

GLUING FOR STABLE FAMILIES OF SURFACES IN MIXED CHARACTERISTIC

QUENTIN POSVA

ABSTRACT. Motivated by the question of properness of the moduli space of stable surfaces in mixed characteristic, we study the gluing problem for stable families of lc surfaces over a mixed characteristic DVR. We prove that the quotient exists, and that the fibers of the quotient are the quotient of the fibers.

CONTENTS

1. Introduction	1
1.1. Acknowledgements	3
2. Preliminaries	3
2.1. Conventions and notations	3
2.2. Quotients by finite equivalence relations	3
2.3. Demi-normal varieties and slc pairs	4
3. Gluing for families of surfaces	5
3.1. Existence of the quotient	5
3.2. Descent of the log canonical sheaf	8
4. Fibers of the quotient	8
References	10

1. INTRODUCTION

Semi-log canonical (slc) singularities play a central role in the moduli theory of canonically polarized varieties in characteristic zero: they appear at the boundary of the compact moduli space of stable varieties [Kol21]. It is expected that slc varieties play a similar role in positive and mixed characteristics.

Recent breakthroughs in the Minimal Model Program (MMP) for threefolds in mixed characteristic [BMP⁺21, TY21] have led to advances in the moduli theory of stable surfaces in mixed characteristic. The corresponding moduli stack is known to be a separated Artin stack with finite diagonal and of finite type [BMP⁺21, Theorem I]. Properness is not known yet, except for some specific subspaces [BMP⁺21, Theorem J].

In characteristic zero, properness of the moduli stack of stable varieties is proved through a delicate use of semi-stable reductions and MMP methods: see [Kol21, §2.4] for a presentation of the proof. In addition to semi-stable reduction and MMP, that proof needs a technical gluing statement. Indeed, we consider a pointed curve $(t \in T)$ and a stable family $X^0 \rightarrow T^0 = T \setminus \{t\}$, and we need to complete the family over T , possibly after a finite base-change $T' \rightarrow T$. The variety X^0 is only demi-normal in general, and the MMP might fail for demi-normal varieties [Fuj14, Example 5.4]. Thus we normalize X^0 and try to complete the family $(X^0)^n \rightarrow T^0$. If we succeed, we need to de-normalize the completed family. In characteristic zero, this is achieved through Kollár's gluing theory [Kol13, §5], that gives a dictionary between slc stable varieties and their normalizations.

In [Pos21] I have extended this dictionary to surfaces and threefolds in positive characteristic, and given applications to the properness of the moduli space of stable surfaces in positive characteristic (see also [Pat17] for related results about this moduli space). In this paper, I study the gluing statement necessary for the proof of properness of the moduli stack of stable

surfaces in mixed characteristic along the lines sketched above. The tailor-made statement is our main result:

Theorem 1 (Proposition 3.1.8). *Let R be a DVR of mixed characteristic with maximal ideal πR . Then normalization gives a bijection*

$$\left(\begin{array}{l} \text{Threefold pairs } (X, \Delta) \\ \text{flat and proper over } R \\ \text{such that } (X, \Delta + X_\pi) \text{ is slc} \\ \text{and } K_X + \Delta \text{ is ample} \end{array} \right) \xrightarrow{1:1} \left(\begin{array}{l} \text{Threefold pairs } (\bar{X}, \bar{D} + \bar{\Delta}) \\ \text{flat and proper over } R \\ \text{such that } (\bar{X}, \bar{D} + \bar{\Delta} + \bar{X}_\pi) \text{ is lc} \\ \text{and } K_{\bar{X}} + \bar{D} + \bar{\Delta} \text{ is ample} \\ \text{plus a generically fixed point free} \\ \text{involution } \tau \text{ of } (\bar{D}^n, \text{Diff}_{\bar{D}^n} \bar{\Delta}). \end{array} \right)$$

Let us comment on the condition that $(X, \Delta + X_\pi)$ is slc. Since we think of the family of surfaces $(X, \Delta) \rightarrow \text{Spec } R$ as an element in moduli stack of stable surfaces, it would be natural to ask for each fiber to be slc. In characteristic zero, this is equivalent to $(X, \Delta + X_\pi)$ being log canonical (lc) [Kol21, Theorem 2.4]. The local-to-global direction relies on inversion of adjunction, which is known in dimension three in positive and mixed characteristics provided that the non-zero characteristics are > 5 ([Pat17, Lemma 3.3] and [BMP⁺21, Corollary 10.1]). On the other hand, there is a difficult step in the proof of the global-to-local direction, namely that $(X, \Delta + X_\pi)$ being slc implies that X_π is S_2 : in characteristic zero this follows from a theorem of Alexeev [Kol13, 7.21]. A similar result is not yet proved in positive and mixed characteristic, so the slc condition on $(X, \Delta + X_\pi)$ is *a priori* weaker.

We prove our gluing statement for this weaker stability assumption. We also prove in Corollary 4.0.5 that if $(\bar{X}_\pi, \text{Diff}_{\bar{X}_\pi}(\bar{D} + \bar{\Delta}))$ is slc then $(X_\pi, \text{Diff}_{X_\pi} \Delta)$ is slc. However the converse statement seems more difficult without an Alexeev-type statement (see Remark 4.0.6). Thus we cannot yet formulate an analog of Theorem 1 for the stronger stability assumption.

Let us say a word about the proof of Theorem 1. We construct an inverse map to the normalization process: \bar{X} should be the normalization of an slc quotient $\bar{X}/R(\tau)$. It turns out that the main difficulty is the existence of the quotient, for then it is easy to show that it has an slc structure, see Proposition 3.2.1.

To show that the quotient exists, thanks to a result of Witaszek [Wit20, Theorem 1.4] and to Kollár's theory [Kol13, 5.13], we only need to show that the equivalence relations induced by τ on the special fiber X_π is finite. To achieve this, we relate \bar{D}_π^n to a divisor contained in the round-down of $\text{Diff}_{X_\pi^n}(\bar{D} + \bar{\Delta})$, and show that the relation on \bar{X}_π comes from a log involution on the reduced boundary of the $k(\pi)$ -pair $(\bar{X}_\pi^n, \text{Diff}_{X_\pi^n}(\bar{D} + \bar{\Delta}))$. This is done using adjunction and classification of codimension two lc singularities. Then we are in position to apply the results from [Pos21] and obtain finiteness. We also obtain a conditional finiteness result in case $\dim \bar{X} = 4$: see Proposition 3.1.8 for the precise statement.

In the second part of the paper, we study the fibers of $\bar{X}/R(\tau)$ over $\text{Spec } R$. We are able to prove that these fibers are the quotients of the fibers of $\bar{X} \rightarrow \text{Spec } R$. More precisely:

Theorem 2 (Theorem 4.0.3). *Let $(\bar{X}, \bar{D} + \bar{\Delta})$ be a proper flat pair over $\text{Spec } R$ such that:*

- (a) $(\bar{X}, \bar{D} + \bar{\Delta} + \bar{X}_\pi)$ is lc,
- (b) $K_{\bar{X}} + \bar{D} + \bar{\Delta}$ is ample,
- (c) there is a log involution τ on $(\bar{D}^n, \text{Diff}_{\bar{D}^n} \bar{\Delta})$ over $\text{Spec } R$, and
- (d) there exists a geometric finite quotient $\bar{X} \rightarrow X := \bar{X}/R(\tau)$.

Then the fibers of $X \rightarrow \text{Spec } R$ are the geometric quotients of the fibers of $\bar{X} \rightarrow \text{Spec } R$ by the equivalence relations induced by τ on each of them.

The proof relies on an analysis of the sections of $K_{\bar{X}_\pi} + \text{Diff}_{\bar{X}_\pi}(\bar{D} + \bar{\Delta})$ that descends to the quotient.

We emphasize that all our results are independent of $\dim \bar{X}$ and of the properties of the positive characteristic residue field $k(\pi)$ — except for Proposition 3.1.8, which shows that the quotient indeed exists in several situations.

1.1. Acknowledgements

Financial support was provided by the European Research Council under grant number 804334.

2. PRELIMINARIES

2.1. Conventions and notations

We will work with schemes that are not necessarily defined over a field, but we keep most of the usual terminology.

A *variety* is a connected separated reduced equidimensional excellent Noetherian scheme. Note that a variety in our sense might be reducible. A *curve* (resp. a *surface*, resp. a *threefold*) is a variety of dimension one (resp. two, resp. three).

Let X be a variety and \mathcal{F} a coherent \mathcal{O}_X -module. We let $\mathcal{F}^* := \mathcal{H}om_X(\mathcal{F}, \mathcal{O}_X)$ be the dual of \mathcal{F} and write $\mathcal{F}^{[m]} := (\mathcal{F}^{\otimes m})^{**}$. We say that \mathcal{F} is S_i if $\text{depth}_{\mathcal{O}_{X,x}} \mathcal{F}_x \geq \min\{i, \dim \mathcal{F}_x\}$ for every $x \in X$.

If X is a variety, we define \mathbb{Q} -divisors and \mathbb{Q} -Cartier divisors the usual way. We usually use these notions when X is (demi-)normal.

If R is a ring that is the quotient of a regular one, and if X is a proper scheme over $\text{Spec } R$, then X has a dualizing complex ω_X^\bullet and a dualizing sheaf $\omega_X := h^{-i}(\omega_X^\bullet)$ where $i := \max\{j \mid h^{-j}(\omega_X^\bullet) \neq 0\}$. See [BMP⁺21, §2.1] for more details. If X is a variety and ω_X is invertible in codimension one, it defines a canonical divisor K_X .

A *pair* is a data (X, Δ) where X is a demi-normal variety, Δ a \mathbb{Q} -divisor with coefficients in $[0, 1]$ without component in common with $\text{Sing}(X)$, such that $K_X + \Delta$ is \mathbb{Q} -Cartier. If $f: Y \rightarrow X$ is a proper birational morphism from a normal variety, then there is a \mathbb{Q} -divisor Δ_Y on Y such that

$$K_Y + \Delta_Y = f^*(K_X + \Delta)$$

and Δ_Y is uniquely defined if we assume $f_*K_Y = K_X$, which we always do. Running through all such $f: Y \rightarrow X$, we define the singularity of the pair (X, Δ) (e.g. *lc*, *klt*) the usual way as in [KM98]. See [BMP⁺21, §2.5] for details.

Let R be a discrete valuation ring (DVR) with maximal ideal πR . If X is an R -scheme, we denote by X_π the scheme-theoretic special fiber. We usually assume that X is flat over $\text{Spec } R$: if in addition X is pure-dimensional, then every irreducible component of X_π has dimension $\dim X - 1$ by [GW20, 14.97].

We will use many times the coarse classification of codimension two lc singularities [Kol13]: if (X, Δ) is an lc pair and $x \in X$ a codimension two point, then $[\Delta]$ is either regular or nodal at x . Notice that the proof given in [Kol13] holds in the generality of normal excellent Noetherian schemes.

Let $(X, D + \Delta)$ be a pair where D is a reduced divisor. Then adjunction theory produces canonically a pair $(D^n, \text{Diff}_{D^n} \Delta)$: see [Kol13, §4.1]. In some cases adjunction already produces a different divisor on D , which will be important for us at a few places. See [Kol13, 4.2] for minimalistic assumptions necessary to perform adjunction.

2.2. Quotients by finite equivalence relations

The theory of quotients by finite equivalence relations is developed in [Kol12] and [Kol13, §9]. For convenience, we recall the basic definitions and constructions that we will need.

Let S be a base scheme, and X, R two reduced S -schemes. An S -morphism $\sigma = (\sigma_1, \sigma_2): R \rightarrow X \times_S X$ is a **set theoretic equivalence relation** (over S) if, for every geometric point $\text{Spec } K \rightarrow S$, the induced map

$$\sigma(\text{Spec } K): \text{Hom}_S(\text{Spec } K, R) \rightarrow \text{Hom}_S(\text{Spec } K, X) \times \text{Hom}_S(\text{Spec } K, X)$$

is injective and an equivalence relation of the set $\text{Hom}_S(\text{Spec } K, X)$. We say in addition that $\sigma: R \rightarrow X \times_S X$ is **finite** if both $\sigma_i: R \rightarrow X$ are finite morphisms.

Suppose that $\sigma: R \hookrightarrow X \times_S X$ is a reduced closed subscheme. Then there is a minimal set theoretic equivalence relation generated by R : see [Kol13, 9.3]. Even if both $\sigma_i: R \rightarrow X$ are finite morphisms, the resulting pro-finite relation may not be finite: achieving transitivity can create infinite equivalence classes.

Let $f: X' \rightarrow X$ be a finite S -morphism, and τ an involution of X' . We denote by $R_X(\tau) \rightrightarrows X$ the set theoretic equivalence relation induced by the closed subscheme $(f \times f)(\Delta_{X'/S} \cup \Gamma_\tau)$ of $X \times_S X$, where $\Delta_{X'/S} \subset X' \times_S X'$ is the diagonal and $\Gamma_\tau \subset X' \times_S X'$ is the graph of τ . If there is no risk of confusion, we drop the subscript and simply write $R(\tau)$.

Let $(\sigma_1, \sigma_2): R \rightarrow X \times_S X$ be a finite set theoretic equivalence relation. A **geometric quotient** of this relation is an S -morphism $q: X \rightarrow Y$ such that

- (a) $q \circ \sigma_1 = q \circ \sigma_2$,
- (b) $(Y, q: X \rightarrow Y)$ is initial in the category of algebraic spaces for the property above, and
- (c) q is finite.

The most important result for us is the following theorem of Witaszek [Wit20, Theorem 1.4]. Assume X is separated of finite type over an excellent scheme S (in our applications, S will be the spectrum of a DVR), and $R \rightarrow X \times_S X$ a finite set-theoretic equivalence relation over S . If the geometric quotient $X_{\mathbb{Q}}/R_{\mathbb{Q}}$ exists, then the geometric quotient X/R also exists (as an algebraic space) and is of finite type over S . If X is projective over S , then X has the Chevalley–Kleiman property, so X/R is a scheme by [Kol12, Corollary 48].

Lemma 2.2.1. *Let X be a reduced Noetherian pure-dimensional scheme, $D \subset X$ a divisor, $\tau: D^n \cong D^n$ an involution such that there exists a finite quotient $q: X \rightarrow Y := X/R(\tau)$. Let L be a line bundle on Y . Then L is equal to the subsheaf of q_*L_X formed by the sections whose pullbacks to D^n are τ -invariant.*

Proof. Let $\sigma_1, \sigma_2: R(\tau) \rightrightarrows X$ be the finite equivalence induced by τ . By [Kol13, 9.10] we have

$$\mathcal{O}_Y = \ker \left[q_*\mathcal{O}_X \xrightarrow{\sigma_1^* - \sigma_2^*} (q \circ \sigma_i)_*\mathcal{O}_{R(\tau)} \right].$$

We claim that the kernel of the map $\sigma_1^* - \sigma_2^*: q_*\mathcal{O}_X \rightarrow (q \circ \sigma_i)_*\mathcal{O}_{R(\tau)}$ is equal to the kernel of $\iota^* - (\iota \circ \tau)^*: q_*\mathcal{O}_X \rightarrow (q \circ \iota)_*\mathcal{O}_{D^n}$, where $\iota: D^n \rightarrow X$ is the structural morphism. Indeed, the inclusion \supseteq is clear. For the converse one, observe that we can find a dense open subset $U \subset D$ which is τ -invariant and contained in the normal locus of D . So for $s \in q_*\mathcal{O}_X$ the condition $(\sigma_1^* - \sigma_2^*)(s) = 0$ implies that $\iota^*(s) - (\iota \circ \tau)^*(s)$ is zero over the dense open subset $U \subset D^n$. Since D^n is S_1 , we obtain that $\iota^*(s) - (\iota \circ \tau)^*(s) = 0$ as desired.

Thus we have an exact sequence

$$0 \rightarrow \mathcal{O}_Y \rightarrow q_*\mathcal{O}_X \xrightarrow{\iota^* - (\iota \circ \tau)^*} (q \circ \iota)_*\mathcal{O}_{D^n}$$

and tensoring it with L we obtain the result. \square

2.3. Demi-normal varieties and slc pairs

We give the definitions and the basic properties of demi-normal varieties and of slc pairs over a DVR.

Definition 2.3.1. A one-dimensional Noetherian local ring (R, \mathfrak{m}) is called a **node** if there exists a ring isomorphism $R \cong S/(f)$, where (S, \mathfrak{n}) is a regular two-dimensional local ring and $f \in \mathfrak{n}^2$ is an element that is not a square in $\mathfrak{n}^2/\mathfrak{n}^3$.

A locally Noetherian reduced scheme (or ring) is called **nodal** if its codimension one local rings are regular or nodal. It is called **demi-normal** if it is S_2 and nodal.

Let X be a reduced scheme with normalization $\pi: \bar{X} \rightarrow X$. The conductor ideal of the normalization is defined as

$$\mathfrak{I} := \mathcal{H}om_X(\pi_*\mathcal{O}_{\bar{X}}, \mathcal{O}_X).$$

It is an ideal in both \mathcal{O}_X and $\mathcal{O}_{\bar{X}}$. We let

$$D := \text{Spec}_X \mathcal{O}_X/\mathfrak{I}, \quad \bar{D} := \text{Spec}_{\bar{X}} \mathcal{O}_{\bar{X}}/\mathfrak{I},$$

and call them the *conductor subschemes*.

Lemma 2.3.2. *Notations as above. Assume that X is the spectrum of a local ring that is essentially of finite type over a DVR. Then:*

- (a) D and \bar{D} are reduced of pure codimension 1.
- (b) If $\eta \in D$ is a generic point such that $\text{char } k(\eta) \neq 2$, the morphism $\bar{D} \rightarrow D$ is étale of degree 2 in a neighborhood of η .
- (c) X has a dualizing sheaf which is invertible in codimension one. In particular, it has a well-defined canonical divisor K_X .
- (d) Let Δ be a \mathbb{Q} -divisor on X with no component supported on D , and such that $K_X + \Delta$ is \mathbb{Q} -Cartier. If $\bar{\Delta}$ denotes the divisorial part of $\pi^{-1}(\Delta)$, then there is a canonical isomorphism

$$\pi^* \omega_X^{[m]}(m\Delta) \cong \omega_{\bar{X}}^{[m]}(m\bar{D} + m\bar{\Delta})$$

for m divisible enough.

Proof. We note that X has a dualizing sheaf, since it is the spectrum of a quotient of a regular ring. The rest is proved as in [Kol13, 5.2, 5.7] (notice that the adjunction property is proved in [Kol13, 4.2, 5.7] in the generality we need). \square

Lemma 2.3.3. *Notations as above. Assume that $\text{char } k(\eta) \neq 2$ for every generic point η of D . Then:*

- (a) the induced morphism of normalizations $\bar{D}^n \rightarrow D^n$ is the geometric quotient by a Galois involution τ ;
- (b) if $K_X + \Delta$ is \mathbb{Q} -Cartier, then τ is a log involution of $(\bar{D}^n, \text{Diff}_{\bar{D}^n} \bar{\Delta})$.

Proof. See [Pos21, 2.2.5]. \square

Definition 2.3.4. We say that $m(X, \Delta)$ is an **semi-log canonical (slc) pair** over a DVR if: X is demi-normal of finite type over a DVR, Δ is a \mathbb{Q} -divisor with no components along D , $K_X + \Delta$ is \mathbb{Q} -Cartier, and the normalization $(\bar{X}, \bar{D} + \bar{\Delta})$ is an lc pair.

3. GLUING FOR FAMILIES OF SURFACES

The goal of this section is the proof of [Theorem 1](#). We consider the following situation (where, compared to [Theorem 1](#), we modify slightly our notations):

Notation 3.0.1. Let R be a DVR of mixed characteristic $(0, p)$, let $X \rightarrow \text{Spec } R$ be a flat proper morphism with connected fibers from an equidimensional normal scheme X of any dimension, let D be an effective reduced Weil divisor on X and Δ an effective \mathbb{Q} -Weil divisor, and assume that $(X, D + \Delta + X_\pi)$ is log canonical. In particular, X_π is reduced and has no component contained in the support of $D + \Delta$.

We denote by $\bar{D} \rightarrow D$ the normalization of D , by \bar{D}_π the special fiber of $\bar{D} \rightarrow \text{Spec } R$ and by \bar{D}_π^n its normalization.

We also assume that there exists an involution τ of the log pair $(\bar{D}, \text{Diff}_{\bar{D}} \Delta)$.

3.1. Existence of the quotient

Lemma 3.1.1. *Let $Z \rightarrow \text{Spec } R$ be a morphism and τ an involution of Z . Then τ is a morphism over $\text{Spec } R$ and restricts to an involution of $\text{red}(Z_\pi)$.*

Proof. Let I be the ideal of $\text{red}(Z_\pi)$. Looking at the exact sequence $0 \rightarrow I \rightarrow \mathcal{O}_Z \rightarrow \mathcal{O}_{\text{red}(Z_\pi)} \rightarrow 0$, we see that τ descends to an involution of $\mathcal{O}_{\text{red}(Z_\pi)}$ if and only if $\tau(I) \subseteq I$.

Since $I = \sqrt{\pi \mathcal{O}_Z}$, we only need to show that $\tau(\pi) \in I$. Since τ is an automorphism, for every $z \in Z$ we have an equality of residue fields $k(z) = k(\tau(z))$, and $\pi \in \mathfrak{m}_{Z,z}$ if and only if $\tau(\pi) \in \mathfrak{m}_{Z,\tau(z)}$. Since $\pi \in \mathfrak{m}_{Z,z}$ if and only if $k(z) = p > 0$, we deduce that $\tau(\pi) \in \mathfrak{m}_{Z,z}$ for all z above the closed point of $\text{Spec } R$. Since $I = \bigcap_{z \in Z_\pi} \mathfrak{m}_{Z,z}$ we obtain that $\tau(\pi) \in I$. \square

Lemma 3.1.2. *Let $(X, D + \Delta) \rightarrow \text{Spec } R$ be as in [Notation 3.0.1](#). If E is a divisor over X whose center $c_X(E)$ belongs to the special fiber X_π , then $a(E; X, D + \Delta) \geq 0$.*

Proof. The proof is the same as [Kol21, 2.14]. Let E be a divisor over X , appearing on a proper birational model $\pi: Y \rightarrow X$. Write $b_E := \text{coeff}_E \pi^* X_\pi$. Since $\pi^* X_\pi$ is Cartier and effective, b_E is a non-negative integer. If $c_X(E) \subset X_\pi$ then b_E is actually a positive integer. Then:

$$-1 \leq a(E; X, D + \Delta + X_\pi) = a(E; X, D + \Delta) - b_E$$

so $a(E; X, D + \Delta) \geq 0$. \square

Lemma 3.1.3. *In the situation of Notation 3.0.1, every irreducible component of D dominates $\text{Spec } R$, and the irreducible components of $D \cap X_\pi$ have dimension $\dim D - 1$.*

Proof. Since X is flat over $\text{Spec } R$, every component of X_π is a divisor by [GW20, 14.97]. Since $(X, D + \Delta + X_\pi)$ is lc, no component of the boundary has coefficient > 1 . Thus D does not contain any component of X_π , and so every component of D dominates $\text{Spec } R$ and so $D \rightarrow \text{Spec } R$ is flat [Har77, III.9.7]. Applying [GW20, 14.97] again yields the result. \square

Lemma 3.1.4. *The special fiber \bar{D}_π of $\bar{D} \rightarrow \text{Spec } R$ is reduced.*

Proof. Since \bar{D} is normal and \bar{D}_π is an hypersurface, \bar{D}_π is S_1 . Thus we only need to show that \bar{D}_π is generically reduced.

The generic points of \bar{D}_π dominates the generic points of the intersection $D \cap X_\pi$, and by Lemma 3.1.3 these points have codimension two in X . Since $(X, D + \Delta + X_\pi)$ is lc, the classification of codimension two lc singularities [Kol13, 2.32] shows that around the generic points of $D \cap X_\pi$, the divisors D and X_π are regular and meet transversally. Hence $\bar{D} \rightarrow D$ is an isomorphism around those generic points, with \bar{D}_π isomorphic to the regular $D \cap X_\pi$. This shows that \bar{D}_π is generically reduced. \square

Lemma 3.1.5. *The involution τ on $(\bar{D}, \text{Diff}_{\bar{D}} \Delta)$ induces an involution σ of the lc pair (\bar{D}_π^n, Γ) , where Γ is defined by the adjunction formula $(K_{\bar{D}} + \text{Diff}_{\bar{D}}(\Delta) + \bar{D}_\pi)|_{\bar{D}_\pi^n} = K_{\bar{D}_\pi^n} + \Gamma$.*

Moreover, the equivalence relation $R_{X_\pi}(\sigma) \rightrightarrows X_\pi$ is equal to the restriction of the equivalence relation $R_X(\tau) \rightrightarrows X$ to X_π .

Proof. By Lemma 3.1.1 and Lemma 3.1.4 the involution τ preserves the Cartier divisor \bar{D}_π^n and descends to an involution σ' on \bar{D}_π . By the universal property of normalization, we obtain an involution σ of the normalization \bar{D}_π^n that makes the diagram

$$(3.1.5.a) \quad \begin{array}{ccc} \bar{D}_\pi^n & \xrightarrow{\sigma} & \bar{D}_\pi^n \\ \downarrow & & \downarrow \\ \bar{D}_\pi & \xrightarrow{\sigma'} & \bar{D}_\pi \end{array}$$

commutative.

Since the \mathbb{Q} -Cartier divisors $K_{\bar{D}} + \text{Diff}_{\bar{D}} \Delta$ and \bar{D}_π are τ -invariant, so is their sum. Hence the pullback of $K_{\bar{D}} + \text{Diff}_{\bar{D}} \Delta + \bar{D}_\pi^n$ to \bar{D}_π^n is σ -invariant.

The pair (\bar{D}_π^n, Γ) is lc by adjunction.

It is clear from the construction that $R(\sigma') \rightrightarrows X_\pi$ is equal to the restriction of $R_X(\tau)$ to X_π , so we only need to compare $R_{X_\pi}(\sigma)$ and $R_{X_\pi}(\sigma')$. Since (3.1.5.a) is commutative, we see that the two equivalence relations are the same. \square

Lemma 3.1.6. *The special fiber X_π is reduced, regular at the generic points of $D \cap X_\pi$ and at worst nodal at other codimension one points. Moreover if $X_\pi^n \rightarrow X_\pi$ is the normalization morphism then:*

- (a) *it is an isomorphism over the generic points of $D \cap X_\pi$,*
- (b) *the pair $(X_\pi^n, \text{Diff}_{X_\pi^n}(D + \Delta))$ is lc, and*
- (c) *the strict transform of $D \cap X_\pi$ is contained in $[\text{Diff}_{X_\pi^n}(D + \Delta)]$.*

Proof. The hypersurface X_π of X is S_1 . It is regular at the generic points of $D \cap X_\pi$ and at worst nodal at other codimension one points by [Kol13, 2.32]. To decide whether the strict transform of $D \cap X_\pi$ appears in $[\text{Diff}_{X_\pi^n}(D + \Delta)]$ is a local question around the generic points of $D \cap X_\pi$. So the first and third points hold. The pair $(X_\pi^n, \text{Diff}_{X_\pi^n}(D + \Delta))$ is lc by adjunction. \square

Lemma 3.1.7. *Denote by $E \subset [\text{Diff}_{X_\pi^n}(D + \Delta)]$ the strict transform of $D \cap X_\pi$, and by (E^n, Θ) the lc pair obtained from $(X_\pi^n, \text{Diff}_{X_\pi^n}(D + \Delta))$ by adjunction. Then there exists a log isomorphism $f: (E^n, \Theta) \cong (\bar{D}_\pi^n, \Gamma)$ such that the diagram*

$$(3.1.7.b) \quad \begin{array}{ccc} E^n & \xrightarrow{f} & \bar{D}_\pi^n \\ & \searrow & \swarrow \\ & X & \end{array}$$

commutes.

Proof. As observed in Lemma 3.1.4 and Lemma 3.1.6, the two morphisms

$$E \rightarrow D \cap X_\pi \leftarrow \bar{D}_\pi$$

are birational and finite. This induces finite birational morphisms

$$E^n \rightarrow (D \cap X_\pi)^n \leftarrow \bar{D}_\pi^n$$

of normal schemes, so these must be isomorphisms. Thus we obtain an isomorphism $f: E^n \cong \bar{D}_\pi^n$ that commutes with the morphisms to X . Since the divisors Θ and Γ are defined by adjunction, we see that f is a log isomorphism. \square

Proposition 3.1.8. *In the situation of Notation 3.0.1, assume that X is proper over $\text{Spec } R$ and that $K_X + D + \Delta$ is ample over $\text{Spec } R$. Assume also that*

- (a) $\dim X = 3$, or
- (b) $\dim X = 4$, and X_π is S_2 in a neighborhood of $D \cap X_\pi$ and the residue field of R is perfect of characteristic > 5 .

Then the quotient $X/R(\tau)$ exists as a demi-normal scheme that is flat and proper over $\text{Spec } R$.

Proof. As explained in Section 2.2, the geometric quotient $X/R(\tau)$ exists as a scheme as soon as $R(\tau)$ is finite and the geometric quotient $X_{\mathbb{Q}}/R(\tau)_{\mathbb{Q}}$ exists.

Lemma 3.1.1 implies that the involution τ on $(\bar{D}, \text{Diff}_{\bar{D}}(\Delta))$ is an involution over $\text{Spec } R$. So it induces an involution $\tau_{\mathbb{Q}}$ on the generic fiber $(\bar{D}_{\mathbb{Q}}, \text{Diff}_{\bar{D}_{\mathbb{Q}}}(\Delta_{\mathbb{Q}}))$ and $R(\tau_{\mathbb{Q}}) = R(\tau)|_{X_{\mathbb{Q}}}$. Since $K_X + D + \Delta$ is ample over $\text{Spec } R$, we see that $(X_{\mathbb{Q}}, D_{\mathbb{Q}} + \Delta_{\mathbb{Q}})$ is a projective lc pair over a field of characteristic zero with ample log canonical divisor and a log involution $\tau_{\mathbb{Q}}$ on $(\bar{D}_{\mathbb{Q}}, \text{Diff}_{\bar{D}_{\mathbb{Q}}}(\Delta_{\mathbb{Q}}))$. By [Kol13, 5.13] the quotient $X_{\mathbb{Q}}/R(\tau)_{\mathbb{Q}}$ exists.

We still have to show that $R(\tau)$ is finite. Since it respects the fibration to $\text{Spec } R$, we only need to show that $R(\tau)|_{X_\pi}$ is finite. By Lemma 3.1.5 it is equivalent to show that $R_{X_\pi}(\sigma) \rightrightarrows X_\pi$ is finite. By Lemma 3.1.7 and the commutativity of (3.1.7.b), we may transport σ to an involution of (E^n, Θ) . We are now in situation to apply [Pos21, 4.1.1, 4.1.3, 5.3.2] which shows that $R_{X_\pi^n}(\sigma)$ is finite.

To go from X_π^n to X_π , observe that the equivalence generated by σ on X_π^n , respectively on X_π , is trivial away from the support of E , respectively away from the support of $D \cap X_\pi$. Now X_π is R_1 in a neighborhood of $D \cap X_\pi$ by Lemma 3.1.7. If it is also S_2 then $X_\pi^n \rightarrow X_\pi$ is an isomorphism in a neighborhood of E , and we deduce that $R_{X_\pi}(\sigma)$ is finite.

If $\dim X = 3$, then we do not need the fact that X_π is S_2 . In this case X_π is a reduced surface that is regular in codimension one and such that $\omega_{X_\pi}^{[m]}(m(D + \Delta)|_{X_\pi})$ is invertible for m sufficiently divisible (this follows from adjunction along $X_\pi \subset X$), hence we can perform adjunction along $E^n \rightarrow X_\pi$ even if X_π is not normal (see [Kol13, 4.2]). The crucial point is that E^n is a curve, so the points of E^n where $E^n \rightarrow D \cap X_\pi$ is not an isomorphism, are contained in $\text{Supp } \Theta$ by [Kol13, 4.5.1]. Thus the the proof of [Pos21, 4.1.1] is also valid in this situation: the finite closed subset $\Sigma := \text{Supp } \Theta$ is σ -invariant and $E^n \setminus \Sigma \rightarrow E \setminus n(\Sigma)$ is an isomorphism. Thus we obtain finiteness.

Flatness of $X/R(\tau)$ over $\text{Spec } R$ follows from [Har77, III.9.7].

It remains to show that $X/R(\tau)$ is demi-normal and proper over $\text{Spec } R$. This is shown as in [Pos21, 3.4.1] (which is formulated for schemes over a field, but the proof also works over a DVR). \square

Remark 3.1.9. More generally, the proof of [Proposition 3.1.8](#) applies to $(X, D + \Delta, \tau)$ as soon as we have a gluing theorems for stable lc varieties of dimension $\dim X - 1$ above the residue field $k(\pi)$ of R , and that the special fiber X_π is S_2 .

3.2. Descent of the log canonical sheaf

We show that in the situation of [Proposition 3.1.8](#), the log canonical \mathbb{Q} -Cartier divisor descends to the quotient. We can actually show it in any dimension, as soon as the quotient exists:

Proposition 3.2.1. *Let $(Y, \Delta_Y) \rightarrow \text{Spec } R$ be a demi-normal flat $\text{Spec } R$ -scheme, with induced normalization $(X, \Delta + D, \tau)$. If $(X, D + \Delta + X_\pi)$ is lc and $\text{Diff}_{D^n}(\Delta)$ is τ -invariant, then $K_Y + \Delta_Y$ is \mathbb{Q} -Cartier.*

Proof. First base-change over the generic point of $\text{Spec } R$. Then [[Kol13](#), 5.38] shows that $K_Y + \Delta_Y$ is \mathbb{Q} -Cartier on the generic fiber.

So the closed locus where $K_Y + \Delta_Y$ is not \mathbb{Q} -Cartier, if not empty, is contained in the special fiber Y_π . By [Lemma 3.1.2](#) and the fact that $(X, D + \Delta) \rightarrow (Y, \Delta_Y)$ is crepant, we see that no lc center of (Y, Δ_Y) is contained in Y_π . Thus we may apply the first part of the proof of [[Kol13](#), 5.38], which is valid in our setting, and conclude that $K_Y + \Delta_Y$ is also \mathbb{Q} -Cartier along the special fiber. \square

Proof of Theorem 1. If (X, Δ) is as in the left-hand side of [Theorem 1](#), we claim its normalization is a triplet $(\bar{X}, \bar{D} + \bar{\Delta}, \tau)$ as on the right-hand side. We indeed have a generically fixed point free involution τ : this follows from [Lemma 2.3.3](#) since by [Lemma 3.1.3](#) the generic points of \bar{D} have characteristic zero residue fields. The other properties clearly hold.

Let $(\bar{X}, \bar{D} + \bar{\Delta}, \tau)$ be as in the right-hand side of [Theorem 1](#). By [Proposition 3.1.8](#) the quotient $X := \bar{X}/R(\tau)$ exists, it is a demi-normal scheme flat and proper over $\text{Spec } R$. By [Proposition 3.2.1](#) we obtain that (X, Δ) is slc.

This defines a map in the opposite direction as the normalization. They are inverse to each other by [[Kol13](#), 5.3] and the fact that \bar{X} is the normalization of X . \square

4. FIBERS OF THE QUOTIENT

In the previous section, we have shown that the quotient exists and has a \mathbb{Q} -Cartier log canonical divisor in several situations. In this section, we study the fibers of the quotient whenever it exists.

Notation 4.0.1. Let $(X, D + \Delta, \tau) \rightarrow \text{Spec } R$ be as in [Notation 3.0.1](#). We assume that the geometric quotient $q: X \rightarrow Y := X/R(\tau)$ exists (so as in [Proposition 3.1.8](#) Y is necessarily demi-normal).

If $\Delta_Y := q_*\Delta$ then $K_Y + \Delta_Y$ is \mathbb{Q} -Cartier by [Proposition 3.2.1](#).

As in [Lemma 3.1.7](#), we let E be the strict transform of $D \cap X_\pi$ in X_π^n , (E^n, Θ) the log pair obtained by adjunction and σ the log involution of (E^n, Θ) .

Lemma 4.0.2. *Assumptions as in [Notation 4.0.1](#). Then the geometric quotient $p: X_\pi \rightarrow Z := X_\pi/R(\tau)|_{X_\pi}$ exists, and $K_{X_\pi^n} + \text{Diff}_{X_\pi^n}(D + \Delta)$ descends to a \mathbb{Q} -Cartier divisor M on Z . Moreover, for a sufficiently divisible integer $e > 0$, the sections of $\mathcal{O}_Z(eM)$ are the sections of $p_*\mathcal{O}_{X_\pi^n}(e(K_X + D + \Delta)|_{X_\pi^n})$ whose restriction to E^n are σ -invariant.*

Proof. As we have seen in the proof of [Lemma 3.1.5](#), the equivalence relation $R(\tau)|_{X_\pi} \rightrightarrows X_\pi$ is equal to the equivalence relation $R(\sigma) \rightrightarrows X_\pi$ and this relation is finite since the quotient exists. So $Z := X_\pi/R(\sigma)$ exists as a $k(\pi)$ -scheme [[Kol12](#), Theorem 6, Corollary 48]. By the universal

property of the geometric quotient, there is a morphism $Z \rightarrow Y_\pi$ that make the diagram

$$(4.0.2.c) \quad \begin{array}{ccc} X_\pi & \hookrightarrow & X \\ \downarrow p & & \downarrow q \\ Z & & Y \\ \downarrow & & \downarrow \\ Y_\pi & \hookrightarrow & Y \end{array}$$

commutative. Since $K_X + D + \Delta$ descends to Y by [Proposition 3.2.1](#), by commutativity we obtain a \mathbb{Q} -Cartier divisor M on Z which pullbacks to $K_{X_\pi^n} + \text{Diff}_{X_\pi^n}(D + \Delta)$ on X_π^n . Therefore the last claim of the statement follows from [Lemma 2.2.1](#). \square

Theorem 4.0.3. *Assumptions as in [Notation 4.0.1](#). We assume in addition that X is projective over $\text{Spec } R$ with relatively ample \mathbb{Q} -Cartier divisor $K_X + D + \Delta$. Then the fibers of Y over $\text{Spec } R$, are the quotients of the fibers of X by τ .*

Proof. The base-change over the generic point of $\text{Spec } R$ is flat and taking the quotient commutes with flat base-change [[Kol13](#), 9.11]. Thus the statement holds for the generic fiber.

Now we consider the special fibers. The divisor $K_Y + \Delta_Y$ is \mathbb{Q} -Cartier and ample over $\text{Spec } R$ by [Proposition 3.2.1](#). Choose an even integer $m > 0$ such that $L := m(K_Y + \Delta_Y)$ is Cartier. We let L_S be the pullback of L to S , for any scheme S that appears in the diagram (4.0.2.c). We may increase m so that L (resp. L_X) is very ample on the fibers of Y (resp. X) over $\text{Spec } R$. Then we have $Y = \text{Proj}_R \bigoplus_{r \geq 0} H^0(Y, rL)$ and furthermore by cohomology and base-change

$$Y_\pi = \text{Proj}_{k(\pi)} \bigoplus_{r \geq 0} H^0(Y, rL) \otimes_R k(\pi) = \text{Proj}_{k(\pi)} \bigoplus_{r \geq 0} H^0(Y_\pi, rL_{Y_\pi}).$$

Since L_Z is also ample we have

$$Z = \text{Proj}_{k(\pi)} \bigoplus_{r \geq 0} H^0(Z, rL_Z).$$

We claim that the section rings of L_{Y_π} and L_Z are equal. By [[Pos21](#), 3.2.5] the global sections of rL_Y are those global sections of rL_X whose restriction to \bar{D} are τ -invariant. By [Lemma 4.0.2](#) the global sections of rL_Z are those global sections of rL_{X_π} whose restriction to E^n are σ -invariant. It is clear that a section of rL_X that is τ -invariant on \bar{D} becomes σ -invariant on $E^n = \bar{D}_\pi^n$. Hence the map

$$H^0(Y, rL) \xrightarrow{\otimes k(\pi)} H^0(Y_\pi, rL_{Y_\pi}) \hookrightarrow H^0(X_\pi, rL_{X_\pi})$$

factors through a map $\alpha_r: H^0(Y, rL) \rightarrow H^0(Z, rL_Z)$ that makes the diagram

$$\begin{array}{ccc} H^0(Y, rL) & \hookrightarrow & H^0(X, rL_X) \\ \downarrow \alpha_r & & \downarrow \beta_r \\ H^0(Z, rL_Z) & \hookrightarrow & H^0(X_\pi, rL_{X_\pi}) \end{array}$$

commutative, where β_r is the restriction map. To conclude, we need to show that α_r is surjective. Pick a section $\bar{s} \in H^0(Z, rL_Z)$. By cohomology and base-change the map β_r is surjective, so we can lift \bar{s} to a section $s \in H^0(X, rL_X)$. We are done if $s|_{\bar{D}}$ is τ -invariant, for then $s \in H^0(Y, rL)$ and so $s \in \text{im}(\alpha_r)$. Let us prove that $t := s|_{\bar{D}}$ is indeed τ -invariant. The restriction $t|_{\bar{D}_\pi^n}$ is σ -invariant by hypothesis, so $t|_{\bar{D}_\pi}$ is σ' -invariant (recall the diagram (3.1.5.a)). Since σ' is the restriction of τ to \bar{D}_π , we see that the support of $t - \tau^*t$ is disjoint from \bar{D}_π . Now $\text{Supp}(t - \tau^*t)$ is closed and its image through the proper morphism $\bar{D} \rightarrow \text{Spec } R$ misses the closed point of $\text{Spec } R$. Hence $\text{Supp}(t - \tau^*t)$ is empty. Therefore $t = \tau^*t$, as claimed. \square

Corollary 4.0.4. *Assumptions as in [Theorem 4.0.3](#). Then Y_π is S_1 , at worst nodal in codimension one, and S_2 at $y = p(x)$ if X_π is S_2 at x .*

Proof. The S_1 property is clear since Y is S_2 .

We have established that $p: X_\pi \rightarrow Y_\pi$ is the quotient by $R(\sigma) \rightrightarrows X_\pi$. By [Lemma 3.1.6](#) we know X_π is regular at the generic points of $D \cap X_\pi$ and at worst nodal at other codimension one points. Since p is an isomorphism away from $D \cap X_\pi$, we only need to consider points of $p(D \cap X_\pi)$. If $y \in p(D \cap X_\pi)$ is a codimension one point of Y_π (resp. the image of an S_2 -point $x \in X_\pi$), we can find an open neighbourhood U of y such that $p^{-1}U$ is normal. Finite quotients of normal varieties (over $k(\pi)$) by involution on normalized divisors are demi-normal [[Pos21](#), 3.4.1]: this proves that Y_π has at worst a nodal singularity at y (resp. that Y_π is S_2 at y). \square

Corollary 4.0.5. *Assumptions as in [Theorem 4.0.3](#). If $(X_\pi, \text{Diff}_{X_\pi}(\Delta + D))$ is slc then $(Y_\pi, \text{Diff}_{Y_\pi} \Delta_Y)$ is also slc.*

Proof. By [Corollary 4.0.4](#) if X_π is S_2 then Y_π is demi-normal. Thus we can perform adjunction along $Y_\pi \subset Y$ (see [[Kol13](#), 4.2]) and we obtain a pair $(Y_\pi, \text{Diff}_{Y_\pi} \Delta_Y)$. From the diagram ([4.0.2.c](#)) we see that its normalization is $(X_\pi^n, \text{Diff}_{X_\pi^n}(\Delta + D))$, which is lc. \square

Remark 4.0.6. The converse direction of [Corollary 4.0.5](#) seems more difficult. The question is whether Y_π being S_2 implies that X_π is also S_2 . The only question is along $D \cap X_\pi$, thus we may localize over a point of $p(D \cap X_\pi)$.

Let W be the quotient of X_π^n by $R(\sigma)$. We have a commutative diagram

$$\begin{array}{ccc} X_\pi^n & \xrightarrow{n} & X_\pi \\ \downarrow q & & \downarrow p \\ W & \xrightarrow{m} & Y_\pi \end{array}$$

where m is a finite morphism. Since X_π is S_2 in codimension one by [Lemma 3.1.6](#), there is a big open subset $U \subset Y_\pi$ such that n_U , and therefore m_U , are isomorphisms. Hence $\mathcal{O}_{Y_\pi} \subset m_*\mathcal{O}_W$ is an equality over U . Since \mathcal{O}_{Y_π} is S_2 and $m_*\mathcal{O}_W$ is torsion-free, we obtain by [[Sta](#), 0AV9] that $\mathcal{O}_{Y_\pi} = m_*\mathcal{O}_W$. In other words $W = Y_\pi$ in a neighbourhood of $p(D \cap X_\pi)$.

However, this equality alone does not imply that X_π is S_2 in a neighbourhood of $D \cap X_\pi$. Here is a counterexample. Consider $S^n = \mathbb{A}_{x,y}^2$ over a field k , the two lines $L_0 = (y = 0)$, $L_1 = (y = 1)$ and the two points $s_0 = (0, 0) \in L_0$, $s_1 = (0, 1) \in L_1$. Let S be the surface obtained from S^n by gluing s_0 and s_1 together, and T be the surface obtained from S^n by gluing L_0 and L_1 along the involution $L_0 \ni (x, 0) \longleftrightarrow (x, 1) \in L_1$. Then T is demi-normal, it is also a quotient of S and we have a commutative diagram

$$\begin{array}{ccc} S^n & \xrightarrow{\quad} & S \\ & \searrow & \swarrow \\ & T & \end{array}$$

However S is R_1 but not S_2 .

REFERENCES

- [BMP⁺21] Bhargav Bhatt, Linqun Ma, Zsolt Patakfalvi, Karl Schwede, Kevin Tucker, Joe Waldron, and Jakub Witaszek. Globally $+$ -regular varieties and the minimal model program for threefolds in mixed characteristic. *ArXiv e-print, arXiv:2012.15801v2*, 2021.
- [Fuj14] Osamu Fujino. Fundamental theorems for semi log canonical pairs. *Algebr. Geom.*, 1(2):194–228, 2014.
- [GW20] Ulrich Görtz and Torsten Wedhorn. *Algebraic geometry I. Schemes—with examples and exercises*. Springer Studium Mathematik—Master. Springer Spektrum, Wiesbaden, [2020] ©2020. Second edition [of 2675155].
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.

- [Kol12] János Kollár. Quotients by finite equivalence relations. In *Current developments in algebraic geometry*, volume 59 of *Math. Sci. Res. Inst. Publ.*, pages 227–256. Cambridge Univ. Press, Cambridge, 2012. With an appendix by Claudiu Raicu.
- [Kol13] János Kollár. *Singularities of the Minimal Model Program*, volume 200 of *Cambridge Tracts in Mathematics*. 2013.
- [Kol21] János Kollár. Families of varieties of general type. Available at <https://web.math.princeton.edu/~kollar/FromMyHomePage/modbook.pdf>, 2021. With the collaboration of K. Altmann and S. Kovács.
- [Pat17] Zsolt Patakfalvi. On the projectivity of the moduli space of stable surfaces in char > 5 . *ArXiv e-print*, *arXiv:1710.03818v3*, 2017.
- [Pos21] Quentin Posva. Gluing theory for slc surfaces and threefolds in positive characteristic. *ArXiv e-print*, *arXiv:2110.09387*, 2021.
- [Sta] The Stacks Project. <https://stacks.math.columbia.edu>.
- [TY21] Teppei Takamatsu and Shou Yoshikawa. Minimal model program for semi-stable threefolds in mixed characteristic. *ArXiv e-print*, *arXiv:2012.07324v2*, 2021.
- [Wit20] Jakub Witaszek. Keel’s base point free theorem and quotients in mixed characteristic. *ArXiv e-print*, *arXiv:2002.11915v1*, 2020.

ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE, SB MATH CAG, MA C3 595 (BÂTIMENT MA), STATION 8, CH-1015 LAUSANNE, SWITZERLAND

Email address: `quentin.posva@epfl.ch`