

# Relativistic formulation of curl force, relativistic Kapitza equation and trapping

Partha Guha\*

Department of Mathematics  
Khalifa University of Science and Technology  
Main Campus, P.O. Box -127788, Abu Dhabi  
United Arab Emirates

Sudip Garai†

Department of Physics  
Diamond Harbour Women's University  
D. H. Road, Sarisha, West-Bengal 743368, India

October 20, 2021

## Abstract

In this present communication the relativistic formulation of the curl forces with saddle potentials has been performed. In particular, we formulated the relativistic version of the Kapitza equation. The dynamics and trapping phenomena of this equation have been studied both theoretically and numerically. The numerical results show interesting characteristics of the charged particles associated with the particle trapping and escaping in the relativistic domain. In addition, the relativistic generalization of the Kapitza equation associated with the monkey saddle has also been discussed.

**PACS numbers:** 05.45.-a, 45.20.-d, 45.20.Dd, 45.50.Jf

**2020 Mathematics Subject Classification:** 01A75, 34A05, 70J25, 70H14

**Keywords:** Curl forces; Kapitza equation; Relativistic Lagrangian; Higher-order saddle potentials; Trapping and escaping

---

\*E-mail: partha.guha@ku.ac.ae

†E-mail: sudip.dhwu@gmail.com

# 1 Introduction

A Newtonian dynamics associated with a non-zero curl force has been a topic of interest to many of the researchers round the globe in the recent past. These forces are known as curl forces which has been introduced to the research community by Berry and Shukla[1, 2]. These forces, in general, are velocity independent in order to satisfy a non-zero curl *viz.* force  $F$  depends only on position  $\mathbf{r}$  and is independent of velocity  $\mathbf{v}$ . The interesting fact is that these curl forces can not be derivable from a scalar potential. The direct applications about the understanding of the system characteristics associated with these curl forces span to many compulsive fields like optics, laser physics, noncentral forces, anisotropic Kepler problem etc.[3, 4, 5, 6, 7, 8, 9, 10, 11, 12]. The generalization of these curl forces have been made with the introduction of dissipation factor present in the system and gyroscopic terms. The application to this directs us to ion trapping, two level atomic state, gain-loss mechanisms etc.

The curl force preserves the volume in phase space domain *viz.*  $(\mathbf{r}, \mathbf{v})$  without any attractors and this can be visualized as follows. Let us take the dynamical equation for a unit mass as  $\ddot{\mathbf{r}} = F(\mathbf{r})$  with a non zero curl i.e.  $\nabla \times F(\mathbf{r}) \neq 0$ . Henceforth one can observe that

$$\nabla_{\mathbf{r}}\mathbf{v} + \nabla_{\mathbf{v}}\dot{\mathbf{v}} = \nabla_{\mathbf{r}}\mathbf{v} + \nabla_{\mathbf{v}}F(\mathbf{r}) = 0, \quad \mathbf{v} = \dot{\mathbf{r}}.$$

Although, in general, the dynamics associated with the curl forces are non-dissipative. Therefore in presence of dissipation the phase space volume is not preserved under the curl force, i.e.  $\nabla \times F(\mathbf{r}) \neq 0$  and  $\nabla_{\mathbf{r}}\mathbf{v} + \nabla_{\mathbf{v}}F(\mathbf{r}) \neq 0$ .

The curl forces are generated by the dynamical motion associated with saddle potentials. A particle remains stable to its equilibrium if the surface rotates around the vertical axis sufficiently fast. To visualize let us take a saddle force field as  $(F_x = x, F_y = -y)$ . We now covert this to a time-dependent one by implementing a rotation to each of the vector counterclockwise with angular velocity  $2\omega$ . This describes the dynamical motion of a particle under this force field with unit mass. From the Earnshaw's theorem we know that a static potential can not trap a charged particle because any static potential, which satisfies the Laplace's law, lacks a potential minimal. Thus the trapping of a charged particle can be achieved by rotating the saddle potential around a suitable axis with a suitable angular frequency. This was the famous idea of Wolfgang Paul[14]. The basic idea of Paul was to stabilize the saddle by means of a 'vibrating' electrostatic field by analogy with so called *Stephenson-Kapitsa*[15, 16] pendulum. On the other hand, in the Penning trap, magnetic fields are used for charged particle's trapping[17, 18, 19]. In recent times, the study regarding the stabilization and other geometric properties of these highly interesting curl forces and the dynamics of nonlinear Hamiltonian curl forces have been reported[13, 20, 21].

The dynamics of the curl forces are related to the theory of Kapitsa-Merkin nonconservative positional forces. The linearized dynamics of a rotating shaft formulated by Kapitsa[22] is given by

$$\ddot{x} + ay + bx = 0, \quad \ddot{y} - ax + by = 0. \quad (1.1)$$

The corresponding characteristics equation reflects that, if we add a non-zero nonconservative curl force ( i.e.  $a \neq 0$ ) to a stable system associated with a stable potential then it becomes unstable. This is associated to the Merkin’s result[23, 24], which states that “the introduction of nonconservative linear forces into a system with a stable potential and with equal frequencies destroys the stability regardless of the form of nonlinear terms”. It is worth mentioning that the positional force, i.e. the terms  $ay$  and  $-ax$  are proportional to  $\omega^2$ , where  $\omega$  is the rotation rate of the shaft. It is also possible to derive Eq.(1.1) associated with a saddle potential via *Euler-Lagrange* method. It is very straightforward to check that the Lagrangian

$$L = \frac{1}{2}(\dot{x}^2 - \dot{y}^2) - \frac{1}{2}a(x^2 - y^2) - bxy, \quad (1.2)$$

yields Eq.(1.1). A symmetric saddle surface can be described by  $\tilde{U} = a(x^2 - y^2)$ , where  $a$  is a geometrical parameter that specifies the curvature of the saddle. It is note worthy to mention that an ion trap potential is a rotating saddle surface on which a ball can be trapped. The pondermotive potential of an ion trap is that of a saddle in  $2D$  but the time evolution of the potential is one that flaps up and down. The potential of the modified rotating system is

$$\mathbb{U}(x, y, t) = \frac{1}{2}(x^2 - y^2) \cos(2\omega t) - xy \sin(2\omega t),$$

where  $\omega$  is the angular drive frequency of the spinning saddle.

The relativistic extension of the nonlinear dynamical equations are not a straightforward job due to the intrinsic extra nonlinearity triggered by the presence of the Lorentz factor[25, 26, 27]. The relativistic harmonic oscillator is a topic of immense interest, although widely discussed, still contains elements of interest. Here the Newtonian kinetic energy is replaced by its special relativistic counterpart. This version of the relativistic harmonic oscillator model has recently been probed experimentally also[28]. Recently, Haas[29] generalized Ermakov systems towards the special relativity domain. In fact relativistic nonlinear dynamics is an open area of research and deserves more attention and intense studies. In this present work we study the relativistic analog of curl force and explore the relativistic Kapitza equation (RKE) and trapping phenomena. One must note that the RKE is the prototypical example of a relativistic curl force.

The paper is organized as follows: In Section 2, we recapitulate, briefly, the construction of the linear curl forces by following Berry and Shukla. We also introduce the Kapitza equation and dynamics related to the higher saddles there. Section 3 is dedicated to the relativistic extension of the curl forces and Kapitza equation. We also describe the trapping phenomena in that section. Lastly, in Section 4, we summarize our findings in the relativistic domain with the comparison to the non-relativistic results along with the possible application towards realistic scenarios.

## 2 Preliminaries: linear curl forces and Hamiltonization

The central fact about curl force theory is that only a small subset of all curl forces are Hamiltonian cases and they find nice applications also. But it is the non-Hamiltonian curl forces that

are really distinctive and exhibit various new features of dynamics.

Let us summarise the classical theory of linear curl forces[1, 2, 9]. We recall from Ref.[1] that the form of the Hamiltonian in the case of curl forces is given by

$$H = \frac{1}{2}\alpha p_x^2 + \beta p_x p_y + \frac{1}{2}\gamma p_y^2 + U(x, y), \quad (2.1)$$

where the potential is defined by

$$U(x, y) = \frac{1}{2}ax^2 + bxy + \frac{1}{2}cy^2. \quad (2.2)$$

The first set of Hamiltonian equations then gives

$$\dot{x} = \alpha p_x + \beta p_y, \quad \dot{y} = \beta p_x + \gamma p_y. \quad (2.3)$$

We can then get the forces from the second set of Hamiltonian equations as

$$a_x = \ddot{x} = \alpha \dot{p}_x + \beta \dot{p}_y = -\alpha \frac{\partial U(x, y)}{\partial x} - \beta \frac{\partial U(x, y)}{\partial y}, \quad (2.4)$$

$$a_y = \ddot{y} = \beta \dot{p}_x + \gamma \dot{p}_y = -\beta \frac{\partial U(x, y)}{\partial x} - \gamma \frac{\partial U(x, y)}{\partial y}. \quad (2.5)$$

The curl is

$$\nabla \times \mathbf{F} = (\alpha - \gamma) \frac{\partial^2 U(x, y)}{\partial x \partial y} \mathbf{i} + \beta \left( \frac{\partial^2 U(x, y)}{\partial x^2} - \frac{\partial^2 U(x, y)}{\partial y^2} \right) \mathbf{j}, \quad (2.6)$$

which yields  $\nabla \times \mathbf{F} = (\alpha - \gamma)b\mathbf{i} + \beta(c - a)\mathbf{j}$ . To ensure that  $\nabla \times F \neq 0$ , it is necessary to take  $\beta = 0$  and  $\alpha = 1 = -\gamma$ . Also the another choice we have is  $\beta \neq 0$  and  $c = -a$ . Henceforth the Hamiltonian turns out to be

$$H = \frac{1}{2}(p_x^2 - p_y^2) + \frac{1}{2}(x^2 - y^2) + bxy, \quad (2.7)$$

which is a Hamiltonian of the Kapitza equation.

## 2.1 Rotating saddle and Kapitza equation related to higher-saddle

Consider the motion of a particle in the rotating saddle potential in the plane

$$U(\mathbf{x}, t) = U_0(R^{-1}\mathbf{x}), \quad U_0 = \frac{1}{2}(x^2 - y^2), \quad \mathbf{x} = (x, y); \quad (2.8)$$

where

$$R = R(\Omega t) = \begin{pmatrix} \cos(\omega t) & \sin(\omega t) \\ -\sin(\omega t) & \cos(\omega t) \end{pmatrix}.$$

The Lagrangian of the rotating saddle potential is given by

$$L_{rotsaddle} = \frac{1}{2}(\dot{x}^2 - \dot{y}^2) - \frac{\Lambda}{2} \left( (x^2 - y^2) \cos(2\omega t) + 2xy \sin(2\omega t) \right), \quad \Lambda = 2 \frac{mgh_0}{r_0^2}, \quad (2.9)$$

and the corresponding equations of motion are

$$\ddot{x} + \Lambda x \cos(2\omega t) + \Lambda y \sin(2\omega t) = 0, \quad \ddot{y} + \Lambda y \cos(2\omega t) - \Lambda x \sin(2\omega t) = 0. \quad (2.10)$$

### 2.1.1 Higher saddle and generalized Kapitza equation

We can generalize the rotating shaft equation formulated by Kapitza using higher-order saddles. The Eq.(1.1) can be manufactured from the Euler-Lagrange equation using simple saddle potential  $g_1(x, y) = x^2 - y^2$  and the rotated version of the same surface,  $g_1^R(x, y) = xy$ . In other words, latter one is obtained from the rotation of the saddle by 90 degree. Let  $z = x + iy$ , then  $x^2 - y^2 = \text{Re}(z^2)$  and  $2xy = \text{Im}(z^2)$ . We follow this guideline to construct equations of rotating shaft associated to monkey saddle

$$g_2(x, y) = x^3 - 3xy^2 = \text{Re}(z^3) \quad (2.11)$$

and its rotated version

$$g_2^R(x, y) = 3x^2y - y^3 = \text{Im}(z^3). \quad (2.12)$$

The generalized rotating shaft pair of equations associated to degree three potential or monkey saddle potential is given by

$$U_3 = \frac{1}{3}(k_1g_2(x, y) + k_2g_2^R(x, y)) = \frac{1}{3}(k_1(x^3 - 3xy^2) + k_2(3x^2y - y^3)).$$

The corresponding equations of motion are given by

$$\ddot{x} + k_1(x^2 - y^2) + 2k_2xy = 0, \quad \ddot{y} + 2k_1xy - k_2(x^2 - y^2) = 0. \quad (2.13)$$

## 3 Relativistic curl force

The relativistic Lagrangian in presence of scalar potential fields is given by

$$L = -mc^2\sqrt{1 - 2\frac{\mathcal{L}}{mc^2}}, \quad \mathcal{L} = \frac{m|\mathbf{v}|^2}{2} - U(x). \quad (3.1)$$

One way to describe the non-relativistic approximation is to apply the limits:  $U(x) \ll mc^2$ ;  $v \ll c$ ,

$$\begin{aligned} L &= -mc^2\sqrt{\left(1 - \frac{|\mathbf{v}|^2}{c^2}\right) + \frac{2U}{mc^2}} = -mc^2\sqrt{1 - \frac{|\mathbf{v}|^2}{c^2}}\sqrt{1 + \frac{2U}{mc^2}\left(1 - \frac{|\mathbf{v}|^2}{c^2}\right)^{-1}} \\ &= -mc^2\gamma^{-1}\sqrt{1 + \frac{2U}{mc^2}\gamma^2}, \quad \gamma = \left(1 - \frac{|\mathbf{v}|^2}{c^2}\right)^{-\frac{1}{2}}. \end{aligned}$$

Expanding the expression within square-root binomially up to second order gives us

$$\tilde{L} = -mc^2\gamma^{-1}\left(1 + \frac{U}{mc^2}\gamma^2\right) = -mc^2\gamma^{-1} - U\gamma. \quad (3.2)$$

Although in *Goldstein's* book the semi-relativistic Lagrangian is given by

$$L_{sr} = mc^2\gamma^{-1} - U. \quad (3.3)$$

If we assume

$$\frac{|\mathbf{v}|}{c} = \alpha \ll 1, \Rightarrow U\gamma \sim U + \frac{U}{2}\alpha^2,$$

for practical purpose; we can drop the second term for small value of  $\alpha$  which immediately yields the semi-relativistic Lagrangian given in *Goldstein*[30].

### 3.1 Relativistic Kapitza equation

The relativistic Lagrangian of the curl force is defined as

$$L = -\frac{c^2}{\Gamma} - U(x, y), \quad \text{with} \quad \Gamma = \left(1 - \frac{\dot{x}^2}{c^2} + \frac{\dot{y}^2}{c^2}\right)^{-\frac{1}{2}}. \quad (3.4)$$

Here the saddle potential is assumed as

$$U(x, y) = \frac{1}{2}k(x^2 - y^2) + bxy. \quad (3.5)$$

**Lemma 3.1** *The Euler-Lagrange equation of the relativistic Lagrangian  $L = -\frac{c^2}{\Gamma} - \frac{1}{2}k(x^2 - y^2) - bxy$  yields*

$$\ddot{x}\left(1 + \frac{\dot{y}^2}{c^2}\right) - \frac{\dot{x}\dot{y}}{c^2}\ddot{y} = -\frac{kx}{\Gamma^3} - \frac{by}{\Gamma^3}, \quad (3.6)$$

$$\ddot{y}\left(1 - \frac{\dot{x}^2}{c^2}\right) + \frac{\dot{x}\dot{y}}{c^2}\ddot{x} = -\frac{ky}{\Gamma^3} + \frac{bx}{\Gamma^3}, \quad (3.7)$$

where  $\Gamma = \left(1 - \frac{\dot{x}^2}{c^2} + \frac{\dot{y}^2}{c^2}\right)^{-\frac{1}{2}}$ .

**Proof** It is straightforward to check that

$$\begin{aligned} \frac{\partial L}{\partial \dot{x}} &= \dot{x}\left(1 - \frac{\dot{x}^2}{c^2} + \frac{\dot{y}^2}{c^2}\right)^{-\frac{1}{2}}, & \frac{d}{dt}\left(\frac{\partial L}{\partial \dot{x}}\right) &= \left[\ddot{x}\left(1 + \frac{\dot{y}^2}{c^2}\right) - \frac{\dot{x}\dot{y}}{c^2}\ddot{y}\right]\left(1 - \frac{\dot{x}^2}{c^2} + \frac{\dot{y}^2}{c^2}\right)^{-\frac{3}{2}}, \\ \frac{\partial L}{\partial \dot{y}} &= -\dot{y}\left(1 - \frac{\dot{x}^2}{c^2} + \frac{\dot{y}^2}{c^2}\right)^{-\frac{1}{2}}, & \frac{d}{dt}\left(\frac{\partial L}{\partial \dot{y}}\right) &= \left[-\ddot{y}\left(1 - \frac{\dot{x}^2}{c^2}\right) - \frac{\dot{x}\dot{y}}{c^2}\ddot{x}\right]\left(1 - \frac{\dot{x}^2}{c^2} + \frac{\dot{y}^2}{c^2}\right)^{-\frac{3}{2}}. \end{aligned}$$

Then using the derivative of  $\frac{\partial L}{\partial \dot{x}}$  and  $\frac{\partial L}{\partial \dot{y}}$  we obtain our result from the Euler-Lagrange equation. We must note here that both the equations (3.6) and (3.7) contain  $\ddot{x}$  and  $\ddot{y}$  terms. The equations of motion are therefore obtained by separating these terms.

**Proposition 3.1** *The Euler-Lagrange equation for the relativistic Kapitza Lagrangian yields the relativistic Kapitza equation*

$$\ddot{x} = \frac{\dot{x}\dot{y}}{\Gamma c^2}(-ky + bx) - \frac{1}{\Gamma \Gamma_x^2}(kx + by), \quad (3.8)$$

$$\ddot{y} = \frac{\dot{x}\dot{y}}{\Gamma c^2}(kx + by) + \frac{1}{\Gamma \Gamma_y^2}(-ky + bx), \quad (3.9)$$

where

$$\Gamma_x = \left(1 - \frac{\dot{x}^2}{c^2}\right)^{-\frac{1}{2}}, \quad \Gamma_y = \left(1 + \frac{\dot{y}^2}{c^2}\right)^{-\frac{1}{2}}. \quad (3.10)$$

**Proof** Multiplying (3.6) by  $(1 - \frac{\dot{x}^2}{c^2})$  we obtain

$$\begin{aligned} \ddot{x}\left(1 + \frac{\dot{y}^2}{c^2}\right)\left(1 - \frac{\dot{x}^2}{c^2}\right) - \frac{\dot{x}\dot{y}}{c^2}\ddot{y}\left(1 - \frac{\dot{x}^2}{c^2}\right) &= -\left(\frac{kx}{\Gamma^3} + \frac{by}{\Gamma^3}\right)\left(1 - \frac{\dot{x}^2}{c^2}\right) \\ \frac{\ddot{x}}{\Gamma^2} &= \frac{\dot{x}\dot{y}^3}{\Gamma}c^2(-ky + bx) - \frac{1}{\Gamma_x^2\Gamma^3}(kx + by). \end{aligned}$$

Similarly, we obtain the second equation via the multiplication of Eq.(3.7) with  $(1 + \frac{\dot{y}^2}{c^2})$ .

**Corollary 3.1** *The equations (3.16) and (3.16) for non-relativistic limit reduce to the Kapitza equation*

$$\ddot{x} + kx + by = 0, \quad \ddot{y} - ky + bx = 0. \quad (3.11)$$

The relativistic Kapitza equation can also be formulated using Hamiltonian method.

**Proposition 3.2** *The (effective) Hamiltonian of the relativistic Kapitza equation is*

$$H = \frac{p_x^2}{2\Gamma} - \frac{p_y^2}{2\Gamma} + \frac{1}{2}k(x^2 - y^2) + bxy, \quad (3.12)$$

and corresponding Hamiltonian equations are given by

$$\dot{x} = \frac{\partial H}{\partial p_x} = \frac{p_x}{\Gamma}, \quad \dot{p}_x = -\frac{\partial H}{\partial x} = -kx - by, \quad (3.13)$$

$$\dot{y} = \frac{\partial H}{\partial p_y} = -\frac{p_y}{\Gamma}, \quad \dot{p}_y = -\frac{\partial H}{\partial y} = ky - bx. \quad (3.14)$$

One can clearly see that  $p_x = \dot{x}\Gamma$ ,  $p_y = -\dot{y}\Gamma$  and the rest follows from lemma 3.1. It can be easily shown that the relativistic Kapitza equation follows from Eqns.(3.13) and (3.14).

It is clear that for the non-relativistic limit the Hamiltonian equation yields the Kapitza equation. One must note that non-relativistic Kapitza equation has two integrals of motion, the Hamiltonian  $H = \frac{p_x^2}{2} - \frac{p_y^2}{2} + \frac{1}{2}k(x^2 - y^2) + bxy$  and the Fradkin tensor  $I = p_x p_y + \frac{1}{2}b(x^2 - y^2) - kxy$ . Unfortunately, for the relativistic case we have just one conserved quantity due to the presence of the Lorentz factor.

### 3.2 Relativistic rotation of saddle

The Lagrangian of the rotating saddle potential is given by

$$L_{\text{rel saddle}} = -\frac{c^2}{\Gamma} - \frac{\Lambda}{2}\left((x^2 - y^2)\cos(2\Gamma\omega t) + 2xy\sin(2\Gamma\omega t)\right), \quad (3.15)$$

where the constant coefficients  $b$  and  $k$  are replaced by the sinusoidal terms, the corresponding equations of motion are

$$\ddot{x} = \frac{\dot{x}\dot{y}}{\Gamma c^2}\Lambda(-\cos(2\Gamma\omega t)y + \sin(2\Gamma\omega t)x) - \frac{1}{\Gamma_x^2}\Lambda(\cos(2\Gamma\omega t)x + \sin(2\Gamma\omega t)y), \quad (3.16)$$




Figure 1: Phase space diagrams (with normalized parameters) via Eq.(3.16) and Eq.(3.17) with  $\Lambda = 0.1$  and  $\omega = 0.5$ . The top two figures are for non-relativistic cases with  $v_x = 0 = v_y$  and the bottom two are for the relativistic cases with  $v_x = 0.5c = v_y$ . The initial conditions are  $x(0) = 0$ ,  $\dot{x}(0) = 0.0$ ,  $y(0) = 0.0$ ,  $\dot{y}(0) = 0.001$ . The normalized parameters are as follows:  $x \rightarrow x\omega_c/c$ ,  $y \rightarrow y\omega_c/c$ ,  $\dot{x} \rightarrow \dot{x}/c$ ,  $\dot{y} \rightarrow \dot{y}/c$ ,  $\ddot{x} \rightarrow \ddot{x}\omega_c c$ ,  $\ddot{y} \rightarrow \ddot{y}\omega_c c$ ,  $\Lambda \rightarrow \Lambda/\omega_c^2$ ,  $\omega \rightarrow \omega/\omega_c$  and  $t \rightarrow t\omega_c$  where  $\omega_c$  can be taken as the cyclotron frequency in the typical microwave range.

$$\ddot{y} = \frac{\dot{x}\dot{y}}{\Gamma_x^2} \Lambda (\cos(2\Gamma\omega t)x + \sin(2\Gamma\omega t)y) + \frac{1}{\Gamma\Gamma_y^2} \Lambda (-\cos(2\Gamma\omega t)y + \sin(2\Gamma\omega t)x), \quad (3.17)$$

where

$$\Gamma_x = (1 - \frac{\dot{x}^2}{c^2})^{-\frac{1}{2}}, \quad \Gamma_y = (1 + \frac{\dot{y}^2}{c^2})^{-\frac{1}{2}}. \quad (3.18)$$

The phase space plots for the rotating saddle have been depicted in figure (1) for Eqs.(3.16) & (3.17). In the non-relativistic scenario, it can be seen from figure (1a) that the trajectory corresponds to limit cycle under one specific saddle point (SP) and spanning over smaller to larger amplitudes. The particle can be trapped inside those periodic boundaries for shorter times but not completely. Figure (1b) corresponds to heteroclinic orbit where it moves from one SP to another over time directs to the escaping of particles. In the relativistic case, both figure (1c) and (1d) correspond heteroclinic orbit and particle escaping.






Figure 2: Phase space diagrams (with normalized parameters) via Eq.(3.19) and Eq.(3.20) with  $\Lambda = 0.1$ ,  $\omega = 0.5$  and  $\alpha_1 = \alpha_2 = 1$ . The top two figures are for non-relativistic cases with  $v_x = 0 = v_y$  and the bottom two are for the relativistic cases with  $v_x = 0.5c = v_y$ . The initial conditions are  $x(0) = 0$ ,  $\dot{x}(0) = 0.0$ ,  $y(0) = 0.0$ ,  $\dot{y}(0) = 0.001$ . The normalization scheme is same as mentioned in figure (1) along with  $\alpha_1 \rightarrow \alpha_1 c/\omega_c$  and  $\alpha_2 \rightarrow \alpha_2 c/\omega_c$ .

In the non-relativistic limit Eqns.(3.16) and (3.18) reduce to

$$\ddot{x} + \Lambda x \cos(2\omega t) - \Lambda y \sin(2\omega t) = 0, \quad \ddot{y} + \Lambda y \cos(2\omega t) - \Lambda x \sin(2\omega t) = 0.$$

The relativistic generalized Kapitza equation corresponding monkey saddle is the extension of the monkey saddle equation,

$$\ddot{x} = \frac{\dot{x}\dot{y}}{\Gamma c^2} (-2k_1xy + k_2(x^2 - y^2)) - \frac{1}{\Gamma\Gamma_x^2} (k_1(x^2 - y^2) + 2k_2xy), \quad (3.19)$$

$$\ddot{y} = \frac{\dot{x}\dot{y}}{\Gamma c^2} (k_1(x^2 - y^2) + 2k_2xy) + \frac{1}{\Gamma\Gamma_y^2} (-2k_1xy + k_2(x^2 - y^2)). \quad (3.20)$$

The phase space plots for the relativistic rotating monkey saddle have been depicted in figure (2) for Eqs.(3.19) & (3.20). Figure (2a) and (2c) show heteroclinic trajectories which remove the possibility of particle trapping for both relativistic and non-relativistic cases. In contrary

to this figure (2b) and (2d) show limit cycle trajectories under one SP for both relativistic and non-relativistic cases. As one can see that there are periodic boundaries which are changing very slowly over time and closely spaced, henceforth particle can be trapper over shorter time scale.

The *relativistic rotating monkey saddle equation* can be obtained by replacing  $k_1 = \Lambda\alpha_1 \cos(\Gamma 2\omega t)$  and  $k_2 = \Lambda\alpha_2 \sin(\Gamma 2\omega t)$ .

### 3.3 Flapping and spinning of relativistic saddle potential and trap

Suppose we allow the potential  $U = g(\frac{1}{2}(x^2 - y^2))$  to vary sinusoidally with time. this becomes

$$\mathcal{U}(x, y, t) = A_{RF} \cos(\omega t) g\left(\frac{1}{2}(x^2 - y^2)\right), \quad (3.21)$$

where  $A_{RF}$  is the amplitude of the AC component of the electric field applied to create the trap, and  $\omega$  is the frequency. This can be viewed as a *nonlinear* generalization of ‘flapping saddle’. Latter occurs in a quadrupole potential in which extreme oscillate between peaks and valleys. Thus Newton’s law yields the following pair of equations

$$\ddot{x} = -A_{RF} \cos(\omega t) x g\left(\frac{1}{2}(x^2 - y^2)\right), \quad \ddot{y} = -A_{RF} y \cos(\omega t) g\left(\frac{1}{2}(x^2 - y^2)\right). \quad (3.22)$$

The linear case coincides with the form of the Mathieu differential equations. Our motivation is to study this new set of equations. In general it is difficult to create flapping saddle on which the ball can move. So we bang on the time-dependent spinning saddle at angular frequency  $\omega$ , which is similar to the flapping potential

$$\hat{\mathcal{U}}(x, y, t) = A(g(x^2 - y^2) \cos(2\omega t) + 2f(xy) \sin(2\omega t)), \quad (3.23)$$


where  $A$  is some constant. We now apply again the relativistic Lagrangian as

$$L_{relflap} = -\frac{c^2}{\Gamma} - \hat{\mathcal{U}}(x, y, t), \quad \text{where } \hat{\mathcal{U}}(x, y, t) = A(g(x^2 - y^2) \cos(2\omega t) + 2f(xy) \sin(2\omega t)), \quad (3.24)$$


to get the relativistic coupled differential equations as

$$\ddot{x} = \frac{\dot{x}\dot{y}}{\Gamma c^2} \Lambda (-yg'(x^2 - y^2) \cos(2\Gamma\omega t) + xf'(xy) \sin(2\Gamma\omega t)) - \frac{1}{\Gamma\Gamma_x^2} (xg'(x^2 - y^2) \cos(2\Gamma\omega t) + yf'(xy) \sin(2\Gamma\omega t)), \quad (3.25)$$


$$\ddot{y} = \frac{\dot{x}\dot{y}}{\Gamma c^2} \Lambda (xg'(x^2 - y^2) \cos(2\Gamma\omega t) + yf'(xy) \sin(2\Gamma\omega t)) + \frac{1}{\Gamma\Gamma_y^2} (-yg'(x^2 - y^2) \cos(2\Gamma\omega t) + xf'(xy) \sin(2\Gamma\omega t)). \quad (3.26)$$




(a) Plot of  $\dot{x}$  vs  $x$ .



(b) Plot of  $\dot{y}$  vs  $y$ .



(c) Plot of  $\dot{x}$  vs  $x$ .



(d) Plot of  $\dot{y}$  vs  $y$ .

Figure 3: Phase space diagrams (with normalized parameters) via Eq.(3.25) and Eq.(3.26) with  $\Lambda = 0.1$ ,  $\omega = 0.5$  and  $\beta_1 = \beta_2 = \beta_3 = \beta_4 = 1$ . The top two figures are for non-relativistic cases with  $v_x = 0 = v_y$  and the bottom two are for the relativistic cases with  $v_x = 0.5c = v_y$ . The initial conditions are  $x(0) = 0$ ,  $\dot{x}(0) = 0.0$ ,  $y(0) = 0.0$ ,  $\dot{y}(0) = 0.001$ . The normalization scheme is same as mentioned in figure (1) along with  $\beta_1 \rightarrow \beta_1 \omega_c^2 / c^2$ ,  $\beta_2 \rightarrow \beta_2 \omega_c^4 / c^4$ ,  $\beta_3 \rightarrow \beta_3 \omega_c^2 / c^2$  and  $\beta_4 \rightarrow \beta_4 \omega_c^4 / c^4$ . It must be noted here that we have chosen  $g(x^2 - y^2) = 1 + \beta_1(x^2 - y^2) + \beta_2(x^2 - y^2)^2$  and  $f(xy) = 1 + \beta_3(xy) + \beta_4(xy)^2$ .

If we take  $\tau = \omega t$  and  $x' = \frac{dx}{d\tau}$  we obtain

$$x'' = \frac{x'y'}{\Gamma c^2} \Lambda (-yg'(x^2 - y^2) \cos(2\Gamma\omega\tau) + xf'(xy) \sin(2\Gamma\omega\tau)) - \frac{1}{\omega^2 \Gamma \Gamma_x^2} (xg'(x^2 - y^2) \cos(2\Gamma\omega\tau) + yf'(xy) \sin(2\Gamma\omega\tau)), \quad (3.27)$$

$$y'' = \frac{x'y'}{\Gamma c^2} \Lambda (xg'(x^2 - y^2) \cos(2\Gamma\omega\tau) + yf'(xy) \sin(2\Gamma\omega\tau)) + \frac{1}{\omega^2 \Gamma \Gamma_y^2} (-yg'(x^2 - y^2) \cos(2\Gamma\omega\tau) + xf'(xy) \sin(2\Gamma\omega\tau)). \quad (3.28)$$

The phase space plots for the relativistic flapping saddle have been depicted in figure (3) for Eqs.(3.25) & (3.26). Like in the previous case of relativistic rotating monkey saddle here also we see that both trapping in shorter time scale and escaping is possible in both relativistic and non-relativistic limits.

## 4 Summary

The relativistic analog of the Kapitza equation has been investigated and associated trapping phenomena of the charged particles have been discussed. We have presented an example of a relativistic curl force and the generalization of the curl force associated with saddles potentials formulated by Berry and Shukla. In case of relativistic, rotating saddle the phase space plots correspond some heteroclinic orbits and particle escaping. With the inclusion of the monkey saddle to the generalized Kapitza equation in relativistic domain the phase plots also show heteroclinic trajectories with a possibility of particle trapping in specific parametric domain in the shorter time scale. In case of relativistic flapping and spinning saddle there is also a possibility of trapping and escaping same as discussed in the previous one. At last we have discussed about the relativistic generalization of the Kapitza equation associated with the monkey saddle. The theoretical outcomes along with the numerical results may direct and or give an idea about the possible mechanism of particle trapping with finite curl forces in the relativistic domain. In the earlier observations[11, 12], we see that there is a possibility of trapping the charged particles under the curl force dynamics with saddle potentials completely, whereas, in this work we have observed that the trapping is not complete in the relativistic domain. As the results in the relativistic domain direct us of incomplete trapping in a shorter time scale therefore it becomes a field of relative interest to further investigate this with the choice of different saddle potentials and to apply the outcomes towards astrophysical scenarios.

## Acknowledgments

PG is immensely grateful to *Professor Sir Michael Berry* for his critical and meticulous reading of this draft and for making invaluable remarks and or comments. PG and SG also convey their sincere thanks to *Professor Haret Rosu* for his suggestions and comments and *Professors Praghya Shukla, Stefan Mancas and Anindya Ghose-Choudhury* for various discussions and correspondences. PG thanks *Khalifa University of Science and Technology* for its continued support towards this research work under the grant number FSU-2021-014 and SG thanks *Diamond Harbour Women's University* for providing the necessary research environment with constant encouragement and support.

## Author contributions

All the authors contributed equally to this work

## Conflicts of interest

The authors declare that they have no conflict of interest.

## Data transparency

Not applicable.

## References

- [1] M V Berry and Shukla P, Classical dynamics with curl forces, and motion driven by time-dependent flux, *J. Phys. A*, 45:305201, 2012.
- [2] M V Berry and Shukla P, Hamiltonian curl forces. *Proc. R. Soc. A*, 471:20150002 (13pp), 2015.
- [3] M V Berry and Shukla P, Physical curl forces: dipole dynamics near optical vortices. *J. Phys. A*, 46:422001, 2013.
- [4] Chaumet P C and Nieto-Vesperinas M, Time-averaged total force on a dipolar sphere in an electromagnetic field, *Opt. Lett.*, 25:1065–1067, 2013.
- [5] Albaladejo S., Marqués M. I., Laroche M., and Sáenz J. J. Scattering forces from the curl of the spin angular momentum. *Phys. Rev. Lett.*, 102:113602, 2009.
- [6] Shimizu Y. and Sasada H, Mechanical force in laser cooling and trapping, *Am. J. Phys.*, 66:960–967, 1998.
- [7] Gutzwiller M. C, The anisotropic kepler problem in two dimensions, *J. Math. Phys.*, 14:139–152, 1973.
- [8] Devaney R. L, Nonregularizability of the anisotropic kepler problem, *J. Diff. Eqns.*, 29:253, 1978.
- [9] Ghose-Choudhury A., Guha P., Paliathanasis A., and Leach P. G. L, Noetherian symmetries of noncentral forces with drag term. *Int. J. Geom. Methods Mod. Phys.*, 14:1750018, 2017.
- [10] Guha P, Generalized Emden-Fowler equations in noncentral curl forces and first integrals, *Acta Mech.*, 231:815–825, 2020.
- [11] S. Garai and P. Guha, Higher-order saddle potentials, nonlinear curl forces, trapping and dynamics, *Nonlinear Dyn* (2021), <https://doi.org/10.1007/s11071-021-06212-w>
- [12] P. Guha and S. Garai, Integrable modulation, curl forces and parametric Kapitza equation with trapping and escaping, (2021), <https://arxiv.org/abs/2104.06319>

- [13] Ghose-Choudhury A. and Guha P., Hamiltonian description of nonlinear curl forces from cofactor systems. *Acta Mech*, 230:2267–2277, 2019.
- [14] Paul W, 1990, Electromagnetic traps for charged and neutral particles, *Reviews of Modern Physics*, Vol. 62, No. 3, 531.
- [15] Stephenson A., Xx. on induced stability, *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, doi: 10.1080/14786440809463763, 15(86):233–236, 1908.
- [16] Kapitsa P. L., Dynamic stability of a pendulum when its point of suspension vibrates., *Soviet Phys. JETP.*, 21:588–592, 1951.
- [17] Byrne J. and Farago P. S., On the production of polarized electrons by spin exchange collisions., *Proc. Phys. Soc.*, 86:801–815, 1965.
- [18] Grff B G, Klempt E, 1967, *Z. Naturforschung* 22a, 1960-1962.
- [19] Sokolov A A, Pavlenko Yu G, 1967, *Optics and Spectroscopy* 22, 1-3.
- [20] Kirillov O, Levi M, 2016, Rotating saddle trap as Foucault’s pendulum, *Am. J. Phys.* 84, 26.
- [21] Kirillov O, Levi M, 2017, Rotating saddle trap: A Coriolis force in an inertial frame, *Nonlinearity* 30, 1109-1119.
- [22] P L Kapitsa, *Stability and transition through the critical speed of fast rotating shafts with friction* . *Zhur. Tekhn. Fiz.* 9 (1939) 124-147.
- [23] D R Merkin, *Gyroscopic systems*, Nauka, Moscow, in Russian (first edition - 1956) 1974.
- [24] D R Merkin, *Introduction to the theory of stability*, 1997 Springer-Verlag, New York.
- [25] Harvey, A.L. *Relativistic harmonic oscillator*, *Phys. Rev. D* 6 (1972), 1474-1475.
- [26] Llibre, J. Makhlof, A. *On the periodic solutions of the relativistic driven harmonic oscillator*, *J. Math. Phys.*61(2020) 012501-012507.
- [27] Li, Z, Liu, J, Lucha, W, Ma, W, Schöberl, F.F, *Relativistic harmonic oscillator*, *J. Math.Phys.*46 (2005) 103514-103525
- [28] Fujiwara, K.M, Geiger, Z.A, Singh, K, Senaratne, R, Rajagopal, S.V, Lipatov, M, Shimasaki, T, Weld, D.M, *Experimental realization of a relativistic harmonic oscillator*, *New J. Phys.* 2018, 20, 063027, 7 pages.
- [29] Haas F, *Relativistic Ermakov-Milne-Pinney Systems and First Integrals*, *Physics* 2021, 3(1), 59-70.
- [30] Goldstein, H, Poole, C, Safko, J. *Classical Mechanics*, 3rd ed. Addison Wesley: San Francisco,USA, 2002; pp. 316.