

A Unified and Refined Convergence Analysis for Non-Convex Decentralized Learning

Sulaiman A. Alghunaim

Kuwait University

sulaiman.alghunaim@ku.edu.kw

Kun Yuan

DAMO Academy, Alibaba Group

kun.yuan@alibaba-inc.com

October 20, 2021

Abstract

We study the consensus decentralized optimization problem where the objective function is the average of n agents private non-convex cost functions; moreover, the agents can only communicate to their neighbors on a given network topology. The stochastic online setting is considered in this paper where each agent can only access a noisy estimate of its gradient. Many decentralized methods can solve such problems including EXTRA, Exact-Diffusion/ D^2 , and gradient-tracking. Unlike the famed DSGD algorithm, these methods have been shown to be robust to the heterogeneity of the local cost functions. However, the established convergence rates for these methods indicate that their sensitivity to the network topology is worse than DSGD. Such theoretical results imply that these methods can perform much worse than DSGD over sparse networks, which, however, contradicts empirical experiments where DSGD is observed to be more sensitive to the network topology.

In this work, we study a general stochastic unified decentralized algorithm (SUDA) that includes the above methods as special cases. We establish the convergence of **SUDA** under both non-convex and the Polyak-Lojasiewicz condition settings. Our results provide improved network topology dependent bounds for these methods (such as Exact-Diffusion/ D^2 and gradient-tracking) compared with existing literature. Moreover, our result shows that these method are less sensitive to the network topology compared to DSGD, which agrees with numerical experiments.

1. Introduction

In a distributed multi-agent optimization problem, the inputs (*e.g.*, functions, variables, data) are spread over multiple computing agents (*e.g.*, nodes, processors) that are connected over some network, and the agents are required to communicate with each other to solve this problem. Distributed optimization have attracted a lot of attention due to the need of

developing efficient methods to solve large-scale optimization problems [1] such as in deep neural networks applications [2]. Decentralized optimization methods are algorithms where the agents seek to find a solution through local interactions (dictated by the network connection) with their neighboring agents. Decentralized methods have several advantages over centralized methods, which require all the agents to communicate with a central coordinator, such as their robustness to failure and privacy. Moreover, decentralized methods have been shown to enjoy lower communication cost compared to centralized methods under certain practical scenarios [3–5].

In this work, we consider a network (graph) of n collaborative agents (nodes) that are interested in solving the following distributed stochastic optimization problem:

$$\underset{x \in \mathbb{R}^d}{\text{minimize}} \quad f(x) = \frac{1}{n} \sum_{i=1}^n f_i(x), \quad f_i(x) \triangleq \mathbb{E}_{\xi_i}[F_i(x; \xi_i)]. \quad (1)$$

In the above formulation, $F_i : \mathbb{R}^d \rightarrow \mathbb{R}$ is a smooth non-convex function privately known by agent i . The notation \mathbb{E}_{ξ_i} is the expected value of the random variable ξ_i (*e.g.*, random data samples) taken with respect to some local distribution. The above formulation is known as the consensus formulation since the agents share a common variable, which they need to agree upon [1]. We consider *decentralized methods* where the agents aim to find a solution of (1) through local interactions (each agent can only send and receive information to its immediate neighbors).

Two important measures of the performance of distributed (or decentralized) methods are the linear speedup and transient time. A decentralized method is said to achieve *linear speedup* if the gradient computational complexity needed to reach certain accuracy reduces linearly with n . The *transient time* of a distributed method is the number of iterations needed to achieve linear speedup. A common method for solving problem (1) is the decentralized/distributed stochastic gradient descent (DSGD) method [6, 7], where each agent employs a local stochastic gradient descent update and a local gossip step (there are several variations based on the order of the gossip step such as diffusion or consensus methods [7–9]). DSGD is simple to implement; moreover, it has been shown to achieve linear speedup asymptotically [3]. This implies that the convergence rate of DSGD asymptotically achieves the same network independent rate as the centralized (also known as parallel) stochastic gradient descent (PSGD) with a central coordinator. While being attractive, DSGD suffers from an error or bias term caused by the heterogeneity of the local cost functions minimizers (*e.g.*, heterogeneous data distributions across the agents) [8, 10]. The existence of such bias term will slow down the convergence of DSGD, and hence enlarge its transient time.

Several bias-correction methods have been proposed to remove the bias of DSGD such as EXTRA [11], Exact-Diffusion (ED) (a.k.a D^2 or NIDS) [12–15], and gradient-tracking (GT) methods [16–19]. While there has been an extensive study on these methods, their convergence properties have not been fully understood as we now explain. Under convex stochastic settings, ED/ D^2 is theoretically shown to improve upon the transient time of DSGD [20, 21], especially under sparse topologies. However, existing non-convex results imply that ED/ D^2 has worse transient time compared to DSGD for sparse networks [15]. Moreover,

the transient time of GT-methods are theoretically worse than DSGD under sparse networks even under convex settings [22]. These existing theoretical results imply that under non-convex settings, bias-correction methods can suffer from worse transient time compared to DSGD. However, empirical results suggest that both ED/D² and GT methods outperform DSGD under sparse topologies (without any acceleration) [15, 23, 24]. This phenomenon is yet to be explained.

In this work, we provide a novel unified convergence analysis of several decentralized bias-correction methods including both ED/D² and GT methods under *non-convex* settings. We establish refined and improved convergence rate bounds over existing results. Moreover, our results show that bias-correction methods such as Exact-Diffusion/D² and GT methods have better network topology dependent bounds compared to DSGD. We also study these methods under the Polyak-Lojasiewicz (PL) condition [25, 26] and provide refined bounds over existing literature. Before we state our main contributions, we will go over the related works.

1.1 Related Works

There exists many works that study decentralized optimization methods under deterministic settings (full knowledge of gradients) – see [9, 11, 19, 27–32] and references therein. For example, the works [31] and [32] provide a unified analysis of several decentralized methods, albeit under deterministic and convex settings. Our focus in this work is on the *stochastic (online)* setting where each agent has access to a random estimate of its gradient at each iteration. For this setting, DSGD is the most widely studied and understood method [3, 4, 33–44]. Under non-convex settings, the transient time of DSGD is on the order of $O(n^3/(1-\lambda)^4)$ [3, 4, 43], where $1-\lambda \in (0, 1)$ denotes the network spectral gap that measures the connectivity of the network topology (*e.g.*, it goes to zero for sparse networks). As a result, improving the convergence rate dependence on the network topology quantity $1-\lambda$ is crucial to enhance the transient time of decentralized methods.

The severe dependence on the network topology in DSGD is caused by the data heterogeneity between different agents [14, 43]. Consequently, the dependence on the network topology can be ameliorated by removing the bias caused by data heterogeneity. For example, the transient time of ED/D² has been shown to have enhanced dependence on network topology compared to DSGD [20, 21]. However, similar results have not been established for non-convex scenarios. In fact, the established transient time of bias-correction methods such as ED/D² and GT in literature are even worse than that of DSGD [15, 24, 45–47]. For instance, the best known transient time for both ED/D² and GT is on the order of $O(n^3/(1-\lambda)^6)$ [15, 24], which is worse than DSGD with transient time $O(n^3/(1-\lambda)^4)$. These counter-intuitive results naturally motivates us to study whether ED/D² and GT can enjoy an enhanced dependence on network topology in the non-convex setting. It is also worth noting that the dependence on network topology established in existing GT references are worse than DSGD even for convex scenarios [22]. This work provides refined and enhanced convergence rates for both ED/D² and GT (as well as other methods such as EXTRA) under the non-convex setting.

In this work, we also study the convergence properties of decentralized methods under the Polyak-Łojasiewicz (PL) condition [25]. The PL condition can hold for non-convex costs, yet it can be used to establish similar convergence rates to strong-convexity rates [26]. For strongly-convex settings, the works [20,21] showed that the transient time of ED/D² is on the order of $O(n/(1-\lambda))$. These are the best available network bounds for decentralized methods so far for strongly-convex settings. It is still unclear, whether bias-correction methods can achieve similar bounds to DSGD under the PL condition. For example, the work [24] shows that under the PL condition, GT methods have transient time on the order $O(n/(1-\lambda)^3)$.

We remark that this work only considers decentralized methods with a *single* gossip round per iteration. It has been shown that combining GT methods with *multiple* gossip rounds can further improve the dependence on network topology [23,48], and this technique can also be incorporated into our studied algorithm and its analysis. However, it is worth noting that the utilization of multiple gossip rounds in decentralized stochastic methods might suffer from several limitations. First, it requires the knowledge of the quantity λ to decide the number of gossip rounds per iteration, which, however, might not be available in practice. Second, the multiple gossip rounds update can take even more time than a global average operation. For example, the experiments provided in [5, Table 17] indicate that, under a certain practical scenario, one gossip step requires half or third the communication overhead of a centralized RING-ALLREDUCE operation [49], which conducts global averaging. This implies that decentralized methods with as much as two or three gossip rounds per iteration can be more costly than global averaging. Third, the theoretical improvements brought by multiple gossip rounds rely heavily on gradient accumulation. Such gradient accumulation can easily cause large batch-size which are empirically and theoretically found to be harmful for generalization performance on unseen dataset [50,51].

1.2 Main Contributions

Our main contributions are formally listed below.

- We unify the analysis of several well-known decentralized methods under *non-convex* and *stochastic* settings. In particular, we study the convergence properties of a general primal-dual algorithmic framework, called stochastic unified decentralized algorithm (**SUDA**), which includes several existing methods such as EXTRA, ED/D², and GT methods as special cases.
- We provide a novel analysis technique for these type of methods. In particular, we employ several novel transformations to **SUDA** that are key to establish our refined convergence rates bounds. Our analysis provides improved network dependent bounds for the special cases of **SUDA** such as ED/D² and GT methods compared to existing best known results. In addition, the established transient time of ED/D² and ATC-GT have improved network topology dependence compared to DSGD – see Table 1.
- We also study the convergence properties of **SUDA** under the PL condition. When specifying **SUDA** to ED/D², we achieve network dependent bound matching the best

Table 1: Comparison with existing *non-convex* convergence rates highlighting the network quantities. Here, $\varsigma_0^2 = \frac{1}{n} \sum_{i=1}^n \|\nabla f_i(0) - \nabla f(0)\|^2$ and ς^2 satisfies $\frac{1}{n} \sum_{i=1}^n \|\nabla f_i(x) - \nabla f(x)\|^2 \leq \varsigma^2$ for all $x \in \mathbb{R}^d$ for DSGD. The quantity $\lambda = \rho(W - \frac{1}{n}\mathbf{1}\mathbf{1}^\top)$ is the mixing rate of the network where W is the network combination matrix. Compared with GT methods our result assumes that W is symmetric and positive-semidefinite.

METHOD	WORK	CONVERGENCE RATE	TRANSIENT TIME
DSGD	[43]	$O\left(\frac{1}{\sqrt{nK}} + \frac{\lambda^{2/3}}{(1-\lambda)^{1/3}K^{2/3}} + \frac{\lambda^{2/3}\varsigma^{2/3}}{(1-\lambda)^{2/3}K^{2/3}}\right)$	$O\left(\frac{n^3}{(1-\lambda)^4}\right)$
ED/D ²	[15]	$O\left(\frac{1}{\sqrt{nK}} + \frac{n\lambda^2}{(1-\lambda)^3K} + \frac{n\varsigma_0^2}{(1-\lambda)^2K^2}\right)$	$O\left(\frac{n^3}{(1-\lambda)^6}\right)$
	This work	$O\left(\frac{1}{\sqrt{nK}} + \frac{n\lambda^2}{(1-\lambda)K} + \frac{n\lambda^2\varsigma_0^2}{(1-\lambda)^2K^2}\right)$	$O\left(\frac{n^3}{(1-\lambda)^2}\right)$
ATC-GT	[24]	$O\left(\frac{1}{\sqrt{nK}} + \frac{n\lambda^2}{(1-\lambda)^3K} + \frac{\lambda^4 \sum_{i=1}^n \ \nabla f_i(0)\ ^2}{(1-\lambda)^3K^2}\right)$	$O\left(\frac{n^3}{(1-\lambda)^6}\right)$
	This work	$O\left(\frac{1}{\sqrt{nK}} + \frac{n\lambda^4}{(1-\lambda)K} + \frac{n\lambda^4}{(1-\lambda)^4K^2} + \frac{n\lambda^4\varsigma_0^2}{(1-\lambda)^3K^2}\right)$	$O\left(\max\left\{\frac{n^3}{(1-\lambda)^2}, \frac{n}{(1-\lambda)^{8/3}}\right\}\right)$

known bounds established under strongly-convex settings. When specifying **SUDA** to GT methods, we achieve an improved network dependent bound compared to current results even under strong-convexity. Table 2 compares the transient times network dependent bounds under the PL or strongly-convex setting.

Notation. Vectors and scalars are denoted by lowercase letters. Matrices are denoted using upper case letters. We use $\text{col}\{a_1, \dots, a_n\}$ (or $\text{col}\{a_i\}_{i=1}^n$) to denote the vector that stacks the vectors (or scalars) a_i on top of each other. We use $\text{diag}\{d_1, \dots, d_n\}$ (or $\text{diag}\{d_i\}_{i=1}^n$) to denote a diagonal matrix with diagonal elements d_i . We also use $\text{blkdiag}\{D_1, \dots, D_n\}$ (or $\text{blkdiag}\{D_i\}_{i=1}^n$) to denote a block diagonal matrix with diagonal blocks D_i . The vector of all ones with size n is denoted by $\mathbf{1}_n$ (or $\mathbf{1}$ and size is known from context). The inner product of two vectors a and b is denoted by $\langle a, b \rangle$. The Kronecker product operation is denoted by \otimes . For a square matrix A , we let $\rho(A)$ denote the spectral radius of A , which is the largest absolute value of its eigenvalues. Upright bold symbols (*e.g.*, \mathbf{x} , \mathbf{f} , \mathbf{W}) are used to denote augmented network quantities.

2. General Algorithm Description

In this section, we describe the deterministic form of the studied algorithm and list several specific instances of interest to us.

Table 2: Comparison with existing network dependent transient times under both strongly-convex and PL condition settings. Here, the quantity $\lambda = \rho(W - \frac{1}{n}\mathbf{1}\mathbf{1}^\top)$ is the mixing rate of the network where W is the network combination matrix. Compared with GT methods our result assumes that W is symmetric and positive-semidefinite.

METHOD	WORK	ASSUMPTION	TRANSIENT TIME
DSGD	[43]	Strongly-convex	$O\left(\frac{n}{(1-\lambda)^2}\right)$
ED/D ²	[20, 21]	Strongly-convex	$O\left(\frac{n}{1-\lambda}\right)$
	This work	PL condition	$O\left(\frac{n}{1-\lambda}\right)$
GT	[22]	Strongly-convex	$O\left(\frac{n}{(1-\lambda)^3}\right)$
	[24]	PL condition	$O\left(\frac{n}{(1-\lambda)^3}\right)$
	This work	PL condition	$O\left(\max\left\{\frac{n}{1-\lambda}, \frac{n^{1/3}}{(1-\lambda)^{4/3}}\right\}\right)$

2.1 General Algorithm

To describe the algorithm, we introduce the network quantities:

$$\mathbf{x} \triangleq \text{col}\{x_1, \dots, x_n\} \in \mathbb{R}^{dn}, \quad (2a)$$

$$\mathbf{f}(\mathbf{x}) \triangleq \sum_{i=1}^n f_i(x_i). \quad (2b)$$

We also introduce the matrix $\mathbf{B} \in \mathbb{R}^{dn \times dn}$ that satisfies

$$\mathbf{B}\mathbf{x} = \mathbf{0} \iff x_1 = x_2 = \dots = x_n. \quad (3)$$

Using the previous definitions, the general algorithmic framework can be described as follows. Set an arbitrary initial estimate $\mathbf{x}^0 \in \mathbb{R}^{dn}$ and set $\mathbf{y}^0 = \mathbf{0}$. Repeat for $k = 0, 1, \dots$

$$\mathbf{x}^{k+1} = \mathbf{A}(\mathbf{C}\mathbf{x}^k - \alpha \nabla \mathbf{f}(\mathbf{x}^k)) - \mathbf{B}\mathbf{y}^k, \quad (4a)$$

$$\mathbf{y}^{k+1} = \mathbf{y}^k + \mathbf{B}\mathbf{x}^{k+1}. \quad (4b)$$

Here, $\alpha > 0$ is the step size (learning rate), and the matrices $\mathbf{A} \in \mathbb{R}^{dn \times dn}$ and $\mathbf{C} \in \mathbb{R}^{dn \times dn}$ are doubly stochastic matrices that are chosen according to the network combination matrix introduced next.

2.2 Network Combination matrix

We let $W = [w_{ij}] \in \mathbb{R}^{n \times n}$ denote the network combination (weighting) matrix assumed to be symmetric. Here, the (i, j) th entry $w_{ij} \geq 0$ is used by agent i to scale information received

from agent j . We consider a decentralized setup where $w_{ij} = 0$ if $j \notin \mathcal{N}_i$ where \mathcal{N}_i is the neighborhood of agent i . If we introduce the augmented combination matrix:

$$\mathbf{W} = W \otimes I_d \in \mathbb{R}^{dn \times dn}.$$

The matrices $\mathbf{A}, \mathbf{B}, \mathbf{C}$ can be chosen as a function of \mathbf{W} to recover several existing decentralized methods. Note that if $\mathbf{u} = \text{col}\{u_i\}_{i=1}^n$ where $u_i \in \mathbb{R}^d$ is local to agent i , then, the i -th block of $\mathbf{W}\mathbf{u} = \text{col}\{\sum_{j \in \mathcal{N}_i} w_{ij} u_j\}_{i=1}^n$ can be computed by agent i through local interactions with its neighbors.

2.3 Relation to Existing Decentralized Methods

We next list the relation of algorithm (4) to several important well-known decentralized algorithms. The details are provided in Appendix A.

- *Exact-Diffusion/D² and EXTRA.* If we select $\mathbf{A} = \mathbf{W}$, $\mathbf{B} = (\mathbf{I} - \mathbf{W})^{1/2}$, and $\mathbf{C} = \mathbf{I}$. Then, algorithm (4) becomes equivalent to ED/D² [12, 15]:

$$x_i^{k+1} = \sum_{j \in \mathcal{N}_i} w_{ij} \left(2x_j^k - x_j^{k-1} - \alpha (\nabla f_j(x_j^k) - \nabla f_j(x_j^{k-1})) \right). \quad (5)$$

If we instead select $\mathbf{A} = \mathbf{I}$, $\mathbf{B} = (\mathbf{I} - \mathbf{W})^{1/2}$, and $\mathbf{C} = \mathbf{W}$, then algorithm (4) is equivalent to EXTRA [11]:

$$x_i^{k+1} = \sum_{j \in \mathcal{N}_i} w_{ij} (2x_j^k - x_j^{k-1}) - \alpha (\nabla f(x_i^k) - \nabla f(x_i^{k-1})). \quad (6)$$

- *Gradient-Tracking (GT) methods.* Consider the adapt-then-combine gradient-tracking (ATC-GT) method [16]:

$$x_i^{k+1} = \sum_{j \in \mathcal{N}_i} w_{ij} (x_j^k - \alpha g_j^k) \quad (7a)$$

$$g_i^{k+1} = \sum_{j \in \mathcal{N}_i} w_{ij} (g_j^k + \nabla f_j(x_j^{k+1}) - \nabla f_j(x_j^k)). \quad (7b)$$

The above method is equivalent to (4) with $\mathbf{A} = \mathbf{W}^2$, $\mathbf{B} = \mathbf{I} - \mathbf{W}$, and $\mathbf{C} = \mathbf{I}$. We can also recover other gradient-tracking variants. For example, if we select $\mathbf{A} = \mathbf{I}$, $\mathbf{B} = \mathbf{I} - \mathbf{W}$, and $\mathbf{C} = \mathbf{W}^2$, then (49) becomes equivalent to the non-ATC-GT method [18]:

$$x_i^{k+1} = \sum_{j \in \mathcal{N}_i} w_{ij} x_j^k - \alpha g_i^k \quad (8a)$$

$$g_i^{k+1} = \sum_{j \in \mathcal{N}_i} w_{ij} g_j^k + \nabla f_i(x_i^{k+1}) - \nabla f_i(x_i^k). \quad (8b)$$

Notice that in (8) the communication (gossip) step only involves the terms x_j^k and g_j^k in the update of each vector. This is in contrast to the ATC structure (7) where the communication (gossip) step involves all terms. We remark that the semi-ATC-GT variations where only the update of x_i^k or g_i^k uses the ATC structure [17] are also equivalent to (4) when $\mathbf{A} = \mathbf{W}$, $\mathbf{B} = \mathbf{I} - \mathbf{W}$, $\mathbf{C} = \mathbf{W}$.

- *UDA*. Consider the unified decentralized algorithm proposed in [31]:

$$\mathbf{z}^{k+1} = \mathbf{C}\mathbf{x}^k - \alpha\nabla\mathbf{f}(\mathbf{x}^k) - \mathbf{B}\mathbf{y}^k \quad (9a)$$

$$\mathbf{y}^{k+1} = \mathbf{y}^k + \mathbf{B}\mathbf{z}^{k+1} \quad (9b)$$

$$\mathbf{x}^{k+1} = \mathbf{A}\mathbf{z}^{k+1}. \quad (9c)$$

In the above description, the matrix \mathbf{C} is equivalent to $\mathbf{I} - \mathbf{C}$ in [31]. The above method is equivalent to (4) if $\mathbf{A}\mathbf{B}^2 = \mathbf{B}^2\mathbf{A}$, which holds under the previous special instances.

3. Stochastic UDA and Assumptions

In this section, we describe the stochastic version of algorithm (4) and list the assumptions used to analyze it.

As stated in problem (1), we consider stochastic settings where each agent may only have access to a stochastic gradient $\nabla F_i(x_i, \xi_i^k)$ at each iteration k instead of the true gradient. This scenario arises in online stochastic settings, where the data are not known in advance; hence, we do not have access to the actual gradient. Moreover, even if the all the data is available, the true gradient might be expensive to compute for large datasets and can be replaced by a gradient at one sample or a mini-batch.

Replacing the actual gradient by its stochastic approximation in (4), we get the **Stochastic Unified Decentralized Algorithm (SUDA)**:

$$\mathbf{x}^{k+1} = \mathbf{A}(\mathbf{C}\mathbf{x}^k - \alpha\nabla\mathbf{F}(\mathbf{x}^k, \boldsymbol{\xi}^k)) - \mathbf{B}\mathbf{y}^k, \quad (10a)$$

$$\mathbf{y}^{k+1} = \mathbf{y}^k + \mathbf{B}\mathbf{x}^{k+1}, \quad (10b)$$

where

$$\nabla\mathbf{F}(\mathbf{x}, \boldsymbol{\xi}^k) \triangleq \text{col}\{\nabla F_1(x_1, \xi_1^k), \dots, \nabla F_n(x_n, \xi_n^k)\}.$$

We next list the Assumptions used in our analyses. Our first assumption is about the network combination matrix given next.

Assumption 1 (Combination matrix). *The combination matrix \mathbf{W} is assumed to be doubly stochastic, symmetric, and primitive. Moreover, we assume that the matrices \mathbf{A} , \mathbf{B}^2 , \mathbf{C} are chosen as a polynomial function of \mathbf{W} :*

$$\mathbf{A} = \sum_{l=0}^p a_l \mathbf{W}^l, \quad \mathbf{B}^2 = \sum_{l=0}^p b_l \mathbf{W}^l, \quad \mathbf{C} = \sum_{l=0}^p c_l \mathbf{W}^l, \quad (11)$$

where $p \geq 1$. The constants $\{a_l, b_l, c_l\}_{l=0}^p$ are chosen such that \mathbf{A} and \mathbf{C} are doubly stochastic and the matrix \mathbf{B} satisfies equation (3). \square

Under Assumption 1, the combination matrix W has a single eigenvalues at one, denoted by $\lambda_1 = 1$. Moreover, all other eigenvalues, denoted by $\{\lambda_i\}_{i=2}^n$, are strictly less than one in magnitude [36], and the mixing rate of the network is:

$$\lambda \triangleq \rho\left(W - \frac{1}{n}\mathbf{1}\mathbf{1}^\top\right) = \max_{i \in \{2, \dots, n\}} |\lambda_i| < 1. \quad (12)$$

Note that the assumptions on $\mathbf{A}, \mathbf{B}^2, \mathbf{C}$ are mild and hold for all the special cases described before.

We now introduce the main assumption on the objective function.

Assumption 2 (Objective function). *Each function $f_i : \mathbb{R}^d \rightarrow \mathbb{R}$ is L -smooth:*

$$\|\nabla f_i(z) - \nabla f_i(y)\| \leq L\|z - y\|, \quad \forall z, y \in \mathbb{R}^d, \quad (13)$$

for some $L > 0$. We also assume that the aggregate function $f(x) = \frac{1}{n} \sum_{i=1}^n f_i(x)$ is bounded below, i.e., $f(x) \geq f^* > -\infty \forall x \in \mathbb{R}^d$ where f^* denote the optimal value of f . \square

The above assumption is standard to establish convergence under non-convex settings. Note that we do not impose the strong assumption of bounded gradient dissimilarity, which is required to establish convergence of DSGD – see [3, 4, 43].

We next list our assumption on the stochastic gradient. To do that, we define the filtration generated by the random process (10):

$$\mathcal{F}^k \triangleq \{\mathbf{x}^0, \mathbf{x}^2, \dots, \mathbf{x}^k\}. \quad (14)$$

The filtration \mathcal{F}^k can be interpreted as the collection of all information available on the past iterates up to time k .

Assumption 3 (Gradient noise). *For all $\{i\}_{i=1}^n$ and $k = 0, 1, \dots$, we assume that the following holds*

$$\mathbb{E} [\nabla F_i(x_i^k; \xi_i^k) - \nabla f_i(x_i^k) \mid \mathcal{F}^k] = 0, \quad (15a)$$

$$\mathbb{E} [\|\nabla F_i(x_i^k; \xi_i^k) - \nabla f_i(x_i^k)\|^2 \mid \mathcal{F}^k] \leq \sigma^2, \quad (15b)$$

for some $\sigma^2 \geq 0$. We also assume that conditioned on \mathcal{F}^k , the random data $\{\xi_i^t\}$ are independent of each other for all $\{i\}_{i=1}^n$ and $\{t\}_{t \leq k}$. \square

The previous assumptions will be used to analyze **SUDA** (10) for general non-convex costs. In the sequel, we will also study **SUDA** under the following additional assumption.

Assumption 4 (PL condition). *The aggregate function $f(x) = \frac{1}{n} \sum_{i=1}^n f_i(x)$ satisfies the PL inequality:*

$$2\mu(f(x) - f^*) \leq \|\nabla f(x)\|^2, \quad \forall x \in \mathbb{R}^d, \quad (16)$$

for some $\mu > 0$ where f^* denote the optimal value of f . \square

The above condition implies that every stationary point is a global minimizer, which is weaker than many assumptions used to establish linear convergence without strong-convexity [26]. Note that the PL condition is also referred to as the gradient dominated condition [52].

4. Fundamental Transformations

The updates of **SUDA** (10), while useful for implementation purposes, are not helpful for analysis purposes. In this section, we will transform **SUDA** (10) into an equivalent recursion that is fundamental to arrive at our results.

4.1 Transformation I

Using the change of variable

$$\mathbf{z}^k \triangleq \mathbf{y}^k - \mathbf{B}\mathbf{x}^k \quad (17)$$

in (10a)–(10b), we can describe (10) by the following equivalent non-incremental form:

$$\mathbf{x}^{k+1} = (\mathbf{A}\mathbf{C} - \mathbf{B}^2)\mathbf{x}^k - \alpha\mathbf{A}(\nabla\mathbf{f}(\mathbf{x}^k) + \mathbf{w}^k) - \mathbf{B}\mathbf{z}^k, \quad (18a)$$

$$\mathbf{z}^{k+1} = \mathbf{z}^k + \mathbf{B}\mathbf{x}^k, \quad (18b)$$

where \mathbf{w}^k is the gradient noise defined as:

$$\mathbf{w}^k \triangleq \nabla\mathbf{F}(\mathbf{x}^k, \boldsymbol{\xi}^k) - \nabla\mathbf{f}(\mathbf{x}^k). \quad (19)$$

If we introduce the quantities

$$\bar{x}^k \triangleq \frac{1}{n}(\mathbf{1}_n^\top \otimes I_d)\mathbf{x}^k = \frac{1}{n} \sum_{i=1}^n x_i^k, \quad (20a)$$

$$\bar{\mathbf{x}}^k \triangleq \mathbf{1}_n \otimes \bar{x}^k, \quad (20b)$$

$$\mathbf{s}^k \triangleq \mathbf{B}\mathbf{z}^k + \alpha\mathbf{A}\nabla\mathbf{f}(\bar{\mathbf{x}}^k), \quad (20c)$$

then recursion (18) can be rewritten as:

$$\mathbf{x}^{k+1} = (\mathbf{A}\mathbf{C} - \mathbf{B}^2)\mathbf{x}^k - \alpha\mathbf{A}(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\bar{\mathbf{x}}^k) + \mathbf{w}^k) - \mathbf{s}^k, \quad (21a)$$

$$\mathbf{s}^{k+1} = \mathbf{s}^k + \mathbf{B}^2\mathbf{x}^k + \alpha\mathbf{A}(\nabla\mathbf{f}(\bar{\mathbf{x}}^{k+1}) - \nabla\mathbf{f}(\bar{\mathbf{x}}^k)). \quad (21b)$$

Remark 1 (Motivation behind \mathbf{s}^k). Let (\mathbf{x}, \mathbf{z}) be a fixed point of the deterministic form of (18) (gradient noise $\mathbf{w}^k = 0$) where $\mathbf{z} \in \mathbb{R}^{dn}$ and $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^{dn}$. Then, it holds that:

$$\mathbf{0} = \alpha\mathbf{A}\nabla\mathbf{f}(\mathbf{x}) + \mathbf{B}\mathbf{z},$$

$$\mathbf{0} = \mathbf{B}\mathbf{x} \iff x_1 = \dots = x_n = x.$$

Now suppose that the variable \mathbf{s}^k defined in (20c) converges to zero for some consensual $\bar{\mathbf{x}} = \mathbf{1} \otimes x$. It follows that

$$\mathbf{0} = \alpha\mathbf{A}\nabla\mathbf{f}(\bar{\mathbf{x}}) + \mathbf{B}\mathbf{z} \Rightarrow 0 = \frac{1}{n}(\mathbf{1}^\top \otimes I_n)(\alpha\mathbf{A}\nabla\mathbf{f}(\bar{\mathbf{x}}) + \mathbf{B}\mathbf{z}) = \frac{\alpha}{n} \sum_{i=1}^n \nabla f(x).$$

The above implies that x is a stationary point of problem (1). □

4.2 Transformation II

We next exploit the structure of the matrices \mathbf{A} , \mathbf{B} , and \mathbf{C} to further transform recursion (21) into a more useful form. Under Assumption 1, the combination matrix W can be decomposed as

$$W = U\Lambda U^\top = [\mathbf{1} \quad \hat{U}] \begin{bmatrix} 1 & 0 \\ 0 & \hat{\Lambda} \end{bmatrix} \begin{bmatrix} \frac{1}{n}\mathbf{1}^\top \\ \hat{U}^\top \end{bmatrix},$$

where $\hat{\Lambda} = \text{diag}\{\lambda_i\}_{i=2}^n$. The matrix U is an orthogonal matrix ($UU^\top = U^\top U = I$) and \hat{U} is an $n \times (n-1)$ matrix that satisfies $\hat{U}\hat{U}^\top = I_n - \frac{1}{n}\mathbf{1}\mathbf{1}^\top$ and $\mathbf{1}^\top\hat{U} = 0$. It follows that

$$\mathbf{W} = \mathbf{U}\Lambda\mathbf{U}^\top = [\mathbf{1} \otimes I_d \quad \hat{U}] \begin{bmatrix} I_d & 0 \\ 0 & \hat{\Lambda} \end{bmatrix} \begin{bmatrix} \frac{1}{n}\mathbf{1}^\top \otimes I_d \\ \hat{U}^\top \end{bmatrix},$$

where $\hat{\Lambda} \triangleq \hat{\Lambda} \otimes I_d \in \mathbb{R}^{d(n-1) \times d(n-1)}$, $\mathbf{U} \in \mathbb{R}^{dn \times dn}$ is an orthogonal matrix, and $\hat{\mathbf{U}} \triangleq \hat{U} \otimes I_d \in \mathbb{R}^{dn \times d(n-1)}$ satisfies:

$$\hat{U}^\top\hat{U} = \mathbf{I}, \quad \hat{\mathbf{U}}\hat{\mathbf{U}}^\top = \mathbf{I} - \frac{1}{n}\mathbf{1}\mathbf{1}^\top \otimes I_d, \quad (\mathbf{1}^\top \otimes I_d)\hat{\mathbf{U}} = 0. \quad (22)$$

Now, since \mathbf{A} , \mathbf{B}^2 , \mathbf{C} are chosen as polynomial function of \mathbf{W} as described in Assumption 1, it holds that

$$\mathbf{A} = \mathbf{U}\Lambda_a\mathbf{U}^\top = [\mathbf{1} \otimes I_d \quad \hat{U}] \begin{bmatrix} I_d & 0 \\ 0 & \hat{\Lambda}_a \end{bmatrix} \begin{bmatrix} \frac{1}{n}\mathbf{1}^\top \otimes I_d \\ \hat{U}^\top \end{bmatrix}, \quad (23a)$$

$$\mathbf{C} = \mathbf{U}\Lambda_c\mathbf{U}^\top = [\mathbf{1} \otimes I_d \quad \hat{U}] \begin{bmatrix} I_d & 0 \\ 0 & \hat{\Lambda}_c \end{bmatrix} \begin{bmatrix} \frac{1}{n}\mathbf{1}^\top \otimes I_d \\ \hat{U}^\top \end{bmatrix}, \quad (23b)$$

$$\mathbf{B}^2 = \mathbf{U}\Lambda_b^2\mathbf{U}^\top = [\mathbf{1} \otimes I_d \quad \hat{U}] \begin{bmatrix} 0 & 0 \\ 0 & \hat{\Lambda}_b^2 \end{bmatrix} \begin{bmatrix} \frac{1}{n}\mathbf{1}^\top \otimes I_d \\ \hat{U}^\top \end{bmatrix}, \quad (23c)$$

where

$$\hat{\Lambda}_a = \text{diag}\{\lambda_{a,i}\}_{i=2}^n \otimes I_d, \quad \hat{\Lambda}_b^2 = \text{diag}\{\lambda_{b,i}^2\}_{i=2}^n \otimes I_d, \quad \hat{\Lambda}_c = \text{diag}\{\lambda_{c,i}\}_{i=2}^n \otimes I_d, \quad (24)$$

with $\lambda_{a,i} \triangleq \sum_{l=0}^p a_l \lambda_i^l$, $\lambda_{b,i}^2 \triangleq \sum_{l=0}^p b_l \lambda_i^l$, and $\lambda_{c,i} \triangleq \sum_{l=0}^p c_l \lambda_i^l$. Moreover, $\hat{\Lambda}_b$ is positive definite due to the null space condition (3). Multiplying both sides of (21) by \mathbf{U}^\top and using the structure (23), we get

$$\mathbf{U}^\top \mathbf{x}^{k+1} = (\Lambda_a \Lambda_c - \Lambda_b^2) \mathbf{U}^\top \mathbf{x}^k - \alpha \Lambda_a \mathbf{U}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k) + \mathbf{w}^k) - \mathbf{U}^\top \mathbf{s}^k, \quad (25a)$$

$$\mathbf{U}^\top \mathbf{s}^{k+1} = \mathbf{U}^\top \mathbf{s}^k + \Lambda_b^2 \mathbf{U}^\top \mathbf{x}^k + \alpha \Lambda_a \mathbf{U}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1}) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)). \quad (25b)$$

Note that

$$\left(\frac{1}{n}\mathbf{1}^\top \otimes I_d\right) \mathbf{s}^k \stackrel{(20c)}{=} \left(\frac{1}{n}\mathbf{1}^\top \otimes I_d\right) (\mathbf{B}\mathbf{z}^k + \alpha \mathbf{A} \nabla \mathbf{f}(\bar{\mathbf{x}}^k)) = \frac{\alpha}{n} \sum_{i=1}^n \nabla f_i(\bar{\mathbf{x}}^k). \quad (26)$$

Moreover, utilizing the structure of \mathbf{U} , we have

$$\begin{aligned} \mathbf{U}^\top \mathbf{x}^k &= \begin{bmatrix} \bar{x}^k \\ \hat{\mathbf{U}}^\top \mathbf{x}^k \end{bmatrix}, & \mathbf{U}^\top \mathbf{s}^k &= \begin{bmatrix} \alpha \bar{\nabla} \mathbf{f}(\bar{\mathbf{x}}^k) \\ \hat{\mathbf{U}}^\top \mathbf{s}^k \end{bmatrix} \\ \mathbf{U}^\top \nabla \mathbf{f}(\mathbf{x}) &= \begin{bmatrix} \bar{\nabla} \mathbf{f}(\mathbf{x}^k) \\ \hat{\mathbf{U}}^\top \nabla \mathbf{f}(\mathbf{x}) \end{bmatrix}, & \mathbf{U}^\top \mathbf{w}^k &= \begin{bmatrix} \bar{\mathbf{w}}^k \\ \hat{\mathbf{U}}^\top \mathbf{w}^k \end{bmatrix}, \end{aligned} \quad (27)$$

where

$$\bar{\nabla} \mathbf{f}(\mathbf{x}^k) \triangleq \frac{1}{n} (\mathbf{1}_n^\top \otimes I_d) \nabla \mathbf{f}(\mathbf{x}^k) = \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_i^k), \quad (28a)$$

$$\bar{\mathbf{w}}^k \triangleq \frac{1}{n} (\mathbf{1}_n^\top \otimes I_d) \mathbf{w}^k = \frac{1}{n} \sum_{i=1}^n (\nabla F_i(x_i^k, \xi_i^k) - \nabla f_i(x_i^k)). \quad (28b)$$

Hence, using the previous quantities and the structure of $\Lambda_a, \Lambda_b^2, \Lambda_c$ given in (23) and (24), we find that

$$\bar{x}^{k+1} = \bar{x}^k - \alpha \bar{\nabla} \mathbf{f}(\mathbf{x}^k) - \alpha \bar{\mathbf{w}}^k, \quad (29a)$$

$$\hat{\mathbf{U}}^\top \mathbf{x}^{k+1} = (\hat{\Lambda}_a \hat{\Lambda}_c - \hat{\Lambda}_b^2) \hat{\mathbf{U}}^\top \mathbf{x}^k - \alpha \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k) + \mathbf{w}^k) - \hat{\mathbf{U}}^\top \mathbf{s}^k, \quad (29b)$$

$$\hat{\mathbf{U}}^\top \mathbf{s}^{k+1} = \hat{\mathbf{U}}^\top \mathbf{s}^k + \hat{\Lambda}_b^2 \hat{\mathbf{U}}^\top \mathbf{x}^k + \alpha \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1}) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)). \quad (29c)$$

Multiplying the third equation by $\hat{\Lambda}_b^{-1}$ and rewriting the previous recursion in matrix notation, we obtain:

$$\bar{x}^{k+1} = \bar{x}^k - \alpha \bar{\nabla} \mathbf{f}(\mathbf{x}^k) - \alpha \bar{\mathbf{w}}^k, \quad (30a)$$

$$\begin{bmatrix} \hat{\mathbf{U}}^\top \mathbf{x}^{k+1} \\ \hat{\Lambda}_b^{-1} \hat{\mathbf{U}}^\top \mathbf{s}^{k+1} \end{bmatrix} = \begin{bmatrix} \hat{\Lambda}_a \hat{\Lambda}_c - \hat{\Lambda}_b^2 & -\hat{\Lambda}_b \\ \hat{\Lambda}_b & \mathbf{I} \end{bmatrix} \begin{bmatrix} \hat{\mathbf{U}}^\top \mathbf{x}^k \\ \hat{\Lambda}_b^{-1} \hat{\mathbf{U}}^\top \mathbf{s}^k \end{bmatrix} - \alpha \begin{bmatrix} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k) + \mathbf{w}^k) \\ \hat{\Lambda}_b^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})) \end{bmatrix}. \quad (30b)$$

The convergence of (30) is governed by the matrix:

$$\mathbf{G} \triangleq \begin{bmatrix} \hat{\Lambda}_a \hat{\Lambda}_c - \hat{\Lambda}_b^2 & -\hat{\Lambda}_b \\ \hat{\Lambda}_b & \mathbf{I} \end{bmatrix} \in \mathbb{R}^{2d(n-1) \times 2d(n-1)}. \quad (31)$$

We next introduce a fundamental factorization of \mathbf{G} that will be used to transform (30) into our final key recursion. The next result is proven in Appendix B.1.

Lemma 1 (Fundamental factorization). *Suppose that the eigenvalues of \mathbf{G} are strictly less than one in magnitude. Then, there exists an invertible matrix $\hat{\mathbf{V}}$ such that the matrix \mathbf{G} admits the similarity transformation*

$$\mathbf{G} = \hat{\mathbf{V}} \mathbf{\Gamma} \hat{\mathbf{V}}^{-1}, \quad (32)$$

where $\mathbf{\Gamma}$ satisfies $\|\mathbf{\Gamma}\| < 1$. □

For the convergence of (30), it is necessary that \mathbf{G} is a stable matrix (has eigenvalues strictly less than one in magnitude). To see this suppose that $\nabla \mathbf{f}(\mathbf{x}) = \mathbf{w}^k = \mathbf{0}$, then the convergence is dictated by (30b), which diverges for unstable \mathbf{G} if $\hat{\mathbf{U}}^\top \mathbf{x}^0 \neq \mathbf{0}$ or $\hat{\mathbf{U}}^\top \mathbf{s}^0 \neq \mathbf{0}$. Thus, we implicitly assume that \mathbf{G} is a stable matrix. Explicit expressions for $\mathbf{\Gamma}$ and $\hat{\mathbf{V}}$ for ED/D², EXTRA, and GT methods are derived in Appendix B.2, where we also find exact expressions for the eigenvalues of \mathbf{G} for these methods.

Finally, multiplying both sides of (30b) by $\frac{1}{v} \hat{\mathbf{V}}^{-1}$ for any $v > 0$ and using the structure (32), we arrive at the following key result.

Lemma 2 (Final transformed recursion). *Under Assumption 1, there exists an invertible matrix $\hat{\mathbf{V}}$ such that recursion (30) can be transformed into*

$$\bar{x}^{k+1} = \bar{x}^k - \alpha \overline{\nabla \mathbf{f}}(\mathbf{x}^k) - \alpha \bar{\mathbf{w}}^k, \quad (33a)$$

$$\hat{\mathbf{e}}^{k+1} = \mathbf{\Gamma} \hat{\mathbf{e}}^k - \alpha \hat{\mathbf{V}}^{-1} \begin{bmatrix} \frac{1}{v} \hat{\mathbf{\Lambda}}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k) + \mathbf{w}^k) \\ \frac{1}{v} \hat{\mathbf{\Lambda}}_b^{-1} \hat{\mathbf{\Lambda}}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})) \end{bmatrix}, \quad (33b)$$

where $\mathbf{\Gamma}$ was introduced in (32), v is an arbitrary strictly positive constant, and

$$\hat{\mathbf{e}}^k \triangleq \frac{1}{v} \hat{\mathbf{V}}^{-1} \begin{bmatrix} \hat{\mathbf{U}}^\top \mathbf{x}^k \\ \hat{\mathbf{\Lambda}}_b^{-1} \hat{\mathbf{U}}^\top \mathbf{s}^k \end{bmatrix}. \quad (34)$$

□

Remark 2. Recall that $\bar{\mathbf{x}}^k = \mathbf{1} \otimes \bar{x}^k$ where $\bar{x}^k = \frac{1}{n} \sum_{i=1}^n x_i^k$. Since $\hat{\mathbf{U}}^\top \hat{\mathbf{U}} = \mathbf{I}$, it holds that

$$\|\hat{\mathbf{U}}^\top \mathbf{x}\|^2 = \mathbf{x}^\top \hat{\mathbf{U}} \hat{\mathbf{U}}^\top \hat{\mathbf{U}} \hat{\mathbf{U}}^\top \mathbf{x} = \|\hat{\mathbf{U}} \hat{\mathbf{U}}^\top \mathbf{x}^k\|^2 \stackrel{(22)}{=} \|\mathbf{x}^k - \bar{\mathbf{x}}^k\|^2.$$

Therefore, the quantity $\|\hat{\mathbf{U}}^\top \mathbf{x}^k\|^2$ measures the deviation of \mathbf{x}^k from the average $\bar{\mathbf{x}}^k$. Similarly, $\|\hat{\mathbf{U}}^\top \mathbf{s}^k\|^2$ measures the deviation of \mathbf{s}^k from the average $(\frac{1}{n} \mathbf{1} \mathbf{1}^\top \otimes I_n) \mathbf{s}^k$. Now, using (34), we have

$$\|v \hat{\mathbf{V}} \hat{\mathbf{e}}^k\|^2 = \|\hat{\mathbf{U}}^\top \mathbf{x}^k\|^2 + \|\hat{\mathbf{\Lambda}}_b^{-1} \hat{\mathbf{U}}^\top \mathbf{s}^k\|^2. \quad (35)$$

Thus, the vector $\hat{\mathbf{e}}^k$ can be interpreted as a measure of the deviation of \mathbf{x}^k and a weighted \mathbf{s}^k from the averages $\bar{\mathbf{x}}^k$ and $\frac{1}{n} (\mathbf{1} \mathbf{1}^\top \otimes I_n) \mathbf{s}^k$, respectively. □

5. Convergence Results

The convergence results are stated and discussed in this section.

5.1 Convergence Under General Non-Convex Costs

The following result establishes the convergence under general non-convex smooth costs.

Theorem 1 (Convergence of SUDA). *Suppose that Assumptions 1–3 hold and the step size satisfies $\alpha \leq \min \left\{ \frac{1}{2L}, \frac{1-\gamma}{2Lv_1v_2\lambda_a}, \frac{\sqrt{\lambda_b(1-\gamma)}}{2L\sqrt{v_1v_2\lambda_a}} \right\}$. Then, the iterates $\{\mathbf{x}^k\}$ of SUDA with $\mathbf{x}^0 = \mathbf{1} \otimes x^0$ ($x^0 \in \mathbb{R}^d$) satisfy*

$$\begin{aligned} & \frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\nabla f(\bar{x}^k)\|^2 + \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 \\ & \leq \frac{8(f(x^0) - f^*)}{\alpha K} + \frac{4\alpha L\sigma^2}{n} + \frac{12\alpha^2 L^2 v_1^2 v_2^2 \zeta_0^2}{\lambda_b^2 (1-\gamma) K} + \frac{16\alpha^2 L^2 v_1^2 v_2^2 \lambda_a^2 \sigma^2}{1-\gamma} + \frac{16\alpha^4 L^4 v_1^2 v_2^2 \lambda_a^2 \sigma^2}{\lambda_b^2 (1-\gamma)^2 n}, \end{aligned} \quad (36)$$

where $v_1 \triangleq \|\hat{\mathbf{V}}\|$, $v_2 \triangleq \|\hat{\mathbf{V}}^{-1}\|$ and

$$\begin{aligned} \gamma & \triangleq \|\mathbf{\Gamma}\| < 1, \quad \lambda_b \triangleq \frac{1}{\|\mathbf{\Lambda}_b^{-1}\|}, \quad \lambda_a \triangleq \|\mathbf{\Lambda}_a\|, \\ \zeta_0 & \triangleq \frac{1}{n} \|(\mathbf{A} - \frac{1}{n} \mathbf{1} \mathbf{1}^\top \otimes I_d)(\nabla \mathbf{f}(\mathbf{x}^0) - \mathbf{1} \otimes \nabla f(x^0))\|. \end{aligned}$$

Consequently, if we set $\alpha = \frac{1}{2L\beta + \sigma\sqrt{K/n}}$ where $\beta \triangleq 1 + \frac{v_1 v_2 \lambda_a}{1-\gamma} + \frac{\sqrt{v_1 v_2 \lambda_a}}{\sqrt{\lambda_b(1-\gamma)}}$. Then, we obtain the convergence rate

$$\frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 \leq O\left(\frac{\sigma}{\sqrt{nK}} + \frac{1}{K} + \frac{n\sigma^2}{n + \sigma^2 K} + \frac{n\zeta_0^2}{nK + \sigma^2 K^2}\right). \quad (37)$$

□

The proof of Theorem 1 is established in Appendix C. The convergence rate (37) shows that SUDA enjoys linear speedup since the dominating term is $O(1/\sqrt{nK})$ for sufficiently large K [3]. Note that the rate given in (37) treats the network quantities $\{\gamma, \lambda_a, \lambda_b, v_1, v_2\}$ as constants. However, these quantities can have a significant influence on the convergence rate as explained in the introduction. While the above result holds for EXTRA, ED/D² and GT methods, it is still unclear whether these methods can achieve enhanced transient time compared to DSGD. To show the network affect and see the implication of Theorem 1 in terms of network quantities, we will specialize Theorem 1 to ED/D² (5) and ATC-GT (7) and reflect the values of the $\{\gamma, \lambda_a, \lambda_b, v_1, v_2\}$ in the convergence rate.

Corollary 1 (ED/D² convergence). *Suppose that all the conditions given in Theorem 1 hold and assume further that W is positive semi-definite. If we set $\alpha = \sqrt{n/K}$ and $K \geq \max \left\{ 4nL^2, \frac{32L^2\lambda^2 n}{(1-\sqrt{\lambda})^2 \lambda}, \frac{\sqrt{32}L^2\lambda n}{\sqrt{1-\lambda}(1-\sqrt{\lambda})\sqrt{\lambda}} \right\}$, then ED/D² (5) with $\mathbf{x}^0 = \mathbf{1} \otimes x^0$ ($x^0 \in \mathbb{R}^d$) has convergence rate*

$$\frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 \leq O\left(\frac{f(x^0) - f^*}{\sqrt{nK}} + \frac{\sigma^2}{\sqrt{nK}} + \frac{n\lambda^2\sigma^2}{(1-\lambda)\lambda K} + \frac{n\lambda^2\sigma^2}{(1-\lambda)^3\lambda K^2} + \frac{nL^2\zeta_0^2}{(1-\lambda)^2 K^2}\right), \quad (38)$$

where $\underline{\lambda}$ is the minimum non-zero eigenvalue of W . Moreover, we have

$$\zeta_0^2 = \frac{1}{n} \sum_{i=1}^n \left\| \sum_{j \in \mathcal{N}_i} w_{ij} \nabla f_j(x^0) - \nabla f(x^0) \right\|^2 \leq \lambda^2 \zeta_0^2,$$

where $\zeta_0^2 \triangleq \frac{1}{n} \sum_{i=1}^n \left\| \nabla f_i(x^0) - \nabla f(x^0) \right\|^2$.

Proof. The proof follows by substituting the bounds (61) established in Appendix B.2 into (36). \square

Corollary 2 (ATC-GT convergence). *Suppose that all the conditions given in Theorem 1 hold and assume further that W is positive semi-definite. If we let $\alpha = \sqrt{n/K}$ and $K \geq \max \left\{ 4nL^2, \frac{432L^2\lambda^4n}{(1-\lambda)^2}, \frac{72L^2\lambda^2n}{(1-\lambda)^2} \right\}$, then ATC-GT (7) with $\mathbf{x}^0 = \mathbf{1} \otimes x^0$ ($x^0 \in \mathbb{R}^d$) has convergence rate*

$$\frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E} \left\| \overline{\nabla \mathbf{f}}(\mathbf{x}^k) \right\|^2 \leq O \left(\frac{f(x^0) - f^*}{\sqrt{nK}} + \frac{\sigma^2}{\sqrt{nK}} + \frac{n\lambda^4\sigma^2}{(1-\lambda)K} + \frac{n\lambda^4\sigma^2}{(1-\lambda)^4K^2} + \frac{n\zeta_0^2}{(1-\lambda)^3K^2} \right). \quad (39)$$

Moreover, we have

$$\zeta_0^2 = \frac{1}{n} \sum_{i=1}^n \left\| \sum_{j \in \mathcal{N}_i} [w_{ij}]^2 \nabla f_j(x^0) - \nabla f(x^0) \right\|^2 \leq \lambda^4 \zeta_0^2,$$

where $\zeta_0^2 \triangleq \frac{1}{n} \sum_{i=1}^n \left\| \nabla f_i(x^0) - \nabla f(x^0) \right\|^2$.

Proof. The proof follows by substituting the bounds (62) established in Appendix B.2 into Theorem 1. \square

Note that linear speedup is achieved when the dominating term is $O(\frac{1}{\sqrt{nK}})$. This is case when K is sufficiently large so that the higher order terms are less than or equal to $O(\frac{1}{\sqrt{nK}})$. Table 1 compares the above results with existing works. In Table 1 we treat $\underline{\lambda}$ for ED/D² as constant since, for example, if we set $W \leftarrow (1-\theta)W + \theta I$ for constant $\theta > 0$, then it holds that $\underline{\lambda} \in (\theta, 1)$. It is clear that our bounds are tighter in terms of the spectral gap $1-\lambda$. Moreover, ED/D² and GT methods have enhanced transient time compared to DSGD.

Remark 3 (Step size selection). We remark that in Corollaries 1 and 2, the step size is chosen as $\alpha = \sqrt{n/K}$ to simplify the expressions. This choice is not optimal and tighter rates can be obtained if we meticulously select the step size. For example, we can select the step size as in Theorem 1 and obtain a rate similar to (37) where the dominating term is $O(\sigma/\sqrt{nK})$ instead of $O(\sigma^2/\sqrt{nK})$. We can even get a tighter rate by carefully selecting the step size similar to [43]. However, such choices does not affect the transient time order in terms of network quantities, which is the main conclusion of our results. \square

Remark 4 (EXTRA and other GT variations). We remark that the matrix \mathbf{G} defined in (31) is identical for both EXTRA (6) and ED/D². Hence, the convergence rate of ED/D² given in (38) also holds for EXTRA with the exception of the value of λ^2 in the *numerators*, which should be replaced by one for EXTRA ($\lambda^2 \rightarrow 1$ in numerators). Likewise, for the non-ATC-GT (8) and semi-ATC-GT (see Appendix A.1), the matrix \mathbf{G} is identical to ATC-GT and the convergence rate of ATC-GT given (39) holds for these other variations except for the value of λ^2 in the *numerators*, which is one for non-ATC-GT ($\lambda^2 \rightarrow 1$ in numerators) and λ for the semi-ATC-GT ($\lambda^2 \rightarrow \lambda$ in numerators). Please see Appendix B.2 for more details. \square

5.2 Convergence Under PL Condition

We next state the convergence of SUDA under the PL condition given in Assumption 4.

Theorem 2 (PL case). *Suppose that Assumptions 1–4 hold and the step size satisfies:*

$$\alpha \leq \min \left\{ \frac{1-\gamma}{3L}, \frac{\lambda_b}{2L}, \frac{1-\gamma}{\sqrt{6}Lv_1v_2\lambda_a}, \left(\frac{\mu\lambda_b^2(1-\gamma)}{8L^4v_1^2v_2^2} \right)^{1/3} \right\}. \quad (40)$$

Then, the iterates $\{\mathbf{x}^k\}$ of SUDA with $\mathbf{x}^0 = \mathbf{1} \otimes x^0$ ($x^0 \in \mathbb{R}^d$) satisfy

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[f(x_i^k) - f^*] \leq (1 - \frac{\alpha\mu}{2})^k r_0 + O \left(\frac{\alpha L \sigma^2}{\mu n} + \frac{\alpha^2 L^2 v_1^2 v_2^2 \lambda_a^2 \sigma^2}{\mu(1-\gamma)} + \frac{\alpha^4 L^4 v_1^2 v_2^2 \lambda_a^2 \sigma^2}{\mu \lambda_b^2 (1-\gamma)^2 n} \right), \quad (41)$$

where $r_0 = 2\mathbb{E}[f(x^0) - f^*] + \alpha^2 L v_1^2 v_2^2 \zeta_0^2 / \lambda_b^2$ and the quantities $v_1, v_2, \gamma, \lambda_a, \lambda_b$ and ζ_0 are defined as in Theorem 1. \square

The proof of Theorem 2 is given in Appendix C.3. To discuss the implication of the above result, we specialize it to ED/D² and ATC-GT.

Corollary 3 (ED/D² convergence under PL condition). *Let the same conditions as in Theorem 2 hold and suppose further that W is positive semi-definite. If the step size satisfies*

$$\alpha \leq \min \left\{ \frac{1-\sqrt{\lambda}}{3L}, \frac{\sqrt{1-\lambda}}{2L}, \frac{(1-\sqrt{\lambda})\lambda}{2\sqrt{12}L\lambda}, \left(\frac{\mu(1-\lambda)(1-\sqrt{\lambda})\lambda}{18L^4} \right)^{1/3} \right\}, \text{ then ED/D}^2 \text{ (5) with } \mathbf{x}^0 = \mathbf{1} \otimes x^0 \text{ (} x^0 \in \mathbb{R}^d \text{) has convergence rate}$$

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[f(x_i^k) - f^*] \leq (1 - \frac{\alpha\mu}{2})^k r_0 + O \left(\frac{\alpha L \sigma^2}{\mu n} + \frac{\alpha^2 L^2 \lambda^2 \sigma^2}{\mu(1-\lambda)\lambda} + \frac{\alpha^4 L^4 \lambda^2 \sigma^2}{\mu(1-\lambda)^3 \lambda n} \right), \quad (42)$$

where $r_0 = 2\mathbb{E}[f(x^0) - f^*] + 8\alpha^2 L \zeta_0^2 / (1-\lambda)\lambda$, $\zeta_0^2 = \frac{1}{n} \sum_{i=1}^n \left\| \sum_{j \in \mathcal{N}_i} w_{ij} \nabla f_j(x^0) - \nabla f(x^0) \right\|^2$, and λ is the minimum non-zero eigenvalue of W . Hence, selecting $\alpha = 2 \ln(K^2) / \mu K$, we obtain

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[f(x_i^k) - f^*]$$

$$\leq \frac{2\mathbb{E}[f(x^0) - f^*]}{K^2} + \tilde{O}\left(\frac{\sigma^2}{Kn} + \frac{\lambda^2\sigma^2}{K^2(1-\lambda)\underline{\lambda}} + \frac{\zeta_0^2}{K^4(1-\lambda)\underline{\lambda}} + \frac{\lambda^2\sigma^2}{\mu^4(1-\lambda)^3\underline{\lambda}n}\right), \quad (43)$$

where $\tilde{O}(\cdot)$ hides logarithmic factors.

Proof. Equation (42) follows from Theorem 2 and the bounds (61) derived in Appendix B.2. Now, if we set $\alpha = 2\ln(K^2)/\mu K$, then $1 - \frac{\alpha\mu}{2} \leq \exp(-\alpha\mu K/2) = \frac{1}{K^2}$ where $\exp(\cdot)$ denote the exponential function. Hence, for $\alpha = 2\ln(K^2)/\mu K$ and large enough K , inequality (42) can be upper bounded by (43). \square

Corollary 4 (ATC-GT convergence under PL condition). *Let the same conditions as in Theorem 2 hold and assume further that W is positive semi-definite. Then, ATC-GT (7) with $\mathbf{x}^0 = \mathbf{1} \otimes x^0$ ($x^0 \in \mathbb{R}^d$) and $\alpha \leq \min\left\{\frac{1-\lambda}{6L}, \frac{1-\lambda}{6\sqrt{18}L\lambda^2}, \left(\frac{\mu(1-\lambda)^3}{432L^4}\right)^{1/3}\right\}$ has convergence rate:*

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[f(x_i^k) - f^*] \leq (1 - \frac{\alpha\mu}{2})^k r_0 + O\left(\frac{\alpha L\sigma^2}{\mu n} + \frac{\alpha^2 L^2 \lambda^4 \sigma^2}{\mu(1-\lambda)} + \frac{\alpha^4 L^4 \lambda^4 \sigma^2}{\mu(1-\lambda)^4 n}\right). \quad (44)$$

where $r_0 = 2\mathbb{E}[f(x^0) - f^*] + 27\alpha^2 L \zeta_0^2 / (1-\lambda)^2$ and $\zeta_0^2 = \frac{1}{n} \sum_{i=1}^n \left\| \sum_{j \in \mathcal{N}_i} [w_{ij}]^2 \nabla f_j(x^0) - \nabla f(x^0) \right\|^2$. Hence, if we set $\alpha = 2\ln(K^2)/\mu K$, then it holds that

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[f(x_i^k) - f^*] \\ & \leq \frac{2\mathbb{E}[f(x^0) - f^*]}{K^2} + \tilde{O}\left(\frac{\sigma^2}{Kn} + \frac{\lambda^4\sigma^2}{K^2(1-\lambda)} + \frac{\zeta_0^2}{K^4(1-\lambda)^2} + \frac{\lambda^4\sigma^2}{K^4(1-\lambda)^4 n}\right), \end{aligned} \quad (45)$$

where $\tilde{O}(\cdot)$ hides logarithmic factors.

Proof. Equation (44) follows from Theorem 2 and the bounds (62) derived in Appendix B.2. Inequality (45) follows by substituting $\alpha = 2\ln(K^2)/\mu K$, into (62) and using $1 - \frac{\alpha\mu}{2} \leq \exp(-\alpha\mu K/2) = \frac{1}{K^2}$. \square

We note that as in Remark 3, the selected step sizes in Corollaries 3 and 4 are clearly not optimal. The hidden log factors can be removed if adopt decaying step-sizes techniques (e.g., [39]). However, the main conclusion we want to emphasize is the network dependent bounds, which do not change if select better step size choice. Under the PL condition, linear speedup is achieved when K is large enough such that the dominating term is $O(\frac{1}{nK})$. Table 2 lists the transient times implied by the above result and compares them with existing results. It is clear that our results significantly improves upon existing GT results. Moreover, our bound for ED/D² matches the existing bound, which are under the stronger assumption of strong-convexity.

Remark 5 (Steady-state error). For constant step size α independent of k , we can let k goes to ∞ in (42) and (44) to arrive at the following steady state results.

- For ED/D² (5), we have

$$\limsup_{k \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \mathbb{E}[f(x_i^k) - f^*] \leq O\left(\frac{\alpha L \sigma^2}{\mu n} + \frac{\alpha^2 L^2 \lambda^2 \sigma^2}{\mu(1-\lambda)} + \frac{\alpha^4 L^4 \lambda^2 \sigma^2}{\mu(1-\lambda)^3 n}\right). \quad (46)$$

- For ATC-GT (7), we have

$$\limsup_{k \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \mathbb{E}[f(x_i^k) - f^*] \leq O\left(\frac{\alpha L \sigma^2}{\mu n} + \frac{\alpha^2 L^2 \lambda^4 \sigma^2}{\mu(1-\lambda)} + \frac{\alpha^4 L^4 \lambda^4 \sigma^2}{\mu(1-\lambda)^4 n}\right). \quad (47)$$

The bound (46) for ED/D² has the same network dependent bounds as in the strongly-convex case [14]. Moreover, the bound (47) for ATC-GT improves upon existing bounds for both strongly-convex [22] and PL settings [24], which are on the order of $O(\alpha^2 \lambda^2 \sigma^2 / (1-\lambda)^3)$. \square

6. Simulation Results

In this section, we validate the established theoretical results with numerical simulations.

6.1 Simulation for non-convex problems

The problem. We consider the logistic regression problem with a non-convex regularization term [24, 53]. The problem formulation is given by $\min_{x \in \mathbb{R}^d} \frac{1}{n} \sum_{i=1}^n f_i(x) + \rho r(x)$, where

$$f_i(x) = \frac{1}{L} \sum_{\ell=1}^L \ln(1 + \exp(-y_{i,\ell} h_{i,\ell}^\top x)) \quad \text{and} \quad r(x) = \sum_{j=1}^d \frac{x(j)^2}{1 + x(j)^2}. \quad (48)$$

In the above problem, $x = \text{col}\{x(j)\}_{j=1}^d \in \mathbb{R}^d$ is the unknown variable to be optimized, $\{h_{i,\ell}, y_{i,\ell}\}_{\ell=1}^L$ is the training dataset held by agent i in which $h_{i,\ell} \in \mathbb{R}^d$ is a feature vector while $y_{i,\ell} \in \{-1, +1\}$ is the corresponding label. The regularization $r(x)$ is a smooth but non-convex function and the regularization constant $\rho > 0$ controls the influence of $r(x)$.

Experimental settings. In our experiments, we set $d = 20$, $L = 2000$ and $\rho = 0.001$. To control data heterogeneity across the agents, we first let each agent i be associated with a local solution x_i^* , and such x_i^* is generated by $x_i^* = x^* + v_i$ where $x^* \sim \mathcal{N}(0, I_d)$ is a randomly generated vector while $v_i \sim \mathcal{N}(0, \sigma_h^2 I_d)$ controls the similarity between each local solution. Generally speaking, a large σ_h^2 result in local solutions $\{x_i^*\}$ that are vastly different from each other. With x_i^* at hand, we can generate local data that follows distinct distributions. At agent i , we generate each feature vector $h_{i,\ell} \sim \mathcal{N}(0, I_d)$. To produce the corresponding label $y_{i,\ell}$, we generate a random variable $z_{i,\ell} \sim \mathcal{U}(0, 1)$. If $z_{i,\ell} \leq 1 + \exp(-y_{i,\ell} h_{i,\ell}^\top x_i^*)$, we set $y_{i,\ell} = 1$; otherwise $y_{i,\ell} = -1$. Clearly, solution x_i^* controls the distribution of the labels. In this way, we can easily control data heterogeneity by adjusting σ_h^2 . Furthermore, to easily control the influence of gradient noise, we will achieve the stochastic gradient by imposing




Figure 1: Performance of different stochastic algorithms to solve problem (48) with homogeneous data. Constant and learning decaying rates are used in the left and right plots, respectively. All algorithms in both plots are over the ring topology with $\lambda = 0.99$.

a Gaussian noise to the real gradient, *i.e.*, $\widehat{\nabla}f_i(x) = \nabla f_i(x) + s_i$ in which $s_i \sim \mathcal{N}(0, \sigma_n^2 I_d)$. We can control the magnitude of the gradient noise by adjusting σ_n^2 . The metric for all simulations is $\mathbb{E} \|\nabla f(\bar{x})\|^2$ where $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$.

Performances of SUDA with and without data heterogeneity. In this set of simulations, we will test the performance of ED/D² and ATC-GT (which are covered by the SUDA framework) with constant and decaying step size, and compare them with DSGD. In the simulation, we organize $n = 32$ agents into an undirected ring topology.

Fig. 1 shows the performances of these algorithms with homogeneous data, *i.e.*, $\sigma_h^2 = 0$. The gradient noise magnitude is set as $\sigma_n^2 = 0.001$. In the left plot, we set up a constant step size $\alpha = 0.01$. In the right plot, we set an initial step size as 0.01, and then scale it by 0.5 after every 100 iterations. It is observed in Fig. 1 that all stochastic algorithms perform similarly to each other with homogeneous data. Fig. 2 shows the performance under heterogeneous data settings with $\sigma_h^2 = 0.2$. The gradient noise and the step size values are the same as in Fig. 1. It is clear from Fig. 2 that ED/D² and ATC-GT are more robust to data heterogeneity compared to DSGD. We see that ED/D² can converge as well as PSGD while ATC-GT performs slightly worse than ED/D².

Influence of network topology. In this set of simulations, we will test the influence of the spectral gap $1 - \lambda$ on various decentralized stochastic algorithms. We generate four topologies: the Erdos-Renyi graph with probability 0.8, the Ring topology, the Grid topology, and the scaled Ring topology with $\lambda = (9 + \lambda_{\text{Ring}})/10$. The value of the mixing rate λ for each topology is listed in the caption in Fig. 3. We utilize a constant step size 0.01 for each plot. It is observed in Fig. 3 that each decentralized algorithm will converge to a less accurate solution as $\lambda \rightarrow 1$ while PSGD is immune to the network topology. In addition, it is also observed that ED/D² is least sensitive to the network topology while DSGD is most sensitive (under heterogeneous setting), which is consistent with the our results listed in Table 1.




Figure 2: Performance of different stochastic algorithms to solve problem (48) with heterogeneous data. Constant and learning decaying rates are used in the left and right plots, respectively. All algorithms in both plots are over the ring topology with $\lambda = 0.99$.




Figure 3: Performance of different stochastic algorithms to solve problem (48) with different topologies. Top-left: Erdos-Renyi random graph with $\lambda = 0.32$; Top-right: Grid with $\lambda = 0.94$; Bottom-left: Ring with $\lambda = 0.99$; Bottom-right: Scaled Ring with $\lambda = 0.999$




Figure 4: Performance of different stochastic algorithms to solve non-convex problem under PL condition with different topologies. Left: Erdos-Renyi random graph with $\lambda = 0.28$; Right: Erdos-Renyi random graph with $\lambda = 0.87$

6.2 Simulation for problems under PL condition

The problem. In this section we examine the performance of ED/D² and GT algorithms for the non-convex problem under PL condition. We consider the same setup used in [24] where the problem formulation is given by $\min_{x \in \mathbb{R}^d} \frac{1}{n} \sum_{i=1}^n f_i(x)$ with $f_i(x) = x^2 + 3 \sin^2(x) + a_i x \cos(x)$. By letting $\sum_{i=1}^n a_i = 0$, we have $f(x) = \frac{1}{n} \sum_{i=1}^n f_i(x) = x^2 + 3 \sin^2(x)$ which is a non-convex cost function that satisfies the PL condition [26].

Experimental settings. We set $n = 32$ in all simulations. To generate a_i , we let $a_i = \sigma_h^2 \cdot i$ and $a_{n-i} = -a_i$ for $i \in \{1, \dots, n/2\}$ where σ_h^2 is used to control data heterogeneity. In this way, we can guarantee $\sum_{i=1}^n a_i = 0$. Similar to Sec. 6, we will achieve the stochastic gradient by imposing a Gaussian noise to the real gradient, *i.e.*, $\widehat{\nabla} f_i(x) = \nabla f_i(x) + s_i$ in which $s_i \sim \mathcal{N}(0, \sigma_n^2 I_d)$. The metric for all simulations is $\mathbb{E} f(\bar{x}) - f^*$ where $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$.

Influence of network topology. We test the influence of the network topology on various stochastic decentralized methods. In simulations, we set data heterogeneity $\sigma_h^2 = 2$ and gradient noise $\sigma_n^2 = 0.1$. We generate two types of topologies: an Erdos-Renyi random graph with $\lambda = 0.28$, and an Erdos-Renyi random graph with $\lambda = 0.87$. We employ a constant step size 0.008 for all tested algorithms. It is observed in Fig. 4 that the performance of all algorithms can be deteriorated by the badly-connected network topology when $\lambda \rightarrow 1$. We see that ED/D² is least sensitive to the network topology while DSGD is most sensitive (under heterogeneous setting), which is consistent with the our results listed in Table 2.

7. Conclusion

In this work, we analyzed the convergence properties of SUDA (10) for decentralized stochastic non-convex optimization problems. SUDA is a general algorithmic framework that includes several state of the art decentralized methods as special case such as EXTRA, Exact-

Diffusion/ D^2 , and gradient-tracking methods. We established the convergence of **SUDA** under both general non-convex and PL condition settings. Explicit convergence rate bounds are provided in terms of the problem parameters and the network topology. When specializing **SUDA** to the particular instances of ED/ D^2 , EXTRA, and GT-methods, we achieve improved network topology dependent rates compared to existing results under non-convex settings. Moreover, our rate shows that ED/ D^2 , EXTRA, and GT-methods are less sensitive to network topology compared to DSGD under heterogeneous data setting.

References

- [1] S. Boyd, N. Parikh, E. Chu, B. Peleato, and J. Eckstein, “Distributed optimization and statistical learning via alternating direction method of multipliers,” *Found. Trends Mach. Lear.*, vol. 3, pp. 1–122, Jan. 2011.
- [2] M. Li, D. G. Andersen, J. W. Park, A. J. Smola, A. Ahmed, V. Josifovski, J. Long, E. J. Shekita, and B.-Y. Su, “Scaling distributed machine learning with the parameter server,” in *11th USENIX Symposium on Operating Systems Design and Implementation (OSDI 14)*, (Broomfield, CO), pp. 583–598, Oct. 2014.
- [3] X. Lian, C. Zhang, H. Zhang, C.-J. Hsieh, W. Zhang, and J. Liu, “Can decentralized algorithms outperform centralized algorithms? A case study for decentralized parallel stochastic gradient descent,” in *Advances in Neural Information Processing Systems (NIPS)*, (Long Beach, CA, USA), pp. 5330–5340, 2017.
- [4] M. Assran, N. Loizou, N. Ballas, and M. Rabbat, “Stochastic gradient push for distributed deep learning,” in *International Conference on Machine Learning*, pp. 344–353, 2019.
- [5] Y. Chen, K. Yuan, Y. Zhang, P. Pan, Y. Xu, and W. Yin, “Accelerating gossip SGD with periodic global averaging,” in *International Conference on Machine Learning*, pp. 1791–1802, PMLR, 2021. Also available as arXiv preprint:2105.09080.
- [6] S. S. Ram, A. Nedic, and V. V. Veeravalli, “Distributed stochastic subgradient projection algorithms for convex optimization,” *J. Optim. Theory Appl.*, vol. 147, no. 3, pp. 516–545, 2010.
- [7] F. S. Cattivelli and A. H. Sayed, “Diffusion LMS strategies for distributed estimation,” *IEEE Trans. Signal Process.*, vol. 58, no. 3, p. 1035, 2010.
- [8] J. Chen and A. H. Sayed, “Distributed pareto optimization via diffusion strategies,” *IEEE J. Sel. Topics Signal Process.*, vol. 7, pp. 205–220, April 2013.
- [9] A. Nedic and A. Ozdaglar, “Distributed subgradient methods for multi-agent optimization,” *IEEE Transactions on Automatic Control*, vol. 54, no. 1, pp. 48–61, 2009.
- [10] K. Yuan, Q. Ling, and W. Yin, “On the convergence of decentralized gradient descent,” *SIAM Journal on Optimization*, vol. 26, no. 3, pp. 1835–1854, 2016.

- [11] W. Shi, Q. Ling, G. Wu, and W. Yin, “EXTRA: An exact first-order algorithm for decentralized consensus optimization,” *SIAM Journal on Optimization*, vol. 25, no. 2, pp. 944–966, 2015.
- [12] K. Yuan, B. Ying, X. Zhao, and A. H. Sayed, “Exact diffusion for distributed optimization and learning-Part I: Algorithm development,” *IEEE Transactions on Signal Processing*, vol. 67, pp. 708–723, Feb. 2019.
- [13] Z. Li, W. Shi, and M. Yan, “A decentralized proximal-gradient method with network independent step-sizes and separated convergence rates,” *IEEE Transactions on Signal Processing*, vol. 67, pp. 4494–4506, Sept. 2019.
- [14] K. Yuan, S. A. Alghunaim, B. Ying, and A. H. Sayed, “On the influence of bias-correction on distributed stochastic optimization,” *IEEE Transactions on Signal Processing*, vol. 68, pp. 4352–4367, 2020.
- [15] H. Tang, X. Lian, M. Yan, C. Zhang, and J. Liu, “D²: Decentralized training over decentralized data,” in *International Conference on Machine Learning*, (Stockholm, Sweden), pp. 4848–4856, 2018.
- [16] J. Xu, S. Zhu, Y. C. Soh, and L. Xie, “Augmented distributed gradient methods for multi-agent optimization under uncoordinated constant stepsizes,” in *Proc. 54th IEEE Conference on Decision and Control (CDC)*, (Osaka, Japan), pp. 2055–2060, 2015.
- [17] P. Di Lorenzo and G. Scutari, “Next: In-network nonconvex optimization,” *IEEE Transactions on Signal and Information Processing over Networks*, vol. 2, no. 2, pp. 120–136, 2016.
- [18] G. Qu and N. Li, “Harnessing smoothness to accelerate distributed optimization,” *IEEE Transactions on Control of Network Systems*, vol. 5, pp. 1245–1260, Sept. 2018.
- [19] A. Nedic, A. Olshevsky, and W. Shi, “Achieving geometric convergence for distributed optimization over time-varying graphs,” *SIAM Journal on Optimization*, vol. 27, no. 4, pp. 2597–2633, 2017.
- [20] K. Yuan and S. A. Alghunaim, “Removing data heterogeneity influence enhances network topology dependence of decentralized SGD,” *arXiv preprint:2105.08023*, 2021.
- [21] K. Huang and S. Pu, “Improving the transient times for distributed stochastic gradient methods,” *arXiv preprint:2105.04851*, 2021.
- [22] S. Pu and A. Nedić, “Distributed stochastic gradient tracking methods,” *Mathematical Programming*, vol. 187, no. 1, pp. 409–457, 2021.
- [23] Y. Lu and C. De Sa, “Optimal complexity in decentralized training,” in *International Conference on Machine Learning*, pp. 7111–7123, PMLR, 2021.
- [24] R. Xin, U. A. Khan, and S. Kar, “An improved convergence analysis for decentralized online stochastic non-convex optimization,” *IEEE Transactions on Signal Processing*, vol. 69, pp. 1842–1858, 2021.
- [25] B. T. Polyak, “Gradient methods for minimizing functionals,” *USSR Comput. Math. Math. Phys.*, vol. 3, no. 4, pp. 864–878, 1963.

- [26] H. Karimi, J. Nutini, and M. Schmidt, “Linear convergence of gradient and proximal-gradient methods under the Polyak-Lojasiewicz condition,” in *Joint European Conference on Machine Learning and Knowledge Discovery in Databases*, (Riva del Garda, Italy), pp. 795–811, Springer, 2016.
- [27] G. Scutari and Y. Sun, “Distributed nonconvex constrained optimization over time-varying digraphs,” *Mathematical Programming*, vol. 176, no. 1-2, pp. 497–544, 2019.
- [28] K. Scaman, F. Bach, S. Bubeck, Y. Lee, and L. Massoulié, “Optimal convergence rates for convex distributed optimization in networks,” *Journal of Machine Learning Research*, vol. 20, pp. 1–31, 2019.
- [29] S. A. Alghunaim, K. Yuan, and A. H. Sayed, “A linearly convergent proximal gradient algorithm for decentralized optimization,” in *Advances in Neural Information Processing Systems (NeurIPS)*, (Vancouver, Canada), pp. 2844–2854, Dec. 2019.
- [30] Y. Arjevani, J. Bruna, B. Can, M. Gurbuzbalaban, S. Jegelka, and H. Lin, “IDEAL: Inexact DEcentralized Accelerated Augmented Lagrangian method,” in *Advances in Neural Information Processing Systems (NeurIPS)*, vol. 33, pp. 20648–20659, 2020.
- [31] S. A. Alghunaim, E. K. Ryu, K. Yuan, and A. H. Sayed, “Decentralized proximal gradient algorithms with linear convergence rates,” *IEEE Transactions on Automatic Control*, vol. 66, pp. 2787–2794, June 2021.
- [32] J. Xu, Y. Tian, Y. Sun, and G. Scutari, “Distributed algorithms for composite optimization: Unified framework and convergence analysis,” *IEEE Transactions on Signal Processing*, vol. 69, pp. 3555–3570, June 2021.
- [33] P. Bianchi and J. Jakubowicz, “Convergence of a multi-agent projected stochastic gradient algorithm for non-convex optimization,” *IEEE Transactions on Automatic Control*, vol. 58, no. 2, pp. 391–405, 2013.
- [34] J. Chen and A. H. Sayed, “On the learning behavior of adaptive networks - Part I: Transient analysis,” *IEEE Transactions on Information Theory*, vol. 61, pp. 3487–3517, June 2015.
- [35] J. Chen and A. H. Sayed, “On the learning behavior of adaptive networks - Part II: Performance analysis,” *IEEE Transactions on Information Theory*, vol. 61, pp. 3518–3548, June 2015.
- [36] A. H. Sayed, “Adaptation, learning, and optimization over networks,” *Foundations and Trends in Machine Learning*, vol. 7, no. 4-5, pp. 311–801, 2014.
- [37] T. Tatarenko and B. Touri, “Non-convex distributed optimization,” *IEEE Transactions on Automatic Control*, vol. 62, no. 8, pp. 3744–3757, 2017.
- [38] B. Swenson, R. Murray, S. Kar, and H. V. Poor, “Distributed stochastic gradient descent: Nonconvexity, nonsmoothness, and convergence to local minima,” *arXiv preprint:2003.02818*, 2020.
- [39] S. Pu, A. Olshevsky, and I. C. Paschalidis, “A sharp estimate on the transient time of distributed stochastic gradient descent,” *arXiv preprint:1906.02702*, 2019.

- [40] S. Vlaski and A. H. Sayed, “Distributed learning in non-convex environments—part II: Polynomial escape from saddle-points,” *IEEE Transactions on Signal Processing*, vol. 69, pp. 1257–1270, 2021.
- [41] Z. Jiang, A. Balu, C. Hegde, and S. Sarkar, “Collaborative deep learning in fixed topology networks,” in *Advances in Neural Information Processing Systems*, (Long Beach, CA, USA), pp. 5904–5914, 2017.
- [42] A. Koloskova, S. Stich, and M. Jaggi, “Decentralized stochastic optimization and gossip algorithms with compressed communication,” in *International Conference on Machine Learning*, pp. 3478–3487, 2019.
- [43] A. Koloskova, N. Loizou, S. Boreiri, M. Jaggi, and S. Stich, “A unified theory of decentralized SGD with changing topology and local updates,” in *Proc. International Conference on Machine Learning*, vol. 119, pp. 5381–5393, PMLR, Jul. 2020.
- [44] J. Wang and G. Joshi, “Cooperative SGD: A unified framework for the design and analysis of local-update SGD algorithms,” *Journal of Machine Learning Research*, vol. 22, no. 213, pp. 1–50, 2021.
- [45] J. Zhang and K. You, “Decentralized stochastic gradient tracking for non-convex empirical risk minimization,” *arXiv preprint:1909.02712*, 2019.
- [46] S. Lu, X. Zhang, H. Sun, and M. Hong, “GNSD: A gradient-tracking based nonconvex stochastic algorithm for decentralized optimization,” in *IEEE Data Science Workshop (DSW)*, pp. 315–321, 2019.
- [47] X. Yi, S. Zhang, T. Yang, T. Chai, and K. H. Johansson, “A primal-dual SGD algorithm for distributed nonconvex optimization,” *arXiv preprint:2006.03474*, 2020.
- [48] R. Xin, S. Das, U. A. Khan, and S. Kar, “A stochastic proximal gradient framework for decentralized non-convex composite optimization: Topology-independent sample complexity and communication efficiency,” *arXiv preprint:2110.01594*, 2021.
- [49] P. Patarasuk and X. Yuan, “Bandwidth optimal all-reduce algorithms for clusters of workstations,” *Journal of Parallel and Distributed Computing*, vol. 69, no. 2, pp. 117–124, 2009.
- [50] Y. You, I. Gitman, and B. Ginsburg, “Large batch training of convolutional networks,” *arXiv preprint arXiv:1708.03888*, 2017.
- [51] M. Gurbuzbalaban, U. Simsekli, and L. Zhu, “The heavy-tail phenomenon in SGD,” in *International Conference on Machine Learning*, pp. 3964–3975, PMLR, 2021.
- [52] Y. Tang, J. Zhang, and N. Li, “Distributed zero-order algorithms for nonconvex multiagent optimization,” *IEEE Transactions on Control of Network Systems*, vol. 8, no. 1, pp. 269–281, 2020.
- [53] A. Antoniadis, I. Gijbels, and M. Nikolova, “Penalized likelihood regression for generalized linear models with non-quadratic penalties,” *Annals of the Institute of Statistical Mathematics*, vol. 63, no. 3, pp. 585–615, 2011.

- [54] R. A. Horn and C. R. Johnson, *Matrix Analysis*. Cambridge University Press, 2012.
- [55] Y. Nesterov, *Introductory Lectures on Convex Optimization: A Basic Course*, vol. 87. Springer, 2013.
- [56] B. Polyak, *Introduction to Optimization*. New York: Optimization Software, 1987.

Appendices

A. Relation of (4) to Existing Methods

In this section, we describe how different existing methods are related to algorithm (4). First, note that we can equivalently describe the updates in (4a) by noting that

$$\begin{aligned} \mathbf{x}^{k+1} - \mathbf{x}^k &= \mathbf{A}\mathbf{C}(\mathbf{x}^k - \mathbf{x}^{k-1}) - \alpha\mathbf{A}(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1})) - \mathbf{B}(\mathbf{y}^k - \mathbf{y}^{k-1}) \\ &\stackrel{(4b)}{=} \mathbf{A}\mathbf{C}(\mathbf{x}^k - \mathbf{x}^{k-1}) - \alpha\mathbf{A}(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1})) - \mathbf{B}^2\mathbf{x}^k. \end{aligned}$$

Hence,

$$\mathbf{x}^{k+1} = (\mathbf{I} - \mathbf{B}^2 + \mathbf{A}\mathbf{C})\mathbf{x}^k - \mathbf{A}\mathbf{C}\mathbf{x}^{k-1} - \alpha\mathbf{A}(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1})). \quad (49)$$

We next show that by choosing specific $\mathbf{A}, \mathbf{B}, \mathbf{C}$ as a function of the combination matrix \mathbf{W} , we can recover several important state-of-the-art methods.

A.1 Specific Instances

In the following, we do not assume that \mathbf{W} is symmetric unless otherwise stated.

- **Exact-Diffusion/D².** For Exact-Diffusion (ED) [12] (a.k.a. D² [15]), we assume \mathbf{W} to be symmetric and positive-semidefinite. If $\mathbf{A} = \mathbf{W}$, $\mathbf{B} = (\mathbf{I} - \mathbf{W})^{1/2}$, and $\mathbf{C} = \mathbf{I}$, we get ED/D² [12, 15]:

$$\mathbf{x}^{k+1} = \mathbf{W}\left(2\mathbf{x}^k - \mathbf{x}^{k-1} - \alpha(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1}))\right). \quad (50)$$

The above methods is also known as NIDS [13].

- **EXTRA.** For EXTRA [11], we also assume \mathbf{W} to be symmetric and positive-semidefinite. If we choose $\mathbf{A} = \mathbf{I}$, $\mathbf{B} = (\mathbf{I} - \mathbf{W})^{1/2}$, and $\mathbf{C} = \mathbf{W}$, we get EXTRA [11]:

$$\mathbf{x}^{k+1} = \mathbf{W}(2\mathbf{x}^k - \mathbf{x}^{k-1}) - \alpha(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1})). \quad (51)$$

- **ATC-GT.** Consider the adapt-then-combine GT method (ATC-GT):

$$\mathbf{x}^{k+1} = \mathbf{W}(\mathbf{x}^k - \alpha\mathbf{g}^k) \quad (52a)$$

$$\mathbf{g}^{k+1} = \bar{\mathbf{W}}(\mathbf{g}^k + \nabla\mathbf{f}(\mathbf{x}^{k+1}) - \nabla\mathbf{f}(\mathbf{x}^k)), \quad (52b)$$

where \mathbf{W} and $\bar{\mathbf{W}}$ are doubly-stochastic combination matrices such that $\bar{\mathbf{W}}\mathbf{W} = \mathbf{W}\bar{\mathbf{W}}$. Subtracting $\bar{\mathbf{W}}\mathbf{x}^k$ from both sides of the first equation, we have

$$\begin{aligned} \mathbf{x}^{k+1} - \bar{\mathbf{W}}\mathbf{x}^k &= \mathbf{W}\mathbf{x}^k - \bar{\mathbf{W}}\mathbf{W}\mathbf{x}^{k-1} - \alpha(\mathbf{W}\mathbf{g}^k - \bar{\mathbf{W}}\mathbf{W}\mathbf{g}^{k-1}) \\ &= \mathbf{W}\mathbf{x}^k - \bar{\mathbf{W}}\mathbf{W}\mathbf{x}^{k-1} - \alpha\mathbf{W}(\mathbf{g}^k - \bar{\mathbf{W}}\mathbf{g}^{k-1}). \end{aligned}$$

Using the second equation and rearranging, we get

$$\mathbf{x}^{k+1} = \mathbf{W}\mathbf{x}^k + \bar{\mathbf{W}}\mathbf{x}^k - \bar{\mathbf{W}}\mathbf{W}\mathbf{x}^{k-1} - \alpha\bar{\mathbf{W}}\mathbf{W}(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1})).$$

The above is the same as (49) when $\mathbf{A} = \bar{\mathbf{W}}\mathbf{W}$, $\mathbf{B}^2 = (\mathbf{I} - \bar{\mathbf{W}})(\mathbf{I} - \mathbf{W})$, and $\mathbf{C} = \mathbf{I}$. Note that when $\bar{\mathbf{W}} = \mathbf{W}$, then (52) becomes the GT method from [16].

- **Non-ATC-GT.** Consider the gradient-tracking algorithm

$$\mathbf{x}^{k+1} = \mathbf{W}\mathbf{x}^k - \alpha\mathbf{g}^k \quad (53a)$$

$$\mathbf{g}^{k+1} = \bar{\mathbf{W}}\mathbf{g}^k + \nabla\mathbf{f}(\mathbf{x}^{k+1}) - \nabla\mathbf{f}(\mathbf{x}^k), \quad (53b)$$

where \mathbf{W} and $\bar{\mathbf{W}}$ are doubly-stochastic combination matrices. Eliminating \mathbf{g}^k , we get:

$$\mathbf{x}^{k+1} = (\bar{\mathbf{W}} + \mathbf{W})\mathbf{x}^k - \bar{\mathbf{W}}\mathbf{W}\mathbf{x}^{k-1} - \alpha(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1})).$$

The above is exactly (49) when $\mathbf{A} = \mathbf{I}$, $\mathbf{B}^2 = (\mathbf{I} - \bar{\mathbf{W}})(\mathbf{I} - \mathbf{W})$, and $\mathbf{C} = \bar{\mathbf{W}}\mathbf{W}$. If $\bar{\mathbf{W}} = \mathbf{W}$ then (8) is the GT method studied in [18, 19].

- **Semi-ATC-GT.** Consider the following variation of GT methods:

$$\mathbf{x}^{k+1} = \mathbf{W}(\mathbf{x}^k - \alpha\mathbf{g}^k) \quad (54a)$$

$$\mathbf{g}^{k+1} = \bar{\mathbf{W}}\mathbf{g}^k + \nabla\mathbf{f}(\mathbf{x}^{k+1}) - \nabla\mathbf{f}(\mathbf{x}^k), \quad (54b)$$

where the ATC structure is used for the update of \mathbf{x}^k only. Following the same argument as before, we can show that (54) is equivalent to

$$\mathbf{x}^{k+1} = \mathbf{W}\mathbf{x}^k + \bar{\mathbf{W}}\mathbf{x}^k - \bar{\mathbf{W}}\mathbf{W}\mathbf{x}^{k-1} - \alpha\mathbf{W}(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1})),$$

The above is the same as (49) when $\mathbf{A} = \mathbf{W}$, $\mathbf{B}^2 = (\mathbf{I} - \bar{\mathbf{W}})(\mathbf{I} - \mathbf{W})$, and $\mathbf{C} = \bar{\mathbf{W}}$.

We can also consider the GT variant:

$$\mathbf{x}^{k+1} = \mathbf{W}\mathbf{x}^k - \alpha\mathbf{g}^k \quad (55a)$$

$$\mathbf{g}^{k+1} = \bar{\mathbf{W}}(\mathbf{g}^k + \nabla\mathbf{f}(\mathbf{x}^{k+1}) - \nabla\mathbf{f}(\mathbf{x}^k)). \quad (55b)$$

where the adapt-then-combine structure is used for the gradient-tracking variable \mathbf{g}^k only. Eliminating the gradient-tracking variable gives:

$$\mathbf{x}^{k+1} = \mathbf{W}\mathbf{x}^k + \bar{\mathbf{W}}\mathbf{x}^k - \bar{\mathbf{W}}\mathbf{W}\mathbf{x}^{k-1} - \alpha\bar{\mathbf{W}}(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1})),$$

which is exactly (49) when $\mathbf{A} = \bar{\mathbf{W}}$, $\mathbf{B}^2 = (\mathbf{I} - \bar{\mathbf{W}})(\mathbf{I} - \mathbf{W})$, and $\mathbf{C} = \mathbf{W}$. Note that when $\bar{\mathbf{W}} = \mathbf{W}$, then (54) and (55) become the GT variants from [17].

A.2 UDA

Here, we show that UDA (9) from [31] is equivalent to (4) if \mathbf{A} and \mathbf{B}^2 commute (*i.e.*, $\mathbf{A}\mathbf{B}^2 = \mathbf{B}^2\mathbf{A}$). We can eliminate the variable \mathbf{y}^k in (9) by noting that

$$\begin{aligned} \mathbf{z}^{k+1} - \mathbf{z}^k &= \mathbf{C}(\mathbf{x}^k - \mathbf{x}^{k-1}) - \alpha(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1})) - \mathbf{B}(\mathbf{y}^k - \mathbf{y}^{k-1}) \\ &= \mathbf{C}(\mathbf{x}^k - \mathbf{x}^{k-1}) - \alpha(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1})) - \mathbf{B}^2\mathbf{z}^k. \end{aligned}$$

Rearranging, we have

$$\mathbf{z}^{k+1} = (\mathbf{I} - \mathbf{B}^2)\mathbf{z}^k + \mathbf{C}(\mathbf{x}^k - \mathbf{x}^{k-1}) - \alpha(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1})).$$

We now multiply both sides by \mathbf{A} and use $\mathbf{A}\mathbf{B}^2 = \mathbf{B}^2\mathbf{A}$, to get

$$\mathbf{x}^{k+1} = (\mathbf{I} - \mathbf{B}^2 + \mathbf{A}\mathbf{C})\mathbf{x}^k - \mathbf{A}\mathbf{C}\mathbf{x}^{k-1} - \alpha\mathbf{A}(\nabla\mathbf{f}(\mathbf{x}^k) - \nabla\mathbf{f}(\mathbf{x}^{k-1})),$$

which is exactly (49).

B. Fundamental Factorization

B.1 Proof of Lemma 1

Recall that $\hat{\Lambda}_a = \text{diag}\{\lambda_{a,i}\}_{i=2}^n \otimes I_d$, $\hat{\Lambda}_b = \text{diag}\{\lambda_{b,i}\}_{i=2}^n \otimes I_d$, $\hat{\Lambda}_c = \text{diag}\{\lambda_{c,i}\}_{i=2}^n \otimes I_d$. Hence, the matrix \mathbf{G} defined in (31) can be rewritten as

$$\mathbf{G} = \begin{bmatrix} \text{diag}\{\lambda_{a,i}\lambda_{c,i} - \lambda_{b,i}^2\}_{i=2}^n & -\text{diag}\{\lambda_{b,i}\}_{i=2}^n \\ \text{diag}\{\lambda_{b,i}\}_{i=2}^n & I_{n-1} \end{bmatrix} \otimes I_d.$$

Utilizing the structure of \mathbf{G} , we can exchange the columns and rows of \mathbf{G} through some permutation matrix $\mathbf{P} \in \mathbb{R}^{2d(n-1) \times 2d(n-1)}$ such that

$$\mathbf{P}\mathbf{G}\mathbf{P}^\top = \text{blkdiag}\{G_i\}_{i=2}^n \otimes I_d,$$

where

$$G_i \triangleq \begin{bmatrix} \lambda_{a,i}\lambda_{c,i} - \lambda_{b,i}^2 & -\lambda_{b,i} \\ \lambda_{b,i} & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}. \quad (56)$$

Let us denote the eigenvalues of G_i by $\gamma_{1,i}$ and $\gamma_{2,i}$. If the eigenvalues of G_i are distinct ($\gamma_{1,i} \neq \gamma_{2,i}$), then there exists a 2×2 invertible V_i such that [54]:

$$G_i = V_i \Gamma_i V_i^{-1}, \quad \Gamma_i \triangleq \text{diag}\{\gamma_{1,i}, \gamma_{2,i}\}.$$

It follows that \mathbf{G} is similar to a diagonal matrix $\mathbf{G} = \hat{\mathbf{V}}\mathbf{\Gamma}\hat{\mathbf{V}}^{-1}$, where

$$\hat{\mathbf{V}} \triangleq \mathbf{P}^\top \mathbf{V}, \quad \mathbf{V} \triangleq \text{blkdiag}\{V_i\}_{i=2}^n \otimes I_d, \quad \mathbf{\Gamma} \triangleq \text{blkdiag}\{\Gamma_i\} \otimes I_d, \quad (57)$$

with $\|\mathbf{\Gamma}\| = \max_{i \in \{2, \dots, n\}} \|\Gamma_i\| = \max_{i \in \{2, \dots, n\}} \{|\gamma_{1,i}|, |\gamma_{2,i}|\} < 1$. Now, suppose that G_i has repeated eigenvalues $\gamma_{1,i} = \gamma_{2,i} = \gamma_i$. Then, using Jordan canonical form [54], there exists an invertible matrix T_i such that

$$G_i = T_i J_i T_i^{-1}, \quad J_i \triangleq \begin{bmatrix} \gamma_i & 1 \\ 0 & \gamma_i \end{bmatrix}.$$

If we let $E_i \triangleq \text{diag}\{1, \epsilon_i\}$ where $\epsilon_i > 0$ is an arbitrary constant, then we can rewrite G_i as

$$G_i = V_i \Gamma_i V_i^{-1}, \quad V_i \triangleq T_i E_i, \quad \Gamma_i \triangleq E_i^{-1} J_i E_i = \begin{bmatrix} \gamma_i & \epsilon_i \\ 0 & \gamma_i \end{bmatrix}.$$

It follows that $\mathbf{G} = \hat{\mathbf{V}}\mathbf{\Gamma}\hat{\mathbf{V}}^{-1}$ where $\hat{\mathbf{V}} = \mathbf{P}^\top \mathbf{V}$, $\mathbf{V} = \text{blkdiag}\{T_i E_i\}_{i=2}^n \otimes I_d$, and $\mathbf{\Gamma} = \text{blkdiag}\{\Gamma_i\} \otimes I_d$. Since the spectral radius of any matrix is upper bounded by any norm [54], it holds that

$$\|\Gamma_i\|^2 = \rho(\Gamma_i \Gamma_i^*) \leq \|\Gamma_i \Gamma_i^*\|_1 = |\gamma_i|^2 + \epsilon_i |\gamma_i| + \epsilon_i^2 \leq (|\gamma_i| + \epsilon_i)^2. \quad (58)$$

Here, $\|\cdot\|_1$ denote the maximum-absolute-column-sum matrix norm. Therefore, $\|\mathbf{\Gamma}\| < 1$ for any $\epsilon_i < 1 - |\gamma_i|$.

B.2 Special Cases of Lemma 1

In this section, we specify the results of Section B.1 to the instances discussed in Appendix A.1. First, we note that for any matrix

$$G = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbb{R}^{2 \times 2},$$

the eigenvalues are:

$$\gamma_{1,2} = \frac{(a+d) \pm \sqrt{(a+d)^2 - 4(ad-bc)}}{2}. \quad (59)$$

Moreover, if $\gamma_1 \neq \gamma_2$, then

$$G = V \text{diag}\{\gamma_1, \gamma_2\} V^{-1}, \quad V = [v_1 \quad v_2], \quad (60)$$

where $v_{1,2} = \frac{1}{r} \text{col}\{b, \gamma_{1,2} - a\}$ (or $v_{1,2} = \frac{1}{r} \text{col}\{\gamma_{1,2} - d, c\}$) if $b \neq 0$ (or $c \neq 0$) for any $r \neq 0$. Here, v_1 and v_2 are eigenvectors G corresponding to eigenvalues γ_1 and γ_2 , respectively.

B.2.1 Exact-Diffusion/D² and EXTRA

ED/D². We will consider the case for ED/D² (50) first. Recall from Appendix A.1 that for ED/D², we have $\mathbf{A} = \mathbf{W}$, $\mathbf{B} = (\mathbf{I} - \mathbf{W})^{1/2}$, and $\mathbf{C} = \mathbf{I}$. For this case, we have $\lambda_{a,i} = \lambda_i$, $\lambda_{c,i} = 1$, and $\lambda_{b,i} = \sqrt{1 - \lambda_i}$ where $\{\lambda_i\}$ denote the eigenvalues of W . Thus, the matrix (56) becomes:

$$G_i = \begin{bmatrix} 2\lambda_i - 1 & -\sqrt{1 - \lambda_i} \\ \sqrt{1 - \lambda_i} & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}.$$

Using (59), the eigenvalues of G_i ($i = 2, \dots, n$) are:

$$\gamma_{(1,2),i} = \lambda_i \pm \sqrt{\lambda_i^2 - \lambda_i}.$$

Note that $|\gamma_{(1,2),i}| < 1$ if $-\frac{1}{3} < \lambda_i < 1$. This implies that for the convergence of ED/D², we require $W > -\frac{1}{3}I$. It is sufficient for our purposes to assume that $W \geq 0$. Thus, we have two cases for the decomposition of G_i derived next.

- If $0 < \lambda_i < 1$, then the eigenvalues are complex and distinct:

$$\gamma_{(1,2),i} = \lambda_i \pm j\sqrt{\lambda_i - \lambda_i^2}, \quad |\gamma_{(1,2),i}| = \sqrt{\lambda_i} < 1,$$

where $j^2 = -1$. Using (60) with $r = \sqrt{1 - \lambda_i}$, we can factor G_i as $G_i = V_i \text{diag}\{\gamma_{1,i}, \gamma_{2,i}\} V_i^{-1}$ where

$$V_i = \begin{bmatrix} -1 & -1 \\ \sqrt{1 - \lambda_i} + j\sqrt{\lambda_i} & \sqrt{1 - \lambda_i} - j\sqrt{\lambda_i} \end{bmatrix}.$$

The inverse of V_i is

$$V_i^{-1} = \frac{1}{2j\sqrt{\lambda_i}} \begin{bmatrix} \sqrt{1 - \lambda_i} - j\sqrt{\lambda_i} & 1 \\ -\sqrt{1 - \lambda_i} - j\sqrt{\lambda_i} & -1 \end{bmatrix} = \frac{1}{2\sqrt{\lambda_i}} \begin{bmatrix} -\sqrt{\lambda_i} - j\sqrt{1 - \lambda_i} & -j \\ -\sqrt{\lambda_i} + j\sqrt{1 - \lambda_i} & j \end{bmatrix}.$$

Note that

$$V_i V_i^* = \begin{bmatrix} 2 & -2\sqrt{1-\lambda_i} \\ -2\sqrt{1-\lambda_i} & 2 \end{bmatrix}.$$

Since the spectral radius of matrix is upper bounded by any of its norm, it holds that $\|V_i\|^2 = \rho(V_i V_i^*) \leq \|V_i V_i^*\|_1 \leq 4$. Following a similar argument for V_i^{-1} , we have

$$\begin{aligned} (V_i^{-1})(V_i^{-1})^* &= \frac{1}{4\lambda_i} \begin{bmatrix} -\sqrt{\lambda_i} - j\sqrt{1-\lambda_i} & -j \\ -\sqrt{\lambda_i} + j\sqrt{1-\lambda_i} & j \end{bmatrix} \begin{bmatrix} -\sqrt{\lambda_i} + j\sqrt{1-\lambda_i} & -\sqrt{\lambda_i} - j\sqrt{1-\lambda_i} \\ j & -j \end{bmatrix} \\ &= \frac{1}{4\lambda_i} \begin{bmatrix} 2 & 2\lambda_i - 2 + 2j\sqrt{\lambda_i - \lambda_i^2} \\ 2\lambda_i - 2 + 2j\sqrt{\lambda_i - \lambda_i^2} & 2 \end{bmatrix}. \end{aligned}$$

Hence, $\|V_i^{-1}\|^2 = \rho(V_i^{-1}(V_i^{-1})^*) \leq \|V_i^{-1}(V_i^{-1})^*\|_1 \leq \frac{1}{\lambda_i}$.

- If $\lambda_i = 0$, then the eigenvalues of G_i are both zero $\gamma_{1,2} = 0$. Using Jordan canonical form, it holds that

$$G_i = \begin{bmatrix} -1 & -1 \\ 1 & 1 \end{bmatrix}, \quad T_i^{-1} G_i T_i = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad T_i = \begin{bmatrix} 1 & -\frac{1}{2} \\ -1 & -\frac{1}{2} \end{bmatrix}.$$

If we let $E = \text{diag}\{1, \epsilon\}$ for $0 < \epsilon < 1$ then

$$G_i = V_i \Gamma_i V_i^{-1}, \quad V_i = T_i E, \quad \Gamma_i = \begin{bmatrix} 0 & \epsilon \\ 0 & 0 \end{bmatrix}$$

Simple calculations show that $\|\Gamma_i\| = \epsilon < 1$ and

$$\|V_i\|^2 \leq \|T_i\|^2 \|E\|^2 = 2, \quad \|V_i^{-1}\|^2 \leq \|T_i^{-1}\|^2 \|E^{-1}\|^2 = \frac{2}{\epsilon^2}.$$

We can choose $\epsilon^2 = \lambda = \max_{i \in \{2, \dots, n\}} \lambda_i$.

Putting things together, we find

$$\|\hat{\mathbf{V}}\|^2 = \|\mathbf{P}^T \mathbf{V}\|^2 \leq 4, \quad \|\hat{\mathbf{V}}^{-1}\|^2 = \|\mathbf{V}^{-1} \mathbf{P}\|^2 \leq \frac{2}{\underline{\lambda}}, \quad \gamma = \|\mathbf{\Gamma}\| = \sqrt{\lambda}, \quad (61a)$$

$$\lambda_a = \|\hat{\mathbf{A}}_a\| = \lambda, \quad \lambda_b = \frac{1}{\|\hat{\mathbf{A}}_b^{-1}\|} = \sqrt{1-\lambda}, \quad (61b)$$

where $\lambda = \max_{i \in \{2, \dots, n\}} \lambda_i$ and $\underline{\lambda}$ is the minimum non-zero eigenvalue of W .

EXTRA. Observe that for EXTRA (51), the matrix G_i is identical to ED/D². Hence, the same bounds (61) hold for EXTRA except for λ_a , which is equal to one, *i.e.*, $\lambda_a = 1$, for EXTRA.

B.2.2 GT methods

ATC-GT. Consider the ATC-GT method (7) (or (52) with $\bar{\mathbf{W}} = \mathbf{W}$). In this case, we have $\lambda_{a,i} = \lambda_i^2$, $\lambda_{b,i} = 1 - \lambda_i$, and $\lambda_{c,i} = 1$. Thus, the matrix G_i ($i = 2, \dots, n$) given in (56) is

$$G_i = \begin{bmatrix} 2\lambda_i - 1 & -(1 - \lambda_i) \\ 1 - \lambda_i & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}.$$

Using (59), we find that the eigenvalues of G_i are identical $\gamma_{1,i} = \gamma_{2,i} = \lambda_i$. Clearly, the eigenvalues are strictly less than one by assumption since $0 \leq \lambda_i < 1$ ($i = 2, \dots, n$). If we let

$$T_i = \begin{bmatrix} -1 & 0 \\ 1 & \frac{1}{1-\lambda_i} \end{bmatrix}, \quad T_i^{-1} = \begin{bmatrix} -1 & 0 \\ 1 - \lambda_i & 1 - \lambda_i \end{bmatrix},$$

then, it holds that

$$G_i = T_i \begin{bmatrix} \lambda_i & 1 \\ 0 & \lambda_i \end{bmatrix} T_i^{-1}.$$

If we further let $E_i = \text{diag}\{1, \epsilon_i\}$ where $\epsilon_i > 0$ is an arbitrary constant, and define

$$V_i \triangleq T_i E_i = \begin{bmatrix} -1 & 0 \\ 1 & \frac{\epsilon_i}{1-\lambda_i} \end{bmatrix}, \quad V_i^{-1} = E_i^{-1} T_i^{-1} = \begin{bmatrix} -1 & 0 \\ \frac{1-\lambda_i}{\epsilon_i} & \frac{1-\lambda_i}{\epsilon_i} \end{bmatrix}.$$

Then, we have $G_i = V_i \Gamma_i V_i^{-1}$ where

$$\Gamma_i \triangleq E_i^{-1} \begin{bmatrix} \lambda_i & 1 \\ 0 & \lambda_i \end{bmatrix} E_i = \begin{bmatrix} \lambda_i & \epsilon_i \\ 0 & \lambda_i \end{bmatrix}.$$

Choosing $\epsilon = (1 - |\lambda_i|)/2$, it can be verified that $\|\mathbf{\Gamma}\| < 1$.

The above derivations are sufficient for our convergence analysis to cover GT methods with $I < W \leq I$. However, we shall assume in the following that $W \geq 0$ in order to get refined network dependent bounds for GT methods. Under this additional condition, we have $\lambda_i \geq 0$ and if we set $\epsilon = (1 - \lambda_i)/2$, it holds that

$$\Gamma_i = \begin{bmatrix} \lambda_i & \frac{1-\lambda_i}{2} \\ 0 & \lambda_i \end{bmatrix}, \quad V_i = \begin{bmatrix} -1 & 0 \\ 1 & \frac{1}{2} \end{bmatrix}, \quad V_i^{-1} = \begin{bmatrix} -1 & 0 \\ 2 & 2 \end{bmatrix}.$$

Hence, for GT methods with $W \geq 0$, we have

$$\|\hat{\mathbf{V}}\|^2 = \|\mathbf{P}^T \mathbf{V}\|^2 \leq 3, \quad \|\hat{\mathbf{V}}^{-1}\|^2 = \|\mathbf{V}^{-1} \mathbf{P}\|^2 \leq 9 \quad \gamma = \|\mathbf{\Gamma}\| \leq \frac{1+\lambda}{2}, \quad (62a)$$

$$\lambda_a = \|\hat{\mathbf{\Lambda}}_a\| = \lambda^2, \quad \lambda_b = \frac{1}{\|\hat{\mathbf{\Lambda}}_b^{-1}\|} = 1 - \lambda, \quad (62b)$$

where $\lambda = \max_{i \in \{2, \dots, n\}} \lambda_i$.

Other GT variants. Note that for the other variations of GT methods considered in Appendix A.1 with $\bar{\mathbf{W}} = \mathbf{W}$, the matrix G_i is identical to the previous case. Hence, the same bounds (62) holds for these other variations except for the value of λ_a , which is $\lambda_a = 1$ for the non-ATC-GT (53) and $\lambda_a = \lambda$ for the semi-ATC-GT variants (54) and (55).

C. Convergence Proof

In this section, we prove Theorems 1 and 2. We will first list some useful inequalities and facts that are used in our analysis.

Useful inequalities

- Under the L -smoothness condition (13) given in Assumption 2, the aggregate cost f is also L -smooth. It follows that [55]:

$$f(y) \leq f(z) + \langle \nabla f(z), y - z \rangle + \frac{L}{2} \|y - z\|^2, \quad \forall z, y \in \mathbb{R}^n. \quad (63)$$

- Recall from (19) that $\mathbf{w}^k = \nabla \mathbf{F}(\mathbf{x}^k, \boldsymbol{\xi}^k) - \nabla \mathbf{f}(\mathbf{x}^k)$ and $\bar{\mathbf{w}}^k = \frac{1}{n} \sum_{i=1}^n \nabla F_i(x_i^k, \xi_i^k) - \nabla f_i(x_i^k)$. Hence, under Assumption 3, we have

$$\mathbb{E}[\mathbf{w}^k | \mathcal{F}^k] = \mathbf{0}, \quad \mathbb{E}[\|\mathbf{w}^k\|^2 | \mathcal{F}^k] \leq n\sigma^2, \quad \mathbb{E}[\bar{\mathbf{w}}^k | \mathcal{F}^k] = \mathbf{0}, \quad \mathbb{E}[\|\bar{\mathbf{w}}^k\|^2 | \mathcal{F}^k] \leq \frac{\sigma^2}{n}. \quad (64)$$

- Using (35), it holds that

$$\|\mathbf{x}^k - \bar{\mathbf{x}}^k\|^2 = \|\hat{\mathbf{U}}^\top \mathbf{x}^k\|^2 = \|v \hat{\mathbf{V}} \hat{\mathbf{e}}^k\|^2 - \|\hat{\Lambda}_b^{-1} \hat{\mathbf{U}}^\top \mathbf{s}^k\|^2 \leq v^2 \|\hat{\mathbf{V}}\|^2 \|\hat{\mathbf{e}}^k\|^2. \quad (65)$$

Moreover, we have

$$\begin{aligned} \|\bar{\nabla \mathbf{f}}(\mathbf{x}^k) - \nabla f(\bar{x}^k)\|^2 &= \left\| \frac{1}{n} \sum_{i=1}^n (\nabla f_i(x_i^k) - \nabla f_i(\bar{x}^k)) \right\|^2 \\ &\leq \frac{1}{n} \sum_{i=1}^n \|\nabla f_i(x_i^k) - \nabla f_i(\bar{x}^k)\|^2 \\ &\stackrel{(13)}{\leq} \frac{L^2}{n} \|\mathbf{x}^k - \bar{\mathbf{x}}^k\|^2 \stackrel{(65)}{\leq} \frac{L^2 v^2 \|\hat{\mathbf{V}}\|^2}{n} \|\hat{\mathbf{e}}^k\|^2. \end{aligned} \quad (66)$$

- Let $v = \sqrt{n}v_2$ where $v_2 = \|\hat{\mathbf{V}}^{-1}\|$. Using the definition of $\hat{\mathbf{e}}^0$ in (34), we have

$$\|\hat{\mathbf{e}}^0\|^2 \leq \frac{v_2^2}{v^2} \left(\|\hat{\mathbf{U}}^\top \mathbf{x}^0\|^2 + \|\hat{\Lambda}_b^{-1} \hat{\mathbf{U}}^\top \mathbf{s}^0\|^2 \right) = \frac{1}{n} \|\hat{\mathbf{U}}^\top \mathbf{x}^0\|^2 + \frac{1}{n} \|\hat{\Lambda}_b^{-1} \hat{\mathbf{U}}^\top \mathbf{s}^0\|^2,$$

If the initialization is identical $\mathbf{x}^0 = \mathbf{1} \otimes x^0$ (for some $x^0 \in \mathbb{R}^d$), then $\|\hat{\mathbf{U}}^\top \mathbf{x}^0\| = 0$ and $\mathbf{x}^0 = \bar{\mathbf{x}}^0$; moreover, $\mathbf{z}^0 = \mathbf{0}$ since $\mathbf{y}^0 = \mathbf{0}$. Hence, we have

$$\begin{aligned} \|\hat{\mathbf{e}}^0\|^2 &\leq \frac{1}{\lambda_b^2 n} \|\hat{\mathbf{U}}^\top \mathbf{s}^0\|^2 \stackrel{(20c)}{=} \frac{\alpha^2}{\lambda_b^2 n} \|\hat{\mathbf{U}}^\top \mathbf{A} \nabla \mathbf{f}(\bar{\mathbf{x}}^0)\|^2 \\ &= \frac{\alpha^2}{\lambda_b^2 n} \|(\mathbf{A} - \frac{1}{n} \mathbf{1} \mathbf{1}^\top \otimes I_d) \nabla \mathbf{f}(\mathbf{x}^0)\|^2 = \frac{\alpha^2 \zeta_0^2}{\lambda_b^2}. \end{aligned} \quad (67)$$

where $\zeta_0 \triangleq \frac{1}{n} \|(\mathbf{A} - \frac{1}{n} \mathbf{1} \mathbf{1}^\top \otimes I_d) (\nabla \mathbf{f}(\mathbf{x}^0) - \mathbf{1} \otimes \nabla f(x^0))\|$.

- For any $0 \leq \eta < 1$, it holds that

$$\sum_{\ell=0}^{k-1} \eta^{k-1-\ell} \leq \sum_{\ell=0}^{\infty} \eta^\ell = \frac{1}{1-\eta}. \quad (68)$$

Moreover, for a non-negative sequence $\{w_\ell\}$ it holds that:

$$\sum_{k=1}^K \sum_{\ell=0}^{k-1} \eta^{k-1-\ell} w_\ell \leq \left(\sum_{\ell=0}^{\infty} \eta^\ell \right) \sum_{k=0}^{K-1} w_k = \frac{1}{1-\eta} \sum_{k=0}^{K-1} w_k. \quad (69)$$

C.1 Descent and Consensus Inequalities

In this section, we establish two inequalities given in Lemmas 3 and 4 that are essential to prove Theorems 1 and 2. We start with the following lemma regarding the iterates average.

Lemma 3 (Descent inequality). *Suppose that Assumptions 2 and 3 holds. If $\alpha \leq \frac{1}{2L}$, then for any $k \geq 0$, we have*

$$\mathbb{E} f(\bar{x}^{k+1}) \leq \mathbb{E} f(\bar{x}^k) - \frac{\alpha}{2} \mathbb{E} \|\nabla f(\bar{x}^k)\|^2 - \frac{\alpha}{4} \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 + \frac{\alpha L^2 v^2 v_1^2}{2n} \mathbb{E} \|\mathbf{e}^k\|^2 + \frac{\alpha^2 L \sigma^2}{2n}, \quad (70)$$

where $v_1 = \|\hat{\mathbf{V}}\|$ and v is an arbitrary strictly positive constant.

Proof. Recall that $\bar{x}^{k+1} = \bar{x}^k - \alpha \bar{\nabla} \mathbf{f}(\mathbf{x}^k) - \alpha \bar{\mathbf{w}}^k$ where $\bar{\nabla} \mathbf{f}(\mathbf{x}^k) = \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_i^k)$. Setting $y = \bar{x}^{k+1}$ and $z = \bar{x}^k$ in inequality (63), taking conditional expectation, and using (64), it holds that

$$\begin{aligned} \mathbb{E}[f(\bar{x}^{k+1}) \mid \mathcal{F}^k] &\leq f(\bar{x}^k) - \alpha \langle \nabla f(\bar{x}^k), \bar{\nabla} \mathbf{f}(\mathbf{x}^k) \rangle + \frac{\alpha^2 L}{2} \mathbb{E} [\|\bar{\nabla} \mathbf{f}(\mathbf{x}^k) + \bar{\mathbf{w}}^k(\mathbf{x}^k)\|^2 \mid \mathcal{F}^k] \\ &\leq f(\bar{x}^k) - \alpha \langle \nabla f(\bar{x}^k), \bar{\nabla} \mathbf{f}(\mathbf{x}^k) \rangle + \frac{\alpha^2 L}{2} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 + \frac{\alpha^2 L \sigma^2}{2n}. \end{aligned} \quad (71)$$

Since $2\langle a, b \rangle = \|a\|^2 + \|b\|^2 - \|a - b\|^2$, we have

$$-\langle \nabla f(\bar{x}^k), \bar{\nabla} \mathbf{f}(\mathbf{x}^k) \rangle = -\frac{1}{2} \|\nabla f(\bar{x}^k)\|^2 - \frac{1}{2} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 + \frac{1}{2} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k) - \nabla f(\bar{x}^k)\|^2. \quad (72)$$

Combining the last two equations, we get

$$\begin{aligned} \mathbb{E}[f(\bar{x}^{k+1}) \mid \mathcal{F}^k] &\leq f(\bar{x}^k) - \frac{\alpha}{2} \|\nabla f(\bar{x}^k)\|^2 - \frac{\alpha}{2} (1 - \alpha L) \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 \\ &\quad + \frac{\alpha}{2} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k) - \nabla f(\bar{x}^k)\|^2 + \frac{\alpha^2 L \sigma^2}{2n}. \end{aligned} \quad (73)$$

Substituting the bound (66) into inequality (73), letting $\alpha \leq \frac{1}{2L}$, and taking the expectation yields (70). \square

We next establish an inequality regarding the consensus iterates $\{\mathbf{e}^k\}$.

Lemma 4 (Consensus inequality). *Suppose that Assumptions 1-3 hold. Then, for all $k \geq 0$, we have*

$$\begin{aligned} \mathbb{E} \|\mathbf{e}^{k+1}\|^2 &\leq \left(\gamma + \frac{2\alpha^2 L^2 v_1^2 v_2^2 \lambda_a^2}{1 - \gamma} \right) \mathbb{E} \|\mathbf{e}^k\|^2 + \frac{2\alpha^4 L^2 v_2^2 \lambda_a^2 n}{\lambda_b^2 (1 - \gamma) v^2} \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 \\ &\quad + \frac{2\alpha^2 v_2^2 \lambda_a^2 n \sigma^2}{v^2} + \frac{2\alpha^4 L^2 v_2^2 \lambda_a^2 \sigma^2}{\lambda_b^2 (1 - \gamma) v^2}, \end{aligned} \quad (74)$$

where $v_1 \triangleq \|\hat{\mathbf{V}}\|$, $v_2 \triangleq \|\hat{\mathbf{V}}^{-1}\|$, $\lambda_a \triangleq \|\mathbf{\Lambda}_a\|$, $\gamma \triangleq \|\mathbf{\Gamma}\| < 1$, and $\lambda_b \triangleq \frac{1}{\|\mathbf{\Lambda}_b^{-1}\|}$.

Proof. From (33b), we have

$$\begin{aligned} \|\mathbf{e}^{k+1}\|^2 &= \left\| \mathbf{\Gamma} \mathbf{e}^k - \frac{\alpha}{v} \hat{\mathbf{V}}^{-1} \begin{bmatrix} \hat{\mathbf{\Lambda}}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)) \\ \hat{\mathbf{\Lambda}}_b^{-1} \hat{\mathbf{\Lambda}}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})) \end{bmatrix} - \frac{\alpha}{v} \hat{\mathbf{V}}^{-1} \begin{bmatrix} \hat{\mathbf{\Lambda}}_a \hat{\mathbf{U}}^\top \mathbf{w}^k \\ \mathbf{0} \end{bmatrix} \right\|^2 \\ &= \left\| \mathbf{\Gamma} \mathbf{e}^k - \frac{\alpha}{v} \hat{\mathbf{V}}^{-1} \begin{bmatrix} \hat{\mathbf{\Lambda}}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)) \\ \hat{\mathbf{\Lambda}}_b^{-1} \hat{\mathbf{\Lambda}}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})) \end{bmatrix} \right\|^2 + \frac{\alpha^2}{v^2} \|\hat{\mathbf{V}}^{-1} \hat{\mathbf{\Lambda}}_a \hat{\mathbf{U}}^\top \mathbf{w}^k\|^2 \end{aligned}$$

$$- \frac{2\alpha}{v} \langle \hat{\mathbf{V}}_l^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top \mathbf{w}^k, \mathbf{\Gamma} \hat{\mathbf{e}}^k \rangle + \frac{2\alpha^2}{v^2} \left\langle \hat{\mathbf{V}}_l^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top \mathbf{w}^k, \hat{\mathbf{V}}^{-1} \left[\begin{array}{c} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)) \\ \hat{\Lambda}_b^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})) \end{array} \right] \right\rangle,$$

where $\hat{\mathbf{V}}_l^{-1}$ is the left part of $\hat{\mathbf{V}}^{-1} = [\hat{\mathbf{V}}_l^{-1} \hat{\mathbf{V}}_r^{-1}]$. Using Cauchy–Schwarz inequality and $ab \leq \frac{a^2}{2} + \frac{b^2}{2}$ for any non-negative scalars a and b , the last term can be upper bounded by:

$$\begin{aligned} & \frac{2\alpha^2}{v^2} \left\langle \hat{\mathbf{V}}_l^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top \mathbf{w}^k, \hat{\mathbf{V}}^{-1} \left[\begin{array}{c} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)) \\ \hat{\Lambda}_b^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})) \end{array} \right] \right\rangle \\ & \leq \frac{\alpha^2}{v^2} \|\hat{\mathbf{V}}_l^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top \mathbf{w}^k\|^2 + \frac{\alpha^2}{v^2} \left\| \hat{\mathbf{V}}^{-1} \left[\begin{array}{c} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)) \\ \hat{\Lambda}_b^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})) \end{array} \right] \right\|^2. \end{aligned}$$

Combining the last two equations and taking conditional expectation, we have

$$\begin{aligned} \mathbb{E} [\|\hat{\mathbf{e}}^{k+1}\|^2 | \mathcal{F}^k] & \leq \mathbb{E} \left[\left\| \mathbf{\Gamma} \hat{\mathbf{e}}^k - \frac{\alpha}{v} \hat{\mathbf{V}}^{-1} \left[\begin{array}{c} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)) \\ \hat{\Lambda}_b^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})) \end{array} \right] \right\|^2 \middle| \mathcal{F}^k \right] \\ & \quad + \frac{2\alpha^2}{v^2} \mathbb{E} [\|\hat{\mathbf{V}}_l^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top \mathbf{w}^k\|^2 | \mathcal{F}^k] - \frac{2\alpha}{v} \mathbb{E} [\langle \hat{\mathbf{V}}_l^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top \mathbf{w}^k, \mathbf{\Gamma} \hat{\mathbf{e}}^k \rangle | \mathcal{F}^k] \\ & \quad + \frac{\alpha^2 v_2^2}{v^2} \mathbb{E} \left[\left\| \left[\begin{array}{c} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)) \\ \hat{\Lambda}_b^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})) \end{array} \right] \right\|^2 \middle| \mathcal{F}^k \right] \\ & \leq \mathbb{E} \left[\left\| \mathbf{\Gamma} \hat{\mathbf{e}}^k - \frac{\alpha}{v} \hat{\mathbf{V}}^{-1} \left[\begin{array}{c} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)) \\ \hat{\Lambda}_b^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})) \end{array} \right] \right\|^2 \middle| \mathcal{F}^k \right] + \frac{2\alpha^2 \|\hat{\mathbf{V}}_l^{-1}\|^2 \lambda_a^2 n \sigma^2}{v^2} \\ & \quad + \frac{\alpha^2 v_2^2}{v^2} \mathbb{E} \left[\left\| \left[\begin{array}{c} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)) \\ \hat{\Lambda}_b^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})) \end{array} \right] \right\|^2 \middle| \mathcal{F}^k \right], \end{aligned}$$

where $v_2 \triangleq \|\hat{\mathbf{V}}^{-1}\|$ and the last inequality holds from (64) and $\|\hat{\mathbf{U}}^\top\| \leq 1$. Taking the expectation and expanding the last term, we get

$$\begin{aligned} \mathbb{E} \|\hat{\mathbf{e}}^{k+1}\|^2 & \leq \mathbb{E} \left\| \mathbf{\Gamma} \hat{\mathbf{e}}^k - \frac{\alpha}{v} \hat{\mathbf{V}}^{-1} \left[\begin{array}{c} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)) \\ \hat{\Lambda}_b^{-1} \hat{\Lambda}_a \hat{\mathbf{U}}^\top (\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})) \end{array} \right] \right\|^2 + \frac{2\alpha^2 \|\hat{\mathbf{V}}_l^{-1}\|^2 \lambda_a^2 n \sigma^2}{v^2} \\ & \quad + \frac{\alpha^2 v_2^2 \lambda_a^2}{v^2} (\mathbb{E} \|\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)\|^2 + (\lambda_b^{-1})^2 \mathbb{E} \|\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})\|^2). \end{aligned} \quad (75)$$

where we used $\|\hat{\mathbf{U}}\| \leq 1$ and defined $\lambda_b^{-1} \triangleq \|\hat{\Lambda}_b^{-1}\|$. We can bound the first term on the right hand side by using the inequality $\|\mathbf{a} + \mathbf{b}\|^2 \leq \frac{1}{t} \|\mathbf{a}\|^2 + \frac{1}{1-t} \|\mathbf{b}\|^2$ for any $t \in (0, 1)$. Doing so with $t = \gamma = \|\mathbf{\Gamma}\| < 1$, we obtain

$$\begin{aligned} & \mathbb{E} \|\hat{\mathbf{e}}^{k+1}\|^2 \\ & \leq \gamma \mathbb{E} \|\hat{\mathbf{e}}^k\|^2 + \frac{\alpha^2 v_2^2 \lambda_a^2}{(1-\gamma)v^2} (\mathbb{E} \|\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)\|^2 + \frac{1}{\lambda_b^2} \mathbb{E} \|\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})\|^2) + \frac{2\alpha^2 \|\hat{\mathbf{V}}_l^{-1}\|^2 \lambda_a^2 n \sigma^2}{v^2} \\ & \quad + \frac{\alpha^2 v_2^2 \lambda_a^2}{v^2} (\mathbb{E} \|\nabla \mathbf{f}(\mathbf{x}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^k)\|^2 + \frac{1}{\lambda_b^2} \mathbb{E} \|\nabla \mathbf{f}(\bar{\mathbf{x}}^k) - \nabla \mathbf{f}(\bar{\mathbf{x}}^{k+1})\|^2) \\ & \leq \gamma \mathbb{E} \|\hat{\mathbf{e}}^k\|^2 + \frac{2\alpha^2 L^2 v_2^2 \lambda_a^2}{(1-\gamma)v^2} \mathbb{E} \|\mathbf{x}^k - \bar{\mathbf{x}}^k\|^2 + \frac{2\alpha^2 L^2 v_2^2 \lambda_a^2 n}{\lambda_b^2 (1-\gamma)v^2} \mathbb{E} \|\bar{\mathbf{x}}^k - \bar{\mathbf{x}}^{k+1}\|^2 + \frac{2\alpha^2 v_2^2 \lambda_a^2 n \sigma^2}{v^2} \\ & \stackrel{(65)}{\leq} \left(\gamma + \frac{2\alpha^2 L^2 v_2^2 \lambda_a^2}{(1-\gamma)v^2} \right) \mathbb{E} \|\hat{\mathbf{e}}^k\|^2 + \frac{2\alpha^2 L^2 v_2^2 \lambda_a^2 n}{\lambda_b^2 (1-\gamma)v^2} \mathbb{E} \|\bar{\mathbf{x}}^k - \bar{\mathbf{x}}^{k+1}\|^2 + \frac{2\alpha^2 v_2^2 \lambda_a^2 n \sigma^2}{v^2}, \end{aligned} \quad (76)$$

where last step holds due to smoothness condition (13), $\|\hat{\mathbf{V}}_l^{-1}\| \leq v_2$, and $1 \leq 1/(1-\gamma)$. The third term can be bounded as follows

$$\mathbb{E}[\|\bar{\mathbf{x}}^k - \bar{\mathbf{x}}^{k+1}\|^2 | \mathcal{F}^k] \stackrel{(33a)}{=} \mathbb{E}[\|\alpha \bar{\nabla} \mathbf{f}(\mathbf{x}^k) + \alpha \bar{\mathbf{w}}^k\|^2 | \mathcal{F}^k] \stackrel{(64)}{\leq} \alpha^2 \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 + \frac{\alpha^2 \sigma^2}{n}.$$

Taking the expectation and substituting the resulting bound into (76) yields inequality (74). \square

C.2 Proof of Theorem 1 (Non-convex Case)

Proof of Theorem 1. We start by deriving an ergodic bound on the consensus iterates $\{\hat{\mathbf{e}}^k\}$. Setting $v = \sqrt{n}v_2$ in (74), we get

$$\mathbb{E} \|\hat{\mathbf{e}}^{k+1}\|^2 \leq \left(\gamma + \frac{2\alpha^2 L^2 v_1^2 v_2^2 \lambda_a^2}{1-\gamma} \right) \mathbb{E} \|\hat{\mathbf{e}}^k\|^2 + \frac{2\alpha^4 L^2 \lambda_a^2}{\lambda_b^2 (1-\gamma)} \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 + 2\alpha^2 \lambda_a^2 \sigma^2 + \frac{2\alpha^4 L^2 \lambda_a^2 \sigma^2}{\lambda_b^2 (1-\gamma)n}. \quad (77)$$

If the step size α satisfies

$$\gamma + \frac{2\alpha^2 L^2 v_1^2 v_2^2 \lambda_a^2}{1-\gamma} \leq \frac{1+\gamma}{2} \Rightarrow \alpha \leq \frac{1-\gamma}{2Lv_1v_2\lambda_a}, \quad (78)$$

then the previous bound can be upper bounded by

$$\mathbb{E} \|\hat{\mathbf{e}}^{k+1}\|^2 \leq \frac{1+\gamma}{2} \mathbb{E} \|\hat{\mathbf{e}}^k\|^2 + \frac{2\alpha^4 L^2 \lambda_a^2}{\lambda_b^2 (1-\gamma)} \left(\mathbb{E} \|\nabla f(\bar{\mathbf{x}}^k)\|^2 + \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 \right) + 2\alpha^2 \lambda_a^2 \sigma^2 + \frac{2\alpha^4 L^2 \lambda_a^2 \sigma^2}{\lambda_b^2 (1-\gamma)n}. \quad (79)$$

Recursively applying the previous inequality, it holds (for any $k = 1, 2, \dots$) that

$$\begin{aligned} \mathbb{E} \|\hat{\mathbf{e}}^k\|^2 &\leq \left(\frac{1+\gamma}{2} \right)^k \|\hat{\mathbf{e}}^0\|^2 + \sum_{\ell=0}^{k-1} \left(\frac{1+\gamma}{2} \right)^{k-1-\ell} \left(2\alpha^2 \lambda_a^2 \sigma^2 + \frac{2\alpha^4 L^2 \lambda_a^2 \sigma^2}{\lambda_b^2 (1-\gamma)n} \right) \\ &\quad + \frac{2\alpha^4 L^2 \lambda_a^2}{\lambda_b^2 (1-\gamma)} \sum_{\ell=0}^{k-1} \left(\frac{1+\gamma}{2} \right)^{k-1-\ell} \left(\mathbb{E} \|\nabla f(\bar{\mathbf{x}}^\ell)\|^2 + \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^\ell)\|^2 \right) \\ &\leq \left(\frac{1+\gamma}{2} \right)^k \|\hat{\mathbf{e}}^0\|^2 + \frac{2}{1-\gamma} \left(2\alpha^2 \lambda_a^2 \sigma^2 + \frac{2\alpha^4 L^2 \lambda_a^2 \sigma^2}{\lambda_b^2 (1-\gamma)n} \right) \\ &\quad + \frac{2\alpha^4 L^2 \lambda_a^2}{\lambda_b^2 (1-\gamma)} \sum_{\ell=0}^{k-1} \left(\frac{1+\gamma}{2} \right)^{k-1-\ell} \left(\mathbb{E} \|\nabla f(\bar{\mathbf{x}}^\ell)\|^2 + \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^\ell)\|^2 \right), \end{aligned} \quad (80)$$

where in the last inequality we used (68). Taking the average of both sides of (80) over $k = 1, 2, \dots, K$, and using (68), we have

$$\begin{aligned} \frac{1}{K} \sum_{k=1}^K \mathbb{E} \|\hat{\mathbf{e}}^k\|^2 &\leq \frac{2\|\hat{\mathbf{e}}^0\|^2}{(1-\gamma)K} + \frac{4\alpha^2 \lambda_a^2 \sigma^2}{1-\gamma} + \frac{4\alpha^4 L^2 \lambda_a^2 \sigma^2}{\lambda_b^2 (1-\gamma)^2 n} \\ &\quad + \frac{2\alpha^4 L^2 \lambda_a^2}{\lambda_b^2 (1-\gamma)K} \sum_{k=1}^K \sum_{\ell=0}^{k-1} \left(\frac{1+\gamma}{2} \right)^{k-1-\ell} \left(\mathbb{E} \|\nabla f(\bar{\mathbf{x}}^\ell)\|^2 + \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^\ell)\|^2 \right) \\ &\leq \frac{2\|\hat{\mathbf{e}}^0\|^2}{(1-\gamma)K} + \frac{4\alpha^2 \lambda_a^2 \sigma^2}{1-\gamma} + \frac{4\alpha^4 L^2 \lambda_a^2 \sigma^2}{\lambda_b^2 (1-\gamma)^2 n} \end{aligned}$$

$$+ \frac{4\alpha^4 L^2 \lambda_a^2}{\lambda_b^2 (1-\gamma)^2 K} \sum_{k=0}^{K-1} \left(\mathbb{E} \|\nabla f(\bar{x}^k)\|^2 + \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 \right), \quad (81)$$

where in the last inequality we used the bound (69) on the last term. Adding $\frac{\|\hat{\mathbf{e}}^0\|^2}{(1-\gamma)K}$ to both sides of the previous inequality and using $\frac{\|\hat{\mathbf{e}}^0\|^2}{K} \leq \frac{\|\hat{\mathbf{e}}^0\|^2}{(1-\gamma)K}$, we get

$$\begin{aligned} \frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\hat{\mathbf{e}}^k\|^2 &\leq \frac{3\|\hat{\mathbf{e}}^0\|^2}{(1-\gamma)K} + \frac{4\alpha^2 \lambda_a^2 \sigma^2}{1-\gamma} + \frac{4\alpha^4 L^2 \lambda_a^2 \sigma^2}{\lambda_b^2 (1-\gamma)^2 n} \\ &\quad + \frac{4\alpha^4 L^2 \lambda_a^2}{\lambda_b^2 (1-\gamma)^2 K} \sum_{k=0}^{K-1} \left(\mathbb{E} \|\nabla f(\bar{x}^k)\|^2 + \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 \right). \end{aligned} \quad (82)$$

Inequality (82) will be used in the upcoming bound (84), which will be established next. Subtracting f^* from both sides of (70), letting $v = \sqrt{n}v_2$, using $1/2 \geq 1/4$, and rearranging gives

$$\mathbb{E} \|\nabla f(\bar{x}^k)\|^2 + \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 \leq \frac{4}{\alpha} \left(\mathbb{E} \tilde{f}(\bar{x}^k) - \mathbb{E} \tilde{f}(\bar{x}^{k+1}) \right) + 2L^2 v_1^2 v_2^2 \mathbb{E} \|\hat{\mathbf{e}}^k\|^2 + \frac{2\alpha L \sigma^2}{n}, \quad (83)$$

where $\tilde{f}(\bar{x}^k) \triangleq f(\bar{x}^k) - f^*$. Summing over $k = 0, \dots, K-1$, dividing by $K \geq 1$, and using $-\tilde{f}(\bar{x}^k) \leq 0$, it holds that

$$\frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\nabla f(\bar{x}^k)\|^2 + \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 \leq \frac{4\tilde{f}(\bar{x}^0)}{\alpha K} + \frac{2\alpha L \sigma^2}{n} + \frac{2L^2 v_1^2 v_2^2}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\hat{\mathbf{e}}^k\|^2. \quad (84)$$

Substituting inequality (82) into (84) and rearranging, we obtain

$$\begin{aligned} &\left(1 - \frac{8\alpha^4 L^4 v_1^2 v_2^2 \lambda_a^2}{\lambda_b^2 (1-\gamma)^2} \right) \frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\nabla f(\bar{x}^k)\|^2 + \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 \\ &\leq \frac{4\tilde{f}(\bar{x}^0)}{\alpha K} + \frac{2\alpha L \sigma^2}{n} + \frac{6L^2 v_1^2 v_2^2 \|\hat{\mathbf{e}}^0\|^2}{(1-\gamma)K} + \frac{8\alpha^2 L^2 v_1^2 v_2^2 \lambda_a^2 \sigma^2}{1-\gamma} + \frac{8\alpha^4 L^4 v_1^2 v_2^2 \lambda_a^2 \sigma^2}{\lambda_b^2 (1-\gamma)^2 n}. \end{aligned} \quad (85)$$

If we set

$$\frac{1}{2} \leq 1 - \frac{8\alpha^4 L^4 v_1^2 v_2^2 \lambda_a^2}{\lambda_b^2 (1-\gamma)^2} \implies \alpha \leq \frac{\sqrt{\lambda_b (1-\gamma)}}{2L\sqrt{v_1 v_2 \lambda_a}}, \quad (86)$$

then we find

$$\begin{aligned} &\frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\nabla f(\bar{x}^k)\|^2 + \mathbb{E} \|\bar{\nabla} \mathbf{f}(\mathbf{x}^k)\|^2 \\ &\leq \frac{8\tilde{f}(\bar{x}^0)}{\alpha K} + \frac{4\alpha L \sigma^2}{n} + \frac{12L^2 v_1^2 v_2^2 \|\hat{\mathbf{e}}^0\|^2}{(1-\gamma)K} + \frac{16\alpha^2 L^2 v_1^2 v_2^2 \lambda_a^2 \sigma^2}{1-\gamma} + \frac{16\alpha^4 L^4 v_1^2 v_2^2 \lambda_a^2 \sigma^2}{\lambda_b^2 (1-\gamma)^2 n}. \end{aligned} \quad (87)$$

Using (67) in the above inequality, we arrive at (36). The step size conditions in the theorem follows from $\alpha \leq \frac{1}{2L}$ (from Lemma 3) and the conditions (78) and (86). \square

C.3 Proof of Theorem 2 (PL Condition Case)

To prove Theorem 2, we need the following result.

Lemma 5. *Under Assumption 2, the following holds:*

$$\|\nabla f(x)\|^2 \leq 2L(f(x) - f^*), \quad \forall x \in \mathbb{R}^d \quad (88)$$

$$\frac{1}{n} \sum_{i=1}^n (f(x_i^k) - f^*) \leq 2(f(\bar{x}^k) - f^*) + \frac{Lv^2v_1^2}{n} \|\hat{\mathbf{e}}^k\|, \quad \forall k. \quad (89)$$

Proof. Inequality (88) is proven in [56]. We include the proof here for convenience. Since $f^* \leq f(z)$ for any $z \in \mathbb{R}^d$, we have

$$f^* \leq f\left(x - \frac{1}{L}\nabla f(x)\right) \leq f(x) - \frac{1}{2L}\|\nabla f(x)\|^2, \quad (90)$$

where the last inequality holds due to (63) with $y = x - \frac{1}{L}\nabla f(x)$ and $z = x$. Rearranging the above we arrive at (88). We now establish inequality (89). The argument adjust the proof [24, Lemma 13] to our case. Substituting $y = x_i^k$ and $z = \bar{x}^k$ in (63), we get

$$f(x_i^k) \leq f(\bar{x}^k) + \langle \nabla f(\bar{x}^k), x_i^k - \bar{x}^k \rangle + \frac{L}{2}\|x_i^k - \bar{x}^k\|^2.$$

Using Cauchy-Schwarz and Young's inequalities, we have

$$\begin{aligned} \langle \nabla f(\bar{x}^k), x_i^k - \bar{x}^k \rangle &\leq \|\nabla f(\bar{x}^k)\| \|x_i^k - \bar{x}^k\| \leq \frac{1}{2L}\|\nabla f(\bar{x}^k)\|^2 + \frac{L}{2}\|x_i^k - \bar{x}^k\|^2 \\ &\stackrel{(88)}{\leq} f(\bar{x}^k) - f^* + \frac{L}{2}\|x_i^k - \bar{x}^k\|^2. \end{aligned}$$

Combining the last two bounds, then subtracting f^* from both sides and averaging over i , we get

$$\frac{1}{n} \sum_{i=1}^n (f(x_i^k) - f^*) \leq 2(f(\bar{x}^k) - f^*) + \frac{L}{n}\|\mathbf{x}^k - \bar{\mathbf{x}}^k\|.$$

Using (65), we arrive at (89). \square

Proof of Theorem 2. Setting $v = \sqrt{nv_2}$ in (70), we get

$$\mathbb{E} f(\bar{x}^{k+1}) \leq \mathbb{E} f(\bar{x}^k) - \frac{\alpha}{2}\mathbb{E}\|\nabla f(\bar{x}^k)\|^2 - \frac{\alpha}{4}\mathbb{E}\|\overline{\nabla \mathbf{f}}(\mathbf{x}^k)\|^2 + \frac{\alpha L^2 v_1^2 v_2^2}{2}\mathbb{E}\|\hat{\mathbf{e}}^k\|^2 + \frac{\alpha^2 L \sigma^2}{2n}. \quad (91)$$

Subtracting f^* from both sides of (91), using the PL inequality (16) and the bound $-\frac{\alpha}{4}\mathbb{E}\|\overline{\nabla \mathbf{f}}(\mathbf{x}^k)\|^2 \leq 0$, we get

$$\mathbb{E} \tilde{f}(\bar{x}^{k+1}) \leq (1 - \mu\alpha)\mathbb{E} \tilde{f}(\bar{x}^k) + \frac{\alpha L^2 v_1^2 v_2^2}{2}\mathbb{E}\|\hat{\mathbf{e}}^k\|^2 + \frac{\alpha^2 L \sigma^2}{2n}, \quad (92)$$

where $\tilde{f}(\bar{x}^k) = f(\bar{x}^k) - f^*$. To establish our result, we will combine the above bound with a later bound given in (97), which we establish next. Letting $v = \sqrt{nv_2}$ in (74), gives

$$\mathbb{E}\|\hat{\mathbf{e}}^{k+1}\|^2 \leq \left(\gamma + \frac{2\alpha^2 L^2 v_1^2 v_2^2 \lambda_a^2}{1-\gamma}\right)\mathbb{E}\|\hat{\mathbf{e}}^k\|^2 + \frac{2\alpha^4 L^2 \lambda_a^2}{\lambda_b^2(1-\gamma)}\mathbb{E}\|\overline{\nabla \mathbf{f}}(\mathbf{x}^k)\|^2 + 2\alpha^2 \lambda_a^2 \sigma^2 + \frac{2\alpha^4 L^2 \lambda_a^2 \sigma^2}{\lambda_b^2(1-\gamma)n}. \quad (93)$$

Note that

$$\|\overline{\nabla \mathbf{f}}(\mathbf{x}^k)\|^2 = \|\overline{\nabla \mathbf{f}}(\mathbf{x}^k) - \nabla f(\bar{x}^k) + \nabla f(\bar{x}^k)\|^2 \leq 2\|\overline{\nabla \mathbf{f}}(\mathbf{x}^k) - \nabla f(\bar{x}^k)\|^2 + 2\|\nabla f(\bar{x}^k)\|^2$$

$$\stackrel{(88)}{\leq} 2\|\overline{\nabla \mathbf{f}}(\mathbf{x}^k) - \nabla f(\bar{x}^k)\|^2 + 4L\tilde{f}(\bar{x}^k) \stackrel{(66)}{\leq} 2v_1^2v_2^2L^2\|\hat{\mathbf{e}}^k\|^2 + 4L\tilde{f}(\bar{x}^k).$$

Substituting the above into (93), we get

$$\mathbb{E}\|\hat{\mathbf{e}}^{k+1}\|^2 \leq \frac{8\alpha^4L^3\lambda_a^2}{\lambda_b^2(1-\gamma)}\mathbb{E}\tilde{f}(\bar{x}^k) + \left(\gamma + \frac{2\alpha^2L^2v_1^2v_2^2\lambda_a^2}{1-\gamma} + \frac{4\alpha^4L^4v_1^2v_2^2\lambda_a^2}{\lambda_b^2(1-\gamma)}\right)\mathbb{E}\|\hat{\mathbf{e}}^k\|^2 + 2\alpha^2\lambda_a^2\sigma^2 + \frac{2\alpha^4L^2\lambda_a^2\sigma^2}{\lambda_b^2(1-\gamma)n}. \quad (94)$$

To simplify later expressions, we choose α such that

$$\frac{2\alpha^2L^2v_1^2v_2^2\lambda_a^2}{1-\gamma} + \frac{4\alpha^4L^4v_1^2v_2^2\lambda_a^2}{\lambda_b^2(1-\gamma)} \leq \frac{3\alpha^2L^2v_1^2v_2^2\lambda_a^2}{1-\gamma}, \quad (95a)$$

$$\gamma + \frac{3\alpha^2L^2v_1^2v_2^2\lambda_a^2}{1-\gamma} \leq \frac{1+\gamma}{2}, \quad (95b)$$

$$\frac{8\alpha^4L^3\lambda_a^2}{\lambda_b^2(1-\gamma)} \leq \frac{\alpha\mu\lambda_a^2}{Lv_1^2v_2^2}. \quad (95c)$$

The above inequalities are satisfied if

$$\alpha \leq \min \left\{ \frac{\lambda_b}{2L}, \frac{1-\gamma}{\sqrt{6}Lv_1v_2\lambda_a}, \left(\frac{\mu\lambda_b^2(1-\gamma)}{8L^4v_1^2v_2^2} \right)^{1/3} \right\}. \quad (96)$$

Under the previous conditions on α , inequality (94) is upper bounded by

$$\mathbb{E}\|\hat{\mathbf{e}}^{k+1}\|^2 \leq \frac{\alpha\mu\lambda_a^2}{Lv_1^2v_2^2}\mathbb{E}\tilde{f}(\bar{x}^k) + \left(\frac{1+\gamma}{2}\right)\mathbb{E}\|\hat{\mathbf{e}}^k\|^2 + 2\alpha^2\lambda_a^2\sigma^2 + \frac{2\alpha^4L^2\lambda_a^2\sigma^2}{\lambda_b^2(1-\gamma)n}. \quad (97)$$

Note that from (89) with $v = \sqrt{n}v_2$, we have

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[f(x_i^k) - f^*] \leq 2\mathbb{E}[f(\bar{x}^k) - f^*] + Lv_1^2v_2^2\mathbb{E}\|\hat{\mathbf{e}}^k\|. \quad (98)$$

Multiplying inequality (92) by 2 and inequality (97) by $Lv_1^2v_2^2$ and rewriting them in matrix form, we have

$$\left[\begin{array}{c} 2\mathbb{E}\tilde{f}(\bar{x}^{k+1}) \\ Lv_1^2v_2^2\mathbb{E}\|\hat{\mathbf{e}}^{k+1}\|^2 \end{array} \right] \leq \underbrace{\left[\begin{array}{cc} 1-\mu\alpha & \alpha L \\ \frac{\alpha\mu\lambda_a^2}{2} & \frac{1+\gamma}{2} \end{array} \right]}_{\triangleq H} \left[\begin{array}{c} 2\mathbb{E}\tilde{f}(\bar{x}^k) \\ Lv_1^2v_2^2\mathbb{E}\|\hat{\mathbf{e}}^k\|^2 \end{array} \right] + \underbrace{\left[\begin{array}{c} \frac{\alpha^2L\sigma^2}{n} \\ 2\alpha^2Lv_1^2v_2^2\lambda_a^2\sigma^2 + \frac{2\alpha^4L^3v_1^2v_2^2\lambda_a^2\sigma^2}{\lambda_b^2(1-\gamma)n} \end{array} \right]}_{\triangleq h}. \quad (99)$$

We will establish the convergence of $\frac{1}{n} \sum_{i=1}^n \mathbb{E}[f(x_i^k) - f^*]$ through the convergence of (99). Note that

$$\rho(H) \leq \|H\|_1 = \max \left\{ 1 - \mu\alpha + \frac{\alpha\mu\lambda_a^2}{2}, \frac{1+\gamma}{2} + \alpha L \right\} \leq 1 - \frac{\mu\alpha}{2}. \quad (100)$$

where the last inequality holds under the step size condition:

$$\alpha \leq \frac{1-\gamma}{3L} \leq \frac{1-\gamma}{2L+\mu}. \quad (101)$$

Hence, $\rho(H) < 1$ and iterating inequality (99), we get

$$\begin{aligned} \begin{bmatrix} 2\mathbb{E}\tilde{f}(\bar{x}^k) \\ Lv_1^2v_2^2\mathbb{E}\|\hat{\mathbf{e}}^k\|^2 \end{bmatrix} &\leq H^k \begin{bmatrix} 2\mathbb{E}\tilde{f}(\bar{x}^0) \\ Lv_1^2v_2^2\mathbb{E}\|\hat{\mathbf{e}}^0\|^2 \end{bmatrix} + \sum_{\ell=0}^{k-1} H^\ell h \\ &\leq H^k \begin{bmatrix} 2\mathbb{E}\tilde{f}(\bar{x}^0) \\ Lv_1^2v_2^2\mathbb{E}\|\hat{\mathbf{e}}^0\|^2 \end{bmatrix} + (I - H)^{-1}h. \end{aligned} \quad (102)$$

Taking the 1-norm, using (98), and using the submultiplicative and triangle inequality properties of the 1-norm, it holds that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[f(x_i^k) - f^*] \leq \|H^k\|_1 c_0 + \|(I - H)^{-1}h\|_1 \leq \|H\|_1^k c_0 + \|(I - H)^{-1}h\|_1. \quad (103)$$

where $c_0 = 2\mathbb{E}\tilde{f}(\bar{x}^0) + Lv_1^2v_2^2\mathbb{E}\|\hat{\mathbf{e}}^0\|^2$. We now bound the last term on the right. To do that, we compute the inverse:

$$(I - H)^{-1} = \begin{bmatrix} \mu\alpha & -\alpha L \\ -\frac{\alpha\mu\lambda_a^2}{2} & \frac{1-\gamma}{2} \end{bmatrix}^{-1} = \frac{1}{\det(I - H)} \begin{bmatrix} \frac{1-\gamma}{2} & \alpha L \\ \frac{\alpha\mu\lambda_a^2}{2} & \mu\alpha \end{bmatrix},$$

where $\det(I - H)$ is the determinant of $I - H$:

$$\det(I - H) = \frac{1}{2}(\alpha\mu(1 - \gamma) - \alpha^2\mu L\lambda_a^2) = \frac{\alpha\mu}{2}(1 - \gamma - \alpha L\lambda_a^2).$$

Hence,

$$\begin{aligned} (I - H)^{-1}h &= \frac{2}{\alpha\mu(1 - \gamma - \alpha L\lambda_a^2)} \begin{bmatrix} \frac{1-\gamma}{2} & \alpha L \\ \frac{\alpha^2\mu\lambda_a^2}{2} & \mu\alpha \end{bmatrix} \begin{bmatrix} \frac{\alpha^2 L\sigma^2}{n} \\ 2\alpha^2 Lv_1^2v_2^2\lambda_a^2\sigma^2 + \frac{2\alpha^4 L^3 v_1^2v_2^2\lambda_a^2\sigma^2}{\lambda_b^2(1-\gamma)n} \end{bmatrix} \\ &= \begin{bmatrix} \frac{\alpha L\sigma^2(1-\gamma)/n + 4\alpha^2 L^2 v_1^2v_2^2\lambda_a^2\sigma^2 + 4\alpha^4 L^4 v_1^2v_2^2\lambda_a^2\sigma^2/(\lambda_b^2(1-\gamma)n)}{\mu(1-\gamma-\alpha L\lambda_a^2)} \\ \frac{\alpha^3\mu L\lambda_a^2\sigma^2/n + 4\alpha^2\mu L v_1^2v_2^2\lambda_a^2\sigma^2 + 4\alpha^4\mu L^3 v_1^2v_2^2\lambda_a^2\sigma^2/(\lambda_b^2(1-\gamma)n)}{\mu(1-\gamma-\alpha L\lambda_a^2)} \end{bmatrix}. \end{aligned} \quad (104)$$

Note that since $\lambda_a \leq 1$; hence, under step size condition (101), we have

$$1 - \gamma - \alpha L\lambda_a^2 \geq (1 - \gamma)/2. \quad (105)$$

Combining the last two equations and using $L \geq \mu$, we get

$$\begin{aligned} \|(I - H)^{-1}h\|_1 &\leq \frac{\alpha L\sigma^2(1-\gamma)/n + 8\alpha^2 L^2 v_1^2v_2^2\lambda_a^2\sigma^2 + 8\alpha^4 L^4 v_1^2v_2^2\lambda_a^2\sigma^2/(\lambda_b^2(1-\gamma)n)}{\mu(1-\gamma)/2} + \frac{2\alpha^3 L\lambda_a^2\sigma^2}{n(1-\gamma)} \\ &= \frac{2\alpha L\sigma^2}{\mu n} + \frac{16\alpha^2 L^2 v_1^2v_2^2\lambda_a^2\sigma^2}{\mu(1-\gamma)} + \frac{16\alpha^4 L^4 v_1^2v_2^2\lambda_a^2\sigma^2}{\mu\lambda_b^2(1-\gamma)^2 n} + \frac{2\alpha^3 L\lambda_a^2\sigma^2}{n(1-\gamma)} \\ &= O\left(\frac{\alpha L\sigma^2}{\mu n}\right) + O\left(\frac{\alpha^2 L^2 v_1^2v_2^2\lambda_a^2\sigma^2}{\mu(1-\gamma)}\right) + O\left(\frac{\alpha^4 L^4 v_1^2v_2^2\lambda_a^2\sigma^2}{\mu\lambda_b^2(1-\gamma)^2 n}\right). \end{aligned}$$

Substituting the above into (103) and using (100), we obtain

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} \tilde{f}(x_i^k) \leq (1 - \frac{\alpha\mu}{2})^k c_0 + O\left(\frac{\alpha L \sigma^2}{\mu n}\right) + O\left(\frac{\alpha^2 L^2 v_1^2 v_2^2 \lambda_a^2 \sigma^2}{\mu(1-\gamma)}\right) + O\left(\frac{\alpha^4 L^4 v_1^2 v_2^2 \lambda_a^2 \sigma^2}{\mu \lambda_b^2 (1-\gamma)^2 n}\right). \quad (106)$$

Note that if $\mathbf{x}^0 = \mathbf{1} \otimes \mathbf{x}^0$, then

$$c_0 = 2\mathbb{E} \tilde{f}(\bar{x}^0) + L v_1^2 v_2^2 \mathbb{E} \|\hat{\mathbf{e}}^0\|^2 \stackrel{(67)}{\leq} 2\mathbb{E} \tilde{f}(\bar{x}^0) + \frac{\alpha^2 L v_1^2 v_2^2 \zeta_0^2}{\lambda_b^2}. \quad (107)$$

Combining the last two inequalities we get (41); moreover, combining the step size conditions (96) and (101), we get the step size condition in Theorem 2. □