

# Hurwitz groups, maximal reducible groups and maximal handlebody groups

Bruno P. Zimmermann

Università degli Studi di Trieste  
Dipartimento di Matematica e Geoscienze  
34127 Trieste, Italy

**Abstract.** A *Hurwitz group* is a finite group of orientation-preserving diffeomorphisms of maximal possible order  $84(g - 1)$  of a closed orientable surface of genus  $g > 1$ . A *maximal handlebody group* instead is a group of orientation-preserving diffeomorphisms of maximal possible order  $12(g - 1)$  of a 3-dimensional handlebody of genus  $g > 1$ . We consider the question of when a Hurwitz group acting on a surface of genus  $g$  contains a subgroup of maximal possible order  $12(g - 1)$  extending to a handlebody (or, more generally, a *maximal reducible group* extending to a product with handles), and show that such Hurwitz groups are closely related to the smallest Hurwitz group of order 168 acting on Klein's quartic of genus 3. We discuss simple groups of small order which are maximal handlebody groups and, more generally, maximal reducible groups. We discuss also the problem of which Hurwitz actions bound geometrically, and in particular whether Klein's quartic bounds geometrically: does there exist a compact hyperbolic 3-manifold with totally geodesic boundary isometric to Klein's quartic?

## 1. Introduction

All finite group-actions in the present paper will be orientation-preserving, all manifolds will be orientable.

### 1.1 *Hurwitz groups*

By the formula of Riemann-Hurwitz, the maximum order of a finite group  $H$  of diffeomorphisms of a closed surface  $\Sigma$  of genus  $g > 1$  is  $84(g - 1)$ ; such a group  $H$  is called a *Hurwitz groups of genus  $g$* . The quotient orbifold  $\Sigma/H$  of such a *Hurwitz action* is the 2-sphere with three branch points of orders 2, 3 and 7, and the Hurwitz groups are exactly the finite quotients of its orbifold fundamental group which is isomorphic to the hyperbolic triangle group of type  $(2,3,7)$ , with presentation

$$\langle x, y, z \mid x^2 = y^3 = z^7 = xyz = 1 \rangle .$$

This triangle group acts by isometries on the hyperbolic plane  $\mathbb{H}^2$ , there is an exact sequence

$$1 \rightarrow K \hookrightarrow (2, 3, 7) \rightarrow H \rightarrow 1,$$

and the factor group  $H \cong (2, 3, 7)/K$  acts by isometries on the hyperbolic surface  $\mathbb{H}^2/K$ .

## 1.2 Reducible and irreducible actions; products with handles

An action of a finite group  $G$  on a surface  $\Sigma$  is *reducible* if there is a nontrivial simple closed curve on  $\Sigma$  which is mapped either to itself or to a disjoint curve by every element of  $G$ , and otherwise *irreducible*. For a reducible action, one can equivariantly attach thickened 2-disks along the curve and its  $G$ -images to the boundary component  $\Sigma \times 1$  of the product  $\Sigma \times I$  to obtain a 3-manifold which is a product with handles of genus  $g$  to which the  $G$ -action extends. Iterating the construction one finishes with a product with handles  $\mathcal{P}$  with a  $G$ -action such that the action of each stabilizer of a boundary component of  $\mathcal{P}$  arising from  $\Sigma \times 1$  is irreducible.

In general, a *product with handles*  $\mathcal{P}$  is defined as follows. Considering the product  $\Sigma \times I$  of a closed surface  $\Sigma$  of genus  $g$  with the unit interval, one attaches 2-handles  $D^2 \times I$  (where  $D^2$  denotes the 2-disk) along the boundary parts  $S^1 \times I$  to a finite collection of disjoint simple closed curves on the surface  $\Sigma \times 1$ , then one closes eventually created 2-sphere boundary components by attaching 3-balls to obtain a 3-manifold  $\mathcal{P}$  whose boundary consists of the surface  $\partial_0\mathcal{P} = \Sigma \times 0$  of genus  $g$  (the *outer boundary* of  $\mathcal{P}$ ) and a collection of surfaces  $\partial_1\mathcal{P}$  (the *inner boundary* which may be empty). We call  $\mathcal{P}$  a *product with handles of genus  $g$*  (see section 4 for an explicit example).

A  $G$ -action on a surface  $\Sigma$  is irreducible if and only if the quotient orbifold  $\Sigma/G$  is the 2-sphere with exactly three branch points. In particular, all Hurwitz actions are irreducible. By the formula of Riemann-Hurwitz, the maximal possible order of a reducible action of  $G$  on a surface of genus  $g > 1$  is  $12(g - 1)$ , and in this case the quotient orbifold  $\Sigma/G$  is the 2-sphere with four branch points of orders 2, 2, 2 and 3 whose orbifold fundamental group is isomorphic to the hyperbolic quadrangle group of type (2,2,2,3), with presentation

$$\langle x_1, x_2, x_3, x_4 \mid x_1^2 = x_2^2 = x_3^2 = x_4^3 = x_1x_2x_3x_4 = 1 \rangle .$$

As before, there is an exact sequence

$$1 \rightarrow K \hookrightarrow (2, 2, 2, 3) \rightarrow G \rightarrow 1,$$

and the factor group  $G \cong (2, 2, 2, 3)/K$  acts by isometries on the hyperbolic surface  $\mathbb{H}^2/K$ . Let  $n$  denote the order of the image of the element  $x_1x_2$  in the finite group  $G$ ; by adding the relation  $(x_1x_2)^n = 1$  to the above presentation, we obtain a presentation of the free product with amalgamation

$$G_n = \mathbb{D}_n *_{\mathbb{Z}_n} (2, 3, n)$$

where  $(2, 3, n)$  denotes the triangle group of type  $(2, 3, n)$  and  $\mathbb{D}_n = (2, 2, n)$  the dihedral group of order  $2n$ . So the finite groups  $G$  admitting a reducible action of maximal

possible order  $12(g-1)$  on a surface of genus  $g > 1$  are exactly the finite factor groups with torsionfree kernel of the quadrangle group  $(2,2,2,3)$ , or equivalently of one of the groups  $G_n$ ,  $n \geq 2$ ; we call such a group a *maximal reducible group*, and specifically a  $G_n$ -group. So the maximal reducible groups are exactly the  $G_n$ -groups, for some  $n \geq 2$ .

A quotient of a product with handles by a finite group action is an *orbifold product with handles*; such an orbifold can be uniformized by a function group, i.e. by a Kleinian groups with an invariant component of the regular set. An approach to orbifold product with handles and the construction of the uniformizing Kleinian groups is given in [R], using the language of fiite graphs of groups.

### 1.3 Completely reducible actions, maximal handlebody groups and maximal bounded surface groups

An action of a finite group  $G$  on a surface  $\Sigma_g$  is *completely reducible* if it extends to a handlebody  $V_g$  with  $\partial V_g = \Sigma_g$ . By the equivariant Dehn Lemma/loop theorem [MY], a completely reducible action is reducible, and the maximum order of a completely reducible action of a finite group  $G$  acting on a 3-dimensional handlebody  $V_g$  of genus  $g > 1$  is again  $12(g-1)$  (as first observed in [Z1], prior to the appearance of the equivariant Dehn lemma/loop theorem, see also [MMZ, Theorem 7.2]); such a group  $G$  is called a *maximal handlebody group of genus  $g$* . By [Z2] (see also [Z5], [MMZ] for the following), the maximal handlebody groups  $G$  are exactly the finite quotients with torsionfree kernel of one of the four groups

$$G_2 = \mathbb{D}_2 *_{\mathbb{Z}_2} \mathbb{D}_3, \quad G_3 = \mathbb{D}_3 *_{\mathbb{Z}_3} \mathbb{A}_4, \quad G_4 = \mathbb{D}_4 *_{\mathbb{Z}_4} \mathbb{S}_4, \quad G_5 = \mathbb{D}_5 *_{\mathbb{Z}_5} \mathbb{A}_5$$

where  $\mathbb{A}_4 = (2, 3, 3)$  denotes the tetrahedral group of order 12,  $\mathbb{S}_4 = (2, 3, 4)$  the octahedral group of order 24 and  $\mathbb{A}_5 = (2, 3, 5)$  the dodecahedral group of order 60; together with the dihedral groups  $\mathbb{D}_n = (2, 2, n)$  of order  $2n$  these are exactly the finite or spherical triangle groups and, together with the cyclic groups, the finite subgroups of  $\text{SO}(3)$ .

The quotient orbifold  $V_g/G$  of a maximal handlebody group  $G$  of genus  $g$  by a finite group action is a *handlebody orbifold* whose orbifold fundamental group is one of the four groups  $G_2, G_3, G_4$  or  $G_5$ ; the underlying topological space is the 3-disk, its boundary  $\partial V_g/G$  is the 2-sphere with four branch points of orders 2, 2, 2 and 3. The handlebody orbifolds are uniformized by virtual Schottky groups (i.e., virtually free function groups with a single component of the regular set), cf. [MMZ], [Z5].

An upper bound for the maximal order of a finite group of homeomorphisms of a compact surface with nonempty boundary (orientable or not) of algebraic genus  $g$  (the rank of its free fundamental group) is also  $12(g-1)$ . This can be seen by taking the product of the surface with a closed interval (twisted if the surface is nonorientable); the result is an orientable handlebody of genus  $g$  to which the finite group action extends orientation-preservingly, so the upper bound for handlebodies applies, and such groups of order

$12(g - 1)$  are called *maximal bounded surface groups*. The maximal bounded surface groups are exactly the finite quotients with torsionfree kernel of the group

$$G_2 = \mathbb{D}_2 *_{\mathbb{Z}_2} \mathbb{D}_3$$

(cf. [MZ]). An interesting example of a maximal handlebody group which is not a maximal bounded surface group is the second Mathieu group  $M_{12}$ .

#### 1.4 Hurwitz actions versus maximal reducible actions

The smallest Hurwitz group is the linear fractional group  $\mathrm{PSL}_2(7)$  of order 168, acting on Klein's quartic  $\mathcal{Q}_3$  of genus 3. The smallest maximal handlebody group is the dihedral group  $\mathbb{D}_6$  of genus 2, the second smallest the octahedral or symmetric group  $\mathbb{S}_4$  of genus 3 (see [Z5, Proposition 7]). The maximal handlebody group  $\mathbb{S}_4$  is a subgroup of index 7 in the Hurwitz group  $\mathrm{PSL}_2(7)$ , and its preimage under the associated surjection  $\psi : (2, 3, 7) \rightarrow \mathrm{PSL}_2(7)$  is the quadrangle group  $(2,2,2,3)$  (by an easy application of the formula of Riemann Hurwitz). Since the restriction  $\psi : (2, 2, 2, 3) \rightarrow \mathbb{S}_4$  factors through one of the three groups  $G_2, G_3$  or  $G_4$  (in fact through all of them), the action of  $\mathbb{S}_4$  on Klein's quartic  $\mathcal{Q}_3$  (unique up to conjugation) extends to a handlebody  $V_3$  of genus 3. Our first main result is the following.

**Theorem 1.** *Let  $H$  be Hurwitz group of genus  $g > 1$  acting on a surface  $\Sigma = \Sigma_g$ .*

*i) If the action of  $H$  on  $\Sigma$  has a reducible subgroup  $G$  of maximal possible order  $12(g - 1)$  then  $\Sigma$  is a finite regular covering of Klein's quartic and the actions of  $H$  and  $G$  are obtained by lifting the actions of  $\mathrm{PSL}_2(7)$  and its subgroup  $\mathbb{S}_4$  to  $\Sigma$ ; in particular,  $H$  surjects onto  $\mathrm{PSL}_2(7)$ . There are infinitely many Hurwitz actions with a maximal reducible subgroup.*

*ii) If  $H$  is simple and not isomorphic to a subgroup of the alternating group  $\mathbb{A}_{24}$  then the action of every proper subgroup of  $H$  on  $\Sigma$  is reducible.*

A lift of a subgroup  $\mathbb{S}_4$  of  $\mathrm{PSL}_2(7)$  to a finite regular covering of Klein's quartic is always reducible but in general not completely reducible; at present we don't know if one can obtain infinitely many completely reducible actions in this way. Considering just the types of groups and not the actions, the following holds.

**Theorem 2.** *A Hurwitz group which does not surject onto  $\mathrm{PSL}_2(7)$  is a maximal reducible group; in particular,  $\mathrm{PSL}_2(7)$  is the only simple Hurwitz group which is not a maximal reducible group.*

### 1.5 Bounding and nonbounding Hurwitz actions

The action of the subgroup  $\mathbb{S}_4$  of  $\mathrm{PSL}_2(7)$  on Klein's quartic  $\mathcal{Q}_3$  extends to a handlebody  $V_3$  of genus 3; on the other hand, the action of  $\mathrm{PSL}_2(7)$  on  $\mathcal{Q}_3$  *does not bound*, i.e. does not extend to any compact 3-manifold  $M$  with exactly one boundary component  $\partial M = \mathcal{Q}_3$  ([Z4, Corollary 1]): the quotient orbifold  $\mathcal{Q}_3/\mathrm{PSL}_2(7)$  is the 2-sphere with three branch points of orders 2, 3 and 7 which does not occur as the unique boundary component of a compact 3-orbifold (since a singular axis starting in the boundary point of order 7 can end only in a dihedral point of type  $\mathbb{D}_7$  but  $\mathrm{PSL}_2(7)$  has no dihedral subgroup  $\mathbb{D}_7$ ).

The second smallest Hurwitz group is the linear fractional group  $\mathrm{PSL}_2(8)$  of order 504 and genus 7, and it is shown in [Z4, Corollary 2] that the action by isometries of  $\mathrm{PSL}_2(8)$  on the unique hyperbolic surface  $\mathcal{Q}_7$  *bounds geometrically*, i.e. extends to a group of isometries of a compact hyperbolic 3-manifold  $M$  with totally geodesic boundary  $\partial M = \mathcal{Q}_7$ . In particular, forgetting about the group action, the hyperbolic Hurwitz surface  $\mathcal{Q}_7$  bounds geometrically. This leads naturally to the following:

*Question* (see [Z6]): Does Klein's quartic bound geometrically? (Since Klein's quartic is the most symmetric surface of genus 3, the suspected answer should be affirmative.)

Up to conjugation by homeomorphisms, there is a unique action of  $\mathbb{S}_4$  on a surface of genus 3; we will show in section 3 that this  $\mathbb{S}_4$ -action bounds geometrically, for some hyperbolic structure on a surface of genus 3. But, in contrast to the Hurwitz actions, the isometric action of  $\mathbb{S}_4$  on a hyperbolic surface of genus 3 does not determine the hyperbolic structure of the surface (since the deformation space of a hyperbolic quadrangle of type (2,2,2,3) has positive dimension whereas a hyperbolic triangle of type (2,3,7) is unique up to isometry). So there are uncountably many hyperbolic structures on a surface of genus 3 such that  $\mathbb{S}_4$  acts by isometries; only countably many of these can bound geometrically, and at present we do not know if one can obtain Klein's quartic by a construction as in section 3.

The Hurwitz groups of linear fractional type are the groups  $\mathrm{PSL}_2(q)$ , for  $q = 7$  or  $q = p$  prime with  $p \equiv \pm 1 \pmod{7}$  (each with a unique Hurwitz action), and for  $q = p^3$  with  $p \equiv \pm 2, \pm 3 \pmod{7}$  (each with three different Hurwitz actions). For a proof of the following theorem see [GZ, Corollary 3.14] (and [RZ, Proposition 2] for a proof that  $\mathrm{PSL}_2(27)$  does not bound).

**Theorem 3.** *The only non-bounding Hurwitz actions of linear fractional type  $\mathrm{PSL}_2(q)$ , with  $q \leq 1000$ , are the unique Hurwitz actions of the groups  $\mathrm{PSL}_2(7)$  and  $\mathrm{PSL}_2(27)$ ; in fact all other Hurwitz actions with  $q \leq 1000$  bound geometrically, and in particular the corresponding Hurwitz surfaces bound geometrically.*

So there is some analogy between the Hurwitz actions of  $\mathrm{PSL}_2(7)$  and  $\mathrm{PSL}_2(27)$ : both do not bound, and in particular, similar as for  $\mathrm{PSL}_2(7)$ , it remains open whether the

Hurwitz surface of the action of  $\mathrm{PSL}_2(27)$  bounds geometrically. Supported by some strong evidence from [GZ], we state the following:

*Conjecture.* With the exceptions of  $q = 7$  and  $27$ , all Hurwitz actions of linear fractional type  $\mathrm{PSL}_2(q)$  bound geometrically.

### 1.6 Maximal reducible groups and maximal handlebody groups

Examples of small maximal handlebody groups which are not maximal bounded surface groups are given in [Z3], and it is shown in [CZ] by extensive computational methods that 161 and 3781 are the two smallest genera for which there exists a maximal handlebody group but not a maximal bounded surface group (answering a question in [MZ]). In the next theorem, we confront maximal reducible groups and maximal handlebody groups; the first claim is proved in [PZ, Corollary 3.3].

**Theorem 4.** *i) The linear fractional group  $\mathrm{PSL}_2(q)$  is a maximal handlebody group exactly for all  $q$  different from 7, 9, 11 and  $3^{2m+1}$ . The group  $\mathrm{PSL}_2(27)$  is the smallest simple group which is a maximal reducible group but not a maximal handlebody group. There are infinitely many maximal reducible groups which are not maximal handlebody groups.*

*ii) The smallest simple groups which are maximal reducible of order less than  $10^6$  and not of linear fractional type  $\mathrm{PSL}_2(q)$  are the second Mathieu group  $M_{12}$ , the Janko group  $J_1$ , the alternating group  $\mathbb{A}_9$  and the Hall-Janko group  $J_2$ .*

### 1.7 $G_7$ -groups and bounding Hurwitz groups

An interesting class of maximal reducible groups closely related to Hurwitz groups are the  $G_7$ -groups,

$$G_7 = \mathbb{D}_7 *_{\mathbb{Z}_7} (2, 3, 7).$$

A surjection  $\varphi : G_7 \rightarrow G$  with torsionfree kernel to a finite group  $G$  determines a  $G$ -action of maximal possible order  $12(g-1)$  on a product with handles  $\mathcal{P}$  of genus  $g$  (obtained as the regular orbifold covering of the orbifold product with handles with fundamental group  $G_7$  associated to the kernel of  $\varphi$ ). Let  $n$  be the index of the image  $\varphi(2, 3, 7)$  of the triangle group  $(2, 3, 7)$  in  $G$ .

If  $n = 1$ , the inner boundary  $\partial_1 \mathcal{P}$  of  $\mathcal{P}$  consists of a single surface of genus  $g'$ , with  $84(g'-1) = 12(g-1)$ , on which  $G$  restricts to a Hurwitz action. For arbitrary  $n$ ,  $\partial_1 \mathcal{P}$  consists of  $n$  copies of a Hurwitz surface, and the stabilizer in  $G$  of each of these surfaces acts as a Hurwitz group (isomorphic to  $\varphi(2, 3, 7)$ ); see section 4 for an explicit example.

We say that a Hurwitz group is *bounding* if admits a bounding Hurwitz action; similarly, a  $G_7$ -group is *bounding* if it admits a bounding maximal reducible action.

**Theorem 5.** *i) All bounding Hurwitz groups are  $G_7$ -groups. The groups  $\mathrm{PSL}_2(7)$  and  $\mathrm{PSL}_2(27)$  are the smallest non-bounding Hurwitz groups, and  $\mathrm{PSL}_2(27)$  is the smallest simple non-bounding  $G_7$ -group.*

*ii) With the exception of  $\mathrm{PSL}_2(7)$ , all Hurwitz groups of linear fractional type are  $G_7$ -groups. The simple  $G_7$ -groups of order less than  $10^6$  which are not Hurwitz groups of linear fractional type are the groups  $\mathrm{PSL}_2(49)$ ,  $J_1$ ,  $\mathbb{A}_9$  and  $J_2$ . The groups  $\mathrm{PSL}_2(49)$  and  $\mathbb{A}_9$  are the smallest simple  $G_7$ -groups which are not Hurwitz groups.*

The simple Hurwitz groups of to order less than  $10^6$  are of linear fractional type, and in addition the Janko groups  $J_1$  and  $J_2$  (cf. [C]).

Let  $H$  be a Hurwitz group with a Hurwitz action associated to a surjection  $\varphi : (2, 3, 7) \rightarrow H$ . We denote by  $[2,3,7]$  the *extended triangle group* generated by the reflections in the sides of a hyperbolic triangle with angles  $2\pi/2$ ,  $2\pi/3$  and  $2\pi/7$  (containing the triangle group  $(2,3,7)$  as the subgroup of index 2 of orientation-preserving elements). The following follows from [Z4, Theorem 3b].

**Theorem 6.** *Let  $H$  be a Hurwitz group with a Hurwitz action associated to a surjection  $(2, 3, 7) \rightarrow H$ . If the surjection extends to a surjection  $[2, 3, 7] \rightarrow H$  then the Hurwitz action of  $H$  bounds geometrically.*

The Hurwitz groups  $H$  as in Theorem 6 are exactly the "non-orientable Hurwitz groups" (also called  $H^*$ -groups), of maximal possible order  $84(g-2)$  acting on a non-orientable surface of genus  $g > 2$  (see [C2]). An interesting example of such a group is the first Janko group  $J_1$ : some Hurwitz action of  $J_1$  bounds geometrically; this remains open for the Hall-Janko group  $J_2$  which is also a Hurwitz group.

## 2. Proofs of Theorems 1 and 2

In the notation of Theorem 1, the reducible group  $G$  of order  $12(g-1)$  has index 7 in the Hurwitz group  $H$  of order  $84(g-1)$ . There is a surjection

$$\pi : (2, 3, 7) \rightarrow H$$

whose kernel is the universal covering group of the surface  $\Sigma_g$ . Since the triangle group is a perfect group (has trivial abelianization), also the Hurwitz group  $H$  is perfect. The preimage  $\pi^{-1}(G)$  of  $G$  in the triangle group  $(2,3,7)$  is a quadrangle group  $(2,2,2,3)$  (by an easy application of the formula of Riemann-Hurwitz, this is the only signature of a subgroup of index 7 in the triangle group  $(2,3,7)$ ).

The permutation representation of  $H$  by right multiplication on the seven right cosets of  $G$  in  $H$  defines a nontrivial homomorphism

$$\phi : H \rightarrow \mathbb{A}_7$$

of  $G$  to the alternating group  $\mathbb{A}_7$  of degree 7; the image of  $\phi$  is perfect and also a Hurwitz group (the kernel of the composition  $\phi \circ \pi : (2, 3, 7) \rightarrow \mathbb{A}_7$  is torsionfree since any normal subgroup of  $(2, 3, 7)$  containing a torsion element is equal to  $(2, 3, 7)$ ). The perfect subgroups of the alternating group  $\mathbb{A}_7$  are the simple groups  $\mathbb{A}_5$ ,  $\mathbb{A}_6$ ,  $\mathbb{A}_7$  and  $\mathrm{PSL}_2(7)$  (see [CCN]). Since the only Hurwitz group among these groups is  $\mathrm{PSL}_2(7)$  (by [C] or by a direct computation using [GAP]), we have surjections

$$\phi : H \rightarrow \mathrm{PSL}_2(7), \quad \phi \circ \pi : (2, 3, 7) \rightarrow \mathrm{PSL}_2(7),$$

and the second one gives the unique Hurwitz action of  $\mathrm{PSL}_2(7)$  on Klein's quartic  $\mathcal{Q}_3$ .

The image  $\phi(G)$  is a subgroup of index 1 or 7 in  $\mathrm{PSL}_2(7)$ . Since  $\mathrm{PSL}_2(7)$  is not a maximal handlebody group (not a surjective image of the quadrangle group  $(2, 2, 2, 3)$  as can easily be checked),  $\phi(G)$  has index 7 in  $\mathrm{PSL}_2(7)$ . Up to conjugation in  $\mathrm{PGL}_2(7)$ , the only subgroup of index 7 in  $\mathrm{PSL}_2(7)$  is the symmetric group  $\mathbb{S}_4$  of degree 4 (see [CCN]).

The kernel  $U$  of the surjection  $\phi : H \rightarrow \mathrm{PSL}_2(7)$  acts freely on  $\Sigma_g$  (since its preimage  $\pi^{-1}(U)$  in the triangle group  $(2, 3, 7)$  is torsionfree), and the actions of  $H$  and  $G$  on  $\Sigma_g$  project to the actions of  $\mathrm{PSL}_2(7) \cong H/U$  and  $\mathbb{S}_4 \cong G/U$  on Klein's quartic  $\mathcal{Q}_3 = \Sigma_g/U$ .

In order to obtain infinitely many examples of such Hurwitz groups, one considers finite-index characteristic subgroups of the fundamental group of Klein's quartic and lifts the actions of  $\mathrm{PSL}_2(7)$  and its subgroup  $\mathbb{S}_4$  to the corresponding finite regular coverings.

This concludes the proof of part i) of Theorem 1. For the proof of part ii), assume that  $G$  is a proper subgroup of  $H$ , of index  $n > 1$ , which acts irreducibly on  $\Sigma$ . Then the extension of  $\pi_1(\Sigma)$  associated to  $G$  is a triangle group of index  $n$  in the triangle group  $(2, 3, 7)$ . The triangle group of maximal possible index in  $(2, 3, 7)$  is the triangle group  $(7, 7, 7)$  of index 24, which implies  $n \leq 24$ . The action by right multiplication of  $G$  on the right cosets of  $H$  in  $G$  defines a nontrivial, hence injective homomorphism of the simple group  $H$  to the alternating group  $\mathbb{A}_n$ , in particular  $H$  is isomorphic to a subgroup of  $\mathbb{A}_{24}$ .

For the *proof of Theorem 2*, suppose that the Hurwitz group  $H$  does not surject onto  $\mathrm{PSL}_2(7)$ . There is a surjection of the triangle group  $(2, 3, 7)$  onto  $H$ , and the image of the quadrangle subgroup  $(2, 2, 2, 3)$  of index 7 in  $(2, 3, 7)$  has index 1 or 7 in  $H$ . If it has index 7 then, by the proof of Theorem 1,  $H$  surjects onto  $\mathrm{PSL}_2(7)$ , otherwise the quadrangle group  $(2, 2, 2, 3)$  surjects onto  $H$  and  $H$  is a maximal reducible group.

Example. The second largest Hurwitz group is the linear fractional group  $\mathrm{PSL}_2(8)$  of order 504 and genus 7, and also the product  $\mathrm{PSL}_2(7) \times \mathrm{PSL}_2(8)$  is a Hurwitz group acting on a unique surface of genus 1009 (by a direct computation using [GAP], see also [C]). In fact, up to isomorphism there is a unique surjection  $\pi : (2, 3, 7) \rightarrow \mathrm{PSL}_2(7) \times \mathrm{PSL}_2(8)$ .

The preimage  $\pi^{-1}(\mathbb{S}_4 \times \mathrm{PSL}_2(8))$  has index 7 in the triangle group  $(2,3,7)$  and is a quadrangle group  $(2,2,2,3)$ . We checked by [GAP] that, for the corresponding surjection  $\pi : (2, 2, 2, 3) \rightarrow \mathbb{S}_4 \times \mathrm{PSL}_2(8)$ , the image of the element  $x_1x_2 \in (2, 2, 2, 3)$  has order 36 and 36 is the minimal  $n$  such that the surjection factors through  $G_n$ . Hence the action of  $\mathbb{S}_4 \times \mathrm{PSL}_2(8)$  on the surface of genus 1009 is reducible but not completely reducible. On the other hand, there are various surjections  $(2, 2, 2, 3) \rightarrow \mathbb{S}_4 \times \mathrm{PSL}_2(8)$  which factor through  $G_2$  or  $G_4$ , so  $\mathbb{S}_4 \times \mathrm{PSL}_2(8)$  is a maximal handlebody group.

### 3. Geometrically bounding actions

Up to isomorphisms there is a unique surjection of the quadrangle group  $(2,2,2,3)$  onto  $\mathbb{S}_4$ ; equivalently, up to conjugation by homeomorphisms there is a unique action of  $\mathbb{S}_4$  on a surface  $\Sigma_3$  of genus 3, with quotient orbifold of type  $(2,2,2,3)$  (cf. [B]). There are infinitely many hyperbolic structures on the quotient orbifold  $\mathbb{H}^2/(2, 2, 2, 3)$ ; lifting such a hyperbolic structure to the surface  $\Sigma_3$ , the group  $\mathbb{S}_4$  acts by isometries.

**Proposition.** *The unique  $\mathbb{S}_4$ -action on a surface of genus 3 (unique up to conjugation by homeomorphisms) bounds geometrically, for some hyperbolic structure on the surface such that  $\mathbb{S}_4$  acts by isometries.*

*Proof.* By Andreev's theorem ([T, chapter 13], [V, p.111]) there exists a unique finite hyperbolic polyhedron  $\mathcal{C} = \mathcal{C}(3, 3, 3)$  in hyperbolic 3-space  $\mathbb{H}^3$  which has the combinatorial structure of a cube as follows. The four edges of a face  $F$  of  $\mathcal{C}$  have right dihedral angles (hence this face is orthogonal to the four adjacent faces), the four edges connecting  $F$  to the opposite face  $F'$  have three right dihedral angles and one dihedral angle  $\pi/3$ , and the face  $F'$  has two consecutive edges with right dihedral angles and two with dihedral angles  $\pi/3$  such that the three edges of  $\mathcal{C}$  with angles  $\pi/3$  form a segment (note that the polyhedron  $\mathcal{C}$  has no incompressible Euclidean 2-suborbifolds). Let  $C$  denote the orientation-preserving subgroup of index two in the group  $C$  generated by the reflections in the five faces of  $\mathcal{C}$  different from  $F$  (so  $C$  is generated by rotations around the eight edges of  $\mathcal{C}$  not in the boundary of  $F$ ); it is easy to check then that there exists a surjection from  $C$  to  $\mathbb{S}_4$ , with torsionfree kernel.

Let  $\mathcal{D}$  denote the double of  $\mathcal{C}$  along its face  $F$  (again of the combinatorial type of a cube, with five dihedral angles  $\pi/3$ ), and by  $D$  the orientation-preserving subgroup of index 2 in the Coxeter group generated by the reflections in the faces of  $\mathcal{D}$ . There exists a surjection  $\pi : D \rightarrow \mathbb{S}_4$  with torsionfree kernel  $K$ , and hence  $D/K \cong \mathbb{S}_4$  acts on the closed hyperbolic 3-manifold  $M = \mathbb{H}^3/K$ . The face  $F$  of  $\mathcal{C}$  lies in a hyperbolic plane on which a quadrangle subgroup  $(2,2,2,3)$  of  $D$  acts, and the restriction of  $\pi$  to this group  $(2,2,2,3)$  is also surjective. The preimage of the face  $F$  of  $\mathcal{C}$  is a closed totally geodesic surface  $\Sigma_3$  of genus 3 in  $M$  which is invariant under the action of  $\mathbb{S}_4$ . The surface  $\Sigma_3$  separates  $M$  into two isometric hyperbolic 3-manifolds, each with an isometric action

of  $\mathbb{S}_4$ , which have the hyperbolic surface  $\Sigma_3$  as their common totally geodesic boundary, and hence the isometric  $\mathbb{S}_4$ -action on the hyperbolic surface  $\Sigma_3$  bounds geometrically.

We don't know if the hyperbolic surface  $\Sigma_3$  is isometric to Klein's quartic (for example, one might try to compute the lengths of the edges of the face  $F$  of  $Q$  and compare with the edges of the corresponding quadrangle for a subgroup  $(2,2,2,3)$  of the triangle group  $(2,3,7)$ ). Note that the construction can be modified by choosing finite cubical polyhedra  $\mathcal{C}$  with other small dihedral angles for the face  $F'$ , obtaining other hyperbolic surfaces  $\Sigma_3$ .

#### 4. Proofs of Theorems 4 and 5

As noted before, the first claim of Theorem 4 is proved in [PZ, Corollary 3.3] (note that the case  $q = 11$  was overlooked in [PZ].) In particular,  $\mathrm{PSL}_2(27)$  is not a maximal handlebody group (which can be checked easily also by a direct computation). The unique Hurwitz action of  $\mathrm{PSL}_2(27)$  is associated to a surjection  $(2, 3, 7) \rightarrow \mathrm{PSL}_2(27)$ ; since  $\mathrm{PSL}_2(27)$  has subgroups  $\mathbb{D}_7$ , this extends to a surjection  $G_7 = \mathbb{D}_7 *_{\mathbb{Z}_7} (2, 3, 7) \rightarrow \mathrm{PSL}_2(27)$ , hence  $\mathrm{PSL}_2(27)$  is a  $G_7$ -group.

The group  $G = \mathrm{PSL}_2(27)$  is a Hurwitz group in a unique way: up to equivalence there is a unique surjection  $(2, 3, 7) \rightarrow G$ ; let  $K$  denote the kernel of this surjection which is a surface group of genus  $g' = 118$ . By abelianization and reduction of coordinates mod  $p$ , for some prime  $p$ , we get a surjection  $K \rightarrow (\mathbb{Z}_p)^{2g'}$  whose kernel is a characteristic subgroup  $\tilde{K}$  of  $K$ , in particular normal in  $(2, 3, 7)$ , so we have an exact sequence

$$1 \rightarrow \tilde{K} \hookrightarrow (2, 3, 7) \rightarrow (\mathbb{Z}_p)^{2g'} \rtimes \mathrm{PSL}_2(27) \rightarrow 1.$$

Since  $G = \mathrm{PSL}_2(27)$  has dihedral subgroups  $\mathbb{D}_7$  (of order 14), the surjection  $(2, 3, 7) \rightarrow (\mathbb{Z}_p)^{2g'} \rtimes G$  extends to a surjection with torsionfree kernel of  $G_7 = \mathbb{D}_7 *_{\mathbb{Z}_7} (2, 3, 7)$  to  $(\mathbb{Z}_p)^{2g'} \rtimes G$ , hence  $(\mathbb{Z}_p)^{2g'} \rtimes G$  is a maximal reducible  $G_7$ -group of Hurwitz type. On the other hand,  $(\mathbb{Z}_p)^{2g'} \rtimes G$  is not a maximal handlebody group for  $p > |G|$  since a surjection with torsionfree kernel  $G_n \rightarrow (\mathbb{Z}_p)^{2g'} \rtimes G$  would induce a surjection  $G_n \rightarrow G$  with torsionfree kernel which does not exist for  $2 \leq n \leq 5$  (since  $G = \mathrm{PSL}_2(27)$  is not a maximal handlebody group).

This completes the proof of Theorem 4 i). Theorem 4 ii) and Theorem 5 ii) can easily be checked by case-by-case computations (using e.g. [GAP]), going over the list of the simple groups of small order, their maximal subgroups and the orders and conjugacy classes of elements (see [CNN]). The first statement of Theorem 5 ii) follows from the well-known classification of the subgroups of the linear fractional groups  $\mathrm{PSL}_2(q)$ ; in particular, with the exception of  $\mathrm{PSL}_2(7)$ , an element of order 7 in a linear fractional Hurwitz group  $H$  lies in a dihedral subgroup  $\mathbb{D}_7$  of order 14, hence  $H$  is a  $G_7$ -group. Theorem 5 i) follows from the argument given in section 1.5 for the case of  $\mathrm{PSL}_2(7)$ .

Example. The Hurwitz group  $\mathrm{PSL}_2(7)$  of genus 3 is not a  $G_7$ -group since an element of order 7 does not lie in a dihedral subgroup  $\mathbb{D}_7$ . We finish with an example of a maximal reducible action of  $\mathrm{PGL}_2(7)$  on a surface  $\Sigma$  of genus 29 and the induced action on a product with handles  $\mathcal{P}$  with outer boundary  $\partial_0\mathcal{P} = \Sigma$  whose inner boundary  $\partial_1\mathcal{P}$  consists of two copies of Klein's quartic with the Hurwitz action of  $\mathrm{PSL}_2(7)$ .

The kernel of the composition of surjections

$$(2, 2, 2, 3) \rightarrow \mathbb{D}_7 *_{\mathbb{Z}_7} (2, 3, 7) \rightarrow \mathrm{PGL}_2(7)$$

is a surface group  $\pi_1(\Sigma)$  of genus  $g = 29$  (note that, in contrast to  $\mathrm{PSL}_2(7)$ ,  $\mathrm{PGL}_2(7)$  has dihedral subgroups  $\mathbb{D}_7$ ). Let  $\bar{\mathcal{P}}$  denote the orbifold product with handles with orbifold fundamental group  $\mathbb{D}_7 *_{\mathbb{Z}_7} (2, 3, 7)$  (cf. [R]), and let  $\mathcal{P}$  be the product with handles of genus  $g = 29$  which is the covering of  $\bar{\mathcal{P}}$  associated to the kernel of the surjection  $\mathbb{D}_7 *_{\mathbb{Z}_7} (2, 3, 7) \rightarrow \mathrm{PGL}_2(7)$ . The outer boundary  $\partial_0\mathcal{P}$  is the surface  $\Sigma$ ; since the triangle group  $(2, 3, 7)$  maps onto the subgroup  $\mathrm{PSL}_2(7)$  of index 2 in  $\mathrm{PGL}_2(7)$ , the inner boundary  $\partial_1\mathcal{P}$  consists of two copies of Klein's quartic of genus 3 with the Hurwitz action of  $\mathrm{PSL}_2(7)$ .

## References

- [B] S.A. Broughton, *Classifying finite group actions on surfaces of low genus*. J. Pure Appl. Alg. 69 (1990), 233-270
- [C1] M. Conder, *The genus of compact Riemann surfaces with maximal automorphism group*. J. Algebra 108 (1987), 204-247
- [C2] M. Conder, *Maximal automorphism groups of symmetric Riemann surfaces with small genus*. J. Algebra 114 (1988), 16-28
- [CZ] M.D.E. Conder, B. Zimmermann, *Maximal bordered surface groups versus maximal handlebody groups*. Contemporary Math. 629 (2014), 99-105
- [CCN] J.H. Conway, R.T. Curtis, S.P. Norton, R.A. Parker, R.A. Wilson, *Atlas of Finite Groups*. Oxford University Press 1985
- [GAP] The GAP Group, GAP - Groups, Algorithms, and Programming, Computational algebra system available under <http://www.gap-system.org>
- [GZ] M. Gradolato, B. Zimmermann, *Extending finite group actions on surfaces to hyperbolic 3-manifolds*. Math. Proc. Cambridge Phil. Soc. 117 (1995), 137-151
- [MMZ] D. McCullough, A. Miller, B. Zimmermann, *Group actions on handlebodies*. Proc. Lond. Math. Soc. 59 (1989), 373-415
- [MY] W.H. Meeks, S.T. Yau, *The equivariant Dehn's lemma and loop theorem*. Comment. Math. Helv. 56 (1981), 225-239
- [MZ] A. Miller, B. Zimmermann, *Large groups of symmetries of handlebodies*. Proc. Amer. Math. Soc. 106 (1989), 829-838

- [PZ] L. Paoluzzi, B. Zimmermann *On finite quotients of the Picard group and related hyperbolic tetrahedral groups*. Rend. Istit. Mat. Univ. Trieste Suppl. 1 Vol. 32, 257-288 (2001)
- [R] M. Reni, *A graph-theoretical approach to Kleinian groups*. Proc. London Math. Soc. 67 (1993), 200-224
- [RZ] M. Reni, B. Zimmermann, *Extending finite group actions from surfaces to handlebodies*. Proc. Amer. Math. Soc. 124 (1996), 2877-2887
- [T] W. Thurston, *The geometry and topology of 3-manifolds*. Lecture Notes, Dept. of Math., Princeton Univ., Princeton 1977
- [V] E.B. Vinberg (Ed.), *Geometry II*. Encyclopaedia of Math. Sciences vol. 29, Springer-Verlag 1993
- [Z1] B. Zimmermann, *Über Abbildungsklassen von Henkelkörpern*. Arch. Math. 33 (1979), 379-382
- [Z2] B. Zimmermann, *Über Homöomorphismen  $n$ -dimensionaler Henkelkörper und endliche Erweiterungen von Schottky-Gruppen*. Comm. Math. Helv. 56 (1981), 474-486
- [Z3] B. Zimmermann, *Finite group actions on handlebodies and equivariant Heegaard genus for 3-manifolds*. Topol. Appl. 43, 263-274 (1992)
- [Z4] B. Zimmermann, *Hurwitz groups and finite group actions on hyperbolic 3-manifolds*. J. London Math. Soc. 52 (1995), 199-208
- [Z5] B. Zimmermann, *Genus actions of finite groups on 3-manifolds*. Michigan Math. J. 43 (1996), 593-610
- [Z6] B. Zimmermann, *A note on surfaces bounding hyperbolic 3-manifolds*. Monatsh. Math. 142 (2004), 267-273