

Characterization of Nonhomogeneous Wavelet Bi-frames for Reducing Subspaces of Sobolev Spaces over Local Fields of Prime Characteristics

M. YOUNUS BHAT

*Department of Mathematical Sciences, Islamic University of Science and Technology
Awantipora, Pulwama, Jammu and Kashmir 192122, India. E-mail:
gyounusg@gmail.com*

Abstract: In this paper, we provide the characterization of nonhomogeneous wavelet bi-frames. First of all we introduce the reducing subspaces of Sobolev spaces over local fields of prime characteristics and then characterize the nonhomogeneous wavelet bi-frames over such fields. **Keywords:** Nonhomogeneous, Local field, Wavelet Bi-frames, Sobolev Spaces.

Mathematics Subject Classification: 42C40; 42C15; 43A70; 11S85; 47A25.

1 Introduction

During the last two decades, there is a substantial body of work that has been concerned with the construction of wavelets on local fields. Even though the structures and metrics of local fields of zero and positive characteristics are similar, their wavelet and MRA (multiresolution analysis) theory are quite different. The theory of wavelets, wavelet frames, multiresolution analysis, Gabor frames on local fields of positive characteristics (LFPC) are extensively studied by many researchers including Benedetto, Behera and Jahan, Ahmad and Shah, Jiang, Li and Ji, Shukla and Mittal in the references [1, 2, 3, 4, 5, 6, 27, 28, 29, 30, 31, 32] but still more concepts required to be studied for its enhancement on LFPC. Continuing our study of frames on local fields, we introduce a comprehensive theory of discrete wave packet systems on local fields by exploiting the machinery of Fourier transforms and we also give the definition of discrete periodic wave packet transform. A characterization of the system to be a Parseval frame and discrete periodic wave packet frame for $\ell^2(\mathcal{Z})$ are obtained. Furthermore, we construct discrete periodic wave packet frame by filter sequence iteration.

We in this paper, provide the characterization of nonhomogeneous wavelet bi-frames. First of all we introduce the reducing subspaces of Sobolev spaces over local fields of prime characteristics and then characterize the nonhomogeneous wavelet bi-frames over such fields.

The rest of the paper is tailored as follows. In Section 2, we recall some basic Fourier analysis on local fields and also some results which are required in the subsequent sections. In section 3, Characterization of NWBFs over Local Field. In Section 4, we construct the J^{th} -stage discrete periodic wave packet frame for $\ell^2(\mathcal{Z})$.

2 Preliminaries on Local Fields

2.1 Local Fields

A local field K is a locally compact, non-discrete and totally disconnected field. If it is of characteristic zero, then it is a field of p -adic numbers \mathbb{Q}_p or its finite extension. If K is of positive characteristic, then K is a field of formal Laurent series over a finite field $GF(p^c)$. If $c = 1$, it is a p -series field, while for $c \neq 1$, it is an algebraic extension of degree c of a p -series field. Let K be a fixed local field with the ring of integers $\mathfrak{D} = \{x \in K : |x| \leq 1\}$. Since K^+ is a locally compact Abelian group, we choose a Haar measure dx for K^+ . The field K is locally compact, non-trivial, totally disconnected and complete topological field endowed with non-Archimedean norm $|\cdot| : K \rightarrow \mathbb{R}^+$ satisfying

- (a) $|x| = 0$ if and only if $x = 0$;
- (b) $|xy| = |x||y|$ for all $x, y \in K$;
- (c) $|x + y| \leq \max\{|x|, |y|\}$ for all $x, y \in K$.

Property (c) is called the ultrametric inequality. Let $\mathfrak{B} = \{x \in K : |x| < 1\}$ be the prime ideal of the ring of integers \mathfrak{D} in K . Then, the residue space $\mathfrak{D}/\mathfrak{B}$ is isomorphic to a finite field $GF(q)$, where $q = p^c$ for some prime p and $c \in \mathbb{N}$. Since K is totally disconnected and \mathfrak{B} is both prime and principal ideal, so there exist a prime element \mathfrak{p} of K such that $\mathfrak{B} = \langle \mathfrak{p} \rangle = \mathfrak{p}\mathfrak{D}$. Let $\mathfrak{D}^* = \mathfrak{D} \setminus \mathfrak{B} = \{x \in K : |x| = 1\}$. Clearly, \mathfrak{D}^* is a group of units in K^* and if $x \neq 0$, then can write $x = \mathfrak{p}^n y, y \in \mathfrak{D}^*$. Moreover, if $\mathcal{U} = \{a_m : m = 0, 1, \dots, q-1\}$ denotes the fixed full set of coset representatives of \mathfrak{B} in \mathfrak{D} , then every element $x \in K$ can be expressed uniquely as $x = \sum_{\ell=k}^{\infty} c_\ell \mathfrak{p}^\ell$ with $c_\ell \in \mathcal{U}$. Recall that \mathfrak{B} is compact and open, so each fractional ideal $\mathfrak{B}^k = \mathfrak{p}^k \mathfrak{D} = \{x \in K : |x| < q^{-k}\}$ is also compact and open and is a subgroup of K^+ . We use the notation in Taibleson's book [33]. In the rest of this paper, we use the symbols \mathbb{N}, \mathbb{N}_0 and \mathbb{Z} to denote the sets of natural, non-negative integers and integers, respectively.

Let χ be a fixed character on K^+ that is trivial on \mathfrak{D} but non-trivial on \mathfrak{B}^{-1} . Therefore, χ is constant on cosets of \mathfrak{D} so if $y \in \mathfrak{B}^k$, then $\chi_y(x) = \chi(y, x), x \in K$. Suppose that χ_u is any character on K^+ , then the restriction $\chi_u|_{\mathfrak{D}}$ is a character on \mathfrak{D} . Moreover, as characters on \mathfrak{D} , $\chi_u = \chi_v$ if and only if $u - v \in \mathfrak{D}$. Hence, if $\{u(n) : n \in \mathbb{N}_0\}$ is a complete list of distinct coset representative of \mathfrak{D} in K^+ , then, as it was proved in [13], the set $\{\chi_{u(n)} : n \in \mathbb{N}_0\}$ of distinct characters on \mathfrak{D} is a complete orthonormal system on \mathfrak{D} .

We now impose a natural order on the sequence $\{u(n)\}_{n=0}^{\infty}$. We have $\mathfrak{D}/\mathfrak{B} \cong GF(q)$ where $GF(q)$ is a c -dimensional vector space over the field $GF(p)$. We choose a set $\{1 = \zeta_0, \zeta_1, \zeta_2, \dots, \zeta_{c-1}\} \subset \mathfrak{D}^*$ such that $\text{span}\{\zeta_j\}_{j=0}^{c-1} \cong GF(q)$. For $n \in \mathbb{N}_0$ satisfying

$$0 \leq n < q, \quad n = a_0 + a_1 p + \dots + a_{c-1} p^{c-1}, \quad 0 \leq a_k < p, \quad \text{and } k = 0, 1, \dots, c-1,$$

we define

$$u(n) = (a_0 + a_1\zeta_1 + \cdots + a_{c-1}\zeta_{c-1}) \mathfrak{p}^{-1}.$$

Also, for $n = b_0 + b_1q + b_2q^2 + \cdots + b_sq^s$, $n \in \mathbb{N}_0$, $0 \leq b_k < q$, $k = 0, 1, 2, \dots, s$, we set

$$u(n) = u(b_0) + u(b_1)\mathfrak{p}^{-1} + \cdots + u(b_s)\mathfrak{p}^{-s}.$$

This defines $u(n)$ for all $n \in \mathbb{N}_0$. In general, it is not true that $u(m+n) = u(m) + u(n)$. But, if $r, k \in \mathbb{N}_0$ and $0 \leq s < q^k$, then $u(rq^k + s) = u(r)\mathfrak{p}^{-k} + u(s)$. Further, it is also easy to verify that $u(n) = 0$ if and only if $n = 0$ and $\{u(\ell) + u(k) : k \in \mathbb{N}_0\} = \{u(k) : k \in \mathbb{N}_0\}$ for a fixed $\ell \in \mathbb{N}_0$. Hereafter we use the notation $\chi_n = \chi_{u(n)}$, $n \geq 0$.

Let the local field K be of characteristic $p > 0$ and $\zeta_0, \zeta_1, \zeta_2, \dots, \zeta_{c-1}$ be as above. We define a character χ on K as follows:

$$\chi(\zeta_\mu \mathfrak{p}^{-j}) = \begin{cases} \exp(2\pi i/p), & \mu = 0 \text{ and } j = 1, \\ 1, & \mu = 1, \dots, c-1 \text{ or } j \neq 1. \end{cases}$$

2.2 Fourier Transforms on Local Fields

The Fourier transform of $f \in L^1(K)$ is denoted by $\widehat{f}(\omega)$ and defined by

$$F\{f(x)\} = \widehat{f}(\omega) = \int_K f(x) \overline{\chi_\omega(x)} dx.$$

It is noted that

$$\widehat{f}(\omega) = \int_K f(x) \overline{\chi_\omega(x)} dx = \int_K f(x) \chi(-\omega x) dx.$$

The properties of Fourier transforms on local field K are much similar to those of on the classical field \mathbb{R} . In fact, the Fourier transform on local fields of positive characteristic have the following properties:

- The map $f \rightarrow \widehat{f}$ is a bounded linear transformation of $L^1(K)$ into $L^\infty(K)$, and $\|\widehat{f}\|_\infty \leq \|f\|_1$.
- If $f \in L^1(K)$, then \widehat{f} is uniformly continuous.
- If $f \in L^1(K) \cap L^2(K)$, then $\|\widehat{f}\|_2 = \|f\|_2$.

The Fourier transform of a function $f \in L^2(K)$ is defined by

$$\widehat{f}(\omega) = \lim_{k \rightarrow \infty} \widehat{f}_k(\omega) = \lim_{k \rightarrow \infty} \int_{|x| \leq q^k} f(x) \overline{\chi_\omega(x)} dx,$$

where $f_k = f \Phi_{-k}$ and Φ_k is the characteristic function of \mathfrak{B}^k . Furthermore, if $f \in L^2(\mathfrak{D})$, then we define the Fourier coefficients of f as

$$\widehat{f}(u(n)) = \int_{\mathfrak{D}} f(x) \overline{\chi_{u(n)}(x)} dx.$$

The series $\sum_{n \in \mathbb{N}_0} \widehat{f}(u(n)) \chi_{u(n)}(x)$ is called the Fourier series of f . From the standard L^2 -theory for compact Abelian groups, we conclude that the Fourier series of f converges to f in $L^2(\mathfrak{D})$ and Parseval's identity holds:

$$\|f\|_2^2 = \int_{\mathfrak{D}} |f(x)|^2 dx = \sum_{n \in \mathbb{N}_0} |\widehat{f}(u(n))|^2.$$

Let $\mathcal{Z} = \{u(n) : n \in \mathbb{N}_0\}$, where $\{u(n) : n \in \mathbb{N}_0\}$ is a complete list of (distinct) coset representation of \mathfrak{D} in K^+ . Then we define

$$\ell^2(\mathcal{Z}) = \left\{ z : \mathcal{Z} \rightarrow \mathbb{C} : \sum_{n \in \mathbb{N}_0} |z(u(n))|^2 < \infty \right\}$$

is a Hilbert space with inner product

$$\langle z, w \rangle = \sum_{n \in \mathbb{N}_0} z(u(n)) \overline{w(u(n))}.$$

The following definitions are natural which are used in the subsequent sections:

Definition 2.1 *The Fourier transform on $\ell^2(\mathcal{Z})$ is the map $\widehat{\cdot} : \ell^2(\mathcal{Z}) \rightarrow L^2(\mathfrak{D})$ defined for $z \in \ell^2(\mathcal{Z})$ by*

$$\widehat{z}(\omega) = \sum_{n \in \mathbb{N}_0} z(u(n)) \chi_{u(n)}(\omega),$$

and the Inverse Fourier transform on $L^2(\mathfrak{D})$ is the map $\vee : L^2(\mathfrak{D}) \rightarrow \ell^2(\mathcal{Z})$ defined for $f \in L^2(\mathfrak{D})$ by

$$f^\vee(u(n)) = \langle f, \chi_{u(n)} \rangle = \int_{\mathfrak{D}} f(x) \overline{\chi_{u(n)}(x)} dx.$$

For $z \in \ell^2(\mathcal{Z})$, we have

$$\begin{aligned} (\widehat{z})^\vee(u(n)) &= \langle \widehat{z}, \chi_{u(n)} \rangle \\ &= \left\langle \sum_{m \in \mathbb{N}_0} z(u(m)) \chi_{u(m)}, \chi_{u(n)} \right\rangle \\ &= \sum_{m \in \mathbb{N}_0} z(u(m)) \langle \chi_{u(m)}, \chi_{u(n)} \rangle \\ &= z(u(n)). \end{aligned}$$

Since $\{\chi_{u(n)} : n \in \mathbb{N}_0\}$ is an orthonormal basis for $L^2(\mathfrak{D})$. It is also clear that the function \widehat{z} is an integral periodic function because for $m \in \mathbb{N}_0$, we have

$$\begin{aligned}\widehat{z}(\omega + u(m)) &= \sum_{n \in \mathbb{N}_0} z(u(n)) \chi_{u(n)}(\omega), \chi_{u(n)}(u(m)). \\ &= \sum_{n \in \mathbb{N}_0} z(u(n)) \chi_{u(n)}(\omega) \\ &= \widehat{z}(\omega).\end{aligned}$$

For $z, w \in \ell^2(\mathcal{Z})$, we have *Parseval's relation*:

$$\langle z, w \rangle = \sum_{n \in \mathbb{N}_0} z(u(n)) \overline{w(u(n))} = \int_{\mathfrak{D}} \widehat{z}(\omega) \overline{\widehat{w}(\omega)} d\omega = \langle \widehat{z}, \widehat{w} \rangle,$$

and *Plancherel's formula*:

$$\|z\|^2 = \sum_{n \in \mathbb{N}_0} |z(u(n))|^2 = \int_{\mathfrak{D}} |\widehat{z}(\omega)|^2 d\omega = \|\widehat{z}\|^2.$$

Definition 2.2 Let \mathcal{H} denotes a separable Hilbert space and \mathcal{P} be countable index set. A countable set $\{\varphi_p : p \in \mathcal{P}\}$ in \mathcal{H} is called a *frame* \mathcal{H} if there exists constants A and B , $0 < C \leq D < \infty$ such that

$$C\|f\|^2 \leq \sum_{p \in \mathcal{P}} |\langle f, \varphi_p \rangle|^2 \leq D\|f\|^2 \quad (1)$$

holds for all $f \in \mathcal{H}$. The greatest possible such C is the *lower frame bound* and the least possible such D is the *upper frame bound*. We say that $\{\varphi_p : p \in \mathcal{P}\}$ is a *tight frame* for \mathcal{H} with *frame bound* C if $C = D$, and a *normalized tight (Parseval) frame* if $C = D = 1$.

Definition 2.3 For $m, k \in \mathbb{N}_0$, the *translation operator* $T_{u(m)} : \ell^2(\mathcal{Z}) \rightarrow \ell^2(\mathcal{Z})$ is defined by

$$T_{u(m)} z(u(n)) = z(u(n) - u(m))$$

and the *modulation operator* $M_{u(k)} : \ell^2(\mathcal{Z}) \rightarrow \ell^2(\mathcal{Z})$ by

$$M_{u(k)} z(u(n)) = z(u(n)) \overline{\chi_{u(k)}}.$$

Definition 2.4 For $v_\ell^j \in \ell^2(\mathcal{Z})$, $0 \leq \ell \leq L, 0 \leq j \leq J$, we call $\mathcal{W}(V)$ a J^{th} -stage *wave packet system* associated with $V = \{v_\ell : 0 \leq \ell \leq L\}$ if

$$\mathcal{W}(V) = \{\mathcal{W}_{k,m}^j v_\ell^j = T_{\mathbf{p}^j u(m)} M_{u(k)} v_\ell^j : k \in \mathbb{N}_0, m \in \mathbb{N}_0, 0 \leq \ell \leq L, 0 \leq j \leq J\}. \quad (2)$$

For $J = 0$, the wave packet system in (2) becomes the *Gabor system*

$$\{\overline{\chi_{u(k)}(u(m))} v_\ell(\cdot - u(m)) : k \in \mathbb{N}_0, 0 \leq \ell \leq L\}.$$

On taking $k = 0$ in (2), the wave packet system degenerates into the *wavelet system*

$$\{v_\ell^j(\cdot - \mathbf{p}^j u(m)) : m \in \mathbb{N}_0, 0 \leq \ell \leq L, j = 1, 2, \dots, J\}.$$

Thus the Gabor system and wavelet system are the particular cases of wave packet systems.

3 Characterization of NWBFs over Local Field

In this section, we provide the characterization of NWBFs in $(FH^s(\Omega), FH^{-s}(\Omega))$. Firstly we need following two lemmas.

Lemma 3.1 *Given $s \in K$, let $\{T_{u(k)}\psi_0 : k \in \mathbb{N}_0\} \cup \{T_{u(k)}\psi_\ell : k \in \mathbb{N}_0, 1 \leq \ell \leq L\}$ be a Bessel sequence in $H^s(K)$. Then*

$$\begin{aligned}
& \sum_{k \in \mathbb{N}_0} |\langle g, \psi_{0,0,k} \rangle|^2 + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \sum_{k \in \mathbb{N}_0} |\langle g, \psi_{\ell,j,k}^s \rangle|^2 \\
&= \int_K |\hat{g}(\omega)|^2 \left(|\hat{\psi}_0(\omega)|^2 + \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{-2js} |\hat{\psi}_\ell(\mathfrak{p}^{-j}\omega)|^2 \right) d\omega \\
&+ \int_K \overline{\hat{g}(\omega)} \sum_{k \in \mathbb{N}_0} \hat{g}(\omega + u(k)) \\
&\times \left(\hat{\psi}_0(\omega) \overline{\hat{\psi}_0(\omega + u(k))} + \sum_{\ell=1}^L \sum_{j=0}^k m^{-2js} \hat{\psi}_\ell(\mathfrak{p}^{-j}\omega) \overline{\hat{\psi}_\ell(\omega + u(k))} \right) \quad (3)
\end{aligned}$$

for $g \in \mathcal{D}$

Proof. From Lemma 2.3, we have

$$\begin{aligned}
& \sum_{k \in \mathbb{N}_0} |\langle g, \psi_{0,0,k} \rangle|^2 + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \sum_{k \in \mathbb{N}_0} |\langle g, \psi_{\ell,j,k}^s \rangle|^2 \\
&= \int_{\mathfrak{D}} \left| \sum_{k \in \mathbb{N}_0} \hat{g}(\omega + u(k)) \overline{\hat{\psi}_0(\omega + u(k))} \right|^2 d\omega \\
&+ \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{j(d-2s)} \int_{\mathfrak{D}} \left| \sum_{k \in \mathbb{N}_0} \hat{g}(\mathbf{p}^j(\omega + u(k))) \overline{\hat{\psi}_\ell(\omega + u(k))} \right|^2 d\omega \\
&= \int_{\mathfrak{D}} \left(\sum_{k \in \mathbb{N}_0} \hat{\psi}_0(\omega + u(k)) \overline{\hat{g}(\omega + u(k))} \right) \left(\sum_{k \in \mathbb{N}_0} \hat{g}(\omega + u(k)) \overline{\hat{\psi}_0(\omega + u(k))} \right) d\omega \\
&+ \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{j(d-2s)} \int_{\mathfrak{D}} \left(\sum_{k \in \mathbb{N}_0} \hat{\psi}_\ell(\omega + u(k)) \overline{\hat{g}(\mathbf{p}^j(\omega + u(k)))} \right) \\
&\times \left(\sum_{k \in \mathbb{N}_0} \hat{g}(\mathbf{p}^j(\omega + u(k))) \overline{\hat{\psi}_\ell(\omega + u(k))} \right) \\
&= \int_{\mathfrak{D}} \left(\sum_{k \in \mathbb{N}_0} \hat{\psi}_0(\omega + u(k)) \overline{\hat{g}(\omega + u(k))} \right) E_0(\omega) d\omega \\
&+ \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{j(d-2s)} \int_{\mathfrak{D}} \left(\sum_{k \in \mathbb{N}_0} \hat{\psi}_\ell(\omega + u(k)) \overline{\hat{g}(\mathbf{p}^j(\omega + u(k)))} \right) E_{\ell,j} d\omega \\
&= R_1 + R_2, \tag{4}
\end{aligned}$$

where $E_0(\cdot) = \sum_{k \in \mathbb{N}_0} \hat{g}(\omega + u(k)) \overline{\hat{\psi}_0(\omega + u(k))}$ and $E_{\ell,j}(\cdot) = \sum_{k \in \mathbb{N}_0} \hat{g}(\mathbf{p}^j(\omega + u(k))) \overline{\hat{\psi}_\ell(\omega + u(k))}$. Since $\{T_{u(k)}\psi_0 : k \in \mathbb{N}_0\}$ is a Bessel sequence in $H^s(K)$ and $g \in \mathcal{D}$, it follows that $|E_0(\cdot)| \leq [\hat{g}, \hat{g}]_{-s}^{\frac{1}{2}}(\cdot) [\hat{\psi}_0, \hat{\psi}_0]_{-s}^{\frac{1}{2}}(\cdot) < \infty$ by Lemma 2.2 and Lemma 2.5. Therefore

$$\int_{\mathfrak{D}} \left| \sum_{k \in \mathbb{N}_0} \hat{g}(\omega + u(k)) \overline{\hat{\psi}_0(\omega + u(k))} E_0(\omega) \right| \leq \|E_0\|_{L^\infty(\mathfrak{D})} \int_{\mathfrak{D}} [\hat{g}, \hat{g}]_{-s}^{\frac{1}{2}}(\omega) [\hat{\psi}_0, \hat{\psi}_0]_{-s}^{\frac{1}{2}}(\omega) < \infty$$

, hence by Fubini-Tonelli theorem

$$\begin{aligned}
& \int_{\mathfrak{D}} \left(\sum_{k \in \mathbb{N}_0} \hat{\psi}_0(\omega + u(k)) \overline{\hat{g}(\omega + u(k))} \right) \left(\sum_{k \in \mathbb{N}_0} \hat{g}(\omega + u(k)) \overline{\hat{\psi}_0(\omega + u(k))} \right) d\omega \\
&= \int_{\mathfrak{D}} \hat{\psi}_0(\omega) \overline{\hat{g}(\omega)} \sum_{k \in \mathbb{N}_0} \hat{g}(\omega + u(k)) \overline{\hat{\psi}_0(\omega + u(k))} d\omega
\end{aligned}$$

Moreover

$$\begin{aligned}
& \int_K |\hat{\psi}_0(\omega) \overline{\hat{g}(\omega)}| \sum_{k \in \mathbb{N}_0} |\hat{g}(\omega + u(k)) \overline{\hat{\psi}_0(\omega + u(k))}| d\omega \\
& \leq \int_{\text{supp}(\hat{g})} \left(\sum_{k \in \mathbb{N}_0} |\hat{g}(\omega + u(k)) \overline{\hat{\psi}_0(\omega + u(k))}| \right)^2 d\omega \\
& \leq \int_{\text{supp}(\hat{g})} [\hat{g}, \hat{g}]_{-s}(\omega) [\hat{\psi}_0, \hat{\psi}_0]_s(\omega) d\omega \\
& < \infty
\end{aligned}$$

since $[\hat{g}, \hat{g}]_{-s}(\omega) [\hat{\psi}_0, \hat{\psi}_0]_s(\cdot)$ is essentially bounded by Lemma 2.2, we have

$$\begin{aligned}
R_1 &= \int_K \hat{\psi}_0(\omega) \overline{\hat{g}(\omega)} \sum_{k \in \mathbb{N}_0} \hat{g}(\omega + u(k)) \overline{\hat{\psi}_0(\omega + u(k))} d\omega \\
&= \int_K |\hat{\psi}_0(\omega)|^2 |\hat{g}(\omega)|^2 d\omega + \int_K \hat{\psi}_0(\omega) \overline{\hat{g}(\omega)} \sum_{k \in \mathbb{N}_0} \hat{g}(\omega + u(k)) \overline{\hat{\psi}_0(\omega + u(k))} d\omega \quad (5)
\end{aligned}$$

In order to complete the proof, we need to calculate R_2 . Let us define \tilde{g} by $\hat{\tilde{g}}(\cdot) = \hat{g}((\mathbf{p})^j \cdot)$. Then as $g \in \mathcal{D}$ and Lemma 2.5, we have

$$[\hat{\tilde{g}}((\mathbf{p})^j \cdot), \hat{\tilde{g}}((\mathbf{p})^j \cdot)](\cdot) \leq C$$

Thus

$$|E_{i,j}(\cdot)| \leq [\hat{\tilde{g}}((\mathbf{p})^j \cdot), \hat{\tilde{g}}((\mathbf{p})^j \cdot)]_{-s}^{\frac{1}{2}}(\cdot) [\hat{\psi}_\ell, \hat{\psi}_\ell]_s^{\frac{1}{2}} < \infty.$$

Hence

$$R_2 = \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{j(d-2s)} \int_K \hat{\psi}_\ell(\omega) \overline{g(\mathbf{p}^j \omega)} \sum_{k \in \mathbb{N}_0} \hat{g}(\mathbf{p}^j(\omega + u(k))) \overline{\hat{\psi}_\ell(\omega + u(k))} d\omega$$

Taking A a bounded set in K such that $\text{supp}(\hat{g}) \subset A$. So by Lemma 2.6, we get

$$A \cap (A + \mathbf{p}^j u(k)) = \emptyset \quad \text{for } (j, k) \notin A_1 \times A_2 \quad \text{with } k \neq 0,$$

where $A_1 \subset \mathbb{N}_0$ and $A_2 \subset \mathbb{N}_0 \setminus \{0\}$ are two finite sets. Therefore

$$R_2 = \sum_{\ell=1}^L \sum_{j \in A_1} m^{j(d-2s)} \int_K \hat{\psi}_\ell(\omega) \overline{g(\mathbf{p}^j \omega)} \sum_{k \in A_2} \hat{g}(\mathbf{p}^j(\omega + u(k))) \overline{\hat{\psi}_\ell(\omega + u(k))} d\omega$$

Denote $S = \bigcup_{k \in A_2 \cap \{0\}} (\bigcup_{j \in A_1} (\mathbf{p}^j A + u(k)))$. Therefore, for each $(j, k) \in A_1 \times A_2$, we have

$$\begin{aligned}
& \int_K \left| \hat{\psi}_\ell(\omega) \overline{g(\mathbf{p}^j \omega)} \hat{g}(\mathbf{p}^j(\omega + u(k))) \overline{\hat{\psi}_\ell(\omega + u(k))} \right| d\omega \\
& \leq \|\hat{g}\|_{L^\infty(K)}^2 \int_{\mathbf{p}^{-j}A} |\hat{\psi}_\ell(\omega) \hat{\psi}_\ell(\omega + u(k))| d\omega \\
& \leq \|\hat{g}\|_{L^\infty(K)}^2 \left(\int_{\mathbf{p}^{-j}A} |\hat{\psi}_\ell(\omega)|^2 d\omega \right)^{\frac{1}{2}} \left(\int_{\mathbf{p}^{-j}A} |\hat{\psi}_\ell(\omega + u(k))|^2 d\omega \right)^{\frac{1}{2}} \\
& \leq \|\hat{g}\|_{L^\infty(K)}^2 \int_S |\hat{\psi}_\ell(\omega)|^2 d\omega
\end{aligned}$$

Note that $1 \leq (\max_{\omega \in A} (1 + |\omega|^2)^{-s})(1 + |\omega|^2)^s$ for $\omega \in A$. Thus

$$\begin{aligned} & \int_K \left| \hat{\psi}_\ell(\omega) \overline{g(\mathbf{p}^j \omega)} \hat{g}(\mathbf{p}^j(\omega + u(k))) \overline{\hat{\psi}_\ell(\omega + u(k))} \right| d\omega \\ & \leq \left(\max_{\omega \in A} (1 + |\omega|^2)^{-s} \right) \|\hat{g}\|_{L^\infty(K)}^2 \int_A |\hat{\psi}_\ell(\omega)|^2 (1 + |\omega|^2)^s d\omega \\ & \leq \left(\max_{\omega \in A} (1 + |\omega|^2)^{-s} \right) \|\hat{g}\|_{L^\infty(K)}^2 \|\psi_\ell\|_{H^s(K)}^2 \\ & < \infty \end{aligned}$$

On combining the formula given above, we get

$$\begin{aligned} R_2 &= \int_K \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{j(d-2s)} |\hat{\psi}_\ell(\omega)|^2 |\hat{g}(\mathbf{p}^j \omega)|^2 d\omega \\ &+ \int_K \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{j(d-2s)} \overline{\hat{g}(\mathbf{p}^j \omega)} \hat{\psi}_\ell(\omega) \sum_{k \in \mathbb{N}_0} \hat{g}(\mathbf{p}^j(\omega + u(k))) \overline{\hat{\psi}_\ell(\omega + u(k))} d\omega \\ &= \int_K \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{-2js} |\hat{\psi}_\ell(\mathbf{p}^{-j} \omega)|^2 |\hat{g}(\omega)|^2 d\omega \\ &+ \int_K \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{-2js} \overline{\hat{g}(\omega)} \hat{\psi}_\ell(\mathbf{p}^{-j} \omega) \sum_{k \in \mathbb{N}_0} \hat{g}(\omega + \mathbf{p}^{-j} u(k)) \overline{\hat{\psi}_\ell(\mathbf{p}^{-j} \omega + u(k))} d\omega \\ &= \int_K \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{-2js} |\hat{\psi}_\ell(\mathbf{p}^{-j} \omega)|^2 |\hat{g}(\omega)|^2 d\omega \\ &+ \int_K \overline{\hat{g}(\omega)} \sum_{k \in \mathbb{N}_0} \hat{g}(\omega + u(k)) \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{-2js} \hat{\psi}_\ell(\mathbf{p}^{-j} \omega) \overline{\hat{\psi}_\ell(\mathbf{p}^{-j} \omega + u(k))} d\omega \quad (6) \end{aligned}$$

using the definition of $\kappa(k)$. Hence using (4), (5) and (6), we get (3). This completes the proof of the lemma.

Lemma 3.2 *Given $s \in K$, let $X^s(\psi_0; \Psi)$ and $X^{-s}(\tilde{\psi}_0; \tilde{\Psi})$ be a Bessel sequences in $H^s(K)$ and $H^{-s}(K)$, respectively. Then for all $f, g \in \mathcal{D}$, we have*

$$\begin{aligned} & \sum_{k \in \mathbb{N}_0} \langle f, \tilde{\psi}_{0,0,k} \rangle \langle \psi_{0,0,k}, g \rangle + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \sum_{k \in \mathbb{N}_0} \langle f, \tilde{\psi}_{\ell,j,k}^{-s} \rangle \langle \psi_{\ell,j,k}^s, g \rangle \\ &= \int_K \hat{f}(\omega) \overline{\hat{g}(\omega)} \left(\hat{\psi}_0(\omega) \overline{\hat{\psi}_0(\omega)} + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \hat{\psi}_\ell(\mathbf{p}^{-j} \omega) \overline{\hat{\psi}_\ell(\mathbf{p}^{-j} \omega)} \right) d\omega \\ &+ \int_K \overline{\hat{g}(\omega)} \sum_{k \in \mathbb{N}_0} \hat{f}(\omega + u(k)) \\ &\times \left(\hat{\psi}_0(\omega) \overline{\hat{\psi}_0(\omega + u(k))} + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \hat{\psi}_\ell(\mathbf{p}^{-j} \omega) \overline{\hat{\psi}_\ell(\mathbf{p}^{-j} \omega + u(k))} \right) d\omega \quad (7) \end{aligned}$$

proof: As $X^s(\psi_0; \Psi)$ and $X^{-s}(\tilde{\psi}_0; \tilde{\Psi})$ are Bessel sequences in $H^s(K)$ and $H^{-s}(K)$, respectively, the expression in (7) is meaningful. Proceeding in a similar fashion as in Lemma 3.1, we have

$$\begin{aligned}
& \sum_{k \in \mathbb{N}_0} \langle f, \tilde{\psi}_{0,0,k} \rangle \langle \psi_{0,0,k}, g \rangle + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \sum_{k \in \mathbb{N}_0} \langle f, \tilde{\psi}_{\ell,j,k}^{-s} \rangle \langle \psi_{\ell,j,k}^s, g \rangle \\
&= \int_K \hat{\psi}_0(\omega) \overline{\hat{g}(\omega)} \overline{\sum_{k \in \mathbb{N}_0} \hat{f}(\omega + u(k))} \hat{\psi}_0(\omega + u(k)) d\omega \\
&+ \sum_{\ell=1}^L \sum_{j=0}^{\infty} q^j \int_K \hat{\psi}_\ell(\omega) \overline{\hat{g}(\mathbf{p}^j \omega)} \overline{\sum_{k \in \mathbb{N}_0} \hat{f}(\mathbf{p}^j(\omega + u(k)))} \hat{\psi}_\ell(\omega + u(k)) d\omega \\
&= I_1 + I_2.
\end{aligned} \tag{8}$$

Note that

$$|\hat{\psi}_0(\cdot) \overline{\hat{g}(\cdot)}| \sum_{k \in \mathbb{N}_0} |\hat{f}(\cdot + u(k)) \overline{\hat{\psi}_0(\cdot + u(k))}| \leq [\hat{f}, \hat{f}]_s^{\frac{1}{2}}(\cdot) [\hat{\psi}_0, \tilde{\psi}_0]_{-s}^{\frac{1}{2}}(\cdot) [\hat{g}, \hat{g}]_{-s}^{\frac{1}{2}}(\cdot) [\hat{\psi}_0, \psi_0]_s^{\frac{1}{2}}(\cdot),$$

is bounded due to Lemma 2.2. Hence

$$\begin{aligned}
& \int_K |\hat{\psi}_0(\omega) \overline{\hat{g}(\omega)}| \overline{\sum_{k \in \mathbb{N}_0} \hat{f}(\omega + u(k))} \hat{\psi}_0(\omega + u(k)) d\omega \\
& \leq \int_{\text{supp}(\hat{g})} |\hat{\psi}_0(\omega) \overline{\hat{g}(\omega)}| \overline{\sum_{k \in \mathbb{N}_0} \hat{f}(\omega + u(k))} \hat{\psi}_0(\omega + u(k)) d\omega < \infty
\end{aligned}$$

Therefore

$$I_1 = \int_K \hat{f}(\omega) \overline{\hat{g}(\omega)} \hat{\psi}_0(\omega) \overline{\hat{\psi}_0(\omega)} + \int_K \hat{\psi}_0(\omega) \overline{\hat{g}(\omega)} \overline{\sum_{k \in \mathbb{N}_0} \hat{f}(\omega + u(k))} \hat{\psi}_0(\omega + u(k)) d\omega. \tag{9}$$

To complete the proof of the lemma, we need to discuss I_2 . Let's break it into two parts, for $k = 0$ and $k \neq 0$. Hence by Cauchy-Schwartz inequality and Lemma 2.4, we have

$$\begin{aligned}
& \sum_{\ell=1}^L \sum_{j=0}^{\infty} |\hat{\psi}_\ell(\mathbf{p}^{-j} \omega) \overline{\hat{\psi}_\ell(\mathbf{p}^{-j} \omega)}| \\
& \leq \left(\sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{-2js} |\hat{\psi}_\ell(\mathbf{p}^{-j} \omega)|^2 \right)^{\frac{1}{2}} \left(\sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{2js} |\hat{\psi}_\ell(\mathbf{p}^{-j} \omega)|^2 \right)^{\frac{1}{2}} \\
& \leq B_1 B_2.
\end{aligned} \tag{10}$$

Therefore

$$\begin{aligned}
& \int_K |\hat{f}(\omega) \overline{\hat{g}(\omega)}| \sum_{\ell=1}^L \sum_{j=0}^{\infty} |\hat{\psi}_\ell(\mathbf{p}^{-j} \omega) \overline{\hat{\psi}_\ell(\mathbf{p}^{-j} \omega)}| \\
& \leq B_1 B_2 \left| \text{supp}(\hat{f}) \cap \text{supp}(\hat{g}) \right| \|\hat{f}\|_{L^\infty(K)} \|\hat{g}\|_{L^\infty(K)} \\
& < \infty
\end{aligned}$$

Fix a compact set $A \in K$ such that $\text{supp}(\hat{f}) \cap \text{supp}(\hat{g}) \subset A$. Using Lemma 2.6, it follows that

$$A \cap (A + \mathbf{p}^j u(k)) = \emptyset \quad \text{for } (j, k) \notin A_1 \times A_2 \quad \text{with } k \neq 0 \quad (11)$$

where $A_1 \subset \mathbb{N}_0$ and $A_2 \subset \mathbb{N}_0 \setminus \{0\}$ are two finite sets. With the same argument as applied to R_2 , we have

$$\begin{aligned} & \int_K \overline{|\hat{g}(\mathbf{p}^j \omega) \hat{f}(\mathbf{p}^j(\omega + u(k))) \hat{\psi}_\ell(\omega) \hat{\psi}_\ell(\omega + u(k))|} d\omega \\ & \leq \|\hat{g}\|_{L^\infty(K)} \|\hat{f}\|_{L^\infty(K)} \left(\int_{\mathbf{p}^{-j}A} |\hat{\psi}_\ell(\omega)|^2 \right)^{\frac{1}{2}} \left(\int_{\mathbf{p}^{-j}A} |\hat{\psi}_\ell(\omega + u(k))|^2 \right)^{\frac{1}{2}} \\ & \leq \|\hat{g}\|_{L^\infty(K)} \|\hat{f}\|_{L^\infty(K)} \left(\int_T |\hat{\psi}_\ell(\omega)|^2 \right)^{\frac{1}{2}} \left(\int_T |\hat{\psi}_\ell(\omega + u(k))|^2 \right)^{\frac{1}{2}} \\ & \leq \|\hat{g}\|_{L^\infty(K)} \|\hat{f}\|_{L^\infty(K)} \left(\max_{\omega \in T} (1 + |\omega|^2) \right)^{-s/2} \left(\max_{\omega \in T} (1 + |\omega|^2) \right)^{s/2} \|\psi_\ell\|_{H^s(K)} \|\tilde{\psi}_\ell\|_{H^{-s}(K)} \\ & < \infty \end{aligned} \quad (12)$$

for $(j, k) \in A_1 \times A_2$, where $T = \bigcup_{k \in A_2 \cup \{0\}} (\bigcup_{j \in A_1} \mathbf{p}^{-j}A + u(k))$. Using (10) and (12), we get

$$\begin{aligned} I_2 &= \sum_{\ell=1}^L \sum_{j=0}^{\infty} \int_K \overline{\hat{g}(\omega) \hat{\psi}_\ell(\mathbf{p}^{-j} \omega)} \sum_{k \in \mathbb{N}_0} \hat{f}(\omega + \mathbf{p}^j u(k)) \overline{\hat{\psi}_\ell(\mathbf{p}^{-j}(\omega + u(k)))} d\omega \\ &= \int_K \sum_{\ell=1}^L \sum_{j=0}^{\infty} \overline{\hat{g}(\omega) \hat{\psi}_\ell(\mathbf{p}^{-j} \omega)} \sum_{k \in \mathbb{N}_0} \hat{f}(\omega + \mathbf{p}^j u(k)) \overline{\hat{\psi}_\ell(\mathbf{p}^{-j}(\omega + u(k)))} d\omega \\ &= \int_K \overline{\hat{g}(\omega)} \sum_{k \in \mathbb{N}_0} \hat{f}(\omega + u(k)) \sum_{\ell=1}^L \sum_{j=0}^{\kappa(k)} \overline{\hat{\psi}_\ell(\mathbf{p}^{-j} \omega) \hat{\psi}_\ell(\mathbf{p}^{-j}(\omega + u(k)))} d\omega \end{aligned} \quad (13)$$

On combining (8), (9) and (13), we get (7), which completes the proof of the lemma. We now present a characterization of NWBFs in $(FH^s(\Omega), FH^{-s}(\Omega))$ in the form of the following theorem.

Theorem 3.3 *Given $s \in K$, let $FH^s(\Omega)$ and $FH^{-s}(\Omega)$ be reducing subspaces of $H^s(K)$ and $H^{-s}(K)$, respectively, $\psi_0 \in H^s(K)$, $\tilde{\psi}_0 \in H^{-s}(K)$ and $\Psi \in H^s(K)$, $\tilde{\Psi} \in H^{-s}(K)$. Suppose that $X^s(\psi_0, \Psi)$ and $X^{-s}(\tilde{\psi}_0, \tilde{\Psi})$ are Bessel sequences in $FH^s(\Omega)$ and $FH^{-s}(\Omega)$, respectively. Then $X^s(\psi_0, \Psi); X^{-s}(\tilde{\psi}_0, \tilde{\Psi})$ is an NWBFs in $(FH^s(\Omega), FH^{-s}(\Omega))$ if and only if*

$$\overline{\hat{\psi}_0(\cdot) \hat{\tilde{\psi}}_0(\cdot + u(k))} \sum_{\ell=1}^L \sum_{j=0}^{\kappa(k)} \overline{\hat{\psi}_\ell(\mathbf{p}^{-j} \cdot) \hat{\tilde{\psi}}_\ell(\mathbf{p}^{-j}(\cdot + u(k)))} = \delta_{0,k} \quad \text{a.e. on } \Omega. \quad (14)$$

Proof: As $\mathcal{D} \cap FH^s(\Omega)$ is dense in $FH^s(\Omega)$, then

$$X^s(\psi_0, \Psi); X^{-s}(\tilde{\psi}_0, \tilde{\Psi})$$

is an NWBF's in $(FH^s(\Omega), FH^{-s}(\Omega))$ iff for $f \in \mathcal{D} \cap FH^s(\Omega)$ and $g \in \mathcal{D} \cap FH^{-s}(\Omega)$

$$\sum_{k \in \mathbb{N}_0} \langle f, \tilde{\psi}_{0,0,k} \rangle \langle \psi_{0,0,k}, g \rangle + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \sum_{k \in \mathbb{N}_0} \langle f, \tilde{\psi}_{\ell,j,k}^{-s} \rangle \langle \psi_{\ell,j,k}^s, g \rangle = \langle f, g \rangle$$

This is equivalent to

$$\begin{aligned} & \sum_{k \in \mathbb{N}_0} \langle (\hat{f}\chi_\Omega), \tilde{\psi}_{0,0,k} \rangle \langle \psi_{0,0,k}, (\hat{g}\chi_\Omega) \rangle + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \sum_{k \in \mathbb{N}_0} \langle (\hat{f}\chi_\Omega), \tilde{\psi}_{\ell,j,k}^{-s} \rangle \langle \psi_{\ell,j,k}^s, (\hat{g}\chi_\Omega) \rangle \\ & = \langle (\hat{f}\chi_\Omega), (\hat{g}\chi_\Omega) \rangle \end{aligned} \quad (15)$$

with $f, g \in \mathcal{D}$ as $\mathcal{D} \cap FH^s(\Omega) = \{(\hat{h}\chi_\Omega) : h \in \mathcal{D}\}$. The expression of (15) is well defined as $X^s(\psi_0, \Psi)$ and $X^{-s}(\tilde{\psi}_0, \tilde{\Psi})$ are Bessel sequences in $H^s(K)$ and $H^{-s}(K)$. Hence using Lemma 3.2, we can write expression (15) as

$$\begin{aligned} & \int_K \hat{f}(\omega) \overline{\hat{g}(\omega)} \chi_\Omega(\omega) \left(\hat{\psi}_0(\omega) \overline{\hat{\psi}_0(\omega)} + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \hat{\psi}_\ell(\mathbf{p}^{-j}\omega) \overline{\hat{\psi}_\ell(\mathbf{p}^{-j}\omega)} \right) d\omega \\ & + \int_K \overline{\hat{g}(\omega)} \chi_\Omega(\omega) \sum_{k \in \mathbb{N}_0} (\hat{f}\chi_\Omega)(\omega + u(k)) \\ & \times \left(\hat{\psi}_0(\omega) \overline{\hat{\psi}_0(\omega + u(k))} + \sum_{\ell=1}^L \sum_{j=0}^{\kappa(k)} \hat{\psi}_\ell(\mathbf{p}^{-j}\omega) \overline{\hat{\psi}_\ell(\mathbf{p}^{-j}\omega + u(k))} \right) d\omega \\ & = \int_K \hat{f}(\omega) \overline{\hat{g}(\omega)} \chi_\Omega(\omega) \end{aligned} \quad (16)$$

with $f, g \in \mathcal{D}$. Hence the expression (14) leads (16). It remains only to prove the converse statement. Suppose (16) hold. Using Cauchy-Schwartz inequality, we have

$$\begin{aligned} & |\hat{\psi}_0(\cdot) \overline{\hat{\psi}_0(\cdot + u(k))}| + \sum_{\ell=1}^L \sum_{j=0}^{\kappa(k)} |\hat{\psi}_\ell(\mathbf{p}^{-j}\cdot) \overline{\hat{\psi}_\ell(\mathbf{p}^{-j}\cdot + u(k))}| \\ & \leq \left(|\hat{\psi}_0(\cdot)|^2 + \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{-2js} |\hat{\psi}_\ell(\mathbf{p}^{-j}\cdot)|^2 \right)^{\frac{1}{2}} \\ & + \left(|\hat{\psi}_0(\cdot + u(k))|^2 + \sum_{\ell=1}^L \sum_{j=0}^{\infty} m^{-2js} |\hat{\psi}_\ell(\mathbf{p}^{-j}(\cdot + u(k)))|^2 \right)^{\frac{1}{2}} \\ & \leq B_1 B_2 (1 + |\cdot|^2)^{-s} (1 + |\cdot + u(k)|^2)^s \\ & = C_k < \infty \end{aligned}$$

for each $k \in \mathbb{N}_0$ using Lemma 2.4. Hence the series

$\hat{\psi}_0(\cdot) \overline{\hat{\psi}_0(\cdot + u(k))} + \sum_{\ell=1}^L \sum_{j=0}^{\kappa(k)} \hat{\psi}_\ell(\mathbf{p}^{-j}\cdot) \overline{\hat{\psi}_\ell(\mathbf{p}^{-j}\cdot + u(k))}$ converges absolutely a. e. on K and is contained in $L^\infty(K)$. Therefore almost all points in K are its Lebesgue

points. Next we consider two cases. When $k = 0$. Let $\omega_0 \neq 0$ be a Lebesgue point of $\hat{\psi}_0(\cdot)\overline{\hat{\psi}_0(\cdot)} + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \hat{\psi}_\ell(\mathbf{p}^{-j}\cdot)\overline{\hat{\psi}_\ell(\mathbf{p}^{-j}\cdot)}$ and $\chi_\Omega(\cdot)$. Fix f and g for $0 < \epsilon < u(1)$, we have

$$\hat{f}(\cdot) = \hat{g}(\cdot) = \frac{\chi_{B(\omega_0, \epsilon)}}{\sqrt{|B(\omega_0, \epsilon)|}}$$

in (16), where $B(\omega_0, \epsilon)$ is an open ball centred at ω_0 and radius ϵ . Therefore

$$\begin{aligned} & \frac{1}{|B(\omega_0, \epsilon)|} \int_{B(\omega_0, \epsilon)} \chi_\Omega(\omega) \left(\hat{\psi}_0(\omega)\overline{\hat{\psi}_0(\omega)} + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \hat{\psi}_\ell(\mathbf{p}^{-j}\omega)\overline{\hat{\psi}_\ell(\mathbf{p}^{-j}\omega)} \right) d\omega \\ &= \frac{1}{|B(\omega_0, \epsilon)|} \int_{B(\omega_0, \epsilon)} \chi_\Omega(\omega) d\omega \end{aligned}$$

letting $\epsilon \rightarrow 0$, we have

$$\hat{\psi}_0(\omega_0)\overline{\hat{\psi}_0(\omega_0)} + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \hat{\psi}_\ell(\mathbf{p}^{-j}\omega_0)\overline{\hat{\psi}_\ell(\mathbf{p}^{-j}\omega_0)}$$

For $k \neq 0$, we fix $k_0 \in \mathbb{N}_0$ and take f and g

$$\hat{f}(\cdot + u(k)) = \hat{g}(\cdot) = \frac{\chi_{B(\omega_0, \epsilon)}}{\sqrt{|B(\omega_0, \epsilon)|}}$$

in (16), with $0 < \epsilon < \frac{1}{2}$. Therefore

$$\begin{aligned} & \frac{1}{|B(\omega_0, \epsilon)|} \int_{B(\omega_0, \epsilon)} \chi_\Omega(\omega) \\ & \times \left(\hat{\psi}_0(\omega)\overline{\hat{\psi}_0(\omega + u(k))} + \sum_{\ell=1}^L \sum_{j=0}^{\kappa(k)} \hat{\psi}_\ell(\mathbf{p}^{-j}\omega)\overline{\hat{\psi}_\ell(\mathbf{p}^{-j}\omega + u(k))} \right) d\omega = 0 \end{aligned}$$

letting $\epsilon \rightarrow 0$ and using Lebesgue differentiation theorem, we get

$$\hat{\psi}_0(\omega_0 + u(k_0))\overline{\hat{\psi}_0(\omega_0)} + \sum_{\ell=1}^L \sum_{j=0}^{\infty} \hat{\psi}_\ell(\mathbf{p}^{-j}\omega_0)\overline{\hat{\psi}_\ell(\mathbf{p}^{-j}\omega_0 + u(k_0))}$$

Hence we obtain (14) by using the arbitrariness of ω_0 and k_0 , which completes the proof of the theorem.

References

- [1] Ahmad O, Sheikh N.A, Ali M. A, Nonuniform nonhomogeneous dual wavelet frames in Sobolev spaces in $L^2(\mathbb{K})$, *Afrika Math.* (2020) doi.org/10.1007/s13370-020-00786-1.
- [2] Ahmad O, Sheikh N.A, On Characterization of nonuniform tight wavelet frames on local fields, *Anal. Theory Appl.*, **34** (2018) 135-146.

- [3] Ahmad O, Shah F. A and Sheikh N. A. *Gabor frames on non-Archimedean fields* International Journal of Geometric Methods in Modern Physics **15** (2018) 1850079 (17 pages).
- [4] Behera B and Jahan Q, Multiresolution analysis on local fields and characterization of scaling functions, *Adv. Pure Appl. Math.* **3** (2) (2012) 181-202.
- [5] Behera B and Jahan Q, Characterization of wavelets and MRA wavelets on local fields of positive characteristic, *Collect. Math.* **66** (1) (2015) 33-53.
- [6] Benedetto J.J and . Benedetto R. L, A wavelet theory for local fields and related groups. *J. Geom. Anal.* **14** (2004) 423-456.
- [7] Christensen O. *An introduction to frames and Riesz bases*. New York: Springer; 2003.
- [8] Chui C. K, Shi X. *Orthonormal wavelets and tight frames with arbitrary real dilations*. Appl. Comput. Harmon. Anal. (2000) **9** 243-264.
- [9] Christensen O, Rahimi A. *Frame properties of wave packet systems in $L^2(\mathbb{R}^d)$* . Adv. Comp. Math. (2008) **29** 101-111.
- [10] Christensen O. *Frames, bases, and discrete Gabor/wavelet expansions*. Bull. Amer. Math. Soc. (2001) **38** 273-291.
- [11] Cordoba A, Fefferman C. *Wave packets and Fourier integral operators* Comm. Partial Differential Equations. (1978) **3** 979-1005.
- [12] Daubechies I. *Ten lecture on wavelets*. Philadelphia(PA):SIAM; (1992).
- [13] Daubechies I, Grossman A, Meyer Y. *Painless non orthogonal expansions*. J. Math. Phys. (1986) **27** 1271-1283.
- [14] Dou X, Jia J , Liu Y. *Spline wavelets in $\ell^2(\mathbb{Z})$* . J. Math. Anal. Appl. **321** (2006) 59-74.
- [15] Duffin and Schaeffer A.C. *A class of nonharmonic Fourier series*. Trans. Amer. Math. Soc. (1952) **72** 341-366.
- [16] Gabor D. *Theory of communications*. J. Inst. Elect. Engn. **93** (1946), 429-457.
- [17] Han B. *Properties of discrete framelet transforms*. Math. Model. Nat. Phenom, **81** (2013) 18-47.
- [18] Ibratovich V. K. *Nonspreading wave packets in quantum mechanics*. Fond. Phys. (1978) **8** 564-571.
- [19] Kaur M, Arora B, Mian M. *Wave packet construction in three-dimensional quantum billiards: Visualizing the closed orbit, collapse and revival of wave packets in the cubical billiard*. J. Phys. (2016) **86** 31-48.
- [20] Labate D, Weiss G, Wilson E. *An approach to the study of wave packet systems*. Contemp. Math., Wavelets, Frames Operator Theory. (2004) **345** 215-235.
- [21] Li. D and Jiang H. K. *The necessary condition and sufficient conditions for wavelet frame on local fields*. J. Math. Anal. Appl. **345** (2008) 500-510.

- [22] Li D. F, Jiang H. K. *Basic results Gabor frame on local fields* Chinese Ann. Math. **28 B** (2) (2007) 165-176.
- [23] Lu D. and Li D. *Frame properties of generalized shift-invariant systems in discrete setting.* Appl. Anal. **95** (2016) 2535-2552.
- [24] Pan A. K. *Understanding the spreading of a Gaussian wave packet using the Bohmian machinery.* J. Phys. (2010) **74** 867-874.
- [25] Ron A, Shen Z. W. *Generalized shift-invariant systems.* Constr. Approx. (2005) **22** 1-45.
- [26] Ron A, Shen Z. W. *Affine systems in $L^2(\mathbb{R}^d)$: the analysis of the analysis operator.* J. Functional Anal. Appl. (1997) **148** 408-447.
- [27] Shah F. A. and Ahmad O. Wave packet systems on local fields, Journal of Geometry and Physics, **120** (2017) 5-18.
- [28] Shah F. A, Ahmad, O, Rahimi A, Frames Associated with Shift Invariant Spaces on Local Fields, *Filomat* **32** (9) (2018) 3097-3110.
- [29] Shah F.A, Ahmad O, Jan H, Sufficient Conditions for Nonuniform Wavelet Frames on Local Fields, *Facta Universtatis*, University of Nis, Serbia, **32** (4) (2017) 551-563.
- [30] Shah F. A, Ahmad O, Sheikh N. A, Some New Inequalities for Wavelet Frames on Local Fields, *Anal. Theory Appl.*, **33** (2) (2017) 134-148.
- [31] Shukla N. K, Maury S. C, Mittal S, Semi-orthogonal Parseval Wavelets Associated with GMRA's on Local Fields of Positive Characteristic, *Mediterr. J. Math.* (2019) 16: 120 <https://doi.org/10.1007/s00009-019-1383-1>.
- [32] Shukla N. K, Maury S. C, Super-wavelets on local fields of positive characteristic. *Math. Nachr.* **291** (4) (2018) 704-719.
- [33] Taibleson M. H (1975). *Fourier Analysis on Local Fields*, Princeton University Press, Princeton, NJ.
- [34] Xu M, Lu D and Fan Q. *Construction of J^{th} -stage discrete periodic wave packet frames* Appl. Anal. DOI: 10.1080/00036811.2017.1343465.