

RESOLVING SINGULARITIES OF CURVES WITH ONE TORIC MORPHISM

ANA BELÉN DE FELIPE, PEDRO D. GONZÁLEZ PÉREZ, AND HUSSEIN MOURTADA

This paper is dedicated to Bernard Teissier.

ABSTRACT. We give an explicit positive answer, in the case of reduced curve singularities, to a question of B. Teissier about the existence of a toric embedded resolution after reembedding. In the case of a curve singularity (C, O) contained in a non singular surface S such a reembedding may be defined in terms of a sequence of maximal contact curves associated to C . We prove that there exists a toric modification, after reembedding, which provides an embedded resolution of C . We use properties of the semivaluation space of S at O to describe how the the dual graph of the minimal embedded resolution of C may be seen on the local tropicalization of S associated to this reembedding.

CONTENTS

| | |
|--|----|
| Introduction | 1 |
| 1. Toric resolutions of reduced curves singularities | 5 |
| 1.1. Toric modifications, their critical and discriminant locus | 5 |
| 1.2. Sufficient conditions for the existence of a toric embedded resolution | 6 |
| 1.3. Toric resolutions after reembedding | 9 |
| 2. Resolving a plane curve singularity with one toric morphism | 13 |
| 2.1. Background on plane curve singularities and divisorial valuations | 13 |
| 2.2. The embedding defined by the maximal contact curves | 18 |
| 3. Embedded resolution and local tropicalization | 22 |
| 3.1. The local tropicalization of S defined by the maximal contact curves of C | 22 |
| 3.2. Initial ideals associated to the local tropicalization and toric resolutions | 30 |
| 3.3. Minimal embedded resolution with one toric morphism | 33 |
| References | 37 |

INTRODUCTION

In [GT00], Goldin and Teissier proved that one can resolve the singularities of a plane branch (i.e., an analytically irreducible plane curve singularity) with one toric morphism, after reembedding it in a possibly higher dimensional affine space. Teissier asked then the following question in [Tei04, Section 5], see also [Tei03].

Date: October 22, 2021.

2010 Mathematics Subject Classification. 3A80,14E15,14E18,14M25.

Key words and phrases. divisorial valuations, curve singularities, generating sequences, resolution of singularities, toric geometry, local tropicalization, toric embedding.

Question. Given a reduced and equidimensional algebraic or formal space X over an algebraically closed field k , is it true that for every point $x \in X$ there is a local formal embedding of (X, x) into an affine space $(\mathbb{A}^m, 0)$ and a toric structure on \mathbb{A}^m such that $(X, x) \subset (\mathbb{A}^m, 0)$ can be resolved by one toric morphism ?

This means that there exist local coordinates u_1, \dots, u_m centered at x and an open (étale or formal) neighborhood U of $x \in \mathbb{A}^m(k)$, such that there is a proper birational toric map $\pi : Z \rightarrow U$ with respect to the coordinates u_1, \dots, u_m with Z non singular and such that the strict transform X' of $X \cap U$ is non singular and transversal to the non dense orbits at every point of $\pi^{-1}(x) \cap X'$. An embedding satisfying the property of the question will be called *torific*, a terminology which combines toric and terrific.

We consider also the problem of existence of *torific embeddings of a triple* $(x, X, \mathbb{A}^n(k))$ where $x \in X \subset \mathbb{A}^n(k)$, and n is the embedding dimension of $x \in X$, that is, if there exists a formal embedding of $\mathbb{A}^n(k)$ into an affine space $\mathbb{A}^m(k)$ endowed with a toric structure, such that a toric morphism of $\mathbb{A}^m(k)$ induces an embedded resolution of the triple $(x, X, \mathbb{A}^n(k))$.

Besides the case of plane branches there are some partial answers to Teissier's question:

- Aroca, Gómez-Morales, and Shabbir considered a notion of *Newton non-degenerate* ideals of the ring of polynomials $k[X_0, \dots, X_n]$ (see [AGS13]). This is a condition on the initial ideals with respect to a weight vector lying in the *tropicalization* of I . In this case, the original embedding of the singularity is already torific. The notion of Newton non-degenerate ideal is related to that of schön variety in [Tev07]. It is a generalization of the notions of hypersurface and complete intersection singularities which are non degenerate with respect to their Newton polyhedra, which was introduced by Khovanskii and Kouchnirenko [Kou76, Kho78]. Recently, Cueto, Popescu-Pampu and Stepanov have proven that the ideals defining the natural embeddings of splice type surface singularities are Newton non-degenerate (see [CPS21]).

- Lejeune-Jalabert and Reguera proved in [LJR99] that sandwiched surface singularities admit natural torific embeddings, which were called *toric environments*.

- González Pérez generalized Goldin and Teissier's result to the case of an irreducible germ of quasi-ordinary hypersurface singularity (see [GP03]).

- More generally, Tevelev proved that if k is an algebraically closed field of characteristic zero and $X \subset \mathbb{P}^n(k)$ is a projective algebraic variety, then there exists an integer $m \gg 0$ such that if $X \subset \mathbb{P}^m(k)$ is the Veronese reembedding of order m , there exist projective coordinates $(z_0 : \dots : z_m)$ on $\mathbb{P}^m(k)$ such that the intersection of X with the torus $T^m(k) = (k^*)^m$ defined by this choice of coordinates, is dense in X and there exists an equivariant map of toric varieties $\pi : Z \rightarrow \mathbb{P}^m(k)$, with Z non singular, such that the strict transform of X is non singular and transversal to the non dense toric orbits in Z (see [Tev14]). The proof of this result uses embedded resolution of singularities which is not known to be available in general when the field k has positive characteristic.

The main result of [GT00] motivates Teissier's strategy towards the proof of local uniformization, which is a very local version of resolution, by a method based on the comparison of a given singular germ by deformation with a space whose resolution is easy and blind to the characteristic, namely using toric methods (see [Tei09, Tei03, Tei14]). Note here that the valuative machinery that makes things work in [GT00], still works to some extent in higher dimension [CMT21] but does not lead to similar conclusions on resolution of singularities.

In addition, the resolution obtained by a torific embedding is often easier to use than classical Hironaka-type resolutions for computing subtle invariants of singularities like motivic or topological zeta functions, monodromy zeta functions, log canonical thresholds and jumping numbers of

multiplier ideals (see [M_VV21, G_GGR21, G_G14]). The resolutions of singularities obtained by the classical approach are often difficult to handle and it is a difficult problem to link invariants of resolution of singularities like Hironaka's order of ideals to subtle invariants of singularities such as those that we have just mentioned.

Let us explain the main contributions of this paper. Along this article, we assume that we are working over an algebraically closed field k of arbitrary characteristic.

In Section 1 we prove the existence of functions defining a toric embedding of a reduced curve singularity of arbitrary embedding dimension in two different ways (see Corollary 1.19 and Theorem 1.22).

In Section 2 we consider a reduced plane curve singularity C at a point O . We take a minimal generating sequence (x_0, \dots, x_m) of the divisorial valuations defined by those prime exceptional divisors of the minimal embedded resolution ψ of C which intersect the strict transform of C . We assume that the strict transform of the branch L_i defined by $x_i = 0$ does not intersect the strict transform of C by ψ , for $i = 0, \dots, m$. Then, we prove that the tuple of functions (x_0, x_1, \dots, x_m) defines a toric embedding of $(O, C, \mathbb{A}^{m+1}(k))$ (see Theorem 2.26).

In Section 3 we prove that the tuple of functions (x_0, \dots, x_m) considered above, defines toric embedding of the triple $(O, C, \mathbb{A}^2(k))$, that is, there exists a toric modification of $\mathbb{A}^{m+1}(k)$ which induces an embedded resolution of $(O, C, \mathbb{A}^2(k))$ (see Theorem 3.32). In addition, we show how the dual graph of the total transform of $\cup_{j=0}^m L_j$ by the minimal embedded resolution ψ of C appears on the support of the fan defining this modification.

Our main result realises the program of the geometric approach to resolution of singularities explained in [Mou20, Mou17]. This program suggests that one can find a toric embedding of a singularity (X, x) from first detecting a finite number of special divisorial valuations which may be called essential and then finding an embedding of X which is toric along each of these divisorial valuations. Given an essential valuation ν , finding an embedding of X which is toric along ν is a valuation theoretical issue; then a toric embedding may be found by a simple procedure (concatenation of the packages of functions defining the toric embeddings along each essential valuation); such a toric embedding is far from being minimal in general.

In this article, for a plane curve singularity, we find these essential valuations on the minimal embedded resolution of the curve and we characterize them by using the Eggers-Wall tree. We use the geometric characterization of the generating sequences of tuples of divisorial valuations by Spivakovskiy [Spi90, Section 8] in the case of one valuation and Delgado, Galindo, and Nuñez [DGN08] in the case of several valuations (see Proposition 2.11). The embedding that we find has no superfluous information as in the concatenation procedure, it is actually minimal for the triple (O, C, \mathbb{A}^2) . One can also detect the essential valuations on a graph which is associated with the jet schemes of the curve singularity [Mou11, LJMR13]. This latter graph makes sense also for higher dimensional singularities [CM21] and the *loc.cit.* program suggests that one can detect essential valuations on it [Mou20].

In order to prove the main results in Section 3 we consider the *local tropicalization* of $\mathbb{A}^2(k)$ associated with the tuple of functions (x_0, \dots, x_m) . The definition and properties of local tropicalizations in a general setting was developed in [PS13] by Popescu-Pampu and Stepanov. Further applications of local tropicalization are obtained recently by Cueto, Popescu-Pampu, and Stepanov in the case of surface singularities of splice type (see [CPS21]).

The *finite local tropicalization* of $\mathbb{A}^2(k)$ associated with the tuple of functions (x_0, \dots, x_m) consists of the tuples $(\nu(x_0), \dots, \nu(x_m)) \in \mathbb{R}_{\geq 0}^m$, where ν runs through the semivaluations of $\mathbb{A}^2(k)$ at the closed point O of C , which take finite values on x_0, \dots, x_m . We show that this set is the

support of a fan \mathcal{T} in \mathbb{R}^{m+1} consisting of at most two dimensional cones (see Theorem 3.17). For instance, if E is an exceptional prime divisor in the minimal embedded resolution of ψ of C and if ν_E denotes its associated divisorial valuation then $(\nu_E(x_0), \dots, \nu_E(x_m)) \in \mathbb{Z}_{>0}^{m+1}$ belongs to the support of \mathcal{T} . Then, we study the initial ideals associated to the ideal I defining the embedding $\mathbb{A}^2(k) \hookrightarrow \mathbb{A}^{m+1}(k)$ with respect to weight vectors w in the support of \mathcal{T} . The key step is to prove that the ideal I is Newton non degenerate (see Proposition 3.31). This is shown by using the properties of expansions in terms of generating sequences. The main theorem of this section is obtained by applying the result of [AGS13] mentioned above. We show that if Y denotes the image of $\mathbb{A}^2(k)$ after reembedding, then there exists a toric modification π of $\mathbb{A}^{m+1}(k)$, defined by a regular fan supported on $\mathbb{R}_{\geq 0}^{m+1}$ which induces a regularization of the fan \mathcal{T} , and such that the restriction of π to the strict transform Y' of Y is an embedded resolution of C (see Theorem 3.32). If in addition the regularization of the fan \mathcal{T} is the minimal one, then the restriction of π to Y' is the minimal embedded resolution of C (see Theorem 3.39). In particular, we prove that the projectivization of the minimal regularization of the fan \mathcal{T} is isomorphic to the dual graph of the total transform of $\cup_{j=0}^m L_j$ by the minimal embedded resolution ψ of C (see Proposition 3.38).

The description of the fan \mathcal{T} is very explicit in terms of the embedding of the *Eggers-Wall tree* of C with respect to the smooth branch L_0 defined by $x_0 = 0$ in the *space of normalized semivaluations* \mathcal{V}_{L_0} with respect to L_0 . This embedding was described by García Barroso, Popescu-Pampu, and González Pérez (see [GGP19a, Section 8] and [GGP19b, Remark 5.37]). The space \mathcal{V}_{L_0} is one of the valuative trees considered in Favre and Jonsson's book [FJ04], see also the presentation of this theory in Jonsson's survey [Jon15, section 7].

Besides providing an answer to Teissier's question for plane curve singularities, our results exhibit the beautiful interplay between semivaluation spaces, resolution of singularities, toric geometry and tropical geometry, which may shed some light on higher dimensional generalizations.

Notation. A curve C is an affine scheme $\text{Spec } \mathcal{O}$, where \mathcal{O} is a complete noetherian local ring of dimension one over k with residue field k . We call the ring \mathcal{O} the local ring of C and denote by \mathcal{M} the maximal ideal of \mathcal{O} . The curve C is reduced if the local ring \mathcal{O} is reduced. If \mathcal{O} is an integral domain, the curve C is said to be a branch. The local ring \mathcal{O} of a reduced curve C has a finite number of minimal prime ideals, say \mathcal{P}_j for $j = 1, \dots, r$, each one defining a branch C_j with local ring $\mathcal{O}/\mathcal{P}_j$. Then we write $C = \cup_{j=1}^r C_j$ and say that C is a curve with r branches. The embedding dimension of a curve C is the dimension of $\mathcal{M}/\mathcal{M}^2$ as k -vector space. A plane curve is a curve of embedding dimension at most two.

Acknowledgment. We are grateful to Javier Fernández de Bobadilla for sharing some ideas which were helpful in the formulation of Theorem 1.22. We thank Bernard Teissier and Alicia Dickenstein for useful discussions and suggestions, and Patrick Popescu-Pampu for his comments on a preliminary version of the paper.

The first author was supported by ERCEA Consolidator Grant 615655-NMST, the Basque Government through the BERC 2018-2021 program, BCAM Severo Ochoa excellence accreditation SEV-2017-0718, grant MTM2016-80659-P, and grant PID2019-105896GB-I00 funded by MCIN/AEI/10.13039/501100011033. The second author was supported by the Spanish grants PID2020-114750GB-C32 and MTM2016-76868-C2-1-P. The third author was supported by the French grant Projet ANR LISA, ANR-17-CE40-0023. We also thank the hospitality of the Institute of Mathematics of the University of Barcelona (IMUB) and the Institut de Mathématiques de Jussieu-Paris Rive Gauche (IMJ-PRG).

1. TORIC RESOLUTIONS OF REDUCED CURVES SINGULARITIES

1.1. Toric modifications, their critical and discriminant locus. In this section we describe the modification induced by an ambient toric modification on an orbit closure in a toric variety. See the textbooks [Ful93, CLS11, Ewa96, Oda88] for standard notions about toric geometry.

Let Σ be a fan with respect to a rank n lattice N , with dual lattice M . If $\sigma \subset N_{\mathbb{R}} := N \otimes \mathbb{R}$ is a cone we denote by $\check{\sigma} \subset M_{\mathbb{R}} := M \otimes \mathbb{R}$ the dual cone, and by σ^{\perp} the orthogonal cone.

If $\sigma \in \Sigma$ then the semigroup $\check{\sigma} \cap M$ is finitely generated and the semigroup algebra

$$k[\check{\sigma} \cap M] = \left\{ \sum_{finite} a_v \chi^v \mid a_v \in k, v \in \check{\sigma} \cap M \right\}$$

is a k -algebra of finite type. It is the coordinate ring of the affine toric variety X_{σ} .

In addition, if the cone σ is of dimension n , then there is a unique minimal system of generators $v_1, \dots, v_{m_{\sigma}}$ of the semigroup $\check{\sigma} \cap M$. Setting, $x_i := \chi^{v_i}$ for $i = 1, \dots, m_{\sigma}$, defines an embedding of the affine toric variety

$$(1.1) \quad X_{\sigma} \hookrightarrow k^{m_{\sigma}},$$

which sends the 0-dimensional orbit to the origin of $k^{m_{\sigma}}$. In this case, the defining ideal of this 0-dimensional orbit is generated by the monomials χ^v , for $v \in \check{\sigma} \cap M \setminus \{0\}$. The completion of $k[\check{\sigma} \cap M]$ with respect to this ideal is the formal power series ring

$$k[[\check{\sigma} \cap M]] = \left\{ \sum a_v \chi^v \mid a_v \in k, v \in \check{\sigma} \cap M \right\}.$$

We denote by $X_{\Sigma, N}$ the normal toric variety associated with the fan Σ and the lattice N , or simply by X_{Σ} , if the lattice N is clear from the context. If $\sigma \in \Sigma$ we denote by X_{σ} the affine open toric subvariety of X_{Σ} , and by $O(\sigma)_{\Sigma}$, or simply by $O(\sigma)$, the corresponding orbit by the action on X_{Σ} of the torus $T_N = \text{Hom}(M, k^*)$. Abstractly, the orbit $O(\sigma)$ is the torus $\text{Hom}(M(\sigma), k^*)$, where $M(\sigma)$ denotes the lattice spanned by $\sigma^{\perp} \cap M$. In particular, we have $T_N = O(\{0\})$. The map which sends a cone $\sigma \in \Sigma$ to its corresponding orbit $O(\sigma)$ defines a bijection of Σ with the set of orbits of the torus action on X_{Σ} .

The closure of the orbit $O(\sigma)$ in X_{Σ} is a normal toric variety. In order to describe its associated fan, we consider the lattices N_{σ} spanned by $\sigma \cap N$ and the quotient $N(\sigma) := N/N_{\sigma}$. The dual lattice of $N(\sigma)$ is precisely the lattice $M(\sigma)$ introduced before. Denote by $\text{Star}(\sigma)_{\Sigma}$ the set of cones in $N(\sigma)_{\mathbb{R}}$ which are images by the canonical projection $N_{\mathbb{R}} \rightarrow N(\sigma)_{\mathbb{R}} = N_{\mathbb{R}}/(N_{\sigma})_{\mathbb{R}}$ of the cones of the fan Σ which contain σ as a face. The set $\text{Star}(\sigma)_{\Sigma}$ is a fan with respect to the lattice $N(\sigma)$.

Lemma 1.1. (see [Ful93, Section 3.1]) *The closure of the orbit $O(\sigma)$ in X_{Σ} is the normal toric variety $X_{\text{Star}(\sigma)_{\Sigma}, N(\sigma)}$ associated with the fan $\text{Star}(\sigma)_{\Sigma}$ and the lattice $N(\sigma)$.*

Remark 1.2. The torus $T_{N(\sigma)} = \text{Hom}(M(\sigma), k^*)$ of the toric variety $X_{\text{Star}(\sigma)_{\Sigma}, N(\sigma)}$ is equal to $O(\sigma)$. More generally, the orbits of the action of the torus $T_{N(\sigma)}$ on $X_{\text{Star}(\sigma)_{\Sigma}, N(\sigma)}$ are orbits of the action of T_N on X_{Σ} .

Notation 1.3. We denote by $\Sigma_{0, n}$ the fan consisting of the faces of the cone $\sigma_{0, n} := \mathbb{R}_{\geq 0}^n$, with respect to the lattice $N := \mathbb{Z}^n$. We often write simply $\sigma_0 := \sigma_{0, n}$ and $\Sigma_0 := \Sigma_{0, n}$, if n is clear from the context. We denote by e_1, \dots, e_n the canonical basis of the lattice N , which spans the cone $\sigma_{0, n}$. We denote by $\check{e}_1, \dots, \check{e}_n$ the dual basis of M , which spans the dual cone $\check{\sigma}_{0, n}$. The toric variety $X_{\Sigma_{0, n}}$ is isomorphic to k^n by (1.1). It is equipped with coordinates $(X_1, \dots, X_n) := (\chi^{\check{e}_1}, \dots, \chi^{\check{e}_n})$.

Example 1.4. Let us fix an integer s with $1 \leq s \leq n-1$. The cone $\sigma := \mathbb{R}_{\geq 0}e_{s+1} + \cdots + \mathbb{R}_{\geq 0}e_n$ belongs to the fan $\Sigma_{0,n}$ (see Notation 1.3). The lattice N_σ has basis e_{s+1}, \dots, e_n . Let us denote by $\bar{u} \in N(\sigma)$ the image of a vector $u \in N$ by the canonical projection $N \rightarrow N(\sigma)$. Then, the lattice $N(\sigma)$ has basis $\bar{e}_1, \dots, \bar{e}_s$. Thus, the closure of the orbit $O(\sigma)$ is the coordinate subspace $k^s \hookrightarrow k^n$ with coordinates (X_1, \dots, X_s) , and defined by $X_{s+1} = \cdots = X_n = 0$.

Let Σ' be a fan subdividing Σ with respect to the lattice N . We denote by $\pi_{\Sigma'}^{\Sigma'}: X_{\Sigma'} \rightarrow X_\Sigma$ the associated toric *modification*. The *exceptional locus* of the modification $\pi_{\Sigma'}^{\Sigma'}$ consists of those subvarieties of $X_{\Sigma'}$ which are mapped to subvarieties of X_Σ of smaller dimension. The *discriminant locus* of the modification $\pi_{\Sigma'}^{\Sigma'}$ is the image of the exceptional locus. These loci can be described in terms of orbits as follows (see [GP03, Section 1.1] and also [GSLJ91, Page 100]).

Proposition 1.5. *The exceptional locus $\text{Exc}(\pi_{\Sigma'}^{\Sigma'})$ of $\pi_{\Sigma'}^{\Sigma'}$ is the union of orbits $O(\sigma)_{\Sigma'}$, for $\sigma \in \Sigma' \setminus \Sigma$. The discriminant locus of $\pi_{\Sigma'}^{\Sigma'}$ is the union of orbits $O(\tau)_\Sigma$, for $\tau \in \Sigma \setminus \Sigma'$.*

Let us fix a cone $\sigma \in \Sigma$. If σ belongs also to Σ' then the orbit $O(\sigma)_\Sigma \subset X_\Sigma$ is not contained in the discriminant locus of $\pi_{\Sigma'}^{\Sigma'}$. We have also an orbit $O(\sigma)_{\Sigma'} \subset X_{\Sigma'}$ and the restriction of $\pi_{\Sigma'}^{\Sigma'}$ defines an isomorphism of orbits $O(\sigma)_{\Sigma'} \rightarrow O(\sigma)_\Sigma$. The fan $\text{Star}(\sigma)_{\Sigma'}$ is a subdivision of $\text{Star}(\sigma)_\Sigma$. The following lemma is a particular case of Lemma 3.3.21 of [CLS11].

Lemma 1.6. *Let Σ' be a subdivision of the fan Σ . If $\sigma \in \Sigma' \cap \Sigma$ then the restriction of $\pi_{\Sigma'}^{\Sigma'}$ to the closure of $O(\sigma)_{\Sigma'}$ in the source and to the closure $O(\sigma)_\Sigma$ in the target is the toric modification*

$$(1.2) \quad \pi_{\text{Star}(\sigma)_\Sigma}^{\text{Star}(\sigma)_{\Sigma'}}: X_{\text{Star}(\sigma)_{\Sigma'}} \rightarrow X_{\text{Star}(\sigma)_\Sigma}.$$

Example 1.7. Let us specialize to the case $n = 3$ in Notation 1.3. Consider the subdivision Σ of Σ_0 at the vector $w = (4, 6, 13)$. It has three-dimensional cones $\sigma_{i,j} = \mathbb{R}_{\geq 0}e_i + \mathbb{R}_{\geq 0}e_j + \mathbb{R}_{\geq 0}w$, for $1 \leq i < j \leq 3$. The exceptional locus of $\pi_{\Sigma_0}^\Sigma$ is the union of orbits $O(\sigma)_\Sigma$ such that the relative interior of the cone σ is contained in $\mathbb{R}_{> 0}^3$, while the discriminant locus of $\pi_{\Sigma_0}^\Sigma$ is reduced to the origin, that is, the 0-dimensional orbit. Let us consider the cone $\sigma = \mathbb{R}_{\geq 0}e_3$. The closure of the orbit $O(\sigma)_{\Sigma_0}$ is the coordinate plane of k^3 defined by $X_3 = 0$. Abstractly, it is the toric variety defined by the cone $\mathbb{R}_{\geq 0}^2 = \mathbb{R}_{\geq 0}\bar{e}_1 + \mathbb{R}_{\geq 0}\bar{e}_2$, with respect to the lattice $N(\sigma) = \mathbb{Z}\bar{e}_1 + \mathbb{Z}\bar{e}_2$, where we use the notation of Example 1.4. The only three-dimensional cones of Σ which contain σ are $\sigma_{1,3}$ and $\sigma_{2,3}$. These two cones intersect along the cone $\mathbb{R}_{\geq 0}w + \mathbb{R}_{\geq 0}e_3$. The fan $\text{Star}(\sigma)_\Sigma$ is the subdivision of the cone $\mathbb{R}_{\geq 0}^2$ along the ray spanned by $2\bar{e}_1 + 3\bar{e}_2$. This ray is the image of $\mathbb{R}_{\geq 0}w + \mathbb{R}_{\geq 0}e_3$ under the canonical projection $N_\mathbb{R} = \mathbb{R}^3 \rightarrow N(\sigma)_\mathbb{R} = \mathbb{R}^2$, which sends $(a_1, a_2, a_3) \mapsto (a_1, a_2)$.

1.2. Sufficient conditions for the existence of a toric embedded resolution. Let C be a reduced curve, not necessarily a plane one. We denote by \mathcal{O} its local ring, with maximal ideal \mathcal{M} . If x_1, \dots, x_n is a set of generators of the maximal ideal \mathcal{M} , then we have a surjection

$$(1.3) \quad k[[X_1, \dots, X_n]] \rightarrow \mathcal{O}, \text{ such that } X_i \mapsto x_i, \text{ for } i = 1, \dots, n.$$

This surjection defines an embedding of germs $C \hookrightarrow k^n$, which maps the special point of C to the origin of k^n . In this section, we give sufficient conditions on the functions x_1, \dots, x_n in order to guarantee that an embedded resolution of a branch $C \subset k^n$ can be obtained by a toric modification of k^n .

In what follows we consider k^n as the affine toric variety X_{Σ_0} and use Notation 1.3. The following definition is slightly more general than the one considered in [GT00].

Definition 1.8. Let C be a reduced curve embedded in k^n by the map (1.3). A toric modification $\pi_{\Sigma_0}^{\Sigma} : X_{\Sigma} \rightarrow X_{\Sigma_0} = k^n$ defined by a regular subdivision Σ of Σ_0 is a *toric embedded resolution* of C if no branch of C is contained in the discriminant locus of $\pi_{\Sigma_0}^{\Sigma}$, and the strict transform \tilde{C} of C is smooth and transversal to the orbit stratification of the exceptional locus of $\pi_{\Sigma_0}^{\Sigma}$.

Recall that the strict transform \tilde{C} of C by $\pi_{\Sigma_0}^{\Sigma}$ is the closure in X_{Σ} of $(\pi_{\Sigma_0}^{\Sigma})^{-1}(C) \setminus \text{Exc}(\pi_{\Sigma_0}^{\Sigma})$.

Remark 1.9. In the paper [GT00] the case when the curve C is contained in a coordinate hyperplane was not considered. In order to deal with this case we introduced the hypothesis about the discriminant locus of the modification in Definition 1.8. The transversality condition in Definition 1.8 should be considered with respect to the modifications of the minimal coordinate subspaces containing the branches of C , which are toric morphisms by Lemma 1.6. Notice that the minimal coordinate subspace of $X_{\Sigma_0} = k^n$ which contains a branch C_j of C is the closure $X_{\text{Star}(\sigma)_{\Sigma_0}}$ of an orbit $O(\sigma)_{\Sigma_0}$, for some $\sigma \in \Sigma_0$ (see Example 1.4). By Proposition 1.5 this coordinate subspace is not contained in the discriminant locus of $\pi_{\Sigma_0}^{\Sigma}$ if and only if $\sigma \in \Sigma$. Then, the toric modification $\pi_{\Sigma_0}^{\Sigma}$ is a toric embedded resolution of C_j according to Definition 1.8 means that the map $\pi_{\text{Star}(\sigma)_{\Sigma_0}}^{\text{Star}(\sigma)_{\Sigma}}$, defined in (1.2), is a toric embedded resolution of $C_j \subset X_{\text{Star}(\sigma)_{\Sigma_0}}$ according to the definition considered in [GT00].

Before stating our results we need to introduce preliminary materials on arcs. See [Ish04] for more on arcs on toric varieties.

An *arc* on X_{Σ_0} is a morphism $\eta : \text{Spec } k[[t]] \rightarrow X_{\Sigma_0}$, which corresponds to a homomorphism of k -algebras $\eta^* : k[\check{\sigma}_0 \cap M] \rightarrow k[[t]]$. The arc η has its *generic point in the torus* of k^n if the series $\chi^v \circ \eta := \eta^*(\chi^v)$ is a nonzero element of $k[[t]]$ for any $v \in \check{\sigma}_0 \cap M$. The arc is *centered at the origin* of k^n if $\chi^v \circ \eta$ belongs to $(t)k[[t]]$, for any nonzero $v \in \check{\sigma}_0 \cap M$. In this case, the arc η induces a map of local k -algebras $k[[\check{\sigma}_0 \cap M]] \rightarrow k[[t]]$ which we denote also by η^* .

If η has its generic point in the torus T_N , then it has an associated *order vector* $\text{ord}(\eta) \in N$. It is defined as the lattice homomorphism $M \rightarrow \mathbb{Z}$ which sends $v \in M$ to the order of the Laurent series $\chi^v \circ \eta \in k((t))$. Notice that $\chi^v \circ \eta \in k[[t]]$, for any $v \in \check{\sigma}_0 \cap M$. This implies that $\text{ord}(\eta) \in \sigma_0$. The coordinates of the order vector $\text{ord}(\eta)$ with respect to the basis e_1, \dots, e_n are equal to

$$(\text{ord}_t(X_1 \circ \eta), \dots, \text{ord}_t(X_n \circ \eta)).$$

Definition 1.10. Let C be a branch embedded in k^n by the map (1.3). We associate to C an arc $\eta_C : \bar{C} \rightarrow k^n$ which is the composition a normalization $\bar{C} \rightarrow C$ of C with the inclusion $C \hookrightarrow k^n$.

In terms of the coordinates (X_1, \dots, X_n) of k^n and a regular parameter t of \bar{C} , the arc η_C in Definition 1.10 corresponds to a local homomorphism of k -algebras

$$\eta_C^* : k[[X_1, \dots, X_n]] = k[[\check{\sigma}_0 \cap M]] \rightarrow k[[t]],$$

whose kernel is the *defining ideal* I_C of the branch C . By definition, the arc η_C is centered at the origin of k^n . If in addition, η_C has its generic point in the torus of k^n , then $\text{ord}(\eta_C)$ belongs to the interior of σ_0 .

1.2.1. *The case when η_C has its generic point in the torus.* We start by discussing what happens to the order vector of an arbitrary arc η on X_{Σ_0} with generic point in the torus, when a toric modification is made.

Remark 1.11. Let Σ be a regular subdivision of Σ_0 . By the valuative criterion of properness there is a unique arc $\tilde{\eta} : \text{Spec } k[[t]] \rightarrow X_{\Sigma}$ such that $\pi_{\Sigma_0}^{\Sigma} \circ \tilde{\eta} = \eta$. Since the modification $\pi_{\Sigma_0}^{\Sigma}$ is the identity on the torus, the lifted arc $\tilde{\eta}$ has also its generic point in the torus T_N and by

definition we get $\text{ord}(\eta) = \text{ord}(\tilde{\eta})$. Let $\sigma \in \Sigma$ be a cone of dimension n which contains the order vector $\text{ord}(\eta)$. Denote by v_1, \dots, v_n the basis of the lattice N which spans the regular cone σ . The affine toric variety X_σ is isomorphic to k^n with coordinates $(U_1, \dots, U_n) = (\chi^{\check{v}_1}, \dots, \chi^{\check{v}_n})$, where $\check{v}_1, \dots, \check{v}_n$ is the dual basis of v_1, \dots, v_n . Then, it follows that the coordinates of the vector $\text{ord}(\tilde{\eta})$ with respect to the basis v_1, \dots, v_n of N are equal to $(\text{ord}_t(U_1 \circ \tilde{\eta}), \dots, \text{ord}_t(U_n \circ \tilde{\eta}))$.

The following proposition provides a sufficient condition to guarantee the existence of a toric embedded resolution of the branch C , when η_C has generic point in the torus of k^n .

Proposition 1.12. *Let C be a branch embedded in k^n by the map (1.3). If the arc η_C has its generic point in the torus of k^n and $\text{ord}(\eta_C)$ is primitive with respect to the lattice N , then any regular subdivision Σ of the fan Σ_0 such that the ray ρ_C spanned by $\text{ord}(\eta_C)$ belongs to Σ defines a toric embedded resolution of C .*

Proof. Let us denote the toric modification $\pi_{\Sigma_0}^\Sigma$ simply by π and the arc η_C by η . Let $\tilde{\eta}$ be the unique lifting of η to X_Σ . Then, the strict transform \tilde{C} of C is equal to the image of $\tilde{\eta}$, and $\tilde{\eta}$ is the composition of the normalization of \tilde{C} with the inclusion $\tilde{C} \hookrightarrow X_\Sigma$.

Since the order vector $\text{ord}(\eta)$ is contained in the interior of the cone σ_0 , Proposition 1.5 implies that the orbit $O(\rho_C)$ is contained in the exceptional locus of π .

Let σ be a regular cone of Σ spanned by a basis v_1, \dots, v_n of N such that $v_1 = \text{ord}(\eta)$. Notice that we use here the assumption that $\text{ord}(\eta)$ is primitive in the lattice N . The affine toric variety X_σ is isomorphic with k^n with coordinates (U_1, \dots, U_n) . Then, it follows from Remark 1.11, that $\text{ord}(\eta) = \text{ord}(\tilde{\eta})$ and

$$(1.4) \quad (\text{ord}_t(U_1 \circ \tilde{\eta}), \dots, \text{ord}_t(U_n \circ \tilde{\eta})) = (1, 0, \dots, 0),$$

since $(1, 0, \dots, 0)$ is the vector of coordinates of $\text{ord}(\eta)$ with respect to the basis v_1, \dots, v_n . The parametrization $\tilde{\eta}$ of \tilde{C} , in terms of the coordinates (U_1, \dots, U_n) , is given by $(U_1 \circ \tilde{\eta}, \dots, U_n \circ \tilde{\eta})$. Therefore, the condition (1.4) implies that the strict transform \tilde{C} of C is smooth and transversal to the orbit $O(\rho_C)$. \square

1.2.2. *The case when η_C does not have generic point in the torus.* We now deal with the generalization of Proposition 1.12 to the case when η_C does not have generic point in the torus of k^n . Then, some of the series $X_j \circ \eta$ are identically zero, say for $j = s + 1, \dots, n$. The smaller coordinate subspace of k^n which contains the branch C is $X_{s+1} = \dots = X_n = 0$. We are in the situation of Remark 1.9. If $\sigma = \mathbb{R}_{\geq 0}e_{s+1} + \dots + \mathbb{R}_{\geq 0}e_n$, then the arc η_C has its generic point in the torus $O(\sigma)_{\Sigma_0}$, η_C factors through the orbit closure $X_{\text{Star}(\sigma)_{\Sigma_0}} = k^s$, and it is centered at the origin of k^s . That is, we have $X_j \circ \eta_C \in (t)k[[t]] \setminus \{0\}$, for $j = 1, \dots, s$. The order vector $\text{ord}(\eta_C)$ is an element of the lattice $N(\sigma)$ which belongs to the interior of the image of the cone σ_0 under the canonical projection $N_{\mathbb{R}} \rightarrow N(\sigma)_{\mathbb{R}}$ (see Example 1.4 and Remark 1.11).

Proposition 1.13. *With the previous hypothesis and notation, let Σ be a regular fan subdividing Σ_0 such that $\sigma \in \Sigma$. Assume that $\text{ord}(\eta_C)$ is a primitive vector in the lattice $N(\sigma)$ and it spans a ray ρ_C of the fan $\text{Star}(\sigma)_{\Sigma}$. Then, $\pi_{\Sigma_0}^\Sigma$ is a toric embedded resolution of C .*

Proof. Since $\sigma \in \Sigma$ the image of η_C , which is equal to the branch C , is not contained in the discriminant locus of π (see Proposition 1.5). This implies that the strict transform \tilde{C} of C exists. Recall that η_C factors through $X_{\text{Star}(\sigma)_{\Sigma_0}} = k^s$, has generic point in the torus of this variety, and it is centered at the origin of k^s . Proposition 1.12 implies that the modification (1.2) is a toric embedded resolution of $C \subset X_{\text{Star}(\sigma)_{\Sigma_0}}$. By Remark 1.9 this implies the result. \square

Example 1.14. We keep the notations of Example 1.7. Let C_1, C_2 be the branches in k^3 given by $\eta_{C_1}^*(X_1) = t^4$, $\eta_{C_1}^*(X_2) = t^6$, $\eta_{C_1}^*(X_3) = t^{13}$, and $\eta_{C_2}^*(X_1) = t^2$, $\eta_{C_2}^*(X_2) = t^3$, $\eta_{C_2}^*(X_3) = 0$.

If Σ' is any regular subdivision of Σ , then by Proposition 1.12 the toric modification defined by Σ' is an embedded resolution of C_1 . The branch C_2 is contained in the closure of the orbit of k^3 associated with $\sigma = \mathbb{R}_{\geq 0}e_3$. The cone spanned by $2\bar{e}_1 + 3\bar{e}_2 \in N(\sigma)$ belongs to the fan $\text{Star}(\sigma)_{\Sigma'}$. The cone $\sigma \in \Sigma_0$ belongs also to Σ' . Since the order vector of C_2 is the primitive vector $2\bar{e}_1 + 3\bar{e}_2$, the modification Σ' induces also an embedded resolution of C_2 by Proposition 1.13. Therefore, in this case $\pi_{\Sigma'_0}^{\Sigma'}$ is a toric embedded resolution of $C_1 \cup C_2$.

1.3. Toric resolutions after reembedding. In this section we prove that certain choices of functions define suitable *torific embeddings* for a reduced curve singularity. We mean by this that the curve singularity can be resolved by one toric modification, after reembedding it in an affine space of possibly higher dimension. This result is a generalization of a theorem of Goldin and Teissier, see [GT00, Th. 6.1, Corollary].

Definition 1.15. Let C be a branch embedded in k^n by the map (1.3). The semigroup Γ_C of the branch C is $\Gamma_C = \{\text{ord}_t(h \circ \eta_C) \mid h \in k[[X_1, \dots, X_n]], h \notin I_C\}$.

The semigroup Γ_C is a finitely generated semigroup of $(\mathbb{Z}_{\geq 0}, +)$ (see [Zar06]).

Lemma 1.16. Let C be a branch embedded in k^n by the map (1.3). Let y_1, \dots, y_m be elements of the maximal ideal of $k[[X_1, \dots, X_n]]$ such that:

- (1) $y_1, \dots, y_m \notin I_C$,
- (2) the numbers $\text{ord}_t(y_j \circ \eta_C) \in \mathbb{Z}_{>0}$, $j = 1, \dots, m$, generate the semigroup Γ_C .

Then, the vector $(\text{ord}_t(y_1 \circ \eta_C), \dots, \text{ord}_t(y_m \circ \eta_C))$ is primitive in the lattice \mathbb{Z}^m .

Proof. Since the local homomorphism $\mathcal{O} \rightarrow k[[t]]$ corresponding to the normalization $\bar{C} \rightarrow C$ induces an isomorphism between the fields of fractions, we must have that the group generated by Γ_C is equal to \mathbb{Z} . This is equivalent to the condition $\text{gcd}(\text{ord}_t(y_1 \circ \eta_C), \dots, \text{ord}_t(y_m \circ \eta_C)) = 1$. \square

Let y_1, \dots, y_m be generators of the maximal ideal of $k[[X_1, \dots, X_n]]$. Then, the surjection

$$k[[Y_1, \dots, Y_m]] \rightarrow k[[X_1, \dots, X_n]] \text{ such that } Y_j \mapsto y_j, \text{ for } j = 1, \dots, m.$$

defines an embedding

$$(1.5) \quad k^n \hookrightarrow k^m.$$

If C is a reduced curve embedded in k^n by the map (1.3) then, by composing the embedding $C \hookrightarrow k^n$ with (1.5), we get an embedding

$$(1.6) \quad C \hookrightarrow k^m.$$

If C_j is a branch of $C = \cup_{l=1}^r C_l$, then $C \hookrightarrow k^m$ induces an embedding $C_j \hookrightarrow k^m$. We denote by $\eta_j: \text{Spec } k[[t_j]] \rightarrow k^m$ the composition of η_{C_j} with the map (1.5). It corresponds to a local homomorphism of k -algebras

$$\eta_j^*: k[[Y_1, \dots, Y_m]] \rightarrow k[[t_j]], \text{ such that } Y_s \mapsto y_s \circ \eta_{C_j}, \text{ for } s = 1, \dots, m.$$

Corollary 1.17. Let $C = \cup_{j=1}^r C_j$ be a reduced curve with $r \geq 1$ branches embedded in k^n by the map (1.3). Let $y_1, \dots, y_m \in k[[X_1, \dots, X_n]]$ be such that:

- (1) y_1, \dots, y_m generate the maximal ideal of $k[[X_1, \dots, X_n]]$,
- (2) $y_1, \dots, y_m \notin I_{C_j}$, for $1 \leq j \leq r$,
- (3) the numbers $\text{ord}_{t_j}(y_s \circ \eta_{C_j}) \in \mathbb{Z}_{>0}$, $s = 1, \dots, m$, generate the group \mathbb{Z} , for $1 \leq j \leq r$.

Let $C \hookrightarrow k^m$ be the embedding defined as (1.6) with respect to y_1, \dots, y_m . Let Σ be a regular subdivision of the fan $\Sigma_{0,m}$ such that the ray ρ_j spanned by $\text{ord}(\eta_j)$ belongs to Σ , for $1 \leq j \leq r$. Denote by \tilde{C}_j the strict transform of C_j under $\pi_{\Sigma_{0,m}}^{\Sigma}$. If the special points of \tilde{C}_j , for $j = 1, \dots, r$, are pairwise different, then the toric modification $\pi_{\Sigma_{0,m}}^{\Sigma}$ is a toric embedded resolution of $C \subset k^m$.

Proof. For $1 \leq j \leq r$, by condition (2), the arc η_j has its generic point in the torus of k^m . Note that the coordinates of the vector $\text{ord}(\eta_j)$ with respect to the canonical basis e_1, \dots, e_m of $N = \mathbb{Z}^m$ (see Notation 1.3) are equal to

$$(\text{ord}_{t_j}(Y_1 \circ \eta_j), \dots, \text{ord}_{t_j}(Y_m \circ \eta_j)) = (\text{ord}_{t_j}(y_1 \circ \eta_{C_j}), \dots, \text{ord}_{t_j}(y_m \circ \eta_{C_j})).$$

The vector $\text{ord}(\eta_j)$ is primitive in the lattice N by condition (3). Then, we apply Proposition 1.12 to the branch C_j embedded in k^m by (1.6) with respect to y_1, \dots, y_m , for $1 \leq j \leq r$. It follows that the strict transform \tilde{C}_j of C_j is smooth and transversal to the orbit $O(\rho_j)_\Sigma$, which is contained in the exceptional locus of $\pi_{\Sigma_{0,m}}^\Sigma$. Since the special points of \tilde{C}_j are pairwise different, the branches \tilde{C}_j and $\tilde{C}_{j'}$ do not intersect if $j \neq j'$. \square

Remark 1.18. Let y_1, \dots, y_m be generators of the maximal ideal of $k[[X_1, \dots, X_n]]$ such that for any $1 \leq j \leq r$ the semigroup Γ_{C_j} is generated by the numbers $\text{ord}_{t_j}(y_\ell \circ \eta_{C_j})$, for $1 \leq \ell \leq m$ such that $\text{ord}_{t_j}(y_\ell \circ \eta_{C_j}) \neq \infty$. We can choose such functions in such a way that the three conditions of Corollary 1.17 are satisfied. The condition (1) holds by assumption. Let us choose a function y in the maximal ideal of $k[[X_1, \dots, X_n]]$ such that $y \notin I_{C_j}$ for $1 \leq j \leq r$ and take an integer $p > 0$. If p is large enough then for any $1 \leq j \leq r$ we have that:

$$\text{ord}_{t_j}((y_i + y^p) \circ \eta_{C_j}) = \begin{cases} \text{ord}_{t_j}(y_i \circ \eta_{C_j}) & \text{if } \text{ord}_{t_j}(y_i \circ \eta_{C_j}) \in \mathbb{N} \\ \text{ord}_{t_j}(y^p \circ \eta_{C_j}) & \text{if } \text{ord}_{t_j}(y_i \circ \eta_{C_j}) = \infty. \end{cases}$$

Thus, if we replace y_ℓ by $y_\ell + y^p$, for $1 \leq \ell \leq m$, conditions (2) and (3) of Corollary 1.17 are also satisfied (see Lemma 1.16).

Corollary 1.19. *Let $C = \cup_{j=1}^r C_j$ be a reduced curve with $r \geq 1$ branches embedded in k^n by the map (1.3). There exist $m \geq 1$ and functions y_1, \dots, y_m such that the conditions (1), (2) and (3) of Corollary 1.17 are satisfied and in addition, the vectors*

$$(1.7) \quad (\text{ord}_{t_j}(y_1 \circ \eta_{C_j}), \dots, \text{ord}_{t_j}(y_m \circ \eta_{C_j})), \text{ for } j = 1, \dots, r,$$

are pairwise different. Then, the modification $\pi_{\Sigma_{0,m}}^\Sigma$ of Corollary 1.17 is a toric embedded resolution of $C \subset k^m$.

Proof. Assume first that we are given functions y_1, \dots, y_m verifying the conditions of Corollary 1.17 and such that the vectors (1.7) are pairwise different. This implies that the cones ρ_1, \dots, ρ_r , and their orbits, are also pairwise different. Since the special point of \tilde{C}_j belongs to the orbit $O(\rho_j)_\Sigma$, for $j = 1, \dots, r$, these special points are also pairwise different. By Corollary 1.17 the modification $\pi_{\Sigma_{0,m}}^\Sigma$ is a toric embedded resolution of $C \subset k^m$.

Let us take functions $y_1, \dots, y_{m'}$ such that the conditions (1), (2) and (3) of Corollary 1.17 are satisfied (see Remark 1.18). Up to relabelling the branches of C we can assume that

$$(1.8) \quad \text{ord}_{t_1}(y_1 \circ \eta_{C_1}) \geq \dots \geq \text{ord}_{t_r}(y_1 \circ \eta_{C_r}).$$

For $i = 1, \dots, r-1$ we take $f_i \in I_{C_i}$ such that $f_i \notin I_{C_j}$ for any $1 \leq j \leq r$ and $j \neq i$. Then we choose an integer ℓ_i large enough such that

$$(1.9) \quad \text{ord}_{t_j}(f_i \circ \eta_{C_j}) < \text{ord}_{t_j}(y_1^{\ell_i} \circ \eta_{C_j}), \text{ for } j = i+1, \dots, r.$$

Set $m = m' + r - 1$ and consider the functions:

$$y_1, \dots, y_{m'}, y_{m'+1} := f_1 + y_1^{\ell_1}, \dots, y_{m'+r-1} := f_{r-1} + y_1^{\ell_{r-1}}.$$

Then, the vectors (1.7) associated with the branches of C , with respect to the functions y_1, \dots, y_m , are pairwise different. This follows from the inequalities

$$\text{ord}_{t_i}(y_{m'+i} \circ \eta_{C_i}) = \text{ord}_{t_i}(y_1^{\ell_i} \circ \eta_{C_i}) \stackrel{(1.8)}{\geq} \text{ord}_{t_j}(y_1^{\ell_i} \circ \eta_{C_j}) \stackrel{(1.9)}{>} \text{ord}_{t_j}(f_i \circ \eta_{C_j}) = \text{ord}_{t_j}(y_{m'+i} \circ \eta_{C_j}),$$

for $i = 1, \dots, r-1$ and $j = i+1, \dots, r$. \square

Example 1.20. The integers $\bar{\beta}_0 := 8$, $\bar{\beta}_1 := 12$, $\bar{\beta}_2 := 26$ and $\bar{\beta}_3 := 53$ define the minimal system of generators of the semigroup of a plane branch (see [Tei06, Sec 3.2]). By the method explained in *loc.cit.* we get that a defining equation of such a plane branch C is

$$((Y^2 - X^3)^2 - X^5 Y)^2 - X^{10}(Y^2 - X^3) = 0.$$

Then, $y_0 = X$, $y_1 = Y$, $y_2 = Y^2 - X^3$, and $y_3 = (Y^2 - X^3)^2 - X^5 Y$ verify the conditions of Corollary 1.17 (see Remark 1.18). But observe that y_0 , y_1 and y_3 verify also these conditions because $\gcd(8, 12, 53) = 1$. By Corollary 1.17, any regular fan subdividing $\Sigma_{0,3}$ which contains the ray spanned by $(8, 12, 53)$ defines a toric embedded resolution of C . Next we study the behaviour of the branch $C \subset k^3$, and also of the plane $k^2 \subset k^3$, under the monomial map $\pi_{\sigma_{0,3}}^\sigma$ corresponding to the regular cone $\sigma \subset \sigma_{0,3}$ generated by the vectors $v_1 = (1, 1, 4)$, $v_2 = (2, 3, 13)$, and $v_3 = (8, 12, 53)$. Consider $X_{\Sigma_{0,3}} = k^3$ with coordinates (X_1, X_2, X_3) as in Notation 1.3, and X_σ with coordinates (U_1, U_2, U_3) , where $U_i = \chi^{\tilde{v}_i}$. The equation that defines $k^2 \subset k^3$ is $X_3 - (X_2^2 - X_1^3)^2 + X_1^5 X_2 = 0$. In addition, the points of $C \subset k^3$ must satisfy the following equation $X_3^2 - X_1^{10}(X_2^2 - X_1^3) = 0$. The points in the strict transform \tilde{C} of the curve C by the monomial map $\pi_{\sigma_{0,3}}^\sigma$ satisfy the following two equations:

$$(1.10) \quad U_2 U_3^5 - (1 - U_1)^2 + U_1^2 U_2 U_3^4 = 0,$$

$$(1.11) \quad U_3^2 - U_1^4 (1 - U_1) = 0,$$

where (1.10) is the equation of the strict transform of k^2 . Starting with (1.11) and then substituting in (1.10), we obtain a parametrization of \tilde{C} of the form

$$U_1 = 1 - t^2 + \dots, \quad U_2 = 1 + \dots, \quad U_3 = t + \dots,$$

where we have omitted higher order terms (compare with (1.4)). As expected, \tilde{C} is smooth and transverse to $U_3 = 0$. Notice that the intersection of the surfaces defined by (1.11) and (1.10), contains also an exceptional component given by $U_1 = 1, U_2 = t, U_3 = 0$, and the strict transform of the plane (1.10) is singular at the point $(1, 0, 0)$.

One may consider this toric resolution as a sort of *toric embedded normalization* of C , since we do not care about the effect of this modification on the original plane $k^2 \supset C$.

In Corollary 1.19 we showed the existence of functions defining a suitable embedding. We consider a different choice of functions in Theorem 1.22 below. We are grateful to Javier Fernández de Bobadilla for sharing some ideas which were helpful in the formulation of the result. We will need the following elementary lemma:

Lemma 1.21. *Denote by N' the sublattice of N spanned by v_1, \dots, v_{s-1} . If v_1, \dots, v_{s-1} are part of a basis of the lattice N and if the canonical image of a vector $w \in N$ is a primitive vector in the lattice N/N' , then v_1, \dots, v_{s-1}, w are part of a basis of the lattice N .*

Proof. Let $v_1, \dots, v_{s-1}, v_s, \dots, v_n$ be a basis of the lattice N . Denote by \bar{u} the class of a vector $u \in N$ in the quotient lattice N/N' . We have that $\bar{v}_s, \dots, \bar{v}_n$ is a basis of N/N' . By hypothesis \bar{w} is a primitive vector in the lattice N/N' , hence there are elements $w_s := w, w_{s+1}, \dots, w_n \in N$ such that $\bar{w}_s = \bar{w}, \bar{w}_{s+1}, \dots, \bar{w}_n$ is a basis of N/N' . If we expand $\bar{w}_j = \sum_{l=s}^n a_{l,j} \bar{v}_l$, for $j = s, \dots, n$, it follows that the matrix $A = (a_{l,j})$ has determinant ± 1 . It is enough to check that $v_1, \dots, v_{s-1}, w_s, w_{s+1}, \dots, w_n$ is a basis of N . The matrix whose columns are the coordinates of these vectors in terms of v_1, \dots, v_n has a block structure of the form

$$\begin{pmatrix} I_{s-1} & * \\ 0 & A \end{pmatrix}$$

where I_{s-1} denotes the identity matrix of size $s-1$. This matrix is unimodular, since A is. Therefore $v_1, \dots, v_{s-1}, w_s, w_{s+1}, \dots, w_n$ is a basis of N as claimed. \square

Theorem 1.22. *Let $C = \cup_{j=1}^r C_j$ be a reduced curve with r branches embedded in k^n by the map (1.3). Let $y_1, \dots, y_m, y_{m+1}, \dots, y_{m+r} \in k[[X_1, \dots, X_n]]$ be such that:*

- (1) y_1, \dots, y_m generate the maximal ideal of $k[[X_1, \dots, X_n]]$,
- (2) $y_1, \dots, y_m \notin I_{C_j}$, for $1 \leq j \leq r$,
- (3) the numbers $\text{ord}_{t_j}(y_s \circ \eta_{C_j}) \in \mathbb{Z}_{>0}$, $s = 1, \dots, m$, generate the group \mathbb{Z} , for $1 \leq j \leq r$,
- (4) $y_{m+j} \in \bigcap_{1 \leq i \leq r}^{i \neq j} I_{C_i}$ and $y_{m+j} \notin I_{C_j}$, for $1 \leq j \leq r$.

Let $C \hookrightarrow k^{m+r}$ be the embedding defined as (1.6) with respect to y_1, \dots, y_{m+r} . Then, there exists a toric modification of $X_{\Sigma_{0,m+r}} = k^{m+r}$ which is a toric embedded resolution of C .

Proof. We denote by Σ_0 the fan $\Sigma_{0,m+r}$ and use Notation 1.3. Condition (4) implies that for $1 \leq j \leq r$, the branch $C_j \subset k^{m+r}$ is contained in the coordinate subspace:

$$Y_{m+1} = \dots = Y_{m+j-1} = Y_{m+j+1} = \dots = Y_{m+r} = 0.$$

This subspace is the closure of the orbit $O(\sigma_j)_{\Sigma_0}$, where the cone $\sigma_j \in \Sigma_0$ is spanned by the vectors $e_{m+1}, \dots, e_{m+j-1}, e_{m+j+1}, \dots, e_{m+r}$, of the canonical basis $N = \mathbb{Z}^{m+r}$. Denote by $\bar{e}_{j,i}$ the image of e_i under the canonical projection $N \rightarrow N(\sigma_j)$, for $i = 1, \dots, m+r$. By conditions (2)-(3), the arc η_j (i.e., the composition of η_{C_j} with $k^n \hookrightarrow k^{m+r}$) has its generic point in the torus $O(\sigma_j)_{\Sigma_0}$, and the order vector $\text{ord}(\eta_j)$ is primitive in the lattice $N(\sigma_j)$. We have that

$$(1.12) \quad \text{ord}(\eta_j) = w_{j,1}\bar{e}_{j,1} + \dots + w_{j,m}\bar{e}_{j,m} + w_{j,m+j}\bar{e}_{j,m+j},$$

in terms of the basis of $\bar{e}_{j,1}, \dots, \bar{e}_{j,m}, \bar{e}_{j,m+j}$ of $N(\sigma_j)$, with all the $w_{j,i}$ in $\mathbb{Z}_{>0}$. Let us consider the vectors:

$$w_0 := w_{1,m+1}e_{m+1} + \dots + w_{r,m+r}e_{m+r}, \quad w_j := w_{j,1}e_1 + \dots + w_{j,m}e_m, \quad \text{for } 1 \leq j \leq r.$$

Notice that the canonical image of $w_0 + w_j$ in $N(\sigma_j)$ is precisely the primitive vector $\text{ord}(\eta_j) \in N(\sigma_j)$. The cone

$$\theta_j := \mathbb{R}_{\geq 0}(w_0 + w_j) + \sigma_j \subset N_{\mathbb{R}}$$

is regular for the lattice N by Lemma 1.21, and its canonical projection in $N(\sigma_j)_{\mathbb{R}}$ is equal to the ray $\rho_j := \mathbb{R}_{\geq 0}\text{ord}(\eta_j) \subset N(\sigma_j)_{\mathbb{R}}$.

Let us prove that the cones θ_j for $j = 1, \dots, r$ are pairwise different. We assume that there are integers $1 \leq s < l \leq r$ such that $\theta := \theta_s = \theta_l$, and proceed by contradiction. By construction, the cone θ is of dimension r . Since σ_s and σ_l are faces of θ , it follows that

$$\sigma_s + \sigma_l = \mathbb{R}_{\geq 0}e_{m+1} + \dots + \mathbb{R}_{\geq 0}e_{m+r}$$

is an r dimensional cone contained in θ . Assume that there is a vector $w \in \theta \setminus (\sigma_s + \sigma_l)$. Since θ is contained in the cone $\mathbb{R}_{\geq 0}e_1 + \dots + \mathbb{R}_{\geq 0}e_{m+r}$, this implies that $w, e_{m+1}, \dots, e_{m+r}$ are linearly independent, contradicting the assumption that θ has dimension r . Hence, we get that $\theta = \mathbb{R}_{\geq 0}e_{m+1} + \dots + \mathbb{R}_{\geq 0}e_{m+r}$. But, then the canonical projection of θ onto $N(\sigma_l)_{\mathbb{R}}$ is equal to the ray spanned by $\bar{e}_{l,m+l}$. This ray is different from the ray ρ_l because the coefficients $w_{l,i}$ appearing in the expansion (1.12) of $\text{ord}(\eta_l)$ are non zero. This contradiction shows that the cones θ_j , for $1 \leq j \leq r$ are pairwise different, as claimed.

Assume that Σ is a regular subdivision of Σ_0 which contains the cones θ_j , for $j = 1, \dots, r$. Then, we have $\sigma_j \in \Sigma$ since σ_j is a face of θ_j , and ρ_j belongs to the fan $\text{Star}(\sigma_j)_{\Sigma}$. This implies that the toric modification $\pi_{\Sigma_0}^{\Sigma}$ is a toric embedded resolution of the branches C_j for $j = 1, \dots, r$, by Proposition 1.13. The strict transform of C_j by this modification is smooth and transversal to the orbit associated with θ_j , for $1 \leq j \leq r$. Since these cones are pairwise different, it follows that the strict transform of C by $\pi_{\Sigma_0}^{\Sigma}$ is smooth.

In order to finish the proof it remains to show that such a regular fan Σ exists. It is enough to show the existence of a fan Σ' subdividing Σ_0 such that the regular cone θ_j belongs to Σ' , for $1 \leq j \leq r$, and then obtain Σ as a regular subdivision of Σ' .

Next, we show that the set consisting of the cones θ_j , for $j = 1, \dots, r$, and their faces is a fan, which contains cones of dimension at most r . Then, there exists a fan Σ' subdividing Σ_0 and containing it, and this ends the proof. Set $1 \leq j, j' \leq r$ and let us check that $\theta_j \cap \theta_{j'}$ is a common face of θ_j and $\theta_{j'}$. If $w_j = w_{j'}$ then $\theta_j \cap \theta_{j'} = \sigma_j \cap \sigma_{j'} + \mathbb{R}_{\geq 0}(w_j + w_0)$. If $w_j \neq w_{j'}$ then $\theta_j \cap \theta_{j'} = \sigma_j \cap \sigma_{j'}$. In both cases the assertion holds by a direct computation when we express a vector $v \in \theta_j \cap \theta_{j'}$ in terms of the basis e_1, \dots, e_{m+r} . \square

In the proof of Theorem 1.22, we show the existence of a fan Σ' subdividing Σ_0 , which can be obtained explicitly as follows.

Remark 1.23. Let N_1 (resp. N_2) be the lattice spanned by e_1, \dots, e_m (resp. e_{m+1}, \dots, e_{m+r}), and let Σ_1 (resp. Σ_2) be the fan of faces of the cone spanned by e_1, \dots, e_m (resp. e_{m+1}, \dots, e_{m+r}). The lattice N is equal to the direct sum of its sublattices N_1 and N_2 . The fan Σ_0 is the *direct* Minkowski sum $\Sigma_1 \oplus \Sigma_2$ of the fans Σ_1 and Σ_2 , in the sense that any cone $\tau \in \Sigma_0$ is the Minkowski sum of unique cones $\tau_1 \in \Sigma_1$ and $\tau_2 \in \Sigma_2$. Let Σ'_1 be a regular subdivision of Σ_1 containing the rays spanned by w_j , for $j = 1, \dots, r$. Let us denote by Σ'_2 the star subdivision of Σ_2 at the vector w_0 . The r -dimensional cones of Σ'_2 are $\sigma_j + \mathbb{R}_{\geq 0}w_0$, for $j = 1, \dots, r$. The $(r+1)$ -dimensional cone $\tau_j := \mathbb{R}_{\geq 0}w_j + \mathbb{R}_{\geq 0}w_0 + \sigma_j$ belongs to the direct Minkowski sum of fans $\Sigma'_1 \oplus \Sigma'_2$ and contains the cones σ_j and $\mathbb{R}_{\geq 0}w_j + \mathbb{R}_{\geq 0}w_0$ as faces. Apply, for $1 \leq j \leq r$, a star subdivision at the vector $\mathbb{R}_{\geq 0}(w_j + w_0)$. This combinatorial operation only affects the cones which contain $\mathbb{R}_{\geq 0}w_j + \mathbb{R}_{\geq 0}w_0$ as a face (see [CLS11, Lemma 11.1.3]). In particular, it replaces the cone τ_j by the cones $\mathbb{R}_{\geq 0}w_j + \mathbb{R}_{\geq 0}(w_0 + w_j) + \sigma_j$, $\mathbb{R}_{\geq 0}w_0 + \mathbb{R}_{\geq 0}(w_0 + w_j) + \sigma_j$, and $\theta_j = \mathbb{R}_{\geq 0}(w_0 + w_j) + \sigma_j$, and their faces. Therefore, the fan Σ' constructed by this process, starting from $\Sigma'_1 \oplus \Sigma'_2$, contains the cones θ_j , for $1 \leq j \leq r$.

2. RESOLVING A PLANE CURVE SINGULARITY WITH ONE TORIC MORPHISM

In this section, we prove that a singular curve $C = \cup_{j=1}^r C_j$ with $r \geq 1$ branches admits a non degenerate reembedding, in such a way that an embedded resolution is obtained by one toric morphism. This result is a generalization of theorems of Goldin and Teissier in [GT00].

2.1. Background on plane curve singularities and divisorial valuations. In this section we explain the basic notation and conventions used in the rest of the paper about plane curve singularities and divisorial valuations. We denote by S the affine space $\text{Spec } k[X, Y]$ and by O its closed point corresponding to the maximal ideal (X, Y) . We denote by R the ring $k[[X, Y]]$ and by m_R its maximal ideal as before.

Let $C = \cup_{j=1}^r C_j$ be a plane curve with $r \geq 1$ branches. Denote by \mathcal{O} its local ring. If x, y generate the maximal ideal of \mathcal{O} , then the embedding of $C \subset \text{Spec } R$ is defined by a local homomorphism of k -algebras

$$(2.1) \quad R \longrightarrow \mathcal{O}, \text{ such that } X \mapsto x, Y \mapsto y.$$

If D is a closed subscheme of $\text{Spec } R$ defined by a principal ideal of R , then we call any generator $f \in R$ of this ideal a *defining function* of D and we write $D = Z(f)$. If $f \in m_R$ generates the kernel of the map (2.1), then f is a defining function of C and $\{C_j\}_{j=1}^r$ is in bijection with the set of irreducible factors of f in R .

If $D_1 = Z(f_1)$ and $D_2 = Z(f_2)$, we denote by $(D_1 \cdot D_2) := \dim_k R/(f_1, f_2)$ the *intersection multiplicity* of D_1 and D_2 . If D_1 is a branch, then $(D_1 \cdot D_2)$ is equal to the order of the series

$f_2 \circ \eta_{D_1} \in k[[t]]$, where $\eta_{D_1} : \text{Spec } k[[t]] \rightarrow \text{Spec } R$ is the arc associated to D_1 (see Definition 1.10).

Remark 2.1. A pair $(x, y) \in R^2$ is a *local coordinate system* on $\text{Spec } R$ if x, y generate the maximal ideal m_R . Once the local coordinate system (x, y) is fixed, the Weierstrass Preparation Theorem allows us to express any branch $A \neq Z(x)$ in the form $A = Z(x_A)$ for a monic element $x_A \in k[[x]][[y]]$ such that $\deg_y x_A = (Z(x) \cdot A)$.

If A is a plane branch on S then the semigroup Γ_A is finitely generated. We mention below some classical properties (see [GBP15, Lemma 1.2 and Theorem 3.2]).

Notation 2.2. Let A be a plane branch on S . Denote by $\bar{\beta}_0, \bar{\beta}_1, \dots, \bar{\beta}_{g(A)}$ the minimal generating system of the semigroup Γ_A , where $\bar{\beta}_0 < \bar{\beta}_1 < \dots < \bar{\beta}_{g(A)}$. We set $e_0 := \bar{\beta}_0$, $e_j = \gcd(e_{j-1}, \bar{\beta}_j)$, and $n_j = e_{j-1}/e_j$ for $j = 1, \dots, g(A)$. We set $n_0 := 0$.

Proposition 2.3.

(1) For every $j \in \{1, \dots, g(A)\}$ one has $n_j > 1$ and

$$(2.2) \quad n_j \bar{\beta}_j = b_0^j \bar{\beta}_0 + b_1^j \bar{\beta}_1 + \dots + b_{j-1}^j \bar{\beta}_{j-1} \text{ with } 0 \leq b_i^j < n_i, \text{ for } i = 1, \dots, j-1.$$

(2) We have the inequalities:

$$(2.3) \quad n_j \bar{\beta}_j < \bar{\beta}_{j+1}, \text{ for } j = 1, \dots, g(A) - 1.$$

(3) There exist a local coordinate system (x_0, x_1) on S and irreducible elements $x_j \in k[[x_0]][[x_1]]$ with $\deg_{x_1} x_j = n_1 \dots n_{j-1}$, for $j \in \{2, \dots, g(A)\}$, such that $(Z(x_j) \cdot A) = \bar{\beta}_j$ for $j \in \{0, \dots, g(A)\}$.

A *model* of (S, O) is a proper birational morphism $\psi: (S(\psi), E(\psi)) \rightarrow (S, O)$, obtained as a composition of blow ups of infinitely near points of O . The preimage $E(\psi) = \psi^{-1}(O)$, seen as a reduced divisor on $S(\psi)$, is called the *exceptional divisor* of ψ . It has simple normal crossings and its irreducible components are projective lines which are called the *exceptional prime divisors* of ψ . We denote by E_O the prime exceptional divisor created by blowing up O in S .

The *dual graph* $G(\psi)$ of the model ψ is a tree whose set of vertices is in bijection with the set of exceptional prime divisors of ψ , and where two vertices are joined by an edge if and only if the corresponding exceptional prime divisors intersect. Recall that the *valency* of a vertex in a tree is the number of edges incident to it. A vertex of a tree is an *end* (resp. a *ramification vertex*) if it has valency equal to one (resp. ≥ 3). We denote by \mathcal{E}_ψ the set of exceptional prime divisors of ψ which correspond to ends of $G(\psi)$.

Notation 2.4. If C is a curve on S we denote by C^ψ its strict transform on $S(\psi)$. If the model ψ is clear from the context we often denote by the same letter a curve C and its strict transform C^ψ on this model. This applies also in the case of an exceptional prime divisor E which was created in another model dominated by ψ and its strict transform E^ψ . For instance, we will often denote E_O^ψ simply by E_O .

Definition 2.5. Let E be an irreducible component of $E(\psi)$. A plane branch C on S is a *curveta* of E at the model ψ if C^ψ is smooth and transversal to E at a smooth point of $E(\psi)$. We denote by $\mathcal{C}_E(\psi)$ the set of curvetas of E at the model ψ . We call ν_E the *divisorial valuation* on (S, O) defined by E , that is, ν_E maps $h \in R \setminus \{0\}$ to the order of vanishing along E of $h \circ \psi$.

Remark 2.6. The value of ν_E at $h \in R \setminus \{0\}$ is characterized in terms of intersection multiplicity of curvetas as follows:

$$(2.4) \quad \nu_E(h) = \min\{(Z(h) \cdot D) \mid D \in \mathcal{C}_E(\psi)\}.$$

Furthermore, the minimum in (2.4) is attained whenever the strict transforms $Z(h)^\psi$ and D^ψ intersect E at different points (see [Spi90, Section 7]).

Definition 2.7. A model ψ is an *embedded resolution* of a curve C on S if the divisor $C^\psi + E(\psi)$ has simple normal crossings. If $C = \cup_{j=1}^r C_j$ is a curve with r branches, we denote by E_{C_j} the unique exceptional prime divisor of ψ that intersects C_j^ψ , for $j = 1, \dots, r$. The *dual graph of the total transform* of C under ψ is the tree $G(\psi, C)$ obtained by adding to $G(\psi)$, a new vertex corresponding to C_j^ψ , and an edge joining it to the vertex of $G(\psi)$ corresponding to E_{C_j} , for $j = 1, \dots, r$.

Definition 2.8. A model ψ is the *minimal resolution* of the divisorial valuation ν_E if ψ is the composition of a finite sequence of point blow-ups above O ,

$$\psi: S(\psi) = S_n \xrightarrow{\psi_n} S_{n-1} \xrightarrow{\psi_{n-1}} \dots \xrightarrow{\psi_2} S_1 \xrightarrow{\psi_1} S_0 = S$$

such that E is the exceptional divisor of ψ_n and the center of ψ_{i+1} in S_i belongs to the exceptional divisor of ψ_i for $i = 1, \dots, n-1$.

If ψ is the minimal resolution of ν_E and A is any curvetta in $\mathcal{C}_E(\psi)$, then by definition ψ is an embedded resolution of A . However, it may not be the minimal one. Denote by $\pi: (S(\pi), E(\pi)) \rightarrow (S, O)$ the minimal embedded resolution of A and by E_A the irreducible component of $E(\pi)$ which intersects A^π . Since by definition ψ dominates π , the strict transform E_A^ψ is a component of $E(\psi)$. We will distinguish the following two cases:

- (1) $E_A^\psi = E$, that is, the minimal resolution of ν_E equals the minimal embedded resolution of any curvetta of E . In this case we set $\ell_E := 0$.
- (2) $E_A^\psi \neq E$, in this case the model $S(\psi)$ is obtained from $S(\pi)$ after blowing up $\ell_E > 0$ additional infinitely near points of A over O . We indicate in Figure 1 the shapes of the dual graph of $G(\psi)$ in both cases.



FIGURE 1. The shape of the dual graph in the case $\ell_E = 0$ on the left (resp. $\ell_E > 0$ on the right). The vertex corresponding to E is marked by a bigger bullet.

We introduce the notion of generating sequence of a finite set of divisorial valuations according to Delgado, Campillo, Galindo and Núñez (see [CG03, DGN08]), based upon the work of Spivakovsky who described the case of one valuation (see [Spi90]).

Let E_1, \dots, E_s be exceptional prime divisors on a model ψ of (S, O) . We denote by $V = (\nu_{E_1}, \dots, \nu_{E_s})$ be the associated tuple of divisorial valuations. Given a nonzero $h \in R$, we denote $V(h) = (\nu_{E_1}(h), \dots, \nu_{E_s}(h)) \in \mathbb{Z}_{\geq 0}^s$. For any element $\gamma \in \mathbb{Z}_{\geq 0}^s$, we consider the *valuation ideal* \mathcal{P}_γ^V of R defined by $\mathcal{P}_\gamma^V = \{h \in R \mid V(h) \geq \gamma\} \cup \{0\}$, where \leq is the partial ordering over \mathbb{Z}^s given by $\delta \leq \gamma$ if and only if $\gamma - \delta \in \mathbb{Z}_{\geq 0}^s$.

Definition 2.9. (See [Spi90, CG03, DGN08]). The sequence $x_0, \dots, x_m \in m_R$ is a *generating sequence* of V if for each $\gamma \in \mathbb{Z}_{\geq 0}^s$ the ideal \mathcal{P}_γ^V is generated by those monomials in x_0, \dots, x_m which belong to \mathcal{P}_γ^V . We say that it is *minimal* if no proper subsequence of it is a generating sequence of V . The *minimal resolution* of V is a model ψ of (S, O) such that ψ dominates the minimal resolution of ν_{E_i} for all $i \in \{1, \dots, s\}$, and ψ is minimal with this property.

Observe that $\mathcal{P}_{\underline{1}}^V = m_R$ where $\underline{1} = (1, \dots, 1)$, and thus a generating sequence x_0, \dots, x_m generates the maximal ideal m_R . If $s = 1$, that is, when $V = (\nu_E)$ we speak simply about generating sequences of the divisorial valuation ν_E .

Remark 2.10. A generating sequence of V is also a generating sequence of ν_{E_i} , for $1 \leq i \leq s$. Indeed, if $p_i: \mathbb{Z}_{\geq 0}^s \rightarrow \mathbb{Z}_{\geq 0}$ denotes the canonical projection onto the i^{th} -coordinate and $\phi \in \mathbb{Z}_{\geq 0}$, then $p_i^{-1}(\phi)$ has a unique minimal element and $\mathcal{P}_{\gamma}^V = \mathcal{P}_{\phi}^{\nu_i}$ with $\gamma = \min p_i^{-1}(\phi)$.

The following result gives a geometric characterization of a minimal generating sequence of a finite set of divisorial valuations (see [Spi90, Section 8] for the case of one valuation and [DGN08] for the general case).

Proposition 2.11. *Let $\psi : (S(\psi), E(\psi)) \rightarrow (S, O)$ be the minimal resolution of a tuple of divisorial valuations $V = (\nu_{E_1}, \dots, \nu_{E_s})$. Assume that $V \neq (\nu_{E_O})$. Write $\mathcal{E}_{\psi} = \{E_0, \dots, E_{g(V)}\}$, where $E_0 = E_O$ and the indices are compatible with the order in which the divisors were created, and take $x_i \in m_R$ such that the branch $L_i = Z(x_i)$ is a curvetta of E_i , for $i = 0, \dots, g(V)$. Then, $x_0, \dots, x_{g(V)}$ is a minimal generating sequence of V and any minimal generating sequence of V is of this form.*

It follows that if E is an exceptional prime divisor then any minimal generating sequence of ν_E has the same length, that is, it is of the form $x_0, x_1, \dots, x_{g(E)}$, for some $g(E) \geq 1$.

Example 2.12. If $E = E_O$ then, for all $h \neq 0$, $\nu_E(h)$ is the largest $n \in \mathbb{Z}_{\geq 0}$ such that $h \in m_R^n$. Any minimal generating sequence of ν_E is of the form $x_0, x_1 \in m_R$ where (x_0, x_1) defines a local coordinate system on S . In this case $\ell_E = 1$, and we have $g(E) = 1$.

Proposition 2.13 (see [Spi90]). *Let us consider a minimal generating sequence $x_0, \dots, x_{g(E)}$ of ν_E . Denote $L_j = Z(x_j)$, for $j \in \{0, \dots, g(E)\}$. Let A be any curvetta in $\mathcal{C}_E(\psi)$. Denote by $\bar{\beta}_0, \dots, \bar{\beta}_{g(A)}$ the minimal system of generators of the semigroup Γ_A (see Notation 2.2). Then:*

$$\nu_E(x_j) = \bar{\beta}_j, \text{ for } j = 0, \dots, g(A), \text{ and } \nu_E(x_A) = n_{g(A)} \bar{\beta}_{g(A)} + \ell_E.$$

In addition, if $\ell_E = 0$ then one has $g(E) = g(A)$, while if $\ell_E > 0$ then $g(E) = g(A) + 1$ and in this case $L_{g(E)}$ is a curvetta in $\mathcal{C}_E(\psi)$.

Notation 2.14. If $\ell_E > 0$, we set $n_{g(E)} := 1$ and $\bar{\beta}_{g(E)} := n_{g(A)} \bar{\beta}_{g(A)} + \ell_E$. We get the formulas $\nu_E(x_A) = n_{g(E)} \bar{\beta}_{g(E)}$ and $\deg_{x_1} x_A = n_1 \dots n_{g(E)}$, which hold also when $\ell_E = 0$.

The next definition is classical.

Definition 2.15. Let $C = \cup_{j=1}^r C_j$ be a reduced singular curve at (S, O) . Let ψ be the minimal embedded resolution of C and let $V = (\nu_{E_1}, \dots, \nu_{E_s})$ be a tuple of divisorial valuations defined by the exceptional prime divisors E_1, \dots, E_s of ψ which intersect the strict transform of C . Let x_0, x_1, \dots, x_m , ($m = g(V)$), be a minimal generating sequence of V and denote $L_i = Z(x_i)$, for $i \in \{1, \dots, m\}$. We say then that L_0, \dots, L_m is a sequence of *maximal contact curves* of C .

By Proposition 2.11 the sequence L_0, \dots, L_m is in bijection with the ends of the dual graph $G(\psi)$, each L_i is a curvetta of a unique exceptional prime divisor of ψ defining an end of $G(\psi)$.

We often require sequences of maximal contact curves of C such that their strict transforms do not intersect the strict transform of C . That is the following genericity condition (see Remark 3.21 below for some consequences of it):

$$(2.5) \quad (C^{\psi} \cdot L_i^{\psi}) = 0, \text{ for } 1 \leq i \leq m.$$

Lemma 2.16. *Let $V = (\nu_{E_1}, \dots, \nu_{E_s})$ be a tuple of divisorial valuations on (S, O) . Take $x_0, \dots, x_{g(V)} \in m_R$ a minimal generating sequence of V and set $L_i = Z(x_i)$ for $i \in \{0, \dots, g(V)\}$. For every $j \in \{0, \dots, g(V)\}$ we can extract from $L_0, \dots, L_g(V)$ a sequence of maximal contact curves of L_j .*

Proof. By Remark 2.10 one can extract from the sequence $x_0, \dots, x_{g(V)}$ a minimal generating sequence of the divisorial valuation $\nu_{E_{i_j}}$, By definition 2.15, this provides a sequence of maximal contact curves of L_j . \square

Example 2.17. Let C be the plane curve defined by $f = f_1 f_2$, where $f_1 = Y^2 - X^3 - X^4$ and $f_2 = Y^2 - X^3 - X^5$. Denote by C_i the branch defined by f_i , for $i = 1, 2$. Let ψ be the minimal embedded resolution of C . We obtain ψ , after blowing up the origin, by successively blowing up the point where the strict transforms of C_1 and C_2 intersect. The exceptional divisor $E(\psi)$ has 5 irreducible components. We call them D_0, \dots, D_4 , where the indices are compatible with the order in which the divisors were created. The divisors corresponding to the ends of $G(\psi)$ are $E_0 := D_0$, $E_1 := D_1$, and $E_2 := D_4$. The strict transforms C_1^ψ and C_2^ψ intersect E_2 . Set $E := E_2$. By construction, the minimal resolution of ν_E is ψ . We have $g(E) = 2$ and $\ell_E = 2$. For any $a \in k$, the plane branch defined by $h_a = Y^2 - X^3 + aX^4$ is a curvetta of E . We may take as a minimal generating sequence of ν_E the functions $x_0 = X$, $x_1 = Y$ and $x_2 = h_a$.

On the one hand, if $a = -1$ then $h_a = f_1$ and $(C_1^\psi \cdot Z(h_a)^\psi) = \infty$. On the other hand, if $a = 0$ then $(C_2^\psi \cdot Z(h_a)^\psi) = 1$. For $a \in k \setminus \{0, -1\}$, Hence, $x_0 := X$, $x_1 := Y$, and $x_2 := Y^2 - X^3 + aX^4$ define a sequence of maximal contact curves of $V = (\nu_E)$ for any $a \in k$, while the conditions (2.5). are satisfied just for $a \in k \setminus \{0, -1\}$.

Example 2.18. Let C be the plane curve defined by $f = f_1 f_2$, where $f_1 = (Y^2 - X^3)^2 - X^5 Y$ and $f_2 = (Y^2 - X^3)^2 - X^6 Y$. Denote by C_i the branch defined by f_i , for $i = 1, 2$. In order to obtain the minimal embedded resolution ψ of C , we start with the minimal embedded resolution of C_1 and then we successively blow up the points at which the total transform of C_2 fails to have simple normal crossing support. The exceptional divisor $E(\psi)$ has 7 irreducible components. We call them E_0, \dots, E_6 , where the indices are compatible with the order in which the divisors were created. With the notation of Definition 2.7, we have $EC_1 = E_4$ and $EC_2 = E_6$.

Let us set $V = (\nu_1, \nu_2)$, where ν_i is the divisorial valuation defined by EC_i , for $i = 1, 2$. By construction, the minimal resolution of V is ψ . The divisors E_0 , E_1 , and E_5 are those which correspond to ends of $G(\psi)$. For any $a \in k$, the branches defined by $x_0 = X$, $x_1 = Y$, and $x_2 = Y^2 - X^3 + aX^4$ form a sequence of maximal contact curves of V .

Consider now C' be the curve defined by $f = f_1 f_2 f_3$ with $f_3 = Y^2 - X^3$. Then, for any $a \in k^*$, $x_0 = X$, $x_1 = Y$, and $x_2 = Y^2 - X^3 + aX^4$ define a sequence of maximal contact curves of C' .

2.1.1. *Expansions in terms of generating sequences.* We recall how to expand a function in terms of a minimal generating sequence $x_0, \dots, x_{g(E)}$ of a divisorial valuation ν_E , and how to determine from it the value of ν_E on this function. The elements $x_2, \dots, x_{g(E)}$ can be chosen as monic polynomials in the ring $k[[x_0]][[x_1]]$ (see Remark 2.1).

Proposition 2.19. (see [PP04, Lemma 7.2]) *Let us consider positive integers $G \geq 1$, and $N_j > 1$ for $j = 1, \dots, G - 1$, and let $f_j \in k[[x]][[y]]$ be monic polynomials in y with*

$$\deg_y f_1 = 1 \text{ and } \deg_y f_j = N_1 \dots N_{j-1}, \text{ for } j = 2, \dots, G.$$

Then, any polynomial $f \in k[[x]][[y]]$ has a unique finite expansion in the form:

$$(2.6) \quad f = \sum_{I=(i_1, \dots, i_G)} c_I(x) \cdot f_1^{i_1} \cdots f_G^{i_G},$$

where $c_I(x) \in k[[x]]$, $0 \leq i_G \leq \left\lfloor \frac{\deg_y(f)}{\deg_y(f_G)} \right\rfloor$, and $0 \leq i_j < N_j$ for $j = 1, \dots, G-1$.

We call the expansion (2.6) the (x, f_1, \dots, f_G) -adic expansion of f .

Next we apply Proposition 2.19 with respect to the minimal generating sequence $x_0, \dots, x_{g(E)}$ of ν_E taking into account the properties of the values $\nu_E(x_0) = \bar{\beta}_0, \dots, \nu_E(x_{g(E)}) = \bar{\beta}_{g(E)}$ (see Proposition 2.13, Notation 2.14 and Proposition 2.3). See also [Spi90, Rem. 8.16] for details.

Proposition 2.20. *Let A be a curveta in $\mathcal{C}_E(\psi)$. The $(x_0, \dots, x_{g(E)})$ -adic expansion of x_A is of the form:*

$$(2.7) \quad x_A = x_{g(E)}^{n_{g(E)}} - \theta_{I_0}^A \cdot x_0^{b_0} x_1^{b_1} \cdots x_{g(E)-1}^{b_{g(E)-1}} + p_A(x_0, \dots, x_{g(E)}),$$

where $\theta_{I_0}^A \in k$, $I_0 = (b_0, b_1, \dots, b_{g(E)-1}, 0)$, the equality

$$(2.8) \quad n_{g(E)} \bar{\beta}_{g(E)} = b_0 \bar{\beta}_0 + \cdots + b_{g(E)-1} \bar{\beta}_{g(E)-1}, \text{ with } 0 \leq b_j < n_j \text{ for } 1 \leq j < g(E) - 1,$$

holds, where $p_A(x_0, \dots, x_{g(E)}) = \sum_{I=(i_0, \dots, i_{g(E)})} \theta_I^A \cdot x_0^{i_0} x_1^{i_1} \cdots x_{g(E)}^{i_{g(E)}}$, with $\theta_I^A \in k$, and whenever $\theta_I^A \neq 0$ we have

$$n_{g(E)} \bar{\beta}_{g(E)} < i_0 \bar{\beta}_0 + \cdots + i_{g(E)} \bar{\beta}_{g(E)}, \text{ with } 0 \leq i_j < n_j \text{ for } j = 1, \dots, g(E).$$

In addition, if $\ell_E = 0$ then $\theta_{I_0}^A \in k^*$.

We have also a similar result for the expansion of an element x_{j+1} of the generating sequence in terms of the previous ones.

Proposition 2.21. *The (x_0, \dots, x_j) -adic expansion of x_{j+1} , for $1 \leq j \leq g(E) - 1$, is of the form:*

$$(2.9) \quad x_{j+1} = x_j^{n_j} - \theta_j \cdot x_0^{b_0^j} x_1^{b_1^j} \cdots x_{j-1}^{b_{j-1}^j} + p_j(x_0, \dots, x_j),$$

where $\theta_j \in k^*$, the relation (2.2) holds, where $p_j(x_0, \dots, x_j) = \sum_{I=(i_0^j, \dots, i_j^j)} \theta_I^j \cdot x_0^{i_0^j} x_1^{i_1^j} \cdots x_j^{i_j^j}$ with $\theta_I^j \in k$, and whenever $\theta_I^j \neq 0$ we have $n_j \bar{\beta}_j < \sum_{t=0}^j i_t^j \bar{\beta}_t$ and $0 \leq i_t^j < n_t$ for $t = 1, \dots, j$.

Remark 2.22. One can use the expansion of $h \in k[[x_0]][[x_1]]$ in terms of the generating sequence of ν_E to determine the value of $\nu_E(h)$. Indeed, if $h \in k[[x_0]][[x_1]]$ and if

$$h = \sum_{I=(i_1, \dots, i_{g(E)})} c_I(x_0) \cdot x_1^{i_1} \cdots x_{g(E)}^{i_{g(E)}},$$

is the $(x_0, \dots, x_{g(E)})$ -adic expansion of h then $\nu_E(h) = \min_I \{ \nu_E(c_I(x_0) \cdot x_1^{i_1} \cdots x_{g(E)}^{i_{g(E)}}) \}$, where $\nu_E(c_I(x_0) \cdot x_1^{i_1} \cdots x_{g(E)}^{i_{g(E)}}) = \text{ord}_{x_0}(c_I) \cdot \bar{\beta}_0 + i_1 \bar{\beta}_1 + \cdots + i_{g(E)} \bar{\beta}_{g(E)}$. This may be seen as a consequence of [Ro19, Th. 4.125].

2.2. The embedding defined by the maximal contact curves. We keep the notations of section 2.1. The curve C is embedded in $S = k^2$ by the map (2.1). Recall that for any branch C_j of C we consider an arc η_{C_j} on k^2 defined in terms of the normalization of C_j and the given embedding of $C_j \subset k^2$ (see Definition 1.10).

Definition 2.23. Let us fix L_0, \dots, L_m a sequence of maximal contact curves of C such that satisfies the genericity conditions (2.5) (see Definition 2.15). We set $D := \cup_{j=0}^m L_j$. Choose a defining function $x_i \in m_R$ of the branch L_i , for $i = 0, \dots, m$. The sequence x_0, \dots, x_m generates

the maximal ideal of R . Hence it defines an embedding $k^2 \hookrightarrow k^{m+1}$ (see (1.5)) corresponding to the surjective homomorphism

$$(2.10) \quad k[[X_0, \dots, X_m]] \longrightarrow R, \text{ such that } X_i \mapsto x_i, \text{ for } i = 0, \dots, m.$$

We get from this an embedding $C \hookrightarrow k^{m+1}$ as in (1.6).

With the notation of Section 1.3, for any branch C_j of C we have an arc η_j on k^{m+1} that corresponds to the local homomorphism of k -algebras

$$k[[X_0, \dots, X_m]] \rightarrow k[[t_j]], \text{ such that } X_i \mapsto x_i \circ \eta_{C_j}, \text{ for } i = 0, \dots, m.$$

By (2.5), we have that $x_0, \dots, x_m \notin I_{C_j}$, and as a consequence the arc η_j has its generic point in the torus of k^{m+1} .

Proposition 2.24. *Keep the previous notation. We have:*

- (1) *If C_j is a branch of C , then the positive integers $\text{ord}_{t_j}(x_0 \circ \eta_{C_j}), \dots, \text{ord}_{t_j}(x_m \circ \eta_{C_j})$, generate the semigroup Γ_{C_j} .*
- (2) *Let $\psi: (S(\psi), E(\psi)) \rightarrow (S, O)$ be the minimal embedded resolution of C and let C_j and $C_{j'}$ be two branches of C . If $\text{ord}(\eta_j) = \text{ord}(\eta_{j'})$, then C_j^ψ and $C_{j'}^\psi$ intersect the same irreducible component of $E(\psi)$.*

Proof. The branch C_j belongs to $\mathcal{C}_{E_{C_j}}(\psi)$, where we recall that E_{C_j} is the irreducible component of $E(\psi)$ which intersects C_j^ψ . By Remark 2.6 and (2.5), we deduce that

$$(2.11) \quad \nu_{E_{C_j}}(x_i) = (L_i \cdot C_j) = \text{ord}_{t_j}(x_i \circ \eta_{C_j}), \text{ for } 0 \leq i \leq m.$$

By Proposition 2.11, x_0, \dots, x_m is a generating sequence of the divisorial valuation $\nu_{E_{C_j}}$, thus this sequence contains a minimal generating sequence of $\nu_{E_{C_j}}$. By Proposition 2.11, the set $\{\text{ord}_{t_j}(x_i \circ \eta_{C_j}) \mid 0 \leq i \leq m\}$ contains the minimal generating system of Γ_{C_j} . This proves (1).

If C_j and $C_{j'}$ are two branches of C and $\text{ord}(\eta_j) = \text{ord}(\eta_{j'})$, then $\nu_{E_{C_j}}(x_i) = \nu_{E_{C_{j'}}}(x_i)$ for $i = 0, \dots, m$, by (2.11). Since x_0, \dots, x_m is a generating sequence of both valuations $\nu_{E_{C_j}}$ and $\nu_{E_{C_{j'}}}$, and their values on this sequence coincide, the valuation ideals must be equal. This implies that $\nu_{E_{C_j}} = \nu_{E_{C_{j'}}}$ and thus $E_{C_j} = E_{C_{j'}}$. \square

The following proposition provides a geometric interpretation of a coefficient appearing on a $(x_0, \dots, x_{g(E)})$ -adic expansion of the defining function x_A , of a curvetta $A \in \mathcal{C}_E(\psi)$.

Proposition 2.25. *Let $\psi: (S(\psi), E(\psi)) \rightarrow (S, O)$ be the minimal resolution of ν_E and let A and B be two curvetas in $\mathcal{C}_E(\psi)$. Let $\theta_{I_0}^A$ and $\theta_{I_0}^B$ be the coefficients of the term $x_0^{b_0} x_1^{b_1} \dots x_{g(E)-1}^{b_{g(E)-1}}$ in the expansions (2.7) of x_A and x_B , respectively. The following are equivalent:*

- (1) *The strict transforms of A and B on $S(\psi)$ intersect E at different points.*
- (2) *The coefficients $\theta_{I_0}^A$ and $\theta_{I_0}^B$ are different.*

Proof. • **The case $\ell_E > 0$.** In this case $L_{g(E)} = Z(x_{g(E)})$ is a curvetta of E . Take local coordinates (w, s) at the point of $S(\psi)$ where $L_{g(E)}$ and E meet, such that $Z(w) = E$, and $x_{g(E)} \circ \psi = w^{\nu_E(x_{g(E)})} \cdot s$, that is, $Z(s) = L_{g(E)}^\psi$. Recall that $\nu_E(x_{g(E)}) = \bar{\beta}_{g(E)}$ (see Notation 2.14). Let us compute the total transform of A . First take $j \in \{0, \dots, g(E) - 1\}$. By construction, the exceptional prime divisor E and the strict transform L_j^ψ do not intersect.

Recall that $\nu_E(x_j) = \bar{\beta}_j$. Therefore, there exists $\epsilon_{0,j} \in k^*$ such that $x_j \circ \psi = w^{\bar{\beta}_j}(\epsilon_{0,j} + w(\dots))$. Denoting $\epsilon_{I_0} = \prod_{j=0}^{g(E)-1} \epsilon_{0,j}^{b_j} \in k^*$, from the expansion (2.7) of x_A we obtain the following:

$$x_A \circ \psi = w^{\bar{\beta}_g(E)} (s - \theta_{I_0}^A \epsilon_{I_0} + w(\dots)).$$

Then, the strict transform of A is $A^\psi = Z(s - \theta_{I_0}^A \epsilon_{I_0} + w(\dots))$ and its intersection with E is the point of coordinates $(0, s(A))$ where $s(A) = \theta_{I_0}^A \epsilon_{I_0}$. If we apply this to B we get that $s(B) = \theta_{I_0}^B \epsilon_{I_0}$. Since $\epsilon_{I_0} \neq 0$, it follows that (1) and (2) are equivalent.

• **The case $\ell_E = 0$.** In what follows, we use g instead of $g(E)$. In this case $L_g = Z(x_g)$ is not a curvetta of E . The model ψ dominates the minimal embedded resolution $\varphi: (S(\varphi), E(\varphi)) \rightarrow (S, O)$ of L_g .

We address first the case where $g > 1$. We denote by R_g the exceptional prime divisor of φ which intersects L_g^φ . Take local coordinates (w, s) at the point of $S(\varphi)$ where L_g^φ and R_g meet, such that $Z(w) = R_g$, and $x_g \circ \varphi = w^{\nu_{R_g}(x_g)} \cdot s$, and then, $Z(s) = L_g^\varphi$. According to Proposition 2.11, the minimal generating system of the semigroup of L_g is $\nu_{R_g}(x_0), \nu_{R_g}(x_1), \dots, \nu_{R_g}(x_{g-1})$. By [GBP15, Proposition 5.4], it is related to $\bar{\beta}_0, \bar{\beta}_1, \dots, \bar{\beta}_{g-1}$ as follows:

$$\nu_{R_g}(x_0) = \bar{\beta}_0/n_g, \nu_{R_g}(x_1) = \bar{\beta}_1/n_g, \dots, \nu_{R_g}(x_{g-1}) = \bar{\beta}_{g-1}/n_g.$$

Using this and Proposition 2.13, we get that $\nu_{R_g}(x_g) = \frac{n_g-1}{n_g} \bar{\beta}_{g-1}$ and thus $x_g \circ \varphi = w^{n_g-1} \bar{\beta}_{g-1}/n_g \cdot s$.

As in the case $\ell_E > 0$, for $j = 0, \dots, g-1$, we have $x_j \circ \varphi = w^{\bar{\beta}_j/n_g}(\delta_{0,j} + w(\dots))$ with $\delta_{0,j} \in k^*$. Set $\delta_{I_0} = \prod_{j=0}^{g-1} \delta_{0,j}^{b_j} \in k^*$. Taking into account that $n_{g-1} \bar{\beta}_{g-1} < \bar{\beta}_g$ (see (2.2)) and denoting by m_g the difference $\bar{\beta}_g - n_{g-1} \bar{\beta}_{g-1}$, from the expansion (2.7) of x_A we obtain that

$$x_A \circ \varphi = w^{n_g-1} \bar{\beta}_{g-1} (s^{n_g} - w^{m_g} (\theta_{I_0}^A \delta_{I_0} + w(\dots))).$$

Hence $A^\varphi = Z(h_A)$, where $h_A = s^{n_g} - w^{m_g} (\theta_{I_0}^A \delta_{I_0} + w(\dots)) \in k[[w, s]]$. Notice that the integers n_g and m_g must be coprime (otherwise A^φ would not be a branch) and the compact edge of the Newton polygon of h_A is the line segment that joins $(0, n_g)$ and $(m_g, 0)$. Now, let us consider the fan $\Sigma_0 := \Sigma_{0,2}$ (see Notation 1.3) and the vector $u = (n_g, m_g)$. Let $\pi: S(\pi) \rightarrow S(\varphi)$ be the toric modification, defined with respect to the local coordinates (w, s) , associated to the minimal regular subdivision Σ of Σ_0 containing the ray $\mathbb{R}_{\geq 0}u$. This morphism is the minimal embedded resolution of A^φ and $\psi = \varphi \circ \pi$. We refer to [GP10] for details. In the chart corresponding to the cone $\mathbb{R}_{\geq 0}u + \mathbb{R}_{\geq 0}v \in \Sigma$, where $v = (c, d)$ and $cm_g - dn_g = 1$, this morphism is given by $w = w_1^c s_1^{n_g}$, and $s = w_1^d s_1^{m_g}$.

We get that $h_A(w_1^c s_1^{n_g}, w_1^d s_1^{m_g}) = w_1^{dn_g} s_1^{n_g m_g} (1 - \theta_{I_0}^A \delta_{I_0} w_1 + s_1(\dots))$. The exceptional divisor E on this chart is $Z(s_1)$. Thus, A^ψ intersects E at the point with coordinates $(w_1, s_1) = (w_1(A), 0)$, where $w_1(A) = 1/\theta_{I_0}^A \delta_{I_0}$. If we apply the previous construction to $B \in \mathcal{C}_E(\psi)$, we get that $w_1(B)$ equals $1/\theta_{I_0}^B \delta_{I_0}$. It follows that (1) and (2) are equivalent.

Assume now that $g = 1$. The expansion (2.7) of x_A is of the form

$$x_A = x_1^{\bar{\beta}_0} - \theta_{I_0}^A x_0^{\bar{\beta}_1} + \sum_{\bar{\beta}_0 i_0 + \bar{\beta}_1 i_1 > \bar{\beta}_0 \bar{\beta}_1} \theta_{I_0}^A x_0^{i_0} x_1^{i_1}.$$

Let Σ_0 be as above and $u = (\bar{\beta}_0, \bar{\beta}_1)$. Then, ψ is the toric modification, defined with respect to the local coordinates (x_0, x_1) , associated to the minimal regular subdivision of Σ_0 containing $\mathbb{R}_{\geq 0}u$. Choosing the chart as above, we see that (1) and (2) are also equivalent in this case. \square

We prove now the announced generalization a theorem of Goldin and Teissier:

Theorem 2.26. *Let $C = \cup_{j=1}^r C_j$ be a reduced singular plane curve with $r \geq 1$ branches embedded in k^2 by the map (2.1). Let x_0, \dots, x_m be defining functions of a sequence of maximal contact curves L_0, \dots, L_m of C which satisfies (2.5). Let $C \hookrightarrow k^{m+1}$ be the embedding defined by the functions x_0, \dots, x_m . Then, any regular subdivision Σ of the fan $\Sigma_{0,m+1}$ such that Σ contains the ray ρ_j spanned by $\text{ord}(\eta_j)$, for $1 \leq j \leq r$, defines a toric embedded resolution of C .*

Proof. The functions x_0, \dots, x_m satisfy conditions (1), (2), and (3) of Corollary 1.17 (see the discussion at the beginning of section 2.2 and Proposition 2.24.(1)). Therefore, the theorem follows directly from Corollary 1.17 if $r = 1$, or $r > 1$ and the order vectors $\text{ord}(\eta_j)$, for $1 \leq j \leq n$, are pairwise different (see Corollary 1.19). So let us assume that $\text{ord}(\eta_j) = \text{ord}(\eta_{j'})$ for some $1 \leq j < j' \leq r$. It was shown in the proof of Corollary 1.17 that the strict transforms \tilde{C}_j and $\tilde{C}_{j'}$ are smooth and transversal to the orbit $O(\rho)_\Sigma$, where $\rho := \rho_j = \rho_{j'}$. Next we show that the special points of \tilde{C}_j and $\tilde{C}_{j'}$ are different. This is enough to end the proof.

Denote by $\tilde{\eta}_j$ (resp. $\tilde{\eta}_{j'}$) the lifting of the arc η_j (resp. $\eta_{j'}$) to X_Σ . Take $\sigma \in \Sigma$ spanned by a basis $v_0 \in \rho, v_1, \dots, v_m$ of $N = \mathbb{Z}^{m+1}$. The affine toric variety X_σ is isomorphic to k^{m+1} with coordinates (U_0, \dots, U_m) , where $U_i = \chi^{v_i}$ for $i = 0, \dots, m$. The parametrization $\tilde{\eta}_j$ of \tilde{C}_j , in terms of these coordinates, is $(U_0 \circ \tilde{\eta}_j, \dots, U_m \circ \tilde{\eta}_j)$, and we have an analogous expression for the parametrization $\tilde{\eta}_{j'}$ of $\tilde{C}_{j'}$. Since $\text{ord}(\eta_j) = \text{ord}(\eta_{j'})$, we have the identities for $0 \leq i \leq m$:

$$X_i \circ \eta_j = x_i \circ \eta_{C_j} = d_i t_j^{\alpha_i} + \dots, \quad X_i \circ \eta_{j'} = x_i \circ \eta_{C_{j'}} = d'_i t_{j'}^{\alpha_i} + \dots,$$

where $d_i, d'_i \in k^*$, we have omitted higher order terms, and $(\alpha_0, \dots, \alpha_{m+1})$ are the coordinates of $\text{ord}(\eta_j)$ with respect to the canonical basis of $\mathbb{Z}^{m+1} = N$.

Let us assume that the special points of \tilde{C}_j and $\tilde{C}_{j'}$ coincide and proceed by contradiction. By assumption, there exist $c_0, c'_0, c_1, \dots, c_m \in k^*$ such that:

$$\begin{aligned} U_0 \circ \tilde{\eta}_j &= c_0 t_j + \dots, & U_i \circ \tilde{\eta}_j &= c_i + \dots, \text{ for } 1 \leq i \leq m; \\ U_0 \circ \tilde{\eta}_{j'} &= c'_0 t_{j'} + \dots, & U_i \circ \tilde{\eta}_{j'} &= c_i + \dots, \text{ for } 1 \leq i \leq m, \end{aligned}$$

where we have omitted higher order terms (see (1.4)).

For $1 \leq i \leq m$, denote by $(a_{0,i}, \dots, a_{m,i})$ the coordinates of v_i with respect to the canonical basis of $\mathbb{Z}^{m+1} = N$. Taking into account the monomial expression of the toric modification in the chart of X_Σ defined by the cone σ , we obtain:

$$d_i = c_0^{\alpha_i} c_1^{a_{i,1}} \dots c_m^{a_{i,m}}, \quad d'_i = (c'_0)^{\alpha_i} c_1^{a_{i,1}} \dots c_m^{a_{i,m}}, \text{ for } 0 \leq i \leq m.$$

Set $\kappa_i := c_1^{a_{i,1}} \dots c_m^{a_{i,m}} \in k^*$, for $i = 0, \dots, m$. Without loss of generality, we may assume that x_0, \dots, x_g is a minimal generating sequence of $\nu := \nu_{E_{C_j}}$ as in Proposition 2.11. In comparison with Notation 2.14, we have that $\alpha_i = \bar{\beta}_i$, for $i = 0, \dots, g$. We consider a (x_0, \dots, x_g) -adic expansion (2.7) of a defining function x_{C_j} of C_j . Substituting x_i by the series $x_i \circ \eta_{C_j}$, for $i = 0, \dots, g$, in the expansion (2.7), provides an expression

$$0 = x_{C_j} \circ \eta_{C_j} = (d_g^{n_g} - \theta_{I_0}^{C_j} \cdot d_0^{b_0} d_1^{b_1} \dots d_{g-1}^{b_{g-1}}) t_j^{n_g \bar{\beta}_g} + \dots,$$

where we have omitted the terms of higher order on the right side. It follows that:

$$(2.12) \quad d_g^{n_g} - \theta_{I_0}^{C_j} \cdot d_0^{b_0} d_1^{b_1} \dots d_{g-1}^{b_{g-1}} = 0.$$

Now substitute $d_i = \kappa_i c_0^{\alpha_i}$ in (2.12). Taking into account the condition (2.8) we get the equality:

$$(\kappa_g^{n_g} - \theta_{I_0}^{C_j} \cdot \kappa_0^{b_0} \kappa_1^{b_1} \dots \kappa_{g-1}^{b_{g-1}}) c_0^{n_g \bar{\beta}_g} = 0.$$

Since $c_0 \neq 0$, we conclude that $\theta_{I_0}^{C_j} = \kappa_g^{n_g} / \kappa_0^{b_0} \kappa_1^{b_1} \cdots \kappa_{g-1}^{b_{g-1}}$. According to Proposition 2.24.(2), we have that $E_{C_j} = E_{C_{j'}}$. Hence, by the same argument applied to $C_{j'}$, it follows that $\theta_{I_0}^{C_j} = \theta_{I_0}^{C_{j'}}$. This gives a contradiction with Proposition 2.25. \square

Example 2.27. Let C be the curve of Example 2.17. The branches L_0, L_1, L_2 defined by $x_0 = X$, $x_1 = Y$, and $x_2 = Y^2 - X^3 + X^4$, respectively, form a sequence of maximal contact curves of C which satisfies (2.5). Let $C \hookrightarrow k^3$ be the embedding defined by this sequence. With the notation of Example 2.17, for $1 \leq i \leq 2$, we have $(L_0, C_i) = 2$, $(L_1, C_i) = 3$, and $(L_2, C_i) = 8$. Hence $\text{ord}(\eta_1) = \text{ord}(\eta_2) = (2, 3, 8)$. By Theorem 2.26, any regular subdivision of $\Sigma_{0,3}$ containing the ray spanned by $v_3 := (2, 3, 8)$ defines a toric embedded resolution of C . Let us study the strict transforms of $C \subset k^3$ and $S = k^2 \subset k^3$ under the monomial map $\pi := \pi_{\sigma_{0,3}}^\sigma$ associated to the regular cone $\sigma \subset \sigma_{0,3}$ generated by the vectors $v_1 = (1, 1, 2)$, $v_2 = (2, 3, 7)$, and v_3 . We take coordinates in $X_{\Sigma_{0,3}} = k^3$, and also in X_σ , as in Example 1.20.

On the one hand, the surface $S \subset k^3$ has equation $X_3 - (X_2^2 - X_1^3 + X_1^4) = 0$. On the other hand, the points of $C \subset k^3$ satisfy the previous equation and $(X_3 - 2X_1^4)(X_3 - X_1^4 - X_1^5) = 0$. Thus the points in the strict transform \tilde{C} of C under π satisfy the following two equations:

$$(2.13) \quad \begin{aligned} 1 - U_1 - U_2 U_3^2 + U_1^2 U_2^2 U_3^2 &= 0, \\ (1 - 2U_1^2 U_2)(1 - U_1^2 U_2 - U_1^3 U_2^3 U_3^2) &= 0. \end{aligned}$$

One checks easily that the strict transform \tilde{S} of S , which is given by equation (2.13), is smooth. The points $(1, t, 0) \in \tilde{S}$, for $t \in k$, are those in the intersection with the exceptional locus of π . For $t = 1/2$ (resp. $t = 1$) we find the intersection with \tilde{C}_1 (resp. \tilde{C}_2).

3. EMBEDDED RESOLUTION AND LOCAL TROPICALIZATION

In this section we consider a singular curve $C \subset S = k^2$ and the embedding $S = k^2 \subset k^{m+1}$ defined by the map (2.10). The goal of this section is to prove that there exists a regular subdivision Σ of the positive quadrant $\Sigma_{0,m+1}$ such that the restriction of π_Σ to the strict transform of S is the minimal embedded resolution of the given plane curve singularity $C \subset S$. In order to prove this we describe the *local tropicalization* of S associated to this embedding.

Throughout this section we keep the notation introduced at the beginning of section 2.2. In particular, $L_0 = Z(x_0), \dots, L_m = Z(x_m)$ is a sequence of maximal contact curves of C satisfying (2.5). We call I_S the kernel of the surjective homomorphism (2.10), that is, the defining ideal of $S \subset k^{m+1}$.

3.1. The local tropicalization of S defined by the maximal contact curves of C . We start by recalling the definition of a semivaluation of a local domain with values in the ordered set $\overline{\mathbb{R}}_{\geq 0} := \mathbb{R}_{\geq 0} \cup \{\infty\}$.

Definition 3.1. Let R be a local domain containing the field k . A *semivaluation* of R is a function $\nu: R \rightarrow \overline{\mathbb{R}}_{\geq 0}$ such that:

- (1) $\nu(fg) = \nu(f) + \nu(g)$ for all $f, g \in R$;
- (2) $\nu(f + g) \geq \min(\nu(f), \nu(g))$ for all $f, g \in R$;
- (3) $\nu(0) = \infty$ and $\nu(\lambda) = 0$ for all $\lambda \in k^*$.

A semivaluation ν of R is *centered in R* if $\nu(m_R) \subseteq (0, \infty]$, where m_R denotes the maximal ideal of R . The semivaluation ν is a *valuation* if it takes the value ∞ only at 0, and it is *trivial* if $\nu(R) = \{0, \infty\}$.

Let us come back to the case $R = k[[X, Y]]$. If $A = Z(h)$ is a plane branch defined by a function $h \in R$, then we set $\nu(A) := \nu(h)$ for every semivaluation ν of R . Associated to a plane branch A , we have the *intersection semivaluation* I^A , defined by $I^A(h) = (A \cdot Z(h))$ for any $h \in R$, the *vanishing order semivaluation* ord^A , whose value on a function $h \in R$ is the order of vanishing of h along the branch A , and the *trivial valuation* triv^A which takes the value ∞ on any function vanishing on A and zero otherwise. The trivial valuations triv^O and triv^S associated to the point O and the surface S respectively, are defined analogously to triv^A .

The *semivaluation space* \mathcal{V} of R (or of the smooth surface germ S) is the set of semivaluations of R , endowed with the topology of pointwise convergence, that is, with the restriction of the product topology of $[0, \infty]^R$. We denote by $\mathcal{V}_{L_0} = \{\nu \in \mathcal{V} \mid \nu(x_0) = 1\}$ the subspace of *normalized semivaluations with respect to L_0* . The semivaluation space \mathcal{V} has been intensively studied after Favre and Jonsson's book [FJ04], see also [Jon15, GGP19a]. The normalized semivaluation space \mathcal{V}_{L_0} , which is also called the *(relative) valuative tree*, is an \mathbb{R} -tree rooted at ord^{L_0} (see [FJ04, section 3.9] and [GGP19a]). One has the following property:

Lemma 3.2. (see [GGP19a, section 6]) *Let $\nu \in \mathcal{V}_{L_0}$ and let A be a plane branch. If $\nu(A) = \infty$ then we have that $\nu = I_{L_0}^A := \frac{1}{(L_0 \cdot A)} \cdot I^A$. If $\nu(A) = 0$, then one has $A = L_0$ and $\nu = \text{ord}^{L_0}$.*

We consider the following notion of *local tropicalization* introduced in [PS13] (see also [St17]).

Definition 3.3. The *local tropicalization* $\text{Trop}_{\geq 0}(I)$ of an ideal $I \subset k[[X_0, \dots, X_m]]$ is the set of tuples $(\mu(X_0), \dots, \mu(X_m)) \in \overline{\mathbb{R}}_{\geq 0}^{m+1}$, for μ running through the semivaluations μ of $k[[X_0, \dots, X_m]]$ such that $\mu|_I = \infty$.

We have also a notion of tropicalization map which is defined in terms of $x_0, \dots, x_m \in m_R$:

Definition 3.4. The *tropicalization map* defined by the tuple (x_0, \dots, x_m) is

$$\begin{aligned} \text{trop}: \mathcal{V} &\rightarrow \overline{\mathbb{R}}_{\geq 0}^{m+1} \\ \nu &\mapsto (\nu(x_0), \dots, \nu(x_m)) \end{aligned}$$

Lemma 3.5. *With the previous notation, we have $\text{Trop}_{\geq 0}(I_S) = \text{trop}(\mathcal{V})$.*

Proof. If μ is a semivaluation of $k[[X_0, \dots, X_m]]$ verifying that $\mu|_{I_S} = \infty$, then one has a semivaluation $\bar{\mu}$ of the quotient ring $k[[X_0, \dots, X_m]]/I_S = R$ by setting $\bar{\mu}(\phi + I_S) := \mu(\phi)$ for every $\phi \in k[[X_0, \dots, X_m]]$, and every non-trivial semivaluation of R can be obtained in this way. \square

The aim of this section is to describe the *finite local tropicalization* of I_S , which is defined as $\text{Trop}_{\geq 0}(I_S) \cap \mathbb{R}_{\geq 0}^{m+1}$. By Lemma 3.5, we have the equality $\text{Trop}_{\geq 0}(I_S) \cap \mathbb{R}_{\geq 0}^{m+1} = \text{trop}(\mathcal{U})$, where \mathcal{U} denotes the set of semivaluations $\nu \in \mathcal{V}$ such that $\nu(x_j) < \infty$, for $j = 0, \dots, m$.

In what follows we denote by e_0, \dots, e_m the canonical basis of \mathbb{R}^{m+1} . Given $w \in \mathbb{R}^{m+1}$, we write $w = \sum_{0 \leq j \leq m} w_j e_j$. If $X \subset \mathbb{R}^{m+1}$, we denote by $\mathbb{R}_{>0}X$ the set $\{ax \mid a \in \mathbb{R}_{>0}, x \in X\}$, and similarly $\mathbb{R}_{\geq 0}X := \mathbb{R}_{>0}X \cup \{0\}$.

Lemma 3.6. *We have the following equalities:*

- (1) $\text{trop}(\mathcal{U}) = (\text{trop}(\mathcal{U}) \cap \mathbb{R}_{>0}^{m+1}) \cup \mathbb{R}_{\geq 0}e_0 \cup \dots \cup \mathbb{R}_{\geq 0}e_m$.
- (2) $\text{trop}(\mathcal{U}) \cap \mathbb{R}_{\geq 0}^{m+1} = \mathbb{R}_{>0} \left(\text{trop} \left(\mathcal{V}_{L_0} \setminus \left\{ \text{ord}^{L_0}, I_{L_0}^{L_1}, \dots, I_{L_0}^{L_m} \right\} \right) \right)$.

Proof. If A is a plane branch, then $\text{ord}^A \in \mathcal{U}$. In addition, the vector $\text{trop}(\text{ord}^A)$ is equal to e_j if $A = L_j$ for some $0 \leq j \leq m$, and zero otherwise. If $\nu \in \mathcal{U}$ is a trivial semivaluation then $\text{trop}(\nu) = 0$. The formula in (1) follows from these observations and Lemma 3.2.

By direct inspection we get that $\text{trop}(\mathcal{U}) \cap \mathbb{R}_{>0}^{m+1} = \mathbb{R}_{>0}(\text{trop}(\mathcal{V}_{L_0}) \cap \mathbb{R}_{>0}^{m+1})$. In order to finish the proof of (2) it is enough to check that the set $\text{trop}(\mathcal{V}_{L_0}) \cap \mathbb{R}_{>0}^{m+1}$ is the image of $\mathcal{V}_{L_0} \setminus \{\text{ord}^{L_0}, I_{L_0}^{L_1}, \dots, I_{L_0}^{L_m}\}$ under the tropicalization map. This follows again from Lemma 3.2. \square

We will study the set $\text{trop}(\mathcal{U})$ by using properties of the semivaluation spaces \mathcal{V} and \mathcal{V}_{L_0} , in connexion with the Eggers-Wall trees which we introduce briefly as follows.

3.1.1. Eggers-Wall trees. We consider a plane curve singularity $C = \cup_{j=1}^r C_j$ with $r \geq 1$ branches and a smooth branch L_0 . We describe some features of the Eggers-Wall tree associated to C with respect to L_0 , see [GGP19a] for precise definitions and details.

Let us introduce first some basic notation about trees (see [GGP19a, section 2]). If a, b and c are three points in a tree T , we denote by $\langle a, b, c \rangle$ the unique point of intersection of the segments $[a, b]$, $[b, c]$ and $[a, c]$. We call the point $\langle a, b, c \rangle$ the *center* of the *tripod* $\{a, b, c\}$. If the tree T is rooted at r , we set $a \leq_r b$ if $[r, a] \subseteq [r, b]$ for $a, b \in T$.

The Eggers-Wall tree $\Theta_{L_0}(C)$ is a rooted tree endowed with an *exponent function* \mathbf{e}_{L_0} and an *index function* \mathbf{i}_{L_0} :

$$\mathbf{e}_{L_0}: \Theta_{L_0}(C) \rightarrow \overline{\mathbb{R}}_{\geq 0}, \quad \mathbf{i}_{L_0}: \Theta_{L_0}(C) \rightarrow \mathbb{Z}_{>0}.$$

If k is of characteristic zero, this tree and these functions are usually defined in terms of the Puiseux expansions with respect to x_0 of the branches of C different from L_0 . If k is of arbitrary characteristic $\Theta_{L_0}(C)$ may be defined as the convex hull of the points labelled by L_0 and the branches of C in the *fan tree* associated to a toroidal pseudo-resolution of C with respect to L_0 . The functions \mathbf{e}_{L_0} and \mathbf{i}_{L_0} are determined then by the *slope function* on the fan tree (see [GGP19b, Remark 5.37]).

A point $P \in \Theta_{L_0}(C)$ is *rational* if $\mathbf{e}_{L_0}(P) \in \mathbb{Q}_{>0}$. The *marked points* of $\Theta_{L_0}(C)$ are its ends (i.e. the root labelled by L_0 and the leaves labelled by C_j , $1 \leq j \leq r$), its ramification points, and the points of discontinuity of the index function. By definition, the index function is constant on any segment of the form $(P, Q]$, where P and Q are two consecutive marked points.

We consider $\Theta_{L_0}(C)$ as a poset with respect to \leq_{L_0} . The exponent function in restriction to $[L_0, C_j]$ is a homeomorphism of posets onto $[0, \infty]$, for $1 \leq j \leq r$ with $C_j \neq L_0$. In addition, the functions \mathbf{e}_{L_0} and \mathbf{i}_{L_0} determine the *contact complexity function*,

$$\mathbf{c}_{L_0}: \Theta_{L_0}(C) \rightarrow \overline{\mathbb{R}}_{\geq 0},$$

defined by $\mathbf{c}_{L_0}(P) = \int_{L_0}^P \frac{de_{L_0}}{i_{L_0}}$. The knowledge of the functions \mathbf{i}_{L_0} and \mathbf{c}_{L_0} allow us to recover the intersection multiplicity of pairs of different branches of $C \cup L_0$. One has that $(L_0 \cdot C_j) = \mathbf{i}_{L_0}(C_j)$ for $j \in \{1, \dots, r\}$ (see [GGP19a, Rem. 3.25]). In addition, by [GGP19a, Cor. 3.26], we have

$$(3.1) \quad (C_i \cdot C_j) = \mathbf{i}_{L_0}(C_i)\mathbf{i}_{L_0}(C_j)\mathbf{c}_{L_0}(\langle L_0, C_i, C_j \rangle) \text{ for } i, j \in \{1, \dots, r\}, \text{ and } i \neq j.$$

For any branch A different from L_0 the *attaching point* of A to the tree $\Theta_{L_0}(C)$ is

$$P_A := \max\{\langle L_0, A, C_j \rangle \mid j = 1, \dots, r\} \in \Theta_{L_0}(C),$$

where the tripods are viewed in $\Theta_{L_0}(C \cup A)$ and the maximum is taken with respect to \leq_{L_0} .

Example 3.7. Assume that L_0, L_1, \dots, L_g is a sequence of maximal contact curves of a branch A of C . Then, the discontinuity locus of the restriction of the index function \mathbf{i}_{L_0} to the segment $[L_0, A]$ is equal to the set of points $\langle L_0, L_j, A \rangle$, for $j = 1, \dots, g$, where the tripods are viewed in the tree $\Theta_{L_0}(C \cup L_1 \cup \dots \cup L_g)$. With Notation 2.2, one has that $\mathbf{i}_{L_0}(\langle L_0, L_1, A \rangle) = 1 = \mathbf{i}_{L_0}(L_1)$ and $\mathbf{i}_{L_0}(\langle L_0, L_j, A \rangle) = n_1 \dots n_{j-1} = \mathbf{i}_{L_0}(L_j)$ for $j = 2, \dots, g$. By (3.1) we get that $\mathbf{c}_{L_0}(\langle L_0, L_j, A \rangle) = \bar{\beta}_0^{-1} \bar{\beta}_1$ and $\mathbf{c}_{L_0}(\langle L_0, L_j, A \rangle) = \bar{\beta}_0^{-1} (n_1 \dots n_{j-1})^{-1} \bar{\beta}_j$ for $j = 2, \dots, g$. In

addition, we get $\mathbf{e}_{L_0}(\langle L_0, L_j, A \rangle) = \beta_j/\beta_0$, for $j = 1, \dots, g$, where $\beta_0 = \bar{\beta}_0$, $\beta_1 = \bar{\beta}_1$ and $\beta_j - \beta_{j-1} = \bar{\beta}_j - n_{j-1}\bar{\beta}_{j-1}$ for $j = 2, \dots, g$. In addition, $\mathbf{i}_{L_0}(\langle L_0, L_1, A \rangle) = 1 = \mathbf{i}_{L_0}(L_1)$ and $\mathbf{i}_{L_0}(\langle L_0, L_1, A \rangle) = n_1 \dots n_{j-1} = \mathbf{i}_{L_0}(L_j)$ for $j = 2, \dots, g$.

If k is of characteristic zero, then the sequence $(\beta_0, \beta_1, \dots, \beta_g)$ is the *characteristic* of the branch A , which can be defined in terms of the Newton Puiseux series of A with respect to L_0 (see Definition 3.2 of [Zar06]).

Example 3.8. In this example we assume for simplicity that the field k is of characteristic zero. Let us consider the branches $L_0 = Z(X)$, $L_1 = Z(Y)$, and L_j , for $j = 2, \dots, 5$, parametrized respectively by the Newton-Puisseux series $\zeta_2 := X^{5/3}$, $\zeta_3 := X^{5/3} + X^2 + X^{5/2}$, $\zeta_4 := X^{3/2}$, and $\zeta_5 := X^{3/2} + X^{7/4}$. We have represented the Eggers-Wall tree $\Theta_{L_0}(D)$ of the curve $D := \cup_{j=1}^5 L_j$ in Figure 2. The marked points are $L_0, \dots, L_5, P_1, \dots, P_4$. In Figure 2 is indicated the constant value of the index function on every interval $(Q, Q']$ between consecutive marked points $Q \leq_{L_0} Q'$. We have that $\mathbf{e}_{L_0}(P_1) = 3/2$, $\mathbf{e}_{L_0}(P_2) = 5/3$, $\mathbf{e}_{L_0}(P_3) = 5/2$, $\mathbf{e}_{L_0}(P_4) = 7/4$, while $\mathbf{e}_{L_0}(L_0) = 0$ and $\mathbf{e}_{L_0}(L_j) = \infty$ for $j = 1, \dots, 5$. By definition, the value of the contact complexity function \mathbf{c}_{L_0} at a point P of the tree is just a finite sum. One has $\mathbf{c}_{L_0}(P_3) = \frac{25}{18}$, $\mathbf{c}_{L_0}(P_1) = \frac{3}{2}$, $\mathbf{c}_{L_0}(P_2) = \frac{5}{3}$, $\mathbf{c}_{L_0}(P_4) = \frac{13}{8}$, and $\mathbf{c}_{L_0}(L_0) = 0$, and $\mathbf{c}_{L_0}(L_j) = \infty$ for $j = 1, \dots, 5$.




FIGURE 2. The Eggers-Wall tree of Example 3.8

Let L_0, \dots, L_m be a sequence of maximal contact curves of C . The following lemma shows a particular property of the Eggers-Wall tree $\Theta_{L_0}(D)$, where we recall that $D = \cup_{j=0}^m L_j$.

Lemma 3.9. *The leaves of the closures of the levels of the index function \mathbf{i}_{L_0} on the tree $\Theta_{L_0}(D)$ are leaves of $\Theta_{L_0}(D)$.*

Proof. The closure of a connected component Ξ of the index function on the tree $\Theta_{L_0}(D)$ is a tree rooted at the point of minimum exponent of Ξ . If P is an end of Ξ which is not an end of $\Theta_{L_0}(D)$ then P , seen on $\Theta_{L_0}(D)$, is a point of discontinuity of \mathbf{i}_{L_0} . Take a branch L of D such that $P <_{L_0} L$. By Lemma 2.16, there exists $1 \leq g \leq m$ such that, up to relabelling L_1, \dots, L_m , we have that L_0, \dots, L_{g-1} is a sequence of maximal contact curves of $L = L_g$. Then, P is also a point of discontinuity of the restriction of \mathbf{i}_{L_0} to the segment $[L_0, L_g]$. By Example 3.7 applied to the branch L_g of D and its sequence of maximal contact curves L_0, \dots, L_{g-1} , there exists $j \in \{1, \dots, g-1\}$ such that $P = \langle L_0, L_j, L_g \rangle$ and then $\mathbf{i}_{L_0}(P) = \mathbf{i}_{L_0}(L_j)$. We get that $P <_{L_0} L_j$ with $L_j \in \Xi$, which is a contradiction. \square

3.1.2. *The finite local tropicalization.* In this section we prove that the finite local tropicalization $\text{trop}(\mathcal{U})$ of I_S is the support of a fan and we describe it in terms of the Eggers-Wall tree. We use the functions \mathbf{e}_{L_0} , \mathbf{i}_{L_0} and \mathbf{c}_{L_0} , which can be expressed in terms of natural "coordinate functions" on the valutive tree \mathcal{V}_{L_0} : the *log-discrepancy*, the *multiplicity* and the *self-interaction* (we use here the terminology of [GGP19a]). These functions are crucial in the work of Favre and Jonsson [FJ04], see also Jonsson's survey [Jon15, section 7].

The embedding (3.2) below is described in [GGP19a, Section 8] if k is a field of characteristic zero, and in [GGP19b, Remark 5.37] if k has arbitrary characteristic.

Proposition 3.10. [GGP19a, Prop. 8.10 and Th. 8.11] *There is an embedding of rooted trees:*

$$(3.2) \quad V_{L_0} : \Theta_{L_0}(D) \hookrightarrow \mathcal{V}_{L_0}, \quad P \mapsto V_{L_0}^P$$

such that for any $P \in \Theta_{L_0}(D)$ and any plane branch A on S we have

$$(3.3) \quad V_{L_0}^P(A) = \begin{cases} (L_0 \cdot A) \cdot \mathbf{c}_{L_0}(\langle L_0, P, A \rangle) & \text{if } A \neq L_0, \\ 1 & \text{if } A = L_0, \end{cases}$$

where $\langle L_0, P, A \rangle$ is the center of the tripod defined by L_0 , P , and A on the Eggers-Wall tree $\Theta_{L_0}(D \cup A)$.

If P is a leaf of $\Theta_{L_0}(D)$, that is, $P = L_j$ for some $1 \leq j \leq m$, then $V_{L_0}^P = I_{L_0}^{L_j}$. In the case of the root $P = L_0$, we have $V_{L_0}^P = \text{ord}^{L_0}$. Finally, if P is an interior point of $\Theta_{L_0}(D)$ then $V_{L_0}^P(A) \in \mathbb{R}_{>0}$ for any branch A , since the function \mathbf{c}_{L_0} restricted to the interior of the tree $\Theta_{L_0}(D)$ has values in $\mathbb{R}_{>0}$.

Definition 3.11. If P is an interior point of the tree $\Theta_{L_0}(D)$, we set $w^P := \text{trop}(V_{L_0}^P) \in \mathbb{R}_{>0}^{m+1}$. For $0 \leq j \leq m$, we define $w^{L_j} := e_j$.

Remark 3.12. Notice that $w^{L_0} = \text{trop}(V_{L_0}^{L_0}) = \text{trop}(\text{ord}^{L_0})$. However, for $1 \leq j \leq m$, we have $w^{L_j} = \text{trop}(\text{ord}^{L_j})$, while $\text{trop}(V_{L_0}^{L_j}) = \text{trop}(I_{L_0}^{L_j}) \notin \mathbb{R}_{\geq 0}^{m+1}$. One has that the semivaluations ord^{L_j} and $I_{L_0}^{L_j}$ are identified in the projective semivaluation space (see [GGP19a]).

Lemma 3.13. *Let P_1, P_2 be two consecutive marked points of $\Theta_{L_0}(D)$. Then, the cone spanned by $\{w^P \mid P \in [P_1, P_2]\}$ is equal to $\mathbb{R}_{\geq 0} w^{P_1} + \mathbb{R}_{\geq 0} w^{P_2}$.*

Proof. We may assume that $P_1 \leq_{L_0} P_2$. Take $j \in \{0, \dots, m\}$ and $P \in [P_1, P_2]$. By definition of the contact complexity function we get the following equalities:

$$(3.4) \quad \mathbf{c}_{L_0}(P) = \int_{L_0}^{P_1} \frac{d\mathbf{e}_{L_0}}{\mathbf{i}_{L_0}} + \int_{P_1}^P \frac{d\mathbf{e}_{L_0}}{\mathbf{i}_{L_0}} = \mathbf{c}_{L_0}(P_1) + \frac{1}{\mathbf{i}_{L_0}(P_2)} (\mathbf{e}_{L_0}(P) - \mathbf{e}_{L_0}(P_1)),$$

where the second equality holds since \mathbf{i}_{L_0} has constant value $\mathbf{i}_{L_0}(P_2)$ on the segment $(P_1, P_2]$.

We discuss the value $V_{L_0}^P(L_j)$ according to the relative position of L_j with respect P_1 and P_2 . Since no ramification point of the tree $\Theta_{L_0}(D)$ belongs to the segment (P_1, P_2) by definition of marked points, then:

- (1) $\langle L_0, L_j, P \rangle = P$ if $P_2 \leq_{L_0} L_j$;
- (2) $\langle L_0, L_j, P \rangle = \langle L_0, L_j, P_1 \rangle$, otherwise.

In case (2), from the definition of V_{L_0} we see that

$$(3.5) \quad V_{L_0}^P(L_j) = V_{L_0}^{P_1}(L_j).$$

In case (1), it follows from the definition of V_{L_0} and the equality (3.4) that

$$(3.6) \quad V_{L_0}^P(L_j) = (L_0 \cdot L_j) \left(\mathbf{c}_{L_0}(P_1) + \frac{1}{\mathbf{i}_{L_0}(P_2)} (\mathbf{e}_{L_0}(P) - \mathbf{e}_{L_0}(P_1)) \right).$$

We distinguish now two cases:

- The point P_2 is not a leaf of $\Theta_{L_0}(D)$. Then we consider the number

$$s(P) := \frac{\mathbf{e}_{L_0}(P) - \mathbf{e}_{L_0}(P_1)}{\mathbf{e}_{L_0}(P_2) - \mathbf{e}_{L_0}(P_1)},$$

which runs through $[0, 1]$ when P runs through $[P_1, P_2]$. From (3.3), (3.5), and (3.6) we obtain the convex combination $w^P = s(P)w^{P_2} + (1 - s(P))w^{P_1}$, which implies the assertion in this case.

- The point P_2 is a leaf of $\Theta_{L_0}(D)$. Then $P_2 = L_i$ for some $1 \leq i \leq m$. We consider the value

$$s(P) := \frac{(L_0 \cdot L_i)}{\mathbf{i}_{L_0}(P_1)} (\mathbf{e}_{L_0}(P) - \mathbf{e}_{L_0}(P_1)),$$

which runs through $[0, \infty]$ when P runs through $[P_1, P_2]$. By (3.3), (3.5), and (3.6) we get that $w^P = w^{P_1} + s(P)w^{P_2}$, for $P \in [P_1, P_2]$. This ends the proof in this case. \square

Lemma 3.14. *The map $\Theta_{L_0}(D) \rightarrow \mathbb{R}_{\geq 0}^{m+1}$, $P \mapsto w^P$ is injective and its restriction to the set of non-leaf points of the tree $\Theta_{L_0}(D)$ is continuous.*

Proof. Take $j \in \{1, \dots, m\}$. The restriction of \mathbf{c}_{L_0} to $[L_0, L_j]$ is an increasing homeomorphism onto $[0, \infty]$. For $P \in [L_0, L_j)$, we have $w_j^P = V_{L_0}^P(L_j) = (L_0 \cdot L_j)\mathbf{c}_{L_0}(P)$. Notice that $w_0^P = 1$ for every non-leaf point P . Hence the assertion on the continuity of this map follows. In addition, in order to prove the injectivity is enough to show that $w^P \neq w^Q$ for any pair of non comparable points $P, Q \in \Theta_{L_0}(D)$ with respect to the order \leq_{L_0} . The special case where one of these points is an end follows from Definition 3.11, using that $w^U = \text{trop}(V_{L_0}^U) \in \mathbb{R}_{>0}^{m+1}$ for any interior point U . Suppose now that P, Q are interior points. Let $1 \leq j \leq m$ be such that $P \in [L_0, L_j]$. Then, if $F := \langle P, Q, L_0 \rangle$ we have that $F <_{L_0} P$ and $F <_{L_0} Q$. Therefore, $w_j^Q = (L_0 \cdot L_j)\mathbf{c}_{L_0}(F) = w_j^F < w_j^P$, since the map \mathbf{c}_{L_0} is strictly increasing on $[L_0, L_j]$. \square

Notation 3.15. The augmented set of marked points $\mathcal{A}_{L_0}^C(D)$ consists of those points $Q \in \Theta_{L_0}(D)$ which are marked points when seen in $\Theta_{L_0}(D \cup C)$. See Remark 3.35 below for a geometrical interpretation of the set $\mathcal{A}_{L_0}^C(D)$.

As a consequence of Lemmas 3.13 and 3.14 we get that:

Proposition 3.16. *There exists a fan \mathcal{T} with respect to the lattice \mathbb{Z}^{m+1} consisting of cones of dimension at most 2, whose support is equal to*

$$|\mathcal{T}| := \mathbb{R}_{\geq 0}\{w^P \mid P \in \Theta_{L_0}(D)\}.$$

Its 2-dimensional cones are $\mathbb{R}_{\geq 0}w^{P_1} + \mathbb{R}_{\geq 0}w^{P_2}$, for $P_1, P_2 \in \mathcal{A}_{L_0}^C(D)$ consecutive points on the tree $\Theta_{L_0}(D)$.

Now, we prove that:

Theorem 3.17. *With the previous notation, we have $\text{trop}(\mathcal{U}) = |\mathcal{T}|$.*

Proof. Since $w^{L_j} = e_j$ for $j = 0, \dots, m$, it is enough to show that

$$(3.7) \quad \text{trop} \left(\mathcal{V}_{L_0} \setminus \left\{ \text{ord}^{L_0}, I_{L_0}^{L_1}, \dots, I_{L_0}^{L_m} \right\} \right) = \{w^P \mid P \text{ is an interior point of } \Theta_{L_0}(D)\}$$

(see Lemma 3.6). For any interior point $P \in \Theta_{L_0}(D)$, we have $w^P = \text{trop}(V_{L_0}^P)$, with $V_{L_0}^P \in \mathcal{V}_{L_0} \setminus \left\{ \text{ord}^{L_0}, I_{L_0}^{L_1}, \dots, I_{L_0}^{L_m} \right\}$. Therefore, the inclusion \supset in the equality (3.7) follows.

Next we show the other inclusion. Take $\nu \in \mathcal{V}_{L_0} \setminus \{\text{ord}^{L_0}, I_{L_0}^{L_1}, \dots, I_{L_0}^{L_m}\}$. Since the valuative tree \mathcal{V}_{L_0} is an \mathbb{R} -tree, and the set $V_{L_0}(\Theta_{L_0}(D))$ is a closed subtree of it, there exists a unique *attaching point* ν' of ν to $V_{L_0}(\Theta_{L_0}(D))$ (see [GGP19a, sections 2, 6 and 8]). The attaching point ν' is the unique point of \mathcal{V}_{L_0} such that the intersection of the interval $[\nu, \nu']$ with the subtree is reduced to $\{\nu'\}$. Notice that the semivaluations $\text{ord}^{L_0}, I_{L_0}^{L_1}, \dots, I_{L_0}^{L_m}$ are the ends of $V_{L_0}(\Theta_{L_0}(D))$, and these semivaluations are also ends of \mathcal{V}_{L_0} . We get that ν' must be of the form $\nu' = V_{L_0}^Q$ for some interior point $Q \in \Theta_{L_0}(D)$. Let us prove that $\text{trop}(\nu) = w^Q$, that is, $\nu(L_j) = w_j^Q$ for $1 \leq j \leq m$, since ν and ν' belong to \mathcal{V}_{L_0} .

We rely on some results of [GGP19a]. There exists a function $\langle \cdot, \cdot \rangle : \mathcal{V}^* \times \mathcal{V}^* \rightarrow [0, \infty]$, called the *bracket*, such that for any branch A on S and any $\mu \in \mathcal{V}^*$ we have

$$(3.8) \quad \langle I^A, \mu \rangle = \mu(A)$$

(see [GGP19a, Prop. 7.5 and 7.11]). In particular, if B is another plane branch we get that

$$(3.9) \quad \langle I^A, I^B \rangle = I^B(A) = (A \cdot B).$$

The *relative self-interaction* function $\mathbf{s}_{L_0} : \mathcal{V}_{L_0} \rightarrow [0, \infty]$ is defined by $\mathbf{s}_{L_0}(\mu) = \langle \mu, \mu \rangle$.

Let us consider $1 \leq j \leq m$. By the *generalized tripod formula* applied to the semivaluations I^{L_j} and ν with respect to the observer L_0 , the following equality holds (see [GGP19a, Prop. 7.18]):

$$(3.10) \quad \langle I^{L_j}, \nu \rangle = \langle I^{L_0}, \nu \rangle \langle I^{L_0}, I^{L_j} \rangle \mathbf{s}_{L_0}(\langle \text{ord}_{L_0}, I_{L_0}^{L_j}, \nu \rangle).$$

By definition of the attaching point, we have that

$$(3.11) \quad \langle \text{ord}_{L_0}, I_{L_0}^{L_j}, \nu \rangle = \langle \text{ord}_{L_0}, I_{L_0}^{L_j}, V_{L_0}^Q \rangle.$$

Taking into account (3.8), (3.9) and (3.11), we can reformulate (3.10) as follows:

$$(3.12) \quad \nu(L_j) = \nu(L_0)(L_0 \cdot L_j) \mathbf{s}_{L_0}(\langle \text{ord}_{L_0}, I_{L_0}^{L_j}, V_{L_0}^Q \rangle) = (L_0 \cdot L_j) \mathbf{s}_{L_0}(\langle \text{ord}_{L_0}, I_{L_0}^{L_j}, V_{L_0}^Q \rangle).$$

On the one hand, we have that $V_{L_0}^{L_0} = \text{ord}^{L_0}$ and $V_{L_0}^{L_j} = I_{L_0}^{L_j}$. On the other hand by [GGP19a, Th. 8.18] one has $\mathbf{s}_{L_0} \circ V_{L_0} = \mathbf{c}_{L_0}$. Combining these facts we reformulate the equality (3.12) as $\nu(L_j) = (L_0 \cdot L_j) \mathbf{c}_{L_0}(\langle L_0, L_j, Q \rangle) = w_j^Q$. \square

We will make use of the following lemma in section 3.2.

Lemma 3.18. *Let P be an interior point of $\Theta_{L_0}(D)$. The point P is rational if and only if $w^P \in \mathbb{Q}_{>0}^{m+1}$.*

Proof. By definition we have that $\mathbf{c}_{L_0}(Q) \in \mathbb{Q}_{>0}$ if and only if Q is a rational point. Taking into account this observation and (3.3), we see that w^P is rational if P is rational. Let us now show the converse. Assume that $w^P \in \mathbb{Q}_{>0}^{m+1}$. Take P_1, P_2 two consecutive marked points of $\Theta_{L_0}(D)$ such that $P \in [P_1, P_2]$ and $j \in \{1, \dots, m\}$ such that $P_2 \leq_{L_0} L_j$. By (3.6), we have that $w_j^P \in \mathbb{Q}_{>0}$ if and only if $\mathbf{e}_{L_0}(P) \in \mathbb{Q}_{>0}$. \square

Let P be a rational point of the Eggers-Wall tree $\Theta_{L_0}(D)$. As it is explained in [GGP19a, Def. 8.14], there exists a unique exceptional prime divisor E_P (up to birational transformation) such that its associated divisorial valuation ν_{E_P} is proportional to the valuation $V_{L_0}^P$. We say that E_P is the *representing divisor* of P (see [GGP19a, Prop. 8.16] for some of its properties). We have

$$w^P = (V_{L_0}^P(x_0), \dots, V_{L_0}^P(x_m)) = \frac{1}{\nu_{E_P}(x_0)} (\nu_{E_P}(x_0), \dots, \nu_{E_P}(x_m)).$$

Our next goal is the description of a generating sequence of the divisorial valuation ν_{E_P} , for P any rational point of $P \in \Theta_{L_0}(D)$. We need some notation and preliminary results.

We denote by $\mathbf{i}_{L_0}^+(P)$ the index of $(\mathbb{Z}, +)$ in the subgroup of $(\mathbb{Q}, +)$ generated by 1, $\mathbf{e}_{L_0}(P)$ and $\mathbf{e}_{L_0}(P')$ for $P' \leq_{L_0} P$ running through the points of discontinuity of the index function (see [GGP19a, Definition 3.14]).

Lemma 3.19. *Let $P \in \Theta_{L_0}(D)$ be a rational point. Let A be a branch such that $P \leq_{L_0} A$ in $\Theta_{L_0}(D \cup A)$ and the limit of $\mathbf{i}_{L_0}(P')$ when $P' \in (P, A)$ tends to P is equal to $\mathbf{i}_{L_0}^+(P)$. Then, A is a curvetta of E_P at the minimal resolution of ν_{E_P} .*

Proof. Let A be any branch such that $P \leq_{L_0} A$ and the limit of $\mathbf{i}_{L_0}(P')$, when $P' \in (P, A)$ tends to P , is $\geq \mathbf{i}_{L_0}^+(P)$. Denote by $\psi: (S(\psi), E(\psi)) \rightarrow (S, O)$ the minimal resolution of ν_{E_P} . We have the following formula for the intersection number at the surface $S(\psi)$:

$$(3.13) \quad (A^\psi \cdot E_P^\psi)_{S(\psi)} = (\mathbf{i}_{L_0}^+(P))^{-1} \cdot \mathbf{i}_{L_0}(A).$$

Formula (3.13) is consequence of the renormalization formulas (see Propositions 1.6.20 (1) and 1.6.22 (1) of [GGP19b], applied to P and the points of discontinuity of the index function \mathbf{i}_{L_0} on the segment $[L_0, P)$). In particular, if $\mathbf{i}_{L_0}^+(P) = \mathbf{i}_{L_0}(A)$ then A^ψ is smooth and intersects transversally E_P^ψ . Then, applying [GGP19b, Lemma 1.6.18] as before, we get that A^ψ does not intersect any other component of $E(\psi)$. Hence A belongs to $\mathcal{C}_{E_P}(\psi)$. \square

Corollary 3.20. *(see [Ro19, Theorem 4.160]) Let $P \in \Theta_{L_0}(D)$ be a rational point. Then, the sequence x_0, \dots, x_m is a generating sequence of the divisorial valuation ν_{E_P} .*

Proof. Take a branch $L \in \{L_1, \dots, L_m\}$ such that $P <_{L_0} L$ and $\mathbf{i}_{L_0}^+(P) = \mathbf{i}_{L_0}(L)$. Such a branch exists by Lemma 3.9. By Lemma 2.16 there exists $1 \leq g \leq m$ such that, up to relabelling L_1, \dots, L_m , one has that L_0, \dots, L_{g-1} is a sequence of maximal contact curves of the branch $L = L_g$. Denote by ψ the minimal resolution of ν_{E_P} . By Lemma 3.19 we have that $L_g \in \mathcal{C}_{E_P}(\psi)$. Since $\mathbf{i}_{L_0}^+(P) = \mathbf{i}_{L_0}(L_g)$ the point P belongs to the segment $[Q, L_g]$ where $Q = \langle L_0, L_{g-1}, L_g \rangle$. In addition, we have that $\mathbf{i}_{L_0}^+(Q) = \mathbf{i}_{L_0}(L_g)$. By Lemma 3.19, the branch L_g belongs to $\mathcal{C}_{E_Q}(\psi')$, where ψ' is the minimal resolution of ν_{E_Q} .

Take a branch B such that $Q \leq_{L_0} B$ and $\mathbf{i}_{L_0}^+(Q) = \mathbf{i}_{L_0}(B)$. Then, $B \in \mathcal{C}_{E_Q}(\psi')$ by Lemma 3.19. Assume in addition that $B^{\psi'} \cap L_g^{\psi'} = \emptyset$. Then, we have that $\nu_{E_Q}(L_g) = (L_g \cdot B)$ by Remark 2.6. We get also that $Q = \langle L_0, L_g, B \rangle$ (see the proof of [GGP19a, Prop. 8.16]). By the tripod formula (3.1) applied to L_0, L_g, B and L_0, L_{g-1}, L_g one has that

$$(3.14) \quad \nu_{E_Q}(L_g) = \mathbf{i}_{L_0}(L_g)^2 \mathbf{c}_{L_0}(Q) = \frac{\mathbf{i}_{L_0}(L_g)}{\mathbf{i}_{L_0}(L_{g-1})} (L_g \cdot L_{g-1}) = n_{g-1} \bar{\beta}_{g-1},$$

where $\bar{\beta}_0, \dots, \bar{\beta}_{g-1}$ denotes the minimal generating system of Γ_{L_g} and n_1, \dots, n_{g-1} is the associated sequence of integers as Notation 2.2. Similarly, we obtain that $\nu_{E_P}(L_g) = \mathbf{i}_{L_0}(L_g)^2 \mathbf{c}_{L_0}(P)$. Since the function \mathbf{c}_{L_0} is strictly increasing on $[Q, L_g]$ then

$$(3.15) \quad \nu_{E_P}(L_g) = \mathbf{i}_{L_0}(L_g)^2 \mathbf{c}_{L_0}(P) \geq \mathbf{i}_{L_0}(L_g)^2 \mathbf{c}_{L_0}(Q) = \nu_{E_Q}(L_g), \text{ for } P \in [Q, L_g],$$

with equality if and only if $P = Q$. By (3.14) and (3.15) it follows that $\ell_{E_Q} = 0$ while $\ell_{E_P} > 0$ if $P \in (Q, L_g)$. By Proposition 2.11 we deduce that a minimal generating sequence of ν_{E_P} is equal to x_0, \dots, x_{g-1} if $P = Q$, and x_0, \dots, x_g if $P \in (Q, L_g)$. \square

Remark 3.21. Let E be an exceptional prime divisor of the minimal embedded resolution ψ of C which intersects the strict transform of C . By Lemma 3.19 it follows that $E = E_Q$, where $Q \in \mathcal{A}_{L_0}^C(D)$ is the attaching point of C_j to the tree $\Theta_{L_0}(D)$. Then we have $(L_i \cdot C_j) = \nu_{E_Q}(L_i)$,

for $i = 0, \dots, m$, by Remark 2.6 since we assume the genericity conditions (2.5). This implies the equality $\text{ord}(\eta_j) = (\nu_{E_Q}(x_0), \dots, \nu_{E_Q}(x_m))$. Since $Q \in \mathcal{A}_{L_0}^C(D)$ by definition, the order vector $\text{ord}(\eta_j)$ defines a ray in the support of the fan \mathcal{T} (see Proposition 3.16).

Example 3.22. Let us come back to the case of Example 3.8. We have represented in Figure 3 the dual graph of the total transform of $D \cup L_0$ by its minimal embedded resolution. Take $x_j \in m_R$ such that $L_j = Z(x_j)$, for $j = 0, \dots, 5$. By Corollary 3.20, the sequence x_0, \dots, x_5 is a generating sequence of $(\nu_{E_{P_1}}, \dots, \nu_{E_{P_4}})$. Then, we use formula (3.3) to find the vector $w^{P_i} = (w_0^{P_i}, \dots, w_5^{P_i})$, for $i = 1, \dots, 4$. We obtain $w_0^{P_i} = 1$ and $w_j^{P_i} = (L_0 \cdot L_j) \mathbf{c}_{L_0}(\langle L_0, L_j, P_i \rangle)$, for $j = 1, \dots, 5$. In particular, for $i = 4$, we get that $\langle L_0, L_j, P_4 \rangle = P_1$ for $j = 1, 2, 3$ while $\langle L_0, L_j, P_4 \rangle = P_4$ for $j = 4, 5$. Then, one obtains $w^{P_4} = (1, 3/2, 9/2, 9, 13/4, 13/2)$, and similarly $w^{P_1} = (1, 3/2, 9/2, 9, 3, 6)$, $w^{P_2} = (1, 5/3, 5, 10, 3, 6)$, and $w^{P_3} = (1, 5/3, 25/6, 25/3, 3, 6)$.




FIGURE 3. The dual graph of the total transform of the curve in Example 3.8 under its minimal resolution.

3.2. Initial ideals associated to the local tropicalization and toric resolutions. In this section we consider the embedding $S = k^2 \subset k^{m+1}$ defined by the map (2.10). We describe the initial ideals $in_w(I_S)$ when $w \in \text{Trop}_{\geq 0}(I_S)$. Then, we prove that there exists a regular subdivision Σ of the positive quadrant $\Sigma_{0,m+1}$ such that the restriction of π_Σ to the strict transform of S is an embedded resolution of the given plane curve singularity $C \subset S$.

Given $\alpha = (\alpha_0, \dots, \alpha_m) \in \mathbf{Z}_{\geq 0}^{m+1}$, we denote $X^\alpha = X_0^{\alpha_0} \cdots X_m^{\alpha_m}$. Then we can write a power series $g \in k[[X_0, \dots, X_m]]$ as $g = \sum_{\alpha \in \mathbf{Z}_{\geq 0}^{m+1}} c_\alpha X^\alpha$. Let $w \in \mathbb{R}_{\geq 0}^{m+1}$. If $g \neq 0$, the w -weight of g is

$$\nu_w(g) := \min \{w \cdot \alpha \mid c_\alpha \neq 0\},$$

where \cdot stands for the usual scalar product of \mathbb{R}^{m+1} . The w -initial part of g , denoted $in_w(g)$, is the sum of all terms in g of w -weight $\nu_w(g)$. Set $\nu_w(0) = \infty$ and $in_w(0) = 0$. Actually, the map ν_w defines a valuation that is called the monomial valuation of weight w . Such a valuation is divisorial if and only if $w \in \mathbb{Q}^{m+1}$.

Let I be an ideal of $k[[X_0, \dots, X_m]]$. The w -initial ideal of I is the ideal generated by the set $\{in_w(g) \mid g \in I\}$. We denote this ideal by $in_w(I)$. Note that the w -initial parts of a system of generators g_1, \dots, g_ℓ of I do not generate in general $in_w(I)$. When this occurs, g_1, \dots, g_ℓ is a w -standard basis of I .

Given $w = (w_0, \dots, w_m) \in \mathbb{R}_{\geq 0}^{m+1}$, we call $\text{supp}(w)$ the set of indices i such that $w_i \neq 0$. We define $C_w(I)$ as the closure in the usual Euclidean topology on \mathbb{R}^{m+1} of the following set:

$$\{w' \in \mathbb{R}_{\geq 0}^{m+1} \mid \text{supp}(w) = \text{supp}(w') \text{ and } in_w(I) = in_{w'}(I)\}.$$

Then, the family $\{C_w(I) \mid w \in \mathbb{R}_{\geq 0}^{m+1}\}$ is finite and forms a rational polyhedral fan $\text{GF}(I)$ subdividing the fan $\Sigma_{0,m+1}$ (see [BT07, PS13, Tou05]). This fan is called the *Gröbner fan* of I and is a local version of the notion of Gröbner fan studied in [MR88].

The following result relates the local tropicalization and the Gröbner fan of I . For the proof, we refer the reader to [Tou05, Theorem 5.8] and [PS13, Theorem 11.2].

Proposition 3.23. *Let I be an ideal of $k[[X_0, \dots, X_m]]$.*

- (1) *Given $w \in \mathbb{R}_{\geq 0}^{m+1}$, $w \in \text{Trop}_{\geq 0}(I)$ if and only if $\text{in}_w(I)$ contains no monomial.*
- (2) *$\text{Trop}_{\geq 0}(I) \cap \mathbb{R}_{\geq 0}^{m+1}$ is a union of cones of $\text{GF}(I)$.*

The following definition is based upon that of non-degenerate functions with respect to their Newton polyhedra of Khovanskii and Kouchnirenko [Kou76, Kho78].

Definition 3.24 ([AGS13, Tev07]). *An ideal $I \subset k[[X_0, \dots, X_m]]$ is *Newton non-degenerate* if for every $w \in \mathbb{R}_{\geq 0}^{m+1}$, the variety $Z(\text{in}_w(I))$ does not have singularities in the torus $(k^*)^{m+1}$.*

The notion of Newton non-degenerate given in Definition 3.24 may be seen as a particular case of the notion of schön compactification of a subvariety of a torus introduced in [Tev07].

Remark 3.25. It is a straightforward consequence of Proposition 3.23 and the Nullstellensatz that $\text{Trop}_{\geq 0}(I) \cap \mathbb{R}_{\geq 0}^{m+1}$ can also be described as the set of all $w \in \mathbb{R}_{\geq 0}^{m+1}$ such that $Z(\text{in}_w(I)) \cap (k^*)^{m+1} \neq \emptyset$. Hence in order to decide whether the ideal I is Newton non-degenerate it is enough to verify that the condition given in Definition 3.24 is satisfied for those w in $\text{Trop}_{\geq 0}(I) \cap \mathbb{R}_{\geq 0}^{m+1}$. In addition, it is sufficient to check the condition for every vector in $\text{Trop}_{\geq 0}(I) \cap \mathbb{Q}_{\geq 0}^{m+1}$. Indeed, if $w \in \text{Trop}_{\geq 0}(I) \cap \mathbb{R}_{\geq 0}^{m+1}$ and $w \notin \mathbb{Q}_{\geq 0}^{m+1}$ then, by Proposition 3.23, there exists a cone $\sigma \in \text{GF}(I)$ such that $w \in \overset{\circ}{\sigma} \subset \text{Trop}_{\geq 0}(I)$. Since σ is a rational cone, we can take $w' \in \overset{\circ}{\sigma} \cap \mathbb{Q}_{\geq 0}^{m+1}$ and we get $\text{in}_w(I) = \text{in}_{w'}(I)$.

Theorem 3.26 (See [AGS13, Tev07]). *Let I be a Newton non-degenerate ideal of $k[[X_0, \dots, X_m]]$. Let Σ be a regular subdivision of the Gröbner fan of I and let $\pi_{\Sigma_{0,m+1}}^{\Sigma} : X_{\Sigma} \rightarrow k^{m+1}$ be the associated toric modification. Then, the strict transform of $Z(I)$ by $\pi_{\Sigma_{0,m+1}}^{\Sigma}$ is non-singular and transversal to the orbit stratification of the exceptional locus of $\pi_{\Sigma_{0,m+1}}^{\Sigma}$.*

In what follows we keep the notation of section 3.1. Let P be a rational point of $\Theta_{L_0}(D)$. Then, w^P belongs to $\text{Trop}_{\geq 0}(I_S) \cap \mathbb{Q}_{> 0}^{m+1}$. Let us relabel the functions x_0, \dots, x_m in such a way that x_0, \dots, x_g is a minimal generating sequence of the divisorial valuation ν_{E_P} (see Corollary 3.20). With the notation of Section 2.1, $g = g(E_P)$. Next we give a system of generators H_2, \dots, H_m of I_S that is well adapted to the computation of w^P -initial parts:

- For $j \in \{1, \dots, g-1\}$, we consider the (x_0, \dots, x_j) -expansion of x_{j+1} given in Proposition 2.21 and set

$$H_{j+1} := -X_{j+1} + X_j^{n_j} - \theta_j \cdot X_0^{b_0^j} X_1^{b_1^j} \cdots X_{j-1}^{b_{j-1}^j} + p_j(X_0, \dots, X_j).$$

- For $j \in \{g, \dots, m-1\}$, we set

$$H_{j+1} := -X_{j+1} + \sum_{I=(i_0, \dots, i_g)} c_{j,I} X_0^{i_0} X_1^{i_1} \cdots X_g^{i_g},$$

where $x_{j+1} = \sum_{I=(i_0, \dots, i_g)} c_{j,I} x_0^{i_0} x_1^{i_1} \cdots x_g^{i_g}$ is the (x_0, \dots, x_g) -adic expansion of x_{j+1} (see Remark 2.22).

By definition, all the H_{j+1} are in the kernel I_S of the homomorphism (2.10). They generate I_S because any element of I_S is congruent modulo (H_2, \dots, H_m) to some element of $k[[X_0, X_1]]$ and the restriction of the homomorphism (2.10) to this ring is injective.

Lemma 3.27. *Let P be a rational point of $\Theta_{L_0}(D)$. With the previous notation, we have:*

- (1) For $j \in \{1, \dots, g-1\}$, $in_{w^P}(H_{j+1}) = X_j^{n_j} - \theta_j \cdot X_0^{b_j^0} X_1^{b_j^1} \dots X_{j-1}^{b_j^{j-1}}$.
- (2) For $j \in \{g, \dots, m-1\}$, $in_{w^P}(H_{j+1}) = -X_{j+1} + in_{w^P} \left(\sum_{I=(i_0, \dots, i_g)} c_{j,I} X_0^{i_0} X_1^{i_1} \dots X_g^{i_g} \right)$.

Proof. With the notation of Section 2.1.1, one has that $w^P = (1, \bar{\beta}_1/\bar{\beta}_0, \dots, \bar{\beta}_g/\bar{\beta}_0, w_{g+1}^P, \dots, w_m^P)$, where $w_i^P = \nu_{E_P}(x_j)/\bar{\beta}_0$, for $g+1 \leq i \leq m$. Assertion (1) follows from Proposition 2.21. Assertion (2) follows from Remark 2.22 applied to $h = x_{j+1}$, for $g \leq j \leq m-1$. \square

Remark 3.28. If P is a rational point of $\Theta_{L_0}(D)$, then the vector $\text{trop}(\nu_{E_P}) = \nu_{E_P}(x_0)w^P$ is primitive (see the proof of Lemma 3.27).

Proposition 3.29. *Let P be a rational point of $\Theta_{L_0}(D)$. With the previous notation,*

$$in_{w^P}(I_S) = (in_{w^P}(H_2), \dots, in_{w^P}(H_m)).$$

Proof. Let $<$ be the monomial order on $k[X_0, \dots, X_m]$ defined by $X^\alpha < X^\beta$ if $w^P \cdot \alpha < w^P \cdot \beta$ or $w^P \cdot \alpha = w^P \cdot \beta$ and $X^\alpha <_{lex} X^\beta$, where $<_{lex}$ is the lexicographic order with respect to $X_m < \dots < X_1 < X_0$. By Lemma 3.27, the $<$ -initial parts of H_2, \dots, H_m are pairwise coprime. Therefore, H_2, \dots, H_m is a $<$ -standard basis of I_S . By [KTV, Theorem 2.1], H_2, \dots, H_m is also a w^P -standard basis of I_S . \square

Proposition 3.30. *Let P be a rational point of $\Theta_{L_0}(D)$. The intersection of the variety $Z(in_{w^P}(I_S)) \subset k^{m+1}$ with the torus $(k^*)^{m+1}$ is an irreducible and smooth surface. In particular, the singular locus of $Z(in_{w^P}(I_S))$ does not meet the torus $(k^*)^{m+1}$.*

Proof. The variety $\mathcal{C} \subset k^g$ defined by the ideal $(in_{w^P}(H_2), \dots, in_{w^P}(H_g))$, seen as an ideal of $k[X_0, \dots, X_{g-1}]$, is a monomial curve with semigroup generated by $\bar{\beta}_0/n_g, \dots, \bar{\beta}_{g-1}/n_g$ (see [Tei06]). By the form of $in_{w^P}(H_{j+1})$, for $j = g, \dots, m-1$, we get that the restriction of the projection $(X_0, \dots, X_m) \mapsto (X_0, \dots, X_g)$ to the variety $Z(in_{w^P}(I_S))$ is an isomorphism whose image is the irreducible surface $\mathcal{C} \times k \subset k^g \times k$ defined by the ideal $(in_{w^P}(H_2), \dots, in_{w^P}(H_g))$ of $k[X_0, \dots, X_g]$. Note that for every $j \in \{g, \dots, m-1\}$, we cannot have that all the coefficients $c_{j,I}$ in the expansion of H_{j+1} vanish for all I , because this would imply that $x_{j+1} = 0$. Since the equations defined by $in_{w^P}(H_{j+1}) = 0$, are non-trivial, for $j \in \{g, \dots, m-1\}$, it follows that the intersection of the variety $Z(in_{w^P}(I_S)) \subset k^{m+1}$ with the torus $(k^*)^{m+1}$ is a surface which is irreducible and smooth. \square

Next we prove the following:

Proposition 3.31. *The ideal $I_S \subset k[[X_0, \dots, X_m]]$ is Newton non-degenerate.*

Proof. By Remark 3.25, it is enough to prove that the singular locus of $Z(in_w(I))$ and the torus $(k^*)^{m+1}$ have empty intersection for every $w \in \text{Trop}_{\geq 0}(I_S) \cap \mathbb{Q}_{\geq 0}^{m+1}$.

Take $w \in \text{Trop}_{\geq 0}(I_S) \cap \mathbb{Q}_{\geq 0}^{m+1}$. By Theorem 3.17, it must be of the form $w = w^P$ for some interior point P of $\Theta_{L_0}(D)$. Moreover, P is a rational point (see Lemma 3.18). According to Proposition 3.30, $\text{Sing}(Z(in_w(I))) \cap (k^*)^{m+1} = \emptyset$.

Now take $w \in \text{Trop}_{\geq 0}(I_S) \cap \mathbb{Q}_{\geq 0}^{m+1}$ such that $w_j = 0$ for some $j \in \{0, \dots, m\}$. Then $w = e_i$ by Lemma 3.6.(1). We can extract from L_0, \dots, L_m a sequence of maximal contact curves of L_i (see Lemma 2.16 and Definition 2.15). Suppose that such a sequence has g elements. Up to relabelling the indices $0, \dots, m$, we can assume that L_0, \dots, L_{g-1} is this sequence, $w =$

e_g , and then L_i becomes L_g . We define H_2, \dots, H_m as we did in the case of w^P where P was a rational point of $\Theta_{L_0}(D)$. The w -initial parts are obtained by substituting $X_g = 0$ on H_2, \dots, H_m . Let us denote by $\bar{\beta}_0, \dots, \bar{\beta}_{g-1}$ the minimal sequence of generators of Γ_{L_g} and set $w' = (\bar{\beta}_0, \dots, \bar{\beta}_{g-1}, 0, \dots, 0) \in \mathbb{Z}^{m+1}$. Let $<'$ be the monomial order on $k[X_0, \dots, X_m]$ defined by $X^\alpha <' X^\beta$ if $w' \cdot \alpha < w' \cdot \beta$ or $w' \cdot \alpha = w' \cdot \beta$ and $X^\alpha <_{lex} X^\beta$, where $<_{lex}$ is the lexicographic order with respect to $X_m < \dots < X_1 < X_0$. Arguing as in Proposition 3.29, we can show that H_2, \dots, H_m is a w -standard basis of I_S by considering the monomial order $<$ defined by $X^\alpha < X^\beta$ if $w \cdot \alpha < w \cdot \beta$ or $w \cdot \alpha = w \cdot \beta$ and $X^\alpha <' X^\beta$. Finally, in order to show that $\text{Sing}(Z(\text{in}_w(I))) \cap (k^*)^{m+1} = \emptyset$, we may apply the arguments given in Proposition 3.30, where in this case \mathcal{C} is a plane branch with semigroup Γ_{L_g} . \square

Theorem 3.32. *Let $C = \cup_{j=1}^r C_j$ be a reduced plane curve singularity with maximal contact curves L_0, \dots, L_m . Let Σ be a regular subdivision of $\text{GF}(I_S)$ inducing a regular subdivision \mathcal{T}_{reg} of the fan \mathcal{T} . Then the restriction of $\pi_{\Sigma_{0,m+1}}^\Sigma$ to the strict transform of $Z(I_S)$ induces an embedded resolution of $C \subset S$.*

Proof. By Remark 3.21, the fan \mathcal{T} contains the rays spanned by $\text{ord}(\eta_j)$ for all the branches C_j of C . Let us denote by $\pi : S(\pi) \rightarrow S$ the restriction of $\pi_{\Sigma_{0,m+1}}^\Sigma$ to the strict transform $S(\pi)$ of $Z(I_S)$. By Proposition 3.31 and Theorem 3.26, $S(\pi)$ is non-singular and transversal to the orbit stratification of the exceptional locus. In addition, by Theorem 2.26 the modification $\pi_{\Sigma_{0,m+1}}^\Sigma$ is a toric embedded resolution of $C \subset k^{m+1}$. Since $C \subset S$ it follows that the strict transform C^π of C is contained in $S(\pi)$ and the transversality properties above imply that C^π is transversal to the exceptional divisor of π . \square

3.3. Minimal embedded resolution with one toric morphism. In this section we describe the divisors of the minimal embedded resolution of C in terms of the Eggers-Wall tree $\Theta_{L_0}(D)$. Then, we prove that when the regular subdivision \mathcal{T}_{reg} in Theorem 3.32 is the minimal one, then the restriction of $\pi_{\Sigma_{0,m+1}}^\Sigma$ to the strict transform of $Z(I_S)$ induces the minimal embedded resolution of $C \subset S$.

We characterize first the set of exceptional divisors of the minimal embedded resolution of C in terms of the Eggers-Wall tree $\Theta_{L_0}(D)$, and we compare it with the set of rays in the minimal regularization of the fan \mathcal{T} .

Let P be a rational point of $\Theta_{L_0}(D)$ and set $Q := \min_{\leq L_0} \{P' \in [L_0, P] \mid \mathbf{i}_{L_0}^+(P') = \mathbf{i}_{L_0}(P)\}$. The point Q is L_0 or a point of discontinuity of the index function \mathbf{i}_{L_0} . By Lemma 3.9 there is a branch L' of D such that $Q \in [L_0, L']$ and $\mathbf{i}_{L_0}^+(Q) = \mathbf{i}_{L_0}(L')$. Denote by ψ the minimal resolution of ν_{E_Q} . If $Q = L_0$ we set $E_Q := L_0$, ψ is the identity map on S and (L_0, L') is a cross at O . If $Q \neq L_0$ then L' belongs to $\mathcal{C}_\psi(E_Q)$ by Lemma 3.19. Let (u, v) be a local coordinate system on $S(\psi)$ defining the cross $(E_Q, (L')^\psi)$, that is, $E_Q = Z(u)$ and $(L')^\psi = Z(v)$ on this model. Then, the divisorial valuation ν_{E_P} is *monomial* in terms of the cross $(E_Q, (L')^\psi)$. This is a particular case of the *quasi-monomial* valuations considered in [FJ04]. There exists a pair of coprime integers (n_P, m_P) , such that for any $0 \neq h \in R$, if one has the expansion $h \circ \psi = \sum a_{s,t} u^s v^t$ in $k[[u, v]]$, then $\nu_{E_P}(h) = \min\{sn_P + tm_P \mid a_{s,t} \neq 0\}$. The pair (n_P, m_P) is determined by the value of *renormalized exponent function* \mathbf{e}_{E_Q} associated to the triple $(S(\psi), E_Q, O')$, where $\{O'\} = E_Q \cap (L')^\psi$ (see [GGP19b, section 5.3 and 5.4]). One has that

$$(3.16) \quad \mathbf{e}_{E_Q}(P) = \mathbf{i}_{L_0}(L') (\mathbf{e}_{L_0}(P) - \mathbf{e}_{L_0}(Q)) \in \mathbb{Q}, \text{ and } \mathbf{e}_{E_Q}(P) = \frac{m_P}{n_P}.$$

We set $(n_{L_0}, m_{L_0}) := (1, 0)$ if $P_1 = L_0$ and $(n_{L_j}, m_{L_j}) := (0, 1)$ if $P_2 = L_j$ for $j \in \{1, \dots, m\}$.

We associate to a pair of points $P_1, P_2 \in \Theta_{L_0}(D)$ the rational cone:

$$\sigma_{P_1, P_2} := \mathbb{R}_{\geq 0}(n_{P_1}, m_{P_1}) + \mathbb{R}_{\geq 0}(n_{P_2}, m_{P_2}) \subset \mathbb{R}^2.$$

In [GGP19b] the toroidal embedded resolution process of a plane curve singularity C is studied with respect to a fixed smooth branch L_0 . One associates to such process a *completion* \hat{C} of C , by adding to the branches of C all the auxiliary curvetas considered in the process to define local coordinates. By using Lemma 3.9 and the presentation of toric embedded resolutions in [GGP19b] it is easy to see that one has a resolution ψ of C , such that $\hat{C} = C \cup D$ (see also [GGGR21]). Since L_0 is transversal to the branches of C , one has that ψ is the *minimal* embedded resolution of C (see [GP03, Section 3.4, Cor. 28]). In [GGP19b, Section 1.6.5] the toroidal embedded resolution ψ is compared with the Eggers-Wall tree $\Theta_{L_0}(D)$. Then, we have a bijection which sends an exceptional prime divisor E of ψ to the interior point $P \in \Theta_{L_0}(D)$ distinguished by C such that $E = E_P$. We can describe the set of points of $\Theta_{L_0}(D)$ whose representing divisors are the exceptional divisors of the minimal embedded resolution of C :

Notation 3.33. A point Q of $\Theta_{L_0}(D)$ is *distinguished by C* if $Q \in \mathcal{A}_{L_0}^C(D)$ or $Q \in (Q_1, Q_2)$ where $Q_1 \leq_{L_0} Q_2$ is a pair of consecutive points of $\mathcal{A}_{L_0}^C(D)$ and (n_Q, m_Q) defines a ray of the minimal regularization of the fan of the cone σ_{Q_1, Q_2} with respect to the lattice \mathbb{Z}^2 (see Notation 3.15).

The facts indicated above imply the following result:

Proposition 3.34. *The dual graph of the total transform of D by the minimal embedded resolution of C is isomorphic to the tree $\Theta_{L_0}(D)$ with vertices running through the set of points distinguished by C , by an isomorphism which preserves the labels by the branches of D .*

Remark 3.35. We have also a similar notion of *toroidal pseudo-resolutions* of C with respect to L_0 . By [GGP19b] one has a toroidal pseudo-resolution ϕ of C such that the dual graph of the total transform of D by ϕ is isomorphic to the tree $\Theta_{L_0}(D)$ with vertices running through the set of augmented marked points $\mathcal{A}_{L_0}^C(D)$ (see Notation 3.15).

Our aim now is to show that some linear relations between the vectors (n_Q, m_Q) associated with certain exceptional divisors E_Q are preserved when we consider the tropicalizations of the associated valuations. We start with a preparatory lemma:

Lemma 3.36. *Let $P_1, P_2 \in \mathcal{A}_{L_0}^C(D)$ be consecutive points in $\Theta_{L_0}(D)$. Assume that $P_1 \leq_{L_0} P_2$ and let $P \in [P_1, P_2]$ be a rational point. Denote by (n_P, m_P) the pair of coprime integers such that $e_{E_{P_1}}(P) = \frac{m_P}{n_P}$. For $0 \leq j \leq m$, there exists non negative integers a_j, b_j such that $\nu_{E_P}(L_j) = a_j n_P + b_j m_P$.*

Proof. On the one hand, for any rational point $P \in [P_1, P_2]$ one has $V_{L_0}^P = \frac{1}{\nu_{E_P}(L_0)} \nu_{E_P}$. On the other hand, by [GGP19a, Prop. 8.16 (3) and Def. 3.14] we deduce that $\nu_{E_P}(L_0) = n_P \mathbf{i}_{L_0}(P_2)$ since P_1 and P_2 belong to $\mathcal{A}_{L_0}^C(D)$, hence the index function \mathbf{i}_{L_0} is constant on $(P_1, P_2]$. In particular, $\nu_{E_{P_1}}(L_0) = \mathbf{i}_{L_0}(P_2)$. Take $j \in \{1, \dots, m\}$. One has that:

$$(3.17) \quad \nu_{E_P}(L_j) = n_P \mathbf{i}_{L_0}(P_2) V_{L_0}^P(L_j).$$

Next, the argument follows closely the discussion of the cases (1) and (2) in the proof of Lemma 3.13. If L_j is in case (1), using (3.17), (3.6), and (3.16) we get that $\nu_{E_P}(L_j) = n_P \mathbf{i}_{L_0}(P_2) \mathbf{i}_{L_0}(L_j) \mathbf{c}_{L_0}(P_1) + m_P \frac{\mathbf{i}_{L_0}(L_j)}{\mathbf{i}_{L_0}(P_2)}$, where $\frac{\mathbf{i}_{L_0}(L_j)}{\mathbf{i}_{L_0}(P_2)}$ is an integer since in this case $P_2 \leq_{L_0} L_j$, hence $\mathbf{i}_{L_0}(P_2) \mathbf{i}_{L_0}(L_j) \mathbf{c}_{L_0}(P_1)$ is also an integer. If L_j is in case (2), using (3.17) and (3.5) we

get that $\nu_{E_P}(L_j) = n_P \mathbf{i}_{L_0}(P_2) V_{L_0}^{P_1}(L_j) = n_P \nu_{E_{P_1}}(L_j)$, where the last equality in the previous formula follows using that $V_{L_0}^{P_1} = \frac{1}{\nu_{E_{P_1}}(L_0)} \nu_{E_{P_1}}$ and $\nu_{E_{P_1}}(L_0) = \mathbf{i}_{L_0}(P_2)$. \square

Proposition 3.37. *Let $P_1, P_2 \in \mathcal{A}_{L_0}^C(D)$ be consecutive points in $\Theta_{L_0}(D)$. Then, the map*

$$\phi_{P_1, P_2}: \sigma_{P_1, P_2} \cap \mathbb{Z}^2 \longrightarrow (\mathbb{R}_{\geq 0} w^{P_1} + \mathbb{R}_{\geq 0} w^{P_2}) \cap \mathbb{Z}^{m+1}, \quad (p, q) \mapsto (a_0 p + b_0 q, \dots, a_m p + b_m q),$$

is an isomorphism.

Proof. The map ϕ_{P_1, P_2} is well-defined and it is a group homomorphism thanks to Lemma 3.36. Note that $a_0 \neq 0$ and there exists an index $j \in \{1, \dots, m\}$ such that $P_2 \leq_{L_0} L_j$, and, then we have $b_j \neq 0$. This implies that ϕ_{P_1, P_2} is injective. By definition and Lemma 3.36 one has that w^{P_1} is a positive multiple of $\phi_{P_1, P_2}(n_{P_1}, m_{P_1})$ and the same happens when we compare w^{P_2} with $\phi_{P_1, P_2}(n_{P_2}, m_{P_2})$. In order to prove the surjectivity of this map it is enough to show that the image of a basis u_1, u_2 of the lattice \mathbb{Z}^2 , with $u_1, u_2 \in \sigma_{P_1, P_2}$, is part of a basis of \mathbb{Z}^{m+1} . This condition holds if and only if the two minors of the $2 \times (m+1)$ matrix G_{u_1, u_2} , whose rows are $\phi_{P_1, P_2}(u_i)$, for $i = 1, 2$, have greatest common divisor equal to one.

Take a branch $L \in \{L_1, \dots, L_m\}$ such that $P_2 <_{L_0} L$ and $\mathbf{i}_{L_0}(P_2) = \mathbf{i}_{L_0}(L)$. This condition implies that $\mathbf{i}_{L_0}(P) = \mathbf{i}_{L_0}(L)$ for $P \in (P_1, L]$, hence, arguing as in Corollary 3.20 we see that $L \in \mathcal{C}_{E_{P_1}}(\psi)$ for some suitable model ψ . By Lemma 2.16, there exists $1 \leq g \leq m$ such that, up to relabelling L_1, \dots, L_m , one has $L = L_g$ and L_0, \dots, L_{g-1} is a sequence of maximal contact curves of the branch L_g .

Take $P \in [P_1, P_2]$ such that $\mathbf{e}_{E_{P_1}}(P) = 1/d$ for some integer $d \geq 1$. Then, the vector $(d, 1)$ belongs to the cone σ_{P_1, P_2} and together with $(n_{P_1}, m_{P_1}) = (1, 0)$, they form a basis of \mathbb{Z}^2 .

By [GGP19a, Prop. 8.16 (1)] we have that $\nu_{E_P}(L_0) = d \mathbf{i}_{L_0}(P_2)$ and $\nu_{E_{P_1}}(L_0) = \mathbf{i}_{L_0}(P_2)$, where we use that the function \mathbf{i}_{L_0} is continuous on (P_1, P_2) . By Lemma 3.36 and the discussion of the cases (1) and (2) in the proof of Lemma 3.13, we get that:

$$\nu_{E_P}(L_j) = d \mathbf{i}_{L_0}(P_2) V_{L_0}^{P_1}(L_j) = d \nu_{E_{P_1}}(L_j), \quad \text{for } j \in 1, \dots, g-1, \text{ and}$$

$$\nu_{E_P}(L_g) = d \mathbf{i}_{L_0}(P_2) \mathbf{i}_{L_0}(L_j) \mathbf{c}_{L_0}(P_1) + 1 = d \nu_{E_{P_1}}(L_g) + 1.$$

Among the two minors of the matrix $G_{(1,0),(d,1)}$ we obtain the values, $\nu_{E_{P_1}}(L_0), \dots, \nu_{E_{P_1}}(L_{g-1})$, which have greatest common divisor equal to one, since they form a system of generators of the semigroup Γ_{L_g} by Proposition 2.11. \square

Let us denote by \mathcal{T}_{reg}^{min} the minimal regularization of the two dimensional fan \mathcal{T} . Its projectivization is the set of images of its non-zero cones in the projective space $\mathbb{P}^m(\mathbb{R})$. The projectivization of \mathcal{T}_{reg}^{min} is a graph since since the fan \mathcal{T}_{reg}^{min} consist of cones of dimension at most two contained in $\mathbb{R}_{\geq 0}^{m+1}$. The following result is a consequence of Propositions 3.34 and 3.37, taking into account Notation 3.33 and the definition of \mathcal{T} (see Proposition 3.16).

Proposition 3.38. *Let ψ be the minimal embedded resolution of C . The projectivization of the fan \mathcal{T}_{reg}^{min} is isomorphic to the dual graph of the total transform of D under ψ , by an isomorphism which respect the labels by the components of D .*

Keep the notation of Theorem 3.32. Let us study the intersection of $S(\pi)$ with the orbits defined by cones of \mathcal{T}_{reg} . This will allow us to describe when $\pi : S(\pi) \rightarrow S$ is the minimal embedded resolution of $C \subset S$.

If e_j is a vector of the canonical basis of \mathbb{Z}^{m+1} , then the orbit $O(\mathbb{R}_{\geq 0} e_j)_{\Sigma_{0, m+1}}$ is not contained in the discriminant locus of $\pi_{\Sigma_{0, m+1}}^{\Sigma}$ and the intersection of $O(\mathbb{R}_{\geq 0} e_j)_{\Sigma}$ with $S(\pi)$ is equal to the strict transform L_j^{π} of L_j . The orbits defined by other rays of \mathcal{T}_{reg} are contained in the critical

locus of $\pi_{\Sigma_{0,m+1}}^{\Sigma}$. Any such ray is of the form $\rho = \mathbb{R}_{\geq 0}w^P$ for a unique rational point P of $\Theta_{L_0}(D)$ (see Lemma 3.14).

Theorem 3.39. *With the hypothesis of Theorem 3.32, assume that $\mathcal{T}_{reg} = \mathcal{T}_{reg}^{min}$. Then:*

- (1) *If P is a rational point of $\Theta_{L_0}(D)$, and $\rho = \mathbb{R}_{\geq 0}w^P \in \mathcal{T}_{reg}^{min}$, then the closure of $S(\pi) \cap O(\rho)$ is equal to E_P .*
- (2) *The restriction of $\pi_{\Sigma_{0,m+1}}^{\Sigma}$ to the strict transform of $Z(I_S)$ induces the minimal embedded resolution of $C \subset S$.*

Proof. We prove first that $S(\pi) \cap O(\rho)$ is irreducible. In order to study this intersection, we can see the orbit $O(\rho)$ in an affine open toric subvariety X_{σ} of X_{Σ} , where $\sigma \in \Sigma$ is spanned by a basis v_0, \dots, v_m of $N = \mathbb{Z}^{m+1}$ and $v_0 := \text{trop}(\nu_{E_P}) = \nu_{E_P}(x_0)w^P$. Then, the affine toric variety X_{σ} is isomorphic to k^{m+1} with coordinates (U_0, \dots, U_m) , where $U_i = \chi^{v_i}$ for $i = 0, \dots, m$. In addition, the restriction π_{σ} of the toric modification $\pi_{\Sigma_{0,m+1}}^{\Sigma}$ to X_{σ} is given by

$$X_i \circ \pi_{\sigma} = U_0^{v_{0,i}} U_1^{v_{1,i}} \dots U_m^{v_{m,i}}, \text{ for } 0 \leq i \leq m,$$

where $v_j = \sum_{i=0}^m v_{j,i}e_i$, and the orbit $O(\rho)$ is defined by $U_0 = 0$ and $U_1 \cdots U_m \neq 0$. Given $g \in k[[X_0, \dots, X_m]]$, we write $g \circ \pi_{\sigma} = U_0^{\nu_{v_0}(g)} \cdot g_{\rho}$, where $g_{\rho} \in k[[U_0, \dots, U_m]]$ and U_0 does not divide g_{ρ} . If $h = \text{in}_{v_0}(g)$, then $h_{\rho} \in k[U_1, \dots, U_m]$ and $h_{\rho} = g_{\rho}(0, U_1, \dots, U_m)$. This fact and Proposition 3.29 imply that $S(\pi) \cap O(\rho)$ is defined by the ideal of $k[U_1^{\pm 1}, \dots, U_m^{\pm 1}]$ generated by the set $\{(\text{in}_{v_0}(H_2))_{\rho}, \dots, (\text{in}_{v_0}(H_m))_{\rho}\}$. Since the polynomials $(\text{in}_{v_0}(H_j))_{\rho}$ belong to $k[U_1, \dots, U_m]$ for $j = 2, \dots, m$, we get that the variety $S(\pi) \cap O(\rho)$ is irreducible if and only if the variety $V \subset (k^*)^{m+1}$ defined by these polynomials, seen in $k[U_0^{\pm 1}, U_1^{\pm 1}, \dots, U_m^{\pm 1}]$, is irreducible. The variety V is also defined by the ideal $(\text{in}_{v_0}(H_2) \circ \pi_{\sigma}, \dots, \text{in}_{v_0}(H_m) \circ \pi_{\sigma})$. The monomial map π_{σ} induces an isomorphism of tori, therefore the variety V is irreducible by Proposition 3.30. This shows that $S(\pi) \cap O(\rho)$ is irreducible.

We can take the regular cone $\sigma \in \Sigma$ in such a way that $v_1 \in |\mathcal{T}|$ and $v_2, \dots, v_m \notin |\mathcal{T}|$. Denote by ρ_i the ray $\mathbb{R}_{\geq 0}v_i$ for $i = 1, \dots, m$ and set $\rho_0 = \rho$. Take $i \in \{2, \dots, m\}$. The condition $v_i \notin |\mathcal{T}|$ implies that there exists a monomial in $k[X_0, \dots, X_m]$ which belongs to the v_i -initial ideal of I_S (see Proposition 3.23). Hence the intersection $S(\pi) \cap O(\rho_i)$ is empty.

Denote by E_i the closure of $S(\pi) \cap O(\rho_i)$ on X_{σ} for $i = 0, 1$. As the map $\pi : S(\pi) \rightarrow S$ is a model of the smooth surface (S, O) , its dual graph $G(\pi)$ is a tree. This implies that the divisors E_0 and E_1 intersect at at most one point, which belongs to the orbit $O(\rho_0 + \rho_1)$. We show now that this intersection consists of exactly one point o_1 .

Assume by contradiction that $E_0 \cap E_1 = \emptyset$. Then as the dual graph $G(\pi)$ is connected, there exists a sequence of rays ρ'_0, \dots, ρ'_s of \mathcal{T}_{reg}^{min} , with $s > 1$, $\rho_0 = \rho'_0$, $\rho_1 = \rho'_s$, such that $\rho'_{i-1} + \rho'_i \in \mathcal{T}_{reg}^{min}$, and $\rho'_{i-1} + \rho'_i \neq \rho_0 + \rho_1$, for $i = 1, \dots, s$. Then, the projectivization of the nonzero elements of the set $\rho_0 + \rho_1 \cup \bigcup_{i=1}^s \rho'_{i-1} + \rho'_i$ contains a non trivial cycle. We get a contradiction by Proposition 3.38, since the projectivization of fan \mathcal{T}_{reg}^{min} is a tree.

Denote by u_j , for $j = 0, \dots, m$, the restriction of the function U_j to the surface $S(\pi)$. We obtain that (u_0, u_1) are local coordinates at o_1 . It follows that the order of vanishing of $x_j \circ \pi_{\sigma}$ along the divisor E_0 is equal to $v_{0,j}$, for $0 \leq j \leq m$. Therefore, $\nu_{E_P}(x_j) = \nu_{E_0}(x_j)$, for $0 \leq j \leq m$.

By the transversality of $S(\pi)$ to the toric stratification of the exceptional divisor of π we have that any irreducible component of $E(\pi)$ must be the closure of $S(\pi) \cap O(\rho')$, for ρ' running through the rays of \mathcal{T}_{reg} which intersect $\mathbb{R}_{>0}^{m+1}$. By Proposition 3.38 we have a bijection between the set of irreducible components of the exceptional divisor of the minimal embedded resolution of C and the rays of \mathcal{T}_{reg} which intersect $\mathbb{R}_{>0}^{m+1}$, which is given by $E_{P'} \mapsto \mathbb{R}_{\geq 0}w^{P'}$, where P' runs

through the set of rational points of $\Theta_{L_0}(D)$ which are distinguished by C (see Notation 3.33). It follows that π is the minimal embedded resolution of C . This implies also that $E_0 = E_P$. \square

REFERENCES

- [AGS13] F. Aroca, M. Gómez-Morales, and K. Shabbir, *Toric modification of Newton non-degenerate ideals*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM **107** (2013), no. 1, 221–239.
- [BT07] R. Bahloul and N. Takayama, *Local Gröbner fans*, Comptes Rendus Mathématique **344** (2007), no. 3, 147–152.
- [CG03] A. Campillo and C. Galindo, *The Poincaré series associated with finitely many monomial valuations*, Math. Proc. Cambridge Philos. Soc. **134** (2003), no. 3, 433–443.
- [CM21] H. Cobo, H. Mourtada, *Jet schemes of quasi-ordinary surface singularities*, Nagoya Math. J. **242** (2021), 77–164.
- [CLS11] D. A. Cox, J. B. Little, and H. K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011.
- [CPS21] M. A. Cueto, P. Popescu-Pampu, D. Stepanov, *Local tropicalizations of splice type surface singularities*, preprint available at [arXiv\[math\]:2108.05912](https://arxiv.org/abs/2108.05912).
- [CMT21] D. Cutkosky, H. Mourtada, B. Teissier, *On the construction of valuations and generating sequences on hypersurface singularities*, Algebraic Geometry **8** (6) (2021) 705–748.
- [DGN08] F. Delgado, C. Galindo, and A. Núñez, *Generating sequences and Poincaré series for a finite set of plane divisorial valuations*, Adv. Math. **219** (2008), no. 5, 1632–1655.
- [Ewa96] G. Ewald, *Combinatorial convexity and algebraic geometry*, Graduate Texts in Mathematics, vol. 168, Springer-Verlag, New York, 1996.
- [FJ04] C. Favre and M. Jonsson, *The valuative tree*, Lecture Notes in Mathematics, vol. 1853, Springer-Verlag, Berlin, 2004.
- [Ful93] W. Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993, The William H. Roever Lectures in Geometry.
- [GGP19a] E. R. García Barroso, P. D. González Pérez, and P. Popescu-Pampu, *The valuative tree is the projective limit of Eggers-Wall trees*, Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas (2019).
- [GGP19b] ———, *The combinatorics of plane curve singularities. How Newton polygons blossom into lotuses*, Handbook of Geometry and Topology of Singularities I (José Luis Cisneros-Molina, Lê Dũng Tráng, and José Seade, eds.), Springer International Publishing, 2020, pp. 1–150.
- [GBP15] E. R. García Barroso and A. Płoski, *An approach to plane algebroid branches*, Rev. Mat. Complut. **28** (2015), no. 1, 227–252.
- [GP03] P. D. González Pérez, *Toric embedded resolutions of quasi-ordinary hypersurface singularities*, Ann. Inst. Fourier (Grenoble) **53** (2003), no. 6, 1819–1881.
- [GP10] ———, *Approximate roots, toric resolutions and deformations of a plane branch*, J. Math. Soc. Japan **62** (2010), no. 3, 975–1004.
- [GG14] P. D. González Pérez, M. González Villa, *Motivic Milnor fiber of a quasi-ordinary hypersurface*, J. Reine Angew. Math. no. 68, pag. 159–205, 2014.
- [GGGR21] P. D. González Pérez, M. González Villa, C. R. Guzmán Durán, M. Robredo Buces, *Multiplier ideals of plane curve singularities via Newton polygons*, preprint available at [arXiv:2109.13294 \[math.AG\]](https://arxiv.org/abs/2109.13294).
- [GSLJ91] G. Gonzalez-Sprinberg and M. Lejeune-Jalabert, *Modèles canoniques plongés. I*, Kodai Math. J. **14** (1991), no. 2, 194–209.
- [GT00] R. Goldin and B. Teissier, *Resolving singularities of plane analytic branches with one toric morphism*, Resolution of singularities (Obergrugl, 1997), Progr. Math., vol. 181, Birkhäuser, Basel, 2000, pp. 315–340.
- [Ish04] S. Ishii, *The arc space of a toric variety*, J. Algebra **278** (2004), no. 2, 666–683.
- [Jon15] M. Jonsson, *Dynamics of Berkovich spaces in low dimensions*, Berkovich spaces and applications, Lecture Notes in Math., vol. 2119, Springer, Cham, 2015, pp. 205–366.
- [KTV] G. Kemper, N.V. Trung, and N.T. Van Anh, *Toward a Theory of Monomial Preorders*, Math. of Computation, **87**, (2018), 313, 2513–2537.
- [Kou76] A. G. Kouchnirenko, *Polyèdres de Newton et nombres de Milnor*, Invent. Math. **32** (1976), no. 1, 1–31.
- [Kho78] A. G. Khovanskii, *Newton polyhedra and the genus of complete intersections*, Functional Anal. Appl. **12** (1978), no. 1, 38–46.

- [LJMR13] M. Lejeune-Jalabert, H. Mourtada, and A. J. Reguera, *Jet schemes and minimal embedded desingularization of plane branches*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM **107** (2013), no. 1, 145–157.
- [LJR99] M. Lejeune-Jalabert and A. J. Reguera, *Arcs and wedges on sandwiched surface singularities*, Amer. J. Math. **121** (1999), no. 6, 1191–1213.
- [MR88] T. Mora and L. Robbiano, *The Gröbner fan of an ideal*, Journal of Symbolic Computation **6** (1988), no. 2-3, 183–208.
- [Mou11] H. Mourtada, *Jet schemes of complex plane branches and equisingularity*, Annales de l’Institut Fourier, Tome **61**, numéro 6 (2011), p. 2313–2336.
- [Mou17] ———, *Jet schemes and generating sequences of divisorial valuations in dimension two*, Michigan Math. J. **66** (2017), no. 1, 155–174.
- [Mou20] ———, *Approches géométriques de la résolution des singularités et des partitions des nombres entiers*, Mémoire d’habilitation à diriger des recherches, (2020), Université de Paris.
- [MVV21] H. Mourtada, W. Veys, L. Vos, *The motivic Igusa zeta function of a space monomial curve with a plane semigroup*, Advances in Geometry, vol. 21, no. 3, 2021, pp. 417–442.
- [Oda88] T. Oda, *Convex bodies and algebraic geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 15, Springer-Verlag, Berlin, 1988, An introduction to the theory of toric varieties, Translated from the Japanese.
- [PP04] P. Popescu-Pampu, *On the analytical invariance of the semigroups of a quasi-ordinary hypersurface singularity*, Duke Math. J. **124** (2004), no. 1, 67–104.
- [PS13] P. Popescu-Pampu and D. Stepanov, *Local tropicalization*, Algebraic and combinatorial aspects of tropical geometry, 253–316, Contemp. Math., 589, Amer. Math. Soc., Providence, RI, 2013.
- [Ro19] M. Robredo Buces, *Invariants of singularities, generating sequences and toroidal structures* PhD Thesis at Univ. Complutense de Madrid, (2019), available at [E-prints Complutense](#).
- [Spi90] M. Spivakovsky, *Valuations in function fields of surfaces*, Amer. J. Math. **112** (1990), no. 1, 107–156.
- [St17] D.A. Stepanov, *Universal valued fields and lifting points in local tropical varieties*, Comm. Algebra **45** (2017), no. 2, 469–480.
- [Tei03] B. Teissier, *Valuations, deformations, and toric geometry*. Valuation theory and its applications, Vol. II (Saskatoon, SK, 1999), 361–459, Fields Inst. Commun., 33, Amer. Math. Soc., Providence, RI, 2003.
- [Tei04] ———, *Monomial ideals, binomial ideals, polynomial ideals*, Trends in commutative algebra, 211–246, Math. Sci. Res. Inst. Publ., 51, Cambridge Univ. Press, Cambridge, 2004.
- [Tei06] ———, Appendix to [\[Zar06\]](#).
- [Tei09] ———, *A viewpoint on local resolution of singularities*, Oberwolfach Workshop on Singularities, September 2009. Oberwolfach Reports, Vol. 6, No. 3, 2009. European Math. Soc. Publications.
- [Tei14] ———, *Overweight deformations of affine toric varieties and local uniformization*, Valuation theory in interaction", Proceedings of the second international conference on valuation theory, Segovia–El Escorial, 2011. Edited by A. Campillo, F-V. Kuhlmann and B. Teissier. European Math. Soc. Publishing House, Congress Reports Series, Sept. 2014, 474–565.
- [Tev07] J. Tevelev, *Compactifications of subvarieties of tori*, Amer. J. Math. **129** (2007), no. 4, 1087–1104.
- [Tev14] ———, *On a question of B. Teissier*, Collectanea Math. **65** (1) (2014), 61–66.
- [Tou05] N. Touda, *Local Tropical Variety*, preprint available at [arXiv:math/0511486 \[math.AG\]](#).
- [Zar06] O. Zariski, *The moduli problem for plane branches*, University Lecture Series, vol. 39, American Mathematical Society, Providence, RI, 2006.

DEPARTAMENTO DE MATEMÁTICAS, ESTADÍSTICA E I.O. SECCIÓN DE MATEMÁTICAS, UNIVERSIDAD DE LA LAGUNA. APARTADO DE CORREOS 456. 38200 LA LAGUNA, TENERIFE, ESPAÑA.

Email address: afelipe@ull.edu.es

INSTITUTO DE MATEMÁTICA INTERDISCIPLINAR, DEPARTAMENTO DE ÁLGEBRA, GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE CIENCIAS MATEMÁTICAS, UNIVERSIDAD COMPLUTENSE DE MADRID, PLAZA DE LAS CIENCIAS 3, MADRID 28040, ESPAÑA.

Email address: pgonzalez@mat.ucm.es

UNIVERSITÉ DE PARIS, SORBONNE UNIVERSITÉ, CNRS, INSTITUT MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, F-75013, PARIS, FRANCE.

Email address: houssein.mourtada@imj-prg.fr