

A ZOO OF DUALITIES

S. ARTSTEIN-AVIDAN, S. SADOVSKY, K. WYCZESANY

ABSTRACT. In this note we study order reversing quasi involutions and their properties. These maps are dualities (order reversing involutions) on their image. We prove that any order reversing quasi involution is induced by a cost. We characterize when an order reversing quasi involution on a subclass can be extended to the whole space and discuss the uniqueness of the extension. We describe ways for constructing order reversing quasi involutions and study their invariant sets. Finally, we present a “zoo of examples” where many transforms are represented as order reversing quasi involutions. Some of the examples are classical, some have recently attracted attention of the convexity community and some are new. We present a unifying point of view deepening the understanding of the underlying principles and structures and offering a new perspective on the topic.

1. DEFINITIONS AND RESULTS

We study abstract dualities on classes of partially ordered sets. A duality is defined as a mapping from a class \mathcal{C} (which has a partial order on it) to itself which is order reversing and which satisfies $T \circ T = I$ (where I is the identity operator). In this setting, several characterization theorems have been proven (see, among others, [1–4, 7–10, 14, 16, 17]).

Of particular significance in geometry are \mathcal{K}_0^n , the class of closed, convex sets in \mathbb{R}^n with the origin in the interior together with the order given by inclusion, and the following two classes of functions: The class $\text{Cvx}(\mathbb{R}^n)$ of lower semi-continuous convex functions from \mathbb{R}^n to $(-\infty, \infty]$, with the order given by point-wise inequality (which corresponds to the inclusion of the epi-graphs,) and the subclass $\text{Cvx}_0(\mathbb{R}^n) \subset \text{Cvx}(\mathbb{R}^n)$ of non-negative lower semi-continuous convex functions attaining value zero at the origin. In each of these examples the order reversing transform can be extended to be defined on a much larger class, and on it, the transform satisfies a slightly modified definition which we call order reversing quasi involution which is given in the following definition. We denote $\mathcal{P}(X)$ to be the power set of X , that is, all possible subsets of X (including the empty set and all of X), sometimes denoted by 2^X .

Definition 1.1. *Let X be some set, and let $T : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$. We will say that T is an order reversing quasi involution if for every $K, L \subset X$, the following hold*

- (i) $K \subset TTK$, (quasi involution)
- (ii) if $L \subset K$ then $TK \subset TL$. (order reversion)

When $T : \mathcal{C} \rightarrow \mathcal{C}$ is only defined on subsets of X in some class $\mathcal{C} \subset \mathcal{P}(X)$, but still satisfies (i) and (ii), we say that T is an order reversing quasi involution on \mathcal{C} .

In this paper, we study order reversing quasi involutions and their structural properties. It turns out that order reversing quasi involutions have many interesting properties in common. In fact, they can all be identified as a form of “cost dualities for sets”, introduced in [6] and used there in the context of optimal transport. These cost dualities are order reversing quasi involutions induced by a function, which we call a cost.

Definition 1.2 (Cost duality for sets). *Let $c : X \times X \rightarrow (-\infty, \infty]$ satisfy $c(x, y) = c(y, x)$. For $K \subset X$ define the c -dual set of K as*

$$K^c = \{y \in X : \inf_{x \in K} c(x, y) \geq 0\}.$$

Our first main theorem reads as follows.

Theorem 1.3. *Let $T : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ be an order reversing quasi involution. Then there exists a cost function $c : X \times X \rightarrow \{\pm 1\}$ such that for all $K \subset X$, $TK = K^c$.*

In the paper we discuss some inherent properties of order reversing quasi involutions. For example, any order reversing quasi involution is always an order reversing involution on its image and this image has a natural lattice structure. Other useful properties follow from the cost representation. When an order reversing quasi involution is defined on a subclass $\mathcal{C} \subset \mathcal{P}(X)$, we determine necessary and sufficient conditions for it to have an extension to $\mathcal{P}(X)$. We say that T respects inclusions if $K \subset \cup_{i \in I} K_i$ implies $TK \supset \cap_{i \in I} TK_i$ for any $K, K_i \in \mathcal{C}$, $i \in I$. It is easy to see that this condition is necessary and the next theorem shows that it is in fact sufficient.

Theorem 1.4. *Let $\mathcal{C} \subset \mathcal{P}(X)$ a family of sets and $T : \mathcal{C} \rightarrow \mathcal{C}$ be an order reversing quasi involution which respects inclusions. Then T can be extended to an order reversing quasi involution $\hat{T} : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$, such that $\hat{T}|_{\mathcal{C}} = T$.*

When the class \mathcal{C} is closed under intersection, any order reversing quasi involution T on it automatically respects inclusions (see Lemma 2.5).

Order reversing quasi involutions can be intersected and composed in various ways producing other order reversing quasi involutions which may inherit some of their properties. Of particular interest are invariant sets of a given order reversing quasi involution and we prove, for instance, the following theorem.

Theorem 1.5. *Let $c : X \times X \rightarrow \{-1, 1\}$ be a symmetric cost function and denote $X_0 = \{x : c(x, x) \geq 0\}$. Let $S_0 \subset X_0$ satisfy $c(x, y) \geq 0$ for all $x, y \in S_0$. Then there exists some $K \subset X$ with $S_0 \subset K$ and such that $K^c \cap X_0 = K$.*

The rest of the paper is devoted to presenting a “zoo of examples” that includes, in particular, dualities on some important classes of convex bodies as well as star-shaped sets, reciprocal bodies and flowers [11, 12]. We further present functional examples, where a function is identified with its epigraph or hypograph. These include the Legendre and Polarity transforms (see e.g. [4]), which were a main motivation for this project. This vast array of examples shows how fundamental the notion of an order reversing quasi involution is, and

offers a unifying point of view which deepens our understanding of the underlying principles and structures.

The paper is organized as follows: In Section 2 we investigate some basic properties of order reversing quasi involutions. In Section 3 we study cost dualities and prove Theorem 1.3. In Section 4 we discuss uniqueness and existence of order reversing quasi involutions defined on subsets of $\mathcal{P}(X)$, and prove Theorem 1.4. In Section 5 we discuss ways of producing new order reversing quasi involutions based on given ones, in particular by using restriction and conjugation. To this end we utilize order preserving transformations and a modification of property (i) in Definition 1.1, where the order of inclusion is reversed. We further address the question of fixed points of the transform $K \mapsto K^c$. Section 6 is the “petting zoo” with several examples in it that we take a closer look at.

Acknowledgment. The authors were supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 770127). The second named author is grateful to the Azrieli foundation for the award of an Azrieli fellowship.

2. GENERAL PROPERTIES OF QUASI INVOLUTIONS ON SETS

We begin by gathering a few simple facts about order reversing quasi involutions. First, every order reversing quasi involution gives rise to an order reversing involution on a subset of $\mathcal{P}(X)$.

Lemma 2.1. *If $T : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ is an order reversing quasi involution then for $\mathcal{C} = \text{Im}(T) \subset \mathcal{P}(X)$, that is, $\mathcal{C} = \{TK : K \subset X\}$, we have that $T : \mathcal{C} \rightarrow \mathcal{C}$ is an abstract duality, that is, an order reversing involution.*

Proof. The fact that T reverses order on \mathcal{C} is inherited from the same property on $\mathcal{P}(X)$. As for the involution property, it amounts to showing $TTTK = TK$ for any K . Using the first property on K , we see that $TTK \supset K$, so applying T and using order reversion we get $TTTK \subset TK$. On the other hand, we can use the order reversion property on TK instead, which yields, $TK \subset TTTK$, and the claim is proven. \square

Secondly we mention that the direction of the inclusion in the quasi involution property (i) can be reversed, and there is a one-to-one correspondence between order reversing quasi involutions as defined in Definition 1.1, and those that one could define with the reverse inclusion, as is apparent from the following lemma.

Lemma 2.2. *Let $T : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ be an order reversing quasi-involution. Then the mapping $S : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ given by $SK = X \setminus (T(X \setminus K))$ satisfies*

- (i) $K \supset SSK$, (complemented quasi involution)
- (ii) if $L \subset K$ then $SK \subset SL$. (order reversion)

Proof. Indeed, for the order reversion property, if $L \subset K$ then $X \setminus L \supset X \setminus K$ and so $T(X \setminus L) \subset T(X \setminus K)$, from which it follows that

$$SL = X \setminus T(X \setminus L) \supset X \setminus T(X \setminus K) = SK.$$

For the quasi involution property, we want to show $K \supset SSK$ which can be rewritten as $X \setminus K \subset X \setminus SSK$, which in turn means $X \setminus K \subset T(X \setminus SK) = TT(X \setminus K)$, which follows from the order reversion property given for T on the subset $X \setminus K$. \square

Remark 2.3. As the reader will readily notice, the transform S amounts to the composition of the given transform T on both sides with the order reversing *involution* given by $K \mapsto X \setminus K$. For other properties of the conjugation of an order reversing quasi involution with specific classes of transformations (such as another order reversing quasi involution, for example) see Section 5.2.

The next property we list is trivial, and amounts to the fact that under an order reversing transformations, maximal elements are mapped to minimal ones (in the image) and vice versa. Joined with the quasi involution property (i), this gives the following lemma.

Lemma 2.4. *If $T : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ is an order reversing quasi involution then $TTX = X$ and $T\emptyset = X$.*

Proof. Indeed, $TTX \supset X$ by the quasi involution condition, so $TTX = X$. Letting $L_0 = TX$, since $\emptyset \subset L_0$ we have by order reversion that $T\emptyset \supset TL_0 = X$ and so $T\emptyset = X$. \square

A fourth useful fact is that given an order reversing quasi involution on a subset \mathcal{C} of $\mathcal{P}(X)$, which is closed under intersection, and with $X \in \mathcal{C}$, we may extend it to the full space $\mathcal{P}(X)$.

Lemma 2.5. *Let $\mathcal{C} \subset \mathcal{P}(X)$ be a class of sets which is closed under intersection, with $X \in \mathcal{C}$. Given an order reversing quasi involution $T : \mathcal{C} \rightarrow \mathcal{C}$, there exists an order reversing quasi involution $\hat{T} : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ with $\hat{T}|_{\mathcal{C}} = T$.*

Remark 2.6. In Proposition 4.1 we show that if we require the image of \hat{T} to be \mathcal{C} , then the extension \hat{T} is unique. Furthermore, Example 4.3 demonstrates that without the property that \mathcal{C} is closed under intersection, one may easily produce an order reversing involution on \mathcal{C} with no extension to the whole $\mathcal{P}(X)$. However, in Theorem 1.4 we show that adding a different, but still necessary condition on T , one can extend also order reversing quasi involutions defined on classes \mathcal{C} which are not closed under intersection.

Proof of Lemma 2.5. Let $\mathcal{C}_0 \subset \mathcal{C}$ denote $\{TK : K \in \mathcal{C}\}$, that is, $T\mathcal{C} = \mathcal{C}_0$. Let $L_0 \in \mathcal{C}$ denote the intersection of all elements in \mathcal{C}_0 (it may be an empty set). Since $X \in \mathcal{C}$, and since for all $K \in \mathcal{C}$, $K \subset X$, we know that $TX \subset TK$ for all $K \in \mathcal{C}$ so that $TX \subset \bigcap_{L \in \mathcal{C}_0} L = L_0$, and as it participates in the intersection, we see $TX = L_0$.

For $K \subset X$ we define

$$\hat{T}(K) = T(\bigcap \{L \in \mathcal{C} : K \subset L\}).$$

First note that as \mathcal{C} is closed under intersection, the set on which we apply T (which we call the “ \mathcal{C} -hull” of K) is in \mathcal{C} as well. We take into account here the fact that $X \in \mathcal{C}$, so that

the intersection is over a non-empty set of elements L . Therefore, T is well-defined. Next, note that for $K \in \mathcal{C}$, this “ \mathcal{C} -hull” is K itself, and therefore $\hat{T}|_{\mathcal{C}} = T$. Let us show that \hat{T} is an order reversing quasi involution. For the order reversion, let $K_1 \subset K_2$, in which case the “ \mathcal{C} -hulls” satisfy the same inclusion $\cap\{L \in \mathcal{C} : K_1 \subset L\} \subset \cap\{L \in \mathcal{C} : K_2 \subset L\}$, and by order reversion of T on \mathcal{C} we get

$$\hat{T}(K_1) = T(\cap\{L \in \mathcal{C} : K_1 \subset L\}) \supset T(\cap\{L \in \mathcal{C} : K_2 \subset L\}) = \hat{T}(K_2).$$

For the quasi involution note that a set is always included in its “ \mathcal{C} -hull”, $K \subset \cap\{L \in \mathcal{C} : K \subset L\}$, and since T is a quasi involution we get

$$K \subset \cap\{L \in \mathcal{C} : K \subset L\} \subset TT(\cap\{L \in \mathcal{C} : K \subset L\}) = \hat{T}T(\cap\{L \in \mathcal{C} : K \subset L\}) = \hat{T}\hat{T}(K).$$

where we used (for the second equality from the right) that $T = \hat{T}$ on \mathcal{C} , and the image of T is in \mathcal{C} . \square

As a corollary we get that the image of an order reversing quasi involution on $\mathcal{P}(X)$ is always closed under intersection (or we can show this directly, see Proposition 2.7).

Examples for sets \mathcal{C} which can be taken in Lemma 2.5 are closed sets, convex sets, sets which include the origin, bounded sets (together with the whole space), sets which are sub-graphs of functions with values in $(-\infty, \infty]$, any intersection of the above, and many more.

If we change the quasi involution property (i) in Definition 1.1 to be in the opposite direction, $TTK \subset K$, then (by Lemma 2.2) the condition of being closed under intersection becomes a condition on being closed under unions (and the condition of containing X becomes the condition of containing \emptyset).

Finally, any order reversing quasi involution defined on all the subsets of the space X respects the “lattice structure” in that it maps unions to intersections. The following simple proposition will be of much use to us in what follows.

Proposition 2.7. *Let $T : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ be an order reversing quasi involution. Then for any collection of sets $K_i \subset X$, $i \in I$,*

$$(1) \quad T(\cup_{i \in I} K_i) = \cap_{i \in I} T(K_i).$$

Proof. We have $K_j \subset \cup_{i \in I} K_i$ for all $j \in I$, so by the order reversion property (i), $T(\cup_{i \in I} K_i) \subset T(K_j)$ for any j , and in particular $T(\cup_{i \in I} K_i) \subset \cap_{j \in I} T(K_j)$.

For the other direction, $\cap_{i \in I} T(K_i) \subset T(K_j)$ for any $j \in I$. Applying property (ii) again, $TT(K_j) \subset T(\cap_{i \in I} TK_i)$. Using property (i), this implies $K_j \subset T(\cap_{i \in I} TK_i)$, so in particular $\cup_{i \in I} K_i \subset T(\cap_{i \in I} TK_i)$. Applying T duality once more

$$T(\cup_{i \in I} K_i) \supset TT(\cap_{i \in I} TK_i) \supset \cap_{i \in I} TK_i$$

where for the rightmost include we used property (i) once more. \square

Remark 2.8. In fact, as is apparent from the proof, when T is defined on a subclass $\mathcal{C} \subset \mathcal{P}(X)$, and is an order reversing quasi involution on \mathcal{C} , property (1) still holds whenever $\cup_{i \in I} K_i \in \mathcal{C}$.

Remark 2.9. A *complete lattice* is a partially ordered set (S, \leq) in which all subsets have both a supremum (join) and an infimum (meet). Proposition 2.7 implies that for an order reversing quasi involution $T : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$, the image $\mathcal{C}_0 = T(\mathcal{P}(X))$ is a complete lattice. For $M \subset \mathcal{C}_0$ the set $\cap\{K : K \in M\} \in \mathcal{C}_0$ is the meet of M , and the set $TT(\cup\{K : K \in M\})$ is the join.

3. COST DUALITIES

Our motivation for studying abstract dualities for subsets arose from our study of optimal transport maps and potentials in [6]. The setting is of a space X given together with a symmetric cost function $c : X \times X \rightarrow (-\infty, \infty]$. When X is a measure space, and a probability measure π on $X \times X$ is given, the “total cost” which π induces is given by $\int_{X \times X} c(x, y) d\pi(x, y)$. When this expression is finite, we say that π is a finite cost plan between its two marginals, μ and ν , defined by $\mu(A) = \pi(A \times X)$ and $\nu(B) = \pi(X \times B)$, where A and B are any measurable subsets of X .

For this total cost to be finite, a necessary condition is for π to be concentrated on the set $S \subset X \times X$ where $c(x, y) < \infty$. In studying restrictions for the existence of finite cost plans between given measures, a natural construction of the “ c -dual” of a subset of X arose, which, roughly speaking, is as follows: Given a subset $A \subset X$, we look at all of the elements $y \in X$ which have finite cost with *at least* one element of A . The complement of this set is then given by

$$TA = \{y \in X : \forall x \in A, \quad c(x, y) = \infty\}.$$

A necessary condition on probability measures μ and ν on X , for the existence of a finite cost plan between them, is that $\mu(A) + \nu(TA) \leq 1$ for all $A \subset X$. It is not hard to check (and we do this below) that $T : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ is an order reversing quasi involution. What is slightly more surprising, and we prove in this section, is that this is in fact the general form of an order reversing quasi involution for sets. In fact, it suffices to take very simple cost functions, which attain only two values (some finite value and the value $+\infty$).

There is no significance in the use of the value $+\infty$, and for the purpose of this note we use Definition 1.1 which we recall here including an extra parameter.

Definition 1.2. *Let $c : X \times X \rightarrow (-\infty, \infty]$ satisfy $c(x, y) = c(y, x)$. Fix $t \in (-\infty, \infty]$ (which will be omitted in the notation as it is a fixed parameter). For $K \subset X$ define the c -dual set of K as*

$$K^c = \bigcap_{x \in K} \{y \in X : c(x, y) \geq t\} = \{y \in X : \inf_{x \in K} c(x, y) \geq t\}.$$

We first remark that the notion of c -duality may be generalized further. One may associate with a cost function $c : X \times Y \rightarrow (-\infty, \infty]$ the corresponding transform from $\mathcal{P}(X)$ to $\mathcal{P}(Y)$, and its counterpart, from $\mathcal{P}(Y)$ to $\mathcal{P}(X)$. For simplicity we restrict here to the case $X = Y$ and symmetric c .

Second, very little of the information given by c is used for the definition of the transform. The choice of t is immaterial as we may exchange c with $c - t$ and take $t = 0$, or, if $t = \infty$,

we can take $-\exp(-c)$, $t = 0$. Moreover, two costs c_1, c_2 , with parameters t_1, t_2 , produce the same transform if and only if

$$\{(x, y) : c_1(x, y) \geq t_1\} = \{(x, y) : c_2(x, y) \geq t_2\}.$$

In other words, the transform is actually defined by this characteristic subset of $X \times X$, and the main reason to keep in mind the cost function and the parameter t is that for a fixed cost function and a varying parameter t we get a family of transforms which is sometimes of interest to study as a whole. In what follows, when t is not specified we always mean $t = 0$.

Lemma 3.1. *The transform $K \mapsto K^c$ is an order reversing quasi involution.*

Proof. Order reversion is immediate from the definition, as if $K \subset L$ and $c(x, y) \geq t$ for all $x \in L$ then $c(x, y) \geq t$ for all $x \in K$. For the property $K \subset TTK$ note that if $x \in K$ and $y \in K^c$ then $c(x, y) \geq t$ so that $x \in K^{cc}$. \square

Let us point out (see also Section 6.2) that in the case of the classical cost $c(x, y) = -\langle x, y \rangle$ and $t = -1$, we get the well known polar set K° , as is obviously the case also for the polar cost $c(x, y) = -\ln(\langle x, y \rangle - 1)_+$ and $t = \infty$. The classical cost $c(x, y) = -\langle x, y \rangle$ and $t = -5$ gives $K \mapsto 5K^\circ$. With $t = 1$, however, we get

$$K \mapsto \{x : \forall x \in K, \langle x, y \rangle \geq 1\}$$

which is very different from duality, and we will see many other examples of cost dualities in Section 6.

Our main observation in this section is, as stated above, that all order reversing quasi involutions on sets are induced by costs, and in fact by costs into $\{\pm 1\}$.

Theorem 1.3. *Let $T : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ be an order reversing quasi involution. Then there exists a cost function $c : X \times X \rightarrow \{\pm 1\}$ such that for all $K \subset X$, $TK = K^c$.*

Proof. We shall construct a cost c as required. To do so, define the following subset of $X \times X$

$$S = \{(x, y) : y \in T(\{x\})\},$$

and define $c(x, y) = 1$ if $(x, y) \in S$ and -1 elsewhere. Note that S is a symmetric set (that is, $(x, y) \in S$ implies $(y, x) \in S$): Indeed, $y \in T(\{x\})$ implies $T(\{y\}) \supset TT(\{x\}) \supset \{x\}$ so that $x \in T(\{y\})$.

Let $K \subset X$, we claim that $TK = K^c$. Note that

$$K^c = \{y : \forall x \in K \ c(x, y) \geq 0\} = \{y : \forall x \in K \ (x, y) \in S\} = \{y : \forall x \in K \ y \in T(\{x\})\}$$

Thus

$$K^c \subset \bigcap_{x \in K} T(\{x\}) = T(\bigcup_{x \in K} \{x\}) = TK.$$

Next let $y \in TK$, then $T(\{y\}) \supset TTK \supset K$ so that for every $x \in K$ we have $x \in T(\{y\})$ which, by the symmetry we mentioned, means that for every $x \in K$, $y \in T(\{x\})$ so that for every $x \in K$, $c(x, y) \geq 0$ namely $y \in K^c$ as required. \square

As the proof above demonstrates, to every symmetric subset $S \subset X \times X$ there corresponds a unique order reversing quasi involution. Indeed, given S we define a cost by $c(x, y) = 1$ if $(x, y) \in S$ and -1 elsewhere. Then

$$T\{x\} = \{x\}^c = \{y : c(x, y) \geq 0\} = \{y : (x, y) \in S\} =: S_x$$

is the fiber of X . For any $K \subset X$

$$TK = T(\cup_{x \in K} \{x\}) = \cap_{x \in K} S_x.$$

Clearly different sets S produce different order reversing quasi involutions T , since at least one of the fibers is different.

Remark 3.2. Let us note that the set S defined in the proof of Theorem 1.3 can be equivalently written as

$$S = \cup\{TK \times TTK : K \subset X\} \subset X \times X,$$

a representation which will resurface in the proof of Theorem 1.4 in Section 4.

4. EXTENSIONS OF ORDER REVERSING QUASI INVOLUTIONS

In this section we study when and how can an order reversing quasi involution T on a subset $\mathcal{C} \subset \mathcal{P}(X)$ be extended to the whole of $\mathcal{P}(X)$. We begin by showing uniqueness of the extension, namely that an order reversing quasi involution on $\mathcal{C} \subset \mathcal{P}(X)$ cannot have more than one extension to $\mathcal{P}(X)$ whose image is $T(\mathcal{C})$.

Proposition 4.1. *Let $T_1, T_2 : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ be order reversing quasi involutions, and assume $T_1(\mathcal{P}(X)) = T_2(\mathcal{P}(X)) =: \mathcal{C}_0 \subset \mathcal{P}(X)$, and further assume $T_1|_{\mathcal{C}_0} = T_2|_{\mathcal{C}_0}$. Then $T_1 = T_2$.*

Proof. We let T stand for either T_1 or T_2 and show that $TL = T(\cap\{K \in \mathcal{C}_0 : L \subset K\})$, which in particular implies that T_1 and T_2 are equal. Note that \mathcal{C}_0 is closed under intersections, and includes X as $TTX = X$. Moreover, as T is an involution on \mathcal{C}_0 , we know that letting $K_0 = \cap_{K \in \mathcal{C}_0} K$ we have $TK_0 = X$ and $TX = K_0$. Clearly all subsets of K_0 are mapped to X as well.

Let $L \in \mathcal{P}(X) \setminus \mathcal{C}_0$, and let $L_0 = \cap\{K \in \mathcal{C}_0 : L \subset K\}$. The intersection is non-empty as $L \subset X$, and by the fact that \mathcal{C}_0 is closed under intersection, $L_0 \in \mathcal{C}_0$. By order reversion, as $L \subset L_0$, we know $TL \supset TL_0$ (and $TTL \subset TTL_0 = L_0$). On the other hand, as $L \subset TTL$ and $TTL \in \mathcal{C}_0$, we know that TTL participates in the intersection defining L_0 , namely $L_0 \subset TTL$. By order reversion, and the involutive property of T on \mathcal{C}_0 , we get $TL_0 \supset TTTL = TL$. That is, $TL = TL_0$. \square

Remark 4.2. Within the proof we have shown the useful fact that for an order reversing quasi involution with image \mathcal{C}_0 ,

$$TL = T(\cap\{K \in \mathcal{C}_0 : L \subset K\}).$$

Proposition 4.1 complements Lemma 2.5, which tells us that given a set \mathcal{C} closed under intersections which includes X , and an order reversing quasi involution on it, we can extend it to the whole set $\mathcal{P}(X)$ with the same image.

We mention that as the example of Legendre transform \mathcal{L} and polarity transform \mathcal{A} on, say, 2-homogeneous convex functions shows, an order reversing quasi involution may have more than one extension to the whole space if we do not insist on keeping the same image. For details see Remark 6.14.

It turns out that if one is given an order reversing quasi involution on a subset $\mathcal{C} \subset \mathcal{P}(X)$ which is not closed under intersection, it may be the case that there is no extension to an order reversing quasi involution on $\mathcal{P}(X)$, as is the case in the following example.

Example 4.3. Consider the following sets $A = \{1, 2\}, B = \{1, 3\}, C = \{2, 3\}$, as subsets of $X = \{1, 2, 3\}$. Note that no two of these sets are comparable with respect to inclusion and hence any involution on these three sets will be order-reversing (in an empty sense). Set $TA = B, TB = A$ and $TC = C$. We then have that $A \subset B \cup C$ but $TA = B \not\supset A \cap C = TB \cap TC$. However, any extension \hat{T} of T would have to satisfy (by Proposition 2.7) that

$$\{1, 3\} = T(A) = \hat{T}(A) \supset \hat{T}(B \cup C) = \hat{T}(B) \cap \hat{T}(C) = TB \cap TC = \{2\}$$

which is not true, therefore, an extension \hat{T} does not exist.

The reason that in this this example one fails to have an extension is quite simple. For an order reversing quasi involution on $\mathcal{C} \subset \mathcal{P}(X)$ to admit an extension to $\mathcal{P}(X)$, Proposition 2.7 implies that it must satisfy the following property:

Definition 4.4. Let $\mathcal{C} \subset \mathcal{P}(X)$ and let $T : \mathcal{C} \rightarrow \mathcal{C}$ be an order reversing quasi involution. We say that T **respects inclusions** if for any $K, K_i \in \mathcal{C}, i \in I$ such that $K \subset \cup_{i \in I} K_i$ one has that $TK \supset \cap_{i \in I} TK_i$.

Remark 4.5. By Lemma 2.5 we know that if an order reversing quasi involution T is defined on a family \mathcal{C} which is closed under intersections then it can be extended to the whole space, and in particular, Proposition 2.7 implies that T respects inclusions.

It turns out that the condition of respecting inclusion is not only necessary, but sufficient for the existence of an extension, as we show next.

Theorem 1.4. Let $\mathcal{C} \subset \mathcal{P}(X)$ a family of sets and $T : \mathcal{C} \rightarrow \mathcal{C}$ be an order reversing quasi involution which respects inclusions. Then T can be extended to an order reversing quasi involution $\hat{T} : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$, such that $\hat{T}|_{\mathcal{C}} = T$.

Proof. Given the transform T define $S = \cup\{TK \times TTK : K \subset X\} \subset X \times X$ as in Remark 3.2. We may then define a cost function c such that $c(x, y) = 1$ if $(x, y) \in S$ and -1 otherwise (this is the same as in the proof of Theorem 1.3). Clearly, the corresponding cost transform is defined on the whole of $\mathcal{P}(X)$ and we must only check that for any $K \in \mathcal{C}$ we have that $TK = K^c$.

To show that $TK \subset K^c$ note that by definition of c , for any $(x, y) \in TTK \times TK$ we have $c(x, y) = 1$. In particular $c(x, y) = 1$ for all $(x, y) \in K \times TK$, as $K \subset TTK$ by property (i). Hence,

$$K^c = \{y : \forall x \in K c(x, y) \geq 0\} \supset \{y : (x, y) \in K \times TK\} = TK$$

which gives $TK \subset K^c$.

For the other direction, note that

$$\begin{aligned} K^c &= \{y : \forall x \in K c(x, y) \geq 0\} = \{y : \forall x \in K (x, y) \in S\} \\ &= \{y : \forall x \in K \exists L_{(x,y)} \in \mathcal{C} \text{ with } (x, y) \in L_{(x,y)} \times TL_{(x,y)}\}. \end{aligned}$$

Fix some $y_0 \in K^c$, and note that the above implies that we must have $K \subset \cup_{x \in K} L_{(x,y_0)}$ and that $y_0 \in \cap_{x \in K} TL_{(x,y_0)}$. By the assumption that T respects inclusions we conclude that $TK \supset \cap_{x \in K} TL_{(x,y_0)}$ and hence $y_0 \in TK$. Since $y_0 \in K^c$ was arbitrary, it follows that $K^c \subset TK$ as required. \square

5. FURTHER OBSERVATIONS

In this section we gather some methods for obtaining new order reversing quasi involutions from some given transforms, and we discuss the existence of invariant subsets under the mapping $K \mapsto K^c$.

5.1. Restrictions. We begin by collecting a few results on how to generate new order reversing quasi involutions.

Proposition 5.1. *Assume T_1 and T_2 are two order reversing quasi involutions. Then also $T_3K = T_1K \cap T_2K$ is an order reversing quasi involution.*

Proof. If T_1 is induced by c_1 and T_2 by c_2 , then letting $c_3 = \min(c_1, c_2)$ we get the transform T_3 . \square

Remark 5.2. Of course, one can also check directly that $L \subset K$ implies $T_3L \supset T_3K$ and $L \subset T_1T_1L \cap T_2T_2L$ so

$$T_3T_3L = T_3(T_1L \cap T_2L) = T_1(T_1L \cap T_2L) \cap T_2(T_1L \cap T_2L) \supset T_1T_1L \cap T_2T_2L \supset L.$$

Secondly, if an order reversing quasi involution is defined on a class $\mathcal{C} \subset \mathcal{P}(X)$ which is closed under intersections, we can define a new order reversing quasi involution by restricting the original transform to some fixed subset. More precisely, we have the following proposition.

Proposition 5.3. *Let $T : \mathcal{C} \rightarrow \mathcal{C}$ be an order reversing quasi involution on \mathcal{C} , where $\mathcal{C} \subset \mathcal{P}(X)$ is closed under intersections. Then for any fixed $M_0 \in \mathcal{C}$, denoting $\mathcal{C}_0 = \{K \cap M_0 : K \in \mathcal{C}\} \subset \mathcal{C}$, we have that $S : \mathcal{C}_0 \rightarrow \mathcal{C}_0$ defined by $SK = TK \cap M_0$ is an order reversing quasi involution.*

Proof. To see that S is order reversing, consider $K, L \in \mathcal{C}_0$ such that $K \subset L$. Then since T is order reversing we get that $TK \supset TL$. Since intersecting with M_0 preserves that inclusion, we get that S is indeed order reversing. To see that for any $K \in \mathcal{C}_0$ we have $K \subset SSK$ note that $SK = TK \cap M_0 \subset TK$. Since \mathcal{C} is closed under intersections we may apply T to both sides again and by the order reversing property of T we get $T(TK \cap M_0) \supset TTK \supset K$. Finally note that since $K \in \mathcal{C}_0$ it follows that $K \subset T(TK \cap M_0) \cap M_0 = SSK$. \square

Remark 5.4. Instead of a direct proof one can use the cost associated with T and restrict it to $M_0 \times M_0$. Such a cost exists by Lemma 2.5 and Theorem 1.3.

One can use Proposition 5.3 to construct order reversing quasi involutions. For example, considering the polarity transform on subsets of \mathbb{R}^n , and letting $M_0 = RB_2^n$, a multiple of the Euclidean ball, the new transform, which is given by

$$SK = K^\circ \cap RB_2^n,$$

is an order reversing quasi involution on subsets of RB_2^n . Its image, namely the class on which it is an involution, is the class of all subsets of RB_2^n which include $\frac{1}{R}B_2^n$. If we instead chose M_0 to be a linear subspace, we would simply get polarity within the subspace.

5.2. Composition and conjugation. There is a close relation between order reversion and order preservation. We collect a few facts.

Lemma 5.5. *Let $S, T : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ be two order reversing quasi involutions with the same image \mathcal{C} . Then $T \circ S : \mathcal{P}(X) \rightarrow \mathcal{C}$ is order preserving, and restricted to \mathcal{C} it is an order preserving bijection on \mathcal{C} .*

Proof. Order preservation is immediate, the fact that the image is included in \mathcal{C} is immediate, and bijectivity follows since if $A \in \mathcal{C}$ then $TS(STA) = TTA = A$. \square

Conjugating an order reversing quasi involution T by an order preserving bijection on $\text{Im}(T)$, one gets an order reversing involution on the image.

Lemma 5.6. *Let $T : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ be an order reversing quasi involution, let $\mathcal{C} = T(\mathcal{P}(X))$ its image, and let $R : \mathcal{C} \rightarrow \mathcal{C}$ be an order preserving bijection. Then $S = R^{-1} \circ T \circ R$ is an order reversing involution on \mathcal{C} .*

Proof. Let $A \subset B \in \mathcal{C}$. Then $RA \subset RB \in \mathcal{C}$ and so $TRA \supset TRB$, and using order preservation of R^{-1} we get $SA = R^{-1}TRA \supset R^{-1}TRB = SB$. The involution property follows from $R^{-1}TRR^{-1}TRA = R^{-1}TTRA = R^{-1}RA = A$. \square

Remark 5.7. Clearly \mathcal{C} is closed under intersections as it is the image of an order reversing quasi involution. Using Lemma 2.5 we may thus extend S to be an order reversing quasi involution on $\mathcal{P}(X)$.

Recall Lemma 2.2, where by conjugating an order reversing quasi involution with the order reversing involution $K \mapsto X \setminus K$ we obtained a complemented order reversing quasi involution

S , namely an order reversing map satisfying $K \supset SSK$. Whereas it may seem that in the specific choice of $RK = X \setminus K$ we only used that it is an order reversing involution, one quite easily checks that it is the *only* order reversing bijection on $\mathcal{P}(X)$. However, this construction can be generalized as follows.

Lemma 5.8. *Let T be an order reversing quasi involution and R be a complemented order reversing quasi involution. Then the composition TRT is an order reversing quasi involution while RTR is a complemented order reversing quasi involution.*

Proof. Assume that T, R are order reversing and that for any $K \subset X$ we have $K \subset TTK$ and $RRK \subset K$. It can be easily checked that a composition of an odd number of order reversing transforms remains order reversing. To see that TRT is a quasi involution note that

$$TT(RTK) \supset RTK \Rightarrow RTTRTK \subset RRTK \subset TK \Rightarrow TRTTRTK \supset TTK \supset K.$$

Similarly, we get that the composition RTR is a complemented quasi involution

$$RR(TRK) \subset TRK \Rightarrow TRRTRK \supset TTRK \supset RK \Rightarrow RTRRTRK \subset RRK \subset K.$$

□

5.3. Invariant sets. For the polarity transform in \mathbb{R}^n the only invariant set is the Euclidean unit ball B_2^n . Similarly, for the Legendre transform (see Example 6.12) the function $\|x\|_2^2/2$ is the unique fixed point. However, for the \mathcal{A} -transform (see Example 6.13), for instance, many self dual functions exist [13].

In this subsection we state some facts regarding invariant sets for arbitrary order reversing quasi involutions. As above, we will denote the transform on $\mathcal{P}(X)$ by $K \mapsto K^c$, where c is a symmetric cost function on X and $K^c = \{y : \forall x \in K, c(x, y) \geq 0\}$. Recall that by Theorem 1.3 any order reversing quasi involution is of this form. If one prefers to omit the cost notation, using $T : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ to denote the order reversing quasi involution at hand, one should replace the set $\{x : c(x, x) \geq 0\}$ defined below, by the set $\{x : x \in T(\{x\})\}$.

Fact 5.9. If a set K is such that $K = K^c$, then $K \subset \{x : c(x, x) \geq 0\}$.

Proof. Let $x \in K$ then $x \in K^c$ so $c(x, y) \geq 0$ for all $y \in K$ in particular for $y = x$. □

We will denote $X_0 = \{x \in X : c(x, x) \geq 0\}$, and the above fact means that if $K = K^c$ then $K \subset X_0$.

Our first result has to do with the special case where all points “stand a chance” of belonging to an invariant set.

Lemma 5.10. *Assume $X = X_0$. Then for any $x_0 \in X$ there exists an invariant set K with $x_0 \in K$.*

Proof. Consider the class \mathcal{S} of subsets $S \subset X$ with $c(x, y) \geq 0$ for all $x, y \in S$. If $S \in \mathcal{S}$ then $S \subset S^c$ by definition. The class \mathcal{S} is non-empty, as it includes $\{x_0\}$ for any $x_0 \in X$. We will show that for any fixed $S_0 \in \mathcal{S}$ there exists an invariant set K with $S_0 \subset K$. Indeed, let \mathcal{S}_0

stand for the class of sets $S \in \mathcal{S}$ such that $S_0 \subset S$. This is again a non-empty class as it includes S_0 itself. Moreover, given a chain (with respect to inclusion) in \mathcal{S}_0 , the union of its elements still belongs to \mathcal{S}_0 and so the conditions of Zorn's lemma are satisfied, that is, every chain has an upper bound. Therefore one may find a maximal element in \mathcal{S}_0 . Denote it by K . Clearly $K \subset K^c$ (since $K \in \mathcal{S}$). However, were it the case that some $z \in K^c \setminus K$, then $K_1 = K \cup \{z\}$ would also belong to \mathcal{S}_0 . Indeed, to this end one uses the fact that $c(z, z) \geq 0$ for all $x \in K$ since by assumption we have $c(z, z) \geq 0$ since $z \in X_0 = X$. But this contradicts maximality. Therefore, $K = K^c$. \square

In the proof we used $X = X_0$ in precisely one place, and the exact same proof gives the following

Theorem 1.5. *Let $S_0 \subset X$ satisfy $c(x, y) \geq 0$ for all $x, y \in S_0$. Then there exists some $K \subset X$ with $S_0 \subset K$ and such that $K^c \cap X_0 = K$.*

We cannot expect much more without extra assumptions, since we may always artificially add to a space X a point z such that $c(z, z) = -1$ and $c(x, z) = 1$ for all $x \neq z$. In such a case, for any K with $z \notin K$ we will have $z \in K^c$ and for any K with $z \in K$ we will have $z \notin K^c$.

Remark 5.11. Let us analyze an important example with multiple invariant sets, namely the \mathcal{A} transform on $\text{Cvx}_0(\mathbb{R}^+)$, see Example 6.13. The invariant sets for this transform were analyzed by Rotem in [13]. In this example, the set X is the upper half space, so a point can be written as $\bar{x} = (x, t)$ with $x \in \mathbb{R}^+$ and $t \geq 0$. The cost is given by $c((x, t), (y, s)) = st + 1 - xy$ and so X_0 is the set (x, t) with $t^2 + 1 - x^2 \geq 0$, or $t \geq \sqrt{x^2 - 1}$. If we fix a point (x_0, t_0) on the boundary of X_0 , then for points outside X_0 , say (y, s) with $s^2 < y^2 - 1$, we have (as clearly $x, y \geq 1$) that

$$c((x, t), (y, s)) = st + 1 - xy = s\sqrt{x^2 - 1} + 1 - xy < \sqrt{y^2 - 1}\sqrt{x^2 - 1} + 1 - xy \leq 0.$$

In other words, having chosen some \bar{x} on the boundary of X_0 , and taken K to be as given in Theorem 1.5, we get that $K^c \subset X_0$ automatically and so $K^c = K$. So for this specific example, for any point in the boundary of X_0 , there exists an invariant set which includes this point. Moreover, no two different points on the boundary of X_0 can belong to the same invariant set (as the inequality above remains valid, where the strictness of the inequality follows from strictness of the arithmetic geometric inequality when the points are different). This is a reason for the multitude of invariant sets in this specific case.

While a complete characterization of invariant sets for order reversing quasi involutions seems difficult, because of the various scenarios which can occur, we can nevertheless gather several more useful facts in this direction. In two of our favorite examples, namely \mathbb{R}^n with the polarity transform and $\text{Cvx}(\mathbb{R}^n)$ with the Legendre transform, there is a unique invariant set, which in the first case is $X_0 = B_2^n$ and $X_0 = \{(x, t) \in \mathbb{R}^n \times \mathbb{R} : t \geq \|x\|^2/2\}$ in the second. These correspond to the instance described in the first item of the following lemma.

Lemma 5.12. *As above, let X be some set and $K \mapsto K^c$ an order reversing quasi involution associated to some symmetric cost function c . Let $X_0 = \{x : c(x, x) \geq 0\}$.*

- (1) *If $X_0^c = X_0$, then X_0 is the unique invariant set for the transform.*
- (2) *If $X_0^c \not\subseteq X_0$ then there is no invariant set for the transform.*
- (3) *If $X_0^c \subsetneq X_0$ then there are examples where no invariant set exists, examples where only one invariant set exists, and examples where more than one invariant set exists.*

Proof. For (1), if $K = K^c$, as we already know $K \subset X_0$ by Fact 5.9, and by order reversion we get that $K = K^c \supset X_0^c = X_0$ and so $K = X_0$.

To see that (2) holds, by the same reasoning, if $K = K^c$ then we have that $X_0 \supset K = K^c \supset X_0^c$, and so if this inclusion does not hold, there will be no invariant sets.

For (3), one case of many invariant sets was given in Remark 5.11. A simpler case is when $c(x, x) = 1$ and for $x \neq y$ we let $c(x, y) = -1$. Then any $\{x\}$ is an invariant set, $X_0 = X$, and $X_0^c = \emptyset$ as soon as X has more than one point in it. An example with just one invariant set, despite $X_0^c \subsetneq X_0$, can be given with a three point space. Let $X = \{1, 2, 3\}$ and set $c(1, 1) = c(2, 2) = c(1, 3) = c(3, 1) = 1$ and all other pairs to have cost -1 . In this case it is easy to check that $X_0 = \{1, 2\}$ and $X_0^c = \emptyset$. The only invariant set is $\{2\}$, and the other subsets of $\{1, 2\}$ satisfy $\{1\} \mapsto \{1, 3\}$ and $\{1, 2\} \mapsto \emptyset$ whereas $\emptyset \mapsto \{1, 2, 3\}$ of course. Finally, to give an example where no invariant set exists despite $X_0^c \subset X_0$, we consider a four point space $X = \{1, 2, 3, 4\}$. We let the cost be given by $c(1, 1) = c(2, 2) = c(1, 3) = c(3, 1) = c(2, 4) = c(4, 2) = 1$, and the cost is -1 for any other pair. If there was some invariant set, it would have to be included in $X_0 = \{1, 2\}$. However $\{1\} \mapsto \{1, 3\}$ and $\{2\} \mapsto \{2, 4\}$, whereas $X_0^c = \emptyset$ (and as usual $\emptyset \mapsto \{1, 2, 3, 4\}$). \square

6. A ZOO OF EXAMPLES

The theory of order reversing quasi involutions has many implications for geometry, as many of the classical operations considered in the theory are in fact of this form. In this section we gather many of these examples, as well as some less standard ones which have received attention in recent years and several new examples which may be of interest.

6.1. Some trivial examples. As a first example, we consider order reversing quasi involutions whose image consists of a finite number of sets. In this case, determining the various order reversing quasi involutions reduces to a question of combinatorial nature. We list below the first few examples.

- Example 6.1.**
- (1) *When there is a single element in the class, it must be all of X and then the order reversing quasi involution is such that all sets are mapped to X .*
 - (2) *When the image includes two sets they must be $K \neq X$ and X , and the only possible duality takes all subsets of K to X , and anything which is not a subset of K to K .*
 - (3) *An order reversing quasi involution whose image has three elements must have as these three elements the set X , and two comparable sets $K_1 \subset K_2$, where T switches*

X and K_1 , while K_2 is a fixed point for T . The pre-image of X is precisely all L such that $L \subset K_1$. The pre-image of K_2 consists of subsets of K_2 which are not subsets of K_1 . All other sets are mapped to K_1 .

- (4) A similarly trivial example is when $\mathcal{C} = \{X, K_0, K_1, K_2\}$ where $K_0 \subset K_1 \subset K_2$, and the mapping switches the two, namely $TK_1 = K_2$ and $TK_2 = K_1$. If $K \subset K_0$ then $T(K) = X$, if $K \subset K_1$ and $K \not\subset K_0$ then $T(K) = K_2$, if $K \subset K_2$ and $K \not\subset K_1$ then $TK = K_1$ and if $K \not\subset K_2$ then $TX = K_0$.

In the next example we consider X which is a metric space, and the transform maps a set to the complement of its t -neighborhood.

Example 6.2. Consider a metric-induced cost, that is, let $X = Y$ be some metric space, and $c(x, y) = d(x, y)$ (more generally one can take $f(d(x, y))$ where $f : [0, \infty) \rightarrow [0, \infty)$ is increasing; for example, $f(x) = x^p$, and the same phenomenon described below occurs. Note that this case includes the classical quadratic cost on \mathbb{R}^n). In such a case,

$$A^c = X \setminus A_t \quad \text{and} \quad A^{cc} = X \setminus (X \setminus A_t)_t.$$

It is clear that $A \subset A^{cc}$, and that for a closed convex subset K of the Euclidean space \mathbb{R}^n , $K^{cc} = K$. In general, there is equality $A = A^{cc}$ if and only if the complement of the set A can be written as a union of balls of radius t .

In the next two sections we will restrict our geometry further, first looking at the class of star shaped sets, and then at classes of convex sets and other related families.

6.2. Examples from convex geometry. Recall once more the class \mathcal{K}_0^n of closed convex bodies including the origin with the usual polarity transform

$$K^\circ = \{y : \forall x \in K \langle x, y \rangle \leq 1\}$$

(for an overview of classical convexity see [15]). This class arises from the classical cost $c(x, y) = -\langle x, y \rangle + 1$. Indeed, $K^c = \{y : \forall x \in K -\langle x, y \rangle + 1 \geq 0\} = K^\circ$. Böröczky and Schneider [7] showed that polarity is essentially the only order reversing involution on \mathcal{K}_0^n . Slightly modifying the cost we get an order reversing quasi involution whose image is the class of unconditional convex bodies.

Example 6.3. Let $X = \mathbb{R}^n$ and consider the cost $c(x, y) = \sum_{i=1}^n |x_i||y_i|$. Then the image of the transform induced by this cost is the family of unconditional convex bodies, namely those convex bodies symmetric with respect to all coordinate hyperplanes.

Another example where the image is a subset of the class of convex sets is as follows.

Example 6.4. Let $X = \mathbb{R}^n$ and consider the cost function $c(x, y) = \langle x, y \rangle - 1$. Then the image of the cost transform consists of intersections of affine half-spaces that do not include the origin. In particular, these are unbounded, closed and convex sets.

Our next examples consist of two classes of bodies studied by Milman, Milman and Rotem [11]: The first are reciprocal bodies and they are a subclass of convex bodies.

Example 6.5. Consider $X = \mathbb{R}^n$ and the following cost function

$$c(x, y) = \sup_{\theta \in S^{n-1}} (\langle x, \theta \rangle \langle y, \theta \rangle) - 1.$$

We claim that the image of this class is reciprocal bodies, namely the Wulff shape (or Alexandrov body) associated with the function $1/h_K$ (on the sphere), i.e. the set

$$\{y \in \mathbb{R}^n : \forall \theta \in S^{n-1} \langle y, \theta \rangle \leq 1/h_K(\theta)\}.$$

We compute the c -transform for any set K :

$$\begin{aligned} K^c &= \{y \in \mathbb{R}^n : \forall x \in K \ c(x, y) \geq 0\} \\ &= \{y \in \mathbb{R}^n : \forall x \in K \ \theta \in S^{n-1} (\langle x, \theta \rangle \langle y, \theta \rangle) \leq 1\} \\ &= \{y \in \mathbb{R}^n : \forall \theta \in S^{n-1} \sup_{x \in K} (\langle x, \theta \rangle \langle y, \theta \rangle) \leq 1\} \\ &= \{y \in \mathbb{R}^n : \forall \theta \in S^{n-1} \ h_K(\theta) \langle y, \theta \rangle \leq 1\} \\ &= \{y \in \mathbb{R}^n : \forall \theta \in S^{n-1} \ \langle y, \theta \rangle \leq 1/h_K(\theta)\}. \end{aligned}$$

Recalling that h_K is always a convex support function, even if K was not convex, we conclude that indeed the image of the transform is reciprocal bodies.

Remark 6.6. Our theory of cost dualities gives, as an immediate corollary, some properties of the reciprocal transform shown in [11]. First, $K''' = K'$, as by Lemma 2.1 any order reversing quasi involution is a duality on its image.

Second, $K' \subset K^\circ$, which we can show by comparing the following cost transforms

$$c(x, y) = -\ln[\sup_{\theta \in S^{n-1}} (\langle x, \theta \rangle \langle y, \theta \rangle)] \leq -\ln[(\langle x, y/|y| \rangle \langle y, y/|y| \rangle)] = -\ln(\langle x, y \rangle) =: c_2(x, y).$$

Indeed, we have shown that, given a set K , $K^c = K'$, and one may easily check that $K^{c_2} = K^\circ$, and thus

$$K' = \{y : \inf_{x \in A} c(x, y) \geq 0\} \subset \{y : \inf_{x \in A} c_2(x, y) \geq 0\} = K^\circ.$$

In fact, in view of Proposition 5.1, one can think of the previous example as an intersection of a continuum of simple order reversing quasi involutions:

Example 6.7. Let $c_\theta(x, y) = 1 - \langle x, \theta \rangle \langle y, \theta \rangle$. In such a case one gets

$$K^{c_\theta} = \{y : \forall x \in K, \langle y, \theta \rangle \leq 1/\langle x, \theta \rangle\},$$

which simply means that $K^{c_\theta} = (P_{\theta\mathbb{R}}K)^\circ$. In particular, the image consists of all slabs in direction θ , a very simple one dimensional class.

As $\inf_\theta c_\theta$ gives the cost in the previous example, we see that the reciprocal body has an elegant representation as

$$K' = \cap_\theta (P_\theta K)^\circ.$$

Our final example for this subsection consists of a relatively new class of bodies, promoted by V. Milman (see [11, 12]), called flowers. These are intimately related to reciprocals and polars, and are not convex but in fact star shaped. A flower is defined as a union of the form

$A = \bigcup_{x \in C} B\left(\frac{x}{2}, \frac{|x|}{2}\right)$, where $C \subset \mathbb{R}^n$ is some closed set. The class of flowers is therefore closed under unions (not intersections). Hence, we present an example of a cost which generates the class of complements of flowers (for the relation between a class and the class of complements recall Lemma 2.2).

The flower transform $K \mapsto K^\clubsuit$ was defined in [11] to be the star shaped set with radial function $r_{K^\clubsuit} = h_K$.

Example 6.8. Consider $X = \mathbb{R}^n$, the set $S \subset X \times X$ given by

$$S = \{(x, y) : \langle x, y \rangle < \frac{1}{2}|x|^2|y|^2\}.$$

Then, the cost

$$c(x, y) = \begin{cases} 1 & x \in S \\ -1 & x \notin S \end{cases}$$

will generate the class of complements of flowers, namely sets of the form $X \setminus \bigcup_{x \in C} B\left(\frac{x}{2}, \frac{|x|}{2}\right)$, where $C \subset \mathbb{R}^n$ is some closed set. Indeed, the c -dual of a single point $x_0 \in X$ is given by

$$\{y : 2\langle z_0, y \rangle < |y|^2\} = \mathbb{R}^n \setminus B(z_0, |z_0|)$$

where $z_0 = x_0/|x_0|^2$. This implies that all sets in the class are complements of flowers. Since the spherical inversion mapping $\Phi(x) = x/|x|^2$ is bijective, by Proposition 2.7 we conclude that all complements of flowers are in the class: $A = X \setminus \bigcup_{x \in C} B\left(\frac{x}{2}, \frac{|x|}{2}\right)$ is the c -dual of the set $\{\Phi^{-1}(x/2) : x \in C\}$. In other words, $K^\clubsuit = X \setminus (\Phi[K/2])^c$.

6.3. Examples with star shaped sets. In the examples that follow, we consider the space $X = \mathbb{R}^n$ and the subset $\mathcal{C} \subset \mathcal{P}(X)$ of closed star shaped sets, namely those sets $A \subset \mathbb{R}^n$ which satisfy $A = \{\lambda x : x \in A, 0 \leq \lambda \leq 1\}$. Given a closed star shaped set A , denote by g_A its gauge function which uniquely determines it. We discuss cost dualities whose image is this class. As it turns out, there are various different dualities on the class.

Example 6.9. Consider the cost

$$c(x, y) = \begin{cases} -\langle x, y \rangle + 1 & y \in \mathbb{R}_+ x \\ +\infty & \text{otherwise} \end{cases}.$$

The image of the c -transform is the class of star shaped bodies, and the c -duality on it takes the gauge function g to $1/g$.

Indeed, consider a closed star shaped A Then

$$\begin{aligned} A^c &= \{y \in \mathbb{R}^n : \forall x \in A, c(x, y) \geq 0\} \\ &= \{y \in \mathbb{R}^n : \forall x \in [0, y/g_A(y)], \langle x, y \rangle \leq 1\} \\ &= \{y \in \mathbb{R}^n : \langle y/g_A(y), y \rangle \leq 1\} \\ &= \{y \in \mathbb{R}^n : |y|^2 \leq g_A(y)\}. \end{aligned}$$

This set is clearly closed, star shaped, and in fact for $u \in S^{n-1}$, if $g_A(u) = \lambda$ then

$$g_{A^c}(u) = \inf\{r : u/r \in A^c\} = \inf\{r : |u/r|^2 \leq g_A(u/r)\} = \inf\{r : 1/r \leq \lambda\} = 1/\lambda = 1/g_A(u).$$

In particular $A \mapsto A^c$ is an involution on the class of closed star shaped domains (not necessarily bounded, and 0 can be on the boundary). It is also easy to check that A^c is always closed and star shaped.

Example 6.10. We may generalize the cost from Example 6.9 to get a large family of dualities on star shaped domains as follows. For any $u \in S^{n-1}$ let $f_u : [0, \infty) \rightarrow [0, \infty)$ be an increasing bijective function, such that f_u depend continuously on u . Let $F : S^{n-1} \rightarrow S^{n-1}$ be continuous and satisfy $F \circ F = Id$. Then

$$c(x, y) = \begin{cases} -f_{y/|y|}(\langle x, |y|F(y/|y|) \rangle) + 1 & y/|y| = F(x/|x|) \\ +\infty & \text{otherwise} \end{cases}$$

is a cost whose image is, again, closed star shaped sets. Indeed, for A a closed star shaped set,

$$\begin{aligned} A^c &= \{y \in \mathbb{R}^n : \forall x \in A, c(x, y) \geq 0\} \\ &= \{y \in \mathbb{R}^n : \forall x \in [0, F(y/|y|)/g_A(F(y/|y|))], \langle x, |y|F(y/|y|) \rangle \leq f_{y/|y|}^{-1}(1)\} \\ &= \{y \in \mathbb{R}^n : \langle F(y/|y|)/g_A(F(y/|y|)), |y|F(y/|y|) \rangle \leq f_{y/|y|}^{-1}(1)\} \\ &= \{y \in \mathbb{R}^n : |y| \leq f_{y/|y|}^{-1}(1)g_A(F(y/|y|))\}. \end{aligned}$$

The resulting set is clearly closed and star shaped. One may easily check that K^c is closed and star shaped even if K is not. Finally, we again see that for $u \in S^{n-1}$, if we have $g_A(u) = \lambda$ then

$$g_{A^c}(u) = \inf\{r : u/r \in A^c\} = \inf\{r : |u/r| \leq f_u^{-1}(1)g_A(F(u))\} = f_u^{-1}(1)/\lambda.$$

This implies that the image of the c -duality is the class of closed star shaped sets.

Remark 6.11. Before moving to the next family of examples, let us mention that while many geometric families of sets appear as images of order reversing quasi involutions or their complement, some classical families of convex bodies cannot arise in this way. For example, the classes of Zonotopes (Minkowski sums of a finite number of segments) or Zonoids (limits of Zonotopes) as they are not closed under intersection or unions.

6.4. Examples of function classes. When considering an order reversing quasi involution on a class of sets, if these sets are epi-graphs of functions we can view the transform on them as a functional transform. The inclusion order on epigraphs translates to a pointwise inequality of the functions. We say that T is an order reversing quasi involution on a class \mathcal{C} of functions $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$ if for any $f, g \in \mathcal{C}$ we have

- (i) $TTf \leq f$ (quasi involution)
- (ii) $f \leq g$ implies $Tf \geq Tg$ (order reversion)

In other words, a functional duality T corresponds to an order reversing quasi involution on $P(X \times \mathbb{R})$, which we also denote by T , whose image class is the set of epi-graphs of functions in the image of T . We define it on sets which are epi-graphs by $T(\text{epi}(f)) := \text{epi}(Tf)$.

Example 6.12. Consider the class of $\text{Cvx}(\mathbb{R}^n)$ consisting of convex functions $\phi : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ which are lower semi continuous. The Legendre transform,

$$\mathcal{L}\phi(y) = \sup_x (\langle x, y \rangle - \phi(x))$$

is an order reversing quasi involution on this class. The same definition applied to all functions, can be shown to satisfy $\mathcal{L}\mathcal{L}\phi \leq \phi$ for all ϕ .

The class of epi-graphs of functions in $\text{Cvx}(\mathbb{R}^n)$ is closed under intersection and in particular, the transform can thus be expressed as a cost-induced transform on subsets of \mathbb{R}^{n+1} . It is easily checked that considering the cost function on \mathbb{R}^{n+1} given by $c((x, t), (y, s)) = t + s - \langle x, y \rangle$ we recover, as the c -transform, the order reversing quasi involution whose restriction to the class of epi-graphs is the one corresponding to the Legendre transform.

Indeed, for some (y, s) to belong to $\text{epi}(\mathcal{L}\phi)$ one needs

$$s \geq \mathcal{L}\phi(y) \Leftrightarrow \forall x, s \geq \langle x, y \rangle - \phi(x) \Leftrightarrow s \geq \langle x, y \rangle - t, \forall (x, t) \in \text{epi}(\phi),$$

and thus

$$T(\text{epi}(\phi)) = \{w = (y, s) : \forall z = (x, t) \in \text{epi}(\phi), c(z, w) \geq 0\} = \text{epi}(\mathcal{L}\phi).$$

We can apply this transform to any subset of \mathbb{R}^{n+1} , and always end up with the graph of a convex lower semi continuous function.

Example 6.13. Consider the class $\text{Cvx}_0(\mathbb{R}^n)$ of geometric convex functions on \mathbb{R}^n (non-negative lower semi-continuous convex functions with minimum 0 at the origin) and on it the order reversing involution \mathcal{A} , given by

$$\mathcal{A}(\phi) = \sup_{y \in \mathbb{R}^n} \frac{\langle x, y \rangle - 1}{\phi(y)}.$$

For discussion of this transform and its many properties, see [4]. Again, the family of epi-graphs of functions in $\text{Cvx}_0(\mathbb{R}^n)$ is closed under intersection. Further, the transform on it can be geometrically described as a composition of standard duality in \mathbb{R}^{n+1} with a reflection (see again [4]). Therefore, the transform is induced by the following cost

$$c((x, t), (y, s)) = st + 1 - \langle x, y \rangle.$$

Remark 6.14. It is well known that \mathcal{L} and \mathcal{A} agree on the subclass of $\text{Cvx}_0(\mathbb{R}^n)$ consisting of 2-homogeneous functions. However, on $\text{Cvx}_0(\mathbb{R}^n)$ they \mathcal{L} and \mathcal{A} differ substantially, and therefore so do their extensions to $\mathcal{P}(\mathbb{R}^{n+1})$. We thus have an example of an order reversing quasi involution (on 2-homogeneous convex functions in \mathbb{R}^n) which admits very different extensions. Of course, each of these extensions has a different image ($\text{Cvx}(\mathbb{R}^n)$ and $\text{Cvx}_0(\mathbb{R}^n)$, respectively).

The two above examples, up to composition with a monotone one-dimension function, belong to a special class of functional order reversing quasi involutions (complemented, in fact) which are used in optimal transport theory. These are the so-called cost-transforms (see [18]), defined as follows.

Definition 6.15. *Let $c : X \times X \rightarrow (-\infty, \infty]$ be some cost function. For $\psi : X \rightarrow \mathbb{R} \cup \{+\infty\}$ its **c -transform** is given by*

$$\psi^c(x) = \inf_{y \in X} (c(x, y) - \psi(y)),$$

In general, such a cost transform, induces a transform on hypo-graphs of functions, namely the $K \subset \mathbb{R}^{n+1}$ such that for any fixed $x \in \mathbb{R}^n$, the set $\{t : (x, t) \in K\}$ is of the form $(-\infty, a]$ for some a , or \mathbb{R} , or empty, a class closed under intersection. Call $\{(x, t) : t \leq \psi(x)\} = \text{hypo}(\psi)$. Given $T : \mathcal{C} \rightarrow \mathcal{C}$ where \mathcal{C} is a class of functions $\psi : \mathbb{R}^n \rightarrow [-\infty, \infty]$, which satisfies $TT\psi \geq \psi$ and $\psi \leq \varphi$ implies $T\psi \geq T\varphi$ we let

$$T(\text{hypo}(\psi)) := \text{hypo}(T\psi).$$

For example, $\psi \mapsto \psi^c$ satisfies these properties (see, e.g., [5]).

As before, since $\psi^{cc} \geq \psi$ we have that $\text{hypo}(\psi^{cc}) \supset \text{hypo}(\psi)$ so that T is an order reversing quasi involution on the class of hypo-graphs of functions in $\text{Cvx}_0(\mathbb{R}^n)$.

It is easily checked that the cost function corresponding to the order reversing quasi involution on sets is given by

$$c((x, t), (y, s)) = c(x, y) - s - t.$$

REFERENCES

- [1] S. Artstein-Avidan, D. Florentin, and V. Milman. Order isomorphisms on convex functions in windows. In *Geometric aspects of functional analysis*, pages 61–122. Springer, 2012.
- [2] S. Artstein-Avidan and V. Milman. A new duality transform. *Comptes Rendus Mathematique*, 346(21-22):1143–1148, 2008.
- [3] S. Artstein-Avidan and V. Milman. The concept of duality in convex analysis, and the characterization of the Legendre transform. *Annals of mathematics*, pages 661–674, 2009.
- [4] S. Artstein-Avidan and V. Milman. Hidden structures in the class of convex functions and a new duality transform. *Journal of the European Mathematical Society*, 13(4):975–1004, 2011.
- [5] S. Artstein-Avidan, S. Sadovskiy, and K. Wyczesany. A Rockafellar-type theorem for non-traditional costs. *arXiv:2011.13263 (submitted)*, 2020.
- [6] S. Artstein-Avidan, S. Sadovskiy, and K. Wyczesany. Optimal measure transportation with respect to non-traditional costs. *arXiv:2104.04838 (submitted)*, 2021.
- [7] K. Böröczky and R. Schneider. A characterization of the duality mapping for convex bodies. *Geometric and Functional Analysis*, 18(3):657–667, 2008.
- [8] D. Florentin and A. Segal. Stability of order preserving transforms. In *Geometric Aspects of Functional Analysis*, pages 205–225. Springer, 2012.
- [9] P. Gruber. The endomorphisms of the lattice of convex bodies. In *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, volume 61, pages 121–130. Springer, 1991.
- [10] A. Iusem, D. Reem, and B. Svaiter. Order preserving and order reversing operators on the class of convex functions in Banach spaces. *Journal of Functional Analysis*, 268(1):73–92, 2015.

- [11] E. Milman, V. Milman, and L. Rotem. Reciprocals and Flowers in Convexity. In *Geometric Aspects of Functional Analysis*, pages 199–227. Springer, 2020.
- [12] V. Milman and L. Rotem. Novel view on classical convexity theory. *arXiv:2005.11253*, 2020.
- [13] L. Rotem. Characterization of self-polar convex functions. *Bulletin des Sciences Mathématiques*, 136(7):831–838, 2012.
- [14] R. Schneider. The endomorphisms of the lattice of closed convex cones. *Beitr. Algebra Geom*, 49:541–547, 2008.
- [15] R. Schneider. *Convex bodies: the Brunn–Minkowski theory*. Number 151. Cambridge university press, 2014.
- [16] A. Segal and B. Slomka. Duality on convex sets in generalized regions. In *Asymptotic Geometric Analysis*, pages 289–298. Springer, 2013.
- [17] B. Slomka. On duality and endomorphisms of lattices of closed convex sets. 2011.
- [18] Cédric Villani. *Optimal transport: old and new*, volume 338. Springer Science & Business Media, 2008.