

The Charged Zipoy - Voorhees Metric

Ozay Gurtug*

*T. C. Maltepe University, Faculty of Engineering
and Natural Sciences, 34857, Istanbul -Turkey*

Mustafa Halilsoy[†] and Mert Mangut[‡]

*Physics Department, Eastern Mediterranean University,
Famagusta, 99628 North Cyprus via Mersin 10, Turkey.*

Abstract

Starting from an integral of the interaction region of colliding Einstein-Maxwell waves and by applying a coordinate transformation, we obtain the charged version of the static Zipoy-Voorhees metric. In Schwarzschild coordinates we investigate the effect of the charge in the newly found spacetime and discuss some of its features.

*Electronic address: ozaygurtug@maltepe.edu.tr

†Electronic address: mustafa.halilsoy@emu.edu.tr

‡Electronic address: mert.mangut@emu.edu.tr

I. INTRODUCTION

The existence of non-spherical stellar objects brings into focus the geometry of metrics with intricate topologies known as the γ -metric or Zipoy-Voorhees (ZV) class of metrics [1–3]. Although there is a growing interest in such planetary objects when searched in the literature, for their charged version, one comes across with only a few references [4, 5]. In their approach [4], test field solution in a background geometry is found by the Killing symmetry and from the energy-momentum of the resulting electromagnetic (em) field, the authors proceed to establish the Einstein-Maxwell (EM) spacetime. Expectedly, the final part is technically difficult and for that reason we follow an alternative procedure in this article. We approach the problem from the spacetime of colliding (EM) waves formulated long ago [6, 7] and transform the metric into the static, non-spherical form of ZV . More clearly, the field equations are solved in prolate coordinates which are found to be very useful in the description of colliding gravitational waves. However, the resulting solution is interpreted in some other coordinates. For this purpose, we use the line element of Chandrasekhar and Xanthopoulos (CX) [8, 9] as the starting point and employ the method of [10]. The spacetime of colliding EM waves is described by the line element

$$ds^2 = \sqrt{\Delta} e^N \left(\frac{d\eta^2}{\Delta} - \frac{d\mu^2}{\delta} \right) - \sqrt{\Delta\delta} (\chi dx^2 + \chi^{-1} dy^2) \quad (1)$$

in which the metric functions (N, χ) depend on (η, μ) alone, $\Delta = 1 - \eta^2$ and $\delta = 1 - \mu^2$. The coupled Ernst equation [11, 12] for EM fields are given by the pair

$$(\xi\bar{\xi} + \eta\bar{\eta} - 1) \nabla^2 \xi = 2\nabla\xi (\bar{\xi}\nabla\xi + \bar{\eta}\nabla\eta) \quad (2)$$

$$(\xi\bar{\xi} + \eta\bar{\eta} - 1) \nabla^2 \eta = 2\nabla\eta (\bar{\xi}\nabla\xi + \bar{\eta}\nabla\eta) \quad (3)$$

where ξ and η represent the gravitational and em complex potentials, respectively. Note that the bar denotes complex conjugation and the operators ∇ and ∇^2 are defined on an appropriate base manifold. We also record that an alternative parametrization for the Ernst potentials are

$$Z = \frac{1 + \xi}{1 - \xi} \quad (4)$$

$$H = \frac{\eta}{1 - \xi} \quad (5)$$

where Z and H become the gravitational and *em* potentials, respectively. The rest of the paper is organized as follows. In section II, we shall solve the Ernst equation, transform the metric into Schwarzschild coordinates and in section III, we shall discuss its main properties. Our conclusion will be presented in section IV.

II. SOLUTION FOR THE ERNST SYSTEM

A particular, real solution to the foregoing set of equations (2, 3), is given by

$$\xi = p\xi_0 \quad (6)$$

$$\eta = q\xi_0 \quad (7)$$

where $\xi_0 = \bar{\xi}_0$ and the real constants (p, q) satisfy

$$p^2 + q^2 = 1. \quad (8)$$

The choice $\xi_0 = \tanh X$, where $X(\eta, \mu)$ satisfies the Euler-Darboux equation

$$(\Delta X_{,\eta})_{,\eta} - (\delta X_{,\mu})_{,\mu} = 0 \quad (9)$$

in which a comma denotes partial derivative, solves the Ernst system in the real domain.

We now make the choice

$$e^{2X} = \left(\frac{1 - \eta}{1 + \eta} \right)^\gamma \quad (10)$$

where γ is a constant to be identified in the sequel as the ZV - parameter. From the referances [8–10], it is kown that the metric function χ is given by

$$\chi = \frac{\sqrt{\Delta\delta}}{\Psi} \quad (11)$$

where the function Ψ is determined from

$$Z = \Psi + H^2. \quad (12)$$

The remaining metric function N is obtained from the integrability equations [10]

$$(N + \ln \Psi)_{,\eta} = \frac{2\eta}{\delta - \Delta} + \frac{\eta}{\Delta} + \frac{2\delta}{\delta - \Delta} [2\mu \Delta X_{,\eta} X_{,\mu} - \eta (\Delta X_{,\eta}^2 + \delta X_{,\mu}^2)]. \quad (13)$$

$$(N + \ln \Psi)_{,\mu} = \frac{2\mu}{\Delta - \delta} + \frac{2\Delta}{\Delta - \delta} [2\eta \Delta X_{,\eta} X_{,\mu} - \mu (\Delta X_{,\eta}^2 + \delta X_{,\mu}^2)]. \quad (14)$$

Upon substitution for X from (10) and integrating for the metric function N , we obtain the final line element

$$ds^2 = M^2(\eta) \left[\Delta^{\gamma^2} (\delta - \Delta)^{1-\gamma^2} \left(\frac{d\eta^2}{\Delta} - \frac{d\mu^2}{\delta} \right) - \Delta \delta dx^2 \right] - \frac{dy^2}{M^2(\eta)} \quad (15)$$

in which

$$M(\eta) = \cosh X - p \sinh X. \quad (16)$$

At this point, it is worthwhile that this line element must satisfy the appropriate boundary conditions of the incoming and interaction regions in order to be considered as a solution to the problem of colliding EM waves. In this regard (15) is not a promising candidate as the factor $(\delta - \Delta)$ in the metric fails to satisfy the boundary conditions. As a solution, however, in the interaction region alone it is useful, as we shall show below.

Having applied the coordinate transformation

$$\begin{aligned} p\eta + 1 &= \frac{r}{m} \quad , \quad x = \varphi \\ \mu &= \cos\theta \quad , \quad y = \tau \end{aligned} \quad (17)$$

supplemented with the condition $(-1)^\gamma = -1$, and appropriate rescaling of time ($\tau \rightarrow t$), we cast the metric into the form

$$ds^2 = \frac{\Delta^\gamma}{K^2} dt^2 - \frac{K^2}{\Delta^\gamma} \left[\Delta^{\gamma^2} \Sigma^{1-\gamma^2} \left(\frac{dr^2}{\Delta} + r^2 d\theta^2 \right) + r^2 \Delta \sin^2 \theta d\varphi^2 \right]. \quad (18)$$

Our notation here is as follows

$$\Delta(r) = 1 - \frac{2m}{r} + \frac{m^2 q^2}{r^2} \quad (19)$$

$$\Sigma(r, \theta) = 1 - \frac{2m}{r} + \frac{m^2}{r^2} (q^2 + p^2 \sin^2 \theta) \quad (20)$$

and

$$K(r) = (1+p) \left(1 - \frac{m(1-p)}{r}\right)^\gamma - (1-p) \left(1 - \frac{m(1+p)}{r}\right)^\gamma. \quad (21)$$

It is checked that line element (18) solves the *EM* equations and for $q = 0$ ($p = 1$), it reduces to the *ZV*-metric. Furthermore, although we assumed the condition $(-1)^\gamma = -1$ for the parameter γ in the beginning, we noticed that this condition can be released and the metric becomes valid for all γ 's. The reason is that the transformation for $\gamma = \text{even}$ also can be incorporated through an analytic continuation $y \rightarrow i\tau$ and $x \rightarrow i\phi$, letting the other coordinates as in the previous transformation. In the next section, we investigate some of the features of our new metric (18).

III. PROPERTIES OF THE SOLUTION

In this section we shall investigate some of the properties of our metric (18). Firstly, we shall expand the time component of the metric to see the asymptotic behaviour for large r values. This will show the Newtonian potential in the presence of both mass and charge. Secondly, we show that static magnetic and electric fields can separately be considered as the source to our spacetime. And thirdly, we shall consider the singularity distribution in our metric.

A. Newtonian Limit

As usual, in order to see the Newtonian limit, we take the g_{tt} component of the metric and express it in the form

$$g_{tt} = \frac{\Delta^\gamma}{K^2} \approx 1 + 2\phi \quad (22)$$

where $\phi(r)$ is the asymptotic expression of the Newtonian potential. Upon expansion to the order r^{-3} , we obtain

$$\phi(r) \approx -\frac{m\gamma}{r} + \frac{\gamma m^2}{2r^2} [\gamma q^2 + 2(\gamma - 1)] - \frac{\gamma(\gamma - 1)}{3} \left(\frac{m}{r}\right)^3 [(1 + 4\gamma)q^2 + 2(\gamma - 2)] + \mathcal{O}\left(\frac{1}{r^4}\right). \quad (23)$$

It is observed that the physical mass, $M = m\gamma$ and charge $Q = mq$ play roles at the monopole, dipole and the quadrupole orders. It will be in order at this point to comment that without charge ($q = 0$), the dipole term $\sim 1/r^2$ can be removed by two successive transformations on the radial coordinate [18]. We shall ignore such a procedure since it will be a repetition. Since we have charge in the present case, the $1/r^2$ term can be attributed to the charge. For the simplest case of spherical symmetry ($\gamma = 1$), we recover the asymptotic form of the Reissner-Nordstrom (*RN*) potential. It is observed also that the case of charge without mass is not available. However, for the pure electromagnetic limit in the case of $\gamma = 1$, which yields the Bertotti-Robinson (*BR*) metric [13, 14], one must consult [10].

B. Electromagnetic Sources for the Metric

1. Pure magnetic case

The vector potential is chosen as

$$A_\mu = (0, 0, 0, C_0 \cos\theta) \quad (24)$$

where C_0 is the magnetic charge proportional to q . The magnetic field becomes $F_{\theta\varphi} = -C_0 \sin\theta$ from $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, which solves the only relevant Maxwell equation

$$\nabla_\mu F^{\mu\nu} = 0. \quad (25)$$

The invariant of the magnetic field is given by

$$I = F_{\mu\nu} F^{\mu\nu} = \frac{2C_0^2}{r^4 K^4} \Delta^{-(\gamma-1)^2} \Sigma^{\gamma^2-1} \quad (26)$$

which trivially reduces to the case of *RN* for $\gamma = 1$. Any divergence in the electromagnetic field can easily be identified from this invariant, which is highly dependent on γ .

It is interesting to see that although the em invariant diverges at $r = 0$; when $\gamma = 1$, it becomes regular for $\gamma \neq 1$. To see this, we express K in the form $K = r^{-\gamma}K_0$ and upon substituting $r = 0$, the invariant becomes $I = \frac{2C_0^2}{K_0^4}(m^2q^2)^{-(\gamma-1)^2}(m^2q^2 + m^2p^2\sin^2\theta)^{\gamma^2-1} < \infty$ since $K_0 \neq 0$. We also add that from the Einstein equations $R_{\mu\nu} = -T_{\mu\nu} = \frac{1}{4}g_{\mu\nu}F_{\alpha\beta}F^{\alpha\beta} - F_{\mu\alpha}F_{\nu}^{\alpha}$, the constant C_0 can be fixed as $C_0^2 = 8m^2\gamma^2p^2q^2$.

2. Pure electric case

In this case, we choose the vector potential as

$$A_{\mu} = (f(r), 0, 0, 0) \quad (27)$$

where $f(r)$ is a function of r which is to be determined from the satisfaction of the Maxwell equation $\nabla_r F^{rt} = 0$. Up to a trivial additive constant, we obtain

$$f(r) = C_1 \int^r \frac{\Delta^{\gamma-1} dr}{r^2 K^2} \quad (28)$$

where C_1 is an integration constant proportional to q . It is seen that finding the exact form of $f(r)$ depends entirely on the γ - term and this expression is rather complicated in general. By choosing the spherical symmetric case $\gamma = 1$ one easily observes that the potential of the pure electric RN solution is recovered.

The invariant of the field is computed in this case as

$$I = F_{\mu\nu}F^{\mu\nu} = -\frac{2C_1^2}{r^4 K^4} \Delta^{-(\gamma-1)^2} \Sigma^{\gamma^2-1} \quad (29)$$

which is in conform (up to an expected sign change) with the magnetic invariant. Just as the magnetic case, the constant C_1 is found as $C_1^2 = 8m^2\gamma^2p^2q^2$.

C. Singularity Detection

Our metric contains singularities. In order to see this we must compute the Kretschmann scalar $R_{\mu\nu\alpha\beta}R^{\mu\nu\alpha\beta}$, however, due to the technical difficulties we follow an alternative way to investigate the Newman-Penrose (NP) component [15] Ψ_2 of null-tetrad formalism. In general, the singularities of Ψ_2 are indicative to the singularities of the overall spacetime. It

will be seen that even this will not be an easy task at all. For this purpose, we make the choice of null-tetrad basis 1– forms

$$\begin{aligned}
\sqrt{2}l &= A(r)dt - C(r)\sin\theta d\varphi = \sqrt{2}l_\mu dx^\mu \\
\sqrt{2}n &= A(r)dt + C(r)\sin\theta d\varphi = \sqrt{2}n_\mu dx^\mu \\
\sqrt{2}m &= B(r, \theta) \left(\frac{dr}{\Delta} + i r d\theta \right) = \sqrt{2}m_\mu dx^\mu \\
\sqrt{2}\bar{m} &= B(r, \theta) \left(\frac{dr}{\Delta} - i r d\theta \right) = \sqrt{2}\bar{m}_\mu dx^\mu
\end{aligned} \tag{30}$$

where $A(r)$, $B(r, \theta)$ and $C(r)$ are abbreviated as

$$\begin{aligned}
A(r) &= \Delta^{\gamma/2} K^{-1} \\
C(r) &= K \Delta^{\frac{1-\gamma}{2}} \\
B(r, \theta) &= K \Delta^{\frac{\gamma^2-\gamma}{2}} \Sigma^{\frac{1-\gamma^2}{2}}
\end{aligned} \tag{31}$$

In the null-tetrad $(l_\mu, n_\mu, m_\mu, \bar{m}_\mu)$ of NP we find the following expression for Ψ_2

$$\begin{aligned}
\Psi_2(r, \theta) &= \frac{\Delta^{1+\gamma-\gamma^2}}{4K^2} \Sigma^{\gamma^2-1} \left\{ \frac{\Delta'}{2r\Delta} - 3 \left(\frac{K'}{K} \right)^2 + \frac{1}{4} \gamma(\gamma-1)(\gamma-\gamma^2-2) \left(\frac{\Delta'}{\Delta} \right)^2 \right. \\
&\quad + \frac{1}{4} (1-\gamma^2)(\gamma^2-2) \left(\frac{\Sigma'}{\Sigma} \right)^2 + \gamma(2-\gamma) \frac{K' \Delta'}{K \Delta} + (\gamma^2-1) \frac{K' \Sigma'}{K \Sigma} \\
&\quad + \frac{1}{4} (1-\gamma^2)(1+2\gamma-2\gamma^2) \frac{\Sigma' \Delta'}{\Sigma \Delta} - \frac{2K'}{rK} + \frac{\gamma^2 \Delta'}{2r\Delta} + \frac{K''}{K} + \frac{1}{2} \gamma(\gamma-1) \frac{\Delta''}{\Delta} \\
&\quad \left. \frac{1}{2} (1-\gamma^2) \frac{\Sigma''}{\Sigma} + \frac{(1-\gamma^2)p^2 m^2}{r^4 \Sigma^2} \left(\cos 2\theta - \frac{p^2 m^2}{\Delta r^2} \sin^2 \theta \right) \right\}
\end{aligned} \tag{32}$$

in which prime denotes derivative with respect to r of the functions given in (18).

We observed that further substitution of the derivatives will add little other than the already wild expression for the Weyl curvature. Even the study of this expression for $\Psi_2(r, \theta)$ reveals the occurrence of singularities at $r = 0$, and the roots of $\Delta(r) = 0$ and $\Sigma(r, \theta) = 0$. It is interesting to observe that the outermost singularity is due to the root of $\Delta(r) = 0$, which gives

$$r_\Delta = m(1+p). \tag{33}$$

The larger root of $\Sigma(r, \theta) = 0$, gives

$$r_{\Sigma} = m(1 + p\cos\theta). \quad (34)$$

which lies inside of r_{Δ} . Thus, any probe to the singularities of the metric must encounter first with $r_{\Delta} = m(1 + p)$. Another point of interest is to comment that for $r > 0$ and $0 < p < 1$, we have $K(r) \neq 0$, so that no additional singularity arises due to the presence of $K(r)$ in the expression for curvature. We remark that the function $K(r)$ is the metric function that arises when Maxwell field is added to the gravitational ZV - metric. $K(r)$ becomes a constant when the electromagnetic source vanishes. A final observation from $\Psi_2(r, \theta)$ about singularities is that directional singularities occur for $\gamma^2 < 3$, since beyond this interval the power of $\Sigma(r, \theta)$ becomes positive to avoid any divergence.

IV. CONCLUSION

Most planets and stellar objects have magnetic fields. Our Earth has a rather weak field, of the order $B \sim 0.5 G$, which is yet crucial for life on Earth. Most of the other planets also have magnetic fields, stronger or weaker than that of Earth. Gravastars, however, have extremely high fields of the order $\sim 10^{13} G$. When the existence of these magnetic fields is combined with the non-spherical topology of planets/stars, we encounter automatically with the case of the charged ZV metric. This has three parameters, m , q and γ . For $\gamma < 1$, the object is oblate and for it $\gamma > 1$, is prolate. Obviously, $\gamma = 1$ gives the spherically symmetric Reissner-Nordstrom (RN) solution and its uniqueness dictates that any other class of EM solutions must all agree at $\gamma = 1$. The source of our metric can both be pure electric and pure magnetic. Their combination, which contains both electric and magnetic fields, sounds more realistic and it can be considered as a separate study. In the light of the existence of directional singularities, i.e. dependence on the angle θ , the singularity structure can further be investigated by the quantum probes [16]. The integrability or non-integrability of the geodesics, chaotic behaviours [17] and the analysis of the outermost singularity $r_{\Delta} = m(1+p)$ can further be analysed.

Finally, by resorting to the Ernst formalism of the colliding EM spacetime, one can charge the stationary version [18] of the ZV metric. The class of ZV - metrics is known to emerge from interaction of aligned, static rods [3]. Our approach goes beyond the finite rods to the infinite plane waves. In that limit due to symmetry, it becomes possible to find

further exact solutions, all by making use of the power of the Ernst formalism.

Acknowledgment:

We extend our sincere gratitude to I. Gullu and S. H. Mazharimousavi for their helpful comments.

-
- [1] D. M. Zipoy, *J. Math. Phys. (N.Y)*, **7**, 1137, (1966).
 - [2] B. H. Voorhees, *Phys. Rev.* , **D 2**, 2119, (1970).
 - [3] F. P. Esposito and L. Witten, *Phys. Lett.* , **58 B**, 357, (1975).
 - [4] J. Horsky and N. V. Mitskievich, *Czech. J. Phys.* , **B 39**, 957, (1989).
 - [5] L. Richterek, J. Novotny and J. Horsky, *Czech. J. Phys.* , **52**, 1021, (2002).
 - [6] P. Bell and P. Szekeres, *Gen. Rel. Grav.* , **5**, 275, (1974).
 - [7] J. B. Griffith, *Colliding Plane Waves in General Relativity* (Oxford University Press, 1991).
 - [8] S. Chandrasekhar and B. C. Xanthopoulos, *Proc. Roy. Soc. Lond.* , **A, 398**, 209, (1985).
 - [9] S. Chandrasekhar and B. C. Xanthopoulos, *Proc. Roy. Soc. Lond.* , **A, 410**, 311, (1987).
 - [10] M. Halilsoy, *Gen. Rel. Grav.* , **25**, 275, (1993).
 - [11] F. J. Ernst, *Phys. Rev.* , **167**, 1175, (1968).
 - [12] F. J. Ernst, *Phys. Rev.* , **168**, 1415, (1968).
 - [13] I. Robinson, *Bull. Acad. Pol. Sci.* , **7**, 351, (1959).
 - [14] B. Bertotti, *Phys. Rev.* , **116**, 1331, (1959).
 - [15] E. Newman and R. Penrose, *J. Math. Phys.*, **3**, 566, (1962).
 - [16] G. T. Horowitz and D. Marolf, *Phys. Rev. D* **52**, 5670 (1995).
 - [17] G. Lukes-Gerakopoulos, *Phys. Rev. D* **86**, 044013, (2012).
 - [18] M. Halilsoy, *J. Math. Phys. (N.Y)*, **33**, 4225, (1992).