

Thermal nature of a generic null surface

Surojit Dalui^{a,*}, Bibhas Ranjan Majhi^{a,†} and T. Padmanabhan^{b,‡}

^a*Department of Physics, Indian Institute of Technology Guwahati, Guwahati 781039, Assam, India.*

^b*IUCAA, Pune University Campus, Ganeshkhind, Pune - 411 007, India.*

(Dated: October 26, 2021)

Dynamical properties of a generic null surface are known to have a thermodynamic interpretation. Such an interpretation is completely based on an analogy between the usual law of thermodynamics and structure of gravitational field equation on the surface. Here we materialise this analogy and show that assigning a temperature on the null surface for a local observer is indeed physically relevant. We find that for a local frame, chosen as outgoing massless chargeless particle (or field mode), perceives a “*local unstable Hamiltonian*” very near to the surface. This is shown both at the field as well as particle descriptions. Due to this it has finite quantum probability to escape through acausal null path which is given by Maxwell-Boltzmann distribution, thereby providing a temperature on the surface.

I. INTRODUCTION AND MOTIVATION

The intimate relationship between gravitational dynamics of the black hole horizon and classical thermodynamics uncovered the fact that black holes possess thermodynamic attributes like entropy [1, 2] and temperature [3–5]. Hawking had shown [3, 4] that the radiating photons are thermal in nature, and the temperature for the corresponding radiating particles was predicted as $T = \hbar\kappa/2\pi$, where κ is the surface gravity of the black hole. Another phenomenon, parallel to Hawking effect, has been predicted theoretically, known as Unruh effect [6]. Such effect can be observed in any local arbitrary gravitational background. To make this perception clearer, we start with the *principle of equivalence* which allows one to construct a local inertial frame around any event in an arbitrary curved spacetime. Given the local inertial frame, one can construct a local Rindler frame, and the observer at rest in the local Rindler frame will perceive a patch of null surface as horizon with a temperature. This result allows us to associate thermodynamical attributions with the null surfaces, which the local Rindler observers perceive as horizons. Such a notable fact leads one to introduce observer-dependent thermodynamic variables [7–9] around any event in spacetime and reinterpret the gravitational field equations near any null surface in the language of thermodynamics [10] or vis versa [11]. The equality between the field equations on the horizon and the thermodynamic identity has been exhibited for a wide class of models like the cases of stationary axisymmetric horizons and evolving spherically symmetric horizons in Einstein gravity [12], static spherically symmetric horizons [13]. In Lanczos-Lovelock gravity sector it has been studied for dynamical apparent

horizons [14] and for generic static horizon [15].

Incidentally, one can provide a thermodynamical interpretation of a gravitational field equation by suitably projecting it on a generic null surface also [16–21]. For instance, Einstein’s equation, contracted with null generator and corresponding auxiliary vector, yields a thermodynamical identity of the form $T\delta_\lambda S = \delta_\lambda E + P\delta_\lambda V$ [16–19], where symbols have their usual meanings and the variation can be interpreted as the change due to virtual displacement of the null surface along an auxiliary null vector, parametrized by the parameter λ (generalization to Lanczos-Lovelock gravity [22] and scalar-tensor theory [23] has been done as well). Therefore, it has been realized that null surfaces which act as one-way membranes to a certain class of observers also possess thermodynamical attributions. This indispensable relation between gravity and thermodynamics led to the idea that *the dynamics of gravity is not fundamental in nature; rather it has emerged from the dynamics of a more fundamental theory*, just like the laws of thermodynamics of a system emerges out from the statistical dynamics of its molecules (see [24] for more details on the concept and aspects of emergent nature of gravity).

However, all these thermodynamical attributions associated with the generic null hypersurface are standing on the platform of complete analogy between the laws of thermodynamics and the structure of the gravitational field equation on the surface. Therefore it is mandatory to provide a clear physical justification in order to call a geometrical quantity on the null surface as a particular thermodynamical entity. To make the concern clear, we point out that Hawking’s calculation [3, 4] on the emission spectrum from the black hole horizon clearly indicates the concept of the temperature of the horizon. Similarly, Unruh effect [6] points out the appearance of temperature on the local Rindler horizon. The lack of such robust justification in the case of generic null surface does not give us complete confidence in assigning temperature or entropy on the surface. In this work, our particular aim is to find out the appearance of temperature without stating any analogical point of view.

In this connection, a recent development related to the

* suroj176121013@iitg.ac.in

† bibhas.majhi@iitg.ac.in

‡ Prof. T. Padmanabhan (Paddy) passed away on 17th September, 2021. The whole idea and calculations were thoroughly discussed among all the authors before his decease, except the final drafting of the manuscript.

thermalization mechanism of the black hole horizon is worth mentioning. It is known that the thermodynamics of a system arises from the underlying statistical theory of microstates. However, in the case of spacetime, which microscopic degrees of freedom is responsible for the emergence of temperature into the system in the presence of the horizon is not known properly. Therefore, the hunt for finding out a concrete explanation for such a feature is still under progress. The general consciousness is - to illuminate the underlying microstructure, it is mandatory to understand the physical mechanism for thermalization of the horizon. In pursuit of achieving to find a unified reason for the origin of horizon thermodynamics, two of the authors have recently found that the thermal nature of the horizon has some connection with instability [25–30] in the near-horizon region. At first glance, these two characteristics may seem to be different to each other, but their unification may become a strong candidate to answer the long-standing question about the origination of horizon thermodynamics. It has been shown through considering a model that in the presence of static spherically symmetric (SSS) black hole [27, 28] or in case of a Kerr black hole [30], an outgoing massless and chargeless particle experiences instability in the near-horizon region. In those works, it was found that a class of observers see a particle following an outgoing null path, is driven by a Hamiltonian xp kind for its near horizon motion [27, 28, 30]. It may be noted that such a Hamiltonian is unstable in nature. Moreover, in the quantum scale, this unstable Hamiltonian provides the temperature into the system, which exactly matches with Hawking’s expression. This noticeable fact leads to a conjecture – “*the local instability in the vicinity region of the horizon acts as the source of the temperature of the system*” (see also [27, 28]). The noticeable feature of the model is it requires mainly the information of a suitable near horizon null path of the particle, which does not crucially consider the underlying symmetry of the spacetime, like the presence of a timelike Killing vector.

Therefore, keeping the above feature in mind, we want to extend the spirit of the aforesaid model in the case of any generic null surface. We will see that it works very well in this case. Let us now summarize the main outcomes and features of our present investigation.

- First, we start with a massless scalar field and, using Klein-Gordon (KG) equation in the field-theoretic approach, we obtain that in the semiclassical limit, the system Hamiltonian for the outgoing mode in the near null hypersurface region comes out to be of xp kind. Also, applying the particle dynamical approach, we obtain the same Hamiltonian for the outgoing particle.
- Using the conventional idea of tunnelling mechanism [31–35] for this Hamiltonian, we obtain that quantum mechanically, the particle can see the null surface as a thermal system. The importance of this observation is that it not only shows an application of tunnelling methodology for a more general

background, but also verifies the fact that the association of temperature on a generic null surface is very much physically acceptable.

- Therefore, our results shed light on the intrinsic reason why temperature is associated with the null surface. The local unstable feature can be responsible for such thermalization.
- Finally, in the present work, unlike the previous ones (SSS and Kerr), we deal with a metric where any intrinsic symmetry, like time translational invariance of spacetime, is absent. Therefore it shows a wide applicability and generality of our developed approach.

The paper is organised as follows: In section II, we first introduce the metric adapted to Gaussian null coordinates (GNC) in the neighbourhood region of any generic null hypersurface and some precursory properties of the line element. Next, in section III we construct the Hamiltonian of the real scalar field mode using Klein-Gordon (KG) equation in the near null hypersurface region. In section IV we construct the Hamiltonian of the outgoing particle but using the Lagrangian formalism this time. The purpose of this formalism is to find out the exact form of the canonical forms of the momenta components for further calculations. In section V we calculate the transverse coordinate average of the Hamiltonian of the system for the implementation of tunneling approach. In the next part in section VI, we apply the tunneling formalism for our system. In the final section VII we conclude.

II. NULL HYPERSURFACES IN GNC COORDINATES

The intent of this paper is to investigate the possible thermodynamics properties of an arbitrary null surface. Therefore, our first objective is to describe the neighbourhood region of that null-hypersurface. A preferable choice of coordinate system exists in order to narrate in this context which is known as Gaussian null coordinates (GNC), in analogy with Gaussian normal coordinates. Usually the Gaussian normal coordinates are constructed by extending the coordinates on a non-null hypersurface to a spacetime neighbourhood using geodesics normal to the surface. However, this construction is not applicable for a null surface because the normal geodesics lies itself on the null surface. Therefore, an uniquely defined auxiliary null geodesics are introduced with some certain conditions in order to construct the Gaussian null coordinates which we shall discuss in the later part of this section. An elaborate description and detailed discussion on the construction of this coordinate system and how metric is constructed in this coordinate system can be found in [36–39]. Here, in order to keep the clarity of our work, we shall briefly describe about some precursory

construction of this coordinate system in a more intuitive way.

The primary objective is to construct a coordinate system around any null surface in any spacetime. Consider a smooth null surface N in a four-dimensional spacetime manifold M , where g_{ab} represents the metric on M . We start with a spacelike 2-surface ζ , on N and the coordinates on ζ are introduced as (x^A) for $A = 1, 2$, i.e. (x^1, x^2) . Now, in order to construct the null surface N using null geodesics, one cannot have this null geodesics lying on the spacelike surface as any vector lies on the spacelike surface has to be spacelike in nature. Therefore, the next coordinate parameter should be introduced in such a way that one must move away from this spacelike surface along any of these null geodesic. Here, we introduce that particular parameter v , not necessarily affine, along the null geodesics increasing in the future direction with the condition $v = 0$ on ζ . Therefore, any point on N in the neighbourhood of ζ can be labelled by the coordinates (v, x^1, x^2) , where (x^1, x^2) corresponds to the label given to the null geodesic passing through that point and v is the chosen parameter at that point. The ‘‘neighbourhood’’ corresponds to that sufficiently small region where the geodesics do not cross each others or do not form any caustics. Let us call the future directed vector field tangent to the null geodesics, $\partial/\partial v$, as \mathbf{l} . Now, our spacetime is four-dimensional, so, it is time for us to introduce the fourth coordinate. With the help of a new set of null geodesics we can construct the coordinate chart in the surrounding of the null-hypersurface. Introducing a unique null vector k^a which is situated at each point on the null surface and satisfying these conditions (i) $k^a k_a = 0$, as it is a null vector; (ii) $l^a k_a = -1$ which suggests that k^a sticks out from each point of the null surface instead of lying on the surface; and (iii) $X_A^a k_a = 0$ where $X_A = \partial/\partial x^A$ are the basis vectors correspond to the coordinates (x^1, x^2) . The choice of the third condition shows that our auxiliary null vector is uniquely defined. As this vector k^a points out off the null surface, it can be used to go off the null surface. The null geodesics emitted from each point on the null surface in the direction of k^a are labelled by the coordinates (v, x^1, x^2) , of that point. We choose a parameter r along this null geodesic and $r = 0$ (we choose again) represents the null surface and $\mathbf{k} = -\partial/\partial r$. Therefore, the chosen affine parameter r can be assigned in the coordinate chart (v, r, x^1, x^2) in the neighbourhood of the null hypersurface means upto the regions where geodesics do not reach to a caustic.

After the introduction of this coordinate system we shall introduce the metric adapted to this coordinate system in the neighbourhood region of any arbitrary null hypersurface. The construction of this metric has been detailed in [37, 38]. However, we shall recall here some of the essential properties of this metric. The line element adapted in this context takes the following form

$$ds^2 = -2r\alpha dv^2 + 2dvdr - 2r\beta_A dv dx^A + \mu_{AB} dx^A dx^B \quad (1)$$

Now, one can see from (1) that there are two null sur-

faces which are at $r = 0$ and at $v = \text{constant}$. Additionally, $r = \text{constant}$ where the constant is non-zero represents the timelike surfaces. Besides, the metric components α , β_A and μ_{AB} are the smooth functions of all the coordinates and μ_{AB} is the transverse $(D - 2)$ -dimensional Riemannian metric on the spacelike surface ζ . We have here a set of null vectors as we mentioned earlier $l^a = (l^v, l^r, l^{x^A}) = (1, 0, \mathbf{0})$ and the unique auxiliary one, i.e. $k^a = (0, -1, \mathbf{0})$. Among these two we can think of l^a as the future-outgoing null vector and k^a as the future-ingoing null vector. The covariant components of these two vectors are $l_a = (-2r\alpha, 1, -r\beta_A)$ and $k_a = (-1, 0, \mathbf{0})$. Therefore, the normal $l_a = \partial_a r$ to the $r = 0$ surface will be a null vector. Moreover this generates the $r = 0$ null surface around which we will do our all analysis.

In order to strengthen the motivation for choosing our desired null hypersurface at $r = 0$, let us first write the static spherically symmetric (SSS) metric in analogous to the GNC form [17]

$$ds^2 = -f(r)dv^2 + 2dvdr + \mu_{AB} dx^A dx^B. \quad (2)$$

One can see $\beta_A = 0$ in this case comparing it with Eq. (1). Now, in the near horizon region we have $f(r) \simeq 2\kappa(r - r_H)$ where $\kappa = f'(r_H)/2$ where the prime denotes the derivative with respect to r . Therefore, comparing (2) with Eq. (1) we obtain $\alpha \equiv \kappa$ and the position of the horizon $r = r_H$ is equivalent to the position of the null surface at $r = 0$ in case of GNC metric (1). Therefore, in this context the null surface at $r = 0$ is reminiscent to the black hole horizon. In our earlier works [27, 28] we studied the particle dynamics near the horizon and obtained that particle trajectory becomes unstable near it. Following that essence our objective in this work is to extend this idea in case of any generic null hypersurface and thereby we particularly choose the null surface which is at $r = 0$. The null normal l_a which is at $r = 0$ is given by $l_a = (0, 1, \mathbf{0})$ is the generator of the null surface providing $l^a l_a = 0$ at $r = 0$. Therefore, for our examination in the near null hypersurface region we shall consider the limit $r \rightarrow 0$ whenever necessary.

In the present context our prime aim is to explain how temperature is physically associated with any generic null hypersurface. In order to do so, the first objective is to find out the emission probability of a particle through the null hypersurface. So, we adopt one of the familiar techniques known as the tunneling method [31–35]. The tunneling formalism predicts the probability of escaping the particle through the null hypersurface. Therefore, to evaluate the tunneling probability first we need to find out the path of the particle for both ingoing and the outgoing cases. Hence, we acquire the Hamiltonian formalism in this case just like our earlier works [27, 28, 30]. We shall calculate the Hamiltonian in the near null hypersurface region for two cases. The first one is for the field modes represented by the massless real scalar fields. The second one using the particle description method. For both the cases the intend is to find out the Hamilto-

nian in the near null hypersurface region for both ingoing and the outgoing cases and proceed for the evaluation of the tunneling probability through the null surface.

III. HAMILTONIAN: FIELD DESCRIPTION

Considering the massless real scalar field ϕ , from the Klein-Gordon (KG) equation $\square\phi = 0$ under the background of metric (1) yields

$$\begin{aligned} & \partial_v(\sqrt{\mu}\partial_r\phi) + \partial_r(\sqrt{\mu}\partial_v\phi) + \partial_r[\sqrt{\mu}(2r\alpha + r^2\beta^2)\partial_r\phi] \\ & + \partial_r(\sqrt{\mu}r\beta^A\partial_A\phi) + \partial_A(\sqrt{\mu}r\beta^A\partial_r\phi) \\ & + \partial_A(\sqrt{\mu}\mu^{AB}\partial_B\phi) = 0, \end{aligned} \quad (3)$$

where μ is the determinant of the induced metric μ_{AB} . Now, we start with the standard ansatz for the scalar field of a particle as (see also [31])

$$\phi = \mathcal{A}(v, r, x^A)e^{-\frac{i}{\hbar}S(v, r, x^A)}, \quad (4)$$

where $S(v, r, x^A)$ is the Hamilton-Jacobi action and with respect to the Hamilton-Jacobi action we define the four-momentum as

$$\frac{\partial S}{\partial x^a} = p_a. \quad (5)$$

Now, expanding $S(v, r, x^A)$ in the powers of \hbar we find,

$$\begin{aligned} S(v, r, x^A) &= S_0(v, r, x^A) + \hbar S_1(v, r, x^A) + \hbar^2 S_2(v, r, x^A) \\ &+ \dots \\ &= S_0(v, r, x^A) + \sum_i \hbar^i S_i(v, r, x^A), \end{aligned} \quad (6)$$

where $i = 1, 2, 3, \dots$. The terms from $\mathcal{O}(\hbar)$ onward are treated as the quantum corrections over the semi-classical value S_0 . However, our analysis is restricted only upto the semi-classical limit, i.e. $\hbar \rightarrow 0$. Therefore, the higher order terms of \hbar can be neglected in the semi-classical limit. At this point, we define $-\partial S_0/\partial v = -p_v = H$, where H is the (semi-classical) Hamiltonian of the system.

Now, the main interest lies in the near $r \rightarrow 0$ region because that is where the dynamics of our massless scalar modes will be studied. The probability of crossing the mode across the null surface will be our main quantity to find. One of the well-known techniques to study such quantity in this region is tunneling formalism. One of the principal ways of implementing the tunneling formalism is the Hamilton-Jacobi method [31]. To implement the inherent idea of tunneling mechanism first, we need to identify the ingoing and the outgoing modes near the null hypersurface region. Here we define the mode moving from $r < 0$ region to $r > 0$ as the outgoing one and that moving from $r > 0$ region to $r < 0$ is the ingoing one. Now, applying $\hat{p} \equiv -i\hbar\partial/\partial r$ on (4) we shall have the momentum eigenvalue as $-\partial S_0/\partial r$ in the semi-classical limit. For the outgoing case we shall have the positive momentum eigenvalue, which means one must have $\partial S_0/\partial r < 0$. Similarly, for the ingoing case we shall have the negative momentum eigenvalue,

i.e. $\partial S_0/\partial r > 0$. After this identification, we need to calculate both the Hamilton-Jacobi actions (outgoing and ingoing) and implement these expressions to calculate the tunneling probability for the particles to cross the null hypersurface.

The tunneling formalism mainly based on the semi-classical approximation and in order to implement it in the present setting we shall consider the limit $\hbar \rightarrow 0$ which we have already mentioned before. In this limit, keeping only the leading order terms, we obtain the following form of the Eq. (3):

$$\begin{aligned} & 2(\partial_v S_0)(\partial_r S_0) + (2r\alpha + r^2\beta^2)(\partial_r S_0)^2 + 2r\beta^A(\partial_A S_0)(\partial_r S_0) \\ & + \mu^{AB}(\partial_B S_0)(\partial_A S_0) = 0. \end{aligned} \quad (7)$$

As we can see from Eq. (7) that $\partial_r S_0$ has two solutions which are

$$\begin{aligned} \partial_r S_0 = & -\frac{\partial_v S_0 + r\beta^A(\partial_A S_0)}{2r\alpha + r^2\beta^2} \pm \left[\left(\frac{\partial_v S_0 + r\beta^A(\partial_A S_0)}{2r\alpha + r^2\beta^2} \right)^2 \right. \\ & \left. - \frac{\mu^{AB}(\partial_A S_0)(\partial_B S_0)}{2r\alpha + r^2\beta^2} \right]^{\frac{1}{2}}. \end{aligned} \quad (8)$$

Among these two solutions one solution corresponds to the outgoing mode and the other one corresponds to the ingoing one. Now, let us get going to identify them.

First we need to find out the leading order solutions of $\partial_r S_0$ in the near null hypersurface region, as we are interested to examine the physics near the null hypersurface which is at $r = 0$. Now, considering the negative sign solution of Eq. (8) first, we obtain the leading order term in $r \rightarrow 0$ limit as (for details please see Appendix A)

$$\partial_r S_0 \Big|_- = -\frac{\partial_v S_0}{\alpha^{(0)}(v, x^A)r}. \quad (9)$$

According to our definition of the Hamiltonian of the system (which we have defined earlier below Eq. (6)) we can write Eq. (9) as

$$\partial_r S_0 \Big|_- = \frac{H}{\alpha^{(0)}(v, x^A)r}. \quad (10)$$

Next, if we consider the initial position is at $r < 0$, then for $H > 0$, $\partial_r S_0 < 0$. Therefore, according to our earlier definition of identifying the directions of the modes, the momentum direction of the mode in this case is in the outward direction, i.e. from $r < 0$ to $r > 0$ region. Hence, the negative sign solution of Eq. (8) corresponds to the outgoing mode. So, we can write the Hamiltonian for the outgoing mode in the near null hypersurface region ($r \rightarrow 0$) as

$$H = \alpha^{(0)}(v, x^A)r p_{r_{out}} \quad (11)$$

where $p_{r_{out}}$ is the momentum in r direction.

Similarly, considering the positive sign solution of Eq. (8) we obtain the leading order term that survives in $r \rightarrow 0$ limit is

$$\partial_r S_0 \Big|_+ = -\frac{1}{2} \frac{\mu^{(0)AB} (\partial_A S_0) (\partial_B S_0)}{\partial_v S_0}, \quad (12)$$

where $\mu^{(0)AB}$ is the first term of the expansion of μ^{AB} about $r = 0$. Therefore, in terms of the Hamiltonian, we can write the above equation as

$$\partial_r S_0 \Big|_+ = \frac{1}{2} \frac{\mu^{(0)AB} (\partial_A S_0) (\partial_B S_0)}{H}. \quad (13)$$

Now, from Eq. (13) we can see for $H > 0$, $\partial_r S_0 > 0$. Therefore, with the similar argument as earlier, the momentum direction of the mode in this case is in the inward direction, i.e. from $r > 0$ to $r < 0$. So, the positive sign solution of $\partial_r S_0$ corresponds to the ingoing mode and the expression of Hamiltonian in this case is

$$H = \frac{1}{2} \frac{\mu^{(0)AB} p_A p_B}{p_{r_{in}}}, \quad (14)$$

where $p_{r_{in}}$ is the momentum in r direction for this ingoing mode.

In the next part we shall show that the form of the outgoing Hamiltonian (Eq. (11)) can also be achieved if one follows the Lagrangian formalism of a massless particle system.

IV. HAMILTONIAN: PARTICLE DESCRIPTION IN LAGRANGIAN FORMALISM

Before proceeding for the thermalization analysis, let us look for a moment for the expression of the Hamiltonian obtained using the Lagrangian of the system. In the previous section using Hamilton-Jacobi formalism we land up to a particular Hamiltonian structure (Eq. (11)) of the outgoing scalar mode in the near null hypersurface region. In the present section we like to find out whether the same Hamiltonian structure can be obtained using the Lagrangian of a particle. Further, we would also like to find out the expressions of the canonical momenta and their dependence on the coordinate variables for further calculations.

Consider the Lagrangian $L = \sqrt{-g_{ab} \dot{x}^a \dot{x}^b}$ where $\dot{x}^a = dx^a/dv$. Since we are considering massless particle, for convenience v has been chosen here as the affine parameter for the geodesics of the particle. Therefore, under the background of metric (1) we obtain the form of the Lagrangian of the system as

$$L = [2r\alpha - 2\dot{r} + 2r\beta_A \dot{x}^A - \mu_{AB} \dot{x}^A \dot{x}^B]^{\frac{1}{2}} \quad (15)$$

where the expressions of the corresponding momentum components are

$$p_r = -\frac{1}{[2r\alpha - 2\dot{r} + 2r\beta_A \dot{x}^A - \mu_{AB} \dot{x}^A \dot{x}^B]^{\frac{1}{2}}}; \quad (16)$$

$$p_A = \frac{r\beta_A - \mu_{AB} \dot{x}^B}{[2r\alpha - 2\dot{r} + 2r\beta_A \dot{x}^A - \mu_{AB} \dot{x}^A \dot{x}^B]^{\frac{1}{2}}}. \quad (17)$$

Therefore, we obtain the Hamiltonian of the system as

$$H = \frac{1}{2p_r} [(2r\alpha + r^2\beta^2)p_r^2 + 2r\beta^A p_A p_r + (1 + p_A^2)] \quad (18)$$

The above expression of the Hamiltonian (18) reveals that there are two solutions of p_r in terms of H and p_A . It is evident that one solution of p_r corresponds to the outgoing particle while the other one corresponds to the ingoing one. Since our interest lies in the near null hypersurface region, our next aim is to find out the expressions of p_r for both the outgoing and the ingoing one in the limit $r \rightarrow 0$. Therefore, implementing the limit $r \rightarrow 0$ and considering the leading order terms in these two solutions of p_r we obtain

$$p_r \Big|_- = \frac{H}{\alpha^{(0)}(v, x^A)r} \quad \text{and} \quad p_r \Big|_+ = \frac{1}{2} \left(\frac{1 + p_A^2}{H} \right) \quad (19)$$

where $(-)$ and $(+)$ sign represents the -ve and the +ve sign solutions of p_r of the quadratic equation (18) respectively. Therefore, we can see that the expression of the -ve sign solution of p_r exactly matches with Eq. (10) which we identified as the momentum in r direction for the outgoing particle, i.e. $p_{r_{out}}$. So, it is evident that the -ve sign solution of p_r corresponds to the momentum of the outgoing particle and we obtain the similar structure of the outgoing Hamiltonian in the near null hypersurface region (see Eq. (11)).

So, from both the analyses, i.e. studying the field mode description and the particle description we obtain that the outgoing Hamiltonian in the near null hypersurface region has the similar expression for both the cases (see Eq. (11)). From the expression of the outgoing Hamiltonian (11) one can see that the outgoing particle suffers a singularity at $r = 0$. Whereas for the ingoing particle the Hamiltonian structure in the near null hypersurface region may differ in those two descriptions (see (14) and Eq. (19)) but their natures are same as the ingoing particle does not suffer any singularity at $r = 0$. This interesting observation has very important implication in the calculation of the tunneling probability which we shall see in the later part of our calculations.

V. TRANSVERSE COORDINATE AVERAGE OF THE HAMILTONIAN

In the previous two sections applying two formalism, i.e. Hamilton-Jacobi and Lagrangian formalism we obtained the Hamiltonian for the outgoing object Eq. (11). Our next aim is to know the consequences of this classical Hamiltonian in the quantum picture. In order to gain knowledge about that, here we are about to implement the tunneling formalism in this case which can tell us about the tunneling probability of the object through the null hypersurface. It may be worth to point out here that in earlier calculations [27, 28, 30] the spacetime metric was static or stationary and hence $\alpha^{(0)}$ was constant. But this is not the case here.

However before proceeding for implementing the tunneling formalism let us prepare the stage for implement-

ing it on our Hamiltonian. The structure of the Hamiltonian for the outgoing particle in the near null hypersurface region is multidimensional in this case due to the presence of $\alpha^{(0)}(v, x^A)$. Therefore, we have reached to a situation where we have to deal with the case of a multidimensional tunneling. Multidimensional tunnelling event has been discussed in [40] for the usual physical systems. One of the proposals to calculate tunnelling probability is to do calculation on an average potential by considering averaging over directions except one. Here we are interested to motion of our object radially just across the null surface and hence following the idea of [40] we read the transverse coordinates average of the Hamiltonian: $\bar{H} = \int H \sqrt{\mu} d^2 x^A / \int \sqrt{\mu} d^2 x^A$. This yields

$$\bar{H} = \bar{\alpha}(v) r p_{r_{out}} \equiv \bar{E}, \quad (20)$$

where $\bar{\alpha}(v)$ is defined as

$$\bar{\alpha}(v) = \frac{\int \alpha^{(0)}(v, x^A) \sqrt{\mu} d^2 x^A}{\int \sqrt{\mu} d^2 x^A}. \quad (21)$$

Here one thing is to be mentioned that during the consideration of the average of the Hamiltonian H , the integrating average applied only on $\alpha^{(0)}(v, x^A)$ because both r and $p_{r_{out}}$ are independent of x^A in a Hamiltonian. We shall use this average Hamiltonian (20) in the next section in order to investigate the thermalization of our null surface through tunneling formalism.

VI. TUNNELING AND THERMALITY

We start by calculating the Hamilton-Jacobi action for the outgoing particle. In this case, the outgoing particle is going from $r < 0$ to $r > 0$ by crossing the $r = 0$ region. Hence, choosing the integration limit from $r = -\epsilon$ to $r = \epsilon$ where $\epsilon > 0$ and is a very small number, suggests that the outgoing particle crosses the null hypersurface from just inside to just outside in the vicinity of the null hypersurface. Therefore, we obtain the Hamilton-Jacobi action for the outgoing particle

$$S_{out} = \frac{\bar{E}}{\bar{\alpha}(v)} \int_{-\epsilon}^{\epsilon} \frac{dr}{r} + \int p_{v_{out}} dv + \int p_{A_{out}} dx^A. \quad (22)$$

From the above integration it can be seen that the first integration term will contain the imaginary part as there exists a singularity at $r = 0$ and other two integrations will contribute in the real part of the total integration. Also in the first integration we have pulled out the term $\bar{E}/\bar{\alpha}(v)$ as it is constant of motion (please see Appendix B). Hence, after performing the integration in Eq. (22) we obtain

$$S_{out} = -\frac{i\pi\bar{E}}{\bar{\alpha}(v)} + \text{Real part}. \quad (23)$$

In a similar way, we can calculate the Hamilton-Jacobi action for the ingoing particle. However, in this case, the action does not contain any singularity at $r = 0$ (see Eq. (14) and Eq. (19)); thus, it turns out to be

$$S_{in} = \text{Real quantity}. \quad (24)$$

Accordingly, the probability for the outgoing particle crossing the null hypersurface turns out to be

$$P_{out} \sim \left| e^{-\frac{i}{\hbar} S_{out}} \right|^2 \propto \exp\left(-\frac{2\pi\bar{E}}{\hbar\bar{\alpha}(v)}\right), \quad (25)$$

whereas the probability of crossing the null hypersurface for the ingoing particle is $P_{in} \sim 1$. Therefore, the tunneling probability comes out to be

$$\Gamma(v) = \frac{P_{out}}{P_{in}} \sim \exp\left(-\frac{2\pi\bar{E}}{\hbar\bar{\alpha}(v)}\right). \quad (26)$$

This particular expression of the tunneling probability is similar to Boltzmann factor and therefore can be considered as thermal in nature. Then, we can identify the temperature of the system as

$$T(v) = \frac{\hbar\bar{\alpha}(v)}{2\pi}. \quad (27)$$

However, one important thing to notice is that this very expression of the temperature is not a constant; rather, it is a function of v . It means at every other $v = \text{constant}$ null hypersurface near the $r = 0$ region, the observer will feel different values of temperature of the system for every different values of v .

VII. DISCUSSION

Let us summarise the results obtained in the present work. We started this work by addressing the fact that gravitational field equations near any null surface in an arbitrary space-time reduces to a thermodynamic identity, and it generalises the results previously available in the context of the horizon. Our prime motive was to find out the underlying reason for this noticeable fact in order to convey the cause why thermodynamical attributions are associated with any arbitrary null hypersurface. We started our calculations with a probed massless and chargeless particle and tried to construct the Hamiltonian of the system in the near null hypersurface region. First, we attempted using the KG equation in the field-theoretic approach and in the semi-classical limit, we obtain that the system Hamiltonian for the outgoing mode in the near null hypersurface region comes out to be of xp kind. The same has been explored in the Lagrangian formalism for a massless outgoing particle as well. In the context of thermality we proceed with the conventional idea of tunneling mechanism, and after implementing the tunneling formalism in the near null hypersurface region, we obtain that our system is thermal in nature. However, the system temperature we found, in this case, is not constant; rather it is a function of the timelike coordinate, unlike the previous results of the black hole horizons (SSS BH and the Kerr one).

To summarise, our results justify the fact how temperature is associated with any generic null hypersurface. Some earlier works predicted that the emergence of

thermality into the system has close connection with the *local* instability of the system in the context of horizon [26–28, 30]. This connection previously showed that if the Hamiltonian of the system turns out to be an unstable one in the classical scale, this instability may lead to the thermality of the system in the quantum scale [27, 28, 30]. Here, we came across the Hamiltonian which consists of a probed massless and chargeless object near any generic null hypersurface and the structure of the outgoing Hamiltonian in this case turns out to be of xp kind (see Eq. (11)). This specific Hamiltonian is unstable in nature which is apparent from the very structure of the Hamiltonian. The importance of this unstable nature of the Hamiltonian is visible when we turn our attention towards the calculation of the tunneling effect in the quantum regime. The tunneling probability of the outgoing particle for crossing the null surface comes out to be of Boltzmann factor which shows that the system is thermal in nature in the quantum scale. Therefore, this result clearly represents the fact that the instability in

the near null hypersurface region provides the the automatic emergence of thermal fluctuations into the system. Our results in this work also represents one of the important applications of tunneling mechanism for a more general background. One of the important aspects of our methodology is to show how tunneling formalism can be implemented for the time dependent background.

Finally, our methodology clearly displays how thermality is associated with any generic null hypersurface without the analogical point of view. Our results also show that the local instability plays crucial role for the emergence of the temperature into the system. Previously, this connection between instability and thermality was established only in specific cases containing horizons. But the results we obtained in this paper simply breaks the illusion when we generalise this connection for any arbitrary null hypersurface in any space-time. Therefore, we feel that the present discussion may unfold the deeper reason for having the thermodynamical quantities of not only horizon but also for any generic null surface at the quantum level.

APPENDICES

Appendix A: Derivation of Eq. (9) and Eq. (12)

In the near null hypersurface region, i.e. $r \rightarrow 0$ limit, $\alpha(v, r, x^A)$ can be expanded (using Taylor series expansion)

$$\alpha(v, r, x^A) = \alpha^{(0)}(v, x^A) + \alpha^{(1)}(v, x^A)r + \mathcal{O}(r^2). \quad (\text{A1})$$

Now, looking back to Eq. (8) we can rewrite it as

$$\partial_r S = -\frac{\partial_v S + r\beta^A(\partial_A S)}{2r\alpha + r^2\beta^2} \pm \left(\frac{\partial_v S + r\beta^A(\partial_A S)}{2r\alpha + r^2\beta^2} \right) \left[1 - \frac{\mu^{AB}(\partial_A S)(\partial_B S)}{(\partial_v S + r\beta^A(\partial_A S))^2} (2r\alpha + r^2\beta^2) \right]^{\frac{1}{2}}. \quad (\text{A2})$$

Now, at $r \rightarrow 0$ the above equation turns into

$$\begin{aligned} \partial_r S &\simeq -\frac{\partial_v S + r\beta^A(\partial_A S)}{2r\alpha + r^2\beta^2} \pm \left(\frac{\partial_v S + r\beta^A(\partial_A S)}{2r\alpha + r^2\beta^2} \right) \left[1 - \frac{1}{2} \frac{\mu^{AB}(\partial_A S)(\partial_B S)}{(\partial_v S + r\beta^A(\partial_A S))^2} (2r\alpha + r^2\beta^2) \right] \\ &\simeq -\frac{\partial_v S + r\beta^A(\partial_A S)}{2r\alpha + r^2\beta^2} \pm \left[\left(\frac{\partial_v S + r\beta^A(\partial_A S)}{2r\alpha + r^2\beta^2} \right) - \frac{1}{2} \frac{\mu^{AB}(\partial_A S)(\partial_B S)}{(\partial_v S + r\beta^A(\partial_A S))} \right]. \end{aligned} \quad (\text{A3})$$

Considering the negative sign solution of $\partial_r S$ we obtain from Eq. (A3)

$$\partial_r S \Big|_- = -2 \frac{\partial_v S + r\beta^A(\partial_A S)}{2r\alpha + r^2\beta^2} + \frac{1}{2} \frac{\mu^{AB}(\partial_A S)(\partial_B S)}{(\partial_v S + r\beta^A(\partial_A S))}. \quad (\text{A4})$$

The first term of Eq. (A4), using the expansion of $\alpha(v, r, x^A)$, in the near null hypersurface region reduces to

$$\begin{aligned} \frac{\partial_v S + r\beta^A(\partial_A S)}{2r\alpha + r^2\beta^2} &\simeq \frac{\partial_v S + r\beta^A(\partial_A S)}{2r\alpha^{(0)} \left(1 + \frac{r\alpha^{(1)}}{\alpha^{(0)}} + \frac{r\beta^2}{2\alpha^{(0)}} \right)} \\ &= \frac{\partial_v S + r\beta^A(\partial_A S)}{2r\alpha^{(0)}} \left(1 - \frac{r\alpha^{(1)}}{\alpha^{(0)}} - \frac{r\beta^2}{2\alpha^{(0)}} \right). \end{aligned} \quad (\text{A5})$$

Similarly, from the second term of Eq. (A4), in the near null hypersurface region we obtain

$$\frac{\mu^{AB}(\partial_A S)(\partial_B S)}{(\partial_v S + r\beta^A(\partial_A S))} \simeq \frac{\mu^{AB}(\partial_A S)(\partial_B S)}{\partial_v S} \left(1 - \frac{r\beta^A(\partial_A S)}{\partial_v S} \right). \quad (\text{A6})$$

Hence, putting the approximated values of these two terms in Eq. (A4) we obtain the only leading order term in the near null hypersurface region

$$\partial_r S \Big|_{-} = -\frac{\partial_v S}{2r\alpha^{(0)}(v, x^A)}, \quad (\text{A7})$$

i.e. Eq. (9) in our main text. In the similar manner we can also obtain the expression of $\partial_r S|_{+}$. Taking the positive sign solution in Eq. (A3) and considering the leading order term at $r \rightarrow 0$ limit we end up getting Eq. (12).

Appendix B: Conserved quantity $\bar{H}/\bar{\alpha}(v)$

Now, we have the near null hypersurface Hamiltonian for the outgoing particle, i.e.

$$H = \alpha^{(0)}(v, x^A) r p_{r_{out}} \quad (\text{B1})$$

and after averaging out the transverse coordinates we have

$$\bar{H} = \bar{\alpha}(v) r p_{r_{out}}. \quad (\text{B2})$$

Now, let us check the variation of $\bar{H}/\bar{\alpha}(v)$ with respect to some affine parameter λ , i.e.

$$\begin{aligned} \frac{d}{d\lambda} \left(\frac{\bar{H}}{\bar{\alpha}(v)} \right) &= \frac{d}{d\lambda} (r p_{r_{out}}) \\ &= \dot{r} p_{r_{out}} + r \dot{p}_{r_{out}} \end{aligned} \quad (\text{B3})$$

where $\dot{} \equiv \frac{d}{d\lambda}$. Now, from the Hamilton's equations of motion we obtain

$$\dot{r} = \frac{\partial \bar{H}}{\partial p_{r_{out}}} = \bar{\alpha}(v) r \quad (\text{B4})$$

and the other one is

$$\dot{p}_{r_{out}} = \frac{\partial \bar{H}}{\partial r} = -\bar{\alpha}(v) p_{r_{out}}. \quad (\text{B5})$$

Now, putting the values of \dot{r} and $\dot{p}_{r_{out}}$ in Eq. (B3) we obtain

$$\begin{aligned} \frac{d}{d\lambda} \left(\frac{\bar{H}}{\bar{\alpha}(v)} \right) &= \bar{\alpha}(v) r p_{r_{out}} + (-\bar{\alpha}(v) r p_{r_{out}}) \\ &= 0. \end{aligned} \quad (\text{B6})$$

It tells that the quantity $\bar{H}/\bar{\alpha}(v) = \bar{E}/\bar{\alpha}$ is conserved during the motion of the particle under the average Hamiltonian \bar{H} .

- [1] J. D. Bekenstein, "Black holes and entropy," *Phys. Rev. D* **7** (1973), 2333-2346
- [2] J. D. Bekenstein, "Generalized second law of thermodynamics in black hole physics," *Phys. Rev. D* **9** (1974), 3292-3300
- [3] S. W. Hawking, "Black hole explosions," *Nature* **248**, 30 (1974).
- [4] S. W. Hawking, "Particle Creation by Black Holes," *Commun. Math. Phys.* **43** (1975), 199-220
- [5] P. C. W. Davies, S. A. Fulling and W. G. Unruh, "Energy Momentum Tensor Near an Evaporating Black Hole," *Phys. Rev. D* **13** (1976), 2720-2723
- [6] W. G. Unruh, "Notes on black hole evaporation," *Phys. Rev. D* **14**, 870 (1976).
- [7] B. R. Majhi and T. Padmanabhan, "Noether current from the surface term of gravitational action, Virasoro algebra and horizon entropy," *Phys. Rev. D* **86**, 101501 (2012) [[arXiv:1204.1422](https://arxiv.org/abs/1204.1422) [gr-qc]].
- [8] B. R. Majhi, "Noether current of the surface term of Einstein-Hilbert action, Virasoro algebra and entropy," *Adv. High Energy Phys.* **2013**, 386342 (2013) [[arXiv:1210.6736](https://arxiv.org/abs/1210.6736) [gr-qc]].
- [9] B. R. Majhi, "Noncommutativity in near horizon symmetries in gravity," *Phys. Rev. D* **95**, no.4, 044020 (2017)

- [arXiv:1701.07952 [gr-qc]].
- [10] T. Padmanabhan, “A Dialogue on the Nature of Gravity,” [arXiv:0910.0839 [gr-qc]].
- [11] T. Jacobson, “Thermodynamics of space-time: The Einstein equation of state,” *Phys. Rev. Lett.* **75** (1995), 1260-1263 [arXiv:gr-qc/9504004 [gr-qc]].
- [12] D. Kothawala, S. Sarkar and T. Padmanabhan, “Einstein’s equations as a thermodynamic identity: The Cases of stationary axisymmetric horizons and evolving spherically symmetric horizons,” *Phys. Lett. B* **652** (2007), 338-342 [arXiv:gr-qc/0701002 [gr-qc]].
- [13] A. Paranjape, S. Sarkar and T. Padmanabhan, “Thermodynamic route to field equations in Lancos-Lovelock gravity,” *Phys. Rev. D* **74** (2006), 104015 [arXiv:hep-th/0607240 [hep-th]].
- [14] R. G. Cai, L. M. Cao, Y. P. Hu and S. P. Kim, “Generalized Vaidya Spacetime in Lovelock Gravity and Thermodynamics on Apparent Horizon,” *Phys. Rev. D* **78** (2008), 124012 [arXiv:0810.2610 [hep-th]].
- [15] D. Kothawala and T. Padmanabhan, “Thermodynamic structure of Lanczos-Lovelock field equations from near-horizon symmetries,” *Phys. Rev. D* **79** (2009), 104020 [arXiv:0904.0215 [gr-qc]].
- [16] K. Parattu, B. R. Majhi and T. Padmanabhan, “Structure of the gravitational action and its relation with horizon thermodynamics and emergent gravity paradigm,” *Phys. Rev. D* **87** (2013) no.12, 124011 [arXiv:1303.1535 [gr-qc]].
- [17] S. Chakraborty, K. Parattu and T. Padmanabhan, “Gravitational field equations near an arbitrary null surface expressed as a thermodynamic identity,” *JHEP* **10** (2015), 097 [arXiv:1505.05297 [gr-qc]].
- [18] S. Chakraborty and T. Padmanabhan, “Thermodynamical interpretation of the geometrical variables associated with null surfaces,” *Phys. Rev. D* **92** (2015) no.10, 104011 [arXiv:1508.04060 [gr-qc]].
- [19] S. Dey and B. R. Majhi, “Covariant approach to the thermodynamic structure of a generic null surface,” *Phys. Rev. D* **102** (2020) no.12, 124044 [arXiv:2009.08221 [gr-qc]].
- [20] S. Chakraborty, S. Bhattacharya and T. Padmanabhan, “Entropy of a generic null surface from its associated Virasoro algebra,” *Phys. Lett. B* **763** (2016), 347-351 [arXiv:1605.06988 [gr-qc]].
- [21] K. Bhattacharya and B. R. Majhi, “Noncommutative Heisenberg algebra in the neighbourhood of a generic null surface,” *Nucl. Phys. B* **934** (2018), 557-577 [arXiv:1802.02862 [gr-qc]].
- [22] S. Chakraborty, “Lanczos-Lovelock gravity from a thermodynamic perspective,” *JHEP* **08** (2015), 029 [arXiv:1505.07272 [gr-qc]].
- [23] S. Dey, K. Bhattacharya and B. R. Majhi, “Thermodynamic structure of a generic null surface in scalar-tensor theory and the zeroth law,” [arXiv:2105.07787 [gr-qc]].
- [24] T. Padmanabhan, “Thermodynamical Aspects of Gravity: New insights,” *Rept. Prog. Phys.* **73** (2010), 046901 [arXiv:0911.5004 [gr-qc]].
- [25] M. Srednicki, “Chaos and Quantum Thermalization,” *Phys. Rev. E* **50**, 888 (1994) [arXiv:cond-mat/9403051].
- [26] T. Morita, “Thermal Emission from Semi-classical Dynamical Systems,” *Phys. Rev. Lett.* **122**, no. 10, 101603 (2019) [arXiv:1902.06940 [hep-th]].
- [27] S. Dalui, B. R. Majhi and P. Mishra, “Horizon induces instability locally and creates quantum thermality,” *Phys. Rev. D* **102** (2020) no.4, 044006 [arXiv:1910.07989 [gr-qc]].
- [28] S. Dalui and B. R. Majhi, “Near horizon local instability and quantum thermality,” *Phys. Rev. D* **102** (2020) no.12, 124047 [arXiv:2007.14312 [gr-qc]].
- [29] B. R. Majhi, “Is randomness near a black hole key for thermalization of its horizon?,” [arXiv:2101.04458 [gr-qc]].
- [30] S. Dalui and B. Ranjan Majhi, “Horizon thermalization of Kerr black hole through local instability,” [arXiv:2103.11613 [gr-qc]].
- [31] K. Srinivasan and T. Padmanabhan, “Particle production and complex path analysis,” *Phys. Rev. D* **60**, 024007 (1999) [arXiv:gr-qc/9812028 [gr-qc]].
- [32] M. K. Parikh and F. Wilczek, “Hawking radiation as tunneling,” *Phys. Rev. Lett.* **85**, 5042-5045 (2000) [arXiv:hep-th/9907001 [hep-th]].
- [33] R. Banerjee and B. R. Majhi, “Connecting anomaly and tunneling methods for Hawking effect through chirality,” *Phys. Rev. D* **79**, 064024 (2009) [arXiv:0812.0497 [hep-th]].
- [34] R. Banerjee and B. R. Majhi, “Hawking black body spectrum from tunneling mechanism,” *Phys. Lett. B* **675**, 243 (2009) [arXiv:0903.0250 [hep-th]].
- [35] B. R. Majhi, “Quantum Tunneling in Black Holes,” [arXiv:1110.6008 [gr-qc]].
- [36] V. Moncrief and J. Isenberg, “Symmetries of cosmological Cauchy horizons,” *Commun. Math. Phys.* **89** (1983) no.3, 387-413
- [37] S. Hollands, A. Ishibashi and R. M. Wald, “A Higher dimensional stationary rotating black hole must be axisymmetric,” *Commun. Math. Phys.* **271** (2007), 699-722 [arXiv:gr-qc/0605106 [gr-qc]].
- [38] E. M. Morales, “On a Second Law of Black Hole Mechanics in a Higher Derivative Theory of Gravity” (2008). URL <https://www.theorie.physik.uni-goettingen.de/forschung/qft/theses/dipl/Morfa-Morales.pdf>
- [39] K. Parattu, S. Chakraborty, B. R. Majhi and T. Padmanabhan, “A Boundary Term for the Gravitational Action with Null Boundaries,” *Gen. Rel. Grav.* **48** (2016) no.7, 94 [arXiv:1501.01053 [gr-qc]].
- [40] M. Razavy, “Quantum Theory of Tunneling,” 2nd ed. (World Scientific, Singapore, 2014)