

Local supersymmetry: Variations on a theme by Volkov and Soroka

Sergei M. Kuzenko

*Department of Physics M013, The University of Western Australia
35 Stirling Highway, Perth W.A. 6009, Australia*

Email: `sergei.kuzenko@uwa.edu.au`

Abstract

As is well known, supergravity was discovered in 1976 by Ferrara, Freedman and van Nieuwenhuizen, and independently by Deser and Zumino. In 1994, Volkov, one of the co-creators of supersymmetry, posted two preprints to the hep-th archive containing “supergravity before 1976” in the title. He argued that the key ingredients of $\mathcal{N} = 1$ supergravity in four dimensions had appeared in his earlier work with Soroka on the super-Higgs effect. In order to better appreciate his point of view, in this paper we first revisit the Volkov-Soroka construction and review it. Then we demonstrate that, for specially chosen parameters of the theory, the Volkov-Soroka action is invariant under two different local supersymmetries. One of them is present for arbitrary values of the parameters and acts on the Goldstino, while the other supersymmetry emerges only in a special case and leaves the Goldstino invariant. The former can be used to gauge away the Goldstino, and then the resulting action coincides with that proposed by Deser and Zumino for consistent supergravity. In this sense, pure $\mathcal{N} = 1$ supergravity is a special case of the Volkov-Soroka theory, and it could have been discovered before 1976.

1 Introduction

Dmitry V. Volkov was one of the co-discoverers of supersymmetry. His most prominent results on rigid and local supersymmetry are the Goldstino model (jointly with Akulov) [1,2] and the super-Higgs mechanism (jointly with Soroka) [3,4]. His approach to nonlinear realisations of internal and space-time symmetries [5], which paved the way to [1–4], has also been highly influential. However, it is less known that Ref. [2] also pioneered the following fundamental concepts of modern theoretical physics: (i) the \mathcal{N} -extended super-Poincaré group and, hence, the \mathcal{N} -extended super-Poincaré algebra¹ for $\mathcal{N} > 1$; and (ii) \mathcal{N} -extended Minkowski superspace.² It is quite remarkable that all these results had appeared before the first paper by Wess and Zumino on supersymmetry [7] was published on 18 February, 1974.

As is well known, supergravity was discovered in 1976 [9, 10]. In 1994, less than two years before his death, Volkov posted two preprints to the hep-th archive [11, 12], both of which contained “supergravity before 1976” in the title. He argued that the crucial ingredients of $\mathcal{N} = 1$ supergravity in four dimensions had appeared in [3, 4].³ In order to better appreciate his point of view, in this paper we first provide a detailed review of the Volkov-Soroka construction. Then we demonstrate that for specially chosen parameters the theory possesses two local supersymmetries. One of them can be used to gauge away the Goldstino, and then the resulting action coincides with that proposed by Deser and Zumino to describe consistent supergravity. This shows that pure $\mathcal{N} = 1$ supergravity is a special case of the Volkov-Soroka theory, and it could have been discovered before 1976.

2 A review of the Volkov-Soroka construction

This section is devoted to a pedagogical review of the Volkov-Soroka construction [3, 4].

Let $\mathfrak{P}(4|\mathcal{N})$ denote the four-dimensional \mathcal{N} -extended super-Poincaré group introduced in [2]. Any element $g \in \mathfrak{P}(4|\mathcal{N})$ is a $(4 + \mathcal{N}) \times (4 + \mathcal{N})$ supermatrix of the form⁴

$$g = S(b, \varepsilon) h(M, U) \equiv Sh , \tag{2.1a}$$

¹The $\mathcal{N} = 1$ super-Poincaré algebra was discovered in 1971 by Golfand and Likhtman [6].

²The Akulov-Volkov paper [2] was submitted to the journal *Theoretical and Mathematical Physics* on 8 January 1973 and published in January 1974, before the publication of the first paper on supersymmetry by Wess and Zumino [7] and long before the work by Salam and Strathdee [8] devoted to the $\mathcal{N} = 1$ superspace approach. It remains largely unknown, perhaps because it was published in a Russian journal.

³Similar thoughts were also expressed by Soroka a few years later [13, 14], see also [15, 16].

⁴Our parametrisation of the elements of $\mathfrak{P}(4|\mathcal{N})$ follows [17].

$$S(b, \varepsilon) := \left(\begin{array}{c|c|c} \mathbb{1}_2 & 0 & 0 \\ \hline -i\tilde{b}_{(+)} & \mathbb{1}_2 & 2\epsilon^\dagger \\ \hline 2\epsilon & 0 & \mathbb{1}_{\mathcal{N}} \end{array} \right) = \left(\begin{array}{c|c|c} \delta_{\alpha\beta} & 0 & 0 \\ \hline -i\tilde{b}_{(+)}^{\dot{\alpha}\beta} & \delta_{\dot{\beta}}^{\dot{\alpha}} & 2\bar{\epsilon}^{\dot{\alpha}j} \\ \hline 2\epsilon_i^\beta & 0 & \delta_i^j \end{array} \right), \quad (2.1b)$$

$$h(M, U) := \left(\begin{array}{c|c|c} M & 0 & 0 \\ \hline 0 & (M^{-1})^\dagger & 0 \\ \hline 0 & 0 & U \end{array} \right) = \left(\begin{array}{c|c|c} M_{\alpha}^{\beta} & 0 & 0 \\ \hline 0 & (\bar{M}^{-1})_{\dot{\beta}}^{\dot{\alpha}} & 0 \\ \hline 0 & 0 & U_i^j \end{array} \right), \quad (2.1c)$$

where $M = (M_{\alpha}^{\beta}) \in \mathbf{SL}(2, \mathbb{C})$, $U = (U_i^j) \in \mathbf{U}(\mathcal{N})$, and

$$\tilde{b}_{(\pm)} = \tilde{b} \pm 2i\epsilon^\dagger\epsilon. \quad (2.2)$$

The tilde notation in (2.1b) and (2.2) reflects the fact that there are two types of relativistic Pauli matrices, σ_a and $\tilde{\sigma}_a$, see the appendix. The group element $S(b, \varepsilon)$ is labelled by 4 commuting real parameters b^a and $4\mathcal{N}$ anti-commuting complex parameters $\varepsilon = (\epsilon, \epsilon^\dagger)$, where $\epsilon = (\epsilon_i^\alpha)$ and $\epsilon^\dagger = (\bar{\epsilon}^{\dot{\alpha}i})$, $\bar{\epsilon}^{\dot{\alpha}i} := \overline{\epsilon_i^\alpha}$. In the vector notation, eq. (2.2) reads

$$b_{(\pm)}^a := b^a \pm i\epsilon_i\sigma^a\bar{\epsilon}^i = b^a \pm i\epsilon_i\alpha^\alpha(\sigma^a)_{\alpha\dot{\alpha}}\bar{\epsilon}^{\dot{\alpha}i}. \quad (2.3)$$

Introduce Goldstone fields $Z^A(x) = (X^a(x), \Theta_i^\alpha(x), \bar{\Theta}^{\dot{\alpha}i}(x))$ for spacetime translations (X^a) and supersymmetry transformations ($\Theta_i^\alpha, \bar{\Theta}^{\dot{\alpha}i}$). They parametrise the homogeneous space (\mathcal{N} -extended Minkowski superspace)

$$\mathbb{M}^{4|4\mathcal{N}} = \frac{\mathfrak{P}(4|\mathcal{N})}{\mathbf{SL}(2, \mathbb{C}) \times \mathbf{U}(\mathcal{N})} \quad (2.4)$$

according to the rule:

$$\mathfrak{S}(Z) = \left(\begin{array}{c|c|c} \mathbb{1}_2 & 0 & 0 \\ \hline -i\tilde{X}_{(+)} & \mathbb{1}_2 & 2\Theta^\dagger \\ \hline 2\Theta & 0 & \mathbb{1}_{\mathcal{N}} \end{array} \right) \implies \mathfrak{S}^{-1}(Z) = \left(\begin{array}{c|c|c} \mathbb{1}_2 & 0 & 0 \\ \hline i\tilde{X}_{(-)} & \mathbb{1}_2 & -2\Theta^\dagger \\ \hline -2\Theta & 0 & \mathbb{1}_{\mathcal{N}} \end{array} \right), \quad (2.5)$$

where

$$\tilde{X}_{(\pm)} = \tilde{X} \pm 2i\Theta^\dagger\Theta. \quad (2.6)$$

We define gauge super-Poincaré transformations by

$$g(x) : Z(x) \rightarrow Z'(x), \quad g\mathfrak{S}(Z) = \mathfrak{S}(Z')h, \quad (2.7)$$

with $g = Sh$. This is equivalent to the following transformations of the Goldstone fields:

$$S(b, \varepsilon) : \quad \tilde{X}' = \tilde{X} + \tilde{b} + 2i(\epsilon^\dagger\Theta - \Theta^\dagger\epsilon), \quad (2.8a)$$

$$\Theta' = \Theta + \epsilon \quad (2.8b)$$

and

$$h(M, U) : \quad \tilde{X}' = (M^\dagger)^{-1} \tilde{X} M^{-1} , \quad (2.9a)$$

$$\Theta' = U \Theta M^{-1} . \quad (2.9b)$$

Introduce a connection $\mathfrak{A} = dx^m \mathfrak{A}_m$ taking its values in the super-Poincaré algebra,

$$\mathfrak{A} := \left(\begin{array}{c|c|c} \Omega & 0 & 0 \\ \hline -i \tilde{e} & -\Omega^\dagger & 2\psi^\dagger \\ \hline 2\psi & 0 & iV \end{array} \right) = \left(\begin{array}{c|c|c} \Omega_{\alpha\beta} & 0 & 0 \\ \hline -i e^{\dot{\alpha}\beta} & -\bar{\Omega}^{\dot{\alpha}\beta} & 2\bar{\psi}^{\dot{\alpha}j} \\ \hline 2\psi_{i\beta} & 0 & iV_i^j \end{array} \right) , \quad (2.10)$$

and possessing the gauge transformation law

$$\mathfrak{A}' = g \mathfrak{A} g^{-1} + g dg^{-1} . \quad (2.11)$$

Here the one-forms $\Omega_{\alpha\beta}$ and $\bar{\Omega}^{\dot{\alpha}\beta}$ are the spinor counterparts of the Lorentz connection $\Omega^{ab} = dx^m \Omega_m^{ab} = -\Omega^{ba}$ such that

$$\Omega_{\alpha\beta} = \frac{1}{2} (\sigma^{ab})_{\alpha\beta} \Omega_{ab} , \quad \bar{\Omega}^{\dot{\alpha}\beta} = -\frac{1}{2} (\tilde{\sigma}^{ab})^{\dot{\alpha}\beta} \Omega_{ab} . \quad (2.12)$$

The Lorentz connection is an independent field. It is expressed in terms of the other fields on the mass shell. The one-form $e^{\dot{\alpha}\beta}$ is the spinor counterpart of the vierbein $e^a = dx^m e_m^a$. The fermionic one-forms $\psi_{i\beta}$ and $\bar{\psi}^{\dot{\alpha}j}$ describe \mathcal{N} gravitini. Finally, the anti-Hermitian one-form iV is the $U(\mathcal{N})$ gauge field.

Associated with \mathfrak{S} and \mathfrak{A} is another connection

$$\mathbb{A} := \mathfrak{S}^{-1} \mathfrak{A} \mathfrak{S} + \mathfrak{S}^{-1} d\mathfrak{S} , \quad (2.13)$$

which is characterised by the gauge transformation law

$$\mathbb{A}' = h \mathbb{A} h^{-1} + h dh^{-1} , \quad (2.14)$$

for an arbitrary gauge parameter $g = Sh$. This transformation law tells us that \mathbb{A} is invariant under all gauge transformations of the form $g = S(b, \epsilon)$ which describe local spacetime translations and supersymmetry transformations. The connection \mathbb{A} is the main object of the Volkov-Soroka construction. It has the form

$$\mathbb{A} := \left(\begin{array}{c|c|c} \Omega & 0 & 0 \\ \hline -i \tilde{E} & -\Omega^\dagger & 2\Psi^\dagger \\ \hline 2\Psi & 0 & iV \end{array} \right) , \quad (2.15)$$

where we have defined

$$\Psi := \psi + \mathcal{D}\Theta , \quad (2.16a)$$

$$\Psi^\dagger := \psi^\dagger + \mathcal{D}\Theta^\dagger , \quad (2.16b)$$

$$\tilde{E} := \tilde{e} + \mathcal{D}\tilde{X} + 4i(\Psi^\dagger - \frac{1}{2}\mathcal{D}\Theta^\dagger)\Theta - 4i\Theta^\dagger(\Psi - \frac{1}{2}\mathcal{D}\Theta) , \quad (2.16c)$$

and \mathcal{D} denotes the covariant derivative,

$$\mathcal{D}\Theta = d\Theta - \Theta\Omega + iV\Theta , \quad (2.17a)$$

$$\mathcal{D}\Theta^\dagger = d\Theta^\dagger - \Omega^\dagger\Theta^\dagger - i\Theta^\dagger V , \quad (2.17b)$$

$$\mathcal{D}\tilde{X} = d\tilde{X} - \Omega^\dagger\tilde{X} - \tilde{X}\Omega . \quad (2.17c)$$

Equation (2.14) is equivalent to the following gauge transformation laws:

$$\Omega' = M\Omega M^{-1} + M dM^{-1} , \quad (2.18a)$$

$$iV' = U(iV)U^{-1} + U dU^{-1} \quad (2.18b)$$

and

$$\tilde{E}' = (M^\dagger)^{-1}\tilde{E}M^{-1} , \quad (2.19a)$$

$$\Psi' = U\Psi M^{-1} , \quad (2.19b)$$

We see that the supersymmetric one-forms E^a and Ψ_i^β transform as tensors with respect to the Lorentz and $U(\mathcal{N})$ gauge groups.

Making use of (2.8), we deduce the local supersymmetry transformation of the gravitini and the vielbein

$$\psi' = \psi - \mathcal{D}\epsilon , \quad (2.20a)$$

$$\tilde{e}' = \tilde{e} + 4i(\epsilon^\dagger\psi - \psi^\dagger\epsilon) + i(\mathcal{D}\epsilon^\dagger\epsilon - \epsilon^\dagger\mathcal{D}\epsilon) . \quad (2.20b)$$

In the infinitesimal case, this transformation can be rewritten in the form

$$\delta_\epsilon\psi = -\mathcal{D}\epsilon , \quad \delta_\epsilon e^a = 2i \operatorname{tr}[\sigma^a(\psi^\dagger\epsilon - \epsilon^\dagger\psi)] . \quad (2.21)$$

As pointed out by Volkov [11], these transformation laws coincide with those used by Deser and Zumino in their construction of $\mathcal{N} = 1$ supergravity [10]. We should remark that the supersymmetry transformations of the Goldstone fields X^a and Θ_i^β are given by the relations (2.8).

Let us consider a local Poincaré translation, $S(b, 0)$. It only acts on the Goldstone vector field X^a and the vierbein e^a ,

$$X'^a = X^a + b^a, \quad e'^a = e^a - \mathcal{D}b^a. \quad (2.22)$$

We have two types of gauge transformations with vector-like parameters, the general coordinates transformations and the local Poincaré translations. The latter gauge freedom can be fixed by imposing the condition $X^a = 0$, and then we stay only with the general coordinate invariance. However, in what follows we will keep X^a intact.

The curvature tensor is given by

$$\mathbb{R} = d\mathbb{A} - \mathbb{A} \wedge \mathbb{A}, \quad \mathbb{R}' = h\mathbb{R}h^{-1}. \quad (2.23)$$

Its explicit form is

$$\mathbb{R} := \left(\begin{array}{c|c|c} R & 0 & 0 \\ \hline -i\tilde{\mathbb{T}} & -R^\dagger & 2\mathcal{D}\Psi^\dagger \\ \hline 2\mathcal{D}\Psi & 0 & iF \end{array} \right), \quad (2.24)$$

where $R = (R_\alpha^\beta)$ and $R^\dagger = (\bar{R}^{\dot{\alpha}}_{\dot{\beta}})$ form the Lorentz curvature, $F = (F_i^j)$ is the Yang-Mills field strength,

$$\mathcal{D}\Psi = d\Psi - \Psi \wedge \Omega - iV \wedge \Psi, \quad (2.25a)$$

$$\mathcal{D}\Psi^\dagger = d\Psi^\dagger + \Omega^\dagger \wedge \Psi^\dagger - i\Psi^\dagger \wedge V \quad (2.25b)$$

are the gravitino field strengths, and

$$\tilde{\mathbb{T}} = d\tilde{E} - \tilde{E} \wedge \Omega + \Omega^\dagger \wedge \tilde{E} - 4i\Psi^\dagger \wedge \Psi = \mathcal{D}\tilde{E} - 4i\Psi^\dagger \wedge \Psi \quad (2.26)$$

is the supersymmetric torsion tensor. In the vector notation, the torsion tensor reads

$$\mathbb{T}^a = \mathcal{D}E^a + 2i\Psi \wedge \sigma^a \bar{\Psi}. \quad (2.27)$$

It should be pointed out that the exterior derivative is defined to obey the property

$$d(\Sigma_p \wedge \Sigma_q) = \Sigma_p \wedge d\Sigma_q + (-1)^q d\Sigma_p \wedge \Sigma_q, \quad (2.28)$$

which is used for superforms [18].

The above results allow one to construct gauge-invariant functionals that can be used to construct a locally supersymmetric action. The invariants proposed in [3, 4] are the following:

- The Einstein-Hilbert action

$$S_{\text{EH}} = \frac{1}{4} \int \varepsilon_{abcd} E^a \wedge E^b \wedge R^{cd} ; \quad (2.29)$$

- The Rarita-Schwinger action

$$S_{\text{RS}} = \frac{1}{2} \int \left(\Psi_i \wedge E^a \wedge \sigma_a \mathcal{D} \bar{\Psi}^i - \mathcal{D} \Psi_i \wedge e^a \wedge \sigma_a \bar{\Psi}^i \right) ; \quad (2.30)$$

- The cosmological term

$$S_{\text{cosmological}} = \frac{1}{24} \int \varepsilon_{abcd} E^a \wedge E^b \wedge E^c \wedge E^d . \quad (2.31)$$

- $\text{O}(\mathcal{N})$ -invariant mass term

$$S_{\text{mass}} = \frac{i}{4} \int E^a \wedge E^b \wedge \left(\delta^{ij} \Psi_i \wedge \sigma_{ab} \Psi_j - \delta_{ij} \bar{\Psi}^i \wedge \tilde{\sigma}_{ab} \bar{\Psi}^j \right) . \quad (2.32)$$

The functionals S_{EH} , S_{RS} and $S_{\text{cosmological}}$ are $\text{U}(\mathcal{N})$ invariant. The cosmological term, eq. (2.31), also contains the kinetic term for the Goldstini [1, 2]. The mass term (2.32) is invariant under local internal transformations only if the group $\text{U}(\mathcal{N})$ is replaced with $\text{O}(\mathcal{N})$, and the gauge connection iV takes its values in the Lie algebra $\mathfrak{so}(\mathcal{N})$. The Yang-Mills action associated with V is obviously supersymmetric, but it will not be used in what follows.

In the $\mathcal{N} = 1$ case, the Volkov-Soroka theory is described by the general action

$$S = S_{\text{EH}} + 4c S_{\text{RS}} + 4m S_{\text{mass}} + \lambda S_{\text{cosmological}} , \quad (2.33)$$

with c , m and λ coupling constants.

3 Second local supersymmetry

In this section our consideration is restricted to the $\mathcal{N} = 1$ case, and the $\text{U}(1)$ gauge field is switched off, $V = 0$. We are going to demonstrate that the action

$$S_{\text{SUGRA}} = S_{\text{EH}} + 4S_{\text{RS}} \quad (3.1)$$

is invariant under a new local supersymmetry transformation described by the parameter $\varepsilon = (\epsilon^\alpha, \bar{\epsilon}^{\dot{\alpha}})$. It acts on the composite fields Ψ^α and E^a , defined by eq. (2.16), and the Lorentz connection as follows

$$\delta_\varepsilon \Psi^\alpha = -\mathcal{D} \epsilon^\alpha , \quad \delta_\varepsilon E^a = 2i (\Psi \sigma^a \bar{\epsilon} - \epsilon \sigma^a \Psi) , \quad (3.2a)$$

$$\frac{1}{4}\varepsilon_{abcd}\delta_{\boldsymbol{\varepsilon}}\Omega^{bc}\wedge E^d = -\boldsymbol{\varepsilon}\sigma_a\mathcal{D}\bar{\Psi} - \mathcal{D}\Psi\sigma_a\bar{\boldsymbol{\varepsilon}}. \quad (3.2b)$$

The Goldstone fields are inert under this transformation,

$$\delta_{\boldsymbol{\varepsilon}}X^a = 0, \quad \delta_{\boldsymbol{\varepsilon}}\Theta^\alpha = 0. \quad (3.2c)$$

The elementary field ψ^α and e^a transform as follows:

$$\delta_{\boldsymbol{\varepsilon}}\psi^\alpha = -\mathcal{D}\boldsymbol{\varepsilon}^\alpha + \Theta^\beta\delta_{\boldsymbol{\varepsilon}}\Omega_{\beta}{}^\alpha, \quad (3.2d)$$

$$\begin{aligned} \delta_{\boldsymbol{\varepsilon}}e^a &= 2i(\Psi\sigma^a\bar{\boldsymbol{\varepsilon}} - \boldsymbol{\varepsilon}\sigma^a\bar{\Psi}) + 2i(\Theta\sigma^a\mathcal{D}\bar{\boldsymbol{\varepsilon}} - \mathcal{D}\boldsymbol{\varepsilon}\sigma^a\bar{\Theta}) - \delta_{\boldsymbol{\varepsilon}}\Omega^a{}_bX^b \\ &\quad + \frac{1}{2}\varepsilon^{abcd}\delta_{\boldsymbol{\varepsilon}}\Omega_{bc}\Theta\sigma_d\bar{\Theta}. \end{aligned} \quad (3.2e)$$

The dependence on $\delta_{\boldsymbol{\varepsilon}}\Omega$ in (3.2d) and (3.2e) is such that the composite fields Ψ^α and E^a remain unchanged when the connection gets the displacement $\Omega \rightarrow \Omega + \delta_{\boldsymbol{\varepsilon}}\Omega$. We should point out that the action (3.1) involves only the one-forms Ψ^α , $\bar{\Psi}^{\dot{\alpha}}$, E^a and Ω^{ab} and their descendants. We also remark that the transformations (3.2a) and (3.2b) reduce to those given by Deser and Zumino [10] if the Goldstone fields X^a and Θ^α are switched off.

It should be pointed out that eq. (3.2b) uniquely determines $\delta_{\boldsymbol{\varepsilon}}\Omega^{bc}$. Indeed, given a vector-valued two-form

$$\Sigma_a = \frac{1}{2}E^c \wedge E^b \Sigma_{a,bc}, \quad (3.3)$$

the following equation

$$\frac{1}{2}\varepsilon_{abcd}\omega^{bc}\wedge E^d \equiv \tilde{\omega}_{ab}\wedge E^b = \Sigma_a, \quad \tilde{\omega}_{ab} = E^c\tilde{\omega}_{c,ab} \quad (3.4)$$

on the one-form $\omega^{ab} = E^c\omega_c{}^{ab}$, which takes its values in the Lorentz algebra, has the unique solution

$$\tilde{\omega}_{c,ab} = \frac{1}{2}\left(\Sigma_{a,bc} - \Sigma_{b,ac} - \Sigma_{c,ab}\right). \quad (3.5)$$

Now we turn to demonstrating that the action (3.1) is invariant under the supersymmetry transformation (3.2). Let us first take into account the variations in (3.2a). Varying the Einstein-Hilbert action (2.29) for $\boldsymbol{\varepsilon} \neq 0$ and $\bar{\boldsymbol{\varepsilon}} = 0$ gives

$$\delta_{\boldsymbol{\varepsilon}}^{(1)}S_{\text{EH}} = -i \int \varepsilon_{abcd}R^{ab}\wedge E^c \wedge \boldsymbol{\varepsilon}\sigma^d\bar{\Psi}. \quad (3.6)$$

Varying the Rarita-Schwinger action (2.30) gives

$$\delta_{\boldsymbol{\varepsilon}}^{(1)}S_{\text{RS}} = \int \left\{ \frac{1}{2}\mathcal{D}E^a \wedge \boldsymbol{\varepsilon}\sigma_a\mathcal{D}\bar{\Psi} + i\boldsymbol{\varepsilon}\sigma^a\bar{\Psi}\Psi\sigma_a\mathcal{D}\bar{\Psi} \right\}$$

$$+\frac{1}{2} \int \left\{ E^a \wedge \epsilon \sigma_a R^\dagger \bar{\Psi} - E^a \wedge \epsilon R \wedge \sigma^a \bar{\Psi} \right\}, \quad (3.7)$$

where we have used the relations

$$\mathcal{D}\mathcal{D}\epsilon = -\epsilon R, \quad \mathcal{D}\mathcal{D}\Psi^\dagger = -R^\dagger \wedge \Psi^\dagger. \quad (3.8)$$

Here $\delta^{(1)}$ means that we have taken into account only the variations (3.2a).

Making use of the identities (A.13), the curvature-dependent contributions in (3.7) can be rearranged as

$$\frac{1}{2} \int \left\{ E^a \wedge \epsilon \sigma_a R^\dagger \bar{\Psi} - E^a \wedge \epsilon R \wedge \sigma^a \bar{\Psi} \right\} = \frac{i}{4} \int \varepsilon_{abcd} R^{ab} \wedge E^c \wedge \epsilon \sigma^d \bar{\Psi}. \quad (3.9)$$

In order for the curvature contributions in (3.6) and (3.7) to cancel each other, we must consider the action (3.1), for which we obtain

$$\delta_{\epsilon}^{(1)}(S_{\text{EH}} + 4S_{\text{RS}}) = 2 \int \left(\mathcal{D}E^a + 2i\Psi\sigma^a\bar{\Psi} \right) \wedge \epsilon \sigma_a \mathcal{D}\bar{\Psi} = 2 \int \epsilon \sigma_a \mathcal{D}\bar{\Psi} \wedge \mathbb{T}^a, \quad (3.10)$$

with \mathbb{T}^a being the supersymmetric torsion tensor (2.27). Adding the complex conjugate part gives

$$\delta_{\bar{\epsilon}}^{(1)}(S_{\text{EH}} + 4S_{\text{RS}}) = 2 \int \left(\epsilon \sigma_a \mathcal{D}\bar{\Psi} + \mathcal{D}\Psi \sigma_a \bar{\epsilon} \right) \wedge \mathbb{T}^a. \quad (3.11)$$

Next, let us vary the action (3.1) with respect to the Lorentz connection Ω^{ab} . We give the Lorentz connection a small disturbance, $\Omega \rightarrow \Omega + \delta_{\epsilon}\Omega$, with $\delta_{\epsilon}\Omega$ to be determined below, and assume that the fields ψ^α and e^a also acquire $\delta_{\epsilon}\Omega$ -dependent variations given in (3.2d) and (3.2e). We denote $\delta^{(2)}$ the corresponding variation. Direct calculations give

$$\delta_{\epsilon}^{(2)}(S_{\text{EH}} + 4S_{\text{RS}}) = \frac{1}{2} \int \varepsilon_{abcd} \delta_{\epsilon} \Omega^{bc} \wedge E^d \wedge \mathbb{T}^a. \quad (3.12)$$

Combining the results (3.11) and (3.12), we end up with

$$\delta_{\epsilon}(S_{\text{EH}} + 4S_{\text{RS}}) = 2 \int \left(\epsilon \sigma_a \mathcal{D}\bar{\Psi} + \mathcal{D}\Psi \sigma_a \bar{\epsilon} + \frac{1}{4} \varepsilon_{abcd} \delta_{\epsilon} \Omega^{bc} \wedge E^d \right) \wedge \mathbb{T}^a. \quad (3.13)$$

This variation vanishes if $\delta_{\epsilon}\Omega$ is given by eq. (3.2b).

We have demonstrated that the theory (3.1) has two types of local supersymmetry. The original Volkov-Soroka supersymmetry is described by the relations

$$\delta_{\epsilon} \psi^\alpha = -\mathcal{D}\epsilon^\alpha, \quad \delta_{\epsilon} e^a = 2i(\psi\sigma^a\bar{\epsilon} - \epsilon\sigma^a\bar{\psi}), \quad (3.14a)$$

$$\delta_\epsilon \Theta^\alpha = \epsilon^\alpha , \quad \delta_\epsilon X^a = i(\Theta \sigma^a \bar{\epsilon} - \epsilon \sigma^a \bar{\Theta}) . \quad (3.14b)$$

The second supersymmetry, introduced in this work, is given by eq. (3.2). The gauge transformations (2.22) and (3.14) allow us to gauge away the Goldstone fields $Z^A(x) = (X^a(x), \Theta^\alpha(x), \bar{\Theta}^{\dot{\alpha}}(x))$ by imposing the conditions

$$X^a = 0 , \quad \Theta^\alpha = 0 . \quad (3.15)$$

As a result, the action (3.1) turns into the supergravity action proposed by Deser and Zumino [10], and the local supersymmetry transformation laws (3.2a) turn into those given in [10].

4 Conclusion

A few years ago, it was shown [19] that the Volkov-Soroka theory (2.33), with non-vanishing parameters c , m and λ , is equivalent to spontaneously broken $\mathcal{N} = 1$ supergravity [20, 21] which was called de Sitter supergravity in [22].

In this paper we have demonstrated that the action (3.1) is a Stückelberg-type extension of the $\mathcal{N} = 1$ supergravity action proposed by Deser and Zumino. Equivalently, the latter theory is a gauged-fixed version of (3.1). Therefore, pure $\mathcal{N} = 1$ supergravity is a special case of the Volkov-Soroka theory, and it could have been discovered before 1976.

The new local supersymmetry (3.2) of the action (3.1) is the main original result of this paper.

Acknowledgements:

I thank Evgeny Buchbinder, Ian McArthur and Dmitri Sorokin for discussions, and Darren Grasso and Michael Ponds for comments on the manuscript. This work was supported in part by the Australian Government through the Australian Research Council, project No. DP200101944.

A Two-component spinor formalism

In this appendix we collect the key formulae of the two-component spinor formalism. Our notation and two-component spinor conventions correspond to those used in [18, 23].

In particular, the Minkowski metric is $\eta_{ab} = \text{diag}(-1, +1, +1, +1)$, and the Levi-Civita tensor ε^{abcd} is normalised by $\varepsilon^{0123} = 1$.

Given a four-vector p^a , it can equivalently be described as an Hermitian 2×2 matrix with lower spinor indices

$$p := p^a \sigma_a = p^\dagger = (p_{\alpha\dot{\beta}}) , \quad \sigma_a = (\mathbb{1}_2, \vec{\sigma}) , \quad (\text{A.1})$$

or as an Hermitian 2×2 matrix with upper spinor indices

$$\tilde{p} := p^a \tilde{\sigma}_a = \tilde{p}^\dagger = (p^{\dot{\alpha}\beta}) , \quad \tilde{\sigma}_a = (\mathbb{1}_2, -\vec{\sigma}) , \quad (\text{A.2})$$

with $\vec{\sigma}$ being the Pauli matrices. The two sets of the relativistic Pauli matrices, σ_a and $\tilde{\sigma}_a$, are related to each other by the rule

$$(\tilde{\sigma}_a)^{\dot{\alpha}\alpha} = \varepsilon^{\dot{\alpha}\dot{\beta}} \varepsilon^{\alpha\beta} (\sigma_a)_{\beta\dot{\beta}} , \quad (\text{A.3})$$

where $\varepsilon^{\alpha\beta}$ and $\varepsilon_{\alpha\beta}$, $\varepsilon^{\dot{\alpha}\dot{\beta}}$ and $\varepsilon_{\dot{\alpha}\dot{\beta}}$ are antisymmetric spinor metrics normalised as $\varepsilon^{12} = \varepsilon_{21} = 1$ and $\varepsilon^{\dot{1}\dot{2}} = \varepsilon_{\dot{2}\dot{1}} = 1$. These are used to raise and lower the spinor indices,

$$\psi^\alpha = \varepsilon^{\alpha\beta} \psi_\beta , \quad \psi_\alpha = \varepsilon_{\alpha\beta} \psi^\beta , \quad (\text{A.4})$$

and similarly for the dotted spinors.

Let $\mathfrak{P}(4)$ be the universal covering group of the restricted Poincaré group $\text{ISO}_0(3, 1)$. It is usually realised as the group of linear inhomogeneous transformations (M, b) acting on the space of 2×2 Hermitian matrices $x = x^a \sigma_a = x^\dagger$ as follows:

$$x \rightarrow x' = x'^a \sigma_a = M x M^\dagger + b , \quad b = b^a \sigma_a = b^\dagger , \quad M = (M_\alpha^\beta) \in \text{SL}(2, \mathbb{C}) . \quad (\text{A.5})$$

Here $M^\dagger := \bar{M}^T$ is the Hermitian conjugate of M , and $\bar{M} = (\bar{M}_{\dot{\alpha}}^{\dot{\beta}})$ the complex conjugate of M , with $\bar{M}_{\dot{\alpha}}^{\dot{\beta}} := \overline{M_\alpha^\beta}$. The group $\mathfrak{P}(4)$ is equivalently realised as the group of linear inhomogeneous transformations acting on the space of 2×2 Hermitian matrices $\tilde{x} := x^a \tilde{\sigma}_a = \tilde{x}^\dagger$ as follows:

$$\tilde{x} \rightarrow \tilde{x}' = x'^a \tilde{\sigma}_a = (M^{-1})^\dagger \tilde{x} M^{-1} + \tilde{b} , \quad \tilde{b} = b^a \tilde{\sigma}_a . \quad (\text{A.6})$$

In Minkowski space $\mathbb{M}^4 \equiv \mathbb{R}^{3,1}$, the transformation (A.5) or, equivalently, (A.6) looks like

$$x'^a = (\Lambda(M))^a_b x^b + b^a , \quad (\Lambda(M))^a_b = -\frac{1}{2} \text{tr}(\tilde{\sigma}^a M \sigma_b M^\dagger) . \quad (\text{A.7})$$

It is also possible to realise $\mathfrak{P}(4)$ as a subgroup of $\mathrm{SU}(2, 2)$ consisting of all block triangular matrices of the form:

$$(M, b) := \left(\begin{array}{c|c} M & 0 \\ \hline -i\tilde{b}M & (M^{-1})^\dagger \end{array} \right) = (\mathbb{1}_2, b) (M, 0) , \quad (\text{A.8})$$

with M and \tilde{b} as in (A.5) and (A.6). Minkowski space is the homogeneous space

$$\mathbb{M}^4 = \mathfrak{P}(4)/\mathrm{SL}(2, \mathbb{C}) , \quad (\text{A.9})$$

compare with eq. (2.4) defining the \mathcal{N} -extended Minkowski superspace. Its points are naturally parametrised by the Cartesian coordinates x^a corresponding to the coset representative:

$$(\mathbb{1}_2, x) = \left(\begin{array}{c|c} \mathbb{1}_2 & 0 \\ \hline -i\tilde{x} & \mathbb{1}_2 \end{array} \right) . \quad (\text{A.10})$$

Given an antisymmetric tensor field $F_{ab} = -F_{ba}$, it can be equivalently described by a symmetric rank-two spinor $F_{\alpha\beta} = F_{\beta\alpha}$ and its conjugate $\bar{F}_{\dot{\alpha}\dot{\beta}}$. The precise correspondence $F_{ab} \longleftrightarrow (F_{\alpha\beta}, \bar{F}_{\dot{\alpha}\dot{\beta}})$ is given by

$$F_{ab} = (\sigma_{ab})^{\alpha\beta} F_{\alpha\beta} - (\tilde{\sigma}_{ab})^{\dot{\alpha}\dot{\beta}} \bar{F}_{\dot{\alpha}\dot{\beta}} , \quad F_{\alpha}{}^{\beta} := \frac{1}{2}(\sigma^{ab})_{\alpha}{}^{\beta} F_{ab} , \quad \bar{F}^{\dot{\alpha}}{}_{\dot{\beta}} := -\frac{1}{2}(\tilde{\sigma}^{ab})^{\dot{\alpha}}{}_{\dot{\beta}} F_{ab} .$$

Here the matrices $\sigma_{ab} = \left((\sigma_{ab})_{\alpha}{}^{\beta} \right)$ and $\tilde{\sigma}_{ab} = \left((\tilde{\sigma}_{ab})^{\dot{\alpha}}{}_{\dot{\beta}} \right)$ are defined by

$$\sigma_{ab} = -\frac{1}{4}(\sigma_a \tilde{\sigma}_b - \sigma_b \tilde{\sigma}_a) , \quad \tilde{\sigma}_{ab} = -\frac{1}{4}(\tilde{\sigma}_a \sigma_b - \tilde{\sigma}_b \sigma_a) . \quad (\text{A.11})$$

These matrices are (anti) self-dual,

$$\frac{1}{2}\varepsilon^{abcd}\sigma_{cd} = -i\sigma^{ab} , \quad \frac{1}{2}\varepsilon^{abcd}\tilde{\sigma}_{cd} = i\tilde{\sigma}^{ab} . \quad (\text{A.12})$$

The important identities involving σ_{ab} and $\tilde{\sigma}_{ab}$ are:

$$\sigma_{ab}\sigma_c = -\frac{1}{2}(\eta_{ac}\sigma_b - \eta_{bc}\sigma_a) - \frac{i}{2}\varepsilon_{abcd}\sigma^d , \quad (\text{A.13a})$$

$$\sigma_c\tilde{\sigma}_{ab} = \frac{1}{2}(\eta_{ac}\sigma_b - \eta_{bc}\sigma_a) - \frac{i}{2}\varepsilon_{abcd}\sigma^d . \quad (\text{A.13b})$$

References

- [1] D. V. Volkov and V. P. Akulov, “Possible universal neutrino interaction,” JETP Lett. **16**, 438 (1972) [Pisma Zh. Eksp. Teor. Fiz. **16**, 621 (1972)]; “Is the neutrino a Goldstone particle?,” Phys. Lett. B **46**, 109 (1973).
- [2] V. P. Akulov and D. V. Volkov, “Goldstone fields with spin 1/2,” Theor. Math. Phys. **18**, 28 (1974) 28 [Teor. Mat. Fiz. **18**, 39 (1974)].
- [3] D. V. Volkov and V. A. Soroka, “Higgs effect for Goldstone particles with spin 1/2,” JETP Lett. **18**, 312 (1973) [Pisma Zh. Eksp. Teor. Fiz. **18**, 529 (1973)].
- [4] D. V. Volkov and V. A. Soroka, “Gauge fields for symmetry group with spinor parameters,” Theor. Math. Phys. **20**, 829 (1974) [Teor. Mat. Fiz. **20**, 291(1974)].
- [5] D. V. Volkov, “Phenomenological Lagrangians,” Sov. J. Particles Nucl. **4**, 1 (1973) [Fiz. Elem. Chast. Atom. Yadra **4**, 3 (1973)].
- [6] Yu. A. Golfand and E. P. Likhtman, “Extension of the algebra of Poincaré group generators and violation of P invariance,” JETP Lett. **13**, 323 (1971) [Pisma Zh. Eksp. Teor. Fiz. **13**, 452 (1971)].
- [7] J. Wess and B. Zumino, “Supergauge transformations in four dimensions,” Nucl. Phys. B **70**, 39 (1974).
- [8] A. Salam and J. A. Strathdee, “Super-gauge transformations,” Nucl. Phys. B **76**, 477 (1974).
- [9] D. Z. Freedman, P. van Nieuwenhuizen and S. Ferrara, “Progress toward a theory of supergravity,” Phys. Rev. D **13**, 3214 (1976).
- [10] S. Deser and B. Zumino, “Consistent supergravity,” Phys. Lett. B **62**, 335 (1976).
- [11] D. V. Volkov, “Supergravity before and after 1976,” in: *Concise Encyclopedia of Supersymmetry and Noncommutative Structures in Mathematics and Physics*, S. Duplij, W. Siegel and J. Bagger (Eds.), Kluwer Academic Publishers (2004), pp. 6–9 [arXiv:hep-th/9404153 [hep-th]].
- [12] D. V. Volkov, “Supergravity before 1976,” in: *History of Original Ideas and Basic Discoveries in Particle Physics*, H. B. Newman and T. Ypsilantis (Eds.), Plenum Press, New York (1996), pp. 663-675 [arXiv:hep-th/9410024 [hep-th]].
- [13] V. A. Soroka, “Starting point of supergravity,” [arXiv:hep-th/0111271 [hep-th]].
- [14] V. A. Soroka, “The Sources of supergravity,” in *The supersymmetric World: The beginning of the Theory*, G. Kane and M. Shifman (Eds.), World Scientific, Singapore (2000), pp. 88–92 [arXiv:hep-th/0203171 [hep-th]].
- [15] A. J. Nurmagambetov, “How old supergravity is: Thirty five years or more?,” Prob. Atomic Sci. Technol. **2011N5**, 3 (2011).
- [16] S. Duplij, “Supergravity was discovered by D.V. Volkov and V.A. Soroka in 1973, wasn’t it?,” Eur. J. Phys. **3**, 81 (2019) [arXiv:1910.03259 [physics.hist-ph]].
- [17] S. M. Kuzenko, “Lectures on nonlinear sigma-models in projective superspace,” J. Phys. A **43**, 443001 (2010) [arXiv:1004.0880 [hep-th]].

- [18] J. Wess and J. Bagger, *Supersymmetry and Supergravity*, Princeton University Press, Princeton, 1992.
- [19] I. Bandos, L. Martucci, D. Sorokin and M. Tonin, “Brane induced supersymmetry breaking and de Sitter supergravity,” *JHEP* **1602**, 080 (2016) [arXiv:1511.03024 [hep-th]].
- [20] S. Deser and B. Zumino, “Broken supersymmetry and supergravity,” *Phys. Rev. Lett.* **38**, 1433 (1977).
- [21] U. Lindström and M. Roček, “Constrained local superfields,” *Phys. Rev. D* **19**, 2300 (1979).
- [22] E. A. Bergshoeff, D. Z. Freedman, R. Kallosh and A. Van Proeyen, “Pure de Sitter supergravity,” *Phys. Rev. D* **92**, no. 8, 085040 (2015) Erratum: [*Phys. Rev. D* **93**, no. 6, 069901 (2016)] [arXiv:1507.08264 [hep-th]].
- [23] I. L. Buchbinder and S. M. Kuzenko, *Ideas and Methods of Supersymmetry and Supergravity or a Walk Through Superspace*, IOP, Bristol, 1998.