

# On two-dimensional Hamiltonian systems with sixth-order integrals of motion

E.O. Porubov, A.V. Tsiganov

*St.Petersburg State University, St.Petersburg, Russia*

*e-mail: st077283@student.spbu.ru, andrey.tsiganov@gmail.com*

## Abstract

We obtain 21 two-dimensional natural Hamiltonian systems with sextic invariants, which are polynomial of the sixth order in momenta. Following to Bertrand, Darboux, and Drach these results of the standard brute force experiments can be applied to construct a new mathematical theory.

**Keywords:** integrable systems

## 1 Introduction

A Hamiltonian system is called natural if Hamiltonian is the sum of positive-definite kinetic energy and potential

$$H = T(q, p) + V(q).$$

The search of natural Hamiltonian systems admitting integral of motion that is a polynomial of order  $N$  in the momenta is a classical problem [4, 5], which is currently intensively studied, see [2, 10, 11, 12, 15, 23] and references within.

There is few two-dimensional natural Hamiltonian systems with invariants (constants of motion or integrals of motion), which are polynomials of the sixth order in momenta. According to [13] the Hamiltonian vector field with Hamiltonian

$$H = p_1^2 + p_2^2 + V(q_1, q_2),$$

admits sextic invariants for a potential of the Holt-type system

$$V = 12q_2^{4/3} + (q_1^2 + a)q_2^{-2/3} + bq_1^{-2},$$

Toda-type system

$$V_1 = e^{\frac{(\sqrt{3}q_1 - q_2)}{2}} + e^{q_2} + e^{-\sqrt{3}q_1}, \quad V_2 = e^{\frac{(\sqrt{3}q_1 - q_2)}{2}} + e^{q_2} + e^{\frac{-\sqrt{3}q_1}{3}},$$

superintegrable Calogero-type system

$$V = v(2q_2) + v(\sqrt{3}q_1 + q_2) + v(-\sqrt{3}q_1 + q_2), \quad v(z) = z^{-2} + \frac{a^2}{2}z^{-2},$$

and a few superintegrable systems associated with the Chebychev theorem on binomial differentials.

In [9] we obtained one more system with sextic invariant at

$$V = (q_1^{-2/3} + a)q_2^{-2/3},$$

which is a generalization of the Fokas-Lagerström system having a cubic invariant [7, 13].

On the pseudo-Euclidean space, there are more natural Hamiltonian systems with sextic invariant [14]

$$H = p_1 p_2 + q_1^a q_2^b,$$

where  $a$  and  $b$  are rational numbers associated with a special set of parameters of the hypergeometric function, for instance

$$V = q_1^2 q_2^{-10/7}, \quad V = q_1^{-2/3} q_2^{-5/6}, \quad V = q_1^{-2/3} q_2^{-7/3}.$$

Similar to the Holt and Fokas-Lagerstrom potentials we have singular potentials which are well-defined functions only on part of the plane.

In [18] we construct one more Hamiltonian system with invariant of degree six on the two-dimensional space with metric depending on local coordinates

$$H = \frac{p_1^2}{2m_1(q_1, q_2)} + \frac{p_2^2}{2m_2(q_1, q_2)} + V(q_1, q_2),$$

i.e. a natural Hamiltonian system with position-dependent mass (effective mass). This system has been obtained using divisor arithmetic on an elliptic curve [18, 19]. Below we present 20 similar natural Hamiltonian systems with effective mass and sextic invariant.

Dynamical systems with position-dependent mass were first introduced in the theory of semiconductor physics, and now these systems can be found in many fields, such as classical Hamiltonian and non-Hamiltonian mechanics, quantum mechanics, relativistic mechanics, nuclear physics, molecular physics, neutrino mass oscillations, quantum information and so on, see references in recent papers [1, 3, 8, 16, 22].

Our aim is not to construct integrable systems with a sixth degree invariant or to study the applications of these systems in physics. Similar to Darboux [5] and Drach [6] we use the direct method of constructing invariants to collect a sufficient number of examples that allow us to build a mathematical theory explaining the existence of such sextic invariants.

## 1.1 Two-dimensional geodesic motion with cubic invariant

Let us consider two-dimensional metric

$$g = \begin{pmatrix} \frac{(kq_1 + q_2)q_1^m}{q_1 - q_2} & 0 \\ 0 & \frac{(kq_2 + q_1)q_2^m}{q_2 - q_1} \end{pmatrix}, \quad q_1 < 0 < q_2,$$

and the corresponding kinetic energy

$$T = g_{11}p_1^2 + g_{22}p_2^2 = \frac{(kq_1 + q_2)q_1^m}{q_1 - q_2} p_1^2 + \frac{(kq_2 + q_1)q_2^m}{q_2 - q_1} p_2^2, \quad (1.1)$$

derived in [18] in the framework of divisor arithmetic on an elliptic curve. Here  $q_{1,2}$  are the position variables and  $p_{1,2}$  are the conjugate momenta for canonical Poisson brackets

$$\{q_1, p_1\} = \{q_2, p_2\} = 1, \quad \{q_1, q_2\} = \{q_1, p_2\} = \{q_2, p_1\} = \{p_1, p_2\} = 0.$$

At  $k = \pm 1$  Hamiltonian system with geodesic Hamiltonian  $T$  (1.1) admits linear invariant and, therefore, sextic invariant may be obtained by using the Chebyshev theorem on binomial differentials [9]. Here, we do not consider this construction of integrals of motion.

**Proposition 1** *At special values of parameters  $m$  and  $k$*

$$\bullet m = 1, k = 2; \quad \bullet m = 3, k = \frac{1}{2}, 3; \quad \bullet m = 4, k = \pm 3, -\frac{3}{5}, -\frac{1}{7}, \frac{1}{5}, \frac{1}{2};$$

geodesic Hamiltonian  $T$  (1.1) commutes

$$\{T, K\} = 0$$

with the reducible cubic polynomial

$$K = K_1 K_2, \quad K_1 = (q_1^2 p_1 - q_2^2 p_2),$$

where  $K_2$  is a polynomial of the second order in momenta.

These special values of parameters are related to divisor arithmetic on an elliptic curve, which allows us to get separated variables for the corresponding Hamilton-Jacobi equation  $T = E$  and prove that these Hamiltonian systems are bi-Hamiltonian.

**Proposition 2** *At special values of parameters  $m$  and  $k$*

$$\bullet m = 1, k = 2; \quad \bullet m = 3, k = \frac{1}{2}, 2; \quad \bullet m = 4, k = -3, \frac{1}{3}, \frac{1}{5}, \frac{3}{5}, 2;$$

geodesic Hamiltonian  $T$  (1.1) commutes

$$\{T, L\} = 0$$

with the reducible quartic polynomial

$$L = K_1^2 L_2, \quad K_1 = (q_1^2 p_1 - q_2^2 p_2),$$

where  $L_2$  is a polynomial of the second order in momenta.

Thus, we have superintegrable geodesic motion at

$$\bullet m = 1, k = 2; \quad \bullet m = 3, k = \frac{1}{2}; \quad \bullet m = 4, k = -3, \frac{1}{5};$$

and in all these cases

$$\{L, K\} = K^2.$$

Following the example from [18], below we consider integrable Hamiltonian systems with integrals of motion

$$\begin{aligned} H_1 &= T + V(q_1, q_2), \\ H_2 &= \left( K_1^2 + U_1(q_1, q_2) \right) \left( K_2 + U_2(q_1, q_2) \right)^2 \\ &= K^2 + \left( U_1 K_2^2 + 2U_2 K_1 K_2 \right) + \left( U_2^2 K_1^2 + 2U_1 U_2 K_2 \right) + U_1 U_2^2. \end{aligned} \quad (1.2)$$

where  $V(q_1, q_2)$  and  $U_{1,2}(q_1, q_2)$  are the solution of partial differential equations obtained from the standard equation

$$\{H_1, H_2\} = 0, \quad (1.3)$$

which must be identically satisfied for all admissible values of  $p_1$  and  $p_2$ .

If reducible cubic polynomial  $K$  admits a few decompositions

$$K = K_1 K_2 = \hat{K}_1 \hat{K}_2 = \tilde{K}_1 \tilde{K}_2,$$

we obtain several integrable systems with integrals of motion (1.2) and

$$\begin{aligned} \hat{H}_1 &= T + \hat{V}(q_1, q_2), \quad \hat{H}_2 = \left( \hat{K}_1^2 + \hat{U}_1(q_1, q_2) \right) \left( \hat{K}_2 + \hat{U}_2(q_1, q_2) \right)^2, \\ \tilde{H}_1 &= T + \tilde{V}(q_1, q_2), \quad \tilde{H}_2 = \left( \tilde{K}_1^2 + \tilde{U}_1(q_1, q_2) \right) \left( \tilde{K}_2 + \tilde{U}_2(q_1, q_2) \right)^2. \end{aligned} \quad (1.4)$$

In the next section, we present formal solutions of the partial differential equations (1.3) associated with all parameters  $m$  and  $k$  in Proposition 1.

## 2 Invariants of sixth order in momenta

In [18] we supposed that  $q_{1,2}$  are parabolic coordinates on the plane

$$q_1 = y - \sqrt{x^2 + y^2}, \quad q_2 = y + \sqrt{x^2 + y^2}$$

so that the corresponding momenta are

$$p_1 = \frac{p_y}{2} + \frac{\sqrt{-q_1 q_2} p_x}{2q_1}, \quad p_2 = \frac{p_y}{2} + \frac{\sqrt{-q_1 q_2} p_x}{2q_2}.$$

In these variables, kinetic energy is equal to

$$\begin{aligned} m = 1, \quad T &= \frac{yp_x^2}{2} + \frac{x(k-1)p_x p_y}{2} + \frac{ky p_y^2}{2}, \\ m = 3, \quad T &= \frac{kyx^2 p_x^2}{2} + \frac{((k-1)x^2 + 4ky^2) x p_x p_y}{2} + \left( \frac{(2k-1)x^2}{2} + 2ky^2 \right) y p_y^2, \\ m = 4, \quad T &= \frac{x^2((k-1)x^2 + 4ky^2) p_x^2}{4} + xy((2k-1)x^2 + 4ky^2) p_x p_y \\ &+ \left( \frac{(k-1)x^4}{4} + (3k-1)x^2 y^2 + 4ky^4 \right) p_y^2. \end{aligned}$$

Below we also present potentials in these  $x, y$ -variables in two dimensional configuration space.

### 2.1 Case $m = 1$ and $k = 2$

In this case geodesic Hamiltonian

$$T = \frac{q_1(2q_1 + q_2)p_1^2}{q_1 - q_2} + \frac{q_2(q_1 + 2q_2)p_2^2}{q_2 - q_1}$$

commutes with cubic and quartic polynomials

$$K = \frac{q_1 q_2 (q_1^2 p_1 - q_2^2 p_2)(p_1 - p_2)^2}{(q_1 - q_2)^3}, \quad L = \frac{(q_1^2 p_1 - q_2^2 p_2)^2 (q_1 p_1^2 - q_2 p_2^2)}{(q_1 - q_2)^3}, \quad \{K, L\} = K^2.$$

Cubic polynomial  $K$  has the form  $K = K_1 K_2 = \hat{K}_1 \hat{K}_2$ , where

$$K_1 = (q_1^2 p_1 - q_2^2 p_2), \quad \text{and} \quad \hat{K}_1 = \frac{q_1 q_2 (p_1 - p_2)}{(q_1 - q_2)^3}.$$

The corresponding functions in (1.2) and (1.4) are

$$\begin{aligned} V &= \frac{a(4q_1^4 + 14q_1^3 q_2 + 19q_1^2 q_2^2 + 14q_1 q_2^3 + 4q_2^4)}{(q_1 + q_2)^2} + b(q_1 + q_2) + c \left( \frac{1}{q_1} + \frac{1}{q_2} \right), \\ U_1 &= 2a(q_1^2 + q_1 q_2 + q_2^2)(q_1 + q_2)(q_1 - q_2)^2 + \frac{b(q_1 - q_2)^2 (q_1 + q_2)^2}{2}, \\ U_2 &= \frac{2aq_1 q_2}{(q_1 + q_2)(q_1 - q_2)} - \frac{c}{q_1 q_2 (q_1 - q_2)}, \end{aligned}$$

and

$$\hat{V} = \frac{a(q_1^4 - 4q_1^3q_2 - 14q_1^2q_2^2 - 4q_1q_2^3 + q_2^4)}{(q_1 + q_2)^2} + \frac{b(q_1^2 + 3q_1q_2 + q_2^2)}{(q_1 + q_2)^{3/2}} + c(q_1 + q_2),$$

$$\hat{U}_1 = \frac{8aq_2^2q_1^2}{(q_1^2 - q_2^2)^4} - \frac{2bq_1q_2}{\sqrt{q_1 + q_2}(q_1 - q_2)^4},$$

$$\hat{U}_2 = \frac{2a(q_1^4 - q_2^4)}{q_1 + q_2} + \frac{b(q_1 - q_2)^2}{2\sqrt{q_1 + q_2}} + c(q_1 - q_2)^2.$$

In  $x, y$ -variables potentials read as

$$V = -\frac{a(x^4 - 8x^2y^2 - 64y^4)}{4y^2} + 2by - \frac{2cy}{x^2}, \quad (2.1)$$

$$\hat{V} = -\frac{a(x^4 - 8x^2y^2 - 4y^4)}{y^2} - \frac{\sqrt{2}b(x^2 - 4y^2)}{4y^{3/2}} + 2cy. \quad (2.2)$$

Using the method of undetermined coefficients we can prove that these Hamiltonian systems do not have quartic invariants. Indeed, substituting quartic polynomial

$$H_3 = L + \sum_{k=0}^4 \sum_{i=0}^k f_{ik}(q_1, q_2) p_1^{k-i} p_2^i$$

into the equations

$$\{T + V, H_3\} = 0 \quad \{T + \hat{V}, H_3\} = 0$$

we obtain two inconsistent systems of partial differential equations on functions  $f_{ik}(q_1, q_2)$ . It means that the corresponding Hamiltonian vector fields admit quartic invariants only at  $V = 0$  and  $\hat{V} = 0$ .

## 2.2 Case $m = 3$ and $k = 3$

Geodesic Hamiltonian

$$T = \frac{q_1^3(3q_1 + q_2)p_1^2}{q_1 - q_2} + \frac{q_2^3(q_1 + 3q_2)p_2^2}{q_2 - q_1},$$

commutes with cubic polynomial

$$K = \frac{q_1^{3/2} q_2^{3/2}}{(q_1 - q_2)^2} (q_1^2 p_1 - q_2^2 p_2) (q_1^{3/2} p_1 - q_2^{3/2} p_2) (q_1^{3/2} p_1 + q_2^{3/2} p_2).$$

So, we can take

$$K_1 = q_1^{3/2} q_2^{3/2} (q_1^2 p_1 - q_2^2 p_2), \quad \hat{K}_1 = q_1^{3/2} q_2^{3/2} (q_1^{3/2} p_1 - q_2^{3/2} p_2).$$

and

$$\tilde{K}_1 = q_1^{3/2} q_2^{3/2} (q_1^{3/2} p_1 + q_2^{3/2} p_2).$$

Solution of the equation (1.3) for  $V$  and  $U_{1,2}$  has the following form

$$V = \frac{a(q_1 + q_2)(q_1^2 + 3q_1q_2 + q_2^2)}{q_1^3 q_2^3} + b(q_1 + q_2) + c \left( \frac{1}{q_1} + \frac{1}{q_2} \right),$$

$$U_1 = \frac{a(q_1 - q_2)^2 (q_1 + q_2)^2}{4} + \frac{bq_1^3 q_2^3 (q_1 - q_2)^2}{3},$$

$$U_2 = \frac{a(q_1^2 + q_1q_2 + q_2^2)}{q_1^3 q_2^3 (q_1 - q_2)} + \frac{b}{3(q_1 - q_2)} + \frac{c}{q_1 q_2 (q_1 - q_2)}.$$

Other decompositions give rise to potentials

$$\hat{V} = \frac{a(13q_1^2 + 4\sqrt{q_1^3 q_2} + 46q_1 q_2 + 4\sqrt{q_1 q_2^3} + 13q_2^2)(\sqrt{q_1} + \sqrt{q_2})^2}{q_1^3 q_2^3} + \frac{bq_1 q_2 (\sqrt{q_1} + \sqrt{q_2})^2}{(3q_1 - 2\sqrt{q_1 q_2} + 3q_2)^2} + \frac{c(\sqrt{q_1} + \sqrt{q_2})^2}{q_1 q_2},$$

$$\tilde{V} = \frac{a(13q_1^2 - 4\sqrt{q_1^3 q_2} + 46q_1 q_2 - 4\sqrt{q_1 q_2^3} + 13q_2^2)(\sqrt{q_1} - \sqrt{q_2})^2}{q_1^3 q_2^3} + \frac{bq_1 q_2 (\sqrt{q_1} - \sqrt{q_2})^2}{(3q_1 + 2\sqrt{q_1 q_2} + 3q_2)^2} + \frac{c(\sqrt{q_1} - \sqrt{q_2})^2}{q_1 q_2},$$

and functions

$$\hat{U}_1 = -8a(\sqrt{q_1} - \sqrt{q_2})^2(\sqrt{q_1} + \sqrt{q_2})^4 + \frac{8bq_1^4 q_2^4 (\sqrt{q_1} - \sqrt{q_2})^2}{3(3q_1 - 2\sqrt{q_1 q_2} + 3q_2)^2},$$

$$\tilde{U}_1 = -8a(\sqrt{q_1} - \sqrt{q_2})^4(\sqrt{q_1} + \sqrt{q_2})^2 - \frac{8bq_1^4 q_2^4 (\sqrt{q_1} + \sqrt{q_2})^2}{3(3q_1 + 2\sqrt{q_1 q_2} + 3q_2)^2},$$

$$\hat{U}_2 = -\frac{a(5q_1 + 2\sqrt{q_1 q_2} + 5q_2)(q_1 - 6\sqrt{q_1 q_2} + q_2)}{(\sqrt{q_1} - \sqrt{q_2})q_1^3 q_2^3} - \frac{bq_1 q_2}{3(\sqrt{q_1} - \sqrt{q_2})(3q_1 - 2\sqrt{q_1 q_2} + 3q_2)^2} + \frac{c}{q_1 q_2(\sqrt{q_1} - \sqrt{q_2})},$$

$$\tilde{U}_2 = -\frac{a(5q_1 - 2\sqrt{q_1 q_2} + 5q_2)(q_1 + 6\sqrt{q_1 q_2} + q_2)}{(\sqrt{q_1} + \sqrt{q_2})q_1^3 q_2^3} + \frac{bq_1 q_2}{3(\sqrt{q_1} + \sqrt{q_2})(3q_1 + 2\sqrt{q_1 q_2} + 3q_2)^2} - \frac{c}{q_1 q_2(\sqrt{q_1} + \sqrt{q_2})},$$

In  $x, y$ -variables potentials have the form

$$V = \frac{2ay(x^2 - 4y^2)}{x^6} + 2by - \frac{2cy}{x^2}, \quad (2.3)$$

$$\hat{V} = i \left( -\frac{8a(iy - 5x - 8y)(iy - 5x + 8y)(iy - x)}{5x^6} - \frac{b(iy - x)x^2}{2(3iy + x)^2} - \frac{2c(iy - x)}{x^2} \right), \quad (2.4)$$

$$\tilde{V} = i \left( -\frac{8a(iy + 5x - 8y)(iy + 5x + 8y)(iy + x)}{5x^6} - \frac{b(iy + x)x^2}{2(3iy - x)^2} - \frac{2c(iy + x)}{x^2} \right), \quad (2.5)$$

where  $i = \sqrt{-1}$ . In two last cases canonical transformation of coordinates

$$y \rightarrow iy \quad \text{and} \quad p_y \rightarrow -ip_y,$$

changes kinetic energy  $T$  and potentials  $\hat{V}, \tilde{V}$  so that the corresponding Hamiltonians

$$\hat{H}_1 = i(T + \hat{V}), \quad \text{and} \quad \tilde{H}_1 = i(T + \tilde{V})$$

become real functions.

### 2.3 Case $m = 3$ and $k = 1/2$

Quadratic polynomial

$$T = \frac{q_1^3 \left(\frac{q_1}{2} + q_2\right) p_1^2}{q_1 - q_2} + \frac{q_2^3 \left(q_1 + \frac{q_2}{2}\right) p_2^2}{q_2 - q_1}$$

commutes with cubic and quartic polynomials

$$K = \frac{q_1^2 q_2^2 (q_1^2 p_1 - q_2^2 p_2)^2 (p_1 - p_2)}{(q_1 - q_2)^3}, \quad L = \frac{(q_1^2 p_1 - q_2^2 p_2)^3 (q_1^2 (q_1 + 3q_2) p_1 + q_2^2 (3q_1 + q_2) p_2)}{4(q_1 - q_2)^3}$$

so that  $\{L, K\} = K^2$ . Because cubic polynomial  $K$  admits two decompositions

$$K = K_1 K_2 = \hat{K}_1 \hat{K}_2,$$

where

$$K_1 = q_1^2 q_2^2 (q_1^2 p_1 - q_2^2 p_2), \quad \text{and} \quad \hat{K}_1 = \frac{p_1 - p_2}{(q_1 - q_2)^3},$$

we can construct two different integrable systems with sextic invariants so that

$$V = \frac{a(q_1^4 - 4q_1^3q_2 - 14q_1^2q_2^2 - 4q_1q_2^3 + q_2^4)}{q_1^2q_2^2(q_1 + q_2)^2} + \frac{b(q_1^2 + 3q_1q_2 + q_2^2)}{\sqrt{q_1q_2}(q_1 + q_2)^{3/2}} + c\left(\frac{1}{q_1} + \frac{1}{q_2}\right),$$

$$U_1 = -\frac{16aq_1^3q_2^3(q_1 - q_2)^2}{q_1 + q_2} + \frac{4bq_1^{7/2}q_2^{7/2}(q_1 - q_2)^2}{\sqrt{q_1 + q_2}},$$

$$U_2 = -\frac{4a(q_1^2 + q_2^2)}{q_1^3q_2^3(q_1^2 - q_2^2)} - \frac{b}{q_1^{3/2}q_2^{3/2}\sqrt{q_1 + q_2}(q_1^2 - q_2^2)} - \frac{2c}{q_1^2q_2^2(q_1 - q_2)},$$

and

$$\hat{V} = \frac{a(4q_1^4 + 14q_1^3q_2 + 19q_1^2q_2^2 + 14q_1q_2^3 + 4q_2^4)}{q_1^2q_2^2(q_1 + q_2)^2} + b(q_1 + q_2) + c\left(\frac{1}{q_1} + \frac{1}{q_2}\right),$$

$$\hat{U}_1 = -\frac{4a(q_1 + q_2)(q_1^2 + q_1q_2 + q_2^2)}{q_1^5q_2^5(q_1 - q_2)^4} - \frac{c(q_1 + q_2)^2}{q_1^4q_2^4(q_1 - q_2)^4},$$

$$\hat{U}_2 = -\frac{4aq_1q_2(q_1 - q_2)^2}{q_1 + q_2} + 2bq_1^2q_2^2(q_1 - q_2)^2.$$

In this case

$$V = -\frac{a(x^4 - 8x^2y^2 - 4y^4)}{x^4y^2} + \frac{ib\sqrt{2}(x^2 - 4y^2)}{4xy^{3/2}} - \frac{2cy}{x^2}, \quad (2.6)$$

$$\hat{V} = -\frac{a(x^4 - 8x^2y^2 - 64y^4)}{4x^4y^2} + 2by - \frac{2cy}{x^2}. \quad (2.7)$$

After changing parameter  $b \rightarrow -ib$  we obtain real potential  $V$  (2.6).

## 2.4 Case $m = 4$ and $k = -3$

In this case geodesic Hamiltonian

$$T = \frac{q_1^4(q_2 - 3q_1)p_1^2}{q_1 - q_2} + \frac{q_2^4(q_1 - 3q_2)p_2^2}{q_2 - q_1}$$

commutes with the reducible cubic polynomial

$$K = K_1K_2 = \hat{K}_1\hat{K}_2 = \frac{q_1q_2(q_1^2p_1 - q_2^2p_2)(q_1^2p_1 + q_2^2p_2)^2}{q_1 - q_2},$$

where

$$K_1 = q_1^2p_1 - q_2^2p_2 \quad \text{and} \quad \hat{K}_1 = \frac{q_1q_2(q_1^2p_1 + q_2^2p_2)}{q_1 - q_2}.$$

The corresponding quartic invariant is equal to

$$L = \frac{(q_1^2p_1 - q_2^2p_2)^2 \left( q_1^4(3q_1^2 - 2q_1q_2 + q_2^2)p_1^2 + 2q_1^2q_2^2(q_1^2 + q_2^2)p_1p_2 + q_2^4(q_1^2 - 2q_1q_2 + 3q_2^2)p_2^2 \right)}{4(q_1 - q_2)^2},$$

so that  $\{L, K\} = K^2$ . The corresponding two integrable systems with sextic invariants (1.2-1.4) are determined by functions

$$\begin{aligned} V &= \frac{a(19q_1^4 - 68q_1^3q_2 + 82q_1^2q_2^2 - 68q_1q_2^3 + 19q_2^4)}{q_1^4q_2^4} + \frac{bq_1^2q_2^2}{(q_1 + q_2)^2} + \frac{c(q_1 - q_2)^2}{q_1^2q_2^2}, \\ U_1 &= -\frac{8a(3q_1^2 - 2q_1q_2 + 3q_2^2)(q_1 - q_2)^2}{q_1^4q_2^4} - \frac{c(q_1 - q_2)^2}{q_1^2q_2^2}, \\ U_2 &= -\frac{4a(q_1 + q_2)^2(q_1 - q_2)}{q_1^3q_2^3} - \frac{bq_1^3q_2^3}{(q_1 + q_2)^2(q_1 - q_2)}, \end{aligned}$$

and

$$\begin{aligned} \hat{V} &= \frac{a(q_1^4 - 2q_1^3q_2 - 2q_1^2q_2^2 - 2q_1q_2^3 + q_2^4)}{q_1^4q_2^4} + \frac{b(q_1^2 - 6q_1q_2 + q_2^2)}{4q_1q_2(q_1 - q_2)} + \frac{c(q_1 - q_2)^2}{q_1^2q_2^2}, \\ \hat{U}_1 &= -\frac{a(q_1 + q_2)^2}{q_1^2q_2^2} - \frac{bq_1q_2}{q_1 - q_2}, \\ \hat{U}_2 &= \frac{a(q_1 + q_2)(q_1 - q_2)(q_1^2 + q_2^2)}{q_1^4q_2^4} + \frac{b(q_1 + q_2)}{4q_1q_2} + c\left(\frac{1}{q_1^2} - \frac{1}{q_2^2}\right). \end{aligned}$$

In  $x, y$ -variables potentials have the form

$$V = \frac{16a(16x^4 + 36x^2y^2 + 19y^4)}{x^8} + \frac{bx^4}{4y^2} + \frac{4c(x^2 + y^2)}{x^4}, \quad (2.8)$$

$$\hat{V} = \frac{4a(x^4 + 6x^2y^2 + 4y^4)}{x^8} + \frac{b(2x^2 + y^2)}{2x^2\sqrt{x^2 + y^2}} + \frac{4c(x^2 + y^2)}{x^4}. \quad (2.9)$$

We can prove that the Hamiltonians  $H_1$  (1.2) and  $\hat{H}_1$  (1.4) do not commute with the following polynomial of fourth order in momenta

$$H_3 = L + \sum_{k=0}^4 \sum_{i=0}^k f_{ik}(q_1, q_2) p_1^{k-i} p_2^i,$$

for any functions  $f_{ik}(q_1, q_2)$ . It means that the corresponding Hamiltonian vector fields admit quartic invariants only for geodesic motion at  $V = 0$  and  $\hat{V} = 0$ .

## 2.5 Case $m = 4$ and $k = 3$

In this case, we have

$$T = \frac{q_1^4(3q_1 + q_2)p_1^2}{q_1 - q_2} + \frac{q_2^4(q_1 + 3q_2)p_2^2}{q_2 - q_1}$$

and

$$K = K_1K_2 = \hat{K}_1\hat{K}_2 = \frac{q_1^2q_2^2}{(q_1 - q_2)^2}(q_1^2p_1 - q_2^2p_2)^2(q_1^2p_1 + q_2^2p_2),$$

where

$$K_1 = q_1^2q_2^2(q_1^2p_1 - q_2^2p_2) \quad \text{and} \quad \hat{K}_1 = (q_1^2p_1 + q_2^2p_2).$$

The first decomposition yields the following solutions of (1.3)

$$\begin{aligned} V &= \frac{a(q_1^4 + 4q_1^3q_2 + 4q_1^2q_2^2 + 4q_1q_2^3 + q_2^4)}{q_1^4q_2^4} + \frac{b(q_1^2 + 4q_1q_2 + q_2^2)}{q_1^2q_2^2} + c \left( \frac{1}{q_1} + \frac{1}{q_2} \right), \\ U_1 &= -\frac{a(3q_1^2 + 2q_1q_2 + 3q_2^2)(q_1 - q_2)^2}{4} - bq_2^2q_1^2(q_1 - q_2)^2, \\ U_2 &= \frac{a(q_1 + q_2)(q_1^2 + q_2^2)}{q_1^4q_2^4(q_1 - q_2)} + \frac{b(q_1 + q_2)}{q_1^2q_2^2(q_1 - q_2)} + \frac{c}{q_1q_2(q_1 - q_2)}. \end{aligned}$$

The second decomposition leads to the solutions

$$\begin{aligned} \hat{V} &= \frac{a(q_1^2 + 6q_1q_2 + q_2^2)(q_1 - q_2)^2}{q_1^4q_2^4} + \frac{bq_1^2q_2^2}{(q_1 + q_2)^2} + \frac{c(q_1 - q_2)^2}{q_1^2q_2^2}, \\ \hat{U}_1 &= -\frac{a(q_1 - q_2)^4}{q_1^4q_2^4} + \frac{2bq_1^2q_2^2}{(q_1 + q_2)^2}, \\ \hat{U}_2 &= -\frac{2a(3q_1^2 + 2q_1q_2 + 3q_2^2)}{q_1^2q_2^2} + 2c. \end{aligned}$$

In  $x, y$ -variables potentials have the form

$$V = -\frac{2a(x^4 - 8y^4)}{x^8} - \frac{2b(x^2 - 2y^2)}{x^4} - \frac{2cy}{x^2}, \quad (2.10)$$

$$\hat{V} = -\frac{16a(x^4 - y^4)}{x^8} + \frac{bx^4}{4y^2} + \frac{4c(x^2 + y^2)}{x^4}. \quad (2.11)$$

## 2.6 Case $m = 4$ and $k = -3/5$

Geodesic Hamiltonian

$$T = \frac{q_1^4 \left( q_2 - \frac{3q_1}{5} \right) p_1^2}{q_1 - q_2} + \frac{q_2^4 \left( q_1 - \frac{3q_2}{5} \right) p_2^2}{q_2 - q_1}$$

commutes with reducible polynomial

$$K = \frac{(q_1^2p_1 - q_2^2p_2)(q_1^2p_1 + q_2^2p_2)(q_1^2(5q_1^2 - 10q_1q_2 + q_2^2)p_1 + q_2^2(q_1^2 - 10q_1q_2 + 5q_2^2)p_2)}{q_1q_2(q_1 - q_2)},$$

from which we can extract three different linear polynomials in momenta

$$K_1 = (q_1^2p_1 - q_2^2p_2), \quad \hat{K}_1 = \frac{(q_1^2p_1 + q_2^2p_2)}{q_1q_2(q_1 - q_2)}$$

and

$$\tilde{K}_1 = q_1^2(5q_1^2 - 10q_1q_2 + q_2^2)p_1 + q_2^2(q_1^2 - 10q_1q_2 + 5q_2^2)p_2.$$

The corresponding potentials in (1.2-1.4) are equal to

$$\begin{aligned} V &= \frac{a(19q_1^2 - 26q_1q_2 + 19q_2^2)}{q_1^2q_2^2} + \frac{bq_1^2q_2^2}{(q_1 + q_2)^2}, \\ U_1 &= -\frac{40a(q_1 - q_2)^2}{q_1^2q_2^2}, \\ U_2 &= \frac{5a(5q_1^2 - 6q_1q_2 + 5q_2^2)(q_1 + q_2)^2}{2q_1^3q_2^3(q_1 - q_2)} - \frac{10bq_1q_2(q_1 - q_2)}{(q_1 + q_2)^2} \end{aligned}$$

and

$$\hat{V} = \frac{a(31q_1^4 - 68q_1^3q_2 + 58q_1^2q_2^2 - 68q_1q_2^3 + 31q_2^4)}{q_1q_2(q_1 - q_2)^3} - \frac{b\sqrt{q_1q_2}(7q_1^2 - 18q_1q_2 + 7q_2^2)}{(q_1 - q_2)^{5/2}} + \frac{cq_1q_2}{q_1 - q_2},$$

$$\hat{U}_1 = -\frac{40a(q_1 + q_2)^2}{q_1^3q_2^3(q_1 - q_2)^3} + \frac{40b}{q_1^{3/2}q_2^{3/2}(q_1 - q_2)^{5/2}},$$

$$\hat{U}_2 = \frac{5a(q_1 + q_2)(3q_1 - q_2)^2(q_1 - 3q_2)^2}{q_1q_2(q_1 - q_2)^2} + \frac{5b\sqrt{q_1q_2}(3q_1 - q_2)(q_1 - 3q_2)(q_1 + q_2)}{(q_1 - q_2)^{3/2}} - 5cq_1q_2(q_1 + q_2).$$

Third decomposition  $K = \tilde{K}_1\tilde{K}_2$  yields the third integrable system  $\tilde{H}_1 = T + \tilde{V}$  with potential

$$\tilde{V} = \frac{a(5q_1^2 - 6q_1q_2 + 5q_2^2)}{q_1q_2(q_1 - q_2)}, \quad \tilde{U}_1 = 0, \quad \tilde{U}_2 = \frac{5a(q_1 + q_2)}{q_1q_2}.$$

Thus, in this case, we have three integrable systems with sextic invariants and potentials

$$V = \frac{4a(16x^2 + 19y^2)}{x^4} + \frac{bx^4}{4y^2}, \quad (2.12)$$

$$\hat{V} = \frac{2a(16x^4 + 48x^2y^2 + 31y^4)}{x^2(x^2 + y^2)^{3/2}} + \frac{\sqrt{2}bx(8x^2 + 7y^2)}{2(x^2 + y^2)^{5/4}} - \frac{cx^2}{2\sqrt{x^2 + y^2}}, \quad (2.13)$$

$$\tilde{V} = \frac{2a(4x^2 + 5y^2)}{x^2\sqrt{x^2 + y^2}}. \quad (2.14)$$

## 2.7 Case $m = 4$ and $k = -1/7$

As above geodesic Hamiltonian

$$T = \frac{q_1^4(q_2 - \frac{q_1}{7})p_1^2}{q_1 - q_2} + \frac{q_2^4(q_1 - \frac{q_2}{7})p_2^2}{q_2 - q_1}$$

commutes with reducible cubic polynomial

$$K = \frac{(q_1^2p_1 - q_2^2p_2)(q_1^2(q_1 - 3q_2)p_1 + q_2^2(3q_1 - q_2)p_2)}{q_1q_2(q_1 - q_2)^3} \\ \times \left( q_1^2(3q_1^3 - 27q_1^2q_2 + 33q_1q_2^2 - q_2^3)p_1 + q_2^2(q_1^3 - 33q_1^2q_2 + 27q_1q_2^2 - 3q_2^3)p_2 \right)$$

from which we can extract three different linear polynomials

$$K_1 = q_1^2p_1 - q_2^2p_2, \quad \hat{K}_1 = q_1^2(q_1 - 3q_2)p_1 + q_2^2(3q_1 - q_2)p_2$$

and

$$\tilde{K}_1 = q_1^2(3q_1^3 - 27q_1^2q_2 + 33q_1q_2^2 - q_2^3)p_1 + q_2^2(q_1^3 - 33q_1^2q_2 + 27q_1q_2^2 - 3q_2^3)p_2.$$

The corresponding potentials in (1.2-1.4) are equal to

$$V = \frac{a(11q_1^2 - 42q_1q_2 + 11q_2^2)}{q_1^2q_2^2} + \frac{bq_1^2q_2^2}{(q_1 + q_2)^2}, \quad U_1 = \frac{56a(q_1 - q_2)^2}{q_1^2q_2^2},$$

$$U_2 = \frac{14a(3q_1 - q_2)(q_1 - 3q_2)(q_1 + q_2)^2}{q_1^3q_2^3(q_1 - q_2)} - \frac{28bq_1q_2(q_1^2 - 6q_1q_2 + q_2^2)}{(q_1 + q_2)^2(q_1 - q_2)},$$

and

$$\hat{V} = \frac{a(11q_1^4 - 84q_1^3q_2 + 66q_1^2q_2^2 - 84q_1q_2^3 + 11q_2^4)}{q_1q_2(q_1^2 - 6q_1q_2 + q_2^2)^{3/2}} - \frac{2bq_1q_2}{\sqrt{q_1^2 - 6q_1q_2 + q_2^2}},$$

$$\hat{U}_1 = -\frac{112a(q_1 - q_2)^2(q_1 + q_2)^2}{q_1q_2\sqrt{q_1^2 - 6q_1q_2 + q_2^2}},$$

$$\hat{U}_2 = \frac{14a(q_1 + q_2)(q_1^2 - 14q_1q_2 + q_2^2)^2}{(q_1^2 - 6q_1q_2 + q_2^2)^{3/2}(q_1 - q_2)q_1^2q_2^2} + \frac{28b(q_1 + q_2)}{\sqrt{q_1^2 - 6q_1q_2 + q_2^2}(q_1 - q_2)}.$$

Third decomposition  $K = \tilde{K}_1\tilde{K}_2$  gives rise to the third integrable system so that

$$\tilde{V} = \frac{a(3q_1 - q_2)(q_1 - 3q_2)}{q_1q_2\sqrt{q_1^2 - 6q_1q_2 + q_2^2}}, \quad \tilde{U}_1 = 0, \quad \tilde{U}_2 = \frac{14a(q_1 + q_2)}{q_1^2q_2^2(q_1 - q_2)\sqrt{q_1^2 - 6q_1q_2 + q_2^2}}.$$

In  $x, y$ -variables these potentials have the form

$$V = \frac{4a(16x^2 + 11y^2)}{x^4} + \frac{bx^4}{4y^2}, \quad (2.15)$$

$$\hat{V} = -\frac{2a(16x^4 + 32x^2y^2 + 11y^4)}{x^2(2x^2 + y^2)^{3/2}} + \frac{bx^2}{\sqrt{2x^2 + y^2}}, \quad (2.16)$$

$$\tilde{V} = -\frac{2a(4x^2 + 3y^2)}{x^2\sqrt{2x^2 + y^2}}. \quad (2.17)$$

## 2.8 Case $m = 4$ and $k = 1/5$

In this case geodesic Hamiltonian

$$T = \frac{q_1^4 \left(\frac{q_1}{5} + q_2\right) p_1^2}{q_1 - q_2} + \frac{q_2^4 \left(q_1 + \frac{q_2}{5}\right) p_2^2}{q_2 - q_1}$$

commutes with the cubic and quartic polynomials

$$K = \frac{q_1q_2(q_1^2p_1 - q_2^2p_2) \left( q_1^2(q_1 + 3q_2)p_1 - q_2^2(3q_1 + q_2)p_2 \right)^2}{(q_1 - q_2)^3},$$

and

$$L = \frac{(q_1^2p_1 - q_2^2p_2)^2 \left( q_1^4(q_1^3 + 9q_1^2q_2 + 21q_1q_2^2 + q_2^3)p_1^2 - 2q_1^2q_2^2(q_1 - q_2)(q_1^2 + 4q_1q_2 + q_2^2)p_1p_2 - q_2^4(q_1^3 + 21q_1^2q_2 + 9q_1q_2^2 + q_2^3)p_2^2 \right)}{4(q_1 - q_2)^3}.$$

so that  $\{L, K\} = K^2$ .

This cubic polynomial can be decomposed as  $K = K_1K_2$  or  $K = \hat{K}_1\hat{K}_2$ , where

$$K_1 = (q_1^2p_1 - q_2^2p_2) \quad \text{and} \quad \hat{K}_1 = q_1^2(q_1 + 3q_2)p_1 - q_2^2(3q_1 + q_2)p_2.$$

Integrals of motion (1.2-1.4) involve the following functions

$$V = \frac{a(11q_1^4 + 156q_1^3q_2 + 546q_1^2q_2^2 + 156q_1q_2^3 + 11q_2^4)}{q_1^4q_2^4} + \frac{b(q_1^2 + 6q_1q_2 + q_2^2)}{q_1^2q_2^2} + \frac{cq_1^2q_2^2}{(q_1 + q_2)^2},$$

$$U_1 = \frac{40a(q_1 - q_2)^2(q_1^2 + 10q_1q_2 + q_2^2)}{q_1^4q_2^4} + \frac{5b(q_1 - q_2)^2}{q_1^2q_2^2},$$

$$U_2 = \frac{20a(q_1^2 + 6q_1q_2 + q_2^2)(q_2 + q_1)^2}{q_1^3q_2^3(q_1 - q_2)} - \frac{5cq_1^3q_2^3}{(q_1 - q_2)(q_2 + q_1)^2},$$

and

$$\begin{aligned}\hat{V} &= \frac{a(q_1^4 + 6q_1^3q_2 + 6q_1^2q_2^2 + 6q_1q_2^3 + q_2^4)}{q_1^4q_2^4} + \frac{b(3q_1^2 + 14q_1q_2 + 3q_2^2)}{q_1q_2\sqrt{q_1^2 + 6q_1q_2 + q_2^2}} + \frac{c(q_1^2 + 6q_1q_2 + q_2^2)}{q_1^2q_2^2}, \\ \hat{U}_1 &= -\frac{5a(q_1 - q_2)^2(q_1^2 + 6q_1q_2 + q_2^2)(q_1 + q_2)^2}{q_1^4q_2^4} - \frac{20b(q_1 - q_2)^2\sqrt{q_1^2 + 6q_1q_2 + q_2^2}}{q_1q_2}, \\ \hat{U}_2 &= \frac{5a(q_1 + q_2)}{q_1^2q_2^2(q_1 - q_2)} - \frac{5b(q_1 + q_2)}{2\sqrt{q_1^2 + 6q_1q_2 + q_2^2}(q_1 - q_2)} - \frac{5c(q_1 + q_2)}{2q_1q_2(q_1 - q_2)}.\end{aligned}$$

In  $x, y$ -variables potentials are equal to

$$V = \frac{16a(16x^4 - 28x^2y^2 + 11y^4)}{x^8} - \frac{4b(x^2 - y^2)}{x^4} + \frac{cx^4}{4y^2}, \quad (2.18)$$

$$\hat{V} = -\frac{4a(x^4 + 2x^2y^2 - 4y^4)}{x^8} + \frac{2b(2x^2 - 3y^2)}{x^2\sqrt{y^2 - x^2}} - \frac{4c(x^2 - y^2)}{x^4}. \quad (2.19)$$

We can prove that Hamiltonians  $H_1$  (1.2) and  $\hat{H}_1$  (1.4) do not commute with polynomial of fourth order in momenta

$$H_3 = L + \sum_{k=0}^4 \sum_{i=0}^k f_{ik}(q_1, q_2) p_1^{k-i} p_2^i$$

for any  $f_{ik}(q_1, q_2)$ . As above it means that the corresponding Hamiltonian vector fields admit quartic invariants only for geodesic motion.

## 2.9 Case $m = 4$ and $k = 1/2$

In this case geodesic Hamiltonian

$$T = \frac{q_1^4 \left(\frac{q_1}{2} + q_2\right) p_1^2}{q_1 - q_2} + \frac{q_2^4 \left(q_1 + \frac{q_2}{2}\right) p_2^2}{q_2 - q_1}$$

commute with cubic polynomial

$$K = K_1 K_2 = \hat{K}_1 \hat{K}_2 = \frac{q_1^2 q_2^2 (q_1^2 p_1 - q_2^2 p_2)^2 (q_1^2 (2q_1 + 3q_2) p_1 - q_2^2 (3q_1 + 2q_2) p_2)}{(q_1 - q_2)^3},$$

where

$$K_1 = q_1^2 (2q_1 + 3q_2) p_1 - q_2^2 (3q_1 + 2q_2) p_2, \quad \text{and} \quad \hat{K}_1 = (q_1^2 p_1 - q_2^2 p_2).$$

As a result, we obtain two integrable systems (1.2-1.4) with

$$\begin{aligned}V &= \frac{a(q_1^2 + 3q_1q_2 + q_2^2)(3q_1^2 + 8q_1q_2 + 3q_2^2)}{q_1^4q_2^4} + \frac{b(q_1^2 + 3q_1q_2 + q_2^2)}{q_1^2q_2^2} + \frac{cq_1^2q_2^2}{(q_1 + q_2)^2}, \\ U_1 &= -\frac{2a(q_1 - q_2)^2(q_1^2 + 3q_1q_2 + q_2^2)^2}{q_1^4q_2^4} - \frac{2cq_1^2q_2^2(q_1 - q_2)^2}{(q_1 + q_2)^2}, \\ U_2 &= \frac{2a(q_1^2 + 4q_1q_2 + q_2^2)}{q_1^2q_2^2(q_1 - q_2)} + \frac{2b}{q_1 - q_2},\end{aligned}$$

and

$$\begin{aligned}\hat{V} &= \frac{a(17q_1^4 + 108q_1^3q_2 + 198q_1^2q_2^2 + 108q_1q_2^3 + 17q_2^4)}{q_1^4q_2^4} + \frac{b(7q_1^2 + 18q_1q_2 + 7q_2^2)}{q_1^2q_2^2} + \frac{c(q_1 + q_2)}{q_1q_2}, \\ \hat{U}_1 &= \frac{16a(q_1 - q_2)^2(q_1^2 + 4q_1q_2 + q_2^2)}{q_1^4q_2^4} + \frac{8b(q_1 - q_2)^2}{q_1^2q_2^2}, \\ \hat{U}_2 &= \frac{8a(q_1 + 3q_2)(3q_1 + q_2)(q_1 + q_2)}{q_1^2q_2^2(q_1 - q_2)} + \frac{4b(q_1 + q_2)}{q_1 - q_2} + \frac{2cq_2q_1}{q_1 - q_2}.\end{aligned}$$

In  $x, y$ -variables potentials are equal to

$$V = \frac{2a(x^2 - 4y^2)(x^2 - 6y^2)}{x^8} - \frac{b(x^2 - 4y^2)}{x^4} - \frac{cx^4}{4y^2}, \quad (2.20)$$

$$\hat{V} = \frac{16a(x^4 - 10x^2y^2 + 17y^4)}{x^8} - \frac{4b(x^2 - 7y^2)}{x^4} - \frac{2cy}{x^2}. \quad (2.21)$$

Thus, we present 21 integrable systems with sextic invariants associated with metric (1.1) depending on coordinates.

Other nontrivial families of integrable systems with quartic invariants can be obtained similarly. Indeed, we can consider integrals of motion

$$H_1 = T + V(q_1, q_2), \quad H_2 = \left(K_1^2 + U_1(q_1, q_2)\right)\left(L_2 + U_2(q_1, q_2)\right) \quad (2.22)$$

where  $V$  and  $U_{1,2}$  are solutions of the equation (1.3). For instance, at  $m = 1$  and  $k = 2$  if we substitute Hamiltonian

$$H_1 = \frac{q_1(2q_1 + q_2)p_1^2}{q_1 - q_2} + \frac{q_2(q_1 + 2q_2)p_2^2}{u_2 - u_1} + V(q_1, q_2)$$

and

$$H_2 = \left((q_1^2p_1 - q_2^2p_2)^2 + U_1(q_1, q_2)\right) \left(\frac{(q_1p_1^2 - q_2p_2^2)}{(q_1 - q_2)^3} + U_2(q_1, q_2)\right)$$

into  $\{H_1, H_2\} = 0$  and solve the resulting equation, then we obtain the following solution

$$V = a(q_1 + q_2)(2q_1^2 + q_1q_2 + 2q_2^2) + b(2q_1^2 + 3q_1q_2 + 2q_2^2) + c(q_1 + q_2) + \frac{d(q_1 + q_2)}{q_1q_2} + \frac{e(q_1 + q_2)^3}{q_1^3q_2^3}$$

and integrable Hamiltonian

$$H_1 = \frac{y(p_x^2 + 2p_y^2)}{2} + \frac{xp_xp_y}{2} + 2ay(3x^2 + 8y^2) + b(x^2 + 8y^2) + 2cy - \frac{2dy}{x^2} + \frac{8ey^3}{x^6}$$

depending on five parameters  $a, b, c, d$ , and  $e$ . The corresponding Hamiltonian vector fields has the quartic invariant  $H_2$ .

### 3 Conclusion

In 1910 Darboux studied two-dimensional systems in the Euclidean plane with quadratic invariants [5] and his calculations became one of the foundations of the modern theory of separation of variables in orthogonal curvilinear coordinates.

In 1935 Drach studied two-dimensional systems in the pseudo-Euclidean plane with cubic invariants [6]. Seven of the ten Drach potentials are superintegrable with two quadratic additional integrals [17] and, therefore, cubic Drach invariants exist due to the Abel and Riemann-Roch theorems [20, 21].

In [14] we studied three families of superintegrable two-dimensional systems in the pseudo-Euclidean planes, for which one additional first integral is quadratic, and the second one can be arbitrarily polynomials of a high degree in momenta. Some of these systems were obtained using degenerate deformations of the Poisson brackets but we do not know of a mathematical justification for the existence of these systems till now.

In this note, we start with two-dimensional systems with a position-dependent mass having cubic invariants. Some of these systems are also superintegrable. In contrast with the Drach systems, these cubic invariants can be used to construct systems with sextic invariants. From the technical point of view, it is a consequence of dependence of metrics on the coordinates.

In [18] we obtain a triad of bivectors  $\Pi_{\pm}$  and  $\pi$ , which form a Poisson bivector

$$\Pi' = \frac{\Pi_-}{a} + \pi + a\Pi_+$$

compatible with canonical Poisson bivector  $\Pi$ , so that

$$X = \Pi dH_1 = \Pi' dH_2$$

for a system with sextic invariant at  $m = 4$  and  $k = 3$ . In the future, we plan to study similar triads of bivectors  $\Pi_{\pm}$  and  $\pi$  and the corresponding bi-Hamiltonian vector fields  $X$  for other integrable cases obtained in this note. We hope that this mathematical experiment allows us to get new results for natural Hamiltonian systems with other metrics.

The work was supported by the Russian Science Foundation (project 21-11-00141).

## References

- [1] Ballesteros A., Gutiérrez-Sagredo I., Pedro Naranjo P., *On Hamiltonians with position-dependent mass from Kaluza–Klein compactifications*, Physics Letters A, v. 381 (7), pp. 701-706, 2017.
- [2] Bialy M., Mironov A.E., *A survey on polynomial in momenta integrals for billiard problems*, Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences, v. 376, n.2131, 20170418, 2018.
- [3] da Costa, Bruno G., Ignacio S., *Information-theoretic measures for a position-dependent mass system in an infinite potential well*, Physica A: Statistical Mechanics and its Applications, v. 541, 123698, 2020.
- [4] Darboux G., *Leçons sur la théorie générale des surfaces*, v. 3, Chelsea Publishing 1894.
- [5] Darboux G., *Leçons sur les Systems Orthogonaux et les Coordinées Curvilignes*, Paris: Gauthier-Villars, 1910.
- [6] Drach J., *Sur l'intégration logique et la transformation des équations de la dyanamique á deux Forces conservatives. Intégrales cubiques*, Comptes Rendus, (Paris), v.200, pp.22-26, 1935.
- [7] Fokas S., Lagerström P.A., *Quadratic and cubic invariants in classical mechanics*, J. Math. Anal. Appl., v.74, pp. 325-341, 1980.
- [8] Grigoryev Yu.A., Sozonov A.P., Tsiganov A.V., *Integrability of nonholonomic Heisenberg type systems*, SIGMA, v.12, 112, 14 pp, 2016.
- [9] Grigoriev Yu.A., Tsiganov A. V., *On superintegrable systems separable in Cartesian coordinates* Physics Letters A, v.382, n.32, pp.2092-2096, 2018.
- [10] Kiyohara K., *Two-dimensional geodesic flows having first integrals of higher degree*, Math. Ann., v.320, n.3, pp.487-505, 2001.

- [11] Kozlov, V.V., Denisova, N.V., *Polynomial integrals of geodesic flows on a two-dimensional torus*, Mat. Sb., v.185, n.1, pp.49-64, 1994.
- [12] Kruglikov, B.S., Matveev, V.S., *Nonexistence of an integral of the 6th degree in momenta for the Zipoy-Voorhees metric*, Phys. Rev. D, v.85, 124057, 2012.
- [13] Hietarinta J., *Direct methods for the search of the second invariant*, Phys.Report, v.147, n.2, pp. 87-154, 1987.
- [14] Maciejewski A.J., Przybylska M., Tsiganov A.V., *On algebraic construction of certain integrable and super-integrable systems*, Physica D, v. 240, p.1426-1448, 2011.
- [15] Post S., Winternitz P., *General  $N$ -th order integrals of the motion in the Euclidean plane*, J. Phys. A: Math. Theor., v.48, 405201 (24pp), 2015.
- [16] Rath B., Mallick P., Mohapatra P., Asad J., Shanak H., Jarrar R., *Position-dependent finite symmetric mass harmonic like oscillator: Classical and quantum mechanical study*, Open Physics, v.19, no.1, pp. 266-276, 2021.
- [17] Tsiganov A. V., *On the Drach superintegrable systems*, J. Phys.A., v.33, p.7407-7423, 2000.
- [18] Tsiganov A. V., *New bi-Hamiltonian systems on the plane*, Journal of Mathematical Physics, v.58, 062901, 2017.
- [19] Tsiganov A.V., *Elliptic curve arithmetic and superintegrable systems*, Physica Scripta, v.94, n.8, 085207, 2019.
- [20] Tsiganov A.V., *Discretization and superintegrability all rolled into one*, Nonlinearity, v.33, pp.4924-4939, 2020.
- [21] Tsiganov A.V., *Superintegrable systems and Riemann-Roch theorem*, Journal of Mathematical Physics, v.61, 012701, 2020.
- [22] Tsiganov A.V., *On a time-dependent nonholonomic oscillator*, Russ. J. Math. Phys., v.27, pp.399-409, 2020.
- [23] Valent G., *Superintegrable geodesic flows on the hyperbolic plane*, Preprint arXiv: 2110.02703, 2021.