

Bending of solitonic beams modelled by coupled KMN equation in optical birefringent fibers

Abhik Mukherjee¹

¹*Fundamental Quantum Science Division, Tsung-Dao Lee Institute,
Shanghai Jiao Tong University, Shanghai, Republic of China*

The dynamics of $(2 + 1)$ dimensional optical solitonic beams in birefringent fibers without four wave mixing terms are considered using the recently introduced coupled Kundu Mukherjee Naskar (KMN) equation. The arbitrary bending of solitonic beams in this coupled optical fiber system has been discussed with various two and three dimensional plots. Such analytical results on the bending of optical pulse may pave new directions of research in this field.

I. INTRODUCTION

The upsurge of research in nonlinear fiber optics has started in early seventies leading extensive research on Raman and Brillouin Scattering, optically induced birefringence, parametric four-wave mixing, self phase modulation etc. [1]. An important contribution was made when it was suggested that optical fibers can support soliton like pulses which was experimentally verified in 1980 [1]. After its discovery in nonlinear optics, the dynamics of soliton pulses has been investigated extensively in $(1+1)$ and higher dimensions using both numerical and analytical techniques. The main specialty of soliton, which is the exact solution of integrable systems, is its stability [1]. The amplitude and velocity of solitons remain constant during its propagation. On the other hand, the presence of lossy or time dependent terms in the evolution equation makes it non-integrable causing the velocity of solitary waves to change. Recently, a deep research interest has been developed on some special intricate exact solutions of nonlinear systems like accelerating solitons [7–9], topological solitons [10–14], rogue wave solutions [15, 16] etc.

In the last decade, a new completely integrable $(2+1)$ dimensional nonlinear evolution equation was derived from the basic equations of hydrodynamics by Anjan Kundu, Abhik Mukherjee and Tapan Naskar to describe the dynamics of two-dimensional oceanic rogue wave phenomena [15]. This equation was also re-derived in magnetized plasma system [17] to model nonlinear ion acoustic waves. The detailed exploration of integrable properties and soliton solutions of the equation were also carried out [15, 17, 18]. The equation was derived in nonlinear optical system by Kundu and Naskar in [19] to explain the arbitrary bending phenomena of optical solitonic beam. Such bending is controlled by the boundary population inversion of atoms in an Erbium doped medium. A first order rogue wave solution of this equation was obtained by one fold Darboux Transformation in [20] where the authors have named this new equation as Kundu-Mukherjee-Naskar (KMN) equation. Thereafter, an extensive amount of research has been done on KMN equation exploring its various interesting properties. Many kinds of solutions of KMN equation were derived like higher order rational solutions [21], optical dromions and domain walls [22, 23], various optical soliton solutions having different features [24–35], different analytic solutions [36–43], power series solutions [44], complex wave solutions [45, 46], periodic solutions (via variational principle) [49], solitons in birefringent fiber system [52–54] etc. Integrable properties such as Lax pair, conservation laws, higher order soliton solutions via Hirota method, symmetry analysis, nonlinear self-adjointness property etc. of KMN equation has been explored in [15, 18, 30, 46, 47]. Time fractional KMN equation has been introduced in [50] to explore periodic properties.

One of the important properties of KMN equation is that it supports both bright as well as dark soliton solutions [29] that actually comes from the current like nonlinearity (which depends on both wave envelope and its spatial derivative) present in the equation. This property is different from the standard Nonlinear Schrodinger (NLS) equation which admits only bright and dark soliton solution for focusing and defocusing nonlinearities respectively. In real experimental conditions, the localized wave structures may have various shapes depending on the physical system. Hence, the search for exact solitary wave solutions with variable amplitude or velocity is also an important problem to investigate. In [51], we have derived a special lump, snoidal wave and topological soliton solution that can get curved in $x - y$ plane arbitrarily due to presence of an arbitrary function of space(x) and time(t) in their analytic expressions. Thus, this important bending feature which comes from the Galilean co-variance property and current like nonlinearity present in KMN equation, is rare for a completely integrable equation having constant coefficients.

The interaction of two modes of shorter wavelengths in optical fiber system has been discussed using a system of coupled Nonlinear Schrodinger Equations in $(1+1)$ dimensions [2–4] as well as in higher dimensions [5]. It is also important to state that the single modes in the optical fiber are actually bimodal due to the presence of birefringence [6]. The coupled KMN equation was introduced in [52, 53] to model $(2+1)$ dimensional optical solitons in birefringent fiber without four wave mixing terms. In this paper, our aim is to explore the bending phenomena of optical solitonic beams in coupled KMN system using Hirota bilinearization method which has not been explored till now. Arbitrary

bending light has been discussed in [19] where the boundary value of population inversion of the medium becomes non uniform. On the other hand, we will show that the bending property for coupled KMN system is intrinsic i.e, does not depend on any external function.

The paper is outlined as follows: the coupled KMN equation is introduced in section-II and its Galilean co-variance is derived in section-III. Exact bright and dark solitons are derived using Hirota method in section IV and V respectively. The bending of optical solitonic beams is discussed in section-VI. Conclusive remarks are stated in section-VII followed by acknowledgments and bibliography.

II. COUPLED KMN EQUATION

The Kundu Mukherjee Naskar (KMN) equation which was first derived in modelling oceanic rogue waves [15] is given as

$$iu_t + u_{xy} + 2iu(uu_x^* - u^*u_x) = 0, \quad (1)$$

where the wave profile is given by $u(x, y, t)$ and the subscripts denote partial derivatives w. r. to space (x, y) and time (t) coordinates. The asterisk denotes complex conjugation. The coupled Kundu Mukherjee Naskar (KMN) equation was first introduced by Yildirim [52, 53] in modelling the dynamics of (2+1) dimensional optical solitonic beams in birefringent fibers without four wave mixing terms. In this work, we take the same system of equations taken in [52, 53] with $a_i = 1, b_i = c_i = 2, i = 1, 2$ for simplicity which can be written as :

$$\begin{aligned} iu_t + u_{xy} + 2iu\{(uu_x^* - u^*u_x) + (vv_x^* - v^*v_x)\} &= 0, \\ iv_t + v_{xy} + 2iv\{(vv_x^* - v^*v_x) + (uu_x^* - u^*u_x)\} &= 0, \end{aligned} \quad (2)$$

where $u(x, y, t), v(x, y, t)$ are two wave profiles. Due to the presence of the current like nonlinearities in Eq. (2), it shows few interesting features which will be discussed below.

III. GALILEAN CO-VARIANCE (GC)

The Galilean Co-variance (GC) property of KMN equation has been derived in [51]. Generalizing the same mathematical procedure to (2), we can also find the GC of coupled KMN equation. The equations (2) remains co-variant as

$$\begin{aligned} iU_T + U_{XY} + 2iU\{(UU_X^* - U^*U_X) + (VV_X^* - V^*V_X)\} &= 0, \\ iV_T + V_{XY} + 2iV\{(VV_X^* - V^*V_X) + (UU_X^* - U^*U_X)\} &= 0, \end{aligned} \quad (3)$$

under the Galilean transformation:

$$X = x - at, \quad Y = y, \quad T = t, \quad u(x, y, t) = U(X, Y, T) e^{iaY}, \quad v(x, y, t) = V(X, Y, T) e^{iaY}, \quad (4)$$

where a is the constant frame velocity. This co-variance property is one of the crucial feature for the bending of optical solitonic beams which will be discussed below.

IV. BRIGHT SOLITONS

Optical solitons of the coupled KMN equation (2) has been derived in [52, 53] using trial equation approach and modified simple equation approach. Now using Hirota bilinearization method, we will derive one bright and one dark solitons of (2) which has not been done before as far as our knowledge goes. The Hirota bilinear transformation [2-4] is defined as

$$u(x, y, t) = \frac{g(x, y, t)}{f(x, y, t)}, \quad v(x, y, t) = \frac{h(x, y, t)}{f(x, y, t)}, \quad (5)$$

where g, h are complex functions and f is a real function. Using (5) in (2) we can form the following bilinear equations :

$$\begin{aligned} (iD_t + D_x D_y)(g.f) &= 0, \\ (iD_t + D_x D_y)(h.f) &= 0, \\ D_x(f.f_y) + iD_x(g^*.g) + iD_x(h^*.h) &= 0, \end{aligned} \quad (6)$$

where the Hirota bilinear operator [2] is defined as

$$D_x^m D_t^n g(x, t) \cdot f(x, t) = \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'} \right)^m \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'} \right)^n g(x, t) f(x', t') \Big|_{x=x', t=t'}. \quad (7)$$

Now we will proceed in the standard way to find the bright and dark soliton solutions. In order to find one bright soliton, we assume

$$g = \epsilon g_1, \quad h = \epsilon h_1, \quad f = 1 + \epsilon^2 f_2, \quad (8)$$

where ϵ is the standard expansion parameter and g_1, h_1, f_2 are new variables. Now using (8) in (6), we can get the following differential equations at different orders of ϵ as

$$\begin{aligned} O(\epsilon) : (iD_t + D_x D_y)(g_1 \cdot 1) &= 0, \quad (iD_t + D_x D_y)(h_1 \cdot 1) = 0, \\ O(\epsilon^2) : -f_{2xy} + iD_x(g_1^* \cdot g_1) + iD_x(h_1^* \cdot h_1) &= 0, \\ O(\epsilon^3) : (iD_t + D_x D_y)(g_1 \cdot f_2) &= 0, \quad (iD_t + D_x D_y)(h_1 \cdot f_2) = 0, \\ O(\epsilon^4) : D_x(f_2 \cdot f_{2y}) &= 0. \end{aligned} \quad (9)$$

Solving the systems of equations (9) one can find the solutions as

$$\begin{aligned} g_1(x, y, t) &= \alpha e^{\eta_1}, \quad h_1(x, y, t) = \beta e^{\eta_1}, \quad \eta_1(x, y, t) = kx + my + ikmt + \eta_0, \\ f_2(x, y, t) &= \left\{ \frac{i(k^* - k)}{(k^* + k)(m^* + m)} \right\} \{ |\alpha|^2 + |\beta|^2 \} e^{(\eta_1 + \eta_1^*)}, \quad e^{\delta_1} = \epsilon^{2 \frac{k_I(|\alpha|^2 + |\beta|^2)}{2k_R m_R}} \end{aligned} \quad (10)$$

where $k, m, \eta_0, \alpha, \beta$ are complex constants. Simplifying above expressions, we can obtain the analytical expressions of two modes as

$$\begin{aligned} u &= \frac{\epsilon g_1}{1 + \epsilon^2 f_2} = \mu_1 \operatorname{Sech}[k_R x + m_R y - (k_R m_I + k_I m_R)t] e^{i[k_I x + m_I y + (k_R m_R - k_I m_I)t]}, \\ v &= \frac{\epsilon h_1}{1 + \epsilon^2 f_2} = \mu_2 \operatorname{Sech}[k_R x + m_R y - (k_R m_I + k_I m_R)t] e^{i[k_I x + m_I y + (k_R m_R - k_I m_I)t]}, \\ \text{where,} \quad \mu_1 &= \frac{\alpha}{2} \sqrt{\frac{2k_R m_R}{k_I(|\alpha|^2 + |\beta|^2)}}, \quad \mu_2 = \frac{\beta}{2} \sqrt{\frac{2k_R m_R}{k_I(|\alpha|^2 + |\beta|^2)}}, \quad \alpha = |\alpha| e^{i\theta}, \quad \beta = |\beta| e^{i\phi} \end{aligned} \quad (11)$$

and “R” and “I” in the subscripts denote the real and imaginary parts of the complex constants. In the above simplification, we have chosen the arbitrary constants $\eta_{0R}, \eta_{0I}, \phi, \theta, \delta_1$ in such a way that they get cancelled and we get the expressions (11). The 3 - dimensional plots of the two bright solitonic modes u and v are shown in FIG.1.




FIG. 1: The modulus of the two optical bright solitonic modes u, v respectively are plotted in $x - y$ plane at $t = 0$. The constants are chosen as $k_R = 1, k_I = 2, m_R = 1, m_I = 2, |\alpha| = 1, |\beta| = 2$.

V. DARK SOLITONS

The importance of the KMN model lies in the fact that it supports both bright and dark soliton solutions. In case of NLSE with Kerr like nonlinearity, bright and dark solitons occur for focusing and defocusing nonlinearities respectively. In order to find one dark solitons of (2) we assume

$$g = g_0(1 + \epsilon g_1), \quad h = h_0(1 + \epsilon h_1), \quad f = 1 + \epsilon f_1, \quad (12)$$

where ϵ is the same expansion parameter as discussed before. Thus using this expansion (12) in (6) we would get the following differential equations at different orders of ϵ .

At $O(\epsilon^0)$, we would get

$$(iD_t + D_x D_y)(g_0,1) = 0, (iD_t + D_x D_y)(h_0,1) = 0, D_x(g_0^*.g_0) + D_x(h_0^*.h_0) = 0. \quad (13)$$

In order to solve (13), we assume

$$g_0 = \tau_1 e^{i\psi_1}, h_0 = \tau_2 e^{i\psi_2}, \psi_i = k_i x + m_i y + n_i t + \psi_i^0, \quad (14)$$

with $i = 1, 2$, where ψ_i s are real and τ_i s are complex quantities. Solving (13), we would get

$$n_1 = -k_1 m_1, n_2 = -k_2 m_2, k_1 |\tau_1|^2 + k_2 |\tau_2|^2 = 0. \quad (15)$$

At $O(\epsilon)$, we would get the following differential equations :

$$\begin{aligned} (g_{1t} - f_{1t}) + k_1(g_{1y} - f_{1y}) + m_1(g_{1x} - f_{1x}) - i(g_{1xy} + f_{1xy}) &= 0, \\ (h_{1t} - f_{1t}) + k_2(h_{1y} - f_{1y}) + m_2(h_{1x} - f_{1x}) - i(h_{1xy} + f_{1xy}) &= 0, \\ f_{1xy} = 2k_1 |\tau_1|^2 (g_1 + g_1^* - h_1 - h_1^*) - i|\tau_1|^2 (g_{1x} - g_{1x}^*) - i|\tau_2|^2 (h_{1x} - h_{1x}^*). \end{aligned} \quad (16)$$

Equations (16) admit the following solutions :

$$g_1 = Z_g e^{\xi_1}, h_1 = Z_h e^{\xi_1}, f_1 = e^{\xi_1}, \xi_1 = P_1 x + Q_1 y + R_1 t + \xi_1^0, \quad (17)$$

where Z_g, Z_h are complex constants and P_1, Q_1, R_1, ξ_1^0 are real constants that are related by following constraints :

$$\begin{aligned} Z_g &= \frac{[(R_1 + k_1 Q_1 + m_1 P_1) + i P_1 Q_1]^2}{[(R_1 + k_1 Q_1 + m_1 P_1)^2 + P_1^2 Q_1^2]}, Z_h = \frac{[(R_1 + k_2 Q_1 + m_2 P_1) + i P_1 Q_1]^2}{[(R_1 + k_2 Q_1 + m_2 P_1)^2 + P_1^2 Q_1^2]}, \\ Q_1 &= \left(\frac{2k_1}{P_1}\right) |\tau_1|^2 (Z_g + Z_g^* - Z_h - Z_h^*) - i|\tau_1|^2 (Z_g - Z_g^*) - i|\tau_2|^2 (Z_h - Z_h^*). \end{aligned} \quad (18)$$

Absorbing ϵ in e^{ξ_1} , the one dark solitons can be expressed as

$$\begin{aligned} u &= (\tau_1/2) e^{i\psi_1} [(1 + Z_g) - (1 - Z_g) \tanh(\xi_1/2)], \\ v &= (\tau_2/2) e^{i\psi_2} [(1 + Z_h) - (1 - Z_h) \tanh(\xi_1/2)]. \end{aligned} \quad (19)$$

The 3 - dimensional plots of the two dark solitonic modes u and v are shown in FIG.2.




FIG. 2: The modulus of the two optical dark solitonic modes u, v respectively are plotted in $x - y$ plane at $t = 0$. The constants are chosen as $k_1 = 1, k_2 = -1, P_1 = 1, Q_1 = 2, R_1 = 2, m_2 = 2, m_1 = \sqrt{5} - 5, \tau_1 = \frac{1+i}{\sqrt{2}}, \tau_2 = \frac{1-i}{\sqrt{2}}$.

VI. BENDING OF OPTICAL SOLITONIC BEAMS

One important property of KMN model (1) is that it processes curved solitons that is explored in [51]. Due to the presence of the current like nonlinearity and Galilean co-variance in (1), this special dark soliton can get curved arbitrarily in $x - y$ plane. In [51], one curved soliton has been derived using ansatz approach. Now, we will rederive such curved one soliton solution of Eq. (1) using Hirota bilinearization method that has not been done before.

A. Derivation of one curved dark soliton solution of KMN equation (1) via Hirota bilinearization method

Here, we will derive one curved dark soliton solution of KMN equation (1) via Hirota bilinearization technique. We consider the KMN equation in the Galilean frame ($X - Y - T$) that is discussed in section - III as

$$iU_T + U_{XY} + 2iU(UU_X^* - U^*U_X) = 0, \quad (20)$$

where $X = x - at$, $Y = y$, $T = t$, $u(x, y, t) = U(X, Y, T)e^{iaY}$. We use the bilinear transformation as

$$U(X, Y, T) = \frac{G(X, Y, T)}{F(X, Y, T)}, \quad (21)$$

where G, F are complex and real functions respectively, to get the pair of bilinear equations :

$$\begin{aligned} [iD_T + D_X D_Y] (G.F) &= 0, \\ D_X [(F.F_Y) + i (G^*.G)] &= 0. \end{aligned} \quad (22)$$

In order to find one dark curved soliton of (20) we assume

$$G = G_0(1 + \epsilon G_1), \quad F = 1 + \epsilon F_1, \quad (23)$$

where ϵ is the same expansion parameter as discussed before. At this stage, we would assume that G, F are independent of time T . That will not make the solution static, the reason will be explored later. Thus using this expansion (23) in (22) we would get the following differential equations at different orders of ϵ .

At $O(\epsilon^0)$, we would get

$$G_{0XY} = 0, \quad (G_{0X}^* G_0 - G_0^* G_{0X}) = 0 \quad (24)$$

In order to solve (24), we assume

$$G_0 = \tau e^{i\psi}, \quad \psi = kX + mY + \psi^0, \quad (25)$$

where k, m, ψ^0 are real constants and τ is complex constant. Solving (24), we would get $k = 0$. The the solution is

$$G_0 = \tau e^{i(mY + \psi^0)}. \quad (26)$$

At $O(\epsilon)$, we would get

$$\begin{aligned} G_{1XY} + F_{1XY} + im(G_{1X} - F_{1X}) &= 0, \\ F_{1XY} = i|G_0|^2(G_{1X}^* - G_{1X}). \end{aligned} \quad (27)$$

Equations (27) admit the following solutions :

$$G_1 = Z_g e^\xi, \quad F_1 = e^\xi, \quad \xi = A(X) + m_1 Y + \xi^0, \quad (28)$$

where Z_g is complex constant and m_1, ξ^0 are real constants. We have introduced an arbitrary real function $A(X)$ which will ultimately cause the bending of solitons. The constants are related as

$$Z_g = \frac{im - m_1}{im + m_1}, \quad |\tau|^2 = \frac{(m_1^2 + m^2)}{4m}. \quad (29)$$

If the function $A(X)$ is nonlinear in X , the soliton will bend in $X - Y$ plane making curvature. The final soliton solution can be written as

$$U = \frac{\tau}{2} [(1 + Z_g) - (1 - Z_g) \tanh(\xi/2)] e^{i(mY)}. \quad (30)$$

Now moving back to the original frame $x - y - t$ and old variable u , we would get

$$\begin{aligned} u &= \frac{\tau}{2} [(1 + Z_g) - (1 - Z_g) \tanh(\xi/2)] e^{i(my + ay)}, \quad \xi = A(X) + m_1 y, \quad X = x - at. \\ Z_g &= \frac{im - m_1}{im + m_1}, \quad \tau = (m + im_1) / \sqrt{4m}. \end{aligned} \quad (31)$$

Hence the soliton bends due to the presence of $A(X)$, which is a nonlinear function of $X = (x - at)$. For the different nonlinear choices of $A(X)$, $X = x - at$, we would get different types of bending of solitons which is discussed in [51].

B. Curved solitons for coupled KMN equations

Now we will concentrate on the main findings of this work. Similar to the previous case, we consider the coupled KMN system (3) in the Galilean frame $X - Y - T$ and take $U = G/F$, $V = H/F$, where H, G, F are independent of T . The functions G, H are complex and F is a real function.

Expanding as :

$$G = G_0(1 + \epsilon G_1), \quad H = H_0(1 + \epsilon H_1), \quad f = 1 + \epsilon F_1, \quad (32)$$

where ϵ is the same expansion parameter as discussed before, we would get the following differential equations at different orders of ϵ .

At $O(\epsilon^0)$, we would get

$$G_{0XY} = 0, \quad H_{0XY} = 0, \quad (G_{0X}^* G_0 - G_0^* G_{0X}) + (H_{0X}^* H_0 - H_0^* H_{0X}) = 0 \quad (33)$$

with following solutions

$$G_0 = \tau_1 e^{i(m_1 Y + \psi_1^0)}, \quad H_0 = \tau_2 e^{i(m_2 Y + \psi_2^0)}, \quad (34)$$

where $m_{1,2}, \psi_{1,2}^0$ are real constants and $\tau_{1,2}$ are complex constants.

At $O(\epsilon)$, we would get the following differential equations :

$$\begin{aligned} G_{1XY} + F_{1XY} + im_1(G_{1X} - F_{1X}) &= 0, \quad H_{1XY} + F_{1XY} + im_2(H_{1X} - F_{1X}) = 0 \\ F_{1XY} &= i|\tau_1|^2(G_{1X}^* - G_{1X}) + i|\tau_2|^2(H_{1X}^* - H_{1X}) \end{aligned} \quad (35)$$

with the solutions :

$$G_1 = Z_g e^{\xi_1}, \quad H_1 = Z_h e^{\xi_1}, \quad F_1 = e^{\xi_1}, \quad \xi_1 = B(X) + PY + \xi_1^0, \quad (36)$$

where Z_g, Z_h are complex constants and P, ξ_1^0 are real constants that are related by following constraints :

$$Z_g = \frac{im_1 - P}{im_1 + P}, \quad Z_h = \frac{im_2 - P}{im_2 + P}, \quad P = 2(Z_{gI}|\tau_1|^2 + Z_{hI}|\tau_2|^2), \quad (37)$$

where I in the subscript denotes the imaginary part. The function $B(X)$ is the arbitrary function of $X = (x - at)$. For different nonlinear choices of $B(X)$, we would get different kinds of bending of u, v . Absorbing ϵ in ξ_1^0 , the one dark solitons in the old frame $x - y - t$ can be expressed as

$$\begin{aligned} u &= \frac{\tau_1}{2} [(Z_g - 1) \tanh \frac{\xi_1}{2} + (Z_g + 1)] e^{i(m_1 y + ay + \psi_{10})}, \quad v = \frac{\tau_2}{2} [(Z_h - 1) \tanh \frac{\xi_1}{2} + (Z_h + 1)] e^{i(m_2 y + ay + \psi_{20})}, \\ Z_g &= \frac{im_1 - P}{im_1 + P}, \quad Z_h = \frac{im_2 - P}{im_2 + P}, \quad P = 2(Z_{gI}|\tau_1|^2 + Z_{hI}|\tau_2|^2), \quad \xi_1 = B(X) + PY + \xi_1^0, \quad X = x - at. \end{aligned} \quad (38)$$

The bending of such dark solitons u, v are shown as contour plots in $x - y$ plane for different choices of $B(X)$ in FIG. 3, 4, 5 respectively.

Thus we can form different shapes of the dark soliton solutions (38) by choosing different functional forms of the arbitrary function $B(X)$, $X = x - at$ present in the argument of the \tanh function. This property may be useful in modeling the bending of light beam [19] in nonlinear optics. Previously, the bending of an optical beam was shown in [55, 56] through the Airy function solution of linear Schrodinger equation. It can preserve its parabolic form over a finite distance. The beam acceleration along an arbitrary curve was shown in [19] by Kundu and Naskar where the shape was not preserved. Such bending is controlled by the boundary population inversion of atoms in an Erbium doped medium. In that situation, the boundary value of population inversion of the medium becomes non uniform. On the other hand, the bending feature of this work is more general and arbitrary. Due to the presence of the arbitrary function $B(X)$, any shape can be obtained which is important for Mathematical modelling of experimental data. Also, the amplitude of the soliton remains constant. Such property, that is the existence of the arbitrary function $B(X)$ causing arbitrary bending, originates from the current like nonlinearity and Galilean Co-variance of the coupled KMN system (2). It is a very rare property which is absent in NLSE with Kerr like nonlinearity.

Now it is important to discuss the condition for larger bending, it would be interesting to derive the amount of curvature due to function $B(X)$. For the static case ($t = 0$), the locus of the maximum amplitude of the curved soliton solution (38) is of the form

$$B(x) + Py + \xi_1^0 = 0. \quad (39)$$




FIG. 3: The contour plots of the curved dark solitons u and v given by (38) in $x - y$ plane at $t = 0$ for the choice of $B(X) = X^2 = (x - at)^2$. The constants are chosen as $m_1 = 1, m_2 = 2, a = P = 1, |\tau_1|^2 = 1/10, |\tau_2|^2 = 1/2$. Due to the presence of nonlinear function $B(X)$ in the argument, the solitons (38) get bend in $x - y$ plane.




FIG. 4: The contour plots of the curved dark solitons u and v given by (38) in $x - y$ plane at $t = 0$ for the choice of $B(X) = \sin X = \sin(x - at)$. The constants are chosen as $m_1 = 1, m_2 = 2, a = P = 1, |\tau_1|^2 = 1/10, |\tau_2|^2 = 1/2$. Due to the presence of nonlinear function $B(X)$ in the argument, the solitons (38) get bend in $x - y$ plane.

Now taking the derivative with respect to x twice, we get

$$\frac{dS}{dx} = -\frac{d^2B}{dx^2}. \quad (40)$$

where the slope is defined as $S = \frac{dy}{dx}$. We see from (40) that for a higher value of $-\frac{d^2B}{dx^2}$, the rate of variation of the slope S is higher. Hence, the slope of the maximum amplitude curve will vary large for traveling the unit distance in x . Larger rate of variation of S describes larger bending. Hence, for large bending to take place, $-\frac{d^2B}{dx^2}$ should be high which can be seen from FIG. 3,4,5 respectively.




FIG. 5: The contour plots of the curved dark solitons u and v given by (38) in $x - y$ plane at $t = 0$ for the choice of $B(X) = X^3 = (x - at)^3$. The constants are chosen as $m_1 = 1, m_2 = 2, a = P = 1, |\tau_1|^2 = 1/10, |\tau_2|^2 = 1/2$. Due to the presence of nonlinear function $B(X)$ in the argument, the solitons (38) get bend in $x - y$ plane.

VII. CONCLUSIVE REMARKS

In this work, we have considered the coupled Kundu Mukherjee Naskar system which was introduced by Yildirim in [52, 53] to model (2+1) dimensional optical solitons in birefringence fiber without four wave mixing terms. The Galilean co-variance property of the system has been derived. The dark and bright one solitons of the system has been calculated using Hirota bilinearization method. It has been shown that the dark solitons of the two modes u and v can bend arbitrarily due to the presence of an arbitrary function of space (x) and time (t) which is shown by the contour plots. Such arbitrary curvature originates from the current like nonlinearity and Galilean co-variance present in the system (2). The condition of larger bending has been derived and explained. Such analytical solutions may pave new directions of research in the field of Nonlinear Optics.

VIII. ACKNOWLEDGMENTS

The work is sponsored by Shanghai Pujiang Program and Tsung Dao Lee Institute starting grant. Author is indebted to Ms. M. Mukherjee for the thoughtful discussions and unconditional support during the progress of the work.

IX. REFERENCES

-
- [1] G.P Agarwal, Nonlinear Fiber Optics, Academic Press, USA, 2007.
 - [2] R. Radhakrishnan, M. Lakshmanan, J. Phys. A : Math.Gen. 28 (1995) 2683.
 - [3] R. Radhakrishnan, M. Lakshmanan, J. Hietarinta, Phys. Rev. E 56 (1997) 2213.
 - [4] R. Radhakrishnan, K. Aravintan, J. Phys. A : Math. Theor. 40 (2007) 13023.
 - [5] X. Liu et.al, Journal of Advanced Research 24 (2020) 167.
 - [6] C. R. Menyuk, IEEE Journal of Quantum Electronics 23 (1987) 174.
 - [7] A.V.Gorbach, D.V Skryabin, Phys.Rev.A 76 (2007) 053803.
 - [8] A. Mukherjee, A. Kundu, Phys Lett A 383 (2019) 985
 - [9] Zaqlao, Nonlinear Dynamics 99 (2020) 2945
 - [10] M.O.Katanaev, Theoretical and Mathematical Physics 132(2) (2004) 163.
 - [11] A. Biswas, D. Milovic, Int. J Theor. Phys 48 (2009) 1104.
 - [12] A. Biswas, Commun Nonlinear Sci Numer Simulat 14 (2009) 2845.

- [13] B. Sturdevant, D. A. Lott, A. Biswas, Progress In Electromagnetics Research Letters 10 (2009) 69.
- [14] M. Mirzazadeh, M. Eslami, D. Milovic, A. Biswas, Optik, 125 (2014) 5480
- [15] A.Kundu, A.Mukherjee, T.Naskar, Proc. Soc. A 470 (2014) 20130576.
- [16] C.Kharif, E.Pelinovsky, A.Slunyaev, Rogue Waves in Ocean, Berlin, Germany, Springer, 2009.
- [17] A. Mukherjee, M. S. Janaki, and A. Kundu, Phys Plasmas 22 (2015) 072302.
- [18] R. Constantinescu, A Florian, Symmetry 12 (2020) 1819.
- [19] A. Kundu, T. Naskar, Physica D 276 (2014) 21
- [20] D. Qiu, Y. Zhang, J. He, Commun Nonlinear Sci Numer Simulat 30 (2016) 307-315.
- [21] X. Wen, Proc Romanian Acad A, 18 (2017) 191-198.
- [22] A. Biswas et.al, Results in Physics 16 (2020) 102850.
- [23] O.G. Gaxiola et.al, Regular and Chaotic Dynamics 25 (2020) 338.
- [24] M. Ekici, A. Sonmezoglu, A. Biswas, M. R. Belice, Chin J Phys 57 (2019) 72-77.
- [25] Y. Yildirim, Optik, 183 (2019) 1061.
- [26] Y. Yildirim, Optik, 184 (2019) 247.
- [27] S.T.R. Rizvi, I. Afzal, K. Ali, Mod Phys Lett B 34 (2020) 2050074-1.
- [28] T.A Sulaiman, H. Bulut, Applied Mathematics and Nonlinear sciences 4(2)(2019)513
- [29] S.Singh, A. Mukherjee, K.Sakkaravarthi, K.Murugesan, Waves in Random and Complex media, (<https://doi.org/10.1080/17455030.2021.1874621>)
- [30] S.Singh, R. Sakthivel, M.Inc, A. Yusuf, K.Murugesan, Mod Phys Lett B 34 (2020) 2050068.
- [31] A.I. Aliyu, Y. Li, D. Baleanu, Chin J Phys 63 (2020) 410.
- [32] H. Rezazadeh et.al, Optical and Quantum Electronics 53 (2021) 317.
- [33] M. Bilal, S.U. Rehman, J. Ahmed, Optical and Quantum Electronics 53 (2021) 283.
- [34] H. Triki et.al, Physics Letters A 401 (2021) 127341.
- [35] E.M.E Zayed, R. Shohib, M.E.M Alngar, Optical and Quantum Electronics (under review).
- [36] R.A Talarposhti et.al, Int. Journal of Modern Physics B, 34 (2020) 2050102
- [37] H.Gunerhan, Mod Phys Lett B (2020) 2050225-1.
- [38] D. Kumar et.al, Physica Scripta 96 (2021) 025218.
- [39] G. Dang, Demonstratio Mathematica 54 (2021) 129.
- [40] S.S. Singh, International Journal of Physical Research 8(2)(2020)40.
- [41] H.M Baskonus et.al, Modern Physics Letters B 35 (2021) 2150217.
- [42] M. Ren, Y. Zhou, Q. Liu, arxiv : 2105.08538v1.
- [43] N.A. Kudryashov, Optik 186 (2019) 22.
- [44] W.Q. Peng, S. F. Tian, T.T Zhang, Modern Physics Letters B 32 (2018) 1850336
- [45] A. Jhangeer et.al, Results in Physics 16 (2020) 102816.
- [46] R. Cimpoiasu et.al, Results in Physics, 28 (2021) 104583.
- [47] S. Kumar, S. Malik, A. Biswas, Chaos Solitons and Fractals 137 (2020) 109855.
- [48] N. A. Kudryashov, Optik 186 (2019) 22-27.
- [49] J. H. He, Results in Physics 17 (2020) 103031.
- [50] J.H. He, Y.O.El-Dib, Results in Physics 19 (2020) 103345.
- [51] A. Mukherjee, Optik 219 (2020) 165194.
- [52] Y. Yildirim, Optik 184 (2019) 121.
- [53] Y. Yildirim, Optik, 183 (2019) 1026.
- [54] Y. Yildirim, M. Mirzazadeh, Chin J Phys 64 (2020) 183.
- [55] G.A. Siviloglou, D.N. Christodoulides, Opt. Lett. 32 (2007) 979.
- [56] G.A. Siviloglou, J. Broky, A. Dogariu, D.N. Chistodoulides, Phys. Rev. Lett. 99 (2007) 213901.