

# Relative subsystems and quantum reference frame transformations

Esteban Castro-Ruiz<sup>1,2,3</sup> and Ognjan Oreshkov<sup>1</sup>

<sup>1</sup>*QuIC, Ecole polytechnique de Bruxelles, C.P. 165,  
Université libre de Bruxelles, 1050 Brussels, Belgium*

<sup>2</sup>*Institute for Quantum Optics and Quantum Information (IQOQI),  
Austrian Academy of Sciences, Boltzmannngasse 3, A-1090 Vienna, Austria*

<sup>3</sup>*Institute for Theoretical Physics, ETH Zurich, Switzerland*

Recently, there has been much effort in developing a quantum generalisation of reference frame transformations. Despite important progress, a complete understanding of their principles and physical meaning is still lacking. Here, we develop a framework for quantum reference frame transformations that focuses on the algebra of relative observables between a system and a reference frame. We show that this perspective sheds light on important conceptual issues regarding reference frame changes. In our framework, a quantum reference frame viewpoint is a preferred partition of the full invariant system into subsystems. A transformation between quantum reference frames is a specific change of preferred partition. Interestingly, the full invariant system contains not only the algebra of relative observables between the system and frame but also an “extra particle,” related to the invariant degrees of freedom of the frame itself. The extra particle contains information about the “quantumness” of the frame and is essential to the unitarity of the transformations. Our approach can be applied to a vast set of symmetry groups and systems, among which we study the centrally extended Galilei group as a particular case.

## I. INTRODUCTION

Transformations between reference frames play a crucial role in our understanding of physical processes. In practice, reference frames are realised by physical systems, which are standardly treated as classical. However, assuming that every physical system is ultimately quantum, it is interesting to ask how a theory of transformations with respect to quantum reference frames (QRFs) would look like, and what implications it would have for our description of the physical world.

The study of QRFs is broad in scope. It includes the connection between QRFs and superselection rules [1–5], the study of quantum mechanics with respect to finite-mass QRFs [6–9], quantum tasks and operations under symmetry constraints [3, 4, 10–22],

QRFs as resources of asymmetry [23–30], and QRFs as a means to define physical observables in quantum gravity [31–36].

Recently, attention has turned towards understanding transformations between QRFs [37–54]. Given the description of a physical process with respect to a QRF A, how do we obtain the description from the point of view of QRF B? A formalism for QRF transformations was introduced in Ref. [37], showing that entanglement and superposition are frame-dependent notions. Subsequently, Ref. [39] showed that some transformations of [37] can be obtained by imposing a symmetry constraint on a "perspective neutral" state. The perspective-neutral point of view was further developed in [40–42, 44–46]. A group-theoretic treatment of QRFs and their transformations was introduced in [49], where reference frames are associated with symmetry groups. These recent developments suggest that a precise formulation and answer to the question of how to transform between the descriptions of different QRFs has the potential to generalise the notion of symmetry and covariance [37, 39, 48, 50, 51]. It can also provide an operational understanding of spin for relativistic particles [38, 47], lead to a new understanding of the physics of gravitating quantum systems [43, 54, 57], and to quantum extensions of the general relativistic equivalence principle [55–57].

Despite the important progress done in this line of research, it is safe to say that the principles and interpretation of "jumping" from one quantum reference frame to another are not yet fully understood. Given the description of a physical process with respect to QRF a A, is there a way to *derive*, using standard quantum mechanical arguments as applied to an external observer, what is "seen" from the perspective of another QRF B? When we talk about "the system" from the point of view of A and that of B, do we refer to the same degrees of freedom? If not, how are these degrees of freedom related to each other? Are the systems considered in a QRF scenario supposed to include (perhaps a coarse graining of) all the degrees of freedom of the universe, or are we allowed to assume that there could be other degrees of freedom out there, which can potentially act as reference frames? In the latter case, is it possible to have a framework for quantum reference frame transformations that captures this possibility in its most general way?

Here, we develop a framework for QRF transformations that focuses on the relations between algebras of relative observables, defined in connection to a symmetry group. Our framework leads naturally to a "mixed-state approach" to quantum reference frames, where symmetry invariance of states and operators is formulated as conjugation with unitary representations of the group. That is, we focus on invariance under transformations of the form  $\rho \rightarrow U(g)\rho U(g)^\dagger$ , rather than invariance of pure states under multiplication of the form  $|\psi\rangle \rightarrow U(g)|\psi\rangle$ . We argue that this approach is better suited to deal with questions like the ones posed above. On the one hand, it clearly identifies the relevant

quantum subsystems from the point of view of A and B, clarifying the QRF-dependence of superpositions and entanglement. On the other hand, the mixed-state approach leads to a more general notion of symmetry transformations. Being broader than invariance under multiplication, invariance under conjugation leads to the possibility of applying QRF transformations to subsystems of the whole universe in full generality.

In our framework, a QRF viewpoint is simply a specific partition of the invariant algebra into subsystems, as defined by a set of subalgebras, and the relation between different QRFs can be viewed as a network of subalgebras partitioning the full invariant space. Essential to this network is the algebra of an "extra particle," which emerges as a consequence of the invariant degrees of freedom of the reference frame. We argue that the extra particle should be included in the relative description of quantum systems in a standard way. The reason why its importance has not been noticed so far is that we normally deal with sharply-defined, classical reference frames, for which, as we show, the extra particle is always in a maximally mixed state. However, when considering general QRFs, the extra particle should be included, because it is essential for obtaining reversible QRF transformations.

As an illustration of the physical meaning of our framework, we analyse quantum reference frame transformations with respect to the (centrally extended) Galilei group.

## II. MODELLING A QUANTUM REFERENCE FRAME

Let us now introduce the basic ingredients of our framework. In particular, we define the notion of quantum reference frame that we will use throughout this work.

Consider the situation of Fig. 1. An observer, Alice, possesses a reference frame A, associated with the symmetry group  $G$ . She uses it to perform quantum operations on a system S, which transforms under some unitary representation of  $G$ . We treat both A and S quantum mechanically. To do this, we imagine an external observer, Eve, with a reference frame E, who has full access to both systems. Eve assigns a Hilbert space to the composite system

$$\mathcal{H}_{AS|E} = \mathcal{H}_{A|E} \otimes \mathcal{H}_{S|E}. \quad (1)$$

The reason for the notation  $|E$  in Eq. (1) is that the quantum mechanical description of A and S is defined with respect to Eve. In the remaining of this section, we will omit this label, as we will be concerned with Eve's description only. However, in section III, this point will be important and we shall introduce the notation again to distinguish it from the "internal" perspective of Alice, who has only access to operators that are invariant under the action of  $G$ . Eventually, we will do away with the external observer by con-

sidering only operators living in the invariant subspace. At this level of description, Eve regards the degrees of freedom of Alice’s measurement apparatus (and Alice herself) as implicit. They lie on the “other side” of Heisenberg’s cut. If desired, the cut can be moved to include such degrees of freedom explicitly.




FIG. 1. In our setup, an observer, Alice, has only access to degrees of freedom that are invariant under the action of the group  $G$ . The latter is assumed defined relative to some external observer. As we show in Section III, the invariant degrees of freedom are independent of any external observer or reference frame. These invariant degrees of freedom include, in particular, degrees of freedom of the system defined relative to the reference frame  $A$ . The latter are described by operators that form an algebra, called  $S|A$ . Importantly, Alice’s apparatus, by means of which these degrees of freedom are accessed are *not* part of the quantum system under consideration. They lie on the “other side” of Heisenberg’s cut.

To make contact with the standard situation in quantum mechanics, where reference frames are assumed to be classical and are treated implicitly, we assume that the QRF  $A$  is perfect. That is, it can be prepared in a basis of states that break the symmetry of  $G$  maximally [3]. Therefore, the Hilbert space of  $A$ ,  $\mathcal{H}_A$ , is the span of a fully distinguishable basis of “classical” states labeled by group elements,  $|g\rangle_A$ . Because basis states are fully distinguishable, we have  $\langle g|g'\rangle = \delta(g^{-1}g')$ . Here,  $\delta(g)$  denotes the Dirac delta distribution for continuous groups, where the group identity element  $e$  plays the role of the real number 0, or the (single-argument) Kronecker delta for discrete groups. Thus,  $\mathcal{H}_A$  consists of square-integrable functions on  $G$  with respect to the invariant measure  $dg$ . (In this work, we consider only unimodular groups, that is, groups for which the left-invariant and the

right-invariant measure are the same.)  $\mathcal{H}_A$  carries the left- and right-regular representations of  $G$ . The left-regular representation,  $L_A$ , acts as  $L_A(g) |g'\rangle_A = |gg'\rangle_A$  for all  $g$  and  $g'$  in  $G$ . The right-regular representation,  $R_A$ , acts as  $R_A(g) |g'\rangle_A = |g'g^{-1}\rangle_A$  for all  $g$  and  $g'$  in  $G$ . Both  $L_A$  and  $R_A$  are unitary representations. The only assumption we make on  $S$  is that it transforms under a unitary representation,  $U_S$ , of  $G$ . Mathematically, this setup closely resembles that of Ref. [4], where the regular representation is used as a token in a quantum communication scheme.

Importantly, the regular representation is highly reducible — it contains all irreducible representations (irreps) of the group. We can write the basis states  $|g\rangle_A$  as [2]

$$|g\rangle_A = \int dq dx dy \sqrt{\frac{\dim(q)}{|G|}} D_{xy}^{(q)}(g) |q; x, y\rangle_A, \quad (2)$$

where  $q$  is the "charge" labelling a specific irrep. For compact groups,  $\dim(q)$  denotes the dimension of the irrep labeled by  $q$ , and  $|G|$  denotes the order of  $G$ . The complex numbers  $D_{xy}^{(q)}(g)$  are matrix elements of the irrep  $q$  for  $g \in G$ . The left-regular representation  $L_A(g)$  acts on the "colour" degrees of freedom, labeled by  $x$ , whereas the right-regular representation  $R_A(g)$  acts on the "flavour" or multiplicity degrees of freedom, labeled by  $y$  [2]. For the regular representation, the dimension of the multiplicity degrees of freedom for a given irrep  $q$  equals the dimension of  $q$ .

Although Eq. (2) is written under the assumption that both  $\dim(q)$  and  $|G|$  are finite, a similar equation holds more generally, not only for compact groups. For example, Eq. (2) reduces to the well-known Fourier transform relation between position eigenvectors  $|x\rangle$  and momentum eigenvectors  $|p\rangle$ :  $|x\rangle = (1/\sqrt{2\pi}) \int dp \exp(-ipx) |p\rangle$ . As we will see in Section V, Eq. (2) is useful in the case of the centrally extended Galilei group, where the quotient  $\dim(q)/|G|$  is replaced by the mass parameter,  $m$ , labeling the irrep.

For an example in the case of compact groups, suppose  $G$  is the rotation group  $SU(2)$ . In this case,  $q$  corresponds to the total angular momentum, and the integral with respect to  $q$  is replaced by a sum that runs over all values of total angular momentum, or equivalently, all irreps of  $SU(2)$ . As the labels  $x$  and  $y$  are discrete, the integral in Eq. (2) is also replaced by a sum running over all possible projections for a given irrep. From Eve's point of view,  $G$  acts physically on the colour degrees of freedom of  $A$ , leaving the multiplicity degrees of freedom untouched. For  $SU(2)$ , the action of  $G$  corresponds to physically rotating the reference frame  $A$ . In this case, the label  $x$  corresponds to all the projections of the angular momentum along a specific axis, say  $\hat{z}$ .

The previous discussion implies that  $\mathcal{H}_A$  has the following associated decomposition:

$$\mathcal{H}_A = \bigoplus_q \mathcal{H}_{A_L}^{(q)} \otimes \mathcal{H}_{A_R}^{(q)}, \quad (3)$$

where the direct sum runs over all possible values of the charge  $q$ . The charge could take discrete or continuous values, where in the latter case the states pertaining to the subspaces labeled by  $q$  need to be properly normalised as elements of the full Hilbert space. For the time being we will ignore this technicality, and revisit it again in Section V and Appendix G.

The left ( $\mathcal{H}_{A_L}^{(q)}$ ) and right ( $\mathcal{H}_{A_R}^{(q)}$ ) tensor factors in each subspace labeled by the charge correspond, respectively, to the colour and flavour degrees of freedom of A. With respect to this decomposition, the left-regular representation has the form  $L_A(g) = \bigoplus_q D_{A_L}^{(q)}(g) \otimes \mathbb{1}_{A_R}^{(q)}$ , where  $D_{A_L}^{(q)}(g)$  is an irrep of  $G$  corresponding to the charge  $q$ . Similarly, the right-regular representation has the form  $R_A(g) = \bigoplus_q \mathbb{1}_{A_L}^{(q)} \otimes D_{A_R}^{(q^*)}(g)$ , where  $D_{A_R}^{(q^*)}$  denotes the conjugate representation corresponding to the charge  $q$ . Given a choice of basis as defined in Eq. (2),  $D_{A_R}^{(q^*)}$  is obtained by complex-conjugating the matrix elements of  $D_{A_R}^{(q)}$ .

In general, a Hilbert space decomposing as a direct sum of tensor products, like in Eq. (3), is said to decompose into subsystems [59, 60]. Here, we will use a slightly more general terminology, associating a subsystem with a subalgebra of operators [61, 62]. In particular, we will speak about the left subsystem, which is associated with the subalgebra of operators of the form  $T_L = \bigoplus_q T_{A_L}^{(q)} \otimes \mathbb{1}_{A_R}^{(q)}$ , and about the right subsystem, which is the commutant of the left, and consists of operators of the form  $T_R = \bigoplus_q \mathbb{1}_{A_L}^{(q)} \otimes T_{A_R}^{(q)}$ . A given subalgebra (equivalently, its commutant) always induces a decomposition of the Hilbert space of the form (3) [59, 60]. Note that the basis vectors  $|g\rangle_A$  generally involve nontrivial superpositions of vectors belonging to the subspaces corresponding to different charges.

What is the physical realisation of an ideal quantum reference frame as defined above? The answer generally depends on the group. In section V, we will discuss reference frames for the centrally extended Galilei group. We will show that for this group a reference frame is physically equivalent to two particles — one that serves as a reference for position and the other one as a reference for velocity.

### III. RELATIVE SUBSYSTEMS

In this section, we construct the description of the setup in Fig. 1 from Alice's reference frame. First, we find the subsystem of the full A and S system that Alice has access to. Afterwards, we construct a map from the Hilbert space associated to the external observer, Eve (see Section II), to a Hilbert space with a tensor product structure that is natural from the point of view A. This map entails a refactorisation of the Hilbert space, which can be interpreted as "jumping" onto Alice's reference frame. We study how the representation of the invariant subsystem changes under this refactorisation. Importantly, we find that the full invariant subsystem is larger than the algebra of relative observables between the

system and frame. It contains an extra subsystem, which we call the “extra particle,” due to its physical realisation in the case of the Galilei group, discussed in Section V.

### A. The invariant subsystem

From Eve’s perspective, the Hilbert space of A and S factorises as  $\mathcal{H}_{AS|E} = \mathcal{H}_{A|E} \otimes \mathcal{H}_{S|E}$ . We call this tensor product factorisation the standard partition. In the standard partition,  $G$  acts transversally on operators  $T$ , as  $T \mapsto L_A(g) \otimes U_S(g) T L_A^\dagger(g) \otimes U_S^\dagger(g)$ , for  $g \in G$ .

What are the degrees of freedom that Alice has access to, and how would she describe them? By assumption, Alice has no access to the external reference frame E. Therefore, she has only access to  $G$ -invariant degrees of freedom on the AS system. That is, operators on  $\mathcal{H}_{AS|E}$  that are invariant under the transversal action of  $G$ :  $T = L_A(g) \otimes U_S(g) T L_A^\dagger(g) \otimes U_S^\dagger(g)$ , for all  $g \in G$ . The set of all  $G$ -invariant operators forms an algebra, which we call the invariant subsystem. We assume that Alice has access to all of these (and only these) operators.

Note that any unitary representation of a group  $G$  on a given Hilbert space  $\mathcal{H}$  induces an analogous decomposition to that in Eq. (3),  $\mathcal{H} = \bigoplus_q \mathcal{J}^{(q)} \otimes \mathcal{K}^{(q)}$ , such that  $G$  acts irreducibly on each  $\mathcal{J}^{(q)}$  and trivially on each  $\mathcal{K}^{(q)}$ . In general, the labels  $q$  need not go over all possible irreps, like in the case of the regular representation, and the Hilbert spaces  $\mathcal{K}^{(q)}$  need not be of the same dimension as  $\mathcal{J}^{(q)}$ . This decomposition is a consequence of the fact that a generally reducible representation splits into a direct sum of irreps, some of which might have nontrivial multiplicities. By Shur’s lemma, all invariant operators are proportional to the identity on  $\mathcal{J}^{(q)}$  for all  $q$  and are possibly nontrivial on the multiplicity factors  $\mathcal{K}^{(q)}$ . These operators form the invariant algebra, or the invariant subsystem. Its commutant – which is the algebra with trivial action on the multiplicity factors  $\mathcal{K}^{(q)}$  – is what we call the gauge subsystem. For example, in the case of the Galilei group for a system of particles, the gauge subsystem corresponds to the centre of mass degrees of freedom [7–9].

In our case, any operator on the gauge subsystem is physically irrelevant for Alice – it is redundant. There is a standard way of removing this redundancy from any operator and extracting only the physically relevant degrees of freedom [3], by applying the  $G$ -twirl operation  $\mathcal{T}_{AS}$ :

$$\mathcal{T}_{AS} = \int dg L_A(g) \otimes U_S(g) \cdot L_A^\dagger(g) \otimes U_S^\dagger(g). \quad (4)$$

As shown in Ref. [3], this operation is equivalent to first projecting the operator into a block-diagonal form over the charge sectors (i.e., killing off-diagonal elements between subspaces corresponding to different charges), followed by applying fully depolarising



channels in the left tensor factors. In the standard partition, the space of physically relevant operators from the point of view of Alice, denoted by  $\mathcal{L}_{\text{inv}}(\mathcal{H}_{AS|E})$ , is defined by those operators which are invariant under the  $G$ -twirl,  $T_{\text{inv}} = \mathcal{T}_{AS}[T_{\text{inv}}]$ .  $\mathcal{L}_{\text{inv}}(\mathcal{H}_{AS|E})$  is a strict subspace of the Hilbert space of operators on  $\mathcal{H}_{AS|E}$ , called  $\mathcal{L}(\mathcal{H}_{AS|E})$ .

Importantly,  $\mathcal{L}_{\text{inv}}(\mathcal{H}_{AS|E})$  is independent of Eve's external reference frame,  $E$ , with respect to which the systems  $A$  and  $S$ , and the action of  $G$  were defined. More precisely, as we show in Appendix A, the invariant algebra of a given system (in this case  $AS$ ) is the largest common subalgebra of the relative algebras  $AS|E$  for all conceivable external reference frames  $E$ . The invariant algebra  $\mathcal{L}_{\text{inv}}(\mathcal{H}_{AS|E})$  can thus be regarded as meaningful on its own. We can imagine external reference frames being "out there" or not; our framework is agnostic to their existence.

Let us now turn to Alice's perspective on  $S$ . Imagine that Alice describes an operator  $T$  acting on the system from her point of view. What would be the corresponding operator in the standard tensor product decomposition? We denote the operator  $T$  on  $S$  relative to  $A$  by  $T_{S|A}$ . All operations on  $S$  from Alice's viewpoint correspond to elements of the algebra of system  $S$  relative to reference frame  $A$ , denoted  $S|A$ . In the standard partition, elements  $T \in S|A$  are of the form [3, 4, 13, 14]

$$T = \int dg |g\rangle\langle g|_A \otimes U_S(g) T_S U_S^\dagger(g), \quad (5)$$

where  $T_S$  is an operator on  $\mathcal{H}_{S|E}$ . As a mathematical object,  $S|A$  is independent of the choice of tensor product decomposition, pretty much in the same way as a point or a tangent vector on a manifold is independent of the choice of coordinates. As we will see below,  $S|A$  can have different representations, which are natural to the viewpoint of different reference frames. A rough analogy is that of a point or a tangent vector to a manifold, which can be represented in different coordinate systems, which are natural from the viewpoint of different observers.

Note that  $S|A$  is not the full algebra of  $G$ -invariant operators. This is because the reference frame  $A$  lives in a Hilbert space that carries the regular representation of  $G$ , which is reducible (see Eq. (3)). As such, it has multiplicity subspaces that are invariant under the action of  $G$  [2, 3]. The multiplicity degrees of freedom that are invariant under the transversal action of  $G$ , as this action is defined in terms of the left-regular representation. As a consequence, any operator  $T_R$  on  $\mathcal{H}_{AS|E}$  of the form

$$T_R = \bigoplus_q \mathbb{1}_{A_L}^{(q)} \otimes T_{A_R}^{(q)} \otimes \mathbb{1}_S \quad (6)$$

is  $G$ -invariant. Here, the first tensor factor denotes the subsystem of  $A$  where  $L_A(g)$  acts, the second denotes the the subsystem of  $A$  where  $R_A(g)$  acts (see Eq. (3)) and the third one



denotes  $S'$  degrees of freedom (all in Eve's standard partition). Note that operators of the form (6) generally overlap with  $S|A$ , but do not belong to it. Therefore, the full invariant system is strictly larger than  $S|A$ . This fact will be very important for the next subsection, where we shall introduce an "extra particle" belonging to the full invariant system.

### B. Change of preferred tensor product factorisation

We now construct a representation of the invariant subsystem that captures Alice's perspective in a natural way. Namely, a representation that *i*) contains only degrees of freedom accessible to Alice (i.e. it is gauge-free), *ii*) contains  $S|A$  as an explicit tensor factor. We call this representation "Alice's perspective." This term is motivated by the conventional treatment of subsystems in quantum mechanics, where each subsystem has a tensor factor of its own (more generally, as noted in Section II, a subsystem is associated with a subalgebra). Thus, when Alice refers to "the system," she is implicitly referring to the system relative to her reference frame. Alice's perspective makes this fact explicit. Moreover, it is justified from an operational perspective (see for example [58]), where Hilbert space operators represent experimental procedures defined with respect to laboratory instruments – Alice's reference frame,  $A$ , in this case.

The first step is to note that there exists an alternative factorisation of  $\mathcal{H}_{AS|E}$  that is induced by the algebra  $S|A$  and its commutant,  $C$ :

$$\mathcal{H}_{AS|E} \cong \mathcal{H}_C \otimes \mathcal{H}_{S|A} =: \mathcal{H}_{C,S|A}. \quad (7)$$

The tensor refactorisation is implemented by a Hilbert space isomorphism

$$|g\rangle_C \otimes |\alpha\rangle_{S|A} \cong |g\rangle_{A|E} \otimes |\alpha\rangle_{S|E}, \quad (8)$$

where  $|\alpha\rangle_{S|E}$  and  $|\alpha\rangle_{S|A}$  are fixed yet arbitrary bases of  $\mathcal{H}_S$  and  $\mathcal{H}_{S|A}$ , respectively. The isomorphism can be written as a map  $V_{E \rightarrow A} : \mathcal{H}_{AS|E} \rightarrow \mathcal{H}_{C,S|A}$ , defined by  $V_{E \rightarrow A} = F_{E \rightarrow A} \circ U_S^\dagger(\hat{g}_A)$ , where  $F_{E \rightarrow A} |g\rangle_A \otimes |\alpha\rangle_S = |g\rangle_C \otimes |\alpha\rangle_{S|A}$ , and

$$U_S^\dagger(\hat{g}_A) = \int dg |g\rangle\langle g|_A \otimes U_S^\dagger(g), \quad (9)$$

Because  $U_S^\dagger(\hat{g}_A)$  is a unitary operator on  $\mathcal{H}_{AS|E}$ , it follows that  $\mathcal{H}_C$  carries the left- and right-regular representations of  $G$ , and  $\mathcal{H}_{S|A}$  carries a representation  $U_{S|A}$  of  $G$  which is isomorphic to  $U_S$ .

A straightforward calculation shows that the super-operator  $\mathcal{V}_{E \rightarrow A} = V_{E \rightarrow A} \cdot V_{E \rightarrow A}^\dagger$  maps the representation of  $S|E$  in  $\mathcal{H}_{AS|E}$  to the tensor factor  $\mathcal{H}_{S|A}$ ,

$$\mathcal{V}_{E \rightarrow A} \left[ \int dg |g\rangle\langle g|_A \otimes U_S(g) T_S U_S^\dagger(g) \right] = \mathbb{1}_C \otimes T_{S|A}, \quad (10)$$

where  $\langle \alpha |_S T_S | \beta \rangle_S = \langle \alpha |_{S|A} T_{S|A} | \beta \rangle_{S|A}$ .

Note that, from Alice's perspective, operators on  $\mathcal{H}_{C,S|A}$  are not redundancy-free. This is because we have not projected out the gauge subsystem as in Eq. (4). To do so, we use that  $\mathcal{V}_{E \rightarrow A}$  maps the gauge subsystem to the left-regular representation of  $\mathcal{H}_C$ :

$$\mathcal{V}_{E \rightarrow A}[L_A(g) \otimes U_S(g)] = L_C(g) \otimes \mathbb{1}_{S|A}. \quad (11)$$

We prove Eq. (11) in Appendix B. Therefore, we can eliminate the redundancy coming from the gauge subsystem by twirling with respect to the left-regular representation in  $\mathcal{H}_C$ . Define

$$\mathcal{T}_C = \int dg L_C(g) \otimes \mathbb{1}_{S|A} \cdot L_C^\dagger(g) \otimes \mathbb{1}_{S|A}. \quad (12)$$

In Alice's perspective, the space of physically relevant operators, denoted by  $\mathcal{L}_{\text{inv}}(\mathcal{H}_{C,S|A})$ , is given by those operators which are invariant under the  $G$ -twirl on the  $C$  subsystem,  $T_{\text{inv}} = \mathcal{T}_C[T_{\text{inv}}]$ . Thus,  $\mathcal{L}_{\text{inv}}(\mathcal{H}_{C,S|A})$  contains the effective, redundancy free description of the physics from the perspective of Alice. The full procedure of refactorising the Hilbert space and eliminating the redundancy is captured by the map  $\mathcal{E}_A = \mathcal{T}_C \circ \mathcal{V}_{E \rightarrow A}$ . In Appendix C, we show that  $\mathcal{E}_A = \mathcal{T}_C \circ \mathcal{V}_{E \rightarrow A} = \mathcal{V}_{E \rightarrow A} \circ \mathcal{T}_{AS}$ , where  $\mathcal{T}_{AS}$  is defined in Eq. (4). This means that removing the redundancy and changing the factorisation commute in a natural way.

Following the reasoning leading to Eq. (3) and the discussion below it, we see that all operators in  $\mathcal{L}_{\text{inv}}(\mathcal{H}_{C,S|E})$  are of the form

$$T_{\text{inv}} = \bigoplus_q \mathbb{1}_{C_L}^{(q)} \otimes T_{C_R,S|A}^{(q)}, \quad (13)$$

where  $\mathcal{T}_{C_R,S|A}^{(q)}$  is an operator on  $\mathcal{H}_{C_R}^{(q)} \otimes \mathcal{H}_{S|A}$ , with a notation analogous to that of Eq. (3). Clearly, the identity operators  $\mathbb{1}_{C_L}^{(q)}$  are not physically meaningful for Alice, as she cannot access the gauge subsystem. For this reason, we could define Alice's perspective by projecting Eq. (13) on each charge sector  $q$  and then tracing out the corresponding  $\mathcal{H}_{C_L}^{(q)}$  Hilbert space. However, we will keep the operators  $\mathbb{1}_{C_L}^{(q)}$  as in Eq. (13) for mathematical convenience, as will be clear in Section IV.

To summarise, in the perspective of A, the full Hilbert space is associated with the following decomposition:

$$\mathcal{H}_{C,S|A} = \left( \bigoplus_q \mathcal{H}_{C_L}^q \otimes \mathcal{H}_{C_R}^q \right) \otimes \mathcal{H}_{S|A}, \quad (14)$$

where the left subsystem of C contains the gauge degrees of freedom.

### C. The extra particle

What is the physical meaning of the right-regular subsystem of  $C$ ? To answer this question, consider a general operator on  $C$ ,  $\int dg' dg T(g', g) |g'\rangle\langle g|_C \otimes \mathbb{1}_{S|A}$ , and act on it with  $\mathcal{T}_C$ . The result is

$$T_{\text{inv}} = \int dg' dg T(g', g) R_C^\dagger(g') R_C(g) \otimes \mathbb{1}_{S|A}. \quad (15)$$

$T_{\text{inv}}$  is  $G$ -invariant, and therefore represents a physically meaningful operator, expressed in Alice's perspective. We call the set of these operators the algebra  $\overline{S|A}$ . It is the complement of  $S|A$  in the full invariant subsystem,  $\mathcal{L}_{\text{inv}}(\mathcal{H}_{C,S|A})$ , in the sense that its tensor product with  $S|A$  gives the full invariant subsystem,  $\mathcal{L}_{\text{inv}}(\mathcal{H}_{C,S|A}) = S|A \otimes \overline{S|A}$ .

In the standard partition,  $\overline{S|A}$  corresponds to a subsystem which is non-trivial in both the right-regular representation and the system, as can be seen by applying the inverse of Eq. (9) to a general operator on  $\overline{S|A}$ . Explicitly, in the standard partition,  $\overline{S|A}$  consists of operators of the form

$$T_{\overline{S|A}} = \int dg' dg |g'\rangle\langle g'|_{A|E} T_{A|E}^R |g'\rangle\langle g'|_{A|E} \otimes U_{S|E}(g') U_{S|E}^\dagger(g), \quad (16)$$

where  $T_{A|E}^R$  is left-invariant.

We call the algebra  $\overline{S|A}$  the "extra particle," because it formally satisfies (in a single mass sector) the algebra of a single particle in the case of the centrally extended Galilei group, as we show in Section V. As we will see in Section IV,  $\overline{S|A}$  is essential to the unitarity of quantum reference frame transformations at the level of algebra of observables. For this reason, we argue that, in a fully relative formulation of quantum mechanics, the "extra particle" has to be considered standardly when we refer to a quantum system. In this way, the relative nature of quantum objects with respect to a reference frame, which is normally considered implicit, becomes *explicit* in our formalism.

One might wonder under which circumstances the extra particle does not play a significant role and can be considered implicitly. This is the case when the state of the reference frame  $A$  in  $E$ 's factorisation is classical, that is, for states on  $\mathcal{H}_{AS}$  of the form  $|g\rangle\langle g|_A \otimes \rho_S$  for  $g \in G$  and  $\rho_S$  a state on  $\mathcal{H}_S$ , or any convex combination (probabilistic mixture) of such states. Applying  $\mathcal{T}_C \circ \mathcal{V}_{E \rightarrow A}$  to any such state, we immediately see that the extra particle  $\overline{S|A}$  is in the maximally mixed state and in a tensor product with the state of  $S|A$ . In this sense, the extra particle carries information about the "quantumness" of the reference frame state. Remarkably, this "quantumness" is independent of any potential external observer, as  $\overline{S|A}$  is part of the invariant subsystem.

#### IV. QUANTUM REFERENCE FRAME TRANSFORMATIONS

Consider now 2 observers, Alice and Bob, with QRFs A and B, respectively. The total Hilbert space in the standard partition is  $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_S$  (we omit the explicit reference to Eve's reference frame for simplicity). As before, A and B are perfect reference frames, so  $\mathcal{H}_A$  and  $\mathcal{H}_B$  each carry the left- and right-regular representation of  $G$ .  $\mathcal{H}_S$  carries an arbitrary unitary representation of  $G$ . Following the procedure of Section III, we can




FIG. 2. The full invariant system can be decomposed in a way that is natural to A (vertical, orange "threads") and in a way that is natural to B (horizontal, green threads). A QRF is a preferred factorisation of the invariant system, and QRF transformation is a change from one preferred factorisation to another. In this illustration, when 2 different subsystems overlap it means that their corresponding operators don't commute in general. In this way, when A refers to "the system," she is actually referring to the subsystem  $A|B$ , which overlaps with  $S|B$  and  $A|B$  from the point of view of B. Importantly, the inclusion of the subsystems  $\overline{SB|A}$  and  $\overline{SA|B}$  is essential to find a unitary relation between A and B's tensor product factorisations.

express the invariant subsystem of the joint system ABS in the perspective of Alice. This gives rise to the invariant subalgebra  $\mathcal{L}_{\text{inv}}(\mathcal{H}_{C,BS|A})$ , where  $\mathcal{H}_{C,BS|A} = \mathcal{H}_C \otimes \mathcal{H}_{BS|A} = \mathcal{H}_C \otimes \mathcal{H}_{B|A} \otimes \mathcal{H}_{S|A}$ , with obvious notation. The space  $\mathcal{H}_C$  decomposes into a left- and a right-invariant part. The left-invariant part is the subsystem  $\overline{BS|A}$  and the right-invariant part is the gauge subsystem.

An analogous procedure gives rise to Bob's perspective, corresponding to the algebra  $\mathcal{L}_{\text{inv}}(\mathcal{H}_{D,S|B})$ . As in the case of Alice,  $\mathcal{H}_D$  decomposes into a left- and a right-invariant

parts, which are the extra particle  $\overline{AS|B}$  and the gauge subsystem from B's perspective, respectively. In this Section, we construct a unitary map that relates Alice's and Bob's perspectives. To this end, we note that both perspectives are unitarily related to the standard decomposition (E). Then, to "jump" between the perspective of A and B, we can map the representation of A to that of E and then map the representation of E to that of B. This same logic is used to relate different quantum reference frames in the "perspective neutral" approach ([39]).

We define a quantum reference frame transformation from Alice to Bob,  $\mathcal{S}_{A \rightarrow B} : \mathcal{L}_{\text{inv}}(\mathcal{H}_{C,BS|A}) \longrightarrow \mathcal{L}_{\text{inv}}(\mathcal{H}_{D,AS|B})$ , as

$$\mathcal{S}_{A \rightarrow B} = \mathcal{V}_{E \rightarrow B} \circ \mathcal{V}_{E \rightarrow A}^\dagger. \quad (17)$$

Following the same logic as in Subsection III B, we define  $\mathcal{V}_{E \rightarrow B} = V_{E \rightarrow B} \cdot V_{E \rightarrow B}^\dagger$  and  $\mathcal{V}_{E \rightarrow A}^\dagger = V_{E \rightarrow A}^\dagger \cdot V_{E \rightarrow A}$ . Here,  $\mathcal{V}_{E \rightarrow B} = F_{E \rightarrow B} \circ U_{AS}^\dagger(\hat{g}_B)$  and  $\mathcal{V}_{E \rightarrow A} = F_{E \rightarrow A} \circ U_{BS}^\dagger(\hat{g}_A)$ , where

$$U_{BS}(\hat{g}_A) = \int dg |g\rangle\langle g|_A \otimes L_B(g) \otimes U_S(g), \quad (18a)$$

$$U_{AS}(\hat{g}_B) = \int dg L_A(g) \otimes |g\rangle\langle g|_B \otimes U_S(g). \quad (18b)$$

$F_{E \rightarrow A}$  acts as  $F_{E \rightarrow A} |g\rangle_A |h\rangle_B |\alpha\rangle_S = |g\rangle_C |h\rangle_{B|A} |\alpha\rangle_{S|A}$ , and an analogous equation holds for  $F_{E \rightarrow B}$ .

We can now use Eq. (17) to compute the transformation of operators from the perspective of A to that of B. Let us divide the set of operators in the reference frame of A into 3 classes. Class 1 is made of operators of the form  $\mathbb{1}_C \otimes \mathbb{1}_{B|A} \otimes T_{S|A}$ , i.e. elements of  $S|A$ ; class 2 is made of operators of the form  $\mathbb{1}_C \otimes T_{B|A} \otimes \mathbb{1}_{S|A}$ , i.e. elements of  $B|A$ . Finally, class 3 is made of operators of the form  $T_C^R \otimes \mathbb{1}_{B|A} \otimes \mathbb{1}_{S|A}$ , where  $T^R$  is left-invariant, i.e. elements of  $\overline{BS|A}$ . The transformation of each of these 3 classes of operators is computed explicitly in Appendix D. The result is

$$\mathcal{S}_{A \rightarrow B}[\mathbb{1}_C \otimes \mathbb{1}_{B|A} \otimes T_{S|A}] = \int dg |g\rangle\langle g|_{A|B} \otimes \mathbb{1}_D \otimes U_{S|B}(g) T_{S|B} U_{S|B}^\dagger(g) \quad (19a)$$

$$\mathcal{S}_{A \rightarrow B}[\mathbb{1}_C \otimes T_{B|A} \otimes \mathbb{1}_{S|A}] = \int dh dg |h^{-1}\rangle\langle h|_{T_{A|B}} |g\rangle\langle g^{-1}|_{A|B} \otimes R_D(h^{-1}g) \otimes U_{S|B}(h^{-1}g) \quad (19b)$$

$$\mathcal{S}_{A \rightarrow B}[T_C^R \otimes \mathbb{1}_{B|A} \otimes \mathbb{1}_{S|A}] = \int dg |g\rangle\langle g|_{A|B} \otimes R_D(g) T_D^R R_D^\dagger(g) \otimes \mathbb{1}_{S|B}. \quad (19c)$$

Eqs. (19) fully characterise the relation between A's natural tensor product factorisation and B's. We thus see that a quantum reference frame is a preferred tensor factorisation of the invariant subsystem. Alice and Bob have 2 such partitions, natural to their relative degrees of freedom. This fact is at the heart of the relativity of entanglement under QRF transformations [37]. As we show in Appendix E, in the zero-charge sector Eq. (17) reduces to the QRF transformation found in Ref. [49], which is equivalent to that of Ref. [37] for the case of translations.

Note that  $S|A$  and  $S|B$  partially overlap but are not equal. The same is true for the subalgebras  $BS|A$  and  $AS|B$ . For this reason, we cannot expect these subalgebras to be unitarily related. However, the extra particle comes to the rescue, as it complements each of  $BS|A$  and  $AS|B$  to the full invariant subsystem. This is why the extra particle is essential for unitarity.

It is worth emphasising the generality of the transformations in Eqs. (19). They do not merely allow us to say how to "jump" between two fixed reference frames, but also how the description from the point of view of one reference frame would change if that reference frame is subjected to an arbitrary active transformation from the perspective of another. For instance, if Alice applies an active unitary transformation on  $B$ ,  $U_{B|A}$ , the state of the invariant subsystem in the perspective of Bob would undergo a corresponding passive unitary transformation, whose form can be computed from Eq. (19b) by plugging  $U_{B|A}$  in the place of  $T_{B|A}$ . The transformation seen by Bob would generally spread over the system, Alice's frame, as well as the extra particle, where the latter is again essential for recovering unitarity (see Appendix F). The transformations of Eq. (19) are obviously not restricted to scenarios involving two reference frames, as additional frames can be included in  $S$ .

In conclusion, our framework decomposes the full invariant subsystem as a network of subsystems, whose "threads" represent the viewpoints of  $A$  and  $B$ . A QRF transformation is a change from a decomposition which is natural to Alice to a decomposition which is natural to Bob. Figure IV depicts how each subalgebra in Alice's reference frame commutes or fails to commute with each subalgebra in Bob's partition. The vertical "threads" correspond to Alice's QRF, whereas the horizontal ones correspond to Bob's QRF. More generally, we can imagine multiple reference frames and the corresponding network of relative subalgebras related via analogous principles. A remarkable feature of these algebraic relations is that, as commented earlier (see Appendix A), they concern algebras that are independent of external reference frames, yet compatible with any potential external reference frame in the sense that they would automatically embed as subalgebras of the corresponding larger invariant algebra entailed by the existence of such a frame. This unveils a tantalising mathematical landscape of nested subalgebras that may represent both actual and potential scenarios.

## V. CENTRALLY EXTENDED GALILEI GROUP

In this Section, we apply our framework to the case of the centrally extended Galilei group. We start by briefly introducing the Galilei group and its central extension. Then, we compute the algebras  $S|A$  and  $\overline{S|A}$ , and give a physical interpretation of the regular

representation as a quantum reference frame. For simplicity, we treat the case of 1 spatial dimension and focus only on spatial translations and boosts, leaving time translations to further work. Although our treatment is formal, glossing over issues regarding normalisation, we can extract the essential physics and obtain interesting insights about the physical realisation of the regular representation as a QRF. For a rigorous construction of covariant "screen observables" for the Galilei and Poincaré groups, see Ref. [64].




FIG. 3. Physical interpretation of the regular representation of the centrally extended Galilei Group. For a given mass sector  $m$ , the regular representation can be seen as a system of 2 particles. Here we depict the case where each particle has a mass  $m/2$ . In this interpretation, the left regular representation corresponds to the degrees of freedom of the centre of mass, CM, of the 2-particle system. The right regular representation corresponds to the distance of any of the 2 particles to the centre of mass, or half their relative distance, REL. Imagine that  $A$  describes an operation on  $S$  using the regular representation of the centrally extended Galilei Group as a reference. We can then ask how this operation "looks like" from the standard partition viewpoint. Roughly speaking, in this viewpoint,  $A$  uses one of the particles,  $A_{m_1}$ , as a reference frame for position, and uses the *other* particle,  $A_{m_2}$  as a reference frame for velocity (see Eqs. (25)).

### A. Introducing the group

In 1 spatial dimension, the Galilei group consist in elements  $(a, v)$ , labeled by a translation parameter  $a \in \mathbb{R}$  and a boost parameter  $v \in \mathbb{R}$ . Physically, the transformation  $(a, v)$  means



changing to a reference frame which is displaced in space by a distance  $a$  and moving with a constant velocity  $v$  with respect to the original reference frame. The composition rule of the Galilei group is  $(a', v') \cdot (a, v) = (a' + a, v' + v)$ .

Galilean transformations on a quantum particle of mass  $m$  are generated by the momentum operator  $\hat{p}$  (translations) and by the boost operator  $\hat{k} = \hat{p}t - m\hat{x}$  (boosts), where  $t$  is the time and  $\hat{x}$  is the position operator. The commutation relation of the group is  $[\hat{p}, \hat{k}] = im$ . The non-commutativity of the Galilean generators in quantum mechanics implies the well known fact that the Galilean group has a projective representation in Hilbert space

$$U^{(m)}(a', v')U^{(m)}(a, v) = e^{i\frac{m}{2}(av' - a'v)}U^{(m)}(a' + a, v' + v), \quad (20)$$

where  $U(a, v) = \exp(-i(a\hat{p} + v\hat{k}))$ . In order to apply our framework, we consider the central extension of the Galilei group,  $\tilde{G}$  (see, for example, [65, 66]).  $\tilde{G}$  has group elements  $(\theta, a, v)$  and group multiplication rule  $(\theta', a', v') \cdot (\theta, a, v) = (\theta' + \theta + \varphi(a', v'; a, v), a' + a, v' + v)$ , where  $\varphi(a', v'; a, v) = (av' - a'v)/2$ . For a given mass  $m$ , we define the (irreducible) representation of  $\tilde{G}$  by  $\tilde{U}^{(m)}(\theta, a, v) = e^{im\theta}U^{(m)}(a, v)$ . It is easy to check that Eq. (20) is an ordinary (i.e. not projective) representation of the centrally extended Galilei group.

The centrally extended Galilei group involves an additional parameter,  $\theta$ , whose physical meaning as the conjugate variable to a dynamical mass variable has been discussed in the literature [66–70]. Regardless the specific meaning of  $\theta$ , we do not miss any physics by conceiving a (possibly fictitious) reference frame for it, since the physically accessible projective representations of the Galilei group are naturally recovered in the case of reference frames of fixed mass. What is more, this treatment highlights the interesting possibility of having an explicit QRF for dynamical mass, allowing for coherences between different mass sectors.

## B. QRFs for the centrally extended Galilei group

Suppose that  $A$  is a QRF carrying the regular representation of  $\tilde{G}$ . This representation is spanned by vectors of the form

$$|(\theta, a, v)\rangle = \int^{\oplus} dm dp \sqrt{m} (\tilde{U}^{(m)}(\theta, a, v) |m; p\rangle_L) \otimes |m; p\rangle_R, \quad (21)$$

with  $\theta, a, v \in \mathbb{R}$  [65], and has inner product (see Appendix G)

$$\langle(\theta', a', v')|(\theta, a, v)\rangle = \delta(\theta' - \theta)\delta(a' - a)\delta(v' - v). \quad (22)$$

Consider a system  $S$  carrying an irreducible representation of  $\tilde{G}$ , labeled by the mass

$m_S$ . This is equivalent to a particle of mass  $m_S$ <sup>1</sup>. Using Eq. (5) we can compute the generators of Galilean transformations,  $\hat{p}_{S|A}$  and  $\hat{k}_{S|A}$ , in the standard partition. As shown in Appendix H, the result is

$$\hat{p}_{S|A} = \mathbb{1}_A \otimes \hat{p}_S - m_S \int^{\oplus} \frac{dm}{m} (\hat{p}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} - \mathbb{1}_{A_L}^{(m)} \otimes \hat{p}_{A_R}^{(m)}) \otimes \mathbb{1}_S, \quad (23a)$$

$$\hat{k}_{S|A} = \mathbb{1}_A \otimes \hat{k}_S - m_S \int^{\oplus} \frac{dm}{m} (\hat{k}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} + \mathbb{1}_{A_L}^{(m)} \otimes \hat{k}_{A_R}^{(m)}) \otimes \mathbb{1}_S, \quad (23b)$$

where  $\mathbb{1}_A = \int^{\oplus} dm \mathbb{1}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)}$ , and we have used the same notation as in Section III.

We can use Eqs. (23) to compute the algebra of the extra particle in the standard partition (see Appendix H):

$$\hat{p}_{S|A}^R = \hat{p}_A^R \otimes \mathbb{1}_S + \mathbb{1}_A \otimes \hat{p}_S - m_S \int^{\oplus} \frac{dm}{m} (\hat{p}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} - \mathbb{1}_{A_L}^{(m)} \otimes \hat{p}_{A_R}^{(m)}) \otimes \mathbb{1}_S, \quad (24a)$$

$$\hat{k}_{S|A}^R = \hat{k}_A^R \otimes \mathbb{1}_S + \mathbb{1}_A \otimes \hat{k}_S - m_S \int^{\oplus} \frac{dm}{m} (\hat{k}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} + \mathbb{1}_{A_L}^{(m)} \otimes \hat{k}_{A_R}^{(m)}) \otimes \mathbb{1}_S, \quad (24b)$$

where  $\hat{p}_A^R = \int^{\oplus} dm \mathbb{1}_{A_L}^{(m)} \otimes \hat{p}_{A_R}^{(m)}$ .  $\hat{p}_{S|A}^R$  and  $\hat{k}_{S|A}^R$  satisfy the commutation relations of the centrally extended Galilei group for a mass of  $\hat{M}_A + m_S$ , where  $\hat{M}_A = \int^{\oplus} dm m \mathbb{1}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)}$ . More explicitly,  $[\hat{p}_{S|A}^R, \hat{k}_{S|A}^R] = i(\hat{M}_A \otimes \mathbb{1}_S + \mathbb{1}_A \otimes m_S \mathbb{1}_S)$ .

Although  $m$  can take, in principle, values over all  $\mathbb{R}$ , we can focus on the positive mass case by restricting the set of states on which our operators act. Let us now focus on a single mass sector of the regular representation, corresponding to mass  $m > 0$ . In Appendix G, we discuss normalisation issues that arise when restricting to a single mass sector. In what follows, it will be more instructive to deal with position operators instead of boost operators, so we write the boost operators in terms of position ones in Eq. (23). Thus, we focus on the momentum operator  $\hat{p}_{S|A}^{(m)} = \mathbb{1}_A^{(m)} \otimes \hat{p}_S - (m_S/m)(\hat{p}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} - \mathbb{1}_{A_L}^{(m)} \otimes \hat{p}_{A_R}^{(m)}) \otimes \mathbb{1}_S$  and the position operator  $\hat{x}_{S|A}^{(m)} = \mathbb{1}_A^{(m)} \otimes \hat{x}_S - \hat{x}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} - \mathbb{1}_{A_L}^{(m)} \otimes \hat{x}_{A_R}^{(m)} \otimes \mathbb{1}_S$ , where  $\mathbb{1}_A^{(m)} = \mathbb{1}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)}$ .

We will now show that the system A can be seen as consisting of two particles, called  $A_{m_1}$  and  $A_{m_2}$  of respective masses  $m_1$  and  $m_2$ , such that  $m_1 + m_2 = m$ , where  $A_{m_1}$  serves as a reference for position and  $A_{m_2}$  as a reference for velocity. We define  $\hat{x}_{A_{m_1}} = \hat{x}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} + \mathbb{1}_{A_L}^{(m)} \otimes \hat{x}_{A_R}^{(m)}$ , and  $\hat{x}_{A_{m_2}} = \hat{x}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} - (m_1/m_2) \mathbb{1}_{A_L}^{(m)} \otimes \hat{x}_{A_R}^{(m)}$ . The momenta  $\hat{p}_{A_{m_1}}$  and  $\hat{p}_{A_{m_2}}$  are the conjugate variables to  $\hat{x}_{A_{m_1}}$  and  $\hat{x}_{A_{m_2}}$ , respectively. In this way,  $\hat{x}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)}$  (the left-regular

<sup>1</sup> While we are considering a single such system, our results apply automatically to a system of multiple particles, where the role of S would be played by the centre of mass. Indeed, for a system of particles with Galilei symmetry the Hilbert space decomposes as [9]  $\mathcal{H}_S \cong \mathcal{H}_{S_{CM}} \otimes \mathcal{H}_{S_{rel}}$ , where the  $\mathcal{H}_{S_{CM}}$  is the gauge subsystem corresponding to the centre of mass, on which the group acts irreducibly, and  $\mathcal{H}_{S_{rel}}$  is the invariant subsystem containing relational degrees of freedom, on which the group acts trivially. Due to the transversal action of the group, when we bring in the reference frame A, it would be as if the group only acts on  $AS_{CM}$ , where  $S_{CM}$  behaves as single particle, while  $S_{rel}$  would remain invariant and would tensor-multiply the invariant subsystem of  $AS_{CM}$  to give the full invariant subsystem of AS.

representation) can be seen as the position operator for the centre of mass of a system of our two particles,  $A_{m_1}$  and  $A_{m_2}$ . That is,  $\hat{x}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} = (m_1 \hat{x}_{A_{m_1}} + m_2 \hat{x}_{A_{m_2}})/m$ . Similarly, the operator  $\hat{p}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)}$  is the momentum of the centre of mass,  $\hat{p}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} = \hat{p}_{A_{m_1}} + \hat{p}_{A_{m_2}}$ . On the other hand, the operator  $\mathbb{1}_{A_L}^{(m)} \otimes \hat{x}_{A_R}^{(m)}$  (the right-regular representation) is proportional to the relative distance between  $A_{m_1}$  and  $A_{m_2}$ ,  $\mathbb{1}_{A_L} \otimes \hat{x}_{A_R}^{(m)} = (m_2/m)(\hat{x}_{A_{m_1}} - \hat{x}_{A_{m_2}})$ , whereas  $\mathbb{1}_{A_L}^{(m)} \otimes \hat{p}_{A_R}^{(m)}$  corresponds to the relative momentum,  $\mathbb{1}_{A_L}^{(m)} \otimes \hat{p}_{A_R}^{(m)} = \hat{p}_{A_{m_1}} - (m_1/m_2)\hat{p}_{A_{m_2}}$ .

Going back to the full space, considering all mass sectors, it follows that Eqs. (23) can be rewritten as

$$\hat{x}_{S|A} = \mathbb{1}_A \otimes \hat{x}_S - \int^{\oplus} dm \hat{x}_{A_{m_1}} \otimes \mathbb{1}_S \quad (25a)$$

$$\hat{p}_{S|A} = \mathbb{1}_A \otimes \hat{p}_S - m_S \int^{\oplus} \frac{dm}{m_2} \hat{p}_{A_{m_2}} \otimes \mathbb{1}_S, \quad (25b)$$

where we have kept  $m_2$  inside the direct integral over  $m$  because it can be different for different mass sectors.

If we have 2 QRFs, A and B, for the centrally extended Galilei group, the natural tensor product decompositions associated to A is related to the decomposition of B via Eqs. (19). In Appendix H, we compute explicitly the QRF transformation connecting the infinitesimal generators of the group "as seen" from QRF A to those "as seen" from QRF B.

In conclusion, the regular representation of the centrally extended Galilei Group can be seen as a system of variable mass, which under a properly normalised restriction to a fixed mass sector, consists of 2 particles, one of them serving as a QRF for position and the other as a QRF for velocity. These particles transform under the usual projective representation of the Galilei group. The case of a single mass sector with  $m_1 = m_2 = m/2$  is depicted in Fig. 3.

### C. Comparison with other frameworks

It is instructive to compare our framework in a given mass sector with other proposals for the relational description of multi-particle systems under Galilei and translation symmetries [7, 9, 37, 39]. Assume that our reference frame A in the given mass sector is realised by particles 1 and 2 (we drop the label A for simplicity) serving as references for position and velocity, respectively, and let the system S consist of  $N - 2$  particles, labeled by  $i = 3, \dots, N$ . Denote the mass of particle  $i$  by  $m_i$  and the pair of its position and momentum operators in the standard partition by  $(\hat{x}_i, \hat{p}_i)$ ,  $i = 1, \dots, N$ . As noted in footnote (1), the Hilbert space of such an  $N$ -particle system defined relative to a hypothetical external observer decomposes as [9]  $\mathcal{H} \cong \mathcal{H}_{CM} \otimes \mathcal{H}_{rel}$ , where  $\mathcal{H}_{CM}$  is the gauge subsystem corresponding to the centre of mass, defined by the position and momentum operators

$\hat{x}_{CM} = \sum_i m_i \hat{x}_i / M$ ,  $P_{CM} = \sum_i \hat{p}_i$ , where  $M = \sum_i m_i$ , and  $\mathcal{H}_{rel}$  is the invariant subsystem containing relational degrees of freedom.

In our framework, the choice of particles 1 and 2 as a QRF gives rise to a decomposition of the invariant subsystem into a tensor product of the  $N - 2$  "system" particles defined relative to the QRF, plus the corresponding extra particle. The relative particles are given by the canonically conjugate pairs of relative position and momentum operators  $(\hat{x}_{i|1}, \hat{p}_{i|2})$ , where  $\hat{x}_{i|1} = \hat{x}_i - \hat{x}_1$ , and  $\hat{p}_{i|2} = \hat{p}_i - \frac{m_i}{m_2} \hat{p}_2$ , for  $i = 3, \dots, N$ , and the extra particle by the canonically conjugate pair (24) restricted to the respective mass sector.

In comparison, Ref. [9] considers only a single particle as a reference for either the position or velocity of the remaining particles. For example, if particle 1 is used as a reference for position, this is associated with a decomposition of the invariant subsystem into  $N - 1$  relational particles, defined by the relative position operators  $\hat{x}_{i|1} = \hat{x}_i - \hat{x}_1$ ,  $i = 2, \dots, N$  and canonically conjugate momenta  $\hat{p}_{i|1} = \hat{p}_i - \frac{m_i}{M} \hat{p}_{CM}$ . Note that, as seen from an external observer, the momenta in this case do not have an interpretation as the relative momenta of one particle relative to another, as the centre of mass is not a separate subsystem from such a perspective but rather it depends on the positions and masses of the whole collection of particles. In contrast, the relative momenta defined here depend only on the momenta of two particles: the momentum  $\hat{p}_i$ ,  $i = 3, \dots, N$  and the momentum of the reference frame for velocity, particle 2.

Ref. [37] has a completely internal treatment, where one "jumps" from the QRF of one internal observer to that of another one without invoking an external observer. It treats translations and Galilean boosts in 1 dimension as 2 separate cases, introducing a QRF transformation for translations and a different QRF transformation for boosts. Similar to Refs. [7, 9], Ref. [37] uses a single-particle model of QRF. A single particle of finite mass  $m$  can be either a perfect reference frame for the translation group, or a perfect reference frame for the group of Galilean boosts in one dimension, but not for both. In contrast, here we consider, in a fixed mass sector, a system of 2 particles serving as a QRF for both translations and boosts (which combined form the Galilei group in 1 dimension). Note that, in the limit  $m \rightarrow \infty$ , a single particle can serve as a perfect reference frame for both position and velocity. It would be interesting to investigate the connection between this limit and the QRF model presented here.

Ref. [39] obtains the QRF transformation for translations of Ref. [37] by means of a gravity-inspired momentum constraint, which forces the centre-of-mass momentum of a "perspective neutral" state to vanish,  $\hat{p}_{CM} |\Psi\rangle = 0$ . Within the  $\hat{p}_{CM} = 0$  subspace, the relational variables of Refs. ([37, 39]) are equivalent to that of [9]. However, the perspective-neutral state of Ref. [39] does not have an immediate operational interpretation, as there is no external observer "out there" to measure such state. Our framework is agnostic

to whether such external observer exists or not, and the constraint state  $|\Psi\rangle$  can be interpreted as a state whose centre-of-mass momentum vanishes "as seen" by the external observer. This can be modelled in our framework by introducing an external reference frame for velocity, aligned with the velocity of the centre of mass. In this case, we would have a total of  $N+1$  particles, where particle  $N + 1$  serves as a reference for velocity (and thereby momentum), while one of the other particles, say particle 1, serves as a reference for position.

Ignoring the extra particle, and assuming that the total momentum of all particles from 1 to  $N$  is zero relative to particle  $N + 1$ , we recover the description of Refs. [37, 39]. In particular, the jumping transformations derived there can be understood as corresponding to changing which particle from 1 to  $N$  serves as a reference for position, while keeping the reference for velocity fixed. As shown in Appendix E, our framework restricted to the zero-charge sector for a general group reduces to the QRF transformation found in Ref. [49]. For the case of translations, this recovers formally the perspective-neutral computation of the QRF transformation developed in Ref. [37].

Finally, the work of Angelo et al., Ref. [7], proposed a relational description of particles within the invariant subsystem that uses a single particle, e.g. particle 1, as a reference for both position and velocity of the other particles, leading to a notion of relational particles with position and momentum operators  $(\hat{x}_{i|1}, \hat{p}_{r_i})$ ,  $i = 2, \dots, N$ , where  $\hat{p}_{r_i} = \frac{m_1 m_r}{m_1 + m_r} (\frac{\hat{p}_i}{m_i} - \frac{\hat{p}_1}{m_1})$  (note that this notion of relative momentum is not equal to the relative velocity of the respective particle times its mass, but times the reduced mass of the particle and the reference, which is needed to ensure the canonical commutation relations for each particle). As emphasised in Ref. [7], these particles are not separate systems, since their algebras do not commute with each other, and the canonical commutation relations are only recovered in the limit  $m_1 \rightarrow \infty$ .

The fact that  $(\hat{x}_{i|1}, \hat{p}_{r_i})$ , as defined by Angelo et al., is not a separate subsystem has drastic consequences, as Ref. [7] illustrate by introducing the "paradox of the 3rd particle". In short, the paradox concerns the observation that, if one uses a single particle as a reference frame for both position and velocity in the context of Galilei symmetry, one arrives to the conclusion that the state of particle  $S_2$  defined relative to particle  $S_1$  depends on whether, relative to an external classical reference frame  $E$  (which can be modeled by a very heavy particle), there exists another particle  $S_3$ , separate from  $S_1$  and  $S_2$ .

The resolution of the paradox proposed in [7] is that two systems that are separate relative to  $E$  (in this case  $S_2$  and  $S_3$ ) may be overlapping when described relative to  $S_1$ , and therefore one cannot trace out  $S_3$  from the state relative to  $S_1$ . Note, however, that this conclusion is obtained for a different model of QRF than the one we consider. In our framework, two separate systems are always separate relative to any QRF, and one can

trace them out in any reference frame. Nevertheless, one should do this with care. As we have seen, when two observers using different QRFs refer to the same “system”, they are referring to DOFs that belong to two different, albeit overlapping, subalgebras. Thus, in general their descriptions of the “system” would not contain the same information. Moreover, even if two observers Alice and Bob each describe the reference frame of the other in addition to the system  $S$ , the description of  $BS$  relative to Alice is given by an algebra  $BS|A$  that is not equal to the algebra  $AS|B$  describing  $AS$  relative to  $B$ . Thus, even in this case their descriptions would not contain the same information. The full invariant information, which is accessible by both observers, is only obtained when the extra particle is included in the description.

Recently, Refs. [50, 51] proposed an alternative analysis of the paradox of the third particle. They introduced quantum reference frame transformations as generalised transformations on the gauge subsystem, which are quantum-controlled by classical configurations. In this framework, a “relational partial trace” was defined, as a means to resolve the paradox in a gauge-independent way.

## VI. DISCUSSION

Symmetry transformations between QRFs can lead to a more general notion of symmetry in quantum mechanics, potentially sharpening our operational understanding of space-time at the quantum level. For this reason, it is very important to understand what is at the root of the key differences between classical and quantum reference frame transformations. In this work, we have developed an approach to QRF transformations that focuses on the algebra of relative observables between a system and a reference frame. From this point of view, a QRF transformation is a change from a preferred tensor factorisation to another one. As the notion of subsystem is key to the definition of entanglement, our approach explains why entanglement and superposition are relative notions in QRFs. Moreover, given a set of QRFs, our approach fully characterises how different subsystem decompositions are connected to each other. This leads to a picture of the full invariant system of a quantum system as being composed by a network of subalgebras, with different parts of the network corresponding to different QRF viewpoints.

Our framework is naturally compatible with an incoherent-twirling approach to QRFs rather than a coherent twirling approach. This feature makes our approach a good candidate for studying QRF symmetries in a strict subsystem of the universe in the most general way. Approaches that restrict to a given charge subspace restrict the possible states in which the subsystem of the universe under study can be with respect to potential new systems out there. For this reason, a framework that focuses on a given charge subspace



fails to capture the potential relation that the subsystem of interest might have with external degrees of freedom. Our approach, developed at the level of the full invariant subsystem of a given system, is compatible with extending the system we are interested in an arbitrary way. In this sense, our framework supports the view that, in some situations, incoherent twirling should be preferred to coherent twirling in the quantisation of systems with gauge symmetries [3, 63].

There are several new research avenues that our work opens. On the one hand, it would be very interesting to study QRF transformations with respect to relativistic groups, i.e. the Lorentz and Poincaré groups, and ask what operational notion of space-time arises from such reference frames. We believe our approach is general and powerful enough to make such a study feasible. On the other hand, we have focused on a very restrictive notion of quantum reference frame, namely, that corresponding to the regular representation of the group. The reason for doing this is to make explicit contact with our more familiar classical notion of reference frame viewpoints and transformations. Admittedly, the regular representation is a highly idealised object, and it would be very important to learn how to treat situations in which our reference frame is bounded in resources [3, 7, 8, 63]. We believe the solution to this open problem can yield important insights beyond the approximation of superpositions of semiclassical causal structures and spacetimes, as for example the case considered in Ref. [71].

**Note added:** During the completion of this work, we became aware of related work by Anne-Catherine de la Hamette, Thomas D. Galley, Philipp A. Höhn, Leon Loveridge, and Markus P. Müller, which appears on the arXiv today [73].

## ACKNOWLEDGEMENTS

This publication was made possible through the support of the ID # 61466 grant from the John Templeton Foundation, as part of the “The Quantum Information Structure of Spacetime (QISS)” Project (qiss.fr). The opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the John Templeton Foundation. This work was supported by the Program of Concerted Research Actions (ARC) of the Université libre de Bruxelles. O. O. is a Research Associate of the Fonds de la Recherche Scientifique (F.R.S.–FNRS). E.C-R. acknowledges financial support from the Austrian Science Fund (FWF) through BeyondC (F7103-N48), the European Commission via Testing the Large-Scale Limit of Quantum Mechanics (TEQ) (No. 766900) project, the Founda-



tional Questions Institute (FQXi), the Swiss National Science Foundation (SNSF) via the National Centers of Competence in Research QSIT and SwissMAP, as well as the project No. 200021\_188541.

---

- [1] Aharonov, Y., & Susskind, L. *Physical Review*, **155(5)**, (1967).
- [2] Kitaev, A., Mayers, D., & Preskill, J. *Physical Review A*, **69(5)**, (2004).
- [3] Bartlett, S. D., Rudolph, T., & Spekkens, R. W. *Reviews of Modern Physics*, **79(2)**, (2007).
- [4] Bartlett, S. D., Rudolph, T., Spekkens, R. W., & Turner, P. S. *New Journal of Physics*, **11(6)**, (2009).
- [5] Bartlett, S. D., Rudolph, T., & Spekkens, R. W. *International Journal of Quantum Information*, **4(01)**, (2006).
- [6] Aharonov, Y., & Kaufherr, T. *Physical Review D*, **30(2)**, (1984).
- [7] Angelo, R. M., Brunner, N., Popescu, S., Short, A. J., & Skrzypczyk, P. *Journal of Physics A: Mathematical and Theoretical*, **44(14)**, (2011).
- [8] Angelo, R. M., & Ribeiro, A. D. *Journal of Physics A: Mathematical and Theoretical*, **45(46)**, (2012).
- [9] Smith, A. R., Piani, M., & Mann, R. B. *Physical Review A*, **94(1)**, (2016).
- [10] Loveridge, L., Miyadera, T., & Busch, P. *Foundations of Physics*, **48(2)**, (2018).
- [11] Loveridge, L., & Busch, P. *The European Physical Journal D*, **62(2)**, (2011).
- [12] Loveridge, L., Busch, P., & Miyadera, *EPL (Europhysics Letters)*, **117(4)**, (2017).
- [13] Miyadera, T., Loveridge, L., & Busch, P. *Journal of Physics A: Mathematical and Theoretical*, **49(18)**, (2016).
- [14] Loveridge, L. Quantum measurements in the presence of symmetry (Doctoral dissertation, University of York) (2012).
- [15] Busch, P., & Loveridge, L. *Physical review letters*, **106(11)**, (2011).
- [16] Loveridge, L. *Journal of Physics: Conference Series* **1638, No. 1**, (2020).
- [17] Miyadera, T., & Loveridge, L. *Journal of Physics: Conference Series* **1638, No. 1**, (2020).
- [18] Loveridge, L. In *Journal of Physics: Conference Series* (Vol. 1638, No. 1, p. 012009). IOP Publishing. (2020).
- [19] Bartlett, S. D., Rudolph, T., Spekkens, R. W., & Turner, P. S. *New Journal of Physics*, **8(4)**, (2006).
- [20] Marvian, I., & Spekkens, R. W. Preprint at arXiv:1212.3378. (2012).
- [21] Bartlett, S. D., Rudolph, T., & Spekkens, R. W. *Physical Review A*, **70(3)**, (2004).
- [22] Bartlett, S. D., Rudolph, T., & Spekkens, R. W. *Physical review letters*, **91(2)**, (2003).
- [23] Marvian, I., & Spekkens, R. W. *Physical Review A*, **94(5)**, (2016).
- [24] Marvian, I., & Spekkens, R. W. *Nature communications*, **5(1)**, (2014).

- [25] Marvian, I., & Spekkens, R. W. *New Journal of Physics*, **15(3)**, (2013).
- [26] Marvian, I., Spekkens, R. W., & Zanardi, P. *Physical Review A*, **93(5)**, (2016).
- [27] Marvian, I., & Spekkens, R. W. *Physical Review A*, **90(6)**, (2014).
- [28] Marvian, I., & Spekkens, R. W. *Physical review letters*, **123(2)**, (2019).
- [29] Gour, G., & Spekkens, R. W. *New Journal of Physics*, **10(3)**, (2008).
- [30] Gour, G., Marvian, I., & Spekkens, R. W. *Physical Review A*, **80(1)**, (2009).
- [31] Rovelli, C. *Classical and Quantum Gravity*, **8(2)**, 297. (1991).
- [32] Rovelli, C. *Quantum gravity*. Cambridge university press. (2004).
- [33] Rovelli, C. *Classical and Quantum Gravity*, **8(2)**, 317 (1991).
- [34] Gambini, R., Porto, R. A., & Pullin, J. *New Journal of Physics*, **6(1)**, (2004).
- [35] Bojowald, M., Hoehn, P. A., & Tsobanjan, A. *Classical and quantum gravity*, **28(3)**, (2011).
- [36] Bojowald, M., Hoehn, P. A., & Tsobanjan, A. *Physical Review D*, **83(12)**, (2011).
- [37] Giacomini, F., Castro-Ruiz, E., & Brukner, Č. *Nature communications*, **10(1)**, (2019).
- [38] Giacomini, F., Castro-Ruiz, E., & Brukner, Č. *Physical review letters*, **123(9)**, (2019).
- [39] Vanrietvelde, A., Hoehn, P. A., Giacomini, F., & Castro-Ruiz, E. *Quantum*. **4**, **225**, (2020).
- [40] Hoehn, P. A. *Universe*, **5(5)**, (2019).
- [41] Vanrietvelde, A., Hoehn, P. A., & Giacomini, F. (2018). Preprint at: arXiv:1809.05093. (2018).
- [42] Hoehn, P. A., & Vanrietvelde, A. *New Journal of Physics*, **22(12)**, (2020).
- [43] Castro-Ruiz, E., Giacomini, F., Belenchia, A., & Brukner, Č. *Nature communications*, **11(1)**, (2020).
- [44] Hoehn, P. A., Smith, A. R., & Lock, M. P. Preprint at: arXiv:1912.00033. (2019).
- [45] Hoehn, P. A., Smith, A. R., & Lock, M. P. Preprint at: arXiv:2007.00580. (2020).
- [46] Hoehn, P. A., Lock, M. P., Ahmad, S. A., Smith, A. R., & Galley, T. D. Preprint at: arXiv:2103.01232. (2021).
- [47] Streiter, L. Giacomini, F. & Brukner, Č. *Physical Review Letters*, **126.23**, (2021).
- [48] Ballesteros, A., Giacomini, F. & Gubitosi, G. *Quantum*, **5**, (2021).
- [49] de la Hamette, A. C., & Galley, T. D. *Quantum*, **4**, (2020).
- [50] Krumm, M., Hoehn, P. A., & Mueller, M. P. Preprint at: arXiv:2011.01951. (2020).
- [51] Hoehn, P. A., Krumm, M & Mueller, M. P. Preprint at: arXiv:2107.07545 (2021).
- [52] Smith, A. R., & Ahmadi, M. *Nature communications*, **11(1)**, (2020).
- [53] Mikusch, M., Barbado, L. C., & Brukner, Č. Preprint at: arXiv:2103.05022. (2021).
- [54] Giacomini, F. Preprint at: arXiv:2101.11628 (2021).
- [55] Hardy, L. Preprint at: arxiv:1807.10980 (2018).
- [56] Hardy, L. Progress and Visions in Quantum Theory in view of Gravity, 189-220. (2020).
- [57] Giacomini, F., & Brukner, Č. Preprint at: arXiv:2012.13754. (2020).

- [58] Peres, A., Quantum theory: concepts and methods. *Springer Science & Business Media*. (2006).
- [59] Zanardi, P. *Physical Review Letters*, **87(7)**, (2001).
- [60] Zanardi, P., Lidar, D. A., & Lloyd, S. *Physical Review Letters*, **92(6)**, (2004).
- [61] Del Rio, L., Kraemer, L., & Renner, R. Preprint at: arXiv:1511.08818 (2015).
- [62] Chiribella, G. *Entropy* **20(5)**, (2018).
- [63] Poulin, D. *International Journal of Theoretical Physics*, **45(7)**, (2006).
- [64] Werner, R. *Journal of mathematical physics*, **27(3)**, (1986).
- [65] Giulini, D. *Annals of Physics*, **249(1)**, (1996).
- [66] Zych, M. & Greenberger, D. M. Preprint at: arXiv:1906.03725, (2019).
- [67] Greenberger, D. M. *Journal of Mathematical Physics*, **11(8)**, 2329-2340, (1970).
- [68] Greenberger, D. M. *Journal of Mathematical Physics*, **11(8)**, 23411-2347, (1970).
- [69] Greenberger, D. M. *Journal of Mathematical Physics*, **15(4)**, 395-405, (1974).
- [70] Hernandez-Coronado, H. *Foundations of Physics*, **42(10)**, (2012).
- [71] Zych, M. Costa, F. Pikovski, I. & Brukner, Č. *Nature Communications*, **10(1)**, (2019).
- [72] Budko, N.V. Samokhin, A. B. *Journal of Physics A: Mathematical and Theoretical*, **40** (2007).
- [73] de la Hamette, A-C., Galley, T. D., Höhn, P. A., Loveridge, L. & Markus P. Müller "Perspective-neutral approach to quantum frame covariance for general symmetry groups" (2021).

## Appendix A

Let  $Q$  be a generic quantum system (in the context of Section II,  $Q = AS$ ). Let  $E$  be an external reference frame with respect to which  $Q$  is described. We now show that, for any other possible external reference frame  $F$  describing  $Q$ , the invariant subalgebra  $\mathcal{L}_{\text{inv}}(\mathcal{H}_Q)$  is the largest subalgebra common to both  $\mathcal{L}(\mathcal{H}_{Q|E})$  and  $\mathcal{L}(\mathcal{H}_{Q|F})$ . This implies that  $\mathcal{L}_{\text{inv}}(\mathcal{H}_Q)$  is independent of any potential external reference frame and at the same time compatible with any such reference frame, in the sense that it is automatically a subalgebra of the larger invariant subalgebra arising from the addition of such a frame<sup>2</sup>.

Consider a Hilbert space containing all systems  $Q$ ,  $E$  and  $F$ ,  $\mathcal{H}_{EFQ} = \mathcal{H}_E \otimes \mathcal{H}_F \otimes \mathcal{H}_Q$  (defined with respect to a yet more powerful observer). By definition,  $\mathcal{L}(\mathcal{H}_{Q|E})$  is formed by operators  $T_{Q|E}$  on  $\mathcal{H}_{EFQ}$  of the form

$$T_{Q|E} = \int dg |g\rangle\langle g|_E \otimes \mathbb{1}_F \otimes U_Q(g) T_Q^{(E)} U_Q^\dagger(g). \quad (\text{A1})$$

<sup>2</sup> It is easy to see that due to the assumed transversal action of the symmetry group, enlarging a given system by tensor-multiplying it with a new system always leads to a larger invariant subsystem that contains the old one as a subsystem.

Analogously,  $\mathcal{L}(\mathcal{H}_{Q|F})$  is made of operators of the form

$$T_{Q|F} = \int dg \mathbb{1}_E \otimes |g\rangle\langle g|_F \otimes U_Q(g) T_Q^{(F)} U_Q^\dagger(g). \quad (\text{A2})$$

On the other hand, operators  $T$  on  $\mathcal{L}_{\text{inv}}(\mathcal{H}_Q)$  have the form

$$T = \mathbb{1}_E \otimes \mathbb{1}_F \otimes T_Q^{\text{inv}}, \quad (\text{A3})$$

where  $T_Q^{\text{inv}}$  is an invariant operator,  $U_Q(g) T_Q^{\text{inv}} U_Q^\dagger(g) = T_Q^{\text{inv}}$  for all  $g \in G$ . From these definitions, it is clear that  $\mathcal{L}_{\text{inv}}(\mathcal{H}_Q)$  is a common subalgebra of  $\mathcal{L}(\mathcal{H}_{Q|E})$  and  $\mathcal{L}(\mathcal{H}_{Q|F})$ , as Eq. (A3) is a particular case of both Eq. (A1) and Eq. (A2).

Now, let  $T$  be a common element of both  $\mathcal{L}(\mathcal{H}_{Q|E})$  and  $\mathcal{L}(\mathcal{H}_{Q|F})$ . We want to show that  $T \in \mathcal{L}_{\text{inv}}(\mathcal{H}_Q)$ . For any Hilbert space carrying the left- and right-regular representations of  $G$ , define  $|\Omega\rangle = \int dg |g\rangle$ . If we set Eq. (A1) equal to Eq. (A2), multiply each side of the equality by  $|\Omega\rangle\langle g|_E \otimes |\Omega\rangle\langle h|_F \otimes \mathbb{1}_Q$  for arbitrary  $g$  and  $h$ , and take the partial trace on E and F, we find that

$$U_Q(g) T_Q^{(E)} U_Q^\dagger(g) = U_Q(h) T_Q^{(F)} U_Q^\dagger(h), \quad (\text{A4})$$

for arbitrary  $g$  and  $h$ . Setting  $g = h$  gives  $T_Q^{(E)} = T_Q^{(F)} = T_Q$ . Setting  $g = e$  and  $h$  arbitrary gives  $T_Q = U_Q(h) T_Q U_Q^\dagger(h)$  for all  $h$ . This shows that  $T \in \mathcal{L}_{\text{inv}}(\mathcal{H}_Q)$ .

## Appendix B

Here we prove Eq. (11). By direct calculation, we find

$$\begin{aligned} \mathcal{V}_{E \rightarrow A}(\hat{g}_A)[L_A(g) \otimes U_S(g)] &= \int dh' dh |h'\rangle\langle h'| L_C(g) |h\rangle\langle h|_C \otimes U_{S|A}^\dagger(h') U_{S|A}(g) U_{S|A}(h) \\ &= \int dh' dh |h'\rangle_C \langle h'| gh \rangle \langle h|_C \otimes U_{S|A}^\dagger(h') U_{S|A}(g) U_{S|A}(h) \\ &= \int dh L_C(g) |h\rangle\langle h|_C \otimes U_{S|A}^\dagger(gh) U_{S|A}(g) U_{S|A}(h) \\ &= L_C(g) \otimes \mathbb{1}_{S|A}, \end{aligned} \quad (\text{B1})$$

Where we have used that  $F_{E \rightarrow A}$  is its own inverse to change the labels in the first step. This proves Eq. (11).

### Appendix C

Here we prove the identity  $\mathcal{T}_C \circ \mathcal{V}_{E \rightarrow A} = \mathcal{V}_{E \rightarrow A} \circ \mathcal{T}_{AS}$ . Let  $T_{AS}$  be an operator on  $\mathcal{H}_A \otimes \mathcal{H}_S$ . Then,

$$\begin{aligned}
\mathcal{E}_A[T_{AS}] &= \mathcal{V}_{E \rightarrow A} \circ \mathcal{T}_{AS}[T_{AS}] \\
&= U_C^\dagger(\hat{g}_{S|A}) \int dg L_C(g) \otimes U_{S|A}(g) T_{C,S|A} L^\dagger(g)_C \otimes U_{S|A}^\dagger(g) U_C(\hat{g}_{S|A}) \\
&= \int dg L_C(g) \otimes \mathbb{1}_{S|A} U_{S|A}^\dagger(\hat{g}_C) T_{C,S|A} U_{S|A}(\hat{g}_C) L_C^\dagger(g) \otimes \mathbb{1}_{S|A} \\
&= \mathcal{T}_C \circ \mathcal{V}_{E \rightarrow A}[T_{AS}].
\end{aligned} \tag{C1}$$

To pass from the second to the third line, we have multiplied by the identity in the form  $U_C(\hat{g}_{S|A}) U_C^\dagger(\hat{g}_{S|A})$  on the left of  $T$  and in the form  $U_C^\dagger(\hat{g}_{S|A}) U_C(\hat{g}_{S|A})$  on the right. Then we have used Eq. (11). This proves our claim.

For any operator  $T_{AS} = \int dg' dg |g'\rangle\langle g|_A \otimes T_S(g', g)$  in the standard partition, we can find its  $G$ -invariant version in the reference frame of Alice by applying the map  $\mathcal{E}_A$ . The answer is

$$\mathcal{E}_A[T] = \int dg' dg R_C^\dagger(g') R_C(g) \otimes U_{S|A}^\dagger(g') T_{S|A}(g', g) U_{S|A}(g). \tag{C2}$$

Proof:

$$\begin{aligned}
\mathcal{E}_A[T_{AS}] &= \mathcal{T}_C \circ \mathcal{V}_{E \rightarrow A}[T_{AS}] \\
&= \int dg' dg R_C^\dagger(g') \int dh |h\rangle\langle h|_C R_C(g) \otimes U_{S|A}^\dagger(g') T_{S|A}(g', g) U_{S|A}(g) \\
&= \int dg' dg R_C^\dagger(g') R_C(g) \otimes U_{S|A}^\dagger(g') T_{S|A}(g', g) U_{S|A}(g).
\end{aligned} \tag{C3}$$

### Appendix D

Here we compute the transformation of observables from A to B. For operators in class 1, we have

$$\begin{aligned}
\mathcal{S}_{A \rightarrow B}[\mathbb{1}_C \otimes \mathbb{1}_{B|A} \otimes T_{S|A}] &= \mathcal{V}_{E \rightarrow B} \circ \mathcal{V}_{E \rightarrow A}^\dagger[\mathbb{1}_C \otimes \mathbb{1}_{B|A} \otimes T_{S|A}] \\
&= \int dg' dg |g'\rangle\langle g'|_{A|B} \otimes |g'\rangle\langle g'|_D \otimes U_{S|B}(g') T_{S|B} U_{S|B}^\dagger(g') \\
&= \int dg |g\rangle\langle g|_{A|B} \otimes \mathbb{1}_D \otimes U_{S|B}(g) T_{S|B} U_{S|B}^\dagger(g).
\end{aligned} \tag{D1}$$

For operators in class 2, we have

$$\begin{aligned}
\mathcal{S}_{A \rightarrow B}[\mathbb{1}_C \otimes T_{B|A} \otimes \mathbb{1}_{S|A}] &= \mathcal{V}_{E \rightarrow B} \left[ \int dg |g\rangle\langle g|_A \otimes L_B(g) T_B L_B^\dagger(g) \otimes \mathbb{1}_S \right] \\
&= \int dh dg dh' |h^{-1}h'\rangle\langle g^{-1}h'|_{A|B} \otimes |h\rangle\langle (h')^{-1}h|_{T_D} |(h')^{-1}g\rangle\langle g|_D \otimes U_{S|B}^\dagger(h'h) U_{S|B}(h'g) \\
&= \int dh dg |h^{-1}\rangle\langle h|_{T_{A|B}} |g\rangle\langle g^{-1}|_{A|B} \otimes R_D(h^{-1}g) \otimes U_{S|B}(h^{-1}g),
\end{aligned} \tag{D2}$$

where we have done the changes of variables  $(h')^{-1}h \rightarrow h$  and  $(h')^{-1}g \rightarrow g$  to pass from the second equality to the third one. Finally, for operators in class 3, we have

$$\begin{aligned} \mathcal{S}_{A \rightarrow B}[T_C^R \otimes \mathbb{1}_{B|A} \otimes \mathbb{1}_{S|A}] &= \mathcal{V}_{E \rightarrow B} \left[ \int dh' dh |h'\rangle\langle h'| T_A^R |h\rangle\langle h|_A \otimes L_B(h') L_B^\dagger(h) \otimes U_S(h') U_S^\dagger(h) \right] \\ &= \int dg dh' dh |g^{-1}\rangle\langle g^{-1}|_{A|B} \otimes R_D^\dagger(g) |h'\rangle\langle h'| T_D^R |h\rangle\langle h| R_D(g) \otimes \mathbb{1}_{S|B} \\ &= \int dg |g\rangle\langle g|_{A|B} \otimes R_D(g) T_D^R R_D^\dagger(g) \otimes \mathbb{1}_{S|B}. \end{aligned} \quad (\text{D3})$$

This proves Eqs. (19).

## Appendix E

In this appendix we consider the special case of the zero-charge sector of the invariant subspace, and show how to formally obtain the transformation rule of [49] from our framework. The derivation follows the perspective neutral framework [39] applied to a general group  $G$ .

Consider 2 reference frames A and B and a quantum system S. As in the main text, the total Hilbert space decomposes into a sum of charge sectors. Suppose we have a quantum state  $|\Psi\rangle$  in the zero-charge sector of the total Hilbert space. In the standard partition, such a state satisfies  $L_A(g) \otimes L_B(g) \otimes U_S(g) |\Psi\rangle = |\Psi\rangle$  for all  $g \in G$ . Note that this condition is strictly stronger than requiring the invariance of the density matrix  $\rho$  under the action of  $G$ :  $L_A(g) \otimes L_B(g) \otimes U_S(g) \rho L_A^\dagger(g) \otimes L_B^\dagger(g) \otimes U_S^\dagger(g) = \rho$ . The state  $|\Psi\rangle$  can be obtained by "coherent group averaging" over an arbitrary state  $|\varphi\rangle = \int dg_A dg_B |g_A\rangle_A \otimes |g_B\rangle_B \otimes |\varphi(g_A, g_B)\rangle_S$ . Then we have

$$|\Psi\rangle = \int dg L_A(g) \otimes L_B(g) \otimes U_S(g) |\varphi\rangle. \quad (\text{E1})$$

As in the main text, the state in the partition natural to A is found by applying  $U_{BS}^\dagger(\hat{g}_{sfA})$  on  $|\Psi\rangle$ . The result is that the state of the reference frame A factors out for any initial state  $|\varphi\rangle$ . That is

$$U_{BS}^\dagger(\hat{g}_A) |\Psi\rangle = |\Omega\rangle_C \otimes \int dg L_{B|A}^\dagger(g) \otimes U_{S|A}^\dagger(g) |\varphi(g)\rangle_{B|A, S|A}, \quad (\text{E2})$$

where  $|\Omega\rangle = \int dg |g\rangle$ , as in Appendix A, and  $|\varphi(g)\rangle_{B|A, S|A} = \int dg' |g'\rangle_{B|A} \otimes |\varphi(g, g')\rangle_{S|A}$ . We interpret  $\int dg L_{B|A}^\dagger(g) \otimes U_{S|A}^\dagger(g) |\varphi(g)\rangle_{B|A, S|A}$  as the state of B and S "as seen" from A.

By construction, applying the operator  $S_{A \rightarrow B} = U_{AS}^\dagger(\hat{g}_B) U_{BS}(\hat{g}_A)$  to  $U_{BS}^\dagger(\hat{g}_A) |\Psi\rangle$  gives

$$U_{AS}^\dagger(\hat{g}_B) |\Psi\rangle = \int dg dg' L_{A|B}(g)^\dagger \otimes \mathbb{1}_D \otimes U_{B|S}^\dagger(g) |g'\rangle_{A|B} \otimes |\Omega\rangle_D \otimes |\varphi(g', g)\rangle_{S|B}, \quad (\text{E3})$$

Which is the analogue of Eq. (E2) with B playing the role of A. Now define

$$\hat{D} = \text{SWAP}_{AB} \circ \mathbb{1}_C \otimes \int dh |h^{-1}\rangle\langle h|_{B|A} \otimes U_{S|A}^\dagger(h), \quad (\text{E4})$$

where  $\text{SWAP}_{AB}$  is the operator that swaps A and B's Hilbert spaces. A straightforward calculation gives

$$\hat{D}U_{BS|A}^\dagger(\hat{g}_A)|\Psi\rangle = U_{AS|B}^\dagger(\hat{g}_B)|\Psi\rangle, \quad (\text{E5})$$

showing that  $S_{A \rightarrow B}$  and  $\hat{D}$  coincide in the zero-charge subspace. The operator  $\hat{D}$  is the one found in [49] up to an arbitrary exchange of the roles between the left- and right-regular representations. In [49], the state associated to the reference frame whose perspective we "jump" into is the neutral element of the group  $e$ . This can be fixed in the present perspective, up to normalisation, by conditioning the state of the reference frame to be  $|e\rangle$  [39]. In conclusion, we have shown that our results formally reduce to those of [49] in the zero-charge subspace.

## Appendix F

To appreciate the importance of the extra particle for obtaining a unitary (passive) transformation in Bob's description when Bob's reference frame is subject to a unitary (active) transformation relative to Alice, consider a simple scenario. Let A be in a classical state and let B be in the state  $|e\rangle\langle e|_{B|A}$  in the perspective of A. This means that A is also in the state  $|e\rangle\langle e|_{A|B}$  in the perspective of B (i.e., the two reference frames are aligned).

Let S be in some pure state  $|\psi\rangle\langle\psi|$ , which would be the same in both perspectives, i.e., we have  $|\psi\rangle\langle\psi|_{S|A}$  and  $|\psi\rangle\langle\psi|_{S|B}$ . If now a unitary is applied in B|A, taking the state of B relative to A to a nontrivial superposition of group states,  $|\phi\rangle\langle\phi|_{B|A}$ , where  $|\phi\rangle_{B|A} = \int dg \phi(g) |g\rangle_{B|A}$ , such that this state is not invariant under the action of the group, it is easy to see that Bob would describe the system and reference frame of Alice by the mixed state

$$\rho_{AS|B} = \int dg |\phi(g)|^2 |g^{-1}\rangle\langle g^{-1}|_{A|B} \otimes U^\dagger(g)_{S|B} |\psi\rangle\langle\psi|_{S|B} U(g)_{S|B}, \quad (\text{F1})$$

which cannot be unitarily related to the initial pure state  $|e\rangle\langle e|_{A|B} \otimes |\psi\rangle\langle\psi|_{S|B}$ . In other words, some information has been lost.

A naive attempt to recover this information by searching for it in the rest of the universe outside of A and B can be immediately seen to fail since any system S outside of A and B will be in an analogous classical correlation with A from the perspective of Bob. The resolution to this apparent paradox is that the state of AS|B is purified on the extra particle  $\overline{AS|B}$ , which is inside the invariant subsystems of ABS. This should come as no surprise since the active unitary transformation we considered was confined within this invariant subsystem.

One may nevertheless ask how come any other system in the rest of the universe gets correlated with A in the perspective of Bob if the transformation is so confined. The answer is that even though other systems in the perspective of Bob correspond to separate



subsystems, such  $A|B$ ,  $S|B$ , etc., each of these subsystems overlaps with the subsystem  $B|A$  on which the unitary acts (in the sense that the corresponding algebras do not commute), hence they would generally all be affected.

## Appendix G

Here we compute the inner product between the basis vectors  $|(\theta, a, v)\rangle$  of the regular representation of the centrally extended Galilei Group, and discuss the normalisation of states belonging to a single mass sector. For simplicity of notation, we omit the subscripts referring to the QRF perspective.

The Hilbert space of the regular representation of the centrally extended Galilei group is of the form

$$\mathcal{H} = \int^{\oplus} dm \mathcal{H}_L^{(m)} \otimes \mathcal{H}_R^{(m^*)}, \quad (\text{G1})$$

where  $\mathcal{H}_L^{(m)}$  ( $\mathcal{H}_R^{(m^*)}$ ) corresponds to the colour (flavour) degrees of freedom for mass  $m$ . The left-regular (right-regular) action of the group is trivial in  $\mathcal{H}_R^{(m^*)}$  ( $\mathcal{H}_L^{(m)}$ ) for all  $m$ . For all  $m$ , the subspace  $\mathcal{H}_L^{(m)} \otimes \mathcal{H}_R^{(m^*)}$  is spanned by vectors of the form  $|m; p, q\rangle$ , satisfying  $\langle m; p, q | m; p', q' \rangle = \delta(p-p')\delta(q-q')$ . The labels  $p$  and  $q$  are eigenvalues of the momentum operator on  $\mathcal{H}_L^{(m)}$  ( $\mathcal{H}_R^{(m^*)}$ ). The inner product of 2 states,  $|\varphi\rangle = \int^{\oplus} dm |\varphi_m\rangle$  and  $|\psi\rangle = \int^{\oplus} dm |\psi_m\rangle$  is defined by

$$\langle \varphi | \psi \rangle = \int dm \langle \varphi_m | \psi_m \rangle. \quad (\text{G2})$$

A normalised state  $|\psi\rangle$  satisfies  $\int dm \langle \psi_m | \psi_m \rangle = 1$ .

By analogy with the compact group case of Eq. (2), vectors corresponding to a fixed group element  $(\theta, a, v)$  are given by

$$|(\theta, a, v)\rangle = \int^{\oplus} dm dp \sqrt{m} (\tilde{U}^{(m)}(\theta, a, v) |m; p\rangle_L) \otimes |m; p\rangle_R, \quad (\text{G3})$$

where  $\tilde{U}^{(m)}(\theta, a, v) = e^{i\theta} e^{-i((a+vt)\hat{p}-mv\hat{x})}$ . We show that they are orthonormal in a generalised sense.

Using the Baker-Campbell-Hausdorff formula, we have

$$\tilde{U}^{(m)}(\theta, a, v) = e^{im(\theta + \frac{v}{2}(a+vt))} e^{-i(a+vt)\hat{p}} e^{imv\hat{x}}. \quad (\text{G4})$$

With the help of Eq. (G3) and Eq. (G4), we can compute straightforwardly the inner product between 2 basis elements

$$\begin{aligned} \langle (\theta', a', v') | (\theta, a, v) \rangle &= \int dm dp m \langle m, p | \tilde{U}^{(m)}(\theta - \theta' - \frac{1}{2}(av' - a'v), a - a', v - v') | m, p \rangle \\ &= \int dm dp m e^{im(s + \frac{\beta}{2}(\alpha + \beta t))} e^{-i(\alpha + \beta t)p} \delta(-m\beta), \end{aligned} \quad (\text{G5})$$

where  $s = \theta - \theta' - \frac{1}{2}(av' - a'v)$ ,  $\alpha = a - a'$  and  $\beta = v - v'$ . Because for any normalised state there will always be an integral over  $\beta$ , we can use the identity  $\delta(-m\beta) = \delta(\beta)/m$ . Performing the integral over  $p$ , and going back to the original variables, the end result is

$$\langle(\theta', a', v') | (\theta, a, v)\rangle = \delta(\theta - \theta')\delta(a - a')\delta(v - v'), \quad (\text{G6})$$

as we wanted to show.

Given the full Hilbert space  $\mathcal{H}$ , how do we represent normalisable states on a single mass sector labeled by  $m$ ? Because  $m$  is a continuous parameter, we will only be able to do this in an approximate way. Consider the states  $|\varphi_m^i\rangle = \int^\oplus dm' \sqrt{\Delta(m - m')} |\varphi_{m,m'}^i\rangle \in \mathcal{H}$ , for  $i$  running over a set of indices  $\mathcal{I}$ . Let  $\{|\varphi_{m,m'}^i\rangle\}_{i \in \mathcal{I}}$  be an orthonormal basis on  $\mathcal{H}_L^{(m')} \otimes \mathcal{H}_R^{(m'^*)}$  for each  $m'$ , and  $\Delta(m - m')$  be a sharply peaked function around  $m$ , such that we can approximate it by a Dirac delta,  $\delta(m - m')$ . Then, in this limit, we say that a normalisable state  $|\psi_m\rangle \in \mathcal{H}$  belongs to the sector of mass  $m$  if it is of the form  $|\psi_m\rangle = \sum_i \psi_m^i |\varphi_m^i\rangle$ , with  $\sum_i |\psi_m^i|^2 = 1$ . Then, formally, we can write the normalised basis states of a subspace of definite mass  $m$  as

$$|\varphi_m^i\rangle = \int^\oplus dm' \sqrt{\delta(m - m')} |\varphi_{m,m'}^i\rangle, \quad (\text{G7})$$

and the projector onto the mass  $m$  sector is given by

$$\Pi_m = \sum_i |\varphi_m^i\rangle\langle\varphi_m^i|. \quad (\text{G8})$$

Using the properties of the Dirac delta function, one can check that  $\Pi_m^2 = \Pi_m$ . For an explicit constructions of a set of functions converging to the square root of the delta function on three dimensions, see [72].

## Appendix H

Here we derive Eqs. (23) of the main text. We work in the standard partition. By definition (Eq. (5)),

$$\hat{p}_{S|A} = \int d\theta da dv |(\theta, a, v)\rangle\langle(\theta, a, v)|_A \otimes \tilde{U}_S^{(ms)}(\theta, a, v) \hat{p}_S \tilde{U}_S^{(ms)\dagger}(\theta, a, v), \quad (\text{H1a})$$

$$\hat{k}_{S|A} = \int d\theta da dv |(\theta, a, v)\rangle\langle(\theta, a, v)|_A \otimes \tilde{U}_S^{(ms)}(\theta, a, v) \hat{k}_S \tilde{U}_S^{(ms)\dagger}(\theta, a, v). \quad (\text{H1b})$$

A straightforward calculation of the second tensor factor in Eqs. (H1) gives

$$\hat{p}_{S|A} = \mathbb{1}_A \otimes \hat{p}_S - m_S \hat{v}_A^{\text{reg.}} \otimes \mathbb{1}_S, \quad (\text{H2a})$$

$$\hat{k}_{S|A} = \mathbb{1}_A \otimes \hat{x}_S + m_S \hat{a}_A^{\text{reg.}} \otimes \mathbb{1}_S, \quad (\text{H2b})$$

where

$$\hat{a}_A^{\text{reg.}} = \int d\theta da dv a |(\theta, a, v)\rangle\langle(\theta, a, v)|_A, \quad (\text{H3})$$

$$\hat{v}_A^{\text{reg.}} = \int d\theta da dv v |(\theta, a, v)\rangle\langle(\theta, a, v)|_A. \quad (\text{H4})$$

(The superscript reg. stands for "regular", as in the regular representation.) Therefore, it all amounts to calculating  $\hat{a}_A^{\text{reg.}}$  and  $\hat{v}_A^{\text{reg.}}$ .

Let us start by expressing the projector  $|(\theta, a, v)\rangle\langle(\theta, a, v)|_A$  in the basis of irreducible representations of  $\tilde{G}$ . As in the main text, we denote the left-regular subsystem of  $A$  by  $A_L$  and the right-regular subsystem by  $A_R$ . For the case of the (noncompact) extended Galilei group, the analogue to Eq. (2) is

$$|(\theta, a, v)\rangle_A = \int^{\oplus} dm dp \sqrt{m} (\tilde{U}^{(m)}(\theta, a, v) |m; p\rangle_{A_L}) \otimes |m; p\rangle_{A_R}, \quad (\text{H5})$$

where  $\hat{p}_{A_L} |m; p\rangle_{A_L} = p |m; p\rangle_{A_L}$ . Note the presence of the factor  $\sqrt{m}$ , analogue to  $\dim(q)/|\mathcal{G}|$ , which ensures that the normalisation condition is fulfilled (see Appendix G). With this identity at hand, together with  $\hat{k} = t\hat{p} - m\hat{x}$ , we can write

$$|(\theta, a, v)\rangle\langle(\theta, a, v)|_A = \int^{\oplus} dm dm' (m e^{i\theta(m-m')} e^{-i(a+vt)(p+\frac{1}{2}mv-p'-\frac{1}{2}m'v)} |m; p + mv\rangle\langle m'; p' + mv|_{A_L} \otimes |m; p\rangle\langle m'; p'|_{A_R}). \quad (\text{H6})$$

The only dependence on  $\theta$  in both  $\hat{a}_{\text{reg.}}$  and  $\hat{v}_{\text{reg.}}$  comes from the first exponential in Eq. (H6). This means we can perform the integral over  $\theta$  straight away, leading to a superselection on the mass. (We neglect factors of  $\pi$  when using the Fourier transform of the Dirac delta function.)

Doing the change of variable  $a + vt \rightarrow a$ , it follows by direct calculation that  $\hat{a}_A^{\text{reg.}} = \hat{a}_A^{*\text{reg.}} - t\hat{v}_A^{\text{reg.}}$ , where

$$\hat{a}_A^{*\text{reg.}} = \int^{\oplus} da dv dmdpd p' m a e^{-ia(p-p')} |p + mv\rangle\langle p' + mv|_{A_L} \otimes |p\rangle\langle p'|_{A_R} \quad (\text{H7})$$

At this point, the integral  $\hat{v}_A^{\text{reg.}}$  follows immediately, resulting in

$$\hat{v}_A^{\text{reg.}} = \int^{\oplus} \frac{dm}{m} (\hat{p}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} - \mathbb{1}_{A_L}^{(m)} \otimes \hat{p}_{A_R}^{(m)}). \quad (\text{H8})$$

For each irrep, labeled by  $m$ , both the left-regular and the right-regular operators  $\hat{p}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)}$  and  $\mathbb{1}_{A_L}^{(m)} \otimes \hat{p}_{A_R}^{(m)}$  are present in Eq. (H8). Importantly, this integral has a block diagonal structure due to superselection on the mass.

Finally, using  $|m; p + mv\rangle_{A_L} = e^{-i\frac{p}{m}\hat{k}} |m; mv\rangle_{A_L}$ , we can compute  $\hat{a}_A^{*\text{reg.}}$ . The result is

$$\hat{a}_A^{*\text{reg.}} = - \int^{\oplus} \frac{dm}{m} (\hat{k}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} + \mathbb{1}_{A_L}^{(m)} \otimes \hat{k}_{A_R}^{(m)}). \quad (\text{H9})$$

Putting all the pieces together, we arrive at

$$\hat{p}_{S|A} = \mathbb{1}_A \otimes \hat{p}_S - m_S \int^{\oplus} \frac{dm}{m} (\hat{p}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} - \mathbb{1}_{A_L}^{(m)} \otimes \hat{p}_{A_R}^{(m)}) \otimes \mathbb{1}_S, \quad (\text{H10a})$$

$$\hat{k}_{S|A} = \mathbb{1}_A \otimes \hat{k}_S - m_S \int^{\oplus} \frac{dm}{m} (\hat{k}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} + \mathbb{1}_{A_L}^{(m)} \otimes \hat{k}_{A_R}^{(m)}) \otimes \mathbb{1}_S, \quad (\text{H10b})$$

which are Eqs. (23).

Let us now compute the generators of the extra particle,  $\hat{p}_{\overline{S|A}}$  and  $\hat{k}_{\overline{S|A}}$ . To do this, we use a trick that is valid for quantum reference frames associated to arbitrary Lie groups  $G$ . Let  $R_A(\delta) = \int^{\oplus} dq \mathbb{1}_{A_L}^{(q)} \otimes D_{A_R}^{(q^*)}(\delta)$  be the right-regular representation of a group element  $\delta$  (for simplicity of notation, we write  $R_A(\delta)$  instead of  $R_{A|E}(\delta)$ ). Assume  $\delta$  is such that, for every value of the charge  $q$ , we can write  $D_{A_R}^{(q^*)}(\delta) = e^{i\epsilon_\delta \cdot X_{A_R}^{(q^*)}}$  for a parametrisation of  $\delta$  given by  $\epsilon_\delta$  and an infinitesimal generator  $X_{A_R}^{(q^*)}$ . By the orthogonality of the subspaces corresponding to different  $q$ 's, we can write

$$R_A(\delta) = \int^{\oplus} dq \mathbb{1}_{A_L}^{(q)} \otimes e^{i\epsilon_\delta \cdot X_{A_R}^{(q^*)}} = e^{i\epsilon_\delta \cdot X_A^R}, \quad (\text{H11})$$

where  $X_A^R = \int^{\oplus} dq \mathbb{1}_{A_L}^{(q)} \otimes X_{A_R}^{(q^*)}$ . Let us expand  $\delta$  be an element  $R_A(\delta)$  to first order in a Taylor series around  $\epsilon_\delta$ , so that  $R_A(\delta) = \mathbb{1}_A + i\epsilon_\delta \cdot X_A^R + \dots$ . Now we can use this representation of  $R_A(\delta)$  to compute  $X_{\overline{S|A}}$  from Eq. (16). There are 2 ways in which we can compute the right-hand side of Eq. (16) for the case of  $R_A(\delta)$ . We can expand to first order in  $\epsilon_\delta$  and then compute the integral, or we can compute the integral first and then expand to first order in  $\epsilon_\delta$ . Equating the order  $\epsilon_\delta$  of both Taylor series gives

$$X_{\overline{S|A}} = X_A^R \otimes \mathbb{1}_S + \mathbb{1}_A \otimes X_S. \quad (\text{H12})$$

Note that both  $X_A^R \otimes \mathbb{1}_S$  and  $\mathbb{1}_A \otimes X_S$  have an overall positive sign in Eq. (H12). This is because the right-regular representation is defined in terms of the complex-conjugate representations  $D_{A_R}^{(q^*)} = e^{i\epsilon_\delta \cdot X_{A_R}^{(q^*)}}$ , whereas  $U_S = e^{-I\epsilon_\delta \cdot X_S}$ .

Applying Eq. (H12) to the case of the centrally extended Galilei group gives immediately

$$\hat{p}_{\overline{S|A}} = \hat{p}_A^R \otimes \mathbb{1}_S + \mathbb{1}_A \otimes \hat{p}_S - m_S \int^{\oplus} \frac{dm}{m} (\hat{p}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} - \mathbb{1}_{A_L}^{(m)} \otimes \hat{p}_{A_R}^{(m)}) \otimes \mathbb{1}_S, \quad (\text{H13a})$$

$$\hat{k}_{\overline{S|A}} = \hat{k}_A^R \otimes \mathbb{1}_S + \mathbb{1}_A \otimes \hat{k}_S - m_S \int^{\oplus} \frac{dm}{m} (\hat{k}_{A_L}^{(m)} \otimes \mathbb{1}_{A_R}^{(m)} + \mathbb{1}_{A_L}^{(m)} \otimes \hat{k}_{A_R}^{(m)}) \otimes \mathbb{1}_S, \quad (\text{H13b})$$

which are Eqs. (24).

For completeness, let us write down explicitly Eqs. (19) for the infinitesimal generators of the centrally Extended Galilei group.

The case of Eq. (19a) is straightforward from the computation of the algebra  $S|A$ , as one only needs to add an extra identity operator in the Hilbert space of  $B$ . We have

already computed this algebra for the centrally extended Galilei group (see Eqs. (23) and Appendix H). The result is

$$\mathcal{S}_{A \rightarrow B}[\mathbb{1}_C \otimes \mathbb{1}_{B|A} \otimes \hat{p}_{S|A}] = \mathbb{1}_{A|B} \otimes \mathbb{1}_D \otimes \hat{p}_{S|B} - m_S \int^{\oplus} \frac{dm}{m} (\hat{p}_{A|B_L}^{(m)} \otimes \mathbb{1}_{A|B_R}^{(m)} - \mathbb{1}_{A|B_L}^{(m)} \otimes \hat{p}_{A|B_R}^{(m)}) \otimes \mathbb{1}_D \otimes \mathbb{1}_{S|B}, \quad (\text{H14a})$$

$$\mathcal{S}_{A \rightarrow B}[\mathbb{1}_C \otimes \mathbb{1}_{B|A} \otimes \hat{k}_{S|A}] = \mathbb{1}_{A|B} \otimes \mathbb{1}_D \otimes \hat{k}_{S|B} - m_S \int^{\oplus} \frac{dm}{m} (\hat{k}_{A|B_L}^{(m)} \otimes \mathbb{1}_{A|B_R}^{(m)} + \mathbb{1}_{A|B_L}^{(m)} \otimes \hat{k}_{A|B_R}^{(m)}) \otimes \mathbb{1}_D \otimes \mathbb{1}_{S|B}. \quad (\text{H14b})$$

Expressing the right-regular action in exponential form, as we did in the derivation that led to Eq. (H12), the case of Eq. (19c) follows in essentially the same way as the case of Eq. (19a), giving

$$\mathcal{S}_{A \rightarrow B}[\hat{p}_C^R \otimes \mathbb{1}_{B|A} \otimes \mathbb{1}_{S|A}] = \mathbb{1}_{A|B} \otimes \hat{p}_D^R \otimes \mathbb{1}_{S|B} - \int^{\oplus} \frac{dm}{m} (\hat{p}_{A|B_L}^{(m)} \otimes \mathbb{1}_{A|B_R}^{(m)} - \mathbb{1}_{A|B_L}^{(m)} \otimes \hat{p}_{A|B_R}^{(m)}) \otimes \hat{M}_D \otimes \mathbb{1}_{S|B}, \quad (\text{H15a})$$

$$\mathcal{S}_{A \rightarrow B}[\hat{k}_C^R \otimes \mathbb{1}_{B|A} \otimes \mathbb{1}_{S|A}] = \mathbb{1}_{A|B} \otimes \hat{k}_D^R \otimes \mathbb{1}_{S|B} - \int^{\oplus} \frac{dm}{m} (\hat{k}_{A|B_L}^{(m)} \otimes \mathbb{1}_{A|B_R}^{(m)} + \mathbb{1}_{A|B_L}^{(m)} \otimes \hat{k}_{A|B_R}^{(m)}) \otimes \hat{M}_D \otimes \mathbb{1}_{S|B}. \quad (\text{H15b})$$

Finally, we can compute the case of Eq. (19b) by means of a similar trick to that leading to Eq. (H12). That is, we can compute Eq. (19b) in 2 equivalent ways and equate the results. In the first way, we solve the integrals in Eq. (19b) for an infinitesimal transformation and then expand the result to first order in the parameter multiplying the generator. In the second way, we expand first and write down the integrals afterwards. Following this technique for the generators of the centrally extended Galilei group yields

$$\mathcal{S}_{A \rightarrow B}[\mathbb{1}_C^R \otimes \hat{p}_{B|A}^R \otimes \mathbb{1}_{S|A}] = -\hat{p}_{A|B}^L \otimes \mathbb{1}_D \otimes \mathbb{1}_{S|B} + \mathbb{1}_{A|B} \otimes \hat{p}_D^L \otimes \mathbb{1}_{S|B} - \mathbb{1}_{A|B} \otimes \mathbb{1}_D \otimes \hat{p}_{S|B}, \quad (\text{H16a})$$

$$\mathcal{S}_{A \rightarrow B}[\mathbb{1}_C^R \otimes \hat{k}_{B|A}^R \otimes \mathbb{1}_{S|A}] = -\hat{k}_{A|B}^L \otimes \mathbb{1}_D \otimes \mathbb{1}_{S|B} + \mathbb{1}_{A|B} \otimes \hat{k}_D^L \otimes \mathbb{1}_{S|B} - \mathbb{1}_{A|B} \otimes \mathbb{1}_D \otimes \hat{k}_{S|B}, \quad (\text{H16b})$$

$$\mathcal{S}_{A \rightarrow B}[\mathbb{1}_C^R \otimes \hat{M}_{B|A} \otimes \mathbb{1}_{S|A}] = -\hat{M}_{A|B} \otimes \mathbb{1}_D \otimes \mathbb{1}_{S|B} + \mathbb{1}_{A|B} \otimes \hat{M}_D \otimes \mathbb{1}_{S|B} - \mathbb{1}_{A|B} \otimes \mathbb{1}_D \otimes m_S \mathbb{1}_{S|B}, \quad (\text{H16c})$$

and

$$\begin{aligned}
\mathcal{S}_{A \rightarrow B}[\mathbb{1}_C^R \otimes \hat{p}_{B|A}^L \otimes \mathbb{1}_{S|A}] &= -\hat{p}_{A|B}^R \otimes \mathbb{1}_D \otimes \mathbb{1}_{S|B} + \mathbb{1}_{A|B} \otimes \hat{p}_D^R \otimes \mathbb{1}_{S|B} - \mathbb{1}_{A|B} \mathbb{1}_D \otimes \hat{p}_{S|B} \\
&\quad - \int^{\oplus} \frac{dm}{m} (\hat{p}_{A|B_L}^{(m)} \otimes \mathbb{1}_{A|B_R}^{(m)} - \mathbb{1}_{A|B_L}^{(m)} \otimes \hat{p}_{A|B_R}^{(m)}) \otimes \hat{M}_D \otimes \mathbb{1}_{S|B} \\
&\quad + \int^{\oplus} \frac{dm}{m} (\hat{p}_{A|B_L}^{(m)} \otimes \mathbb{1}_{A|B_R}^{(m)} - \mathbb{1}_{A|B_L}^{(m)} \otimes \hat{p}_{A|B_R}^{(m)}) \otimes \mathbb{1}_D \otimes m_S \mathbb{1}_{S|B}, \quad (\text{H17a})
\end{aligned}$$

$$\begin{aligned}
\mathcal{S}_{A \rightarrow B}[\mathbb{1}_C^R \otimes \hat{k}_{B|A}^L \otimes \mathbb{1}_{S|A}] &= -\hat{k}_{A|B}^R \otimes \mathbb{1}_D \otimes \mathbb{1}_{S|B} + \mathbb{1}_{A|B} \otimes \hat{k}_D^R \otimes \mathbb{1}_{S|B} - \mathbb{1}_{A|B} \mathbb{1}_D \otimes \hat{k}_{S|B} \\
&\quad - \int^{\oplus} \frac{dm}{m} (\hat{k}_{A|B_L}^{(m)} \otimes \mathbb{1}_{A|B_R}^{(m)} + \mathbb{1}_{A|B_L}^{(m)} \otimes \hat{k}_{A|B_R}^{(m)}) \otimes \hat{M}_D \otimes \mathbb{1}_{S|B} \\
&\quad + \int^{\oplus} \frac{dm}{m} (\hat{k}_{A|B_L}^{(m)} \otimes \mathbb{1}_{A|B_R}^{(m)} + \mathbb{1}_{A|B_L}^{(m)} \otimes \hat{k}_{A|B_R}^{(m)}) \otimes \mathbb{1}_D \otimes m_S \mathbb{1}_{S|B}. \quad (\text{H17b})
\end{aligned}$$