

Gauge-invariant perturbation theory on the Schwarzschild background spacetime Part I : — Formulation and odd-mode perturbations —

Kouji Nakamura [†]

*Gravitational-Wave Science Project, National Astronomical Observatory of Japan,
2-21-1, Osawa, Mitaka, Tokyo 181-8588, Japan*

28/10/2021

.....
This is the Part I paper of our series of papers on a gauge-invariant perturbation theory on the Schwarzschild background spacetime. After reviewing our general framework of the gauge-invariant perturbation theory, we propose the strategy of the gauge-invariant perturbation theory on the Schwarzschild spacetime. In the above general framework, the “zero-mode problem” was a remaining important problem to develop gauge-invariant perturbation theories on generic background spacetime. In perturbation theories on the Schwarzschild background spacetime, $l = 0, 1$ -mode problem corresponds to the above “zero-mode problem.” The above strategy proposed in this paper is a resolution of this $l = 0, 1$ -mode problem in perturbations on the Schwarzschild background spacetime. Following this proposal, we derive the linearized Einstein equations for any modes of $l \geq 0$ in gauge-invariant manner. We discuss the solutions to the odd-mode perturbation equations in the linearized Einstein equations and show that these perturbations include the Kerr parameter perturbation in these odd-mode perturbation, which is physically reasonable. In the Part II and Part III papers [K. Nakamura, arXiv:2110.13512 [gr-qc]; arXiv:2110.13519 [gr-qc].] of this series of papers, we will show that the even-mode solutions to the linearized Einstein equations obtained through our proposal are also physically reasonable. Then, we conclude that our proposal of the resolution of the $l = 0, 1$ -mode problem is also physically reasonable.

1. Introduction

Gravitational-wave astronomy has begun from the first event GW150914 of the direct observation of gravitational waves in 2015 [1]. This event was also the beginning of the multi-messenger astronomy including gravitational waves [2]. We are now on the stage where we can directly measure gravitational waves and we can carry out scientific research through these gravitational-wave events. We can also expect that one future direction of gravitational-wave astronomy is the development as a precise science by the detailed studies of source science, the tests of general-relativity, and the developments of the global network of gravitational-wave detectors [2–5]. In addition to the current network of ground-based detectors, as future ground-based gravitational-wave detectors, the projects of Einstein Telescope [6] and Cosmic Explorer [7] are also progressing to achieve more sensitive detections.

Besides these ground-based detectors, some projects of space gravitational-wave antenna are also progressing [8–11]. Among them, the Extreme-Mass-Ratio-Inspiral (EMRI), which is a source of

[†]E-mail address: kouji.nakamura@nao.ac.jp

gravitational waves from the motion of a stellar mass object around a supermassive black hole, is a promising target of the Laser Interferometer Space Antenna [8]. To describe the gravitational wave from EMRIs, black hole perturbations are used [12]. Furthermore, the sophistication of higher-order black hole perturbation theories is required to support these gravitational-wave physics as a precise science. Recently, the backaction effect of mass and angular momentum accretion on the Schwarzschild black hole due to the Blandford-Znajek process [13] is also discussed [14], which are higher-order effects of two-parameter perturbations [15, 16]. The motivation of this paper is in the theoretical sophistication of black hole perturbation theories toward higher-order perturbations for very wide physical situations including the topic in Ref. [14].

In the current situation of black hole perturbation theories, we may say that further sophistications are possible even in perturbation theories on the Schwarzschild background spacetime, although realistic black holes have their angular momentum, and we have to consider the perturbation theory of a Kerr black hole for direct applications to the EMRI. From the pioneering works by Regge and Wheeler [17] and Zerilli [18–20], there have been many studies on the perturbations in the Schwarzschild background spacetime [20–33]. They usually decompose the perturbations on the Schwarzschild spacetime using the spherical harmonics Y_{lm} and classify them into odd- and even-modes based on their parity, because the Schwarzschild spacetime has a spherical symmetry. However, in the current situations, $l = 0$ and $l = 1$ modes should be separately treated [30–33]. Furthermore, “*gauge-invariant*” treatments for $l = 0$ and $l = 1$ even-modes remain unknown.

On the other hand, toward unambiguous sophisticated nonlinear general-relativistic perturbation theories, we have been developing the general formulation of a higher-order gauge-invariant perturbation theory on a generic background spacetime [15, 16, 34–37] and have been applying it to cosmological perturbations [38–45]. This paper is the Part I paper of the series of papers on the application of our gauge-invariant perturbation theory on generic background spacetime to that on the Schwarzschild background spacetime. This series of papers is the full paper version of our short paper [46].

We review our framework of the gauge-invariant perturbation theory on generic background spacetime [15, 16] in Sec. 2 of this paper. This framework is based on a conjecture (Conjecture 2.1 below), which roughly states that *we already know the procedure to find gauge-invariant variables for linear-order metric perturbations*. A proof of Conjecture 2.1 was already discussed in Ref. [34–36]. In this proof, we have to introduce some Green functions for some elliptic derivative operators and ignored the kernel modes of these elliptic derivative operators due to a technical reason. We call these kernel modes “*zero modes*,” and the treatment of these zero modes remains unclear. We also called the problem of finding a treatment to these zero modes as the “*zero-mode problem*.”

In the case of the perturbations on the Schwarzschild background spacetime, as we will see in Sec. 2, these zero modes correspond to the above the $l = 0, 1$ modes. From the viewpoint of the application of our higher-order gauge-invariant perturbation theory, the above special treatments of these modes become an obstacle when we develop nonlinear perturbation theory because of the mode couplings owing to the nonlinear effects produce higher-order $l = 0, 1$ modes [47]. For this reason, the finding of a gauge-invariant treatment of $l = 0, 1$ modes in the perturbations on Schwarzschild background spacetime is quite crucial and is a resolution of the above zero-mode problem in a specific background spacetime. This resolution is an important step of the development of the higher-order gauge-invariant perturbation theory on the Schwarzschild background spacetime. In addition to the perturbation theory on a specific background spacetime, this resolution will become a clue to the perturbation theory on a generic background spacetime.

In this Part I paper, we propose a gauge-invariant treatment of the $l = 0, 1$ -mode perturbations on the Schwarzschild background spacetime and show that Conjecture 2.1 is true even for these modes. In addition, we also derive the solutions to the linearized Einstein equation for odd-mode perturbations. Following the strategy proposed in this paper, in the Part II paper [48], the strategy to solve the even-mode perturbations following Proposal 3.1 and their $l = 0, 1$ -mode solutions are discussed. Furthermore, in the Part III paper [49], we will discuss the realization of two exact solutions in terms of the linear perturbations on the Schwarzschild background spacetime. A brief discussion on the extension to the higher-order perturbations are given in Ref. [50].

The organization of this Part I paper is as follows. In Sec. 2, we briefly review the framework of the general-relativistic gauge-invariant perturbation theory within the linear perturbation theory, as mentioned above. This framework can be easily extended to higher-order perturbations [15, 16, 34–37]. In Sec. 3, we explain the situation in many studies why the special treatments of $l = 0, 1$ modes are required. Then, we propose a strategy for gauge-invariant treatments of $l = 0, 1$ modes. In Sec. 4, we construct gauge-invariant variables including $l = 0, 1$ modes through the proposal described in Sec. 3. This is a complete proof of Conjecture 2.1 for perturbations on the background spacetimes with spherical symmetry. In Sec. 5, we derive the Einstein equations for any mode perturbations following the proposal in Sec. 3. In Sec. 6, we show the strategy to solve the odd-mode perturbations and derive the explicit solutions for $l = 0, 1$ mode perturbations through the component treatment of gauge-invariant variables in the Einstein equations derived in Sec. 5. The final section 7 is devoted to the summary and discussions within this Part I paper.

Throughout this paper, we use the unit $G = c = 1$, where G is Newton’s constant of gravitation, and c is the velocity of light.

2. Review of our general-relativistic gauge-invariant perturbation theory

In this section, we briefly review our general framework of the gauge-invariant perturbation theory [15, 16]. Although the main purpose of the framework of the gauge-invariant perturbation theory developed in Refs. [15, 16] is the extension to the higher-order perturbation theory, in this review, we concentrate only on the linear perturbations. This is because we treat only the linear perturbations within this paper. Since we want to explain the gauge-invariant perturbation theory in general relativity, first of all, we have to explain the notion of “gauge” in general relativity [40, 43, 45].

General relativity is a theory with general covariance. This general covariance intuitively states that there is no preferred coordinate system in nature. This general covariance also introduces the notion of “gauge” in the theory. In the theory with general covariance, these “gauges” give rise to the unphysical degree of freedom and we have to fix the “gauges” or to extract some invariant quantities to obtain physical results. Therefore, treatments of “gauges” are crucial in general relativity and this situation becomes more delicate in general relativistic perturbation theories.

In 1964, Sachs [51] pointed out that there are two kinds of “gauges” in general relativity. Sachs called these two “gauges” as the first- and the second-kind gauges, respectively. Here, we review these concepts of “gauge,” which are different from each other. Furthermore, the distinction of these “gauges” is important to understand the ingredients of this paper. In Sec. 2.1, we first explain the notion of the first kind gauge. Second, we explain the notion of the second-kind gauge in Sec. 2.2. We expect that the reader can distinguish these two different notions of gauges in general relativity through these explanations. Then, we review our general framework of the general-relativistic gauge-invariant perturbation theory on generic background spacetimes in Sec. 2.3. We have to note




Fig. 1 The first kind gauge is a coordinate system of a single manifold.

that the aim of our general formulation of general-relativistic gauge-invariant perturbation theory is to exclude the degree of freedom of the second-kind gauge, completely.

2.1. First kind gauge

The first kind gauge is a coordinate system on a single manifold \mathcal{M} . This first kind gauge is not the “gauge” of our “gauge-invariant perturbation theory.” However, we have to explain this first kind gauge to distinguish the notions of the first-kind gauge and the second-kind gauge, as emphasized above.

In standard textbooks of manifolds (for example, see [52]), the following property of a manifold is written, “On a manifold, we can always introduce a coordinate system as a diffeomorphism ψ_α from an open set $O_\alpha \subset \mathcal{M}$ to an open set $\psi_\alpha(O_\alpha) \subset \mathbb{R}^n$ ($n = \dim \mathcal{M}$).” This diffeomorphism ψ_α , i.e., coordinate system of the open set O_α , is called *gauge choice* (of the first kind). If we consider another open set in $O_\beta \subset \mathcal{M}$, we have another gauge choice $\psi_\beta : O_\beta \mapsto \psi_\beta(O_\beta) \subset \mathbb{R}^n$ for O_β . If these two open sets O_α and O_β have the intersection $O_\alpha \cap O_\beta \neq \emptyset$, we can consider the diffeomorphism $\psi_\beta \circ \psi_\alpha^{-1}$. This diffeomorphism $\psi_\beta \circ \psi_\alpha^{-1}$ is just a coordinate transformation: $\psi_\alpha(O_\alpha \cap O_\beta) \subset \mathbb{R}^n \mapsto \psi_\beta(O_\alpha \cap O_\beta) \subset \mathbb{R}^n$, which is called *gauge transformation* (of the first kind) in general relativity. These are depicted in Fig. 1 which is a famous figure in many textbooks of the theory of manifolds.

According to the theory of a manifold, coordinate systems are not on a manifold itself but we can always introduce a coordinate system as a map from an open set in the manifold \mathcal{M} to an open set of \mathbb{R}^n . Furthermore, we may choose an different coordinate system through the different map from an open set in the manifold \mathcal{M} to an open set of \mathbb{R}^n . We can always change the coordinate system as we want. This is a realization of the statement of the general covariance that “there is no preferred coordinate system in nature.” For this reason, general covariance in general relativity is automatically included in the premise that our spacetime is regarded as a single manifold. The first kind gauge does arise due to this general covariance. The gauge issue of the first kind is usually represented by the question, “Which coordinate system is convenient?” The answer to this question depends on the problem which we are addressing, i.e., what we want to clarify. In some cases, this gauge issue of the first kind is important. On the other hand, in many cases, this gauge issue becomes harmless if we apply a covariant theory on the manifold. However, this notion of the first kind gauge appears when we interpret our results in Sec. 6 of this paper.

2.2. Second kind gauge

The *second kind gauge* appears in perturbation theories in a theory with general covariance. To explain this, we have to remind what we are doing in perturbation theories.

First, in any perturbation theories, we always treat two spacetime manifolds. One is the *physical spacetime* \mathcal{M}_{ph} . We want to describe the properties of this physical spacetime \mathcal{M}_{ph} through perturbative analyses. This physical spacetime \mathcal{M}_{ph} is usually identified with our nature itself. The other is the *background spacetime* \mathcal{M} . This background spacetime have nothing to do with our nature and is a fictitious manifold which is introduced as a reference to carry out perturbative analyses by us. We emphasize that these two spacetime manifolds \mathcal{M}_{ph} and \mathcal{M} are distinct. Let us denote the physical spacetime by $(\mathcal{M}_{\text{ph}}, \bar{g}_{ab})$ and the background spacetime by (\mathcal{M}, g_{ab}) , where \bar{g}_{ab} is the metric on the physical spacetime manifold, \mathcal{M}_{ph} , and g_{ab} is the metric on the background spacetime manifold, \mathcal{M} . Further, we formally denote the spacetime metric and the other physical tensor fields on \mathcal{M}_{ph} by Q and its background value on \mathcal{M} by Q_0 .

Second, in any perturbation theory, we always write equations for the perturbation of the variable Q as follows:

$$Q(\text{“}p\text{”}) = Q_0(p) + \delta Q(p). \quad (2.1)$$

Equation (2.1) gives a relation between variables on different manifolds. Actually, $Q(\text{“}p\text{”})$ in Eq. (2.1) is a variable on \mathcal{M}_{ph} , whereas $Q_0(p)$ and $\delta Q(p)$ are variables on \mathcal{M} . Because we regard Eq. (2.1) as a field equation, Eq. (2.1) includes an implicit assumption of the existence of a point identification map $\mathcal{M} \rightarrow \mathcal{M}_{\text{ph}} : p \in \mathcal{M} \mapsto \text{“}p\text{”} \in \mathcal{M}_{\text{ph}}$. This identification map is a *gauge choice* in general-relativistic perturbation theories (see Fig. 2). This is the notion of the *second-kind gauge* pointed out by Sachs [51]. Note that this second-kind gauge is a different notion from the degree of freedom of the coordinate transformation on a single manifold which is explained above.

To develop this understanding of the “gauge,” we introduce an infinitesimal parameter ε for perturbations and 4 + 1-dimensional manifold $\mathcal{N} = \mathcal{M}_{\text{ph}} \times \mathbb{R}$ ($4 = \dim \mathcal{M}$) such that $\mathcal{M} = \mathcal{N}|_{\varepsilon=0}$ and $\mathcal{M}_{\text{ph}} = \mathcal{M}_\varepsilon = \mathcal{N}|_{\mathbb{R}=\varepsilon}$. On \mathcal{N} , the point-identification choice is regarded as a diffeomorphism $\mathcal{X}_\varepsilon : \mathcal{N} \rightarrow \mathcal{N}$ such that $\mathcal{X}_\varepsilon : \mathcal{M} \rightarrow \mathcal{M}_\varepsilon$. This point-identification is a gauge choice of the second kind [40, 43, 45, 51, 53–55]. Furthermore, we introduce a gauge choice \mathcal{X}_ε as an exponential map with a generator $\mathcal{X} \eta^a$, which is chosen such that its integral curve in \mathcal{N} is transverse to each \mathcal{M}_ε




Fig. 2 The second kind gauge is a point-identification between the physical spacetime $\mathcal{M}_{\text{ph}} = \mathcal{M}_\varepsilon$ and the background spacetime \mathcal{M} on the extended manifold \mathcal{N} . Through Eq. (2.1), we implicitly assume the existence of a point-identification map between \mathcal{M}_ε and \mathcal{M} . However, this point-identification is not unique by virtue of the general covariance in the theory. We may choose the gauge of the second kind so that $p \in \mathcal{M}$ and “ p ” $\in \mathcal{M}_\varepsilon$ is same (\mathcal{X}_ε). We may also choose the gauge so that $q \in \mathcal{M}_0$ and “ p ” $\in \mathcal{M}_\varepsilon$ is same (\mathcal{Y}_ε). These are different gauge choices. The gauge transformation $\mathcal{X}_\varepsilon \rightarrow \mathcal{Y}_\varepsilon$ is given by the diffeomorphism $\Phi = \mathcal{X}_\varepsilon^{-1} \circ \mathcal{Y}_\varepsilon$.

everywhere on \mathcal{N} . Points lying on the same integral curve are regarded as the “same point” by the gauge choice \mathcal{X}_ε .

The first-order perturbation of the variable Q on \mathcal{M}_ε is defined as the pulled-back $\mathcal{X}_\varepsilon^* Q$ on \mathcal{M} , which is induced by \mathcal{X}_ε , and is expanded as

$$\mathcal{X}_\varepsilon^* Q = Q_0 + \varepsilon \mathcal{L}_{\mathcal{X}_\varepsilon} Q \Big|_{\mathcal{M}} + O(\varepsilon^2), \quad (2.2)$$

where $Q_0 = Q|_{\mathcal{M}}$ is the background value of Q and all terms in Eq. (2.2) are evaluated on the background spacetime \mathcal{M} . Because Eq. (2.2) is the perturbative expansion of $\mathcal{X}_\varepsilon^* Q_\varepsilon$, the first-order perturbation of Q is given by ${}^{(1)}Q := \mathcal{L}_{\mathcal{X}_\varepsilon} Q \Big|_{\mathcal{M}}$.

When we have two gauge choices \mathcal{X}_ε and \mathcal{Y}_ε with the generators $\mathcal{X}_\varepsilon \eta^a$ and $\mathcal{Y}_\varepsilon \eta^a$, respectively, and when these generators have different tangential components to each \mathcal{M}_ε , \mathcal{X}_ε and \mathcal{Y}_ε are regarded as *different gauge choices*. A *gauge-transformation* is regarded as the change of the point-identification $\mathcal{X}_\varepsilon \rightarrow \mathcal{Y}_\varepsilon$, which is given by the diffeomorphism $\Phi_\varepsilon := (\mathcal{X}_\varepsilon)^{-1} \circ \mathcal{Y}_\varepsilon : \mathcal{M} \rightarrow \mathcal{M}$. The diffeomorphism Φ_ε does change the point-identification. Here, Φ_ε induces a pull-back from the representation $\mathcal{X}_\varepsilon^* Q_\varepsilon$ to the representation $\mathcal{Y}_\varepsilon^* Q_\varepsilon$ as $\mathcal{Y}_\varepsilon^* Q_\varepsilon = \Phi_\varepsilon^* \mathcal{X}_\varepsilon^* Q_\varepsilon$. From general arguments of the Taylor

expansion [56], the pull-back Φ_ε^* is expanded as

$$\mathcal{Y}_\varepsilon^* Q_\varepsilon = \mathcal{X}_\varepsilon^* Q_\varepsilon + \varepsilon \mathcal{L}_{\xi_{(1)}} \mathcal{X}_\varepsilon^* Q_\varepsilon + O(\varepsilon^2), \quad (2.3)$$

where $\xi_{(1)}^a$ is the generator of Φ_ε . From Eqs. (2.2) and (2.3), the linear-order gauge-transformation is given as

$${}^{(1)}\mathcal{Y}Q - {}^{(1)}\mathcal{X}Q = \mathcal{L}_{\xi_{(1)}} Q_0. \quad (2.4)$$

We also employ the *order by order gauge invariance* as a concept of gauge invariance [41]. We call the k th-order perturbation ${}^{(k)}\mathcal{X}Q$ as gauge invariant if and only if ${}^{(k)}\mathcal{X}Q = {}^{(k)}\mathcal{Y}Q$ for any gauge choice \mathcal{X}_ε and \mathcal{Y}_ε .

2.3. The general-relativistic gauge-invariant linear perturbation theory

Based on the above setup, we proposed a procedure to construct gauge-invariant variables of higher-order perturbations [15, 16]. In this paper, we concentrate only on the explanations of the linear perturbations. First, we expand the metric on the physical spacetime \mathcal{M}_ε , which was pulled back to the background spacetime \mathcal{M} through a gauge choice \mathcal{X}_ε as

$$\mathcal{X}_\varepsilon^* \bar{g}_{ab} = g_{ab} + \varepsilon \mathcal{X}h_{ab} + O(\varepsilon^2). \quad (2.5)$$

Although the expression (2.5) depends entirely on the gauge choice \mathcal{X}_ε , henceforth, we do not explicitly express the index of the gauge choice \mathcal{X}_ε in the expression if there is no possibility of confusion. The important premise of our proposal was the following conjecture [15, 16] for the linear metric perturbation h_{ab} :

Conjecture 2.1. *If the gauge-transformation rule for a tensor field h_{ab} is given by ${}^{(1)}\mathcal{Y}h_{ab} - {}^{(1)}\mathcal{X}h_{ab} = \mathcal{L}_{\xi_{(1)}} g_{ab}$ with the background metric g_{ab} , there then exist a tensor field \mathcal{F}_{ab} and a vector field Y^a such that h_{ab} is decomposed as $h_{ab} =: \mathcal{F}_{ab} + \mathcal{L}_Y g_{ab}$, where \mathcal{F}_{ab} and Y^a are transformed as ${}^{(1)}\mathcal{Y}\mathcal{F}_{ab} - {}^{(1)}\mathcal{X}\mathcal{F}_{ab} = 0$ and ${}^{(1)}\mathcal{Y}Y^a - {}^{(1)}\mathcal{X}Y^a = \xi_{(1)}^a$ under the gauge transformation, respectively.*

We call \mathcal{F}_{ab} and Y^a as the *gauge-invariant* and *gauge-variant* parts of h_{ab} , respectively.

The proof of Conjecture 2.1 is highly nontrivial [34], and it was found that the gauge-invariant variables are essentially non-local. Despite this non-triviality, once we accept Conjecture 2.1, we can construct gauge-invariant variables for the linear perturbation of an arbitrary tensor field ${}^{(1)}\mathcal{X}Q$, whose gauge-transformation is given by Eq. (2.4), through the gauge-variant part of the metric perturbation Y_a in Conjecture 2.1 as

$${}^{(1)}\mathcal{Q} := {}^{(1)}\mathcal{X}Q - \mathcal{L}_Y Q_0. \quad (2.6)$$

This definition implies that the linear perturbation ${}^{(1)}\mathcal{X}Q$ of an arbitrary tensor field $\mathcal{X}_\varepsilon^* Q$ is always decomposed into its gauge-invariant part ${}^{(1)}\mathcal{Q}$ and gauge-variant part $\mathcal{L}_Y Q_0$ as

$${}^{(1)}\mathcal{X}Q = {}^{(1)}\mathcal{Q} + \mathcal{L}_Y Q_0. \quad (2.7)$$

As examples, the linearized Einstein tensor ${}^{(1)}\mathcal{X}G_a^b$ and the linear perturbation of the energy-momentum tensor ${}^{(1)}\mathcal{X}T_a^b$ are also decomposed as

$${}^{(1)}\mathcal{X}G_a^b = {}^{(1)}\mathcal{G}_a^b[\mathcal{F}] + \mathcal{L}_Y G_a^b, \quad {}^{(1)}\mathcal{X}T_a^b = {}^{(1)}\mathcal{T}_a^b + \mathcal{L}_Y T_a^b, \quad (2.8)$$

where G_{ab} and T_{ab} are the background values of the Einstein tensor and the energy-momentum tensor, respectively. The gauge-invariant part ${}^{(1)}\mathcal{G}_a^b$ of the linear-order perturbation of the Einstein

tensor is given by

$${}^{(1)}\mathcal{G}_a{}^b[A] := {}^{(1)}\Sigma_a{}^b[A] - \frac{1}{2}\delta_a{}^b{}^{(1)}\Sigma_c{}^c[A], \quad (2.9)$$

$${}^{(1)}\Sigma_a{}^b[A] := -2\nabla_{[a}H_{d]}{}^{bd}[A] - A^{cb}R_{ac}, \quad H_{ba}{}^c[A] := \nabla_{(a}A_{b)}{}^c - \frac{1}{2}\nabla^c A_{ab}. \quad (2.10)$$

Then, using the background Einstein equation $G_a{}^b = 8\pi T_a{}^b$, the linearized Einstein equation ${}^{(1)}\mathcal{G}_a{}^b = 8\pi {}^{(1)}\mathcal{T}_a{}^b$ is automatically given in the gauge-invariant form

$${}^{(1)}\mathcal{G}_a{}^b[\mathcal{F}] = 8\pi {}^{(1)}\mathcal{T}_a{}^b \quad (2.11)$$

even if the background Einstein equation is nontrivial. We also note that, in the case of a vacuum background case, i.e., $G_a{}^b = 8\pi T_a{}^b = 0$, Eq. (2.8) shows that the linear perturbations of the Einstein tensor and the energy-momentum tensor are automatically gauge-invariant.

We can also derive the perturbation of the divergence of $\bar{\nabla}_a \bar{T}_b{}^a$ of the second-rank tensor $\bar{T}_b{}^a$ on $(\mathcal{M}_{\text{ph}}, \bar{g}_{ab})$. Through the gauge choice \mathcal{X}_ε , the tensor $\bar{T}_b{}^a$ is pulled back to $\mathcal{X}_\varepsilon^* \bar{T}_b{}^a$ on the background spacetime (\mathcal{M}, g_{ab}) , and the covariant derivative operator $\bar{\nabla}_a$ on $(\mathcal{M}_{\text{ph}}, \bar{g}_{ab})$ is pulled back to a derivative operator

$$\bar{\nabla}_a := \mathcal{X}_\varepsilon^* \bar{\nabla}_a (\mathcal{X}_\varepsilon^{-1})^* \quad (2.12)$$

on (\mathcal{M}, g_{ab}) . Note that the derivative $\bar{\nabla}_a$ is the covariant derivative associated with the metric $\mathcal{X}_\varepsilon^* \bar{g}_{ab}$, whereas the derivative ∇_a on the background spacetime (\mathcal{M}, g_{ab}) is the covariant derivative associated with the background metric g_{ab} . Bearing in mind the difference in these derivatives, the first-order perturbation of $\bar{\nabla}_a \bar{T}_b{}^a$ is given by

$${}^{(1)}(\bar{\nabla}_a \bar{T}_b{}^a) = \nabla_a {}^{(1)}\mathcal{T}_b{}^a + H_{ca}{}^a[\mathcal{F}]T_b{}^c - H_{ba}{}^c[\mathcal{F}]T_c{}^a + \mathcal{L}_Y \nabla_a T_b{}^a. \quad (2.13)$$

The derivation of the formula (2.13) is given in Ref. [16]. If the tensor field $\bar{T}_b{}^a$ is the Einstein tensor $\bar{G}_a{}^b$, Eq. (2.13) yields the linear-order perturbation of the Bianchi identity

$$\nabla_a {}^{(1)}\mathcal{G}_b{}^a[\mathcal{F}] + H_{ca}{}^a[\mathcal{F}]G_b{}^c - H_{ba}{}^c[\mathcal{F}]G_c{}^a = 0 \quad (2.14)$$

and if the background Einstein tensor vanishes $G_a{}^b = 0$, we obtain the identity

$$\nabla_a {}^{(1)}\mathcal{G}_b{}^a[\mathcal{F}] = 0. \quad (2.15)$$

By contrast, if the tensor field $\bar{T}_b{}^a$ is the energy-momentum tensor, Eq. (2.13) yields the continuity equation of the energy-momentum tensor

$$\nabla_a {}^{(1)}\mathcal{T}_b{}^a + H_{ca}{}^a[\mathcal{F}]T_b{}^c - H_{ba}{}^c[\mathcal{F}]T_c{}^a = 0, \quad (2.16)$$

where we used the background continuity equation $\nabla_a T_b{}^a = 0$. If the background spacetime is vacuum $T_{ab} = 0$, Eq. (2.16) yields a linear perturbation of the energy-momentum tensor given by

$$\nabla_a {}^{(1)}\mathcal{T}_b{}^a = 0. \quad (2.17)$$

Thus, starting from the Conjecture 2.1, we can develop the gauge-invariant perturbation theory through the above framework. Furthermore, this formulation can be extended to any order perturbations [15, 16, 34, 37] from Conjecture 2.1. In this sense, the proof of the Conjecture 2.1 is crucial to this framework.




Fig. 3 A second kind gauge transformation induces a coordinate transformation. The diffeomorphism $\Psi_\alpha \circ \mathcal{X}_\epsilon^{-1}$ maps the open set $\mathcal{X}_\epsilon(O_\alpha) \subset \mathcal{M}_{\text{ph}}$ to a open set on \mathbb{R}^4 . If we change the gauge choice from \mathcal{X}_ϵ to \mathcal{Y}_ϵ , this change induces the coordinate transformation $\Psi_\alpha \circ \mathcal{X}_\epsilon^{-1}$ to $\Psi_\alpha \circ \mathcal{Y}_\epsilon^{-1}$.

We should also note that the decomposition of the metric perturbation h_{ab} into its gauge-invariant part \mathcal{F}_{ab} and into its gauge-variant part Y^a is not unique [41, 43, 45]. For example, the gauge-invariant part \mathcal{F}_{ab} has six components and we can create the gauge-invariant vector field Z^a through the component \mathcal{F}_{ab} such that the gauge-transformation of the vector field Z^a is given by $\mathcal{Y}Z^a - \mathcal{X}Z^a = 0$. Using this gauge-invariant vector field Z^a , the original metric perturbation can be expressed as follows:

$$h_{ab} = \mathcal{F}_{ab} - \mathcal{L}_Z g_{ab} + \mathcal{L}_{Z+Y} g_{ab} =: \mathcal{H}_{ab} + \mathcal{L}_X g_{ab}. \quad (2.18)$$

The tensor field $\mathcal{H}_{ab} := \mathcal{F}_{ab} - \mathcal{L}_Z g_{ab}$ is also regarded as the gauge-invariant part of the perturbation h_{ab} because $\mathcal{Y}\mathcal{H}_{ab} - \mathcal{X}\mathcal{H}_{ab} = 0$. Similarly, the vector field $X^a := Z^a + Y^a$ is also regarded as the gauge-variant part of the perturbation h_{ab} because $\mathcal{Y}X^a - \mathcal{X}X^a = \xi_{(1)}^a$. This non-uniqueness appears in the solutions derived in Sec. 6.

Finally, we comment on the relation between the gauge-transformation Φ_ϵ and the coordinate transformation [43, 45, 57, 58]. As mentioned above, the notion of the second-kind gauges above is different from the notion of the degree of freedom of the coordinate transformation on a single manifold which is called first-kind gauge. However, the gauge-transformation Φ_ϵ of the second kind induces the coordinate transformations. To see this, we introduce the coordinate system $\{O_\alpha, \Psi_\alpha\}$ on the background spacetime \mathcal{M} , where O_α are open sets on the background spacetime and Ψ_α

are diffeomorphisms from O_α to \mathbb{R}^4 ($4 = \dim \mathcal{M}$) as depicted in Fig. 3. The coordinate system $\{O_\alpha, \psi_\alpha\}$ is the set of collections of the pair of open sets O_α and diffeomorphism $O_\alpha \mapsto \mathbb{R}^4$. If we employ a gauge choice \mathcal{X}_ε of the second kind, we have the correspondence of the physical spacetime $\mathcal{M}_\varepsilon = \mathcal{M}_{\text{ph}}$ and the background spacetime \mathcal{M} . Together with the coordinate system ψ_α on \mathcal{M} , this correspondence between \mathcal{M}_ε and \mathcal{M} induces the coordinate system on \mathcal{M}_ε . Actually, $\mathcal{X}_\varepsilon(O_\alpha)$ for each α is an open set of \mathcal{M}_ε . Then, $\psi_\alpha \circ \mathcal{X}_\varepsilon^{-1}$ becomes a diffeomorphism from an open set $\mathcal{X}_\varepsilon(O_\alpha) \subset \mathcal{M}_\varepsilon$ to $\mathbb{R}^4(\{x^\mu\})$. This diffeomorphism $\psi_\alpha \circ \mathcal{X}_\varepsilon^{-1}$ induces a coordinate system of an open set on \mathcal{M}_ε . When we have two different gauge choices \mathcal{X}_ε and \mathcal{Y}_ε of the second kind, $\psi_\alpha \circ \mathcal{X}_\varepsilon^{-1} \mapsto \mathbb{R}^4(\{x^\mu\})$ and $\psi_\alpha \circ \mathcal{Y}_\varepsilon^{-1} \mapsto \mathbb{R}^4(\{y^\mu\})$ become different coordinate systems on \mathcal{M}_ε . We can also consider the coordinate transformation from the coordinate system $\psi_\alpha \circ \mathcal{X}_\varepsilon^{-1}$ to another coordinate system $\psi_\alpha \circ \mathcal{Y}_\varepsilon^{-1}$. Because the gauge transformation $\mathcal{X}_\varepsilon \rightarrow \mathcal{Y}_\varepsilon$ is induced by the diffeomorphism $\Phi_\varepsilon := (\mathcal{X}_\varepsilon)^{-1} \circ \mathcal{Y}_\varepsilon$, this diffeomorphism Φ_ε induces the coordinate transformation as

$$y^\mu(q) := x^\mu(p) = ((\Phi_\varepsilon^{-1})^* x^\mu)(q) \quad (2.19)$$

in the passive point of view [15, 56], where $p \in \mathcal{M}$, $\mathcal{X}_\varepsilon(p) = \text{“}p\text{”} \in \mathcal{M}_{\text{ph}}$ and $q \in \mathcal{M}$, $\mathcal{Y}_\varepsilon(q) = \text{“}p\text{”} \in \mathcal{M}_{\text{ph}}$. If we represent this coordinate transformation in terms of the Taylor expansion (2.3), we have the coordinate transformation

$$y^\mu(q) = x^\mu(q) - \varepsilon \xi_{(1)}^\mu(q) + O(\varepsilon^2). \quad (2.20)$$

We should emphasize that the coordinate transformation (2.20) is not the starting point of the gauge-transformation but a result of the above framework. Because our above framework of the gauge-invariant perturbation theory is constructed without a coordinate transformation (2.20), we avoid the use of the coordinate transformation (2.20) as much as possible.

3. Linear perturbations on spherically symmetric background

Here, we consider the 2+2 formulation of perturbations of a spherically symmetric background spacetime, which originally proposed by Gerlach and Sengupta [25–28]. In this formulation, we pay attention to the symmetry of the background spacetime. Spherically symmetric spacetimes are characterized by the direct product $\mathcal{M} = \mathcal{M}_1 \times S^2$ and the metric on this spacetime is given by

$$g_{ab} = y_{ab} + r^2 \gamma_{ab}, \quad (3.1)$$

$$y_{ab} = y_{AB}(dx^A)_a(dx^B)_b, \quad \gamma_{ab} = \gamma_{pq}(dx^p)_a(dx^q)_b, \quad (3.2)$$

where $x^A = (t, r)$, $x^p = (\theta, \phi)$, and γ_{pq} is the metric on the unit sphere. In the Schwarzschild spacetime, the metric (3.1) is given by

$$y_{ab} = -f(dt)_a(dt)_b + f^{-1}(dr)_a(dr)_b, \quad f := 1 - \frac{2M}{r}, \quad (3.3)$$

$$\gamma_{ab} = (d\theta)_a(d\theta)_b + \sin^2 \theta (d\phi)_a(d\phi)_b = \theta_a \theta_b + \phi_a \phi_b, \quad (3.4)$$

$$\theta_a = (d\theta)_a, \quad \phi_a = \sin \theta (d\phi)_a. \quad (3.5)$$

In Sec. 3.1, we review the conventional decomposition of the metric perturbation and its inverse relation and show that the conventional decomposition is essentially non-local and the two Green functions for the derivative operators are necessary to derive its inverse relation. The kernel modes of these derivative operators are $l = 0, 1$ modes. This is the reason why $l = 0, 1$ modes in the perturbations on the spherically symmetric background spacetime should be treated, separately. In Sec. 3.2,

we discuss the treatment in which the special treatments of these kernel modes are not necessary. To develop such treatment, we use the different scalar harmonic functions from the conventional spherical harmonic functions. We also summarize the conditions for the harmonic functions should be satisfied. In Sec. 3.3, we derive the explicit form of the mode functions. In Sec. 3.4, we propose a treatment of $l = 0, 1$ modes in perturbations on spherically symmetric background spacetime.

3.1. Conventional perturbation decomposition and its inverse relation

On the above background spacetime (\mathcal{M}, g_{ab}) , the components of the metric perturbation are given by

$$h_{ab} = h_{AB}(dx^A)_a(dx^B)_b + 2h_{Ap}(dx^A)_{(a}(dx^p)_{b)} + h_{pq}(dx^p)_a(dx^q)_b. \quad (3.6)$$

Here, we note that the components h_{AB} , h_{Ap} , and h_{pq} are regarded as components of scalar, vector, and tensor on S^2 , respectively. In many literatures, these components are decomposed through the decomposition [59–61] using the spherical harmonics $S = Y_{lm}$ as follows:

$$h_{AB} = \sum_{l,m} \tilde{h}_{AB} S, \quad (3.7)$$

$$h_{Ap} = r \sum_{l,m} [\tilde{h}_{(e1)A} \hat{D}_p S + \tilde{h}_{(o1)A} \varepsilon_{pq} \hat{D}^q S], \quad (3.8)$$

$$h_{pq} = r^2 \sum_{l,m} \left[\frac{1}{2} \gamma_{pq} \tilde{h}_{(e0)} S + \tilde{h}_{(e2)} \left(\hat{D}_p \hat{D}_q - \frac{1}{2} \gamma_{pq} \hat{D}^r \hat{D}_r \right) S + 2\tilde{h}_{(o2)} \varepsilon_{r(p} \hat{D}_{q)} \hat{D}^r S \right], \quad (3.9)$$

where \hat{D}_p is the covariant derivative associated with the metric γ_{pq} on S^2 , $\hat{D}^p = \gamma^{pq} \hat{D}_q$, $\varepsilon_{pq} = \varepsilon_{[pq]} = 2\theta_{[p} \phi_{q]}$ is the totally antisymmetric tensor on S^2 . Here, we note that the covariant derivatives of the basis θ_p and ϕ_p on S^2 are given by

$$\hat{D}_p \theta_q = \cot \theta \phi_p \phi_q, \quad \hat{D}_p \phi_q = -\cot \theta \phi_p \theta_q. \quad (3.10)$$

Through these formulae, we can check $\hat{D}_r \varepsilon_{pq} = 0$. We also note that the curvature tensors ${}^{(2)}R_{pqrs}$ and ${}^{(2)}R_{pr}$ associated with the metric γ_{pq} are given by

$${}^{(2)}R_{pqrs} = 2\gamma_{p[r} \gamma_{s]q}, \quad {}^{(2)}R_{pr} = \gamma_{pr}. \quad (3.11)$$

Although the matrix representations of the independent harmonic functions are used in the pioneer papers [17–20], these are equivalent to the covariant form (3.7)–(3.9) with the choice $S = Y_{lm}$. The choice $S = Y_{lm}$ is the starting point of the original 2+2 formulation proposed by Gerlach and Sengupta [25–28]. They showed the constructions of gauge-invariant variables for $l \geq 2$ modes and derived Einstein equations. If we apply the decomposition (3.7)–(3.9) with $S = Y_{lm}$ to the metric perturbation h_{ab} , special treatments for $l = 0, 1$ modes are required [17–28, 30–33]. This is due to the fact that the set of harmonic functions

$$\left\{ S, \hat{D}_p S, \varepsilon_{pq} \hat{D}^q S, \frac{1}{2} \gamma_{pq} S, \left(\hat{D}_p \hat{D}_q - \frac{1}{2} \gamma_{pq} \right) S, 2\varepsilon_{r(p} \hat{D}_{q)} \hat{D}^r S \right\} \quad (3.12)$$

loses its linear independence in $l = 0, 1$ cases. To clarify this situation, we consider the inverse relation of the decomposition formula (3.7)–(3.9). Furthermore, we see that the inverse-relation of the decomposition formulae (3.7)–(3.9) requires the Green functions of the derivative operators $\hat{\Delta} := \hat{D}^r \hat{D}_r$ and $\hat{\Delta} + 2 := \hat{D}^r \hat{D}_r + 2$, respectively. The eigen mode of these operators are $l = 0$ and $l = 1$, respectively. This is the reason why the special treatments for these modes are required in many literatures.

In the derivation of the inverse relation of the decomposition formulae (3.7)–(3.9), we use the orthogonality

$$\int_{S^2} d\Omega Y_{lm} Y_{l'm'} = \delta_{ll'} \delta_{mm'} \quad (3.13)$$

of the spherical harmonic function $S = Y_{lm}$, where $d\Omega = \sin\theta d\theta d\phi$. Therefore, we do not show the final expressions as the results of the application of Eq. (3.13).

First, we consider the inverse relation of the decomposition (3.8). Taking the divergence of Eq. (3.8), we obtain

$$\hat{D}^p h_{Ap} = r \sum_{l,m} \tilde{h}_{(e1)A} \hat{D}^p \hat{D}_p S = r \sum_{l,m,(l \neq 0)} \tilde{h}_{(e1)A} \hat{\Delta} S. \quad (3.14)$$

Thus, we should regard that the mode coefficient $\tilde{h}_{(e1)A}$ in Eq. (3.8) does not include $l = 0$ mode. Using the Green function $\hat{\Delta}^{-1}$, we obtain

$$\sum_{l,m,(l \neq 0)} \tilde{h}_{(e1)A} S = \frac{1}{r} \hat{\Delta}^{-1} \hat{D}^p h_{Ap}. \quad (3.15)$$

Furthermore, using the orthogonal property (3.13) of the $S = Y_{lm}$ with $l \neq 0$, we obtain the mode coefficient $\tilde{h}_{(e1)A}$ for each mode, except for $l = 0$ mode. Similarly, taking the rotation of Eq. (3.8), we obtain

$$\sum_{l,m,(l \neq 0)} \tilde{h}_{(o1)A} S = \frac{1}{r} \hat{\Delta}^{-1} \hat{D}_r (\varepsilon^{rq} h_{Aq}) \quad (3.16)$$

and the mode coefficient $\tilde{h}_{(o1)A}$ for each mode, except for $l = 0$ mode, through the orthogonal property (3.13) of the $S = Y_{lm}$ with $l \neq 0$.

The explicit form of the Green function is given by Refs. [62, 63]. The expressions (3.15) and (3.16) indicates that the decomposition (3.8) is meaningless for the modes which belongs to the kernel $\hat{\Delta} := \hat{D}^r \hat{D}_r$, i.e., $l = 0$ mode.

Next, we consider the inverse relation of (3.9). First, we note that the trace of Eq. (3.9) yields

$$\sum_{l,m} \tilde{h}_{(e0)} S = \frac{1}{r^2} \gamma^{pq} h_{pq}, \quad (3.17)$$

and the traceless part of Eq. (3.9) yields

$$\mathbb{H}_{pq}[h_{tu}] := h_{pq} - \frac{1}{2} \gamma_{pq} \gamma^{rs} h_{rs}, \quad (3.18)$$

$$= r^2 \sum_{l,m} \left[\tilde{h}_{(e2)} \left(\hat{D}_p \hat{D}_q - \frac{1}{2} \gamma_{pq} \hat{\Delta} \right) S + 2 \tilde{h}_{(o2)} \varepsilon_{r(p} \hat{D}_{q)} \hat{D}^r S \right]. \quad (3.19)$$

The mode coefficient $\tilde{h}_{(e0)}$ for each mode is obtained through the orthogonal property (3.13) of the spherical harmonics $S = Y_{lm}$ from the trace part (3.17) of h_{pq} . Therefore, we may concentrate on the traceless part (3.19) of h_{pq} . Taking the divergence of Eq. (3.19), we obtain

$$\hat{D}^p \mathbb{H}_{pq}[h_{tu}] = r^2 \sum_{l,m} \left[\tilde{h}_{(e2)} \frac{1}{2} \hat{D}_q (\hat{\Delta} + 2) S + \tilde{h}_{(o2)} \varepsilon_{rq} \hat{D}^r (\hat{\Delta} + 2) S \right] \quad (3.20)$$

$$= r^2 \sum_{l,m,(l \neq 1)} \left[\tilde{h}_{(e2)} \frac{1}{2} \hat{D}_q (\hat{\Delta} + 2) S + \tilde{h}_{(o2)} \varepsilon_{rq} \hat{D}^r (\hat{\Delta} + 2) S \right]. \quad (3.21)$$

Equation (3.21) indicates that the mode coefficients $\tilde{h}_{(e2)}$ and $\tilde{h}_{(o2)}$ do not include $l = 1$ mode if $S = Y_{lm}$ because the $l = 1$ spherical harmonic function Y_{1m} is in the kernel of the derivative operator

$\hat{\Delta} + 2$. Furthermore, we take the divergence of Eq. (3.21), and obtain

$$\hat{D}^q \hat{D}^p \mathbb{H}_{pq}[h_{tu}] = r^2 \sum_{l,m,(l \neq 1)} \left[\frac{1}{2} \tilde{h}_{(e2)} \hat{\Delta} (\hat{\Delta} + 2) S \right] = r^2 \sum_{l,m,l \geq 2} \left[\frac{1}{2} \tilde{h}_{(e2)} \hat{\Delta} (\hat{\Delta} + 2) S \right]. \quad (3.22)$$

Equation (3.22) indicates that, in addition to the $l = 1$ mode, the mode coefficient $\tilde{h}_{(e2)}$ does not include the $l = 0$ mode which is the kernel mode of the derivative operator $\hat{\Delta}$. Then, through the Green functions of the derivative operators $\hat{\Delta}$ and $(\hat{\Delta} + 2)$, we obtain the solution to Eq. (3.22) as

$$\sum_{l,m,l \geq 2} \tilde{h}_{(e2)} S = \frac{2}{r^2} [\hat{\Delta} + 2]^{-1} \hat{\Delta}^{-1} \hat{D}^q \hat{D}^p \mathbb{H}_{pq}[h_{tu}]. \quad (3.23)$$

From the orthogonal property (3.13) of the spherical harmonic function $S = Y_{lm}$ with $l \geq 2$, we obtain the mode coefficient $\tilde{h}_{(e2)}$.

On the other hand, multiplying ε^{qs} to Eq. (3.21), we obtain

$$\varepsilon^{qs} \hat{D}^p \mathbb{H}_{pq}[h_{tu}] = r^2 \sum_{l,m,(l \neq 1)} \left[\frac{1}{2} \tilde{h}_{(e2)} \varepsilon^{qs} \hat{D}_q (\hat{\Delta} + 2) S + \tilde{h}_{(o2)} \hat{D}^s (\hat{\Delta} + 2) S \right], \quad (3.24)$$

and then, taking the divergence of Eq. (3.24), we obtain

$$\varepsilon^{qs} \hat{D}_s \hat{D}^p \mathbb{H}_{pq}[h_{tu}] = r^2 \hat{\Delta} (\hat{\Delta} + 2) \sum_{l,m,(l \neq 1)} \tilde{h}_{(o2)} S = r^2 \hat{\Delta} (\hat{\Delta} + 2) \sum_{l,m,(l \geq 2)} \tilde{h}_{(o2)} S. \quad (3.25)$$

Equation (3.25) indicates that, in addition to the $l = 1$ mode, the mode coefficient $\tilde{h}_{(o2)}$ does not include the $l = 0$ mode, which is the kernel mode of the derivative operator $\hat{\Delta}$. Through the Green functions of the derivative operators $\hat{\Delta}$ and $\hat{\Delta} + 2$, we can solve Eq. (3.25) as

$$\sum_{l,m,l \geq 2} \tilde{h}_{(o2)} S = \frac{1}{r^2} [\hat{\Delta} + 2]^{-1} \hat{\Delta}^{-1} \varepsilon^{qs} \hat{D}_s \hat{D}^p \mathbb{H}_{pq}[h_{tu}]. \quad (3.26)$$

From the orthogonality property (3.13) of the spherical harmonic function $S = Y_{lm}$ with $l \neq 0, 1$, we obtain the mode coefficient $\tilde{h}_{(o2)}$.

Since the eigenvalue of the Laplacian operator $\hat{\Delta}$ on S^2 is $-l(l+1)$ with the non-negative integer l , the fact that we have to use the Green function of the operators Δ and $(\Delta + 2)$ implies that the one-to-one correspondence between the set of variables $\{h_{pq}\}$ and the set of the variables $\{h_{(e0)}, h_{(e2)}, h_{(o2)}\}$ is not guaranteed for the kernel modes $l = 0$ and $l = 1$.

We also note that the operators $\hat{\Delta}^{-1} \hat{\Delta}$ and $[\hat{\Delta} + 2]^{-1} [\hat{\Delta} + 2]$ are not identity operators but should be regarded as the projection operators. We regard that the domain of the operators $\hat{\Delta}^{-1} \hat{\Delta}$ and $[\hat{\Delta} + 2]^{-1} [\hat{\Delta} + 2]$ is the L^2 -space which is spanned by the spherical harmonics $\{Y_{lm}\}$. Since the operator $\hat{\Delta}$ eliminates the kernel

$$\mathcal{K}_{\hat{\Delta}} := \{f \in \mathcal{F} | \hat{\Delta} f = 0\}, \quad (3.27)$$

where \mathcal{F} is the function algebra, the range of the operator $\hat{\Delta}^{-1} \hat{\Delta}$ is the L^2 -space which is spanned by the spherical harmonics $\{Y_{lm} | l \neq 0\}$, i.e.,

$$\{Y_{lm} | l \neq 0\} = L^2 \setminus \mathcal{K}_{\hat{\Delta}}. \quad (3.28)$$

Similarly, the domain of the operator $[\hat{\Delta} + 2]^{-1} [\hat{\Delta} + 2]$ is the L^2 -space which is spanned by the spherical harmonics $\{Y_{lm} | l \geq 0\}$, while the kernel

$$\mathcal{K}_{\hat{\Delta}+2} := \{f \in \mathcal{F} | (\hat{\Delta} + 2)f = 0\}. \quad (3.29)$$

is excluded in the range of the operator $[\hat{\Delta} + 2]^{-1} [\hat{\Delta} + 2]$, i.e., the range of this operator is

$$\{Y_{lm} | l \neq 1\} = L^2 \setminus \mathcal{K}_{(\hat{\Delta}+2)}. \quad (3.30)$$

Namely, the operators $\hat{\Delta}^{-1} \hat{\Delta}$ and $[\hat{\Delta} + 2]^{-1} [\hat{\Delta} + 2]$ are regarded as the projection operators as

$$\hat{\Delta}^{-1} \hat{\Delta} \quad : \quad L^2 \quad \mapsto \quad L^2 \setminus \mathcal{K}_{(\hat{\Delta})} \quad (3.31)$$

$$[\hat{\Delta} + 2]^{-1} [\hat{\Delta} + 2] \quad : \quad L^2 \quad \mapsto \quad L^2 \setminus \mathcal{K}_{(\hat{\Delta}+2)}. \quad (3.32)$$

From Eqs. (3.31) and (3.32), we obtain the projection operator

$$\hat{\Delta}^{-1} [\hat{\Delta} + 2]^{-1} [\hat{\Delta} + 2] \hat{\Delta} \quad : \quad L^2 \quad \mapsto \quad L^2 \setminus \left(\mathcal{K}_{(\hat{\Delta})} \oplus \mathcal{K}_{(\hat{\Delta}+2)} \right). \quad (3.33)$$

This is a reason why we should discuss the treatments of the modes $l = 0$ and $l = 1$, separately, if we choose $S = Y_{lm}$.

3.2. Treatments of the kernel modes

As seen in Sec. 3.1, the decomposition formulae (3.7)–(3.9) with $S = Y_{lm}$ does not include the $l = 0, 1$ modes of the perturbations. In the general-relativistic gauge-invariant perturbation theory proposed in Refs. [15, 16], we assumed the separation of the linear-order metric perturbation into its gauge-invariant and gauge-variant parts, i.e., Conjecture 2.1. In Refs. [34, 36], we discuss a scenario of the proof of Conjecture 2.1 on the generic background spacetime. In this scenario of the proof, we had to use the Green functions of some elliptic differential operators. In other words, we ignored the kernel modes of these elliptic differential operators in the scenario of the proof of Conjecture 2.1 in Refs. [34, 36]. The treatment of these kernel modes was unclear at that time. We call these kernel modes as zero modes. Furthermore, we call the problem to find the treatment of these zero modes as the zero-mode problem. In the case of the perturbations on the spherically symmetric background spacetimes, the $l = 0, 1$ modes correspond to the above zero mode in Refs. [34, 36]. This is also the well-known problem as “ $l = 0, 1$ mode problem” in the treatments of perturbations on spherically symmetric background spacetimes.

Here, we consider the resolution of this $l = 0, 1$ mode problem. To carry out this, we re-examine the derivation of the inverse relations of the decomposition formulae (3.7)–(3.9), again. In this re-examination, we use the harmonic function $S = Y_{lm}$ for $l \geq 2$ model, because the set of the harmonic functions (3.12) has the linear independence at least for $l \geq 2$ mode. For $l = 0, 1$ mode, we change the harmonic function S from the spherical harmonic function Y_{00} and Y_{1m} to $k_{(\hat{\Delta})}$ and $k_{(\hat{\Delta}+2)}$, respectively, i.e., we use the harmonic functions S which are given by

$$S = S_{\delta} := \begin{cases} Y_{lm} & (l \geq 2); \\ k_{(\hat{\Delta}+2)} & (l = 1); \\ k_{(\hat{\Delta})} & (l = 0). \end{cases} \quad (3.34)$$

In this paper, we look for the explicit form of functions $k_{(\hat{\Delta})}$ and $k_{(\hat{\Delta}+2)}$ within the constraints

$$k_{(\hat{\Delta})} \in \mathcal{K}_{(\hat{\Delta})}, \quad k_{(\hat{\Delta}+2)} \in \mathcal{K}_{(\hat{\Delta}+2)}, \quad (3.35)$$

respectively. Within these domain (3.35) of the kernel modes, we specify the conditions for the functions $k_{(\hat{\Delta})}$ and $k_{(\hat{\Delta}+2)}$ to realize the independence of the set of the harmonic functions (3.12).

3.2.1. h_{pq} . Here, we first consider the decomposition of the component h_{pq} . Previously, we considered the decomposition of the component h_{pq} as Eq. (3.9):

$$h_{pq} = r^2 \sum_{l,m} \left[\frac{1}{2} \gamma_{pq} \tilde{h}_{(e0)} S + \tilde{h}_{(e2)} \left(\hat{D}_p \hat{D}_q - \frac{1}{2} \gamma_{pq} \hat{D}^r \hat{D}_r \right) S + 2 \tilde{h}_{(e2)} \varepsilon_{r(p} \hat{D}_q) \hat{D}^r S \right]. \quad (3.36)$$

As shown in Eq. (3.17), we can separate the component h_{pq} into the trace part and the traceless part. The trace part of h_{pq} is given by Eq. (3.17), which is also given by

$$\sum_{l,m,l \geq 2} \tilde{h}_{(e0,l \geq 2)} Y_{lm} + \sum_{m=-1,0,1} \tilde{h}_{(e0,l=1)} k_{(\hat{\Delta}+2)} + \tilde{h}_{(e0,l=0)} k_{(\hat{\Delta})} = \frac{1}{r^2} \gamma^{pq} h_{pq}. \quad (3.37)$$

Here, we note the effects (3.31) and (3.32) of the operators $\hat{\Delta}^{-1} \hat{\Delta}$ and $[\hat{\Delta} + 2]^{-1} [\hat{\Delta} + 2]$ as projection operators. If we apply the derivative operator $[\hat{\Delta} + 2]$ to Eq. (3.37), we obtain

$$\sum_{l,m,l \geq 2} \tilde{h}_{(e0,l \geq 2)} [\hat{\Delta} + 2] Y_{lm} + 2 \tilde{h}_{(e0,l=0)} k_{(\hat{\Delta})} = \frac{1}{r^2} [\hat{\Delta} + 2] \gamma^{pq} h_{pq}, \quad (3.38)$$

since we chose the functions $k_{(\hat{\Delta})}$ and $k_{(\hat{\Delta}+2)}$ are eigen-functions through Eqs. (3.35). Furthermore, applying the derivative operator $\hat{\Delta}$ to Eq. (3.38) as

$$\sum_{l,m,l \geq 2} \tilde{h}_{(e0,l \geq 2)} \hat{\Delta} [\hat{\Delta} + 2] Y_{lm} = \frac{1}{r^2} \hat{\Delta} [\hat{\Delta} + 2] \gamma^{pq} h_{pq}. \quad (3.39)$$

The left- and right-hand sides of Eq. (3.39) are in the domain of the Green functions $[\hat{\Delta}]^{-1}$ and $[\hat{\Delta} + 2]^{-1}$. Therefore, we may apply the Green functions $[\hat{\Delta}]^{-1}$ and $[\hat{\Delta} + 2]^{-1}$ to Eq. (3.39) and obtain

$$\sum_{l,m,l \geq 2} \tilde{h}_{(e0,l \geq 2)} Y_{lm} = \frac{1}{r^2} [\hat{\Delta} + 2]^{-1} \hat{\Delta}^{-1} \hat{\Delta} [\hat{\Delta} + 2] \gamma^{pq} h_{pq}. \quad (3.40)$$

Through the orthogonal property (3.13) of the spherical harmonic function, we obtain

$$\tilde{h}_{(e0,l \geq 2)} = \frac{1}{r^2} \int_{S^2} d\Omega Y_{lm}^* [\hat{\Delta} + 2]^{-1} \hat{\Delta}^{-1} \hat{\Delta} [\hat{\Delta} + 2] \gamma^{pq} h_{pq} =: \tilde{h}_{(e0,l \geq 2)} [h_{pq}]. \quad (3.41)$$

Thus, for $l \geq 2$, the mode coefficients $\tilde{h}_{(e0,l \geq 2)}$ is given by the functional of the original metric component h_{pq} .

Substituting Eq. (3.41) into Eq. (3.38), we obtain

$$2 \tilde{h}_{(e0,l=0)} k_{(\hat{\Delta})} = \frac{1}{r^2} [\hat{\Delta} + 2] \gamma^{pq} h_{pq} - \sum_{l,m,l \geq 2} \tilde{h}_{(e0,l \geq 2)} [h_{pq}] [\hat{\Delta} + 2] Y_{lm} =: 2 \tilde{h}_{(e0,l=0)} [h_{pq}] k_{(\hat{\Delta})}. \quad (3.42)$$

Then, the mode coefficient $\tilde{h}_{(e0,l=0)}$ is obtained as a functional of the original metric perturbation h_{pq} if $k_{(\hat{\Delta})} \neq 0$. Furthermore, from Eqs. (3.37), (3.41), and (3.42), we obtain

$$\sum_{m=-1,0,1} \tilde{h}_{(e0,l=1)} k_{(\hat{\Delta}+2)} = \frac{1}{r^2} \gamma^{pq} h_{pq} - \sum_{l,m,l \geq 2} \tilde{h}_{(e0,l \geq 2)} [h_{pq}] Y_{lm} - \tilde{h}_{(e0,l=0)} [h_{pq}] k_{(\hat{\Delta})}. \quad (3.43)$$

To resolve the degeneracy of the modes with $m = 0, \pm 1$ in Eq. (3.43), we choose $k_{(\hat{\Delta}+2)}$ as

$$k_{(\hat{\Delta}+2)} = k_{(\hat{\Delta}+2)m} = \Theta_{1m}(\theta) e^{im\phi}. \quad (3.44)$$

Through the orthogonality condition

$$\frac{1}{2\pi} \int_0^{2\pi} d\phi e^{i(m-m')\phi} = \delta_{mm'}, \quad (3.45)$$

we obtain

$$e^{+im\phi} \frac{1}{2\pi} \int_0^{2\pi} d\phi e^{-im'\phi} k_{(\hat{\Delta}+2)m} = k_{(\hat{\Delta}+2)m} \delta_{mm'}. \quad (3.46)$$

Applying the property (3.46) to Eq. (3.43), we obtain

$$\begin{aligned} \tilde{h}_{(e0,l=1)} k_{(\hat{\Delta}+2)m} &= e^{+im\phi} \frac{1}{2\pi} \int_0^{2\pi} d\phi e^{-im\phi} \\ &\quad \times \left[\frac{1}{r^2} \gamma^{pq} h_{pq} - \sum_{l,m',(l \neq 0,1)} \tilde{h}_{(e0,l \geq 2)} [h_{pq}] Y_{lm'} - \tilde{h}_{(e0,l=0)} [h_{pq}] k_{(\hat{\Delta})} \right] \\ &=: \tilde{h}_{(e0,l=1)} [h_{pq}] k_{(\hat{\Delta}+2)m}. \end{aligned} \quad (3.47)$$

Then, if $\Theta_{1m}(\theta) \neq 0$, i.e., $k_{(\hat{\Delta}+2)} \neq 0$, the mode coefficient $\tilde{h}_{(e0,l=1)}$ is given in the functional form of the original metric perturbation h_{pq} .

Thus, the mode decomposition of the trace part (3.37) of the metric perturbation h_{pq} is invertible. In this argument, we essentially used the equations (3.35) for the eigen functions and the ϕ -dependence (3.44) of the function $k_{(\hat{\Delta}+2)}$.

Next, we consider the traceless part (3.18) of h_{pq} as Eq. (3.19). Taking the divergence of Eq. (3.19), we obtain

$$\begin{aligned} \frac{1}{r^2} \hat{D}^p \mathbb{H}_{pq} [h_{tu}] &= \sum_{l,m} \left[\frac{1}{2} \tilde{h}_{(e2)} \hat{D}_q (\hat{\Delta} + 2) Y_{lm} - \tilde{h}_{(o2)} \varepsilon_{qr} \hat{D}^r (\hat{\Delta} + 2) Y_{lm} \right] \\ &= \sum_{l,m,l \geq 2} \left[\frac{1}{2} \tilde{h}_{(e2,l \geq 2)} \hat{D}_q (\hat{\Delta} + 2) Y_{lm} - \tilde{h}_{(o2,l \geq 2)} \varepsilon_{qr} \hat{D}^r (\hat{\Delta} + 2) Y_{lm} \right] \\ &\quad + \tilde{h}_{(e2,l=0)} \hat{D}_q k_{(\hat{\Delta})} - \tilde{h}_{(o2,l=0)} 2\varepsilon_{qr} \hat{D}^r k_{(\hat{\Delta})}, \end{aligned} \quad (3.48)$$

where we used Eqs. (3.11) and (3.35). We have to emphasize that the $l = 1$ mode does not appear in the expression (3.48). Taking the divergence of Eq. (3.48), again, we have

$$\frac{1}{r^2} \hat{D}^q \hat{D}^p \mathbb{H}_{pq} [h_{tu}] = \frac{1}{2} \sum_{l,m,l \geq 2} \tilde{h}_{(e2,l \geq 2)} (\hat{\Delta} + 2) \hat{\Delta} Y_{lm}, \quad (3.49)$$

where we used the property of the eigen equation for $k_{(\hat{\Delta})}$ in Eqs. (3.35). Through the Green functions $\hat{\Delta}^{-1}$ and $[\hat{\Delta} + 2]^{-1}$ and the orthogonal property (3.13) of the spherical harmonics Y_{lm} , we obtain the same result as Eq. (3.26) and the mode coefficient $\tilde{h}_{(e2,l \geq 2)}$ of each mode is given in a functional form of the original metric perturbation h_{tu} as

$$\tilde{h}_{(e2,l \geq 2)} = \frac{2}{r^2} \int_{S^2} d\Omega Y_{lm}^* [\hat{\Delta}]^{-1} [\hat{\Delta} + 2]^{-1} \hat{D}^q \hat{D}^p \mathbb{H}_{pq} [h_{tu}] =: \tilde{h}_{(e2,l \geq 2)} [h_{tu}]. \quad (3.50)$$

On the other hand, taking the rotation of Eq. (3.48) and use the eigen equation for $k_{(\hat{\Delta})}$ in Eqs. (3.35), Green functions $[\hat{\Delta}]^{-1}$ and $[\hat{\Delta} + 2]$, and the orthogonal properties (3.13) of the spherical harmonics Y_{lm} , we obtain the mode coefficient $\tilde{h}_{(o2,l \geq 2)}$ in the functional form of the original metric perturbation

h_{tu} as

$$\tilde{h}_{(o2,l \geq 2)} = \frac{1}{r^2} \int_{S^2} d\Omega Y_{lm}^* [\hat{\Delta}]^{-1} [\hat{\Delta} + 2]^{-1} \varepsilon^{ps} \hat{D}_s \hat{D}^q \mathbb{H}_{pq} [h_{tu}] =: \tilde{h}_{(o2,l \geq 2)} [h_{tu}]. \quad (3.51)$$

Substituting Eqs. (3.50) and (3.51) into Eq. (3.48), we obtain

$$\begin{aligned} & \tilde{h}_{(e2,l=0)} \hat{D}_q k_{(\hat{\Delta})} - \tilde{h}_{(o2,l=0)} 2\varepsilon_{qr} \hat{D}^r k_{(\hat{\Delta})} \\ &= - \sum_{l,m,l \geq 2} \left[\frac{1}{2} \tilde{h}_{(e2,l \geq 2)} [h_{tu}] \hat{D}_q (\hat{\Delta} + 2) Y_{lm} - \tilde{h}_{(o2,l \geq 2)} [h_{tu}] \varepsilon_{qr} \hat{D}^r (\hat{\Delta} + 2) Y_{lm} \right] \\ & \quad + \frac{1}{r^2} \hat{D}^p \mathbb{H}_{pq} [h_{tu}]. \end{aligned} \quad (3.52)$$

If $\hat{D}_q k_{(\hat{\Delta})} \neq 0$, the vectors $\hat{D}_q k_{(\hat{\Delta})}$ and $\varepsilon_{qr} \hat{D}^r k_{(\hat{\Delta})}$ are orthogonal to each other. Then, we have

$$\begin{aligned} \tilde{h}_{(e2,l=0)} &= \tilde{h}_{(e2,l=0)} [h_{tu}] \\ &:= \frac{\left(\hat{D}^q k_{(\hat{\Delta})} \right)}{\left(\hat{D}_s k_{(\hat{\Delta})} \right) \left(\hat{D}^s k_{(\hat{\Delta})} \right)} \left[\frac{1}{r^2} \hat{D}^p \mathbb{H}_{pq} [h_{tu}] - \sum_{l,m,l \geq 2} \left\{ \frac{1}{2} \tilde{h}_{(e2,l \geq 2)} [h_{tu}] \hat{D}_q (\hat{\Delta} + 2) Y_{lm} \right. \right. \\ & \quad \left. \left. - \tilde{h}_{(o2,l \geq 2)} [h_{tu}] \varepsilon_{qr} \hat{D}^r (\hat{\Delta} + 2) Y_{lm} \right\} \right] \end{aligned} \quad (3.53)$$

and

$$\begin{aligned} \tilde{h}_{(o2,l=0)} &= \tilde{h}_{(o2,l=0)} [h_{tu}] \\ &:= \frac{\left(\varepsilon^{qr} \hat{D}_r k_{(\hat{\Delta})} \right)}{\left(\hat{D}_s k_{(\hat{\Delta})} \right) \left(\hat{D}^s k_{(\hat{\Delta})} \right)} \left[\frac{1}{r^2} \hat{D}^p \mathbb{H}_{pq} [h_{tu}] - \sum_{l,m,l \geq 2} \left\{ \frac{1}{2} \tilde{h}_{(e2,l \geq 2)} [h_{tu}] \hat{D}_q (\hat{\Delta} + 2) Y_{lm} \right. \right. \\ & \quad \left. \left. - \tilde{h}_{(o2,l \geq 2)} [h_{tu}] \varepsilon_{qr} \hat{D}^r (\hat{\Delta} + 2) Y_{lm} \right\} \right]. \end{aligned} \quad (3.54)$$

Now, we return to the original definition (3.19) of the traceless part \mathbb{H}_{pq} . From Eqs. (3.18), (3.50), (3.51), (3.53), and (3.54), we obtain

$$\begin{aligned} & \sum_{m=-1,0,1} \left[\tilde{h}_{(e2,l=1,m)} \left(\hat{D}_p \hat{D}_q - \frac{1}{2} \gamma_{pq} \hat{\Delta} \right) k_{(\hat{\Delta}+2)} + 2\tilde{h}_{(o2,l=1,m)} \varepsilon_{r(p} \hat{D}_q) \hat{D}^r k_{(\hat{\Delta}+2)} \right] \\ &= \frac{1}{r^2} \mathbb{H}_{pq} [h_{tu}] - \left[\sum_{l,m,l \geq 2} \left\{ \tilde{h}_{(e2)} [h_{tu}] \left(\hat{D}_p \hat{D}_q - \frac{1}{2} \gamma_{pq} \hat{\Delta} \right) Y_{lm} + 2\tilde{h}_{(o2)} [h_{tu}] \varepsilon_{r(p} \hat{D}_q) \hat{D}^r Y_{lm} \right\} \right. \\ & \quad \left. + \tilde{h}_{(e2,l=0)} [h_{tu}] \left(\hat{D}_p \hat{D}_q - \frac{1}{2} \gamma_{pq} \hat{\Delta} \right) k_{(\hat{\Delta})} + 2\tilde{h}_{(o2,l=0)} [h_{tu}] \varepsilon_{r(p} \hat{D}_q) \hat{D}^r k_{(\hat{\Delta})} \right] \\ &=: H_{(\hat{\Delta}+2)pq} [h_{tu}]. \end{aligned} \quad (3.55)$$

To simplify the notation, we define

$$K_{(m)pq} := \left(\hat{D}_p \hat{D}_q - \frac{1}{2} \gamma_{pq} \hat{D}^r \hat{D}_r \right) k_{(\hat{\Delta}+2)m}, \quad J_{(m)pq} := 2\varepsilon_{r(p} \hat{D}_q) \hat{D}^r k_{(\hat{\Delta}+2)m}, \quad (3.56)$$

and we evaluate $K_{(m)pq}K_{(m')^{pq}}^{pq}$, $J_{(m)pq}K_{(m')^{pq}}^{pq}$, and $J_{(m)pq}J_{(m')^{pq}}^{pq}$, which are given by

$$K_{(m)pq}K_{(m')^{pq}}^{pq} = \left(\hat{D}^p \hat{D}^q k_{(\hat{\Delta}+2)m} \right) \left(\hat{D}^p \hat{D}^q k_{(\hat{\Delta}+2)m'} \right) - 2 \left(k_{(\hat{\Delta}+2)m} \right) \left(k_{(\hat{\Delta}+2)m'} \right), \quad (3.57)$$

$$J_{(m)pq}K_{(m')^{pq}}^{pq} = 2\varepsilon^{rp} \hat{D}_r \hat{D}_q k_{(\hat{\Delta}+2)m} \hat{D}_p \hat{D}^q k_{(\hat{\Delta}+2)m'}, \quad (3.58)$$

$$J_{(m)pq}J_{(m')^{pq}}^{pq} = 4 \left[\left(\hat{D}^p \hat{D}^q k_{(\hat{\Delta}+2)m} \right) \left(\hat{D}_p \hat{D}_q k_{(\hat{\Delta}+2)m'} \right) - 2 \left(k_{(\hat{\Delta}+2)m} \right) \left(k_{(\hat{\Delta}+2)m'} \right) \right]. \quad (3.59)$$

To carry out the resolution of the degeneracy in Eq. (3.55), we use the property (3.44) of the function $k_{(\hat{\Delta}+2)}$. From the property (3.44), we have

$$\hat{D}_p k_{(\hat{\Delta}+2)m} = \left(\frac{d}{d\theta} \Theta_m(\theta) \right) e^{im\phi} \theta_p + \frac{im}{\sin \theta} \Theta_m(\theta) e^{im\phi} \phi_p \quad (3.60)$$

and

$$\begin{aligned} \hat{D}_p \hat{D}_q k_{(\hat{\Delta}+2)m} &= \left(\frac{d^2}{d\theta^2} \Theta_m(\theta) \right) e^{im\phi} \theta_p \theta_q \\ &+ \left[\left(\frac{d}{d\theta} \Theta_m(\theta) \right) \cot \theta - m^2 \frac{1}{\sin^2 \theta} \Theta_m(\theta) \right] e^{im\phi} \phi_p \phi_q \\ &+ im \frac{1}{\sin \theta} \left[\frac{d}{d\theta} \Theta_m(\theta) - \cot \theta \Theta_m(\theta) \right] e^{im\phi} 2\theta_{(p} \phi_{q)}. \end{aligned} \quad (3.61)$$

From Eq. (3.61), we obtain

$$\begin{aligned} K_{(m)pq} &:= \left(\hat{D}_p \hat{D}_q - \frac{1}{2} \gamma_{pq} \hat{\Delta} \right) k_{(\hat{\Delta}+2)m} \\ &= -(\theta_p \theta_q - \phi_p \phi_q) \left[\cot \theta \frac{d}{d\theta} \Theta_m(\theta) + \left(1 - \frac{m^2}{\sin^2 \theta} \right) \Theta_m(\theta) \right] e^{im\phi} \\ &+ 2\theta_{(p} \phi_{q)} \frac{im}{\sin \theta} \left(\frac{d}{d\theta} \Theta_m(\theta) - \cot \theta \Theta_m(\theta) \right) e^{im\phi}, \end{aligned} \quad (3.62)$$

where we used $(\hat{\Delta} + 2)k_{(\hat{\Delta}+2)m} = 0$, i.e.,

$$\frac{d^2}{d\theta^2} \Theta_m(\theta) + \cot \theta \frac{d}{d\theta} \Theta_m(\theta) + \left(2 - \frac{m^2}{\sin^2 \theta} \right) \Theta_m(\theta) = 0. \quad (3.63)$$

From the expression of the components $K_{(m')pq}$, $J_{(m')pq}$, θ_p , and ϕ_p , we can confirm

$$\frac{1}{2\pi} \int_0^{2\pi} d\phi e^{-im\phi} K_{(m')pq} = K_{(m)pq} e^{-im\phi} \delta_{mm'}, \quad (3.64)$$

$$\frac{1}{2\pi} \int_0^{2\pi} d\phi e^{-im\phi} J_{(m')pq} = J_{(m)pq} e^{-im\phi} \delta_{mm'}. \quad (3.65)$$

Furthermore, straightforward calculations yield

$$K_{(m)pq}K_{(m)}^{pq} = \left(\hat{D}_p \hat{D}_q k_{(\hat{\Delta}+2)m} \right) \left(\hat{D}^p \hat{D}^q k_{(\hat{\Delta}+2)m} \right) - 2 \left(k_{(\hat{\Delta}+2)m} \right)^2, \quad (3.66)$$

$$J_{(m)pq}K_{(m)}^{pq} = 0, \quad (3.67)$$

$$J_{(m)pq}J_{(m)}^{pq} = 4K_{(m)pq}K_{(m)}^{pq}. \quad (3.68)$$

Through Eqs. (3.64) and (3.65), we can consider the resolution of the m -degeneracy of $l = 1$ mode in Eq. (3.55) as follows:

$$\begin{aligned}
& \frac{1}{2\pi} e^{+im\phi} \int_0^{2\pi} d\phi e^{-im\phi} H_{(\hat{\Delta}+2)pq}[h_{tu}] \\
&= \sum_{m'=-1,0,1} \left[\tilde{h}_{(e2,l=1,m)} e^{+im\phi} \frac{1}{2\pi} \int_0^{2\pi} d\phi e^{-im\phi} K_{(m')pq} \right. \\
&\quad \left. + \tilde{h}_{(o2,l=1,m)} e^{+im\phi} \frac{1}{2\pi} \int_0^{2\pi} d\phi e^{-im\phi} J_{(m')pq} \right] \\
&= \sum_{m'=-1,0,1} \left[\tilde{h}_{(e2,l=1,m)} K_{(m)pq} \delta_{mm'} + \tilde{h}_{(o2,l=1,m)} J_{(m)pq} \delta_{mm'} \right] \\
&= \tilde{h}_{(e2,l=1,m)} K_{(m)pq} + \tilde{h}_{(o2,l=1,m)} J_{(m)pq}. \tag{3.69}
\end{aligned}$$

Furthermore, from Eqs. (3.66)–(3.68), we obtain

$$\tilde{h}_{(e2,l=1,m)} = [K_{(m)pq} K_{(m)}^{pq}]^{-1} K_{(m)}^{pq} \frac{1}{2\pi} e^{+im\phi} \int_0^{2\pi} d\phi e^{-im\phi} H_{(\hat{\Delta}+2)pq}[h_{tu}] =: \tilde{h}_{(e2,l=1,m)}[h_{tu}], \tag{3.70}$$

and

$$\tilde{h}_{(o2,l=1,m)} = \frac{1}{4} [K_{(m)pq} K_{(m)}^{pq}]^{-1} J_{(m)}^{pq} \frac{1}{2\pi} e^{+im\phi} \int_0^{2\pi} d\phi e^{-im\phi} H_{(\hat{\Delta}+2)pq}[h_{tu}] =: \tilde{h}_{(o2,l=1,m)}[h_{tu}]. \tag{3.71}$$

Thus, we have obtained the mode coefficients $\tilde{h}_{(e2,l=1,m)}$ and $\tilde{h}_{(o2,l=1,m)}$ in the functional forms of the original metric h_{tu} .

Here, we summarize the conditions for the eigen functions $k_{(\hat{\Delta})}$ and $k_{(\hat{\Delta}+2)}$ to obtain the inverse relation of the metric decomposition (3.36). To obtain the inverse relations of the mode decomposition of the trace and the traceless-part of Eq. (3.36), we use the conditions

$$k_{(\hat{\Delta})} \in \mathcal{K}_{(\hat{\Delta})}, \quad k_{(\hat{\Delta}+2)} \in \mathcal{K}_{(\hat{\Delta}+2)}, \quad k_{(\hat{\Delta}+2)} = k_{(\hat{\Delta}+2)m} = \Theta_{1m}(\theta) e^{im\phi}, \tag{3.72}$$

$$\left(\hat{D}_p k_{(\hat{\Delta})} \right) \left(\hat{D}^p k_{(\hat{\Delta})} \right) \neq 0, \tag{3.73}$$

$$K_{(m)pq} K_{(m)}^{pq} = \left(\hat{D}_p \hat{D}_q k_{(\hat{\Delta}+2)m} \right) \left(\hat{D}^p \hat{D}^q k_{(\hat{\Delta}+2)m} \right) - 2 \left(k_{(\hat{\Delta}+2)m} \right)^2 \neq 0. \tag{3.74}$$

The condition (3.74) implies the nonvanishing $K_{(m)pq}$ and $J_{(m)pq}$.

3.2.2. h_{Ap} . Next, we consider the inversion relation of the decomposition (3.8) taking account of the kernel modes $k_{(\hat{\Delta})}$ and $k_{(\hat{\Delta}+2)}$.

$$h_{Ap} = r \sum_{l,m} \left[\tilde{h}_{(e1)A} \hat{D}_p S + \tilde{h}_{(o1)A} \epsilon_{pq} \hat{D}^q S \right] \tag{3.75}$$

$$\begin{aligned}
&= r \sum_{l,m,l \geq 2} \left[\tilde{h}_{(e1,l \geq 2)A} \hat{D}_p Y_{lm} + \tilde{h}_{(o1,l \geq 2)A} \epsilon_{pq} \hat{D}^q Y_{lm} \right] \\
&\quad + r \sum_m \left[\tilde{h}_{(e1,l=1)A} \hat{D}_p k_{(\hat{\Delta}+2)} + \tilde{h}_{(o1,l=1)A} \epsilon_{pq} \hat{D}^q k_{(\hat{\Delta}+2)} \right] \\
&\quad + r \left[\tilde{h}_{(e1,l=0)A} \hat{D}_p k_{(\hat{\Delta})} + \tilde{h}_{(o1,l=0)A} \epsilon_{pq} \hat{D}^q k_{(\hat{\Delta})} \right]. \tag{3.76}
\end{aligned}$$

Taking the divergence of Eq. (3.76), we obtain

$$\hat{D}^p h_{Ap} = r \sum_{l,m,l \geq 2} \tilde{h}_{(e1,l \geq 2)A} \hat{\Delta} Y_{lm} - 2r \sum_m \tilde{h}_{(e1,l=1,m)A} k_{(\hat{\Delta}+2)}. \tag{3.77}$$

Applying the derivative operator $\hat{\Delta} + 2$ to Eq. (3.77), we obtain

$$[\hat{\Delta} + 2]\hat{D}^p h_{Ap} = r \sum_{l,m,l \geq 2} \tilde{h}_{(e1,l \geq 2)A} [\hat{\Delta} + 2]\hat{\Delta} Y_{lm}. \quad (3.78)$$

Using the Green functions $[\hat{\Delta} + 2]^{-1}$, $\hat{\Delta}^{-1}$, and the orthogonal property (3.13) of the spherical harmonics Y_{lm} , we obtain

$$\tilde{h}_{(e1,l \geq 2)A} = \frac{1}{r} \int_{S^2} Y_{lm}^* \hat{\Delta}^{-1} [\hat{\Delta} + 2]^{-1} [\hat{\Delta} + 2]\hat{D}^p h_{Ap} =: \tilde{h}_{(e1,l \geq 2)A} [h_{Bs}]. \quad (3.79)$$

Thus, the mode coefficient $\tilde{h}_{(e1)A}$ is given in the form of the functional of the original metric component h_{Ap} . Through Eq. (3.79), Eq. (3.77) is expressed as

$$\sum_m \tilde{h}_{(e1,l=1,m)A} k_{(\hat{\Delta}+2)} = \frac{1}{2} \sum_{l,m,l \geq 2} \tilde{h}_{(e1,l \geq 2)A} [h_{Br}] \hat{\Delta} Y_{lm} - \frac{1}{2r} \hat{D}^p h_{Ap}. \quad (3.80)$$

To resolve the m -degeneracy of Eq. (3.80), we use Eq. (3.72) and (3.46). Then, we have

$$\begin{aligned} \tilde{h}_{(e1,l=1,m)A} &= \frac{e^{im\phi}}{k_{(\hat{\Delta}+2)m}} \frac{1}{4\pi} \int_0^{2\pi} d\phi e^{-im'\phi} \left[\sum_{l,m',l \geq 2} \tilde{h}_{(e1,l \geq 2)A} [h_{Br}] \hat{\Delta} Y_{lm'} - \frac{1}{r} \hat{D}^p h_{Ap} \right] \\ &=: \tilde{h}_{(e1,l=1,m)A} [h_{Bs}] \end{aligned} \quad (3.81)$$

On the other hand, taking the rotation of Eq. (3.76), we have

$$\varepsilon^{pq} \hat{D}_q h_{Ap} = r \sum_{l,m,l \geq 2} [\tilde{h}_{(o1,l \geq 2)A} \hat{\Delta} Y_{lm}] - 2r \sum_m [\tilde{h}_{(o1,l=1)A} k_{(\hat{\Delta}+2)}]. \quad (3.82)$$

As in the case of Eq. (3.79), we have

$$\tilde{h}_{(o1,l \geq 2)A} = \frac{1}{r} \int_{S^2} Y_{lm}^* \hat{\Delta}^{-1} [\hat{\Delta} + 2]^{-1} [\hat{\Delta} + 2] \varepsilon^{pq} \hat{D}_q h_{Ap} =: \tilde{h}_{(o1,l \geq 2)A} [h_{Br}], \quad (3.83)$$

$$\begin{aligned} \tilde{h}_{(o1,l=1)A} &= \frac{e^{im\phi}}{k_{(\hat{\Delta}+2)m}} \frac{1}{4\pi} \int_0^{2\pi} d\phi e^{-im'\phi} \left[\sum_{l,m',l \geq 2} \{ \tilde{h}_{(o1,l \geq 2)A} [h_{Br}] \hat{\Delta} Y_{lm'} \} - \frac{1}{r} \varepsilon^{pq} \hat{D}_q h_{Ap} \right] \\ &=: \tilde{h}_{(o1,l=1)A} [h_{Bs}]. \end{aligned} \quad (3.84)$$

Through Eqs. (3.79), (3.81), (3.83), and (3.84), we obtain

$$\begin{aligned} &\tilde{h}_{(e1,l=0)A} \hat{D}_p k_{(\hat{\Delta})} + \tilde{h}_{(o1,l=0)A} \varepsilon_{pq} \hat{D}^q k_{(\hat{\Delta})} \\ &= \frac{1}{r} h_{Ap} - \sum_{l,m,l \geq 2} [\tilde{h}_{(e1,l \geq 2)A} [h_{Bs}] \hat{D}_p Y_{lm} + \tilde{h}_{(o1,l \geq 2)A} [h_{Bs}] \varepsilon_{pq} \hat{D}^q Y_{lm}] \\ &\quad - \sum_m [\tilde{h}_{(e1,l=1)A} [h_{Bs}] \hat{D}_p k_{(\hat{\Delta}+2)} + \tilde{h}_{(o1,l=1)A} [h_{Bs}] \varepsilon_{pq} \hat{D}^q k_{(\hat{\Delta}+2)}] \end{aligned} \quad (3.85)$$

$$=: \mathbb{H}_{Ap} [h_{Bs}]. \quad (3.86)$$

Here, we use the condition (3.73). Then, we have

$$\tilde{h}_{(e1,l=0)A} = \left[\left(\hat{D}^q k_{(\hat{\Delta})} \right) \left(\hat{D}_q k_{(\hat{\Delta})} \right) \right]^{-1} \hat{D}^p k_{(\hat{\Delta})} \mathbb{H}_{Ap} [h_{Bs}] =: \tilde{h}_{(e1,l=0)A} [h_{Bs}], \quad (3.87)$$

$$\tilde{h}_{(o1,l=0)A} = \left[\left(\hat{D}^r k_{(\hat{\Delta})} \right) \left(\hat{D}_r k_{(\hat{\Delta})} \right) \right]^{-1} \varepsilon^{pq} \hat{D}_q k_{(\hat{\Delta})} \mathbb{H}_{Ap} [h_{Bs}] =: \tilde{h}_{(o1,l=0)A} [h_{Bs}]. \quad (3.88)$$

Thus, we have shown that the mode coefficients $\tilde{h}_{(e1)A}$ and $\tilde{h}_{(o1)A}$ for all $l \geq 0$ modes are given in the functional forms (3.79), (3.81), (3.83), (3.84), (3.87), and (3.88) of the original metric h_{Ap} under the conditions (3.72)–(3.74).

3.2.3. h_{AB} . Through the harmonic functions Y_{lm} ($l \geq 2$), $k_{(\hat{\Delta}+2)m}$, and $k_{(\hat{\Delta})}$, the component h_{AB} of the metric perturbation h_{ab} is decomposed as

$$h_{AB} = \sum_{l,m(l \geq 2)} \tilde{h}_{(l \geq 2)AB} S + \sum_{m=-1,0,1} \tilde{h}_{(l=1,m)AB} k_{(\hat{\Delta}+2)m} + \tilde{h}_{(l=0)AB} k_{(\hat{\Delta})}. \quad (3.89)$$

This decomposition has the same form as Eq. (3.37) for the trace part of the component h_{pq} . Then, we obtain the inverse relations

$$\tilde{h}_{(l \geq 2)AB} = \int_{S^2} d\Omega Y_{lm}^* [\hat{\Delta} + 2]^{-1} \hat{\Delta}^{-1} \hat{\Delta} [\hat{\Delta} + 2] h_{AB} =: \tilde{h}_{(l \geq 2)AB} [h_{AB}], \quad l \geq 2, \quad (3.90)$$

$$\begin{aligned} \tilde{h}_{(l=0)AB} &= \frac{1}{2k_{(\hat{\Delta})}} \left[[\hat{\Delta} + 2] h_{AB} - \sum_{l,m,(l \neq 0,1)} \tilde{h}_{(l \geq 2)AB} [h_{AB}] [\hat{\Delta} + 2] Y_{lm} \right] \\ &=: \tilde{h}_{(l=0)AB} [h_{AB}], \end{aligned} \quad (3.91)$$

$$\begin{aligned} \tilde{h}_{(l=1,m)AB} &= \frac{1}{k_{(\hat{\Delta}+2)m}} e^{+im\phi} \frac{1}{2\pi} \int_0^{2\pi} d\phi e^{-im\phi} \\ &\quad \times \left[\frac{1}{r^2} h_{AB} - \sum_{l,m',(l \neq 0,1)} \tilde{h}_{(l \geq 2)AB} [h_{AB}] Y_{lm'} - \tilde{h}_{(l=0)AB} [h_{AB}] k_{(\hat{\Delta})} \right] \\ &=: \tilde{h}_{(l=1)AB} [h_{AB}]. \end{aligned} \quad (3.92)$$

which correspond to Eqs. (3.37), (3.42), and (3.47), respectively.

3.2.4. *Summary of the mode decomposition including $l = 0, 1$ modes.* Here, we summarize the mode decomposition by harmonic functions Y_{lm} ($l \geq 2$), $k_{(\hat{\Delta}+2)m}$, and $k_{(\hat{\Delta})}$. We decompose the components $\{h_{AB}, h_{Ap}, h_{pq}\}$ of the metric perturbation h_{ab} as Eqs. (3.7)–(3.9) with

$$S = \begin{cases} Y_{lm} & \text{for } l \geq 2; \\ k_{(\hat{\Delta}+2)m} & \text{for } l = 1; \\ k_{(\hat{\Delta})} & \text{for } l = 0. \end{cases} \quad (3.93)$$

This decomposition is invertible for any l, m modes including $l = 0, 1$ if the conditions (3.72)–(3.74), i.e.,

$$k_{(\hat{\Delta})} \in \mathcal{K}_{(\hat{\Delta})}, \quad k_{(\hat{\Delta}+2)} \in \mathcal{K}_{(\hat{\Delta}+2)}, \quad k_{(\hat{\Delta}+2)} = k_{(\hat{\Delta}+2)m} = \Theta_{1m}(\theta) e^{im\phi}, \quad (3.94)$$

$$\left(\hat{D}_p k_{(\hat{\Delta})} \right) \left(\hat{D}^p k_{(\hat{\Delta})} \right) \neq 0, \quad (3.95)$$

$$K_{(m)pq} K_{(m)}^{pq} = \left(\hat{D}_p \hat{D}_q k_{(\hat{\Delta}+2)m} \right) \left(\hat{D}^p \hat{D}^q k_{(\hat{\Delta}+2)m} \right) - 2 \left(k_{(\hat{\Delta}+2)m} \right)^2 \neq 0. \quad (3.96)$$

are satisfied. As the inverse relation of Eqs. (3.7)–(3.9), the mode coefficients of these decomposition are given in the functional form of the metric components h_{AB} , h_{Ap} , and h_{pq} as Eqs. (3.41), (3.42), (3.47), (3.50), (3.51), (3.53), (3.54), (3.70), (3.71), (3.79), (3.81), (3.83), (3.84), (3.87), (3.88), and (3.90)–(3.92). From Eqs. (3.7)–(3.9), the components $\{h_{AB}, h_{Ap}, h_{pq}\}$ vanish if all mode coefficients $\{\tilde{h}_{AB}, \tilde{h}_{(e1)A}, \tilde{h}_{(o1)A}, \tilde{h}_{(e0)}, \tilde{h}_{(e2)}, \tilde{h}_{(o2)}\}$ vanish. On the contrary, from the obtained functional forms, all mode coefficients $\{\tilde{h}_{AB}, \tilde{h}_{(e1)A}, \tilde{h}_{(o1)A}, \tilde{h}_{(e0)}, \tilde{h}_{(e2)}, \tilde{h}_{(o2)}\}$ vanish if the components $\{h_{AB}, h_{Ap}, h_{pq}\}$ vanish. This indicates the linear independence of the set of the harmonic functions (3.12). Therefore, the conditions (3.94)–(3.96) guarantee the linear independence of the set of these harmonic functions (3.12).

We also note that the Green functions $\hat{\Delta}^{-1}$ and $[\hat{\Delta} + 2]^{-1}$ which used above do not directly operate to the functions $k_{(\hat{\Delta})}$, nor $k_{(\hat{\Delta}+2)m}$. There, the domain of these Green function $\hat{\Delta}^{-1}$ and $[\hat{\Delta} + 2]^{-1}$ are L^2 -space spanned by $\{Y_{lm}|l \neq 0\}$ and $\{Y_{lm}|l \neq 1\}$, respectively. The explicit form of these Green functions are given in Ref. [62, 63].

3.3. Explicit form of the mode functions

Here, we consider the explicit expression of the mode function $k_{\hat{\Delta}}$ and $k_{(\hat{\Delta}+2)}$ which satisfy the conditions (3.94)–(3.96). In Appendix A, we explicitly see that the choice $S = Y_{lm}$ for $l \geq 0$ does not satisfy these conditions and what is happen in this choice. As the result of this appendix, in the choice $S = Y_{lm}$, any vector and tensor harmonics does not have their values for $l = 0$ mode. On the other hand, for $l = 1$ modes, the vector harmonics have their vector value and the trace parts of the second-rank tensor of each modes have their tensor values, while all traceless even and odd mode harmonics identically vanish. Therefore, in the choice $S = Y_{lm}$, the set of harmonics (3.12) does not play the role of basis of tangent space on S^2 for $l = 0, 1$ mode. This situation already appeared in terms of the Green function $\hat{\Delta}^{-1}$ and $(\hat{\Delta} + 2)^{-1}$ in the inverse relations in Sec. 3.1. For this reason, we seek an alternative choice of S which satisfy the conditions (3.94)–(3.96).

3.3.1. Explicit form of $k_{(\hat{\Delta})}$. Here, we treat the modes which belong to the kernel of the derivative operator $\hat{\Delta}$, i.e.,

$$\hat{\Delta}k_{(\hat{\Delta})} = \frac{1}{\sqrt{\gamma}}\partial_p \left(\sqrt{\gamma}\gamma^{pq}\partial_q k_{(\hat{\Delta})} \right) = 0. \quad (3.97)$$

We look for the function which satisfies the conditions (3.94) and (3.95). We emphasize that we do not impose the regularity on the function $k_{(\hat{\Delta})}$ on S^2 itself in this selection of $k_{(\hat{\Delta})}$. Since the regularity is a kind of boundary conditions for perturbations, this regularity may be imposed on the solutions when we solve the Einstein equations.

Our guiding principle to look for the solution to Eq. (3.97) with a simple modification from the conventional spherical harmonic functions. Although the conditions (3.94) and (3.95) do not restrict the ϕ -dependence for $k_{(\hat{\Delta})}$, we look for the solution to Eq. (3.97) which is independent of ϕ as the original Y_{00} in the conventional spherical harmonics is so. Then, in terms of the coordinate system where γ_{ab} is given by Eq. (3.4), Eq. (3.97) yields

$$\frac{d^2}{dy^2}k_{(\hat{\Delta})} = 0, \quad (3.98)$$

where we introduced an independent variable y by

$$y = \ln \left(\frac{1 - \cos \theta}{1 + \cos \theta} \right)^{1/2}. \quad (3.99)$$

As the solution to Eq. (3.98), we choose

$$k_{(\hat{\Delta})} = 1 + \delta y = 1 + \delta \ln \left(\frac{1 - \cos \theta}{1 + \cos \theta} \right)^{1/2}, \quad \delta \in \mathbb{R}. \quad (3.100)$$

If $\delta \neq 0$, we see that

$$\hat{\Delta}_p k_{(\hat{\Delta})} (dx^p)_a = \delta (dy)_a = \frac{\delta}{\sin \theta} (d\theta)_a \neq 0, \quad (3.101)$$

and

$$(\hat{D}_p k_{(\Delta)}) (\hat{D}^p k_{(\Delta)}) = \frac{\delta^2}{\sin^2 \theta} \neq 0. \quad (3.102)$$

Thus, $\hat{D}_p k_{(\hat{\Delta})}$ given by Eq. (3.101) and $\varepsilon_{pq} \hat{D}^q k_{(\hat{\Delta})}$ span the vector space though their norm is singular at $\theta = 0, \pi$. The solution (3.100) to Eq. (3.98) also yields

$$\left(\hat{D}_p \hat{D}_q - \frac{1}{2} \gamma_{pq} \hat{\Delta} \right) k_{(\hat{\Delta})} = \hat{D}_p \hat{D}_q k_{(\hat{\Delta})} = \delta \frac{\cos \theta}{\sin^2 \theta} (-\theta_p \theta_q + \phi_p \phi_q) \neq 0, \quad (3.103)$$

$$\varepsilon_{r(p} \hat{D}_{q)} \hat{D}^r k_{(\hat{\Delta})} = -2\delta \frac{\cos \theta}{\sin^2 \theta} \theta_{(p} \phi_{q)} \neq 0. \quad (3.104)$$

Together with the trace part

$$\frac{1}{2} \gamma_{pq} k_{(\hat{\Delta})} = \frac{1}{2} \left(1 + \delta \ln \left(\frac{1 - \cos \theta}{1 + \cos \theta} \right)^{1/2} \right) (\theta_p \theta_q + \phi_p \phi_q), \quad (3.105)$$

the tensor (3.103) and (3.104) span the basis of the space of the second-rank tensor field though these are singular at $\theta = 0, \pi$.

3.3.2. Explicit form of $k_{(\hat{\Delta}+2)}$. Here, we consider the kernel mode $k_{(\hat{\Delta}+2)}$ for the operator $\hat{\Delta} + 2$. The condition (3.94) for $k_{(\hat{\Delta}+2)}$ is given by

$$(\hat{\Delta} + 2) k_{(\hat{\Delta}+2)} = \frac{1}{\sqrt{\gamma}} \partial_p \left(\sqrt{\gamma} \gamma^{pq} \partial_q k_{(\hat{\Delta}+2)} \right) + 2k_{(\hat{\Delta}+2)} = 0. \quad (3.106)$$

We look for the function which satisfies the conditions (3.94) and (3.96). We emphasize that we do not impose the regularity on the function $k_{(\hat{\Delta}+2)}$ on S^2 itself as in the case of $k_{(\hat{\Delta})}$. To obtain the solution to Eq. (3.106) which satisfies the conditions (3.94) and (3.96), we first consider the ϕ -dependence from the condition (3.94). Then, Eq. (3.106) is given by

$$\sin \theta \partial_\theta (\sin \theta \partial_\theta \Theta_{1m}(\theta)) - m^2 \Theta_{1m}(\theta) + 2 \sin^2 \theta \Theta_{1m}(\theta) = 0. \quad (3.107)$$

To solve Eq. (3.107), we introduce the independent variable

$$z = \cos \theta, \quad dz = -\sin \theta d\theta. \quad (3.108)$$

In terms of the independent variable z , we obtain

$$\sin \theta \frac{d}{d\theta} = -(1 - z^2) \frac{d}{dz}. \quad (3.109)$$

Then, Eq. (3.107) is given by

$$(1 - z^2) \frac{d^2}{dz^2} \Theta_{1m}(\theta) - 2z \frac{d}{dz} \Theta_{1m}(\theta) + \left(1(1+1) - \frac{m^2}{1-z^2} \right) \Theta_{1m}(\theta) = 0. \quad (3.110)$$

Suppose that we have obtained the solution to Eq. (3.110) as

$$k_{(\hat{\Delta}+2)m} = \Theta_{1m}(\theta) e^{im\phi}. \quad (3.111)$$

Here, we introduce the ladder operator \hat{L}_\pm [64] as

$$\hat{L}_\pm := -ie^{\pm i\phi} (\pm i \partial_\theta - \cot \theta \partial_\phi) \quad (3.112)$$

and examine the function defined by

$$\begin{aligned}\hat{L}_+k_{(\hat{\Delta}+2,m)} &= -ie^{+i\phi} (+i\partial_\theta - \cot\theta\partial_\phi)\Theta_{1m}(\theta)e^{im\phi} \\ &= (\partial_\theta - m\cot\theta)\Theta_{1m}(\theta)e^{i(m+1)\phi}.\end{aligned}\quad (3.113)$$

Evidently, the function given by Eq. (3.113) is the eigenfunction of the operator $-i\partial_\phi$ with the eigenvalue $m+1$:

$$-i\partial_\phi\hat{L}_+k_{(\hat{\Delta}+2,m)} = (m+1)\hat{L}_+k_{(\hat{\Delta}+2,m)}.\quad (3.114)$$

Now, we consider the variable Φ_+ defined by

$$\begin{aligned}\Phi_+ &:= (\partial_\theta - m\cot\theta)\Theta_{1m} \\ &= -(1-z^2)^{-1/2}\left[(1-z^2)\frac{d}{dz}\Theta_{1,m} + mz\Theta_{1m}\right],\end{aligned}\quad (3.115)$$

and straightforward calculations using Eq. (3.110) yields

$$(1-z^2)\frac{d^2}{dz^2}\Phi_+ - 2z\frac{d}{dz}\Phi_+ + \left(1(1+1) - \frac{(m+1)^2}{1-z^2}\right)\Phi_+ = 0.\quad (3.116)$$

This indicates

$$\Phi_+ = \Theta_{1,m+1}(\theta).\quad (3.117)$$

Therefore, we conclude that

$$\hat{L}_+k_{(\hat{\Delta}+2,m)} = k_{(\hat{\Delta}+2,m+1)}.\quad (3.118)$$

On the other hand, we consider the operator \hat{L}_- defined by

$$\begin{aligned}\hat{L}_-k_{(\hat{\Delta}+2,m)} &= -ie^{-i\phi} (-i\partial_\theta - \cot\theta\partial_\phi)\Theta_{1,m}e^{im\phi} \\ &= (-\partial_\theta - m\cot\theta)\Theta_{1,m}e^{i(m-1)\phi}\end{aligned}\quad (3.119)$$

Evidently, the function given by Eq. (3.119) is an eigenfunction of the operator $-i\partial_\phi$ with the eigenvalue $m-1$:

$$-i\partial_\phi\hat{L}_-k_{(\hat{\Delta}+2,m)} = (m-1)\hat{L}_-k_{(\hat{\Delta}+2,m)}.\quad (3.120)$$

Now, we consider

$$\Phi_- := (-\partial_\theta - m\cot\theta)\Theta_{1,m}\quad (3.121)$$

and straightforward calculations using Eq. (3.110) yields

$$(1-z^2)\frac{d^2}{dz^2}\Phi_- - 2z\frac{d}{dz}\Phi_- + \left[1(1+1) - \frac{(m-1)^2}{1-z^2}\right]\Phi_- = 0.\quad (3.122)$$

This indicates

$$\Phi_- = \Theta_{1,m-1}(\theta).\quad (3.123)$$

Therefore, we conclude that

$$\hat{L}_-k_{(\hat{\Delta}+2)m} = k_{(\hat{\Delta}+2)m-1}.\quad (3.124)$$

From the above operator \hat{L}_\pm and

$$\hat{L}_\pm k_{(\hat{\Delta}+2)m} = k_{(\hat{\Delta}+2)m\pm 1}, \quad (3.125)$$

we may concentrate only to solve $m = 0$ case. Corresponding $m = \pm 1$ modes with $l = 1$ can be derived from Eq. (3.125). Since $k_{(\hat{\Delta}+2)m=0} = \Theta_{10}(\theta)$, the equation for $\Theta_{10}(\theta)$ is given by

$$(1-z^2) \frac{d^2}{dz^2} \Theta_{10}(\theta) - 2z \frac{d}{dz} \Theta_{10}(\theta) + 1(1+1) \Theta_{10}(\theta) = 0. \quad (3.126)$$

Here, we note that $\Theta_{10} = z \propto Y_{10}$ should be a solution to Eq. (3.126). To obtain the other independent solution, we consider the solution in the form $\Theta_{10} = \Psi(z)z$. Substituting this into Eq. (3.126) we can solve Eq. (3.126) as

$$\Theta_{10} = z + \delta \left(\frac{1}{2} z \ln \frac{1+z}{1-z} - 1 \right), \quad (3.127)$$

where we choose one of constant of integration as 1 and δ is another integration constant. Then, we obtain

$$k_{(\hat{\Delta}+2)m=0} = z + \delta \left(\frac{1}{2} z \ln \frac{1+z}{1-z} - 1 \right) = P_1(z) + \delta Q_1(z), \quad (3.128)$$

where $P_1(z)$ is the Legendre polynomial and $Q_1(z)$ is the first order and the second kind Legendre function.

Since we have the explicit form (3.128) of $k_{(\hat{\Delta}+2)m=0}$ as

$$k_{(\hat{\Delta}+2)m} = \Theta_{1m}(\theta) e^{im\phi}, \quad (3.129)$$

we can derive the $m = \pm 1$ modes by applying the ladder operators \hat{L}_\pm 3.112 as

$$k_{(\hat{\Delta}+2)m=\pm 1} = \hat{L}_\pm k_{(\hat{\Delta}+2)m=0} = \left[\sqrt{1-z^2} + \delta \left(\frac{1}{2} \sqrt{1-z^2} \ln \frac{1+z}{1-z} + \frac{z}{\sqrt{1-z^2}} \right) \right] e^{\pm i\phi}. \quad (3.130)$$

Equations (3.128) and (3.130) are summarized as

$$k_{(\hat{\Delta}+2,m=0)} = \cos \theta + \delta \left(\frac{1}{2} \cos \theta \ln \frac{1+\cos \theta}{1-\cos \theta} - 1 \right), \quad \delta \in \mathbb{R}, \quad (3.131)$$

$$k_{(\hat{\Delta}+2,m=\pm 1)} = \left[\sin \theta + \delta \left(+\frac{1}{2} \sin \theta \ln \frac{1+\cos \theta}{1-\cos \theta} + \cot \theta \right) \right] e^{\pm i\phi}. \quad (3.132)$$

Here, we check the non-vanishing properties of $\hat{D}_p k_{(\hat{\Delta}+2)}$ and $\hat{D}_p \hat{D}_q k_{(\hat{\Delta}+2)}$. For $m = 0$ modes, the vector $\hat{D}_p k_{(\hat{\Delta}+2,m=0)}$ is given by

$$\hat{D}_p k_{(\hat{\Delta}+2,m=0)} = - \left[1 + \frac{1}{2} \delta \left(\ln \frac{1+\cos \theta}{1-\cos \theta} + \frac{2 \cos \theta}{\sin^2 \theta} \right) \right] \sin \theta \theta_p. \quad (3.133)$$

Then $\hat{D}_p k_{(\hat{\Delta}+2,m=0)}$ and $\varepsilon_{pq} \hat{D}^q k_{(\hat{\Delta}+2),m=0}$ span the basis of the tangent space on S^2 .

Next, we consider the tensor $\hat{D}_q \hat{D}_p k_{(\hat{\Delta}+2, m=0)}$ as

$$\begin{aligned} & \hat{D}_q \hat{D}_p k_{(\hat{\Delta}+2, m=0)} \\ = & - \left[\cos \theta + \frac{1}{2} \delta \left(+ \cos \theta \ln \frac{1 + \cos \theta}{1 - \cos \theta} - 4 - 2 \cot^2 \theta \right) \right] \theta_p \theta_q \\ & - \left[\cos \theta + \frac{1}{2} \delta \cos \theta \left(+ \ln \frac{1 + \cos \theta}{1 - \cos \theta} + \frac{2 \cos \theta}{\sin^2 \theta} \right) \right] \phi_p \phi_q. \end{aligned} \quad (3.134)$$

This does not proportional to γ_{ab} . Therefore, we should have nonvanishing $K_{(m)pq}$ and $J_{(m)pq}$. To confirm this, we evaluate the condition (3.74) as

$$\left(\hat{D}_p \hat{D}_q k_{(\hat{\Delta}+2)} \right) \left(\hat{D}^p \hat{D}^q k_{(\hat{\Delta}+2)} \right) - 2 \left(k_{(\hat{\Delta}+2)} \right)^2 = \frac{2\delta}{\sin^4 \theta}. \quad (3.135)$$

This indicates that we have nonvanishing $K_{(m)pq}$ and $J_{(m)pq}$ if $\delta \neq 0$. However, we should note that these tensor singular at $\theta = 0, \pi$.

For $m = \pm 1$ modes, the vector $\hat{D}_p k_{(\hat{\Delta}+2, m=\pm 1)}$ is given by

$$\begin{aligned} \hat{D}_p k_{(\hat{\Delta}+2, m=\pm 1)} & = \left[\cos \theta + \delta \left(+ \frac{1}{2} \cos \theta \ln \frac{1 + \cos \theta}{1 - \cos \theta} - 1 - \frac{1}{\sin^2 \theta} \right) \right] e^{\pm i\phi} \theta_p \\ & + (\pm i) \left[1 + \delta \left(+ \frac{1}{2} \sin \theta \ln \frac{1 + \cos \theta}{1 - \cos \theta} + \frac{\cos \theta}{\sin^2 \theta} \right) \right] e^{\pm i\phi} \phi_p. \end{aligned} \quad (3.136)$$

Finally, we evaluate the condition (3.74) as

$$\begin{aligned} \hat{D}_q \hat{D}_p k_{(\hat{\Delta}+2, m=\pm 1)} & = \left[-\sin \theta + \delta \left(-\frac{1}{2} \sin \theta \ln \frac{1 + \cos \theta}{1 - \cos \theta} - \frac{\cos \theta}{\sin \theta} + \frac{2 \cos \theta}{\sin^3 \theta} \right) \right] e^{\pm i\phi} \theta_p \theta_q \\ & + \left[-\sin \theta + \delta \left(-\frac{1}{2} \sin \theta \ln \frac{1 + \cos \theta}{1 - \cos \theta} - \frac{\cos \theta}{\sin \theta} - \frac{2 \cos \theta}{\sin^3 \theta} \right) \right] e^{\pm i\phi} \phi_p \phi_q \\ & \mp \frac{4i\delta}{\sin^3 \theta} e^{\pm i\phi} \theta_p \phi_q. \end{aligned} \quad (3.137)$$

This does not proportional to γ_{pq} . Therefore, we should have nonvanishing $K_{(m)pq}$ and $J_{(m)pq}$. This is confirmed by the check of the condition (3.74) as

$$\left(\hat{D}_p \hat{D}_q k_{(\hat{\Delta}+2)} \right) \left(\hat{D}^p \hat{D}^q k_{(\hat{\Delta}+2)} \right) - 2 \left(k_{(\hat{\Delta}+2)} \right)^2 = -\frac{8\delta^2}{\sin^4 \theta} e^{\pm 2i\phi}. \quad (3.138)$$

Then, we have seen that if $\delta \neq 0$, the condition (3.74) is satisfied, though this norm is singular at $\theta = 0, \pi$. We also note that $K_{(m)pq}$ is orthogonal to $J_{(m)pq}$ as shown in Eqs. (3.66)–(3.68). Therefore, γ_{pq} , $K_{(m)pq}$, and $J_{(m)pq}$ span the basis of the second-rank tensor field on S^2 .

3.4. Proposal of the treatment of $l = 0, 1$ -mode perturbations

As shown in above, it is shown that the harmonic decomposition (3.7)–(3.9) have the one-to-one correspondence between the original metric perturbations $\{h_{AB}, h_{Ap}, h_{pq}\}$ and the mode coefficients $\{\tilde{h}_{AB}, \tilde{h}_{(e1)A}, \tilde{h}_{(o1)A}, \tilde{h}_{(e0)}, \tilde{h}_{(e2)}, \tilde{h}_{(o2)}\}$ for any modes $l \geq 0$ through the employment of the scalar

harmonic functions

$$S_\delta = \begin{cases} Y_{lm} & \text{for } l \geq 2; \\ k_{(\hat{\Delta}+2)m} & \text{for } l = 1; \\ k_{(\hat{\Delta})} & \text{for } l = 0, \end{cases} \quad (3.139)$$

where $k_{(\hat{\Delta})}$ is given by Eq. (3.100), i.e.,

$$k_{(\hat{\Delta})} = 1 + \delta \ln \left(\frac{1 - \cos \theta}{1 + \cos \theta} \right)^{1/2}, \quad \delta \in \mathbb{R} \quad (3.140)$$

and $k_{(\hat{\Delta}+2)m}$ are given by Eqs. (3.128) and (3.130), i.e.,

$$k_{(\hat{\Delta}+2)m=0} = \cos \theta + \delta \left(\frac{1}{2} \cos \theta \ln \frac{1 + \cos \theta}{1 - \cos \theta} - 1 \right), \quad (3.141)$$

$$k_{(\hat{\Delta}+2)m=\pm 1} = \left[\sin \theta + \delta \left(\frac{1}{2} \sin \theta \ln \frac{1 + \cos \theta}{1 - \cos \theta} + \cot \theta \right) \right] e^{\pm i\phi}. \quad (3.142)$$

These mode functions $k_{(\hat{\Delta}+2)m}$ and $k_{(\hat{\Delta})}$ are parametrized by the single parameter δ . This choice satisfies the conditions (3.94)–(3.96) but singular at $\theta = 0, \pi$ if $\delta \neq 0$. When $\delta = 0$, we have $k_{(\hat{\Delta})} \propto Y_{00}$ and $k_{(\hat{\Delta}+2)m} \propto Y_{1m}$. In this decomposition, for each mode of any $l \geq 0$, the set of harmonic functions (3.12) are a linear-independent set in the both senses of the second-rank tensor field and the function on S^2 .

Using the above harmonics functions S_δ in Eq. (3.139), we propose the following strategy:

Proposal 3.1. *We decompose the metric perturbation h_{ab} on the background spacetime with the metric (3.1)–(3.4) through Eqs. (3.7)–(3.9) with the harmonic function S_δ given by Eq. (3.139). Then, Eqs. (3.7)–(3.9) become invertible including $l = 0, 1$ modes. After deriving the field equations such as linearized Einstein equations by using the harmonic functions S_δ , we choose $\delta = 0$ when we solve these field equations as regularities of solutions.*

Since the set of the mode functions (3.12) with $S = S_\delta$ have the linear-independence including $l = 0, 1$ modes, we can construct gauge-invariant variables and evaluate the field equations through the mode-by-mode analyses including $l = 0, 1$ modes through the choice of these mode functions.

4. Construction of gauge-invariant variables

In this section, we construct the gauge invariant variables for perturbations on spherically symmetric background with the metric (3.1) through Proposal 3.1. To construct the gauge-invariant variables, we first discuss the gauge-transformation rule for the metric perturbation h_{ab} . In the derivation of the gauge-transformation rules for the mode coefficient in the decomposition (3.7)–(3.9) with the harmonic function $S = S_\delta$ given by Eq. (3.139). In this section, we use the relations of the covariant derivatives associated with the metrics g_{ab} , y_{ab} , and γ_{ab} , which are summarized in Appendix B. In Sec. 4.1, we derive the gauge-transformation rules for the mode coefficients of the metric perturbation in the decomposition (3.7)–(3.9) with the harmonic function $S = S_\delta$. In Sec. 4.2, we explicitly construct gauge-invariant variables for the metric perturbations through the mode-by-mode analyses. In Sec. 4.2.3, we summarize gauge-invariant and gauge-variant variables in the four-dimensional form.

4.1. Gauge-transformation rules

Here, we consider the gauge-transformation rules for the linear-order metric perturbation h_{ab} following to Proposal 3.1. The gauge-transformation rule for linear-order metric perturbation is given by

$$\mathcal{L}h_{ab} - \mathcal{X}h_{ab} = \mathcal{L}\xi g_{ab} = 2\nabla_{(a}\xi_{b)}. \quad (4.1)$$

We rewrite this gauge-transformation rule in terms of 2+2 formulation. To do this, the generator of gauge-transformation rules is decomposed as

$$\xi_a = \xi_A(dx^A)_a + \xi_p(dx^p)_a. \quad (4.2)$$

Through the component-representations (3.6) and (4.2), the gauge-transformation rules (4.1) are given by

$$\mathcal{L}h_{AB} - \mathcal{X}h_{AB} = \nabla_A\xi_B + \nabla_B\xi_A = \bar{D}_A\xi_B + \bar{D}_B\xi_A, \quad (4.3)$$

$$\mathcal{L}h_{Ap} - \mathcal{X}h_{Ap} = \nabla_A\xi_p + \nabla_p\xi_A = \bar{D}_A\xi_p + \hat{D}_p\xi_A - \frac{2}{r}\bar{D}_A r\xi_p, \quad (4.4)$$

$$\mathcal{L}h_{pq} - \mathcal{X}h_{pq} = \nabla_p\xi_q + \nabla_q\xi_p = \hat{D}_p\xi_q + \hat{D}_q\xi_p + 2r\bar{D}^A r\gamma_{pq}\xi_A. \quad (4.5)$$

Furthermore, through the mode-decomposition (3.7)–(3.9) and

$$\xi_A =: \sum_{l,m} \zeta_A S_\delta, \quad (4.6)$$

$$\xi_p =: r \sum_{l,m} [\zeta_{(e)} \hat{D}_p S_\delta + \zeta_{(o)} \epsilon_{pq} \hat{D}^q S_\delta] \quad (4.7)$$

with the harmonic function S_δ , we can carry out the mode-by-mode analyses, since the set of the harmonic functions (3.12) has the linear-independence due to the choice $S = S_\delta$. From Eq. (4.3), we obtain

$$\mathcal{L}\tilde{h}_{AB} - \mathcal{X}\tilde{h}_{AB} = 2\bar{D}_{(A}\zeta_{B)}. \quad (4.8)$$

From Eq. (4.4), we obtain

$$\mathcal{L}\tilde{h}_{(e1)A} - \mathcal{X}\tilde{h}_{(e1)A} = \frac{1}{r}\zeta_A + \bar{D}_A\zeta_{(e)} - \frac{1}{r}\bar{D}_A r\zeta_{(e)}, \quad (4.9)$$

$$\mathcal{L}\tilde{h}_{(o1)A} - \mathcal{X}\tilde{h}_{(o1)A} = \bar{D}_A\zeta_{(o)} - \frac{1}{r}\bar{D}_A r\zeta_{(o)}. \quad (4.10)$$

Finally, the gauge-transformation rules (4.5) yield

$$\mathcal{L}\tilde{h}_{(e0)} - \mathcal{X}\tilde{h}_{(e0)} = \frac{4}{r} \left(-\frac{1}{2}l(l+1)\zeta_{(e)} + \bar{D}^A r\zeta_A \right), \quad (4.11)$$

$$\mathcal{L}\tilde{h}_{(e2)} - \mathcal{X}\tilde{h}_{(e2)} = \frac{2}{r}\zeta_{(e)}, \quad (4.12)$$

$$\mathcal{L}\tilde{h}_{(o2)} - \mathcal{X}\tilde{h}_{(o2)} = -\frac{1}{r}\zeta_{(o)}. \quad (4.13)$$

We note that these gauge-transformation rules (4.8)–(4.13) are not only that for $l \geq 2$ modes but also $l = 0, 1$ modes.

4.2. Gauge-invariant and gauge-variant variables

Inspecting gauge-transformation rules (4.8)–(4.13), we can define gauge-invariant variables.

4.2.1. *Odd modes.* From gauge-transformation rules (4.10) and (4.13), we easily find that the following combination is gauge-invariant:

$$\tilde{h}_{(o1)A} - \bar{D}_A(-r\tilde{h}_{(o2)}) + \frac{1}{r}\bar{D}_{Ar}(-r\tilde{h}_{(o2)}) = \tilde{h}_{(o1)A} + r\bar{D}_A\tilde{h}_{(o2)} =: \tilde{F}_A. \quad (4.14)$$

We also note that the gauge-transformation rule (4.13) implies that

$$-r^2\vartheta\tilde{h}_{(o2)} + r^2\mathcal{X}\tilde{h}_{(o2)} = r\zeta_{(o)}. \quad (4.15)$$

4.2.2. *Even modes.* Now, we note that the gauge-transformation rule (4.12) implies that

$$\frac{r^2}{2}\vartheta\tilde{h}_{(e2)} - \frac{r^2}{2}\mathcal{X}\tilde{h}_{(e2)} = r\zeta_{(e)}. \quad (4.16)$$

Inspecting gauge-transformation rules (4.9) and (4.12), we define the variable \tilde{Y}_A as

$$\begin{aligned} \tilde{Y}_A &:= r\tilde{h}_{(e1)A} - r\bar{D}_A\left(\frac{r}{2}\tilde{h}_{(e2)}\right) + \bar{D}_{Ar}\left(\frac{r}{2}\tilde{h}_{(e2)}\right) \\ &= r\tilde{h}_{(e1)A} - \frac{r^2}{2}\bar{D}_A\tilde{h}_{(e2)}. \end{aligned} \quad (4.17)$$

We easily check that the gauge-transformation rules for the variable \tilde{Y}_A is given by

$$\vartheta\tilde{Y}_A - \mathcal{X}\tilde{Y}_A = \zeta_A. \quad (4.18)$$

From the gauge-transformation rules (4.16) and (4.18), we easily define the gauge-invariant variables as follows: First, from the gauge-transformation rules (4.8) and (4.18), the following combination is gauge-invariant:

$$\tilde{F}_{AB} := \tilde{h}_{AB} - 2\bar{D}_{(A}\tilde{Y}_{B)}. \quad (4.19)$$

Second, from the gauge-transformation rules (4.11), (4.16), and (4.18), we can define the gauge-invariant variables \tilde{F} as follows:

$$\begin{aligned} \tilde{F} &:= \tilde{h}_{(e0)} - \frac{4}{r}\tilde{Y}_A\bar{D}^Ar + \frac{2}{r}\frac{r}{2}\tilde{h}_{(e2)}l(l+1) \\ &= \tilde{h}_{(e0)} - \frac{4}{r}\tilde{Y}_A\bar{D}^Ar + \tilde{h}_{(e2)}l(l+1). \end{aligned} \quad (4.20)$$

4.2.3. *Summary of gauge-invariant and gauge-variant variables.* In summary, we have defined gauge-invariant variables as follows:

$$\tilde{F}_A := \tilde{h}_{(o1)A} + r\bar{D}_A\tilde{h}_{(o2)}, \quad (4.21)$$

$$\tilde{F} := \tilde{h}_{(e0)} - \frac{4}{r}\tilde{Y}_A\bar{D}^Ar + \tilde{h}_{(e2)}l(l+1), \quad (4.22)$$

$$\tilde{F}_{AB} := \tilde{h}_{AB} - 2\bar{D}_{(A}\tilde{Y}_{B)}, \quad (4.23)$$

where we defined the variable \tilde{X}_A by

$$\tilde{Y}_A := r\tilde{h}_{(e1)A} - \frac{r^2}{2}\bar{D}_A\tilde{h}_{(e2)} \quad (4.24)$$

The gauge-transformation rules for the variable \tilde{Y}_A is given by

$$\vartheta\tilde{Y}_A - \mathcal{X}\tilde{Y}_A = \zeta_A. \quad (4.25)$$

We also note that the gauge-transformation rules (4.15) and (4.16), i.e.,

$$-r^2 \vartheta \tilde{h}_{(o2)} + r^2 \mathcal{X} \tilde{h}_{(o2)} = r \zeta_{(o)}. \quad (4.26)$$

$$\frac{r^2}{2} \vartheta \tilde{h}_{(e2)} - \frac{r^2}{2} \mathcal{X} \tilde{h}_{(e2)} = r \zeta_{(e)}. \quad (4.27)$$

Therefore, it is reasonable to define the variables $\tilde{Y}_{(o)}$ and $\tilde{Y}_{(e)}$ as follows:

$$\tilde{Y}_{(o1)} := -r^2 \tilde{h}_{(o2)}, \quad (4.28)$$

$$\tilde{Y}_{(e1)} := \frac{r^2}{2} \tilde{h}_{(e2)} \quad (4.29)$$

so that their gauge-transformation rules are given by

$$\vartheta \tilde{Y}_{(o1)} - \mathcal{X} \tilde{Y}_{(o1)} = r \zeta_{(o)}, \quad (4.30)$$

$$\vartheta \tilde{Y}_{(e1)} - \mathcal{X} \tilde{Y}_{(e1)} = r \zeta_{(e)}. \quad (4.31)$$

Furthermore, we define the variable

$$Y_a := \sum_{l,m} \tilde{Y}_A S_\delta (dx^A)_a + \sum_{l,m} (\tilde{Y}_{(e1)} \hat{D}_p S_\delta + \tilde{Y}_{(o1)} \varepsilon_{pq} \hat{D}^q S_\delta) (dx^p)_a. \quad (4.32)$$

The gauge transformation rule for the variable Y_a is given by

$$\begin{aligned} \vartheta Y_a - \mathcal{X} Y_a &= \sum_{l,m} (\vartheta \tilde{Y}_A - \mathcal{X} \tilde{Y}_A) S_\delta (dx^A)_a \\ &\quad + \sum_{l,m} ((\vartheta \tilde{Y}_{(e)} - \mathcal{X} \tilde{Y}_{(e)}) \hat{D}_p S_\delta + (\vartheta \tilde{Y}_{(o)} - \mathcal{X} \tilde{Y}_{(o)}) \varepsilon_{pq} \hat{D}^q S_\delta) (dx^p)_a \\ &= \sum_{l,m} \zeta_A S_\delta (dx^A)_a + \sum_{l,m} (r \zeta_{(e)} \hat{D}_p S_\delta + r \zeta_{(o)} \varepsilon_{pq} \hat{D}^q S_\delta) (dx^p)_a \\ &= \zeta_A (dx^A)_a + \zeta_p (dx^p)_a \\ &= \xi_a, \end{aligned} \quad (4.33)$$

where we used Eqs. (4.6) and (4.7).

In terms of the gauge-invariant variables $\{\tilde{F}_A, \tilde{F}, \tilde{F}_{AB}\}$ defined by Eqs. (4.21)–(4.23) and gauge-variant variables Y_a defined by (4.32), we can express the original components $\{h_{AB}, h_{Ap}, h_{pq}\}$. First, we consider the component h_{AB} :

$$\begin{aligned} h_{AB} &= \sum_{l,m} (\tilde{h}_{AB}) S_\delta = \sum_{l,m} (\tilde{F}_{AB} + 2\bar{D}_{(A} \tilde{Y}_{B)}) S_\delta, \\ &= F_{AB} + 2\bar{D}_{(A} Y_{B)}, \end{aligned} \quad (4.34)$$

where we defined the gauge-invariant variable F_{AB} by

$$F_{AB} := \sum_{l,m} \tilde{F}_{AB} S_\delta. \quad (4.35)$$

Next, we consider the component h_{Ap} :

$$\begin{aligned} h_{Ap} &= r \sum_{l,m} [(\tilde{h}_{(e1)A}) \hat{D}_p S_\delta + (\tilde{h}_{(o1)A}) \varepsilon_{pq} \hat{D}^q S_\delta] \\ &= r F_{Ap} + \hat{D}_p Y_A + \bar{D}_A Y_p - \frac{2}{r} \bar{D}_A r Y_p, \end{aligned} \quad (4.36)$$

where we defined

$$F_{Ap} := \sum_{l,m} [\tilde{F}_A \varepsilon_{pq} \hat{D}^q S_\delta], \quad \hat{D}^p F_{Ap} = 0. \quad (4.37)$$

Finally, we consider the component h_{pq} :

$$\begin{aligned} h_{pq} &= r^2 \sum_{l,m} \left[\tilde{h}_{(e0)} \frac{1}{2} \gamma_{pq} S_\delta + (\tilde{h}_{(e2)}) \left(\hat{D}_p \hat{D}_q - \frac{1}{2} \gamma_{pq} \hat{D}^r \hat{D}_r \right) S_\delta \right. \\ &\quad \left. + 2 (\tilde{h}_{(o2)}) \varepsilon_{r(p} \hat{D}_{q)} \hat{D}^r S_\delta \right] \\ &= \frac{1}{2} \gamma_{pq} r^2 F + 2r \gamma_{pq} \bar{D}^A r Y_A + \hat{D}_p Y_q + \hat{D}_q Y_p, \end{aligned} \quad (4.38)$$

where we have defined

$$F := \sum_{l,m} \tilde{F} S_\delta. \quad (4.39)$$

Then, we have obtained

$$h_{AB} = F_{AB} + 2\bar{D}_{(A} Y_{B)}, \quad (4.40)$$

$$h_{Ap} = r F_{Ap} + \hat{D}_p Y_A + \bar{D}_A Y_p - \frac{2}{r} \bar{D}_A r Y_p, \quad (4.41)$$

$$h_{pq} = +\frac{1}{2} \gamma_{pq} r^2 F + 2r \gamma_{pq} \bar{D}^A r Y_A + \hat{D}_p Y_q + \hat{D}_q Y_p. \quad (4.42)$$

Comparing with the gauge-transformation rules (4.3)–(4.5), the expression (4.40)–(4.42) are summarized as

$$h_{ab} =: \mathcal{F}_{ab} + \mathcal{L}_Y g_{ab}, \quad (4.43)$$

where \mathcal{F}_{ab} is the gauge-invariant part in the 2+2 formulation. The components of \mathcal{F}_{ab} is given by

$$\mathcal{F}_{AB} = F_{AB} = \sum_{l,m} \tilde{F}_{AB} S_\delta, \quad (4.44)$$

$$\mathcal{F}_{Ap} = r F_{Ap} = r \sum_{l,m} \tilde{F}_A \varepsilon_{pq} \hat{D}^q S_\delta, \quad \hat{D}^p \mathcal{F}_{Ap} = 0, \quad (4.45)$$

$$\mathcal{F}_{pq} = \frac{1}{2} \gamma_{pq} r^2 F = \frac{1}{2} \gamma_{pq} r^2 \sum_{l,m} \tilde{F} S_\delta. \quad (4.46)$$

Here, we note that the above arguments include not only $l \geq 2$ modes but also $l = 0, 1$ modes of metric perturbations. Equations (4.43)–(4.46) is complete proof of the Conjecture 2.1 for the perturbations on the spherically symmetric background spacetime. Therefore, our general arguments on the gauge-invariant perturbation theory reviewed in Sec. 2 are applicable to perturbations on the Schwarzschild background spacetime without special treatment of $l = 0, 1$ modes. Thus, we have resolved the zero-mode problem in the perturbations on the Schwarzschild background spacetime.

We also note that we only used the forms (3.1) and (3.4) of the background metric and did not used the specific forms of the Schwarzschild metric (3.3). Therefore, our construction of the gauge-invariant and gauge-variant part of the metric perturbation is also valid for the metric perturbations on any spherically symmetric spacetime. Thus, if we accept Proposal 3.1, we reached to the following statement:

Theorem 4.1. *If the gauge-transformation rule for a tensor field h_{ab} is given by $\mathcal{Y}h_{ab} - \mathcal{X}h_{ab} = \xi_{(1)}^a g_{ab}$ with the background metric g_{ab} with spherical symmetry, there then exist a tensor field \mathcal{F}_{ab} and a vector field Y^a such that h_{ab} is decomposed as $h_{ab} =: \mathcal{F}_{ab} + \xi_Y g_{ab}$, where \mathcal{F}_{ab} and Y^a are transformed into $\mathcal{Y}\mathcal{F}_{ab} - \mathcal{X}\mathcal{F}_{ab} = 0$ and $\mathcal{Y}Y^a - \mathcal{X}Y^a = \xi_{(1)}^a$ under the gauge transformation, respectively.*

5. Einstein equations

Here, we consider the linearized Einstein equations (2.11) on the spherically symmetric background spacetime with the metric (3.1). The gauge-invariant part of the linearized Einstein tensor ${}^{(1)}\mathcal{G}_a{}^b[\mathcal{F}]$ is given by Eqs. (2.9) and (2.10). The components of the tensor fields $H_{abc}[\mathcal{F}]$, $H_{ab}{}^c[\mathcal{F}]$, and $H_a{}^{bc}[\mathcal{F}]$ in terms of the variables F_{AB} , F_{Ap} and F in Eqs. (4.44)–(4.46) are summarized in Appendix C. Through these formulae and the mode decomposition in Eqs. (4.44)–(4.46) with the harmonic functions S_δ defined by Eq. (3.139), the components of the tensor ${}^{(1)}\mathcal{G}_a{}^b[\mathcal{F}]$ are given by

$$\begin{aligned}
{}^{(1)}\mathcal{G}_A{}^B &= \frac{1}{2} \sum_{l,m} \left[\left(-\bar{D}_D \bar{D}^D + \frac{l(l+1)}{r^2} - \frac{2}{r} (\bar{D}_D r) \bar{D}^D \right) \tilde{F}_A{}^B + \left(\bar{D}_D \bar{D}_A + \frac{2}{r} (\bar{D}_D r) \bar{D}_A \right) \tilde{F}^{BD} \right. \\
&\quad + \left(\bar{D}^D \bar{D}^B + \frac{2}{r} (\bar{D}^D r) \bar{D}^B \right) \tilde{F}_{AD} - \bar{D}_A \bar{D}^B \tilde{F}_D{}^D \\
&\quad \left. - \left(\bar{D}_A \bar{D}^B + \frac{1}{r} (\bar{D}_A r) \bar{D}^B + \frac{1}{r} (\bar{D}^B r) \bar{D}_A \right) \tilde{F} \right] S_\delta \\
&\quad + \frac{1}{2} \delta_A{}^B \sum_{l,m} \left[\left(\bar{D}_E \bar{D}^E - \frac{l(l+1)+1}{r^2} + \frac{2}{r} (\bar{D}_E r) \bar{D}^E + \frac{1}{r^2} (\bar{D}^E r) (\bar{D}_E r) \right) \tilde{F}_D{}^D \right. \\
&\quad \left. - \left(\bar{D}_D \bar{D}_E + \frac{4}{r} (\bar{D}_D r) \bar{D}_E + \frac{2}{r^2} (\bar{D}_E r) (\bar{D}_D r) \right) \tilde{F}^{ED} \right. \\
&\quad \left. + \left(\bar{D}_D \bar{D}^D - \frac{l(l+1)-2}{2r^2} + \frac{3}{r} (\bar{D}^D r) \bar{D}_D \right) \tilde{F} \right] S_\delta, \tag{5.1}
\end{aligned}$$

$$\begin{aligned}
{}^{(1)}\mathcal{G}_A{}^q &= \frac{1}{2r^2} \sum_{l,m} \left[\left(-\bar{D}_A + \frac{1}{r} (\bar{D}_A r) \right) \tilde{F}_D{}^D + \bar{D}^D \tilde{F}_{AD} - \frac{1}{2} \bar{D}_A \tilde{F} \right] \hat{D}^q S_\delta \\
&\quad + \frac{1}{2r} \sum_{l,m} \left[\left(-\bar{D}^D \bar{D}_D + \frac{l(l+1)}{r^2} - \frac{2}{r} (\bar{D}^D r) \bar{D}_D + \frac{3}{2r^2} \{ (\bar{D}^D r) (\bar{D}_D r) - 1 \} \right) \tilde{F}_A \right. \\
&\quad \left. + \left(\bar{D}^D \bar{D}_A + \frac{3}{r} (\bar{D}^D r) \bar{D}_A - \frac{1}{r} (\bar{D}_A r) \bar{D}^D - \frac{2}{r^2} (\bar{D}_A r) (\bar{D}^D r) \right) \tilde{F}_D \right] \varepsilon^{qt} \hat{D}_t S_\delta, \tag{5.2}
\end{aligned}$$

$$\begin{aligned}
{}^{(1)}\mathcal{G}_p{}^B &= \frac{1}{2} \sum_{l,m} \left[\left(-\bar{D}^B + \frac{1}{r} (\bar{D}^B r) \right) \tilde{F}_D{}^D + \bar{D}_D \tilde{F}^{BD} - \frac{1}{2} \bar{D}^B \tilde{F} \right] \hat{D}_p S_\delta \\
&\quad + \frac{r}{2} \sum_{l,m} \left[\left(-\bar{D}_D \bar{D}^D + \frac{l(l+1)}{r^2} - \frac{2}{r} (\bar{D}_D r) \bar{D}^D + \frac{3}{2r^2} \{ (\bar{D}^D r) (\bar{D}_D r) - 1 \} \right) \tilde{F}^B \right. \\
&\quad \left. + \left(\bar{D}_D \bar{D}^B + \frac{3}{r} (\bar{D}_D r) \bar{D}^B - \frac{1}{r} (\bar{D}^B r) \bar{D}_D - \frac{2}{r^2} (\bar{D}^B r) (\bar{D}_D r) \right) \tilde{F}^D \right] \varepsilon_{pq} \hat{D}^q S_\delta, \tag{5.3}
\end{aligned}$$

$$\begin{aligned}
{}^{(1)}\mathcal{G}_p^q &= \sum_{l,m} \left[\frac{1}{2r^2} \bar{D}_D (r^2 \bar{D}^D \tilde{F}) - \frac{1}{r^2} \bar{D}_D (r^2 \bar{D}_E \tilde{F}^{ED}) \right. \\
&\quad \left. + \left(\bar{D}_E \bar{D}^E + \frac{1}{r} (\bar{D}_E r) \bar{D}^E - \frac{l(l+1)}{2r^2} \right) \tilde{F}_D^D \right] \frac{1}{2} \gamma_p^q S_\delta \\
&\quad + \frac{1}{2r^2} \sum_{l,m} \left[-\tilde{F}_D^D \left(\hat{D}_p \hat{D}^q - \frac{1}{2} \gamma_p^q \hat{D}^s \hat{D}_s \right) S_\delta - \bar{D}_D (r \tilde{F}^D) (\varepsilon^{sq} \hat{D}_p \hat{D}_s + \varepsilon_{sp} \hat{D}^q \hat{D}^s) S_\delta \right], \quad (5.4)
\end{aligned}$$

where we used the fact that the background Ricci curvature vanishes and the background Einstein equations (B67) and (B68).

We also decompose the components of the linearized energy-momentum tensor ${}^{(1)}\mathcal{T}_a^b$ as follows:

$${}^{(1)}\mathcal{T}_A^B = \sum_{l,m} \tilde{T}_A^B S_\delta, \quad (5.5)$$

$${}^{(1)}\mathcal{T}_A^q = \frac{1}{r} \sum_{l,m} \left\{ \tilde{T}_{(e1)A} \hat{D}^q S_\delta + \tilde{T}_{(o1)A} \varepsilon^{qr} \hat{D}_r S_\delta \right\}, \quad (5.6)$$

$${}^{(1)}\mathcal{T}_p^B = r \sum_{l,m} \left\{ \tilde{T}_{(e1)}^B \hat{D}_p S_\delta + \tilde{T}_{(o1)}^B \varepsilon_{pr} \hat{D}^r S_\delta \right\}, \quad (5.7)$$

$$\begin{aligned}
{}^{(1)}\mathcal{T}_p^q &= \sum_{l,m} \left\{ \tilde{T}_{(e0)} \frac{1}{2} \gamma_p^q S_\delta + \tilde{T}_{(e2)} \left(\hat{D}_p \hat{D}^q S_\delta - \frac{1}{2} \gamma_p^q \hat{D}_r \hat{D}^r S_\delta \right) \right. \\
&\quad \left. + \tilde{T}_{(o2)} (\varepsilon_{sp} \hat{D}^q \hat{D}^s S_\delta + \varepsilon^{sq} \hat{D}_p \hat{D}_s S_\delta) \right\}. \quad (5.8)
\end{aligned}$$

The linearized continuity equation (2.17) for the energy-momentum tensor \mathcal{T}_a^b is summarized as

$$\bar{D}^C \tilde{T}_C^B + \frac{2}{r} (\bar{D}^D r) \tilde{T}_D^B - \frac{1}{r} l(l+1) \tilde{T}_{(e1)}^B - \frac{1}{r} (\bar{D}^B r) \tilde{T}_{(e0)} = 0, \quad (5.9)$$

$$\bar{D}^C \tilde{T}_{(e1)C} + \frac{3}{r} (\bar{D}^C r) \tilde{T}_{(e1)C} + \frac{1}{2r} \tilde{T}_{(e0)} - \frac{1}{2r} (l-1)(l+2) \tilde{T}_{(e2)} = 0, \quad (5.10)$$

$$\bar{D}^C \tilde{T}_{(o1)C} + \frac{3}{r} (\bar{D}^D r) \tilde{T}_{(o1)D} + \frac{1}{r} (l-1)(l+2) \tilde{T}_{(o2)} = 0. \quad (5.11)$$

Through the components (5.1)–(5.4) for the linearized Einstein tensor and the components (5.5)–(5.8) for the linearized energy-momentum tensor, we evaluate the linearized Einstein equation (2.11). Due to the linear-independence of the set of harmonics (3.12), we can carry out the mode-by-mode analyses including $l = 0, 1$ modes. Since the odd-mode perturbations and the even-mode perturbations are decoupled with each other, we consider these perturbations, separately.

5.1. Odd mode perturbation equations

From the linearized Einstein equation (2.11) through Eqs. (5.1)–(5.4) and Eq. (5.5)–(5.8), the odd-mode part in the linearized Einstein equations are simplified as the constraint equation

$$\bar{D}_D (r \tilde{F}^D) = -16\pi r^2 \tilde{T}_{(o2)}, \quad (5.12)$$

and the evolution equation

$$\begin{aligned}
& - \left[\bar{D}^D \bar{D}_D - \frac{l(l+1)}{r^2} \right] (r \tilde{F}_A) - \frac{2}{r^2} (\bar{D}^D r) (\bar{D}_A r) (r \tilde{F}_D) + \frac{2}{r} (\bar{D}^D r) \bar{D}_A (r \tilde{F}_D) \\
& = 16\pi r (\tilde{T}_{(o1)A} + r \bar{D}_A \tilde{T}_{(o2)}). \quad (5.13)
\end{aligned}$$

Furthermore, we have the continuity equation (5.11) for the odd-mode matter perturbation which is derived from the divergence of the first-order perturbation of the energy-momentum tensor. The

explicit strategy to solve these odd-mode perturbations and $l = 0, 1$ mode solutions will be discussed in Sec. 6 in this paper.

5.2. Even mode perturbation equations

Here, we consider the even-mode perturbations from Eqs. (5.1)–(5.4) and (5.5)–(5.8). The traceless even part of the (p, q) -component of the linearized Einstein equation (2.11) is given by

$$\tilde{F}_D{}^D = -16\pi r^2 \tilde{T}_{(e2)}. \quad (5.14)$$

Using this equation, the even part of (A, q) -component, equivalently (p, B) -component, of the linearized Einstein equation (2.11) yields

$$\bar{D}^D \tilde{\mathbb{F}}_{AD} - \frac{1}{2} \bar{D}_A \tilde{F} = 16\pi \left[r \tilde{T}_{(e1)A} - \frac{1}{2} r^2 \bar{D}_A \tilde{T}_{(e2)} \right] =: 16\pi S_{(ec)A} \quad (5.15)$$

through the definition of the traceless part $\tilde{\mathbb{F}}_{AB}$ of the variable \tilde{F}_{AB} defined by

$$\tilde{\mathbb{F}}_{AB} := \tilde{F}_{AB} - \frac{1}{2} y_{AB} \tilde{F}_C{}^C. \quad (5.16)$$

Using Eqs. (5.14), (5.15), and the component (B67) of background Einstein equation, the trace part of (p, q) -component of the linearized Einstein equation (2.11) is given by

$$\bar{D}^D \tilde{T}_{(e1)D} + \frac{3}{r} (\bar{D}^D r) \tilde{T}_{(e1)D} + \frac{1}{2r} \tilde{T}_{(e0)} - \frac{(l-1)(l+2)}{2r} \tilde{T}_{(e2)} = 0 \quad (5.17)$$

This coincides with the component (5.10) of the continuity equation for the linearized energy-momentum tensor.

Next, we consider the (A, B) -components of the linearized Einstein equation (2.11). Through Eqs. (5.14) and (5.15), the trace part of the (A, B) -component of the linearized Einstein equation (2.11) is given by

$$\left(\bar{D}_D \bar{D}^D + \frac{2}{r} (\bar{D}^D r) \bar{D}_D - \frac{(l-1)(l+2)}{r^2} \right) \tilde{F} - \frac{4}{r^2} (\bar{D}_C r) (\bar{D}_{Dr}) \tilde{\mathbb{F}}^{CD} = 16\pi S_{(F)}, \quad (5.18)$$

$$S_{(F)} := \tilde{T}_C{}^C + 4(\bar{D}_D r) \tilde{T}_{(e1)}^D - 2r(\bar{D}_{Dr}) \bar{D}^D \tilde{T}_{(e2)} - (l(l+1) + 2) \tilde{T}_{(e2)}. \quad (5.19)$$

On the other hand, the traceless part of the (A, B) -component of the linearized Einstein equation (2.11) is given by

$$\begin{aligned} & \left[-\bar{D}_D \bar{D}^D - \frac{2}{r} (\bar{D}_D r) \bar{D}^D + \frac{4}{r} (\bar{D}^D \bar{D}_{Dr}) + \frac{l(l+1)}{r^2} \right] \tilde{\mathbb{F}}_{AB} + \frac{4}{r} (\bar{D}^D r) \bar{D}_{(A} \tilde{\mathbb{F}}_{B)D} - \frac{2}{r} (\bar{D}_{(Ar}) \bar{D}_{B)} \tilde{F} \\ & = 16\pi S_{(\mathbb{F})AB}, \end{aligned} \quad (5.20)$$

$$\begin{aligned} S_{(\mathbb{F})AB} & := T_{AB} - \frac{1}{2} y_{AB} T_C{}^C - 2 \left(\bar{D}_{(A} (r \tilde{T}_{(e1)B)}) - \frac{1}{2} y_{AB} \bar{D}^D (r \tilde{T}_{(e1)D}) \right) \\ & + 2 \left((\bar{D}_{(Ar}) \bar{D}_{B)} - \frac{1}{2} y_{AB} (\bar{D}^D r) \bar{D}_D \right) (r \tilde{T}_{(e2)}) + r \left(\bar{D}_A \bar{D}_B - \frac{1}{2} y_{AB} \bar{D}^D \bar{D}_D \right) (r \tilde{T}_{(e2)}) \\ & + 2 \left((\bar{D}_{Ar}) (\bar{D}_{Br}) - \frac{1}{2} y_{AB} (\bar{D}^C r) (\bar{D}_{Cr}) \right) \tilde{T}_{(e2)} + 2y_{AB} (\bar{D}^C r) \tilde{T}_{(e1)C} - ry_{AB} (\bar{D}^C r) \bar{D}_C \tilde{T}_{(e2)}, \end{aligned} \quad (5.21)$$

where we used the background Einstein equation (B68).

Equations (5.14), (5.15), (5.18), and (5.20) are all independent equations of the linearized Einstein equation for even-mode perturbations. These equations are coupled equations for the variables \tilde{F}_C^C , F , and \tilde{F}_{AB} and the energy-momentum tensor for the matter field. When we solve these equations, we have to take into account of the continuity equations (5.9) and (5.10) for the matter fields. We note that these equations are valid not only for $l \geq 2$ modes but also $l = 0, 1$ modes in our formulation.

The explicit strategy to solve these Einstein equations for even modes, and the explicit solution for $l = 0, 1$ mode perturbations are discussed in the Part II paper [48].

6. Component treatment for the odd-mode perturbations of the Einstein equations

6.1. Strategy to solve odd-mode perturbations

Here, we consider the component treatment for the odd-mode perturbations based on the old paper by Regge and Wheeler [17], and Zerilli [18, 19] and re-derivation by Nakano [20]. We introduce the component of $r\tilde{F}^D$ as

$$r\tilde{F}_D =: X_{(o)}(dt)_D + Y_{(o)}(dr)_D, \quad r\tilde{F}^D = -f^{-1}X_{(o)}(\partial_t)^D + fY_{(o)}(\partial_r)^D, \quad (6.1)$$

where the background metric is given by Eqs. (3.1)–(3.4). In terms of the components (6.1), Eq. (5.12) is given by

$$-\partial_t X_{(o)} + ff'Y_{(o)} + f^2\partial_r Y_{(o)} = -16\pi r^2 f\tilde{T}_{(o)2}, \quad (6.2)$$

where $f' = \partial_r f$. The components of Eq. (5.13) are summarized as follows:

$$\begin{aligned} & \frac{1}{f}\partial_t^2 X_{(o)} - f\partial_r^2 X_{(o)} - \frac{2(1-f)}{r^2}X_{(o)} + \frac{l(l+1)}{r^2}X_{(o)} - \frac{1-3f}{r}\partial_t Y_{(o)} \\ & = 16\pi r (\tilde{T}_{(o)1t} + r\partial_t \tilde{T}_{(o)2}), \end{aligned} \quad (6.3)$$

$$\begin{aligned} & \partial_t^2 Y_{(o)} - f\partial_r(f\partial_r Y_{(o)}) + \frac{2(2f-1)f}{r}\partial_r Y_{(o)} + \frac{(l-1)(l+2)}{r^2}fY_{(o)} + \frac{(1-f)(5f-1)}{r^2}Y_{(o)} \\ & = +16\pi r (f\tilde{T}_{(o)1r} + rf\partial_r \tilde{T}_{(o)2} + (1-f)\tilde{T}_{(o)2}). \end{aligned} \quad (6.4)$$

In addition to these equations, the odd-mode perturbation (5.11) of the divergence of the energy-momentum tensor.

Here, we consider Eqs. (6.4). We define the dependent variable $Z_{(o)}$ by

$$Y_{(o)} =: \frac{r}{f}Z_{(o)} \quad (6.5)$$

and we have obtained the famous equation which is called Regge-Wheeler equation

$$\partial_t^2 Z_{(o)} - f\partial_r(f\partial_r Z_{(o)}) + \frac{1}{r^2}f[l(l+1) - 3(1-f)]Z_{(o)} = 16\pi f [f\tilde{T}_{(o)1r} + r\partial_r(f\tilde{T}_{(o)2})]. \quad (6.6)$$

We can solve Eq. (6.6) with appropriate boundary conditions and obtain the variable $Y_{(o)}$ through Eq. (6.5). For the $l \geq 2$ case, the analytic solutions to Eq. (6.6) are constructed by the formulation proposed by Mano, Suzuki, and Takasugi [65–68] (MST formulation). However, this is a partial solution to the odd-mode Einstein equations. We cannot regard such solutions as the total solution to the Einstein equation for odd-mode perturbations, because we have other two equations of the Einstein equation (6.3) and the constraint equation (6.2). To obtain the solution to the total Einstein equations for odd-mode perturbations, we have to discuss Eqs. (6.2), (6.3), and (5.11), i.e.,

$$-\frac{1}{f}\partial_t \tilde{T}_{(o)1t} + f\partial_r \tilde{T}_{(o)1r} + f'\tilde{T}_{(o)1r} + \frac{3}{r}f\tilde{T}_{(o)1r} + \frac{1}{r}(l-1)(l+2)\tilde{T}_{(o)2} = 0. \quad (6.7)$$

in addition to Eq. (6.6).

To obtain the solution to the total Einstein equations for odd-mode perturbations, it is convenient to introduce the Cunningham-Price-Moncrief variable $\Phi_{(o)}$ [23] by

$$\Phi_{(o)} := 2r \left[r^2 \partial_r \left(\frac{X_{(o)}}{r^2} \right) - \partial_t Y_{(o)} \right] \quad (6.8)$$

$$= 2r \partial_r X_{(o)} - 4X_{(o)} - 2r \partial_t Y_{(o)}. \quad (6.9)$$

Here, we consider the time derivative of $\Phi_{(o)}$ and use Eqs. (6.2), (6.4), and the background Einstein equation (B65) as

$$\begin{aligned} \partial_t \Phi_{(o)} &= 2 \frac{(l-1)(l+2)}{r} f Y_{(o)} - 32\pi r^2 f \tilde{T}_{(o1)r} \\ &= 2(l-1)(l+2)Z_{(o)} - 32\pi r^2 f \tilde{T}_{(o1)r}. \end{aligned} \quad (6.10)$$

The relation (6.10) indicates that the variable $Z_{(o)}$ is related to $\Phi_{(o)}$ for $l \neq 1$ modes, while the time derivative of $\Phi_{(o)}$ is just the matter degree of freedom $\tilde{T}_{(o1)r}$ for the $l = 1$ mode. This relation also gives the relation with the metric perturbation $Y_{(o)}$ as

$$(l-1)(l+2)Y_{(o)} = \frac{r}{2f} \partial_t \Phi_{(o)} + 16\pi r^3 \tilde{T}_{(o1)r}. \quad (6.11)$$

On the other hand, using Eqs. (6.2) and (6.3), the r -derivative of $\Phi_{(o)}$ through Eq. (6.9) is given by

$$\partial_r \Phi_{(o)} = -\frac{1}{r} \Phi_{(o)} + \frac{2}{rf} (l-1)(l+2)X_{(o)} - 32\pi \frac{r^2}{f} \tilde{T}_{(o1)t}. \quad (6.12)$$

Then, we obtain the relation

$$(l-1)(l+2)X_{(o)} = \frac{f}{2} (r \partial_r \Phi_{(o)} + \Phi_{(o)}) + 16\pi r^3 \tilde{T}_{(o1)t}. \quad (6.13)$$

From Eqs. (6.10) and (6.12) and the constraint (6.2), we obtain

$$\begin{aligned} \partial_r \partial_t \Phi_{(o)} - \partial_t \partial_r \Phi_{(o)} &= \partial_r \left[+2 \frac{(l-1)(l+2)}{r} f Y_{(o)} - 32\pi r^2 f \tilde{T}_{(o1)r} \right] \\ &\quad - \partial_t \left[-\frac{1}{r} \Phi_{(o)} + \frac{1}{r} 2(l-1)(l+2) \frac{1}{f} X_{(o)} - 32\pi r^2 \frac{1}{f} \tilde{T}_{(o1)t} \right] \\ &= -32\pi r^2 \left[-\frac{1}{f} \partial_t \tilde{T}_{(o1)t} + f' \tilde{T}_{(o1)r} + f \partial_r \tilde{T}_{(o1)r} \right. \\ &\quad \left. + \frac{3}{r} f \tilde{T}_{(o1)r} + \frac{1}{r} (l-1)(l+2) \tilde{T}_{(o2)} \right] \\ &= 0. \end{aligned} \quad (6.14)$$

The final equality comes from the odd-mode perturbation (6.7) of the energy-momentum tensor. Thus, Eqs. (6.10) and (6.12) are integrable under the constraint (6.2) and the continuity equation (6.7).

We emphasize that the relations (6.11) and (6.13) gives the relations of the metric components $(X_{(o)}, Y_{(o)})$ and the master variable $\Phi_{(o)}$ only for $l \neq 1$ mode. In the case of the $l = 1$ mode, these equations give the constraint of the master variable $\Phi_{(o)}$ and the matter degree of freedom. Furthermore, in the derivation of the relation (6.13), we used Eq. (6.3) and (6.2), which means that the relation (6.13) carries the information of Eq. (6.3).

From Eq. (6.10), we evaluate the second time-derivative of the master variable $\Phi_{(o)}$. On the other hand, from Eq. (6.12) we also evaluate the second derivative of $\Phi_{(o)}$ with respect to the tortoise coordinate $f\partial_r$. Furthermore, using Eqs. (6.9) and (6.13), we obtain

$$\partial_t^2 \Phi_{(o)} - f\partial_r [f\partial_r \Phi_{(o)}] + \frac{1}{r^2} f[l(l+1) - 3(1-f)]\Phi_{(o)} = 32\pi r f [\partial_r (r\tilde{T}_{(o1)t}) - r\partial_t \tilde{T}_{(o1)r}]. \quad (6.15)$$

This has the same form as Eq. (6.6) but we have different source terms from Eq. (6.6). For the $l \geq 2$ case, the analytic solutions to Eq. (6.15) is also constructed by the MST formulation [65–68]. In the vacuum case, Eq. (6.10) with $l \neq 1$ implies that the component $Y_{(o)}$ of the metric perturbation corresponds to the time-derivative of the variable $\Phi_{(o)}$. This indicates that Eq. (6.15) corresponds to the time-integration of Eq. (6.6) in the vacuum case. However, there is no degree of freedom of the integration constant in Eq. (6.15). Therefore, we may say that the initial conditions for Eq. (6.15) is restricted more than that of Eq. (6.6).

Here, we note that Eq. (6.11) is derived from Eqs. (6.2) and (6.6). This means that the relation (6.11) does not includes the information (6.3). On the other hand, the relation (6.13) is derived from Eq. (6.2) and (6.3). This means that the relation (6.13) does not includes the information of Eq. (6.6). In other words, we may regard the relation (6.11) as a result of Eq. (6.6), while Eq. (6.12) as a result of Eq. (6.3). Therefore, we obtain the two equations (6.11) and (6.13) from the three equations (6.2), (6.6), and (6.3). On the other hand, we have derived Eq. (6.15) from Eqs. (6.2), (6.6), and (6.3), which is independent of Eqs. (6.11) and (6.13). Thus, we may regard that all information of the set of three equations (6.2), (6.6), and (6.3) is included in the set of three equations (6.11), (6.13), and (6.15). In addition to these equations, we have to take into account of the continuity equation (6.7) for the odd-mode perturbations of the matter field.

However, as emphasized above, these arguments are not valid for $l = 1$ mode. Therefore, we have to reconsider the derivation of equations in the case of $l = 1$ mode, separately. Here, we examine the $l = 1$ modes. In this case, Eq. (6.10) is also valid, though this equation does not give the component $Y_{(o)}$ of the metric perturbations. In this case, the time-derivative of the variable $\Phi_{(o)}$ is given by

$$\partial_t \Phi_{(o)} = -32\pi r^2 f \tilde{T}_{(o1)r}, \quad (6.16)$$

which indicates that $\partial_t \Phi_{(o)}$ is determined by the matter degree of freedom. Similarly, Equation (6.12) is also valid even in the case of $l = 1$ mode, though this equation does not give the component $X_{(o)}$ of the metric perturbations. In this case, we obtain

$$f\partial_r \Phi_{(o)} = -\frac{1}{r} f \Phi_{(o)} - 32\pi r^2 \tilde{T}_{(o1)t}. \quad (6.17)$$

This equation indicates that the $\partial_r \Phi_{(o)}$ is also determined by the matter degree of freedom. From Eqs. (6.16) and (6.17), we can confirm that the variable $\Phi_{(o)}$ satisfies Eq. (6.15) with $l = 1$. However, we do not have to solve Eq. (6.15) with $l = 1$ in this case, because we can directly integrate Eqs. (6.16) and (6.17). Actually, the integrability condition $\partial_t \partial_r \Phi_{(o)} = \partial_r \partial_t \Phi_{(o)}$ of Eqs. (6.16) and (6.17) can be checked through the continuity equation (6.7) with $l = 1$.

Since we obtain the variable $\Phi_{(o)}$ by the direct integration of Eqs. (6.16) and (6.17), we can obtain the relation between the components $X_{(o)}$ and $Y_{(o)}$ of the metric perturbations through the definition (6.9). In addition to the solution $\Phi_{(o)}$, if we have a solution to $Z_{(o)} = \frac{f}{r} Y_{(o)}$, independently, we obtain the components $X_{(o)}$ and $Y_{(o)}$ of the metric perturbations through the above relation between $X_{(o)}$ and $Y_{(o)}$. Note that $Z_{(o)} = \frac{f}{r} Y_{(o)}$ can be determined through the integration of Eq. (6.6) with $l = 1$ with appropriate boundary conditions. In this case, the continuity equation (6.7) for odd-mode matter perturbations is used as consistency check of the solutions.

6.2. Odd-mode solutions

Since the construction of solutions for $l \geq 2$ mode is accomplished by the MST formulation [65–68], we discuss the $l = 0, 1$ -mode solutions for odd-mode perturbations along Proposal 3.1 and the strategy discussed in Sec. 6.1.

6.2.1. $l = 0$ odd mode. We choose Eq. (3.100) as the harmonic function $k_{(\hat{\Delta})}$ and used the set $\{\hat{D}_p k_{(\hat{\Delta})}, \varepsilon_{pr} \hat{D}^r k_{(\hat{\Delta})}, \hat{D}_p \hat{D}_q k_{(\hat{\Delta})}, 2\varepsilon_{r(p} \hat{D}_{q)} \hat{D}^r k_{(\hat{\Delta})}\}$ as the basis of the vector and tensors on S^2 . The bases of the odd-mode perturbations are $\varepsilon_{pr} \hat{D}^r k_{(\hat{\Delta})}$ and $2\varepsilon_{r(p} \hat{D}_{q)} \hat{D}^r k_{(\hat{\Delta})}$. Following Proposal 3.1, we choose $\delta = 0$ as the regularity of solutions when we solve the linearized Einstein equations. As shown in Eqs. (3.102) and (3.104), $\varepsilon_{pr} \hat{D}^r k_{(\hat{\Delta})} = 0 = 2\varepsilon_{r(p} \hat{D}_{q)} \hat{D}^r k_{(\hat{\Delta})}$. Then, we conclude that there is no nontrivial solution for odd-mode perturbations with $l = 0$.

6.2.2. $l = 1$ odd-mode vacuum solution. Following the strategy to solve the $l = 1$ odd-mode perturbation given in Sec. 6.1, we consider the equations (6.6), (6.7) with $l = 1$, (6.9), (6.16), and (6.17). To derive the non-vacuum solution to the linearized Einstein equations for $l = 1$ odd-mode perturbations, it is instructive to consider the vacuum case in which $\tilde{T}_{(o1)t} = \tilde{T}_{(o1)r} = \tilde{T}_{(o2)} = 0$.

From Eqs. (6.16) and (6.17), we obtain the solution to these equations as

$$\Phi_{(o)} = \frac{\alpha}{r}, \quad (6.18)$$

where α is constant of integration. On the other hand, $Y_{(o)}$ is obtained as the solution to the $l = 1$ version of the Regge-Wheeler equation (6.6) without source terms through Eq. (6.5). Here, we first consider the case $Y_{(o)} = 0$. Through the definition (6.8) of the variable $\Phi_{(o)}$ and Eq. (6.18), we obtain

$$\frac{\alpha}{r} = 2r \left[r^2 \partial_r \left(\frac{X_{(o)}}{r^2} \right) \right]. \quad (6.19)$$

The solution to Eq. (6.19) together with the assumption $Y_{(o)} = 0$ is a solution to the linearized Einstein equations for $l = 1$ odd-mode perturbations as follows:

$$X_{(o)} = -\frac{\alpha}{6r} + \partial_t \beta(t) r^2, \quad Y_{(o)} = 0. \quad (6.20)$$

From Eqs. (4.45) and (6.1), we can derive the gauge-invariant metric perturbation \mathcal{F}_{Ap} which corresponds to the solution (6.20). In the $l = 1$ modes, there are $m = 0, \pm 1$ modes. In this paper, we only consider the $m = 0$ -mode perturbation, since the generalization to $m = \pm 1$ modes is straightforward. If we choose $\delta = 0$ in the mode function (3.133), we obtain

$$\mathcal{F}_{Ap} = rF_{Ap} = r\tilde{F}_A \varepsilon_{pq} \hat{D}^q k_{(\hat{\Delta}+2, m=0)}, \quad \varepsilon_{pq} \hat{D}^q k_{(\hat{\Delta}+2, m=0)} = \sin^2 \theta (d\phi)_p. \quad (6.21)$$

Then, we have

$$\begin{aligned} 2\mathcal{F}_{Ap} (dx^A)_{(a} (dx^p)_{b)} &= 2rF_{(Ap)} (dx^A)_{(a} (dx^p)_{b)} \\ &= 2X_{(o)} \sin^2 \theta (dt)_{(a} (d\phi)_{b)} + 2Y_{(o)} \sin^2 \theta (dr)_{(a} (d\phi)_{b)} \end{aligned} \quad (6.22)$$

$$= \left(-\frac{\alpha}{3r} + 2\partial_t \beta(t) r^2 \right) \sin^2 \theta (dt)_{(a} (d\phi)_{b)}. \quad (6.23)$$

Here, the term $\partial_t \beta(t) r^2$ is diverge as $r \rightarrow \infty$. At this moment, we choose $\partial_t \beta(t) = 0$. Then, we have obtained

$$2\mathcal{F}_{Ap} (dx^A)_{(a} (dx^p)_{b)} = -\frac{\alpha}{3r} \sin^2 \theta (dt)_{(a} (d\phi)_{b)}. \quad (6.24)$$

Eq. (6.24) is the linearized Kerr solution. Actually, the Kerr solution with the Kerr parameter a is expressed as [69, 70]

$$g_{ab} = - \left[1 - \frac{2Mr}{\Sigma} \right] (dt)_a (dt)_b - \frac{2aMr \sin^2 \theta}{\Sigma} (dt)_{(a} (d\phi)_{b)} + \frac{\Sigma}{\Delta} (dr)_a (dr)_b + \Sigma (d\theta)_a (d\theta)_b + \left[r^2 + a^2 + \frac{2Mr}{\Sigma} a^2 \sin^2 \theta \right] \sin^2 \theta (d\phi)_a (d\phi)_b, \quad (6.25)$$

where

$$\Sigma := r^2 + a^2 \cos^2 \theta, \quad \Delta := r^2 + a^2 - 2Mr \quad (6.26)$$

In the metric (6.25), we replace $a \rightarrow \varepsilon a$, where ε is the parameter for the perturbative expansion. Then, when the Kerr metric (6.25) is expressed as follows:

$$g_{ab} = y_{ab} + r^2 \gamma_{ab} + \varepsilon \left(-\frac{2aM}{r} \sin^2 \theta (dt)_{(a} (d\phi)_{b)} \right) + O(\varepsilon^2), \quad (6.27)$$

Comparing Eqs. (6.24) and (6.27), the constant of integration α in Eq. (6.24) is identified as the angular momentum perturbation in Kerr solution by choosing

$$\frac{\alpha}{3} = 2aM =: 2a_{10}M. \quad (6.28)$$

Thus, we have seen that the solution (6.23) is given using the Kerr parameter a_{10} as follows:

$$2\mathcal{F}_{Ap}(dx^A)_{(a}(dx^P)_{b)} = 2 \left(-\frac{a_{10}M}{r} + \partial_t \beta(t) r^2 \right) \sin^2 \theta (dt)_{(a} (d\phi)_{b)}. \quad (6.29)$$

Next, we consider the physical meaning of the arbitrary function $\beta(t)$ in the solution (6.23). If we consider the frame with the rigid rotation

$$t = t', \quad \phi = \varphi + \varepsilon \omega(t') t' \quad (6.30)$$

In terms of (t', φ) , the background metric (3.1) with Eqs. (3.3) and (3.4).

$$g_{ab} = -f(dt')_a (dt')_b + f^{-1} (dr)_a (dr)_b + r^2 [(d\theta)_a (d\theta)_b + \sin^2 \theta (d\varphi)_a (d\varphi)_b] + 2\varepsilon (\partial_{t'} \omega(t') t' + \omega(t')) r^2 \sin^2 \theta (dt')_{(a} (d\varphi)_{b)} + O(\varepsilon^2). \quad (6.31)$$

Comparing Eq. (6.31) and Eq. (6.29), we can see that the arbitrary function $\beta(t)$ corresponds to

$$\partial_t \beta(t) = \partial_t \omega(t) t + \omega(t). \quad (6.32)$$

Thus, we may interpret the integration constant $\partial_t \beta(t)$ as non-inertia term due to the rigidly rotating frame with the angular velocity $\omega(t)$. We also note that this angular velocity $\omega(t)$ may depend on t .

Finally, we consider the solution $Y_{(o)}$ to Eq. (6.5) and (6.6). Here, we consider the situation $Y_{(o)} \neq 0$ and introduce the variable $W_{(o)}$ as follows:

$$Y_o =: r^2 \partial_r W_{(o)}, \quad Z_{(o)} = \frac{f}{r} Y_{(o)} = r f \partial_r W_{(o)}. \quad (6.33)$$

Through the solution (6.18) with Eq. (6.28) and the definition (6.8) of the variable $\Phi_{(o)}$, we obtain the equation

$$\frac{6a_{10}M}{r} = 2r \left[r^2 \partial_r \left(\frac{X_{(o)}}{r^2} \right) - r^2 \partial_t \partial_r W_{(o)} \right]. \quad (6.34)$$

Integrating this equation, we obtain

$$X_{(o)} = -\frac{a_{10}M}{r} + \partial_t \beta(t) r^2 + r^2 \partial_t W_{(o)}. \quad (6.35)$$

Through Eqs. (4.45) and (6.1), we obtain

$$\begin{aligned} 2\mathcal{F}_{Ap}(dx^A)_{(a)}(dx^p)_{(b)} &= 2 \left(-\frac{a_{10}M}{r} + r^2 \partial_t \beta(t) + r^2 \partial_t W_{(o)} \right) \sin^2 \theta (dt)_{(a)}(d\phi)_{(b)} \\ &\quad + 2r^2 \partial_r W_{(o)} \sin^2 \theta (dr)_{(a)}(d\phi)_{(b)}. \end{aligned} \quad (6.36)$$

Note again that the variable $Z_{(o)} = r f \partial_r W_{(o)}$ satisfy the Regge-Wheeler equation (6.6) with $l = 1$.

The above interpretation of the arbitrary function $\beta(t)$ as the inertia force on the rigidly rotation frame is instructive to consider the interpretation of the odd-mode vacuum solution (6.36). To see this, we consider the component expression of $\mathcal{L}_V g_{ab}$, where V^a is constructed from gauge-invariant variables, which is discussed in Sec. 2. To obtain the components of $\mathcal{L}_V g_{ab}$, the explicit components of the Christoffel symbol Γ_{ab}^c for the background metric (3.1) with Eqs. (3.3) and (3.4) are convenient, which are summarized in Eqs. (B71). Here, we assume that $V_a = V_\phi (d\phi)_a$, then the non-vanishing components of $\mathcal{L}_V g_{ab}$ are given by

$$\mathcal{L}_V g_{t\phi} = \partial_t V_\phi, \quad \mathcal{L}_V g_{r\phi} = \partial_r V_\phi - \frac{2}{r} V_\phi, \quad \mathcal{L}_V g_{\theta\phi} = \partial_\theta V_\phi - 2 \cot \theta V_\phi. \quad (6.37)$$

Comparing Eqs. (6.36) and (6.37), we obtain

$$V_a = (\beta(t) + W_{(o)}(t, r)) r^2 \sin^2 \theta (d\phi)_a, \quad (6.38)$$

$$\begin{aligned} \mathcal{L}_V g_{ab} &= \partial_t (\beta(t) + W_{(o)}(t, r)) r^2 \sin^2 \theta 2(dt)_{(a)}(d\phi)_{(b)} \\ &\quad + (\partial_r W_{(o)}(t, r)) r^2 \sin^2 \theta 2(dr)_{(a)}(d\phi)_{(b)}. \end{aligned} \quad (6.39)$$

This coincides with the perturbation (6.36) with the condition of the vanishing Kerr parameter $a_{10} = 0$. Then, we have

$$2\mathcal{F}_{Ap}(dx^A)_{(a)}(dx^p)_{(b)} = -\frac{2a_{10}M}{r} \sin^2 \theta (dt)_{(a)}(d\phi)_{(b)} + \mathcal{L}_V g_{ab}, \quad (6.40)$$

$$V_a = (\beta(t) + W_{(o)}(t, r)) r^2 \sin^2 \theta (d\phi)_a. \quad (6.41)$$

Here, we note that the vector field V_a and $\mathcal{L}_V g_{ab}$ are gauge-invariant.

6.3. Odd mode non-vacuum $l = 1$ solution

Inspecting the derivation of the vacuum solution for $l = 1$ modes in Sec. 6.2.2, we consider the non-vacuum solution for $l = 1$ modes. For $l = 1$ modes, the linearized Einstein equations for the master variable $\Phi_{(o)}$ defined by Eq. (6.8) are given by Eqs. (6.16) and (6.17). As mentioned in Sec. 6.1, the

integrability condition for these equations is guaranteed by the continuity equation (6.7) with $l = 1$. Inspecting Eqs. (6.18) and (6.28), we consider the solution in the form

$$\Phi_{(o)} = \frac{6Ma_1(t,r)}{r}. \quad (6.42)$$

Substituting Eq. (6.42) into Eqs. (6.16) and (6.17), we obtain

$$\partial_t a_1(t,r) = -\frac{16\pi}{3M} r^3 f \tilde{T}_{(o1)r}, \quad \partial_r a_1(t,r) = -\frac{16\pi}{3M} r^3 \frac{1}{f} \tilde{T}_{(o1)t}. \quad (6.43)$$

The integrability of Eqs. (6.43) is equivalent to the integrability of Eqs. (6.16) and (6.17) which is guaranteed by the continuity equation (6.7) with $l = 1$. Then, we may integrate Eqs. (6.43) as follows:

$$\begin{aligned} a_1(t,r) &= -\frac{16\pi}{3M} r^3 f \int dt \tilde{T}_{(o1)r} + a_{10} \\ &= -\frac{16\pi}{3M} \int dr r^3 \frac{1}{f} \tilde{T}_{(o1)t} + a_{10}, \end{aligned} \quad (6.44)$$

where a_{10} is the constant which corresponds to the Kerr parameter a in Eq. (6.25) as shown in the vacuum case.

Similar arguments to those in Sec. 6.2.2, which lead the results (6.40) and (6.41), also leads

$$2\mathcal{F}_{Ap}(dx^A)_{(a)}(dx^p)_{(b)} = 6Mr^2 \left[\int dr \frac{a_1(t,r)}{r^4} \right] \sin^2 \theta (dt)_{(a)}(d\phi)_{(b)} + \mathcal{L}_V g_{ab}, \quad (6.45)$$

$$V_a = (\beta(t) + W_{(o)}(t,r)) r^2 \sin^2 \theta (d\phi)_a. \quad (6.46)$$

Here, we note that the vector field V_a and $\mathcal{L}_V g_{ab}$ are gauge-invariant. However, we can eliminate the term $\mathcal{L}_V g_{ab}$ by an gauge-fixing procedure. From this situation, we may say that the concept of “the complete gauge fixing” is stronger restriction of the metric perturbation than the concept of “gauge-invariant”.

7. Summary and Discussions

In summary, after reviewing our general framework of the gauge-invariant perturbation theory, we discussed the resolution of the “zero-mode problem” in perturbations on the Schwarzschild background spacetime. The “zero-mode problem” in the context of our general framework of the gauge-invariant perturbation theory corresponds to the $l = 0, 1$ mode problem in perturbations of the Schwarzschild background spacetime. In the review of our general framework of the gauge invariant perturbation theory, we emphasize the importance of the distinction of the first- and the second-kind gauge in general relativity. It should be emphasized that our general framework for the gauge-invariant perturbation theory is the formulation to exclude the second-kind gauge in general relativistic perturbation theories.

As emphasize in Sec. 2, Conjecture 2.1 is the non-trivial and important premise of our general framework of gauge-invariant perturbation theories. If Conjecture 2.1 is actually true, we can develop gauge-invariant perturbation theory on general background spacetime and we can also extend this gauge-invariant perturbation theory to higher-order perturbation theory. For this reason, the gauge-invariant treatment of the $l = 0, 1$ modes in perturbations of the Schwarzschild background spacetime is important not only for the development of the linear perturbations but also for the development of the higher-order perturbation theory on the Schwarzschild background spacetime.

To find the gauge-invariant treatment of the $l = 0, 1$ mode perturbations on the Schwarzschild background spacetime, we first reviewed 2+2 formulation in which the decomposition formulae (3.7)–(3.9) with the spherical harmonic functions Y_{lm} as the scalar harmonic function S and explained why $l = 0, 1$ modes should be separately treated in conventional perturbation theory on the Schwarzschild background spacetime. The special treatment in the conventional formulation caused by the loss of the linear independence of the set (3.12) of the tensor harmonic functions on S^2 , i.e., vector and/or tensor harmonic functions vanishes in $l = 0, 1$ modes and does not play a role of the bases of tangent space on S^2 .

To recover this situation, instead of the spherical harmonics Y_{00} and Y_{1m} for $l = 0, 1$ modes, we introduce the mode functions $k_{(\hat{\Delta})}$ and $k_{(\hat{\Delta}+2)m}$, which belongs to the kernel of the derivative operator $\hat{\Delta}$ and $\hat{\Delta} + 2$, respectively. We also derive the sufficient condition for which the decomposition formulae (3.7)–(3.9) with the harmonic function $S = S_\delta$ defined by Eq. (3.34) is invertible not only for $l \geq 2$ modes but also $l = 0, 1$ modes. As the result, we showed that the mode functions (3.140)–(3.142) with the parameter δ for $l = 0, 1$ modes satisfy this sufficient condition. These mode functions realize the conventional spherical harmonic functions Y_{00} and Y_{1m} when $\delta = 0$, but the set of harmonic functions (3.34) loses the linear independence as the bases of the tangent space on S^2 as the conventional case. On the other hand, when $\delta \neq 0$, the set of the mode functions (3.34) has the linear-independence as the bases of the tangent space on S^2 . However, the mode functions $k_{(\hat{\Delta})}$ and $k_{(\hat{\Delta}+2)m}$ with $\delta \neq 0$ are singular functions.

Due to this situation, we proposed Proposal 3.1 as the strategy to define the gauge-invariant variables for $l = 0, 1$ modes and to derive and solve the linearized Einstein equation. Following Proposal 3.1, we can construct gauge-invariant and gauge-variant variables for linear metric perturbation through the similar manner to the case of the $l \geq 2$ modes. This construction is a proof of Conjecture 2.1 for the perturbations on the spherically symmetric background spacetime. Then, we reach to the statement Theorem 4.1. Owing to Theorem 4.1, we can develop gauge-invariant perturbation theory on spherically symmetric background spacetimes including $l = 0, 1$ modes. Furthermore, Theorem 4.1 yields that we can develop higher-order gauge-invariant perturbation theory on spherically symmetric background spacetimes, although this development is beyond the current scope of this paper. A brief discussion of this development to higher-order perturbations is already given in Ref. [50].

Besides the discussion on the extension to the higher-order perturbation theory, it is also true that we are proposing unusual procedure as Proposal 3.1. Therefore, we have to confirm that the solutions to the Einstein equation derived by Proposal 3.1 are physically reasonable. To check this, we derived the linearized Einstein equations on the Schwarzschild background spacetime following Proposal 3.1. We consider the mode decomposition of the general expression of the linearized energy-momentum tensor as the source term of the linearized Einstein equations. To solve the derived linearized Einstein equations, the linearized perturbation of the continuity equation of the energy-momentum tensor should be taken into account. The metric perturbations on the Schwarzschild spacetime are classified into the odd-mode and the even-mode perturbations. In this Part I paper, we concentrate only on the odd-mode perturbations and derive the $l = 0, 1$ -mode solutions following Proposal 3.1.

For odd-mode perturbations, we examined the strategy to solve the linearized Einstein equations for any l modes following the Proposal 3.1, through we take care of the structure of equations for $l = 1$ mode perturbations. As well-known, to solve the odd-mode perturbations, Einstein equations

for the $l \geq 2$ odd-mode perturbations are reduced to the Regge-Wheeler equation. Furthermore, the solutions to the Regge-Wheeler equation for $l \geq 2$ modes are constructed through the MST formulation [65–68]. Therefore, we concentrated on the $l = 0, 1$ mode perturbations.

Following Proposal 3.1, for $l = 0$ odd-mode perturbations, we reached to the conclusion that there is no non-trivial solution to the linearized Einstein equation as expected. Then, we carefully examined the solutions to the Einstein equations for $l = 1$ odd-mode perturbations. We first consider the vacuum solution to the linearized Einstein equation in which the linear perturbation of the energy-momentum tensor vanishes. Then, we obtain the linearized Kerr parameter perturbation with the term given in the form of the Lie derivative of the background metric g_{ab} . Through the variation of constant, we derived the general solutions to non-vacuum linearized Einstein equations for the $l = 1$ odd-mode perturbations. Since we use the constant Kerr parameter in the variation of constant, we can expect that the obtained general solution describes the spin-up or the spin-down of the black hole due to the effect of the linearized energy-momentum tensor.

In addition to the Kerr parameter perturbations, we obtain the term which has the form of the Lie derivative of the background metric g_{ab} in the derived solution. The appearance of such term is natural due to the diffeomorphism symmetry of the linearized Einstein equations. We also note that the appearance of such terms is related to the non-uniqueness of the gauge-invariant variables which was emphasized by Eq. (2.18) in Sec. 2. Since our formulation completely exclude the gauge degree of freedom of the second kind, we have to regard that such terms are the gauge degree of freedom of the first kind which was explained in Sec. 2. Similar results are also obtained in even-mode perturbations which will be shown in the Part II paper.

Apart from these terms of the Lie derivative of the background metric g_{ab} , in vacuum case, the only non-trivial solution in $l = 1$ odd-mode perturbation is the Kerr parameter perturbations. This will be related to the uniqueness of the Kerr solution in the vacuum Einstein equations in the local sense [14], though the assertion of the uniqueness theorem of the Kerr solution includes topological statement. Besides the relation of the uniqueness theorem of Kerr black hole, at least, we may say that the derived vacuum solution for $l = 0, 1$ odd-mode perturbations is physically reasonable. In this sense, we may say that Proposal 3.1 is also physically reasonable.

Acknowledgements

The author deeply acknowledged to Professor Hiroyuki Nakano for various discussions and suggestions.

Appendix

A. Explicit form of conventional spherical harmonics on S^2

First, we summarize the properties of the conventional spherical harmonic functions Y_{lm} . The spherical harmonic functions $Y_{lm}(\theta, \phi)$ satisfy the equations

$$[\hat{\Delta} + l(l+1)]Y_{lm} = 0, \quad (\text{A1})$$

$$\partial_\phi Y_{lm} = imY_{lm}. \quad (\text{A2})$$

To be explicit, they are expressed in terms of the Legendre functions as

$$Y_{lm}(\theta, \phi) = \sqrt{\frac{(2l+1)(l-m)!}{4\pi(l+m)!}} P_l^m(\cos\theta) e^{im\phi}. \quad (\text{A3})$$

For $l = 0, 1$ modes, the spherical harmonic functions $Y_{lm} = Y_{l,m}$ are explicitly given by

$$Y_{00} = \sqrt{\frac{1}{4\pi}}, \quad (A4)$$

$$Y_{10} = \sqrt{\frac{3}{4\pi}} \cos \theta, \quad Y_{11} = \sqrt{\frac{3}{8\pi}} \sin \theta e^{i\phi}, \quad Y_{1-1} = -\sqrt{\frac{3}{8\pi}} \sin \theta e^{-i\phi}. \quad (A5)$$

Employing these spherical harmonic functions (A3) as the scalar harmonics, we construct the set of the tensor harmonics on S^2 . Since the dimension of S^2 is two, we have enough number of tensor harmonic functions as bases of tangent space on S^2 .

On the unit sphere any vector field v^p is written in terms of two scalar functions v and w as

$$v^p = \hat{D}^p v + \varepsilon^{pq} \hat{D}_q w. \quad (A6)$$

Here, $\hat{D}^p v$ is even part and $\varepsilon^{pq} \hat{D}_q w$ is the odd part, which corresponds to $\hat{D}_p S$ and $\varepsilon_{pq} \hat{D}^q S$ in Eq. (3.8), respectively. If we choose $S = Y_{lm}$, these vectors are given by

$$\hat{D}_p Y_{lm}, \quad \varepsilon_{pq} \hat{D}^q Y_{lm}. \quad (A7)$$

For $l = 0$ modes, the spherical harmonic function Y_{00} is constant as in Eq. (A4) and corresponding vector harmonics vanish:

$$\hat{D}_p Y_{00} = 0, \quad \varepsilon_{pq} \hat{D}^q Y_{00} = 0. \quad (A8)$$

On the other hand, for $l = 1$ modes, vector harmonics has the vector values as

$$\hat{D}_p Y_{10} = -\sqrt{\frac{3}{4\pi}} \sin \theta \theta_p, \quad (A9)$$

$$\hat{D}_p Y_{11} = \sqrt{\frac{3}{8\pi}} e^{i\phi} (\cos \theta \theta_p + i\phi_p), \quad \hat{D}_p Y_{1-1} = \sqrt{\frac{3}{8\pi}} e^{-i\phi} (-\cos \theta \theta_p + i\phi_p) \quad (A10)$$

and

$$\varepsilon_{pq} \hat{D}^q Y_{10} = \sqrt{\frac{3}{4\pi}} \sin \theta \phi_p, \quad (A11)$$

$$\varepsilon_{pq} \hat{D}^q Y_{11} = \sqrt{\frac{3}{8\pi}} e^{i\phi} (-\cos \theta \phi_p + i\theta_p), \quad \varepsilon_{pq} \hat{D}^q Y_{1-1} = \sqrt{\frac{3}{8\pi}} e^{-i\phi} (\cos \theta \phi_p + i\theta_p). \quad (A12)$$

Thus, vector harmonics has its vector value for $l = 1$ modes, while does not for $l = 0$ mode.

Any smooth symmetric second-rank tensor field t^{pq} on the unit sphere can be expressed in terms of its trace $t = t^p_p$ and two scalar fields v and w as

$$t^{pq} = \frac{1}{2} t \gamma^{pq} + \left(\hat{D}^p \hat{D}^q - \frac{1}{2} \gamma^{pq} \hat{\Delta} \right) v + 2\varepsilon^{r(q} \hat{D}^p) D_r w. \quad (A13)$$

These three terms correspond to the terms proportional to $\frac{1}{2} \gamma_{pq} S$, $\left(\hat{D}_p \hat{D}_q - \frac{1}{2} \gamma^{pq} \hat{\Delta} \right) S$, and $2\varepsilon_{r(p} \hat{D}_{q)} \hat{D}^r S$ in Eq. (3.9). As in the case of vector harmonics above, for $l = 0$ modes, the spherical harmonic function Y_{00} is constant as in Eq. (A4) and the only non-vanishing harmonics is its

trace part

$$\frac{1}{2}\gamma_{pq}Y_{00} = \frac{1}{2}\gamma_{pq}\sqrt{\frac{1}{4\pi}} \quad (\text{A14})$$

and the other traceless even and odd parts vanish. For $l = 1$ modes, from Eqs. (A5), the trace parts are trivially given by

$$\frac{1}{2}\gamma_{pq}Y_{10} = \frac{1}{2}\sqrt{\frac{3}{4\pi}}\cos\theta\gamma_{pq}, \quad (\text{A15})$$

$$\frac{1}{2}\gamma_{pq}Y_{11} = \frac{1}{2}\sqrt{\frac{3}{8\pi}}\sin\theta e^{i\phi}\gamma_{pq}, \quad (\text{A16})$$

$$\frac{1}{2}\gamma_{pq}Y_{1-1} = -\frac{1}{2}\sqrt{\frac{3}{8\pi}}\sin\theta e^{-i\phi}\gamma_{pq}. \quad (\text{A17})$$

On the other hand, the traceless even and odd parts for $\left(\hat{D}_p\hat{D}_q - \frac{1}{2}\gamma^{pq}\hat{\Delta}\right)Y_{1m}$, and $2\varepsilon_{r(p}\hat{D}_{q)}\hat{D}^rY_{1m}$ identically vanish for all $m = -1, 0, 1$.

As a summary of $S = Y_{lm}$ cases, for $l = 0$ mode, any vector and tensor harmonics does not have their values, and these do not play roles of bases of the tangent space on S^2 . On the other hand, for $l = 1$ modes, the vector harmonics have their vector value and play roles of bases of the tangent space on S^2 . The trace parts of the second-rank tensor of each modes have their tensor values, while all traceless even and odd mode harmonics identically vanish and does not play roles of bases of the tangent space on S^2 .

B. Covariant derivatives in 2+2 formulation and background curvatures

In this Appendix, we summarize the relation between the covariant derivatives ∇_a associated with the metric g_{ab} , \bar{D}_A associated with the metric y_{ab} , and \hat{D}_p associated with the metric γ_{ab} . These formulae are convenient to derive the gauge-transformation rules, linearized Einstein equations, and so on. Here, the metrics g_{ab} , y_{ab} , and γ_{ab} are given by Eq. (3.1). We assume that y_{ab} depends on $\{x^A\}$ and $r = r(x^A)$. We also assume that γ_{ab} depends only on $\{x^p\}$. Under these assumptions, the Christoffel symbol $\Gamma_{ab}{}^c$ are given by

$$\Gamma_{ab}{}^c = \frac{1}{2}g^{cd}(\partial_a g_{db} + \partial_b g_{da} - \partial_d g_{ab}), \quad (\text{B1})$$

$$\Gamma_{AB}{}^C = \frac{1}{2}y^{CD}(\partial_A y_{DB} + \partial_B y_{DA} - \partial_D y_{AB}) =: \bar{\Gamma}_{AB}{}^C, \quad (\text{B2})$$

$$\Gamma_{pB}{}^C = 0, \quad (\text{B3})$$

$$\Gamma_{pq}{}^C = -r(\bar{D}^C r)\gamma_{pq}, \quad (\text{B4})$$

$$\Gamma_{AB}{}^P = 0, \quad (\text{B5})$$

$$\Gamma_{qA}{}^P = \frac{1}{r}(\bar{D}_A r)\gamma_q{}^P, \quad (\text{B6})$$

$$\Gamma_{qr}{}^P = \frac{1}{2}\gamma^{pd}(\partial_q \gamma_{dr} + \partial_r \gamma_{dq} - \partial_d \gamma_{qr}) =: \hat{\Gamma}_{qr}{}^P. \quad (\text{B7})$$

Here, we note that

$$\hat{D}_p \bar{D}_A t_B = \bar{D}_A \hat{D}_p t_B, \quad (\text{B8})$$

and

$$\hat{D}_p \bar{D}_A t_q = \partial_p \bar{D}_A t_q - \hat{\Gamma}_{qp}{}^r \bar{D}_A t_q = \bar{D}_A \hat{D}_p t_q, \quad (\text{B9})$$

since

$$\partial_p \bar{\Gamma}_{AB}{}^C = 0, \quad \partial_A \hat{\Gamma}_{pq}{}^r = 0. \quad (\text{B10})$$

Then, we obtain the formulae for the covariant derivatives $\nabla_a v_b$ and $\nabla_a t^b$ as

$$\nabla_A v_B = \bar{D}_A v_B, \quad (\text{B11})$$

$$\nabla_A v_p = \bar{D}_A v_p - \frac{1}{r} \bar{D}_A r v_p, \quad (\text{B12})$$

$$\nabla_p v_A = \hat{D}_p v_A - \frac{1}{r} \bar{D}_A r v_p, \quad (\text{B13})$$

$$\nabla_p v_q = \hat{D}_p v_q + r \bar{D}^A r \gamma_{pq} v_A, \quad (\text{B14})$$

$$\nabla_A t^B = \bar{D}_A t^B, \quad (\text{B15})$$

$$\nabla_A t^p = \partial_A t^p + \frac{1}{r} \bar{D}_A r t^p, \quad (\text{B16})$$

$$\nabla_p t^A = \hat{D}_p t^A - r \bar{D}^A r \gamma_{pq} t^q, \quad (\text{B17})$$

$$\nabla_p t^q = \hat{D}_p t^q + \frac{1}{r} \bar{D}_A r \gamma_p{}^q t^A. \quad (\text{B18})$$

Here, we also summarize the expression of $\nabla_a T_{bc}$ for an arbitrary tensor T_{bc} in terms of the covariant derivatives \bar{D}_A and \hat{D}_p which are associated with the metric y_{AB} and γ_{pq} , respectively, from

$$\nabla_a T_{bc} = \partial_a T_{bc} - \Gamma_{ba}{}^d T_{dc} - \Gamma_{ca}{}^d T_{bd}. \quad (\text{B19})$$

These are given by

$$\nabla_A T_{BC} = \bar{D}_A T_{BC}, \quad (\text{B20})$$

$$\nabla_A T_{Bp} = \bar{D}_A T_{Bp} - \frac{1}{r} \bar{D}_A r T_{Bp}, \quad (\text{B21})$$

$$\nabla_A T_{pC} = \bar{D}_A T_{pC} - \frac{1}{r} \bar{D}_A r T_{pC}, \quad (\text{B22})$$

$$\nabla_p T_{BC} = \hat{D}_p T_{BC} - \frac{1}{r} \bar{D}_B r T_{pC} - \frac{1}{r} \bar{D}_C r T_{Bp}, \quad (\text{B23})$$

$$\nabla_p T_{qC} = \hat{D}_p T_{qC} + r \bar{D}^D r \gamma_{qp} T_{DC} - \frac{1}{r} \bar{D}_C r T_{qp}, \quad (\text{B24})$$

$$\nabla_p T_{Bq} = \hat{D}_p T_{Bq} - \frac{1}{r} \bar{D}_B r T_{pq} + r \bar{D}^D r \gamma_{qp} T_{BD}, \quad (\text{B25})$$

$$\nabla_A T_{pq} = \bar{D}_A T_{pq} - \frac{2}{r} \bar{D}_A r T_{pq}, \quad (\text{B26})$$

$$\nabla_p T_{qr} = \hat{D}_p T_{qr} + r \bar{D}^D r \gamma_{qp} T_{Dr} + r \bar{D}^D r \gamma_{rp} T_{qD}. \quad (\text{B27})$$

Furthermore, to derive the linearized Einstein equation, we have to derive the components of

$$\nabla_a H_c{}^{bd} = \partial_a H_c{}^{bd} - \Gamma_{ca}{}^e H_e{}^{bd} + \Gamma_{ea}{}^b H_c{}^{ed} + \Gamma_{ea}{}^d H_c{}^{be}. \quad (\text{B28})$$

Then, these are summarized as

$$\nabla_A H_C^{BD} = \bar{D}_A H_C^{BD}, \quad (\text{B29})$$

$$\nabla_A H_C^{Bs} = \bar{D}_A H_C^{Bs} + \frac{1}{r} \bar{D}_A r H_C^{Bs}, \quad (\text{B30})$$

$$\nabla_A H_C^{qD} = \bar{D}_A H_C^{qD} + \frac{1}{r} \bar{D}_A r H_C^{qD}, \quad (\text{B31})$$

$$\nabla_A H_C^{qs} = \bar{D}_A H_C^{qs} + \frac{2}{r} \bar{D}_A r H_C^{qs}, \quad (\text{B32})$$

$$\nabla_A H_r^{BD} = \bar{D}_A H_r^{BD} - \frac{1}{r} \bar{D}_A r H_r^{BD}, \quad (\text{B33})$$

$$\nabla_A H_r^{Bs} = \bar{D}_A H_r^{Bs}, \quad (\text{B34})$$

$$\nabla_A H_r^{qD} = \bar{D}_A H_r^{qD}, \quad (\text{B35})$$

$$\nabla_A H_r^{qs} = \bar{D}_A H_r^{qs} + \frac{1}{r} \bar{D}_A r H_r^{qs}, \quad (\text{B36})$$

$$\nabla_p H_C^{BD} = \hat{D}_p H_C^{BD} - \frac{1}{r} \bar{D}_C r H_p^{BD} - r \bar{D}^B r \gamma_{tp} H_C^{tD} - r \bar{D}^D r \gamma_{tp} H_C^{Bt}, \quad (\text{B37})$$

$$\nabla_p H_C^{Bs} = \hat{D}_p H_C^{Bs} - \frac{1}{r} \bar{D}_C r H_p^{Bs} + \frac{1}{r} \bar{D}_E r \gamma_p^s H_C^{BE} - r \bar{D}^B r \gamma_{tp} H_C^{ts}, \quad (\text{B38})$$

$$\nabla_p H_C^{qD} = \hat{D}_p H_C^{qD} - \frac{1}{r} \bar{D}_C r H_p^{qD} + \frac{1}{r} \bar{D}_E r \gamma_p^q H_C^{ED} - r \bar{D}^D r \gamma_{tp} H_C^{qt}, \quad (\text{B39})$$

$$\nabla_p H_C^{qs} = \hat{D}_p H_C^{qs} - \frac{1}{r} \bar{D}_C r H_p^{qs} + \frac{1}{r} \bar{D}_E r \gamma_p^q H_C^{Es} + \frac{1}{r} \bar{D}_E r \gamma_p^s H_C^{qE}, \quad (\text{B40})$$

$$\nabla_p H_r^{BD} = \hat{D}_p H_r^{BD} + r \bar{D}^E r \gamma_{rp} H_E^{BD} - r \bar{D}^B r \gamma_{tp} H_r^{tD} - r \bar{D}^D r \gamma_{tp} H_r^{Bt}, \quad (\text{B41})$$

$$\nabla_p H_r^{Bs} = \hat{D}_p H_r^{Bs} + r \bar{D}^E r \gamma_{rp} H_E^{Bs} - r \bar{D}^B r \gamma_{tp} H_r^{ts} + \frac{1}{r} \bar{D}_E r \gamma_p^s H_r^{BE}, \quad (\text{B42})$$

$$\nabla_p H_r^{qD} = \hat{D}_p H_r^{qD} + r \bar{D}^E r \gamma_{rp} H_E^{qD} - r \bar{D}^D r \gamma_{tp} H_r^{qt} + \frac{1}{r} \bar{D}_E r \gamma_p^q H_r^{ED}, \quad (\text{B43})$$

$$\nabla_p H_r^{qs} = \hat{D}_p H_r^{qs} + r \bar{D}^E r \gamma_{rp} H_E^{qs} + \frac{1}{r} \bar{D}_E r \gamma_p^q H_r^{Es} + \frac{1}{r} \bar{D}_E r \gamma_p^s H_r^{qE}. \quad (\text{B44})$$

Next, we summarize the components of the background curvatures induced by the metric Eq. (3.1). We derive these components through the components of the connection (B2)–(B7) and the formula of the Riemann curvature

$$R_{abc}{}^d = \partial_b \Gamma_{ac}{}^d - \partial_a \Gamma_{bc}{}^d + \Gamma_{ac}{}^e \Gamma_{eb}{}^d - \Gamma_{bc}{}^e \Gamma_{ea}{}^d. \quad (\text{B45})$$

To derive the components of this curvature, we use

$$\bar{D}_A \gamma_{pq} = 0 = \hat{D}_p \gamma_{AB}, \quad \hat{D}_p r = 0. \quad (\text{B46})$$

The components of the non-vanishing Riemann curvature are summarized as

$$R_{ABC}{}^D = {}^{(2)}\bar{R}_{ABC}{}^D, \quad (\text{B47})$$

$$R_{pBr}{}^D = -r(\bar{D}_B \bar{D}^D r) \gamma_{pr}, \quad (\text{B48})$$

$$R_{pBC}{}^s = \frac{1}{r} (\bar{D}_B \bar{D}_C r) \gamma_p^s, \quad (\text{B49})$$

$$R_{pqr}{}^s = {}^{(2)}\hat{R}_{pqr}{}^s - 2(\bar{D}^E r)(\bar{D}_E r) \gamma_{r[p} \gamma_{q]}^s. \quad (\text{B50})$$

The components of the Ricci curvature are summarized as

$$R_{AC} = {}^{(2)}\bar{R}_{AC} - \frac{2}{r}(\bar{D}_A \bar{D}_C r), \quad (\text{B51})$$

$$R_{Ar} = 0, \quad (\text{B52})$$

$$R_{pr} = {}^{(2)}\hat{R}_{pr} - [r(\bar{D}_E \bar{D}^E r) + (\bar{D}^E r)(\bar{D}_E r)] \gamma_{pr}. \quad (\text{B53})$$

The Ricci scalar curvature is given by

$$R = g^{ac} R_{ac} = {}^{(2)}\bar{R} + \frac{1}{r^2} {}^{(2)}\hat{R} - \frac{4}{r} (\bar{D}^C \bar{D}_C r) - \frac{2}{r^2} (\bar{D}^E r)(\bar{D}_E r). \quad (\text{B54})$$

Next, we derive the components of the Einstein tensor

$$G_{ab} := R_{ab} - \frac{1}{2} g_{ab} R \quad (\text{B55})$$

and its components are summarized as

$$G_{AB} = -\frac{2}{r} (\bar{D}_A \bar{D}_B r) + \frac{1}{r^2} y_{AB} [-1 + 2r(\bar{D}^C \bar{D}_C r) + (\bar{D}^E r)(\bar{D}_E r)], \quad (\text{B56})$$

$$G_{Aq} = 0, \quad (\text{B57})$$

$$G_{pq} = \gamma_{pq} \left[r(\bar{D}^C \bar{D}_C r) - \frac{1}{2} r^2 {}^{(2)}\bar{R} \right], \quad (\text{B58})$$

where we used the two-dimensional Einstein tensors are identically vanish and the fact that the metric γ_{pq} is the maximally symmetric space with positive curvature, i.e.,

$${}^{(2)}\hat{R}_{pqrs} = 2\gamma_{p[r} \gamma_{s]q}, \quad {}^{(2)}\hat{R}_{pr} = \gamma_{pr}, \quad {}^{(2)}\hat{R} = 2. \quad (\text{B59})$$

Here, we consider the static solution whose metric is given by

$$y_{AB} = -f(dt)_A(dt)_B + f^{-1}(dr)_A(dr)_B, \quad (\text{B60})$$

where $f = f(r)$. Due to the Birkhoff theorem, the vacuum solution with the spherically symmetric spacetime must be the Schwarzschild spacetime. We check this fact from Eqs. (B56)–(B58) with the substitution (B60). Actually, we obtain

$$\bar{D}_B r = (dr)_B, \quad \bar{D}^B r = f \left(\frac{\partial}{\partial r} \right)^B, \quad \bar{D}_A \bar{D}_B r = \frac{f'}{2} y_{AB}. \quad (\text{B61})$$

Then, we have

$$(\bar{D}^B r)(\bar{D}_B r) = f, \quad \bar{D}^A \bar{D}_B r = \frac{f'}{2} \delta_B^A, \quad \bar{D}^C \bar{D}_C r = f'. \quad (\text{B62})$$

From Eq. (B56) as

$$y^{AB} G_{AB} = \frac{2}{r} \left(f' - \frac{1-f}{r} \right) = 0, \quad G_{AB} - \frac{1}{2} y_{AB} G_{AB} = 0. \quad (\text{B63})$$

The solution to Eq. (B63) is given by

$$f = 1 - \frac{2M}{r}, \quad (\text{B64})$$

$$f' = \frac{1-f}{r}, \quad (\text{B65})$$

where M is the constant of integration. This is the Schwarzschild metric. We also evaluate the component $G_{pq} = 0$ through Eq. (B58) using Eq. (B64) as

$${}^{(2)}\bar{R} = \frac{2}{r}(\bar{D}^C \bar{D}_C r) \quad (\text{B66})$$

As the summary of the background vacuum Einstein equations, we have

$$r(\bar{D}^C \bar{D}_C r) + (\bar{D}^E r)(\bar{D}_E r) = 1, \quad (\text{B67})$$

$$(\bar{D}_A \bar{D}_B r) = \frac{1}{2} \gamma_{AB} (\bar{D}^C \bar{D}_C r), \quad (\text{B68})$$

$${}^{(2)}\bar{R} = \frac{2}{r}(\bar{D}^C \bar{D}_C r). \quad (\text{B69})$$

Eq. (B67) is equivalent to Eq. (B65). Since the two-dimensional curvature ${}^{(2)}\bar{R}_{DAEC}$ has only one independent component, ${}^{(2)}\bar{R}_{DAEC}$ is written as

$${}^{(2)}\bar{R}_{DAEC} = \frac{2}{r}(\bar{D}^F \bar{D}_F r) \gamma_{D[E} \gamma_{C]A}, \quad {}^{(2)}\bar{R}_{DE} = \frac{1}{r}(\bar{D}^F \bar{D}_F r) \gamma_{DE}, \quad {}^{(2)}\bar{R} = \frac{2}{r}(\bar{D}^F \bar{D}_F r). \quad (\text{B70})$$

The above formulae are expressed the covariant form of the 2+2 formulation. However, the explicit components of $\Gamma_{ab}{}^c$ are also convenient to leads the results in Sec. 6.2. From Eqs. (B2)–(B7) and the background metric (3.1) with Eqs. (3.3) and (3.4), non-vanishing components of $\Gamma_{ab}{}^c$ are summarized as

$$\begin{aligned} \Gamma_{tr}{}^t &= \frac{f'}{2f}, & \Gamma_{tt}{}^r &= \frac{1}{2} f f', & \Gamma_{rr}{}^r &= -\frac{f'}{2f}, & \Gamma_{\theta\theta}{}^r &= -r f, \\ \Gamma_{\phi\phi}{}^r &= -r f \sin^2 \theta, & \Gamma_{r\theta}{}^\theta &= \frac{1}{r}, & \Gamma_{\phi\phi}{}^\theta &= -\sin \theta \cos \theta, \\ \Gamma_{r\phi}{}^\phi &= \frac{1}{r}, & \Gamma_{\phi\theta}{}^\phi &= \cot \theta. \end{aligned} \quad (\text{B71})$$

C. Summary of the 2+2 representations of the tensor $H_{abc}[\mathcal{F}]$, $H_{ab}{}^c[\mathcal{F}]$, $H_a{}^{bc}[\mathcal{F}]$

Here, we summarize the components of $H_{abc}[\mathcal{F}]$ through the expressions (4.44)–(4.46):

$$H_{ABC} = \bar{D}_{(A} F_{B)C} - \frac{1}{2} \bar{D}_C F_{AB}, \quad (\text{C1})$$

$$H_{pBC} = \frac{1}{2} (\hat{D}_p F_{BC} + r \bar{D}_B F_{Cp} - r \bar{D}_C F_{Bp} - (\bar{D}_B r) F_{Cp} - (\bar{D}_C r) F_{Bp}), \quad (\text{C2})$$

$$H_{pqC} = \frac{1}{2} \left(2r \hat{D}_{(p} F_{q)C} - \frac{1}{2} \gamma_{pq} r^2 \bar{D}_C F - r (\bar{D}_C r) \gamma_{pq} F + 2r (\bar{D}^D r) \gamma_{pq} F_{DC} \right), \quad (\text{C3})$$

$$H_{ABr} = r \bar{D}_{(A} F_{B)r} + (\bar{D}_{(A} r) F_{B)r} - \frac{1}{2} \hat{D}_r F_{AB}, \quad (\text{C4})$$

$$H_{pBr} = \frac{1}{2} \left(r \hat{D}_p F_{rB} - r \hat{D}_r F_{pB} + \frac{1}{2} r^2 \gamma_{pr} \bar{D}_B F \right), \quad (\text{C5})$$

$$H_{pqr} = \frac{1}{2} r^2 \gamma_{r(q} \hat{D}_{p)} F - \frac{1}{4} r^2 \gamma_{pq} \hat{D}_r F + r^2 \bar{D}^D r \gamma_{pq} F_{Dr}. \quad (\text{C6})$$

Next, we summarize the components of $H_{ab}{}^c[\mathcal{F}]$ through the expressions (4.44)–(4.46):

$$H_{AB}{}^C = \bar{D}_{(A}F_{B)}{}^C - \frac{1}{2}\bar{D}^C F_{AB}, \quad (C7)$$

$$H_{pB}{}^C = \frac{1}{2}(\hat{D}_p F_B{}^C + r\bar{D}_B F_p{}^C - r\bar{D}^C F_{Bp} - (\bar{D}_B r)F_p{}^C - (\bar{D}^C r)F_{Bp}), \quad (C8)$$

$$H_{pq}{}^C = \frac{1}{2}\left(2r\hat{D}_{(p}F_{q)}{}^C - \frac{1}{2}\gamma_{pq}r^2\bar{D}^C F - r(\bar{D}^C r)\gamma_{pq}F + 2r(\bar{D}^D r)\gamma_{pq}F_D{}^C\right), \quad (C9)$$

$$H_{AB}{}^r = \frac{1}{r}\bar{D}_{(A}F_{B)}{}^r + \frac{1}{r^2}(\bar{D}_{(A}r)F_{B)}{}^r - \frac{1}{2r^2}\hat{D}^r F_{AB}, \quad (C10)$$

$$H_{pB}{}^r = \frac{1}{2r}\hat{D}_p F_B{}^r - \frac{1}{2r}\hat{D}^r F_{Bp} + \frac{1}{4}\gamma_p{}^r \bar{D}_B F, \quad (C11)$$

$$H_{pq}{}^r = \frac{1}{2}\gamma_{(q}{}^r \hat{D}_{p)}F - \frac{1}{4}\gamma_{pq}\hat{D}^r F + (\bar{D}^D r)\gamma_{pq}F_D{}^r. \quad (C12)$$

Finally, we summarize the component $H_a{}^{bc}[\mathcal{F}]$ through the expression (4.44)–(4.46):

$$H_A{}^{BC} = \frac{1}{2}(\bar{D}_A F^{BC} + \bar{D}^B F_A{}^C - \bar{D}^C F_A{}^B), \quad (C13)$$

$$H_A{}^{Br} = \frac{1}{2r}\bar{D}_A F^{Br} + \frac{1}{2r}\bar{D}^B F_A{}^r + \frac{1}{2r^2}(\bar{D}_A r)F^{Br} + \frac{1}{2r^2}(\bar{D}^B r)F_A{}^r - \frac{1}{2r^2}\hat{D}^r F_A{}^B, \quad (C14)$$

$$H_A{}^{qC} = \frac{1}{2r^2}(\hat{D}^q F_A{}^C + r\bar{D}_A F^{qC} - r\bar{D}^C F_A{}^q - (\bar{D}_A r)F^{qC} - (\bar{D}^C r)F_A{}^q), \quad (C15)$$

$$H_A{}^{qr} = \frac{1}{2r^3}\left[\hat{D}^q F_A{}^r - \hat{D}^r F_A{}^q + \frac{1}{2}r\gamma^{qr}\bar{D}_A F\right], \quad (C16)$$

$$H_p{}^{BC} = \frac{1}{2}(\hat{D}_p F^{BC} + r\bar{D}^B F_p{}^C - r\bar{D}^C F_p{}^B - (\bar{D}^B r)F_p{}^C - (\bar{D}^C r)F_p{}^B), \quad (C17)$$

$$H_p{}^{Br} = \frac{1}{2r}\hat{D}_p F^{Br} - \frac{1}{2r}\hat{D}^r F_p{}^B + \frac{1}{4}\gamma_p{}^r \bar{D}^B F, \quad (C18)$$

$$H_p{}^{qC} = \frac{1}{2r^2}\left(r\hat{D}_p F^{qC} + r\hat{D}^q F_p{}^C - \frac{1}{2}\gamma_p{}^q r^2\bar{D}^C F - r(\bar{D}^C r)\gamma_p{}^q F + 2r(\bar{D}^D r)\gamma_p{}^q F_D{}^C\right), \quad (C19)$$

$$H_p{}^{qr} = \frac{1}{r^2}\left(\frac{1}{4}\gamma^{qr}\hat{D}_p F + \frac{1}{4}\gamma_p{}^r \hat{D}^q F - \frac{1}{4}\gamma_p{}^q \hat{D}^r F + (\bar{D}^D r)\gamma_p{}^q F_D{}^r\right). \quad (C20)$$

References

- [1] B. P. Abbot et al., (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. Lett. **116** (2016), 061102.
- [2] LIGO Scientific Collaboration 2021 home page: <https://ligo.org>
- [3] Virgo 2021 home page : <https://virgo-gw.eu>
- [4] KAGRA 2021 home page: <https://gwcenter.icrr.u-tokyo.ac.jp>
- [5] LIGO INDIA 2021 home page : <https://ligo-india.in>
- [6] Einstein Telescope 2021 home page : <https://www.et-gw.eu>
- [7] Cosmic Explorer 2021 home page : <https://cosmicexplorer.org>
- [8] LISA home page : <https://lisa.nasa.gov>
- [9] S. Kawamura, et al., Prog. Theor. Exp. Phys. 2021 (2021), 05A105.
- [10] J. Mei, et al., Prog. Theor. Exp. Phys. 2020 (2020), 05A107.
- [11] Z. Luo, et al., Prog. Theor. Exp. Phys. 2020 (2020),05A108.
- [12] L. Barack and A. Pound, Rep. Prog. Phys. **82** (2019) 016904.
- [13] R. D. Blandford and R. L. Znajek, Mon. Not. Roy. Astron. Soc. **179** (1977), 433.
- [14] M. Kimura, T. Harada, A. Naruko, K. Toma, Prog. Theor. Exp. Phys. (2021) in press.
- [15] K. Nakamura, Prog. Theor. Phys. **110**, (2003), 723.
- [16] K. Nakamura, Prog. Theor. Phys. **113** (2005), 481.
- [17] T. Regge and J. A. Wheeler, Phys. Rev. **108** (1957), 1063.

-
- [18] F. Zerilli, Phys. Rev. Lett. **24** (1970), 737.
- [19] F. Zerilli, Phys. Rev. D **2** (1970), 2141.
- [20] H. Nakano, Private note on “Regge-Wheeler-Zerilli formalism” (2019).
- [21] V. Moncrief, Ann. Phys. (N.Y.) **88** (1974), 323.
- [22] V. Moncrief, Ann. Phys. (N.Y.) **88** (1974), 343.
- [23] C. T. Cunningham, R. H. Price, and V. Moncrief, Astrophys. J. **224** (1978), 643.
- [24] S. Chandrasekhar, *The mathematical theory of black holes* (Oxford: Clarendon Press, 1983).
- [25] U.H. Gerlach and U.K. Sengupta, Phys. Rev. D **19** (1979), 2268.
- [26] U.H. Gerlach and U.K. Sengupta, Phys. Rev. D **20** (1979), 3009.
- [27] U.H. Gerlach and U.K. Sengupta, J. Math. Phys. **20** (1979), 2540.
- [28] U.H. Gerlach and U.K. Sengupta, Phys. Rev. D **22** (1980), 1300.
- [29] T. Nakamura, K. Oohara, Y. Kojima, Prog. Theor. Phys. Suppl. No. **90** (1987), 1.
- [30] C. Gundlach and J.M. Martín-García, Phys. Rev. D **61** (2000), 084024.
- [31] J.M. Martín-García and C. Gundlach, Phys. Rev. D **64** (2001), 024012.
- [32] A. Nagar and L. Rezzolla, Class. Quantum Grav. **22** (2005), R167; Erratum *ibid.* **23** (2006), 4297.
- [33] K. Martel and E. Poisson, Phys. Rev. D **71** (2005), 104003.
- [34] K. Nakamura, Class. Quantum Grav. **28** (2011), 122001.
- [35] K. Nakamura, Int. J. Mod. Phys. D **21** (2012), 124004.
- [36] K. Nakamura, Prog. Theor. Exp. Phys. **2013** (2013), 043E02.
- [37] K. Nakamura, Class. Quantum Grav. **31** (2014), 135013.
- [38] K. Nakamura, Phys. Rev. D **74** (2006), 101301(R).
- [39] K. Nakamura, Prog. Theor. Phys. **117** (2007), 17.
- [40] K. Nakamura, ““Gauge” in General Relativity: – Second-order general relativistic gauge-invariant perturbation theory –”, in *Lie Theory and its Applications in Physics VII* ed. V. K. Dobrev et al, (Heron Press, Sofia, 2008)
- [41] K. Nakamura, Phys. Rev. D **80** (2009), 124021.
- [42] K. Nakamura, Prog. Theor. Phys. **121** (2009), 1321.
- [43] K. Nakamura, Advances in Astronomy, **2010** (2010), 576273.
- [44] A. J. Christopherson, K. A. Malik, D. R. -Matravers, K. Nakamura, Class. Quantum Grav. **28** (2011), 225024.
- [45] K. Nakamura, “Second-order Gauge-invariant Cosmological Perturbation Theory: Current Status Updated in 2019”, Chapter 1 in *Theory and Applications of Physical Science, Vol.3* Ed. M. Rafatullah, (Book Publisher International, 2020), ISBN 978-93-89816-24-2 (Print); ISBN 978-93-89816-25-9 (eBook), Preprint arXiv:1912.12805 [gr-qc].
- [46] K. Nakamura, Class. Quantum Grav. **38** (2021), 145010.
- [47] D. Brizuela, J. M. Martín Garcia, and G. A. Mena Marugan, Phys. Rev. D **76** (2007), 024004.
- [48] K. Nakamura, “Gauge-invariant perturbation theory on the Schwarzschild background spacetime Part II : — Even-mode perturbations —,” Preprint arXiv:2110.13512 [gr-qc].
- [49] K. Nakamura, “Gauge-invariant perturbation theory on the Schwarzschild background spacetime Part III : — Realization of exact solutions —,” Preprint arXiv:2110.13519 [gr-qc].
- [50] K. Nakamura, “Formal solution of the any-order mass, angular-momentum, dipole perturbations on the Schwarzschild background spacetime,” Letters in High Energy Physics in press. Preprint arXiv:2102.10650v3 [gr-qc].
- [51] R. K. Sachs, “Gravitational Radiation”, in *Relativity, Groups and Topology* ed. C. DeWitt and B. DeWitt, (New York: Gordon and Breach, 1964).
- [52] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry, volume I*, (Wiley Classic Library Edition (1996), John Wiley & Sons, Inc.)
- [53] J. M. Stewart and M. Walker, Proc. R. Soc. London A **341** (1974), 49.
- [54] J. M. Stewart, Class. Quantum Grav. **7** (1990), 1169.
- [55] J. M. Stewart, *Advanced General Relativity* (Cambridge University Press, Cambridge, 1991).
- [56] M. Bruni, S. Matarrese, S. Mollerach and S. Sonego, Class. Quantum Grav. **14** (1997), 2585.
- [57] M. Bruni and S. Sonego, Class. Quantum Grav. **16** (1999), L29.
- [58] S. Sonego and M. Bruni, Commun. Math. Phys. **193** (1998), 209.
- [59] J. W. York, Jr. J. Math. Phys. **14** (1973), 456.
- [60] J. W. York, Jr. Ann. Inst. H. Poincaré **21** (1974), 319.
- [61] S. Deser, Ann. Inst. H. Poincaré **7** (1967), 149.
- [62] R. Szmytkowski, J. Math. Phys. **47** (2006), 063506.
- [63] R. Szmytkowski, J. Phys. A: Math. Theor. **40** (2007), 995-1009; and references therein.
- [64] J.J. Sakurai, *Modern Quantum mechanics, Revised Edition*, edited by S. F. Tuan, (Addison-Wesley, 1994).
- [65] S. Mano, H. Suzuki, and E. Takasugi, Prog. Theor. Phys. **95** (1996), 1079.
- [66] S. Mano, H. Suzuki, and E. Takasugi, Prog. Theor. Phys. **96** (1996), 549.
- [67] S. Mano and E. Takasugi, Prog. Theor. Phys. **97** (1997), 213.
- [68] H. Tagoshi, S. Mano, and E. Takasugi, Prog. Theor. Phys. **98** (1997), 829.
- [69] R. H. Boyer and R. W. Lindquist, J. Math. Phys. **8** (1967), 265.

[70] R. M. Wald, *General Relativity* (Chicago, IL, University of Chicago Press, 1984).