

Static and Spherically Symmetric Solutions in $f(Q)$ Gravity

Wenyi Wang,^{1,*} Hua Chen,^{1,†} and Taishi Katsuragawa^{1,‡}

¹*Institute of Astrophysics, Central China Normal University, Wuhan 430079, China*

Abstract

$f(Q)$ gravity is the extension of Symmetric Teleparallel General Relativity (STGR), in which both curvature and torsion vanish, and gravity is attributed to non-metricity. This work performs theoretical analyses of static and spherically symmetric solutions with an anisotropic fluid for general $f(Q)$ gravity. We find that the off-diagonal component of field equation due to coincident gauge leads to stringent restrictions on the functional form of $f(Q)$ gravity, and thereby provides us with ways to construct models of cosmological interest in $f(Q)$ gravity. In addition, although the exact Schwarzschild solution only exists in STGR, we obtain Schwarzschild-like solutions in nontrivial $f(Q)$ gravity and study its asymptotic behavior and deviation from the exact one.

arXiv:2110.13565v1 [gr-qc] 26 Oct 2021

* wangwy@mails.cnu.edu.cn

† huachen@mails.cnu.edu.cn

‡ taishi@mail.cnu.edu.cn

I. INTRODUCTION

Modern observations have confirmed that the expansion of the current Universe is accelerating, which could be sourced by some unknown components, called dark energy (DE) [1, 2]. A possible way to explain DE is to modify pure General Relativity (GR), where gravity is ascribed to curvature, with the torsion and non-metricity assumed to vanish. For instance, Λ CDM model is the minimal modification, where the cosmological constant Λ is responsible for DE. Along this way, one can generalize GR to $f(R)$ gravity by improving the Ricci scalar R to its functional form [3, 4]. On the other hand, choosing the torsion T or the non-metricity Q as a geometric basis provides two different but equivalent descriptions of gravity. These are the so-called Teleparallel Equivalent of General Relativity (TEGR) [5, 6] and Symmetric Teleparallel General Relativity (STGR)¹ [7, 8].

In STGR theory, both the curvature and the torsion vanish, as the non-metricity describes the gravity. In this theory, under the teleparallelism constraint, we can always choose the coincident gauge, which restricts the affine connection to disappear, and makes the metric tensor the only basic variable. In analogy to $f(R)$ gravity, TEGR and STGR can be generalized to $f(T)$ gravity [9] and $f(Q)$ gravity [8, 10]. Although the latter one is relatively less investigated, it has many similar properties to those in $f(T)$ gravity. For instance, as similar to $f(T)$ theory [11–13], the gauge choice breaks the coordinate transformation invariant in $f(Q)$ theory, which predicts different consequences in various coordinate systems [14]. Recently, there have been several applications in $f(Q)$ theory; cosmology [15–18], bouncing model [19], wormhole solutions [20], energy conditions [21], and Newtonian limit [22] have been discussed.

In the analogy to $f(T)$ gravity, it is known that the non-zero off-diagonal component of field equations in $f(T)$ gravity, which originates from the specific gauge choice, restricts the functional form of $f(T)$ [23]. Therefore, it would also put restrictions on the functional form of $f(Q)$ gravity, which potentially gives us a guideline for the model building of $f(Q)$ gravity. This work aims to investigate possible functional forms of $f(Q)$ under the restriction of the static and spherically symmetric geometry with an anisotropic fluid. In particular, we will show that there is no exact Schwarzschild solution for the nontrivial $f(Q)$ function. With the non-metricity scalar Q being constant, we also analyze the deviation of the metric from

¹ This theory is also called Symmetric Teleparallel Equivalent to General Relativity (STTEGR).

the exact Schwarzschild solution.

This paper is organized as follows. In Sec. II, we briefly review $f(Q)$ theory, where the action and the equations of motions are introduced. In Sec. III, we derive the non-metricity scalar Q and the equations of motion for generic static and spherically symmetric geometry with an anisotropic fluid. We will see that the off-diagonal component of field equations leads to two solutions: STGR which recovers GR, and the constant non-metricity scalar. Then, with a focus on the second solution in Sec. IV, we investigate the constraints on the functional form of $f(Q)$ and geometry under various conditions. Finally, Sec. V is devoted to the conclusions and discussions.

II. $f(Q)$ GRAVITY

$f(Q)$ gravity considers a generic metric-affine spacetime, in which the metric tensor $g_{\mu\nu}$ and connection $\Gamma_{\mu\nu}^\lambda$ are treated independently, and the non-metricity of the connection is defined by

$$Q_{\alpha\mu\nu} \equiv \nabla_\alpha g_{\mu\nu} = \partial_\alpha g_{\mu\nu} - \Gamma_{\alpha\mu}^\lambda g_{\lambda\nu} - \Gamma_{\alpha\nu}^\lambda g_{\mu\lambda}. \quad (1)$$

The general form of affine connection can be decomposed into the following three independent components,

$$\Gamma_{\mu\nu}^\lambda = \{\overset{\lambda}{\mu\nu}\} + K_{\mu\nu}^\lambda + L_{\mu\nu}^\lambda, \quad (2)$$

where $\{\overset{\lambda}{\mu\nu}\}$ denotes the Levi-Civita connection determined by the metric $g_{\mu\nu}$

$$\{\overset{\lambda}{\mu\nu}\} \equiv \frac{1}{2}g^{\lambda\beta} (\partial_\mu g_{\beta\nu} + \partial_\nu g_{\beta\mu} - \partial_\beta g_{\mu\nu}), \quad (3)$$

$K_{\mu\nu}^\lambda$ is the contortion written as

$$K_{\mu\nu}^\lambda \equiv \frac{1}{2}T_{\mu\nu}^\lambda + T_{(\mu}^\lambda{}_{\nu)}, \quad (4)$$

with the torsion tensor $T_{\mu\nu}^\lambda$ defined as the anti-symmetric part of the affine connection, $T_{\mu\nu}^\lambda \equiv 2\Gamma_{[\mu\nu]}^\lambda$, and the disformation $L_{\mu\nu}^\lambda$ is defined by

$$L_{\mu\nu}^\lambda \equiv \frac{1}{2}Q_{\mu\nu}^\lambda - Q_{(\mu}^\lambda{}_{\nu)}. \quad (5)$$

Then let us introduce the following non-metricity conjugate,

$$P^\alpha_{\mu\nu} = -\frac{1}{4}Q^\alpha_{\mu\nu} + \frac{1}{2}Q_{(\mu}{}^\alpha{}_{\nu)} + \frac{1}{4}\left(Q^\alpha - \tilde{Q}^\alpha\right)g_{\mu\nu} - \frac{1}{4}\delta^\alpha_{(\mu}Q_{\nu)}, \quad (6)$$

and its two independent traces,

$$Q_\alpha \equiv Q^\mu{}_{\alpha\mu}, \quad \tilde{Q}_\alpha \equiv Q^\mu{}_{\alpha\mu}. \quad (7)$$

Finally, the non-metricity scalar is defined as follows

$$Q = -Q_{\alpha\mu\nu}P^{\alpha\mu\nu}. \quad (8)$$

Supplemented with Lagrange multipliers, one introduces $f(Q)$ gravity given by the following action [24]:

$$S = \int \sqrt{-g} d^4x \left[\frac{1}{2}f(Q) + \lambda_\alpha^{\beta\mu\nu}R^\alpha_{\beta\mu\nu} + \lambda_\alpha^{\mu\nu}T^\alpha_{\mu\nu} + \mathcal{L}_m \right], \quad (9)$$

where g is the determinant of the metric $g_{\mu\nu}$, $f(Q)$ is an arbitrary function of the non-metricity Q , $\lambda_\alpha^{\beta\mu\nu}$ is the Lagrange multipliers, and \mathcal{L}_m is the matter Lagrangian density,

Varying the action (9) with respect to the metric gives the field equation

$$-T_{\mu\nu} = \frac{2}{\sqrt{-g}}\nabla_\alpha(\sqrt{-g}f_Q P^\alpha_{\mu\nu}) + \frac{1}{2}g_{\mu\nu}f + f_Q(P_{\mu\alpha\beta}Q_\nu{}^{\alpha\beta} - 2Q_{\alpha\beta\mu}P^{\alpha\beta}{}_\nu), \quad (10)$$

where a subscript Q stands for a derivative of $f(Q)$ with respect to Q , $f_Q \equiv \partial_Q f(Q)$. The energy-momentum tensor is defined in the standard way,

$$T_{\mu\nu} \equiv -\frac{2}{\sqrt{-g}}\frac{\delta(\sqrt{-g}\mathcal{L}_m)}{\delta g^{\mu\nu}}. \quad (11)$$

Varying Eq. (9) with respect to the connection, one obtains

$$\nabla_\rho \lambda_\alpha^{\nu\mu\rho} + \lambda_\alpha^{\mu\nu} = \sqrt{-g}f_Q P^\alpha_{\mu\nu} + H_\alpha^{\mu\nu}, \quad (12)$$

where the hypermomentum tensor density written as

$$H_\alpha^{\mu\nu} = -\frac{1}{2}\frac{\delta\mathcal{L}_m}{\delta\Gamma^\alpha_{\mu\nu}}. \quad (13)$$

By taking into account the anti-symmetry property of μ and ν in the Lagrangian multiplier coefficients, the equation (12) can be reduced to

$$\nabla_\mu \nabla_\nu (\sqrt{-g}f_Q P^{\mu\nu}{}_\alpha + H_\alpha^{\mu\nu}) = 0. \quad (14)$$

Taking $\nabla_\mu \nabla_\nu H_\alpha^{\mu\nu} = 0$ (see discussion in Ref. [24]), we have

$$\nabla_\mu \nabla_\nu (\sqrt{-g} f_Q P^{\mu\nu}_\alpha) = 0. \quad (15)$$

The curvatureless and torsionless make the affine connection have the following form [8]

$$\Gamma^\alpha_{\mu\nu} = \left(\frac{\partial x^\alpha}{\partial \xi^\lambda} \right) \partial_\mu \partial_\nu \xi^\lambda. \quad (16)$$

We can make a special coordinate choice, the so-called coincident gauge, so that $\Gamma^\alpha_{\mu\nu} = 0$. Then, the non-metricity reduces to

$$Q_{\alpha\mu\nu} = \partial_\alpha g_{\mu\nu}, \quad (17)$$

and thereby largely simplifies the calculation since only the metric is the fundamental variable.

The price is, however, the action no longer remains diffeomorphism invariant, except for STGR [15]. One can utilize the covariant formulation of $f(Q)$ gravity to avoid the problem. Since the affine connection in Eq. (16) is purely inertial, one could utilize the covariant formulation by first determining affine connection in the absence of gravity [14]. As shown in this work, however, the off-diagonal component of the field equations in coincident gauge would put strict constraints on $f(Q)$ gravity, thereby providing us with nontrivial functional forms of $f(Q)$.

III. STATIC AND SPHERICALLY SYMMETRIC GEOMETRY IN POLAR COORDINATE

The metric ansatz for a generic static and spherically symmetric spacetime is written as

$$ds^2 = -e^{a(r)} dt^2 + e^{b(r)} dr^2 + r^2 d\Omega^2, \quad (18)$$

where $d\Omega^2 \equiv d\theta^2 + \sin^2 \theta d\phi^2$. By substituting Eq. (18) into Eq. (8), the non-metricity scalar Q is written in terms of r

$$Q(r) = -\frac{2e^{-b}}{r} \left(a' + \frac{1}{r} \right), \quad (19)$$

where a prime ($'$) denotes a derivative with respect to the radial coordinate r .

Corresponding to the spherically symmetric geometry, the energy-momentum tensor for an anisotropic fluid with spherical symmetry is given by

$$T_{\mu\nu} = (\rho + p_t) u_\mu u_\nu + p_t g_{\mu\nu} + (p_r - p_t) v_\mu v_\nu. \quad (20)$$

Here, u_μ is the four-velocity, v_μ the unitary space-like vector in the radial direction, satisfying $u^\mu u_\mu = -1$, $v^\mu v_\mu = 1$ and $u^\mu v_\mu = 0$. $\rho(r)$ is the energy density, $p_r(r)$ is the pressure in the direction of v_μ (radial pressure), and $p_t(r)$ is the pressure orthogonal to v_μ (tangential pressure).

For the anisotropic fluid (20), independent components of equations of motion (10) are listed as follows:

$$\rho = \frac{f}{2} - f_Q \left(Q + \frac{1}{r^2} + \frac{e^{-b}}{r} (a' + b') \right), \quad (21)$$

$$p_r = -\frac{f}{2} + f_Q \left(Q + \frac{1}{r^2} \right), \quad (22)$$

$$p_t = -\frac{f}{2} + f_Q \left\{ \frac{Q}{2} - e^{-b} \left[\frac{a''}{2} + \left(\frac{a'}{4} + \frac{1}{2r} \right) (a' - b') \right] \right\}, \quad (23)$$

$$0 = \frac{\cot \theta}{2} Q' f_{QQ}. \quad (24)$$

Substituting the non-metricity scalar Eq. (19) into the equations of motion, we have

$$2p'_r + a' (\rho + p_r) = f_Q \frac{e^{-b}}{r} \left[\frac{2}{r} (a' + b') - a' (a' - b') + \frac{4}{r^2} (1 - e^b) - 2a'' \right] = 0, \quad (25)$$

where in the last equality we have assumed isotropic fluid and used $p_r - p_t = 0$ in Eqs. (22) and (23). Therefore, the Tolman-Oppenheimer-Volkov (TOV) equation is satisfied in general $f(Q)$ gravity for static and spherically symmetric metric with isotropic fluids.

Now looking at the off-diagonal component in Eq. (24), one finds that the solutions to $f(Q)$ gravity is constrained to the following two cases:

$$f_{QQ} = 0 \Rightarrow f(Q) = a_0 + a_1 Q, \quad (26)$$

$$Q' = 0 \Rightarrow Q = Q_0, \quad (27)$$

where Q_0 , a_0 , and a_1 are constant. This result is similar to that in $f(T)$ gravity [25], i.e. $f_{TT} = 0$ or $T' = 0$ for the static and spherically symmetric assumption with the diagonal tetrad in $f(T)$ gravity.

The first solution in Eq. (26) is obviously reduced to STGR and is thereby equivalent to GR, with the ratio a_0/a_1 corresponding to cosmological constant Λ . Nevertheless, we shall confirm whether the Schwarzschild solution exists in linear $f(Q)$ gravity.

In the vacuum with $\rho = p_r = p_t = 0$, the equations of motion are reduced to

$$0 = a' + b', \quad (28)$$

$$Q = \frac{a_0}{a_1} - \frac{2}{r^2}, \quad (29)$$

$$0 = \frac{a_0}{2} + a_1 e^{-b} \left[\frac{a''}{2} + \left(\frac{a'}{4} + \frac{1}{2r} \right) (a' - b') \right]. \quad (30)$$

The first equation indicates that $a(r) = -b(r) + c$, where c is an integration constant and can be ignored by rescaling the time coordinate t to $e^{-c/2}t$. As a result, rr -component is the inverse of tt -component in Eq. (18). The second equation implies a cosmological constant term, $2\Lambda = a_0/a_1$. Since $Q = -R$ up to a total derivative or surface term in the action of STGR, the sign of Λ is flipped compared with the case in GR due to our convention of non-metricity scalar Eq. (8).

Using Eqs. (19), (28), and (29), we obtain

$$e^{-b} = 1 + \frac{c_1}{r} - \frac{a_0}{6a_1} r^2. \quad (31)$$

Note that, throughout the paper, we shall use c_i to denote the integration constant. Then, the line element in Eq. (18) is written as

$$ds^2 = - \left(1 + \frac{c_1}{r} - \frac{a_0}{6a_1} r^2 \right) dt^2 + \left(1 + \frac{c_1}{r} - \frac{a_0}{6a_1} r^2 \right)^{-1} dr^2 + r^2 d\Omega^2. \quad (32)$$

One recognizes that it represents the Schwarzschild (Anti-)de Sitter solution, as that in GR, with $\Lambda = a_0/(2a_1)$ and $c_1 = -2M$, where M is a mass parameter.

In the next section, we shall discuss more on the nontrivial solution that $Q = Q_0$ in Eq. (27), which will also lead to strict constraints on the functional forms. In addition, the constant non-metricity scalar Q_0 could be interpreted as cosmological constant Λ , as shown in Eq. (29).

IV. CONSTANT NON-METRICITY SCALAR: $Q = Q_0$

For constant Q , the geometric quantities are related to each other according to Eq. (19)

$$a' = -\frac{Q_0 r}{2} e^b - \frac{1}{r}, \quad (33)$$

and Eqs. (21)-(23) become

$$\rho = \frac{f(Q_0)}{2} - f_Q(Q_0) \left[Q_0 + \frac{1}{r^2} + \frac{e^{-b}}{r} (a' + b') \right], \quad (34)$$

$$p_r = -\frac{f(Q_0)}{2} + f_Q(Q_0) \left(Q_0 + \frac{1}{r^2} \right), \quad (35)$$

$$p_t = -\frac{f(Q_0)}{2} + f_Q(Q_0) \left\{ \frac{Q_0}{2} - e^{-b} \left[\frac{a''}{2} + \left(\frac{a'}{4} + \frac{1}{2r} \right) (a' - b') \right] \right\}. \quad (36)$$

With these four equations in hand, we will try to determine the remaining six quantities, i.e. $f(Q)$, $a(r)$, $b(r)$, $\rho(r)$, $p_r(r)$ and $p_t(r)$, by imposing some conditions.

A. Vacuum Solution

We first focus on the vacuum case with $\rho = p_r = p_t = 0$, then the equations reduce to

$$0 = f_Q(Q_0) \frac{e^{-b}}{r} (a' + b'), \quad (37)$$

$$0 = -\frac{f(Q_0)}{2} + f_Q(Q_0) \left(Q_0 + \frac{1}{r^2} \right), \quad (38)$$

$$0 = f_Q(Q_0) \left\{ \frac{Q_0}{2} + \frac{1}{r^2} + e^{-b} \left[\frac{a''}{2} + \left(\frac{a'}{4} + \frac{1}{2r} \right) (a' - b') \right] \right\}. \quad (39)$$

One immediately finds that, from the second equation, $f_Q(Q_0)$ and $f(Q_0)$ must be vanished. Such strict restrictions imply that a general functional form of $f(Q)$ should be $f(Q) = \sum_n a_n (Q - Q_0)^n$. Therewith one could construct specific models of cosmological interest, for instance DE models of polynomial form

$$f(Q) = Q + nQ_0 \left[1 - \left(\frac{Q}{Q_0} \right)^{1/n} \right] - Q_0, \quad (40)$$

and of exponential form

$$f(Q) = Q - nQ_0 \left\{ 1 - \exp \left[1 - \left(\frac{Q}{Q_0} \right)^{1/n} \right] \right\} - Q_0. \quad (41)$$

Other variants of the above models could be those of arctangent form

$$f(Q) = Q + nQ_0 \tanh \left[1 - \left(\frac{Q}{Q_0} \right)^{1/n} \right] - Q_0, \quad (42)$$

and of logarithmic form

$$f(Q) = Q + nQ_0 \left\{ 1 - \left[1 + \ln \left(\frac{Q}{Q_0} \right) \right]^{1/n} \right\} - Q_0. \quad (43)$$

Our models are similar to the well known DE models in $f(R)$ gravity [3], moreover, all of them satisfy the stabilization conditions, i.e., $f_Q > 0$ and $f_{QQ} > 0$. Furthermore, our polynomial model can be reduced to the models proposed in Ref. [15] with particular parameter choices except for a constant difference. And one of the advantages of our model is that it has only one undetermined parameter n due to the strong constraints on $f(Q)$. Cosmological applications of these models are out of the scope of the current work, so we shall leave it for future work.

Now that we have derived the $f(Q)$ form, we would like to analyze the spacetime structure, which is less constrained due to the triviality of Eqs. (37) and (39) provided that $f(Q_0) = f_Q(Q_0) = 0$.

1. *Case 1: $a' + b' = 0$*

Eq. (37) also allows for a solution that $a' + b' = 0$, which is of particular interest in physics and produce the exact Schwarzschild solution as known in GR. Eq. (33) solves b to

$$e^{-b(r)} = \frac{c_3}{r} - \frac{Q_0}{6}r^2, \quad (44)$$

The line element Eq. (18) is then given by

$$ds^2 = - \left(\frac{c_3}{r} - \frac{Q_0}{6}r^2 \right) dt^2 + \left(\frac{c_3}{r} - \frac{Q_0}{6}r^2 \right)^{-1} dr^2 + r^2 d\Omega^2. \quad (45)$$

It shows that there is no exact Schwarzschild solution for nontrivial $f(Q)$ functions. It is, however, a Schwarzschild (Anti-)de Sitter-like solution with $c_3 = -2M$ and $Q_0 = 2\Lambda$, and its asymptotic behavior shall be the same as Eq. (32) at $r \ll 1$ and $r \gg 1$. Note that inversely, substituting the Schwarzschild metric into Eq. (33), Q_0 cannot be constant, which guarantees the absence of the exact Schwarzschild solution.

Interestingly, Eq. (45) is similar to the Schwarzschild-like solution found by the Noether symmetry approach in $f(T)$ gravity [26], where the line element is expressed as follows

$$ds^2 = -A(r)dt^2 + \frac{1}{d_3^2 A(r)} dr^2 + r^2 d\Omega^2, \quad (46)$$

with

$$A(r) = \frac{2d_1}{3d_3}r^2 - \frac{2d_\mu}{d_3 r}. \quad (47)$$

Here, $d_\mu = d_1 d_4 - d_2 d_3$, and $d_1 \cdots d_4$ are free parameters (in Ref. [26], they are integration constants). Identifying $d_3 = 1$, $d_1 = -Q_0/4$, and $d_\mu = -c_3/2$, we can rewrite Eq. (45) as in the following form:

$$\frac{c_3}{r} - \frac{Q_0}{6} r^2 = \eta \left(1 - \frac{r_\star}{r}\right) R(r). \quad (48)$$

In the above expression we have defined

$$\eta = \left(-\frac{Q_0 c_3^2}{6}\right)^{1/3}, \quad (49)$$

$$r_\star = \left(\frac{6c_3}{Q_0}\right)^{1/3}, \quad (50)$$

$$R(r) = 1 + \frac{r}{r_\star} + \left(\frac{r}{r_\star}\right)^2. \quad (51)$$

The r_\star is a characteristic radius with the restriction $c_3 Q_0 > 0$ so that $r_\star > 0$, and the function $R(r)$ can be viewed as a distortion factor which quantifies the deviation from the exact Schwarzschild solution. By definition of $\eta > 0$, $c_3 < 0$, and thus, $Q_0 < 0$. Thus, at the scale $r \rightarrow r_\star$, Eq. (45) can mimic the Schwarzschild solution, and the $f(Q)$ gravity approximately produces a Schwarzschild spacetime in the vacuum.

Next, we shall verify whether the term inside the braces in Eq. (39) can vanish under the condition that $a' + b' = 0$. Substituting Eq. (33) leads to

$$0 = (e^{-b})'' + \frac{2(e^{-b})'}{r} + \frac{2}{r^2} + Q_0, \quad (52)$$

which solves b to

$$e^{-b(r)} = \left(c_4 + \frac{c_5}{r} - \frac{Q_0 r^2}{6} - 2 \ln r\right). \quad (53)$$

It is apparently in conflict with Eq. (44), which indicates the inconsistency between the two possibilities.

2. Case 2: $a' + b' \neq 0$

Therefore, we could assume $a' + b' \neq 0$ and make the term inside the braces in Eq. (39) vanish, which reads

$$\frac{Q_0}{2} + \frac{1}{r^2} - e^{-b} \left[\frac{a''}{2} + \left(\frac{a'}{4} + \frac{1}{2r}\right) (a' - b') \right] = 0. \quad (54)$$

Substituting Eq. (33) into the above equation gives

$$Q_0^2 r^4 + 2Q_0 r^2(2B + B'r) + 4B(B + 4 + B'r) = 0, \quad (55)$$

where we have defined $B(r) \equiv e^{-b(r)}$.

Although there is no analytic solution, we can still qualitatively analyze its asymptotic behavior. Eq. (55) gives B' in terms of B and r :

$$B' = -\frac{(Q_0 r^2 + 2B)^2 + 16B}{2r(Q_0 r^2 + 2B)}. \quad (56)$$

Let us study the asymptotic form of $B(r)$ at $r \rightarrow 0$ and $r \rightarrow \infty$ by rewriting Eq. (56) as

$$B' = -\frac{Q_0 r}{2} - \frac{B}{r} - \frac{8B}{Q_0 r^3 + 2Br}. \quad (57)$$

First, we consider the case $r \rightarrow 0$. Assuming B converges, Eq. (57) approximates $B' \sim -B/r - 4/r$. The solution is $B(r) \sim -4 + c_6/r$, but it is inconsistent with the assumption. Thus, B should diverge at $r \rightarrow 0$, and then, the second term in Eq. (57) is dominant, $B' \sim -B/r$. Then, we find the asymptotic form of $b(r)$ at $r \sim 0$:

$$e^{-b(r)} \sim \frac{c_7}{r}, \quad (58)$$

Next, we consider the case $r \rightarrow \infty$. In a way similar to the case $r \rightarrow 0$, assuming B converges to a finite value at $r \rightarrow \infty$, the first term in Eq. (57) is dominant, which gives an inconsistent result $B(r) \sim -Q_0 r^2/4 + c_8$. Thus, $B(r)$ should diverge also at $r \rightarrow \infty$. Assuming $B \sim r^n$, the case $0 < n < 2$ suggests the first term in Eq. (57) is dominant, which gives which gives $B(r) \sim -Q_0 r^2/2n$, and it is inconsistent with the assumption. Moreover, the case $n > 2$ suggests the second term is dominant, which also gives the inconsistent result $B \sim c_9/r$. Finally, one finds an consistent form $B(r) \sim B_2 r^2$, where B_0 is a parameter. In this case, Eq. (57) approximates $B' \sim -Q_0 r/2 - B/r$. Therefore, the coefficient B_2 satisfies $2B_2 = -Q_0/2 - B_2$, to find $B_2 = -Q_2/6$. Finally, we find the asymptotic form of $b(r)$ at $r \sim \infty$:

$$e^{-b(r)} \sim -\frac{Q_0}{6} r^2. \quad (59)$$

The above analysis on the asymptotic behavior suggests that the solution $e^{-b(r)}$ to Eq. (54) and Eq. (33) is similar to the Schwarzschild-like solution in Eq. (44). However, $a' + b' \neq 0$, and $a(r)$ is determined by Eq. (33). $e^{-b(r)} \sim c_{10}/r$ at $r \sim 0$, and then, Eq. (33) gives

$a' \sim -Q_0 r^2 / 2c_{10} - 1/r$. Because the second term is dominant, the asymptotic form of $a(r)$ at $r \sim 0$ is $e^{a(r)} \sim c_{11}/r$. In the same manner, $e^{-b(r)} \sim -\frac{Q_0}{6}r^2$ at $r \sim \infty$, and Eq. (33) approximates $a' \sim 2/r$. Thus, the asymptotic form of $a(r)$ at $r \sim \infty$ is $e^{a(r)} \sim c_{12}r^2$.

3. Case 3: $e^a = 1 - \frac{2M}{r}$

In previous case studies, we have confirmed the Schwarzschild-like solution and its asymptotic behavior. Presupposing that $e^a = 1 - \frac{2M}{r}$, we further discuss the deviation from the exact Schwarzschild metric, where the deviation is written only by rr -component of metric $e^{b(r)}$. Under the above assumption, Eq. (33) solves b to

$$e^b = -\frac{2}{Q_0 r (r - 2M)}, \quad (60)$$

where we require $Q_0 < 0$, and the metric reads

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \frac{-2}{Q_0 r^2} \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2. \quad (61)$$

Comparing the above with the Schwarzschild metric, one finds an extra prefactor in the rr -component. Taking the limit $r \gg 2M$, we find the leading term in rr -component is proportional to $1/r^2$, which shows the absence of a conventional Newtonian limit.

B. Non-vacuum solutions

We have seen that the vacuum solution for $Q = Q_0$ is closely related to the solutions that $f(Q_0) = f_Q(Q_0) = 0$, which provide us with nontrivial functional forms of $f(Q)$. In this subsection, however, we shall focus on the non-vacuum case by imposing conditions on geometric quantities. In particular, We shall assume that $f_Q(Q_0) \neq 0$ to look for particular solutions of interest.

1. *Case 1: $a' + b' = 0$*

The solutions for $a' + b' = 0$ are the same as Eqs. (44) and (45), while the equations of motion are constrained to

$$-\rho = p_r = p_t + f_Q(Q_0) \frac{1}{r^2}, \quad (62)$$

$$p_t = -\frac{f(Q_0)}{2} + Q_0 f_Q(Q_0). \quad (63)$$

Note that the energy density and radial pressure obey the inverse-square law with respect to the radius if $f_Q(Q_0) \neq 0$, although the tangential pressure is constant. And since p_t is a constant in terms of Q_0 , we rewrite it as p_{t0} , and obtain a particular solution to $f(Q)$ is

$$f(Q) = c_{13} Q_0 \sqrt{\frac{Q}{Q_0}} - 2p_{t0}. \quad (64)$$

2. *Case 2: $a' = 0$*

When a is a constant, Eq. (33) solves $b(r)$ to

$$e^{-b} = -\frac{Q_0}{2} r^2. \quad (65)$$

Then the metric reads

$$ds^2 = -dt^2 - \frac{2}{Q_0 r^2} dr^2 + r^2 d\Omega^2, \quad (66)$$

where we have absorbed a into time coordinate. The equations of motion are

$$\rho = -p_r - Q_0 f_Q(Q_0), \quad (67)$$

$$p_r = p_t + f_Q(Q_0) \frac{1}{r^2}, \quad (68)$$

$$p_t = -\frac{f(Q_0)}{2} + Q_0 f_Q(Q_0). \quad (69)$$

The values of pressures are the same as case 1, so does the particular solution to $f(Q)$.

3. *Case 3: $b = 0$*

For $b = 0$, Eq. (33) gives

$$e^a = \frac{c_{14}}{r} \exp\left(-\frac{Q_0}{4} r^2\right), \quad (70)$$

with the metric being

$$ds^2 = -\frac{c_{14}}{r} \exp\left(-\frac{Q_0}{4}r^2\right) dt^2 + dr^2 + r^2 d\Omega^2. \quad (71)$$

Then the equations of motion read

$$\rho = \frac{f(Q_0)}{2} - \frac{Q_0 f_Q(Q_0)}{2}, \quad (72)$$

$$p_r = -\rho + f_Q(Q_0) \left(\frac{Q_0}{2} + \frac{1}{r^2}\right), \quad (73)$$

$$p_t = -\rho - f_Q(Q_0) \left(\frac{Q_0}{4} + \frac{Q_0^2 r^2}{16} + \frac{1}{4r^2}\right). \quad (74)$$

Since ρ is a constant in terms of Q_0 , we rewrite it as ρ_0 . Then a particular solution to $f(Q)$ is

$$f(Q) = c_{15}Q + 2\rho_0, \quad (75)$$

which recovers STGR with cosmological constant $\Lambda = \rho_0/c_{15}$.

V. CONCLUSIONS AND DISCUSSION

The choice of different geometric bases allows us to have different descriptions of gravity. Of particular concern in recent years is the STGR, in which gravity is attributed to the non-metricity tensor. As an extension of STGR, $f(Q)$ gravity is an intriguing approach in the study of modified gravity. In this work, we have analyzed the static and spherically symmetric solutions in general $f(Q)$ gravity and shown the two possible solutions due to the non-zero off-diagonal components, that is, linear $f(Q)$ function and constant non-metricity scalar $Q = Q_0$. We confirmed that the Schwarzschild (Anti-)de Sitter solution only exists in linear $f(Q)$ gravity, which recovers STGR and is thereby equivalent to GR.

With a focus on the second solution $Q = Q_0$, we have found that the vacuum solution, especially Eq. (38), is closely related to $f(Q_0) = f_Q(Q_0) = 0$, which puts stringent constraints on the functional form of $f(Q)$ gravity. On the other hand, these solutions make the geometric quantities less constrained and thereby provide us with considerable freedom to investigate the geometric structure. We, therefore, could construct some DE models with one parameter. Examining such models would be intriguing and inspiring in cosmology. As examples, we discuss three cases trying to find the exact Schwarzschild solution, asymptotic

behavior, and deviation. As for the non-vacuum solution for $Q = Q_0$, we start from the geometric assumptions listed as three examples. By proper conditions on tt -component or rr -component, we find some particular solutions to $f(Q)$ with constant energy density or constant tangent pressure.

In the end, we comment on the gauge fixing and possible analogy to the known results in $f(T)$ gravity. The curvatureless and torsionless conditions correspond to the coincident gauge so that the affine connection is always zero. However, the choice of the coincident gauge makes the $f(Q)$ theory no longer invariant under the general coordinate transformation. As in the case of $f(T)$ gravity, the covariant formulation of $f(Q)$ theory [14, 27, 28] can allow us to avoid issues caused by the coincident gauge. In our present analysis, the existence of the constraint equation (24) makes the $f(Q)$ gravity have two choices; a linear function or constant non-metricity scalar. Concerning the gauge fixing and breakdown of diffeomorphism, the different coordinate systems may allow $f(Q)$ gravity to possess the Schwarzschild solution even in the non-covariant approach to $f(Q)$ gravity. As in the $f(T)$ theory, an off-diagonal component of field equations shows up in the polar coordinate, which prohibits the Schwarzschild spacetime as a solution; otherwise, $f(T)$ function should be a linear to the torsion scalar T . On the other hand, $f(T)$ gravity with higher orders of T can possess the Schwarzschild solution in the isotropic coordinate [23]. By relying on the analogy between $f(T)$ and $f(Q)$ theories, it may be worth studying the spherically symmetric solution of $f(Q)$ gravity in the isotropic coordinate. Applying our present analysis to the covariant $f(Q)$ gravity or other coordinate systems, we may be able to obtain the Schwarzschild solution even in the nontrivial $f(Q)$ function.

ACKNOWLEDGMENTS

The authors thank Taotao Qiu for valuable comments on the scientific content. T.K. is grateful to Shin'ichi Nojiri for his advice on the calculations.

[1] A. G. Riess et al. (Supernova Search Team), *Astron. J.* **116**, 1009 (1998), [arXiv:astro-ph/9805201 \[astro-ph\]](#).

- [2] N. Aghanim *et al.* (Planck), *Astron. Astrophys.* **641**, A6 (2020), [Erratum: *Astron.Astrophys.* 652, C4 (2021)], [arXiv:1807.06209 \[astro-ph.CO\]](#).
- [3] A. De Felice and S. Tsujikawa, *Living Rev. Rel.* **13**, 3 (2010), [arXiv:1002.4928 \[gr-qc\]](#).
- [4] S. Nojiri, S. D. Odintsov, and V. K. Oikonomou, *Phys. Rept.* **692**, 1 (2017), [arXiv:1705.11098 \[gr-qc\]](#).
- [5] K. Hayashi and T. Shirafuji, *Phys. Rev. D* **19**, 3524 (1979), [Addendum: *Phys.Rev.D* 24, 3312–3314 (1982)].
- [6] J. W. Maluf, *Annalen Phys.* **525**, 339 (2013), [arXiv:1303.3897 \[gr-qc\]](#).
- [7] M. Adak, O. Sert, M. Kalay, and M. Sari, *Int. J. Mod. Phys. A* **28**, 1350167 (2013), [arXiv:0810.2388 \[gr-qc\]](#).
- [8] J. Beltrán Jiménez, L. Heisenberg, and T. Koivisto, *Phys. Rev. D* **98**, 044048 (2018), [arXiv:1710.03116 \[gr-qc\]](#).
- [9] Y.-F. Cai, S. Capozziello, M. De Laurentis, and E. N. Saridakis, *Rept. Prog. Phys.* **79**, 106901 (2016), [arXiv:1511.07586 \[gr-qc\]](#).
- [10] L. Heisenberg, *Phys. Rept.* **796**, 1 (2019), [arXiv:1807.01725 \[gr-qc\]](#).
- [11] B. Li, T. P. Sotiriou, and J. D. Barrow, *Phys. Rev. D* **83**, 064035 (2011), [arXiv:1010.1041 \[gr-qc\]](#).
- [12] T. P. Sotiriou, B. Li, and J. D. Barrow, *Phys. Rev. D* **83**, 104030 (2011), [arXiv:1012.4039 \[gr-qc\]](#).
- [13] N. Tamanini and C. G. Boehmer, *Phys. Rev. D* **86**, 044009 (2012), [arXiv:1204.4593 \[gr-qc\]](#).
- [14] D. Zhao, (2021), [arXiv:2104.02483 \[gr-qc\]](#).
- [15] J. Beltrán Jiménez, L. Heisenberg, T. S. Koivisto, and S. Pekar, *Phys. Rev. D* **101**, 103507 (2020), [arXiv:1906.10027 \[gr-qc\]](#).
- [16] F. K. Anagnostopoulos, S. Basilakos, and E. N. Saridakis, *Phys. Lett. B* **822**, 136634 (2021), [arXiv:2104.15123 \[gr-qc\]](#).
- [17] B. J. Barros, T. Barreiro, T. Koivisto, and N. J. Nunes, *Phys. Dark Univ.* **30**, 100616 (2020), [arXiv:2004.07867 \[gr-qc\]](#).
- [18] K. F. Dialektopoulos, T. S. Koivisto, and S. Capozziello, *Eur. Phys. J. C* **79**, 606 (2019), [arXiv:1905.09019 \[gr-qc\]](#).
- [19] F. Bajardi, D. Vernieri, and S. Capozziello, *Eur. Phys. J. Plus* **135**, 912 (2020), [arXiv:2011.01248 \[gr-qc\]](#).

- [20] Z. Hassan, S. Mandal, and P. K. Sahoo, *Fortsch. Phys.* **69**, 2100023 (2021), [arXiv:2102.00915 \[gr-qc\]](#).
- [21] S. Mandal, P. K. Sahoo, and J. R. L. Santos, *Phys. Rev. D* **102**, 024057 (2020), [arXiv:2008.01563 \[gr-qc\]](#).
- [22] K. Flathmann and M. Hohmann, *Phys. Rev. D* **103**, 044030 (2021), [arXiv:2012.12875 \[gr-qc\]](#).
- [23] R. Ferraro and F. Fiorini, *Phys. Rev. D* **84**, 083518 (2011), [arXiv:1109.4209 \[gr-qc\]](#).
- [24] J. Beltrán Jiménez, L. Heisenberg, and T. S. Koivisto, *JCAP* **08**, 039 (2018), [arXiv:1803.10185 \[gr-qc\]](#).
- [25] C. G. Boehmer, A. Mussa, and N. Tamanini, *Class. Quant. Grav.* **28**, 245020 (2011), [arXiv:1107.4455 \[gr-qc\]](#).
- [26] A. Paliathanasis, S. Basilakos, E. N. Saridakis, S. Capozziello, K. Atazadeh, F. Darabi, and M. Tsamparlis, *Phys. Rev. D* **89**, 104042 (2014), [arXiv:1402.5935 \[gr-qc\]](#).
- [27] R.-H. Lin and X.-H. Zhai, *Phys. Rev. D* **103**, 124001 (2021), [arXiv:2105.01484 \[gr-qc\]](#).
- [28] F. D'Ambrosio, S. D. B. Fell, L. Heisenberg, and S. Kuhn, (2021), [arXiv:2109.03174 \[gr-qc\]](#).