

# Integrability and solvability of polynomial Liénard differential systems

Maria V. Demina

*HSE University, 34 Tallinskaya Street, 123458, Moscow, Russian Federation, maria\_dem@mail.ru*

---

## Abstract

We provide the necessary and sufficient conditions of Liouvillian integrability for Liénard differential systems describing nonlinear oscillators with a polynomial damping and a polynomial restoring force. We prove that Liénard differential systems are not Darboux integrable excluding subfamilies with certain restrictions on the degrees of the polynomials arising in the systems. We demonstrate that if the degree of a polynomial responsible for the restoring force is greater than the degree of a polynomial producing the damping, then a generic Liénard differential system is not Liouvillian integrable with the exception of linear Liénard systems. However, for any fixed degrees of the polynomials describing the damping and the restoring force we present subfamilies possessing Liouvillian first integrals. As a by-product of our results, we find a number of novel Liouvillian integrable subfamilies. In addition, we study the existence of non-autonomous Darboux first integrals and non-autonomous Jacobi last multipliers with a time-dependent exponential factor.

*Keywords:* Liénard differential systems, Darboux integrability, Liouvillian integrability, invariant algebraic curves, Puiseux series

---

## 1. Introduction

Performing classifications of integrable or solvable subfamilies for a given multi-parameter system of ordinary differential equations is a very difficult problem. The aim of the present article is to solve the integrability problem for the following systems of first-order ordinary differential equations

$$x_t = y, \quad y_t = -f(x)y - g(x). \quad (1.1)$$

We suppose that  $f(x)$  and  $g(x)$  are polynomials

$$f(x) = f_0x^m + \dots + f_m, \quad g(x) = g_0x^n + \dots + g_n, \quad f_0g_0 \neq 0 \quad (1.2)$$

with coefficients from the field  $\mathbb{C}$ . Systems (1.1) are named in honor of the French physicist and engineer Alfred–Marie Liénard [33]. These systems describe oscillators with a polynomial damping  $f(x)$  and a polynomial restoring force  $g(x)$ . In addition, Liénard differential systems have a variety of other applications in physics, chemistry, biology, economics, etc.

For example, these systems (1.1) arise as traveling wave reductions of the following general families of reaction-convection-diffusion equations

$$u_t = Du_{xx} + A(u)u_x + B(u), \quad u = u(x, t), \quad (1.3)$$

where  $D$  is a diffusion coefficient,  $A(u)$  describes a nonlinear convective flux and  $B(u)$  is responsible for a reaction force.

Let the variable  $y$  be privileged with respect to the variable  $x$ , then the function  $y(x)$  satisfies the following family of Abel differential equations of the second kind

$$yy_x + f(x)y + g(x) = 0. \quad (1.4)$$

The integrability properties of Liénard differential systems are investigated by various methods and in the framework of various theories. Let us enumerate the most important studies:

1. local analysis [8, 11, 28, 30, 36];
2. classical Lie symmetry analysis [37, 38] and  $\lambda$  symmetries [40];
3. local [6, 7, 10, 27, 29] and non-local transformations [2, 23, 24, 31, 41, 42];
4. differential Galois theory [1];
5. extended Prolle–Singer method [4] and the Darboux theory of integrability [9, 16, 18, 19, 23–25, 35, 44].

A collection of integrable and solvable subfamilies of Liénard differential systems is presented by A. D. Polyanin and V. F. Zaitsev in [39]. The transformation  $y(x) = 1/w(x)$  brings Abel differential equations of the second kind (1.4) to Abel differential equations of the first kind

$$w_x = g(x)w^3 + f(x)w^2. \quad (1.5)$$

Consequently, certain results available for such Abel differential equations can be transferred to equations (1.4) and related Liénard differential systems [6, 7, 10]. Let us note that many scientific works dealing with the global integrability problem present sufficient conditions of integrability. Thus, these works do not provide classifications of integrable Liénard differential systems and Abel differential equations. Meanwhile, it is an important scientific problem to find all integrable families for fixed degrees of the polynomials  $f(x)$  and  $g(x)$ .

This article is devoted to the study of the general integrability properties of polynomial Liénard differential systems. We focus on the necessary and sufficient conditions of Darboux and Liouvillian integrability. J. Llibre and C. Valls [35] proved that Liénard differential systems (1.1) under the condition  $\deg g \leq \deg f$  do not have Liouvillian first integrals excluding the trivial case  $g(x) = \alpha f(x)$ ,  $\alpha \in \mathbb{C}$ . Consequently, we only need to study systems (1.1) satisfying the restriction  $\deg g > \deg f$ . Throughout this article it is supposed

that  $f(x) \not\equiv 0$ . Any Liénard differential system is Hamiltonian with a polynomial first integral

$$I(x, y) = y^2 + 2 \int_0^x g(s) ds \quad (1.6)$$

whenever the relation  $f(x) \equiv 0$  is valid.

We shall prove that Liénard differential systems satisfying the restrictions  $\deg g > \deg f$  and  $\deg g \neq 2 \deg f + 1$  are not Darboux integrable, while there are Liouvillian integrable subfamilies. In contrast, Liénard differential systems in the case  $\deg g = 2 \deg f + 1$  exhibit a variety of rational, Darboux, and Liouvillian first integrals existing under certain restrictions on the parameters. This fact is also recognized by many scientists [4, 10, 29, 31, 37, 38, 40]. Our main tools include the modern Darboux theory of integrability [13, 43], the method of Puiseux series [17, 20], and the local theory of invariants [22]. We do not impose any non-trivial restrictions on the coefficients of the polynomials  $f(x)$  and  $g(x)$  with the exception of Liénard differential systems from the families  $\deg g = 2 \deg f + 1$ . We mainly study systems that are non-resonant near infinity provided that the following restriction  $\deg g = 2 \deg f + 1$  is valid. To be more precise, we say that a system (1.1) is resonant near infinity whenever the highest-degree coefficients  $f_0$  and  $g_0$  satisfy a resonance condition. This condition is explicitly given in Section 4 and arises only in the case  $\deg g = 2 \deg f + 1$ . Let us note that the subset of resonant systems is of zero Lebesgue measure in the set of all polynomial systems (1.1) with fixed degrees of the polynomials  $f(x)$  and  $g(x)$ . Our results are also valid in the resonant case, but they are not complete. For all other polynomial Liénard differential systems we present a complete classification of Liouvillian integrable subfamilies. In addition, we classify polynomial Liénard differential systems possessing non-autonomous Darboux first integrals and non-autonomous Jacobi last multipliers with a time-dependent exponential factor.

We demonstrate that the integrability properties of Liénard differential systems are substantially different in the following three cases:

- (A)  $\deg f < \deg g < 2 \deg f + 1$ ;
- (B)  $\deg g = 2 \deg f + 1$ ;
- (C)  $\deg g > 2 \deg f + 1$ .

This article is organized as follows. Sections 2, 3, and 4 contain a review of the known results and several preliminary observations on the methods we use in the subsequent part. In Section 2, we describe the Darboux theory of integrability and consider some related questions. Section 3 is devoted to the method of Puiseux series and to the local theory of invariants. In Section 4, the results on invariant algebraic curves of Liénard differential systems are described. In Section 5, we present some integrability properties valid for a generic polynomial Liénard differential system. In Sections 6, 7, 8, we investigate the integrability and solvability of Liénard differential systems from families (A), (B), and (C), respectively. In Section 9, we consider an example: we study the Liénard differential systems satisfying the restrictions  $\deg f = 2$  and  $\deg g = 4$ .

## 2. The Darboux theory of integrability

The main aim of the present section is to describe some basic aspects of the Darboux theory of integrability. We focus on the problem of finding Darboux and Liouvillian first integrals of polynomial differential systems in the plane

$$x_t = P(x, y), \quad y_t = Q(x, y), \quad (2.1)$$

where  $P(x, y)$  and  $Q(x, y)$  are relatively prime elements of the ring  $\mathbb{C}[x, y]$ . By  $\mathbb{C}[x, y]$  we denote the ring of bivariate polynomials with complex-valued coefficients. The vector field related to system (2.1) is defined as

$$\mathcal{X} = P(x, y) \frac{\partial}{\partial x} + Q(x, y) \frac{\partial}{\partial y}. \quad (2.2)$$

**Definition 2.1.** A non-constant function  $I(x, y): D \subset \mathbb{C}^2 \rightarrow \mathbb{C}$  is called a first integral of differential system (2.1) and the related vector field  $\mathcal{X}$  on an open subset  $D \subset \mathbb{C}^2$  if  $I(x(t), y(t)) = C$  with  $C$  being a constant for all values of  $t$  such that the solution  $(x(t), y(t))$  of system (2.1) is defined in  $D$ .

If  $I(x, y)$  is of a class at least  $C^1$  in  $D$ , then  $I(x, y)$  is a first integral of differential system (2.1) if and only if  $\mathcal{X}I = 0$ .

**Definition 2.2.** A non-constant function  $M(x, y): D \subset \mathbb{C}^2 \rightarrow \mathbb{C}$  is called an integrating factor of differential system (2.1) and the related vector field  $\mathcal{X}$  in an open subset  $D \subset \mathbb{C}^2$  if the differential form  $M(x, y)(P(x, y)dy - Q(x, y)dx)$  is exact in  $D$ . In other words, there exists a function  $I(x, y)$  of a class at least  $C^1$  in  $D$  such that the following relation is valid  $M(x, y)(P(x, y)dy - Q(x, y)dx) = dI(x, y)$ .

If an integrating factor  $M(x, y)$  is of a class at least  $C^1$  in  $D$ , then it satisfies the following linear first-order partial differential equation  $\mathcal{X}M = -\text{div } \mathcal{X}M$ , where  $\text{div } \mathcal{X} = P_x + Q_y$  is the divergence of the vector field  $\mathcal{X}$ .

A function  $I(x, y)$  is referred to as a Liouvillian function of two variables  $x$  and  $y$  if it belongs to a Liouvillian extension of the field of rational functions  $\mathbb{C}(x, y)$  over  $\mathbb{C}$ . Generally speaking, any Liouvillian function can be represented as a finite superposition of algebraic functions, antiderivatives, and exponentials [43]. A function  $\Phi(x, y)$  is called a Darboux function of two variables  $x$  and  $y$ , if it can be presented in the form

$$\Phi(x, y) = F_1^{d_1}(x, y) \dots F_K^{d_K}(x, y) \exp\{R(x, y)\}, \quad (2.3)$$

where  $F_1(x, y) \in \mathbb{C}[x, y]$ ,  $\dots$ ,  $F_K(x, y) \in \mathbb{C}[x, y]$ ,  $R(x, y) \in \mathbb{C}(x, y)$ ,  $d_1, \dots, d_K \in \mathbb{C}$ . We see that any Darboux function is a Liouvillian function. The converse is not generally true.

A differential system (2.1) is called Darboux (Liouvillian) integrable if it possesses a Darboux (Liouvillian) first integral. It is known that the problem of establishing Darboux or Liouvillian integrability of a differential system (2.1) can be reduced to the problem of constructing all irreducible invariant algebraic curves of (2.1) and all exponential invariants of (2.1), for more details see [13, 43, 45].

**Definition 2.3.** *The curve  $F(x, y) = 0$  with  $F(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}$  is an invariant algebraic curve of a differential system (2.1) whenever the following condition  $F_t|_{F=0} = (PF_x + QF_y)|_{F=0} = 0$  is valid.*

If the polynomial  $F(x, y)$  producing the invariant algebraic curve  $F(x, y) = 0$  is irreducible in  $\mathbb{C}[x, y]$ , then the ideal generated by  $F(x, y)$  is radical. Consequently, there exists an element  $\lambda(x, y)$  of the ring  $\mathbb{C}[x, y]$  such that  $F(x, y)$  satisfies the partial differential equation  $P(x, y)F_x + Q(x, y)F_y = \lambda(x, y)F$ . The polynomial  $\lambda(x, y)$  is called *the cofactor* of the invariant algebraic curve  $F(x, y) = 0$ . It is straightforward to show that the degree of  $\lambda(x, y)$  is at most  $L - 1$ , where  $L$  is the maximum between the degrees of the polynomials  $P(x, y)$  and  $Q(x, y)$ . We conclude that an invariant algebraic curve of differential system (2.1) is formed from solutions of the latter. A solution of differential system (2.1) has either empty intersection with the zero set of  $F(x, y)$  or it is entirely contained in  $F(x, y) = 0$ . The generating polynomial  $F(x, y)$  of an invariant algebraic curve  $F(x, y) = 0$  is referred to as a *Darboux polynomial* or *an algebraic invariant*.

**Definition 2.4.** *A function  $E(x, y) = \exp[g(x, y)/f(x, y)]$  with the relatively prime polynomials  $g(x, y), f(x, y) \in \mathbb{C}[x, y]$  is called an exponential invariant of a differential system (2.1) whenever the following condition  $\mathcal{X}E = \varrho(x, y)E$  is valid, where  $\varrho(x, y) \in \mathbb{C}[x, y]$ .*

The polynomial  $\varrho(x, y)$  is referred to as *the cofactor* of the exponential invariant  $E(x, y)$ . It is straightforward to show that the product of the exponential invariants  $E_1(x, y)$  and  $E_2(x, y)$  with the cofactors  $\varrho_1(x, y)$  and  $\varrho_2(x, y)$ , respectively, is an exponential invariant possessing the cofactor  $\varrho(x, y) = \varrho_1(x, y) + \varrho_2(x, y)$ . It is known that the polynomial  $f(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}$  arising in an exponential invariant  $E(x, y) = \exp[g(x, y)/f(x, y)]$  produces an invariant algebraic curve  $f(x, y) = 0$  of the system under consideration [13].

It turns out that the study of autonomous first integrals and autonomous integrating factors is sometimes restrictive, even if an autonomous differential system is under consideration. In this article we do not consider the Darboux theory of integrability for non-autonomous systems in the general case, for more details see [3, 15, 34].

A non-autonomous first integral  $I(x, y, t)$  and a non-autonomous integrating factor  $M(x, y, t)$  of a differential system (2.1) are defined similarly to the autonomous case. They satisfy the following linear partial differential equations  $I_t + \mathcal{X}I = 0$  and  $M_t + \mathcal{X}M = -\text{div } \mathcal{X}M$ , respectively, whenever  $I(x, y, t)$  and  $M(x, y, t)$  are function of a class at least  $C^1$  in  $D \subset \mathbb{C}^3$ . Non-autonomous integrating factors are commonly referred to as Jacobi last multipliers or simply Jacobi multipliers.

The following theorems are the essence of the modern Darboux theory of integrability.

**Theorem 2.1.** *A polynomial differential system (2.1) is Darboux integrable if and only if it has a rational integrating factor.*

The fact that a Darboux integrable differential system (2.1) has a rational integrating factor was derived by J. Chavarriga et al., see [5]. The converse statement was established by C. Christopher et al., see [12].

**Theorem 2.2.** *A polynomial differential system (2.1) is Liouvillian integrable if and only if it has a Darboux integrating factor.*

Theorem 2.2 was proved by M. F. Singer [43].

**Theorem 2.3.** *A polynomial differential system (2.1) possesses a first integral of the form*

$$I(x, y, t) = \prod_{j=1}^K F_j^{d_j}(x, y) \exp \left\{ \frac{S(x, y)}{R(x, y)} \right\} \exp(\omega t), \quad \omega, d_1, \dots, d_K \in \mathbb{C}, \quad (2.4)$$

where  $F_1(x, y), \dots, F_K(x, y)$  are pairwise relatively prime irreducible bivariate polynomials from the ring  $\mathbb{C}[x, y]$ ,  $S(x, y)$  and  $R(x, y)$  are relatively prime bivariate polynomials from the ring  $\mathbb{C}[x, y]$ , if and only if  $F_1(x, y) = 0, \dots, F_K(x, y) = 0, R(x, y) = 0$  are invariant algebraic curves of the system and  $E(x, y) = \exp\{S(x, y)/R(x, y)\}$  is an exponential invariant of the system such that the following condition

$$\sum_{j=1}^N d_j \lambda_j(x, y) + \varrho(x, y) + \omega = 0 \quad (2.5)$$

holds identically. In this expression  $\lambda_j(x, y)$  is the cofactor of the invariant algebraic curve  $F_j(x, y) = 0$  and  $\varrho(x, y)$  is the cofactor of the exponential invariant  $E(x, y)$ .

Theorem 2.3 follows from the classical theory of Darboux integrability, see [45]. We name a first integral of Theorem 2.3 as a non-autonomous Darboux first integral provided that  $\omega \neq 0$ . If  $\omega = 0$ , then function (2.4) gives a Darboux first integral.

**Theorem 2.4.** *Under the assumptions of Theorem 2.3 a polynomial differential system (2.1) possesses a Jacobi last multiplier of the form*

$$M(x, y, t) = \prod_{j=1}^K F_j^{d_j}(x, y) \exp \left\{ \frac{S(x, y)}{R(x, y)} \right\} \exp(\omega t), \quad \omega, d_1, \dots, d_K \in \mathbb{C}, \quad (2.6)$$

if and only if  $F_1(x, y) = 0, \dots, F_K(x, y) = 0, R(x, y) = 0$  are invariant algebraic curves of the system and  $E(x, y) = \exp\{S(x, y)/R(x, y)\}$  is an exponential invariant of the system such that the following condition

$$\sum_{j=1}^N d_j \lambda_j(x, y) + \varrho(x, y) + \omega = -\operatorname{div} \mathcal{X} \quad (2.7)$$

is identically valid. In this expression  $\lambda_j(x, y)$  is the cofactor of the invariant algebraic curve  $F_j(x, y) = 0$  and  $\varrho(x, y)$  is the cofactor of the exponential invariant  $E(x, y)$ .

Theorem 2.4 with the restriction  $\omega = 0$  was derived by C. Christopher [13]. The case  $\omega \neq 0$  was considered in article [15]. A Jacobi last multiplier of Theorem 2.4 will be referred to as a non-autonomous Darboux–Jacobi last multiplier whenever  $\omega \neq 0$ .

These theorems suggest the following algorithm for searching autonomous and non-autonomous Darboux first integrals and Jacobi last multipliers:

1. find all relatively prime irreducible invariant algebraic curves and all exponential invariants with linearly independent cofactors;
2. find, or prove the non-existence of, complex numbers  $d_1, \dots, d_K, \omega$  such that condition (2.5) or (2.7) is identically satisfied; the polynomial  $\varrho(x, y)$  arising in conditions (2.5) and (2.7) equals the sum of the cofactors of exponential invariants found at the first step.

Let us note that there exist certain estimates of the number of pairwise distinct invariants which guarantees the existence of rational, Darboux or Liouvillian first integrals in the autonomous case. For more details see books [32, 45] and the references therein.

The first step of this algorithm is extremely difficult. This is due to the absence of *a priori* upper bounds on the degrees of bivariate polynomials giving irreducible invariant algebraic curves. It is shown in articles [14, 17, 20] that the method of Puiseux series, which is described in the next section, can facilitate the first step.

It is straightforward to see that integrating factors and Jacobi last multipliers are defined modulo to the multiplication by a non-zero constant. Two integrating factors or Jacobi last multipliers producing a constant ratio are supposed to be equivalent. We do not distinguish them. The uniqueness of integrating factors and Jacobi last multipliers is understood exactly in this sense.

In general, the existence of only one independent non-autonomous first integral is not sufficient for the complete integrability of the system under study in the framework of the Darboux theory. However, the knowledge of a non-autonomous Darboux first integral can be used to derive the general solutions. Several examples are given in article [19]. Moreover, we demonstrate that some Liénard differential systems from family (B) simultaneously have independent autonomous and non-autonomous Darboux first integrals. Eliminating the variable  $y = x_t$  from these first integrals, one can find explicit expressions of the general solutions; see also article [40], where some similar examples are presented. To conclude this section let us note that trajectories lying in a zero set of inverse Jacobi last multipliers are of importance in the qualitative theory of differential systems [45]. Consequently, the classification of Liénard differential systems possessing Darboux–Jacobi last multipliers seem to be a significant problem.

### 3. The method of Puiseux series and the local theory of invariants

We start with a brief review of the theory of fractional-power (or Puiseux) series. A Puiseux series around a point  $x_0 \in \mathbb{C}$  can be presented as

$$y(x) = \sum_{l=0}^{+\infty} c_l (x - x_0)^{\frac{l_0+l}{n_0}}, \quad c_0 \neq 0, \quad (3.1)$$

where  $l_0 \in \mathbb{Z}$  and  $n_0 \in \mathbb{N}$ . It is without loss of generality to suppose that the number  $n_0$  is relatively prime with the greatest common divisor of the numbers  $\{l_0 + l : c_l \neq 0\}$ ,

$l \in \mathbb{N} \cup \{0\}$ . While a Puiseux series around the point  $x = \infty$  takes the form

$$y(x) = \sum_{l=0}^{+\infty} b_l x^{\frac{l_0-l}{n_0}}, \quad b_0 \neq 0, \quad (3.2)$$

where  $l_0 \in \mathbb{Z}$  and  $n_0 \in \mathbb{N}$ . Again we assume that the number  $n_0$  is relatively prime with the greatest common divisor of the numbers  $\{l_0 - l : b_l \neq 0, l \in \mathbb{N} \cup \{0\}\}$ .

Let us consider the algebraic equation  $F(x, y) = 0$ , where  $F(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}[x]$ . Giving preference to one of the variables with respect to another, a solution  $y$  of this equation viewed as a function of  $x$  can be locally expanded into a convergent Puiseux series. This statement is known as the Newton–Puiseux theorem.

The set of all Puiseux series given by (3.1) or (3.2) forms an algebraically closed field, which we denote by  $\mathbb{C}_{x_0}\{x\}$  or  $\mathbb{C}_\infty\{x\}$ , respectively. The ring of polynomials in one variable  $y$  with coefficients from the fields  $\mathbb{C}_{x_0}\{x\}$  or  $\mathbb{C}_\infty\{x\}$  is denoted as  $\mathbb{C}_{x_0}\{x\}[y]$  or  $\mathbb{C}_\infty\{x\}[y]$ , respectively.

Suppose  $S(x, y)$  is an element of the ring  $\mathbb{C}_\infty\{x\}[y]$ . Let us introduce two operators of projection acting on this ring. The first operator  $\{S(x, y)\}_+$  gives the sum of the monomials of  $S(x, y)$  with non-negative integer powers. In other words,  $\{S(x, y)\}_+$  yields the polynomial part of  $S(x, y)$ . Analogously, the projection  $\{S(x, y)\}_- = S(x, y) - \{S(x, y)\}_+$  produces the non-polynomial part of  $S(x, y)$ . It is straightforward to show that these projections are linear operators. In addition, we see that the action of the projection operators can be extended to the ring of the Puiseux series in  $y$  near the point  $y = \infty$  with coefficients from the field  $\mathbb{C}_\infty\{x\}$ . We denote this ring as  $\mathbb{C}_\infty\{x\}[\{y\}]$ . Thus, we get the relation  $\{S(x, y)\}_+ \in \mathbb{C}[x, y]$ , where  $S(x, y) \in \mathbb{C}_\infty\{x\}[\{y\}]$ . We endow the fields  $\mathbb{C}_{x_0}\{x\}$  and  $\mathbb{C}_\infty\{x\}$  with the differential operator  $\partial_x$ . Analogously, we endow the rings  $\mathbb{C}_{x_0}\{x\}[y]$  and  $\mathbb{C}_\infty\{x\}[y]$  with the differential operators  $\partial_x$  and  $\partial_y$ . The action of these differential operators is standard.

The following basic statements are proved in articles [16, 17, 20].

**Lemma 3.1.** *Let  $y(x)$  be a Puiseux series from one of the fields  $\mathbb{C}_{x_0}\{x\}$  or  $\mathbb{C}_\infty\{x\}$ . Suppose that the series  $y(x)$  satisfies the equation  $F(x, y(x)) = 0$ , where  $F(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}[x]$  gives an invariant algebraic curve  $F(x, y) = 0$  of a system (2.1). Then the series  $y(x)$  solves the equation*

$$P(x, y)y_x - Q(x, y) = 0. \quad (3.3)$$

Using algebraic closeness of the field of Puiseux series  $\mathbb{C}_\infty\{x\}$ , it is straightforward to find the general structure of irreducible invariant algebraic curves and their cofactors.

**Theorem 3.1.** *Let  $F(x, y) = 0$ , where  $F(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}[x]$ , be an irreducible invariant algebraic curve of a differential system (2.1). Then the polynomial  $F(x, y)$  and the cofactor  $\lambda(x, y) \in \mathbb{C}[x, y]$  of the curve  $F(x, y) = 0$  take the form*

$$F(x, y) = \left\{ \mu(x) \prod_{j=1}^N \{y - y_{j,\infty}(x)\} \right\}_+, \quad (3.4)$$

$$\lambda(x, y) = \left\{ \sum_{m=0}^{+\infty} \sum_{j=1}^N \frac{\{Q(x, y) - P(x, y)(y_{j, \infty})_x\} (y_{j, \infty})^m}{y^{m+1}} + P(x, y) \sum_{m=0}^{+\infty} \sum_{l=1}^L \frac{\nu_l x_l^m}{x^{m+1}} \right\}_+, \quad (3.5)$$

where  $y_{1, \infty}(x), \dots, y_{N, \infty}(x)$  are pairwise distinct Puiseux series in a neighborhood of the point  $x = \infty$  that satisfy equation (3.3),  $x_1, \dots, x_L$  are pairwise distinct zeros of the polynomial  $\mu(x) \in \mathbb{C}[x]$  with multiplicities  $\nu_1, \dots, \nu_L \in \mathbb{N}$  and  $L \in \mathbb{N} \cup \{0\}$ . Moreover, the degree of  $F(x, y)$  with respect to  $y$  does not exceed the number of distinct Puiseux series of the form (3.2) satisfying equation (3.3) whenever this number is finite. If  $\mu(x) = \mu_0$ , where  $\mu_0 \in \mathbb{C}$ , then we suppose that  $L = 0$  and the first series is absent in the expression of the cofactor  $\lambda(x, y)$ .

This theorem introduces an algebraic tool enabling one to find invariant algebraic curves explicitly, for more details see article [21].

The method of Puiseux series can be also used whenever one wishes to find exponential invariants with non-polynomial arguments. The following theorem giving necessary conditions for an exponential invariant related to an invariant algebraic curve to exist is proved in article [16].

**Theorem 3.2.** *Let the polynomial  $f(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}[x]$  give an invariant algebraic curve  $f(x, y) = 0$  of a differential system (2.1). The cofactor of the invariant algebraic curve  $f(x, y) = 0$  we denote by  $\lambda(x, y) \in \mathbb{C}[x, y]$ . Suppose that this system admits an exponential invariant  $E = \exp(g/f)$  related to the algebraic curve  $f(x, y) = 0$ , then for each non-constant Puiseux series  $y_{j, \infty}(x)$  in a neighborhood of the point  $x = \infty$  that satisfies the equation  $f(x, y) = 0$  there exists a number  $q \in \mathbb{Q}$  such that the Puiseux series for the function  $\lambda(x, y_{j, \infty}(x))/P(x, y_{j, \infty}(x))$  in a neighborhood of the point  $x = \infty$  is*

$$\frac{\lambda(x, y_{j, \infty}(x))}{P(x, y_{j, \infty}(x))} = \sum_{k=n}^{+\infty} b_k x^{-\frac{k}{n}}, \quad b_n = q. \quad (3.6)$$

Now our aim is to introduce a notion of local invariant curves that are given by elements from one of the rings  $\mathbb{C}_{x_0}\{x\}[y]$  and  $\mathbb{C}_\infty\{x\}[y]$ . Let  $x_0$  be a point of the extended complex plane  $\overline{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ . Elements of the ring  $\mathbb{C}_{x_0}\{x\}[y]$  are referred to as Puiseux polynomials. Note that the polynomials  $P(x, y)$  and  $Q(x, y)$  can be regarded as Puiseux polynomials for any  $x_0 \in \overline{\mathbb{C}}$ . Again giving preference to the variable  $y$ , we shall deal with formal Puiseux series solutions  $y = y_{x_0}(x)$  of algebraic first-order ordinary differential equation (3.3). We can do the same analysis choosing the variable  $x$  as depended.

**Definition 3.1.** *The curve  $F(x, y) = 0$ , where  $F(x, y) \in \mathbb{C}_{x_0}\{x\}[y]$ , is called a local invariant curve of a differential system (2.1) whenever the following condition  $F_t|_{F=0} = (PF_x + QF_y)|_{F=0} = 0$  is valid.*

If the element  $F(x, y) \in \mathbb{C}_{x_0}\{x\}[y]$  gives a local invariant curve  $F(x, y) = 0$  of differential system (2.1), then there exists the Puiseux polynomial  $\lambda(x, y) \in \mathbb{C}_{x_0}\{x\}[y]$  such that the following equation  $\mathcal{X}F(x, y) = \lambda(x, y)F(x, y)$  is satisfied [22]. By analogy with the algebraic

case, the Puiseux polynomials  $F(x, y)$  and  $\lambda(x, y)$  will be called a local invariant and the cofactor of the related local invariant and local curve, respectively. The theory of local invariants is introduced in article [22], where all the statements given below are proved.

In what follows we call local invariants and associated local invariant curves given by the polynomials  $F(x, y) = h_{x_0}(x)$  and  $F(x, y) = y - y_{x_0}(x)$ ,  $h_{x_0}(x), y_{x_0}(x) \in \mathbb{C}_{x_0}\{x\}$  elementary. Any local invariant  $F(x, y)$  can be represented as a product of elementary local invariants and the cofactor of the local invariant  $F(x, y)$  is the sum of the cofactors of all the factors.

**Theorem 3.3.** *Let  $y_{x_0}(x)$  be a Puiseux series near the point  $x = x_0$ . The element  $F(x, y) = y - y_{x_0}(x)$  of the ring  $\mathbb{C}_{x_0}\{x\}[y]$  gives a local invariant curve of differential system (2.1) if and only if the series  $y = y_{x_0}(x)$  satisfies equation (3.3).*

It is straightforward to obtain an explicit expression of the cofactor similar to that presented in Theorem 3.1. We introduce the operator of projection  $\{S(x, y)\}_{+, y}$  yielding the polynomial part with respect to  $y$  of the element  $S(x, y) \in \mathbb{C}_{x_0, \infty}\{x\}[\{y\}]$ . The symbol  $\mathbb{C}_{x_0, \infty}\{x\}[\{y\}]$  denotes the ring of Puiseux series in  $y$  near the point  $y = \infty$  with coefficients from the field  $\mathbb{C}_{x_0}\{x\}$ . In other words, we obtain the relation  $\{S(x, y)\}_{+, y} \in \mathbb{C}_{x_0}\{x\}[y]$ .

**Theorem 3.4.** *Let  $F(x, y) = y - Y_{x_0}(x)$ ,  $Y_{x_0}(x) \in \mathbb{C}_{x_0}\{x\}$  give an elementary local invariant curve  $F(x, y) = 0$  of differential system (2.1), then its cofactor takes the form*

$$\lambda(x, y) = \left\{ \sum_{m=0}^{+\infty} \frac{\{Q(x, y) - P(x, y)(Y_{x_0}(x))_x\} Y_{x_0}^m(x)}{y^{m+1}} \right\}_{+, y}. \quad (3.7)$$

Similarly to the case of local invariant curves, we can introduce a concept of local exponential invariants.

**Definition 3.2.** *A function  $E(x, y) = \exp[g(x, y)/f(x, y)]$  with relatively prime elements  $g(x, y)$  and  $f(x, y)$  of the ring  $\mathbb{C}_{x_0}\{x\}[y]$  is called a local exponential invariant of a differential system (2.1) if  $E(x, y)$  satisfies the partial differential equation  $\mathcal{X}E(x, y) = \varrho(x, y)E(x, y)$ , where  $\varrho(x, y)$  is a Puiseux polynomial.*

Again the Puiseux polynomial  $\varrho(x, y)$  will be referred to as the cofactor of the local exponential invariant  $E(x, y)$ . Note that exponential invariants belong to a differential extension of the field of fractions of the ring  $\mathbb{C}_{x_0}\{x\}[y]$ .

**Lemma 3.2.** *Let  $E(x, y) = \exp[g(x, y)/f(x, y)]$ ,  $f(x, y) \in \mathbb{C}_{x_0}\{x\}[y] \setminus \mathbb{C}_{x_0}\{x\}$  be a local exponential invariant of a differential system (2.1), then  $f(x, y) = 0$  is a local invariant curve of the system in question.*

We shall say that local exponential invariants

$$\begin{aligned} E(x, y) &= \exp [g_l(x)y^l], \quad g_l(x) \in \mathbb{C}_{x_0}\{x\}, \quad l \in \mathbb{N} \cup \{0\}; \\ E(x, y) &= \exp \left[ \frac{u(x)}{\{y - y_{x_0}(x)\}^n} \right], \quad y_{x_0}(x), u(x) \in \mathbb{C}_{x_0}\{x\}, \quad n \in \mathbb{N} \end{aligned} \quad (3.8)$$

are *elementary local exponential invariants*. Since  $E^\alpha(x, y)$  with  $\alpha \in \mathbb{C}$  is a local exponential invariant whenever so does  $E(x, y)$ , the highest-order coefficients of the series  $g_l(x)$  and  $u(x)$  can be fixed in advance. Any local exponential invariant  $E(x, y)$  is as a product of elementary local exponential invariants and the cofactor of the local invariant  $E(x, y)$  is the sum of the cofactors of all the factors.

We conclude that if system (2.1) has a global invariant (algebraic or exponential), then for any  $x_0 \in \overline{\mathbb{C}}$  this invariant must be a product of local elementary invariants in such a way that the sum of their cofactors is a true, not Puiseux, polynomial. This observation is used in subsequent sections.

#### 4. Invariant algebraic curves of polynomial Liénard differential systems

With the aim to study the integrability properties of polynomial Liénard differential systems we need information on the existence of their invariant algebraic curves. The general structure of bivariate polynomials giving irreducible invariant algebraic curves is derived in articles [20, 21].

**Theorem 4.1.** *Let  $F(x, y) = 0$ ,  $F(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}$  be an irreducible invariant algebraic curve of a Liénard differential system (1.1) from family (A). Then the polynomial  $F(x, y)$  and the cofactor  $\lambda(x, y)$  of the invariant algebraic curve  $F(x, y) = 0$  take the form*

$$F(x, y) = \left\{ \prod_{j=1}^{N-k} \left\{ y - y_{j,\infty}^{(1)}(x) \right\} \left\{ y - y_{N,\infty}^{(2)}(x) \right\}^k \right\}_+, \quad (4.1)$$

$$\lambda(x, y) = -Nf - (N - k)q_x - kp_x, \quad (4.2)$$

where  $k = 0$  or  $k = 1$ ,  $N \in \mathbb{N}$ , and  $y_{1,\infty}^{(1)}(x), \dots, y_{N-1,\infty}^{(1)}(x), y_{N,\infty}^{(2)}(x)$  are the series

$$(I) : y_{j,\infty}^{(1)}(x) = q(x) + \sum_{l=0}^{+\infty} b_{m+1+l}^{(j)} x^{-l}, \quad j = 1, \dots, N - k; \quad (4.3)$$

$$(II) : y_{N,\infty}^{(2)}(x) = p(x) + \sum_{l=0}^{+\infty} b_{n-m+l}^{(N)} x^{-l}.$$

The coefficients of the series of type (II) and of the polynomials

$$q(x) = -\frac{f_0}{m+1} x^{m+1} + \sum_{l=1}^m q_{m+1-l} x^l \in \mathbb{C}[x], \quad (4.4)$$

$$p(x) = -\frac{g_0}{f_0} x^{n-m} + \sum_{l=1}^{n-m-1} p_{n-m-l} x^l \in \mathbb{C}[x]$$

are uniquely determined. The coefficients  $b_{m+1}^{(j)}$ ,  $j = 1, \dots, N - k$  are pairwise distinct. All other coefficients  $b_{m+1+l}^{(j)}$ ,  $l \in \mathbb{N}$  are expressible via  $b_{m+1}^{(j)}$ , where  $j = 1, \dots, N - k$ . The corresponding product in (4.1) is unit whenever  $k = 1$  and  $N = 1$ .

**Theorem 4.2.** *A Liénard differential system (1.1) from family (A) with fixed coefficients of the polynomials  $f(x)$  and  $g(x)$  has at most two distinct irreducible invariant algebraic curves simultaneously. If the two distinct irreducible invariant algebraic curves exist, then the first has  $k = 1$  in representation (4.1) and the second is given by a first-degree polynomial with respect to  $y$  and takes the form  $y - q(x) - z_0 = 0$ .*

*Corollary.* Liénard differential systems (1.1) from family (A) are not integrable with a rational first integral.

Theorems 4.1 and 4.2 are proved in article [20].

The structure of polynomials producing invariant algebraic curves for Liénard differential systems from family (B) is in strong correlation with the properties of the following quadratic equation

$$p^2 - \varrho p + (m + 1)\varrho = 0, \quad (4.5)$$

where we have introduced notation

$$\varrho = 4(m + 1) - \frac{f_0^2}{g_0}. \quad (4.6)$$

The set of all positive rational numbers will be denoted as  $\mathbb{Q}^+$ . Let  $p_1$  and  $p_2$  be the roots of equation (4.5).

**Theorem 4.3.** *Suppose  $F(x, y) = 0$ ,  $F(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}$  is an irreducible invariant algebraic curve of a Liénard differential system (1.1) from family (B) and equation (4.5) has no solutions in  $\mathbb{Q}^+$ . One of the following statements holds.*

1. *If  $p_1 p_2 \neq 0$ , then the polynomial  $F(x, y)$  is of degree at most two with respect to  $y$  and*

$$\begin{aligned} F(x, y) &= \left\{ \left\{ y - y_\infty^{(1)}(x) \right\}^{s_1} \left\{ y - y_\infty^{(2)}(x) \right\}^{s_2} \right\}_+, \\ \lambda(x, y) &= -(s_1 + s_2)f(x) - \left\{ s_1 \left( y_\infty^{(1)} \right)_x + s_2 \left( y_\infty^{(2)} \right)_x \right\}_+, \\ y_{+\infty}^{(k)}(x) &= \sum_{l=0}^{\infty} b_l^{(k)} x^{m+1-l}, \quad b_0^{(k)} = \frac{f_0}{p_k - 2(m + 1)}, \quad k = 1, 2, \end{aligned} \quad (4.7)$$

where  $s_1$  and  $s_2$  are either 0 or 1 independently,  $s_1 + s_2 > 0$ . The Puiseux series  $y_\infty^{(k)}(x)$ ,  $k = 1, 2$  are Laurent series and possess uniquely determined coefficients.

2. *If  $p_1 p_2 = 0$ , then  $p_1 = p_2 = 0$ . The polynomial  $F(x, y)$  and its cofactor  $\lambda(x, y)$  take the form*

$$\begin{aligned} F(x, y) &= y + \frac{f_0}{2(m + 1)} x^{m+1} - \sum_{l=1}^{m+1} b_l x^{m+1-l}, \\ \lambda(x, y) &= -f(x) + \frac{f_0}{2} x^m - \sum_{l=1}^m (m + 1 - l) b_l x^{m-l}, \end{aligned} \quad (4.8)$$

where the coefficients  $b_1, \dots, b_{m+1}$  are uniquely determined. In addition, the following relation  $4(m + 1)g_0 - f_0^2 = 0$  is valid.

This theorem is proved in [21]. Note that the Liénard differential systems such that the related equation (4.5) has a positive rational solution are also studied in [21]. We do not reproduce the related results here.

The structure of polynomials giving invariant algebraic curves and their cofactors for Liénard differential systems satisfying the condition  $\deg g > 2 \deg f + 1$  is derived in [21]. The following theorem is valid.

**Theorem 4.4.** *Let  $F(x, y) = 0$ ,  $F(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}$  be an irreducible invariant algebraic curve of a Liénard differential system (1.1) from family (C). Then the polynomial  $F(x, y)$  and the cofactor  $\lambda(x, y)$  of the invariant algebraic curve  $F(x, y) = 0$  take the form*

$$F(x, y) = \left\{ \prod_{j=1}^{N_1} \left\{ y - y_{j,\infty}^{(1)}(x) \right\} \prod_{j=1}^{N_2} \left\{ y - y_{j,\infty}^{(2)}(x) \right\} \right\}_+, \quad (4.9)$$

$$\lambda(x, y) = -(N_1 + N_2)f - \left\{ N_1 h_x^{(1)} + N_2 h_x^{(2)} \right\}_+ \quad (4.10)$$

where the Puiseux series  $y_{j,\infty}^{(1,2)}(x)$  are given by the relations

$$y_{j,\infty}^{(1,2)}(x) = h^{(1,2)}(x) + \sum_{k=2(n+1)}^{+\infty} b_{k,j}^{(1,2)} x^{\frac{n+1}{2} - \frac{k}{2}}, \quad h^{(1,2)}(x) = \sum_{k=0}^{2n+1} b_k^{(1,2)} x^{\frac{n+1}{2} - \frac{k}{2}}, \quad (4.11)$$

$N_1, N_2 \in \mathbb{N} \cup \{0\}$ , and  $N_1 + N_2 \geq 1$ . The coefficients  $b_{2(n+1),j}^{(1,2)}$  with the same upper index are pairwise distinct and all the coefficients  $b_{l,j}^{(1,2)}$  with  $l > 2(n+1)$  are expressible via  $b_{2(n+1),j}^{(1,2)}$ . If  $n$  is an odd number, then the corresponding Puiseux series are Laurent series and  $b_{2l-1}^{(1,2)} = 0$ ,  $b_{2l-1,j}^{(1,2)} = 0$ , when  $l \in \mathbb{N}$ . In addition,  $N_k = 1$  whenever  $n$  is odd and  $N_l = 0$ , where  $k, l = 1, 2$  and  $k \neq l$ . If  $n$  is an even number, then  $N_1 = N_2$ .

All the Puiseux series arising in Theorems 4.1, 4.3, 4.4 solve the equation (1.4) related to the Liénard differential system under consideration. The following theorem provides the necessary and sufficient conditions for a Liénard differential system (1.1) to have an invariant algebraic curve.

**Theorem 4.5.** *The polynomial  $F(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}$  of degree  $N > 0$  with respect to  $y$  gives an invariant algebraic curve  $F(x, y) = 0$  of a Liénard differential system (1.1) if and only if there exists  $N$  Puiseux series  $y_{1,\infty}(x), \dots, y_{N,\infty}(x)$  from the field  $\mathbb{C}_\infty\{x\}$  that solve equation (1.4) and satisfy the conditions*

$$\left\{ \sum_{j=1}^N y_{j,\infty}(x) \right\}_- = 0. \quad (4.12)$$

The theorems presented in this section essentially are based on the method of Puiseux series.

## 5. Integrability of a generic Liénard differential system

We begin this section by investigating the existence of exponential invariants with polynomial arguments.

**Lemma 5.1.** *If a Liénard differential system (1.1) satisfying the condition  $\deg g > \deg f$  has an exponential invariant of the form  $E(x, y) = \exp[h(x, y)]$ ,  $h(x, y) \in \mathbb{C}[x, y]$  with the cofactor  $\varrho(x, y) \in \mathbb{C}[x, y]$  of degree at most  $\deg g - 1$ , then the polynomial  $\varrho(x, y)$  is divisible by  $y$  in the ring  $\mathbb{C}[x, y]$ .*

*Proof.* It is straightforward to show that the polynomial  $h(x, y)$  satisfies the following linear inhomogeneous partial differential equation

$$yh_x - \{f(x)y + g(x)\}h_y = \varrho(x, y). \quad (5.1)$$

Let us represent the functions  $h(x, y)$  and  $\varrho(x, y)$  as polynomials in  $y$  with polynomial in  $x$  coefficients. Substituting the expressions  $h(x, y) = h_0(x) + h_1(x)y + \dots$  and  $\varrho(x, y) = \varrho_0(x) + \varrho_1(x)y + \dots$  into equation (5.1) and selecting the coefficients of  $y^0$ , we get the relation  $g(x)h_1(x) = -\varrho_0(x)$ . Since the degree of the polynomial  $\varrho_0(x)$  is at most  $\deg g - 1$ , we conclude that  $h_1(x) \equiv 0$  and  $\varrho_0(x) \equiv 0$ . As a result the polynomial  $\varrho(x, y)$  is given by the relation  $\varrho(x, y) = (\varrho_1(x) + \dots)y$ . This completes the proof.  $\square$

As a direct consequence of this lemma, we establish that Darboux or Liouvillian integrable Liénard differential systems necessarily have invariant algebraic curves.

**Theorem 5.1.** *If a Liénard differential system (1.1) satisfying the condition  $\deg g > \deg f$  has no invariant algebraic curves, then this system is not integrable with a Darboux or Liouvillian first integral.*

*Proof.* It is a simple result of the Darboux theory of integrability that a differential system (2.1) without invariant algebraic curves cannot have rational first integrals.

Suppose that a Liénard differential system without invariant algebraic curves possesses a Darboux first integral. Then this first integral is given by an exponential invariant with a zero cofactor. Consequently, the argument of the exponential function in the invariant provides a rational first integral. This is a contradiction.

If a Liénard differential system (1.1) without invariant algebraic curves is Liouvillian integrable, then there exists a Darboux integrating factor and this integrating factor equals some exponential invariant of the form  $E(x, y) = \exp[h(x, y)]$ ,  $h(x, y) \in \mathbb{C}[x, y]$ . Calculating the divergence of the vector field

$$\mathcal{X} = y\frac{\partial}{\partial x} - (f(x)y + g(x))\frac{\partial}{\partial y}, \quad (5.2)$$

we get:  $\operatorname{div} \mathcal{X} = -f(x)$ . The divergence is independent of  $y$ , while by Lemma 5.1 the cofactor of the exponential invariant  $E(x, y) = \exp[h(x, y)]$  is divisible by  $y$  provided that  $\deg g > \deg f$ . Consequently, condition (2.5) in the autonomous case is not valid.  $\square$

Thus, we conclude that Liouvillian integrable Liénard differential systems (1.1) with the restriction  $\deg f < \deg g$  must have at least one invariant algebraic curve.

**Theorem 5.2.** *A generic Liénard differential system (1.1) with fixed degrees of the polynomials  $f(x)$  and  $g(x)$  is neither Darboux nor Liouvillian integrable provided that the following restrictions  $\deg g > \deg f$  and  $(\deg f, \deg g) \neq (0, 1)$  are valid.*

*Proof.* Let us denote the set of all Liénard differential systems with fixed degrees of the polynomials  $f(x)$  and  $g(x)$  as  $L_{m,n}$ . Any particular Liénard differential system can be identified with a point in  $\mathbb{C}^{m+n} \times (\mathbb{C} \setminus \{0\})^2$ . In view of Theorem 5.1, we need to prove that the subset of Liénard differential systems (1.1) without invariant algebraic curves is of full Lebesgue measure in the set  $L_{m,n}$ . Note that here we speak about finite invariant algebraic curves. Extending systems (1.1) from the complex plane  $\mathbb{C}^2$  to the complex projective plane  $\mathbb{C}P^2$ , we see that the infinite line is always invariant for Liénard differential systems.

We begin by considering systems from family (A). The subset of Liénard differential systems such that the related equations (1.4) possess a family of formal Puiseux series solutions  $y_\infty^{(1)}(x)$  has Lebesgue measure zero. Indeed, there always arises a compatibility condition enabling this family of series to exist. We only need to show that the compatibility condition cannot be identically satisfied. For this aim we track the appearance of the coefficient  $g_{n-m}$  in the series  $y_\infty^{(1)}(x)$ . We use the following representation

$$y_\infty^{(1)}(x) = \sum_{l=0}^{+\infty} v_l(x) (g_{n-m})^l, \quad (5.3)$$

where  $\{v_l(x)\}$  are elements of the field  $\mathbb{C}_\infty\{x\}$ . The compatibility condition arises, when one tries to find the coefficient of  $x^0$ . Substituting representation (5.3) into equations (1.4) and setting to zero the coefficients of  $g_{n-m}^l$  for  $l = 0, 1$ , we find the ordinary differential equations

$$v_0 v_{0,x} + f(x)v_0 + \hat{g}(x) = 0, \quad v_0 v_{1,x} + (f(x) + v_{0,x})v_1 + x^m = 0, \quad (5.4)$$

where we use the designation  $\hat{g}(x) = g(x) - g_{n-m}x^m$ . The dominant behavior of the series  $v_0(x)$  near the point  $x = \infty$  is

$$v_0(x) = -\frac{f_0}{m+1}x^{m+1} + o(x^{m+1}), \quad x \rightarrow \infty. \quad (5.5)$$

Analyzing the ordinary differential equation for the series  $v_1(x)$ , we see that this equation should have a solution with the dominant behavior  $v_1(x) = e_0x^0$ ,  $e_0 \in \mathbb{C} \setminus \{0\}$ ,  $x \rightarrow \infty$ , but it does not. In addition, the equations for  $v_l(x)$ ,  $l \geq 2$  have a zero solution whenever  $v_1(x)$  is zero. Thus, the compatibility condition enabling the series  $y_\infty^{(1)}(x)$  to exist is given by a first-degree polynomial with respect to  $g_{n-m}$  and cannot hold identically. Consequently, it is necessary to consider systems with invariant algebraic curves given by bivariate polynomials not involving the series  $y_\infty^{(1)}(x)$  into the factorization in the ring  $\mathbb{C}_\infty\{x\}[y]$ . In view of Theorem 4.1, the corresponding irreducible invariant algebraic curve is given by the polynomial  $F(x, y) = y - p(x) - z_1$ , where  $z_1 \in \mathbb{C}$  and the polynomial  $p(x)$  of degree  $n - m$  is

described by expression (4.4). Now we consider the subset of Liénard differential systems with the invariant algebraic curve  $y - p(x) - z_1 = 0$ . Since the related equation (1.4) possesses the polynomial solution  $y(x) = p(x) + z_1$ , we find the polynomial  $g(x)$ . The result is  $g(x) = -(p + z_1)(p_x + f)$ . We see that the dimension of the subset of Liénard differential systems under consideration is less than the dimension of  $L_{m,n}$ .

We turn to systems from family (B). It follows from Theorem 4.3 that if a generic Liénard differential system from family (B) possesses irreducible invariant algebraic curves, then their generating polynomials are of degrees either 1 or 2 with respect to  $y$ . Let us suppose that there exists the irreducible invariant algebraic curve  $y - q_l(x) = 0$ . In this expression  $q_l(x) = \{y_\infty^{(l)}(x)\}_+$ ,  $l = 1, 2$  is a polynomial of degree  $m + 1$ . By analogue with systems from family (A), we can find the polynomial  $g(x)$ . We see from the relation  $g(x) = -q_l(f + q_{l,x})$  that the subset of Liénard differential systems with the invariant algebraic curve  $y - q_l(x) = 0$  is of dimension  $2m + 3$ , while the dimension of  $L_{m,n}$  is  $m + n + 2 = 3(m + 1)$ . Hence the subset in question is of zero Lebesgue measure in  $L_{m,n}$  provided that  $m > 0$ . Now we suppose that Liénard differential systems from family (B) have the hyperelliptic invariant algebraic curve  $(y + u(x))^2 + w(x) = 0$ . By Theorem 4.3 this curve is unique. In addition, we see that the polynomial  $u(x)$  is of degree  $m + 1$  and the polynomial  $w(x)$  is of degree at most  $2m + 2$ . Following H. Żołądec [46], we substitute the bivariate polynomial  $F(x, y) = (y + u(x))^2 + w(x)$  into the partial differential equation

$$yF_x - (f(x)y + g(x))F_y - \lambda(x, y)F = 0. \quad (5.6)$$

Further, we set to zero the coefficients at different powers of  $y$ . Since the cofactor  $\lambda(x, y)$  is independent of  $y$ , we express the polynomials  $f(x)$  and  $g(x)$  via  $u(x)$  and  $w(x)$ . The result is

$$f(x) = u_x + \frac{uw_x}{2w}, \quad g(x) = \frac{w_x}{2} + \frac{u^2w_x}{2w}. \quad (5.7)$$

We see from these expressions that any zero of the polynomial  $w(x)$  is also a zero of the polynomial  $u(x)$ . The polynomial  $u(x)$  is parameterized by at most  $m + 2$  parameters and the polynomial  $w(x)$  adds only one new parameter. Thus, we conclude that the dimension of Liénard differential systems from family (B) with the hyperelliptic invariant algebraic curve  $(y + u(x))^2 + w(x) = 0$  is at most  $m + 3$ . While the dimension of  $L_{m,2m+1}$  is  $3m + 3$ .

Finally, we are left with systems from family (C). Let us track the dependence of the Puiseux series  $y_\infty^{(1)}(x)$  and  $y_\infty^{(2)}(x)$  on the coefficient  $f_m$ . We introduce the representation

$$y_\infty^{(k)}(x) = \sum_{l=0}^{+\infty} v_l^{(k)}(x) (f_m)^l, \quad k = 1, 2, \quad (5.8)$$

where  $\{v_l^{(k)}(x)\}$  are elements of the field  $\mathbb{C}_\infty\{x\}$ . Let us introduce the following designation:  $\hat{f}(x) = f(x) - f_m$ . Substituting these representations into equation (1.4) and setting to zero the coefficients of  $f_m^l$  with  $l = 0, 1$ , we find the ordinary differential equations

$$v_0 v_{0,x} + \hat{f}(x)v_0 + g(x) = 0, \quad v_0 v_{1,x} + (\hat{f}(x) + v_{0,x})v_1 + v_0 = 0. \quad (5.9)$$

Note that we omit the upper index. Let us suppose that  $n$  is even. Puiseux series from the field  $\mathbb{C}_\infty\{x\}$  that satisfy these equations are of the form

$$\begin{aligned} v_0^{(k)}(x) &= \sum_{j=0}^{+\infty} c_j^{(k)} x^{\frac{n+1-k}{2}}, \quad k = 1, 2, \quad c_0^{(1,2)} = \pm \frac{\sqrt{-2(n+1)g_0}}{n+1}; \\ v_1^{(k)}(x) &= \sum_{j=0}^{+\infty} e_j^{(k)} x^{\frac{2-k}{2}}, \quad k = 1, 2, \quad e_0^{(1,2)} = -\frac{2}{n+1}. \end{aligned} \tag{5.10}$$

Theorem 4.5 gives the following necessary condition for an invariant algebraic curve  $F(x, y) = 0$  to exist:  $N_1 b_{n+3}^{(1)} + N_2 b_{n+3}^{(2)} = 0$ . Recall that  $b_{n+3}^{(k)}$  is a coefficient of the family of series  $y_\infty^{(k)}(x)$  and  $N_k$  is the number of times the family of series  $y_\infty^{(k)}(x)$  enters the factorization of  $F(x, y)$  in the ring  $\mathbb{C}_\infty\{x\}[y]$ . Using Theorem 4.4, we get  $N_1 = N_2$ . Calculating the first five coefficients of the series  $v_1^{(k)}(x)$ , we see that  $e_4^{(1)} + e_4^{(2)}$  is not identically zero for any non-zero polynomial  $f(x)$  of degree  $m < (n-1)/2$ . Hence the expression  $N_1(b_{n+3}^{(1)} + b_{n+3}^{(2)})$  is a polynomial with respect to  $f_m$  possessing a non-zero coefficient of  $f_m$  in the generic case. Now we assume that  $n$  is odd. The Puiseux series  $v_0^{(k)}(x)$  and  $v_1^{(k)}(x)$  solving equations (5.9) are of the form (5.10), where the coefficients  $\{c_j^{(k)}\}$  and  $\{e_j^{(k)}\}$  with odd lower indices equal zero. Further, we again consider the necessary condition  $N_1 b_{n+3}^{(1)} + N_2 b_{n+3}^{(2)} = 0$ , but now the numbers  $N_1$  and  $N_2$  may not be equal. Calculating the first three non-trivial coefficients of the series  $v_1^{(k)}(x)$ , we see that the condition  $N_1 e_4^{(1)} + N_2 e_4^{(2)}$  is not identically zero for any numbers  $N_1, N_2 \geq 0, N_1 + N_2 > 0$  and any non-zero polynomial  $f(x)$  of degree  $m < (n-1)/2$ . Consequently, the expression  $N_1 b_{n+3}^{(1)} + N_2 b_{n+3}^{(2)}$ , regarded as a polynomial with respect to  $f_m$ , possesses a non-zero coefficient of  $f_m$  in the generic case. We conclude that a generic Liénard differential system from family (C) has no finite invariant algebraic curves. □

It is straightforward to see that if  $(\deg f, \deg g) = (0, 1)$ , then the associated Liénard differential systems are linear. They always have invariant lines and are Darboux integrable.

## 6. Integrability of Liénard differential systems from family (A)

We have proved in the previous section that if a polynomial Liénard differential system is Liouvillian integrable, then it has at least one invariant algebraic curve. Let us investigate the existence of exponential invariants with non-polynomial arguments. Here and in what follows we use the designations of Theorem 4.1. In particular, the polynomials  $q(x)$  and  $p(x)$  give the initial parts of the Puiseux series near the point  $x = \infty$  that solve equation (1.4) whenever  $\deg f < \deg g < 2 \deg f + 1$ . These polynomials are presented in expression (4.4).

**Lemma 6.1.** *Liénard differential systems (1.1) from family (A) do not have exponential invariants of the form  $E(x, y) = \exp\{h(x, y)/r(x, y)\}$ , where  $h(x, y) \in \mathbb{C}[x, y]$  and  $r(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}$  are coprime polynomials.*

*Proof.* The proof is by contradiction. Let  $E(x, y) = \exp \{h(x, y)/r(x, y)\}$  be an exponential invariant of a Liénard differential system (1.1) from family (A). Since the polynomial  $r(x, y)$  is not a constant, we see that  $r(x, y) = 0$  is an invariant algebraic curve of the differential system under consideration [13]. According to the results of Theorem 4.1 we can represent the polynomial  $r(x, y)$  in the form  $r(x, y) = F_1^{n_1}(x, y)F_2^{n_2}(x, y)$ , where  $n_1, n_2 \in \mathbb{N}_0$ ,  $n_1 + n_2 > 0$ , and the polynomials  $F_1(x, y)$  and  $F_2(x, y)$  produce invariant algebraic curves of the related differential system. These polynomials are given by expression (4.1) with  $N = 1$ ,  $k = 0$  (the invariant algebraic curve  $F_1(x, y) = 0$ ) and by expression (4.1) with  $N \in \mathbb{N}$ ,  $k = 1$  (the invariant algebraic curve  $F_2(x, y) = 0$ ). Relation (4.2) yields the explicit expressions of the cofactors

$$\begin{aligned}\lambda_1(x, y) &= -f(x) - q_x(x) = o(x^m), \quad x \rightarrow \infty, \\ \lambda_2(x, y) &= -Nf(x) - (N-1)q_x(x) - p_x(x) = -f_0x^m + o(x^m), \quad x \rightarrow \infty.\end{aligned}\tag{6.1}$$

related to these invariant algebraic curves, respectively. The cofactor of the invariant algebraic curve  $r(x, y) = 0$  equals  $\lambda(x, y) = n_1\lambda_1(x, y) + n_2\lambda_2(x, y)$ .

If the Liénard differential system under consideration possesses only one irreducible invariant algebraic curve, for example  $F_1(x, y) = 0$ , then we assume that  $n_2 = 0$  and vice versa.

Here and in what follows,  $O$  denotes the zero element of the field  $\mathbb{C}_\infty\{x\}$ . Let us suppose that  $y_\infty^{(2)}(x)$  is a Puiseux series of type (II) such that the following condition  $F_2(x, y_\infty^{(2)}(x)) = O$  is valid. Substituting  $y = y_\infty^{(2)}(x)$  into the function  $\lambda(x, y)/y$ , we can expand it into a Puiseux series near infinity. Supposing that  $n_2 > 0$ , we find the dominant behavior near the point  $x = \infty$  of this series. The result is

$$\frac{\lambda(x, y_\infty^{(2)}(x))}{y_\infty^{(2)}(x)} = \frac{n_2 f_0^2}{g_0} x^{2m-n} + o(x^{2m-n}), \quad x \rightarrow \infty.\tag{6.2}$$

The inequality  $\deg f < \deg g < 2 \deg f + 1$  yields  $2m - n > -1$ . Using Theorem 3.2, we come to a contradiction. Consequently, we should set  $n_2 = 0$ . The polynomial  $r(x, y)$  now takes the form  $r(x, y) = F_1^{n_1}(x, y)$  with  $F_1(x, y) = y - q(x) - z_0$  and  $z_0 \in \mathbb{C}$ .

Equation  $\mathcal{X}E = \varrho(x, y)E$  produces the following linear inhomogeneous partial differential equation

$$yh_x - (g(x)y + f(x))h_y = n_1\lambda_1(x, y)h + \varrho(x, y)F_1^{n_1}(x, y), \quad \varrho(x, y) \in \mathbb{C}[x, y]\tag{6.3}$$

satisfied by the polynomial  $h(x, y)$ . Let us consider the truncated Puiseux series  $y(x) = q(x) + z_0$  that solves equation (1.4) whenever invariant algebraic curve  $F_1(x, y) = 0$  exists. Recall that  $y(x)$  is a zero of the polynomial  $F_1(x, y)$ . Considering the restriction  $H(x) = h(x, y)|_{y=y(x)}$ , we obtain the ordinary differential equation

$$(q(x) + z_0) \frac{dH}{dx} = n_1\lambda_1(x, y)H, \quad \lambda_1(x, y) = -f(x) - q_x(x).\tag{6.4}$$

Since the polynomials  $h(x, y)$  and  $F_1(x, y)$  are relatively prime, we conclude that  $H(x) \not\equiv 0$ . Indeed, assuming the converse and using the Bézout's theorem, we see that the bivariate polynomials  $h(x, y)$  and  $r(x, y)$  have a common factor.

It follows from the relations  $\deg q = m + 1$  and  $\deg \lambda_1 \leq m - 1$  that equation (6.4) does not have non-zero polynomial solutions. This fact contradicts the existence of exponential invariants  $E(x, y) = \exp \{h(x, y)/r(x, y)\}$  with a non-constant polynomial  $r(x, y)$ .  $\square$

Now let us establish that Liénard differential systems (1.1) do not have Darboux first integrals whenever  $\deg f < \deg g < 2 \deg f + 1$ .

**Theorem 6.1.** *Liénard differential systems (1.1) from family (A) are not Darboux integrable.*

*Proof.* Using Theorem 4.2, we see that a Liénard differential system (1.1) with the restriction  $\deg f < \deg g < 2 \deg f + 1$  has at most two distinct irreducible invariant algebraic curves simultaneously. As in the proof of Lemma 6.1, we denote these curves as  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$ . Without loss of generality, we can suppose that the generating polynomials of these algebraic curves are given by expression (4.1) with  $N = 1$ ,  $k = 0$  and  $N \in \mathbb{N}$ ,  $k = 1$ , respectively. If there exists a first integral being a Darboux function, then this first integral should be of the form

$$I(x, y) = F_1^{d_1}(x, y)F_2^{d_2}(x, y), \quad d_1, d_2 \in \mathbb{C}, \quad |d_1| + |d_2| > 0. \quad (6.5)$$

Indeed, Lemma 6.1 forbids the existence of exponential factors given by invariants  $E(x, y) = \exp \{h(x, y)/r(x, y)\}$ , where  $h(x, y) \in \mathbb{C}[x, y]$  and  $r(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}$ . By Lemma 5.1 exponential invariants  $E(x, y) = \exp[h(x, y)]$  that could arise in an expression of the first integral have divisible by  $y$  cofactors, while invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$  have cofactors that are independent of  $y$ . Recall that there are no rational first integrals by the corollary to Theorem 4.2. Consequently, first integrals that are Darboux functions do not have exponential factors.

Let us note that if the Liénard differential system in question has only one irreducible invariant algebraic curve, for example  $F_1(x, y) = 0$ , then we suppose that  $d_2 = 0$  and vice versa.

The cofactors  $\lambda_1(x, y)$  and  $\lambda_2(x, y)$  of invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$  are given in relation (6.1). First integral (6.5) exists provided that the following condition  $d_1\lambda_1(x, y) + d_2\lambda_2(x, y) = 0$  is satisfied. This condition can be rewritten as

$$d_1[f(x) + q_x(x)] + d_2[Nf(x) + (N - 1)q_x(x) + p_x(x)] = 0. \quad (6.6)$$

The highest-degree term in expression (6.5) is  $d_2f_0x^m$ . Since  $f_0 \neq 0$ , we get  $d_2 = 0$ . Consequently, the first integral can be chosen as a rational function with  $d_1 = 1$ . Again we recall the corollary to Theorem 4.2, which excludes the existence of rational first integrals.  $\square$

Further, our aim is to study the existence of non-autonomous Darboux first integrals with a time-dependent exponential factor.

**Lemma 6.2.** *A Liénard differential system (1.1) from family (A) has a non-autonomous Darboux first integral with a time-dependent exponential factor (2.4) if and only if  $\deg g = \deg f + 1$  and the following condition*

$$g(x) = \omega \left( \int_0^x f(s)ds - \omega x - z_0 \right), \quad \omega, z_0 \in \mathbb{C}, \quad \omega \neq 0 \quad (6.7)$$

is valid. A first integral takes the form

$$I(x, y, t) = \left( y + \int_0^x f(s)ds - \omega x - z_0 \right) \exp(\omega t). \quad (6.8)$$

There are no other independent non-autonomous Darboux first integrals with a time-dependent exponential factor.

*Proof.* If the polynomial  $g(x)$  is given by relation (6.7), then it is straightforward to verify that function (6.8) is a non-autonomous first integral of the related differential system.

Let us establish the necessity of condition (6.7). We repeat the reasoning given in the proof of Theorem 6.1. By Theorems 2.3 and 4.2 a non-autonomous first integral (2.4) reads as

$$I(x, y, t) = F_1^{d_1}(x, y)F_2^{d_2}(x, y) \exp(\omega t), \quad d_1, d_2, \omega \in \mathbb{C}, \quad |d_1| + |d_2| > 0, \quad \omega \neq 0. \quad (6.9)$$

Now we need to study the following condition

$$d_1[f(x) + q_x(x)] + d_2[Nf(x) + (N - 1)q_x(x) + p_x(x)] - \omega = 0. \quad (6.10)$$

Similarly to the case of Theorem 6.1, we obtain  $d_2 = 0$ . Further, it is without loss of generality to set  $d_1 = 1$ . Considering relation (6.10) as an ordinary differential equation for the polynomial  $q(x)$  and performing the integration, we find its solution

$$q(x) = - \int_0^x f(s)ds + \omega x \quad (6.11)$$

satisfying the condition  $q(0) = 0$  presented in (4.4). Finally, we note that the invariant algebraic curve  $F_1(x, y) = 0$  given by the polynomial  $F_1(x, y) = y - q(x) - z_0$  exists whenever the series of type (I) defined in (4.3) terminates at the zero term. In this case equation (1.4) has a polynomial solution  $y(x) = q(x) + z_0$ . Substituting our results into equation (1.4), we find expression (6.7). Consequently, we obtain the following equality  $\deg g = \deg f + 1$ . The absence of other independent non-autonomous Darboux first integrals (2.4) follows from the previous considerations and the uniqueness of the invariant algebraic curve  $F_1(x, y) = 0$ .  $\square$

It follows from Lemma 6.2 that ordinary differential equation (1.4) related to a Liénard differential system from family (A) possessing a non-autonomous Darboux first integral has a polynomial solution  $y(x) = q(x) + z_0$ .

We recall that integrating factors and Jacobi last multipliers with a constant ratio belong to the same equivalence class. We do not distinguish between them.

**Theorem 6.2.** *A Liénard differential system (1.1) from family (A) is Liouvillian integrable if and only if the following assertions are valid:*

1. *the system under consideration possesses two distinct irreducible invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$ , where the polynomial  $F_1(x, y) = y - q(x) - z_0$  is given by expression (4.1) with  $N = 1$ ,  $k = 0$  and the polynomial  $F_2(x, y) \in \mathbb{C}[x, y]$  takes the form (4.1) with  $N \in \mathbb{N}$ ,  $k = 1$ ;*
2. *the polynomials  $q(x)$  and  $p(x)$  giving initial parts of the Puiseux series near the point  $x = \infty$  that solve equation (1.4) identically satisfy the condition*

$$(n - m)[f(x) + q_x(x)] + (m + 1)p_x(x) = 0. \quad (6.12)$$

*The related Liénard differential system has the unique Darboux integrating factor*

$$M(x, y) = \frac{(y - q(x) - z_0)^{\frac{N(m+1)-(n+1)}{m+1}}}{F_2(x, y)}. \quad (6.13)$$

*Proof.* Using Theorem 2.2, we see that a Liouvillian integrable differential system should have a Darboux integrating factor. Arguing as in the proof of Theorem 6.1, we conclude that such an integrating factor does not involve exponential invariants and is of the form

$$M(x, y) = F_1^{d_1}(x, y)F_2^{d_2}(x, y), \quad d_1, d_2 \in \mathbb{C}, \quad |d_1| + |d_2| > 0. \quad (6.14)$$

In this expression the polynomials  $F_1(x, y)$  and  $F_2(x, y)$  define invariant algebraic curves. The polynomial  $F_1(x, y)$  is given by expression (4.1) with  $N = 1$  and  $k = 0$ . The polynomial  $F_2(x, y)$  is produced by the same expression with  $N \in \mathbb{N}$  and  $k = 1$ . Again we note that if the Liénard differential system in question has only one irreducible invariant algebraic curve, for example  $F_1(x, y) = 0$ , then we set  $d_2 = 0$  and vice versa. Calculating the divergence of vector field (5.2) yields  $\text{div } \mathcal{X} = -f(x)$ . Hence the necessary and sufficient condition for Darboux integrating factor (6.14) to exist takes the form  $d_1\lambda_1(x, y) + d_2\lambda_2(x, y) = f(x)$ , where  $\lambda_1(x, y)$  and  $\lambda_2(x, y)$  are the cofactors of invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$ , respectively. These cofactors are given in expression (6.1). The condition enabling the existence of the Darboux integrating factor explicitly reads as

$$d_1[f(x) + q_x(x)] + d_2[Nf(x) + (N - 1)q_x(x) + p_x(x)] = -f(x). \quad (6.15)$$

Balancing the highest-degree terms in this expression, we get  $d_2 = -1$ . Further, we recall that the invariant algebraic curve  $F_1(x, y) = 0$  is given by the polynomial  $F_1(x, y) = y -$

$q(x) - z_0$ . If  $d_1 = 0$ , then the curve  $F_1(x, y) = 0$  either does not exist or does not enter the explicit expression (6.14) of the integrating factor.

Now let us consider the representation of the cofactors  $\lambda_1(x, y)$  and  $\lambda_2(x, y)$  in the ring  $\mathbb{C}_\infty\{x\}[y]$ . They are of the form

$$\lambda_1(x, y) = -f(x) - q_x(x), \quad \lambda_2(x, y) = -\sum_{j=1}^{N-1} \left[ f(x) + \left( y_{j,\infty}^{(1)} \right)_x \right] - \left[ f(x) + \left( y_{N,\infty}^{(2)} \right)_x \right]. \quad (6.16)$$

Substituting these expressions into the condition  $d_1\lambda_1(x, y) + d_2\lambda_2(x, y) = f(x)$  with  $d_2 = -1$ , we find

$$d_1[f(x) + q_x(x)] - \sum_{j=1}^{N-1} \left[ f(x) + \left( y_{j,\infty}^{(1)} \right)_x \right] = \left( y_{N,\infty}^{(2)} \right)_x. \quad (6.17)$$

The Puiseux series  $y(x) = y_{j,\infty}^{(1)}(x)$  and the polynomial  $y(x) = q(x) + z_0$  solve equation (1.4). Thus, we obtain

$$f(x) + q_x(x) = -\frac{g(x)}{q(x) + z_0}, \quad f(x) + \left( y_{j,\infty}^{(1)} \right)_x = -\frac{g(x)}{y_{j,\infty}^{(1)}}. \quad (6.18)$$

Again we suppose that if there is no polynomial solution  $y(x) = q(x) + z_0$ , then  $d_1 = 0$ . Using expressions (6.18), we rewrite condition (6.17) in the form

$$g(x) \left[ \sum_{j=1}^{N-1} \frac{1}{y_{j,\infty}^{(1)}} - \frac{d_1}{q(x) + z_0} \right] = \left( y_{N,\infty}^{(2)} \right)_x. \quad (6.19)$$

Let us find the highest-order terms in a neighborhood of the point  $x = \infty$ . Using the asymptotic formulae

$$\begin{aligned} g(x) &= g_0 x^n + o(x^n), & q(x) &= -\frac{f_0}{m+1} x^{m+1} + o(x^{m+1}), & x &\rightarrow \infty \\ y_{j,\infty}^{(1)}(x) &= -\frac{f_0}{m+1} x^{m+1} + o(x^{m+1}), & y_{N,\infty}^{(2)}(x) &= -\frac{g_0}{f_0} x^{n-m} + o(x^{n-m}), & x &\rightarrow \infty, \end{aligned} \quad (6.20)$$

we collect the coefficients of  $x^{n-m-1}$  in expression (6.19). This yields

$$d_1 = N - 1 - \frac{n-m}{m+1}. \quad (6.21)$$

Using inequalities  $m < n < 2m+1$ , we see that the parameter  $d_1$  cannot be zero. Substituting relation (6.21) and  $d_2 = -1$  into expressions (6.14) and (6.15), we find condition (6.12) and the Darboux integrating factor as given in (6.14). □

*Remark.* Condition (6.12) is identically satisfied whenever  $\deg g = \deg f + 1$ . Indeed, if  $n = m + 1$ , then we get

$$f(x) + q_x(x) = \frac{(m+1)g_0}{f_0}, \quad p(x) = -\frac{g_0}{f_0}x. \quad (6.22)$$

Consequently, Liénard differential systems satisfying the restriction  $\deg g = \deg f + 1$  and possessing two distinct irreducible invariant algebraic curves are always Liouvillian integrable.

We see from Theorem 6.2 that equation (1.4) related to a Liouvillian integrable Liénard differential system from family (A) has a polynomial solution given by the expression  $y(x) = q(x) + z_0$ . In addition, using condition (6.12) and equation (1.4), we can represent the polynomials  $f(x)$  and  $g(x)$  in the integrable cases as

$$f(x) = -q_x(x) - \frac{m+1}{n-m}p_x(x), \quad g(x) = \frac{m+1}{n-m}[q(x) + z_0]p_x(x). \quad (6.23)$$

Recall that the polynomial  $q(x)$  is of degree  $m + 1$  and the polynomial  $p(x)$  is of degree  $n - m$ . It follows from Theorem 6.2 that Liénard differential systems (1.1) from family (A) are Liouvillian integrable if and only if the polynomials  $f(x)$  and  $g(x)$  are given by expressions (6.23) and the systems possess an invariant algebraic curve with the generating polynomial taking the form (4.1), where  $N \in \mathbb{N}$  and  $k = 1$ .

Now let us study the integrability of Liénard differential systems such that the related equation (1.4) has two distinct polynomial solutions. The following theorem is valid.

**Theorem 6.3.** *A Liénard differential system (1.1) from family (A) with two distinct invariant algebraic curves given by first-degree polynomials with respect to  $y$  is Liouvillian integrable if and only if the system is of the form*

$$x_t = y, \quad y_t = [k\beta v^{k-1}(x) + (k+l)v^{l-1}(x)]v_x y - k[\beta v^k(x) + v^l(x)]v^{l-1}(x)v_x, \quad (6.24)$$

where  $\beta \in \mathbb{C} \setminus \{0\}$ ,  $v(x)$  is a polynomial of degree  $(n-m)/l$ ,  $k$  and  $l$  are relatively prime natural numbers such that the restriction  $(m+1)l = (n-m)k$  holds. The related Liénard differential system has the unique Darboux integrating factor

$$M(x, y) = \{y - \beta v^k(x) - v^l(x)\}^{-\frac{1}{k}} \{y - v^l(x)\}^{-1} \quad (6.25)$$

and the invariant algebraic curves  $y - \beta v^k(x) - v^l(x) = 0$ ,  $y - v^l(x) = 0$ . A Liouvillian first integral is of the form

$$I(x, y) = \frac{k\beta^{\frac{1}{k}}}{k-l} \{y - \beta v^k(x) - v^l(x)\}^{\frac{k-l}{k}} + \sum_{j=0}^m \exp \left[ -\frac{\pi l(2j+1)i}{k} \right] \times \ln \left\{ \{y - \beta v^k(x) - v^l(x)\}^{\frac{1}{m+1}} - \exp \left[ \frac{\pi(2j+1)i}{m+1} \right] [\beta v^k(x)]^{\frac{1}{m+1}} \right\}. \quad (6.26)$$

*Proof.* We begin the proof by noting that we use novel designations for the polynomial solutions of equation (1.4). We set  $\tilde{q}(x) = q(x) + z_0$  and  $\tilde{p}(x) = p(x) + z_1$ . Condition (6.12) and equation (1.4) provide explicit expressions (6.23) for the polynomials  $f(x)$  and  $g(x)$ . Substituting the relation  $y(x) = \tilde{p}(x)$  and expressions (6.23) into equation (1.4), we integrate the resulting equation with respect to the polynomial  $\tilde{q}(x)$ . This way, we get

$$\tilde{q}(x) = \beta \tilde{p}^{\frac{m+1}{n-m}}(x) + \tilde{p}(x), \quad (6.27)$$

where  $\beta \in \mathbb{C}$  is a constant of integration. Recalling the fact that  $\tilde{q}(x)$  is a polynomial of degree  $m + 1$  and  $\tilde{p}(x)$  is a polynomial of degree  $n - m$ , we find that  $\beta \neq 0$ . Introducing relatively prime natural numbers  $k$  and  $l$  according to the rule  $(m + 1)l = (n - m)k$ , we represent the polynomials  $\tilde{p}(x)$  and  $\tilde{q}(x)$  as

$$\tilde{p}(x) = v^l(x), \quad \tilde{q}(x) = \beta v^k(x) + v^l(x). \quad (6.28)$$

In this expression  $v(x)$  is an arbitrary polynomial of degree  $(n - m)/l$ . Substituting  $N = 1$  into expression (6.13), we find the integrating factor as given in relation (6.25). Finally, we calculate the line integral

$$I(x, y) = \int_{(x_0, y_0)}^{(x, y)} M(x, y) [y dy + \{f(x)y + g(x)\} dx] \quad (6.29)$$

and obtain first integral (6.26), where  $i$  is the imaginary unit.  $\square$

*Remark.* The family of systems (6.24) can be transformed to the following simple form

$$s_\tau = z, \quad z_\tau = [k\beta s^{k-1} + (k + l)s^{l-1}] z - k [\beta s^k + s^l] s^{l-1} \quad (6.30)$$

via the generalized Sundman transformation  $s(\tau) = v(x)$ ,  $z(\tau) = y$ ,  $d\tau = v_x(x)dt$ . Substituting  $v(x) = s$ ,  $y = z$  into (6.26), we find a Liouvillian first integral for systems (6.30).

Let us demonstrate that for any fixed degrees of the polynomials  $f(x)$  and  $g(x)$  there exist Liénard differential systems (1.1) from family (A) that have Liouvillian first integrals. With this aim we suppose that the polynomial  $\tilde{p}(x)$  takes the following form  $\tilde{p}(x) = x^{n-m}$ . We find the polynomial  $\tilde{q}(x)$  using expression (6.27). The result is  $\tilde{q}(x) = \beta x^{m+1} + x^{n-m}$ . Thus, we conclude that the following family of Liénard differential systems

$$x_t = y, \quad y_t = [(m + 1)\beta x^m + (n + 1)x^{n-m-1}] y - (m + 1) (\beta x^n + x^{2n-2m-1}) \quad (6.31)$$

is Liouvillian integrable. The related Darboux integrating factor reads as

$$M(x, y) = (y - x^{n-m})^{-1} (y - \beta x^{m+1} - x^{n-m})^{\frac{m-n}{m+1}}. \quad (6.32)$$

In addition, if the numbers  $n$  and  $m$  are the following  $n = l(k + 1) - 1$  and  $m = lk - 1$ , where  $l, k \in \mathbb{N}$ ,  $k > 1$ , then relations (6.23) and (6.27) produce Liouvillian integrable systems

$$x_t = y, \quad y_t = [k\beta \tilde{p}^{k-1}(x) + k + 1] \tilde{p}_x(x)y - k (\beta \tilde{p}^{k-1}(x) + 1) \tilde{p}(x)\tilde{p}_x(x), \quad (6.33)$$

where  $\tilde{p}(x)$  is a polynomial of degree  $n - m = l$ . Using expression (6.25), we see that systems (6.33) possess the Darboux integrating factor

$$M(x, y) = \{y - \beta\tilde{p}^k(x) - \tilde{p}(x)\}^{-\frac{1}{k}} \{y - \tilde{p}(x)\}^{-1}. \quad (6.34)$$

A related Liouvillian first integral is fairly simple in the case  $k = 2$ . Let us explicitly present another expression of a first integral:

$$I(x, y) = \operatorname{arctanh} \left\{ \sqrt{\frac{\beta\tilde{p}^2(x)}{F(x, y)}} \right\} + \sqrt{\beta F(x, y)}, \quad F(x, y) = \beta\tilde{p}^2(x) + \tilde{p}(x) - y. \quad (6.35)$$

Finally, let us note that there exist Liouvillian integrable Liénard differential systems from family (A) such that the polynomial  $F_2(x, y)$  in expression (6.13) is of degree with respect to  $y$  greater than 1. In fact, the degree with respect to  $y$  of the polynomial  $F_2(x, y)$  can be an arbitrary natural number. This fact was established in article [25].

Our next step is to investigate the existence of non-autonomous Darboux–Jacobi last multipliers. The cases  $\deg g = \deg f + 1$  and  $\deg g \neq \deg f + 1$  will be considered separately.

**Lemma 6.3.** *A Liénard differential system (1.1) satisfying the condition  $\deg f + 1 < \deg g < 2 \deg f + 1$  has a non-autonomous Darboux–Jacobi last multiplier of the form (2.6) if and only if the following assertions are valid:*

1. *the system under consideration possesses two distinct irreducible invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$ , where the polynomial  $F_1(x, y) = y - q(x) - z_0$  is given by expression (4.1) with  $N = 1$ ,  $k = 0$  and the polynomial  $F_2(x, y) \in \mathbb{C}[x, y]$  reads as (4.1) with  $N \in \mathbb{N}$ ,  $k = 1$ ;*
2. *the polynomials  $q(x)$  and  $p(x)$  giving initial parts of the Puiseux series near the point  $x = \infty$  that solve equation (1.4) identically satisfy the condition*

$$(n - m)[f(x) + q_x(x)] + (m + 1)[p_x(x) + \omega] = 0, \quad (6.36)$$

where  $\omega \in \mathbb{C} \setminus \{0\}$  is a constant.

The non-autonomous Darboux–Jacobi last multiplier is unique and takes the form

$$M(x, y, t) = \frac{(y - q(x) - z_0)^{\frac{N(m+1)-(n+1)}{m+1}}}{F_2(x, y)} \exp(\omega t). \quad (6.37)$$

*Proof.* We use Theorem 2.4 and repeat the proof of the previous theorem. The only difference is in condition (6.15). Now this condition reads as

$$d_1[f(x) + q_x(x)] + d_2[Nf(x) + (N - 1)q_x(x) + p_x(x)] - \omega = -f(x), \quad (6.38)$$

where  $\omega \neq 0$ . The related non-autonomous Darboux–Jacobi last multiplier is given by the expression

$$M(x, y, t) = F_1^{d_1}(x, y) F_2^{d_2}(x, y) \exp[\omega t]. \quad (6.39)$$

Similarly to the case of Theorem 6.2, we find the values of  $d_1$  and  $d_2$ . They are  $d_1 = N - 1 - (n - m)/(m + 1)$  and  $d_2 = -1$ . Both of them are non-zero. Consequently, a Liénard differential system (1.1) with a non-autonomous Darboux–Jacobi last multiplier has two distinct irreducible invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$ . Substituting explicit values of the parameters  $d_1$  and  $d_2$  into condition (6.38) yields relation (6.36).

If there exist two distinct non-autonomous Darboux–Jacobi last multipliers (2.6), then their ratio is a Darboux first integral either autonomous or non-autonomous of the form (2.4). By Theorem 6.1 and Lemma 6.2 such a situation is impossible.  $\square$

**Lemma 6.4.** *A Liénard differential system (1.1) satisfying the condition  $\deg g = \deg f + 1$  has a non-autonomous Darboux–Jacobi last multiplier of the form (2.6) if and only if the system under study possesses the irreducible invariant algebraic curve  $F_2(x, y) = 0$  given by relation (4.1) with  $N \in \mathbb{N}$  and  $k = 1$ . A non-autonomous Darboux–Jacobi last multiplier is of the form*

$$M(x, y, t) = \frac{\exp \left[ \frac{g_0}{f_0} \{1 - (N - 1)(m + 1)\} t \right]}{F_2(x, y)}. \quad (6.40)$$

*There are no other non-autonomous Darboux–Jacobi last multipliers whenever the invariant algebraic curve  $F_2(x, y) = 0$  is unique. If, in addition, there exists the invariant algebraic curve  $F_1(x, y) = 0$  presented in expression (4.1) with  $N = 1$  and  $k = 0$ , then the system under consideration has a family of non-autonomous Darboux–Jacobi last multipliers*

$$M(x, y, t) = \frac{F_1^{d_1}(x, y) \exp \left[ \frac{g_0}{f_0} \{1 + \{d_1 - (N - 1)\}(m + 1)\} t \right]}{F_2(x, y)}, \quad d_1 \in \mathbb{C}. \quad (6.41)$$

*Proof.* Similarly to the case of Lemma 6.3, we see that a non-autonomous Darboux–Jacobi last multiplier of the form (2.6) exists if and only if the system under consideration has at least one invariant algebraic curve and condition (6.38) is valid. Again we suppose that  $d_j = 0$  whenever the invariant algebraic curve  $F_j(x, y) = 0$  does not exist. The explicit expression of a non-autonomous Darboux–Jacobi last multiplier is given by (6.39). Balancing the highest-degree terms in condition (6.38) yields  $d_2 = -1$ . This fact proves the existence of the invariant algebraic curve  $F_2(x, y) = 0$ . Further, we find that the polynomials  $f(x) + q_x(x)$  and  $p_x(x)$  are constants, see relation (6.22). Substituting equalities  $d_2 = -1$ ,  $f(x) + q_x(x) = (m + 1)g_0/f_0$  and  $p_x(x) = -g_0/f_0$  into condition (6.38), we obtain

$$(m + 1)g_0d_1 + \{1 - (N - 1)(m + 1)\}g_0 - f_0\omega = 0. \quad (6.42)$$

Setting  $d_1 = 0$ , we find the value of  $\omega$  as given in relation (6.40). If the Liénard differential system in question does not have the invariant algebraic curve  $F_1(x, y) = 0$ , then non-autonomous Darboux–Jacobi last multiplier (6.40) is unique. In the converse case, non-autonomous Darboux–Jacobi last multiplier (6.40) also exists. In addition, we recall that the system has time-dependent Darboux first integral (6.8). It is straightforward to show that the product of a first integral and a non-constant Jacobi last multiplier is another Jacobi last multiplier.  $\square$

It is demonstrated in [18] that Liénard differential systems with non-autonomous Darboux–Jacobi last multipliers (6.40) indeed exist. In addition, let us note that the famous Duffing–van der Pol oscillators belong to family (A). The classification of Liouvillian integrable Duffing–van der Pol oscillators is performed in [16].

## 7. Integrability of Liénard differential systems from family (B)

Let us consider families of Liénard differential systems with fixed degrees of the polynomials  $f(x)$  and  $g(x)$  such that the following relation  $\deg g = 2 \deg f + 1$  is satisfied. If we do not impose restrictions on the highest-degree coefficients  $f_0$  and  $g_0$  of the polynomials  $f(x)$  and  $g(x)$ , then it has been established in [21] that the Fuchs indices of the Puiseux series near the point  $x = \infty$  that solve related equations (1.4) depend on the parameters  $f_0$  and  $g_0$ . Consequently, performing the classification of irreducible invariant algebraic curves and Darboux or Liouvillian first integrals is a very difficult problem whenever only the degrees of the polynomials  $f(x)$  and  $g(x)$  are fixed. The method of Puiseux series can deal with each case of a positive rational Fuchs index separately. Interestingly, such a degeneracy leads to a variety of distinct integrable cases arising in Liénard differential systems from family (B).

This section is mainly devoted to the non-resonant case. We say that a Liénard differential system from family (B) is resonant near infinity if equation (4.5) possesses a solution in  $\mathbb{Q}^+$ . For convenience, we introduce the parameter  $\delta$  according to the rule

$$g_0 = \frac{f_0^2 - \delta^2}{4(m+1)}, \quad \delta \in \mathbb{C}, \quad \delta \neq \pm f_0. \quad (7.1)$$

Using this normalization, we solve equation (4.5). As a result we find the Fuchs indices. They take the form

$$p_1 = \frac{2(m+1)\delta}{\delta - f_0}, \quad p_2 = \frac{2(m+1)\delta}{\delta + f_0}. \quad (7.2)$$

There are no positive rational Fuchs indices if and only if the condition  $\delta/f_0 \notin \mathbb{Q} \setminus \{0\}$  is valid. In addition, we assume that the following inequality  $\deg f > 0$  holds. The integrability problem for Liénard differential systems (1.1) under restrictions  $\deg f = 0$  and  $\deg g = 1$  is simple, see the end of Section 5.

In this section we use the designations of Theorem 4.3. In particular, the Puiseux series near the point  $x = \infty$  that solve equation (1.4) are denoted as  $y_\infty^{(1)}(x)$  and  $y_\infty^{(2)}(x)$ . Introducing the variable  $\delta$  instead of the parameter  $g_0$ , we see that these series have the following dominant behavior

$$\begin{aligned} y_\infty^{(1)}(x) &= \frac{\delta - f_0}{2(m+1)} x^{m+1} + o(x^{m+1}), \quad x \rightarrow \infty; \\ y_\infty^{(2)}(x) &= -\frac{\delta + f_0}{2(m+1)} x^{m+1} + o(x^{m+1}), \quad x \rightarrow \infty. \end{aligned} \quad (7.3)$$

Let us denote the polynomial parts of the series  $y_\infty^{(1)}(x)$  and  $y_\infty^{(2)}(x)$  as  $q_1(x)$  and  $q_2(x)$ , respectively. Thus, we have the equalities

$$q_1(x) = \{y_\infty^{(1)}(x)\}_+, \quad q_2(x) = \{y_\infty^{(2)}(x)\}_+. \quad (7.4)$$

If  $\delta = 0$ , then the series  $y_\infty^{(1)}(x)$  and  $y_\infty^{(2)}(x)$  coincide. Let us omit the indices and set  $q(x) = \{y_\infty(x)\}_+$ .

We begin by investigating the existence of exponential invariants related to invariant algebraic curves.

**Lemma 7.1.** *Let  $h(x, y) \in \mathbb{C}[x, y]$  and  $r(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}$  be relatively prime polynomials. A Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta/f_0 \notin \mathbb{Q} \setminus \{0\}$  has exponential invariants  $E(x, y) = \exp \{h(x, y)/r(x, y)\}$  if and only if the following statements are valid:*

1.  $\delta = 0$ ;
2. there exists the invariant algebraic curve  $F(x, y) = 0$  with the generating polynomial  $F(x, y) = y - q(x)$ ;
3. the ordinary differential equation

$$q(x)u_x(x) + [f(x) + q_x(x)]u(x) = 0 \quad (7.5)$$

has a non-zero polynomial solution  $u(x)$ .

The related exponential invariant is of the form

$$E(x, y) = \exp \left[ \frac{u(x)}{y - q(x)} \right] \quad (7.6)$$

and possesses the cofactor  $\varrho(x, y) = u_x(x)$ .

*Proof.* It follows from Theorem 4.3 that any Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta/f_0 \notin \mathbb{Q} \setminus \{0\}$  has at most two distinct irreducible invariant algebraic curves simultaneously. The degrees with respect to  $y$  of polynomials producing irreducible invariant algebraic curves are either 1 or 2. If there exists an irreducible invariant algebraic curves of degree 2 with respect to  $y$ , then it is unique. Further, there can arise at most two distinct irreducible invariant algebraic curves of degree 1 with respect to  $y$ . Since  $E(x, y) = \exp \{h(x, y)/r(x, y)\}$  is an exponential invariant, we conclude that  $r(x, y) = 0$  is an invariant algebraic curve of the related system. Let us represent exponential invariants in the form

$$E(x, y) = \exp \left[ \frac{h(x, y)}{F_1^{n_1}(x, y)F_2^{n_2}(x, y)} \right], \quad n_1, n_2 \in \mathbb{N}_0, \quad n_1 + n_2 > 0, \quad (7.7)$$

where  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$  are irreducible invariant algebraic curves. Without loss of generality we set  $n_k = 0$  whenever the invariant algebraic curve  $F_k(x, y) = 0$  does not exist. Here  $k = 1$  or  $k = 2$ .

*Case 1.* Let us suppose that there exists only one irreducible invariant algebraic curve  $F_1(x, y) = 0$  of degree 2 with respect to  $y$ . Using relation (4.7), we find the cofactor  $\lambda_1(x, y)$ . The result is

$$\lambda(x, y) = -2f(x) - \left\{ (y_\infty^{(1)})_x + (y_\infty^{(2)})_x \right\}_+ \quad (7.8)$$

The dominant behavior of the cofactor near the point  $x = \infty$  is  $\lambda(x, y) = -f_0x^m + o(x^m)$ . Now let us take one of the Puiseux series, for example, the series  $y_\infty^{(1)}(x)$ . We find the asymptotic relation

$$\frac{\lambda\left(x, y_\infty^{(1)}(x)\right)}{y_\infty^{(1)}(x)} = -\frac{2(m+1)f_0}{(\delta-f_0)x} + o\left(\frac{1}{x}\right), \quad x \rightarrow \infty \quad (7.9)$$

Since  $\delta/f_0 \notin \mathbb{Q} \setminus \{0\}$ , we conclude that the conditions of Theorem 3.2 are not satisfied whenever  $\delta \neq 0$ . The case  $\delta = 0$  will be considered separately. Exponential invariants do not exist provided that  $\delta \neq 0$ .

*Case 2.* Let us suppose that the differential system under consideration has two distinct irreducible invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$ . The polynomials producing the curves are of degree 1 with respect to  $y$  and have the cofactors

$$\lambda_m(x, y) = -f(x) - \left\{ \left( y_\infty^{(m)} \right)_x \right\}_+, \quad m = 1, 2. \quad (7.10)$$

The cofactor of the invariant algebraic curve  $F_1^{n_1}(x, y)F_2^{n_2}(x, y) = 0$  is given by the polynomial  $\lambda(x, y) = n_1\lambda_1(x, y) + n_2\lambda_2(x, y)$  and has the following dominant behavior near the point  $x = \infty$ :

$$\lambda(x, y) = \frac{1}{2} [n_2(\delta - f_0) - n_1(\delta + f_0)] x^m + o(x^m), \quad x \rightarrow \infty. \quad (7.11)$$

Suppose that one of the numbers  $n_1$  and  $n_2$  is non-zero, for example,  $n_1$ . Further, we consider the following asymptotic relation

$$\frac{\lambda\left(x, y_\infty^{(1)}(x)\right)}{y_\infty^{(1)}(x)} = (m+1) \left( n_2 - \frac{(\delta + f_0)n_1}{\delta - f_0} \right) \frac{1}{x} + o\left(\frac{1}{x}\right), \quad x \rightarrow \infty. \quad (7.12)$$

Using Theorem 3.2 and expression  $\delta/f_0 \notin \mathbb{Q} \setminus \{0\}$ , we see that there are no exponential invariants (7.7) whenever  $\delta \neq 0$ .

*Case 3.* Let us suppose that the differential system in question has only one irreducible invariant algebraic curve: either  $F_1(x, y) = 0$  or  $F_2(x, y) = 0$ . If  $\delta \neq 0$ , then arguing as in the previous case, we prove the non-existence of exponential invariants.

*Case 4.* Now let us suppose that  $\delta = 0$ . Recall that the Puiseux series  $y_\infty^{(1)}(x)$  and  $y_\infty^{(2)}(x)$  merge in this case. The unique Puiseux series centered at the point  $x = \infty$  that satisfies equation (1.4) is denoted as  $y_\infty(x)$ . In what follows we use the local theory of invariants considered in Section 3. Let a Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta = 0$  have the irreducible invariant algebraic curve  $F(x, y) = 0$ , where  $F(x, y) = y - q(x)$ . If in addition the system possesses an exponential invariant  $E(x, y) = \exp\{h(x, y)/F^k(x, y)\}$  for some  $k \in \mathbb{N}$ , then it is without loss of generality to suppose that the degree of the polynomial  $h(x, y)$  with respect to  $y$  is at most  $k - 1$ . There exists a finite number of local elementary exponential invariants

$$E_j(x, y) = \exp \left[ \frac{u_j(x)}{\{y - y_\infty(x)\}^{k_j}} \right], \quad u_j(x) \in \mathbb{C}_\infty\{x\}, \quad (7.13)$$

$$k_j \in \mathbb{N}, \quad j = 1, \dots, K, \quad K \in \mathbb{N}$$

such that the exponential invariant  $E(x, y)$  equals the product  $\prod_{j=1}^K E_j(x, y)$ . We assume that the following inequalities  $1 \leq k_1 < k_2 < \dots < k_K \leq k$  are valid. Let us denote the cofactor of the local elementary exponential invariant  $E_j(x, y)$  by  $\varrho_j(x, y) \in \mathbb{C}_\infty\{x\}[y]$ . We see that the following expression  $\sum_{j=1}^K \varrho_j(x, y)$  equals the cofactor  $\varrho(x, y)$  of the invariant  $E(x, y)$ . Note that the cofactor  $\varrho(x, y)$  is an element of the ring  $\mathbb{C}[x, y]$ .

Substituting the explicit representation of  $E_j(x, y)$  into the partial differential equation  $\mathcal{X}E_j(x, y) = \varrho_j(x, y)E_j(x, y)$ , we find

$$yu_{j,x} = k_j \lambda_j(x, y) u_j(x) + \varrho_j(x, y) \{y - y_\infty(x)\}^{k_j}. \quad (7.14)$$

In this expression  $\lambda_j(x, y) \in \mathbb{C}_\infty\{x\}[y]$  is the cofactor of the local elementary invariant  $f(x, y) = y - y_\infty(x)$ . Using Theorem 3.4, we obtain the cofactor  $\lambda_j(x, y)$ . The result is

$$\lambda_j(x, y) = -f(x) - \{y_\infty(x)\}_x. \quad (7.15)$$

Analyzing expression (7.14), we see that  $j = 1$ ,  $k_1 = k = 1$ ,  $\varrho_1(x, y) = u_{1,x}(x)$ , and the series  $u_1(x)$  satisfies the following ordinary differential equation

$$y_\infty(x)u_{1,x}(x) + (f(x) + \{y_\infty(x)\}_x) u_1(x) = 0. \quad (7.16)$$

Since there exists the invariant algebraic curve  $y - q(x) = 0$ , we conclude that the Puiseux series  $y_\infty(x)$  is in fact the polynomial  $q(x)$ :  $y_\infty(x) = q(x)$ . Consequently, the exponential invariant  $E(x, y) = \exp\{h(x, y)/F(x, y)\}$  exists if and only if  $h(x, y) = u_1(x)$  and  $u_1(x)$  is a polynomial. Omitting the index, we get expressions (7.5) and (7.6). □

*Remark.* Balancing the highest-degree terms in equation (7.5), it can be shown that the polynomial  $u(x)$  is of degree  $m + 1$ .

Our next step is to derive the necessary and sufficient conditions of Darboux integrability for non-resonant Liénard differential systems from family (B). The case  $\delta = 0$  will be considered separately.

**Theorem 7.1.** *A Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta/f_0 \notin \mathbb{Q}$  is Darboux integrable if and only if the system is of the form*

$$x_t = y, \quad y_t = \frac{2f_0}{f_0 - \delta} q_{1,x} y - \frac{f_0 + \delta}{f_0 - \delta} q_{1,x} q_1, \quad (7.17)$$

where  $q_1(x)$  is a polynomial of degree  $m + 1$  with the highest-degree coefficient  $(\delta - f_0)/(2\{m + 1\})$ . A related Darboux first integral reads as

$$I(x, y) = [y - q_1(x)]^{\delta - f_0} \left[ y - \frac{(f_0 + \delta)}{(f_0 - \delta)} q_1(x) \right]^{\delta + f_0}. \quad (7.18)$$

*Proof.* It is straightforward to verify that expression (7.18) gives a Darboux first integral whenever all other conditions of the theorem are valid.

Let us prove the converse statement. It follows from Lemmas 5.1 and 7.1 that Darboux first integrals of Liénard differential systems (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta/f_0 \notin \mathbb{Q}$  do not have exponential factors. By Theorem 4.3 the systems under consideration have at most two distinct irreducible invariant algebraic curves simultaneously.

First, we consider a Liénard differential system (1.1) that satisfies the conditions  $\deg g = 2 \deg f + 1$ ,  $\delta/f_0 \notin \mathbb{Q}$  and possesses only one irreducible invariant algebraic curve. If there exists a Darboux first integral, then it can be chosen as a bivariate polynomial  $I(x, y) = F(x, y)$  producing the invariant algebraic curve  $F(x, y) = 0$ . Using Theorem 4.3, we see that the generating polynomial  $F(x, y)$  is of degree at most 2 with respect to  $y$ . The Darboux first integral  $I(x, y) = F(x, y)$  exists if and only if the cofactor  $\lambda(x, y)$  of the invariant algebraic curve  $F(x, y) = 0$  is identically zero. We need to consider three possibilities. Let us write down the related cofactors and their dominant behavior near the point  $x = \infty$ . With the help of relations (7.3) and (7.4), we get

$$\begin{aligned} N = 1 : \lambda(x, y) &= -f(x) - q_{1,x}, \quad \lambda(x, y) = -\frac{1}{2}(\delta + f_0)x^m + o(x^m), \quad x \rightarrow \infty; \\ N = 1 : \lambda(x, y) &= -f(x) - q_{2,x}, \quad \lambda(x, y) = \frac{1}{2}(\delta - f_0)x^m + o(x^m), \quad x \rightarrow \infty; \\ N = 2 : \lambda(x, y) &= -2f(x) - q_{1,x} - q_{2,x}, \quad \lambda(x, y) = -f_0x^m + o(x^m), \quad x \rightarrow \infty. \end{aligned} \quad (7.19)$$

In these expressions  $N$  denotes the degree of the polynomial  $F(x, y)$  with respect to  $y$ . We conclude that the cofactors are not identically zero. Thus, there are no Darboux first integrals.

Now, we consider a Liénard differential system (1.1) that satisfies the conditions  $\deg g = 2 \deg f + 1$ ,  $\delta/f_0 \notin \mathbb{Q}$  and possesses two distinct irreducible invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$ . By Theorem 4.3 the polynomials  $F_1(x, y)$  and  $F_2(x, y)$  are of the form  $F_1(x, y) = y - q_1(x)$  and  $F_2(x, y) = y - q_2(x)$ . If there exists a Darboux first integral, then it can be represented in the form

$$I(x, y) = F_1^{d_1}(x, y)F_2^{d_2}(x, y), \quad d_1, d_2 \in \mathbb{C}, \quad |d_1| + |d_2| > 0 \quad (7.20)$$

This first integral exists if and only if the following condition  $d_1\lambda_1(x, y) + d_2\lambda_2(x, y) = 0$  is satisfied. Finding the dominant behavior near the point  $x = \infty$  of the cofactors

$$\lambda_1(x, y) = -\frac{1}{2}(\delta + f_0)x^m + o(x^m), \quad \lambda_2(x, y) = \frac{1}{2}(\delta - f_0)x^m + o(x^m), \quad (7.21)$$

we obtain  $d_1 = \delta - f_0$ ,  $d_2 = \delta + f_0$  and the expression

$$2\delta f(x) + (\delta - f_0)q_{1,x}(x) + (\delta + f_0)q_{2,x}(x) = 0. \quad (7.22)$$

Recall that one of the parameters  $d_1$  or  $d_2$  can be chosen arbitrary. By Theorem 7.1 equation (1.4) related to the Liénard differential system under consideration possesses two distinct

polynomial solutions  $y(x) = q_1(x)$  and  $y(x) = q_2(x)$ . Thus, we obtain the following relations

$$f(x) + q_{1,x}(x) = -\frac{g(x)}{q_1(x)}, \quad f(x) + q_{2,x}(x) = -\frac{g(x)}{q_2(x)}. \quad (7.23)$$

Substituting these relations and the values of  $d_1$  and  $d_2$  into the condition on the cofactors  $d_1[f(x) + q_{1,x}(x)] + d_2[f(x) + q_{2,x}(x)] = 0$  yields the expression  $q_2(x) = -d_2q_1(x)/d_1$ . Finally, we use relations (7.22) and (7.23) in order to derive the explicit representations of the polynomials  $f(x)$  and  $g(x)$ .  $\square$

*Corollary.* The family of first-order ordinary differential equations

$$yy_x - \frac{2f_0}{f_0 - \delta}q_{1,x}y + \frac{f_0 + \delta}{f_0 - \delta}q_{1,x}q_1 = 0 \quad (7.24)$$

associated with systems (7.17) has two distinct polynomial solutions of the form  $y(x) = q_1(x)$  and  $y(x) = (f_0 + \delta)q_1(x)/(f_0 - \delta)$ .

*Remark 1.* Suppose we are in assumptions of Theorem 7.1 with the exception of the condition  $\delta/f_0 \notin \mathbb{Q} \setminus \{0\}$ . Then function (7.18) is still a Darboux first integral of the related Liénard differential system. In addition, the corollary to Theorem 7.1 is also valid. However, there may exist other resonant Liénard differential systems from family (B) with Darboux first integrals.

*Remark 2.* Expression (7.17) provides a set of systems with rational first integrals (7.18) provided that  $f_0/\delta$  is a rational number.

Next, let us study the Darboux integrability in the case  $\delta = 0$ .

**Theorem 7.2.** *A Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta = 0$  is Darboux integrable if and only if the system can be represented in the form*

$$x_t = y, \quad y_t = 2q_x(x)y - q(x)q_x(x), \quad (7.25)$$

where  $q(x)$  is a polynomial of degree  $m + 1$ . A related Darboux first integral reads as

$$I(x, y) = [y - q(x)] \exp \left[ -\frac{q(x)}{y - q(x)} \right]. \quad (7.26)$$

*Proof.* By direct computations we verify that expression (7.26) gives a Darboux first integral of a Liénard differential system (1.1) with the polynomials  $f(x)$  and  $g(x)$  satisfying relations (7.25) and  $\delta = 0$ . Thus, we have established sufficiency of conditions presented in the theorem.

Let us prove their necessity. Suppose that a Liénard differential system from family (B) with  $\delta = 0$  possesses a Darboux first integral. A Darboux integrable differential system (2.1) has at least one invariant algebraic curve. It follows from Theorem 4.3 that the Liénard differential system in question possesses at most one irreducible invariant algebraic curve. This curve is of the form  $y - q(x) = 0$  and has the cofactor  $\lambda(x, y) = -f(x) - q_x(x)$ . Thus, we have established the existence of the invariant algebraic curve  $y - q(x) = 0$ .

According to Lemma 7.1, the Liénard differential system under consideration may have exponential invariants associated with the invariant algebraic curve  $y - q(x) = 0$ . These invariants take the form (7.6) and possess the cofactor  $\varrho(x, y) = u_x(x)$ , where the polynomial  $u(x)$  satisfies equation (7.5). Using Lemma 5.1, we conclude that exponential invariants with a polynomial argument cannot enter explicit expressions of Darboux first integrals. Consequently, a Darboux first integral can be represented in the form

$$I(x, y) = [y - q(x)]^d \exp \left[ \frac{u(x)}{y - q(x)} \right], \quad d \in \mathbb{C}, \quad (7.27)$$

where we suppose that  $u(x) \equiv 0$  whenever exponential invariants (7.6) do not exist.

If  $d = 0$ , then the related Liénard differential system has an invariant algebraic curve  $u(x) = 0$  independent of  $y$ . It is impossible due to Theorem 4.3. Thus, it is without loss of generality to set  $d = 1$ . The cofactors of all the invariants identically satisfy the relation  $\lambda(x, y) + \varrho(x, y) = 0$  provided that first integral (7.27) exists. As a result, we get the expression  $f(x) = u_x(x) - q_x(x)$ . Substituting this expression into equation (7.5) yields  $u(x) = -q(x)$ . This relation proves existence of exponential invariants (7.6). Since  $y = q(x)$  is a polynomial solution of equation (1.4) and  $f(x) = -2q_x(x)$ , we find the polynomial  $g(x)$ . The result is  $g(x) = q(x)q_x(x)$ . □

Interestingly, Liénard differential systems (7.17) and (7.25) are those characterized by the so-called Chiellini integrability condition

$$\frac{d}{dx} \left[ \frac{g(x)}{f(x)} \right] = \alpha f(x), \quad \alpha \in \mathbb{C} \setminus \{0\}. \quad (7.28)$$

This condition was originally introduced by A. Chiellini [10]. Chiellini integrable Liénard differential systems can be transformed to linear systems  $s_\tau = z$ ,  $z_\tau = -z - \alpha s$  via the generalized Sundman transformation  $s(\tau) = \int f(x)dx$ ,  $z(\tau) = y$ ,  $d\tau = f(x)dt$ , see [2]. Some other properties of Chiellini integrable Liénard differential systems are presented in [29].

Our next step is to study the existence of non-autonomous Darboux first integrals with a time-dependent exponential factor (2.4).

**Lemma 7.2.** *A Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$ ,  $\deg f \neq 0$ , and  $\delta/f_0 \notin \mathbb{Q}$  possesses a non-autonomous Darboux first integral (2.4) if and only if the system reads as*

$$\begin{aligned} x_t = y, \quad y_t = - \left\{ f_0(x - x_0)^m + \frac{(m+2)\omega}{2(m+1)\delta} \right\} y - \frac{f_0^2 - \delta^2}{4(m+1)}(x - x_0)^{2m+1} \\ - \frac{f_0\omega}{2(m+1)\delta}(x - x_0)^{m+1} - \frac{\omega^2}{4(m+1)\delta^2}(x - x_0), \end{aligned} \quad (7.29)$$

where  $x_0 \in \mathbb{C}$  and  $\omega \in \mathbb{C} \setminus \{0\}$ . A related non-autonomous Darboux first integral takes the form

$$\begin{aligned} I(x, y, t) &= \left[ y + \frac{f_0 - \delta}{2(m+1)}(x - x_0)^{m+1} + \frac{\omega}{2(m+1)\delta}(x - x_0) \right]^{\delta - f_0} \\ &\times \left[ y + \frac{\delta + f_0}{2(m+1)}(x - x_0)^{m+1} + \frac{\omega}{2(m+1)\delta}(x - x_0) \right]^{\delta + f_0} \exp(\omega t). \end{aligned} \quad (7.30)$$

*Proof.* It is straightforward to derive that expression (7.30) is a non-autonomous Darboux first integral of systems (7.29). We only need to prove the converse statement. Supposing that a non-resonant Liénard differential system from family (B) possesses a non-autonomous Darboux first integral (2.4), we use the arguments given in the proof of Theorem 7.1 to represent this first integral as

$$I(x, y, t) = [y - q_1(x)]^{\delta - f_0} [y - q_2(x)]^{\delta + f_0} \exp(\omega t), \quad \omega \neq 0, \quad (7.31)$$

where  $q_1(x)$  and  $q_2(x)$  are distinct polynomial solutions of equation (1.4). In addition, we get the following condition

$$2\delta f(x) + (\delta - f_0)q_{1,x}(x) + (\delta + f_0)q_{2,x}(x) - \omega = 0. \quad (7.32)$$

Further, we express the polynomial  $f(x)$  from this condition. Substituting  $y(x) = q_1(x)$  into equation (1.4), we obtain the polynomial  $g(x)$ . Let us introduce the polynomial  $v(x)$  according to the rule  $q_2(x) - q_1(x) = v(x)$ . Requiring that the function  $q_2(x) = q_1(x) + v(x)$  is a solution of equation (1.4), we get the relation

$$[(\delta - f_0)v(x) + 2\delta q_1(x)]v_x(x) + \omega v(x) = 0 \quad (7.33)$$

Since  $q_1(x)$  is a polynomial, we obtain the ordinary differential equation  $\beta(x - x_0)v_x = v$ , where  $\beta, x_0 \in \mathbb{C}$ . Integrating this equation yields  $v(x) = v_0(x - x_0)^{1/\beta}$  with  $v_0 \in \mathbb{C}$  being a constant of integration. It follows from expression  $q_2(x) - q_1(x) = v(x)$  that  $v(x)$  is a polynomial of degree  $m + 1$ . Thus, we obtain  $\beta = 1/(m + 1)$ . As a result the polynomials  $q_1(x)$  and  $q_2(x)$  can be represented in the form

$$\begin{aligned} q_1(x) &= \frac{\delta - f_0}{2(m+1)}(x - x_0)^{m+1} - \frac{\omega}{2(m+1)\delta}(x - x_0), \\ q_2(x) &= -\frac{\delta + f_0}{2(m+1)}(x - x_0)^{m+1} - \frac{\omega}{2(m+1)\delta}(x - x_0). \end{aligned} \quad (7.34)$$

Finally, we find the polynomials  $f(x)$  and  $g(x)$  from condition (7.32) and equation (1.4) recalling the fact that, for example,  $y(x) = q_1(x)$  is a solution of the latter. The uniqueness of independent non-autonomous Darboux first integral (7.30) follows from the uniqueness of the polynomials  $q_1(x)$ ,  $q_2(x)$  and the dominant behavior of the cofactors given by expression (7.21). □

*Remark.* Suppose we are in assumptions of Lemma 7.2 with the exception of the condition  $\delta/f_0 \notin \mathbb{Q} \setminus \{0\}$ . Then function (7.30) is still a non-autonomous Darboux first integral of the related Liénard differential system. However, there may exist other resonant Liénard differential systems from family (B) with non-autonomous Darboux first integrals of the form (2.4).

Let us note that if  $\delta = \pm m f_0 / (m + 2)$ , then any system (7.29) has not only a non-autonomous Darboux first integral (7.30), but also an independent Darboux first integral (7.18), where  $q_1(x)$  is given by the relation

$$q(x) = -\frac{f_0 x^{m+1}}{(m+1)(m+2)} - \frac{(m+2)\omega x}{2(m+1)f_0 m}. \quad (7.35)$$

Let  $I_1(x, y)$  be Darboux first integral (7.18) and  $I_2(x, y, t)$  be non-autonomous Darboux first integral (7.30). Eliminating  $y$  from the relations  $I_1(x, y) = C_1$  and  $I_2(x, y, t) = C_2$ , we can find the general solution of a system (7.29) under the condition  $\delta = \pm m f_0 / (m + 2)$ . Note that such a system is resonant near  $x = \infty$ . The general solution in the case  $m = 2$  previously appeared in [40], see also [24].

**Lemma 7.3.** *A Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$ ,  $\deg f \neq 0$ , and  $\delta = 0$  possesses a non-autonomous Darboux first integral (2.4) if and only if the system is of the form*

$$x_t = y, \quad y_t = -\left\{f_0(x - x_0)^m + \frac{(m+2)\omega}{m+1}\right\}y - \frac{(x - x_0)}{4(m+1)}\{f_0(x - x_0)^m + 2\omega\}^2, \quad (7.36)$$

where  $x_0 \in \mathbb{C}$  and  $\omega \in \mathbb{C} \setminus \{0\}$ . A related non-autonomous Darboux first integral reads as

$$I(x, y, t) = \exp\left[\frac{f_0(x - x_0)^{m+1}}{2(m+1)y + f_0(x - x_0)^{m+1} + 2\omega(x - x_0)}\right] \times \left[y + \frac{f_0(x - x_0)^{m+1}}{2(m+1)} + \frac{\omega(x - x_0)}{m+1}\right] \exp(\omega t). \quad (7.37)$$

*Proof.* By direct computations we verify that expression (7.37) is a time-dependent first integral of system (7.36).

Let us prove the converse statement. We suppose that a Liénard differential system from family (B) satisfies the restriction  $\delta = 0$  and possesses a non-autonomous Darboux first integral (2.4). Repeating the arguments used in the proof of Theorem 7.2, we represent such a first integral in the form

$$I(x, y, t) = [y - q(x)]^d \exp\left[\frac{u(x)}{y - q(x)}\right] \exp(\omega t), \quad d \in \mathbb{C}, \quad \omega \in \mathbb{C} \setminus \{0\}. \quad (7.38)$$

In addition, we note that the related system possesses the invariant algebraic curve  $y - q(x) = 0$  with the cofactor  $\lambda(x, y) = -f(x) - q_x(x)$ . By Lemma 7.1 the cofactor  $\varrho(x, y)$  of the exponential invariant  $E(x, y) = \exp[u(x)/(y - q(x))]$  reads as  $\varrho(x, y) = u_x(x)$ . Condition (2.7) relating these cofactors and the parameter  $\omega$  takes the form

$$d[f(x) + q_x(x)] - u_x(x) - \omega = 0, \quad d \in \mathbb{C}. \quad (7.39)$$

Let us begin with the case  $d = 0$ . We conclude from condition (7.39) that  $u(x)$  is a first-degree polynomial. Using the remark to Lemma 7.1, we find the value of  $m$ . The result is  $m = 0$ . As it was mentioned at the beginning of this section, we do not consider Liénard differential systems with the restriction  $m = 0$  ( $\deg f = 0$ ).

We turn to the case  $d \neq 0$ . Without loss of generality, we set  $d = 1$ . Now let us suppose that the restriction  $u(x) \equiv 0$  is valid. The related system may not have exponential invariants. Finding the dominant behavior of the cofactor  $\lambda(x, y) = -f(x) - q_x(x)$  near the point  $x = \infty$ , we obtain

$$\lambda(x, y) = -\frac{f_0}{2}x^m + o(x^m), \quad x \rightarrow \infty. \quad (7.40)$$

Recalling the inequality  $m > 0$ , we see that condition (7.39) is not satisfied. Thus, we conclude that the exponential invariant exists. Further, we eliminate from relations (7.5) and (7.39) the polynomial  $f(x)$ . As a result we get the following expression

$$q(x) = -u(x) - \omega \frac{u(x)}{u_x(x)}. \quad (7.41)$$

Consequently, the ratio  $u(x)/u_x(x)$  is a polynomial. It is straightforward to see that this polynomial is of the first degree and can be represented as  $\beta(x - x_0)$ , where  $\beta, x_0 \in \mathbb{C}$ . Integrating the ordinary differential equation  $\{\beta(x - x_0)\}u_x(x) = u(x)$ , we obtain  $u(x) = u_0(x - x_0)^{1/\beta}$ , where  $u_0 \in \mathbb{C}$  is a constant of integration. We recall that  $u(x)$  is a polynomial of degree  $m + 1$ . As a result we get  $\beta = 1/(m + 1)$ . Substituting the relation  $u(x) = u_0(x - x_0)^{m+1}$  into expressions (7.41) and (7.39), we find the polynomials  $q(x)$  and  $f(x)$ . In addition, we choose the parametrization  $u_0 = f_0/(2\{m + 1\})$ . The polynomial  $g(x)$  we find recalling the fact that  $y(x) = q(x)$  is the polynomial solution of the related equation (1.4).

Thus, we see that if the Liénard differential system in question has a non-autonomous Darboux first integral (2.4), then there exist the invariant algebraic curve  $y - q(x) = 0$  and the exponential invariant  $E(x, y) = \exp[\alpha u(x)/(y - q(x))]$ , where  $\alpha \in \mathbb{C}$ , the polynomial  $q(x)$  is given by expression (7.41), and the polynomial  $u(x)$  is  $u(x) = f_0(x - x_0)^{m+1}/(2\{m + 1\})$ .  $\square$

Below we shall prove that Liénard differential systems (7.29) and (7.36) are Liouvillian integrable.

Let us study the Liouvillian integrability of Liénard differential systems from family (B). We begin with some partial case characterized by Darboux integrating factors of a special form. Note that in Theorem 7.3 we use novel designations for polynomial solutions of equations (1.4) related to Liénard differential systems. We need novel designations because polynomials  $p_1(x)$  and  $p_2(x)$  may have coinciding dominant terms. After considering this special case, we shall turn to non-resonant systems.

**Theorem 7.3.** *A Liénard differential system (1.1) from family (B) possesses the Darboux integrating factor*

$$M(x, y) = [y - p_1(x)]^{d_1} [y - p_2(x)]^{d_2}, \quad d_1, d_2 \in \mathbb{C} \setminus \{0\}, \quad (7.42)$$

where  $p_1(x)$  and  $p_2(x)$  are distinct polynomials, if and only if one of the following assertions is valid.

1. The system is of the form (7.17) and the polynomials  $p_1(x)$  and  $p_2(x)$  are linearly dependent:  $p_2(x) = (f_0 + \delta)/(f_0 - \delta)p_1(x)$ . In this case the parameters  $d_1$  and  $d_2$  can be chosen as  $d_1 = d_2 = -1$  and the following relations  $p_1(x) = q_1(x)$ ,  $p_2(x) = q_2(x)$  are valid. In fact, there exists a family of Darboux integrating factors (7.42) that are products of the integrating factor  $M_0(x, y) = [y - q_1(x)]^{-1}[y - q_2(x)]^{-1}$  and the Darboux first integrals  $I^\varkappa(x, y)$ , where the function  $I(x, y)$  is given by expression (7.18) and  $\varkappa \in \mathbb{C}$ .
2. The system reads as

$$\begin{aligned} x_t = y, \quad y_t = & \left[ \beta(l+k)u^{k-1} + \frac{\{(2d_1+1)l+k\}l}{k-l}u^{l-1} \right] u_x y \\ & - \left[ l\beta^2 u^{2k-1} + \frac{\{(2d_1+1)l+k\}l\beta}{k-l}u^{k+l-1} + \frac{(ld_1+k)(d_1+1)l^2}{(k-l)^2}u^{2l-1} \right] u_x, \end{aligned} \quad (7.43)$$

where  $k$  and  $l$  are relatively prime both non-unit natural numbers,  $u(x)$  is a polynomial of degree  $(m+1)/\max\{k, l\}$  and  $\beta \in \mathbb{C} \setminus \{0\}$ . The polynomials  $p_1(x)$  and  $p_2(x)$  can be represented as

$$p_1(x) = \beta u^k(x) + \frac{(d_1+1)l}{k-l}u^l(x), \quad p_2(x) = \beta u^k(x) + \frac{(ld_1+k)}{k-l}u^l(x) \quad (7.44)$$

and the parameter  $d_2$  is given by the relation  $d_2 = -(d_1 + 1 + k/l)$ .

*Proof.* Expression (7.42) gives a Darboux integrating factor of a Liénard differential system if and only if the system possesses the invariant algebraic curves  $y - p_1(x) = 0$  and  $y - p_2(x) = 0$  such that the following condition  $d_1\lambda_1(x, y) + d_2\lambda_2(x, y) - f(x) = 0$  is identically satisfied. It is straightforward to find the cofactor  $\lambda_j(x, y)$  of the invariant algebraic curve  $y - p_j(x) = 0$ . The result is  $\lambda_j(x, y) = -f(x) - p_{j,x}(x)$ ,  $j = 1, 2$ . Thus, we arrive at the condition

$$(d_1 + d_2 + 1)f(x) + d_1p_{1,x}(x) + d_2p_{2,x}(x) = 0. \quad (7.45)$$

If the following restriction  $d_2 = -1 - d_1$  is valid, then integrating equation (7.45) with respect to the polynomial  $p_1(x)$ , we obtain  $p_1(x) = (d_1 + 1)p_2(x)/d_1 + \beta$ , where  $\beta \in \mathbb{C}$  is a constant of integration. Recalling the fact that  $y = p_1(x)$  and  $y = p_2(x)$  are polynomial solutions of equation (1.4), we find the polynomials  $f(x)$  and  $g(x)$ . The polynomial  $f(x)$  can be represented in the form

$$f(x) = -\frac{(2d_1+1)p_2(x) + d_1(d_1+1)\beta}{d_1(p_2(x) + d_1\beta)}p_{2,x}(x). \quad (7.46)$$

Analyzing this expression, we conclude that the function on the right-hand side is not a polynomial whenever  $p_2(x)$  is non-constant.

Let us consider the case  $d_2 \neq -1 - d_1$ . We find the polynomials  $f(x)$  and  $g(x)$  from condition (7.45) and equation (1.4), where we set  $y(x) = p_1(x)$ . Substituting the resulting expressions into equation (1.4) and recalling the fact that  $y(x) = p_2(x)$  is a solution of the latter, we obtain the equation

$$\{(d_2 + 1)p_1(x) + d_1 p_2(x)\}p_{1,x}(x) - \{d_2 p_1(x) + (d_1 + 1)p_2(x)\}p_{2,x}(x) = 0. \quad (7.47)$$

Introducing the polynomial  $v(x)$  according to the rule  $v(x) = p_2(x) - p_1(x)$ , we substitute the relation  $p_2(x) = p_1(x) + v(x)$  into equation (7.47). Integrating the result with respect to the polynomial  $p_1(x)$ , we obtain

$$\begin{aligned} d_2 = -2 - d_1 : \quad p_1(x) &= -(d_1 + 1)v(x) \ln v(x) + \beta v(x); \\ d_2 \neq -2 - d_1 : \quad p_1(x) &= \beta v^{-d_2 - d_1 - 1}(x) - \frac{d_1 + 1}{d_2 + d_1 + 2}v(x), \end{aligned} \quad (7.48)$$

where  $\beta \in \mathbb{C}$  is a constant of integration. Analyzing the first possibility, we need to set  $d_1 = -1$ . As a result, we get Darboux integrable family (7.17) of Liénard differential systems. The parameter  $\beta$  can be derived with the help of the dominant behavior of the polynomials  $p_1(x)$  and  $p_2(x)$ , which now coincide with  $q_1(x)$  and  $q_2(x)$ , respectively. Recalling the fact that the product of an integrating factor and a first integral is again an integrating factor, we obtain the family of integrating factors  $M_0(x, y)I^\varkappa(x, y)$ , where  $M_0(x, y) = [y - q_1(x)]^{-1}[y - q_2(x)]^{-1}$ ,  $\varkappa \in \mathbb{C}$ , and the Darboux first integral  $I(x, y)$  is given by expression (7.18).

Now we turn to the case  $d_2 \neq -2 - d_1$ . If  $\beta = 0$ , then we get the equality  $p_2(x) = -(d_2 + 1)p_1(x)/(d_1 + 1)$ . In addition, we obtain the following representations of the polynomials  $f(x)$  and  $g(x)$ :  $f(x) = (d_2 - d_1)p_{1,x}(x)/(d_1 + 1)$  and  $g(x) = -(d_2 + 1)p_1(x)p_{1,x}(x)/(d_1 + 1)$ . Considering these expressions, we again arrive at Darboux integrable systems (7.17).

Thus, it is without loss of generality to set  $\beta \neq 0$ . Recalling the restriction  $d_2 \neq -1 - d_1$ , we introduce relatively prime both non-unit natural numbers  $k$  and  $l$  satisfying the condition  $d_2 + d_1 + 1 = -k/l$ . Analyzing expression (7.48), we conclude that there exists a polynomial  $u(x)$  such that the following relation  $v(x) = u^l(x)$  holds. In this way we express the polynomials  $p_1(x)$  and  $p_2(x)$  via the polynomial  $u(x)$ . The result is given in expression (7.44). By construction the degree of the polynomial  $u(x)$  equals  $(m + 1)/\max\{k, l\}$ .  $\square$

Using integrating factor (7.42), we find the following expression of a Liouvillian first integral

$$I(x, y) = \frac{p_2(x)B\left(\frac{y-p_1(x)}{p_2(x)-p_1(x)}; 1 + d_1, -d_1 - \frac{k}{l}\right)}{\{p_2(x) - p_1(x)\}^{\frac{k}{l}}} - \frac{B\left(\frac{y-p_1(x)}{p_2(x)-p_1(x)}; 1 + d_1, 1 - d_1 - \frac{k}{l}\right)}{\{p_2(x) - p_1(x)\}^{\frac{k}{l}-1}} \quad (7.49)$$

of systems (7.43). The polynomials  $p_1(x)$  and  $p_2(x)$  are given by relation (7.44). Symbol  $B(s; \alpha, \delta)$  denotes the incomplete beta function

$$B(s; \alpha, \delta) = \int_0^s z^{\alpha-1}(1-z)^{\delta-1} dz. \quad (7.50)$$

The family of systems (7.43) can be transformed to the following simple form

$$\begin{aligned} s_\tau = z, \quad z_\tau = & \left[ \beta(l+k)s^{k-1} + \frac{\{(2d_1+1)l+k\}l}{k-l}s^{l-1} \right] z \\ & - \left[ l\beta^2 s^{2k-1} + \frac{\{(2d_1+1)l+k\}l\beta}{k-l}s^{k+l-1} + \frac{(ld_1+k)(d_1+1)l^2}{(k-l)^2}s^{2l-1} \right] \end{aligned} \quad (7.51)$$

via the generalized Sundman transformation  $s(\tau) = u(x)$ ,  $z(\tau) = y$ ,  $d\tau = u_x(x)dt$ . Substituting  $u(x) = s$ ,  $y = z$  into (7.49) and (7.44), we find a Liouvillian first integral for systems (7.51).

The careful examination of expression (7.44) shows that systems (7.43) are resonant near infinity whenever the following inequality  $k > l$  is valid. Indeed, two distinct polynomial solutions of equation (1.4) have the coinciding dominant behavior near the point  $x = \infty$  only in a resonant case. Let us find the necessary and sufficient conditions of the Liouvillian integrability in the non-resonant case.

**Theorem 7.4.** *A Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta/f_0 \notin \mathbb{Q}$  is Liouvillian integrable if and only if the system is either Darboux integrable and reads as (7.17) or takes the form (7.43), where  $k < l$  and the following normalization*

$$d_1 = \frac{(k-l)f_0}{2l\delta} - \frac{l+k}{2l}, \quad u_0 = \left( -\frac{\delta}{m+1} \right)^{\frac{1}{7}} \quad (7.52)$$

is introduced. By  $u_0$  we denote the highest-degree coefficient of the polynomial  $u(x)$ . In the case of systems (7.43) the Darboux integrating factor is the following

$$M(x, y) = \{y - q_1(x)\}^{\frac{(k-l)f_0}{2l\delta} - \frac{l+k}{2l}} \{y - q_2(x)\}^{\frac{(l-k)f_0}{2l\delta} - \frac{l+k}{2l}} \quad (7.53)$$

with the polynomials  $q_j(x) \equiv p_j(x)$ ,  $j = 1, 2$  given by expression (7.44).

*Proof.* By direct computations we verify that systems (7.17) and (7.43) are Liouvillian integrable provided that all other conditions of the theorem are satisfied. This observation proves the sufficiency of these conditions.

Let us prove their necessity. It follows from Lemmas 5.1 and 7.1 that exponential invariants cannot arise in a Darboux integrating factor. Consequently, a Darboux integrating factor is constructed from generating polynomials of invariant algebraic curves. It is straightforward to see that if there are no invariant algebraic curves, then Darboux integrating factors do not exist. In view of Theorem 4.3 we need to consider three distinct cases.

*Case 1.* Let us suppose that a Liouvillian integrable Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta/f_0 \notin \mathbb{Q}$  has only one irreducible invariant algebraic curve with a generating polynomial of the first degree with respect to  $y$ . This curve reads as  $y - q_k(x) = 0$ ,  $k = 1$  or  $k = 2$ , and has the cofactor  $\lambda(x, y) = -f(x) - q_{k,x}(x)$ . A Darboux integrating factor can be represented in the form

$$M(x, y) = [y - q_k(x)]^{d_k}, \quad d_k \in \mathbb{C} \setminus \{0\}. \quad (7.54)$$

This integrating factor exists if and only if the following condition

$$d_k\{f(x) + q_{k,x}(x)\} + f(x) = 0 \quad (7.55)$$

is identically valid. Balancing the terms at  $x^m$  in this relation, we obtain

$$k = 1 : \quad d_1 = -\frac{2f_0}{\delta + f_0}; \quad k = 2 : \quad d_2 = \frac{2f_0}{\delta - f_0}. \quad (7.56)$$

Expressing  $f(x)$  and  $g(x)$  from relation (7.55) and equation (1.4) with  $y(x) = q_k(x)$ , we see that our Liénard differential system is of the form (7.17) and possesses two distinct irreducible invariant algebraic curves. It is a contradiction.

*Case 2.* Let us suppose that a Liouvillian integrable Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta/f_0 \notin \mathbb{Q}$  has an irreducible invariant algebraic curve with a generating polynomial of the second degree with respect to  $y$ . This curve is given by the expression  $\{[y - y_\infty^{(1)}(x)][y - y_\infty^{(2)}(x)]\}_+ = 0$ . Its cofactor reads as  $\lambda(x, y) = -2f(x) - q_{1,x}(x) - q_{2,x}(x)$ . A Darboux integrating factor can be represented in the form

$$M(x, y) = F^d(x, y), \quad F(x, y) = \{[y - y_\infty^{(1)}(x)][y - y_\infty^{(2)}(x)]\}_+, \quad d \in \mathbb{C} \setminus \{0\} \quad (7.57)$$

and exists if and only if the following condition

$$d\{2f(x) + q_{1,x}(x) + q_{2,x}(x)\} + f(x) = 0 \quad (7.58)$$

is identically satisfied. Considering the coefficients of  $x^m$  in this condition yields the value of  $d$ :  $d = -1$ . Further, we represent the polynomial  $F(x, y)$  in the form  $F(x, y) = y^2 + v(x)y + w(x)$ , where  $v(x), w(x) \in \mathbb{C}[x]$ . Substituting expression  $M(x, y) = F^{-1}(x, y)$  into the partial differential equation

$$yM_x - [f(x)y + g(x)]M_y - f(x)M = 0, \quad (7.59)$$

we get rid of the denominator. Setting to zero the coefficients of different powers of  $y$ , we get  $w(x) = \beta v(x)^2$ , where  $\beta \in \mathbb{C} \setminus \{0\}$ . This equality contradicts irreducibility of the polynomial  $F(x, y)$ .

*Case 3.* Now we assume that a Liouvillian integrable Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta/f_0 \notin \mathbb{Q}$  has two distinct irreducible invariant algebraic curves  $y - q_1(x) = 0$  and  $y - q_2(x) = 0$ . Their cofactors are the following  $\lambda_1(x, y) = -f(x) - q_{1,x}(x)$  and  $\lambda_2(x, y) = -f(x) - q_{2,x}(x)$ , respectively. Condition (2.7) enabling the existence of a Darboux integrating factor

$$M(x, y) = [y - q_1(x)]^{d_1}[y - q_2(x)]^{d_2}, \quad d_1, d_2 \in \mathbb{C}, \quad |d_1| + |d_2| > 0 \quad (7.60)$$

is of the form

$$d_1\{f(x) + q_{1,x}(x)\} + d_2\{f(x) + q_{2,x}(x)\} + f(x) = 0. \quad (7.61)$$

All the Liénard differential systems from family (B) with integrating factor of the form (7.60) have been identified in Theorem 7.3. We need to extract non-resonant systems from those given by expression (7.43). Thus, the polynomials  $q_1(x)$  and  $q_2(x)$  necessarily have distinct dominant terms. This fact yields the inequality  $k < l$ . Finally, we need to introduce the normalization adopted at the beginning of this section. Using relations (7.3) and (7.4), we obtain expression (7.52).  $\square$

*Corollary.* Liénard differential systems (7.29) with a non-autonomous Darboux first integral (7.30) are Liouvillian integrable. These systems have the Darboux integrating factor

$$M(x, y) = \frac{\left[ y + \frac{\delta + f_0}{2(m+1)}(x - x_0)^{m+1} + \frac{\omega}{2(m+1)\delta}(x - x_0) \right]^{\frac{mf_0 - (m+2)\delta}{2(m+1)\delta}}}{\left[ y + \frac{f_0 - \delta}{2(m+1)}(x - x_0)^{m+1} + \frac{\omega}{2(m+1)\delta}(x - x_0) \right]^{\frac{mf_0 + (m+2)\delta}{2(m+1)\delta}}}. \quad (7.62)$$

*Proof.* We establish the validity of the statement substituting relations

$$k = 1, l = m + 1, u(x) = u_0(x - x_0), \beta = -\frac{\omega}{2(m+1)\delta u_0}, u_0 = \left\{ -\frac{\delta}{m+1} \right\}^{\frac{1}{m+1}} \quad (7.63)$$

into expressions (7.43) and (7.53). In addition, we recall that the parameter  $d_1$  reads as (7.52).  $\square$

**Theorem 7.5.** *A Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta = 0$  is Liouvillian integrable if and only if the system has the irreducible invariant algebraic curve  $y - q(x) = 0$  and an exponential invariant  $E(x, y) = \exp[u(x)/(y - q(x))]$  such that one of the following assertions is valid.*

1. *The system is Darboux integrable and takes the form (7.25). A related Darboux integrating factor reads as*

$$M(x, y) = \frac{1}{[y - q(x)]^2} \quad (7.64)$$

*The polynomial  $u(x)$  arising in the exponential invariant is  $u(x) = \alpha q(x)$ ,  $\alpha \in \mathbb{C} \setminus \{0\}$ .*

2. *The system is of the form*

$$\begin{aligned} x_t = y, \quad y_t = - \left[ \frac{2l^2}{l-k} v^{l-1} - (l+k)\beta v^{k-1} \right] v_x y \\ + \left[ \frac{2l^2\beta}{l-k} v^{l+k-1} - \frac{l^3}{(l-k)^2} v^{2l-1} - l\beta^2 v^{2k-1} \right] v_x, \end{aligned} \quad (7.65)$$

*where  $\beta \in \mathbb{C} \setminus \{0\}$ ,  $v(x)$  is a non-constant polynomial,  $k$  and  $l$  are relatively prime natural numbers satisfying the inequality  $k < l$ . The associated Darboux integrating factor reads as*

$$M(x, y) = [y - q(x)]^{-\frac{l+k}{l}} \exp \left[ \frac{v^l(x)}{y - q(x)} \right], \quad q(x) = -\frac{l}{l-k} v^l + \beta v^k. \quad (7.66)$$

In addition, the following relation  $m + 1 = l \deg v$  is valid.

*Proof.* It is straightforward to verify that expressions (7.64) and (7.66) are Darboux integrating factors of Liénard differential systems (1.1) satisfying the restrictions  $\deg g = 2 \deg f + 1$  and  $\delta = 0$  whenever all other conditions of the theorem are satisfied.

Let us prove the converse statement. Suppose we consider a Liouvillian integrable Liénard differential system (1.1) such that  $\deg g = 2 \deg f + 1$  and  $\delta = 0$ . By Theorems 2.2, 4.3 and Lemmas 5.1, 7.1 the system has the irreducible invariant algebraic curve  $y - q(x) = 0$  and a Darboux integrating factor that can be represented in the form

$$M(x, y) = [y - q(x)]^d \exp \left[ \frac{u(x)}{y - q(x)} \right], \quad d \in \mathbb{C}, \quad (7.67)$$

where we suppose that  $u(x) \equiv 0$  whenever the exponential invariant  $E(x, y) = \exp[u(x)/(y - q(x))]$  related to the invariant algebraic curve  $y - q(x) = 0$  either does not exist or is not involved into an explicit expression of the integrating factor. Condition (2.7) with  $\omega = 0$  now takes the form

$$d[f(x) + q_x(x)] - u_x(x) + f(x) = 0. \quad (7.68)$$

Further, we shall consider several distinct cases separately.

*Case 1.* Let us begin with the case  $u(x) \equiv 0$ . Substituting the asymptotic relations  $f(x) = f_0 x^m + o(x^m)$  and  $q(x) = -f_0 x^{m+1}/(2\{m+1\}) + o(x^{m+1})$ ,  $x \rightarrow \infty$  into condition (7.68) and setting to zero the coefficient of  $x^m$ , we obtain the equalities  $d = -2$  and  $f(x) = -2q_x(x)$ . Recalling the fact that  $y = q(x)$  is the polynomial solution of equation (1.4), we find the polynomial  $g(x)$  as given in (7.25). Using Theorem 7.2, we conclude that the related Liénard differential system possesses a Darboux first integral (7.26) and the exponential invariants  $E(x, y) = \exp[\alpha q(x)/(y - q(x))]$ ,  $\alpha \in \mathbb{C} \setminus \{0\}$ . Note that integrating the differential form

$$\frac{ydy + (qq_x - 2q_x y)dx}{\{y - q(x)\}^2}. \quad (7.69)$$

yields a first integral in the form  $\ln I(x, y)$  with the function  $I(x, y)$  given by expression (7.26).

*Case 2.* Now let us suppose that the polynomial  $u(x)$  is not identically zero. We see that the system under consideration has the exponential invariant  $E(x, y) = \exp[u(x)/(y - q(x))]$  with the polynomial  $u(x)$  satisfying equation (7.5). Expressing  $f(x)$  from the latter equation and substituting the result into condition (7.68) yields the relation

$$(d + 1)qu_x + q_x u + uu_x = 0. \quad (7.70)$$

This relation viewed as an ordinary differential equation with respect to the polynomial  $q(x)$  can be integrated. Thus, we find the expressions

$$d \neq -2: \quad q(x) = -\frac{1}{d+2}u + \beta u^{-(d+1)}; \quad d = -2: \quad q(x) = (\beta - \ln u)u, \quad (7.71)$$

where  $\beta \in \mathbb{C}$  is a constant of integration. If  $d = -2$ , then  $q(x)$  is not a polynomial. Further, we set  $d \neq -2$ . The case  $\beta = 0$  again leads to a Darboux integrable family of Liénard differential systems given in Theorem 7.2. Thus, we suppose that the constant  $\beta$  is non-zero. Recalling the fact that  $q(x)$  and  $u(x)$  are polynomials with the dominant behavior  $q(x) = -f_0x^{m+1}/(2\{m+1\}) + o(x^{m+1})$  and  $u(x) = u_0x^{m+1} + o(x^{m+1})$ ,  $u_0 \in \mathbb{C} \setminus \{0\}$  near the point  $x = \infty$ , we find two possibilities

$$d = -1 : \quad u(x) = \beta - q(x); \quad d = -\frac{l+k}{l} : \quad u(x) = v^l(x). \quad (7.72)$$

In this expressions  $l$  and  $k$  are relatively prime natural numbers satisfying the restriction  $k < l$  and  $v(x)$  is a non-constant polynomial. Analyzing the possibility  $d = -1$ , we substitute the equality  $u(x) = \beta - q(x)$  into relation (7.5) and find the polynomial  $f(x)$ . The result is

$$f(x) = \frac{(2q(x) - \beta)q_x(x)}{\beta - q(x)}. \quad (7.73)$$

Let  $x_0$  be a zero of the polynomial  $\beta - q(x)$ . Considering the behavior near  $x_0$  of the rational function on the right-hand side of expression (7.73), we see that  $f(x)$  is not a polynomial whenever  $\beta \neq 0$ .

Finally, we suppose that the following relations  $d = -(l+k)/l$  and  $u(x) = v^l(x)$  are valid. We use expressions (7.71), (7.68), and (1.4) to find the polynomials  $f(x)$ ,  $g(x)$ , and  $q(x)$  as given in relations (7.65) and (7.66). In addition, we verify that equation (7.5) is identically satisfied. The proof is completed.  $\square$

*Corollary 1.* Liénard differential systems (7.36) possessing a non-autonomous Darboux first integral (7.37) are Liouvillian integrable with the Darboux integrating factor

$$M(x, y) = \exp \left[ \frac{mf_0(x-x_0)^{m+1}}{(m+1)\{2(m+1)y + f_0(x-x_0)^{m+1} + 2\omega(x-x_0)\}} \right] \times \left[ y + \frac{f_0(x-x_0)^{m+1}}{2(m+1)} + \frac{\omega(x-x_0)}{m+1} \right]^{-\frac{m+2}{m+1}}. \quad (7.74)$$

*Proof.* We prove the validity of the statement substituting relations

$$k = 1, \quad l = m + 1, \quad v(x) = v_0(x - x_0), \quad \beta = -\frac{\omega}{(m+1)v_0}, \quad v_0 = \left\{ \frac{mf_0}{2(m+1)^2} \right\}^{\frac{1}{m+1}} \quad (7.75)$$

into expressions (7.65) and (7.66).  $\square$

*Corollary 2.* If the following inequality  $k > l$  holds, then systems (7.65) are also Liouvillian integrable Liénard differential systems from family (B). But these systems are resonant near infinity. The related Darboux integrating factor again is given by expression (7.66).

A Liouvillian first integral produced by integrating factor (7.66) reads as

$$I(x, y) = v^{l-k}(x) \gamma \left( -\frac{l-k}{l}, \frac{v^l(x)}{q(x)-y} \right) - \frac{q(x)}{v^k(x)} \gamma \left( \frac{k}{l}, \frac{v^l(x)}{q(x)-y} \right), \quad (7.76)$$

where the polynomial  $q(x)$  is given in expression (7.65) and  $\gamma(\delta, s)$  is the lower incomplete Gamma function

$$\gamma(\delta, s) = \int_0^s t^{\delta-1} \exp(-t) dt. \quad (7.77)$$

Note that we need to consider the analytic continuation of this integral for complex or real non-positive values of  $s$ . If  $k = 1$  and  $l = 2$ , then we obtain another representation of a Liouvillian first integral

$$I(x, y) = 2\sqrt{\beta v(x) - 2v^2(x) - y} \exp \left[ \frac{v^2(x)}{y + 2v^2(x) - \beta v(x)} \right] - \sqrt{\pi} \beta \operatorname{erfc} \left[ \frac{v(x)}{\sqrt{\beta v(x) - 2v^2(x) - y}} \right], \quad (7.78)$$

where  $\operatorname{erfc}(s)$  is the complementary error function

$$\operatorname{erfc}(s) = \frac{2}{\sqrt{\pi}} \int_s^\infty \exp(-t^2) dt. \quad (7.79)$$

The family of systems (7.65) can be transformed to the following simple form

$$s_\tau = z, \quad z_\tau = - \left[ \frac{2l^2}{l-k} s^{l-1} - (l+k)\beta s^{k-1} \right] z + \frac{2l^2\beta}{l-k} s^{l+k-1} - \frac{l^3}{(l-k)^2} s^{2l-1} - l\beta^2 s^{2k-1} \quad (7.80)$$

via the generalized Sundman transformation  $s(\tau) = v(x)$ ,  $z(\tau) = y$ ,  $d\tau = v_x(x)dt$ . Substituting  $v(x) = s$ ,  $y = z$  into (7.76), we find a Liouvillian first integral for systems (7.80).

It seems that Liouvillian integrable families of Liénard differential systems given in Theorems 7.3, 7.4, and 7.5 are new with the exception of systems (7.36). The latter are presented by T. Stachowiak [44].

Now let us investigate the existence of Jacobi last multipliers with a time-dependent exponential factor.

**Lemma 7.4.** *A Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta/f_0 \notin \mathbb{Q}$  has a non-autonomous Darboux–Jacobi last multiplier of the form (2.6) if and only if one the following assertions is valid.*

1. *The system under consideration possesses one irreducible invariant algebraic curve  $y - q_k(x) = 0$ , where  $k = 1$  or  $k = 2$ , such that the polynomials  $f(x)$  and  $q_k(x)$  identically satisfy the condition*

$$\begin{aligned} k = 1 : & (f_0 - \delta)f(x) + 2f_0q_{1,x}(x) + (f_0 + \delta)\omega = 0, \quad \omega \in \mathbb{C} \setminus \{0\}, \\ k = 2 : & (f_0 + \delta)f(x) + 2f_0q_{2,x}(x) + (f_0 - \delta)\omega = 0, \quad \omega \in \mathbb{C} \setminus \{0\}. \end{aligned} \quad (7.81)$$

A related Darboux–Jacobi last multiplier reads as

$$\begin{aligned} k = 1 : \quad M(x, y, t) &= [y - q_1(x)]^{-\frac{2f_0}{\delta+f_0}} \exp(\omega t), \\ k = 2 : \quad M(x, y, t) &= [y - q_2(x)]^{\frac{2f_0}{\delta-f_0}} \exp(\omega t). \end{aligned} \quad (7.82)$$

2. The system under consideration possesses one irreducible invariant algebraic curve  $F(x, y) = 0$  with  $F(x, y) = \{[y - y_\infty^{(1)}(x)] [y - y_\infty^{(2)}(x)]\}_+$  such that the polynomials  $f(x)$ ,  $q_1(x) = \{y_\infty^{(1)}(x)\}_+$ , and  $q_2(x) = \{y_\infty^{(2)}(x)\}_+$  identically satisfy the condition

$$f(x) + q_{1,x}(x) + q_{2,x}(x) + \omega = 0, \quad \omega \in \mathbb{C} \setminus \{0\}. \quad (7.83)$$

A related Darboux–Jacobi last multiplier reads as

$$M(x, y, t) = \frac{\exp(\omega t)}{F(x, y)}, \quad F(x, y) = \{[y - y_\infty^{(1)}(x)] [y - y_\infty^{(2)}(x)]\}_+. \quad (7.84)$$

3. The system under consideration possesses two distinct irreducible invariant algebraic curves  $y - q_1(x) = 0$  and  $y - q_2(x) = 0$  such that the polynomials  $f(x)$ ,  $q_1(x)$ , and  $q_2(x)$  identically satisfy the condition

$$\begin{aligned} [(2d_2 + 1)\delta - f_0]f(x) + [(\delta - f_0)d_2 - 2f_0]q_{1,x}(x) + (\delta + f_0)d_2q_{2,x}(x) \\ - (\delta + f_0)\omega = 0, \quad d_2 \in \mathbb{C}, \quad \omega \in \mathbb{C} \setminus \{0\}. \end{aligned} \quad (7.85)$$

A related Darboux–Jacobi last multiplier reads as

$$M(x, y, t) = [y - q_1(x)]^{\frac{(\delta-f_0)d_2-2f_0}{\delta+f_0}} [y - q_2(x)]^{d_2} \exp(\omega t). \quad (7.86)$$

This lemma is proved similarly to Theorem 7.4.

**Lemma 7.5.** A Liénard differential system (1.1) satisfying the conditions  $\deg g = 2 \deg f + 1$  and  $\delta = 0$  has a non-autonomous Darboux–Jacobi last multiplier of the form (2.6) if and only if one the following assertions is valid.

1. The system under consideration possesses the irreducible invariant algebraic curve  $y - q(x) = 0$  such that the polynomials  $f(x)$  and  $g(x)$  can be represented as

$$f(x) = -2q_x(x) - \omega, \quad g(x) = q(x)(q_x(x) + \omega), \quad \omega \in \mathbb{C} \setminus \{0\}. \quad (7.87)$$

A related Darboux–Jacobi last multiplier reads as

$$M(x, y, t) = \frac{\exp(\omega t)}{[y - q(x)]^2}. \quad (7.88)$$

2. The system under consideration possesses the irreducible invariant algebraic curve  $y - q(x) = 0$  and the exponential invariant  $E(x, y) = \exp[u(x)/(y - q(x))]$  such that the polynomials  $f(x)$ ,  $q(x)$ , and  $u(x)$  identically satisfy the condition

$$(d + 1)f(x) + dq_x(x) = u_x(x) + \omega, \quad d \in \mathbb{C}, \quad \omega \in \mathbb{C} \setminus \{0\}. \quad (7.89)$$

A related Darboux–Jacobi last multiplier reads as

$$M(x, y, t) = [y - q(x)]^d \exp \left[ \frac{u(x)}{y - q(x)} \right] \exp(\omega t). \quad (7.90)$$

The proof of this lemma is analogous to the proof of Theorem 7.5.

Concluding this section let us note that autonomous and non-autonomous Darboux first integrals of non-resonant Liénard differential systems (1.1) satisfying the conditions  $\deg f = 1$ ,  $\deg g = 3$  and  $\deg f = 2$ ,  $\deg g = 5$  are classified in [23, 24].

## 8. Integrability of Liénard differential systems from family (C)

We start investigating integrability properties of Liénard differential systems from family (C) by proving the absence of exponential invariants related to invariant algebraic curves. We use designations of Theorem 4.4. In particular, by  $h^{(1)}(x)$  and  $h^{(2)}(x)$  we denote the initial parts of the Puiseux series  $y_\infty^{(1)}(x)$  and  $y_\infty^{(2)}(x)$ , accordingly. These initial parts involve monomials with exponents exceeding  $-(n + 1)/2$ . Recall that we use the designations  $\deg g = n$  and  $\deg f = m$ . Thus, the following inequality  $n > 2m + 1$  is valid.

**Lemma 8.1.** *Suppose a Liénard differential system (1.1) from family (C) is not integrable with a rational first integral. Then this system does not have exponential invariants of the form  $E(x, y) = \exp \{h(x, y)/r(x, y)\}$ , where  $h(x, y) \in \mathbb{C}[x, y]$  and  $r(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}$  are relatively prime polynomials.*

*Proof.* We shall use the local theory presented in Section 3. If a Liénard differential system (1.1) has an exponential invariant  $E(x, y) = \exp \{h(x, y)/r(x, y)\}$  with  $r(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}$ , then  $r(x, y) \in \mathbb{C}[x, y] \setminus \mathbb{C}[x]$  and it is without loss of generality to assume that the degree of the polynomial  $h(x, y)$  with respect to  $y$  is less than the degree of the polynomial  $r(x, y)$  with respect to  $y$ . There exists a finite number of local elementary exponential invariants

$$E_j(x, y) = \exp \left[ \frac{u_j(x)}{\{y - Y_{j,\infty}(x)\}^{n_j}} \right], \quad u_j(x), Y_{j,\infty}(x) \in \mathbb{C}_\infty\{x\}, \quad (8.1)$$

$$n_j \in \mathbb{N}, \quad j = 1, \dots, K, \quad K \in \mathbb{N}$$

such that the exponential invariant  $E(x, y)$  equals the product  $E_1(x, y) \times \dots \times E_K(x, y)$ . It follows from Theorem 3.3 and Lemma 3.2 that each series  $Y_{j,\infty}(x)$  satisfies equation (1.4). In fact, the series  $Y_{j,\infty}(x)$  coincides with one of the series  $y_{j,\infty}^{(1,2)}(x)$  presented in Theorem 4.4. Let us denote the cofactor of the local elementary exponential invariant  $E_j(x, y)$  by  $\varrho_j(x, y) \in \mathbb{C}_\infty\{x\}[y]$ . We see that the following expression  $\varrho_1(x, y) + \dots + \varrho_K(x, y)$

equals the cofactor  $\varrho(x, y)$  of the invariant  $E(x, y)$ . Recall that the cofactor  $\varrho(x, y)$  is an element of the ring  $\mathbb{C}[x, y]$ .

Substituting the explicit representation of  $E_j(x, y)$  into the partial differential equation  $\mathcal{X}E_j(x, y) = \varrho_j(x, y)E_j(x, y)$ , we get

$$yu_{j,x} = n_j\lambda_j(x, y)u_j(x) + \varrho_j(x, y)\{y - Y_{j,\infty}(x)\}^{n_j}. \quad (8.2)$$

In this expression  $\lambda_j(x, y) \in \mathbb{C}_\infty\{x\}[y]$  is the cofactor of the local elementary invariant  $F_j(x, y) = y - Y_{j,\infty}(x)$ . Using Theorem 3.4, we find the cofactor  $\lambda_j(x, y)$ . The result is

$$\lambda_j(x, y) = -f(x) - \{Y_{j,\infty}(x)\}_x. \quad (8.3)$$

Analyzing expression (8.2), we see that  $n_j = 1$ ,  $\varrho_j(x, y) = u_{j,x}(x)$ , and the series  $u_j(x)$  satisfies the following ordinary differential equation

$$Y_{j,\infty}(x)u_{j,x}(x) + (f(x) + [Y_{j,\infty}(x)]_x)u_j(x) = 0. \quad (8.4)$$

Using the dominant behavior of the series  $Y_{j,\infty}(x)$  given by the monomial  $b_{0,j}x^{(n+1)/2}$ , we find the dominant monomial of the series  $u_j(x)$ . Thus, we get the relation  $u_j(x) = \sigma_j x^{-(n+1)/2} + o(x^{-(n+1)/2})$ , where  $\sigma_j \in \mathbb{C}$  and  $x \rightarrow \infty$ . Hence the cofactor  $\varrho_j(x, y) = u_{j,x}(x)$  does not have monomials with non-negative exponents. Consequently, the polynomial  $\varrho(x, y) = \varrho_1(x, y) + \dots + \varrho_K(x, y)$  should be identically zero. We conclude that the argument  $h(x, y)/r(x, y)$  of the exponential invariant  $E(x, y) = \exp\{h(x, y)/r(x, y)\}$  is a rational first integral of the differential system in question. It is a contradiction.  $\square$

*Remark.* It will be shown below that Liénard differential systems (1.1) satisfying the condition  $\deg g > 2 \deg f + 1$  do not have rational first integrals. Thus, this condition can be removed from the statement of Lemma 8.1.

Now our aim is to prove that Liénard differential systems (1.1) do not have Darboux first integrals provided that  $\deg g > 2 \deg f + 1$ .

**Theorem 8.1.** *Liénard differential systems (1.1) from family (C) are not Darboux integrable.*

*Proof.* Let us suppose that a Liénard differential system (1.1) with  $\deg g > 2 \deg f + 1$  has a Darboux first integral. In view of Theorem 2.1 the system possesses a rational integrating factor. Consequently, there exists  $K \in \mathbb{N}$  pairwise distinct irreducible invariant algebraic curves  $F_1(x, y) = 0, \dots, F_K(x, y) = 0$  and  $K$  non-zero integer numbers  $d_1, \dots, d_K$  such that the following condition

$$\sum_{j=1}^K d_j \lambda_j(x, y) = f(x), \quad d_1, \dots, d_K \in \mathbb{Z} \quad (8.5)$$

is valid. In this expression  $\lambda_j(x, y)$  is the cofactor of the invariant algebraic curve  $F_j(x, y) = 0$ . Suppose the family of Puiseux series  $y_\infty^{(l)}(x)$  arises  $N_{l,j}$  times in the factorization of the

polynomial  $F_j(x, y)$  in the ring  $\mathbb{C}_\infty\{x\}[y]$ . Here  $l = 1, 2$  and we use the designations of Theorem 4.4. The cofactor  $\lambda_j(x, y)$  reads as

$$\lambda_j(x, y) = -(N_{1,j} + N_{2,j})f(x) - \{N_{1,j}h_x^{(1)}(x) + N_{2,j}h_x^{(2)}(x)\}_+, \quad (8.6)$$

$$N_{1,j}, N_{2,j} \in \mathbb{N}_0, \quad N_{1,j} + N_{2,j} > 0,$$

where  $h^{(l)}(x)$  is the initial part of Puiseux series  $y_\infty^{(l)}(x)$  introduced in Theorem 4.4. Substituting expression (8.6) into relation (8.5) yields

$$\sum_{j=1}^K d_j N_{1,j} \left( f(x) + \{h_x^{(1)}(x)\}_+ \right) + \sum_{j=1}^K d_j N_{2,j} \left( f(x) + \{h_x^{(2)}(x)\}_+ \right) = -f(x). \quad (8.7)$$

Let us suppose that  $n$  is odd. The dominant behavior of the truncated series  $h^{(1)}(x)$  and  $h^{(2)}(x)$  is  $b_0x^{(n+1)/2}$  and  $-b_0x^{(n+1)/2}$ , respectively. Setting to zero the coefficient of  $x^{(n-1)/2}$  in expression (8.7), we obtain

$$\sum_{j=1}^K d_j N_{1,j} = \sum_{j=1}^K d_j N_{2,j}. \quad (8.8)$$

If  $n$  is even, then by Theorem 4.4 we get  $N_{1,j} = N_{2,j}$ . Consequently, relation (8.8) is identically satisfied. As a result, condition (8.7) takes the form

$$B \left( 2f(x) + \{h_x^{(1)}(x) + h_x^{(2)}(x)\}_+ \right) = -f(x), \quad B = \sum_{j=1}^K d_j N_{1,j}. \quad (8.9)$$

Obviously  $B$  is a non-zero integer number. Now let us consider the equation

$$yy_x + \varepsilon f(x)y + g(x) = 0. \quad (8.10)$$

Puiseux series near the point  $x = \infty$  that satisfy this equation coincide with the Puiseux series  $y_{j,\infty}^{(1,2)}(x)$  presented in Theorem 4.4 provided that  $\varepsilon = 1$ . Analogously to the case  $\varepsilon = 1$ , we find two families of Puiseux series near the point  $x = \infty$  solving equation (8.10). We denote these families as  $Y_\infty^{(1)}(x)$  and  $Y_\infty^{(2)}(x)$ . It is straightforward to see that these series can be represented as

$$Y_\infty^{(1,2)}(x) = \sum_{k=0}^{\infty} \varepsilon^k v_k^{(1,2)}(x), \quad (8.11)$$

where the coefficients  $v_k^{(1,2)}(x)$  are Puiseux series near the point  $x = \infty$ . Substituting expression (8.11) into equation (8.10) and setting to zero the coefficients of different powers of  $\varepsilon$ , we find ordinary differential equations for the series  $v_k^{(1,2)}(x)$ . Two first equations take the form

$$v_0 v_{0,x} + g(x) = 0, \quad v_0 v_{1,x} + v_{0,x} v_1 + f(x) v_0 = 0, \quad (8.12)$$

where the upper index is omitted. Thus, we get

$$v_0^{(2)}(x) = -v_0^{(1)}(x), \quad v_1^{(1,2)}(x) = -\frac{2f_0}{2m+n+3}x^{m+1} + o(x^{m+1}), \quad x \rightarrow \infty. \quad (8.13)$$

Analyzing other ordinary differential equations for the series  $v_k^{(1,2)}(x)$  with  $k \geq 2$ , we find  $v_k^{(1,2)}(x) = o(x^{m+1})$ ,  $x \rightarrow \infty$ ,  $k \geq 2$ . We recall that the Puiseux series  $y_{j,\infty}^{(1,2)}(x)$  with the same upper index have coinciding initial parts involving monomials with exponents exceeding  $-(n+1)/2$ . These initial parts are given by  $h^{(1)}(x)$  and  $h^{(2)}(x)$ . Substituting our results into condition (8.9) and considering the coefficients of the leading term  $x^m$ , we come to the equation

$$B \left( 2f_0 - \frac{4(m+1)f_0}{2m+n+3} \right) = -f_0. \quad (8.14)$$

Solving this equation, we find the expression

$$B = -\frac{m+1}{n+1} - \frac{1}{2}. \quad (8.15)$$

Inequality  $n > 2m + 1$  shows that  $B$  is not an integer. It is a contradiction.  $\square$

*Corollary 1.* A Liénard differential system (1.1) from family (C) has at most one invariant algebraic curve  $F(x, y) = 0$  with the property  $N_1 = N_2$ .

*Proof.* Recall that the variables  $N_1$  and  $N_2$  give the number of distinct Puiseux series  $y_\infty^{(1)}(x)$  and  $y_\infty^{(2)}(x)$  from the field  $\mathbb{C}_\infty\{x\}$  arising in the factorization of the polynomial producing the algebraic curve  $F(x, y) = 0$ , respectively. For more details see Theorem 4.4. Note that we do not require the polynomial  $F(x, y)$  to be irreducible. The cofactor of such an algebraic curve takes the form

$$\lambda(x, y) = -2N_1f(x) - N_1 \{h_x^{(1)}(x) + h_x^{(2)}(x)\}_+. \quad (8.16)$$

If there exists another invariant algebraic curve with the property  $N_1 = N_2$ , then the system under consideration possesses a rational first integral. This fact contradicts Theorem 8.1.  $\square$

*Corollary 2.* A Liénard differential system (1.1) from family (C) cannot have two distinct irreducible invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$  with the property  $N_{1,1}N_{2,2} - N_{1,2}N_{2,1} = 0$ , where  $N_{l,j}$  is the number of times the family of Puiseux series  $y_\infty^{(l)}(x)$  enters the factorization of the polynomial  $F_j(x, y)$  in the ring  $\mathbb{C}_\infty\{x\}[y]$ .

*Proof.* The proof is by contradiction. Suppose a Liénard differential system (1.1) satisfying the condition  $\deg g > 2 \deg f + 1$  possesses two distinct irreducible invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$  with the property  $N_{1,1}N_{2,2} - N_{1,2}N_{2,1} = 0$ .

First of all, let us assume that one of the numbers  $N_{l,j}$ , where  $l, j = 1, 2$ , is zero. Without loss of generality, we choose  $N_{2,1} = 0$ . This gives  $N_{1,1}N_{2,2} = 0$ . The polynomial  $F_j(x, y)$ ,  $j = 1, 2$  should have at least one Puiseux series near the point  $x = \infty$  in its factorization.

Thus, we get  $N_{1,1} > 0$ ,  $N_{2,2} = 0$ , and  $N_{1,2} > 0$ . By Theorem 4.4 there exists at most one irreducible invariant algebraic curve possessing only one family of Puiseux series near the point  $x = \infty$  in the factorization. This yields a contradiction.

Now suppose that all the numbers  $N_{l,j}$ , where  $l, j = 1, 2$ , are non-zero. Introducing the variable

$$\varkappa = \frac{N_{1,1}}{N_{2,1}} = \frac{N_{1,2}}{N_{2,2}}, \quad (8.17)$$

we represent the cofactors  $\lambda_1(x, y)$  and  $\lambda_2(x, y)$  of the invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$  in the form

$$\lambda_j(x, y) = -N_{2,j} \left[ (\varkappa + 1)f(x) + \{ \varkappa h_x^{(1)}(x) + h_x^{(2)}(x) \}_+ \right], \quad j = 1, 2. \quad (8.18)$$

We conclude that the cofactors are dependent over the ring  $\mathbb{Z}$ . Consequently, the Liénard differential system under study has a rational first integral. This is a contradiction.  $\square$

We see from Corollary 1 and Theorem 4.4 that if  $\deg g$  denoted as  $n$  is an even number, then a Liénard differential system (1.1) satisfying the condition  $\deg g > 2 \deg f + 1$  cannot have more than one irreducible invariant algebraic curve simultaneously.

Next, let us investigate the existence of non-autonomous Darboux first integrals. The following lemma is valid.

**Lemma 8.2.** *A Liénard differential system (1.1) from family (C) has a non-autonomous Darboux first integral with a time-dependent exponential factor (2.4) if and only if  $\deg f = 0$  and one of the following assertions is valid.*

1. *There exists an irreducible invariant algebraic curve  $F(x, y) = 0$  such that the family of Puiseux series  $y_\infty^{(1)}(x)$  arises in the factorization of the polynomial  $F(x, y)$  in the ring  $\mathbb{C}_\infty\{x\}[y]$  as many times as so does the family  $y_\infty^{(2)}(x)$ , i.e.  $N_1 = N_2$ . A first integral takes the form*

$$I(x, y, t) = F(x, y) \exp \left[ \frac{2(n+1)f_0 N_1 t}{n+3} \right]. \quad (8.19)$$

2. *There exist two distinct irreducible invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$  such that the following relation  $N_{1,j} \neq N_{2,j}$ ,  $j = 1, 2$  is valid, where  $N_{l,j}$  is the number of times the family of Puiseux series  $y_\infty^{(l)}(x)$  enters the factorization of the polynomial  $F_j(x, y)$  in the ring  $\mathbb{C}_\infty\{x\}[y]$ . A first integral reads as*

$$I(x, y, t) = \frac{\{F_1(x, y)\}^{N_{1,2}-N_{2,2}}}{\{F_2(x, y)\}^{N_{1,1}-N_{2,1}}} \exp \left[ \frac{2(n+1)f_0 \Omega t}{n+3} \right] \quad (8.20)$$

*with the parameter  $\Omega$  given by the relation  $\Omega = N_{1,2}N_{2,1} - N_{1,1}N_{2,2}$ .*

There are no other independent non-autonomous Darboux first integrals with a time-dependent exponential factor (2.4).

*Proof.* By Lemmas 5.1 and 6.1 exponential factors cannot enter an explicit expression of a non-autonomous Darboux first integral (2.4). It follows from Theorem 2.3 that a Liénard differential system (1.1) satisfying the condition  $\deg g > 2 \deg f + 1$  has a first integral (2.4) if and only if there exists  $K \in \mathbb{N}$  pairwise distinct irreducible invariant algebraic curves  $F_1(x, y) = 0, \dots, F_K(x, y) = 0$  and  $K + 1$  non-zero complex numbers  $d_1, \dots, d_K, \omega$  such that the following condition

$$\sum_{j=1}^K d_j \lambda_j(x, y) + \omega = 0, \quad d_1, \dots, d_K, \omega \in \mathbb{C} \setminus \{0\} \quad (8.21)$$

is valid. In this expression  $\lambda_j(x, y)$  is the cofactor of the invariant algebraic curve  $F_j(x, y) = 0$ . The related first integral reads as

$$I(x, y, t) = \prod_{j=1}^K F_j^{d_j}(x, y) \exp[\omega t]. \quad (8.22)$$

Suppose the family of Puiseux series  $y_\infty^{(l)}(x)$  arises  $N_{l,j}$  times in the factorization of the polynomial  $F_j(x, y)$  in the ring  $\mathbb{C}_\infty\{x\}[y]$ . Here  $l = 1, 2$  and again we use the designations of Theorem 4.4. The cofactor  $\lambda_j(x, y)$  is given by relation (8.6). The necessary and sufficient condition for first integral (8.22) to exist reads as

$$\sum_{j=1}^K d_j N_{1,j} \left( f(x) + \{h_x^{(1)}(x)\}_+ \right) + \sum_{j=1}^K d_j N_{2,j} \left( f(x) + \{h_x^{(2)}(x)\}_+ \right) = \omega. \quad (8.23)$$

This condition is not satisfied whenever relation (8.8) is not valid. Further, we rewrite condition (8.23) in the form

$$B \left( 2f(x) + \{h_x^{(1)}(x) + h_x^{(2)}(x)\}_+ \right) = \omega, \quad B = \sum_{j=1}^K d_j N_{1,j}, \quad (8.24)$$

where unlike the case of Theorem 8.1 the parameter  $B$  may be complex-valued. If  $m > 0$ , then we arrive at the expression

$$B \left( 2f_0 - \frac{4(m+1)f_0}{2m+n+3} \right) = 0, \quad (8.25)$$

which is not valid. Consequently, we should set  $m = 0$ . Relations (8.8) and (8.24) now become

$$\sum_{j=1}^K d_j N_{1,j} = \sum_{j=1}^K d_j N_{2,j}, \quad 2f_0(n+1) \sum_{j=1}^K d_j N_{1,j} = (n+3)\omega. \quad (8.26)$$

If the original Liénard differential system has only one irreducible invariant algebraic curve ( $K = 1$ ), then algebraic system (8.26) is satisfied if and only if the following relation  $N_{1,1} = N_{2,1}$  is valid. We note that the parameter  $d_1 \neq 0$  can be chosen arbitrarily. Thus, we set  $d_1 = 1$ . The second equation in (8.26) produces the value of  $\omega$ . As a result, we obtain non-autonomous Darboux first integral (8.19), where the index  $j = 1$  is omitted.

Now let us suppose that the Liénard differential system under consideration has at least two distinct irreducible invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$  satisfying the restriction  $N_{1,j} \neq N_{2,j}$ ,  $j = 1, 2$ . We see that algebraic system (8.26) is always satisfied. Indeed, setting  $K = 2$  and recalling the fact that one of the exponents  $d_1$  and  $d_2$  can be chosen arbitrary, we obtain a solution

$$d_1 = N_{1,2} - N_{2,2}, \quad d_2 = N_{2,1} - N_{1,1}, \quad \omega = \frac{2(n+1)f_0\Omega}{n+3}, \quad (8.27)$$

where we use the designation  $\Omega = N_{1,2}N_{2,1} - N_{1,1}N_{2,2}$ . Hence we have found time-dependent Darboux first integral (8.20). By Corollary 2 to Theorem 8.1, the following relation  $\Omega \neq 0$  is valid.

Suppose a Liénard differential system has two independent non-autonomous Darboux first integrals with a time-dependent exponential factor (2.4), then this system is Darboux integrable. This fact contradicts Theorem 8.1. □

*Remark.* Liénard differential systems (1.1) satisfying the conditions of item 2 can only arise when  $n = \deg g$  is an odd number.

There exist Liénard differential systems (1.1) with non-autonomous Darboux first integrals (8.19) and (8.20). An example was given in article [19]. Finally, we turn to the Liouvillian integrability.

**Theorem 8.2.** *A Liénard differential system (1.1) from family (C) is Liouvillian integrable if and only if the relation*

$$4(m+1)f(x) + (2m+n+3) \{h_x^{(1)}(x) + h_x^{(2)}(x)\}_+ = 0 \quad (8.28)$$

*is identically satisfied and one of the following assertions is valid.*

1. *There exists an irreducible invariant algebraic curve  $F(x, y) = 0$  such that the family of Puiseux series  $y_\infty^{(1)}(x)$  arises in the factorization of the polynomial  $F(x, y)$  in the ring  $\mathbb{C}_\infty\{x\}[y]$  as many times as so does the family  $y_\infty^{(2)}(x)$ , i.e.  $N_1 = N_2$ . In this case the system has the unique Darboux integrating factor*

$$M(x, y) = \{F(x, y)\}^{-\frac{2m+n+3}{2(n+1)N_1}}. \quad (8.29)$$

2. *There exist two distinct irreducible invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$  such that the following relation  $N_{1,j} \neq N_{2,j}$ ,  $j = 1, 2$  is valid, where  $N_{1,j}$  is the number*

of times the family of Puiseux series  $y_\infty^{(l)}(x)$  enters the factorization of the polynomial  $F_j(x, y)$  in the ring  $\mathbb{C}_\infty\{x\}[y]$ . In this case the system has the unique Darboux integrating factor

$$M(x, y) = \frac{\{F_1(x, y)\}^{\frac{(2m+n+3)(N_{2,2}-N_{1,2})}{2(n+1)\Omega}}}{\{F_2(x, y)\}^{\frac{(2m+n+3)(N_{2,1}-N_{1,1})}{2(n+1)\Omega}}}. \quad (8.30)$$

The following designation  $\Omega = N_{1,2}N_{2,1} - N_{1,1}N_{2,2}$  is introduced in expression (8.30).

*Proof.* It follows from Theorem 2.2 that a Liouvillian integrable differential system (2.1) has a Darboux integrating factor. By Lemmas 5.1 and 6.1 exponential invariants cannot enter an explicit expression of a Darboux integrating factor. Thus, the Darboux integrating factor reads as

$$M(x, y) = \prod_{j=1}^K F_j^{d_j}(x, y), \quad d_1, \dots, d_K \in \mathbb{C}, \quad K \in \mathbb{N}, \quad (8.31)$$

where the polynomials  $F_1(x, y), \dots, F_K(x, y)$  give pairwise distinct irreducible invariant algebraic curves  $F_1(x, y) = 0, \dots, F_K(x, y) = 0$  of a Liénard differential system. Without loss of generality, we suppose that the numbers  $d_1, \dots, d_K$  are all non-zero. The cofactor  $\lambda_j(x, y)$  of the invariant algebraic curve  $F_j(x, y) = 0$  is given by relation (8.6). The necessary and sufficient condition  $d_1\lambda_1(x, y) + \dots + d_K\lambda_K(x, y) = -\operatorname{div} \mathcal{X}$  for Darboux integrating factor (8.31) to exist now takes the form

$$\sum_{j=1}^K d_j N_{1,j} \left( f(x) + \{h_x^{(1)}(x)\}_+ \right) + \sum_{j=1}^K d_j N_{2,j} \left( f(x) + \{h_x^{(2)}(x)\}_+ \right) = -f(x). \quad (8.32)$$

This condition is not satisfied provided that relation (8.8) is not valid. Using relation (8.8), we simplify condition (8.32). Thus, we get

$$B \left( 2f(x) + \{h_x^{(1)}(x) + h_x^{(2)}(x)\}_+ \right) = -f(x), \quad B = \sum_{j=1}^K d_j N_{1,j}, \quad (8.33)$$

where unlike the case of Theorem 8.1 the parameter  $B$  may be complex-valued. Doing the same as in the proof of Theorem 8.1, we find the following equality

$$B \left( 2f_0 - \frac{4(m+1)f_0}{2m+n+3} \right) = -f_0, \quad (8.34)$$

which gives the value of  $B$ . The result is

$$B = -\frac{2m+n+3}{2(n+1)}. \quad (8.35)$$

Substituting this equality into condition (8.33) yields relation (8.28). Finally, we are left with the following algebraic system

$$\sum_{j=1}^K d_j N_{1,j} = \sum_{j=1}^K d_j N_{2,j}, \quad \sum_{j=1}^K d_j N_{1,j} = -\frac{2m+n+3}{2(n+1)} \quad (8.36)$$

with respect to the unknowns  $d_1, \dots, d_K$ .

Suppose the Liénard differential system under study has only one irreducible invariant algebraic curve ( $K = 1$ ). Algebraic system (8.36) is satisfied if and only if  $N_{1,1} = N_{2,1}$ . Omitting the index  $j$ , we find the value of  $d$  and Darboux integrating factor (8.29).

Now we assume that the Liénard differential system in question possesses at least two distinct irreducible invariant algebraic curves. Setting  $K = 2$ , we see that the determinant of algebraic system (8.36) equals  $\Omega = N_{1,2}N_{2,1} - N_{1,1}N_{2,2}$ . By Corollary 2 to Theorem 8.1 we get  $\Omega \neq 0$ . Consequently, algebraic system (8.36) has the unique solution. As a result we obtain Darboux integrating factor (8.30).

If there are no invariant algebraic curves or there exists the unique irreducible invariant algebraic curve satisfying the condition  $N_1 \neq N_2$ , then system (8.36) is inconsistent.

Suppose the Liénard differential system has two distinct Darboux integrating factors. Then their ratio is a Darboux first integral. This fact contradicts Theorem 8.1. □

*Corollary.* A Liouvillian integrable Liénard differential system (1.1) from family (C) has at most two distinct irreducible invariant algebraic curves simultaneously provided that  $n = \deg g$  is an odd number.

*Proof.* Assuming that a Liouvillian integrable Liénard differential system has three or more pairwise distinct irreducible invariant algebraic curves, we use Theorem 8.2 to find at least two distinct Darboux integrating factors. Consequently, the system possesses a Darboux first integral. It is a contradiction. □

*Remark 1.* Item 2 can only arise if the number  $n = \deg g$  is odd. Moreover, integrating factor (8.30) transforms into integrating factor (8.29) whenever the following condition  $N_{1,1} + N_{1,2} = N_{2,1} + N_{2,2}$  holds. In this case the polynomial  $F(x, y)$  in (8.29) is reducible:  $F(x, y) = F_1(x, y)F_2(x, y)$ .

*Remark 2.* Relation (8.28) is identically satisfied whenever  $m = 0$  ( $\deg f = 0$ ). This statement follows from relations (8.11) and (8.12).

Let us obtain all Liouvillian integrable Liénard differential systems (1.1) from family (C) with a hyperelliptic invariant algebraic curve  $y^2 + u(x)y + v(x) = 0$ , where  $u(x), v(x) \in \mathbb{C}[x]$ .

**Theorem 8.3.** A Liénard differential system (1.1) from family (C) with a hyperelliptic invariant algebraic curve  $y^2 + u(x)y + v(x) = 0$ , where  $u(x), v(x) \in \mathbb{C}[x]$ , is Liouvillian integrable if and only the system is of the form

$$x_t = y, \quad y_t = -\frac{(k+2l)}{4}w^{l-1}w_x y - \frac{k}{8}(w^{2l-1} + 4\beta w^{k-1})w_x, \quad (8.37)$$

where  $\beta \in \mathbb{C} \setminus \{0\}$ ,  $w(x)$  is a polynomial of degree  $(m+1)/l$ ,  $k$  and  $l$  are relatively prime natural numbers such that the following relation  $(m+1)k = (n+1)l$  is valid. The associated Liénard differential system has the unique Darboux integrating factor

$$M(x, y) = \left\{ y^2 + w^l y + \frac{1}{4}w^{2l} + \beta w^k \right\}^{-\left(\frac{1}{2} + \frac{l}{k}\right)} \quad (8.38)$$

and the hyperelliptic invariant algebraic curve reads as  $4y^2 + 4w^l y + w^{2l} + 4\beta w^k = 0$ . A Liouvillian first integral is of the form

$$I(x, y) = \frac{(2l - k)(2y + w^l)}{4kw^{\frac{k}{2}}\beta^{\frac{1}{2} + \frac{l}{k}}} {}_2F_1\left(\frac{1}{2}, \frac{1}{2} + \frac{l}{k}; \frac{3}{2}; -\frac{(2y + w^l)^2}{4\beta w^k}\right) + \left\{y^2 + w^l y + \frac{1}{4}w^{2l} + \beta w^k\right\}^{\frac{1}{2} - \frac{l}{k}}, \quad (8.39)$$

where  ${}_2F_1(\alpha, \delta; \sigma; s)$  is the hypergeometric function.

*Proof.* It is straightforward to verify that system (8.37) is Liouvillian integrable with an integrating factor and a first integral given by expressions (8.38) and (8.39), respectively. Since  $\beta \neq 0$  and  $w(x)$  is a polynomial of degree  $(m + 1)/l$ , we conclude that system (8.37) is from family (C).

Now our goal is to prove the converse statement. Let us suppose that a Liénard differential system (1.1) from family (C) is Liouvillian integrable and possesses a hyperelliptic invariant algebraic curve  $y^2 + u(x)y + v(x) = 0$ . By Theorem 8.2 condition (8.28) is identically satisfied and the system has integrating factor given by expression (8.29), where the polynomial  $F(x, y)$  can be chosen in the form  $F(x, y) = y^2 + u(x)y + v(x)$ . In addition, we set  $N_1 = 1$ . Substituting integrating factor (8.29) into the partial differential equation  $yM_x - [f(x)y + g(x)]M_y - f(x)M = 0$  and equating to zero the coefficients of different powers of  $y$  yields the relations

$$f(x) = \frac{2m + n + 3}{4(m + 1)}u_x, \quad g(x) = \frac{1}{2}v_x + \frac{n - 2m - 1}{8(m + 1)}uu_x \quad (8.40)$$

and the following equation

$$uv_x - \frac{n + 1}{m + 1}u_x v + \frac{n - 2m - 1}{4(m + 1)}u^2 u_x = 0. \quad (8.41)$$

Let us note that condition (8.28) produces an explicit expression of the polynomial  $f(x)$  similar to that given in relations (8.40). Integrating equation (8.41) with respect to the function  $v(x)$ , we obtain

$$v(x) = \beta u^{\frac{n+1}{m+1}} + \frac{1}{4}u^2, \quad (8.42)$$

where  $\beta \in \mathbb{C}$  is a constant of integration. Using Theorem 4.4 and the arguments given in the proof of Theorem 8.1, we conclude that  $u(x)$  is a polynomial of degree  $m + 1$  and  $v(x)$  is a polynomial of degree  $n + 1$ . Thus, we see that  $\beta$  is non-zero. Further, we introduce relatively prime natural numbers  $k$  and  $l$  satisfying the relation  $(m + 1)k = (n + 1)l$ . It follows from expression (8.42) that there exists a polynomial  $w(x)$  of degree  $(m + 1)/l$  such that the polynomial  $u(x)$  can be represented in the form  $u(x) = w^l(x)$ . Hence we obtain the equality  $v(x) = \beta w^k(x) + w^{2l}(x)$ . Substituting the explicit representations of the polynomials  $u(x)$  and  $v(x)$  into relations (8.40), we find the polynomials  $f(x)$  and  $g(x)$  as given in (8.37). Expressing the number  $n$  from the relation  $(m + 1)k = (n + 1)l$ , we find Darboux integrating factor (8.38) giving Liouvillian first integral (8.39).  $\square$

*Remark 1.* We do not require that the polynomial  $y^2 + u(x)y + v(x)$  is irreducible. See also Remark 1 to Theorem 8.2.

*Remark 2.* The family of systems (8.37) can be transformed to the following simple form

$$s_\tau = z, \quad z_\tau = -\frac{(k+2l)}{4}s^{l-1}z - \frac{k}{8}(s^{2l-1} + 4\beta s^{k-1}) \quad (8.43)$$

via the generalized Sundman transformation  $s(\tau) = w(x)$ ,  $z(\tau) = y$ ,  $d\tau = w_x(x)dt$ . Substituting  $w(x) = s$ ,  $y = z$  into (8.39), we find a Liouvillian first integral for systems (8.43).

It follows from Theorem 4.4 that equation (1.4) related to a Liénard differential system (1.1) from family (C) may have a polynomial solution only if  $n = \deg g(x)$  is an odd number. Such a polynomial solution gives rise to an invariant algebraic curve with the generating polynomial of the first degree with respect to  $y$ . Let us study the Liouvillian integrability of Liénard differential systems (1.1) from family (C) possessing invariant algebraic curves with generating polynomials of the first degree with respect to  $y$ . Since arbitrary coefficients arise in the non-polynomial part of the series  $y_\infty^{(l)}(x)$ ,  $l = 1, 2$ , we conclude that equation (1.4) has at most two distinct polynomial solutions simultaneously provided that the inequality  $\deg g > 2\deg f + 1$  holds. In what follows we denote these polynomial solutions as  $y = p_1(x)$  and  $y = p_2(x)$ . Note that the following relations  $p_l(x) = \{h^{(l)}(x)\}_+$ ,  $l = 1, 2$  are valid, where  $h^{(l)}(x)$  is the initial part of the series  $y_\infty^{(l)}(x)$ .

**Theorem 8.4.** *A Liénard differential system (1.1) from family (C) with two distinct invariant algebraic curves given by first-degree polynomials with respect to  $y$  is Liouvillian integrable if and only if  $n = \deg g(x)$  is an odd number, the system is of the form (8.37) and other conditions of Theorem 8.3 are satisfied with the additional restriction: either  $k$  is an even number or otherwise  $(m+1)/l$  is an even number and the polynomial  $w(x)$  has only double roots. The polynomials  $p_1(x)$  and  $p_2(x)$  producing the invariant algebraic curves  $y - p_1(x) = 0$  and  $y - p_2(x) = 0$  can be represented in the form*

$$p_1(x) = \sqrt{\beta}w^{\frac{k}{2}} + \frac{1}{2}w^l(x), \quad p_2(x) = -\sqrt{\beta}w^{\frac{k}{2}} + \frac{1}{2}w^l(x), \quad \beta \in \mathbb{C} \setminus \{0\}. \quad (8.44)$$

*Proof.* We use item 2 of Theorem 8.2 and the arguments given in the proof of Theorem 8.3. Let us note that the hyperelliptic invariant algebraic curve of Theorem 8.3 with the generating polynomial  $y^2 + w^l y + \frac{1}{4}w^{2l} + \beta w^k = (y + w^l/2)^2 + \beta w^k$  splits into two distinct invariant algebraic curves  $y - p_1(x) = 0$  and  $y - p_2(x) = 0$  if and only if  $n$  is an odd number and either  $k$  is an even number or otherwise  $(m+1)/l$  is an even number and  $w(x)$  is a polynomial with double roots. In addition, recall that the degree of the polynomial  $w(x)$  equals  $(m+1)/l$ . □

*Remark.* This theorem can also be proved directly without using Theorem 8.3. As an example, see Theorem 6.3.

Further, our goal is to demonstrate that there exist Liouvillian integrable Liénard differential systems (1.1) from family (C) for any choice of the numbers  $m = \deg f(x)$  and

$n = \deg g(x)$ . Setting  $u(x) = x^{m+1}$  in expression (8.42), we find the following Liouvillian integrable Liénard differential systems from family (C)

$$x_t = y, \quad y_t = -\frac{2m+n+3}{4}x^m y - \frac{n+1}{8}(4\beta x^n + x^{2m+1}). \quad (8.45)$$

The related Darboux integrating factor reads as

$$M(x, y) = \left( y^2 + x^{m+1}y + \beta x^{n+1} + \frac{1}{4}x^{2(m+1)} \right)^{-\frac{2m+n+3}{2(n+1)}}. \quad (8.46)$$

The numbers  $n = \deg g$  and  $m = \deg f$  can be chosen arbitrarily. In addition, if the following conditions  $n = l(m+1) - 1$ ,  $l \in \mathbb{N}$ , and  $l > 2$  hold, then we obtain another particular family of Liouvillian integrable Liénard differential systems

$$x_t = y, \quad y_t = -\frac{l+2}{4}u_x y - \frac{l}{8}(4\beta u^{l-1} + u)u_x, \quad (8.47)$$

where  $u(x)$  is an arbitrary polynomial of degree  $m+1$ . The associated Darboux integrating factor can be represented in the form

$$M(x, y) = \left( y^2 + uy + \beta u^l + \frac{1}{4}u^2 \right)^{-\frac{l+2}{2l}}. \quad (8.48)$$

Now let us study the the existence of non-autonomous Darboux–Jacobi last multipliers. The case  $\deg f = 0$  is simple. There are families of distinct Jacobi last multipliers arising as products of integrating factors (8.29), (8.30) and non-autonomous first integrals  $I^\varkappa(x, y, t)$ , where  $\varkappa \in \mathbb{C}$  and the function  $I(x, y, t)$  is given by relations (8.19) and (8.20).

**Lemma 8.3.** *A Liénard differential system (1.1) satisfying the conditions  $\deg g > 2 \deg f + 1$  and  $\deg f > 0$  has a non-autonomous Darboux–Jacobi last multiplier of the form (2.6) if and only if there exists a non-zero complex number  $\omega$  such that the relation*

$$4(m+1)f(x) + (2m+n+3) \{h_x^{(1)}(x) + h_x^{(2)}(x)\}_+ + 2(n+1)\omega = 0 \quad (8.49)$$

is identically satisfied and one of the following assertions is valid.

1. *There exists an irreducible invariant algebraic curve  $F(x, y) = 0$  such that the family of Puiseux series  $y_\infty^{(1)}(x)$  arises in the factorization of the polynomial  $F(x, y)$  in the ring  $\mathbb{C}_\infty\{x\}[y]$  as many times as so does the family  $y_\infty^{(2)}(x)$ , i.e.  $N_1 = N_2$ . In this case the system has the unique Darboux–Jacobi last multiplier*

$$M(x, y, t) = \{F(x, y)\}^{-\frac{2m+n+3}{2(n+1)N_1}} \exp[\omega t]. \quad (8.50)$$

2. *There exist two distinct irreducible invariant algebraic curves  $F_1(x, y) = 0$  and  $F_2(x, y) = 0$  such that the following relation  $N_{1,j} \neq N_{2,j}$ ,  $j = 1, 2$  is valid, where  $N_{1,j}$  is the number*

of times the family of Puiseux series  $y_\infty^{(l)}(x)$  enters the factorization of the polynomial  $F_j(x, y)$  in the ring  $\mathbb{C}_\infty\{x\}[y]$ . In this case the system has the unique Darboux–Jacobi last multiplier

$$M(x, y, t) = \frac{\{F_1(x, y)\}^{\frac{(2m+n+3)(N_{2,2}-N_{1,2})}{2(n+1)\Omega}}}{\{F_2(x, y)\}^{\frac{(2m+n+3)(N_{2,1}-N_{1,1})}{2(n+1)\Omega}}} \exp[\omega t], \quad (8.51)$$

where the parameter  $\Omega$  is given by the relation  $\Omega = N_{1,2}N_{2,1} - N_{1,1}N_{2,2}$ .

*Proof.* We repeat the proof of Theorem 8.2. The only difference is in condition (8.32). In the non-autonomous case this condition takes the form

$$\sum_{j=1}^K d_j N_{1,j} \left( f(x) + \{h_x^{(1)}(x)\}_+ \right) + \sum_{j=1}^K d_j N_{2,j} \left( f(x) + \{h_x^{(2)}(x)\}_+ \right) = \omega - f(x), \quad (8.52)$$

where  $\omega$  is a non-zero complex constant. □

We have established that Liénard differential systems (1.1) from family (C) have neither rational nor Darboux first integrals. Let us note that the famous Duffing oscillators belong to family (C). These oscillators are studied in articles [16, 19] in details.

## 9. Quartic Liénard differential equations with a quadratic damping function

The aim of the present section is to demonstrate that the necessary and sufficient conditions of Liouvillian integrability presented in the previous sections can be used to find all Liouvillian integrable subfamilies of Liénard differential systems without performing the classification of irreducible invariant algebraic curves. As an example, we consider Liénard differential systems with the restrictions  $\deg f = 2$  and  $\deg g = 4$ :

$$x_t = y, \quad y_t = -(\zeta x^2 + \beta x + \alpha)y - (\varepsilon x^4 + \xi x^3 + e x^2 + \sigma x + \delta), \quad \zeta \varepsilon \neq 0. \quad (9.1)$$

Introducing suitable rescalings and shifts, it is without loss of generality to set  $\zeta = 3$ ,  $\varepsilon = -3$ , and  $\beta = 0$ . In what follows we work with the following systems

$$x_t = y, \quad y_t = -(3x^2 + \alpha)y + 3x^4 - \xi x^3 - e x^2 - \sigma x - \delta, \quad \zeta \varepsilon \neq 0. \quad (9.2)$$

Let us solve the integrability problem for systems (9.2).

**Theorem 9.1.** *Quartic Liénard differential systems with a quadratic damping function (9.2) are Liouvillian integrable if and only the tuple of the parameters  $(\alpha, \xi, \delta, \sigma, e)$  equals*

$$\begin{aligned} I : \quad (\alpha, \xi, \delta, \sigma, e) &= \left( -\frac{25}{12}, -7, \frac{125}{432}, -\frac{25}{36}, -5 \right); \\ II : \quad (\alpha, \xi, \delta, \sigma, e) &= \left( -\frac{61}{12}, -7, \frac{3905}{432}, \frac{623}{36}, 4 \right). \end{aligned} \quad (9.3)$$

The related Darboux integrating factors can be represented as

$$\begin{aligned}
I : \quad M(x, y) &= \frac{1}{\left(y + x^3 + \frac{3}{2}x^2 + \frac{5}{12}x - \frac{25}{216}\right)^{\frac{2}{3}} \left(y - x^2 - \frac{5}{3}x - \frac{25}{36}\right)}; \\
II : \quad M(x, y) &= \frac{\left(y + x^3 + \frac{3}{2}x^2 - \frac{31}{12}x - \frac{781}{216}\right)^{\frac{1}{3}}}{y^2 + \frac{(6x+5)(6x-13)(6x+11)y}{216} - \frac{(6x-13)(6x+11)^2(6x+5)^2}{7776}}.
\end{aligned} \tag{9.4}$$

*Proof.* Our proof is based on the results of Theorems 4.1 and 6.2. The Puiseux series given in relation (4.3) are now the following

$$\begin{aligned}
y_{\infty}^{(1)}(x) &= -x^3 - \frac{3}{2}x^2 + \left(\xi - \alpha + \frac{9}{2}\right)x + b_3 + \sum_{l=1}^{\infty} b_{l+3}x^{-l}; \\
y_{\infty}^{(2)}(x) &= x^2 - \frac{1}{3}(\xi + 2)x + \frac{1}{3}(\xi + 2 - \alpha - e) + \sum_{l=1}^{\infty} a_{l+2}x^{-l}.
\end{aligned} \tag{9.5}$$

The Puiseux series  $y_{\infty}^{(1)}(x)$  has an arbitrary coefficient  $b_3$  and exists whenever the restriction  $e = 3(27 + 6\xi - 4\alpha)/4$  holds. The Puiseux series  $y_{\infty}^{(2)}(x)$  possesses uniquely determined coefficients. Note that we use novel designations for the coefficients of the Puiseux series  $y_{\infty}^{(2)}(x)$ . The series  $y_{\infty}^{(1)}(x)$  terminates at the zero term under the condition

$$\delta = \frac{1}{24}(2\xi + 9)(18\alpha + 4\xi\alpha - 4\sigma - 4\xi^2 - 36\xi - 81). \tag{9.6}$$

Thus, we see that systems (9.2) possess the invariant algebraic curve  $F_1(x, y) = 0$  of Theorem 6.2 whenever  $e = 3(27 + 6\xi - 4\alpha)/4$  and  $\delta$  is of the form (9.6). The related polynomial and the cofactor can be represented as

$$\begin{aligned}
F_1(x, y) &= y + x^3 + \frac{3}{2}x^2 - \left(\xi - \alpha + \frac{9}{2}\right)x - \frac{1}{3}(\sigma + \xi^2 - \xi\alpha) \\
&\quad - \frac{1}{4}(27 + 12\xi - 6\alpha), \quad \lambda_1(x, y) = 3x - \xi - \frac{9}{2}.
\end{aligned} \tag{9.7}$$

Condition (6.12) gives the following restriction:  $\xi = -7$ . Finally, we use Theorem 4.5 to find an irreducible invariant algebraic curve that exists simultaneously with  $F_1(x, y) = 0$  and is given by expression (4.1) where  $k = 1$  and  $N \in \mathbb{N}$ . As a result, we obtain the values of the parameters as presented in relation (9.3). The related invariant algebraic curves are given by the polynomials

$$\begin{aligned}
(I) : \quad F_2(x, y) &= y - x^2 - \frac{5}{3}x - \frac{25}{36}, \quad \lambda_2(x, y) = -3x^2 - 2x + \frac{5}{12}; \\
(II) : \quad F_2(x, y) &= y^2 + \frac{(6x+5)(6x-13)(6x+11)y}{216} - \frac{(6x-13)}{7776} \\
&\quad \times (6x+11)^2(6x+5)^2, \quad \lambda_2(x, y) = -3x^2 + x + \frac{71}{12}.
\end{aligned} \tag{9.8}$$

We calculate explicit expressions of Darboux integrating factors with the help of expression (6.13).  $\square$

In case (I) a Liouvillian first integral is given by expression (6.26), where one sets

$$l = 2, \quad k = 3, \quad \beta = -1, \quad v(x) = x + \frac{5}{6}. \quad (9.9)$$

In case (II) a Liouvillian first integral reads as

$$\begin{aligned} I(x, y) = & \frac{\sqrt{w(x)}}{w(x)} \sum_{l=0}^2 \left( \left\{ \sqrt{w(x)} - v(x) \right\} U(x) \ln \left( z^{\frac{1}{3}} - U(x) \exp \left( \frac{2\pi li}{3} \right) \right) \right. \\ & \left. + \left\{ \sqrt{w(x)} + v(x) \right\} V(x) \ln \left( z^{\frac{1}{3}} - V(x) \exp \left( \frac{2\pi li}{3} \right) \right) \right) \exp \left( \frac{2\pi li}{3} \right) + 6z^{\frac{1}{3}}, \end{aligned} \quad (9.10)$$

where we have introduced the notation

$$U(x) = \left\{ u(x) - v(x) + \sqrt{w(x)} \right\}^{\frac{1}{3}}, \quad V(x) = \left\{ u(x) - v(x) - \sqrt{w(x)} \right\}^{\frac{1}{3}}, \quad z = y + u(x). \quad (9.11)$$

The polynomials  $u(x)$ ,  $v(x)$ ,  $w(x)$  take the form

$$\begin{aligned} u(x) = & \frac{(6x + 11)(36x^2 - 12x - 71)}{216}, \quad v(x) = \frac{(6x + 5)(6x - 13)(6x + 11)}{432}, \\ w(x) = & \frac{(6x - 13)(6x + 11)^3(6x + 5)^2}{186624}. \end{aligned} \quad (9.12)$$

Concluding this section we note that the method of Puiseux series and the explicit expression (3.5) of the cofactor of an invariant algebraic curve greatly facilitate the classification of integrable multi-parameter planar differential systems.

## 10. Conclusion

This work completely solves the Liouvillian integrability problem for polynomial Liénard differential systems (1.1) satisfying the condition  $\deg g \neq 2 \deg f + 1$ . In the case  $\deg g = 2 \deg f + 1$  our results are complete for the non-resonant systems. We say that a Liénard differential system with the restriction  $\deg g = 2 \deg f + 1$  is resonant near infinity if equation (4.5) possesses a positive rational solution. The resonance condition introduces a restriction on the highest-degree coefficients  $f_0$  and  $g_0$  of the polynomials  $f(x)$  and  $g(x)$ .

We have established that a generic nonlinear polynomial Liénard differential system (1.1) with fixed degrees of the polynomials  $f(x)$  and  $g(x)$  is not Liouvillian integrable provided that the following restriction  $\deg g > \deg f$  is valid. However, as we have demonstrated, Liouvillian integrable subfamilies exist for any degrees of the polynomials  $f(x)$  and  $g(x)$  whenever  $\deg g > \deg f$ . Besides that, we have classified polynomial Liénard differential systems possessing non-autonomous Darboux first integrals and non-autonomous Jacobi last multipliers with a time-dependent exponential factor. Some of our results describing explicit families of Liouvillian integrable Liénard differential systems are gathered in Table 1.

*Remarks to Table 1*

1. Natural numbers  $l$  and  $k$  are both non-unit.
2. Symbols  $D$ ,  $E$ , and  $L$  mean Darboux, elementary, and Liouvillian, respectively. Symbols  $\overline{D}$  and  $\overline{E}$  mean non-Darboux and non-elementary.
3. Family  $(A)_1$  gives all Liouvillian integrable families of Liénard differential systems (1.1) such that the related equation (1.4) possesses two distinct polynomial solutions and the following inequalities  $\deg f < \deg g < 2 \deg f + 1$  are valid.
4. Families  $(B)_1$  and  $(B)_2$  produce all non-resonant Darboux integrable Liénard differential systems (1.1) satisfying the restriction  $\deg g = 2 \deg f + 1$ .
5. Families  $(B)_1$ ,  $(B)_2$ ,  $(B)_3$ , and  $(B)_4$  include all non-resonant Liouvillian integrable Liénard differential systems (1.1) satisfying the restriction  $\deg g = 2 \deg f + 1$ . Note that families  $(B)_3$  and  $(B)_4$  also involve integrable resonant systems. Consequently, additional restrictions should be imposed if one is interested only in the non-resonant case. These restrictions are described in Theorems 7.4 and 7.5.
6. Family  $(C)_1$  gives all Liouvillian integrable families of Liénard differential systems (1.1) with  $\deg g > 2 \deg f + 1$  possessing either a hyperelliptic invariant algebraic curve or two distinct invariant algebraic curves with generating polynomials of the first degree with respect to  $y$ .

Let us note that families  $(B)_1$  and  $(B)_2$  are those given by the Chiellini integrability condition  $\{f(x)/g(x)\}_x = \alpha f(x)$ , see [10]. Remarkably, Chiellini integrable Liénard differential systems can be linearized via generalized Sundman transformations [2]. Other integrable families from Table 1 can also be transformed to a more simple form via generalized Sundman transformations, see remarks and comments to Theorems 6.3, 7.3, 7.5, and 8.3. These systems with the exception of a number of partial cases that appear in [4, 37–39, 44] seem to be new.

Let us enumerate some unsolved problems related to the integrability and solvability of Liénard differential systems. Despite the fact that the subset of resonant Liénard differential systems is of Lebesgue measure zero in the set of all polynomial Liénard differential systems satisfying the condition  $\deg g = 2 \deg f + 1$ , it is an interesting open problem to perform a classification of invariant algebraic curves and integrable subfamilies of particular resonant polynomial Liénard differential systems provided that only a resonant condition is imposed on the parameters of the systems. The method of Puiseux series [17, 21] can deal with each family of resonant Liénard differential systems characterized by a fixed positive rational Fuchs index separately. Note that several novel families of Liouvillian integrable resonant systems are presented in Theorems 7.3 and 7.5, see also Corollary 2 after the latter theorem. In addition, a number of integrable resonant families with  $\deg f = 2$  and  $\deg g = 5$  are found via  $\lambda$  symmetries in [40]. These families possess an exciting property: they simultaneously have autonomous and non-autonomous Darboux first integrals given by expression (2.4) with  $\omega = 0$  and  $\omega \neq 0$ , respectively. This fact allowed A. Ruiz and C. Muriel to obtain nice

Family	$f(x), g(x)$	$M(x, y)$	$I(x, y)$ , type of first integral
(A) <sub>1</sub> Th 6.3	$f(x) = -[k\beta v^{k-1} + (k+l)v^{l-1}]v_x$ $g(x) = k[\beta v^k + v^l]v^{l-1}v_x$ $z = y - \beta v^k - v^l, \frac{m+1}{n-m} = \frac{k}{l}, (l, k) = 1$ $v(x) \in \mathbb{C}[x], \deg v = \frac{n-m}{l}, \beta \in \mathbb{C} \setminus \{0\}$	$\frac{z^{-\frac{1}{k}}}{y-v^l}$	$\frac{k\beta^{\frac{1}{k}}}{k-l}z^{\frac{k-l}{k}} + \sum_{j=0}^m \exp\left[-\frac{\pi l(2j+1)i}{k}\right]$ $\times \ln\left\{z^{\frac{1}{m+1}} - \exp\left[\frac{\pi(2j+1)i}{m+1}\right][\beta v^k]^{\frac{1}{m+1}}\right\}$ $\overline{DEL}$
(B) <sub>1</sub> Th 7.1	$f(x) = -\frac{2f_0}{f_0-\delta}q_{1,x}, g(x) = \frac{f_0+\delta}{f_0-\delta}q_{1,x}q_1$ $q_1(x) \in \mathbb{C}[x], \delta, f_0 \in \mathbb{C} \setminus \{0\}$ $q_1(x) = \frac{\delta-f_0}{2l(m+1)}x^{m+1} + o(x^{m+1})$	$\frac{1}{y-\frac{(f_0+\delta)}{(f_0-\delta)}q_1}$ $\times \frac{1}{y-q_1}$	$[y - q_1]^{\delta-f_0} \left[ y - \frac{(f_0+\delta)}{(f_0-\delta)}q_1 \right]^{\delta+f_0}$ $DEL$
(B) <sub>2</sub> Th 7.2	$f(x) = -2q_x, g(x) = qq_x$ $q(x) \in \mathbb{C}[x], \deg q = m+1$	$\frac{1}{\{y-q\}^2}$	$[y - q(x)] \exp\left[-\frac{q(x)}{y-q(x)}\right]$ $DEL$
(B) <sub>3</sub> Th 7.3	$f(x) = -\left[\frac{\{(2d_1+1)l+k\}l}{k-l}u^{l-1} + \beta(l+k)u^{k-1}\right]u_x$ $g(x) = \left[l\beta^2u^{2k-1} + \frac{\{(2d_1+1)l+k\}l\beta}{k-l} \times u^{k+l-1} + \frac{(ld_1+k)(d_1+1)l^2}{(k-l)^2}u^{2l-1}\right]u_x$ $p_1(x) = \beta u^k(x) + \frac{(d_1+1)l}{k-l}u^l(x)$ $p_2(x) = \beta u^k(x) + \frac{(ld_1+k)}{k-l}u^l(x)$ $(l, k) = 1, d_1, \beta \in \mathbb{C} \setminus \{0\}$ $u(x) \in \mathbb{C}[x], \deg u = \frac{m+1}{\max\{k, l\}}$	$\frac{[y-p_1]^{d_1}}{[y-p_2]^{d_1+1+\frac{k}{l}}}$	$\frac{p_2 B\left(\frac{y-p_1}{p_2-p_1}; 1+d_1, -d_1-\frac{k}{l}\right)}{\{p_2-p_1\}^{\frac{k}{l}}}$ $-\frac{B\left(\frac{y-p_1}{p_2-p_1}; 1+d_1, 1-d_1-\frac{k}{l}\right)}{\{p_2-p_1\}^{\frac{k}{l}-1}}$ $\overline{DEL} (d_1 \notin \mathbb{Q})$
(B) <sub>4</sub> Th 7.5	$f(x) = \left[\frac{2l^2}{l-k}v^{l-1} - (l+k)\beta v^{k-1}\right]v_x$ $g(x) = \left[\frac{l^3}{(l-k)^2}v^{2l-1} + l\beta^2v^{2k-1} - \frac{2l^2\beta}{l-k}v^{l+k-1}\right]v_x$ $q(x) = -\frac{l}{l-k}v^l + \beta v^k, \beta \in \mathbb{C} \setminus \{0\}$ $v(x) \in \mathbb{C}[x], \deg v = \frac{m+1}{\max\{k, l\}}, (l, k) = 1$	$\frac{\exp\left[\frac{v^l}{y-q}\right]}{[y-q]^{\frac{l+k}{l}}}$	$v^{l-k}\gamma\left(-\frac{l-k}{l}, \frac{v^l}{q-y}\right) - \frac{q}{v^k}\gamma\left(\frac{k}{l}, \frac{v^l}{q-y}\right)$ $\overline{DEL}$
(C) <sub>1</sub> Th 8.3 Th 8.4	$f(x) = \frac{(k+2l)}{4}w^{l-1}w_x$ $g(x) = \frac{k}{8}(w^{2l-1} + 4\beta w^{k-1})w_x$ $z = \left[y + \frac{w^l}{2}\right]^2 + \beta w^k, \frac{n+1}{m+1} = \frac{k}{l}$ $w(x) \in \mathbb{C}[x], \deg w = \frac{m+1}{l}, (l, k) = 1$	$z^{-\left(\frac{1}{2}+\frac{l}{k}\right)}$	${}_2F_1\left(\frac{1}{2}, \frac{1}{2} + \frac{l}{k}, \frac{3}{2}, -\frac{(2y+w^l)^2}{4\beta w^k}\right)$ $\times \frac{(2l-k)(2y+w^l)}{4kw^{\frac{k}{2}}\beta^{\frac{1}{2}+\frac{l}{k}}} + z^{\frac{1}{2}-\frac{l}{k}}$ $\overline{DEL}$

Table 1: The explicit Liouvillian integrable families of Liénard differential systems.

expressions of the general solutions. Systems (7.29) with  $\delta = \pm m f_0 / (m + 2)$  have the same property.

Along with this, it is a difficult open problem to perform a classification of integrable polynomial Liénard differential systems with non-Liouvillian first integrals. At the moment only particular examples that can be transformed to linear equations are available, for more details see articles [1, 22, 26, 27].

Another important problem is to study rational Liénard differential systems. If the functions  $f(x)$  and  $g(x)$  in expression (1.1) are rational, then systems (1.1) give rise to the

following polynomial differential systems in the plane

$$x_t = h(x)y, \quad y_t = -\tilde{f}(x)y - \tilde{g}(x), \quad h(x), \tilde{f}(x), \tilde{g}(x) \in \mathbb{C}[x]. \quad (10.1)$$

Investigating the analytic and qualitative properties of these systems with respect to the degrees of polynomials  $h(x)$ ,  $\tilde{f}(x)$ , and  $\tilde{g}(x)$  is a future challenge.

## 11. Acknowledgments

This research was supported by Russian Science Foundation grant 19–71–10003.

## References

- [1] Acosta-Humánez, P. B., Lázaro, J. T., Morales-Ruiz, J. J., and Pantazi, C. (2015). On the integrability of polynomial vector fields in the plane by means of Picard–Vessiot theory. *Discrete Contin. Dyn. Syst. Ser. A*, 35:1767–1800.
- [2] Berkovich, L. M. (1996). The method of an exact linearization of  $n$ -order ordinary differential equations. *Journal of Nonlinear Mathematical Physics*, 3(3–4):341–350.
- [3] Blázquez-Sanz, D. and Pantazi, C. (2012). A note on the Darboux theory of integrability of non-autonomous polynomial differential systems. *Nonlinearity*, 25:2615–2624.
- [4] Chandrasekar, V., Pandey, S., Senthilvelan, M., and Lakshmanan, M. (2006). A simple and unified approach to identify integrable nonlinear oscillators and systems. *J. Math. Phys.*, 47:023508.
- [5] Chavarriga, J., Giacomini, J., Giné, J., and Llibre, J. (2003). Darboux integrability and the inverse integrating factor. *J. Differential Eqs.*, 194:116–139.
- [6] Cheb-Terrab, E. (2003). An Abel ordinary differential equation class generalizing known integrable classes. *European J. Appl. Math.*, 17:217–229.
- [7] Cheb-Terrab, E. (2004). Solutions for the general, confluent and biconfluent Heun equations and their connection with Abel equations. *J. Phys. A*, 37:9923–9949.
- [8] Cherkas, L. A. (1976). Conditions for a Liénard equation to have a centre. *Differ. Equ.*, 12:201–206.
- [9] Chèze, G. and Cluzeau, T. (2013). On the nonexistence of Liouvillian first integrals for generalized Liénard polynomial differential systems. *Journal of Nonlinear Mathematical Physics*, 20(4):475–479.
- [10] Chiellini, A. (1931). Sull'integrazione della equazione differenziale  $y' + Py^2 + Qy^3 = 0$ . *Bollettino della Unione Matematica Italiana*, 10:301–307.
- [11] Christopher, C. (1999a). An algebraic approach to the classification of centers in polynomial Liénard systems. *J. Math. Anal. Appl.*, 229(1):319–329.
- [12] Christopher, C., Llibre, J., Pantazi, C., and Walcher, S. (2019). On planar polynomial vector fields with elementary first integrals. *Journal of Differential Equations*, 267(8):4572–4588.
- [13] Christopher, C. J. (1999b). Liouvillian first integrals of second order polynomial differential equations. *Electron. J. Differential Equations*, 49:1–7.
- [14] Demina, M. V. (2018a). Invariant algebraic curves for Liénard dynamical systems revisited. *Applied Mathematics Letters*, 84:42–48.
- [15] Demina, M. V. (2018b). Invariant surfaces and Darboux integrability for non-autonomous dynamical systems in the plane. *Journal of Physics A: Mathematical and Theoretical*, 51:505202.
- [16] Demina, M. V. (2018c). Novel algebraic aspects of Liouvillian integrability for two-dimensional polynomial dynamical systems. *Physics Letters A*, 382(20):1353–1360.
- [17] Demina, M. V. (2020a). Classifying algebraic invariants and algebraically invariant solutions. *Chaos, Solitons and Fractals*, 140:110219.
- [18] Demina, M. V. (2020b). Integrability and Jacobi last multipliers of cubic Liénard differential equations with quadratic damping. *Discontinuity, Nonlinearity, and Complexity*, 9(4):499–507.

- [19] Demina, M. V. (2021a). Liouvillian integrability of the generalized Duffing oscillators. *Analysis and Mathematical Physics*, 11(1):1–25.
- [20] Demina, M. V. (2021b). The method of Puiseux series and invariant algebraic curves. *Communications in contemporary mathematics*, 24(3):2150007.
- [21] Demina, M. V. (2021c). Necessary and sufficient conditions for the existence of invariant algebraic curves. *Electronic Journal of Qualitative Theory of Differential Equations*, 48:1–22.
- [22] Demina, M. V., Giné, J., and Valls, C. (2022). Puiseux integrability of differential equations. *Qualitative Theory of Dynamical Systems*, 21(2):1–35.
- [23] Demina, M. V. and Sinelshchikov, D. I. (2019). Integrability properties of cubic Liénard oscillators with linear damping. *Symmetry*, 11:1378.
- [24] Demina, M. V. and Sinelshchikov, D. I. (2021). Darboux first integrals and linearizability of quadratic–quintic Duffing–van der Pol oscillators. *Journal of Geometry and Physics*, 165:104215.
- [25] Demina, M. V. and Valls, C. (2020). On the Poincaré problem and Liouvillian integrability of quadratic Liénard differential equations. *Proc. Roy. Soc. Edinburgh Sec. A: Mathematics*, 150(6):3231–3251.
- [26] García, I. A. and Giné, J. (2003). Generalized cofactors and nonlinear superposition principles. *Appl. Math. Lett.*, 26(7):1137–1141.
- [27] García, I. A., Giné, J., and Llibre, J. (2008). Liénard and Riccati differential equations related via Lie algebras. *Discrete Contin. Dyn. Syst. Ser. B*, 10:485–494.
- [28] Gasull, A. and Giné, J. (2017). Integrability of Liénard systems with a weak saddle. *Z. Angew. Math. Phys.*, 68(13):1–13.
- [29] Ghose Choudhury, A. and Guha, P. (2017). Chiellini integrability condition, planar isochronous systems and Hamiltonian structures of Liénard equation. *Discrete Contin. Dyn. Syst. Ser. B*, 22:2465–2478.
- [30] Giné, J. and Llibre, J. (2011). Weierstrass integrability in Liénard differential systems. *Journal of Mathematical Analysis and Applications*, 377:362–369.
- [31] Guha, P. and Ghose Choudhury, A. (2019). Nonlocal transformations of the generalized Liénard type equations and dissipative Ermakov–Milne–Pinney systems. *International Journal of Geometric Methods in Modern Physics*, 16(7):1950107.
- [32] Ilyashenko, Y. and Yakovenko, S. (2008). *Lectures on Analytic Differential Equations*, volume 86. Graduate Studies in Mathematics, American Mathematical Society.
- [33] Liénard, A. (1928). Etude des oscillations entretenues. *Revue générale de l'électricité*, 23:901–912.
- [34] Llibre, J. and Pantazi, C. (2009). Darboux theory of integrability for a class of nonautonomous vector fields. *J. Math. Phys.*, 50:102705.
- [35] Llibre, J. and Valls, C. (2013). Liouvillian first integrals for generalized Liénard polynomial differential systems. *Adv. Nonlinear Stud.*, 13:819–829.
- [36] Llibre, J. and Valls, C. (2016). On the analytic integrability of the Liénard analytic differential systems. *Discrete Contin. Dyn. Syst. Ser. B*, 21(2):557–573.
- [37] Pandey, S., Bindu, P., Senthilvelan, M., and Lakshmanan, M. (2009a). A group theoretical identification of integrable cases of the Liénard–type equation  $\ddot{x} + f(x)\dot{x} + g(x) = 0$ . I: Equations having nonmaximal number of Lie point symmetries. *J. Math. Phys.*, 50(8):082702.
- [38] Pandey, S., Bindu, P., Senthilvelan, M., and Lakshmanan, M. (2009b). A group theoretical identification of integrable equations in the Liénard–type equation  $\ddot{x} + f(x)\dot{x} + g(x) = 0$ . II: Equations having maximal Lie point symmetries. *J. Math. Phys.*, 50(10):102701.
- [39] Polyanin, A. D. and Zaitsev, V. F. (2003). *Handbook of exact solutions for ordinary differential equations, Secod Edition*. Chapman and Hall, Boca Raton.
- [40] Ruiz, A. and Muriel, C. (2018). On the integrability of Liénard I–type equations via  $\lambda$  symmetries and solvable structures. *Appl. Math. Comput.*, 339:888–898.
- [41] Sinelshchikov, D. I. (2020). On linearizability via nonlocal transformations and first integrals for second-order ordinary differential equations. *Chaos, Solitons and Fractals*, 141:110318.
- [42] Sinelshchikov, D. I. (2021). Nonlocal deformations of autonomous invariant curves for liénard equations with quadratic damping. *Chaos, Solitons and Fractals*, 152:111412.
- [43] Singer, M. (1992). Liouvillian first integrals of differential systems. *Trans. Amer. Math. Soc.*, 333:673–

688.

- [44] Stachowiak, T. (2019). Hypergeometric first integrals of the Duffing and van der Pol oscillators. *Journal of Differential Equations*, 266:5895–5911.
- [45] Zhang, X. (2017). *Integrability of Dynamical Systems: Algebra and Analysis*. Springer Singapore.
- [46] Żołądec, H. (1998). Algebraic invariant curves for the Liénard equation. *Trans. Amer. Math. Soc.*, 350:1681–1701.