

# Thermalization without chaos in harmonic systems

Niccolò Cocciaglia,<sup>1</sup> Angelo Vulpiani,<sup>1</sup> and Giacomo Gradenigo<sup>2,3</sup>

<sup>1</sup>*Dipartimento di Fisica, Università di Roma “Sapienza”, Piazzale A. Moro 2, I-00185, Roma, Italy*

<sup>2</sup>*Gran Sasso Science Institute, Viale F. Crispi 7, 67100 L’Aquila, Italy*

<sup>3</sup>*INFN-Laboratori Nazionali del Gran Sasso, Via G. Acitelli 22, 67100 Assergi (AQ), Italy\**

Recent numerical results showed that thermalization of Fourier modes is achieved in short time-scales in the Toda model, despite its integrability and the absence of chaos. Here we provide numerical evidence that the scenario according to which chaos is irrelevant for thermalization is realized even in the simplest of all classical integrable system: the harmonic chain. We study relaxation from an atypical condition given with respect to *random* modes, showing that a thermal state with equilibrium properties is attained in short times. Such a result is independent from the orthonormal base used to represent the chain state, provided it is random.

## INTRODUCTION

Whether chaos is a necessary or just a sufficient condition in order for many-body Hamiltonian systems to exhibit thermalization is an old debate, to which a final assessment has not yet been given [1–3]. A possibility, outlined with interesting mathematical argument first by Khinchin in 1949 [4] and then by Mazur and van der Linden [5], is that basically the applicability of the statistical mechanics description to a macroscopic object relies on the number of its microscopic constituents being very large, irrespective to details of their dynamics. There are in fact examples showing that chaos is not necessary to have good statistical features [6], sometimes being not even sufficient [7, 8]. It is in this spirit that two of us recently investigated how an atypical initial condition relaxes to equipartition in an integrable Hamiltonian system, the Toda chain [9]. It turned out that in order to see fast relaxation to thermal equilibrium it is sufficient to consider appropriate observables, for instance observables almost independent from the conserved quantities of the integrable system [9]. For what concerns the Toda model at not too low energies this is for instance the case of the Fourier modes harmonic energies, which thermalize fast. On the other hand, one could say that thermalization of Fourier modes in the Toda chain is specific to the choice of variables and to its intricate relation with the Toda modes; in fact it is well known that, if one considers the Toda modes, the thermodynamic of the system is well described by the Generalized Gibbs Ensemble [10–13]. The goal of the present work is to give further and more stringent evidences in favour of the scenario tested numerically in [9] and at the basis of the Khinchin approach [4]: in the large- $N$  limit details of the microscopic dynamics are not relevant for statistical mechanics.

In the present paper we discuss an alternative way to study the behaviour of a system which is a textbook example: the harmonic chain. Our goal is to show that the lack of thermalization is a property *specific* of the Fourier modes, in the sense that there are infinitely many other choices of canonical variables to represent the chain state

and that using such variables one can see fast relaxation to a thermal state. In order to complement the results of [9] we will present here the numerical evidence of two important facts: 1) there are *many* (ideally infinitely many) independent sets of canonical variables with respect to which the system shows thermalization and 2) the relaxation to a thermal state is a large- $N$  effect. The fact that the validity of statistical mechanics can be generally claimed for any system *only* in the large- $N$  limit is particularly relevant for integrable systems, for which at small sizes the regular behaviour of the dynamics is manifest. It is therefore definitely worth to recall the enunciate of the Khinchin result, in order to grasp the relevance of the bound imposed by the large- $N$  limit. Given that  $f(q, p) : \mathbb{R}^{2dN} \rightarrow \mathbb{R}$  is an observable of our many-body classical Hamiltonian system, with  $\langle f \rangle$  and  $\bar{f}$  respectively its statistical and time averages, if  $f$  is a sum function, namely something of the kind  $f(q, p) = \sum_{i=1}^N f_i(q_i, p_i)$  (in practice any sort of additive quantity which can be measured as a function of small portions of the system), then under quite general hypothesis [4] one has that the probability to have a difference between  $\langle f \rangle$  and  $\bar{f}$  small in  $N$  is itself small in  $N$ . By denoting as  $\mathcal{P}$  this probability, in formulas we have:

$$\mathcal{P} \left( \left| \frac{\bar{f} - \langle f \rangle}{\langle f \rangle} \right| \geq \frac{C_1}{N^{1/4}} \right) < \frac{C_2}{N^{1/4}}, \quad (1)$$

where  $C_1$  and  $C_2$  are  $\mathcal{O}(1)$  constants which do not depend on  $N$ . Without going in further details of the mathematical results we can notice that its underlying message is in agreement with our results: irrespective to the nature of the microscopic dynamics ensemble and dynamical averages are for practical purposes equivalent in the large- $N$  limit.

## MODEL

Let us consider an harmonic chain with the following Hamiltonian

$$\mathcal{H}(q, p) = \sum_{i=1}^N \frac{p_i^2}{2m} + \frac{k}{2} \sum_{i=0}^N (q_{i+1} - q_i)^2, \quad (2)$$

where we can set masses and the elastic coefficient of the springs to  $m = k = 1$  and fixed boundary conditions  $q_0 = q_{N+1} = 0$ . The Hamiltonian in Eq. (2) can be easily put in a diagonal form with the following change of variables

$$\begin{aligned} Q_k &= \sqrt{\frac{2}{N+1}} \sum_{i=1}^N q_i \sin\left(\frac{\pi i k}{N+1}\right) \\ P_k &= \sqrt{\frac{2}{N+1}} \sum_{i=1}^N p_i \sin\left(\frac{\pi i k}{N+1}\right). \end{aligned} \quad (3)$$

By applying the above transformation, which is a canonical change of coordinates, one gets:

$$\mathcal{H}(Q, P) = \frac{1}{2} \sum_{k=1}^N (P_k^2 + \omega_k^2 Q_k^2) \quad (4)$$

where

$$\omega_k = 2 \sin\left(\frac{\pi k}{2N+2}\right) \quad (5)$$

is the angular frequency of the  $k$ -th normal mode. It is then convenient for our purposes to introduce the semi-canonical complex variables:

$$\begin{aligned} z_k &= \frac{P_k + i\omega_k Q_k}{\sqrt{2\omega_k}} \\ z_k^* &= \frac{P_k - i\omega_k Q_k}{\sqrt{2\omega_k}}, \end{aligned} \quad (6)$$

such that

$$\begin{aligned} \mathcal{H}(z, z^*) &= \sum_{k=1}^N \omega_k |z_k|^2 \\ \{z_k^*, z_q\} &= i \delta_{k,q} \end{aligned} \quad (7)$$

---

It is easy to check that the random rotation does not alter the Poisson parentheses structure between the variables:

$$\{z_p(\theta), z_q^*(\theta)\} = \sum_{ij} M_{pi}(\theta) M_{qj}^*(\theta) \{z_i, z_j^*\} = i \sum_{i=1}^N M_{pi}(\theta) M_{iq}^\dagger(\theta) = i \delta_{pq} \quad (11)$$

Clearly, in terms of these random modes the Hamiltonian is not anymore diagonal:

$$\mathcal{H}(z(\theta), z^*(\theta)) = \sum_{qp=1}^N c_{qp}(\theta) z_q^*(\theta) z_p(\theta), \quad (12)$$

## RANDOM MODES THERMALIZATION

The considered system is such that if only one harmonic is excited at the beginning, i.e., one sets as initial condition  $|z_k|^2 = E$ , for a given  $k$  and  $|z_q|^2 = 0$  for all  $q \neq k$ , energy is never shared among Fourier modes due to the lack of interaction. In other words, Fourier modes are integrability Liouville-Arnol'd theorem variables, so that no signature of thermalization can be found by definition looking at such variables. But let us consider a different point of view. The harmonic chain is a system characterized by a set of  $2N$  real coordinate-momenta variables, or by a set of  $N$  complex coordinates, which can be chosen at will. The complex coordinates  $z_k$  represent just one of the infinitely many choices available. Any random rotation in the  $N$ -dimensional complex space leads to another possible system of coordinates. Let us denote with the symbol

$$\theta = \{\theta_1, \dots, \theta_N\}, \quad (8)$$

the  $N$  random angles which define a random rotation in  $\mathbb{C}^N$  and with  $M(\theta)$  the unitary matrix,  $M(\theta) \in U(N)$ , which represent this rotation. In particular we have

$$M^\dagger M = M M^\dagger = \mathbf{1}, \quad (9)$$

where  $M^\dagger$  means transpose and complex conjugate. We thus have infinitely many choices of “*random modes*”, which define as random rotations of the Fourier modes and indicate as:

$$z_k(\theta) = \sum_{q=1}^N M_{kq}(\theta) z_q. \quad (10)$$

---

where

$$c_{qp} = \sum_{k=1}^N \omega_k \mathcal{M}_{kq}^*(\theta) \mathcal{M}_{kp}(\theta), \quad (13)$$




Figure 1. Panels *a*) and *b*) show two different choices of *random modes*, i.e., two independent choices of the unitary random matrix  $M(\theta)$ . *Main*: Normalized random modes energy spectrum at different times,  $u_k(\theta, t)$  vs  $k$  for increasing values of  $t$ ; *Inset*: effective number of degrees of freedom as a function of time,  $n_{\text{eff}}(\theta, t)$  vs  $t$ .

with  $\mathcal{M}(\theta) = [M^{-1}(\theta)]$ ,  $\mathcal{M}(\theta) \in U(N)$ .

We then define the energy of the random mode  $k$  as

$$\mathcal{E}_k(\theta) = c_{kk}(\theta) |z_k(\theta)|^2, \quad (14)$$

since  $c_{kk}(\theta)$  is real then also  $\mathcal{E}_k(\theta)$  is real. We have in fact by definition  $c_{kk}(\theta) = c_{kk}^*(\theta)$ , see Eq. (13). We can thus write the Hamiltonian by separating the diagonal from the non-diagonal contribution as

$$\mathcal{H}(z(\theta), z^*(\theta)) = \sum_{k=1}^N \mathcal{E}_k(\theta) + \sum_{q \neq p}^N c_{qp}(\theta) z_q^*(\theta) z_p(\theta), \quad (15)$$

and then perform the standard FPUT numerical experiment, which amounts to the numerical study of the relaxational dynamics when energy is initially fed to a single random mode:

$$\mathcal{E}_k = \begin{cases} E_0 & k = 1 \\ 0 & k \neq 1. \end{cases} \quad (16)$$

Clearly the value of  $E_0$  set in the initial condition does not corresponds to the total energy of the system, which comprises the non-diagonal terms of Eq. (15). The configuration of the system is then evolved according to the standard Hamiltonian dynamics in the particles coordinate and momenta representation.

In agreement with the traditional approach to the Fermi-Pasta-Ulam-Tsingou problem [14–18] we have monitored along the dynamics the behaviour of the energy spectrum and of the spectral entropy, which quantifies with a number between zero and one the degree of energy equipartition between modes. A way to quantify the degree of equipartition between modes is obtained by monitoring the energy density

$$u_k(\theta, t) = \frac{\langle \mathcal{E}_k(\theta) \rangle_t}{\langle \mathcal{E}_{\text{tot}}(\theta) \rangle_t}, \quad (17)$$

where  $\langle \mathcal{E}_{\text{tot}}(\theta) \rangle_t = \sum_{k=1}^N \langle \mathcal{E}_k(\theta) \rangle_t$  and we consider cumulative time averages of the kind

$$\langle \mathcal{E}_k(\theta) \rangle_t = \frac{1}{t} \int_0^t ds \mathcal{E}_k(\theta, s). \quad (18)$$

The same information can be represented by studying the behaviour of the effective number of degrees of freedom  $n_{\text{eff}}(\theta, t)$ ,

$$n_{\text{eff}}(\theta, t) = \frac{\exp(S_{\text{sp}}(\theta, t))}{N}. \quad (19)$$

where  $S_{\text{sp}}(\theta, t)$  is the so-called *spectral entropy*:

$$S_{\text{sp}}(\theta, t) = - \sum_{k=1}^N u_k(\theta, t) \log u_k(\theta, t) \quad (20)$$

The parametric dependence of the spectrum  $u_k(\theta, t)$  on the time variable  $t$  is illustrated in Fig. (1) for two different choices of the random rotation  $\theta$ . Two remarkable observations are in order: equipartition among random modes is reached quite fast and this happens on the same time-scale for all choices of the random rotation  $\theta$ . This means that, apart the Fourier basis, fast relaxation to a thermal state is the *typical* phenomenon for random choice of the basis to represent the chain configuration. The inset of each of the four panels of Fig.1 represents the behaviour of the effective number of degrees of freedom  $n_{\text{eff}}(t)$ : the typical sigmoidal shape with a fast convergence to  $n_{\text{eff}}(t) \approx 1$  signals the reaching of equipartition. Despite integrability thermalization is observed even in the harmonic chain, provided the “*right variables*” are considered. And, what is most remarkable, *thermalization is typical*, while the lack of it is specific only to the representation of the chain configuration in Fourier space. The idea that in the large- $N$  limit the relevant thermodynamic properties of a system cannot




Figure 2. Effective number of degrees of freedom as a function of time,  $n_{\text{eff}}(\theta, t)$  vs  $t$ : the continuous (black) line represents the average  $\overline{n_{\text{eff}}}(t)$  over 30 choices of the random modes base, points represent the individual behaviour of 8 different instances.

be tight to a particular choice of coordinates was at the basis of an averaging strategy proposed already in [19] to capture the essential thermodynamic features of the low temperature glassy phase of an optimization problem.

In Fig. 2 the behaviour of the effective number of degrees of freedom averaged over  $M = 30$  instances of the random modes,  $\overline{n_{\text{eff}}}(\theta, t)$  (continuous line), is compared with the behaviour for single instances of the random modes,  $n_{\text{eff}}(\theta, t)$  (points): there is a clear evidence that for a chain of  $N = 1024$  the behaviour of  $n_{\text{eff}}(\theta, t)$  for each single instance is typical.

$$\overline{n_{\text{eff}}}(t) = \frac{1}{M} \sum_{i=1}^M n_{\text{eff}}(\theta_i, t) \quad (21)$$

### EQUILIBRIUM DYNAMICS: IMPORTANCE OF THE LARGE- $N$ LIMIT.

Beside the “standard” FPUT-like numerical experiment discussed in the previous section, probing the relaxation from an atypical initial condition to a thermal state, it is important to investigate the equilibrium dynamics of the system. The first test of thermalization has been the study of the single-mode energy probability distributions, simply defined as the histogram of values taken by  $\mathcal{E}_k$ . In the inset of Fig. 3 it is shown that  $p(\mathcal{E}_k)$ , plotted for different values of  $k$ , has in all cases a nice exponential behaviour, as expected from a thermal ensemble:

$$p(\mathcal{E}_k) \sim \exp(-b\mathcal{E}_k), \quad (22)$$




Figure 3. *Main*: time auto-correlation function  $C_k(t)$  as a function of time for different choices of the random modes index  $k$ , system size  $N = 1024$ ; *Inset*: probability distribution of random-modes energy  $\mathcal{E}_k$  for some values of  $k$ , system size  $N = 1024$ .

where the value of  $b$ , proportional to the inverse of energy per degree of freedom, is identical for all random modes  $k$ . The second important test of thermalization is to check that the system reached a stationary state where, “to all practical purposes”, no memory of a generic initial condition is left.

Clearly, this corresponds to a very pragmatic point of view: we look at a generic random modes basis and simply ignore that the system is integrable. How does look like an time auto-correlation function  $C_k(t)$  of the energy on the random mode  $k$ ? Let us define  $C_k(t)$  as

$$C_k(t) = \frac{\langle \mathcal{E}_k(t)\mathcal{E}_k(0) \rangle - \langle \mathcal{E}_k \rangle^2}{\langle \mathcal{E}_k^2 \rangle - \langle \mathcal{E}_k \rangle^2}. \quad (23)$$

In the main panel of Fig. 3 it is shown the behaviour of  $C_k(t)$  for different  $k$ : the function rapidly decays to values close to zero, around which it then oscillates. How it is possible that an integrable system, moreover characterized by simple linear interactions, presents such good signatures of an equilibrium behaviour? Isn’t it inconsistent with integrability? The way out to this apparent contradiction comes by taking into account the same two points which are the cornerstone of the Khinchin theorem: to consider the appropriate observables and to consider the large- $N$  limit. The dynamics of Fourier modes amplitudes in the harmonic chain is clearly analytically predictable, as done in a very insightful paper of Mazur and Montroll [6].

Already in [6] in fact it was shown that, even in an integrable system like an harmonic chain, if one takes into account a “generic” observable like the energy of a single particle of the chain, the time-autocorrelation function decays fast to a zero. In that case it was then shown that the imprint of the integrable nature of the system was brought by a sort of quasi-periodic oscillations




Figure 4. Time autocorrelation function  $C_k(t)$  as a function of time for different choices of the random-modes index  $k$ , system size  $N = 7$ .

around zero of amplitude  $1/N^{1/2}$  [6]. The picture emerging already from [6] and corroborated by the present results on random modes is therefore clear: if one considers as observables functions which are a (random) combination of *all* the coordinates diagonalizing the Hamiltonian, the “integrability effects” are of order  $\mathcal{O}(N^{-1/2})$ . From this point of view we can already find in the work of Mazur and Montroll [6] the idea that thermalization comes about simply with the thermodynamic limit and does not depend on the details of the micro-

scopic dynamics. For what concerns the present work we have shown that the autocorrelation function of a random mode energy is for all practical purposes equivalent to that of single particle in the harmonic chain, since they both result from the combination of all Fourier modes. In agreement with the results of [6], it is worth showing that even in the present case the fast decay of  $\langle \mathcal{E}_k(t) \mathcal{E}_k(0) \rangle$  is a large- $N$  effect. Without going through a too much detailed finite-size analysis, we show here in Fig. 4 that for small system sizes (e.g.,  $N = 7$  particles) the recurrent nature of the integrable dynamics clearly emerges.

In Fig. 4 is shown the time dependence of the autocorrelation function defined in Eq. (23) for an harmonic chain of  $N = 7$  particles and for some values of  $k$ : it can be clearly seen that at a characteristic time  $\tau \approx 4520$  the system is back to its initial condition.

## OVERLAPS DISTRIBUTION

Let us now represent in a quantitatively precise manner the statement, already stressed by some of us in [9], that the variables allowing to see thermalization in an integrable system must be *almost uncorrelated* to those diagonalizing the Hamiltonian. To this aim we introduce the *overlap* between Fourier modes and random modes,

$$|q_{kp}| = \left| \frac{\langle z_k(\theta) | z_q \rangle}{\langle z_q | z_q \rangle} \right| = |M_{kq}|, \quad (24)$$

which turns out to be nothing but the the matrix element  $M_{kq}$  of the random unitary matrix connecting Fourier modes and random modes (we have to consider its modulus because it is a complex variable). Since  $|q_{kp}| = |M_{kq}|$ , from the properties of random unitary matrices the probability distribution  $\rho(|q|)$  can be computed exactly. For the ease of the reader we report this elementary large deviation estimate in the Appendix, quoting here only the

final result:

$$\rho(|q|) = 2N|q| \exp(-N|q|^2) \quad (25)$$

By recalling that we are dealing with a complex variable,  $q \in \mathbb{C}$ , the linear dependence on  $|q|$  in front  $\rho(|q|)$  has to be understood simply as the Jacobian element for the representation of the complex variable in polar coordinates, so that the probability distribution of the overlap,  $\mathcal{P}(q) : \mathbb{C} \rightarrow \mathbb{R}$  is simply Gaussian:

$$\mathcal{P}(q) \propto \exp(-Nq^2), \quad (26)$$

which completes our discussion about choosing a set of canonical coordinates which are *random combination* of the coordinates which make the integrable nature of the system manifest. In addition to what already observed in [9], in the present work provided sufficient evidence from data in favour of the hypothesis that the set of (semi) canonical coordinates which behaves “thermally” are infinitely many, actually all the random unitary transformations in  $M(\theta) : \mathbb{C}^N \rightarrow \mathbb{C}^N$ . It is remarkable how




Figure 5. *Main*: probability distribution  $P(|q|)$  of the absolute value of the overlap between a Fourier mode and a random mode, for different sizes  $N$  of the system; *Inset*: collapsed data for different  $N$ .

the large- $N$  prediction of Eq. (25) is in good agreement with the numerical estimates of the probability distribution of overlaps at finite  $N$ , which is shown in Fig. 5. The existence of an infinite number of basis of orthonormal functions to represent the chain configuration, each base looking equally random to any other, and the relevance of this property for the possibility to detect equilibrium with respect to each base closely resembles a similar property studied for the choice of eigenfunction basis in quantum systems [24].

## CONCLUSIONS

The results presented in this work aims at clarifying a foundational problem in statistical mechanics: is dynamical chaos really a necessary condition to guarantee the thermalization of systems with many degrees of freedom? Our results generalized and strengthened the conclusions drawn for the Toda model in [9]: thermalization is achieved even in an integrable system, provided that an appropriate choice of the observables is made. The study of the harmonic chain added an insight: we have shown that there are many (ideally infinitely many) different choices of canonical coordinates which allow to detect thermalization in the harmonic chain. This evidence tells us that the lack of equilibrium is really a specific property of the coordinates which diagonalizes the Hamiltonian. Robust thermodynamic properties must be independent from the choice of the coordinates used to write the partition sum, an idea already exploited with success in replica calculations for ordered systems with a glassy phase [19]. This idea, which is substantiated mathematically by the Khinchin approach and of which we presented numerical evidence in this paper, tells us that even for integrable systems there are no

a-priori reasons to not expect the validity of equilibrium ensembles. This of course is not in contradiction with well established frameworks for integrable systems such as the Generalized Gibbs Ensemble [10–13, 20–23], the latter being specific of particular choice of the canonical coordinates. At the same time it is clear that the possibility of unveil specific properties of the dynamics crucially depends on the choice of variables, in particular for system where ergodicity is spontaneously broken as spin glasses [25–28] or systems highly structured and intrinsically characterized by multiple length/time scales like macro-molecules [29].

As a final remark, let us notice that a generalization of the present results to quantum mechanics may provide further insights on the thermalization mechanism in quantum isolated systems. The point made by our findings on the *irrelevance of chaos* for the thermalization of harmonic systems are in fact quite similar to the Von Neumann quantum ergodic theorem approach [31]. According to the latter thermalization is in fact related to an appropriate choice of observables, rather than to the spectral properties of the Hamiltonian. The same scenario, where the choice of observables plays a central role, is the one underlying the Eigenstate Thermalization Hypothesis [32, 33]. Numerical experiments to better explore the analogies between the foundation of statistical mechanics in classical and quantum systems, for instance investigating the scope of thermal ensembles applicability in the presence of an external driving [34], i.e. fluctuation-dissipation relations, are therefore in order.

**Acknowledgments** We thank J. Kurchan, R. Livi, S. Pappalardi and A. Ponso for useful discussions. A.V. acknowledge partial financial support of project MIUR-PRIN2017 *Coarse-grained description for non-equilibrium systems and transport phenomena (CO-NEST)*.

## APPENDIX A

In this Appendix we show how to compute the probability distribution of the generic matrix element  $M_{ki}$  (for simplicity of notation, let us omit the angle  $\theta$ ). To this aim it is possible to exploit the fact that each row/column of the matrix represents one (normalized) element of an eigenvector basis of  $\mathbb{C}^N$ , so that the following condition holds:

$$\sum_{i=1}^N |M_{ki}|^2 = A \quad \forall k, \quad (27)$$

where  $A$  is the normalization that we will fix to 1 at the end of the calculation. Changing the notation so that:  $\zeta_i = M_{ki}$ , we have that the volume spanned by the elements of a row in  $M$  reads:

$$\Omega_N(A) = \int \prod_{i=1}^N d\text{Re}(\zeta_i) d\text{Im}(\zeta_i) \delta\left(A - \sum_{i=1}^N |\zeta_i|^2\right), \quad (28)$$

so that the joint probability density is:

$$\rho(\zeta_1, \dots, \zeta_N | A) = \frac{1}{\Omega_N(A)} \delta\left(A - \sum_{i=1}^N |\zeta_i|^2\right). \quad (29)$$

Since we are looking for the distribution of a single matrix element, we are interested in the marginal:

$$\rho(\zeta_1 | A) = \int \prod_{i=2}^N d\text{Re}(\zeta_i) d\text{Im}(\zeta_i) \rho(\zeta_1, \dots, \zeta_N | A). \quad (30)$$

In order to compute  $\Omega_N(A)$ , let us switch to polar coordinates  $\begin{cases} \text{Re}(\zeta_i) = r_i \cos(\phi_i) \\ \text{Im}(\zeta_i) = r_i \sin(\phi_i) \end{cases}$ :

$$\Omega_N(A) = (2\pi)^N \int_0^\infty \left[ \prod_{i=1}^N dr_i r_i \right] \delta\left(A - \sum_{i=1}^N r_i^2\right). \quad (31)$$

In order to get to the result, we compute its Laplace transform:

$$\begin{aligned} \Omega_N(\mu) &= \int_0^\infty dA e^{-\mu A} \Omega_N(A) \\ &= (2\pi)^N \int_0^\infty \left[ \prod_{i=1}^N dr_i r_i \right] \exp[-\mu \sum_i r_i^2] = \\ &= \left[ 2\pi \int_0^\infty dr r e^{-\mu r^2} \right]^N = \left(\frac{\pi}{\mu}\right)^N \\ &= \pi^N \exp[-N \log \mu] \end{aligned} \quad (32)$$

and then its inverse transform by applying the saddle point approximation, in the large- $N$  limit:

$$\begin{aligned} \Omega_N(A) &= \frac{1}{2\pi i} \int_{\mu_0 - i\infty}^{\mu_0 + i\infty} d\mu e^{\mu A} \Omega_N(\mu) \\ &= \frac{\pi^N}{2\pi i} \int_{\mu_0 - i\infty}^{\mu_0 + i\infty} d\mu e^{N[\mu a - \log \mu]} \\ &= \frac{\pi^N}{2\pi i} \exp\{N[\mu^*(a)a - \log \mu^*(a)]\}, \end{aligned} \quad (33)$$

having put  $a = A/N$  and defined  $\mu^*(a)$  as the saddle point value, which can be explicitly computed using the saddle point equation:

$$\frac{\partial}{\partial \mu} [\mu a - \log(\mu)] = 0 \quad \implies \quad \mu^* = \frac{1}{a} = \frac{N}{A} = N, \quad (34)$$

having finally imposed the normalization condition:  $A = 1 \implies a = 1/N$ . Since we selected the value of  $\mu$  which corresponds to the correct normalization of eigenvectors, we can simply compute the Laplace transform of the unnormalized marginal  $\rho(\zeta|A)$  and plug in the correct value of  $\mu^*$  in order to estimate its leading behaviour. Namely, we first consider:

$$\rho(r|A) \propto \int_0^\infty \left[ \prod_{i=2}^N dr_i r_i \right] r \delta\left(A - r^2 - \sum_{i=2}^N r_i^2\right) \quad (35)$$

and then its Laplace transform, with the same procedure we already employed:

$$\rho(r|\mu) \propto \theta(r) r e^{-\mu r^2} \xrightarrow{\mu^* = N} \rho(r|N) = \mathcal{N} \theta(r) r e^{-Nr^2}, \quad (36)$$

where  $\theta(\cdot)$  is the Heaviside function and  $\mathcal{N}$  the normalization factor. Therefore, considering that  $r_i = |\zeta_i| = |M_{ki}|$ , it's been shown that the probability distribution of the overlap between one of the normal modes and one of the random ones, in the large- $N$  limit, is the product of a linear function and a Gaussian centered in  $r = 0$  with standard deviation:  $\sigma_g \sim 1/\sqrt{N}$ . We can easily compute the normalization factor of the distribution, as well as the mean value and the variance, by solving Gaussian integrals; the results are:

$$\begin{aligned} A &= 2N \\ \langle r \rangle &= \sqrt{\frac{\pi}{4N}} \\ \sigma^2 &= \langle r^2 \rangle - \langle r \rangle^2 = \left(1 - \frac{\pi}{4}\right) \frac{1}{N} \end{aligned} \quad (37)$$

In conclusion, the probability distribution for the overlaps  $|q|$  (dropping the indices) reads:

$$\rho(|q|) = 2N|q|e^{-N|q|^2} \quad (38)$$

---

\* giacomo.gradenigo@gssi.it

- [1] P. Gaspard, “*Chaos, Scattering and Statistical Mechanics*” (Cambridge University Press, 1998)
- [2] N. Zanghì *I fondamenti concettuali dell’ approccio statistico in fisica*, in *La Natura delle Cose V*. Allori, M. Dorato, F. Laudisa, N. Zanghì (Ed.s), pp. 202 (Carocci Editore, 2005).
- [3] P. Castiglione, M. Falcioni, A. Lesne, A. Vulpiani, “*Chaos and Coarse Graining in Statistical Mechanics*” (Cambridge University Press, 2008).
- [4] A. I. Khinchin, “*Mathematical Foundations of Statistical Mechanics*” (Dover, New York, 1949).
- [5] P. Mazur, J. van der Linden, “*Asymptotic form of the structure function for real systems*”, *J. Math. Phys.* **4**, 271 (1963).
- [6] P. Mazur, E. Montroll, “*Poincaré Cycles, Ergodicity, and Irreversibility in Assemblies of Coupled Harmonic Oscillators*”, *J. Math. Phys.* **1**, 70 (1960).
- [7] R. Livi, M. Pettini, S. Ruffo, A. Vulpiani, “*Chaotic behaviour in nonlinear Hamiltonian systems and equilibrium statistical mechanics*”, *J. Stat. Phys.* **48**, 539 (1987).
- [8] M. Falcioni, U. Marini Bettolo Marconi, A. Vulpiani, “*Ergodic properties of high-dimensional symplectic maps*”, *Phys. Rev. A* **44**, 2263 (1991).
- [9] M. Baldovin, A. Vulpiani, G. Gradenigo, “*Statistical mechanics of an integrable system*”, *Journal of Statistical Physics* **183**, 1-16 (2021).
- [10] H. Spohn, “*Generalized Gibbs Ensembles of the Classical Toda Chain*”, *J. Stat. Phys.* **180**, 4 (2020).
- [11] L. Vidmar and M. Rigol, “*Generalized Gibbs ensemble in integrable lattice models*”, *J. Stat. Mech.* 064007 (2016).
- [12] A. De Luca, G. Mussardo, “*Equilibration properties of classical integrable field theories*”, *J. Stat. Mech.* 064011 (2016).
- [13] G. Del Vecchio Del Vecchio, A. Bastianello, A. De Luca, G. Mussardo, “*Exact out-of-equilibrium steady states in the semiclassical limit of the interacting Bose gas*”, *SciPost Phys.* **9**, 002 (2020).
- [14] E. Fermi, J. Pasta, S. Ulam, “*Studies of nonlinear problems. I.*”, Los Alamos Report, LA-1940 (1955).
- [15] R. Livi, M. Pettini, S. Ruffo, M. Sparpaglione, A. Vulpiani, “*Equipartition threshold in nonlinear large Hamiltoniansystems: The Fermi-Pasta-Ulam model*”, *Phys. Rev. A* **31**, 1039 (1985).
- [16] G. Gallavotti (Ed.), “*The Fermi-Pasta-Ulam problem: a status report*” (Springer, 2008).
- [17] S. Flach, M. V. Ivanchenko, O. I. Kanakov, “*q-Breathers and the Fermi-Pasta-Ulam problem*”, *Phys. Rev. Lett.* **95**, 064102 (2005).
- [18] G. Benettin, H. Christodoulidi, A. Ponno, “*The Fermi-Pasta-Ulam problem and its underlying integrable dynamics*”, *Journ. Stat. Phys.* **152**, 195-212 (2013).
- [19] E. Marinari, G. Parisi, F. Ritort, “*Replica field theory for deterministic models: I. Binary sequences with low autocorrelation*”, *J. Phys. A: Math. Gen.* **27** (23), 7615 (1994); E. Marinari, G. Parisi, F. Ritort, “*Replica field theory for deterministic models. II. A non-random spin glass with glassy behaviour*”, *J. Phys. A: Math. Gen.* **27** (23), 7647 (1994).
- [20] D. Rossini, A. Silva, G. Mussardo and G. E. Santoro, “*Effective Thermal Dynamics Following a Quantum Quench in a Spin Chain*”, *Phys. Rev. Lett.* **102**, 127204 (2009).
- [21] L. F. Cugliandolo, G. S. Lozano, N. Nesi, M. Picco, A. Tartaglia, “*Quenched dynamics of classical isolated systems: the spherical spin model with two-body random interactions or the Neumann integrable model*”, *J. Stat. Mech.* 063206 (2018).
- [22] D. Barbier, L. F. Cugliandolo, G. S. Lozano, N. Nesi, M. Picco, A. Tartaglia, “*Pre-asymptotic dynamics of the infinite size Neumann (p=2 spherical)*”, *J. Phys. A: Math. Theor.* **52**, 454002 (2019).
- [23] D. Barbier, L. F. Cugliandolo, G. S. Lozano, N. Nesi “*(Non equilibrium) Thermodynamics of Integrable models: The Generalized Gibbs Ensemble description of the classical Neumann Model*”, *Europhys. Lett.* **132** 50002 (2020).
- [24] S. Goldstein, J. L. Lebowitz, R. Tumulka, N. Zanghì, “*Any Orthonormal Basis in High Dimension is Uniformly Distributed over the Sphere*”, *Ann. Inst. H. Poincaré Probab. Statist.* **53**: 701-717 (2017).
- [25] M. Mézard, G. Parisi, M.-A. Virasoro, “*Spin glass theory and beyond: An Introduction to the Replica Method and Its Applications.*”, World Scientific Publishing Company (1987).
- [26] L.F. Cugliandolo, J. Kurchan, L. Peliti, “*Energy flow, partial equilibration, and effective temperatures in systems with slow dynamics*”, *Phys. Rev. E* **55** (4), 3898 (1997).
- [27] P. Contucci, J. Kurchan, E. Mingione, “*Equilibrium and dynamics of a multi-bath Sherrington-Kirkpatrick model*”, *J. Phys. A: Math. Theor.* **52**, 324001 (2019).
- [28] J. Kurchan, “*Time-reparametrization invariances, multi-thermalization and the Parisi scheme*”, arXiv:2101.12702 (2021).
- [29] F. Piazza, P. De Los Rios, F. Cecconi, “*Temperature Dependence of Normal Mode Reconstructions of Protein Dynamics*”, *Phys. Rev. Lett.* **102**, 218104 (2004).
- [30] M. Baldovin, G. Gradenigo, A. Vulpiani, “*Statistical Features of High-Dimensional Hamiltonian Systems*”, to appear as a contribution in “*EPS Grand Challenges: Physics for Society at the Horizon 2050*”, arXiv:2106.06609 (2021).
- [31] J. Von Neumann , “*Proof of the Ergodic Theorem and H-Theorem in Quantum Mechanics*”, *Europhys. Jour. H* **35**, 201 (2010).
- [32] M. Srednicki, “*The approach to thermal equilibrium in quantized chaotic systems*”, *J. Phys. A: Math. Gen.* **32**, 1163-1175 (1999).
- [33] L. D’Alessio, Y. Kafri, A. Polkovnikov, M. Rigol, “*From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics*”, *Adv. Phys.* **65** 239-362, (2016).
- [34] S. Pappalardi, L. Foini, J. Kurchan, “*Quantum bounds and fluctuation-dissipation relations*”, arXiv:2110.03497 (2021).