

Abstract

We establish frame inequalities for signals in Paley–Wiener spaces on two specific families of graphs consisting of combinations of cubes and cycles. The frame elements are localizations to cubes, regarded as clusters in the graphs, of vertex functions that are eigenvectors of certain spatio–spectral limiting operators on graph signals. **keywords:** Graph product, Dirichlet eigenvector, graph Fourier transform

AMS subject classifications: 94A12, 94A20, 42C10, 65T99

Sampling low-spectrum signals on graphs via cluster-concentrated modes: examples.

Jeffrey A. Hogan

School of Mathematical and Physical Sciences

University of Newcastle

Callaghan NSW 2308 Australia

email: jeff.hogan@newcastle.edu.au

Joseph D. Lakey

Department of Mathematical Sciences

New Mexico State University

Las Cruces, NM 88003–8001

email: jlakey@nmsu.edu

January 12, 2022

1 Introduction

We investigate here, in the context of signals on graphs, extensions of two specific sampling methods that were mentioned in [3], namely generalized sampling of bandlimited signals (loc. cit. p. 68) cf., [2, 10, 11], and time and band limiting (loc. cit. p. 65), also known as spatio-spectral limiting in geometric contexts. The former is now well established in the context of sampling spectrum-limited signals on graphs while the latter has also been considered in relation to sampling on graphs [15], cf., [6, 7].

We study inner products with eigenvectors of spatio-spectral limiting as generalized sampling measurements for two very specific families of finite graphs involving combinations of cubes and cycles, in which the cubes are regarded as *clusters*, and the cycle is regarded as a skeleton that connects the clusters. Of particular interest is how sampling inequalities—frame inequalities of the form $A\|f\|^2 \leq \sum |\langle f, \psi_j \rangle|^2 \leq B\|f\|^2$ where, for fixed $\{\psi_j\}$, $\langle f, \psi_j \rangle$ are regarded as measurements or generalized samples of vertex functions f in the span of low-spectrum eigenvalues of the graph Laplacian—depend on how the cubes and cycles are combined. This study is motivated by recent work of I. and M. Pesenson [12] who established sampling inequalities on graphs involving a single sampling measurement per cluster. Our study seeks corresponding inequalities involving a number of measurements per cluster according to the number of spectral modes that can be concentrated in each cluster. The reason

to study the specific families of graphs examined here is that these graphs allow an analytic apparatus to estimate or count the number of such modes explicitly. Here is an outline of the contents.

In Sect. 2 we provide background, briefly reviewing generalized sampling in Paley–Wiener spaces on \mathbb{R} in Sect. 2.1, and time and band limiting and the role of prolate functions in generalized sampling in Sect. 2.2. Section 2.3 reviews basic concepts of spectral graph theory and a recent result of I. and M. Pesenson concerning cluster-wise sampling on graphs. We review particular properties of cubes and cycle graphs in Sect. 2.4.

Section 3 considers certain graphs that we denote by $\mathcal{B}_N \vdash C_m$ in which a cube \mathcal{B}_N is substituted for each vertex on a cycle C_m . Each inserted cube is regarded as a cluster and is connected to two other cubes, each by a single edge, to form a cycle of clusters. In Sect. 3.1, eigenvectors (of the graph Laplacians) of these graphs are identified as having either *Dirichlet type*, supported on a single copy of \mathcal{B}_N , or *Neumann-type*. The latter are eigenvectors of a certain *augmented Laplacian* on \mathcal{B}_N modulated by eigenvectors of the cycle. The structure of the eigenvectors of this graph is summarized in Thm. 7. Section 3.2 discusses a certain spatial-limiting operator Q that truncates a vertex function f to a single \mathcal{B}_N -cluster and a spectral-limiting operator P that projects onto the span of the low-eigenvalue eigenvectors of the Laplacian on $\mathcal{B}_N \vdash C_m$. Conjecture 8 describes the eigenvectors of PQ and proposes a complete set of eigenvectors of shifted PQ -operators whose eigenvalues should exceed $1/2$, meaning that the corresponding low-spectrum eigenvectors are concentrated in a single copy of \mathcal{B}_N . Section 3.3 provides a specific example fixing the cube parameter $N = 7$ and cycle parameter $m = 21$.

Section 4 considers Cartesian products $\mathcal{B}_N \square C_m$ of cubes and cycles. Here, each \mathcal{B}_N -slice (fixed cycle coordinate) can be regarded as a cluster but, in contrast to $\mathcal{B}_N \vdash C_m$, each vertex in a cluster is now adjacent to a unique corresponding vertex in each of the two adjacent clusters. Theorem 9 describes orthonormal bases (ONBs) for decompositions of Paley–Wiener spaces on these Cartesian products in terms of components localized on a single cubic slice, versus components that are distributed over all cubic slices. Section 4.1 provides a specific example fixing the same cube parameters $N = 7$ and cycle parameter $m = 21$ for the Cartesian case.

Finally, Sect. 5 discusses how our results for specific graph families might extend to other finite graphs. The Appendix establishes a condition for generalized sampling in the context of spatial and spectral limiting on (standard Cayley graphs of) finite abelian groups, which include $\mathcal{B}_N \square C_m$. The observation is not central to the main theme here, but it provides a general context for an important aspect of the examples presented here. Throughout, notation will be introduced as needed to define quantities that use the notation.

2 Background

2.1 Generalized sampling

Generalized sampling on \mathbb{R} in the sense of Papoulis [11] refers to sampling formulas of the form

$$f(t) = \sum_{v=1}^M \sum_{k=-\infty}^{\infty} z_v(kT) p_v(t - kT), \quad f \in \text{PW}_\Omega(\mathbb{R}) \quad (1)$$

where $\widehat{z}_\nu(\xi) = \widehat{f}(\xi)\widehat{h}_\nu(\xi)$, which implies that $z_\nu(kT) = f * h(kT) = \langle f, h(\cdot - kT) \rangle$ for real filter functions h_ν , $\nu = 1, \dots, M$. The uniform recovery condition developed in Brown [2] is a uniform lower bound on the determinant of the matrix $A_{\nu\ell}(\omega) = \widehat{h}_\nu(\omega + \ell\sigma)$, $\nu = 1, \dots, M$; $\ell = 0, \dots, M - 1$ on the interval $-\Omega \leq \xi \leq -\Omega + \sigma$ where $\sigma = 2\Omega/M$. In this event the reconstruction functions $p_\nu(t)$ have the form $p_\nu(t) = \sum_{\ell=0}^{M-1} \int_{-\Omega}^{\sigma-\Omega} b_{\nu\ell}(\xi) e^{2\pi i(\xi+\ell\sigma)t} d\xi$ where $b_{\nu\ell}$ is the (ν, ℓ) th element of the inverse of the matrix A . A critical part of the argument is that the invertibility condition allows uniform reconstruction of the exponential $e^{2\pi i t \xi}$ on $(-\Omega, \sigma - \Omega)$, then on $(-\Omega, \Omega)$, in the form

$$e^{2\pi i t \xi} = \sum_{\nu=1}^M \widehat{h}_\nu(\xi) \sum_{n=-\infty}^{\infty} y_\nu(t - kT) e^{2\pi i k T \xi}; \quad \widehat{y}_\nu(\xi, t) = \sum_{\ell=0}^{M-1} e^{2\pi i k \sigma t} b_{\nu\ell}(\xi) \quad (2)$$

which is a refinement of the original approach of Papoulis.

2.2 Time and band limiting, prolate functions, and prolate shift frames

A particular case of generalized sampling arises when the filter functions h_ν above are eigenfunctions of time- and band-limiting operators. The bandlimiting operator P_Ω projects onto the Paley–Wiener space $\text{PW}_\Omega = \{f \in L^2(\mathbb{R}) : \widehat{f}(\xi) = 0, |\xi| > \Omega\}$ where $\widehat{f}(\xi) = \int_{-\infty}^{\infty} f(t) e^{-2\pi i t \xi} dt$ when $f \in L^1 \cap L^2(\mathbb{R})$. The time-limiting operator is $(Q_T f)(t) = f(t)$ if $|t| \leq T$ and $(Q_T f)(t) = 0$ if $|t| > T$. Up to a dilation factor, the eigenfunctions of the operator $P_\Omega Q_T$ are the *prolate spheroidal wave functions* (PSWFs) ψ_ν with eigenvalues $1 > \lambda_0 > \lambda_1 > \dots$, e.g., [4]. The eigenvalues of $P_\Omega Q_T$ are the same as the concentrations of L^2 -norm on $[-T, T]$, that is, $\|Q_T \psi\|^2 / \|\psi\|^2$. The famous $2\Omega T$ theorem [9] states that the number of eigenvalues of $P_\Omega Q_{T/2}$ that are *close to one* is approximately $2\Omega T - O(\log(2\Omega T))$ for large enough values of ΩT . This observation is tied to the *spectral accumulation property* of prolates, namely that $\sum_{\nu=0}^{\infty} \lambda_\nu |\widehat{\psi}_\nu(\xi)|^2 = 1$ on $[-\Omega, \Omega]$ (and vanishes for $|\xi| > \Omega$). Another useful property of prolates is that they are orthogonal on $[-T, T]$: $\langle Q_T \psi_n, \psi_m \rangle = \lambda_n \delta_{n,m}$.

Theorem 1 *Let ψ_ν denote the ν th prolate bandlimited to $[-\Omega, \Omega]$ and time-concentrated in $[-1, 1]$. If $N = 2\Omega\alpha \in \mathbb{N}$ then the functions $\{\psi_\nu(\cdot - \alpha k)\}_{\nu=0, k \in \mathbb{Z}}^{N-1}$ form a Riesz basis for PW_Ω . That is, there exist constants $0 < A \leq B < \infty$ such that for any sequence $\{c_{\nu k}\}_{\nu=0, k \in \mathbb{Z}}^{N-1} \in \ell^2(\mathbb{Z}^N)$ one has*

$$A \sum_{\nu k} |c_{\nu k}|^2 \leq \left\| \sum_{\nu k} c_{\nu k} \psi_\nu(\cdot - \alpha k) \right\|^2 \leq B \sum_{\nu k} |c_{\nu k}|^2.$$

If $N \geq \lceil 2\Omega\alpha \rceil$ then the functions $\{\psi_\nu(\cdot - \alpha k)\}_{\nu=0, k \in \mathbb{Z}}^{N-1}$ form a frame for PW_Ω : there exist constants $0 < A \leq B < \infty$ such that for any $f \in \text{PW}_\Omega$ one has

$$A \|f\|_{L^2}^2 \leq \sum_{\nu=0}^{N-1} \sum_{k \in \mathbb{Z}} |\langle f, \psi_\nu(\cdot - \alpha k) \rangle|^2 \leq B \|f\|_{L^2}^2.$$

If $N < \lceil 2\Omega\alpha \rceil$ then the lower frame bound fails.

The Riesz basis case follows along lines similar to the generalized sampling case as in (2), specifically by verifying the matrix invertibility criterion for generalized sampling. It is observed further in [5] that the frame or Riesz basis bounds are most snug when $N \approx 4\Omega$, or $\alpha \approx 2$ so the number of sampling functions used is approximately equal to the time–bandwidth product. In this event, $\sum_{v=0}^{N-1} |\psi_v(\xi)|^2$ is smoothly bounded away from zero on $[-\Omega, \Omega]$ and the series approximation of $e^{2\pi i t \xi}$ by shifted prolates along the lines of (2) is numerically effective.

2.3 Generalized sampling in Paley–Wiener spaces on graphs and clusters

Graphs and their Fourier transforms Let \mathcal{G} be a simple unweighted, undirected connected finite graph with n vertices $V(\mathcal{G})$ and edges $E(\mathcal{G}) \subset V(\mathcal{G}) \times V(\mathcal{G})$. Given an ordering v_1, \dots, v_n of $V(\mathcal{G})$, the adjacency matrix A has (i, j) -entry equal to one if $(v_i, v_j) \in E(\mathcal{G})$ and equal to zero otherwise. We write $v \sim w$ if $(v, w) \in E(\mathcal{G})$. When \mathcal{G} is fixed we will simply write V or E for $V(\mathcal{G})$ and $E(\mathcal{G})$ (and similarly for other quantities that refer to the graph \mathcal{G}). The space of functions $f : V(\mathcal{G}) \rightarrow \mathbb{C}$ is denoted $\ell^2(\mathcal{G})$. When referring to a graph Laplacian on \mathcal{G} we always mean the unnormalized Laplacian operator $L(\mathcal{G})$ that maps a vertex function $f \in \ell^2(\mathcal{G})$ to $(Lf)(v) = \sum_{w \sim v} [f(v) - f(w)]$. If D is the diagonal degree matrix with i th diagonal entry $D_{ii} = \#\{w : (v_i, w) \in E\}$ (called the degree $\deg v_i$ of v_i) then $L = D - A$. It is standard to refer to the eigenvalues of $L(\mathcal{G})$ as the eigenvalues or *spectrum* of \mathcal{G} . The operator L is nonnegative and symmetric. It has a single eigenvalue zero (with constant eigenvector) and norm at most $\max\{\deg v\}_{v \in V}$. We may write the eigenvalues of \mathcal{G} as $0 = \lambda_0(\mathcal{G}) < \lambda_1(\mathcal{G}) \leq \dots \leq \lambda_n(\mathcal{G})$. The graph Fourier transform $F(\mathcal{G})$ is represented by the unitary matrix whose columns are the corresponding unit-normalized eigenvectors of L . We assume that the columns are ordered by increasing Laplacian eigenvalue. Given $\Omega > 0$ the space PW_Ω is defined as the span of the Laplacian eigenvectors having eigenvalue at most Ω . If F_Ω is the matrix consisting of the columns of $F(\mathcal{G})$ with Laplacian eigenvalue at most Ω then the orthogonal projection onto PW_Ω is expressed by the matrix $F_\Omega F_\Omega^*$.

A cluster sampling inequality of I. and M. Pesenson In [12], I. and M. Pesenson consider the problem of sampling and interpolation of low-spectrum information associated with clusters in undirected weighted graphs. Some notation is needed to describe a version of the result of [12] of greatest relevance here. Let \mathcal{G} be a connected finite unweighted graph and $\mathcal{S} = \{S_j\}_{j=1}^m$ a partition of $V(\mathcal{G})$. Let L_j be the Laplace operator of the induced graph \mathcal{G}_j on S_j with first nonzero eigenvalue $\lambda_{1,j}$ and $\varphi_{0,j}$ its zero eigenfunction. Assume for each j a *sampling function* $\psi_j \in \ell^2(\mathcal{G})$ is supported in S_j and satisfies $\langle \psi_j, \varphi_{0,j} \rangle \neq 0$. We assume that $\|\psi_j\| = 1$ and set $\theta_j = 1/|\langle \varphi_{0,j}, \psi_j \rangle|^2$, which is generally larger than one (and equal to one precisely when ψ_j is equal to $\varphi_{0,j}$). The Pesensons set $\Lambda = \inf_j \lambda_{1,j}$ and $\Theta = \sup_j \theta_j$ when the partition $\{S_j\}$ and sampling vectors $\{\psi_j\}$ are fixed. As a consequence of certain Poincaré inequalities, I. and M. Pesenson proved a result that we state as follows for the case considered here.

Theorem 2 For every $f \in \text{PW}_\Omega$, where $0 < \Omega < \Lambda/\Theta$, the Plancherel–Polya inequalities

$$\frac{(1-\mu)\epsilon}{(1+\epsilon)\Theta} \|f\|^2 \leq \sum_{j=1}^m |\langle f, \psi_j \rangle|^2 \leq C \|f\|^2 \quad (3)$$

hold for a fixed $C > 0$ provided $\mu = (1+\epsilon)\frac{\Theta}{\Lambda}\Omega < 1$.

It is also proved that the coefficients $\langle f, \psi_j \rangle$ identify $f \in \text{PW}_\Omega$ uniquely.

The condition on Ω which implies that $\Omega < \Lambda/\Theta = \inf \lambda_{1,j} \inf |\langle \psi_j, \varphi_{0,j} \rangle|^2$ is fairly restrictive since it implies that $\Omega < \lambda_{1,j}$. The Pesensons raise the question how big is the admissible interval $[0, \Lambda/\Theta)$. In the limiting case of a disconnected graph, this interval contains only the zero eigenvalue of each cluster. In the weighted case, if one allows for very weak weighted connections between disjoint clusters, and accounts for continuity of the spectrum with respect to the weights, the admissible interval will then contain a number of eigenvalues equal to the number of clusters.

The Pesensons go on to give examples of sampling functionals defined by ψ_j , including cluster averages, pointwise samples, and variational splines, explaining how the constants in (3) depend on these choices. They then raise the question whether similar results can be obtained for a segment of the spectrum strictly larger than the interval $[0, \Lambda/\Theta)$. This raises a broader question, whether sampling inequalities like (3) for clustered graphs can be extended beyond Paley–Wiener spaces that encode a single eigenvalue per cluster. Stated differently: for a given graph \mathcal{G} and vertex partition $\mathcal{S} = \{S_j\}$, is there a frame for PW_Ω in which each frame element is well concentrated in one of the S_j ? In contrast to the case of a homogeneous setting like the real line in Thm. 1, how sampling or frame inequalities apply will depend on the geometry of the clusters relative to the whole graph. We will consider two similar looking, but structurally different examples in Sects. 3 and 4 to elaborate on this point. First, we make some general remarks about how local versus global geometry is reflected in the Laplacian.

Consistent with classical sampling theorems, in order to obtain sampling bounds for larger values of Ω , including values that account for larger eigenvalues of the cluster Laplacians, requires enough sampling functionals per cluster to account for larger cluster eigenmodes. The unnormalized graph Laplacian L can be expressed as $L = \tilde{L} + \sum_j L_j$ where \tilde{L} accounts for connections between clusters. If v is an *insulated* vertex, meaning that all neighbors of v in G lie in S_j , then $(Lf)(v) = (L_j f)(v)$. More generally, if \tilde{L} has small norm, it suggests that an inequality like (3) should hold on a subspace of PW_Ω accounted by eigenvalues of separate L_j smaller than $\Omega - \|\tilde{L}\|$, leaving the question how to account for global behavior encoded in \tilde{L} .

2.4 Cycles and cubes

In Sects. 3 and 4 we study examples of graphs amenable to analytical methods to study concentration of low-spectrum components on relatively densely connected subsets of vertices. These graphs are combinations of cycles and cubes whose properties we outline now.

The m -cycle C_m is the graph whose vertices are the elements of \mathbb{Z}_m (we denote simply by ℓ the vertex of the element $\ell \in \mathbb{Z}_m$) and adjacency is defined by $\ell \sim \ell'$ if and only if $\ell' = \ell \pm 1 \pmod m$. The Laplacian $L(C_m)$ of C_m is usually represented by the $m \times m$ matrix $2I - A_m$ where A_m is the symmetric matrix with entries equal to $A_m(k, k+1) = 1 = A_m(k-1, k)$ and

$A_m(k, \ell) = 0$ if $|(k - \ell) \bmod m| \neq 1$ (here entries are indexed by $0, \dots, m - 1$ with addition modulo m). The eigenvectors of $L(C_m)$ are, up to normalization, $E_k(\ell) = e^{2\pi i k \ell / m} / \sqrt{m}$ with eigenvalue $2 - 2 \cos(2\pi k / m) = 4 \sin^2(\pi k / m)$. When $m = 2$ these eigenvalues are zero and two. The Fourier transform of C_m is represented by the matrix with columns E_k . When $m = 2$ it is the Haar matrix $H = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$.

The Cartesian product of two graphs \mathcal{G} and \mathcal{H} is the graph $\mathcal{G} \square \mathcal{H}$ with $V(\mathcal{G} \square \mathcal{H}) = V(\mathcal{G}) \times V(\mathcal{H})$ and $E(\mathcal{G} \square \mathcal{H})$ defined by $(u_1, v_1) \sim (u_2, v_2)$ if $u_1 \sim u_2$ and $v_1 = v_2$ or $u_1 = u_2$ and $v_1 \sim v_2$. The Laplacian $L(\mathcal{G} \square \mathcal{H})$ is the Kronecker product $L(\mathcal{G}) \otimes L(\mathcal{H})$ of $L(\mathcal{G})$ and $L(\mathcal{H})$ (if A is $m \times n$ then $A \otimes B$ has $m \times n$ blocks, and (k, ℓ) -th block equal to $a_{k\ell} B$) and therefore the eigenvalues of $L(\mathcal{G} \square \mathcal{H})$ are sums of eigenvalues of $L(\mathcal{G})$ and $L(\mathcal{H})$ (with corresponding multiplicities).

The Boolean cube \mathcal{B}_N is the N -fold Cartesian product of C_2 . It has vertices \mathbb{Z}_2^N where two vertices $v = (\epsilon_1, \dots, \epsilon_N)$ and $w = (\epsilon'_1, \dots, \epsilon'_N)$ are adjacent precisely when $v - w = e_i$ (subtraction in \mathbb{Z}_2^N) for some $i \in \{1, \dots, N\}$ where e_i has one in the i th coordinate and zeros in the other coordinates. The unnormalized Laplacian eigenvalues are $0, 2, \dots, 2N$ where $2k$ has multiplicity $\binom{N}{k}$. The Fourier transform of \mathcal{B}_N is a Hadamard matrix H_N of size $2^N \times 2^N$ whose columns, up to reindexing (reordering of rows and columns) are those of the N th Kronecker product of the Haar matrix H . The Fourier transform of \mathbb{Z}_2^N can also be indexed by elements $\gamma \in \mathbb{Z}_2^N$. In this case, if h_γ is the column of H_N corresponding to $\gamma \in \mathbb{Z}_2^N$ then $h_\gamma(v) = 2^{-N/2} (-1)^{\langle v, \gamma \rangle}$. Further elaboration on these basic concepts on cubes relevant to following discussion can be found in [6, 7]

3 Example 1: Sampling on Vertex substitution of cubes on cycles.

3.1 Vertex substitution of cubes on cycles

Spectral properties of edge substitution graphs were studied by Strichartz in [13] and in [14], where sampling properties were also established. In edge substitution, a copy of a fixed graph \mathcal{H} which has two designated vertices v_0 and v_1 , is placed along each edge of a graph \mathcal{G} in such a way that v_0 and v_1 are identified with the terminals of each corresponding edge. In the case of a 2-regular graph (a cycle) it makes sense, instead, to substitute a fixed graph for each vertex of the cycle in such a way that v_0 and v_1 become terminals of successive edges. We denote such a substitution of \mathcal{H} for each vertex of C_m by $\mathcal{H} \vdash C_m$. The symbol “ \vdash ” is used to indicate the asymmetric roles of the component graphs.

Our first set of examples of structured graphs consisting of several connected clusters are the graphs $\mathcal{B}_N \vdash C_m$, in which each vertex of a cycle is replaced by a copy of \mathcal{B}_N . We will denote by \mathcal{B}_N^k the copy of \mathcal{B}_N placed on the k th vertex of C_m ($k = 0, \dots, m - 1$) and by v_0^k and v_1^k the vertices of \mathcal{B}_N^k corresponding to the zero element $\mathbf{0} = (0, \dots, 0)$ and ones-element $\mathbf{1} = (1, \dots, 1)$ of \mathbb{Z}_2^N . Since the transformation that replaces $v = (\epsilon_1, \dots, \epsilon_N) \in \mathbb{Z}_2^N$ by $\tilde{v} = (1 + \epsilon_1, \dots, 1 + \epsilon_N) \in \mathbb{Z}_2^N$ is an isomorphism of \mathcal{B}_N , it is not really critical whether we assign v_0^k or v_1^k as the neighbor of v_1^{k-1} (resp. v_0^{k+1}), but for simplicity we always assign $v_1^{k-1} \sim v_0^k$ and $v_1^k \sim v_0^{k+1}$, see Fig. 1. Generically we will denote by v^k the copy of $v \in V(\mathcal{B}_N)$

lying in $V(\mathcal{B}_N^k)$. We refer to $\mathcal{B}_N \vdash C_m$ as the *vertex substitution* of \mathcal{B}_N on C_m . As in [13], we will use the term *Dirichlet eigenvector* to refer to an eigenvector of the Laplacian $L(\mathcal{B}_N)$ that vanishes at $\mathbf{0}$ and $\mathbf{1}$.

Denote by ι_k the mapping of a vertex function φ on \mathcal{B}_N to one on $\mathcal{B}_N \vdash C_m$ such that $(\iota_k \varphi)(u^k) = \varphi(u)$, $u \in V(\mathcal{B}_N)$ and $(\iota_k \varphi)(u^\ell) = 0$, $\ell \neq k$. Any vertex in $V(\mathcal{B}_N \vdash C_m)$ that is not a copy of v_0 or v_1 will be called an *insulated vertex*. It is clear that, at any insulated vertex $u^k \in V(\mathcal{B}_N^k)$, for any vertex function $f \in \ell^2(\mathcal{B}_N)$, $(L(\mathcal{B}_N \vdash C_m))(\iota_k f)(u^k) = (\iota_k L(\mathcal{B}_N)f)(u^k)$. The following is then clear.

Lemma 3 *If φ is a Dirichlet eigenvector of $L(\mathcal{B}_N)$ then $\iota_k(\varphi)$ is an eigenvector of $L(\mathcal{B}_N \vdash C_m)$ with the same eigenvalue as the $L(\mathcal{B}_N)$ -eigenvalue of φ , and vanishing outside \mathcal{B}_N^k .*

We will also refer to any function of the form $\iota_k \varphi$, where φ is a Dirichlet eigenvector of $L(\mathcal{B}_N)$, as a Dirichlet eigenvector (of $L(\mathcal{B}_N \vdash C_m)$).




Figure 1: Graphic representation of $\mathcal{B}_2 \vdash C_3$

Lemma 4 *For each fixed K ($0 < K < N$) the span of the $2K$ -Dirichlet eigenvectors on $\mathcal{B}_N \vdash C_m$ has dimension $m \binom{N}{K} - 1$.*

Proof. There are $\binom{N}{K}$ linearly independent (mutually orthogonal) Hadamard eigenvectors h_γ with $L(\mathcal{B}_N)$ -eigenvalue $2K$. Recall that $h_\gamma(v) = 2^{-N/2}(-1)^{\langle v, \gamma \rangle}$. In particular, $h_\gamma(\mathbf{0}) = 2^{-N/2}$ for every γ and $h_\gamma(\mathbf{1}) = 2^{-N/2}(-1)^{\#\{j: \gamma_j=1\}}$ for each $\gamma \in \mathbb{Z}_2^N$. Therefore $h_\gamma(\mathbf{1}) = 2^{-N/2}(-1)^K$ when $\sum \gamma_i = K$ as is the case for each $2K$ -eigenvector of $L(\mathcal{B}_N)$. An element of the span of the $2K$ -eigenvectors has the form $\sum_{|\gamma|=K} c_\gamma h_\gamma$. The closed subspace generated by coefficients $\{c_\gamma\}_{|\gamma|=K} : \sum c_\gamma = 0$ has one-dimensional orthocomplement (c_γ constant) and therefore has dimension $\binom{N}{K} - 1$. This subspace of the $2K$ -eigenspace of $L(\mathcal{B}_N)$ evidently contains the Dirichlet eigenvectors of \mathcal{B}_N . There is one isomorphic copy of this subspace associated with block \mathcal{B}_N^k in $\ell^2(\mathcal{B}_N \vdash C_m)$. Since these copies have disjoint supports, the dimension of the direct sum of these copies is $m \binom{N}{K} - 1$. ■

This raises the question of eigenvalues of $L(\mathcal{B}_N \vdash C_m)$ that are not Dirichlet. It is clear from the proof of Lem. 4 that $L(\mathcal{B}_N)$ itself has $\binom{N}{K} - 1$ Dirichlet eigenvalues equal to $2K$ ($0 < K < N$), leaving one eigenvalue for each K that is not Dirichlet. We claim that the remaining $2K$ -eigenvalue of $L(\mathcal{B}_N)$ has an eigenvector of the form $\varphi = c_K \sum_{|\gamma|=K} h_\gamma$. We refer to such φ as the Neumann eigenvector of $L(\mathcal{B}_N)$ with eigenvalue $2K$.

Lemma 5 *For each fixed K ($0 \leq K \leq N$) the vector $h_{n,K} = c_K \sum_{|\gamma|=K} h_\gamma$ (c_K is chosen to make $\|h_{n,K}\| = 1$) is a Neumann eigenvector of $L(\mathcal{B}_N)$. The vectors $h_{n,K}$ are mutually orthogonal for different K and are orthogonal to the Dirichlet eigenvectors of $L(\mathcal{B}_N)$.*

Proof. Since each h_γ with $|\gamma| = K$ is a $2K$ -eigenvector of $L(\mathcal{B}_N)$, the same holds for $\sum_{|\gamma|=K} h_\gamma$. Since the Hadamard vectors are orthogonal ($\langle h_\gamma, h_{\gamma'} \rangle = \delta_{\gamma, \gamma'}$), it follows that $\langle h_{n,K}, h_{n,K'} \rangle = 0$ if $K \neq K'$. Orthogonality between Dirichlet and Neumann eigenvectors for different K follows in the same way. For the same K , if $\varphi = \sum_{|\gamma|=K} c_\gamma h_\gamma$ is such that $\sum_{|\gamma|=K} c_\gamma = 0$ then

$$\langle \varphi, h_{n,K} \rangle = c_K \sum_{|\gamma|=|\gamma'|=K} c_\gamma \langle h_\gamma, h_{\gamma'} \rangle = c_K \sum_{|\gamma|=|\gamma'|=K} c_\gamma \delta_{\gamma, \gamma'} = c_K \sum_{|\gamma|=K} c_\gamma = 0.$$

This proves that $h_{n,K}$ is orthogonal to any Dirichlet eigenvector and completes the proof of the Lemma. ■

Concatenating $2K$ -Dirichlet eigenvectors supported on \mathcal{B}_N^k , $k = 0, \dots, m-1$ results in a $2K$ -eigenvector of $L(\mathcal{B}_N \uplus C_m)$. Remaining eigenvectors of $L(\mathcal{B}_N \uplus C_m)$ are formed by joining elements of Neumann eigenspaces on the \mathcal{B}_N^k in such a way that the resulting vertex function on $\mathcal{B}_N \uplus C_m$ is a global eigenvector of $L(\mathcal{B}_N \uplus C_m)$. We do this by leveraging the exponential structure of eigenvectors of $L(C_m)$ to define *augmented Laplacian* operators on \mathcal{B}_N whose eigenvectors can be concatenated to form eigenvectors of $L(\mathcal{B}_N \uplus C_m)$.

For $\alpha \in \mathbb{C} \setminus 0$, define a *corner matrix* C_α of size $2^N \times 2^N$ such that

$$C_\alpha(1, 1) = 1 = C_\alpha(2^N, 2^N), \quad C_\alpha(1, 2^N) = -\frac{1}{\alpha}, \quad \text{and} \quad C_\alpha(2^N, 1) = -\alpha$$

and $C_\alpha(i, j) = 0$ otherwise. C_α has a single eigenvalue equal to two and all other eigenvalues equal to zero. Recall that we use lexicographic ordering on \mathcal{B}_N so that for a $2^N \times 2^N$ matrix A acting on $\ell^2(\mathcal{B}_N)$ and vertex function f , the first column of Af is $f(\mathbf{0})$ times the first column of A and the last column of Af is $f(\mathbf{1})$ times the last column of A . Therefore, $C_\alpha f$ is the vector whose first entry is $f(\mathbf{0}) - f(\mathbf{1})/\alpha$, whose last entry is $f(\mathbf{1}) - \alpha f(\mathbf{0})$, and whose other entries are equal to zero. We now assume that $|\alpha| = 1$ and define the augmented Laplacian

$$L_\alpha(\mathcal{B}_N) = L(\mathcal{B}_N) + C_\alpha. \quad (4)$$

At any insulated vertex $v \in V(\mathcal{B}_N)$, $(L_\alpha f)(v) = (Lf)(v)$ ($L = L(\mathcal{B}_N)$). In particular, any $2K$ -Dirichlet eigenvector of L is also a $2K$ -eigenvector of L_α . On the other hand, $(L_\alpha f)(\mathbf{0}) = (Lf)(\mathbf{0}) + f(\mathbf{0}) - f(\mathbf{1})/\alpha$ and $(L_\alpha f)(\mathbf{1}) = (Lf)(\mathbf{1}) + f(\mathbf{1}) - \alpha f(\mathbf{0})$. The non-Dirichlet eigenvectors of L_α can be regarded as perturbations of the Neumann eigenvectors of L . Since the norm of C_α is two and since the output vectors of C_α are zero in all but the first and last rows, whereas the Neumann eigenvectors are nonzero in all rows, the norm of C_α on the Neumann subspace of $\ell^2(\mathcal{B}_N)$ is strictly smaller than two, and each non-Dirichlet eigenvector of L_α can be regarded as a nonnegative perturbation of the Neumann eigenvector of L having the closest eigenvalue. We summarize this analysis of Dirichlet and *Neumann-type* eigenvectors of $L_\alpha(\mathcal{B}_N)$ as follows.

Lemma 6 *Let $|\alpha| = 1$. The operator $L_\alpha(\mathcal{B}_N)$ has a complete set of eigenvectors of the following form: (i) h is a Dirichlet eigenvector of $L(\mathcal{B}_N)$ (with the same eigenvalue), or (ii) $h = \sum_{\kappa=0}^N c_\kappa(\alpha) h_{n,\kappa}$. There are $N+1$ mutually orthogonal eigenvectors of $L_\alpha(\mathcal{B}_N)$ of the second type.*

Proof. The different eigenvectors of type (ii) have eigenvalue $2K + \epsilon_K$ ($\epsilon_K \in (0, 2)$) for different $K \in \{0, \dots, N\}$ and are therefore mutually orthogonal. ■

Augmented Laplacians and eigenvectors of $L(\mathcal{B}_N \vdash C_m)$ The eigenvectors of $L(C_m)$ are $E_\nu = \{e^{2\pi i \nu \ell / m} / \sqrt{m}\}_{\ell=0}^{m-1}$. For $f \in \ell^2(\mathcal{B}_N \vdash C_m)$ one has

$$(L(\mathcal{B}_N \vdash C_m)f)(v_0^k) = (L(\mathcal{B}_N^k)f)(v_0^k) + [f(v_0^k) - f(v_1^{k-1})]$$

and similarly

$$(L(\mathcal{B}_N \vdash C_m)f)(v_1^k) = (L(\mathcal{B}_N^k)f)(v_1^k) + [f(v_1^k) - f(v_0^{k+1})],$$

while $(L(\mathcal{B}_N \vdash C_m)f)(v) = (L(\mathcal{B}_N^k)f)(v)$ for any insulated vertex $v \in V(\mathcal{B}_N^k)$. Suppose now that the sample vectors $\{f(v_0^k)\}_{k=0}^{m-1}$ and $\{f(v_1^k)\}_{k=0}^{m-1}$ form eigenvectors of $L(C_m)$. Then there is a value $\nu \in \{0, \dots, m-1\}$ such that $f(v_0^{k+1}) = e^{2\pi i \nu / m} f(v_0^k)$. Assume that, for the same ν , we also have $f(v_1^k) = e^{2\pi i \nu / m} f(v_1^{k-1})$. In this event, for $\alpha = e^{2\pi i \nu / m}$ one then has

$$(L(\mathcal{B}_N \vdash C_m)f)(v_0^k) = (L(\mathcal{B}_N^k)f)(v_0^k) + [f(v_0^k) - e^{-2\pi i \nu / m} f(v_1^k)] = L_\alpha(\mathcal{B}_N^k)f)(v_0^k)$$

and similarly

$$(L(\mathcal{B}_N \vdash C_m)f)(v_1^k) = (L(\mathcal{B}_N^k)f)(v_1^k) + [f(v_1^k) - e^{2\pi i \nu / m} f(v_0^k)] = L_\alpha(\mathcal{B}_N^k)f)(v_1^k).$$

Suppose now that the restriction of φ to \mathcal{B}_N^0 is an eigenvector of $L_\alpha(\mathcal{B}_N^0)$ where $\alpha = e^{2\pi i \nu / m}$, in other words, this restriction has the form $\iota_0 f$ where f is an eigenvector of $L_\alpha(\mathcal{B}_N)$, and suppose that the restriction of φ to \mathcal{B}_N^k is equal to $e^{2\pi i \nu k / m} \iota_k f$ for the same f , for each $k = 0, \dots, m$. We conclude from the observations above then that φ is an eigenvector of $L(\mathcal{B}_N \vdash C_m)$ whose eigenvalue is equal to the $L_\alpha(\mathcal{B}_N)$ -eigenvalue of f for $\alpha = e^{2\pi i \nu / m}$. We can use Lem. 6 to describe a complete family of eigenvectors of $L(\mathcal{B}_N \vdash C_m)$ as follows.

Theorem 7 *The eigenvectors φ of $L(\mathcal{B}_N \vdash C_m)$ have one of the following types:*

(i) *Dirichlet type: φ is supported in \mathcal{B}_N^k for some $k \in \mathbb{Z}_m$ and is a Dirichlet $2K$ -eigenvector of $L(\mathcal{B}_N^k)$ for a fixed value of $K \in \{0, 1, \dots, N\}$, or*

(ii) *Neumann type: the sample vectors $\{\varphi(v_0^k)\}$ and $\{\varphi(v_1^k)\}$, $k = 0, \dots, m-1$, are eigenvectors of $L(C_m)$ and the restriction of φ to $\mathcal{B}_N^{(k)}$ is an eigenvector of $L_\alpha(\mathcal{B}_N)$, $\alpha = e^{2\pi i \nu / m}$ for a fixed $\nu \in \mathbb{Z}_m$.*

The number of linearly independent $2K$ -eigenvectors of Dirichlet type is $m\binom{N}{K} - 1$ for each $K = 0, \dots, N$ and the number of linearly independent eigenvectors of Neumann type is $m(N+1)$, with values indexed by K and ν . Together, the linearly independent sets of eigenvectors of Dirichlet or Neumann type form a complete set of $m2^N$ eigenvectors of $L(\mathcal{B}_N \vdash C_m)$.

It is worth mentioning that for each group $K = 0, \dots, N$ there is a collection of m eigenvectors of Neumann type whose eigenvalues are of the form $2K + \epsilon_{K,\nu}$ where $\epsilon_{K,\nu} \in (0, 2)$, see Fig. 2. These eigenvectors have the form $\sum_{\kappa=0}^N c_\kappa(\alpha) h_{n,\kappa}$ on \mathcal{B}_N^k .

of Thm. 7. By Lem. 4, there are $\binom{N}{K} - 1$ linearly independent Dirichlet $2K$ -eigenvectors of $L(\mathcal{B}_N)$. Since images $\iota_k \varphi$ of such eigenvectors for different k have disjoint supports and since $(L(\mathcal{B}_N \vdash C_m)\iota_k \varphi) = \iota_k(L(\mathcal{B}_N)\varphi)$ for a Dirichlet eigenvector, it follows that there are $m\binom{N}{K} - 1$ linearly independent eigenvectors of Dirichlet type. The total number of linearly independent eigenvectors of Dirichlet type is $\sum_{K=0}^N m\binom{N}{K} - 1 = m(2^N - N - 1)$.

The analysis above shows that if φ is an eigenvector of $L_\alpha(\mathcal{B}_N)$ of norm one with $\alpha = e^{2\pi iv/m}$ then the vector whose values on \mathcal{B}_N^k are of the form $e^{2\pi ivk/m} \iota_k \varphi$ are eigenvectors of $L(\mathcal{B}_N \uplus C_m)$. By Lem. 6, there are $N + 1$ linearly independent Neumann-type eigenvectors of $L_\alpha(\mathcal{B}_N)$ for each such α and, when $\varphi_0, \dots, \varphi_N$ is a linearly independent set of such vectors, the corresponding eigenvectors $e^{2\pi ivk/m} \iota_k \varphi_j$ also form a linearly independent set of $m(N + 1)$ Neumann-type elements of $\ell^2(\mathcal{B}_N \uplus C_m)$. Since Dirichlet and Neumann type eigenvectors are linearly independent, adding the total numbers of eigenvectors of each type one obtains a collection of $m2^N = \dim(\ell^2(\mathcal{B}_N \uplus C_m))$ eigenvectors. Therefore these eigenvectors are complete in $\ell^2(\mathcal{B}_N \uplus C_m)$. ■

3.2 Band limiting and block limiting on $\mathcal{B}_N \uplus C_m$

On $\mathcal{B}_N \uplus C_m$ let P_Ω denote the operator that projects $f \in \ell^2(\mathcal{B}_N \uplus C_m)$ onto the span of the eigenvectors of $L(\mathcal{B}_N \uplus C_m)$ having eigenvalue at most Ω and let Q denote the operator that truncates $f \in \ell^2(\mathcal{B}_N \uplus C_m)$ to \mathcal{B}_N^0 by multiplying by zero outside \mathcal{B}_N^0 . Since P_Ω and Q are orthogonal projections, the norm of $P_\Omega Q$ is at most one. Since any Dirichlet eigenvector of $L(\mathcal{B}_N^0)$ having eigenvalue at most Ω is in the range of P_Ω , there are nontrivial unit eigenvectors of $P_\Omega Q$ when $\Omega \geq 2$. Contrast this to the case of the real line where eigenvalues of the corresponding iterated project operator $P_\Omega(\mathbb{R})Q(\mathbb{R})$ —where $P_\Omega(\mathbb{R})$ bandlimits elements of $L^2(\mathbb{R})$ to $[-\Omega, \Omega]$ and $Q(\mathbb{R})$ truncates elements of $L^2(\mathbb{R})$ to $[-1, 1]$ —has all eigenvalues strictly smaller than one. On the other hand, if $f \in \text{PW}_\Omega(\mathcal{B}_N \uplus C_m)$ is in the span of the Neumann-type eigenvectors of $L(\mathcal{B}_N \uplus C_m)$ identified in Thm. 7, then f is not supported in \mathcal{B}_N^0 so, if it is an eigenvector of $P_\Omega Q$, then the eigenvalue is strictly smaller than one.

The analysis in Thm. 7 indicates that the dimension of PW_{2K} is equal to

$$m \sum_{\kappa=0}^K \left(\binom{N}{\kappa} - 1 \right) + m \sum_{\kappa=0}^{K-1} 1 = m \left(-1 + \sum_{\kappa=0}^K \binom{N}{\kappa} \right) = m(\text{dim}_K - 1) \quad (5)$$

where $\text{dim}_K = \sum_{\kappa=0}^K \binom{N}{\kappa}$ is the volume of the K -ball in \mathcal{B}_N . The first terms in (5) come from Dirichlet eigenvectors that are perfectly localized on \mathcal{B}_N^k for each $k = 0, \dots, m - 1$. Those localized on \mathcal{B}_N^0 are also eigenvectors of PQ having eigenvalue equal to one. Thus PQ has $\sum_{\kappa=0}^K (\binom{N}{\kappa} - 1)$ eigenvalues equal to one. The second terms in (5) correspond to Neumann-type eigenvectors whose norms are equally distributed over all blocks \mathcal{B}_N^k . The example below suggests that there should be K Neumann-type eigenvalues of PQ that are *close to one*, see Conj. 8, whose proof requires showing that there are K Neumann-type eigenvalues of PQ larger than $1/2$. In turn, it requires proving that the vector consisting of the samples $\{\varphi(v^k)\}$ where v^k is the \mathcal{B}_N^k -representative of fixed $v \in \mathcal{B}_M$, is nearly cardinal for the corresponding eigenvectors φ , with the zero-block value equal to at least half the ℓ^2 -norm of the sample vector.

The conjecture implies that when $\Omega = 2K \in 2\mathbb{N}$, there exists a basis of eigenvectors ψ of PQ for $\text{PW}_{2K}(\mathcal{B}_N \uplus C_m)$, most of whose eigenvalues are equal to one, and the rest having eigenvalues close to one. By associating an operator $PQ^{(k)}$ to each separate cluster \mathcal{B}_N^k then, the conjecture implies that there is a basis of $\text{PW}_{2K}(\mathcal{B}_N \uplus C_m)$ of vectors that are either perfectly or predominantly localized on a single cluster. For such a basis one will have inequalities of

the form

$$\mu_K \|Q^{(k)} f\| \leq \sum_{j=1}^{\dim_K - 1} |\langle Q^{(k)} f, \psi_{j,k} \rangle|^2 \leq \|Q^{(k)} f\|^2, \quad f \in \text{PW}_{2K} \quad (6)$$

uniformly in the block parameter k , where μ_K is the minimum eigenvalue of the Neumann-type eigenvectors of PQ .

We compare these observations with Thm. 2, which identifies one measurement per cluster to provide a sampling criterion for PW_Ω on a generic graph \mathcal{G} , but requiring Ω to be sufficiently small. While graphs $\mathcal{B}_N \uplus C_m$ form an extremely specific family of graphs, a consequence of (6) is that on each block \mathcal{B}_N^k there is an orthonormal family of *measurement* vectors $\psi_{j,k}$ in PW_{2K} that are also orthogonal on \mathcal{B}_N^k , such that $\dim_K - (K + 1)$ of the vectors are fully supported in \mathcal{B}_N^k , while the remaining K vectors are concentrated in \mathcal{B}_N^k , and such that most of the ℓ^2 -norm of $f \in \text{PW}_{2K}$ on \mathcal{B}_N^k is accounted for by the measurements $\langle Q^{(k)} f, \psi_{j,k} \rangle$. In the specific case outlined in Sect. 3.3, the value of μ_K in (6) is much larger than $1/2$.

To describe the measurement vectors $\psi_{j,k}$ we introduce the notation $f \uplus g$ for vertex functions f and g defined on C_m and \mathcal{B}_N respectively, to denote the function defined on $\mathcal{B}_N \uplus C$ that takes the value $f(k) \iota_k g(v)$ at the image of v in \mathcal{B}_N^k . Neumann type eigenvectors on $L(\mathcal{B}_N \uplus C_m)$ described in Thm. 7 have the form

$$E_\nu \uplus \sum_{\kappa=0}^N c_\kappa(\alpha, K) h_{n,\kappa} \equiv E_\nu \uplus \varphi_{\nu,K,n} \quad (7)$$

where $\{h_{n,\kappa}\}_{\kappa=0}^N$ forms an ONB for the Neumann eigenvectors of $L(\mathcal{B}_N)$ and $\varphi_{\nu,K,n} = \sum_{\kappa=0}^N c_\kappa(\alpha, K) \varphi_{n,\kappa}$ is an eigenvector of $L_\alpha(\mathcal{B}_N)$ with $\alpha = e^{2\pi i \nu / m}$ as described in Thm. 7. The index K in (7) indexes the $L(\mathcal{B}_N \uplus C_m)$ -eigenvalue of $E_\nu \uplus \varphi_{\nu,K,n}$: as suggested by Thm. 7, for each $K \in \{0, \dots, N\}$, and each ν , there is a unique Neumann-type $L(\mathcal{B}_N \uplus C_m)$ -eigenvalue $2K + \epsilon_\nu$ with $\epsilon_\nu \in (0, 2)$ corresponding to the unique $L_\alpha(\mathcal{B}_N)$ -eigenvalue of the form $2K + \epsilon$. Neumann-type eigenvectors of PQ then have the form

$$\psi_{\nu,K,n} = \sum_\nu \sum_{\kappa=0}^{K-1} \beta_{\nu,\kappa} E_\nu \uplus \varphi_{\nu,\kappa,n} = \sum_\nu E_\nu \uplus \sum_{\kappa=0}^{K-1} \beta_{\nu,\kappa} \varphi_{\nu,\kappa,n} \equiv \sum_\nu E_\nu \uplus \psi_{\nu,K}$$

where P is the orthogonal projection onto $\text{PW}_{2K}(\mathcal{B}_N \uplus C_m)$. Observe that in this case the $\mathcal{B}_N \uplus C_m$ Neumann-type components $\varphi_{\nu,\kappa,n}$ contributing to $\psi_{\nu,K,n}$ themselves lie in $\text{PW}_{2K}(\mathcal{B}_N \uplus C_m)$.

Conjecture 8 *Let $0 < K < N$ and let P, Q be defined as above. There are $\dim_K - 1$ eigenvalues of PQ larger than $1/2$, including $\dim_K - (K + 1)$ eigenvalues equal to one and another K eigenvalues in $(1/2, 1)$. The cyclic shifts of the corresponding eigenvectors are linearly independent in $\ell^2(\mathcal{B}_N \uplus C_m)$. Therefore, these vectors form a basis for PW_{2K} which has dimension $m(\dim_K - 1)$.*

That the Neumann-type PQ eigenvectors should have eigenvalues larger than $1/2$ would rely on a sort of *uniform cardinality* property of the sequences of 2^N th samples of $\psi_{\nu,k,n} = \sum_\nu E_\nu \uplus \psi_{\nu,K}$, see Fig. 5 for the case $N = 7, m = 21$.




Figure 2: First 620 eigenvalues of $L(\mathcal{B}_7 + C_{21})$




Figure 3: Eigenvalues of PQ on $\mathcal{B}_7 + C_{21}$. First 60 eigenvalues equal one, corresponding to Dirichlet eigenvectors of $L(\mathcal{B}_7)$




Figure 4: Eigenvectors of PQ on $\mathcal{B}_7 \vdash C_{21}$ corresponding to eigenvalue indices 32, 62 and 64 whose respective PQ eigenvalues are 1, 0.9982 and 0.0148. The first three cycle lengths ($3 \cdot 128$) are plotted on the left. Their Fourier coefficients are plotted on the right. Eigenvector 32 has its Fourier support in the Dirichlet eigenvalues while the other eigenvectors have their Fourier supports in the Neumann-type eigenvalues.




Figure 5: Sample values of eigenvectors of PQ on $\mathcal{B}_7 \vdash C_{21}$ corresponding to eigenvalue indices 61–64. Each group plots the set of sample values $\varphi_\nu(k + 128 * \ell)$, $\ell = 0, \dots, 20$, $k = 1, \dots, 128$ cyclically shifted by a fixed amount for each $\nu = 61$ (left) to $\nu = 64$ (right). The samples are essentially cardinal (peak at the center and zeros elsewhere) for $\nu = 61$ and become *less cardinal* to the point of containing an oscillation for $\nu = 64$

3.3 Specific case $N = 7, K = 3, m = 21$ of $\mathcal{B}_N \vdash C_m$

When $m = 21, N = 7$ and $K = 3$ there are $m(\dim_K - 1) = 21 \cdot (-1 + \sum_{\kappa=0}^3 \binom{7}{\kappa}) = 21 \cdot 63 = 1323$ eigenvalues of $L(\mathcal{B}_N \vdash C_m)$ with value at most 6. Most of these come from the $21(\binom{7}{1} + \binom{7}{2} + \binom{7}{3} - 3) = 21 \cdot 60$ $2K$ -Dirichlet eigenvalues of $L(\mathcal{B}_7 \vdash C_{21})$ for $K = 0, \dots, 3$. As shown in Fig. 2, the zero eigenvalue and groups of $m(\binom{N}{k} - 1) = 21(\binom{7}{k} - 1)$ Dirichlet eigenvalues equal to $2k$ are punctuated by transition intervals of length 21 between the eigenvalues $2k$ and $2(k + 1)$ ($k = 0, \dots, 2$). The eigenvalues of PQ , where P is the projection onto PW_6 , are plotted in Fig. 3. There are 60 Dirichlet eigenvalues of PQ equal to one, and three additional eigenvalues close to one. Typical eigenvectors for the eigenvalues equal to one, close to one, and close to zero, and their graph Fourier transforms, are plotted in Fig. 4. The eigenvalues close to one have eigenvectors whose samples $\varphi(v^k)$, where $v^k \in \mathcal{B}_N^k$, correspond to fixed $v \in \mathcal{B}_N$, are nearly cardinal. These samples are plotted in Fig. 5.

The matrix whose columns are cyclic shifts of the Neumann-type PQ -eigenvectors numerically has full rank. Since these shifts are orthogonal to corresponding shifts of Dirichlet eigenvectors of PQ , together these shifts form a basis for PW_6 .

4 Example 2: Sampling on the Cartesian product of a cube and a cycle




Figure 6: Graphic representation of $\mathcal{B}_2 \square C_3$

In the Cartesian product $\mathcal{B}_N \square C_m$ of a cube and a cycle, each copy of the cube can be regarded as a cluster (to which we refer as a \mathcal{B}_N slice). In contrast to the graphs formed by substituting cubes for vertices of a cycle, in the Cartesian product with a cycle, each vertex in a single cluster is connected to a vertex in each of two adjacent clusters, see Fig. 6. This prohibits Dirichlet eigenvectors of the Laplacian $L(\mathcal{B}_N \square C_m)$. Nonetheless, $\mathcal{B}_N \square C_m$ is $(N + 2)$ -regular with N neighbors coming from inside a cluster, so if N is large then most of the degree of each vertex is associated with neighbors within the cluster.

The eigenvalues of $\mathcal{B}_N \square C_m$ are sums of eigenvalues of \mathcal{B}_N and of C_m [1], thus have the form $2k_1 + 4 \sin^2(\pi k_2/m)$, $0 \leq k_1 \leq N, 0 \leq k_2 < m$. Suppose that $K \geq 3$. Since the

eigenvalues of C_m are at most four, the $\mathcal{B}_N \square C_m$ -Laplacian eigenvalues that are less than or equal to $2K$ are sums of the form (i) a \mathcal{B}_N -eigenvalue at most $2(K-2)$ plus any C_m -eigenvalue; (ii) a \mathcal{B}_N -eigenvalue at most $2(K-1)$ plus any C_m -eigenvalue with value at most two, or (iii) a \mathcal{B}_N -eigenvalue equal to $2K$ plus the zero eigenvalue of C_m . The $\mathcal{B}_N \square C_m$ -eigenvectors are tensor products of \mathcal{B}_N -eigenvectors and C_m -eigenvectors of the corresponding types, whose eigenvalues add to at most $2K$. Formally, letting $\Lambda_\nu(\mathcal{G})$ denote the λ_ν -eigenspace of $L(\mathcal{G})$, we can write

$$\begin{aligned} \text{PW}_{2K}(\mathcal{B}_N \square C_m) &= [\text{PW}_{2K-4}(\mathcal{B}_N) \otimes \ell^2(C_m)] \\ &\oplus [\Lambda_{2K-2}(\mathcal{B}_N) \otimes \text{PW}_2(C_m)] \\ &\oplus [\Lambda_{2K}(\mathcal{B}_N) \otimes \Lambda_0(C_m)]. \end{aligned} \quad (8)$$

The dimension of the space $\text{PW}_{2K}(\mathcal{B}_N \square C_m)$ is equal to the number of elements of a basis for PW_{2K} . In the next few paragraphs we count the dimensions of the three components in (8), and discuss the behavior of the operator PQ that truncates to a \mathcal{B}_N slice then bandlimits. Specifically, let Q denote the operator that truncates a vertex function f to the zero slice $\mathcal{B}_N \square \{0\}$ of $\mathcal{B}_N \square C_m$. That is, $(Qf)(v) = f(v)$ if $v = (u, 0)$ for some vertex u of \mathcal{B}_N , and $(Qf)(v) = 0$, otherwise. For a fixed $K \in \{1, \dots, N-1\}$, let P denote the operator that bandlimits f to PW_{2K} . That is, $Pf = \sum_{\lambda_\nu \leq 2K} \sum_{j=1}^{\dim \Lambda_\nu} \langle f, h_{\nu,j} \rangle h_{\nu,j}$ where $\{h_{\nu,j}\}_{j=1}^{\dim \Lambda_\nu}$ is an orthonormal basis of eigenvectors of the λ_ν -eigenspace Λ_ν of $L(\mathcal{B}_N \square C_m)$.

Since $L(C_m)$ has norm at most four, $\text{PW}_4(C_m) = \ell^2(C_m)$. In particular, the delta function $\delta(\ell)$ on C_m is in $\text{PW}_4(C_m)$. It follows that if a vertex function $f \in \text{PW}_{2K-4}$ is supported in the zero- \mathcal{B}_N slice of $\mathcal{B}_N \square C_m$ then $PQf = f$. The space $\text{PW}_{2K-4}(\mathcal{B}_N)$ has dimension $\dim_{K-2} = \sum_{k=0}^{K-2} \binom{N}{k}$. Since each \mathcal{B}_N slice contributes such a subspace of $\text{PW}_{2K}(\mathcal{B}_N \square C_m)$, the component $\text{PW}_{2K-4}(\mathcal{B}_N) \otimes \ell^2(C_m)$ of $\text{PW}_{2K}(\mathcal{B}_N \square C_m)$ in (8) has dimension $m(\dim_{K-2})$.

Next, consider the component $\Lambda_{2(K-1)}(\mathcal{B}_N) \otimes \text{PW}_2(C_m)$ of $\text{PW}_{2K}(\mathcal{B}_N \square C_m)$. For explicit calculations below we assume that m has the form $4m' + 1$, but the stated results hold for other values of m . The factor $\text{PW}_2(C_m)$ has a basis of exponentials $\{e^{2\pi i k \ell / m}\}$ such that $|k \bmod m| \leq \lfloor m/4 \rfloor$. There are $\lfloor (m+1)/2 \rfloor$ such terms. Thus the space generated by corresponding tensor products has dimension $\lfloor (m+1)/2 \rfloor \binom{N}{K-1}$.

Consider eigenvectors of PQ that lie in this component. Those having the largest possible eigenvalue of PQ are products of elements of $\Lambda_{2(K-1)}(\mathcal{B}_N)$ and elements of $\text{PW}_2(C_m)$ having the largest possible value at zero. This value is that of the unit-normalized Dirichlet kernel defining $\text{PW}_2(C_m)$ by convolution in $\ell^2(C_m)$. Since $\{e^{2\pi i k \ell / m}\}_{\ell=0}^{m-1}$ has $L(C_m)$ -eigenvalue $4 \sin^2(\pi k / m)$, the finite Dirichlet kernel

$$D_n(\ell) = \sum_{k=-n}^n e^{2\pi i k \ell / m} = \frac{\sin(2\pi(n+1/2)\ell/m)}{\sin(\pi\ell/m)}$$

lies in $\text{PW}_2(C_m)$ provided that $n \leq \lfloor m/4 \rfloor$. The $\ell^2(C_m)$ -norm of D_n is $m(2n+1)$, so $\bar{D}_n = D_n / \sqrt{m(2n+1)}$ has unit norm and $\bar{D}_n(0) = \sqrt{(2n+1)/m}$. When $n = m' = (m-1)/4$, one has $\bar{D}_n \in \text{PW}_2(C_m)$ and $\bar{D}_n(0) = \sqrt{1/2 + 1/(2m)}$. If $f(u, \ell) = \psi(u) \bar{D}_{m'}(\ell)$ is in the $(2K-2)$ -eigenspace of the \mathcal{B}_N -Laplacian on $\mathcal{B}_N \square \{0\}$ then $f \in \text{PW}_{2K}$ and $PQ(\bar{D}_{m'}\psi) = P_{2,C_m}(\delta_{0,\ell} \bar{D}_{m'}(\ell)) P_{2K-2,\mathcal{B}_N} \psi(u) = \bar{D}_{m'}(0) \bar{D}_{m'}(\ell) \psi(u) = \bar{D}_{m'}(0) f$ where $\bar{D}_{m'}(0) = \sqrt{\frac{1}{2} + \frac{1}{2m}}$ and $P_{\lambda,\mathcal{G}}$ denotes the projection of $\ell^2(\mathcal{G})$ onto $\text{PW}_\lambda(\mathcal{G})$. The space of such f has dimension

$\binom{N}{K-1}$. The subspace of $\ell^2(C_m)$ spanned by the cyclic shifts $D_n(\cdot - \ell)$ has dimension $2n + 1$ when $n \leq \lfloor (m-1)/2 \rfloor$. To see this, observe that the Fourier transform of $D_n(\cdot - k)$ is $e^{-2\pi i \xi k/m} \widehat{D}_n(\xi) = e^{-2\pi i \xi k/m} \mathbb{1}_{|\xi \bmod m| \leq n}(\xi)$. Since $2n + 1 \leq m$, the span of the truncated exponentials $\{e^{-2\pi i \xi k/m} \mathbb{1}_{|\xi \bmod m| \leq n}(\xi)\}_k$ has dimension at most $2n + 1$ and so the same holds for the shifts $D_n(\cdot - k)$. In the case $n = m' = (m-1)/4$, $2m' + 1 = \lfloor (m+1)/2 \rfloor$. The space generated by the cyclic shifts $\{D_{m'}(\cdot - \ell)\psi(u) : \psi \in \Lambda_{2(K-1)}(\mathcal{B}_N)\}$ then has dimension $\lfloor (m+1)/2 \rfloor \binom{N}{K-1}$.

Finally, if the restriction of f to $\mathcal{B}_N \square \{0\}$ is in the $2K$ -eigenspace Λ_{2K} of the \mathcal{B}_N -Laplacian then the only way in which f itself can lie in PW_{2K} is if f is the tensor product of its zero slice with the constant function equal to one on C_m : $f(u, \ell) = c \mathbb{1}_{C_m}(\ell)\psi(u)$. The space of such functions has dimension $\binom{N}{K}$. In this case, $PQf = P(\delta_{0,\ell}f) = cP_{0,C_m}(\delta_0)P_{2K,\mathcal{B}_N}\psi = c\frac{1}{\sqrt{m}}\mathbb{1}_{C_m}(\ell)\psi(u) = \frac{1}{\sqrt{m}}f$.

According to (8), combining bases for $\text{PW}_{2(K-2)}(\mathcal{B}_N) \otimes \ell^2(C_m)$, for $\Lambda_{2K-2}(\mathcal{B}_N) \otimes \text{PW}_2(C_m)$, and for $\Lambda_{2K}(\mathcal{B}_N) \otimes \text{PW}_0(C_m)$, produces a basis for PW_{2K} . Its dimension is thus $m(\sum_{k=0}^{K-2} \binom{N}{K-k}) + \lfloor (m+1)/2 \rfloor \binom{N}{K-1} + \binom{N}{K}$. We collect the observations above.

Theorem 9 *For $0 < K < N$ the space $\text{PW}_{2K}(\mathcal{B}_N \square C_m)$ is the orthogonal direct sum of the spaces (i) $\text{PW}_{2K-4}(\mathcal{B}_N) \otimes \ell^2(C_m)$, (ii) $\Lambda_{2K-2}(\mathcal{B}_N) \otimes \text{PW}_2(C_m)$, and (iii) $\Lambda_{2K}(\mathcal{B}_N) \otimes \Lambda_0(C_m)$. One has the following:*

1. *The space $\text{PW}_{2K-4}(\mathcal{B}_N) \otimes \ell^2(C_m)$ has dimension $m \sum_{\kappa=0}^{K-2} \binom{N}{\kappa}$. An orthonormal basis for this space consists of cyclic shifts of the vectors $\delta_0(\ell)\psi_{\kappa,\nu}(u)$ where $\{\psi_{\kappa,\nu}\}_\nu$ is an ONB for $\Lambda_{2\kappa}(\mathcal{B}_N)$. On the zero- \mathcal{B}_N slice, for any f in this space one has*

$$\|Qf\|^2 = \sum_{\kappa=0}^{K-2} \sum_{\nu=1}^{\dim \Lambda_\kappa} |\langle f, \psi_{\kappa,\nu} \rangle|^2.$$

2. *The space $\Lambda_{2K-2}(\mathcal{B}_N) \otimes \text{PW}_2(C_m)$ has dimension $\lfloor (m+1)/2 \rfloor \binom{N}{K-1}$ when m has the form $m = 4m' + 1$. A basis of this space consists of the cyclic shifts $\widehat{D}_{m'}(\cdot - 2\ell)\psi_{K-1,\nu}$, $\ell = 0, \dots, (m-1)/2$, $\nu = 1, \dots, \dim \Lambda_{2(K-1)}$. On the zero- \mathcal{B}_N slice, for any f in this space one has*

$$\left(\frac{1}{2} + \frac{1}{2m}\right)\|Qf\|^2 = \sum_{\nu=1}^{\dim \Lambda_{2K-2}} |\langle f, \psi_{K-1,\nu} \rangle|^2.$$

3. *The space $\Lambda_{2K}(\mathcal{B}_N) \otimes \text{PW}_0(C_m)$ has dimension $\binom{N}{K}$. An orthonormal basis of this space consists of $\frac{1}{\sqrt{m}}\mathbb{1}_{C_m}(\ell)\psi_{K,\nu}(u)$, $\nu = 1, \dots, \binom{N}{K}$ where $\{\psi_{K,\nu}\}$ is an ONB for $\Lambda_{2K}(\mathcal{B}_N)$. On the zero- \mathcal{B}_N slice, for any f in this space one has*

$$\frac{1}{m}\|Qf\|^2 = \sum_{\nu=1}^{\dim \Lambda_{2K}} |\langle f, \psi_{K,\nu} \rangle|^2.$$

In contrast to the reconstruction functions p_ν in (1), the functions $Q\psi_{k,\nu}$ are not themselves in PW_{2K} when $k = K-1$ or $k = K$. However, the $Q\psi_{k,\nu}$ are mutually orthogonal on $\mathcal{B}_N \square \{0\}$ and have unique extensions $\psi_{k,\nu} \in \text{PW}_{2K}$.

In comparison with Thm. 2, Thm. 9 identifies three separate regimes (localized on a slice, concentrated on a slice, and equally distributed over all slices)—in a very specific case, in which a graph \mathcal{G} might satisfy a Plancherel–Polya type inequality on each cluster, in each regime, but the corresponding constants (1, $1/2 + 1/(2m)$ and $1/m$ respectively reflect the nature of the regime.

4.1 Specific case $N = 7, K = 3, m = 21$ of $\mathcal{B}_N \square C_m$

As we did in the case of $\mathcal{B}_N \vdash C_m$, we fix $N = 7$ and $K = 3$. The dimension of $\text{PW}_6(\mathcal{B}_3 \square C_{21})$ is

$$21 \left(\binom{3}{0} + \binom{3}{1} \right) + \lfloor (21+1)/2 \rfloor \binom{7}{2} + \binom{7}{3} = 21 \cdot 8 + 11 \cdot 21 + 35 = 434.$$

The corresponding Laplacian eigenvalues that are at most six are shown in Fig. 7. The 64 nonzero eigenvalues of PQ are plotted in Fig. 8.

Consider, on the other hand, the eigenvalues of PQ as described above. In the case $N = 7, K = 3$ and $m = 21$ there are 8 eigenvalues of PQ equal to one, 21 equal to $1/2 + 1/42$, and 35 equal to $1/21$. The remaining eigenvalues are equal to zero.




Figure 7: First 620 eigenvalues of $L(\mathcal{B}_7 \square C_{21})$. Eigenvalues of $L(\mathcal{B}_7 \vdash C_{21})$ are plotted (dashed) for comparison

5 Discussion and conclusions

We have considered two simple families of graphs, $\mathcal{B}_N \vdash C_m$ and $\mathcal{B}_N \square C_m$ in which cubes \mathcal{B}_N are thought of as clusters connected by a cycle. A very rough measure of *clusterness* of a subgraph determined by a subset of vertices, most of whose neighbors are in the subset, is the ratio of the average number of neighbors within the cluster to the average number of neighbors in the whole graph. In the case of $\mathcal{B}_N \square C_m$ this ratio is $N/(N+2)$ whereas in the case of $\mathcal{B}_N \vdash C_m$ the ratio is $(N2^N)/(N2^N+2) = N/(N+2^{1-N})$, which is much closer to one.




Figure 8: Eigenvalues of PQ on $\mathcal{B}_7 \square C_{21}$. Corresponding eigenvalues of PQ on $\mathcal{B}_7 \vdash C_{21}$ are plotted (dashed) for comparison




Figure 9: Eigenvectors of PQ on $\mathcal{B}_7 \square C_{21}$ corresponding to eigenvalue indices 1 (top), 11 and 30 (bottom) whose respective eigenvalues are 1, $1/2 + 1/42$ and $1/21$. The first three block lengths ($3 * 128$) are plotted. The bottom eigenvector is periodic of period 128.

In the case of $\mathcal{B}_N \vdash C_m$, a majority of Laplacian eigenvectors are completely supported in a cluster. This allows for effective sampling inequalities in Paley–Wiener spaces in which samples are inner products with eigenvectors of spatio–spectral limiting PQ -operators.

In the case of $\mathcal{B}_N \square C_m$ there are three regimes: one in which eigenvectors of PQ are completely localized in clusters, a second in which more than half of the energy of eigenvectors of PQ is concentrated in a cluster, and one in which the corresponding eigenvectors of PQ are equally spread over the full graph. (One could avoid this third regime by limiting the spectrum to Laplacian eigenvalues strictly smaller than $2K$ for some K .) The measurement vectors in Thm. 9, which have the form $Q\psi$ for eigenvectors of PQ , thus are not all themselves in the Paley–Wiener space as they are in [12], but as localizations of eigenvectors of PQ , they are mutually orthogonal on the cluster.

We acknowledge that $\mathcal{B}_N \vdash C_m$ and $\mathcal{B}_N \square C_m$ are far from being graphs that occur in real networks. However, the three regimes governing concentration of elements of Paley–Wiener spaces on clusters in the example of $\mathcal{B}_N \square C_m$ should be a feature in graphs whose clusters have induced Laplacian eigenvalues that are separated in magnitude on the order of the norm of the *skeleton* graph in which each cluster is compressed down to a single vertex.

Appendix: PQ on a finite abelian group and spectral accumulation

In Sect. 2.1 it was observed that generalized sampling expansions involve expansion of the exponential Fourier kernel $e^{2\pi i t \xi}$ in terms of inversion of a matrix with entries indexed by Fourier transforms of sampling convolvers (2). The particular case when these convolvers are eigenfunctions of time and band limiting that we outlined in Sect. 2.2 has an analogous version in the more general setting of locally compact abelian groups, and we outline the case of finite abelian groups here.

Let G be a finite abelian group. By the fundamental theorem of finite abelian groups, G is a product $\prod_{\nu=1}^N \mathbb{Z}_{m_\nu}$ of cycles of lengths that divide the order $|G|$ of G . A Fourier basis consists of tensor products of normalized exponential vectors $\{e^{2\pi i \ell k_\nu / m_\nu} / \sqrt{m_\nu}\}_{\ell=0}^{m_\nu-1}$ indexed by $k_\nu = 0, \dots, m_\nu - 1$. The Fourier transform elements form a dual group \widehat{G} under componentwise multiplication. \widehat{G} and G are isomorphic. We make use of these facts to establish a *spectral accumulation property* of spatio–spectral localization operators on G (see [8] for the case on \mathbb{R} , which is a well-known). In what follow we assume indexings of the vertex elements $s \in G$ and $\sigma \in \widehat{G}$. We denote by $F(s, \sigma)$ the matrix of the Fourier transform of G with respect to these indexings such that the columns (fixed σ) are pairwise orthonormal: $FF^* = I_{|G|}$. Since the entries are products of normalized exponentials, $|F(s, \sigma)|^2 = 1/|G|$.

Suppose that S and Σ are symmetric subsets of G, \widehat{G} (e.g., $S = -S$). Let P be the orthogonal projection onto the span of the vectors $\{F(\cdot, \sigma) : \sigma \in \Sigma\}$ and let Q be pointwise multiplication by the characteristic function of S . Then PQ is self-adjoint and has rank equal to $|\Sigma|$ if $|S| \geq |\Sigma|$. Let $\mu_1 \geq \mu_2 \geq \dots \geq \mu_{|\Sigma|}$ be an ordering of the nonzero eigenvalues PQ with corresponding eigenfunctions φ_ν so that (by Mercer’s theorem) $K(s, t) = \sum_{\nu=1}^{|\Sigma|} \varphi_\nu(s) \overline{\varphi_\nu(t)}$ is the kernel of

PQ . For each $\sigma \in \Sigma$ one has

$$F(s, \sigma) = \sum_{\nu} \langle F(\cdot, \sigma), \varphi_{\nu} \rangle \varphi_{\nu}(s) = \sum_{\nu} \frac{1}{\mu_{\nu}} \langle F(\cdot, \sigma), Q\varphi_{\nu} \rangle \varphi_{\nu}(s)$$

since $PQ\varphi_{\nu} = \mu_{\nu}\varphi_{\nu}$ and $F(\cdot, \sigma)PQ\varphi_{\nu} = F(\cdot, \sigma)Q\varphi_{\nu}$ for $\sigma \in \Sigma$.

Therefore,

$$\begin{aligned} \sum \mu_{\nu} |F\varphi_{\nu}(\sigma)|^2 &= \sum \frac{1}{\mu_{\nu}} |F(PQ\varphi_{\nu})(\sigma)|^2 = \sum \frac{1}{\mu_{\nu}} \mathbb{1}_{\Sigma} |F(Q\varphi_{\nu})(\sigma)|^2 \\ &= \sum \frac{1}{\mu_{\nu}} \mathbb{1}_{\Sigma} F(Q\varphi_{\nu})(\sigma) \overline{F(Q\varphi_{\nu})(\sigma)} \\ &= \sum \frac{1}{\mu_{\nu}} \mathbb{1}_{\Sigma} \langle F(\cdot, \sigma), (Q\varphi_{\nu})(\cdot) \rangle \overline{\langle F(\cdot, \sigma), (Q\varphi_{\nu})(\cdot) \rangle} \\ &= \mathbb{1}_{\Sigma} \left\langle \sum_{\nu} \frac{1}{\mu_{\nu}} \langle F(\cdot, \sigma), Q\varphi_{\nu} \rangle \varphi_{\nu}, QF(\cdot, \sigma) \right\rangle \\ &= \mathbb{1}_{\Sigma} \langle F(\cdot, \sigma), \mathbb{1}_S F(\cdot, \sigma) \rangle = \mathbb{1}_{\Sigma}(\sigma) \sum_{s \in S} F(s, \sigma) \overline{F(s, \sigma)} \\ &= \frac{1}{|G|} \mathbb{1}_{\Sigma}(\sigma) \sum \mathbb{1}_S = \frac{1}{|G|} \mathbb{1}_{\Sigma}(\sigma) |S| \end{aligned}$$

using that Q is self adjoint. The group structure played a critical role in that the kernel of the Fourier matrix is made of elements of modulus equal to one.

We summarize the calculation above in the following.

Proposition 10 *Let G be a finite abelian group with dual \widehat{G} and let S and Σ be symmetric subsets of G and \widehat{G} respectively with $|S| \geq |\Sigma|$. Denote by P the orthogonal projection onto the span of the Fourier vectors indexed by Σ and let $(Qf)(s) = \mathbb{1}_S(s)f(s)$ such that $\mu_1 \geq \dots \geq \mu_{|\Sigma|}$ are the eigenvalues of PQ with eigenvectors $\varphi_1, \dots, \varphi_{|\Sigma|}$.*

$$\sum \mu_{\nu} |F\varphi_{\nu}(\sigma)|^2 = \frac{1}{|G|} \mathbb{1}_{\Sigma}(\sigma) |S|$$

References

- [1] A.E. Brouwer and W.H. Haemers, *Spectra of graphs*, Universitext, Springer, New York, 2012.
- [2] J. Brown, *Multi-channel sampling of low-pass signals*, IEEE Transactions on Circuits and Systems **28** (1981), no. 2, 101–106.
- [3] J.R. Higgins, *Five short stories about the cardinal series*, Bull. Amer. Math. Soc. (N.S.) **12** (1985), 45–89.
- [4] J.A. Hogan and J. Lakey, *Duration and Bandwidth Limiting. Prolate Functions, Sampling, and Applications.*, Birkhäuser, Boston, MA, 2012.
- [5] J.A. Hogan and J. Lakey, *Frame properties of shifts of prolate spheroidal wave functions*, Applied and Computational Harmonic Analysis **39** (2015), 21–32.

- [6] J.A. Hogan and J. Lakey, *An analogue of Slepian vectors on Boolean hypercubes*, J. Fourier Anal. Appl. **25** (2019), no. 4, 2004–2020.
- [7] ———, *Spatio-spectral limiting on Boolean cubes*, J. Fourier Anal. Appl. **27** (2021), 40.
- [8] J.A. Hogan and J.D. Lakey, *Time–Frequency and Time–Scale Methods*, Birkhäuser Boston Inc., Boston, MA, 2005.
- [9] H.J. Landau and H. Widom, *Eigenvalue distribution of time and frequency limiting*, J. Math. Anal. Appl. **77** (1980), 469–481.
- [10] A. Papoulis, *Systems and transforms with application in optics*, McGraw-Hill, New York, 1968.
- [11] A. Papoulis, *Generalized sampling expansion*, IEEE Trans. Circuits and Systems **24** (1977), 652–654.
- [12] I.Z. Pesenson and M.Z. Pesenson, *Graph signal sampling and interpolation based on clusters and averages*, J. Fourier Anal. Appl. **27** (2021), no. 3, Paper No. 39, 28. MR 4248652
- [13] R.S. Strichartz, *Transformation of spectra of graph Laplacians*, Rocky Mountain J. Math. **40** (2010), no. 6, 2037–2062. MR 2764237
- [14] ———, *Half sampling on bipartite graphs*, J. Fourier Anal. Appl. **22** (2016), no. 5, 1157–1173. MR 3547716
- [15] M. Tsitsvero, S. Barbarossa, and P. Di Lorenzo, *Signals on graphs: Uncertainty principle and sampling*, IEEE Trans. Signal Process. **64** (2016), 4845–4860.