

# TESTING SEMIPARAMETRIC

## MODEL–EQUIVALENCE HYPOTHESES BASED ON THE CHARACTERISTIC FUNCTION

FEIFEI CHEN<sup>a</sup>, SIMOS G. MEINTANIS<sup>b,c,1</sup>, LIXING ZHU<sup>a,c</sup>

<sup>a</sup>*Center for Statistics and Data Science, Beijing Normal University, Zhuhai, PRC*

<sup>b</sup>*Department of Economics, National and Kapodistrian University of Athens, Athens,  
Greece*

<sup>c</sup>*Pure and Applied Analytics, North–West University, Potchefstroom, South Africa*

February 7, 2022

**Abstract.** We propose three test criteria each of which is appropriate for testing, respectively, the equivalence hypotheses of symmetry, of homogeneity, and of independence, with multivariate data. All quantities have the common feature of involving weighted–type distances between characteristic functions and are convenient from the computational point of view if the weight function is properly chosen. The asymptotic behavior of the tests under the null hypothesis is investigated, and numerical studies are conducted in order to examine the performance of the criteria in finite samples.

*Keywords.* Neighborhood–of–model validation; Precise hypotheses test; Characteristic function; Independence testing; Symmetry testing; Two–sample problem.

## 1 Introduction

In a parametric framework the term “precise hypothesis” is mostly used in bioequivalence when the null hypothesis  $\vartheta = \vartheta_0$  about a given parameter  $\vartheta$  is being tested against the alternative  $\vartheta \neq \vartheta_0$ . The issue of whether a point null hypothesis such as this is reasonable or if a less precise hypothesis would be more appropriate has a long

---

<sup>1</sup>On sabbatical leave from the University of Athens.

history and is being strongly debated often in connection to the so-called “Lindley’s Paradox”. This paradox stated in rough terms says that as the sample size increases any frequentist approach tends to lean towards the alternative while from the Bayesian perspective the data will become even more inclined towards the null hypothesis; see for instance Sprenger (2013), and Berger and Delampady (1987). In view of this debate some researchers have opted towards reformulation of the problem as one of testing the (im)precise null hypothesis  $|\vartheta - \vartheta_0| \leq \Delta$  against the alternative  $|\vartheta - \vartheta_0| > \Delta$ , for some fixed  $\Delta > 0$ . (Note that  $\Delta$  can be chosen as a shrinking neighborhood of the sample size, analogously to the common practice of power analysis by means of contiguous alternatives; Borovkov, 1998, §49 and §55).

Recently there is even a tendency for interchange of the null and the alternative hypothesis so that rejection of the former indicates that the parameter  $\vartheta$  is “ $\Delta$ -close” to  $\vartheta_0$ . This formulation, which has become standard by now, also has its roots in the area of bioequivalence problems which are distinguished from conventional testing problems by the form of the hypotheses to be established. For example, in the specific case of testing for bioequivalence of a new treatment against an established treatment, an obvious choice is a parameter measuring the margin of improvement. Then a threshold is set on this parameter so that rejection of the null hypothesis, that is when the margin of improvement lies in a neighborhood within the threshold, this implies bioequivalence of the two treatments, while rejection implies that either the new is better than the old or the opposite, with the margin of improvement left unspecified. In this restricted context of inference about a parameter there is considerable literature for which the monograph of Wellek (2010) is a good starting point. On the other hand the problem of testing “model equivalence” has only recently been considered in a more general context, beyond that of testing about a single parameter. We refer to Dette and Munk (1998), Janssen (2000), Freitag et al. (2007), Baringhaus et al. (2017), Dette et al. (2018), Möllenhoff et al. (2019), and Henze and Mayer (2020), for testing model equivalence (or “neighborhood-of-model validation”) in various settings.

In this paper we will restate the three major semiparametric problems of distributional homogeneity (or the two-sample problem), symmetry and independence in the form of model equivalence hypotheses testing, and suggest population and sample

measures by means of which these tests can be performed. Specifically for a given model, say  $\mathcal{M}$ , and an arbitrary random vector  $Z$  we propose the (model-specific) population measure

$$(1.1) \quad \Delta_{w,Z}^{(\mathcal{M})} = \int |D_Z^{(\mathcal{M})}(t)|^2 w(t) dt,$$

which satisfies  $\Delta_{w,Z}^{(\mathcal{M})} = 0$ , if and only if the law of random vector  $Z$  lies within model  $\mathcal{M}$ , where  $D_Z^{(\mathcal{M})}(\cdot)$  is a pointwise distance from the null hypothesis, and  $w > 0$  denotes a weight function to be further discussed along the paper. The composition of the vector  $Z$  will vary depending on the context, consisting of a single vector for the case of testing for symmetry, or of two independent random vectors of equal dimensions in the case of homogeneity, or consisting of a pair of random vectors of potentially different dimensions in the case of independence testing. Also note that herewith we are using the word “model” as being synonymous to symmetry of a given random variable, while for two random variables or pairs of variables signifies distributional homogeneity or independence of these variables, respectively. In this framework we consider the model-equivalence problem of testing the null hypothesis

$$(1.2) \quad \mathcal{H}_0^{(\mathcal{M})} : \Delta_{w,Z}^{(\mathcal{M})} \geq \Delta,$$

against the alternative

$$(1.3) \quad \mathcal{H}_1^{(\mathcal{M})} : \Delta_{w,Z}^{(\mathcal{M})} < \Delta,$$

so that if  $\mathcal{H}_0^{(\mathcal{M})}$  is false this will imply that the law of  $Z$  lies within a neighborhood of “model”  $\mathcal{M}$  of fixed length  $\Delta > 0$ . (Formally, the null hypothesis  $\mathcal{H}_0^{(\mathcal{M})}$  as well as the alternative should be indexed by  $\Delta$ , but we will suppress this dependence for simplicity).

The rest of the paper unfolds as follows. In Section 2 we introduce the null and alternative hypotheses, propose an appropriate population measure  $\Delta_{w,Z}^{(\mathcal{M})}$  in each case, and formulate our testing approach in general terms. In Section 3, the three test criteria are specified along with the weight function  $w(\cdot)$  and asymptotic results are presented. The finite-sample properties of the test criteria are studied in Section 4 by means of Monte Carlo experiments. The paper concludes in Section 5 with discussion and outlook. Technical proofs are postponed to the Appendix.

## 2 Population measures, computations and test criteria

### 2.1 Population measures

We will consider population measures that are formulated as  $L_2$ -type distances, and measure discrepancy between symmetry and asymmetry of a given law, while for pairs of distributions they express distance between these distributions, or distance between independence and arbitrary modes of dependence of corresponding random variables. These population measures are formulated in terms of characteristic functions (CFs). Specifically suppose that  $Z := \{X, Y\}$  where  $X \in \mathbb{R}^p$  and  $Y \in \mathbb{R}^p$  are two independent random vectors, and write  $\varphi_X(\cdot)$  and  $\varphi_Y(\cdot)$  for the corresponding CFs. Then we define the CF-distance between  $X$  and  $Y$  as

$$(2.1) \quad \Delta_{w,Z}^{(\mathcal{H})} = \int_{\mathbb{R}^p} |\varphi_X(t) - \varphi_Y(t)|^2 w(t) dt.$$

We wish to test the null hypothesis figuring in (1.2) against the alternative hypothesis (1.3) by means of an empirical counterpart of  $\Delta_{w,Z}^{(\mathcal{H})}$ . In this setting rejection of  $\mathcal{H}_0^{(\mathcal{M})}$  would imply that the laws of the two random variables are equivalent, or  $\Delta$ -close, in terms of CFs.

Note that the law of a random vector  $X \in \mathbb{R}^p$ ,  $p \geq 1$ , is symmetric around the origin if and only if  $\text{Im}(\varphi_X(t)) = 0, \forall t \in \mathbb{R}^p$ , where  $\text{Im}(z)$  denotes the imaginary part of a complex number  $z$ . Then the next population measure corresponds to testing symmetry is defined by setting  $Z := X$  in (1.1), as

$$(2.2) \quad \Delta_{w,Z}^{(\mathcal{S})} = \int_{\mathbb{R}^p} \{\text{Im}(\varphi_X(t))\}^2 w(t) dt.$$

Again rejection of the null hypothesis would imply that the law of  $X$  belongs to the equivalence class of all centrally symmetric distributions, in the sense of being within a (CF-based) neighborhood of symmetry of size  $\Delta$ .

Our last population measure will be used to measure the significance of dependence between a pair of random vectors  $X \in \mathbb{R}^p$  and  $Y \in \mathbb{R}^q$ . To this end set  $Z := (X^\top, Y^\top)^\top$ , write  $\varphi_{X,Y}(\cdot, \cdot)$  for the CF of  $Z$  and define

$$(2.3) \quad \Delta_{w,Z}^{(\mathcal{I})} = \int_{\mathbb{R}^{p+q}} |\varphi_{X,Y}(t_1, t_2) - \varphi_X(t_1)\varphi_Y(t_2)|^2 w(t_1, t_2) dt_1 dt_2,$$

where  $w(t_1, t_2) = w_p(t_1)w_q(t_2)$ , with each  $w_p(\cdot)$  and  $w_q(\cdot)$  being a weight function in the indicated dimension.

## 2.2 Specific computations

We note that CF-based population measures such as these figuring in (2.1)–(2.3) may be further manipulated to express distance in terms of more conventional quantities. The readers are referred to Chen et al. (2019), §3 for such interpretations. For analogous interpretations based on moments see Székely et al. (2007) and Székely and Rizzo (2013). By way of example we consider here the homogeneity distance. In this connection further below we will assume that the weight function is a density of a spherical distribution but for our present purposes also assume that this density may be written as  $w(t) = c\varphi_V^2(t)$ , where  $c > 0$  is a global constant and  $\varphi_V(t)$  denotes the CF of a random vector  $V$  having a symmetric around zero distribution. A representation like this is possible for a few distributions such as the  $p$ -dimensional Kotz-type distribution with density  $f(x) = c \exp(-2\|x\|^\gamma)$ ,  $0 < \gamma \leq 2$ , and the generalized spherical Laplace distribution with density  $f(x) = c(1 + \|x\|^2)^{-(p+1)/2}$ . Here  $\|\cdot\|$  denotes the Euclidean norm. In the Kotz-case,  $V$  corresponds to a spherical stable distribution with CF  $\exp(-\|t\|^\gamma)$ , while in the latter case to a generalized spherical Laplace distribution with CF  $(1 + \|t\|^2)^{-(p+1)/4}$ . We refer to Nadarajah (2003) and Kozubowski et al. (2013) for the Kotz-type distribution and the Laplace distribution, respectively.

With such a weight function and by using Parseval's identity we have from (2.1),

$$\begin{aligned} \Delta_{w,Z}^{(\mathcal{H})} &= \int_{\mathbb{R}^p} |\varphi_X(t) - \varphi_Y(t)|^2 c \varphi_V^2(t) dt \\ &= c \int_{\mathbb{R}^p} |\varphi_{X*V}(t) - \varphi_{Y*V}(t)|^2 dt \\ &= c(2\pi)^p \int_{\mathbb{R}^p} (f_{X*V}(x) - f_{Y*V}(x))^2 dx, \end{aligned}$$

where  $*$  denotes convolution. Thus apart from a constant, the CF-based homogeneity distance between a pair of distributions is equal to a distance associated with the densities of these distributions, whereby the weight function  $w(t)$  acts on these densities as convolution with the density corresponding to the CF  $\varphi_V(t)$  involved in  $w(t)$ .

We nevertheless note that, for example, measures of independence such as those

based on distribution functions or densities are not always easy to interpret, almost to the same extent that CF-based measures are. On the other hand, correlation, which is arguably the most straightforward measure of independence, has been shown by Székely et al. (2007) to be directly related to distance covariance. We will not repeat the arguments here but simply compute the independence measure for a bivariate zero-mean Gaussian distribution with unit component variances and correlation equal to  $\varrho$ , say  $\Delta_{\varrho,G}^{(\mathcal{I})}$ . Specifically with weight function  $w(t) = \exp(-t_1^2 - t_2^2)$  we obtain from (2.3),

$$\begin{aligned} \Delta_{\varrho,G}^{(\mathcal{I})} &= \int_{\mathbb{R}^2} \left| \exp\left(-\frac{t_1^2 + t_2^2 + 2\varrho t_1 t_2}{2}\right) - \exp\left(-\frac{t_1^2 + t_2^2}{2}\right) \right|^2 \exp(-t_1^2 - t_2^2) dt_1 dt_2 \\ &= \pi \left( \frac{1}{2} + \frac{1}{\sqrt{4 - \varrho^2}} - \frac{4}{\sqrt{16 - \varrho^2}} \right), \end{aligned}$$

which is increasing with  $|\varrho|$  and vanishes at  $\varrho = 0$ . It is also immediate from (2.1) that  $\Delta_{\varrho,G}^{(\mathcal{I})}$  coincides with the population homogeneity distance  $\Delta_{\varrho,G}^{(\mathcal{H})}$  between the aforementioned bivariate Gaussian distribution and a corresponding bivariate zero-mean Gaussian distribution with independent components, i.e. with  $\varrho = 0$ . In both cases the parameter of interest is equal to  $\varrho$ , with smaller (larger)  $|\varrho|$  implying smaller (larger) deviation from the corresponding exact property of independence or homogeneity, and thus values of  $\Delta_{\varrho,G}^{(\mathcal{I})}$  and  $\Delta_{\varrho,G}^{(\mathcal{H})}$  admit a simple interpretation in terms of the corresponding parameter of interest.

As a further example in arbitrary dimension  $p \geq 1$ , we consider a two-component equal mixture of a standard Gaussian distribution with the same Gaussian distribution shifted by  $\delta \in \mathbb{R}^p$ . For weight function  $\exp(-\|t\|^2)$  we compute from (2.2) the resulting symmetry distance for this mixture distribution as

$$\Delta_{\delta,G}^{(\mathcal{S})} = \int_{\mathbb{R}^p} \frac{1 - \cos(2\delta^\top t)}{8} \exp(-2\|t\|^2) dt = \frac{\pi^{\frac{p}{2}}}{2^{\frac{p}{2}+3}} \left( 1 - \exp\left(-\frac{\|\delta\|^2}{2}\right) \right),$$

which is again an easily interpretable function of the parameter of interest  $\delta$ . The corresponding homogeneity distance between this mixture and a standard Gaussian distribution may also be computed as

$$\Delta_{\delta,G}^{(\mathcal{H})} = \frac{\pi^{\frac{p}{2}}}{2^{\frac{p}{2}+1}} \left( 1 - \exp\left(-\frac{\|\delta\|^2}{8}\right) \right),$$

and the same observations apply. Thus, and although these examples are clearly only indicative, it appears that when a certain property (symmetry, homogeneity or inde-

pendence) is expressed via a given parameter, the corresponding distance measure is a function (perhaps complicated) having properties that are in line with intuition regarding the role of this parameter. We close this paragraph by stressing that the range of possible values of  $\Delta_{w,Z}^{(\mathcal{M})}$ , when placed in the appropriate context, is important in order to acquire a feeling about meaningful values of the threshold parameter  $\Delta$ , which is again heavily context-dependent; see for instance Wellek (2010, §1.7). Yet this range is not always as straightforward to compute as in the previous examples, and in fact may require high-dimensional numerical integration.

### 2.3 Test criteria

Now assume that  $\Delta_{n,w}^{(\mathcal{M})}$  is an empirical counterpart of  $\Delta_{w,Z}^{(\mathcal{M})}$  resulting from a sample of size  $n$  on  $Z$ , such that

$$(2.4) \quad \sqrt{n} \left( \Delta_{n,w}^{(\mathcal{M})} - \Delta_{w,Z}^{(\mathcal{M})} \right) \xrightarrow{\mathcal{D}} \mathcal{N} \left( 0, \sigma_w^{2(\mathcal{M})} \right)$$

when  $\Delta_{w,Z}^{(\mathcal{M})} > 0$ , and

$$(2.5) \quad \sqrt{n} \Delta_{n,w}^{(\mathcal{M})} \xrightarrow{\mathbb{P}} 0$$

when  $\Delta_{w,Z}^{(\mathcal{M})} = 0$ . Here  $\xrightarrow{\mathcal{D}}$  and  $\xrightarrow{\mathbb{P}}$  signify convergence in distribution and in probability, respectively. Suppose further that the limit variance  $\sigma_w^{2(\mathcal{M})}$  (again we economize on notation by suppressing dependence of  $\sigma_w^{2(\mathcal{M})}$  on  $Z$ ) can be estimated by, say,  $\sigma_{n,w}^{2(\mathcal{M})}$ , such that

$$(2.6) \quad \sigma_{n,w}^{2(\mathcal{M})} \xrightarrow{\mathbb{P}} \sigma_w^{2(\mathcal{M})}.$$

This limit variance  $\sigma_w^{2(\mathcal{M})}$  as well as its consistent estimator  $\sigma_{n,w}^{2(\mathcal{M})}$  will be discussed in detail in Section 3.

On the basis of the preceding discussion and for some fixed threshold  $\Delta > 0$ , we suggest the test defined by the critical region

$$(2.7) \quad \Delta_{n,w}^{(\mathcal{M})} \leq \Delta + \frac{\sigma_{n,w}^{(\mathcal{M})}}{\sqrt{n}} z_\alpha,$$

where  $\sigma_{n,w}^{(\mathcal{M})}$  is the positive square root of  $\sigma_{n,w}^{2(\mathcal{M})}$ , and  $z_\alpha$  is the  $\alpha$  quantile of the standard normal distribution. Clearly rejection of  $\mathcal{H}_0^{(\mathcal{M})}$  implies model equivalence, or more

precisely that the law of our data is  $\Delta$ -close to model  $\mathcal{M}$  under the corresponding distance measure  $\Delta_{w,Z}^{(\mathcal{M})}$ . In fact it readily follows that the criterion with critical region defined by (2.7) leads to an asymptotic size- $\alpha$  test, which is consistent against fixed alternatives. Specifically for each law of  $Z$  with corresponding population measure  $\Delta_{w,Z}^{(\mathcal{M})} > 0$ , if (2.4) and (2.6) hold true, we have for the probability of rejection of the null hypothesis in (1.2) that

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \mathbb{P} \left( \text{Reject } \mathcal{H}_0^{(\mathcal{M})} \mid \Delta_{w,Z}^{(\mathcal{M})} > 0 \right) \\
&= \lim_{n \rightarrow \infty} \mathbb{P} \left( \Delta_{n,w}^{(\mathcal{M})} \leq \Delta + \frac{\sigma_{n,w}^{(\mathcal{M})}}{\sqrt{n}} z_\alpha \mid \Delta_{w,Z}^{(\mathcal{M})} > 0 \right) \\
&= \lim_{n \rightarrow \infty} \mathbb{P} \left( \frac{\sqrt{n}}{\sigma_{n,w}^{(\mathcal{M})}} \left( \Delta_{n,w}^{(\mathcal{M})} - \Delta_{w,Z}^{(\mathcal{M})} \right) \leq \frac{\sqrt{n}}{\sigma_{n,w}^{(\mathcal{M})}} \left( \Delta - \Delta_{w,Z}^{(\mathcal{M})} \right) + z_\alpha \mid \Delta_{w,Z}^{(\mathcal{M})} > 0 \right) \\
(2.8) \quad &= \begin{cases} 0, & \text{if } \Delta_{w,Z}^{(\mathcal{M})} > \Delta, \\ \alpha, & \text{if } \Delta_{w,Z}^{(\mathcal{M})} = \Delta, \\ 1, & \text{if } \Delta_{w,Z}^{(\mathcal{M})} < \Delta. \end{cases}
\end{aligned}$$

In addition, for each law of  $Z$  with  $\Delta_{w,Z}^{(\mathcal{M})} = 0$ , and by (2.5) and (2.6), we have

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \mathbb{P} \left( \text{Reject } \mathcal{H}_0^{(\mathcal{M})} \mid \Delta_{w,Z}^{(\mathcal{M})} = 0 \right) \\
&= \lim_{n \rightarrow \infty} \mathbb{P} \left( \sqrt{n} \Delta_{n,w}^{(\mathcal{M})} \leq \sqrt{n} \Delta + \sigma_{n,w}^{(\mathcal{M})} z_\alpha \mid \Delta_{w,Z}^{(\mathcal{M})} = 0 \right) \\
(2.9) \quad &= 1.
\end{aligned}$$

**Remark 2.1** Note that the first two cases in (2.8) correspond to null hypotheses, while the last case in (2.8) and the case in (2.9) correspond to alternatives. Therefore, if (2.4)–(2.6) hold true, we have

$$\lim_{n \rightarrow \infty} \mathbb{P}(\text{Reject } \mathcal{H}_0^{(\mathcal{M})} \mid \mathcal{H}_0^{(\mathcal{M})}) \leq \alpha, \quad \lim_{n \rightarrow \infty} \mathbb{P}(\text{Reject } \mathcal{H}_0^{(\mathcal{M})} \mid \mathcal{H}_1^{(\mathcal{M})}) = 1,$$

meaning that under the standing conditions, the proposed test defined by the critical region (2.7) is an asymptotic size- $\alpha$  test which is consistent against all fixed alternatives.



The general theory and specific conditions under which (2.4)–(2.6) hold true have been presented by using a Hilbert space approach, in Baringhaus et al. (2017) and Henze and Mayer (2020), for distributional homogeneity and symmetry–equivalence testing, respectively.

Our contribution herein is to introduce a calculus by means of which the test defined in (2.7) is made feasible. More precisely by utilizing the results in Chen et al. (2019) and the asymptotic behavior of  $\Delta_{n,w}^{(\mathcal{M})}$  obtained within a  $U$ –statistic framework, we determine specific instances of the test which may be carried out in a computationally friendly way, which also facilitates Monte Carlo approximation and the study of small–sample properties of the resulting tests.

### 3 Specification of test statistics

For computational convenience as well as for asymptotics we will impose the following assumption on the weight functions. The generic notation  $d$  for dimension is used below, with  $d$  varying depending on context.

- (A) The weight function  $w(t)$  is the density function of a symmetric around zero distribution on  $\mathbb{R}^d$ , i.e.  $w(t) = w(-t)$ ,  $t \in \mathbb{R}^d$ , and  $\int_{\mathbb{R}^d} w(t)dt = 1$ .

More specifically as weight function we fix the density of a spherically symmetric distribution. The CF of this density will be denoted by  $C_w(\cdot)$ , and will be specified further down the paper.

By taking into account the characterization in Chen et al. (2019) we have that, under assumption (A), the population measures in (2.1)–(2.3) can be rendered as

$$(3.1) \quad \Delta_{w,Z}^{(\mathcal{H})} = \mathbb{E} [C_w(X - X_1) + C_w(Y - Y_1) - 2C_w(X - Y)],$$

$$(3.2) \quad \Delta_{w,Z}^{(\mathcal{S})} = \frac{1}{2} \mathbb{E} [C_w(X - X_1) - C_w(X + X_1)],$$

$$(3.3) \quad \Delta_{w,Z}^{(\mathcal{T})} = \mathbb{E} [C_{w_p}(X - X_1)C_{w_q}(Y - Y_1)] + \mathbb{E} [C_{w_p}(X - X_1)] \mathbb{E} [C_{w_q}(Y - Y_1)] \\ - 2\mathbb{E} [C_{w_p}(X - X_1)C_{w_q}(Y - Y_2)],$$

where  $Z_1$  and  $Z_2$  denote independent copies of the random vector  $Z$ . Thus, it is natural to estimate the population measure  $\Delta_{w,Z}^{(\mathcal{M})}$  by means of suitable  $U$ -statistics, whose corresponding kernels are determined by  $C_w(\cdot)$ ; see, e.g., Hoeffding (1948) and Serfling (1980).

To begin with the symmetry statistic, suppose that  $\{X_1, \dots, X_n\}$  are independent copies of  $X$ , the empirical counterpart of  $\Delta_{w,Z}^{(S)}$  may be obtained as

$$(3.4) \quad \Delta_{n,w}^{(S)} = \frac{1}{2n(n-1)} \sum_{1 \leq i \neq j \leq n} \{C_w(X_i - X_j) - C_w(X_i + X_j)\},$$

i.e. as a  $U$ -statistic with symmetric kernel  $\psi^S(x, x_1) = \{C_w(x - x_1) - C_w(x + x_1)\}/2$ .

Likewise suppose that in addition to  $\{X_1, \dots, X_n\}$ , and independently, we have  $\{Y_1, \dots, Y_n\}$  as independent copies of  $Y$ . Then the empirical counterpart of  $\Delta_{w,Z}^{(\mathcal{H})}$  may be formulated as

$$(3.5) \quad \Delta_{n,w}^{(\mathcal{H})} = \frac{1}{n(n-1)} \sum_{1 \leq i \neq j \leq n} \{C_w(X_i - X_j) + C_w(Y_i - Y_j) - 2C_w(X_i - Y_j)\},$$

which is a  $U$ -statistic with symmetric kernel  $\psi^{\mathcal{H}}(z, z_1) = C_w(x - x_1) + C_w(y - y_1) - C_w(x - y_1) - C_w(x_1 - y)$ . Here,  $z = (x^\top, y^\top)^\top$ .

In turn if  $\{(X_1, Y_1), \dots, (X_n, Y_n)\}$  are independent copies of  $(X, Y)$ , the empirical counterpart of  $\Delta_{w,Z}^{(\mathcal{I})}$  is given by

$$(3.6) \quad \begin{aligned} \Delta_{n,w}^{(\mathcal{I})} &= \frac{1}{n(n-1)} \sum_{1 \leq i \neq j \leq n} C_{w_p}(X_i - X_j) C_{w_q}(Y_i - Y_j) \\ &+ \left\{ \frac{1}{n(n-1)} \sum_{1 \leq i \neq j \leq n} C_{w_p}(X_i - X_j) \right\} \left\{ \frac{1}{n(n-1)} \sum_{1 \leq i \neq j \leq n} C_{w_q}(Y_i - Y_j) \right\} \\ &- \frac{2}{n(n-1)(n-2)} \sum_{1 \leq i \neq j \neq k \leq n} C_{w_p}(X_i - X_j) C_{w_q}(Y_i - Y_k) \\ &:= U_{n,1} + U_{n,2} U_{n,3} - 2U_{n,4}. \end{aligned}$$

Denote  $S_1 = \mathbb{E} [C_{w_p}(X - X_1) C_{w_q}(Y - Y_1)]$ ,  $S_2 = \mathbb{E} [C_{w_p}(X - X_1)]$ ,  $S_3 = \mathbb{E} [C_{w_q}(Y - Y_1)]$ , and  $S_4 = \mathbb{E} [C_{w_p}(X - X_1) C_{w_q}(Y - Y_2)]$ . Thus each  $U_{n,i}$  ( $i = 1, \dots, 4$ ) is a  $U$ -statistic of degree  $m_i$  for estimation of  $S_i$  based on the symmetric kernel  $\psi^{\mathcal{I},i}$ , where  $m_1 = m_2 = m_3 = 2$ ,  $m_4 = 3$ ,  $\psi^{\mathcal{I},1}(z, z_1) = C_{w_p}(x - x_1) C_{w_q}(y - y_1)$ ,  $\psi^{\mathcal{I},2}(z, z_1) = C_{w_p}(x - x_1)$ ,

$\psi^{\mathcal{I},3}(z, z_1) = C_{w_q}(y - y_1)$ , and

$$\begin{aligned} \psi^{\mathcal{I},4}(z, z_1, z_2) &= \frac{1}{6} \left\{ C_{w_p}(x - x_1)C_{w_q}(y - y_2) + C_{w_p}(x - x_2)C_{w_q}(y - y_1) \right. \\ &\quad + C_{w_p}(x - x_1)C_{w_q}(y_1 - y_2) + C_{w_p}(x_1 - x_2)C_{w_q}(y - y_1) \\ &\quad \left. + C_{w_p}(x - x_2)C_{w_q}(y_1 - y_2) + C_{w_p}(x_1 - x_2)C_{w_q}(y - y_2) \right\}. \end{aligned}$$

Here  $\psi^{\mathcal{I},4}$  is a symmetrized version of the asymmetric kernel  $C_{w_p}(x-x_1)C_{w_q}(y-y_2)$  of  $S_4$ . Consequently,  $\Delta_{n,w}^{(\mathcal{I})}$  is a function of the  $U$ -statistics  $U_{n,1}, \dots, U_{n,4}$ . Specifically if we let  $h(u) = u_1 + u_2u_3 - 2u_4$  be a known real-valued function of a four-dimensional vector  $u$ , then  $\Delta_{w,Z}^{(\mathcal{I})} = h(S)$  and  $\Delta_{n,w}^{(\mathcal{I})} = h(U_n)$ , where  $S = (S_1, \dots, S_4)$  and  $U_n = (U_{n,1}, \dots, U_{n,4})$ .

According to the above discussion, the asymptotic behavior of  $\Delta_{n,w}^{(S)}$  and  $\Delta_{n,w}^{(\mathcal{H})}$  can be deduced from the general theory of  $U$ -statistics (see, e.g., Hoeffding, 1948, and Serfling, 1980). Likewise, the asymptotic behavior of  $\Delta_{n,w}^{(\mathcal{I})}$  can be obtained from the asymptotic properties of functions of several  $U$ -statistics (see e.g., Hoeffding, 1948, and Lee, 1990). We now formulate the asymptotic properties of  $\Delta_{n,w}^{(\mathcal{M})}$  in (3.4)–(3.6) in the following theorem uniformly. Again the composition of the random vector  $Z$  below varies depending on the context. Also recall that in the context of homogeneity-equivalence test,  $Z = \{X, Y\}$ , with  $X$  and  $Y$  assumed independent, but we will reconsider this assumption in Remark 3.2.

**Theorem 3.1** *Let  $\{Z_1, \dots, Z_n\}$  be independent copies of  $Z$ . Then under assumption (A), and as  $n \rightarrow \infty$ , we have the following results:*

(i) *If  $\Delta_{w,Z}^{(\mathcal{M})} > 0$ , then*

$$\sqrt{n} \left( \Delta_{n,w}^{(\mathcal{M})} - \Delta_{w,Z}^{(\mathcal{M})} \right) \xrightarrow{\mathcal{D}} \mathcal{N} \left( 0, \sigma_w^{2(\mathcal{M})} \right),$$

*with the limit variance  $\sigma_w^{2(\mathcal{M})}$ , respectively given by,*

$$\sigma_w^{2(S)} = 4 \left\{ \mathbb{E} \left[ \psi^S(X, X_1)\psi^S(X, X_2) \right] - \mathbb{E}^2 \left[ \psi^S(X, X_1) \right] \right\},$$

$$\sigma_w^{2(\mathcal{H})} = 4 \left\{ \mathbb{E} \left[ \psi^{\mathcal{H}}(Z, Z_1)\psi^{\mathcal{H}}(Z, Z_2) \right] - \mathbb{E}^2 \left[ \psi^{\mathcal{H}}(Z, Z_1) \right] \right\},$$

$$\sigma_w^{2(\mathcal{I})} = \sum_{i,j=1}^4 m_i m_j h'_i(S) h'_j(S) \zeta_1(i, j),$$

*where  $h'_i(S) := \frac{\partial h}{\partial u_i}|_{u=S}$  denote the first-order partial derivative of  $h$ , and  $\zeta_1(i, j) = \zeta_1(j, i)$  are defined in the Appendix.*

(ii) If  $\Delta_{w,Z}^{(\mathcal{M})} = 0$ , then

$$\sqrt{n}\Delta_{n,w}^{(\mathcal{M})} \xrightarrow{\mathbb{P}} 0$$

□

Theorem 3.1 presents the specific conditions under which (2.4) and (2.5) hold true. Now we turn to study the consistent estimation of the limit variance  $\sigma_w^{2(\mathcal{M})}$ . Based on the strong law of large numbers (SLLN) for  $U$ -statistics (Theorem 5.4A in Serfling, 1980) and Slutsky's theorem, natural consistent estimators of  $\sigma_w^{2(\mathcal{S})}$  and  $\sigma_w^{2(\mathcal{H})}$  can be defined as

$$(3.7) \quad \sigma_{n,w}^{2(\mathcal{S})} = \frac{4}{n(n-1)(n-2)} \sum_{1 \leq i \neq j \neq k \leq n} \psi^{\mathcal{S}}(X_i, X_j) \psi^{\mathcal{S}}(X_i, X_k) - 4 \left\{ \frac{1}{n(n-1)} \sum_{1 \leq i \neq j \leq n} \psi^{\mathcal{S}}(X_i, X_j) \right\}^2,$$

and

$$(3.8) \quad \sigma_{n,w}^{2(\mathcal{H})} = \frac{4}{n(n-1)(n-2)} \sum_{1 \leq i \neq j \neq k \leq n} \psi^{\mathcal{H}}(Z_i, Z_j) \psi^{\mathcal{H}}(Z_i, Z_k) - 4 \left\{ \frac{1}{n(n-1)} \sum_{1 \leq i \neq j \leq n} \psi^{\mathcal{H}}(Z_i, Z_j) \right\}^2.$$

respectively. Note that the SLLN for  $U$ -statistics applies under the condition that the corresponding kernel has a bounded expectation. In this connection, kernels associated with  $\sigma_{n,w}^{2(\mathcal{S})}$  and  $\sigma_{n,w}^{2(\mathcal{H})}$  only depend on  $C_w(\cdot)$ , and thus in our context this condition translates into the condition that  $\mathbb{E}|C_w(\cdot)|^2 < \infty$ , which is trivially true for any real-valued CF and any underlying law of observations.

However, in order to obtain an estimator of  $\sigma_w^{2(\mathcal{I})}$  analogous to (3.7) and (3.8), the number of terms to be estimated in  $\sigma_w^{2(\mathcal{I})}$  is equal to  $4(4+1)/2 = 10$ . Therefore, an analogous estimator of  $\sigma_w^{2(\mathcal{I})}$  is very complicated to simulate. Instead, we will estimate the variance of  $\Delta_{n,w}^{(\mathcal{I})} = h(U_n)$  by jackknife, which is a widely applicable statistical tool used for estimating variance; see Efron (1982) for a full account of the jackknife. The jackknife estimate of variance, denoted by  $\sigma_{n,j}^2$ , can be calculated without resampling, and its definition is given based on the leave-one-out statistics  $h(U_{n-1}(-i))$ ,

$i = 1, \dots, n$ , which are computed from the sample of  $n - 1$  observations formed by deleting the  $i^{\text{th}}$  observation. Specifically,

$$\sigma_{n,J}^2 = \frac{n-1}{n} \sum_{i=1}^n \left( h(U_{n-1}(-i)) - \bar{h}(U_{n-1}) \right)^2,$$

where  $\bar{h}(U_{n-1}) = n^{-1} \sum_{i=1}^n h(U_{n-1}(-i))$ . In fact,  $n\sigma_{n,J}^2$  is a strongly consistent estimator of the limit variance  $\sigma_w^{2(\mathcal{I})}$ ; see Section 5.1.3 in Lee (1990). Then a consistent estimator of  $\sigma_w^{2(\mathcal{I})}$  is given by

$$(3.9) \quad \sigma_{n,w}^{2(\mathcal{I})} = (n-1) \sum_{i=1}^n \left( h(U_{n-1}(-i)) - \bar{h}(U_{n-1}) \right)^2.$$

**Remark 3.2** As a referee kindly pointed out, even without the assumption of independence of  $X$  and  $Y$ , it holds

$$\Delta_{w,Z}^{(\mathcal{H})} = \mathbb{E}[C_w(X - X_1) + C_w(Y - Y_1) - C_w(X - Y_1) - C_w(Y - X_1)],$$

in the case of homogeneity testing. This has the same symmetric kernel  $\psi^{\mathcal{H}}(z, z_1)$  with that in (3.5), and opens up the possibility of extending the method to a more general two-sample problem with dependent observations. Here however we remain strictly within the classical two-sample problem under which  $X$  and  $Y$  are independent. For more information on the case of dependent observations we refer to Hlávka et al. (2020).

We close this section by specifying the spherical densities which will be employed in our simulation study. In this connection note that in principle any spherical density (or in fact any centrally symmetric density) may be used as weight function. However there exist some choices that are convenient from the computational point of view. Specifically we will employ as weight function the density of the spherical stable distribution and the density of the generalized spherical Laplace distribution. The CFs of these densities will be denoted by  $C_\gamma(\cdot)$ , where  $C_\gamma(t) = \exp(-\|t\|^\gamma)$ ,  $\gamma \in (0, 2]$ , corresponds to the stable density, while  $C_\gamma(t) = (1 + \|t\|^2)^{-\gamma}$ ,  $\gamma > 0$ , corresponds to the Laplace density; see Nolan (2013) and Kozubowski et al. (2013), respectively. Here  $\gamma$  may be termed “characteristic exponent”. With these densities as weight functions, formulae for the sample measures  $\Delta_{n,Z}^{(\mathcal{M})}$  in (3.4)–(3.6) as well as the estimated limit variance  $\sigma_{n,w}^{2(\mathcal{M})}$  in (3.7)–(3.9), readily result by simply replacing  $C_w(\cdot)$  by  $C_\gamma(\cdot)$  in the

corresponding equations. (Recall though that in the case of independence testing and due to the complexity of the resulting formulae, we employ the jackknife in order to estimate the limit variance).

## 4 Numerical studies

In this section, a series of simulation experiments are carried out in order to investigate the finite-sample performance of the proposed tests. As mentioned at the end of Section 3, we employ the stable density and the Laplace density as the weight function, and the corresponding tests are denoted as  $T_{n,S}^{(\mathcal{M})}$  and  $T_{n,L}^{(\mathcal{M})}$  respectively. As for the values of the characteristic exponent, we choose  $\gamma \in \{0.5, 1.0, 1.5, 2.0\}$  for the stable density and  $\gamma \in \{0.1, 0.25, 1.0, 4.0\}$  for the Laplace density. In the following, empirical rejection rates according to the critical regions (2.7) are calculated respectively at 5% significance level using Monte Carlo 2000 trials. Further, the dimensions of  $X$  (and  $Y$ ) considered are  $p$  ( $= q$ )  $\in \{2, 4, 6\}$ , and the sample sizes are  $n \in \{100, 200, 300\}$ .

We also wish to compare the new procedures with competitors. In the standard scenario of testing symmetry, homogeneity or independence there are many procedures, even with multivariate data and the reader is referred to Meintanis and Ngatchou-Wandji (2012), to Ghosh and Biswas (2016), and to Zhang et al. (2018) and Peters et al. (2017, Chapt. 7 and A.2), respectively, for reviews of such testing procedures. In the existing literature however we did not find corresponding methods for testing in the context of multivariate equivalence hypotheses. Nevertheless the energy statistics of Székely and Rizzo (2013) have been adapted herein to the context of model-equivalence testing for comparison purposes. For these tests we write  $T_{n,E}^{(\mathcal{M})}$ . By way of example, Székely and Rizzo (2013) characterized the distance between two independent random vectors  $X$  and  $Y$  by

$$\mathcal{E}_\gamma(X, Y) = \mathbb{E}[2\|X - Y\|^\gamma - \|X - X_1\|^\gamma - \|Y - Y_1\|^\gamma],$$

where  $\gamma \in (0, 2)$ ,  $\mathbb{E}[\|X\|^\gamma] < \infty$ , and  $\mathbb{E}[\|Y\|^\gamma] < \infty$ . Consequently the energy-based homogeneity population measure may formally be obtained by replacing  $C_w(\cdot)$  with  $-\|\cdot\|^\gamma$  in (3.1), and consequently the test  $T_{n,E}^{(\mathcal{H})}$  can be implemented by simply replacing  $C_w(\cdot)$  with  $-\|\cdot\|^\gamma$  in (3.5). With analogous replacements one can obtain from (3.2)

and (3.3) corresponding energy population measures for symmetry and independence, respectively. Furthermore, within a  $U$ -statistic framework, the asymptotic properties of the energy-based test statistics analogous to (2.4)–(2.6) can be obtained when the moment conditions  $\mathbb{E}[\|X\|^{2\gamma}] < \infty$  and  $\mathbb{E}[\|Y\|^{2\gamma}] < \infty$  hold true. However, it is worth noticing that the proposed test statistics in this paper require no moment conditions. In what follows we compare our procedures with their energy counterparts  $T_{n,E}^{(\mathcal{M})}$ , for  $\gamma \in \{0.5, 1.0, 1.5, 2.0\}$  (note that although the value  $\gamma = 2.0$  is excluded from the energy-based characterizations we nevertheless include it in our simulations).

**Example 1.** Symmetry–equivalence test: We consider two scenarios in this example, with  $X$  generated from the multivariate skew-normal distribution  $\mathcal{SN}_p(\xi, \Omega, \delta)$  in Example 1(a) and from the multivariate skew-Cauchy distribution  $\mathcal{SC}_p(\xi, \Omega, \delta)$  in Example 1(b), respectively. Here,  $\xi$  is a  $p$ -dimensional vector representing the location parameter,  $\Omega$  is a symmetric positive definite  $p \times p$  scatter matrix, and  $\delta$  is a  $p$ -dimensional vector which regulates the skewness of the distributions; see the monograph Azzalini (2013) and the R package `sn` (Azzalini, 2021) for additional information. In Example 1(a), we choose  $X \sim \mathcal{SN}_p(0_p, I_p, \theta 1_p)$ , where  $0_p$  and  $1_p$  are  $p$ -dimensional vectors of zeros and ones respectively,  $I_p$  denotes the identity matrix, and  $\theta \geq 0$  is a scalar. With this parametrization, the degree of asymmetry can be measured in terms of  $\theta$ , in the sense that smaller  $\theta$  corresponds to less asymmetry. Particularly,  $\mathcal{SN}_p(0_p, I_p, \theta 1_p)$  reduces to the multivariate standard normal distribution when  $\theta = 0$ . We choose some fixed value  $\theta = \theta_0$  as benchmark, and compute the value of  $\Delta_{w,X}^{(S)}$  in (3.2) for  $X \sim \mathcal{SN}_p(0_p, I_p, \theta_0 1_p)$  as a general threshold  $\Delta$  in (2.7) for significant departure from symmetry. Thus distributions  $\mathcal{SN}_p(0_p, I_p, \theta 1_p)$  with  $\theta \geq \theta_0$  correspond to the null hypotheses and we are in the alternatives when  $\theta < \theta_0$ . A random approximation procedure is used to compute  $\Delta_{w,X}^{(S)}$  for  $\mathcal{SN}_p(0_p, I_p, \theta_0 1_p)$ . Specifically, we generate a large number  $B$  of i.i.d. copies  $\{X_1, \dots, X_B\}$  of  $X \sim \mathcal{SN}_p(0_p, I_p, \theta_0 1_p)$ , and calculate  $\Delta_{n,w}^{(S)}$  in (3.4) based on this sample as the threshold  $\Delta$ . To ensure the accuracy of the approximation,  $B = 5000$  is used, which is a fairly large value; see the closing paragraph of this section for more information on the accuracy of the random approximation method. As benchmark we choose  $\mathcal{SN}_p(0_p, I_p, \theta_0 1_p)$  with  $\theta_0 = 3$ , and also consider  $\theta \in \{5, 4, 3, 2, 1, 0\}$ . In Example 1(b), we repeat Example 1(a) under

identical conditions except that  $X \sim \mathcal{SC}_p(0_p, I_p, \theta 1_p)$ .

The empirical rejection rates of the proposed symmetry–equivalence test with the test  $T_{n,S}^{(\mathcal{S})}$  corresponding to the spherical stable density as weight function are shown in Table 1. We also display the comparative results of all three tests  $T_{n,S}^{(\mathcal{S})}$ ,  $T_{n,L}^{(\mathcal{S})}$ , and  $T_{n,E}^{(\mathcal{S})}$  for Examples 1(a) and 1(b) in Figures 1 and 2, respectively.

Table 1 and Figures 1, 2 about here

In general, the performance of the three tests  $T_{n,S}^{(\mathcal{S})}$ ,  $T_{n,L}^{(\mathcal{S})}$ , and  $T_{n,E}^{(\mathcal{S})}$  are comparable in Example 1(a). However, in Example 1(b), where  $X \sim \mathcal{SC}_p(0_p, I_p, \theta 1_p)$  and the expectation of  $X$  does not exist, the energy–based test  $T_{n,E}^{(\mathcal{S})}$  with  $\gamma = 1.5$  or  $\gamma = 2.0$  can not control the Type–I error rates, even when  $p = 2, n = 300$ . We may attribute this behavior to the fact that in this case the moment condition corresponding to the energy statistic is not satisfied. On the other hand, it is clear that the Type–I error rates of the proposed tests  $T_{n,S}^{(\mathcal{S})}$  and  $T_{n,L}^{(\mathcal{S})}$  can be controlled well for lower dimensions ( $p = 2, 4$ ) with all sample sizes and a higher dimension ( $p = 6$ ) with a larger sample size ( $n = 300$ ). Furthermore, it can be seen that, for all cases, the Type–I error rates in the interior of the null hypothesis (i.e.,  $\theta = 5, 4$ ) are lower than those at the boundary of the null hypothesis (i.e.,  $\theta_0 = 3$ ), which is line with the limit result in (2.8). As for the empirical powers, it may be observed that with the increase of the sample size and/or distance away from the null hypothesis, the empirical powers increase with a few exceptions, which again is expected in view of our consistency results. Finally, the value of  $\gamma$  is seen to affect the empirical powers of all three tests, but only slightly.

**Example 2.** Homogeneity–equivalence test: Two scenarios are considered in this example. In Example 2(a), we consider an independent pair of random vectors such that  $X \sim \mathcal{N}_p(0, 1)$  and  $Y \sim \mathcal{N}_p(\mu, 1)$ , where  $\mathcal{N}_p(\mu, \sigma^2)$  denotes a  $p$ –dimensional Gaussian random vector whose coordinates are independent and identically distributed as univariate normal distributions with mean  $\mu$  and variance equal to  $\sigma^2$ . We also choose some fixed value of  $\mu = \mu_0$  as a benchmark, and set the corresponding value of  $\Delta_{w,Z}^{(\mathcal{H})}$  in (3.1) as a general threshold  $\Delta$  in (2.7) for significant departure from distributional homogeneity. Thus  $\mu \geq \mu_0$  and  $\mu < \mu_0$  correspond to the null hypotheses and alternatives, respectively. Also, the benchmark threshold  $\Delta$  is computed according to a



random approximation procedure. Furthermore, the benchmark and non-benchmark values considered are  $\mu_0 = 2.0$  and  $\mu \in \{2.2, 2.1, 2.0, 1.9, 1.8, 1.7\}$ . In Example 2(b), we repeat Example 2(a) under identical conditions except that the coordinates of  $X$  and  $Y$  are independently generated from  $G(5, 1)$  and  $G(5, \mu)$ , where  $G(a, s)$  denotes the univariate Gamma distribution with shape parameter  $a$  and scale parameter  $s$ .

The resulting empirical rejection rates for  $T_{n,S}^{(\mathcal{H})}$  are reported in Table 2 while corresponding comparative results of all three tests  $T_{n,S}^{(\mathcal{H})}$ ,  $T_{n,L}^{(\mathcal{H})}$ , and  $T_{n,E}^{(\mathcal{H})}$  are displayed in Figures 3 and 4. It may be seen clearly that the Type-I error rates are generally controlled with a few exceptions appearing when  $p$  is higher and  $n$  is smaller, and that empirical rejection rates under alternatives are in line with the fact that the tests are consistent. In addition, the energy-based test  $T_{n,E}^{(\mathcal{H})}$  is more powerful in Example 2(a), whereas in Example 2(b), the opposite is true. Furthermore in Example 2(a), and as it may be seen from Figure 3, the value of  $\gamma$  has a certain effect on the empirical powers of the proposed tests  $T_{n,S}^{(\mathcal{H})}$  and  $T_{n,L}^{(\mathcal{H})}$ . We postpone further discussion on  $\gamma$  till the end of this section.

Table 2 and Figures 3, 4 about here

**Example 3.** Independence-equivalence test: We consider two scenarios in this example, with  $Z \sim \mathcal{MN}(0_{p+q}, \Sigma_\rho)$  in Example 3(a) and  $Z \sim \mathcal{MT}(0_{p+q}, \Sigma_\rho, 5)$  in Example 3(b), respectively. Here,  $Z = (X^\top, Y^\top)^\top$ ,  $\mathcal{MN}(\mu, \Sigma)$  denotes the multivariate normal distribution with mean vector  $\mu$  and covariance matrix  $\Sigma$ , and  $\mathcal{MT}(\delta, \Sigma, \nu)$  denotes the multivariate  $t$  distribution (Kotz and Nadarajah, 2004) with non-centrality parameter  $\delta$ , scale matrix  $\Sigma$  and degrees of freedom  $\nu$ . Furthermore,

$$\Sigma_\rho = \begin{pmatrix} I_p & \Sigma_0 \\ \Sigma_0^\top & I_q \end{pmatrix}$$

is a  $(p+q) \times (p+q)$  matrix, and  $\Sigma_0$  is a  $p \times q$  matrix with the  $(i, j)^{\text{th}}$  element equals to  $\rho \times \delta_{ij}$  for all  $i = 1, \dots, p$  and  $j = 1, \dots, q$ , where  $\delta_{ij} = 1$  if  $i = j$  and  $\delta_{ij} = 0$ , otherwise. Thus  $\rho$  quantifies the dependence between  $X$  and  $Y$ . We again choose some fixed value of  $\rho = \rho_0$  as a benchmark, and set the corresponding value of  $\Delta_{w,Z}^{(\mathcal{I})}$  in (3.3) as  $\Delta$ , with  $\Delta$  being approximated by means of the random approximation procedure.

The reported empirical rejection rates for  $T_{n,S}^{(\mathcal{I})}$ ,  $T_{n,L}^{(\mathcal{I})}$  and  $T_{n,E}^{(\mathcal{I})}$  in Table 3 and Figures 5, 6, correspond to the benchmark value  $\rho_0 = 0.8$  and the non-benchmark values employed  $\rho \in \{0.84, 0.82, 0.8, 0.75, 0.7, 0.65\}$ . Similar conclusions as before may be drawn from these results regarding Type-I errors and powers. At the same time however, the energy-based test  $T_{n,E}^{(\mathcal{I})}$  seems to be less powerful for nearly all cases in Example 3(a) as well as in Example 3(b).

Table 3 and Figures 5, 6 about here

The simulation results of Examples 1–3 suggest that in some cases the weight function used as well as the value of  $\gamma$  may affect the performance of the proposed tests to a certain degree. Nevertheless most research on this topic concentrates on the choice of the weight parameter conditionally on the choice of the weight function, rather than on the choice of weight function itself. For a detailed discussion on this topic we refer to Meintanis (2016). In this connection several researchers and in varying circumstances have asserted that such an optimal choice, if it exists, would require some knowledge about the type of departure from the null hypothesis expected; see for instance Bugni et al. (2009). One approach is to analyze the asymptotic behavior of the test under the null against some “qualitative” classes of alternatives and suggest a compromise value that performs well across these classes of alternatives. In the rather narrow context of testing for multivariate normality, Tenreiro (2009) suggested specific weights that give good powers for long-tailed, short-tailed, symmetric and skewed alternatives to normality, and in a more broad context of testing goodness-of-fit to parametric families of distributions Allison and Santana (2015) and Tenreiro (2019) suggest data-driven ways for choosing the weight parameter. In our current semiparametric context though, proper choice of  $\gamma$  clearly leads to a highly non-trivial analytic problem which needs to be studied separately. We refer to Mélisande et al. (2019) for results on a recent “aggregate” test for independence that employs a collection of weights rather than a single choice. However as these results are preliminary and only apply to independence testing, and not to symmetry or homogeneity testing, and certainly not to equivalence testing, we do not follow this approach. We instead opt for the practical approach of suggesting a compromise value of the weight parameter that seems to work well across

several settings. In this connection and based on our simulation results we suggest the use of a “medium” sized value of  $\gamma$  such as  $\gamma = 1.0$  for both the spherical stable and the Laplace density as an overall good choice. Clearly though this suggestion is conditioned on our Monte Carlo results, and therefore can not be overgeneralized.

This section concludes with a numerical investigation of the random approximation procedure. In this connection, consider for instance the homogeneity–equivalence test in Example 2(a) and write the distance  $\Delta_{w,Z}^{(\mathcal{H})}$ , with  $X \sim \mathcal{N}_p(0, 1)$  and  $Y \sim \mathcal{N}_p(\mu_0, 1)$ , in the following form:

$$\begin{aligned}
 (4.1) \quad \Delta_{w,Z}^{(\mathcal{H})} &= \int_{\mathbb{R}^p} |\varphi_X(t) - \varphi_Y(t)|^2 w(t) dt \\
 &= \int_{\mathbb{R}^p} \left| \exp\left(-\frac{1}{2}t^\top t\right) - \exp\left(i\mu_0 \mathbf{1}_p^\top t - \frac{1}{2}t^\top t\right) \right|^2 w(t) dt \\
 &= 2 \int_{\mathbb{R}^p} \exp(-t^\top t) (1 - \cos(\mu_0 \mathbf{1}_p^\top t)) w(t) dt,
 \end{aligned}$$

(recall that  $\mathbf{1}_p$  denotes the  $p$ -dimensional vector of ones). Note in this connection that the density functions of the spherical stable distribution and the generalized spherical Laplace distribution can be found in Nolan (2013) and Kozubowski et al. (2013), respectively, and hence the values of  $\Delta_{w,Z}^{(\mathcal{H})}$  in Example 2(a) may be computed by numerical integration utilizing (4.1). The corresponding results reported in Table 4 are the values of  $\Delta_{w,Z}^{(\mathcal{H})}$  for the benchmark in Example 2(a) with  $\mu_0 = 2.0$ , first computed by random approximation and also by numerical integration, and show that the two methods are in close agreement, a fact that corroborates the use of the random approximation procedure.

Table 4 about here

## 5 Conclusions

In this paper, we propose three test criteria based on weighted  $L_2$ -type distance measures, each of which is appropriate for testing, respectively, symmetry, distributional homogeneity, and independence, all three hypotheses being formulated as equivalence hypotheses. The asymptotic properties of the proposed tests are proved by utilizing the theory of  $U$ -statistics which properly combined with certain choices of the adopted

weighting scheme lead to computationally convenient, asymptotically normal tests that are consistent against any fixed alternative. Our Monte Carlo results confirm the applicability of the suggested procedures as well as their competitiveness vis-à-vis their energy-based counterparts. In perspective we point out that the techniques employed herein may potentially be applied in the context of testing goodness-of-fit with arbitrary parametric families of distributions.

## Acknowledgments

This research was supported by the National Natural Science Foundation of China [Grant No.12101055, No.12131006]. The authors wish to sincerely thank the Executive Editors for handling our paper in a timely manner, and two anonymous reviewers for their constructive criticism that led to improvement of the article.

## References

- Allison, J. and Santana, L. (2015). On a data-dependent choice of the tuning parameter appearing in certain goodness-of-fit tests. *J. Statist. Comput. Simul.*, 85, 3276–3288.
- Azzalini, A. and Capitanio, A. (1999). Statistical applications of the multivariate skew normal distribution. *J. Roy. Statist. Soc. B.*, 61, 579–602.
- Azzalini, A. (2013). *The Skew-Normal and Related Families (in collaboration with Antonella Capitanio)*. Cambridge University Press, Cambridge.
- Azzalini, A. (2021). *The R Package sn: The Skew-Normal Probability Distributions (and Related Distributions such as the Skew-t and the SUN)*. Available at <http://azzalini.stat.unipd.it/SN/>
- Baringhaus, L., Ebner, B., and Henze, N. (2017). The limit distribution of weighted  $L^2$ -goodness-of-fit statistics under fixed alternatives, with applications. *Ann. Instit. Statist. Math.*, 69, 969–995.
- Berger, J., and Delampady, M. (1987). Testing precise hypotheses (with discussion). *Statist. Scienc.*, 2, 317–352.
- Borovkov, A.A. (1998). *Mathematical Statistics*. Gordon and Breach, Amsterdam.

- Bugni, F.A., Hall, P., Horowitz, J.L. and Neumann, G.R. (2009). Goodness-of-fit tests for functional data. *Econometrics Journal*, 12, 1–18.
- Chen, F., Meintanis, S.G., and Zhu, L. (2019). On some characterizations and multidimensional criteria for testing homogeneity, symmetry and independence. *J. Multivar. Anal.*, 173, 125–144.
- Dette, H., Möllenhoff, K., Volgushev, S., and Bretz, F. (2018). Equivalence of regression curves. *J. Amer. Statist. Assoc.*, 113, 711–729.
- Dette, H., and Munk, A. (1998). Validation of linear regression models. *Ann. Statist.*, 26, 778–800.
- Efron, B. (1982). The jackknife, the bootstrap and other resampling plans. SIAM, Philadelphia.
- Freitag, G., Czado, C., and Munk, A. (2007). A nonparametric test for similarity of marginals—with applications to the assessment of bioequivalence. *J. Statist. Plann. Infer.*, 137, 697–711.
- Ghosh, A.K., and Biswas, M. (2016). Distribution-free high-dimensional two-sample tests based on discriminating hyperplanes. *TEST*, 25, 525–547.
- Henze, N., and Mayer, C. (2020). More good news on the HKM test for multivariate reflected symmetry about an unknown centre. *Ann. Instit. Statist. Math.*, 72, 741–770.
- Hlávka, Z., Hušková, M., and Meintanis, S.G. (2020). Change-point methods for multivariate time-series: paired vectorial observations. *Statist. Pap.*, 61, 1351–1383.
- Hoeffding, W. (1948). A class of statistics with asymptotically normal distribution. *Ann. Math. Statist.*, 19, 293–325.
- Janssen, A. (2000). Nonparametric bioequivalence tests for statistical functionals and their efficient power functions. *Statist. & Decisions*, 18, 49–78.
- Kotz, S., and Nadarajah, S. (2004). *Multivariate t Distributions and Their Applications*. Cambridge University Press, Cambridge.
- Kozubowski, T.J., Podgórski K., and Rychlik, I. (2013). Multivariate generalized Laplace distribution and related random fields. *J. Multivar. Anal.*, 113, 59–72.
- Lee, A.J. (1990). *U-Statistics: Theory and Practice*. Marcel Dekker, Inc., New York.
- Meintanis, S.G. (2016). A review of testing procedures based on the empirical characteristic function (with discussion and rejoinder). *South Afric. J. Statist.*, 50, 1–41.

- Meintanis, S.G., and Ngatchou-Wandji, J. (2012). Recent tests for symmetry with multivariate and structured data: A review. J Jiang et al. (Eds.) *Non-Parametric Statistical Methods and Related Topics*, pp 35-73, World Scientific, New Jersey.
- Mélisande, A., Laurent-Bonneau, B., Marrel, A., and Meynaoui, A. (2019). Adaptive tests of independence based on HSIC measures. <https://arxiv.org/abs/1902.06441>.
- Möllenhoff, K., Bretz, F., and Dette, H. (2019). Equivalence of regression curves sharing common parameters. *Biometrics*, *76*, 518–529.
- Nadarajah, S. (2003). The Kotz-type distribution with applications. *Statistics*, *37*, 341–358.
- Nolan, J.P. (2013). Multivariate elliptically contoured stable distributions: Theory and estimation. *Comput. Statist.*, *28*, 2067–2089.
- Peters, J., Janzing, D., and Schölkopf, B. (2017). *Elements of Causal Inference*. The MIT Press, Cambridge, Massachusetts.
- Serfling, R.J. (1980). *Approximation Theorems of Mathematical Statistics*. John Wiley & Sons, Inc., New York.
- Sprenger, J. (2013). Testing a precise null hypothesis: The case of Lindley’s paradox. *Philosophy of Science*, *80*, 733–744.
- Székely, G.J., and Rizzo, M.L. (2013). Energy statistics: A class of statistics based on distances. *J. Statist. Plann. Infer.*, *143*, 1249–1272.
- Székely, G.J., Rizzo, M.L., and Bakirov, N.K. (2007). Measuring and testing dependence by correlation of distances. *Ann. Statist.*, *35*, 2769–2794.
- Tenreiro, C. (2009). On the choice of the smoothing parameter for the BHEP goodness-of-fit test. *Comput. Statist. Dat. Anal.*, *53*, 1038–1053.
- Tenreiro, C. (2019). On the automatic selection of the tuning parameter appearing in certain families of goodness-of-fit tests. *J. Statist. Comput. Simul.*, *89*, 1780–1797.
- Wellek, S. (2010). *Testing Statistical Hypotheses of Equivalence and Noninferiority*. CRC Press, New York.
- Zhang, Q., Filippi, S., Gretton, A., and Sejdinovic, D. (2018). Large-scale kernel methods for independence testing. *Statist. & Comput.*, *28*, 113–130.

# Appendix

*Proof of Theorem 3.1.*

First, we proof the asymptotic properties of  $\Delta_{n,w}^{(S)}$  using the general theory of  $U$ -statistics. Define  $\psi_1^S(x) = \mathbb{E}[\psi^S(x, X)]$ , where the expected value is taken with respect to the random vector  $X$ . Thus

$$\begin{aligned}
 (5.1) \quad \text{Var}[\psi_1^S(X)] &= \mathbb{E}[\psi_1^S(X)^2] - \mathbb{E}^2[\psi_1^S(X)] \\
 &= \mathbb{E}\{\mathbb{E}[\psi^S(X, X_1)|X]\}^2 - \mathbb{E}^2\{\mathbb{E}[\psi^S(X, X_1)|X]\} \\
 &= \mathbb{E}\{\mathbb{E}[\psi^S(X, X_1)\psi^S(X, X_2)|X]\} - \mathbb{E}^2[\psi^S(X, X_1)] \\
 &= \mathbb{E}[\psi^S(X, X_1)\psi^S(X, X_2)] - \mathbb{E}^2[\psi^S(X, X_1)],
 \end{aligned}$$

where  $X_1$  and  $X_2$  are independent copies of  $X$ . Consider that any characteristic function is uniformly bounded, thus  $\text{Var}[\psi_1^S(X)] < \infty$  can be deduced without any extra assumptions about the distribution of  $X$ . Hence according to the asymptotic property of  $U$ -statistics (see, e.g., Section 5.5.1 in Serfling, 1980), as  $n \rightarrow \infty$ , we have

$$\begin{aligned}
 \sqrt{n}(\Delta_{n,w}^{(S)} - \Delta_{w,X}^{(S)}) &= \frac{2}{\sqrt{n}} \sum_{i=1}^n (\psi_1^S(X_i) - \Delta_{w,X}^{(S)}) + o_p(1) \\
 &\xrightarrow{\mathcal{D}} \mathcal{N}(0, \sigma_w^{2(S)}),
 \end{aligned}$$

where  $\sigma_w^{2(S)} = 4\text{Var}[\psi_1^S(X)]$ , and the limiting distribution is non-singular provided that  $\sigma_w^{2(S)} > 0$ . Notice further that  $\sigma_w^{2(S)} = 0$  if and only if the law of  $X$  is symmetric about the origin, i.e.,  $\Delta_{w,X}^{(S)} = 0$ . Thus when  $\Delta_{w,X}^{(S)} = 0$ ,

$$\sqrt{n}\Delta_{n,w}^{(S)} \xrightarrow{\mathcal{D}} 0, \quad \text{as } n \rightarrow \infty.$$

The second part of the proof for  $\Delta_{n,w}^{(S)}$  can be finished immediately.

Define  $\psi_1^H(z) = \mathbb{E}[\psi^H(z, Z)]$ , where the expected value is taken with respect to the random vector  $Z$ . Similar to (5.1), we have

$$(5.2) \quad \text{Var}[\psi_1^H(Z)] = \mathbb{E}[\psi^H(Z, Z_1)\psi^H(Z, Z_2)] - \mathbb{E}^2[\psi^H(Z, Z_1)].$$

Then the asymptotic properties of  $\Delta_{n,w}^{(H)}$  can be obtained by similar arguments as in the case of  $\Delta_{n,w}^{(S)}$ .

Define  $\psi_1^{\mathcal{I},i}(z) = \mathbb{E}[\psi^{\mathcal{I},i}(z, Z_1, \dots, Z_{m_i-1})]$ ,  $i = 1, \dots, 4$ , where  $m_i$  is the degree of kernel  $\psi^{\mathcal{I},i}$ , and the expected value is taken with respect to the random vectors  $Z_1, \dots, Z_{m_i-1}$ . Further, define

$$\zeta_1(i, j) = \text{Cov} \left( \psi_1^{\mathcal{I},i}(Z), \psi_1^{\mathcal{I},j}(Z) \right), \quad i, j = 1, \dots, 4.$$

Obviously  $\zeta_1(i, j)$  is symmetric in the sense that  $\zeta_1(i, j) = \zeta_1(j, i)$ . Similar to (5.1), for  $i, j = 1, 2, 3$ , we have

$$\begin{aligned} \zeta_1(i, j) &= \mathbb{E} \left[ \psi^{\mathcal{I},i}(Z, Z_1) \psi^{\mathcal{I},j}(Z, Z_2) \right] - S_i S_j, \\ \zeta_1(i, 4) &= \mathbb{E} \left[ \psi^{\mathcal{I},i}(Z, Z_1) \psi^{\mathcal{I},4}(Z, Z_2, Z_3) \right] - S_i S_4, \\ \zeta_1(4, 4) &= \mathbb{E} \left[ \psi^{\mathcal{I},4}(Z, Z_1, Z_2) \psi^{\mathcal{I},4}(Z, Z_3, Z_4) \right] - S_4^2. \end{aligned}$$

Here all pairs of kernels  $\psi^{\mathcal{I},i}$  and  $\psi^{\mathcal{I},j}$  are evaluated on two subsets with only one element in common. Thus  $\zeta_1(i, j) < \infty$ ,  $i, j = 1, \dots, 4$ , can be obtained according to the uniform boundedness of characteristic functions, and the proof can be finished by using Theorem 7.5 in Hoeffding (1948) and the fact that  $\sigma_w^{2(\mathcal{I})} = 0$  holds true if and only if  $X$  and  $Y$  are mutually independent.

□



Table 1: Simulation results for Example 1. Rejection rates of  $T_{n,S}^{(S)}$  at 5% significance level using 2000 trials.

			Example 1(a)						Example 1(b)					
			Null hypotheses			Alternatives			Null hypotheses			Alternatives		
			$\theta = 5$	$\theta = 4$	$\theta = 3$	$\theta = 2$	$\theta = 1$	$\theta = 0$	$\theta = 5$	$\theta = 4$	$\theta = 3$	$\theta = 2$	$\theta = 1$	$\theta = 0$
$n = 100$	$p = 2$	$\gamma = 0.5$	0.0185	0.0300	0.0575	0.1890	0.8615	1	0.0385	0.0520	0.0920	0.2245	0.8320	1
		$\gamma = 1.0$	0.0200	0.0315	0.0585	0.2020	0.8755	1	0.0465	0.0610	0.0945	0.2315	0.8095	1
		$\gamma = 1.5$	0.0215	0.0320	0.0620	0.2100	0.8870	1	0.0495	0.0620	0.1080	0.2410	0.7975	1
		$\gamma = 2.0$	0.0235	0.0330	0.0645	0.2235	0.8930	1	0.0495	0.0675	0.1130	0.2465	0.7960	1
	$p = 4$	$\gamma = 0.5$	0.0410	0.0460	0.0580	0.1070	0.4590	1	0.0405	0.0440	0.0545	0.0950	0.3855	1
		$\gamma = 1.0$	0.0430	0.0450	0.0570	0.1130	0.4835	1	0.0565	0.0625	0.0770	0.1090	0.3480	1
		$\gamma = 1.5$	0.0430	0.0500	0.0630	0.1200	0.5120	1	0.0755	0.0810	0.0910	0.1335	0.3550	1
		$\gamma = 2.0$	0.0505	0.0550	0.0700	0.1310	0.5400	1	0.0870	0.0925	0.1060	0.1480	0.3735	0.9985
	$p = 6$	$\gamma = 0.5$	0.0785	0.0840	0.0985	0.1285	0.3410	1	0.0815	0.0855	0.0910	0.1135	0.2705	1
		$\gamma = 1.0$	0.0790	0.0830	0.0975	0.1335	0.3460	1	0.0860	0.0880	0.0940	0.1140	0.2470	0.9990
		$\gamma = 1.5$	0.0815	0.0855	0.0950	0.1320	0.3535	1	0.1060	0.1085	0.1225	0.1415	0.2725	0.9920
		$\gamma = 2.0$	0.0990	0.1065	0.1205	0.1465	0.3770	1	0.1420	0.1435	0.1515	0.1745	0.2955	0.9765
$n = 200$	$p = 2$	$\gamma = 0.5$	0.0085	0.0145	0.0430	0.2575	0.9825	1	0.0165	0.0255	0.0705	0.2765	0.9705	1
		$\gamma = 1.0$	0.0100	0.0160	0.0500	0.2730	0.9865	1	0.0175	0.0330	0.0800	0.2930	0.9565	1
		$\gamma = 1.5$	0.0105	0.0180	0.0590	0.2940	0.9900	1	0.0200	0.0385	0.0900	0.3005	0.9450	1
		$\gamma = 2.0$	0.0105	0.0210	0.0670	0.3150	0.9900	1	0.0250	0.0440	0.0960	0.3025	0.9375	1
	$p = 4$	$\gamma = 0.5$	0.0230	0.0280	0.0415	0.1035	0.6525	1	0.0165	0.0225	0.0300	0.0680	0.4965	1
		$\gamma = 1.0$	0.0220	0.0275	0.0430	0.1045	0.6855	1	0.0230	0.0275	0.0355	0.0790	0.4405	1
		$\gamma = 1.5$	0.0235	0.0315	0.0480	0.1185	0.7145	1	0.0345	0.0390	0.0530	0.0985	0.4420	1
		$\gamma = 2.0$	0.0300	0.0365	0.0540	0.1350	0.7285	1	0.0425	0.0505	0.0685	0.1130	0.4530	1
	$p = 6$	$\gamma = 0.5$	0.0540	0.0615	0.0750	0.1090	0.4485	1	0.0465	0.0495	0.0540	0.0870	0.3245	1
		$\gamma = 1.0$	0.0490	0.0550	0.0680	0.1055	0.4555	1	0.0455	0.0485	0.0535	0.0810	0.2710	1
		$\gamma = 1.5$	0.0460	0.0530	0.0645	0.1015	0.4285	1	0.0580	0.0615	0.0675	0.0955	0.2595	1
		$\gamma = 2.0$	0.0510	0.0570	0.0630	0.1020	0.4315	1	0.0720	0.0750	0.0845	0.1105	0.2690	1
$n = 300$	$p = 2$	$\gamma = 0.5$	0.0040	0.0120	0.0425	0.3270	1	1	0.0070	0.0225	0.0705	0.3800	0.9960	1
		$\gamma = 1.0$	0.0045	0.0145	0.0520	0.3455	1	1	0.0155	0.0265	0.0795	0.3660	0.9940	1
		$\gamma = 1.5$	0.0055	0.0140	0.0600	0.3715	1	1	0.0165	0.0310	0.0825	0.3765	0.9910	1
		$\gamma = 2.0$	0.0055	0.0135	0.0645	0.3915	1	1	0.0180	0.0320	0.0900	0.3760	0.9895	1
	$p = 4$	$\gamma = 0.5$	0.0205	0.0265	0.0405	0.1135	0.8065	1	0.0115	0.0155	0.0280	0.0695	0.6190	1
		$\gamma = 1.0$	0.0200	0.0255	0.0390	0.1160	0.8290	1	0.0155	0.0210	0.0290	0.0760	0.5170	1
		$\gamma = 1.5$	0.0175	0.0245	0.0430	0.1260	0.8510	1	0.0235	0.0300	0.0420	0.0885	0.5180	1
		$\gamma = 2.0$	0.0185	0.0255	0.0460	0.1400	0.8645	1	0.0310	0.0365	0.0515	0.1065	0.5335	1
	$p = 6$	$\gamma = 0.5$	0.0430	0.0490	0.0655	0.1210	0.5815	1	0.0285	0.0295	0.0355	0.0605	0.3735	1
		$\gamma = 1.0$	0.0395	0.0415	0.0515	0.1080	0.5820	1	0.0275	0.0305	0.0355	0.0575	0.2815	1
		$\gamma = 1.5$	0.0310	0.0340	0.0445	0.0900	0.5485	1	0.0385	0.0415	0.0490	0.0705	0.2625	1
		$\gamma = 2.0$	0.0295	0.0345	0.0465	0.0895	0.5400	1	0.0505	0.0540	0.0600	0.0815	0.2625	1

Table 2: Simulation results for Example 2. Rejection rates of  $T_{n,S}^{(\mathcal{H})}$  at 5% significance level using 2000 trials.

		Example 2(a)						Example 2(b)						
		Null hypotheses			Alternatives			Null hypotheses			Alternatives			
		$\mu = 2.2$	$\mu = 2.1$	$\mu = 2.0$	$\mu = 1.9$	$\mu = 1.8$	$\mu = 1.7$	$\mu = 2.2$	$\mu = 2.1$	$\mu = 2.0$	$\mu = 1.9$	$\mu = 1.8$	$\mu = 1.7$	
$n = 100$	$p = 2$	$\gamma = 0.5$	0.0010	0.0080	0.0510	0.1810	0.4285	0.7090	0.0005	0.0075	0.0545	0.2580	0.6475	0.9215
		$\gamma = 1.0$	0.0020	0.0120	0.0500	0.1390	0.3230	0.5550	0.0005	0.0075	0.0570	0.2655	0.6500	0.9175
		$\gamma = 1.5$	0.0050	0.0185	0.0510	0.1195	0.2490	0.4240	0	0.0080	0.0565	0.2675	0.6475	0.9140
		$\gamma = 2.0$	0.0070	0.0215	0.0525	0.1090	0.2120	0.3735	0	0.0075	0.0605	0.2750	0.6410	0.9130
	$p = 4$	$\gamma = 0.5$	0	0.0030	0.0500	0.2490	0.6325	0.9070	0	0.0005	0.0525	0.3955	0.8840	0.9980
		$\gamma = 1.0$	0.0075	0.0210	0.0565	0.1515	0.3060	0.5395	0	0.0010	0.0660	0.4020	0.8785	0.9950
		$\gamma = 1.5$	0.0335	0.0450	0.0655	0.0995	0.1575	0.2560	0	0.0030	0.0825	0.4120	0.8480	0.9880
		$\gamma = 2.0$	0.0510	0.0585	0.0740	0.0925	0.1340	0.1860	0	0.0085	0.0960	0.4220	0.8360	0.9835
	$p = 6$	$\gamma = 0.5$	0	0.0030	0.0495	0.3225	0.7675	0.9790	0	0	0.0400	0.5260	0.9655	1
		$\gamma = 1.0$	0.0115	0.0270	0.0570	0.1275	0.2580	0.4450	0	0	0.0545	0.5085	0.9405	1
		$\gamma = 1.5$	0.0625	0.0695	0.0755	0.0895	0.1150	0.1485	0	0.0030	0.0795	0.5100	0.9180	0.9975
		$\gamma = 2.0$	0.0955	0.0980	0.1035	0.1070	0.1155	0.1390	0	0.0065	0.1135	0.5345	0.9035	0.9940
$n = 200$	$p = 2$	$\gamma = 0.5$	0	0.0010	0.0450	0.2560	0.6540	0.9315	0	0	0.0515	0.3795	0.8600	0.9985
		$\gamma = 1.0$	0	0.0035	0.0400	0.1940	0.4870	0.8055	0	0	0.0575	0.3935	0.8690	0.9960
		$\gamma = 1.5$	0	0.0060	0.0420	0.1480	0.3570	0.6250	0	0	0.0570	0.3920	0.8615	0.9950
		$\gamma = 2.0$	0.0015	0.0085	0.0405	0.1275	0.3010	0.5405	0	0.0005	0.0570	0.3885	0.8575	0.9935
	$p = 4$	$\gamma = 0.5$	0	0.0015	0.0430	0.3845	0.8740	0.9925	0	0	0.0450	0.6055	0.9920	1
		$\gamma = 1.0$	0.0015	0.0080	0.0495	0.1750	0.4625	0.7780	0	0	0.0490	0.6010	0.9880	1
		$\gamma = 1.5$	0.0170	0.0280	0.0530	0.0980	0.1910	0.3425	0	0	0.0565	0.5800	0.9800	1
		$\gamma = 2.0$	0.0295	0.0405	0.0595	0.0860	0.1350	0.2265	0	0.0005	0.0590	0.5690	0.9700	1
	$p = 6$	$\gamma = 0.5$	0	0	0.0495	0.5100	0.9570	0.9995	0	0	0.0400	0.7715	0.9990	1
		$\gamma = 1.0$	0.0075	0.0185	0.0520	0.1580	0.3700	0.6505	0	0	0.0510	0.7145	0.9980	1
		$\gamma = 1.5$	0.0395	0.0485	0.0600	0.0805	0.1100	0.1625	0	0	0.0645	0.6915	0.9945	1
		$\gamma = 2.0$	0.0635	0.0670	0.0725	0.0785	0.0930	0.1115	0	0.0010	0.0860	0.6925	0.9885	1
$n = 300$	$p = 2$	$\gamma = 0.5$	0	0.0010	0.0390	0.3245	0.8050	0.9890	0	0	0.0405	0.4815	0.9520	1
		$\gamma = 1.0$	0	0.0025	0.0340	0.2280	0.6375	0.9260	0	0.0005	0.0475	0.4905	0.9555	1
		$\gamma = 1.5$	0	0.0040	0.0325	0.1625	0.4565	0.7910	0	0.0005	0.0475	0.4835	0.9465	1
		$\gamma = 2.0$	0.0005	0.0060	0.0305	0.1335	0.3850	0.6990	0	0.0005	0.0475	0.4775	0.9410	1
	$p = 4$	$\gamma = 0.5$	0	0	0.0445	0.5050	0.9600	0.9995	0	0	0.0405	0.7565	1	1
		$\gamma = 1.0$	0.0005	0.0080	0.0415	0.2200	0.5815	0.8945	0	0	0.0480	0.7510	1	1
		$\gamma = 1.5$	0.0115	0.0235	0.0480	0.1090	0.2260	0.4240	0	0	0.0475	0.7200	0.9980	1
		$\gamma = 2.0$	0.0215	0.0305	0.0515	0.0895	0.1550	0.2700	0	0	0.0535	0.6995	0.9960	1
	$p = 6$	$\gamma = 0.5$	0	0.0005	0.0415	0.6205	0.9920	1	0	0	0.0280	0.8925	1	1
		$\gamma = 1.0$	0.0020	0.0115	0.0450	0.1690	0.4485	0.7850	0	0	0.0460	0.8345	1	1
		$\gamma = 1.5$	0.0295	0.0350	0.0450	0.0755	0.1170	0.1735	0	0	0.0615	0.7910	0.9995	1
		$\gamma = 2.0$	0.0470	0.0515	0.0585	0.0670	0.0855	0.1090	0	0	0.0745	0.7755	0.9990	1

Table 3: Simulation results for Example 3. Rejection rates of  $T_{n,S}^{(\mathcal{I})}$  at 5% significance level using 2000 trials with the variance of  $\Delta_{n,w}^{(\mathcal{I})}$  estimated by jackknife.

		Example 3(a)						Example 3(b)						
		Null hypotheses			Alternatives			Null hypotheses			Alternatives			
		$\rho = 0.84$	$\rho = 0.82$	$\rho = 0.8$	$\rho = 0.75$	$\rho = 0.7$	$\rho = 0.65$	$\rho = 0.84$	$\rho = 0.82$	$\rho = 0.8$	$\rho = 0.75$	$\rho = 0.7$	$\rho = 0.65$	
$n = 100$	$p = 2$	$\gamma = 0.5$	0.0025	0.0100	0.0415	0.3825	0.8050	0.9660	0	0.0030	0.0185	0.2470	0.7190	0.9480
		$\gamma = 1.0$	0.0015	0.0090	0.0390	0.4275	0.8505	0.9765	0	0.0040	0.0175	0.2945	0.7760	0.9610
		$\gamma = 1.5$	0.0005	0.0090	0.0430	0.4575	0.8740	0.9825	0.0005	0.0030	0.0240	0.3280	0.7835	0.9555
		$\gamma = 2.0$	0	0.0080	0.0450	0.4775	0.8825	0.9845	0.0005	0.0050	0.0295	0.3360	0.7815	0.9490
	$p = 4$	$\gamma = 0.5$	0	0.0020	0.0240	0.4955	0.9525	0.9995	0	0.0005	0.0120	0.3385	0.8795	0.9960
		$\gamma = 1.0$	0	0.0025	0.0285	0.5500	0.9630	0.9990	0	0.0060	0.0265	0.3345	0.8085	0.9735
		$\gamma = 1.5$	0	0.0040	0.0370	0.4910	0.9270	0.9970	0.0030	0.0140	0.0430	0.3050	0.6680	0.9035
		$\gamma = 2.0$	0.0010	0.0040	0.0450	0.4240	0.8605	0.9835	0.0055	0.0205	0.0575	0.2810	0.5945	0.8300
	$p = 6$	$\gamma = 0.5$	0	0.0005	0.0095	0.5720	0.9795	1	0	0	0.0030	0.2455	0.8430	0.9860
		$\gamma = 1.0$	0	0.0035	0.0270	0.5560	0.9620	1	0.0010	0.0080	0.0410	0.2565	0.6545	0.9020
		$\gamma = 1.5$	0.0015	0.0145	0.0550	0.4155	0.8155	0.9625	0.0130	0.0390	0.0750	0.2545	0.5145	0.7345
		$\gamma = 2.0$	0.0105	0.0320	0.0815	0.3600	0.6500	0.8230	0.0325	0.0595	0.0975	0.2705	0.4670	0.6460
$n = 200$	$p = 2$	$\gamma = 0.5$	0	0.0050	0.0505	0.6450	0.9845	1	0	0.0025	0.0230	0.5350	0.9655	1
		$\gamma = 1.0$	0	0.0030	0.0500	0.7005	0.9915	1	0	0.0025	0.0240	0.5930	0.9785	1
		$\gamma = 1.5$	0	0.0025	0.0485	0.7515	0.9945	0.9995	0	0.0030	0.0280	0.6195	0.9785	0.9990
		$\gamma = 2.0$	0	0.0035	0.0500	0.7680	0.9955	0.9995	0	0.0040	0.0340	0.6095	0.9740	0.9990
	$p = 4$	$\gamma = 0.5$	0	0	0.0310	0.8475	0.9990	1	0	0	0.0085	0.7895	1	1
		$\gamma = 1.0$	0	0	0.0310	0.8910	1	1	0	0	0.0165	0.6515	0.9895	1
		$\gamma = 1.5$	0	0.0010	0.0370	0.8420	0.9995	1	0	0.0025	0.0295	0.4880	0.9400	0.9975
		$\gamma = 2.0$	0	0.0010	0.0410	0.7740	0.9965	1	0	0.0065	0.0370	0.4205	0.8715	0.9835
	$p = 6$	$\gamma = 0.5$	0	0	0.0160	0.9330	1	1	0	0	0.0020	0.7040	0.9985	1
		$\gamma = 1.0$	0	0.0005	0.0320	0.9060	1	1	0	0.0030	0.0240	0.4515	0.9160	0.9970
		$\gamma = 1.5$	0	0.0020	0.0460	0.7410	0.9980	1	0.0010	0.0165	0.0505	0.3425	0.7420	0.9300
		$\gamma = 2.0$	0	0.0065	0.0565	0.5635	0.9430	0.9960	0.0045	0.0250	0.0740	0.3235	0.6530	0.8545
$n = 300$	$p = 2$	$\gamma = 0.5$	0	0.0040	0.0635	0.8110	0.9970	1	0	0.0010	0.0195	0.7030	0.9965	1
		$\gamma = 1.0$	0	0.0020	0.0655	0.8515	0.9990	1	0	0.0005	0.0235	0.7730	0.9990	1
		$\gamma = 1.5$	0	0.0025	0.0645	0.8970	1	1	0	0.0005	0.0290	0.7775	0.9985	1
		$\gamma = 2.0$	0	0.0020	0.0630	0.9135	1	1	0	0.0005	0.0310	0.7730	0.9985	1
	$p = 4$	$\gamma = 0.5$	0	0.0005	0.0375	0.9670	1	1	0	0	0.0125	0.9490	1	1
		$\gamma = 1.0$	0	0	0.0355	0.9790	1	1	0	0.0005	0.0185	0.8230	1	1
		$\gamma = 1.5$	0	0.0010	0.0455	0.9615	1	1	0	0.0020	0.0255	0.6330	0.9870	1
		$\gamma = 2.0$	0	0.0025	0.0530	0.9275	1	1	0.0005	0.0040	0.0315	0.5445	0.9585	0.9995
	$p = 6$	$\gamma = 0.5$	0	0	0.0120	0.9960	1	1	0	0	0.0050	0.9150	1	1
		$\gamma = 1.0$	0	0	0.0265	0.9850	1	1	0	0.0015	0.0290	0.6050	0.9850	1
		$\gamma = 1.5$	0	0	0.0460	0.9100	1	1	0.0015	0.0090	0.0435	0.4310	0.8720	0.9875
		$\gamma = 2.0$	0	0.0020	0.0525	0.7765	0.9945	1	0.0020	0.0165	0.0545	0.3875	0.7920	0.9515




Figure 1: Simulation results for Example 1(a) based on 2000 trials. The horizontal line corresponds to the 5% significance level. The capital letters S, L, and E in panel (a) represent  $T_{n,S}^{(S)}$ ,  $T_{n,L}^{(S)}$ , and  $T_{n,E}^{(S)}$  respectively.




Figure 2: Simulation results for Example 1(b) based on 2000 trials. The horizontal line corresponds to the 5% significance level. The capital letters S, L, and E in panel (a) represent  $T_{n,S}^{(S)}$ ,  $T_{n,L}^{(S)}$ , and  $T_{n,E}^{(S)}$  respectively.




Figure 3: Simulation results for Example 2(a) based on 2000 trials. The horizontal line corresponds to the 5% significance level. The capital letters S, L, and E in panel (a) represent  $T_{n,S}^{(\mathcal{H})}$ ,  $T_{n,L}^{(\mathcal{H})}$ , and  $T_{n,E}^{(\mathcal{H})}$  respectively.




Figure 4: Simulation results for Example 2(b) based on 2000 trials. The horizontal line corresponds to the 5% significance level. The capital letters S, L, and E in panel (a) represent  $T_{n,S}^{(\mathcal{H})}$ ,  $T_{n,L}^{(\mathcal{H})}$ , and  $T_{n,E}^{(\mathcal{H})}$  respectively.




Figure 5: Simulation results for Example 3(a) based on 2000 trials. The horizontal line corresponds to the 5% significance level. The capital letters S, L, and E in panel (a) represent  $T_{n,S}^{(\mathcal{I})}$ ,  $T_{n,L}^{(\mathcal{I})}$ , and  $T_{n,E}^{(\mathcal{I})}$  respectively.






Figure 6: Simulation results for Example 3(b) based on 2000 trials. The horizontal line corresponds to the 5% significance level. The capital letters S, L, and E in panel (a) represent  $T_{n,S}^{(\mathcal{I})}$ ,  $T_{n,L}^{(\mathcal{I})}$ , and  $T_{n,E}^{(\mathcal{I})}$  respectively.

Table 4: Values of  $\Delta_{w,Z}^{(\mathcal{H})}$  for the benchmark configuration in Example 2(a) based on the random approximation (RA) and numerical integration (NI).

Weight function		$p = 2$		$p = 4$		$p = 6$	
		RA	NI	RA	NI	RA	NI
Stable density	$\gamma = 0.5$	0.216287	0.216954	0.177086	0.182100	0.154451	0.150669
	$\gamma = 1.0$	0.312581	0.315284	0.170617	0.171631	0.097647	0.098159
	$\gamma = 1.5$	0.320700	0.324544	0.107838	0.108673	0.034544	0.034622
	$\gamma = 2.0$	0.314935	0.319257	0.076042	0.076714	0.015729	0.015833
Laplace density	$\gamma = 0.1$	0.160431	0.160761	0.167135	0.167664	0.163473	0.161230
	$\gamma = 0.25$	0.307405	0.308412	0.290496	0.291522	0.266015	0.264073
	$\gamma = 1.0$	0.388005	0.391292	0.221240	0.222430	0.141637	0.141809
	$\gamma = 4.0$	0.122471	0.124460	0.014131	0.014279	0.001981	0.002002