

Overlapping and nonoverlapping models

Huan Qing

QINGHUAN@CUMT.EDU.CN

School of Mathematics

China University of Mining and Technology

Xuzhou, 221116, P.R. China

Abstract

Consider a directed network with K_r row communities and K_c column communities. Previous works found that modeling directed networks in which all nodes have overlapping property requires $K_r = K_c$ for identifiability. In this paper, we propose an overlapping and nonoverlapping model to study directed networks in which row nodes have overlapping property while column nodes do not. The proposed model is identifiable when $K_r \leq K_c$. Meanwhile, we provide one identifiable model as extension of ONM to model directed networks with variation in node degree. Two spectral algorithms with theoretical guarantee on consistent estimations are designed to fit the models. A small scale of numerical studies are used to illustrate the algorithms.

Keywords: Community detection, directed networks, spectral clustering, asymptotic analysis, SVD.

1. Introduction

In the study of social networks, various models have been proposed to learn the latent structure of networks. Due to the extremely intensive studies on community detection, we only focus on identifiable models that are closely relevant to our study in this paper. For undirected network, the Stochastic Blockmodel (SBM) (Holland et al., 1983) is a classical and widely used model to generate undirected networks. The degree-corrected stochastic blockmodel (DCSBM) Karrer and Newman (2011) extends SBM by introducing degree heterogeneities. Under SBM and DCSBM, all nodes are pure such that each node only belong to one community. While, in real cases some nodes may belong to multiple communities, and such nodes have overlapping (also known as mixed membership) property. To model undirected networks in which nodes have overlapping property, Airoldi et al. (2008) designs the Mixed Membership Stochastic Blockmodel (MMSB). Jin et al. (2017) introduces the degree-corrected mixed membership model (DCMM) which extends MMSB by considering degree heterogeneities. Zhang et al. (2020) designs the OCCAM model which equals DCMM actually. Spectral methods with consistent estimations under the above models are provided in Rohe et al. (2011); Qin and Rohe (2013); Lei and Rinaldo (2015); Joseph and Yu (2016); Jin (2015); Jin et al. (2017); Mao et al. (2020, 2018). For directed networks in which all nodes have nonoverlapping property, Rohe et al. (2016) proposes a model called Stochastic co-Blockmodel (ScBM) and its extension DCScBM by considering degree heterogeneity, where ScBM (DCScBM) is extension of SBM (DCSBM). ScBM and DCScBM can model nonoverlapping directed networks in which row nodes belong to K_r

row communities and column nodes belong to K_c column communities, where K_r can differ from K_c . Zhou and A.Amini (2019); Qing and Wang (2021b) study the consistency of some adjacency-based spectral algorithms under ScBM. Wang et al. (2020) studies the consistency of the spectral method D-SCORE under DCScBM when $K_r = K_c$. Qing and Wang (2021a) designs directed mixed membership stochastic blockmodel (DiMMSB) as an extension of ScBM and MMSB to model directed networks in which all nodes have overlapping property. Meanwhile, DiMMSB can also be seen as an extension of the two-way blockmodels with Bernoulli distribution of Airoldi et al. (2013). All the above models are identifiable under certain conditions. The identifiability of ScBM and DCScBM holds even for the case when $K_r \neq K_c$. DiMMSB is identifiable only when $K_r = K_c$. Sure, SBM, DCScBM, MMSB, DCMM and OCCAM are identifiable when $K_r = K_c$ since they model undirected networks. For all the above models, row nodes and column nodes have symmetric structural information such that they always have nonoverlapping property or overlapping property simultaneously. As shown by the identifiability of DiMMSB, to model a directed network in which all nodes have overlapping property, the identifiability of the model requires $K_r = K_c$. Naturally, there is a bridge model from ScBM to DiMMSB such that the bride model can model a directed network in which row nodes and column nodes have asymmetric structural information such that they have different overlapping property. In this paper, we introduce this model and name it as overlapping and nonoverlapping model.

Our contributions in this paper are as follows. We propose an identifiable model for directed networks, the overlapping and nonoverlapping model (ONM for short). ONM allows that nodes in a directed network can have different overlapping property. Without loss of generality, in a directed network, we let row nodes have overlapping property while column nodes do not. The proposed model is identifiable when $K_r \leq K_c$. Recall that the identifiability of ScBM modeling nonoverlapping directed networks holds even for the case $K_r \neq K_c$, and DiMMSB modeling overlapping directed networks is identifiable only when $K_r = K_c$, this is the reason we call ONM modeling directed networks in which row nodes have different overlapping property as column nodes as a bridge model from ScBM to DiMMSB. Similar as DCScBM is an extension of ScBM, we propose an identifiable model overlapping and degree-corrected nonoverlapping model (ODCNM) as extension of ONM by considering degree heterogeneity. We construct two spectral algorithms to fit ONM and ODCNM. We show that our method enjoy consistent estimations under mild conditions by delicate spectral analysis. Especially, our theoretical results under ODCNM match those under ONM when ODCNM degenerates to ONM.

Notations. We take the following general notations in this paper. For any positive integer m , let $[m] := \{1, 2, \dots, m\}$. For a vector x and fixed $q > 0$, $\|x\|_q$ denotes its l_q -norm. For a matrix M , M' denotes the transpose of the matrix M , $\|M\|$ denotes the spectral norm, $\|M\|_F$ denotes the Frobenius norm, and $\|M\|_{2 \rightarrow \infty}$ denotes the maximum l_2 -norm of all the rows of M . Let $\sigma_i(M)$ be the i -th largest singular value of matrix M , and $\lambda_i(M)$ denote the i -th largest eigenvalue of the matrix M ordered by the magnitude. $M(i, :)$ and $M(:, j)$ denote the i -th row and the j -th column of matrix M , respectively. $M(S_r, :)$ and $M(:, S_c)$ denote the rows and columns in the index sets S_r and S_c of matrix M , respectively. For any matrix M , we simply use $Y = \max(0, M)$ to represent $Y_{ij} = \max(0, M_{ij})$ for any i, j . For any matrix $M \in \mathbb{R}^{m \times m}$, let $\text{diag}(M)$ be the $m \times m$ diagonal matrix whose i -th

diagonal entry is $M(i, i)$. $\mathbf{1}$ is a column vector with all entries being ones. e_i is a column vector whose i -th entry is 1 while other entries are zero.

2. The overlapping and nonoverlapping model

Consider a directed network $\mathcal{N} = (V_r, V_c, E)$, where $V_r = \{1, 2, \dots, n_r\}$ is the set of row nodes, $V_c = \{1, 2, \dots, n_c\}$ is the set of column nodes, and E is the set of edges from row nodes to column nodes. Note that since row nodes can be different from column nodes, we may have $V_r \cap V_c = \emptyset$, where \emptyset denotes the null set. In this paper, we use subscript r and c to distinguish terms for row nodes and column nodes. Let $A \in \{0, 1\}^{n_r \times n_c}$ be the bi-adjacency matrix of directed network \mathcal{N} such that $A(i_r, i_c) = 1$ if there is a directional edge from row node i_r to column node i_c , and $A(i_r, i_c) = 0$ otherwise.

We propose a new block model which we call overlapping and nonoverlapping model (ONM for short). ONM can model directed networks whose row nodes belong to K_r overlapping row communities while column nodes belong to K_c nonoverlapping column communities.

For row nodes, let $\Pi_r \in \mathbb{R}^{n_r \times K_r}$ be the membership matrix of row nodes such that

$$\Pi_r(i_r, \cdot) \geq 0, \|\Pi_r(i_r, \cdot)\|_1 = 1 \quad \text{for } i_r \in [n_r]. \quad (1)$$

Call row node i_r *pure* if $\Pi_r(i_r, \cdot)$ degenerates (i.e., one entry is 1, all others $K_r - 1$ entries are 0) and *mixed* otherwise. From such definition, row node i_r has mixed membership and may belong to more than one row communities for $i_r \in [n_r]$.

For column nodes, let ℓ be the $n_c \times 1$ vector whose i_c -th entry $\ell(i_c) = k$ if column node i_c belongs to the k -th column community, and $\ell(i_c)$ takes value from $\{1, 2, \dots, K_c\}$ for $i_c \in [n_c]$. Let $\Pi_c \in \mathbb{R}^{n_c \times K_c}$ be the membership matrix of column nodes such that

$$\Pi_c(i_c, k) = 1 \text{ when } \ell(i_c) = k, \text{ and } 0 \text{ otherwise, and } \|\Pi_c(i_c, \cdot)\|_1 = 1 \quad \text{for } i_c \in [n_c], k \in [K_c]. \quad (2)$$

From such definition, column node i_c belongs to *exactly* one of the K_c column communities for $i_c \in [n_c]$. Sure, all column nodes are pure nodes.

In this paper, we assume that

$$K_r \leq K_c. \quad (3)$$

Eq (3) is required for the identifiability of ONM.

Let $P \in \mathbb{R}^{K_r \times K_c}$ be the probability matrix (also known as connectivity matrix) such that

$$0 \leq P(k, l) \leq \rho \leq 1 \quad \text{for } k \in [K_r], l \in [K_c], \quad (4)$$

where ρ controls the network sparsity and is called sparsity parameter in this paper. For convenience, set $P = \rho \tilde{P}$ where $\tilde{P}(k, l) \in [0, 1]$ for $k \in [K_r], l \in [K_c]$, and $\max_{k \in [K_r], l \in [K_c]} \tilde{P}(k, l) = 1$ for model identifiability. For all pairs of (i_r, i_c) with $i_r \in [n_r], i_c \in [n_c]$, our model assumes that $A(i_r, i_c)$ are independent Bernoulli random variables satisfying

$$\Omega := \Pi_r P \Pi_c', \quad A(i_r, i_c) \sim \text{Bernoulli}(\Omega(i_r, i_c)) \quad \text{for } i_r \in [n_r], i_c \in [n_c], \quad (5)$$

where $\Omega = \mathbb{E}[A]$, and we call it population adjacency matrix in this paper.

Definition 1 Call model (1)-(5) the *Overlapping and Nonoverlapping model (ONM)* and denote it by $ONM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c)$.

The following conditions are sufficient for the identifiability of ONM:

- (I1) $\text{rank}(P) = K_r, \text{rank}(\Pi_r) = K_r$ and $\text{rank}(\Pi_c) = K_c$.
- (I2) There is at least one pure row node for each of the K_r row communities.

Here, $\text{rank}(\Pi_r) = K_r$ means that $\sum_{i_r=1}^{n_r} (\Pi_r(i_r, k)) > 0$ for all $k \in [K_r]$; $\text{rank}(\Pi_c) = K_c$ means that each column community has at least one column node. For $k \in [K_r]$, let $\mathcal{I}_r^{(k)} = \{i \in \{1, 2, \dots, n_r\} : \Pi_r(i, k) = 1\}$. By condition (I2), $\mathcal{I}_r^{(k)}$ is non empty for all $k \in [K_r]$. For $k \in [K_r]$, select one row node from $\mathcal{I}_r^{(k)}$ to construct the index set \mathcal{I}_r , i.e., \mathcal{I}_r is the indices of row nodes corresponding to K_r pure row nodes, one from each community. W.L.O.G., let $\Pi_r(\mathcal{I}_r, :) = I_{K_r}$ (Lemma 2.1 Mao et al. (2020) also has similar setting to design their spectral algorithms under MMSB.). \mathcal{I}_c is defined similarly for column nodes such that $\Pi_c(\mathcal{I}_c, :) = I_{K_c}$. Next proposition guarantees that once conditions (I1) and (I2) hold, ONM is identifiable.

Proposition 2 *If conditions (I1) and (I2) hold, ONM is identifiable: For eligible (P, Π_r, Π_c) and $(\check{P}, \check{\Pi}_r, \check{\Pi}_c)$, if $\Pi_r P \Pi_c' = \check{\Pi}_r \check{P} \check{\Pi}_c'$, then $P = \check{P}, \Pi_r = \check{\Pi}_r$, and $\Pi_c = \check{\Pi}_c$.*

Compared to some previous models for directed networks, ONM models different directed networks.

- When all row nodes are pure, our ONM reduces to ScBM with K_r row clusters and K_c column clusters Rohe et al. (2016). However, ONM allows row nodes to have overlapping memberships while ScBM does not. Meanwhile, for model identifiability, ScBM does not require $\text{rank}(P) = K_r$ while ONM requires, and this can be seen as the cost of ONM when modeling overlapping row nodes.
- Though DiMMSB Qing and Wang (2021a) can model directed networks whose row and column nodes have overlapping memberships, DiMMSB requires $K_r = K_c$ for model identifiability. For comparison, our ONM allows $K_r \leq K_c$ at the cost of losing overlapping property of column nodes.

2.1 A spectral algorithm for fitting ONM

The primary goal of the proposed algorithm is to estimate the row membership matrix Π_r and column membership matrix Π_c from the observed adjacency matrix A with given K_r and K_c .

We now discuss our intuition for the design of our algorithm to fit ONM. Under conditions (I1) and (I2), by basic algebra, we have $\text{rank}(\Omega) = K_r$. Let $\Omega = U_r \Lambda U_c'$ be the compact singular value decomposition of Ω , where $U_r \in \mathbb{R}^{n_r \times K_r}, \Lambda \in \mathbb{R}^{K_r \times K_r}, U_c \in \mathbb{R}^{n_c \times K_r}, U_r' U_r = I_{K_r}, U_c' U_c = I_{K_r}$, and I_{K_r} is a $K_r \times K_r$ identity matrix. Let $n_{c,k} = |\{i_c : \ell(i_c) = k\}|$ be the size of the k -th column community for $k \in [K_c]$. Let $n_{c,\max} = \max_{k \in [K_c]} n_{c,k}$ and $n_{c,\min} = \min_{k \in [K_c]} n_{c,k}$. Meanwhile, without causing confusion, let n_{c,K_r} be the K_r -th largest size among all column communities. The following lemma guarantees that U_r enjoys ideal simplex structure and U_c has K_c distinct rows.

Lemma 3 Under $ONM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c)$, there exist an unique $K_r \times K_r$ matrix B_r and an unique $K_c \times K_r$ matrix B_c such that

- $U_r = \Pi_r B_r$ where $B_r = U_r(\mathcal{I}_r, :)$. Meanwhile, $U_r(i_r, :) = U_r(\bar{i}_r, :)$ when $\Pi_r(i_r, :) = \Pi_r(\bar{i}_r, :)$ for $i_r, \bar{i}_r \in [n_r]$.
- $U_c = \Pi_c B_c$. Meanwhile, $U_c(i_c, :) = U_c(\bar{i}_c, :)$ when $\ell(i_c) = \ell(\bar{i}_c)$ for $i_c, \bar{i}_c \in [n_c]$, i.e., U_c has K_c distinct rows. Furthermore, when $K_r = K_c = K$, we have $\|B_c(k, :) - B_c(l, :)\|_F = \sqrt{\frac{1}{n_{c,k}} + \frac{1}{n_{c,l}}}$ for all $1 \leq k < l \leq K$.

Lemma 3 says that the rows of U_c form a K_r -simplex in \mathbb{R}^{K_r} which we call the Ideal Simplex (IS), with the K_r rows of B_r being the vertices. Such IS is also found in Jin et al. (2017); Mao et al. (2020); Qing and Wang (2021a). Meanwhile, Lemma 3 says that U_c has K_c distinct rows, and if two column nodes i_c and \bar{i}_c are from the same column community, then $U_c(i_c, :) = U_c(\bar{i}_c, :)$.

Under ONM, to recover Π_c from U_c , since U_c has K_c distinct rows, applying k-means algorithm on all rows of U_c returns true column communities by Lemma 3. Meanwhile, since U_c has K_c distinct rows, we can set $\delta_c = \min_{k \neq l} \|B_c(k, :) - B_c(l, :)\|_F$ to measure the minimum center separation of B_c . By Lemma 3, $\delta_c \geq \sqrt{\frac{2}{n_{c, \max}}}$ when $K_r = K_c = K$ under $ONM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c)$. However, when $K_r < K_c$, it is challenge to obtain a positive lower bound of δ_c , see the proof of Lemma 3 for detail.

Under ONM, to recover Π_c from U_c , since B_r is full rank, if U_r and B_r are known in advance ideally, we can exactly recover Π_r by setting $\Pi_r = U_r B_r' (B_r B_r')^{-1}$ by Lemma 3. Set $Y_r = U_r B_r' (B_r B_r')^{-1}$, since $Y_r \equiv \Pi_r$ and $\|\Pi_r(i_r, :)\|_1 = 1$ for $i_r \in [n_r]$, we have

$$\Pi_r(i_r, :) = \frac{Y_r(i_r, :)}{\|Y_r(i_r, :)\|_1}, i_r \in [n_r].$$

With given U_r , since it enjoys IS structure $U_r = \Pi_r B_r \equiv \Pi_r U_r(\mathcal{I}_r, :)$, as long as we can obtain the row corner matrix $U_r(\mathcal{I}_r, :)$ (i.e., B_r), we can recover Π_r exactly. As mentioned in Jin et al. (2017); Mao et al. (2020), for such ideal simplex, the successive projection (SP) algorithm Gillis and Vavasis (2015) (for detail of SP, see Algorithm 3) can be applied to U_r with K_r row communities to find $U_r(\mathcal{I}_r, :)$.

Based on the above analysis, we are now ready to give the following algorithm which we call Ideal ONA. Input Ω, K_r, K_c with $K_r \leq K_c$. Output: Π_r and ℓ .

- Let $\Omega = U_r \Lambda U_c'$ be the compact SVD of Ω such that $U_r \in \mathbb{R}^{n_r \times K_r}, U_c \in \mathbb{R}^{n_c \times K_r}, \Lambda \in \mathbb{R}^{K_r \times K_r}, U_r' U_r = I_{K_r}, U_c' U_c = I_{K_r}$.
- For row nodes,
 - Run SP algorithm on all rows of U_r assuming there are K_r row communities to obtain $U_r(\mathcal{I}_r, :)$. Set $B_r = U_r(\mathcal{I}_r, :)$.
 - Set $Y_r = U_r B_r' (B_r B_r')^{-1}$. Recover Π_r by setting $\Pi_r(i_r, :) = \frac{Y_r(i_r, :)}{\|Y_r(i_r, :)\|_1}$ for $i_r \in [n_r]$.

For column nodes,

- Run k-means on U_c assuming there are K_c column communities, i.e., find the solution to the following optimization problem

$$M^* = \operatorname{argmin}_{M \in M_{n_c, K_r, K_c}} \|M - U_c\|_F^2,$$

where M_{n_c, K_r, K_c} denotes the set of $n_c \times K_r$ matrices with only K_c different rows.

- use M^* to obtain the labels vector ℓ of column nodes.

Follow similar proof of Theorem 1 of Qing and Wang (2021a), Ideal ONA exactly recovers row nodes memberships and column nodes labels, and this also verifies the identifiability of ONM in turn. For convenience, call the two steps for column nodes as “run k-means on U_c assuming there are K_c column communities to obtain ℓ ”.

We now extend the ideal case to the real case. Set $\tilde{A} = \hat{U}_r \hat{\Lambda} \hat{U}_c'$ be the top- K_r -dimensional SVD of A such that $\hat{U}_r \in \mathbb{R}^{n_r \times K_r}$, $\hat{U}_c \in \mathbb{R}^{n_c \times K_r}$, $\hat{\Lambda} \in \mathbb{R}^{K_r \times K_r}$, $\hat{U}_r' \hat{U}_r = I_{K_r}$, $\hat{U}_c' \hat{U}_c = I_{K_r}$, and $\hat{\Lambda}$ contains the top K_r singular values of A . For the real case, we use $\hat{B}_r, \hat{B}_c, \hat{Y}_r, \hat{\Pi}_r, \hat{\Pi}_c$ given in Algorithm 1 to estimate $B_r, B_c, Y_r, \Pi_r, \Pi_c$, respectively. Algorithm 1 called overlapping and nonoverlapping algorithm (ONA for short) is a natural extension of the Ideal ONA to the real case. In ONA, we set the negative entries of \hat{Y}_r as 0 by setting $\hat{Y}_r = \max(0, \hat{Y}_r)$ for the reason that weights for any row node should be nonnegative while there may exist some negative entries of $\hat{U}_r \hat{B}_r' (\hat{B}_r \hat{B}_r')^{-1}$. Note that, in a directed network, if column nodes have overlapping property while row nodes do not, to do community detection for such directed network, set the transpose of the adjacency matrix as input when applying our algorithm.

Algorithm 1 Overlapping and Nonoverlapping Algorithm (ONA)

Require: The adjacency matrix $A \in \mathbb{R}^{n_r \times n_c}$ of a directed network, the number of row communities K_r , and the number of column communities K_c with $K_r \leq K_c$.

Ensure: The estimated $n_r \times K_r$ membership matrix $\hat{\Pi}_r$ for row nodes, and the estimated $n_c \times 1$ labels vector $\hat{\ell}$ for column nodes.

- 1: Compute $\hat{U}_r \in \mathbb{R}^{n_r \times K_r}$ and $\hat{U}_c \in \mathbb{R}^{n_c \times K_r}$ from the top- K_r -dimensional SVD of A .
- 2: For row nodes:
 - Apply SP algorithm (i.e., Algorithm 3) on the rows of \hat{U}_r assuming there are K_r row clusters to obtain the near-corners matrix $\hat{U}_r(\hat{\mathcal{I}}_r, :) \in \mathbb{R}^{K_r \times K_r}$, where $\hat{\mathcal{I}}_r$ is the index set returned by SP algorithm. Set $\hat{B}_r = \hat{U}_r(\hat{\mathcal{I}}_r, :)$.
 - Compute the $n_r \times K_r$ matrix \hat{Y}_r such that $\hat{Y}_r = \hat{U}_r \hat{B}_r' (\hat{B}_r \hat{B}_r')^{-1}$. Set $\hat{Y}_r = \max(0, \hat{Y}_r)$ and estimate $\Pi_r(i_r, :)$ by $\hat{\Pi}_r(i_r, :) = \frac{\hat{Y}_r(i_r, :)}{\|\hat{Y}_r(i_r, :)\|_1}, i_r \in [n_r]$.

For column nodes: run k-means on \hat{U}_c assuming there are K_c column communities to obtain $\hat{\ell}$.

2.2 Main results for ONA

In this section, we show the consistency of our algorithm for fitting the ONM as the number of row nodes n_r and the number of column nodes n_c increase. Throughout this paper, $K_r \leq K_c$ are two known integers. First, we assume that

Assumption 4 $\rho \max(n_r, n_c) \geq \log(n_r + n_c)$.

Assumption (4) controls the sparsity of directed network considered for theoretical study. By Lemma 4 of Qing and Wang (2021a), we have below lemma.

Lemma 5 (*Row-wise singular eigenvector error*) *Under $ONM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c)$, when Assumption (4) holds, suppose $\sigma_{K_r}(\Omega) \geq C\sqrt{\rho(n_r + n_c)\log(n_r + n_c)}$, with probability at least $1 - o((n_r + n_c)^{-\alpha})$,*

$$\|\hat{U}_r \hat{U}_r' - U_r U_r'\|_{2 \rightarrow \infty} = O\left(\frac{\sqrt{K_r}(\kappa(\Omega))\sqrt{\frac{\max(n_r, n_c)\mu}{\min(n_r, n_c)} + \sqrt{\log(n_r + n_c)}}}{\sqrt{\rho}\sigma_{K_r}(\tilde{P})\sigma_{K_r}(\Pi_r)\sqrt{n_{c, K_r}}}\right),$$

where μ is the incoherence parameter defined as $\mu = \max\left(\frac{n_r \|U_r\|_{2 \rightarrow \infty}^2}{K_r}, \frac{n_c \|U_c\|_{2 \rightarrow \infty}^2}{K_r}\right)$.

For convenience, set $\varpi = \|\hat{U}_r \hat{U}_r' - U_r U_r'\|_{2 \rightarrow \infty}$ in this paper. To measure the performance of ONA for row nodes memberships, since row nodes have mixed memberships, naturally, we use the l_1 norm difference between Π_r and $\hat{\Pi}_r$. Since column nodes are all pure nodes, we consider the performance criterion defined in Joseph and Yu (2016) to measure estimation error of ONA on column nodes. We introduce this measurement of estimation error as below.

Let $\mathcal{T}_c = \{\mathcal{T}_{c,1}, \mathcal{T}_{c,2}, \dots, \mathcal{T}_{c,K_c}\}$ be the true partition of column nodes $\{1, 2, \dots, n_c\}$ obtained from ℓ such that $\mathcal{T}_{c,k} = \{i_c : \ell(i_c) = k\}$ for $k \in [K_c]$. Let $\hat{\mathcal{T}}_c = \{\hat{\mathcal{T}}_{c,1}, \hat{\mathcal{T}}_{c,2}, \dots, \hat{\mathcal{T}}_{c,K_c}\}$ be the estimated partition of column nodes $\{1, 2, \dots, n_c\}$ obtained from $\hat{\ell}$ of ONA such that $\hat{\mathcal{T}}_{c,k} = \{i_c : \hat{\ell}(i_c) = k\}$ for $k \in [K_c]$. The criterion is defined as

$$\hat{f}_c = \min_{\pi \in S_{K_c}} \max_{k \in [K_c]} \frac{|\mathcal{T}_{c,k} \cap \hat{\mathcal{T}}_{c,\pi(k)}^c| + |\mathcal{T}_{c,k}^c \cap \hat{\mathcal{T}}_{c,\pi(k)}|}{n_{c,k}},$$

where S_{K_c} is the set of all permutations of $\{1, 2, \dots, K_c\}$ and the superscript c denotes complementary set. As mentioned in Joseph and Yu (2016), \hat{f}_c measures the maximum proportion of column nodes in the symmetric difference of $\mathcal{T}_{c,k}$ and $\hat{\mathcal{T}}_{c,\pi(k)}$.

Next theorem gives theoretical bounds on estimations of memberships for both row and column nodes, which is the main theoretical result for ONA.

Theorem 6 *Under $ONM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c)$, suppose conditions in Lemma 5 hold, with probability at least $1 - o((n_r + n_c)^{-\alpha})$,*

- for row nodes, there exists a permutation matrix \mathcal{P}_r such that

$$\max_{i_r \in [n_r]} \|e'_{i_r}(\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O(\varpi \kappa(\Pi_r' \Pi_r) K_r \sqrt{\lambda_1(\Pi_r' \Pi_r)}).$$

- for column nodes,

$$\hat{f}_c = O\left(\frac{K_r K_c \max(n_r, n_c) \log(n_r + n_c)}{\sigma_{K_r}^2(\tilde{P}) \rho \delta_c^2 \sigma_{K_r}^2(\Pi_r) n_{c, K_r} n_{c, \min}}\right).$$

Especially, when $K_r = K_c = K$,

$$\hat{f}_c = O\left(\frac{K^2 \max(n_r, n_c) n_{c, \max} \log(n_r + n_c)}{\sigma_K^2(\tilde{P}) \rho \sigma_K^2(\Pi_r) n_{c, \min}^2}\right).$$

Add conditions similar as Corollary 3.1 in Mao et al. (2020), we have the following corollary.

Corollary 7 Under $ONM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c)$, suppose conditions in Lemma 5 hold, and further suppose that $\lambda_{K_r}(\Pi_r' \Pi_r) = O(\frac{n_r}{K_r})$, $n_{c, \min} = O(\frac{n_c}{K_c})$, with probability at least $1 - o((n_r + n_c)^{-\alpha})$,

- for row nodes, when $K_r = K_c = K$,

$$\max_{i_r \in [n_r]} \|e'_{i_r}(\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O\left(\frac{K^2(\sqrt{\frac{C \max(n_r, n_c)}{\min(n_r, n_c)}} + \sqrt{\log(n_r + n_c)})}{\sigma_K(\tilde{P})\sqrt{\rho n_c}}\right).$$

- for column nodes,

$$\hat{f}_c = O\left(\frac{K_r^2 K_c^3 \max(n_r, n_c) \log(n_r + n_c)}{\sigma_{K_r}^2(\tilde{P}) \rho \delta_c^2 n_r n_c^2}\right).$$

When $K_r = K_c = K$,

$$\hat{f}_c = O\left(\frac{K^4 \max(n_r, n_c) \log(n_r + n_c)}{\sigma_K^2(\tilde{P}) \rho n_r n_c}\right).$$

Especially, when $n_r = O(n)$, $n_c = O(n)$, $K_r = O(1)$ and $K_c = O(1)$,

- for row nodes, when $K_r = K_c$,

$$\max_{i_r \in [n_r]} \|e'_{i_r}(\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O\left(\frac{\sqrt{\log(n)}}{\sigma_{K_r}(\tilde{P})\sqrt{\rho n}}\right).$$

- for column nodes,

$$\hat{f}_c = O\left(\frac{\log(n)}{\sigma_{K_r}^2(\tilde{P}) \rho \delta_c^2 n^2}\right).$$

When $K_r = K_c = K$,

$$\hat{f}_c = O\left(\frac{\log(n)}{\sigma_K^2(\tilde{P}) \rho n}\right).$$

When $K_r \neq K_c$, though it is challenge to obtain the lower bound of δ_c , we can roughly set $\sqrt{\frac{2}{n_{c, \max}}}$ as the lower bound of δ_c since $\delta_c \geq \sqrt{\frac{2}{n_{c, \max}}}$ when $K_r = K_c$.

When ONM degenerates to SBM by setting $\Pi_r = \Pi_c$ and all nodes are pure, applying the separation condition and sharp threshold criterion developed in Qing (2021b) on the upper bounds of error rates in Corollary 7, sure we can obtain the classical separation condition of a balanced network and sharp threshold of the Erdős-Rényi random graph $G(n, p)$ of Erdos and Rényi (2011), and this guarantees the optimality of our theoretical results.

3. The overlapping and degree-corrected nonoverlapping model

Similar as DCSBM Karrer and Newman (2011) is an extension of SBM by introducing node specific parameters to allow for varying degrees, in this section, we propose an extension of ONM by considering degree heterogeneity and build theoretical guarantees for algorithm fitting our model.

Let θ_c be an $n_c \times 1$ vector whose i_c -th entry is the degree heterogeneity of column node i_c , for $i_c \in [n_c]$. Let Θ_c be an $n_c \times n_c$ diagonal matrix whose i_c -th diagonal element is $\theta_c(i_c)$. The extended model for generating A is as follows:

$$\Omega := \Pi_r P \Pi_c' \Theta_c, \quad A(i_r, i_c) \sim \text{Bernoulli}(\Omega(i_r, i_c)) \quad \text{for } i_r \in [n_r], i_c \in [n_c]. \quad (6)$$

Definition 8 Call model (1), (2), (3),(4), (6) the *Overlapping and Degree-Corrected Nonoverlapping model (ODCNM)* and denote it by $ODCNM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c, \Theta_c)$.

Note that, under ODCNM, the maximum element of P can be larger than 1 since $\max_{i_c \in [n_c]} \theta_c(i_c)$ also can control the sparsity of the directed network \mathcal{N} . The following proposition guarantees that ODCNM is identifiable in terms of P, Π_r and Π_c , and such identifiability is similar as that of DCSBM and DCScBM.

Proposition 9 *If conditions (I1) and (I2) hold, ODCNM is identifiable for membership matrices: For eligible $(P, \Pi_r, \Pi_c, \Theta_c)$ and $(\check{P}, \check{\Pi}_r, \check{\Pi}_c, \check{\Theta}_c)$, if $\Pi_r P \Pi_c' \Theta_c = \check{\Pi}_r \check{P} \check{\Pi}_c' \check{\Theta}_c$, then $\Pi_r = \check{\Pi}_r$ and $\Pi_c = \check{\Pi}_c$.*

Remark 10 *By setting $\theta_c(i_c) = \rho$ for $i_c \in [n_c]$, ODCNM reduces to ONM, and this is the reason that ODCNM can be seen as an extension of ONM. Meanwhile, though DC-ScBM Rohe et al. (2016) can model directed networks with degree heterogeneities for both row and column nodes, DCScBM does not allow the overlapping property for nodes. For comparison, our ODCNM allows row nodes have overlapping property at the cost of losing the degree heterogeneities and requiring $K_r \leq K_c$ for model identifiability. Furthermore, another identifiable model extends ONM by considering degree heterogeneity for row nodes with overlapping property is provided in Appendix D, in which we also explain why we do not extend ONM by considering degree heterogeneities for both row and column nodes.*

3.1 A spectral algorithm for fitting ODCNM

We now discuss our intuition for the design of our algorithm to fit ODCNM. Without causing confusion, we also use $U_r, U_c, B_r, B_c, \delta_c, Y_r$, and so on under ODCNM. Let $U_{c,*} \in \mathbb{R}^{n_c \times K_r}$ be the row-normalized version of U_c such that $U_{c,*}(i_c, :) = \frac{U_c(i_c, :)}{\|U_c(i_c, :)\|_F}$ for $i_c \in [n_c]$. Then clustering the rows of $U_{c,*}$ by k-means algorithm can return perfect clustering for column nodes, and this is guaranteed by next lemma.

Lemma 11 *Under $ODCNM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c, \Theta_c)$, there exist an unique $K_r \times K_r$ matrix B_r and an unique $K_c \times K_r$ matrix B_c such that*

- $U_r = \Pi_r B_r$ where $B_r = U_r(\mathcal{I}_r, :)$. Meanwhile, $U_r(i_r, :) = U_r(\bar{i}_r, :)$ when $\Pi_r(i_r, :) = \Pi_r(\bar{i}_r, :)$ for $i_r, \bar{i}_r \in [n_r]$.

- $U_{c,*} = \Pi_c B_c$. Meanwhile, $U_{c,*}(i_c, :) = U_{c,*}(\bar{i}_c, :)$ when $\ell(i_c) = \ell(\bar{i}_c)$ for $i_c, \bar{i}_c \in [n_c]$. Furthermore, when $K_r = K_c = K$, we have $\|B_c(k, :) - B_c(l, :)\|_F = \sqrt{2}$ for all $1 \leq k < l \leq K$.

Recall that we set $\delta_c = \min_{k \neq l} \|B_c(k, :) - B_c(l, :)\|_F$, by Lemma 11, $\delta_c = \sqrt{2}$ when $K_r = K_c = K$ under $ODCNM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c, \Theta_c)$. However, when $K_r < K_c$, it is challenge to obtain a positive lower bound of δ_c , see the proof of Lemma 11 for detail.

Under ODCNM, to recover Π_c from U_c , since $U_{c,*}$ has K_c distinct rows, applying k-means algorithm on all rows of $U_{c,*}$ returns true column communities by Lemma 11; to recover Π_r from U_r , just follow same idea as that of under ONM.

Based on the above analysis, we are now ready to give the following algorithm which we call Ideal ODCNA. Input Ω, K_r, K_c with $K_r \leq K_c$. Output: Π_r and ℓ .

- Let $\Omega = U_r \Lambda U_c'$ be the compact SVD of Ω such that $U_r \in \mathbb{R}^{n_r \times K_r}, U_c \in \mathbb{R}^{n_c \times K_r}, \Lambda \in \mathbb{R}^{K_r \times K_r}, U_r' U_r = I_{K_r}, U_c' U_c = I_{K_r}$. Let $U_{c,*}$ be the row-normalization of U_c .
- For row nodes,
 - Run SP algorithm on all rows of U_r assuming there are K_r row communities to obtain $U_r(\mathcal{I}_r, :)$. Set $B_r = U_r(\mathcal{I}_r, :)$.
 - Set $Y_r = U_r B_r' (B_r B_r')^{-1}$. Recover Π_r by setting $\Pi_r(i_r, :) = \frac{Y_r(i_r, :)}{\|Y_r(i_r, :)\|_1}$ for $i_r \in [n_r]$.

For column nodes: run k-means on $U_{c,*}$ assuming there are K_c column communities to obtain ℓ .

Sure, Ideal ODCNA exactly recoveries row nodes memberships and column nodes labels, and this also supports the identifiability of ODCNM.

We now extend the ideal case to the real case. Let $\hat{U}_{c,*} \in \mathbb{R}^{n_c \times K_r}$ be the row-normalized version of \hat{U}_c such that $\hat{U}_{c,*}(i_c, :) = \frac{\hat{U}_c(i_c, :)}{\|\hat{U}_c(i_c, :)\|_F}$ for $i_c \in [n_c]$. Algorithm 2 called overlapping and degree-corrected nonoverlapping algorithm (ODCNA for short) is a natural extension of the Ideal ODCNA to the real case.

3.2 Main results for ODCNA

Set $\theta_{c,\max} = \max_{i_c \in [n_c]} \theta_c(i_c), \theta_{c,\min} = \min_{i_c \in [n_c]} \theta_c(i_c)$, and $P_{\max} = \max_{k \in [K_r], l \in [n_c]} P(k, l)$. Assume that

Assumption 12 $P_{\max} \max(\theta_{c,\max} n_r, \|\theta_c\|_1) \geq \log(n_r + n_c)$.

By the proof of Lemma 4.3 of Qing (2021a), we have below lemma.

Lemma 13 (Row-wise singular eigenvector error) *Under $ODCNM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c, \Theta_c)$, when Assumption (12) holds, suppose $\sigma_{K_r}(\Omega) \geq C \sqrt{\theta_{c,\max}(n_r + n_c) \log(n_r + n_c)}$, with probability at least $1 - o((n_r + n_c)^{-\alpha})$,*

$$\|\hat{U}_r \hat{U}_r' - U_r U_r'\|_{2 \rightarrow \infty} = O\left(\frac{\sqrt{\theta_{c,\max} K_r} (\kappa(\Omega) \sqrt{\frac{\max(n_r, n_c) \mu}{\min(n_r, n_c)}} + \sqrt{\log(n_r + n_c)})}{\theta_{c,\min} \sigma_{K_r}(P) \sigma_{K_r}(\Pi_r) \sqrt{n_{c, K_r}}}\right).$$

Algorithm 2 Overlapping and Degree-Corrected Nonoverlapping Algorithm (ODCNA)

Require: The adjacency matrix $A \in \mathbb{R}^{n_r \times n_c}$ of a directed network, the number of row communities K_r , and the number of column communities K_c with $K_r \leq K_c$.

Ensure: The estimated $n_r \times K_r$ membership matrix $\hat{\Pi}_r$ for row nodes, and the estimated $n_c \times 1$ labels vector $\hat{\ell}$ for column nodes.

- 1: Compute $\hat{U}_r \in \mathbb{R}^{n_r \times K_r}$ and $\hat{U}_c \in \mathbb{R}^{n_c \times K_r}$ from the top- K_r -dimensional SVD of A . Compute $\hat{U}_{c,*}$ from \hat{U}_c .
- 2: For row nodes:
 - Apply SP algorithm (i.e., Algorithm 3) on the rows of \hat{U}_r assuming there are K_r row clusters to obtain the near-corners matrix $\hat{U}_r(\hat{\mathcal{I}}_r, :) \in \mathbb{R}^{K_r \times K_r}$, where $\hat{\mathcal{I}}_r$ is the index set returned by SP algorithm. Set $\hat{B}_r = \hat{U}_r(\hat{\mathcal{I}}_r, :)$.
 - Compute the $n_r \times K_r$ matrix \hat{Y}_r such that $\hat{Y}_r = \hat{U}_r \hat{B}_r' (\hat{B}_r \hat{B}_r')^{-1}$. Set $\hat{Y}_r = \max(0, \hat{Y}_r)$ and estimate $\Pi_r(i_r, :)$ by $\hat{\Pi}_r(i_r, :) = \frac{\hat{Y}_r(i_r, :)}{\|\hat{Y}_r(i_r, :)\|_1}$, $i_r \in [n_r]$.

For column nodes: run k-means on $\hat{U}_{c,*}$ assuming there are K_c column communities to obtain $\hat{\ell}$.

Next theorem is the main theoretical result for ODCNA, where we also use same measurements as ONA to measure the performances of ODCNA.

Theorem 14 Under $ODCNM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c, \Theta_c)$, suppose conditions in Lemma 13 hold, with probability at least $1 - o((n_r + n_c)^{-\alpha})$,

- for row nodes,

$$\max_{i_r \in [n_r]} \|e'_{i_r} (\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O(\varpi \kappa (\Pi_r' \Pi_r) K_r \sqrt{\lambda_1(\Pi_r' \Pi_r)}).$$

- for column nodes,

$$\hat{f}_c = O\left(\frac{\theta_{c, \max}^2 K_r K_c \max(\theta_{c, \max} n_r, \|\theta_c\|_1) n_{c, \max} \log(n_r + n_c)}{\sigma_{K_r}^2(P) \theta_{c, \min}^4 \delta_c^2 m_{V_c}^2 \sigma_{K_r}^2(\Pi_r) n_{c, K_r} n_{c, \min}}\right),$$

where m_{V_c} is a parameter defined in the proof of this theorem, and it is 1 when $K_r = K_c$. Especially, when $K_r = K_c = K$,

$$\hat{f}_c = O\left(\frac{\theta_{c, \max}^2 K^2 \max(\theta_{c, \max} n_r, \|\theta_c\|_1) n_{c, \max} \log(n_r + n_c)}{\sigma_K^2(P) \theta_{c, \min}^4 \sigma_K^2(\Pi_r) n_{c, \min}^2}\right).$$

Add some conditions on model parameters, we have the following corollary.

Corollary 15 Under $ODCNM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c, \Theta_c)$, suppose conditions in Lemma 13 hold, and further suppose that $\lambda_{K_r}(\Pi_r' \Pi_r) = O(\frac{n_r}{K_r})$, $n_{c, \min} = O(\frac{n_c}{K_c})$, with probability at least $1 - o((n_r + n_c)^{-\alpha})$,

- for row nodes, when $K_r = K_c = K$,

$$\max_{i_r \in [n_r]} \|e'_{i_r}(\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O\left(\frac{K^2 \sqrt{\theta_{c,\max}} \left(\sqrt{\frac{C \max(n_r, n_c)}{\min(n_r, n_c)}} + \sqrt{\log(n_r + n_c)}\right)}{\theta_{c,\min} \sigma_K(P) \sqrt{n_c}}\right).$$

- for column nodes,

$$\hat{f}_c = O\left(\frac{\theta_{c,\max}^2 K_r^2 K_c^2 \max(\theta_{c,\max} n_r, \|\theta_c\|_1) \log(n_r + n_c)}{\sigma_{K_r}^2(P) \theta_{c,\min}^4 \delta_c^2 m_{V_c}^2 n_r n_c}\right).$$

When $K_r = K_c = K$,

$$\hat{f}_c = O\left(\frac{\theta_{c,\max}^2 K^4 \max(\theta_{c,\max} n_r, \|\theta_c\|_1) \log(n_r + n_c)}{\sigma_K^2(P) \theta_{c,\min}^4 n_r n_c}\right).$$

Especially, when $n_r = O(n)$, $n_c = O(n)$, $K_r = O(1)$ and $K_c = O(1)$,

- for row nodes, when $K_r = K_c$,

$$\max_{i_r \in [n_r]} \|e'_{i_r}(\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O\left(\frac{\sqrt{\theta_{c,\max} \log(n)}}{\theta_{c,\min} \sigma_K(P) \sqrt{n}}\right).$$

- for column nodes,

$$\hat{f}_c = O\left(\frac{\theta_{c,\max}^2 \max(\theta_{c,\max} n_r, \|\theta_c\|_1) \log(n)}{\sigma_{K_r}^2(P) \theta_{c,\min}^4 \delta_c^2 m_{V_c}^2 n^2}\right).$$

When $K_r = K_c = K$,

$$\hat{f}_c = O\left(\frac{\theta_{c,\max}^2 \max(\theta_{c,\max} n_r, \|\theta_c\|_1) \log(n)}{\sigma_K^2(P) \theta_{c,\min}^4 n^2}\right).$$

When $K_r \neq K_c$, though it is challenge to obtain the lower bounds of δ_c and m_{V_c} , we can roughly set $\sqrt{2}$ and 1 as the lower bounds of δ_c and m_{V_c} , respectively, since $\delta_c = \sqrt{2}$ and $m_{V_c} = 1$ when $K_r = K_c$. Meanwhile, if we further set $\theta_{c,\max} = O(\rho)$ and $\theta_{c,\min} = O(\rho)$, we have below corollary.

Corollary 16 Under $ODCNM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c, \Theta_c)$, suppose conditions in Lemma 13 hold, and further suppose that $\lambda_{K_r}(\Pi_r \Pi_r) = O(\frac{n_r}{K_r})$, $n_{c,\min} = O(\frac{n_c}{K_c})$ and $\theta_{c,\max} = O(\rho)$, $\theta_{c,\min} = O(\rho)$, with probability at least $1 - o((n_r + n_c)^{-\alpha})$,

- for row nodes, when $K_r = K_c = K$,

$$\max_{i_r \in [n_r]} \|e'_{i_r}(\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O\left(\frac{K^2 \left(\sqrt{\frac{C \max(n_r, n_c)}{\min(n_r, n_c)}} + \sqrt{\log(n_r + n_c)}\right)}{\sigma_K(P) \sqrt{\rho n_c}}\right).$$

- for column nodes,

$$\hat{f}_c = O\left(\frac{K_r^2 K_c^2 \max(n_r, n_c) \log(n_r + n_c)}{\sigma_{K_r}^2(P) \rho \delta_c^2 m_{V_c}^2 n_r n_c}\right).$$

When $K_r = K_c = K$,

$$\hat{f}_c = O\left(\frac{K^4 \max(n_r, n_c) \log(n_r + n_c)}{\sigma_K^2(P) \rho n_r n_c}\right).$$

Especially, when $n_r = O(n)$, $n_c = O(n)$, $K_r = O(1)$ and $K_c = O(1)$,

- for row nodes, when $K_r = K_c$,

$$\max_{i_r \in [n_r]} \|e'_{i_r} (\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O\left(\frac{\sqrt{\log(n)}}{\sigma_K(P) \sqrt{\rho n}}\right).$$

- for column nodes,

$$\hat{f}_c = O\left(\frac{\log(n)}{\sigma_{K_r}^2(P) \rho \delta_c^2 m_{V_c}^2 n}\right).$$

When $K_r = K_c = K$,

$$\hat{f}_c = O\left(\frac{\log(n)}{\sigma_K^2(P) \rho n}\right).$$

By setting $\Theta_c = \rho I$, $ODCNM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c, \Theta_c)$ degenerates to $ONM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c)$. By comparing Corollary 7 and Corollary 16, we see that theoretical results under ODCNM are consistent with those under ONM when ODCNM degenerates to ONM for the case that $K_r = K_c = K$.

4. Simulations

In this section, we present some simulations to investigate the performance of the three proposed algorithms. We measure their performances by Mixed-Hamming error rate (MHamm for short) for row nodes and Hamming error rate (Hamm for short) for column nodes defined below

$$\text{MHamm} = \frac{\min_{\pi \in S_{K_r}} \|\hat{\Pi}_r \pi - \Pi_r\|_1}{n_r} \quad \text{and} \quad \text{Hamm} = \frac{\min_{\pi \in S} \|\hat{\Pi}_c \pi - \Pi_c\|_1}{n_c},$$

where $\hat{\Pi}_c \in \mathbb{R}^{n_c \times K_c}$ is defined as $\hat{\Pi}_c(i_c, k) = 1$ if $\hat{\ell}(i_c) = k$ and 0 otherwise for $i_c \in [n_c]$, $k \in [K_c]$.

For all simulations in this section, the parameters $(n_r, n_c, K_r, K_c, P, \rho, \Pi_r, \Pi_c, \Theta_c)$ are set as follows. Unless specified, set $n_r = 400$, $n_c = 300$, $K_r = 3$, $K_c = 4$. For column nodes, generate Π_c by setting each column node belonging to one of the column communities with equal probability. Let each row community have 100 pure nodes, and let all the mixed

row nodes have memberships (0.6, 0.3, 0.1). $P = \rho \tilde{P}$ is set independently under ONM and ODCNM. Under ONM, ρ is 0.5 in Experiment 1 and we study the influence of ρ in Experiment 2; Under ODCNM, for $z_c \geq 1$, we generate the degree parameters for column nodes as below: let $\theta_c \in \mathbb{R}^{n_c \times 1}$ such that $1/\theta_c(i_c) \stackrel{iid}{\sim} U(1, z_c)$ for $i_c \in [n_c]$, where $U(1, z_c)$ denotes the uniform distribution on $[1, z_c]$. We study the influences of Z_c and ρ under ODCNM in Experiments 3 and 4, respectively. For all settings, we report the averaged MHamm and the averaged Hamm over 50 repetitions.

Experiment 1: Changing n_c under ONM. Let n_c range in $\{50, 100, 150, \dots, 300\}$. For this experiment, P is set as

$$P = \rho \begin{bmatrix} 1 & 0.3 & 0.2 & 0.3 \\ 0.2 & 0.9 & 0.1 & 0.2 \\ 0.3 & 0.2 & 0.8 & 0.3 \end{bmatrix}.$$

Let $\rho = 0.5$ under for this experiment designed under ONM. The numerical results are shown in panels (a) and (b) of Figure 1. The results show that as n_c increases, ONA and ODCNA perform better. Meanwhile, the total run-time for this experiment is roughly 70 seconds. For row nodes, since both ONA and ODCNA apply SP algorithm on \tilde{U} to estimate Π_r , the estimated row membership matrices of ONA and ODCNA are same, and hence MHamm for ONA always equal to that of ODCNA.

Experiment 2: Changing ρ under ONM. P is set same as Experiment 1, and we let ρ range in $\{0.1, 0.2, \dots, 1\}$ to study the influence of ρ on performances of ONA and ODCNA under ONM. The results are displayed in panels (c) and (d) of Figure 1. From the results, we see that both methods perform better as ρ increases since a larger ρ gives more edges generated in a directed network. Meanwhile, the total run-time for this experiment is roughly 136 seconds.


Experiment 3: Change z_c under ODCNM. P is set same as Experiment 1. Let z_c range in $\{1, 2, \dots, 8\}$. Increasing z_c decreases edges generated under ODCNM. Panels (e) and (f) in Figure 1 display simulation results of this experiment. The results show that, generally, increasing the variability of node degrees makes it harder to detect node memberships for both ONA and ODCNA. Though ODCNA is designed under ODCNM, it holds similar performances as ONA for directed networks in which column nodes have various degrees in this experiment, and this is consistent with our theoretical findings in Corollaries 7 and 15. Meanwhile, the total run-time for this experiment is around 131 seconds.

Experiment 4: Change ρ under ODCNM. Set $z_c = 3$, P is set same as Experiment 1, and let ρ range in $\{0.1, 0.2, \dots, 1\}$ under ODCNM. Panels (g) and (h) in Figure 1 displays simulation results of this experiment. The performances of the two proposed methods are similar as that of Experiment 2. Meanwhile, the total run-time for this experiment is around 221 seconds.


5. Discussions

In this paper, we introduced overlapping and nonoverlapping models and its extension by considering degree heterogeneity. The models can model directed network with K_r row communities and K_c column communities, in which row node can belong to multiple row communities while column node only belong to one of the column communities. The


OVERLAPPING AND NONOVERLAPPING MODELS




(a) Changing n_c under ONM: MHamm.




(b) Changing n_c under ONM: Hamm.




(c) Changing ρ under ONM: MHamm.




(d) Changing ρ under ONM: Hamm.




(e) Changing z_c under ODCNM: MHamm.



(f) Changing z_c under ODCNM: Hamm.



(g) Changing ρ under ODCNM: MHamm.



(h) Changing ρ under ODCNM: Hamm.

Figure 1: Estimation errors of ONA and ODCNA.

proposed models are identifiable when $K_r \leq K_c$ and some other popular constraints on the connectivity matrix and membership matrices. For comparison, modeling directed network in which row nodes have overlapping property while column nodes do not with $K_r > K_c$ is unidentifiable. Meanwhile, since previous works found that modeling directed networks in which both row and column nodes have overlapping property with $K_r \neq K_c$ is unidentifiable, our identifiable ONM and ODCNM as well as the DCONM in Appendix D supply a gap in modeling overlapping directed networks when $K_r \neq K_c$. These models provide exploratory tools for studying community structure in directed networks with one side is overlapping while another side is nonoverlapping. Two spectral algorithms are designed to fit ONM and ODCNM. We also showed estimation consistency under mild conditions for our methods. Especially, when ODCNM reduces to ONM, our theoretical results under ODCNM are consistent with those under ONM. But perhaps the main limitation of the models is that the K_r and K_c in the directed network are assumed given, and such limitation also holds for the ScBM and DCScBM of Rohe et al. (2016). In most community problems, the number of row community and the number of column community are unknown, therefore a complete calculation and theoretical study require not only the algorithms and their theoretically consistent estimations described in this paper but also a method for estimating K_r and K_c . We leave studies of this problem to our future work.

Appendix A. Successive Projection algorithm

Algorithm 3 is the Successive Projection algorithm.

Algorithm 3 Successive Projection (SP) (Gillis and Vavasis, 2015)

Require: Near-separable matrix $Y_{sp} = S_{sp}M_{sp} + Z_{sp} \in \mathbb{R}_+^{m \times n}$, where S_{sp}, M_{sp} should satisfy Assumption 1 Gillis and Vavasis (2015), the number r of columns to be extracted.

Ensure: Set of indices \mathcal{K} such that $Y_{sp}(\mathcal{K}, :) \approx S$ (up to permutation)

- 1: Let $R = Y_{sp}, \mathcal{K} = \{\}, k = 1$.
 - 2: **While** $R \neq 0$ and $k \leq r$ **do**
 - 3: $k_* = \operatorname{argmax}_k \|R(k, :)\|_F$.
 - 4: $u_k = R(k_*, :)$.
 - 5: $R \leftarrow (I - \frac{u_k u_k'}{\|u_k\|_F^2})R$.
 - 6: $\mathcal{K} = \mathcal{K} \cup \{k_*\}$.
 - 7: $k = k + 1$.
 - 8: **end while**
-

Appendix B. Proofs under ONM

B.1 Proof of Proposition 2

Proof By Lemma 3, let $U_r \Lambda U_c'$ be the compact SVD of Ω such that $\Omega = U_r \Lambda U_c'$, since $\Omega = \Pi_r P \Pi_c' = \check{\Pi}_r \check{P} \check{\Pi}_c'$, we have $\Omega(\mathcal{I}_r, \mathcal{I}_c) = P = \check{P}$, which gives $P = \check{P}$. By Lemma 3, since $U_r = \Pi_r U_r(\mathcal{I}_r, :) = \check{\Pi}_r U_r(\mathcal{I}_r, :)$, we have $\Pi_r = \check{\Pi}_r$ where we have used the fact that the inverse of $U_r(\mathcal{I}_r, :)$ exists. Since $\Omega = \Pi_r P \Pi_c' = \check{\Pi}_r \check{P} \check{\Pi}_c' = \Pi_r \check{P} \check{\Pi}_c'$, we have $\Pi_r P \Pi_c' = \Pi_r \check{P} \check{\Pi}_c'$. By Lemma 7 of Qing and Wang (2021a), we have $P \Pi_c' = \check{P} \check{\Pi}_c'$, i.e.,

$\Pi_c X = \check{\Pi}_c X$ where we set $X = P' \in \mathbb{R}^{K_c \times K_r}$. Let $\check{\ell}$ be the $n_c \times 1$ vector of column nodes labels obtained from $\check{\Pi}_c$. For $i_c \in [n_c], k \in [K_r]$, from $\Pi_c X = \check{\Pi}_c X$, we have $(\Pi_c X)(i_c, k) = \Pi_c(i_c, :)X(:, k) = X(\ell(i_c), k) = X(\check{\ell}(i_c), k)$, which means that we must have $\ell(i_c) = \check{\ell}(i_c)$ for all $i_c \in [n_c]$, i.e., $\ell = \check{\ell}$ and $\Pi_c = \check{\Pi}_c$. Note that, for the special case $K_r = K_c = K$, $\Pi_c = \check{\Pi}_c$ can be obtained easily: since $P\Pi'_c = P\check{\Pi}'_c$ and $P \in \mathbb{R}^{K \times K}$ is assumed to be full rank, we have $\Pi_c = \check{\Pi}_c$. Thus the proposition holds. \blacksquare

B.2 Proof of Lemma 3

Proof For U_r , since $\Omega = U_r \Lambda U'_c$ and $U'_c U_c = I_{K_r}$, we have $U_r = \Omega U_c \Lambda^{-1}$. Recall that $\Omega = \Pi_r P \Pi'_c$, we have $U_r = \Pi_r P \Pi'_c U_c \Lambda^{-1} = \Pi_r B_r$, where we set $B_r = P \Pi'_c U_c \Lambda^{-1}$. Since $U_r(\mathcal{I}_r, :) = \Pi_r(\mathcal{I}_r, :)B_r = B_r$, we have $B_r = U_r(\mathcal{I}_r, :)$. For $i_r \in [n_r]$, $U_r(i_r, :) = e'_{i_r} \Pi_r B_r = \Pi_r(i_r, :)B_r$, so sure we have $U_r(i_r, :) = U_r(\bar{i}_r, :)$ when $\Pi_r(i_r, :) = \Pi_r(\bar{i}_r, :)$.

For U_c , follow similar analysis as for U_r , we have $U_c = \Pi_c B_c$, where $B_c = P' \Pi'_r U_r \Lambda^{-1}$. Note that $B_c \in \mathbb{R}^{K_c \times K_r}$. Sure, $U_c(i_c, :) = U_c(\bar{i}_c, :)$ when $\ell(i_c) = \ell(\bar{i}_c)$ for $i_c, \bar{i}_c \in [n_c]$.

Now, we focus on the case when $K_r = K_c = K$. For this case, since $B_c \in \mathbb{R}^{K_c \times K_r}$, B_c is full rank when $K_r = K_c$. Since $I_{K_r} = I_K = U'_c U_c = B'_c \Pi'_c \Pi_c B_c$, we have $\Pi'_c \Pi_c = (B_c B'_c)^{-1}$. Since $\Pi'_c \Pi_c = \text{diag}(n_{c,1}, n_{c,2}, \dots, n_{c,K})$, we have $B_c B'_c = \text{diag}(\frac{1}{n_{c,1}}, \frac{1}{n_{c,2}}, \dots, \frac{1}{n_{c,K}})$. When $K_r = K_c = K$, we have $B_c(k, :)B'_c(l, :) = 0$ for any $k \neq l$ and $k, l \in [K]$. Then, we have $B_c B'_c = \text{diag}(\|B_c(1, :)\|_F^2, \|B_c(2, :)\|_F^2, \dots, \|B_c(K, :)\|_F^2) = \text{diag}(\frac{1}{n_{c,1}}, \frac{1}{n_{c,2}}, \dots, \frac{1}{n_{c,K}})$ and the lemma follows.

Note that when $K_r < K_c$, since B_c is not full rank now, we can not obtain $\Pi'_c \Pi_c = (B_c B'_c)^{-1}$ from $I_{K_r} = B'_c \Pi'_c \Pi_c B_c$. Therefore, when $K_r < K_c$, the equality $\|B_c(k, :) - B_c(l, :)\|_F = \sqrt{\frac{1}{n_{c,k}} + \frac{1}{n_{c,l}}}$ does not hold for any $k \neq l$. And we can only know that U_c has K_c distinct rows when $K_r < K_c$, but have no knowledge about the minimum distance between any two distinct rows of U_c . \blacksquare

B.3 Proof of Theorem 6

Proof For row nodes, when conditions in Lemma 5 hold, by Theorem 2 of Qing and Wang (2021a), with probability at least $1 - o((n_r + n_c)^{-\alpha})$ for any $\alpha > 0$, there exists a permutation matrix \mathcal{P}_r such that, for $i_r \in [n_r]$, we have

$$\|e'_{i_r}(\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O(\varpi \kappa(\Pi'_r \Pi_r) K_r \sqrt{\lambda_1(\Pi'_r \Pi_r)}).$$

Next, we focus on column nodes. By the proof of Lemma 2.3 of Qing and Wang (2021b), there exists an orthogonal matrix \hat{O} such that

$$\|\hat{U}_c \hat{O} - U_c\|_F \leq \frac{2\sqrt{2K_r} \|A - \Omega\|}{\sqrt{\lambda_{K_r}(\Omega' \Omega)}}. \quad (7)$$

Under $ONM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c)$, by Lemma 10 of Qing and Wang (2021a), we have

$$\sqrt{\lambda_{K_r}(\Omega' \Omega)} \geq \rho \sigma_{K_r}(\tilde{P}) \sigma_{K_r}(\Pi_r) \sigma_{K_r}(\Pi_c). \quad (8)$$

Since all column nodes are pure, $\sigma_{K_r}(\Pi_c) = \sqrt{n_{c,K_r}}$. By Lemma 3 of Qing and Wang (2021a), when Assumption (4) holds, with probability at least $1 - o((n_r + n_c)^{-\alpha})$, we have

$$\|A - \Omega\| = O(\sqrt{\rho \max(n_r, n_c) \log(n_r + n_c)}). \quad (9)$$

Substitute the two bounds in Eqs (8) and (9) into Eq (7), we have

$$\|\hat{U}_c \hat{O} - U_c\|_F \leq C \frac{\sqrt{K_r \max(n_r, n_c) \log(n_r + n_c)}}{\sigma_{K_r}(\tilde{P}) \sqrt{\rho} \sigma_{K_r}(\Pi_r) \sqrt{n_{c,K_r}}}. \quad (10)$$

Let $\varsigma > 0$ be a small quantity, by Lemma 2 in Joseph and Yu (2016), if

$$\frac{\sqrt{K_c}}{\varsigma} \|U_c - \hat{U}_c \hat{O}\|_F \left(\frac{1}{\sqrt{n_{c,k}}} + \frac{1}{\sqrt{n_{c,l}}} \right) \leq \|B_c(k, :) - B_c(l, :)\|_F, \text{ for each } 1 \leq k \neq l \leq K_c, \quad (11)$$

then the clustering error $\hat{f}_c = O(\varsigma^2)$. Recall that we set $\delta_c = \min_{k \neq l} \|B_c(k, :) - B_c(l, :)\|_F$ to measure the minimum center separation of B_c . Setting $\varsigma = \frac{2}{\delta_c} \sqrt{\frac{K_c}{n_{c,\min}}} \|U_c - \hat{U}_c \hat{O}\|_F$ makes Eq (11) hold for all $1 \leq k \neq l \leq K_c$. Then we have $\hat{f}_c = O(\varsigma^2) = O\left(\frac{K_c \|U_c - \hat{U}_c \hat{O}\|_F^2}{\delta_c^2 n_{c,\min}}\right)$. By Eq (10), we have

$$\hat{f}_c = O\left(\frac{K_r K_c \max(n_r, n_c) \log(n_r + n_c)}{\sigma_{K_r}^2(\tilde{P}) \rho \delta_c^2 \sigma_{K_r}^2(\Pi_r) n_{c,K_r} n_{c,\min}}\right).$$

Especially, when $K_r = K_c = K$, $\delta_c \geq \sqrt{\frac{2}{n_{c,\max}}}$ under $ONM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c)$ by Lemma 3. When $K_r = K_c = K$, we have

$$\hat{f}_c = O\left(\frac{K^2 \max(n_r, n_c) n_{c,\max} \log(n_r + n_c)}{\sigma_K^2(\tilde{P}) \rho \sigma_K^2(\Pi_r) n_{c,\min}^2}\right).$$

■

B.4 Proof of Corollary 7

Proof For row nodes, under conditions of Corollary 7, we have

$$\max_{i_r \in [n_r]} \|e'_{i_r} (\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O(\varpi K_r \sqrt{\frac{n_r}{K_r}}) = O(\varpi \sqrt{K n_r}).$$

Under conditions of Corollary 7, $\kappa(\Omega) = O(1)$ and $\mu \leq C$ for some $C > 0$ by the proof of Corollary 1 Qing and Wang (2021a). Then, by Lemma 5, we have

$$\varpi = O\left(\frac{\sqrt{K} (\kappa(\Omega) \sqrt{\frac{\max(n_r, n_c) \mu}{\min(n_r, n_c)}} + \sqrt{\log(n_r + n_c)})}{\sqrt{\rho} \sigma_K(\tilde{P}) \sigma_K(\Pi_r) \sqrt{n_{c,K_r}}}\right) = O\left(\frac{\sqrt{K} (\sqrt{\frac{C \max(n_r, n_c)}{\min(n_r, n_c)}} + \sqrt{\log(n_r + n_c)})}{\sqrt{\rho} \sigma_K(\tilde{P}) \sigma_K(\Pi_r) \sqrt{n_{c,\min}}}\right)$$

$$= O\left(\frac{K^{1.5}\left(\sqrt{\frac{C\max(n_r, n_c)}{\min(n_r, n_c)}} + \sqrt{\log(n_r + n_c)}\right)}{\sigma_K(\tilde{P})\sqrt{\rho n_r n_c}}\right),$$

which gives that

$$\max_{i_r \in [n_r]} \|e'_{i_r}(\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O\left(\frac{K^2\left(\sqrt{\frac{C\max(n_r, n_c)}{\min(n_r, n_c)}} + \sqrt{\log(n_r + n_c)}\right)}{\sigma_K(\tilde{P})\sqrt{\rho n_c}}\right).$$

Note that, when $K_r < K_c$, we can not draw a conclusion that $\mu \leq C$. Because, when $K_r < K_c$, the inverse of $B_c B'_c$ does not exist since $B_c \in \mathbb{R}^{K_c \times K_r}$. Therefore, Lemma 8 of Qing and Wang (2021a) does not hold, and we can not obtain the upper bound of $\|U_c\|_{2 \rightarrow \infty}$, causing the impossibility of obtaining the upper bound of μ , and this is the reason that we only consider the case when $K_r = K_c$ for row nodes here.

For column nodes, under conditions of Corollary 7, we have

$$\begin{aligned} \hat{f}_c &= O\left(\frac{K_r K_c \max(n_r, n_c) \log(n_r + n_c)}{\sigma_{K_r}^2(\tilde{P}) \rho \delta_c^2 \sigma_{K_r}^2(\Pi_r) n_{c, K_r} n_{c, \min}}\right) = O\left(\frac{K_r K_c \max(n_r, n_c) \log(n_r + n_c)}{\sigma_{K_r}^2(\tilde{P}) \rho \delta_c^2 (n_r / K_r) (n_c / K_c) (n_c / K_c)}\right) \\ &= O\left(\frac{K_r^2 K_c^3 \max(n_r, n_c) \log(n_r + n_c)}{\sigma_{K_r}^2(\tilde{P}) \rho \delta_c^2 n_r n_c^2}\right). \end{aligned}$$

For the special case $K_r = K_c = K$, since $\frac{n_{c, \max}}{n_{c, \min}} = O(1)$ when $n_{c, \min} = O(\frac{n_c}{K})$, we have

$$\hat{f}_c = O\left(\frac{K^4 \max(n_r, n_c) \log(n_r + n_c)}{\sigma_K^2(\tilde{P}) \rho n_r n_c}\right).$$

When $n_r = O(n)$, $n_c = O(n)$, $K_r = O(1)$ and $K_c = O(1)$, the corollary follows immediately by basic algebra. \blacksquare

Appendix C. Proofs under ODCNM

C.1 Proof of Proposition 9

Proof Since $\Omega = \Pi_r P \Pi'_c \Theta_c = \check{\Pi}_r \check{P} \check{\Pi}'_c \check{\Theta}_c = U_r \Lambda U'_c$, we have $U_r = \Pi_r U_r(\mathcal{I}_r, :) = \check{\Pi}_r U_r(\mathcal{I}_r, :)$ by Lemma 11, which gives that $\Pi_r = \check{\Pi}_r$. Since $U_{c, *} = \Pi_c B_c = \Pi_c U_{c, *}(\mathcal{I}_c, :) = \check{\Pi}_c U_{c, *}(\mathcal{I}_c, :)$ by Lemma 11, we have $\Pi_c = \check{\Pi}_c$. \blacksquare

C.2 Proof of Lemma 11

Proof

- For U_r : since $\Omega = U_r \Lambda U'_c$ and $U'_c U_c = I_{K_r}$, we have $U_r = \Omega U_c \Lambda^{-1}$. Recall that $\Omega = \Pi_r P \Pi'_c \Theta_c$ under ODCNM, we have $U_r = \Pi_r P \Pi'_c \Theta_c U_c \Lambda^{-1} = \Pi_r B_r$, where $B_r = P \Pi'_c \Theta_c U_c \Lambda^{-1}$. Sure, $U_r(i_r, :) = U_r(\bar{i}_r, :)$ holds when $\Pi_r(i_r, :) = \Pi_r(\bar{i}_r, :)$ for $i_r, \bar{i}_r \in [n_r]$.

- For U_c : let D_c be a $K_c \times K_c$ diagonal matrix such that $D_c(k, k) = \frac{\|\Theta_c \Pi_c(:, k)\|_F}{\|\theta_c\|_F}$ for $k \in [K_c]$. Let Γ_c be an $n_c \times K_c$ matrix such that $\Gamma_c(:, k) = \frac{\Theta_c \Pi_c(:, k)}{\|\Theta_c \Pi_c(:, k)\|_F}$ for $k \in [K_c]$. For such D_c and Γ_c , we have $\Gamma_c' \Gamma_c = I_{K_c}$ and $\Omega = \Pi_r P \|\theta_c\|_F D_c \Gamma_c'$, i.e., $\Theta_c \Pi_c = \|\theta_c\|_F \Gamma_c D_c$.

Since $\Omega = U_r \Lambda U_c'$ and $U_r' U_r = I_{K_r}$, we have $U_c = \Theta_c \Pi_c P' \Pi_r' U_r \Lambda^{-1}$. Since $\Theta_c \Pi_c = \|\theta_c\|_F \Gamma_c D_c$, we have $U_c = \Gamma_c \|\theta_c\|_F D_c P' \Pi_r' U_r \Lambda^{-1} = \Gamma_c V_c$, where we set $V_c = \|\theta_c\|_F D_c P' \Pi_r' U_r \Lambda^{-1} \in \mathbb{R}^{K_c \times K_r}$. Note that since $U_c' U_c = I_{K_r} = V_c' \Gamma_c' \Gamma_c V_c = V_c' V_c$, we have $V_c' V_c = I_{K_r}$. Now, for $i_c \in [n_c], k \in [K_r]$, we have

$$\begin{aligned} U_c(i_c, k) &= e_{i_c}' U_c e_k = e_{i_c}' \Gamma_c V_c e_k = \Gamma_c(i_c, :) V_c e_k \\ &= \theta_c(i_c) \left[\frac{\Pi_c(i_c, 1)}{\|\Theta_c \Pi_c(:, 1)\|_F} \quad \frac{\Pi_c(i_c, 2)}{\|\Theta_c \Pi_c(:, 2)\|_F} \quad \cdots \quad \frac{\Pi_c(i_c, K_c)}{\|\Theta_c \Pi_c(:, K_c)\|_F} \right] V_c e_k \\ &= \frac{\theta_c(i_c)}{\|\Theta_c \Pi_c(:, \ell(i_c))\|_F} V_c(\ell(i_c), k), \end{aligned}$$

which gives that

$$U_c(i_c, :) = \frac{\theta_c(i_c)}{\|\Theta_c \Pi_c(:, \ell(i_c))\|_F} [V_c(\ell(i_c), 1) \quad V_c(\ell(i_c), 2) \quad \cdots \quad V_c(\ell(i_c), K_r)] = \frac{\theta_c(i_c)}{\|\Theta_c \Pi_c(:, \ell(i_c))\|_F} V_c(\ell(i_c), :).$$

Then we have

$$U_{c,*}(i_c, :) = \frac{V_c(\ell(i_c), :)}{\|V_c(\ell(i_c), :)\|_F}. \quad (12)$$

Sure, we have $U_{c,*}(i_c, :) = U_{c,*}(\bar{i}_c, :)$ when $\ell(i_c) = \ell(\bar{i}_c)$ for $i_c, \bar{i}_c \in [n_c]$. Let $B_c \in \mathbb{R}^{K_c \times K_r}$ such that $B_c(l, :) = \frac{V_c(l, :)}{\|V_c(l, :)\|_F}$ for $l \in [K_c]$. Eq (12) gives $U_{c,*} = \Pi_c B_c$, which guarantees the existence of B_c .

Now we consider the case when $K_r = K_c = K$. Since $V_c \in \mathbb{R}^{K_c \times K_r}$ and $U_c = \Gamma_c V_c \in \mathbb{R}^{n_c \times K_r}$, we have $V_c \in \mathbb{R}^{K \times K}$ and $\text{rank}(V_c) = K$. Since $V_c' V_c = I_{K_r}$, we have $V_c' V_c = I_K$ when $K_r = K_c = K$. Then we have

$$V_c' V_c = I_K \Rightarrow V_c' V_c V_c' = V_c' \Rightarrow V_c' (V_c V_c' - I_K) = 0 \xrightarrow{\text{rank}(V_c)=K} V_c V_c' = I_K. \quad (13)$$

Since $V_c V_c' = V_c' V_c = I_K$, we have $U_{c,*}(i_c, :) = V_c(\ell(i_c), :)$ by Eq (12), and $\|U_{c,*}(i_c, :) - U_{c,*}(\bar{i}_c, :)\|_F = \|V_c(\ell(i_c), :) - V_c(\ell(\bar{i}_c), :)\|_F = \sqrt{2}$ when $\ell(i_c) \neq \ell(\bar{i}_c)$ for $i_c, \bar{i}_c \in [n_c]$, i.e., $\|B_c(k, :) - B_c(l, :)\|_F = \sqrt{2}$ for $k \neq l \in [K]$.

Note that, when $K_r < K_c$, since $\text{rank}(V_c) = K_r$ and $V_c \in \mathbb{R}^{K_c \times K_r}$, the inverse of V_c does not exist, which causes that the last equality in Eq (13) does not hold and $\|B_c(k, :) - B_c(l, :)\| \neq \sqrt{2}$ for all $k \neq l \in [K_c]$. ■

C.3 Proof of Theorem 14

Proof For row nodes, when conditions in Lemma 13 hold, by Theorem 2 of Qing and Wang (2021a), we have

$$\max_{i_r \in [n_r]} \|e'_{i_r}(\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O(\varpi \kappa(\Pi'_r \Pi_r) K_r \sqrt{\lambda_1(\Pi'_r \Pi_r)}).$$

Next, we focus on column nodes. By the proof of Lemma 2.3 of Qing and Wang (2021b), there exists an orthogonal matrix \hat{O} such that

$$\|\hat{U}_c \hat{O} - U_c\|_F \leq \frac{2\sqrt{2K_r} \|A - \Omega\|}{\sqrt{\lambda_{K_r}(\Omega' \Omega)}}. \quad (14)$$

Under $ODCNM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c, \Theta_c)$, by Lemma 4 of Qing (2021a), we have

$$\sqrt{\lambda_{K_r}(\Omega' \Omega)} \geq \theta_{c, \min} \sigma_{K_r}(P) \sigma_{K_r}(\Pi_r) \sqrt{n_{c, K_r}}. \quad (15)$$

By Lemma 4.2 of Qing (2021a), when Assumption (12) holds, with probability at least $1 - o((n_r + n_c)^{-\alpha})$, we have

$$\|A - \Omega\| = O(\sqrt{\max(\theta_{c, \max} n_r, \|\theta_c\|_1) \log(n_r + n_c)}). \quad (16)$$

Substitute the two bounds in Eqs (15) and (16) into Eq (14), we have

$$\|\hat{U}_c \hat{O} - U_c\|_F \leq C \frac{\sqrt{K_r \max(\theta_{c, \max} n_r, \|\theta_c\|_1) \log(n_r + n_c)}}{\sigma_{K_r}(P) \theta_{c, \min} \sigma_{K_r}(\Pi_r) \sqrt{n_{c, K_r}}}. \quad (17)$$

For $i_c \in [n_c]$, by basic algebra, we have

$$\|\hat{U}_{c, *}(i_c, :) \hat{O} - U_{c, *}(i_c, :)\|_F \leq \frac{2\|\hat{U}_c(i_c, :) \hat{O} - U_c(i_c, :)\|_F}{\|U_c(i_c, :)\|_F}.$$

Set $m_c = \min_{1 \leq i_c \leq n_c} \|U_c(i_c, :)\|_F$, we have

$$\|\hat{U}_{c, *}\hat{O} - U_{c, *}\|_F = \sqrt{\sum_{i_c=1}^{n_c} \|\hat{U}_{c, *}(i_c, :) \hat{O} - U_{c, *}(i_c, :)\|_F^2} \leq \frac{2\|\hat{U}_c \hat{O} - U_c\|_F}{m_c}.$$

Next, we provide lower bounds of m_c . By the proof of Lemma 11, we have

$$\begin{aligned} \|U_c(i_c, :)\|_F &= \left\| \frac{\theta_c(i_c)}{\|\Theta_c \Pi_c(:, \ell(i_c))\|_F} V_c(\ell(i_c), :)\right\|_F = \frac{\theta_c(i_c)}{\|\Theta_c \Pi_c(:, \ell(i_c))\|_F} \|V_c(\ell(i_c), :)\|_F \\ &= \frac{\theta_c(i_c)}{\|\Theta_c \Pi_c(:, \ell(i_c))\|_F} \geq \frac{\theta_{c, \min}}{\theta_{c, \max} \sqrt{n_{c, \max}}} m_{V_c}, \end{aligned}$$

where we set $m_{V_c} = \min_{k \in [K_c]} \|V_c(k, :)\|_F$. Note that when $K_r = K_c = K$, by the proof of Lemma 11, we know that $V_c V_c' = I_K$, which gives that $\|V_c(k, :)\|_F = 1$ for $k \in [K]$, i.e.,

$m_{V_c} = 1$ when $K_r = K_c = K$. However, when $K_r < K_c$, it is challenge to obtain a positive lower bound of m_{V_c} . Hence, we have $\frac{1}{m_c} \leq \frac{\theta_{c,\max}\sqrt{n_{c,\max}}}{\theta_{c,\min}m_{V_c}}$. Then, by Eq (17), we have

$$\|\hat{U}_{c,*}\hat{O} - U_{c,*}\|_F = O\left(\frac{\theta_{c,\max}\sqrt{K_r}\max(\theta_{c,\max}n_r, \|\theta_c\|_1)n_{c,\max}\log(n_r + n_c)}{\sigma_{K_r}(P)\theta_{c,\min}^2 m_{V_c}\sigma_{K_r}(\Pi_r)\sqrt{n_{c,K_r}}}\right).$$

Let $\varsigma > 0$ be a small quantity, by Lemma 2 in Joseph and Yu (2016), if

$$\frac{\sqrt{K_c}}{\varsigma}\|U_{c,*} - \hat{U}_{c,*}\hat{O}\|_F\left(\frac{1}{\sqrt{n_{c,k}}} + \frac{1}{\sqrt{n_{c,l}}}\right) \leq \|B_c(k, \cdot) - B_c(l, \cdot)\|_F, \text{ for each } 1 \leq k \neq l \leq K_c, \quad (18)$$

then the clustering error $\hat{f}_c = O(\varsigma^2)$. Setting $\varsigma = \frac{2}{\delta_c}\sqrt{\frac{K_c}{n_{c,\min}}}\|U_{c,*} - \hat{U}_{c,*}\hat{O}\|_F$ makes Eq (18) hold for all $1 \leq k \neq l \leq K_c$. Then we have $\hat{f}_c = O(\varsigma^2) = O\left(\frac{K_c\|U_{c,*} - \hat{U}_{c,*}\hat{O}\|_F^2}{\delta_c^2 n_{c,\min}}\right)$. By Eq (17), we have

$$\hat{f}_c = O\left(\frac{\theta_{c,\max}^2 K_r K_c \max(\theta_{c,\max}n_r, \|\theta_c\|_1)n_{c,\max}\log(n_r + n_c)}{\sigma_{K_r}^2(P)\theta_{c,\min}^4 \delta_c^2 m_{V_c}^2 \sigma_{K_r}^2(\Pi_r)n_{c,K_r}n_{c,\min}}\right).$$

Especially, when $K_r = K_c = K$, $\delta_c = \sqrt{2}$ under $ODCNM_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c, \Theta_c)$ by Lemma 11, and $m_{V_c} = 1$. When $K_r = K_c = K$, we have

$$\hat{f}_c = O\left(\frac{\theta_{c,\max}^2 K^2 \max(\theta_{c,\max}n_r, \|\theta_c\|_1)n_{c,\max}\log(n_r + n_c)}{\sigma_K^2(P)\theta_{c,\min}^4 \sigma_K^2(\Pi_r)n_{c,\min}^2}\right).$$

■

C.4 Proof of Corollary 15

Proof For row nodes, under conditions of Corollary 15, we have

$$\max_{i_r \in [n_r]} \|e'_{i_r}(\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O(\varpi K_r \sqrt{\frac{n_r}{K_r}}) = O(\varpi \sqrt{K n_r}).$$

Under conditions of Corollary 15, $\kappa(\Omega) = O(1)$ and $\mu \leq C \frac{\theta_{c,\max}^2}{\theta_{c,\min}^2} \leq C$ for some $C > 0$ by Lemma 2 of Qing (2021a). Then, by Lemma 13, we have

$$\begin{aligned} \varpi &= O\left(\frac{\sqrt{\theta_{c,\max}K_r}(\kappa(\Omega)\sqrt{\frac{\max(n_r, n_c)\mu}{\min(n_r, n_c)}} + \sqrt{\log(n_r + n_c)})}{\theta_{c,\min}\sigma_{K_r}(P)\sigma_{K_r}(\Pi_r)\sqrt{n_{c,K_r}}}\right) \\ &= O\left(\frac{\sqrt{\theta_{c,\max}K}(\kappa(\Omega)\sqrt{\frac{\max(n_r, n_c)\mu}{\min(n_r, n_c)}} + \sqrt{\log(n_r + n_c)})}{\theta_{c,\min}\sigma_K(P)\sigma_K(\Pi_r)\sqrt{n_{c,\min}}}\right) \\ &= O\left(\frac{K^{1.5}\sqrt{\theta_{c,\max}}(\sqrt{\frac{C\max(n_r, n_c)}{\min(n_r, n_c)}} + \sqrt{\log(n_r + n_c)})}{\theta_{c,\min}\sigma_K(P)\sqrt{n_r n_c}}\right), \end{aligned}$$

which gives that

$$\max_{i_r \in [n_r]} \|e'_{i_r} (\hat{\Pi}_r - \Pi_r \mathcal{P}_r)\|_1 = O\left(\frac{K^2 \sqrt{\theta_{c,\max}} \left(\sqrt{\frac{C_{\max}(n_r, n_c)}{\min(n_r, n_c)}} + \sqrt{\log(n_r + n_c)}\right)}{\theta_{c,\min} \sigma_K(P) \sqrt{n_c}}\right).$$

The reason that we do not consider the case when $K_r < K_c$ for row nodes is similar as that of Corollary 7, and we omit it here.

For column nodes, under conditions of Corollary 15, we have

$$\begin{aligned} \hat{f}_c &= O\left(\frac{\theta_{c,\max}^2 K_r K_c \max(\theta_{c,\max} n_r, \|\theta_c\|_1) n_{c,\max} \log(n_r + n_c)}{\sigma_{K_r}^2(P) \theta_{c,\min}^4 \delta_c^2 m_{V_c}^2 \sigma_{K_r}^2(\Pi_r) n_{c,K_r} n_{c,\min}}\right) \\ &= O\left(\frac{\theta_{c,\max}^2 K_r^2 K_c^2 \max(\theta_{c,\max} n_r, \|\theta_c\|_1) \log(n_r + n_c)}{\sigma_{K_r}^2(P) \theta_{c,\min}^4 \delta_c^2 m_{V_c}^2 n_r n_c}\right). \end{aligned}$$

For the case $K_r = K_c = K$, we have

$$\begin{aligned} \hat{f}_c &= O\left(\frac{\theta_{c,\max}^2 K^2 \max(\theta_{c,\max} n_r, \|\theta_c\|_1) n_{c,\max} \log(n_r + n_c)}{\sigma_K^2(P) \theta_{c,\min}^4 \sigma_K^2(\Pi_r) n_{c,\min}^2}\right) \\ &= O\left(\frac{\theta_{c,\max}^2 K^4 \max(\theta_{c,\max} n_r, \|\theta_c\|_1) \log(n_r + n_c)}{\sigma_K^2(P) \theta_{c,\min}^4 n_r n_c}\right). \end{aligned}$$

When $n_r = O(n)$, $n_c = O(n)$, $K_r = O(1)$ and $K_c = O(1)$, the corollary follows immediately by basic algebra. \blacksquare

Appendix D. The degree-corrected overlapping and nonoverlapping model

Here, we extend ONM by introducing degree heterogeneities for row nodes with overlapping property in the directed network \mathcal{N} . Let θ_r be an $n_r \times 1$ vector whose i_r -th entry is the degree heterogeneity of row node i_r , for $i_r \in [n_r]$. Let Θ_r be an $n_r \times n_r$ diagonal matrix whose i_r -th diagonal element is $\theta_r(i_r)$. The extended model for generating A is as follows:

$$\Omega := \Theta_r \Pi_r P \Pi_c', \quad A(i_r, i_c) \sim \text{Bernoulli}(\Omega(i_r, i_c)) \quad \text{for } i_r \in [n_r], i_c \in [n_c]. \quad (19)$$

Definition 17 Call model (1), (2), (3), (4), (19) the Degree-Corrected Overlapping and Nonoverlapping model (DCONM) and denote it by $\text{DCONM}_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c, \Theta_r)$.

The following conditions are sufficient for the identifiability of DCONM:

- (II1) $\text{rank}(P) = K_r$, $\text{rank}(\Pi_r) = K_r$, $\text{rank}(\Pi_c) = K_c$, and $P(k, k) = 1$ for $k \in [K_r]$.
- (II2) There is at least one pure row node for each of the K_r row communities.

For degree-corrected overlapping models, it is popular to require that P has unit-“diagonal” elements for model identifiability, see the model identifiability requirements on the DCMM model of Jin et al. (2017) and the OCCAM model of Zhang et al. (2020). Follow similar proof as that of Lemma 3, we have the following lemma.

Lemma 18 Under $D\text{CONM}_{n_r, n_c}(K_r, K_c, P, \Pi_r, \Pi_c, \Theta_r)$, there exist an unique $K_r \times K_r$ matrix B_r and an unique $K_c \times K_r$ matrix B_c such that

- $U_r = \Theta_r \Pi_r B_r$ where $B_r = \Theta_r^{-1}(\mathcal{I}_r, \mathcal{I}_r) U_r(\mathcal{I}_r, :)$.
- $U_c = \Pi_c B_c$. Meanwhile, $U_c(i_c, :) = U_c(\bar{i}_c, :)$ when $\ell(i_c) = \ell(\bar{i}_c)$ for $i_c, \bar{i}_c \in [n_c]$, i.e., U_c has K_c distinct rows. Furthermore, when $K_r = K_c = K$, we have $\|B_c(k, :) - B_c(l, :)\|_F = \sqrt{\frac{1}{n_{c,k}} + \frac{1}{n_{c,l}}}$ for all $1 \leq k < l \leq K$.

The following proposition guarantees the identifiability of $D\text{CONM}$.

Proposition 19 If conditions (II1) and (II2) hold, $D\text{CONM}$ is identifiable: For eligible $(P, \Pi_r, \Pi_c, \Theta_r)$ and $(\check{P}, \check{\Pi}_r, \check{\Pi}_c, \check{\Theta}_r)$, if $\Theta_r \Pi_r P \Pi_c' = \check{\Theta}_r \check{\Pi}_r \check{P} \check{\Pi}_c'$, then $P = \check{P}$, $\Pi_r = \check{\Pi}_r$, $\Pi_c = \check{\Pi}_c$, $\Theta_r = \check{\Theta}_r$.

Proof By Lemma 18, since $U_c = \Pi_c B_c = \Pi_c U_c(\mathcal{I}_c, :) = \check{\Pi}_c U_c(\mathcal{I}_c, :)$, we have $\Pi_c = \check{\Pi}_c$. Since $\Omega(\mathcal{I}_r, \mathcal{I}_c) = \Theta_r(\mathcal{I}_r, \mathcal{I}_r) \Pi_r(\mathcal{I}_r, :) P \Pi_c'(\mathcal{I}_c, :) = \Theta_r(\mathcal{I}_r, \mathcal{I}_r) P = U_r(\mathcal{I}_r, :) \Lambda U_c'(\mathcal{I}_c, :)$, we have $\Theta_r(\mathcal{I}_r, \mathcal{I}_r) = \text{diag}(U_r(\mathcal{I}_r, :) \Lambda U_c'(\mathcal{I}_c, :))$ by the condition that $P(k, k) = 1$ for $k \in [K_r]$. Therefore, we also have $\check{\Theta}_r(\mathcal{I}_r, \mathcal{I}_r) = \text{diag}(U_r(\mathcal{I}_r, :) \Lambda U_c'(\mathcal{I}_c, :))$, which gives that $\Theta_r(\mathcal{I}_r, \mathcal{I}_r) = \check{\Theta}_r(\mathcal{I}_r, \mathcal{I}_r)$. Since $\check{\Theta}_r(\mathcal{I}_r, \mathcal{I}_r) \check{P} = U_r(\mathcal{I}_r, :) \Lambda U_c'(\mathcal{I}_c, :) = \Theta_r(\mathcal{I}_r, \mathcal{I}_r) P$, we have $P = \check{P}$. By Lemma 18, since $U_r = \Theta_r \Pi_r \Theta_r^{-1}(\mathcal{I}_r, \mathcal{I}_r) U_r(\mathcal{I}_r, :) = \check{\Theta}_r \check{\Pi}_r \check{\Theta}_r^{-1}(\mathcal{I}_r, \mathcal{I}_r) U_r(\mathcal{I}_r, :) = \check{\Theta}_r \check{\Pi}_r \check{\Theta}_r^{-1}(\mathcal{I}_r, \mathcal{I}_r) U_r(\mathcal{I}_r, :)$ and $U_r(\mathcal{I}_r, :)$ is a nonsingular matrix, we have $\Theta_r \Pi_r = \check{\Theta}_r \check{\Pi}_r$. Since $\|\Pi_r(i_r, :)\|_1 = \|\check{\Pi}_r(i_r, :)\|_1 = 1$ for $i_r \in [n_r]$, we have $\Pi_r = \check{\Pi}_r$ and $\Theta_r = \check{\Theta}_r$.

Remark 20 (The reason that we do not introduce a model as an extension of ONM by considering degree heterogeneities for both row and column nodes) Suppose we propose an extension model (call it nontrivial-extension-of-ONM, and ne-ONM for short) of ONM such that $\mathbb{E}[A] = \Omega = \Theta_r \Pi_r P \Pi_c' \Theta_c$. For model identifiability, we see that if ne-ONM is identifiable, the following should hold: when $\Omega = \Theta_r \Pi_r P \Pi_c' \Theta_c = \check{\Theta}_r \check{\Pi}_r \check{P} \check{\Pi}_c' \check{\Theta}_c$, we have $\Theta_r = \check{\Theta}_r$, $\Pi_r = \check{\Pi}_r$, $P = \check{P}$, $\Pi_c = \check{\Pi}_c$ and $\Theta_c = \check{\Theta}_c$. Now we check the identifiability of ne-ONM. Follow proof of Lemma 19, since $\Omega(\mathcal{I}_r, \mathcal{I}_c) = \Theta_r(\mathcal{I}_r, \mathcal{I}_r) \Pi_r(\mathcal{I}_r, :) P \Pi_c'(\mathcal{I}_c, :)$ $\Theta_c(\mathcal{I}_c, \mathcal{I}_c) = \Theta_r(\mathcal{I}_r, \mathcal{I}_r) P \Theta_c(\mathcal{I}_c, \mathcal{I}_c) = U_r(\mathcal{I}_r, :) \Lambda U_c'(\mathcal{I}_c, :)$, we have $\Theta_r(\mathcal{I}_r, \mathcal{I}_r) P = U_r(\mathcal{I}_r, :) \Lambda U_c'(\mathcal{I}_c, :) \Theta_c^{-1}(\mathcal{I}_c, \mathcal{I}_c)$. If we assume that $P(k, k) = 1$ for $k \in [K_r]$, we have $\Theta_r(\mathcal{I}_r, \mathcal{I}_r) = \text{diag}(U_r(\mathcal{I}_r, :) \Lambda U_c'(\mathcal{I}_c, :) \Theta_c^{-1}(\mathcal{I}_c, \mathcal{I}_c))$. Similarly, we have $\check{\Theta}_r(\mathcal{I}_r, \mathcal{I}_r) = \text{diag}(U_r(\mathcal{I}_r, :) \Lambda U_c'(\mathcal{I}_c, :) \check{\Theta}_c^{-1}(\mathcal{I}_c, \mathcal{I}_c))$, and it is impossible to guarantee the uniqueness of $\Theta_r(\mathcal{I}_r, \mathcal{I}_r)$ such that $\check{\Theta}_r(\mathcal{I}_r, \mathcal{I}_r)$ unless we further assume that $\Theta_c(\mathcal{I}_c, \mathcal{I}_c)$ is a fixed matrix. However, when we fix $\Theta_c(\mathcal{I}_c, \mathcal{I}_c)$ such that ne-ONM is identifiable, ne-ONM is nontrivial due to the fact $\Theta_c(\mathcal{I}_c, \mathcal{I}_c)$ is fixed. And ne-ONM is trivial only when we set $\Theta_c(\mathcal{I}_c, \mathcal{I}_c) = I_{K_c}$, however, for such ne-ONM when $\Theta_c(\mathcal{I}_c, \mathcal{I}_c) = I_{K_c}$, ne-ONM is $D\text{CONM}$ actually. The above analysis proposes the reason that why we do not extend ONM by considering Θ_r and Θ_c simultaneously.

Follow similar idea as Qing (2021a), we can design spectral algorithm with consistent estimation to fit $D\text{CONM}$. Compared with ONM and ODCNM , the identifiability requirement of $D\text{CONM}$ on P is too strict such that $D\text{CONM}$ only model directed network generated from P with diagonal ‘‘unit’’ elements, and this is the reason we do not provide $D\text{CONM}$ in the main text and propose further algorithmic study as well as theoretical study for it. . ■

References

- Edoardo M. Airoldi, David M. Blei, Stephen E. Fienberg, and Eric P. Xing. Mixed membership stochastic blockmodels. *Journal of Machine Learning Research*, 9:1981–2014, 2008.
- Edoardo M. Airoldi, Xiaopei Wang, and Xiaodong Lin. Multi-way blockmodels for analyzing coordinated high-dimensional responses. *The Annals of Applied Statistics*, 7(4):2431–2457, 2013.
- P. Erdos and A. Rényi. On the evolution of random graphs. pages 38–82, 2011.
- Nicolas Gillis and Stephen A. Vavasis. Semidefinite programming based preconditioning for more robust near-separable nonnegative matrix factorization. *SIAM Journal on Optimization*, 25(1):677–698, 2015.
- Paul W. Holland, Kathryn Blackmond Laskey, and Samuel Leinhardt. Stochastic blockmodels: First steps. *Social Networks*, 5(2):109–137, 1983.
- Jiashun Jin. Fast community detection by SCORE. *Annals of Statistics*, 43(1):57–89, 2015.
- Jiashun Jin, Zheng Tracy Ke, and Shengming Luo. Estimating network memberships by simplex vertex hunting. *arXiv: Methodology*, 2017.
- Antony Joseph and Bin Yu. Impact of regularization on spectral clustering. *Annals of Statistics*, 44(4):1765–1791, 2016.
- Brian Karrer and M. E. J. Newman. Stochastic blockmodels and community structure in networks. *Physical Review E*, 83(1):16107, 2011.
- Jing Lei and Alessandro Rinaldo. Consistency of spectral clustering in stochastic block models. *Annals of Statistics*, 43(1):215–237, 2015.
- Xueyu Mao, Purnamrita Sarkar, and Deepayan Chakrabarti. Overlapping clustering models, and one (class) svm to bind them all. In *Advances in Neural Information Processing Systems*, volume 31, pages 2126–2136, 2018.
- Xueyu Mao, Purnamrita Sarkar, and Deepayan Chakrabarti. Estimating mixed memberships with sharp eigenvector deviations. *Journal of the American Statistical Association*, pages 1–13, 2020.
- Tai Qin and Karl Rohe. Regularized spectral clustering under the degree-corrected stochastic blockmodel. *Advances in Neural Information Processing Systems 26*, pages 3120–3128, 2013.
- Huan Qing. Directed degree corrected mixed membership model and estimating community memberships in directed networks. *arXiv preprint arXiv:2109.07826*, 2021a.
- Huan Qing. A useful criterion on studying consistent estimation in community detection. *arXiv preprint arXiv:2109.14950*, 2021b.

- Huan Qing and Jingli Wang. Directed mixed membership stochastic blockmodel. *arXiv preprint arXiv:2101.02307*, 2021a.
- Huan Qing and Jingli Wang. Consistency of spectral clustering for directed network community detection. *arXiv preprint arXiv:2109.10319*, 2021b.
- Karl Rohe, Sourav Chatterjee, and Bin Yu. Spectral clustering and the high-dimensional stochastic blockmodel. *Annals of Statistics*, 39(4):1878–1915, 2011.
- Karl Rohe, Tai Qin, and Bin Yu. Co-clustering directed graphs to discover asymmetries and directional communities. *Proceedings of the National Academy of Sciences of the United States of America*, 113(45):12679–12684, 2016.
- Zhe. Wang, Yingbin. Liang, and Pengsheng. Ji. Spectral algorithms for community detection in directed networks. *Journal of Machine Learning Research*, 21:1–45, 2020.
- Yuan Zhang, Elizaveta Levina, and Ji Zhu. Detecting overlapping communities in networks using spectral methods. *SIAM Journal on Mathematics of Data Science*, 2(2):265–283, 2020.
- Zhixin Zhou and Arash A.Amini. Analysis of spectral clustering algorithms for community detection: the general bipartite setting. *Journal of Machine Learning Research*, 20(47): 1–47, 2019.