

A sharp isoperimetric inequality in metric measure spaces with non-negative Ricci curvature

Bang-Xian Han*

November 5, 2021

Abstract

We prove a sharp dimension-free isoperimetric inequality, involving the volume entropy, in non-compact metric measure spaces with non-negative synthetic Ricci curvature.

Keywords: isoperimetric inequality, curvature-dimension condition, metric measure space, optimal transport, volume entropy.

Contents

1	Introduction	1
2	Main results	3

1 Introduction

The aim of this note is to present a sharp isoperimetric inequality for a class of metric measure spaces with non-negative Ricci curvature in the sense of Lott-Sturm-Villani.

Let (X, d, \mathbf{m}) be a metric measure space. Recall that the Minkowski content of a Borel set $Z \subset X$ with $\mathbf{m}(Z) < +\infty$ is defined by

$$\mathbf{m}^+(Z) := \liminf_{\epsilon \rightarrow 0} \frac{\mathbf{m}(Z^\epsilon) - \mathbf{m}(Z)}{\epsilon}$$

where $Z^\epsilon \subset X$ is the ϵ -neighbourhood of Z defined by $Z^\epsilon := \{x : d(x, Z) < \epsilon\}$. Isoperimetric inequalities for a class \mathcal{M} of metric measure spaces relate the size of the boundary of sets to their measure. Precisely, there is a function $I_{\mathcal{M}}(v)$, called isoperimetric profile, such that

$$\mathbf{m}^+(\Omega) \geq I_{\mathcal{M}}(v)$$

*Wu Wen-Tsun Key Laboratory of Mathematics and School of Mathematical Sciences, University of Science and Technology of China (USTC), 230026, Hefei, China. Email: hanbangxian@ustc.edu.cn

for all $(X, d, \mathbf{m}) \in \mathcal{M}$ and $\Omega \subset X$ with $\mathbf{m}(\Omega) = v$.

Recently, isoperimetric inequalities in the class of non-compact metric measure spaces with synthetic non-negative Ricci curvature is studied in [1] and [3]. In these isoperimetric inequalities, a key number in the isoperimetric profile is a constant called *asymptotic volume ratio* of a metric measure space (X, d, \mathbf{m}) , defined as

$$\text{AVR}(X, d, \mathbf{m}) := \lim_{r \rightarrow +\infty} \frac{\mathbf{m}(B_r(x_0))}{r^N} \in [0, +\infty].$$

When $\text{AVR}(X, d, \mathbf{m}) > 0$, we call that the space (X, d, \mathbf{m}) has Euclidean volume growth.

Since the dimension upper bound N appear in the asymptotic volume ratio, the isoperimetric inequalities obtained in [1] and [3] are all dimension-dependent. So it is natural to find a dimension-free version of the isoperimetric inequality for metric measure spaces with non-negative Ricci curvature. Firstly, recall that a space has non-negative Ricci curvature (without dimension restriction) in the sense of Lott-Sturm-Villani means:

Definition 1.1 (Lott-Sturm-Villani's curvature-dimension condition, cf. [5, 7]). We say that a metric measure space (X, d, \mathbf{m}) has non-negative Ricci curvature, or satisfies $\text{CD}(0, \infty)$ condition, if the entropy functional $\text{Ent}_{\mathbf{m}}$ is displacement convex on the L^2 -Wasserstein space $(\mathcal{P}_2(X), W_2)$. This means, for any two probability measures $\mu_0, \mu_1 \in \mathcal{P}_2(X)$ with $\mu_0, \mu_1 \ll \mathbf{m}$, there is a L^2 -Wasserstein geodesic $(\mu_t)_{t \in [0, 1]}$ such that

$$\text{Ent}_{\mathbf{m}}(\mu_t) \leq t \text{Ent}_{\mathbf{m}}(\mu_1) + (1 - t) \text{Ent}_{\mathbf{m}}(\mu_0) \quad (1.1)$$

where $\text{Ent}_{\mathbf{m}}(\mu_t)$ is defined as $\int \rho_t \ln \rho_t d\mathbf{m}$ if $\mu_t = \rho_t \mathbf{m}$, otherwise $\text{Ent}_{\mathbf{m}}(\mu_t) = +\infty$.

In the study of metric (Riemannian) geometry, there are some important spaces where the asymptotic volume ratio $\text{AVR} = +\infty$. In these cases, we often consider instead the *volume entropy*, which is an important concept in both Riemannian geometry (cf. [2]) and dynamical system (cf. [6]).

Definition 1.2 (Volume entropy). A metric measure space (X, d, \mathbf{m}) admits the *volume entropy* at $x_0 \in X$, denoted by $h(X, d, \mathbf{m})(x_0)$, provided

$$h(X, d, \mathbf{m})(x_0) := \lim_{r \rightarrow +\infty} \frac{\ln \mathbf{m}(B_r(x_0))}{r} \in [0, \infty].$$

It is not hard to prove that a metric measure space with non-negative Ricci curvature in the sense of Definition 1.1, surely admits the volume entropy which is independent of the choice of x_0

The main result of this note is the following sharp isoperimetric inequality involving volume entropy:

Theorem 1.3 (Sharp isoperimetric inequality). *Let (X, d, \mathbf{m}) be a metric measure space with non-negative synthetic Ricci curvature. Then for any $\Omega \subset X$ with finite measure, it holds the following isoperimetric inequality*

$$\mathbf{m}^+(\Omega) \geq \mathbf{m}(\Omega) h(X, d, \mathbf{m}). \quad (1.2)$$

Moreover, the constant $h(X, d, \mathbf{m})$ in (1.2) can not be replaced by any larger ones.

2 Main results

In the following two theorems we will prove a sharp isoperimetric inequality for the class of metric measure spaces with non-negative synthetic Ricci curvature.

Theorem 2.1 (Dimension-free isoperimetric inequality). *Let (X, d, \mathbf{m}) be a metric measure space with non-negative Ricci curvature. Then for any $\Omega \subset X$ with finite measure, it holds the following isoperimetric inequality*

$$\mathbf{m}^+(\Omega) \geq \mathbf{m}(\Omega)h(X, d, \mathbf{m}). \quad (2.1)$$

Proof. Given $x_0 \in X$ and $R > 0$. Define $\mu_0 = \frac{1}{\mathbf{m}(\Omega)}\mathbf{m}|_{\Omega}$ and $\mu_1 = \frac{1}{\mathbf{m}(B_R(x_0))}\mathbf{m}|_{B_R(x_0)}$. By 1.1 there is an L^2 -Wasserstein geodesic (μ_t) connecting μ_0, μ_1 such that

$$\text{Ent}_{\mathbf{m}}(\mu_t) \leq t\text{Ent}_{\mathbf{m}}(\mu_1) + (1-t)\text{Ent}_{\mathbf{m}}(\mu_0) \quad (2.2)$$

Denote the support of μ_t by $\text{supp } \mu_t$ and the set of t -intermediate points

$$Z_t := \left\{ z : \exists x \in \Omega, y \in B_R(x_0), \text{ such that } \frac{d(z, x)}{t} = \frac{d(z, y)}{1-t} = d(x, y) \right\}.$$

It can be seen that $\text{supp } \mu_t \subset Z_t$. Then by (2.2) and Jensen's inequality, we have

$$-\ln(\mathbf{m}(Z_t)) \leq -\ln(\mathbf{m}(\text{supp } \mu_t)) \leq -t \ln(\mathbf{m}(B_R(x_0))) - (1-t) \ln(\mathbf{m}(\Omega)). \quad (2.3)$$

Let $\epsilon := t(\text{diam}(\Omega) + R)$. It is not hard to see that $Z_t \subset \Omega^\epsilon$. If $\mathbf{m}^+(\Omega) = +\infty$, there is nothing to prove. Otherwise, we have $\lim_{\epsilon \rightarrow 0} \mathbf{m}(\Omega^\epsilon) = \mathbf{m}(\Omega)$. Then we have

$$\begin{aligned} \frac{\mathbf{m}^+(\Omega)}{\mathbf{m}(\Omega)} &= \liminf_{\epsilon \rightarrow 0} \frac{1}{\mathbf{m}(\Omega)} \frac{\mathbf{m}(\Omega^\epsilon) - \mathbf{m}(\Omega)}{\epsilon} \\ &= \liminf_{\epsilon \rightarrow 0} \frac{\ln(\mathbf{m}(\Omega^\epsilon)) - \ln(\mathbf{m}(\Omega))}{\mathbf{m}(\Omega^\epsilon) - \mathbf{m}(\Omega)} \frac{\mathbf{m}(\Omega^\epsilon) - \mathbf{m}(\Omega)}{\epsilon} \\ &\geq \liminf_{t \rightarrow 0} \frac{\ln(\mathbf{m}(Z_t)) - \ln(\mathbf{m}(\Omega))}{t(\text{diam}(\Omega) + R)} \\ \text{By (2.3)} \quad &\geq \liminf_{t \rightarrow 0} \frac{t \ln(\mathbf{m}(B_R(x_0))) + (1-t) \ln(\mathbf{m}(\Omega)) - \ln(\mathbf{m}(\Omega))}{t(\text{diam}(\Omega) + R)} \\ &= \frac{\ln(\mathbf{m}(B_R(x_0))) - \ln(\mathbf{m}(\Omega))}{\text{diam}(\Omega) + R}. \end{aligned}$$

Letting $R \rightarrow \infty$, by Definition 1.2, we get

$$\frac{\mathbf{m}^+(\Omega)}{\mathbf{m}(\Omega)} \geq h(X, d, \mathbf{m})$$

which is the thesis. □

Theorem 2.2 (Sharp inequality). *The inequality (2.1) in Theorem 2.1 is sharp. This means, for any $C > h(X, d, \mathbf{m})$, the inequality $\mathbf{m}^+(\Omega) \geq C\mathbf{m}(\Omega)$ does not hold for any Borel set $\Omega \subset X$.*

Proof. We will prove the theorem by contradiction. Assume there is a constant $C > h(X, d, \mathbf{m})$, such that

$$\mathbf{m}^+(\Omega) \geq C\mathbf{m}(\Omega) \quad (2.4)$$

for any Borel set $\Omega \subset X$.

Apply (2.3) with $\Omega = B_{r+\delta}(x_0)$ and $R = \epsilon$. We get the following interpolation inequality

$$\ln(\mathbf{m}(Z_t)) \geq t \ln(\mathbf{m}(B_\epsilon(x_0))) + (1-t) \ln(\mathbf{m}(B_{r+\delta}(x_0))) \quad \forall t \in [0, 1]. \quad (2.5)$$

For any $z \in Z_t$, we have $d(z, x_0) \leq (1-t)(r+\delta) + \epsilon$. Choosing $t = \frac{\delta+\epsilon}{r+\delta}$, we have $Z_t \subset B_r(x_0)$. Thus (2.5) implies

$$\ln(\mathbf{m}(B_r(x_0))) \geq \frac{\delta+\epsilon}{r+\delta} \ln(\mathbf{m}(B_\epsilon(x_0))) + \frac{r-\epsilon}{r+\delta} \ln(\mathbf{m}(B_{r+\delta}(x_0))).$$

Then

$$\begin{aligned} & \frac{r-\epsilon}{r+\delta} \left(\frac{\ln(\mathbf{m}(B_{r+\delta}(x_0))) - \ln(\mathbf{m}(B_r(x_0)))}{\delta} \right) \\ & \leq \frac{\delta+\epsilon}{\delta(r+\delta)} \left(\ln(\mathbf{m}(B_r(x_0))) - \ln(\mathbf{m}(B_\epsilon(x_0))) \right). \end{aligned}$$

From the proof of Theorem 2.1, we can see that $\frac{\ln(\mathbf{m}(B_{r+\delta}(x_0))) - \ln(\mathbf{m}(B_r(x_0)))}{\delta} \geq C$, so

$$\frac{r-\epsilon}{r+\delta} C \leq \frac{\delta+\epsilon}{\delta(r+\delta)} \left(\ln(\mathbf{m}(B_r(x_0))) - \ln(\mathbf{m}(B_\epsilon(x_0))) \right).$$

Letting $r \rightarrow \infty$, we get

$$C \leq \frac{\delta+\epsilon}{\delta} h(X, d, \mathbf{m}).$$

Letting $\epsilon \rightarrow 0$ we get the contradiction. \square

For metric measure spaces satisfying a generalized Bishop-Gromov inequality, such as $CD(0, N)$ spaces or $MCP(0, N)$ spaces with $N < +\infty$, whose volume entropy vanish, it is known that there is no isometric inequality in the form of (2.1) with a positive C . More generally, as a direct consequence of Theorem 2.1 and Theorem 2.2, we have the following result.

Corollary 2.3. *Let (X, d, \mathbf{m}) be a metric measure space with non-negative Ricci curvature $h(X, d, \mathbf{m}) = 0$, then there is no isoperimetric inequality in the form of*

$$\mathbf{m}^+(\Omega) \geq C\mathbf{m}(\Omega) \quad \forall \Omega \subset X$$

from some $C > 0$.

Example 2.4. Consider the 1-dimensional metric measure space $(\mathbb{R}, |\cdot|, e^t \mathcal{L}^1)$. It can be seen that this is a $\text{RCD}(0, \infty)$ space and its volume entropy is 1. Take $\Omega = (-\infty, 0]$, we have $(e^t \mathcal{L}^1)(\Omega) = 1$ and $(e^t \mathcal{L}^1)^+(\Omega) = 1$, so that the equality holds in (2.1).

Inspired by the example above, we propose the following conjecture, which will be studied in a forthcoming paper.

Conjecture 2.5 (Rigidity). *Let (X, d, \mathbf{m}) be a $\text{RCD}(0, \infty)$ metric measure space with $h(X, d, \mathbf{m}) > 0$. Assume there is $\Omega \subset X$ with $\mathbf{m}(\Omega) < +\infty$ such that*

$$\mathbf{m}^+(\Omega) = h(X, d, \mathbf{m})\mathbf{m}(\Omega) > 0.$$

Then

$$(X, d, \mathbf{m}) \cong \left(\mathbb{R}, |\cdot|, e^t dt \right) \times (Y, d_Y, \mathbf{m}_Y)$$

for some $\text{RCD}(0, \infty)$ space (Y, d_Y, \mathbf{m}_Y) with $\mathbf{m}_Y(Y) < +\infty$, and up to change of variables

$$\Omega = (-\infty, e] \times Y \subset \mathbb{R} \times Y$$

where $e \in \mathbb{R}$ satisfies $\int_{-\infty}^e e^s ds = \mathbf{m}(\Omega)$.

References

- [1] Z. M. BALOGH AND A. KRISTÁLY, *Sharp geometric inequalities in spaces with nonnegative Ricci curvature and Euclidean volume growth*. Preprint, arXiv: 2012.11862, 2021.
- [2] G. BESSON, G. COURTOIS, AND S. GALLOT, *Entropies et rigidités des espaces localement symétriques de courbure strictement négative*, *Geom. Funct. Anal.*, 5 (1995), pp. 731–799.
- [3] F. CAVALLETTI AND D. MANINI, *Isoperimetric inequality in noncompact MCP spaces*. Preprint, arXiv: 2110.07528, 2021.
- [4] C. CONNELL, X. DAI, J. NÚÑEZ-ZIMBRÓN, R. PERALES, P. SUÁREZ-SERRATO, AND G. WEI, *Maximal volume entropy rigidity for $\text{RCD}^*(-(N-1), N)$ spaces*, *J. Lond. Math. Soc.*, (2021).
- [5] J. LOTT AND C. VILLANI, *Ricci curvature for metric-measure spaces via optimal transport*, *Ann. of Math. (2)*, 169 (2009), pp. 903–991.
- [6] A. MANNING, *Topological entropy for geodesic flows*, *Ann. of Math. (2)*, 110 (1979), pp. 567–573.
- [7] K.-T. STURM, *On the geometry of metric measure spaces. I*, *Acta Math.*, 196 (2006), pp. 65–131.