

QUASILINEAR WAVE EQUATIONS ON SCHWARZSCHILD–DE SITTER

GEORGIOS MAVROGIANNIS

*University of Cambridge, Department of Pure Mathematics and Mathematical Statistics,
Wilberforce Road, Cambridge CB3 0WB, United Kingdom*

ABSTRACT. We give an elementary new argument for global existence and exponential decay of solutions of quasilinear wave equations on Schwarzschild–de Sitter black hole backgrounds, for appropriately small initial data. The core of the argument is entirely local, based on time translation invariant energy estimates in spacetime slabs of fixed time length. Global existence then follows simply by iterating this local result in consecutive spacetime slabs. We infer that an appropriate future energy flux decays exponentially with respect to the energy flux of the initial data.

CONTENTS

1. Introduction	1
2. Preliminaries and notation	5
3. Metric close to Schwarzschild–de Sitter and the Local well-posedness result	10
4. The main Theorems	11
5. Energy estimates on a spacetime slab of fixed time length	13
6. Proof of Theorem 1	18
7. Proof of Theorem 2	20
8. The semilinear equation on Schwarzschild–de Sitter	22
Appendix A. Proof of the Cauchy stability result of Proposition 3.1	27
References	39

1. INTRODUCTION

We revisit the problem of global existence of solutions of quasilinear wave equations, of the form

$$(1.1) \quad \square_{g(\nabla\psi)}\psi = \partial\psi \cdot \partial\psi,$$

with $g(\nabla\psi) = g_{M,\Lambda} + h(\nabla\psi)$, where $g_{M,\Lambda}$ is the metric of the Schwarzschild–de Sitter black hole spacetime, also $\partial\psi \cdot \partial\psi = a^{ij}\partial_i\psi\partial_j\psi$, where a, h are sufficiently regular tensors, with $h(0) = 0$. Specifically, we are interested in a spacetime region that is slightly larger than that enclosed by the event and cosmological horizons, respectively \mathcal{H}^+ , $\bar{\mathcal{H}}^+$, see the shaded region of Figure 1.

The problem of stability of quasilinear wave equations on such backgrounds has been extensively studied by Hintz and Vasy, see [20, 19, 21, 22], where they arrived at global stability results. Their papers appeal to machinery from microlocal analysis and Nash Moser iteration arguments. These results were preceded by a long list of results on the linear problem, see [39, 40, 2, 11, 8, 9, 32, 15, 14]. Moreover, note the recent remarkable global non-linear stability proof for the slowly rotating Kerr–de Sitter black hole as a solution of the Einstein vacuum equation with $\Lambda > 0$, by Hintz–Vasy [23],

E-mail address: gm615@cam.ac.uk.

Date: November 19, 2021.

based in part on the above works. (For some results on the cosmological region see [36, 35]). Note that the non-linear stability of the pure de Sitter spacetime has been obtained previously by Friedrich [17].

Schwarzschild–de Sitter can be thought of as the $\Lambda > 0$ analogue of the Schwarzschild and Kerr spacetimes, which are celebrated solutions of the vacuum Einstein equation with $\Lambda = 0$. For the study of linear equations on the latter see for instance [10, 38] and the definitive [13]. These results have been used to prove non-linear stability results for equations of type (1.1), see [29, 28]. In general, these non-linear problems are more difficult than the $\Lambda > 0$ case, because the expected decay is only polynomial and one has to assume and exploit suitable null structure (see [26]) for the non-linearities. For results on stability of black hole spacetimes with $\Lambda = 0$ see [6, 37, 18, 1, 27, 7, 5].

In principle, one approach to the study of (1.1) on Schwarzschild or Kerr–de Sitter backgrounds would be to directly adapt the methods from the $\Lambda = 0$ case. Such an approach, however, would not fully exploit the aspects that make the $\Lambda > 0$ problem easier.

The purpose of this paper is to introduce a physical space approach to (1.1) which is well tailored to this setting. Our approach is based entirely on local in time translation invariant energy estimates (Theorem 1) and an iteration argument in consecutive spacetime regions (Theorem 2). This approach will use the results of our accompanying physical space linear paper [30], where we utilized a physical space commutation with a vector field \mathcal{G} , see already (1.6), and proved a *relatively non-degenerate estimate*, using also a Morawetz estimate proved in [9]. Note that Holzegel–Kauffman originally introduced the analogue of the \mathcal{G} vector field in the $\Lambda = 0$ case, see [24]. Our physical space commutation with \mathcal{G} , in the high frequency limit, connects with the work of previous authors on ‘lossless estimates’ and ‘non-trapping estimates’, e.g. see [34, 3, 4, 25, 33, 19, 16].

We will present the rough version of our Theorems, for which the reader may wish to refer to Figure 1 for the Schwarzschild–de Sitter spacetime.




FIGURE 1. The Schwarzschild–de Sitter spacetime

We denote as

$$(1.2) \quad \mathcal{M}_\delta$$

the ‘extended’ exterior region, also see the dark shaded region of Figure 1, where δ is a smallness parameter that parametrises how far the boundaries of \mathcal{M}_δ (which we denote as \mathcal{H}_δ^+ , $\bar{\mathcal{H}}_\delta^+$) are from the event horizon \mathcal{H}^+ and the cosmological horizon $\bar{\mathcal{H}}^+$ respectively. We choose δ in the proof of our Theorem 2 in Section 7. For the precise definition of \mathcal{M}_δ see already Definition 2.1. The spacetime

domain \mathcal{M}_δ , see the dark shaded region of Figure 1, is foliated by the spacelike hypersurfaces

$$(1.3) \quad \{\bar{t} = \tau\},$$

with $\tau \geq 0$, see Figure 1, where

$$(1.4) \quad (\bar{t}, r, \theta, \varphi)$$

are appropriate (non-standard) hyperboloidal coordinates in which the metric takes the form (2.6). The coordinate vector field $\partial_{\bar{t}}$ is Killing. We denote by Ω_α , $\alpha = 1, 2, 3$ the generators of the Lie algebra $so(3)$ associated with the (θ, φ) spheres.

We will consider two types of energies.

The first energy is a non-degenerate energy which we define as

$$(1.5) \quad E_j[\psi](\tau) = \sum_{1 \leq i_1 + i_2 + i_3 \leq j} \sum_{\alpha=1,2,3} \int_{\{\bar{t}=\tau\}} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_\alpha^{i_3} \psi)^2$$

for all $j \geq 1$, with respect to the induced volume form of $g_{M,\Lambda}$ on the $\{\bar{t} = \tau\}$ hypersurface.

Now, we discuss the second energy that we will consider. The first two terms of the second energy, see (1.7), are associated with the $C^{0,1}$ vector field \mathcal{G} of [30] (see the previous [24] for $\Lambda = 0$), which is defined as

$$(1.6) \quad \mathcal{G} \doteq \begin{cases} r \sqrt{1 - \frac{2M}{r} - \frac{\Lambda}{3} r^2} \partial_r, & 1 - \frac{2M}{r} - \frac{\Lambda}{3} r^2 > 0 \\ 0, & 1 - \frac{2M}{r} - \frac{\Lambda}{3} r^2 \leq 0 \end{cases}$$

in appropriate hyperboloidal coordinates $(\bar{t}, r, \theta, \varphi)$, see already Definition 2.4. We define the second energy as

$$(1.7) \quad \begin{aligned} E_{\mathcal{G},j}[\psi](\tau) = & \int_{\{\bar{t}=\tau\}} \sum_{1 \leq i_1 + i_2 + i_3 \leq j-1, i_3 \geq 1} \sum_{\alpha=1,2,3} \left(1 - \frac{2M}{r} - \frac{\Lambda}{3} r^2\right)^{2i_3-1} (\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \partial_r^{i_3} \mathcal{G}\psi)^2 \\ & + \sum_{1 \leq i_1 + i_2 \leq j-1} \sum_{\alpha=1,2,3} (\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \mathcal{G}\psi)^2 \\ & + \sum_{1 \leq i_1 + i_2 + i_3 \leq j-1} \sum_{\alpha=1,2,3} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_\alpha^{i_3} \psi)^2, \end{aligned}$$

for all $j \geq 2$, with respect to the induced volume form of $g_{M,\Lambda}$ on the $\{\bar{t} = \tau\}$ hypersurface. Note that the highest order integrands of the energies in (1.7) are identically zero where $\mathcal{G} \equiv 0$.

Note the inequality

$$(1.8) \quad E_{j-1}[\psi](\tau) \leq E_{\mathcal{G},j}[\psi](\tau) \lesssim E_j[\psi](\tau),$$

also see already Remark 2.5. For the formal definition of the above energies see Definition 2.5.

The rough version of our main Theorem 1 is the following. (See the dark shaded region of Figure 2 for the local spacetime slab we will consider.)

Theorem 1 (rough version). *Let $k \geq 7$. There exists a constant $C > 0$, depending on k, M, Λ and on the tensors a, h , see (1.1), such that, for $L > 0$ an arbitrary positive number, the following holds.*

There exists a $\tau_{step}(L) > 0$ sufficiently large and there exist

$$(1.9) \quad \delta = \delta(\tau_{step}) > 0, \quad \epsilon = \epsilon(\tau_{step}, \delta) > 0$$

sufficiently small such that for all $\tau_1 \geq 0$ and

$$(1.10) \quad \tau_2 = \tau_1 + \tau_{step}$$

if we take initial data for (1.1) on $\{\tau = \tau_1\}$ with

$$(1.11) \quad E_{k+1}[\psi](\tau_1) \leq \epsilon$$




FIGURE 2. A space-time slab of finite time length τ_{step}

then there exists a unique solution to the quasilinear wave equation (1.1) on $\mathcal{M}_\delta \cap \{\tau_1 \leq \tau \leq \tau_2\}$ and the following estimates are satisfied

$$(1.12) \quad E_{\mathcal{G},k}[\psi](\tau_2) \leq \frac{1}{L} E_{\mathcal{G},k}[\psi](\tau_1),$$

and

$$(1.13) \quad \begin{aligned} E_{\mathcal{G},k}[\psi](\tau') &\leq C E_{\mathcal{G},k}[\psi](\tau_1), \\ E_{k+1}[\psi](\tau') &\leq C E_{k+1}[\psi](\tau_1), \\ E_{k+2}[\psi](\tau') &\leq C E_{k+2}[\psi](\tau_1), \end{aligned}$$

for all $\tau' \in [\tau_1, \tau_1 + \tau_{step}]$.

Finally,

$$(1.14) \quad E_{k+1}[\psi](\tau_2) \leq C e^{-\frac{1}{3} \log(L) + \frac{2}{3} \log C} (E_{\mathcal{G},k}[\psi](\tau_1))^{1/3} (E_{k+2}[\psi](\tau_1))^{2/3}.$$

It is instructive to compare the results of Theorem 1 with the main results of our linear theory [30] on a Schwarzschild–de Sitter background. Specifically, the estimate of inequality (1.12) and the first inequality of (1.13) corresponds to the estimate in linear theory that one obtains from the commutation with the \mathcal{G} vector field, see already Theorem 2.1. The remaining inequalities of (1.13) correspond to the uniform boundedness results of [9].

We use the result of Theorem 1, for a well chosen L so that the constant in (1.14) is sufficiently small and, by a completely elementary iteration argument on consecutive spacetime regions, see Figure 2 for such a region, we prove that the solution of the quasilinear wave equation (1.1) exists globally and decays exponentially. The rough version of our main Theorem 2 is the following.

Theorem 2 (rough version). *Let $k \geq 7$. Then, there exist constants $c_d, c_g > 0$ and there exists a $\delta > 0$ and an $\epsilon > 0$ sufficiently small, such that if the initial energy satisfies*

$$(1.15) \quad E_{k+2}[\psi](0) \leq \epsilon,$$

on $\Sigma = \{\bar{t} = 0\}$ then the solution exists globally on \mathcal{M}_δ and the energy $E_{\mathcal{G},k}[\psi]$ decays exponentially

$$(1.16) \quad E_{\mathcal{G},k}[\psi](\tau) \lesssim e^{-c_d \tau} E_{\mathcal{G},k}[\psi](0).$$

We note that the classical Sobolev energies $E_{k-1}[\psi](\tau), E_{k+1}[\psi](\tau)$ decay exponentially

$$(1.17) \quad E_{k-1}[\psi](\tau) \lesssim e^{-c_d \tau} E_{\mathcal{G},k}[\psi](0), \quad E_{k+1}[\psi](\tau) \lesssim e^{-c_d \tau} E_{k+2}[\psi](0)$$

while for the top order energy we only have

$$(1.18) \quad E_{k+2}[\psi](\tau) \lesssim e^{c_g \tau} E_{k+2}[\psi](0).$$

Remark 1.1. Note that the growth constant c_g of (1.18) of Theorem 2 can in fact be made arbitrarily small, restricting to sufficiently small $\epsilon > 0$. See already Remark 4.2.

Remark 1.2. In our Theorem 2, we improve slightly on the regularity assumption of initial data of [22]. Specifically, we only require the initial data to be in the Sobolev space H^9 , see our main Theorems 1, 2 and Remark 4.1.

Remark 1.3. In Section 8, we present Theorems 1' and 2' which give global non-linear stability in the semilinear case, with the tensor $h \equiv 0$, under weaker assumptions than those of Theorem 2. Specifically, we require the initial data only to be in the Sobolev space H^7 .

Remark 1.4. Note that by using the results of our forthcoming [31] on Kerr–de Sitter, the global stability results of the present paper generalize to the slowly rotating Kerr–de Sitter case.

1.1. Acknowledgments. I would like to thank my supervisor Mihalis Dafermos, for his continuous support, for suggesting that the results of [30] may also treat the nonlinear stability problems of the present paper and for carefully reading previous versions of this paper. The author would also like to thank Christoph Kehle for valuable discussions and useful comments.

2. PRELIMINARIES AND NOTATION

2.1. The manifolds, metrics and spacetime domains. The definitions and notation of this Section have already been introduced in our [30].

Fix $M, \Lambda > 0$ such that

$$(2.1) \quad r_+ < \bar{r}_+$$

are the two positive real roots of

$$(2.2) \quad 1 - \mu \doteq 1 - \frac{2M}{r} - \frac{\Lambda}{3} r^2.$$

We need the following definition

Definition 2.1. For $\tau_2 \geq \tau_1 \geq 0$ we define

$$(2.3) \quad D(\tau_1, \tau_2) = [\tau_1, \tau_2]_{\bar{t}} \times [r_+, \bar{r}_+]_r \times \mathbb{S}_{(\theta, \varphi)}^2.$$

We also define, for $\delta > 0$,

$$(2.4) \quad D_\delta(\tau_1, \tau_2) = [\tau_1, \tau_2]_{\bar{t}} \times [r_+ - \delta, \bar{r}_+ + \delta]_r \times \mathbb{S}_{(\theta, \varphi)}^2$$

and

$$(2.5) \quad \mathcal{M} = D(0, \infty) \doteq [0, \infty]_{\bar{t}} \times [r_+, \bar{r}_+]_r \times \mathbb{S}_{(\theta, \varphi)}^2, \quad \mathcal{M}_\delta = D_\delta(0, \infty) \doteq [0, \infty]_{\bar{t}} \times [r_+ - \delta, \bar{r}_+ + \delta]_r \times \mathbb{S}_{(\theta, \varphi)}^2.$$

We refer to the coordinates $(\bar{t}, r, \theta, \varphi)$ as regular hyperboloidal coordinates.

We need the following definition

Definition 2.2. We denote as $(\mathcal{M}_\delta, g_{M, \Lambda})$ the Schwarzschild–de Sitter spacetime, with metric

$$(2.6) \quad g_{M, \Lambda} = - \left(1 - \frac{2M}{r} - \frac{\Lambda}{3} r^2 \right) (d\bar{t})^2 - 2 \frac{1 - \frac{3M}{r}}{\sqrt{1 - 9M^2 \Lambda}} \sqrt{1 + \frac{6M}{r}} d\bar{t} dr + \frac{27M^2}{1 - 9M^2 \Lambda} \frac{1}{r^2} (dr)^2 + r^2 d\sigma_{\mathbb{S}^2},$$

in regular hyperboloidal coordinates $(\bar{t}, r, \theta, \varphi)$, where $d\sigma_{\mathbb{S}^2} = d\theta^2 + \sin^2 \theta d\varphi^2$ is the standard metric of the unit sphere \mathbb{S}^2 . We will often denote the metric $g_{M, \Lambda}$ as

$$(2.7) \quad \mathring{g}.$$

Note the following remarks

Remark 2.1. For a sufficiently small $\delta > 0$, note that the hypersurfaces

$$(2.8) \quad \{\bar{t} = \tau\}$$

are spacelike in \mathcal{M}_δ , with respect to the metric of the following Definition 2.2.

Remark 2.2. Note that δ will be fixed in the proof of Theorem 2 in Section 7. In what follows δ will always be assumed sufficiently small.

Remark 2.3. The inverse metric components of the metric of Definition 2.2 are

$$(2.9) \quad \mathring{g}^{rr} = (1 - \mu), \quad \mathring{g}^{\bar{t}\bar{t}} = -\frac{1 - \xi^2(r)}{1 - \mu}, \quad \mathring{g}^{\bar{t}r} = -\xi(r).$$

for

$$(2.10) \quad \xi(r) = \frac{1 - \frac{3M}{r}}{\sqrt{1 - 9M^2\Lambda}} \sqrt{1 + \frac{6M}{r}}.$$

Remark 2.4. Note the usual expression of the Schwarzschild–de Sitter metric is in coordinates $(t, r, \theta, \varphi) \in \mathbb{R}_t \times (r_+, \bar{r}_+)_r \times \mathbb{S}_{(\theta, \varphi)}^2$, which reads

$$(2.11) \quad \mathring{g} = -(1 - \mu)(dt)^2 + (1 - \mu)^{-1}(dr)^2 + r^2 d\sigma_{\mathbb{S}^2},$$

where note the relation

$$(2.12) \quad \bar{t} = t + H(r), \quad H(r) = \int_{3M}^r \frac{\xi(r)}{1 - \mu} dr.$$

We denote as

$$(2.13) \quad \nabla$$

the covariant derivative of the Riemannian metric $r^2 d\sigma_{\mathbb{S}^2}$. Moreover, we denote the standard generators of the $so(3)$ Lie Algebra

$$(2.14) \quad \Omega_\alpha, \quad \alpha = 1, 2, 3,$$

associated with the (θ, φ) spheres. These can be thought of as vector fields on \mathcal{M}_δ .

Note that

$$(2.15) \quad |\nabla f|^2 \sim \sum_\alpha \frac{1}{r^2} |\Omega_\alpha f|^2$$

for any sufficiently regular f , where the constants in the above only depend on the black hole parameters. (We have included the inessential r factor above for comparison with the asymptotically flat case.)

2.2. The horizons and auxilliary spacelike hypersurfaces. We need the following definition

Definition 2.3. The event and cosmological horizons are defined as

$$(2.16) \quad \mathcal{H}^+ = \{r = r_+\}, \quad \bar{\mathcal{H}}^+ = \{r = \bar{r}_+\}$$

also see [30, 31]. These hypersurfaces are null with respect to the metric $g_{M,\Lambda}$.

For sufficiently small $\delta > 0$, we define the following spacelike hypersurfaces

$$(2.17) \quad \mathcal{H}_\delta^+ = \{r = r_+ - \delta, \bar{t} \geq 0\}, \quad \bar{\mathcal{H}}_\delta^+ = \{r = \bar{r}_+ + \delta, \bar{t} \geq 0\},$$

in the coordinates $(\bar{t}, r, \theta, \varphi)$ of Definition 2.2.

2.3. Volume forms and normals for the metric $g_{M,\Lambda}$. We denote the spacetime volume form of \mathring{g} as

$$(2.18) \quad d\mathring{g} = r^2 \sin \theta d\bar{t} dr d\theta d\varphi$$

with respect to the $(\bar{t}, r, \theta, \varphi)$ coordinates.

By pulling back the spacetime volume form (2.18) into hypersurfaces of constant \bar{t} , we obtain that the $\{\bar{t} = \tau\}$ hypersurfaces admit the volume form

$$(2.19) \quad d\mathring{g}_{\{\bar{t}=\tau\}} = r \sqrt{\frac{27M^2}{1-9M^2\Lambda}} \sin \theta dr d\theta d\varphi.$$

We note that the normal of $\{\bar{t} = \tau\}$, with respect to \mathring{g} , is

$$(2.20) \quad \mathring{n}_{\{\bar{t}=\tau\}} = \sqrt{\frac{27M^2}{1-9M^2\Lambda}} \frac{1}{r^2} \frac{\partial}{\partial \bar{t}} + \frac{\xi(r)}{\sqrt{\frac{27M^2}{1-9M^2\Lambda}} \frac{1}{r^2}} \frac{\partial}{\partial r},$$

which we also simply denote as

$$(2.21) \quad \mathring{n},$$

for $\xi(r)$ see Remark 2.3. We denote the normals of the event and cosmological horizons respectively as

$$(2.22) \quad \mathring{n}_{\mathcal{H}^+} = \partial_{\bar{t}}, \quad \mathring{n}_{\bar{\mathcal{H}}^+} = \partial_{\bar{t}}.$$

With the above choice of normals, the corresponding volume forms of the respective null hypersurfaces take the form

$$(2.23) \quad d\mathring{g}_{\mathcal{H}^+} = r^2 \sin \theta d\bar{t} d\sigma_{\mathbb{S}^2}, \quad d\mathring{g}_{\bar{\mathcal{H}}^+} = r^2 \sin \theta d\bar{t} d\sigma_{\mathbb{S}^2}.$$

Furthermore, for $\delta > 0$ sufficiently small, the vectors

$$(2.24) \quad \mathring{n}_{\mathcal{H}_\delta^+} = \frac{\nabla r}{\sqrt{|\mathring{g}(\nabla r, \nabla r)|}} \Big|_{r=r_+-\delta}, \quad \mathring{n}_{\bar{\mathcal{H}}_\delta^+} = -\frac{\nabla r}{\sqrt{|\mathring{g}(\nabla r, \nabla r)|}} \Big|_{r=\bar{r}_++\delta}$$

are the unit outward normals of the spacelike hypersurfaces $\mathcal{H}_\delta^+, \bar{\mathcal{H}}_\delta^+$, respectively, with respect to the metric \mathring{g} . Note that there exist smooth functions

$$(2.25) \quad c_1 : (r_+ - \delta, r_+) \rightarrow \mathbb{R}, \quad \bar{c}_1 : (\bar{r}_+, \bar{r}_+ + \delta) \rightarrow \mathbb{R},$$

with

$$(2.26) \quad \sqrt{|1-\mu|} c_1(r) = 1 + \mathcal{O}(1-\mu), \quad \sqrt{|1-\mu|} \bar{c}_1(r) = 1 + \mathcal{O}(1-\mu),$$

as $r \rightarrow r_+, r \rightarrow \bar{r}_+$ respectively, and $c_2(r) = -\sqrt{|1-\mu|}, \bar{c}_2(r) = \sqrt{|1-\mu|}$, such that the normals of $\mathcal{H}_\delta^+, \bar{\mathcal{H}}_\delta^+$ can be written respectively as

$$(2.27) \quad \mathring{n}_{\mathcal{H}_\delta^+} = c_1(r) \partial_{\bar{t}} + c_2(r) \partial_r, \quad \mathring{n}_{\bar{\mathcal{H}}_\delta^+} = \bar{c}_1(r) \partial_{\bar{t}} + \bar{c}_2(r) \partial_r.$$

By pulling back the spacetime volume form (2.18) on hypersurfaces of constant $r = r_+ - \delta$, or $r = \bar{r}_+ + \delta$, we obtain the respective volume forms

$$(2.28) \quad d\mathring{g}_{\mathcal{H}_\delta^+}, \quad d\mathring{g}_{\bar{\mathcal{H}}_\delta^+} = r^2 \sqrt{|1-\mu|} \sin \theta d\bar{t} d\sigma_{\mathbb{S}^2},$$

which we both denote simply as $d\mathring{g}_{\mathcal{H}_\delta}$.

2.4. Coarea formula. There exists a constant $C(M, \Lambda) > 0$ such that

$$(2.29) \quad C^{-1} \int_{\tau_1}^{\tau_2} d\tau \int_{\{\bar{t}=\tau\}} f d\mathring{g}_{\{\bar{t}=\tau\}} \leq \int \int_{D_\delta(\tau_1, \tau_2)} f d\mathring{g} \leq C \int_{\tau_1}^{\tau_2} d\tau \int_{\{\bar{t}=\tau\}} f d\mathring{g}_{\{\bar{t}=\tau\}}$$

for any continuous non-negative function f .

2.5. The vector fields and the energies. We need the following

Definition 2.4. We define the following vector field

$$(2.30) \quad \mathcal{G} \doteq r \sqrt{1 - \frac{2M}{r} - \frac{\Lambda}{3} r^2} \frac{\partial}{\partial r},$$

for $r \in [r_+, \bar{r}_+]$ with respect to the hyperboloidal coordinates $(\bar{t}, r, \theta, \varphi)$.

We extend the vector field (2.30) to a $C^{0,1}$ vector field on \mathcal{M}_δ by

$$(2.31) \quad \mathcal{G} \equiv 0 \text{ in } \{r_+ - \delta \leq r \leq r_+\} \cup \{\bar{r}_+ \leq r \leq \bar{r}_+ + \delta\}.$$

We need the following definition

Definition 2.5. On the spacelike hypersurface $\{\bar{t} = \tau\} \subset D_\delta(0, \infty)$, with respect to the metric \mathring{g} , we define the non-degenerate high order energy

$$(2.32) \quad E_j[\psi](\tau) = \sum_{1 \leq i_1 + i_2 + i_3 \leq j} \sum_{\alpha=1,2,3} \int_{\{\bar{t}=\tau\}} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_\alpha^{i_3} \psi)^2 d\mathring{g}_{\{\bar{t}=\tau\}},$$

with $j \geq 1$, where for Ω^α see Section 2.1.

We define the following high order energy

$$(2.33) \quad E_{\mathcal{G},j}[\psi](\tau) = \int_{\{\bar{t}=\tau\}} \left(\sum_{1 \leq i_1 + i_2 + i_3 \leq j-1, i_3 \geq 1} \sum_{\alpha=1,2,3} (1-\mu)^{2i_3-1} (\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \partial_r^{i_3} \mathcal{G}\psi)^2 \right. \\ \left. + \sum_{1 \leq i_1 + i_2 \leq j-1} \sum_{\alpha=1,2,3} (\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \mathcal{G}\psi)^2 \right. \\ \left. + \sum_{1 \leq i_1 + i_2 + i_3 \leq j-1} \sum_{\alpha=1,2,3} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_\alpha^{i_3} \psi)^2 \right) d\mathring{g}_{\{\bar{t}=\tau\}},$$

for $j \geq 2$.

Remark 2.5. For any $j \geq 2$, there exists a constant $C(j, M, \Lambda) > 0$, such that for all $\tau \geq 0$ the following holds

$$(2.34) \quad E_{j-1}[\psi](\tau) \leq E_{\mathcal{G},j}[\psi](\tau) \leq C E_j[\psi](\tau).$$

Moreover, note the interpolation statement

Lemma 2.1. Let $k \geq 0$. Then, there exist a constant

$$(2.35) \quad C_{int}(k, M, \Lambda) > 0$$

such that for ψ a sufficiently regular function we obtain

$$(2.36) \quad E_{k+1}[\psi](\tau) \leq C_{int} (E_{k-1}[\psi](\tau))^{1/3} (E_{k+2}[\psi](\tau))^{2/3},$$

for any $\tau \geq 0$.

2.6. Notation for derivatives. We need the following notations

Definition 2.6. Let X be a Lipschitz vector field on the manifold \mathcal{M}_δ and let ψ be a smooth function on the manifold \mathcal{M}_δ . Then, for all $j \geq 1$ we define

$$(2.37) \quad |\partial\psi| = |\partial_{\bar{t}}\psi| + |\partial_r\psi| + \sum_{\alpha=1,2,3} |\Omega_\alpha\psi|, \\ (\partial^j\psi)^2 = \sum_{1 \leq j_1 + j_2 + j_3 \leq j} \sum_{\alpha} (\partial_{\bar{t}}^{j_1} \partial_r^{j_2} \Omega_\alpha^{j_3} \psi)^2, \\ (X\partial^j\psi)^2 = \sum_{1 \leq j_1 + j_2 + j_3 \leq j} \sum_{\alpha} (X\partial_{\bar{t}}^{j_1} \partial_r^{j_2} \Omega_\alpha^{j_3} \psi)^2, \\ (\partial^j X\psi)^2 = \sum_{1 \leq j_1 + j_2 + j_3 \leq j} \sum_{\alpha} (\partial_{\bar{t}}^{j_1} \partial_r^{j_2} \Omega_\alpha^{j_3} X\psi)^2.$$

Note that if X is only Lipschitz (for instance $X = \mathcal{G}$) then the last expression of (2.37) may not necessarily be finite.

It will be convenient to compare the above expressions with expressions of coordinate derivatives in ambient globally defined Cartesian coordinates. We define the map

$$(2.38) \quad \begin{aligned} \mathcal{M}_\delta &\rightarrow \mathbb{R}^4 \\ (\bar{t}, r, \theta, \varphi) &\mapsto (x^0, x^1, x^2, x^3) \end{aligned}$$

to be a change of coordinates from the coordinates ascribed to the manifold \mathcal{M}_δ , see Definition 2.1, to Cartesian coordinates, where

$$(2.39) \quad x^0 = \bar{t}, \quad x^1 = r \sin \theta \cos \varphi, \quad x^2 = r \sin \theta \sin \varphi, \quad x^3 = r \cos \theta.$$

Define the set

$$(2.40) \quad \text{Cart}_m = \left\{ \underbrace{\partial_{x^1} \partial_{x^2} \dots \partial_{x^1}}_m, \partial_{x^1} \partial_{x^2} \dots \partial_{x^0}, \dots \right\}$$

to have as elements all the operators that are comprised of any collection of m derivatives of the coordinate vector fields of the Cartesian coordinates (x^0, x^1, x^2, x^3) , see (2.38).

For all $j \geq 1$, we note the similarities

$$(2.41) \quad \begin{aligned} |\partial \psi| &\sim |\partial_{x^0} \psi| + |\partial_{x^1} \psi| + |\partial_{x^2} \psi| + |\partial_{x^3} \psi|, \\ (\partial^j \psi)^2 &\sim \sum_{1 \leq i \leq j} \sum_{\text{all } D^i \in \text{Cart}_i} (D^i \psi)^2. \end{aligned}$$

Finally, for an arbitrary smooth Lorentzian metric g , the wave operator is

$$(2.42) \quad \square_g = g^{ab} \partial_a \partial_b + \Gamma^c_{ab}(g) \partial_c,$$

where $\Gamma^c_{ab}(g)$ are the Christoffel symbols, of the Levi-Civita connection, with respect to the metric g .

2.7. The smooth tensor a . We fix a smooth tensor

$$(2.43) \quad a : T\mathcal{M}_\delta \times T\mathcal{M}_\delta \rightarrow \mathbb{R}.$$

Note that for any $m \in \mathbb{N} \cup \{0\}$ there exist constants $A_m < \infty$, such that if a^{ij} are the components of its inverse in Cartesian coordinates (2.38) and D^m any element of the set Cart_m , see (2.40), then

$$(2.44) \quad |D^m a^{ij}| \leq A_m$$

for all m, i, j .

In what follows, if a constant C depends on A_0, A_1, \dots, A_m we denote it as

$$(2.45) \quad C = C(A_{[m]}).$$

2.8. Sobolev inequality. Note the following Sobolev inequality.

Lemma 2.2. *There exists a constant $C = C(M, \Lambda) > 0$ such that for $f \in C^\infty(\{\bar{t} = c\})$, the following holds*

$$(2.46) \quad \|f\|_{L^\infty(\{\bar{t}=c\})}^2 \leq C(M, \Lambda) \sum_{1 \leq i_1 + i_2 \leq 2} \sum_{\alpha=1,2,3} \int_{\{\bar{t}=c\}} (\partial_r^{i_1} \Omega_\alpha^{i_2} f)^2 d\hat{g}_{\{\bar{t}=c\}}.$$

2.9. The \mathcal{G} vector field commutation estimate. In our [30], we proved the following on a Schwarzschild–de Sitter background.

Theorem 2.1. (Theorem 3 of [30]) *Let ψ satisfy the inhomogeneous wave equation*

$$(2.47) \quad \square_{g_{M,\Lambda}} \psi = F$$

on $D_\delta(\tau_1, \tau_2)$. Then, for any $k \geq 3$ there exists a constant $C(k, M, \Lambda) > 0$, such that the following estimate holds

$$(2.48) \quad \begin{aligned} & E_{\mathcal{G},k}[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) \\ & \leq C E_{\mathcal{G},k}[\psi](\tau_1) \\ & \quad + C \int \int_{D_\delta(\tau_1, \tau_2)} d\mathring{g} \sum_{0 \leq i_1 + i_2 + i_3 \leq k-2} \sum_{\alpha} \left(\partial_{\bar{t}}^{i_1} \partial_r^{i_2} (\Omega_\alpha)^{i_3} F \right)^2 + (1-\mu) \sum_{0 \leq i_1 + i_3 \leq k-2} \sum_{\alpha} \left(\partial_{\bar{t}}^{i_1} (\Omega_\alpha)^{i_3} \mathcal{G}F \right)^2 \\ & \quad + C \int_{\{\bar{t}=\tau_2\}} d\mathring{g}_{\{\bar{t}=\tau_2\}} \sum_{0 \leq i_1 + i_2 + i_3 \leq k-3} \sum_{\alpha} \left(\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \partial_r^{i_3} F \right)^2 + (1-\mu)^{2i_3+1} \left(\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \partial_r^{i_3} \mathcal{G}F \right)^2 \end{aligned}$$

where the volume forms are with respect to the Schwarzschild–de Sitter metric $g_{M,\Lambda}$, see Section 2.3.

Proof. This is proved in [30]. \square

Remark 2.6. *Note that the highest order term of the error hypersurface terms on the right hand side of (2.48) is of order $k-2$, while the highest order term of the error bulk terms on the right hand side of (2.48) is of order $k-1$. The weights in $(1-\mu)$ on the error terms on the right hand side of (2.48) ensure that terms on the right hand side of (2.48) are regular for a sufficiently regular function F .*

3. METRIC CLOSE TO SCHWARZSCHILD–DE SITTER AND THE LOCAL WELL-POSEDNESS RESULT

3.1. Metric close to Schwarzschild–de Sitter. We define the class of metrics close to Schwarzschild–de Sitter.

We fix a sufficiently regular tensor

$$(3.1) \quad h \in \Gamma(TM_\delta^* \times TM_\delta^* \times TM_\delta^*)$$

with

$$(3.2) \quad h(\cdot, \cdot, v = 0) = 0,$$

and define

$$(3.3) \quad g(v) = \mathring{g} + h(v).$$

We also define $h^{ij}(v) \doteq g^{ij}(v) - \mathring{g}^{ij}$, where $g^{ij}(v), \mathring{g}^{ij}$ are the components of the inverses of the respective tensors in Cartesian coordinates.

Note that for every $m \in \mathbb{N} \cup \{0\}$ there exist constants $B_m < \infty$, such that, for $m \geq 0$, the following hold in Cartesian coordinates

$$(3.4) \quad |D^m h_{ijk}|, \quad |D^m h^{ij}{}_k| \leq B_m,$$

for all m, i, j, k , where $h^{ij}{}_k v^k = h^{ij}(v)$ and D^m is any element of the set Cart_m , see (2.40).

In what follows, if a constant C depends on B_0, B_1, \dots, B_m , we denote it as

$$(3.5) \quad C = C(B_{[m]}).$$

3.2. The Cauchy stability type result. We present a Cauchy stability type result for the quasilinear wave equation, in the form that we will use, appropriately tailored to display the energies that are used in our main global stability results, see already Theorem 1 and Theorem 2.

Proposition 3.1. *Let $k \geq 7$ and let the tensors a, h be as in Sections 2.7, 3.1 respectively. There exists a constant*

$$(3.6) \quad C_{wp}(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 1,$$

where for $A_{[k+1]}, B_{[k+1]}$ see Sections 2.7, 3.1 respectively, such that the following holds.

Let $\tau_{max} > 0$ be an arbitrary positive number. There exist

$$(3.7) \quad \delta(\tau_{max}, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}), \quad \epsilon(\tau_{max}, \delta, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$$

both sufficiently small, such that if we take initial data for (1.1) on $\{\bar{t} = \tau_1\}$ with

$$(3.8) \quad E_{k+1}[\psi](\tau_1) \leq \epsilon,$$

for some $\tau_1 \geq 0$, then there exists a unique H^{k+1} solution ψ to the quasilinear wave equation (1.1), on $D_\delta(\tau_1, \tau_1 + \tau_{max})$, such that

$$(3.9) \quad E_{k+1}[\psi](\tau') \leq C_{wp} E_{k+1}[\psi](\tau_1),$$

$$(3.10) \quad E_{k+2}[\psi](\tau') \leq C_{wp} E_{k+2}[\psi](\tau_1),$$

for all $\tau' \in [\tau_1, \tau_1 + \tau_{max}]$, where (3.10) holds if $E_{k+2}[\psi](\tau_1) < \infty$, in which case the solution is H^{k+2} .

Proof. See the appendix A. □

Remark 3.1. *The result of Proposition 3.1 is a refinement of the usual Cauchy stability and thus we will have to prove it explicitly. The independence of C_{wp} on τ_{max} is connected to the uniform boundedness result [9] for the linear wave equation.*

4. THE MAIN THEOREMS

4.1. Theorem on an arbitrary slab of length τ_{step} . We give the detailed statement of the Theorem, on a fixed large time domain

Theorem 1. *Let $k \geq 7$ and let the tensors a, h be as in Sections 2.7, 3.1 respectively. There exists a constant*

$$(4.1) \quad C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$$

where for $A_{[k+1]}, B_{[k+1]}$ see Sections 2.7, 3.1 respectively, such that the following holds.

For all $L > 0$ there exists a $\tau_{step}(L, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}, C_{int}, C_{wp})$ sufficiently large, where for C_{int}, C_{wp} see respectively Lemma 2.1 and Proposition 3.1, such that there exist

$$(4.2) \quad \delta(\tau_{step}, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}), \quad \epsilon(\tau_{step}, \delta, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$$

sufficiently small such that if we take initial data for (1.1) in $\{\tau = \tau_1\}$ with

$$(4.3) \quad E_{k+1}[\psi](\tau_1) \leq \epsilon, \quad E_{k+2}[\psi](\tau_1) < \infty$$

and

$$(4.4) \quad \tau_2 = \tau_1 + \tau_{step},$$

then there exists a unique H^{k+2} solution in $D_\delta(\tau_1, \tau_2)$ to the quasilinear wave equation (1.1), and for all $\tau' \in [\tau_1, \tau_2]$ the following inequalities hold

$$(4.5) \quad E_{G,k}[\psi](\tau') \leq C E_{G,k}[\psi](\tau_1),$$

$$(4.6) \quad E_{k+1}[\psi](\tau') \leq C_{wp} E_{k+1}[\psi](\tau_1),$$

$$(4.7) \quad E_{k+2}[\psi](\tau') \leq C_{wp} E_{k+2}[\psi](\tau_1).$$

Moreover, the following holds

$$(4.8) \quad E_{\mathcal{G},k}[\psi](\tau_2) \leq \frac{1}{L} E_{\mathcal{G},k}[\psi](\tau_1).$$

Finally, the following holds

$$(4.9) \quad E_{k+1}[\psi](\tau_2) \leq C_{int} e^{-\frac{1}{3} \log(L) + \frac{2}{3} \log C_{wp}} (E_{\mathcal{G},k}[\psi](\tau_1))^{1/3} (E_{k+2}[\psi](\tau_1))^{2/3}.$$

Remark 4.1. For the requirement $k \geq 7$, see already the computations of inequality (5.19) and Proposition 3.1.

4.2. The global nonlinear stability of the quasilinear wave equation (1.1). Now, we present the main Theorem of our paper, which is in fact a Corollary of Theorem 1.

Theorem 2. Let $k \geq 7$, let the tensors a, h be as in Sections 2.7, 3.1 respectively. Then, there exist constants

$$(4.10) \quad c_d(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}), \quad c_g(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}), \quad C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$$

and there exist

$$(4.11) \quad \delta = \delta(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0, \quad \epsilon = \epsilon(\delta, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$$

sufficiently small, such that for

$$(4.12) \quad E_{k+2}[\psi](0) \leq \epsilon,$$

on $\Sigma = \{\bar{t} = 0\}$, the arising H^{k+2} solution of the quasilinear wave equation (1.1) exists globally on $D_\delta(0, \infty)$ and satisfies the following exponential decay

$$(4.13) \quad E_{\mathcal{G},k}[\psi](\tau) \leq C e^{-c_d \tau} E_{\mathcal{G},k}[\psi](0),$$

for any $\tau \geq 0$.

Moreover, we have the following exponential decay of the lower order energies

$$(4.14) \quad E_{k-1}[\psi](\tau) \leq C e^{-c_d \tau} E_{\mathcal{G},k}[\psi](0), \quad E_{k+1}[\psi](\tau) \leq C e^{-c_d \tau} E_{k+2}[\psi](0),$$

while for the top order energy we have the growth estimate

$$(4.15) \quad E_{k+2}[\psi](\tau) \leq C e^{c_g \tau} E_{k+2}[\psi](0).$$

Remark 4.2. Note that the growth constant c_g of inequality (4.15) of Theorem 2 can be made arbitrarily small, if $\delta > 0$ and $\epsilon > 0$ are sufficiently small. See already the proof of Theorem 2 in Section 7. Although (4.15) allows the top order energy $E_{k+2}[\psi]$ to a priori grow exponentially, one can in fact prove stronger results. Specifically, with the additional use of a top order uniform boundedness estimate, one can prove Theorem 2 assuming only small $E_{k+1}[\psi](0)$ initial data, and also uniformly bound the energy $E_{k+1}[\psi](\tau)$ for all times. (Therefore, we expect to only need the initial energy to lie in the Sobolev space H^8 .) We will not however pursue this here as the weaker estimate provided by Proposition 3.1 leading to (4.15) is sufficient. This improvement is however important in the asymptotically flat case.

4.3. Extension of our results on Kerr–de Sitter. We can extend our results for the quasilinear wave equation to the slowly rotating Kerr–de Sitter case by utilizing the fixed frequency vector field commutation that we introduce in our forthcoming [31]. This will be presented in a future paper.

5. ENERGY ESTIMATES ON A SPACETIME SLAB OF FIXED TIME LENGTH

Before turning to the proofs of Theorems 1 and 2, we need a preliminary Proposition.

Proposition 5.1. *Let $k \geq 7$ and let the tensors a, h be as in Sections 2.7, 3.1 respectively. Let ψ be a solution of the quasilinear wave equation (1.1) on the Schwarzschild–de Sitter domain $D_\delta(\tau_1, \tau_2)$ for $\tau_1 \leq \tau_2$ and any $\delta > 0$ sufficiently small. Then, there exists a positive constant*

$$(5.1) \quad C(k, M, \Lambda, A_{[k]}, B_{[k]}) > 0,$$

where for $A_{[k]}, B_{[k]}$ see Sections 2.7, 3.1 respectively, and an $\epsilon > 0$ sufficiently small, such that if

$$(5.2) \quad \sup_{D_\delta(\tau_1, \tau_2)} \sum_{1 \leq j \leq k-1} \sum_{\partial \in \{\partial_{\bar{t}}, \partial_r, \Omega_1, \Omega_2, \Omega_3\}} |\partial^j \psi| \leq \sqrt{\epsilon}$$

holds, then the following estimate holds

$$(5.3) \quad E_{\mathcal{G}, k}[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G}, k}(\tau) \leq C E_{\mathcal{G}, k}[\psi](\tau_1) + C \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G}, k}[\psi](\tau) E_{k+1}[\psi](\tau),$$

for all $\tau_1 \leq \tau_2$.

Proof of Proposition 5.1. We will use the derivatives notation of Section 2.6. Moreover, for the semilinear term $\partial\psi \cdot \partial\psi$ and for the definition of smooth tensors a, h see Sections 2.7, 3.1 respectively. In this proof, when we write

$$(5.4) \quad h_{ab}$$

it is to be understood as $h_{ab}(\nabla\psi)$ in Cartesian coordinates, see (2.38). When we write

$$(5.5) \quad h^{ab}$$

it is to be understood as $h^{ab}(\nabla\psi) = g^{ab}(\nabla\psi) - \dot{g}^{ab}$, in Cartesian coordinates. Finally, in this proof when we write \lesssim it is to be understood that we omit a constant $C(k, M, \Lambda, A_{[m]}, B_{[m]})$, where for $A_{[m]}, B_{[m]}$ see Sections 2.7, 3.1 respectively, and $m \leq k+1$.

Note that we may rewrite $\square_{g(\nabla\psi)}\psi = \partial\psi \cdot \partial\psi$ as

$$(5.6) \quad \square_{\dot{g}}\psi = \underbrace{(\square_{\dot{g}} - \square_{g(\nabla\psi)})\psi}_{F_1[\psi]} + \underbrace{\partial\psi \cdot \partial\psi}_{F_2[\psi]}.$$

We name, for convenience,

$$(5.7) \quad F[\psi] = F_1[\psi] + F_2[\psi] = (\square_{\dot{g}} - \square_{g(\nabla\psi)})\psi + \partial\psi \cdot \partial\psi.$$

Then, by the integrated energy estimate (2.48) of Theorem 2.1 for the inhomogeneous wave equation (5.6), we obtain the following

$$(5.8) \quad \begin{aligned} & E_{\mathcal{G}, k}[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G}, k}[\psi](\tau) \\ & \lesssim E_{\mathcal{G}, k}[\psi](\tau_1) + \int \int_{D(\tau_1, \tau_2)} \sum_{0 \leq i_1 + i_2 + i_3 \leq k-2} \sum_{\alpha} ((\partial_{\bar{t}})^{i_1} \partial_r^{i_2} (\Omega_\alpha)^{i_3} F)^2 + (1-\mu) \sum_{0 \leq i+j \leq k-2} \sum_{\alpha} ((\partial_{\bar{t}})^i (\Omega_\alpha)^j \mathcal{G}F)^2 \\ & \quad + \int_{\{\bar{t}=\tau_2\}} \sum_{0 \leq i_1 + i_2 + i_3 \leq k-3} \sum_{\alpha} (1-\mu)^{2i_3+1} (\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \partial_r^{i_3} \mathcal{G}F)^2 + \sum_{0 \leq i_1 + i_2 + i_3 \leq k-3} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \partial_r^{i_3} F)^2 \\ & \lesssim E_{\mathcal{G}, k}[\psi](\tau_1) + \int \int_{D(\tau_1, \tau_2)} \sum_{0 \leq i_1 + i_2 + i_3 \leq k-2} \sum_{\alpha} ((\partial_{\bar{t}})^{i_1} \partial_r^{i_2} (\Omega_\alpha)^{i_3} F_1)^2 + \sum_{0 \leq i+j \leq k-2} \sum_{\alpha} ((\partial_{\bar{t}})^i (\Omega_\alpha)^j \mathcal{G}F_1)^2 \\ & \quad + \int \int_{D(\tau_1, \tau_2)} \sum_{0 \leq i_1 + i_2 + i_3 \leq k-2} \sum_{\alpha} ((\partial_{\bar{t}})^i (\Omega_\alpha)^j F_2)^2 + (1-\mu) \sum_{0 \leq i+j \leq k-2} \sum_{\alpha} ((\partial_{\bar{t}})^{i_1} \partial_r^{i_2} (\Omega_\alpha)^{i_3} \mathcal{G}F_2)^2 \\ & \quad + \int_{\{\bar{t}=\tau_2\}} \sum_{0 \leq i_1 + i_2 + i_3 \leq k-3} \sum_{\alpha} (1-\mu)^{2i_3+1} (\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \partial_r^{i_3} \mathcal{G}F)^2 + \sum_{0 \leq i_1 + i_2 + i_3 \leq k-3} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \partial_r^{i_3} F)^2. \end{aligned}$$

We readily bound the term $\int \int_{D_\delta(\tau_1, \tau_2)} (1-\mu) \sum_{i=0}^{k-2} (\partial_t^i \mathcal{G} F_2)^2$ as follows

$$\begin{aligned}
(5.9) \quad & \int \int_{D(\tau_1, \tau_2)} (1-\mu) \sum_{i=0}^{k-2} (\partial_t^i \mathcal{G} F_2)^2 = \int \int_{D(\tau_1, \tau_2)} (1-\mu) \sum_{i=0}^{k-2} (\partial_t^i \mathcal{G} (a^{\alpha\beta} \partial_\alpha \psi \partial_\beta \psi))^2 \\
& = \int \int_{D(\tau_1, \tau_2)} (1-\mu) \sum_{i=0}^{k-2} (\partial_t^i ([\mathcal{G}, a^{\alpha\beta}] \partial_\alpha \psi \partial_\beta \psi + a^{\alpha\beta} \mathcal{G} (\partial_\alpha \psi \partial_\beta \psi)))^2 \\
& = \int \int_{D(\tau_1, \tau_2)} (1-\mu) \sum_{i=0}^{k-2} (\partial_t^i ([\mathcal{G}, a^{\alpha\beta}] \partial_\alpha \psi \partial_\beta \psi + a^{\alpha\beta} \partial_\beta \psi \mathcal{G} \partial_\alpha \psi + \partial_\alpha \psi \mathcal{G} \partial_\beta \psi))^2 \\
& = \int \int_{D(\tau_1, \tau_2)} (1-\mu) \sum_{i=0}^{k-2} \left(\partial_t^i \left([\mathcal{G}, a^{\alpha\beta}] \partial_\alpha \psi \partial_\beta \psi + a^{\alpha\beta} \partial_\beta \psi [\mathcal{G}, \partial_\alpha] \psi + a^{\alpha\beta} \partial_\beta \psi \partial_\alpha \mathcal{G} \psi \right. \right. \\
& \quad \left. \left. + a^{\alpha\beta} \partial_\alpha \psi [\mathcal{G}, \partial_\beta] \psi + a^{\alpha\beta} \partial_\alpha \psi \partial_\beta \mathcal{G} \psi \right) \right)^2 \\
& \lesssim \int \int_{D(\tau_1, \tau_2)} (1-\mu) \sum_{i=0}^{k-2} (\partial_t^i ([\mathcal{G}, a^{\alpha\beta}] \partial_\alpha \psi \partial_\beta \psi))^2 + (1-\mu) \sum_{i=0}^{k-2} (\partial_t^i (a^{\alpha\beta} \partial_\beta \psi [\mathcal{G}, \partial_\alpha] \psi))^2 \\
& \quad + (1-\mu) \sum_{i=0}^{k-2} (\partial_t^i (a^{\alpha\beta} \partial_\alpha \psi [\mathcal{G}, \partial_\beta] \psi))^2 \\
& \quad + (1-\mu) \sum_{i=0}^{k-2} (\partial_t^i (a^{\alpha\beta} \partial_\alpha \psi \partial_\beta \mathcal{G} \psi))^2 + (1-\mu) \sum_{i=0}^{k-2} (\partial_t^i (a^{\alpha\beta} \partial_\beta \psi \partial_\alpha \mathcal{G} \psi))^2 \\
& \lesssim \int \int_{D(\tau_1, \tau_2)} \sum_{1 \leq i+j \leq k-1} (\partial^i \psi)^2 (\partial^j \psi)^2 \\
& \quad + (1-\mu) \sum_{i=0}^{k-2} (\partial_t^i (a^{\alpha\beta} \partial_\alpha \psi \partial_\beta \mathcal{G} \psi))^2 + (1-\mu) \sum_{i=0}^{k-2} (\partial_t^i (a^{\alpha\beta} \partial_\beta \psi \partial_\alpha \mathcal{G} \psi))^2 \\
& \lesssim \int \int_{D(\tau_1, \tau_2)} \sum_{1 \leq i+j \leq k-1} (\partial^i \psi)^2 (\partial^j \psi)^2 + \int \int_{D(\tau_1, \tau_2)} \sum_{1 \leq i+j \leq k-2, j \leq \lfloor \frac{k-2}{2} \rfloor} (1-\mu) (\partial_t^i \partial \mathcal{G} \psi)^2 (\partial^j \psi)^2.
\end{aligned}$$

Therefore, by using the coarea formula (2.29), see Section 2.4, we bound (5.9) from

$$\begin{aligned}
(5.10) \quad & \int_{\tau_1}^{\tau_2} d\tau \sum_{1 \leq i+j \leq k-1, i \leq \lfloor \frac{k-1}{2} \rfloor} \sup_{\{\bar{t}=\tau\}} (\partial^i \psi)^2 \int_{\{\bar{t}=\tau\}} (\partial^j \psi)^2 + \int_{\tau_1}^{\tau_2} d\tau \sup_{\{\bar{t}=\tau\}} \sum_{1 \leq i+j \leq k-2, j \leq \lfloor \frac{k-2}{2} \rfloor} (\partial^j \partial \psi)^2 \int_{\{\bar{t}=\tau\}} (1-\mu) (\partial_t^i \partial \mathcal{G} \psi)^2 \\
& \quad + \int_{\tau_1}^{\tau_2} d\tau \sum_{j \geq \lfloor \frac{k-2}{2} \rfloor + 1, i \leq k-2 - \lfloor \frac{k-2}{2} \rfloor - 1} \sup_{\{\bar{t}=\tau\}} (\partial^{i+2} \psi)^2 \int_{\{\bar{t}=\tau\}} (\partial^{j+1} \psi)^2 \\
& \lesssim \int_{\tau_1}^{\tau_2} d\tau E_{k-1}[\psi](\tau) E_{k-1}[\psi](\tau) + \int_{\tau_1}^{\tau_2} d\tau E_{k-1}[\psi](\tau) E_{\mathcal{G}, k}[\psi](\tau) \\
& \leq C(k, M, \Lambda, A_{[k-1]}, B_{[k-1]}) \int_{\tau_1}^{\tau_2} d\tau E_k[\psi](\tau) E_{\mathcal{G}, k}[\psi](\tau)
\end{aligned}$$

where in the first inequality we use the Sobolev inequality, see Lemma 2.2, and we also used the requirement $k \geq 6$. The remaining terms

$$(5.11) \quad \int \int_{D(\tau_1, \tau_2)} (1-\mu) \sum_{0 \leq i+j \leq k-2} \sum_{\alpha} ((\partial_t)^{i_1} (\Omega_\alpha)^{i_3} \mathcal{G} F_2)^2$$

are similarly bounded by the right hand side of (5.10).

Moreover, we readily bound the term $\int \int_{D(\tau_1, \tau_2)} \sum_{i=0}^{k-2} (\partial_{\bar{t}}^i F_2)^2$ on the right hand side of equation (5.8) by using the coarea formula (2.29), see Section 2.4, by the expression below

$$\begin{aligned}
(5.12) \quad & \int \int_{D(\tau_1, \tau_2)} \sum_{i=0}^{k-2} (\partial_{\bar{t}}^i F_2)^2 = \int \int_{D(\tau_1, \tau_2)} \sum_{i=0}^{k-2} (\partial_{\bar{t}}^i (a^{\alpha\beta} \partial_\alpha \psi \partial_\beta \psi))^2 \\
& \lesssim \int \int_{D(\tau_1, \tau_2)} \sum_{1 \leq i+j \leq k-1} (\partial^i \psi)^2 (\partial^j \psi)^2 \\
& \lesssim \int_{\tau_1}^{\tau_2} \sum_{1 \leq i+j \leq k-1, i \leq \lfloor \frac{k-1}{2} \rfloor} \sup_{\{\bar{t}=\tau\}} (\partial^i \psi)^2 \int_{\{\bar{t}=\tau\}} (\partial^j \psi)^2 \\
& \lesssim \int_{\tau_1}^{\tau_2} d\tau E_{k-1}[\psi](\tau) E_{k-1}[\psi](\tau) \\
& \leq C(k, M, \Lambda, A_{[k-1]}, B_{[k-1]}) \int_{\tau_1}^{\tau_2} d\tau E_{k-1}[\psi](\tau) E_{\mathcal{G}, k}[\psi](\tau),
\end{aligned}$$

where in the second to last inequality we use the Sobolev inequality, see Lemma 2.2, and we also used the requirement $k \geq 4$. The remaining terms

$$(5.13) \quad \int \int_{D(\tau_1, \tau_2)} \sum_{0 \leq i_1+i_2+i_3 \leq k-2} \sum_{\alpha} ((\partial_{\bar{t}})^i (\Omega_\alpha)^j F_2)^2$$

are similarly bounded by the right hand side of (5.12). Note that *we have not yet appealed to the smallness* (5.2), since we have thus far only been discussing semi-linear terms.

We estimate the term $\int \int_{D_\delta(\tau_1, \tau_2)} \sum_{i=0}^{k-2} (\partial_{\bar{t}}^i \mathcal{G} F_1)^2$, on the right hand side of equation (5.8), by using the coarea formula (2.29), by the following expression

$$\begin{aligned}
(5.14) \quad & \int_{\tau_1}^{\tau_2} d\tau \int_{\{\bar{t}=\tau\}} \sum_{i=0}^{k-2} \left(\partial_{\bar{t}}^i \mathcal{G} ((\mathring{g}^{ab} - g^{ab}(\nabla\psi)) \partial_a \partial_b \psi) \right. \\
& \quad \left. + \partial_{\bar{t}}^i \mathcal{G} (\mathring{g}^{ab} \Gamma^c{}_{ab}(\mathring{g}) \partial_c \psi - g^{ab}(\nabla\psi) \Gamma^c{}_{ab}(g(\nabla\psi)) \partial_c \psi) \right)^2.
\end{aligned}$$

Now, we note that

$$(5.15) \quad \Gamma^c{}_{ab}(g(\nabla\psi)) = \Gamma^c{}_{ab}(\mathring{g}) + S^c{}_{ab}(h),$$

where

$$\begin{aligned}
(5.16) \quad & S^c{}_{ab}(h) \doteq \frac{1}{2} \mathring{g}^{ce} (\partial_e h_{ab} + \partial_a h_{eb} + \partial_b h_{ea}) \\
& \quad + \frac{1}{2} h^{ce} (-\partial_e h_{ab} + \partial_a h_{eb} + \partial_b h_{ea}) \\
& \quad + \frac{1}{2} h^{ce} (-\partial_e \mathring{g}_{ab} + \partial_a \mathring{g}_{eb} + \partial_b \mathring{g}_{ea}).
\end{aligned}$$

Therefore, we write (5.14) in the following form

$$\begin{aligned}
(5.17) \quad & \int_{\tau_1}^{\tau_2} d\tau \int_{\{\bar{t}=\tau\}} \sum_{i=0}^{k-2} \left(\partial_{\bar{t}}^i \mathcal{G} (h^{ab} \partial_a \partial_b \psi) + \partial_{\bar{t}}^i \mathcal{G} \left((\dot{g}^{ab} S^c{}_{ab}(h) - h^{ab} \Gamma^c{}_{ab}(\dot{g}) - h^{ab} S^c{}_{ab}(h)) \partial_c \psi \right) \right)^2 \\
& = \int_{\tau_1}^{\tau_2} d\tau \int_{\{\bar{t}=\tau\}} \sum_{i=0}^{k-2} \left(\partial_{\bar{t}}^i \mathcal{G} (h^{ab} \partial_a \partial_b \psi) + \partial_{\bar{t}}^i \mathcal{G} (\dot{g}^{ab} S^c{}_{ab}(h) \partial_c \psi) - \partial_{\bar{t}}^i \mathcal{G} (h^{ab} \Gamma^c{}_{ab}(\dot{g}) \partial_c \psi) - \partial_{\bar{t}}^i \mathcal{G} (h^{ab} S^c{}_{ab}(h) \partial_c \psi) \right)^2 \\
& \lesssim \int_{\tau_1}^{\tau_2} d\tau \int_{\{\bar{t}=\tau\}} \sum_{1 \leq i \leq k-1} \left(\left(\partial^i (h^{ab} \partial_a \partial_b \psi) \right)^2 + \left(\partial^i (\dot{g}^{ab} S^c{}_{ab}(h) \partial_c \psi) \right)^2 + \left(\partial^i (h^{ab} \Gamma^c{}_{ab}(\dot{g}) \partial_c \psi) \right)^2 \right. \\
& \quad \left. + \left(\partial^i (h^{ab} S^c{}_{ab}(h) \partial_c \psi) \right)^2 \right)
\end{aligned}$$

which, by using the smallness (5.2) for a sufficiently small $\epsilon > 0$, the definition of h , see Section 3.1, and by distributing the derivatives (with the Cartesian coordinates of Section 2.6), we bound the above from

$$\begin{aligned}
(5.18) \quad & \lesssim \int_{\tau_1}^{\tau_2} d\tau \int_{\{\bar{t}=\tau\}} \sum_{1 \leq i \leq k-1} \left(\sum_{i_1+i_2=i} (|\partial^{i_1} h^{ab}| |\partial^{i_2} \partial_a \partial_b \psi|)^2 + \sum_{i_1+i_2+i_3=i} (|\partial^{i_1} \dot{g}^{ab}| |\partial^{i_2} S^c{}_{ab}(h)| |\partial^{i_3} \partial_c \psi|)^2 \right. \\
& \quad + \sum_{i_1+i_2+i_3=i} (|\partial^{i_1} h^{ab}| |\partial^{i_2} \Gamma^c{}_{ab}(\dot{g})| |\partial^{i_3} \partial_c \psi|)^2 \\
& \quad \left. + \sum_{i_1+i_2+i_3=i} (|\partial^{i_1} h^{ab}| |\partial^{i_2} S^c{}_{ab}(h)| |\partial^{i_3} \partial_c \psi|)^2 \right) \\
& \lesssim \int_{\tau_1}^{\tau_2} d\tau \int_{\{\bar{t}=\tau\}} \sum_{1 \leq i \leq k-1, i_1 \leq \lfloor \frac{i}{2} \rfloor} (\partial^{i_1+1} \psi)^2 (\partial^{i_2+2} \psi)^2 + \sum_{4 \leq i \leq k-1} \sum_{i_1 = \lfloor \frac{i}{2} \rfloor + 1, i_2 = \lceil \frac{i}{2} \rceil - 1} (\partial^{i_1+1} \psi)^2 (\partial^{i_2+2} \psi)^2 \\
& \quad + \sum_{4 \leq i \leq k-1} \sum_{i_1 = \lfloor \frac{i}{2} \rfloor + 2, i_2 = \lceil \frac{i}{2} \rceil - 2} (\partial^{i_1+1} \psi)^2 (\partial^{i_2+2} \psi)^2 + \sum_{5 \leq i \leq k-1} \sum_{i_2 = \lceil \frac{i}{2} \rceil - 3, i_1 > \lfloor \frac{i}{2} \rfloor + 3} (\partial^{i_1+1} \psi)^2 (\partial^{i_2+2} \psi)^2 \\
& \quad + \sum_{1 \leq i \leq 4} \sum_{i_1+i_2=i} (\partial^{i_1+1} \psi)^2 (\partial^{i_2+2} \psi)^2 \\
& \quad + \sum_{1 \leq i_1+i_2 \leq k-1, i_1 \leq \lceil \frac{k-1}{3} \rceil} (\partial^{i_1+1} \psi)^2 (\partial^{i_2+1} \psi)^2 \\
& \quad + \sum_{1 \leq i_1+i_2+i_3 \leq k-1, i_1 \leq \lceil \frac{k-1}{3} \rceil} (\partial^{i_1} \psi)^2 (\partial^{i_2+1} \psi)^2 (\partial^{i_3+1} \psi)^2
\end{aligned}$$

therefore, by using Sobolev inequalities, see Lemma 2.2, we obtain from (5.18) the following

$$\begin{aligned}
& \lesssim \int_{\tau_1}^{\tau_2} d\tau \sum_{1 \leq i \leq k-1, i_1 \leq \lfloor \frac{i}{2} \rfloor} \sup_{\{\bar{t}=\tau\}} (\partial^{i_1+1} \psi)^2 \int_{\{\bar{t}=\tau\}} (\partial^{i_2+2} \psi)^2 \\
& \quad + \sum_{4 \leq i \leq k-1} \sum_{i_1 = \lfloor \frac{i}{2} \rfloor + 1, i_2 = \lfloor \frac{i}{2} \rfloor - 1} \sup_{\{\bar{t}=\tau\}} (\partial^{i_1+1} \psi)^2 \int_{\{\bar{t}=\tau\}} (\partial^{i_2+2} \psi)^2 \\
& \quad + \sum_{4 \leq i \leq k-1} \sum_{i_1 = \lfloor \frac{i}{2} \rfloor + 2, i_2 = \lfloor \frac{i}{2} \rfloor - 2} \sup_{\{\bar{t}=\tau\}} (\partial^{i_1+1} \psi)^2 \int_{\{\bar{t}=\tau\}} (\partial^{i_2+2} \psi)^2 \\
& \quad + \sum_{5 \leq i \leq k-1} \sum_{i_2 \leq \lfloor \frac{i}{2} \rfloor - 3, i_1} \sup_{\{\bar{t}=\tau\}} (\partial^{i_2+2} \psi)^2 \int_{\{\bar{t}=\tau\}} (\partial^{i_1+1} \psi)^2 \\
(5.19) \quad & \quad + \sum_{1 \leq i \leq 4} \sum_{i_1+i_2=i} \sup_{\{\bar{t}=\tau\}} (\partial^{i_1+1} \psi)^2 \int_{\{\bar{t}=\tau\}} (\partial^{i_2+2} \psi)^2 \\
& \quad + \sum_{1 \leq i_1+i_2 \leq k-1, i_1 \leq \lfloor \frac{k-1}{3} \rfloor} \sup_{\{\bar{t}=\tau\}} (\partial^{i_1+1} \psi)^2 \int_{\{\bar{t}=\tau\}} (\partial^{i_2+1} \psi)^2 \\
& \quad + \sum_{1 \leq i_1+i_2+i_3 \leq k-1, i_1 \leq \lfloor \frac{k-1}{3} \rfloor} \sup_{\{\bar{t}=\tau\}} (\partial^{i_1} \psi)^2 \int_{\{\bar{t}=\tau\}} (\partial^{i_2+1} \psi)^2 (\partial^{i_3+1} \psi)^2 \\
& \lesssim \int_{\tau_1}^{\tau_2} d\tau E_{k-1}[\psi](\tau) E_{k+1}[\psi](\tau) \\
& \leq C(k, M, \Lambda, A_{[k]}, B_{[k]}) \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) E_{k+1}[\psi](\tau)
\end{aligned}$$

where in the second to last inequality we also used that $k = 7$ is the smallest integer such that all of the following hold

$$(5.20) \quad \left(\lfloor \frac{k-1}{2} \rfloor + 1 \right) + 2 \leq k-1, \quad \left(\lfloor \frac{k-1}{2} \rfloor + 1 + 1 \right) + 2 \leq k+1, \quad \left(\lfloor \frac{k-1}{2} \rfloor + 2 + 1 \right) + 2 \leq k+1, \quad \left(\lfloor \frac{k-1}{2} \rfloor - 3 + 2 \right) + 2 \leq k-1$$

where the addition of $+2$ on the above inequalities comes from the Sobolev inequality, see Lemma 2.2.

Now, by similar calculations, we bound the following term of the right hand side of (5.8)

$$(5.21) \quad \int \int_{D(\tau_1, \tau_2)} \sum_{0 \leq i_1+i_2+i_3 \leq k-2} \sum_{\alpha} ((\partial_{\bar{t}})^{i_1} \partial_r^{i_2} (\Omega_{\alpha})^{i_3} F_1)^2 + \sum_{0 \leq i+j \leq k-2} \sum_{\alpha} ((\partial_{\bar{t}})^i (\Omega_{\alpha})^j \mathcal{G} F_1)^2 \leq C \int_{\tau_1}^{\tau_2} E_{\mathcal{G},k}[\psi](\tau) E_k[\psi](\tau).$$

Finally, by using the bounds (5.10) (5.12), (5.19), (5.21), the integrated energy estimate of equation (5.8) implies the following

$$\begin{aligned}
(5.22) \quad & E_{\mathcal{G},k}[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}(\tau) \\
& \leq C(M, \Lambda) E_{\mathcal{G},k}[\psi](\tau_1) + C(k, M, \Lambda, A_{[k]}, B_{[k]}) \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) E_{k+1}[\psi](\tau) \\
& \quad + C(M, \Lambda) \int_{\{\bar{t}=\tau_2\}} \sum_{0 \leq i_1+i_2+i_3 \leq k-3} \sum_{\alpha} (1-\mu)^{2i_3+1} (\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \partial_r^{i_3} \mathcal{G} F)^2 + \sum_{0 \leq i_1+i_2+i_3 \leq k-3} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \partial_r^{i_3} F)^2,
\end{aligned}$$

where recall that $F = (\square_{\hat{g}} - \square_{g(\nabla\psi)}) \psi + \partial\psi \cdot \partial\psi$.

Finally, we want to absorb the boundary terms of the right hand side at $\{\bar{t} = \tau_2\}$ by the relevant boundary term $E_{\mathcal{G},k}[\psi](\tau_2)$ of the left hand side. By using the smallness assumption (5.2), it is evident that we can appropriately distribute derivatives, in view of the requirement $k \geq 7$, to

conclude that

(5.23)

$$\begin{aligned} E_{\mathcal{G},k}[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}(\tau) \\ \leq C(k, M, \Lambda, A_{[k]}, B_{[k]}) E_{\mathcal{G},k}[\psi](\tau_1) + C(k, M, \Lambda, A_{[k]}, B_{[k]}) \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) E_{k+1}[\psi](\tau), \end{aligned}$$

for a constant $C(k, M, \Lambda, A_{[k]}, B_{[k]})$.

The proof of Proposition 5.1 is thus complete. \square

6. PROOF OF THEOREM 1

We are ready to prove Theorem 1.

Proof of Theorem 1. Given $L > 0$, and for a

$$(6.1) \quad \tau_{step}(L, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$$

to be determined later (see already (6.19)), we apply Proposition 3.1 with $\tau_{max} = \tau_{step}$ to find

$$(6.2) \quad \delta(\tau_{step}, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0, \quad \epsilon(\tau_{step}, \delta, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$$

sufficiently small, such that if we take

$$(6.3) \quad E_{k+1}[\psi](\tau_1) \leq \epsilon$$

then the solution ψ of the quasilinear wave equation (1.1) exists in $D_\delta(\tau_1, \tau_2)$, where

$$(6.4) \quad \tau_2 = \tau_1 + \tau_{step}.$$

The two inequalities (4.6), (4.7) are an immediate consequence of the Cauchy stability results (3.9), (3.10) respectively, of Proposition 3.1.

To obtain the remaining statements, we assume the bootstrap

$$(6.5) \quad \sup_{D_\delta(\tau_1, \tau_2)} \sum_{1 \leq j \leq k-1} \sum_{\partial \in \{\partial_{\bar{t}}, \partial_r, \Omega_1, \Omega_2, \Omega_3\}} |\partial^j \psi| \leq C_b \sqrt{\epsilon},$$

for a constant

$$(6.6) \quad C_b(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$$

which will be determined later.

First, we prove inequality (4.5), namely

$$(6.7) \quad \sup_{\tau \in [\tau_1, \tau_2]} E_{\mathcal{G},k}[\psi](\tau) \leq C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) E_{\mathcal{G},k}[\psi](\tau_1).$$

To prove (6.7), we will introduce an additional bootstrap assumption

$$(6.8) \quad \sup_{\tau \in [\tau_1, \tau_2]} E_{\mathcal{G},k}[\psi](\tau) \leq C_{boot} E_{\mathcal{G},k}[\psi](\tau_1).$$

We choose $\epsilon(C_{boot}) > 0$ sufficiently small so that, in view of the bootstrap (6.5), the energy estimate (5.3) of Proposition 5.1 holds, namely

(6.9)

$$\begin{aligned} E_{\mathcal{G},k}[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} E_{\mathcal{G},k}[\psi](\tau') d\tau' \\ \leq C(k, M, \Lambda, A_{[k]}, B_{[k]}) E_{\mathcal{G},k}[\psi](\tau_1) + C(k, M, \Lambda, A_{[k]}, B_{[k]}) \int_{\tau_1}^{\tau_2} E_{\mathcal{G},k}[\psi](\tau') E_{k+1}[\psi](\tau') d\tau'. \end{aligned}$$

Now we use the bootstrap assumption (6.8) and inequality (4.6), which is already proven, to obtain (6.10)

$$\begin{aligned}
E_{\mathcal{G},k}[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} E_{\mathcal{G},k}[\psi](\tau') d\tau' &\leq C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) E_{\mathcal{G},k}[\psi](\tau_1) \\
&\quad + C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) C_{\text{boot}} \tau_{\text{step}} E_{\mathcal{G},k}[\psi](\tau_1) E_{k+1}[\psi](\tau_1) \\
&\leq C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) E_{\mathcal{G},k}[\psi](\tau_1) \\
&\quad + C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) C_{\text{boot}} \tau_{\text{step}} \cdot \epsilon \cdot E_{\mathcal{G},k}[\psi](\tau_1) \\
&\leq C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) (1 + C_{\text{boot}} \tau_{\text{step}} \epsilon) E_{\mathcal{G},k}[\psi](\tau_1).
\end{aligned}$$

Therefore, by choosing

$$(6.11) \quad \frac{C_{\text{boot}}(k, M, \Lambda, A_{[k+1]}, B_{[k+1]})}{2} \gg C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]})$$

we take $\epsilon(\tau_{\text{step}}, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$ sufficiently small in inequality (6.10), and obtain

$$(6.12) \quad \sup_{\tau \in [\tau_1, \tau_2]} E_{\mathcal{G},k}[\psi](\tau) \leq \frac{C_{\text{boot}}}{2} E_{\mathcal{G},k}[\psi](\tau_1).$$

We improved the bootstrap (6.8) and thus proved (6.7). Therefore, we concluded inequality (4.5).

Now, we proceed to prove inequality (4.8). We use the energy estimate (6.9), in view of the now established bound (6.7), and inequality (4.6), namely $E_{k+1}[\psi](\tau') \leq C_{\text{wp}} E_{k+1}[\psi](\tau_1)$ for all $\tau' \in [\tau_1, \tau_2]$, to obtain

$$\begin{aligned}
(6.13) \quad &\int_{\tau_1}^{\tau_2} E_{\mathcal{G},k}[\psi](\tau') d\tau' \leq C E_{\mathcal{G},k}[\psi](\tau_1) + C \tau_{\text{step}} E_{\mathcal{G},k}[\psi](\tau_1) E_{k+1}[\psi](\tau_1) \\
\implies &\int_{\frac{\tau_1+\tau_2}{2}}^{\tau_2} E_{\mathcal{G},k}[\psi](\tau') d\tau' \leq C E_{\mathcal{G},k}[\psi](\tau_1) + C \tau_{\text{step}} E_{\mathcal{G},k}[\psi](\tau_1) E_{k+1}[\psi](\tau_1) \\
\implies &E_{\mathcal{G},k}[\psi](\tau_{1,2}) \leq \frac{2C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]})}{\tau_{\text{step}}} (E_{\mathcal{G},k}[\psi](\tau_1) + \tau_{\text{step}} E_{\mathcal{G},k}[\psi](\tau_1) E_{k+1}[\psi](\tau_1))
\end{aligned}$$

where we used that there exists a $\tau_{1,2} \in [\frac{\tau_1+\tau_2}{2}, \tau_2]$ such that

$$(6.14) \quad \frac{\tau_{\text{step}}}{2} E_{\mathcal{G},k}[\psi](\tau_{1,2}) \leq \int_{\frac{\tau_1+\tau_2}{2}}^{\tau_2} E_{\mathcal{G},k}[\psi](\tau') d\tau',$$

by the mean value theorem. Therefore, from inequality (6.13) we conclude that, if we take

$$(6.15) \quad E_{k+1}[\psi](\tau_1) \leq \epsilon,$$

then there exists a $\tau_{1,2} \in [\frac{\tau_1+\tau_2}{2}, \tau_2]$ such that

$$(6.16) \quad E_{\mathcal{G},k}[\psi](\tau_{1,2}) \leq \left(\frac{2C}{\tau_{\text{step}}} + 2C\epsilon \right) E_{\mathcal{G},k}[\psi](\tau_1).$$

Furthermore, by using the finite in time bound (6.7), for $E_{\mathcal{G},k}[\psi]$, in the time domain

$$(6.17) \quad [\tau_{1,2}, \tau_2],$$

in conjunction with (6.16), we find a, potentially different, constant $C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$ such that

$$(6.18) \quad E_{\mathcal{G},k}[\psi](\tau_2) \leq \left(\frac{2C}{\tau_{\text{step}}} + 2C\epsilon \right) E_{\mathcal{G},k}[\psi](\tau_1).$$

Finally, we choose $\tau_{\text{step}}(L, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$ sufficiently large and conclude

$$(6.19) \quad E_{\mathcal{G},k}[\psi](\tau_2) \leq \frac{1}{L} E_{\mathcal{G},k}[\psi](\tau_1),$$

after taking $\epsilon(\tau_{\text{step}})$ sufficiently small. We have concluded the inequality (4.8).

To conclude (4.9), we use the now established (4.5), (4.6), (4.7), (4.8) and the classical interpolation Lemma 2.1. Namely, by recalling the constant

$$(6.20) \quad C_{int}(k, M, \Lambda) > 0$$

from the interpolation Lemma 2.1, we obtain

$$(6.21) \quad \begin{aligned} E_{k+1}[\psi](\tau_2) &\leq C_{int} (E_{k-1}[\psi](\tau_2))^{1/3} (E_{k+2}[\psi](\tau_2))^{2/3} \\ &\leq C_{int} (E_{\mathcal{G},k}[\psi](\tau_2))^{1/3} (E_{k+2}[\psi](\tau_2))^{2/3} \\ &\leq C_{int} \left(\frac{1}{L}\right)^{1/3} (C_{wp})^{2/3} (E_{\mathcal{G},k}[\psi](\tau_1))^{1/3} (E_{k+2}[\psi](\tau_1))^{2/3} \\ &\leq C_{int} e^{-\frac{1}{3} \log(L) + \frac{2}{3} \log C_{wp}} (E_{\mathcal{G},k}[\psi](\tau_1))^{1/3} (E_{k+2}[\psi](\tau_1))^{2/3}, \end{aligned}$$

where for the constant

$$(6.22) \quad C_{wp}(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$$

see Proposition 3.1. We have concluded (4.9).

Of course our inequalities are still conditional on improving the bootstrap assumption (6.5). We see immediately however that by taking $C_b \gg C_{wp}$ we improve the bootstrap (6.5) by applying a Sobolev inequality on the already proved inequality (4.6).

The proof of Theorem 1 is thus complete. \square

7. PROOF OF THEOREM 2

We are now ready to prove Theorem 2.

Proof of Theorem 2. We will use the results of Theorem 1 in conjunction with an iteration argument in consecutive spacetime regions.

To apply Theorem 1, we choose $L > 0$ sufficiently large so that

$$(7.1) \quad L > \left(2(C_{int} + 1) e^{\frac{2}{3} \log(C_{wp})}\right)^3$$

where for the constants $C_{int}, C_{wp} > 0$ see equation (4.9). Note that with this choice, the constant on the right hand side of (4.9) satisfies

$$(7.2) \quad C_{int} e^{-\frac{1}{3} \log L + \frac{2}{3} \log C_{wp}} < \frac{1}{2}.$$

This gives us τ_{step}, δ and ϵ from the statement of Theorem 1 and our choice of L , namely (7.1). We obtain

$$(7.3) \quad E_{k+1}[\psi](\tau_2) \leq (E_{\mathcal{G},k}[\psi](\tau_1))^{1/3} (E_{k+2}[\psi](\tau_1))^{2/3},$$

for $\tau_2 = \tau_1 + \tau_{step}$.

First, we prove that there exists a strictly increasing sequence of real numbers $\{t_i\}_{i \in \mathbb{N}}$, $t_i \rightarrow \infty$, such that

$$(7.4) \quad t_0 = 0, \quad t_{i+1} - t_i = \tau_{step},$$

the solution exists in $D_\delta(0, t_i)$ and the following hold

$$(7.5) \quad \begin{aligned} E_{\mathcal{G},k}[\psi](t_i) &\leq \left(\frac{1}{L}\right)^i E_{\mathcal{G},k}[\psi](0), \\ E_{k+1}[\psi](t_i) &\leq \epsilon, \\ E_{k+2}[\psi](t_i) &\leq (C_{wp})^i E_{k+2}[\psi](0) \end{aligned}$$

Note that (7.5) holds for t_0 . For the purpose of using induction, we assume that the solution exists in $D_\delta(0, t_i)$ and also assume the iterative step that (7.5) holds for t_i , so we want to prove (7.5) for t_{i+1} . Now, in view of Theorem 1 we obtain that the solution exists in $D_\delta(0, t_{i+1})$ and moreover

in view of inequality (7.3), the condition for L namely (7.1) and by the iterative step assumptions we obtain

$$\begin{aligned}
(7.6) \quad E_{k+1}[\psi](t_{i+1}) &\leq (E_{\mathcal{G},k}[\psi](t_i))^{1/3} (E_{k+2}[\psi](t_i))^{2/3} \\
&\leq \left(\frac{1}{L}\right)^{i/3} e^{\frac{2}{3}i \log(C_{wp})} (E_{\mathcal{G},k}[\psi](0))^{1/3} (E_{k+2}[\psi](0))^{2/3} \\
&\leq e^{i(-\frac{1}{3} \log(L) + \frac{2}{3} \log(C_{wp}))} (E_{\mathcal{G},k}[\psi](0))^{1/3} (E_{k+2}[\psi](0))^{2/3} \\
&\leq (E_{\mathcal{G},k}[\psi](0))^{1/3} (E_{k+2}[\psi](0))^{2/3} \\
&\leq \epsilon.
\end{aligned}$$

Moreover, since $E_{k+1}[\psi](0) \leq \epsilon$ is sufficiently small $\epsilon > 0$, we apply (4.8) of Theorem 1 in conjunction with the inductive step and obtain

$$(7.7) \quad E_{\mathcal{G},k}[\psi](t_{i+1}) \leq \frac{1}{L} E_{\mathcal{G},k}[\psi](t_i) \leq \left(\frac{1}{L}\right)^{i+1} E_{\mathcal{G},k}[\psi](0).$$

Furthermore, we apply inequality (4.7) of Theorem 1 in conjunction with the inductive step and obtain

$$(7.8) \quad E_{k+2}[\psi](t_{i+1}) \leq C_{wp} E_{k+2}[\psi](t_i) \leq (C_{wp})^{i+1} E_{k+2}[\psi](0)$$

Therefore, by (7.6), (7.7), (7.8), it follows that (7.5) holds for t_{i+1} . Therefore, by induction (7.5) hold for all t_j , $j \in \mathbb{N}$.

Now we proceed to prove the the exponential decay (4.13), of $E_{\mathcal{G},k}[\psi]$ for all times $\tau \geq 0$. Note that for any $\tau \in \mathbb{R}$ there exists a $t_l \in \{t_i\}_{i \in \mathbb{N}}$, such that

$$(7.9) \quad |\tau - t_l| \leq \tau_{step}, \quad t_l < \tau.$$

We use the finite in time energy estimate (4.5), of Theorem 1, and the decay of the energy of the sequence $\{t_i\}$, see (7.5), and obtain the following

$$(7.10) \quad E_{\mathcal{G},k}[\psi](\tau) \leq C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) E_{\mathcal{G},k}[\psi](t_l) \leq C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) \left(\frac{1}{L}\right)^l E_{\mathcal{G},k}[\psi](0).$$

Now, by using one final time the finite in time energy estimate (4.5), of Theorem 1, we conclude the desired inequality, by noting

$$\begin{aligned}
(7.11) \quad E_{\mathcal{G},k}[\psi](\tau) &\leq C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) \left(\frac{1}{L}\right)^l E_{\mathcal{G},k}(0) \\
&\leq C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) e^{-l \log L} E_{\mathcal{G},k}[\psi](0) \\
&\leq C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) e^{-c_d \tau} E_{\mathcal{G},k}[\psi](0),
\end{aligned}$$

for $c_d(K, M, \Lambda, A_{[k+1]}, B_{[k+1]}) = \frac{\log L}{\tau_{step}}$, where in the last inequality we utilized that

$$(7.12) \quad t_l = (t_l - t_{l-1}) + (t_{l-1} - t_{l-2}) + (t_{l-2} - t_{l-3}) + \cdots + (t_2 - t_1) + t_1 = \tau_{step} l$$

so

$$(7.13) \quad e^{-l \log L} \leq e^{-\log L \frac{t_l}{\tau_{step}}} \leq e^{\log L - \frac{\log L}{\tau_{step}} \tau},$$

which concludes inequality (4.13).

The exponential decay of the lower order energy $E_{k+1}[\psi](\tau)$ follows by similar considerations, from inequality (7.6), which concludes (4.14), for a certain constant $c_d(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$.

Note that for the top order energy $E_{k+2}[\psi](\tau)$, we use (4.7) of Theorem 1 and (7.5), and we only obtain

$$(7.14) \quad E_{k+2}[\psi](\tau) \leq C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) e^{\frac{\log(C_{wp})}{\tau_{step}} \tau} E_{k+2}[\psi](0),$$

which concludes (4.15), for a constant $c_g(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) = \frac{\log(C_{wp})}{\tau_{step}} > 0$, where note that $C_{wp} > 1$, see Proposition 3.1.

The proof of Theorem 2 is thus complete. \square

8. THE SEMILINEAR EQUATION ON SCHWARZSCHILD–DE SITTER

We here present a simplified proof for the global non-linear stability for the semilinear case

$$(8.1) \quad \square_{g_{M,\Lambda}} \psi = \partial\psi \cdot \partial\psi,$$

with $\partial\psi \cdot \partial\psi = a^{ij} \partial_i \psi \partial_j \psi$, where a^{ij} are sufficiently regular components of a smooth tensor

$$(8.2) \quad a : TM \times TM \rightarrow \mathbb{R}$$

in Cartesian coordinates, see Section 2.7. The results here give a more direct proof and a better regularity for the initial data than the relevant Theorems 1, 2 that treat the more general quasilinear wave equation.

Remark 8.1. *Note that in this Section we study $D(\tau_1, \tau_2)$ instead of $D_\delta(\tau_1, \tau_2)$, as $\delta = 0$ here. One may easily a posteriori extend the results to $D_\delta(\tau_1, \tau_2)$. See Remark A.1.*

We give the statement of our Theorem, the analogue of Theorem 1, on a fixed large time domain.

Theorem 1'. *Let $k \geq 7$ and let the tensor a be as in Section 2.7.*

Then, there exists a $\tau_{step}(k, M, \Lambda, A_{[k-1]}) > 0$ sufficiently large and there exists

$$(8.3) \quad \epsilon(\tau_{step}, k, M, \Lambda, A_{[k-1]}) > 0$$

sufficiently small, where for $A_{[k]}$ see Section 2.7, such that if we take initial data for (8.1) on $\{\tau = \tau_1\}$ with

$$(8.4) \quad E_{\mathcal{G},k}[\psi](\tau_1) \leq \epsilon, \quad E_k[\psi](\tau_1) < \infty$$

and

$$(8.5) \quad \tau_2 = \tau_1 + \tau_{step},$$

then there exists a unique H^k solution in $D_\delta(\tau_1, \tau_2)$ to the semilinear wave equation (8.1), and the following inequality holds

$$(8.6) \quad E_{\mathcal{G},k}[\psi](\tau') \leq \frac{1}{2} E_{\mathcal{G},k}[\psi](\tau_1),$$

for all $\tau' \in [\tau_1, \tau_2]$.

A Corollary of Theorem 1', which is also the analogue of Theorem 2, is the following global existence and exponential decay result.

Theorem 2'. *Let $k \geq 7$ and let the tensor a be as in Section 2.7. Then, there exist positive constants*

$$(8.7) \quad c_d(k, M, \Lambda, A_{[k-1]}), \quad C(k, M, \Lambda, A_{[k-1]}) > 0$$

such that the following holds.

There exists an $\epsilon > 0$ sufficiently small, such that for

$$(8.8) \quad E_{\mathcal{G},k}[\psi](0) \leq \epsilon, \quad E_k[\psi](0) < \infty$$

the solution of the semilinear wave equation (8.1) exists globally on $D(0, \infty)$ and the following holds

$$(8.9) \quad E_{\mathcal{G},k}[\psi](\tau) \leq C e^{-c_d \tau} E_{\mathcal{G},k}[\psi](0),$$

for all $\tau \geq 0$.

Before proving Theorem 1' we need the following energy estimate on a fixed spacetime slab. This is the analogue of Proposition 3.1.

Proposition 8.1. *Let $k \geq 5$ and let the tensor a be as in Section 2.7. There exists a positive constant*

$$(8.10) \quad C(k, M, \Lambda, A_{[k-1]}) > 0,$$

where for $A_{[k-1]}$ see Section 2.7, such that the following holds.

Let ψ be a solution of the semilinear wave equation (8.1), on a Schwarzschild–de Sitter background $D(\tau_1, \tau_2)$ for $\tau_1 < \tau_2$. Then, if the following holds

$$(8.11) \quad \sup_{D(\tau_1, \tau_2)} \sum_{1 \leq i \leq \lfloor \frac{k-1}{2} \rfloor + 1} \sum_{\partial \in \{\partial_{\bar{t}}, \partial_r, \Omega_1, \Omega_2, \Omega_3\}} |\partial^i \psi| \leq \sqrt{\epsilon}$$

for a sufficiently small $\epsilon > 0$, then we obtain

$$(8.12) \quad E_{\mathcal{G},k}[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) \leq C E_{\mathcal{G},k}[\psi](\tau_1) + C \left(\sup_{\tau_1 \leq \tau' \leq \tau_2} E_{k-1}[\psi](\tau') \right) \left(\int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) \right),$$

for all $0 \leq \tau_1 \leq \tau_2$.

Proof of Proposition 8.1. We will use the derivatives notation of Section 2.6. Moreover, for the semilinear term $\partial\psi \cdot \partial\psi$ and for the definition of the tensor a see Section 2.7.

Let

$$(8.13) \quad F = \partial\psi \cdot \partial\psi$$

then by (2.48) we obtain

$$(8.14) \quad \begin{aligned} E_{\mathcal{G},k}[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) \\ \leq C(M, \Lambda) E_{\mathcal{G},k}[\psi](\tau_1) \\ + C(M, \Lambda) \int \int_{D(\tau_1, \tau_2)} \sum_{i+j=0}^{k-2} \sum_{\alpha} (1-\mu) (\partial_{\bar{t}}^i (\Omega_{\alpha})^j \mathcal{G}F)^2 + \sum_{0 \leq i_1+i_2+i_3 \leq k-2} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} (\Omega_{\alpha})^{i_3} F)^2 \\ + C(M, \Lambda) \int_{\{\bar{t}=\tau_2\}} \sum_{0 \leq i_1+i_2+i_3 \leq k-3} \sum_{\alpha} (1-\mu)^{2i_3+1} (\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \partial_r^{i_3} \mathcal{G}F)^2 + \sum_{0 \leq i_1+i_2+i_3 \leq k-3} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \partial_r^{i_3} F)^2. \end{aligned}$$

For the terms

$$(8.15) \quad \int \int_{D(\tau_1, \tau_2)} \sum_{i+j=0}^{k-2} \sum_{\alpha} (1-\mu) (\partial_{\bar{t}}^i (\Omega_{\alpha})^j \mathcal{G}F)^2$$

we have already computed in the proof of Proposition 5.1, see inequality (5.10), that

$$(8.16) \quad \begin{aligned} \int \int_{D(\tau_1, \tau_2)} \sum_{i+j=0}^{k-2} \sum_{\alpha} (1-\mu) (\partial_{\bar{t}}^i (\Omega_{\alpha})^j \mathcal{G}F)^2 \leq C(k, M, \Lambda, A_{[k-1]}) \int_{\tau_1}^{\tau_2} d\tau E_{k-1}[\psi](\tau) E_{\mathcal{G},k}[\psi](\tau) \\ \leq C(k, M, \Lambda, A_{[k-1]}) \sup_{\tau' \in [\tau_1, \tau_2]} E_{k-1}[\psi](\tau') \int_{\tau_1}^{\tau_2} E_{\mathcal{G},k}[\psi](\tau) d\tau. \end{aligned}$$

From the terms

$$(8.17) \quad \int \int_{D(\tau_1, \tau_2)} \sum_{0 \leq i_1+i_2+i_3 \leq k-2} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} (\Omega_{\alpha})^{i_3} F)^2$$

on the right hand side of (8.14), we only discuss the term

$$(8.18) \quad \int \int_{D(\tau_1, \tau_2)} \sum_{i=0}^{k-2} (\partial_{\bar{t}}^i F)^2$$

as the rest will admit the same bound. So, we bound the term (8.18), by using the coarea formula (2.29), as follows

$$\begin{aligned}
(8.19) \quad & \int \int_{D(\tau_1, \tau_2)} \sum_{i=0}^{k-2} (\partial_{\bar{t}}^i F)^2 = \int \int_{D(\tau_1, \tau_2)} \sum_{i=0}^{k-2} (\partial_{\bar{t}}^i (a^{\alpha\beta} \partial_\alpha \psi \partial_\beta \psi))^2 \\
& \leq C(k, M, \Lambda, A_{[k-1]}) \int \int_{D(\tau_1, \tau_2)} \sum_{1 \leq i+j \leq k-1} (\partial^i \psi)^2 (\partial^j \psi)^2 \\
& \leq C(k, M, \Lambda, A_{[k-1]}) \int_{\tau_1}^{\tau_2} d\tau \sum_{1 \leq i+j \leq k-1, i \leq \lfloor \frac{k-1}{2} \rfloor} \sup_{\{\bar{t}=\tau\}} (\partial^i \psi)^2 \int_{\{\bar{t}=\tau\}} (\partial^j \psi)^2 \\
& \leq C(k, M, \Lambda, A_{[k-1]}) \sup_{\tau_1 \leq \tau' \leq \tau_2} (E_{k-1}[\psi](\tau')) \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau),
\end{aligned}$$

where, in the last inequality, we used a Sobolev inequality, see Lemma 2.2, and the assumption that $k \geq 5$.

Therefore, by using (8.16), (8.19), inequality (8.14) implies

$$\begin{aligned}
(8.20) \quad & E_{\mathcal{G},k}[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) \\
& \leq C(k, M, \Lambda) E_{\mathcal{G},k}[\psi](\tau_1) \\
& + C(k, M, \Lambda, A_{[k-1]}) \left(\sup_{\tau_1 \leq \tau' \leq \tau_2} E_{k-1}(\tau') \right) \left(\int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) \right) \\
& + C(k, M, \Lambda) \int_{\{\bar{t}=\tau_2\}} \sum_{0 \leq i_1+i_2+i_3 \leq k-3} \sum_{\alpha} (1-\mu)^{2i_3+1} (\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \partial_r^{i_3} \mathcal{G}F)^2 + \sum_{0 \leq i_1+i_2+i_3 \leq k-3} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \partial_r^{i_3} F)^2,
\end{aligned}$$

where recall that $F = \partial\psi \cdot \partial\psi$.

Finally, we want to absorb the boundary terms of the right hand side at $\{\bar{t} = \tau_2\}$ by the relevant boundary term $E_{\mathcal{G},k}[\psi](\tau_2)$ of the left hand side. By using the smallness assumption (8.11), it is evident that we can appropriately distribute derivatives, in view of the restriction $k \geq 5$, to conclude that

$$(8.21) \quad E_{\mathcal{G},k}[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) \leq C E_{\mathcal{G},k}[\psi](\tau_1) + C \left(\sup_{\tau_1 \leq \tau' \leq \tau_2} E_{k-1}[\psi](\tau') \right) \left(\int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) \right),$$

for a constant $C = C(k, M, \Lambda, A_{[k]})$.

The proof of Proposition 8.1 is thus complete. \square

Now we are ready to prove Theorem 1'.

Proof of Theorem 1'. Note that the semilinear wave equation (8.1) is well posed in H^k , $k \geq 7$, by well-known arguments. The existence of the solution in $D(\tau_1, \tau_2)$, for τ_2 defined below, will follow easily from the estimates that we will prove, so for convenience we here assume existence.

Moreover, we assume the bootstrap assumption

$$(8.22) \quad \sup_{D(\tau_1, \tau_2)} \sum_{1 \leq i \leq \lfloor \frac{k-1}{2} \rfloor + 1} \sum_{\partial \in \{\partial_{\bar{t}}, \partial_r, \Omega_1, \Omega_2, \Omega_3\}} |\partial^i \psi| \leq C_b \sqrt{\epsilon}$$

for any $\tau \geq 0$, for a constant $C_b(k, M, \Lambda, A_{[k-1]}) > 0$ to be determined later.

First, we want to prove that for $\tau_{step} > 0$ to be chosen later, there exists an $\epsilon(\tau_{step}, C_b) > 0$ sufficiently small such that if

$$(8.23) \quad E_{\mathcal{G},k}[\psi](\tau_1) \leq \epsilon$$

for some $\tau_1 \geq 0$, then for

$$(8.24) \quad \tau_2 = \tau_1 + \tau_{step}$$

we obtain

$$(8.25) \quad \begin{aligned} E_{k-1}[\psi](\tau_2) &\leq \epsilon, \\ E_{\mathcal{G},k}[\psi](\tau_2) &\leq \frac{1}{2}E_{\mathcal{G},k}[\psi](\tau_1). \end{aligned}$$

Let $\tau_1 \geq 0$, then in view of the bootstrap assumption (8.22), and for $\epsilon(\tau_{step}, C_b)$ we obtain the result of Proposition 8.1, namely

$$(8.26) \quad E_{\mathcal{G},k}[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) \leq CE_{\mathcal{G},k}[\psi](\tau_1) + C \left(\sup_{\tau_1 \leq \tau' \leq \tau_2} E_{k-1}[\psi](\tau') \right) \left(\int_{\tau_1}^{\tau_2} d\tau E_{\mathcal{G},k}[\psi](\tau) \right)$$

for some constant $C = C(k, M, \Lambda, A_{[k-1]})$. Now, for a sufficiently small $\epsilon(\tau_{step})$ if we assume that

$$(8.27) \quad E_{\mathcal{G},k}[\psi](\tau_1) \leq \epsilon,$$

we also have that $E_{k-1}[\psi](\tau_1) \leq \epsilon$, and from (8.26) we conclude that there exists a constant

$$(8.28) \quad C(k, M, \Lambda, A_{[k-1]}) > 0$$

where for $A_{[k-1]}$ see Section 2.7, such that

$$(8.29) \quad E_{\mathcal{G},k}[\psi](\tau') \leq C(k, M, \Lambda, A_{[k-1]})E_{\mathcal{G},k}[\psi](\tau_1)$$

for all $\tau' \in [\tau_1, \tau_2]$.

Moreover, by using again the result of Proposition 8.1 (also see the relevant computation in the proof of Theorem 1), there exists a constant $C(k, M, \Lambda, A_{[k-1]}) > 0$ such that for any $\tau_{step} > 0$ sufficiently large, there exists an $\epsilon(\tau_{step}) > 0$ sufficiently small, and a value

$$(8.30) \quad \tau_{1,2} \in \left[\frac{\tau_1 + \tau_2}{2}, \tau_2 \right]$$

such that, after taking

$$(8.31) \quad E_{\mathcal{G},k}[\psi](\tau_1) \leq \epsilon$$

and recalling $E_{k-1}[\psi](\tau_1) \leq E_{\mathcal{G},k}[\psi](\tau_1)$, we obtain

$$(8.32) \quad E_{\mathcal{G},k}[\psi](\tau_{1,2}) \leq \frac{1}{\tau_{step}} C(k, M, \Lambda, A_{[k-1]}) E_{\mathcal{G},k}[\psi](\tau_1),$$

where we used the mean value theorem. By using (8.29) and (8.32) we obtain

$$(8.33) \quad E_{\mathcal{G},k}[\psi](\tau_2) \leq \frac{1}{\tau_{step}} C(k, M, \Lambda, A_{[k-1]}) E_{\mathcal{G},k}[\psi](\tau_1)$$

for $\tau_{step} > 0$ and $\epsilon(\tau_{step}, C_b) > 0$ sufficiently small, and for a different constant $C(k, M, \Lambda, A_{[k-1]})$. Now, note that

$$(8.34) \quad E_{k-1}[\psi](\tau_2) \leq E_{\mathcal{G},k}[\psi](\tau_2) \leq \frac{1}{\tau_{step}} C(k, M, \Lambda, A_{[k-1]}) E_{\mathcal{G},k}[\psi](\tau_1).$$

Therefore, we choose a τ_{step} sufficiently large, such that

$$(8.35) \quad \frac{1}{\tau_{step}} C(k, M, \Lambda, A_{[k-1]}) < \frac{1}{2}$$

and we obtain

$$(8.36) \quad \begin{aligned} E_{k-1}[\psi](\tau_2) &\leq E_{\mathcal{G},k}[\psi](\tau_1) \leq \epsilon, \\ E_{\mathcal{G},k}[\psi](\tau_2) &\leq \frac{1}{2}E_{\mathcal{G},k}[\psi](\tau_1). \end{aligned}$$

Of course our inequalities are still conditional on improving the bootstrap assumption (8.22). For the purpose of improving constant $C_b > 0$ of the bootstrap assumption (8.22) we note from (8.29) that

$$(8.37) \quad E_{k-1}[\psi](\tau') \leq C(k, M, \Lambda, A_{[k-1]}) E_{\mathcal{G},k}[\psi](\tau_1) \leq C(k, M, \Lambda, A_{[k-1]})\epsilon,$$

for all $\tau' \in [\tau_1, \tau_2]$. Now, since $k = 7$ is the smallest integer such that

$$(8.38) \quad \left(\lfloor \frac{k-1}{2} \rfloor + 1 \right) + 2 \leq k - 1$$

we use a Sobolev inequality in (8.37) and prove

$$(8.39) \quad \sup_{D(0,\tau)} \sum_{1 \leq i \leq \lfloor \frac{k-1}{2} \rfloor + 1} \sum_{\partial \in \{\partial_t, \partial_r, \Omega_1, \Omega_2, \Omega_3\}} |\partial^i \psi| \leq C(k, M, \Lambda, A_{[k-1]}) \sqrt{\epsilon}$$

which improves the bootstrap for a sufficiently large $C_b \gg C(k, M, \Lambda, A_{[k-1]})$.

This concludes the proof the Theorem. \square

Now, we are ready to prove Theorem 2', which is in fact a Corollary of Theorem 1'.

Proof of Theorem 2'. We want to prove that for sufficiently small initial data

$$(8.40) \quad E_{\mathcal{G},k}[\psi](0) \leq \epsilon,$$

and for the following sequence of ascending real numbers

$$(8.41) \quad t_0 = \tau_{step}, \quad t_{i+1} - t_i = \tau_{step},$$

the solution exists in $D(0, t_i)$ and we obtain that

$$(8.42) \quad \begin{aligned} E_{k-1}[\psi](t_i) &\leq C(k, M, \Lambda)\epsilon, \\ E_{\mathcal{G},k}[\psi](t_i) &\leq \left(\frac{1}{2}\right)^i E_{\mathcal{G},k}[\psi](0), \end{aligned}$$

for all $i \in \mathbb{N}$, where for the constant $C(k, M, \Lambda)$ see Remark 2.5.

Note that the inequalities (8.42) hold for t_0 . Therefore, for the purpose of using induction we assume the inductive step that (8.42) hold for t_i . Then, from Theorem 1', specifically from inequality (8.6) and the inductive step we obtain

$$(8.43) \quad \begin{aligned} E_{k-1}[\psi](t_{i+1}) &\leq C(k, M, \Lambda)\epsilon, \\ E_{\mathcal{G},k}[\psi](t_{i+1}) &\leq \left(\frac{1}{2}\right)^{i+1} E_{\mathcal{G},k}[\psi](0), \end{aligned}$$

and moreover conclude that the solution exists in $D(0, t_{i+1})$.

Therefore, we have concluded global existence and the inequalities (8.42) for all $i \in \mathbb{N}$.

Now, we want to conclude the result of the Theorem, namely we want to prove that for

$$(8.44) \quad E_{\mathcal{G},k}[\psi](0) \leq \epsilon$$

we obtain

$$(8.45) \quad E_{\mathcal{G},k}[\psi](\tau) \leq C e^{-c_d \tau} E_{\mathcal{G},k}[\psi](0),$$

for constants $c_d(k, M, \Lambda, A_{[k-1]})$, $C(k, M, \Lambda, A_{[k-1]}) > 0$. With initial data

$$(8.46) \quad E_{\mathcal{G},k}[\psi](0) \leq \epsilon$$

for a sufficiently small $\epsilon > 0$, we use equation (8.42) and the local in time inequality (8.6) of Theorem 1', to immediately conclude (8.45).

We conclude the Theorem. \square

APPENDIX A. PROOF OF THE CAUCHY STABILITY RESULT OF PROPOSITION 3.1

In this Section we prove the Cauchy stability statement of Proposition 3.1. This proof will require using geometric estimates with respect to the metric $g(\nabla\psi)$, so as to not lose derivatives. We will use some standard definitions and results of the linear theory. Therefore, we first need to discuss some standard notions.

A.1. Energy momentum tensor and the divergence theorem of a metric g . Let g be a sufficiently regular Lorentzian metric, and ψ a sufficiently regular function. Then, we define the energy momentum tensor with respect to that metric as

$$(A.1) \quad \mathbb{T}(g)[\psi] = d\psi \otimes d\psi - \frac{1}{2}g|\nabla\psi|_g^2,$$

where $|\nabla\psi|_g^2 \doteq g^{ab}\partial_a\psi\partial_b\psi$.

Note the following Proposition

Proposition A.1. *Let g be a sufficiently regular Lorentzian metric on \mathcal{M}_δ and $\delta(M, \Lambda) \geq 0$ sufficiently small. Also, let ψ satisfy the equation*

$$(A.2) \quad \square_g\psi = F.$$

Then, we obtain the following identity

$$(A.3) \quad \int_{\{\bar{t}=\tau_2\} \cap D_\delta(\tau_1, \tau_2)} \mathbb{T}(g)(X, n)[\psi] + \int_{\mathcal{H}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} \mathbb{T}(g)(X, n)[\psi] + \int_{\bar{\mathcal{H}}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} \mathbb{T}(g)(X, n)[\psi] \\ + \int_{D_\delta(\tau_1, \tau_2)} K^X(g) = \int_{\{\bar{t}=\tau_1\} \cap D_\delta(\tau_1, \tau_2)} \mathbb{T}(g)(X, n)[\psi] + \int_{D_\delta(\tau_1, \tau_2)} Err^X[\psi]$$

for any $0 \leq \tau_1 \leq \tau_2$ and for any vector field X , where

$$(A.4) \quad K^X(g) = \frac{1}{2} \binom{X}{\pi_{\mu\nu}}(g) \mathbb{T}^{\mu\nu}(g)[\psi], \quad Err^X[\psi] = -(X\psi)F,$$

where note that the raised indices are with the metric g .

The volume forms on the spacetime domains are to be understood with respect to the metric g , and the volume forms on the hypersurfaces are to be understood as the ones induced by the volume form of g . Also

$$(A.5) \quad n$$

is to be understood as the unit normal of each hypersurface with respect to g .

Note that in the case $g = g_{M, \Lambda}$, $\delta \geq 0$, we have explicitly computed the volume forms and the normals, with which the divergence theorem is to be understood, in Section 2.3.

A.2. The redshift vector field N on the Schwarzschild background \dot{g} . We present the following redshift Lemmata, see the Lecture notes [12].

Lemma A.1. *Let $q(M, \Lambda) > 0$ be sufficiently small. Then, for all $\delta(M, \Lambda) > 0$ sufficiently small there exist a timelike vector field*

$$(A.6) \quad N,$$

on $D_\delta(0, \infty)$, such that the following hold

$$(A.7) \quad K^N(\dot{g}) \geq C(M, \Lambda) \mathbb{T}(\dot{g})(N, N)[\psi], \quad \text{in } \{r_+ - \delta \leq r \leq r_+ + q\} \cup \{\bar{r}_+ - q \leq r \leq \bar{r}_+ + \delta\} \\ N = \partial_{\bar{t}}, \quad \text{in } D_\delta(0, \infty) \setminus \{r_+ - \delta \leq r \leq r_+ + q\} \cup \{\bar{r}_+ - q \leq r \leq \bar{r}_+ + \delta\}.$$

where for $K^N(\dot{g})$ see the divergence theorem in Proposition A.1.

The following Lemma, is a higher order manifestation of the redshift effect.

Lemma A.2. *For any $k \geq 0$ there exist positive constants*

$$(A.8) \quad \kappa_k > 0, \quad \bar{\kappa}_k > 0$$

such that for the equation

$$(A.9) \quad \square_{\dot{g}} \psi = F,$$

we obtain that the following holds on \mathcal{H}^+

$$(A.10) \quad \square_{\dot{g}} N^k \psi = \kappa_k N^{k+1} \psi + \sum_{1 \leq i \leq 5} \sum_{0 \leq \sum_i m_i \leq |m| \leq k+1, m_5 \leq k} c_m \Omega_1^{m_1} \Omega_2^{m_2} \Omega_3^{m_3} \partial_{\bar{t}}^{m_4} N^{m_5} \psi + N^k F,$$

where c_m are smooth functions on \mathcal{H}^+ , and the following holds on $\bar{\mathcal{H}}^+$

$$(A.11) \quad \square_{\dot{g}} N^k \psi = \bar{\kappa}_k N^{k+1} \psi + \sum_{1 \leq i \leq 5} \sum_{0 \leq \sum_i m_i \leq |m| \leq k+1, m_5 \leq k} \bar{c}_m \Omega_1^{m_1} \Omega_2^{m_2} \Omega_3^{m_3} \partial_{\bar{t}}^{m_4} N^{m_5} \psi + N^k F,$$

where \bar{c}_m are smooth functions on $\bar{\mathcal{H}}^+$.

A.3. An estimate for metrics close to \dot{g} . Note the following lemma

Lemma A.3. *Let $g(v) = \dot{g} + h(v)$ be a Lorentzian metric that belongs in the class of Section 3.1, where $v \in \Gamma(TM_\delta)$ is considered fixed, and let $\delta(M, \Lambda) > 0$ be sufficiently small.*

Then there exists an $\epsilon(\delta) > 0$ such that for

$$(A.12) \quad |h_{ij}(v)|, \quad |h^{ij}(v)| \leq \sqrt{\epsilon}$$

in Cartesian coordinates, the hypersurface $\{\bar{t} = \tau\}$, is a Cauchy hypersurface, with respect to $g(v)$, in $D_\delta(0, \infty)$. Moreover, the unit normal n of $\{\bar{t} = c\}$ is timelike on $D_\delta(0, \infty)$ with respect to $g(v)$ and the redshift vector field, see Lemma A.1, is timelike with respect to $g(v)$, as well. Furthermore, the hypersurfaces

$$(A.13) \quad \mathcal{H}_\delta^+, \quad \bar{\mathcal{H}}_\delta^+$$

are spacelike hypersurfaces with respect to the metric $g(v)$ in $D_\delta(0, \infty)$.

Finally, we obtain

$$(A.14) \quad \begin{aligned} \int_{\{\bar{t}=\tau\}} \mathbb{T}(g(v))(n, n)[\psi] dg(v)_{\{\bar{t}=\tau\}} &\sim \int_{\{\bar{t}=\tau\}} \mathbb{T}(\dot{g})(\dot{n}, \dot{n})[\psi] d\dot{g}_{\{\bar{t}=\tau\}}, \\ \int_{\mathcal{H}_\delta^+} \mathbb{T}(g(v))(N, n)[\psi] dg(v)_{\mathcal{H}_\delta^+} &\sim \int_{\mathcal{H}_\delta^+} \frac{d\dot{g}_{\mathcal{H}_\delta^+}}{\sqrt{|1-\mu|}} (\delta(\partial_r \psi)^2 + (\partial_{\bar{t}} \psi)^2 + |\nabla \psi|^2), \\ \int_{\bar{\mathcal{H}}_\delta^+} \mathbb{T}(g(v))(N, n)[\psi] dg(v)_{\bar{\mathcal{H}}_\delta^+} &\sim \int_{\bar{\mathcal{H}}_\delta^+} \frac{d\dot{g}_{\bar{\mathcal{H}}_\delta^+}}{\sqrt{|1-\mu|}} (\delta(\partial_r \psi)^2 + (\partial_{\bar{t}} \psi)^2 + |\nabla \psi|^2), \end{aligned}$$

where the constants only depend on the black hole parameters, and $dg(v)_{\{\bar{t}=\tau\}}, dg(v)_{\mathcal{H}_\delta^+}, dg(v)_{\bar{\mathcal{H}}_\delta^+}$ are the induced volume form of the spacetime volume form of the metric $g(v)$ on the hypersurfaces $\{\bar{t} = \tau\}, \mathcal{H}_\delta^+, \bar{\mathcal{H}}_\delta^+$, respectively. For the energy momentum tensor \mathbb{T} see Section A.1, for the redshift vector field N see Lemma A.1.

Proof. The proof of this Lemma is direct by noting that the energy momentum tensor of the metric $g(v)$ is

$$(A.15) \quad \mathbb{T}(g(v))(X, Y)[\psi] = X\psi Y\psi - \frac{1}{2}g(v)(X, Y)|\nabla\psi|_{g(v)}^2,$$

for any two smooth vector fields X, Y .

Specifically, by taking

$$(A.16) \quad |h_{ij}(v)|, \quad |h^{ij}(v)| \leq \sqrt{\epsilon}$$

sufficiently small, we conclude the causal behaviour of the hypersurfaces mentioned, and by pointwise estimates on the integrands we conclude

$$(A.17) \quad \int_{\{\bar{t}=\tau\}} \mathbb{T}(g(v))(n, n)[\psi] dg(v)_{\{\bar{t}=\tau\}} \sim \int_{\{\bar{t}=\tau\}} \mathbb{T}(\hat{g})(\hat{n}, \hat{n})[\psi] d\hat{g}_{\{\bar{t}=\tau\}},$$

$$\int_{\mathcal{H}_\delta^+} \mathbb{T}(g(v))(N, n)[\psi] dg(v)_{\mathcal{H}_\delta^+} \sim \int_{\mathcal{H}_\delta^+} \frac{d\hat{g}_{\mathcal{H}_\delta^+}}{\sqrt{|1-\mu|}} \left(\delta(\partial_r \psi)^2 + (\partial_{\bar{t}} \psi)^2 + |\nabla \psi|^2 + \sqrt{\epsilon(\delta)} \left((\partial_r \psi)^2 + (\partial_{\bar{t}} \psi)^2 + |\nabla \psi|^2 \right) \right),$$

$$\int_{\bar{\mathcal{H}}_\delta^+} \mathbb{T}(g(v))(N, n)[\psi] dg(v)_{\bar{\mathcal{H}}_\delta^+} \sim \int_{\bar{\mathcal{H}}_\delta^+} \frac{d\hat{g}_{\bar{\mathcal{H}}_\delta^+}}{\sqrt{|1-\mu|}} \left(\delta(\partial_r \psi)^2 + (\partial_{\bar{t}} \psi)^2 + |\nabla \psi|^2 + \sqrt{\epsilon(\delta)} \left((\partial_r \psi)^2 + (\partial_{\bar{t}} \psi)^2 + |\nabla \psi|^2 \right) \right).$$

Now, for $\epsilon(\delta) \ll \delta$ we conclude (A.14). \square

A.4. Elliptic estimates and estimates on \mathcal{H}_δ^+ , $\bar{\mathcal{H}}_\delta^+$ for the quasilinear wave equation. Note the following elliptic estimate

Lemma A.4. *Let $k \geq 4$ and let ψ satisfy the quasilinear wave equation (1.1) in $D_\delta(\tau_1, \tau_2)$ for $\tau_1 \leq \tau_2$ and for a sufficiently small $\delta(M, \Lambda) > 0$, where for the tensors a, h see the Sections 2.7, 3.1 respectively. Then, for any $r_+ < r_0 < r_1 < \bar{r}_+$, there exist constants*

$$(A.18) \quad C(r_0, r_1, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}), \quad C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$$

independent of δ , where for $A_{[k+1]}, B_{[k+1]}$ see Sections 2.7, 3.1 respectively, and there exists an $\epsilon = \epsilon(\delta, k, M, \Lambda, A_{[k+1]}, B_{[k+1]})$ sufficiently small such that the following holds.

If

$$(A.19) \quad \sup_{D_\delta(\tau_1, \tau_2)} \sum_{1 \leq j \leq k-1} \sum_{\partial \in \{\partial_{\bar{t}}, \partial_r, \Omega_1, \Omega_2, \Omega_3\}} |\partial^j \psi| \leq \sqrt{\epsilon}$$

holds, we obtain that

$$(A.20) \quad \int_{\{\bar{t}=\tau'\}} \sum_{1 \leq i_1+i_2+i_3 \leq j} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_\alpha^{i_3} \psi)^2 d\hat{g}_{\{\bar{t}=\tau'\}} \leq C \int_{\{\bar{t}=\tau\}} \sum_{1 \leq i \leq j-1} \mathbb{T}(\hat{g})(N, n)[N^i \psi] d\hat{g}_{\{\bar{t}=\tau\}},$$

$$\int_{\{\bar{t}=\tau'\} \cap [r_0, r_1]} \sum_{1 \leq i_1+i_2+i_3 \leq j} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_\alpha^{i_3} \psi)^2 d\hat{g}_{\{\bar{t}=\tau\}} \leq C(r_0, r_1) \int_{\{\bar{t}=\tau\} \cap [r_0, r_1]} \sum_{1 \leq i_1+i_2 \leq j} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \psi)^2 d\hat{g}_{\{\bar{t}=\tau\}},$$

$$\int_{\mathcal{H}_\delta^+} \frac{d\hat{g}_{\mathcal{H}_\delta^+}}{\sqrt{|1-\mu|}} \left(\delta \sum_{1 \leq i_1+i_2+i_3 \leq j, i_2 \geq 1} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_\alpha^{i_3} \psi)^2 + \sum_{1 \leq i_1+i_2 \leq j} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \psi)^2 \right) \leq C \int_{\mathcal{H}_\delta^+} \sum_{1 \leq i \leq j-1} \mathbb{T}(\hat{g})(N, n)[N^i \psi] d\hat{g}_{\mathcal{H}_\delta^+},$$

$$\int_{\bar{\mathcal{H}}_\delta^+} \frac{d\hat{g}_{\bar{\mathcal{H}}_\delta^+}}{\sqrt{|1-\mu|}} \left(\delta \sum_{1 \leq i_1+i_2+i_3 \leq j, i_2 \geq 1} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_\alpha^{i_3} \psi)^2 + \sum_{1 \leq i_1+i_2 \leq j} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \Omega_\alpha^{i_2} \psi)^2 \right) \leq C \int_{\bar{\mathcal{H}}_\delta^+} \sum_{1 \leq i \leq j-1} \mathbb{T}(\hat{g})(N, n)[N^i \psi] d\hat{g}_{\bar{\mathcal{H}}_\delta^+},$$

for $1 \leq j \leq k+2$.

Proof. The proof of this Lemma comes from the estimates of Lemma A.3.

For the first inequality of (A.20), for $j = 1$, we use the smallness (A.19), and Lemma A.3. We prove the estimate for all orders j , by first noting that the energy momentum tensor of $g(\nabla \psi)$ is

$$(A.21) \quad \mathbb{T}(g(\nabla \psi))(X, n)[\tilde{X}^i \psi] = \mathbb{T}(\hat{g})(X, n)[\tilde{X}^i \psi] - \frac{1}{2} h(X, n, \nabla \psi) |\nabla \tilde{X}^i \psi|_{g(\nabla \psi)}^2,$$

for any two smooth vector field X, \tilde{X} . Therefore, for $\epsilon(\delta) > 0$ sufficiently small, we use elliptic estimates on the difference $\square_{\hat{g}} \psi = (\square_{\hat{g}} - \square_{g(\nabla \psi)}) \psi + \partial \psi \cdot \partial \psi$ and conclude. We also used that $k \geq 4$.

For the second inequality of (A.20), we obtain the estimate by arguing as above and by taking $\epsilon(\delta) > 0$ small. We also used that $k \geq 4$.

For the the last two inequalities of (A.20), on the hypersurfaces $\mathcal{H}_\delta^+, \bar{\mathcal{H}}_\delta^+$, to get the estimate for $j = 1$, we use the smallness (A.19), and Lemma A.3. We prove the analogous estimate for all orders j , for a sufficiently small $\epsilon(\delta) > 0$ by noting the form of the energy momentum tensor (A.21), and

by using elliptic estimates on the difference $\square_{\dot{g}}\psi = (\square_{\dot{g}} - \square_{g(\nabla\psi)})\psi + \partial\psi \cdot \partial\psi$. We also used that $k \geq 4$. \square

Finally, note the following estimate

Lemma A.5. *Let $k \geq 4$. Let ψ satisfy the quasilinear wave equation (1.1) in $D_\delta(\tau_1, \tau_2)$ for a sufficiently small $\delta(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$, where for the tensors a, h see the Sections 2.7, 3.1 respectively*

Then, there exists a constant

$$(A.22) \quad C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$$

independent of δ , and an $\epsilon = \epsilon(\delta, k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$ sufficiently small such that the following holds. If

$$(A.23) \quad \sup_{D_\delta(\tau_1, \tau_2)} \sum_{1 \leq j \leq k-1} \sum_{\partial \in \{\partial_{\bar{t}}, \partial_r, \Omega_1, \Omega_2, \Omega_3\}} |\partial^j \psi| \leq \sqrt{\epsilon},$$

then we obtain that

$$(A.24) \quad \begin{aligned} & \left| \int_{\mathcal{H}_\delta^+} \sum_{1 \leq i_1 + i_2 \leq j-1} \sum_{\alpha} \mathbb{T}(\dot{g})(\partial_{\bar{t}}, n) [\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi] d\dot{g}_{\mathcal{H}_\delta} - \int_{\mathcal{H}_\delta^+} \sum_{0 \leq i_1 + i_2 \leq j-1} \sum_{\alpha} (\partial_{\bar{t}} \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi)^2 \frac{1}{\sqrt{|1-\mu|}} d\dot{g}_{\mathcal{H}_\delta} \right| \\ & \leq C\delta \int_{\mathcal{H}_\delta^+} \sum_{1 \leq i \leq j-1} \mathbb{T}(\dot{g})(N, n) [N^i \psi] d\dot{g}_{\mathcal{H}_\delta}, \\ & \left| \int_{\mathcal{H}_\delta^+} \sum_{1 \leq i_1 + i_2 \leq j-1} \sum_{\alpha} \mathbb{T}(\dot{g})(\partial_{\bar{t}}, n) [\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi] d\dot{g}_{\mathcal{H}_\delta} - \int_{\mathcal{H}_\delta^+} \sum_{0 \leq i_1 + i_2 \leq j-1} \sum_{\alpha} (\partial_{\bar{t}} \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi)^2 \frac{1}{\sqrt{|1-\mu|}} d\dot{g}_{\mathcal{H}_\delta} \right| \\ & \leq C\delta \int_{\mathcal{H}_\delta^+} \sum_{1 \leq i \leq j-1} \mathbb{T}(\dot{g})(N, n) [N^i \psi] d\dot{g}_{\mathcal{H}_\delta}, \end{aligned}$$

for $1 \leq j \leq k+2$.

Proof. We note that for for the metric \dot{g} , we obtain

$$(A.25) \quad \begin{aligned} \mathbb{T}(\dot{g})(\partial_{\bar{t}}, \hat{n}_{\mathcal{H}_\delta}) [\partial_{\bar{t}}^j \psi] &= \left((\partial_{\bar{t}}^{j+1} \psi)^2 \left(c_1(r) - \frac{1}{2} \dot{g}_{\bar{t}\bar{t}} \dot{g}^{\bar{t}\bar{t}} - \frac{1}{2} c_2(r) \dot{g}_{\bar{t}r} \dot{g}^{\bar{t}r} \right) \right. \\ & \quad + \partial_{\bar{t}} \partial_{\bar{t}}^j \psi \partial_r \partial_{\bar{t}}^j \psi \left(c_2(r) - c_1(r) \dot{g}_{\bar{t}\bar{t}} \dot{g}^{\bar{t}r} - c_2(r) \dot{g}_{\bar{t}r} \dot{g}^{\bar{t}r} \right) \\ & \quad + |\nabla \partial_{\bar{t}}^j \psi|^2 \left(-\frac{1}{2} c_1(r) g_{\bar{t}\bar{t}} - \frac{1}{2} c_2(r) g_{\bar{t}r} \right) \\ & \quad \left. + (\partial_r \partial_{\bar{t}}^j \psi)^2 \left(-\frac{1}{2} c_1(r) \dot{g}_{\bar{t}\bar{t}} \dot{g}^{rr} - \frac{1}{2} c_2(r) \dot{g}_{\bar{t}r} \dot{g}^{rr} \right) \right), \end{aligned}$$

where for $\hat{n}_{\mathcal{H}_\delta}, c_1(r), c_2(r)$ see Section 2.3, and specifically $c_1(r) = \frac{1}{\sqrt{|1-\mu|}} + \mathcal{O}(\sqrt{|1-\mu|})$ as $r \rightarrow r_+$ and $c_2(r) = -\sqrt{|1-\mu|}$. At $r = r_+ - \delta$ the following hold

$$(A.26) \quad \begin{aligned} & \left| -\frac{1}{2} \dot{g}_{\bar{t}\bar{t}} \dot{g}^{\bar{t}\bar{t}} - \frac{1}{2} c_2(r) \dot{g}_{\bar{t}r} \dot{g}^{\bar{t}\bar{t}} \right| \sim \sqrt{|1-\mu|} \sim \delta^{1/2}, \\ & \left| c_2(r) - c_1(r) \dot{g}_{\bar{t}\bar{t}} \dot{g}^{\bar{t}r} - c_2(r) \dot{g}_{\bar{t}r} \dot{g}^{\bar{t}r} \right| \sim \sqrt{|1-\mu|} \sim \delta^{1/2}, \\ & \left| -\frac{1}{2} c_1(r) g_{\bar{t}\bar{t}} - \frac{1}{2} c_2(r) g_{\bar{t}r} \right| \sim \sqrt{|1-\mu|} \sim \delta^{1/2}, \\ & \left| -\frac{1}{2} c_1(r) \dot{g}_{\bar{t}\bar{t}} \dot{g}^{rr} - \frac{1}{2} c_2(r) \dot{g}_{\bar{t}r} \dot{g}^{rr} \right| \sim |1-\mu|^{3/2} \sim \delta^{3/2}, \end{aligned}$$

see Remark 2.3 for the inverse metric components. A similar estimate holds for the hypersurface $\bar{\mathcal{H}}_\delta^+$, since $\hat{n}_{\bar{\mathcal{H}}_\delta^+}$ has a favorable form, see Section 2.3. We obtain the linear version of the Lemma.

For the quasilinear version we use the smallness (A.23) to obtain

$$\begin{aligned}
 (A.27) \quad & \left| \int_{\mathcal{H}_\delta^+} \sum_{1 \leq i_1+i_2 \leq j-1} \sum_{\alpha} \mathbb{T}(\partial_{\bar{t}}, \dot{n}_{\mathcal{H}_\delta}) [\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi] d\dot{g}_{\mathcal{H}_\delta} - \int_{\mathcal{H}_\delta^+} \sum_{0 \leq i_1+i_2 \leq j-1} \sum_{\alpha} (\partial_{\bar{t}} \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi)^2 \frac{1}{\sqrt{|1-\mu|}} d\dot{g}_{\mathcal{H}_\delta} \right| \\
 & \leq C\sqrt{\delta} \int_{\mathcal{H}_\delta^+} \sum_{1 \leq i \leq j-1} \mathbb{T}(N, \dot{n}_{\mathcal{H}_\delta}) [N^i \psi] d\dot{g}_{\mathcal{H}_\delta} + \sqrt{\epsilon(\delta)} \int_{\mathcal{H}_\delta^+} \sum_{1 \leq i_1+i_2+i_3 \leq j} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_{\alpha}^{i_3} \psi)^2 \frac{d\dot{g}_{\mathcal{H}_\delta}}{\sqrt{|1-\mu|}}, \\
 & \left| \int_{\mathcal{H}_\delta^+} \sum_{1 \leq i_1+i_2 \leq j-1} \sum_{\alpha} \mathbb{T}(\partial_{\bar{t}}, \dot{n}_{\mathcal{H}_\delta}) [\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi] d\dot{g}_{\mathcal{H}_\delta} - \int_{\mathcal{H}_\delta^+} \sum_{0 \leq i_1+i_2 \leq j-1} \sum_{\alpha} (\partial_{\bar{t}} \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi)^2 \frac{1}{\sqrt{|1-\mu|}} d\dot{g}_{\mathcal{H}_\delta} \right| \\
 & \leq C\sqrt{\delta} \int_{\mathcal{H}_\delta^+} \sum_{1 \leq i \leq j-1} \mathbb{T}(N, \dot{n}_{\mathcal{H}_\delta}) [N^i \psi] d\dot{g}_{\mathcal{H}_\delta} + \sqrt{\epsilon(\delta)} \int_{\mathcal{H}_\delta^+} \sum_{1 \leq i_1+i_2+i_3 \leq j} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_{\alpha}^{i_3} \psi)^2 \frac{d\dot{g}_{\mathcal{H}_\delta}}{\sqrt{|1-\mu|}},
 \end{aligned}$$

for all $1 \leq j \leq k+2$, where $C = C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]})$. Therefore, for a sufficiently small δ and $\epsilon(\delta) \ll \delta$, we use Lemma A.4 and conclude. \square

A.5. Proof of Proposition 3.1. Now we proceed to the proof of the Proposition.

Proof. Let $\tau_1 \geq 0$.

We will use the derivatives notation of Section 2.6. Moreover, for the semilinear term $\partial\psi \cdot \partial\psi$ and for the definition of smooth tensors a, h , see Sections 2.7, 3.1 respectively. In this proof, when we write

$$(A.28) \quad h_{ab}$$

it is to be understood $h_{ab}(\nabla\psi)$ in Cartesian coordinates, see (2.38). When we write

$$(A.29) \quad h^{ab}$$

it is to be understood as $h^{ab}(\nabla\psi) = g^{ab}(\nabla\psi) - \dot{g}^{ab}$, in Cartesian coordinates. Finally, in this proof when we write \lesssim it is to be understood that we omit a constant $C(k, M, \Lambda, A_{[m]}, B_{[m]})$, where for $A_{[m]}, B_{[m]}$ see Sections 2.7, 3.1 respectively, where $m \leq k+1$.

Existence and uniqueness.

The existence and uniqueness part of the proof follow readily, by well known arguments, from the energy estimates (3.9), (3.10). Therefore, we only prove (3.9), (3.10), assuming existence.

An auxiliary energy.

We here introduce an auxiliary energy with which we will estimate the $\partial_{\bar{t}}$ flux on $\{\bar{t} = \tau\}$, see already (A.52). We define

$$(A.30) \quad E_{j,q}[\psi](\tau) = \int_{\{\bar{t}=\tau\} \cap \{r_++q \leq r \leq \bar{r}_+-q\}} \sum_{1 \leq i_1+i_2 \leq j} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi)^2 + (1-\mu) \sum_{1 \leq i_1+i_2+i_3 \leq j} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_{\alpha}^{i_3} \psi)^2 d\dot{g}_{\{\bar{t}=\tau\}},$$

for $q > 0$ as in the redshift Lemma A.1.

The linear wave equation analogue

Before proving the Cauchy stability estimate (3.9), (3.10) for the solutions of the quasilinear wave equation, we sketch the proof of (3.9), (3.10) for the solutions of the linear wave equation

$$(A.31) \quad \square_{\dot{g}} \psi = 0,$$

so that the reader can see the similarity between the two proofs. Also see the Lecture notes [12]. The steps of the proof of the relevant quasilinear estimate are similar, *but by also treating the non-linearities appropriately.*

First, we apply Proposition A.1 with multiplier $\partial_{\bar{t}}$ and obtain that there exists a constant $C(M, \Lambda) > 1$ such that

$$(A.32) \quad E_{1,q}[\psi](\tau_2) \leq C(M, \Lambda) E_1[\psi](\tau_1)$$

for all $\tau_1 \leq \tau_2$. By commuting the wave equation appropriately many times with $\partial_{\bar{t}}\Omega_\alpha$ and by using elliptic estimates, then (A.32) implies

$$(A.33) \quad E_{j,q}[\psi](\tau_2) \leq C(j, M, \Lambda)E_j[\psi](\tau_1)$$

for all $j \geq 1$, for some constant $C(j, M, \Lambda) > 1$.

Second, we apply the divergence Theorem with multiplier the red-shift vector field N , see Lemma A.1, and obtain

$$(A.34) \quad E_1[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_1[\psi](\tau) \leq C(j, M, \Lambda) \int_{\tau_1}^{\tau_2} E_{1,q}[\psi](\tau) + C(j, M, \Lambda)E_1[\psi](\tau_1),$$

for all $\tau_1 \leq \tau_2$. By commuting the wave equation $j-1$ times with N , also see the redshift Lemma A.2 and by using elliptic estimates, then (A.34) implies

$$(A.35) \quad E_j[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_j[\psi](\tau) \leq C(j, M, \Lambda) \int_{\tau_1}^{\tau_2} E_{j,q}[\psi](\tau) + C(j, M, \Lambda)E_j[\psi](\tau_1),$$

for all $j \geq 1$.

Now, by using (A.33) and (A.35) we obtain that there exists a constant $C(j, M, \Lambda)$ such that

$$(A.36) \quad E_j[\psi](\tau_2) + \int_{\tau_1}^{\tau_2} E_j[\psi](\tau)d\tau \leq C(j, M, \Lambda)E_j[\psi](\tau_1)(\tau_2 - \tau_1) + C(j, M, \Lambda)E_j[\psi](\tau_1),$$

for all $\tau_1 \leq \tau_2$. Note that (A.34) holds for all $\tau'_1 \geq \tau_1$ in the place of τ_1 . We readily obtain that

$$(A.37) \quad E_j[\psi](\tau) \leq C(j, M, \Lambda)E_j[\psi](\tau_1)$$

for all $\tau \geq \tau_1$, which concludes the proof of the energy estimates (3.9), (3.10) for (A.31).

The bootstrap.

We now return to the proof of the Proposition 3.1.

We introduce the bootstrap assumption

$$(A.38) \quad \sup_{D_\delta(\tau_1, \tau_1 + \tau_{\max})} \sum_{1 \leq j \leq k-1} \sum_{\partial \in \{\partial_{\bar{t}}, \partial_r, \Omega_1, \Omega_2, \Omega_3\}} |\partial^j \psi| \leq C_b \sqrt{\epsilon}$$

for a large constant $C_b(M, \Lambda) > 0$ to be determined later.

Volume form notation.

In this proof in any integrals over spacetime domains or hypersurfaces without explicit volume forms it is to be understood that the volume forms of the integrals over spacetime domains are with respect to the metric $g(\nabla\psi)$. The volume forms of the integrals over hypersurfaces are with respect to the induced volume form of the metric $g(\nabla\psi)$ on those hypersurfaces.

The multiplier estimates.

First, we write the general energy identities (multiplier estimates) we need. We will later use them for the vector fields $\partial_{\bar{t}}, N$.

We apply the divergence Theorem, see Proposition A.1, for the quasilinear wave equation (1.1), for the metric $g(\nabla\psi)$, with multiplier X and appropriately many commutations with a vector field \tilde{X} , on the extended region

$$(A.39) \quad D_\delta(\tau_1, \tau_1 + \tau_{\max}).$$

We consider any

$$(A.40) \quad \tau_2 - \tau_1 \leq \tau_{\max}.$$

We obtain

$$\begin{aligned}
& \text{(A.41)} \\
& \left(\int_{\{\bar{t}=\tau_2\} \cap D_\delta(\tau_1, \tau_2)} + \int_{\mathcal{H}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} \right) \mathbb{T}(g(\nabla\psi))(X, n)[\tilde{X}^i\psi] \\
& + \int_{D_\delta(\tau_1, \tau_2)} \frac{1}{2} {}^{(X)}\pi_{\mu\nu}(g(\nabla\psi))\mathbb{T}^{\mu\nu}(g(\nabla\psi))[\tilde{X}^i\psi] \\
& = \int_{\{\bar{t}=\tau_1\} \cap D_\delta(\tau_1, \tau_2)} \mathbb{T}(g(\nabla\psi))(X, n)[\tilde{X}^i\psi] - \int_{D_\delta(\tau_1, \tau_2)} X \tilde{X}^i \psi \left(\sum_{l=0}^{j-1} \tilde{X}^l [\square_{g(\nabla\psi)}, \tilde{X}] \tilde{X}^{i-1-l} \psi + \tilde{X}^i (a^{\alpha\beta} \partial_\alpha \psi \partial_\beta \psi) \right)
\end{aligned}$$

for all $1 \leq i \leq k+1$. Note the deformation tensor of $g(\nabla\psi)$ is

$$\begin{aligned}
& \text{(A.42)} \\
& \frac{1}{2} {}^{(X)}\pi^{\mu\nu}(g(\nabla\psi))\mathbb{T}_{\mu\nu}(g(\nabla\psi))[\tilde{X}^i\psi] \\
& = \frac{1}{2} (\mathcal{L}_X(\dot{g} + h(\nabla\psi)))_{\mu\nu} g^{\mu\alpha}(\nabla\psi) g^{\nu\beta}(\nabla\psi) \mathbb{T}_{\alpha\beta}(g(\nabla\psi))[\tilde{X}^i\psi] \\
& = \frac{1}{2} (\mathcal{L}_X \dot{g} + \mathcal{L}_X h(\nabla\psi))_{\mu\nu} (\dot{g}^{\mu\alpha} + h^{\mu\alpha}(\nabla\psi)) (\dot{g}^{\nu\beta} + h^{\nu\beta}(\nabla\psi)) \left(\mathbb{T}(\dot{g})(\partial_\alpha, \partial_\beta)[\tilde{X}^i\psi] - \frac{1}{2} h(\partial_\alpha, \partial_\beta, \nabla\psi) |\nabla \tilde{X}^i \psi|_{g(\nabla\psi)}^2 \right) \\
& \doteq \frac{1}{2} ({}^{(X)}\pi_{\mu\nu}(\dot{g}))\mathbb{T}^{\mu\nu}(\dot{g})[\tilde{X}^i\psi] + {}^{(X)}\pi_{non-lin}(\tilde{X}^i\psi),
\end{aligned}$$

with

$$\begin{aligned}
& \text{(A.43)} \\
& {}^{(X)}\pi_{non-lin}(\tilde{X}^i\psi) \\
& = (\mathcal{L}_X \dot{g})_{\mu\nu} \dot{g}^{\mu\alpha} \left(\dot{g}^{\nu\beta} \frac{-1}{2} h_{\alpha\beta}(\nabla\psi) |\nabla \tilde{X}^i \psi|_{g(\nabla\psi)}^2 + h^{\nu\beta} \mathbb{T}(\dot{g})(\partial_\alpha, \partial_\beta)[\tilde{X}^i\psi] + h^{\nu\beta}(\nabla\psi) \frac{-1}{2} h_{\alpha\beta}(\nabla\psi) \right) \\
& + (\mathcal{L}_X \dot{g})_{\mu\nu} h^{\mu\alpha}(\nabla\psi) \left(\dot{g}^{\nu\beta} \mathbb{T}(\dot{g})(\partial_\alpha, \partial_\beta)[\tilde{X}^i\psi] \right. \\
& \quad \left. + \dot{g}^{\nu\beta} \frac{-1}{2} h_{\alpha\beta}(\nabla\psi) |\nabla \tilde{X}^i \psi|_{g(\nabla\psi)}^2 + h^{\nu\beta} \mathbb{T}(\dot{g})(\partial_\alpha, \partial_\beta)[\tilde{X}^i\psi] + h^{\nu\beta}(\nabla\psi) \frac{-1}{2} h_{\alpha\beta}(\nabla\psi) \right) \\
& + (\mathcal{L}_X h(\nabla\psi))_{\mu\nu} \dot{g}^{\mu\alpha} \left(\dot{g}^{\nu\beta} \mathbb{T}(\dot{g})(\partial_\alpha, \partial_\beta)[\tilde{X}^i\psi] \right. \\
& \quad \left. + \dot{g}^{\nu\beta} \frac{-1}{2} h_{\alpha\beta}(\nabla\psi) |\nabla \tilde{X}^i \psi|_{g(\nabla\psi)}^2 + h^{\nu\beta} \mathbb{T}(\dot{g})(\partial_\alpha, \partial_\beta)[\tilde{X}^i\psi] + h^{\nu\beta}(\nabla\psi) \frac{-1}{2} h_{\alpha\beta}(\nabla\psi) \right) \\
& + (\mathcal{L}_X h(\nabla\psi))_{\mu\nu} h^{\mu\alpha}(\nabla\psi) \left(\dot{g}^{\nu\beta} \mathbb{T}(\dot{g})(\partial_\alpha, \partial_\beta)[\tilde{X}^i\psi] \right. \\
& \quad \left. + \dot{g}^{\nu\beta} \frac{-1}{2} h_{\alpha\beta}(\nabla\psi) |\nabla \tilde{X}^i \psi|_{g(\nabla\psi)}^2 + h^{\nu\beta} \mathbb{T}(\dot{g})(\partial_\alpha, \partial_\beta)[\tilde{X}^i\psi] + h^{\nu\beta}(\nabla\psi) \frac{-1}{2} h_{\alpha\beta}(\nabla\psi) \right)
\end{aligned}$$

where we used that the energy momentum tensor of $g(\nabla\psi)$ is

$$\text{(A.44)} \quad \mathbb{T}(g(\nabla\psi))(X, Y)[\psi] = \mathbb{T}(\dot{g})(X, Y)[\psi] - \frac{1}{2} h(X, Y, \nabla\psi) |\nabla \psi|_{g(\nabla\psi)}^2$$

for any two smooth vector field X, Y , with

$$\text{(A.45)} \quad |\nabla \psi|_{g(\nabla\psi)}^2 = g^{ab}(\nabla\psi) \partial_a \psi \partial_b \psi.$$

The $\partial_{\bar{t}}$ multiplier estimate.

We apply the energy identity (A.41) with multiplier $\partial_{\bar{t}}$ and commutators $\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2}$, in view of (A.42) and of

$$(A.46) \quad (\partial_{\bar{t}}) \pi_{\mu\nu}(\mathring{g}) \mathbb{T}^{\mu\nu}(\mathring{g}) = 0.$$

We obtain

$$(A.47) \quad \begin{aligned} & \int_{\{\bar{t}=\tau_2\}} \mathbb{T}(g(\nabla\psi))(\partial_{\bar{t}}, n) [\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi] \\ &= \int_{\{\bar{t}=\tau_1\} \cap D_{\delta}(\tau_1, \tau_2)} \mathbb{T}(g(\nabla\psi))(\partial_{\bar{t}}, n) [\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi] \\ & \quad - \left(\int_{\mathcal{H}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} \right) (\mathbb{T}(g(\nabla\psi))(\partial_{\bar{t}}, n) [\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi]) \\ & \quad - \int \int_{D_{\delta}(\tau_1, \tau_2)} (\partial_{\bar{t}}) \pi_{non-lin}(\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi) \\ & \quad - \int \int_{D_{\delta}(\tau_1, \tau_2)} \partial_{\bar{t}} \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi \left(\sum_{j_1+j_2+j_3+j_4=i_1+i_2-1} \sum_{j_5+j_6=1} \partial_{\bar{t}}^{j_1} \Omega_{\alpha}^{j_2} [\square_{g(\nabla\psi)}, \partial_{\bar{t}}^{j_5} \Omega_{\alpha}^{j_6}] \partial_{\bar{t}}^{j_3} \Omega_{\alpha}^{j_4} \psi + \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} (a^{\alpha\beta} \partial_{\alpha} \psi \partial_{\beta} \psi) \right). \end{aligned}$$

Therefore, by summing over i_1, i_2, α and by adding in the hypersurface terms

$$(A.48) \quad \sum_{1 \leq i_1+i_2 \leq i} \sum_{\alpha} \left(\int_{\mathcal{H}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} \right) (\partial_{\bar{t}}^{i_1+i_2} \Omega_{\alpha}^{i_2} \psi)^2 \frac{1}{\sqrt{|1-\mu|}} d\mathring{g}_{\mathcal{H}_{\delta}}$$

on both sides of (A.47) and recall the the energy momentum tensor of $g(\nabla\psi)$, see (A.44), we obtain

$$(A.49) \quad \begin{aligned} & \sum_{1 \leq i_1+i_2 \leq i} \sum_{\alpha} \int_{\{\bar{t}=\tau_2\}} \mathbb{T}(\mathring{g})(\partial_{\bar{t}}, n) [\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi] \\ & + \sum_{1 \leq i_1+i_2 \leq i} \sum_{\alpha} \left(\int_{\mathcal{H}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} \right) (\partial_{\bar{t}}^{i_1+i_2} \Omega_{\alpha}^{i_2} \psi)^2 \frac{1}{\sqrt{|1-\mu|}} d\mathring{g}_{\mathcal{H}_{\delta}} \\ &= \sum_{1 \leq i_1+i_2 \leq i} \sum_{\alpha} \int_{\{\bar{t}=\tau_1\}} \mathbb{T}(g(\nabla\psi))(\partial_{\bar{t}}, n) [\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi] \\ & \quad + \sum_{1 \leq i_1+i_2 \leq i} \sum_{\alpha} \int_{\{\bar{t}=\tau_2\}} \frac{1}{2} h(\partial_{\bar{t}}, n, \nabla\psi) |\nabla \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi|_{g(\nabla\psi)}^2 \\ & \quad - \sum_{1 \leq i_1+i_2 \leq i} \sum_{\alpha} \left(\int_{\mathcal{H}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} \right) (\mathbb{T}(g(\nabla\psi))(\partial_{\bar{t}}, n) [\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi]) \\ & \quad + \sum_{1 \leq i_1+i_2 \leq i} \sum_{\alpha} \left(\int_{\mathcal{H}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} \right) (\partial_{\bar{t}}^{i_1+i_2} \Omega_{\alpha}^{i_2} \psi)^2 \frac{1}{\sqrt{|1-\mu|}} d\mathring{g}_{\mathcal{H}_{\delta}} \\ & \quad - \sum_{1 \leq i_1+i_2 \leq i} \sum_{\alpha} \int \int_{D_{\delta}(\tau_1, \tau_2)} (\partial_{\bar{t}}) \pi_{non-lin}(\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi) \\ & \quad - \sum_{1 \leq i_1+i_2 \leq i} \sum_{\alpha} \int \int_{D_{\delta}(\tau_1, \tau_2)} \partial_{\bar{t}} \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi \left(\sum_{j_1+j_2+j_3+j_4=i_1+i_2-1} \sum_{j_5+j_6=1} \partial_{\bar{t}}^{j_1} \Omega_{\alpha}^{j_2} [\square_{g(\nabla\psi)}, \partial_{\bar{t}}^{j_5} \Omega_{\alpha}^{j_6}] \partial_{\bar{t}}^{j_3} \Omega_{\alpha}^{j_4} \psi \right. \\ & \quad \left. + \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} (a^{\alpha\beta} \partial_{\alpha} \psi \partial_{\beta} \psi) \right), \end{aligned}$$

Now, note that there exist constants $c(M, \Lambda), C(M, \Lambda) > 0$, such that

$$(A.50) \quad c(M, \Lambda) \left((\partial_{\bar{t}} \psi)^2 + |\nabla \psi|^2 \right) \leq \mathbb{T}(\mathring{g})(\partial_{\bar{t}}, \hat{n}_{\{\bar{t}=\tau\}}) [\psi] + C(M, \Lambda) |1-\mu| (\partial_r \psi)^2,$$

in sufficiently small neighborhoods of the horizons and

$$(A.51) \quad c(M, \Lambda, q) \left((\partial_{\bar{t}} \psi)^2 + (\partial_r \psi)^2 + |\nabla \psi|^2 \right) \leq \mathbb{T}(\mathring{g})(\partial_{\bar{t}}, \hat{n}_{\{\bar{t}=\tau\}}) [\psi]$$

in $(r_+ + q, \bar{r}_+ - q)$. Therefore, in view of Lemma A.3 and the elliptic estimate of Lemma A.4, we note that there exists a $\delta > 0$ sufficiently small, and an $\epsilon(\delta, C_b) > 0$ sufficiently small, such that

$$\begin{aligned}
& \text{(A.52)} \\
& cE_{j,q}(\tau_2) \\
& \leq CE_j(\tau_1) \\
& + C\delta \int_{\{\bar{t}=\tau_2\}} \sum_{1 \leq i_1+i_2+i_3 \leq j} \sum_{\alpha} \left(\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_{\alpha}^{i_3} \psi \right)^2 d\mathring{g}_{\{\bar{t}=\tau\}} \\
& - \sum_{1 \leq i_1+i_2 \leq j-1} \sum_{\alpha} \left(\int_{\mathcal{H}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} \right) (\mathbb{T}(g(\nabla\psi))(\partial_{\bar{t}}, n)[\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi]) \\
& + \sum_{1 \leq i_1+i_2 \leq j-1} \sum_{\alpha} \left(\int_{\mathcal{H}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} \right) \left(\partial_{\bar{t}}^{i_1+1} \Omega_{\alpha}^{i_2} \psi \right)^2 \frac{1}{\sqrt{|1-\mu|}} d\mathring{g}_{\mathcal{H}_{\delta}} \\
& + \sum_{1 \leq i_1+i_2 \leq i} \sum_{\alpha} \int_{\{\bar{t}=\tau_2\}} \frac{1}{2} h(\partial_{\bar{t}}, n, \nabla\psi) |\nabla \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi|_g^2(\nabla\psi) \\
& - \sum_{1 \leq i_1+i_2 \leq j-1} \sum_{\alpha} \int \int_{D_{\delta}(\tau_1, \tau_2)} \overset{(\partial_{\bar{t}})}{\pi_{non-lin}}(\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi) \\
& - \sum_{1 \leq i_1+i_2 \leq j-1} \sum_{\alpha} \int \int_{D_{\delta}(\tau_1, \tau_2)} \partial_{\bar{t}} \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi \left(\sum_{j_1+j_2+j_3+j_4=i_1+i_2-1} \sum_{j_5+j_6=1} \partial_{\bar{t}}^{j_1} \Omega_{\alpha}^{j_2} [\square_{g(\nabla\psi)}, \partial_{\bar{t}}^{j_5} \Omega_{\alpha}^{j_6}] \partial_{\bar{t}}^{j_3} \Omega_{\alpha}^{j_4} \psi \right. \\
& \quad \left. + \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} (a^{\alpha\beta} \partial_{\alpha} \psi \partial_{\beta} \psi) \right) \\
& = CE_j(\tau_1) \\
& + C\delta \int_{\{\bar{t}=\tau_2\}} \sum_{1 \leq i_1+i_2+i_3 \leq j} \sum_{\alpha} \left(\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_{\alpha}^{i_3} \psi \right)^2 d\mathring{g}_{\{\bar{t}=\tau\}} \\
& - \sum_{1 \leq i_1+i_2 \leq j-1} \sum_{\alpha} \left(\int_{\mathcal{H}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} \right) (\mathbb{T}(g(\nabla\psi))(\partial_{\bar{t}}, n)[\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi]) \\
& + \sum_{1 \leq i_1+i_2 \leq j-1} \sum_{\alpha} \left(\int_{\mathcal{H}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_{\delta}^+ \cap D_{\delta}(\tau_1, \tau_2)} \right) \left(\partial_{\bar{t}}^{i_1+1} \Omega_{\alpha}^{i_2} \psi \right)^2 \frac{1}{\sqrt{|1-\mu|}} d\mathring{g}_{\mathcal{H}_{\delta}} \\
& + \sum_{1 \leq i_1+i_2 \leq i} \sum_{\alpha} \int_{\{\bar{t}=\tau_2\}} \frac{1}{2} h(\partial_{\bar{t}}, n, \nabla\psi) |\nabla \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi|_g^2(\nabla\psi) \\
& - \sum_{1 \leq i_1+i_2 \leq j-1} \sum_{\alpha} \int \int_{D_{\delta}(\tau_1, \tau_2)} \overset{(\partial_{\bar{t}})}{\pi_{non-lin}}(\partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi) \\
& - \sum_{1 \leq i_1+i_2 \leq j-1} \sum_{\alpha} \int \int_{D_{\delta}(\tau_1, \tau_2)} \partial_{\bar{t}} \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} \psi \left(\sum_{j_1+j_2+j_3+j_4=i_1+i_2-1} \sum_{j_5+j_6=1} \partial_{\bar{t}}^{j_1} \Omega_{\alpha}^{j_2} [h^{ab}(\nabla\psi) \partial_a \partial_b + S^c{}_{ab}(h) \partial_c, \partial_{\bar{t}}^{j_5} \Omega_{\alpha}^{j_6}] \partial_{\bar{t}}^{j_3} \Omega_{\alpha}^{j_4} \psi \right. \\
& \quad \left. + \partial_{\bar{t}}^{i_1} \Omega_{\alpha}^{i_2} (a^{\alpha\beta} \partial_{\alpha} \psi \partial_{\beta} \psi) \right)
\end{aligned}$$

for all $1 \leq j \leq k+1$, for some constants $c(M, \Lambda), C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$, where for $S^c{}_{ab}$ see (5.16). Note that to get the energies $E_{j,q}, E_j$ on the left hand side and right hand side respectively, we used the bootstrap assumption (A.38) and Lemma A.3, for $\epsilon(\delta, C_b) > 0$ sufficiently small.

We now estimate the hypersurface terms on $\mathcal{H}_{\delta}^+, \bar{\mathcal{H}}_{\delta}^+$ on the right hand side of (A.52) (in view of Lemma A.5) and the two last (non-linear) terms of (A.52). Specifically, in view of the bootstrap (A.38), and the definitions of the smooth tensors a, h , see Sections 2.7, 3.1 respectively, there exists a $\delta > 0$ sufficiently small, and an $\epsilon(\delta, C_b) > 0$ sufficiently small, such that for the two non-linear

terms we use the Lemma A.3 and the coarea formula (2.4), to obtain

$$\begin{aligned}
& E_{k+1,q}(\tau_2) \\
& \leq CE_{k+1}(\tau_1) + C\delta E_{k+1}(\tau_2) \\
\text{(A.53)} \quad & + C\sqrt{\delta} \left(\int_{\mathcal{H}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} \right) \sum_{1 \leq i \leq k} \mathbb{T}(N, n)[N^i \psi] d\dot{g}_{\mathcal{H}_\delta} \\
& + \sqrt{\epsilon(\delta)} C \int_{\tau_1}^{\tau_2} d\tau E_j(\tau),
\end{aligned}$$

for $k \geq 7$, where $C = C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$. We appropriately distributed derivatives and used a Sobolev estimate, see Lemma 2.2, in view of the fact that $k = 7$ is the smallest integer such that

$$\text{(A.54)} \quad \lfloor \frac{k+1}{2} \rfloor + 2 \leq k-1.$$

The N multiplier estimate.

We apply the energy identity (A.41) with multiplier N and commutators N^i , in view of (A.42).

We obtain

$$\begin{aligned}
\text{(A.55)} \quad & \left(\int_{\{\bar{t}=\tau_2\} \cap D_\delta(\tau_1, \tau_2)} + \int_{\mathcal{H}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} \right) \mathbb{T}(g(\nabla\psi))(N, n)[N^i \psi] \\
& + \int_{D_\delta(\tau_1, \tau_2)} \frac{1}{2} {}^{(N)}\pi_{\mu\nu}(\dot{g}) \mathbb{T}^{\mu\nu}(\dot{g})[N^i \psi] \\
& = \int_{\{\bar{t}=\tau_1\} \cap D_\delta(\tau_1, \tau_2)} \mathbb{T}(g(\nabla\psi))(N, n)[N^i \psi] \\
& - \int_{D_\delta(\tau_1, \tau_2)} {}^{(N)}\pi_{non-lin}(N^i \psi) - \int_{D_\delta(\tau_1, \tau_2)} N^{i+1} \psi \left(\sum_{l=0}^{i-1} N^l [\square_{g(\nabla\psi)}, N] N^{i-1-l} \psi + N^i (a^{\alpha\beta} \partial_\alpha \psi \partial_\beta \psi) \right)
\end{aligned}$$

for all $1 \leq i \leq k$.

Therefore, for a sufficiently small δ we obtain

$$\begin{aligned}
\text{(A.56)} \quad & \left(\int_{\{\bar{t}=\tau_2\} \cap D_\delta(\tau_1, \tau_2)} + \int_{\mathcal{H}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} \right) \mathbb{T}(g(\nabla\psi))(N, n)[N^i \psi] \\
& + \int_{D_\delta(\tau_1, \tau_2) \cap (\{r_+ - q \leq r \leq r_+ + q\} \cup \{\bar{r}_+ - q \leq r \leq \bar{r}_+ + q\})} \frac{1}{2} {}^{(N)}\pi_{\mu\nu}(\dot{g}) \mathbb{T}^{\mu\nu}(\dot{g})[N^i \psi] \\
& = \int_{\{\bar{t}=\tau_1\} \cap D_\delta(\tau_1, \tau_2)} \mathbb{T}(g(\nabla\psi))(N, n)[N^i \psi] \\
& - \int_{D_\delta(\tau_1, \tau_2) \setminus (\{r_+ - q \leq r \leq r_+ + q\} \cup \{\bar{r}_+ - q \leq r \leq \bar{r}_+ + q\})} \frac{1}{2} {}^{(N)}\pi_{\mu\nu}(\dot{g}) \mathbb{T}^{\mu\nu}(g(\nabla\psi))[N^i \psi] \\
& - \int_{D_\delta(\tau_1, \tau_2)} {}^{(N)}\pi_{non-lin}(N^i \psi) - \int_{D_\delta(\tau_1, \tau_2)} N^{i+1} \psi \left(\sum_{l=0}^{i-1} N^l [\square_{g(\nabla\psi)}, N] N^{i-1-l} \psi + N^i (a^{\alpha\beta} \partial_\alpha \psi \partial_\beta \psi) \right),
\end{aligned}$$

for all $1 \leq i \leq k$, where for q see Lemma A.1. Now, we estimate the second term on the right hand side of (A.56), and obtain

$$\begin{aligned}
& \text{(A.57)} \\
& \left(\int_{\{\bar{t}=\tau_2\} \cap D_\delta(\tau_1, \tau_2)} + \int_{\mathcal{H}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} \right) \mathbb{T}(g(\nabla\psi))(N, n)[N^i\psi] \\
& + \int_{D_\delta(\tau_1, \tau_2) \cap (\{r_+ - q \leq r \leq r_+ + q\} \cup \{\bar{r}_+ - q \leq r \leq \bar{r}_+ + q\})} \frac{1}{2} {}^{(N)}\pi_{\mu\nu}(\dot{g}) \mathbb{T}^{\mu\nu}(\dot{g})[N^i\psi] \\
& \leq \int_{\{\bar{t}=\tau_1\} \cap D_\delta(\tau_1, \tau_2)} \mathbb{T}(g(\nabla\psi))(N, n)[N^i\psi] \\
& + C\tau_{max} \sup_{\tau \in [\tau_1, \tau_2]} \int_{\{\bar{t}=\tau\} \cap \{r_+ + q \leq r \leq \bar{r}_+ - q\}} \sum_{1 \leq i_1 + i_2 + i_3 = i+1} \sum_{\alpha} (\partial_{\bar{t}}^{i_1} \partial_r^{i_2} \Omega_\alpha^{i_3} \psi)^2 \\
& - \int_{D_\delta(\tau_1, \tau_2)} {}^{(N)}\pi_{non-lin}(N^i\psi) - \int_{D_\delta(\tau_1, \tau_2)} N^{i+1}\psi \left(\sum_{l=0}^{i-1} N^l [\square_{g(\nabla\psi)}, N] N^{i-1-l}\psi + N^i (a^{\alpha\beta} \partial_\alpha \psi \partial_\beta \psi) \right),
\end{aligned}$$

for all $1 \leq i \leq k$, where the constant $C > 0$ depends only on the black hole parameters.

Now, in view of Lemma A.3 and the elliptic estimates of Lemma A.4, there exist $\delta, \epsilon(\delta, C_b)$, sufficiently small, such that the estimate (A.57) implies

$$\begin{aligned}
& cE_j(\tau_2) + c \int_{\tau_1}^{\tau_2} d\tau E_j(\tau) \\
& + c \left(\int_{\mathcal{H}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} \right) \sum_{1 \leq i \leq j-1} \mathbb{T}(N, n)[N^i\psi] d\dot{g}_{\mathcal{H}_\delta} \\
& \leq CE_j(\tau_1) + C\tau_{max} \sup_{\tau \in [\tau_1, \tau_2]} E_{j,q}(\tau) \\
& - \int_{D_\delta(\tau_1, \tau_2)} \sum_{1 \leq i \leq j-1} {}^{(N)}\pi_{non-lin}(N^i\psi) \\
\text{(A.58)} \quad & - \int_{D_\delta(\tau_1, \tau_2)} \sum_{1 \leq i \leq j-1} N^{i+1}\psi \left(\sum_{l=0}^{i-1} N^l [\square_{g(\nabla\psi)}, N] N^{i-1-l}\psi + N^i (a^{\alpha\beta} \partial_\alpha \psi \partial_\beta \psi) \right) \\
& = CE_j(\tau_1) + C\tau_{max} \sup_{\tau \in [\tau_1, \tau_2]} E_{j,q}(\tau) \\
& - \int_{D_\delta(\tau_1, \tau_2)} \sum_{1 \leq i \leq j-1} {}^{(N)}\pi_{non-lin}(N^i\psi) \\
& - \int_{D_\delta(\tau_1, \tau_2)} \sum_{1 \leq i \leq j-1} N^{i+1}\psi \left(\sum_{l=0}^{i-1} N^l [\square_{\dot{g}}, N] N^{i-1-l}\psi \right) \\
& - \int_{D_\delta(\tau_1, \tau_2)} \sum_{1 \leq i \leq j-1} N^{i+1}\psi \left(\sum_{l=0}^{i-1} N^l [h^{ab} \partial_a \partial_b + S^c{}_{ab} \partial_c, N] N^{i-1-l}\psi + N^i (a^{\alpha\beta} \partial_\alpha \psi \partial_\beta \psi) \right),
\end{aligned}$$

for all $1 \leq j \leq k+1$, for some constants $c(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}), C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]})$, where for $S^c{}_{ab}$ see (5.16).

Similarly to (A.52), we estimate the two last (non-linear) terms of (A.58). Note that we estimate the third to last bulk term of (A.58), in view on the linear N -commutation Lemma A.2. Specifically, in view of the bootstrap (A.38), and the definitions of the smooth tensors a, h , see Sections 2.7, 3.1 respectively, there exists a $\delta > 0$ sufficiently small, and an $\epsilon(\delta, C_b) > 0$ sufficiently small, such that for the two non-linear terms we use the Lemma A.3 and the coarea formula (2.4), to obtain

$$\begin{aligned}
\text{(A.59)} \quad & E_{k+1}(\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{k+1}(\tau) + \left(\int_{\mathcal{H}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} \right) \sum_{1 \leq i \leq k} \mathbb{T}(N, n)[N^i\psi] d\dot{g}_{\mathcal{H}_\delta} \\
& \leq CE_{k+1}(\tau_1) + C\tau_{max} \sup_{\tau \in [\tau_1, \tau_2]} E_{j,q}(\tau)
\end{aligned}$$

for $k \geq 7$, where $0 < C = C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]})$. We appropriately distributed derivatives and used a Sobolev injection, see Lemma 2.2, in view of that $k = 7$ is the smallest integer such that

$$(A.60) \quad \lfloor \frac{k+1}{2} \rfloor + 2 \leq k - 1.$$

We obtain

The combination of the $\partial_{\bar{t}}$ and N multiplier estimates.

Now, we combine the $\partial_{\bar{t}}$ -estimate, namely (A.53), and the N -estimate, namely (A.59).

We obtain

$$(A.61) \quad \begin{aligned} & E_{k+1}(\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{k+1}(\tau) + \left(\int_{\mathcal{H}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} \right) \sum_{1 \leq i \leq k} \mathbb{T}(N, n)[N^i \psi] d\dot{g}_{\mathcal{H}_\delta} \\ & \leq C E_{k+1}(\tau_1) + C \tau_{max} \left(C E_{k+1}(\tau_1) + C \delta E_{k+1}(\tau_2) \right. \\ & \quad \left. + C \sqrt{\delta} \left(\int_{\mathcal{H}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} + \int_{\bar{\mathcal{H}}_\delta^+ \cap D_\delta(\tau_1, \tau_2)} \right) \sum_{1 \leq i \leq k} \mathbb{T}(N, n)[N^i \psi] d\dot{g}_{\mathcal{H}_\delta} \right. \\ & \quad \left. + \sqrt{\epsilon(\delta)} C \int_{\tau_1}^{\tau_2} d\tau E_{k+1}(\tau) \right), \end{aligned}$$

for $k \geq 7$.

The integral inequality. Therefore, for $\delta(\tau_{max}), \epsilon(\tau_{max}, \delta) > 0$ sufficiently small, then from inequality (A.61) we obtain

$$(A.62) \quad E_{k+1}(\tau_2) + \int_{\tau_1}^{\tau_2} d\tau E_{k+1}(\tau) \leq C E_{k+1}(\tau_1) + C(\tau_2 - \tau_1) E_{k+1}(\tau_1),$$

for $k \geq 7$, and for a constant $C(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 0$. Note that (A.62) is the quasilinear analogue of the integral inequality (A.36).

Finishing the proof, improving the bootstrap (A.38).

Note that the estimate (A.62) holds for any $\tau_1' \geq \tau_1$ in the place of τ_1 .

Therefore, arguing as in the linear case, see the integral inequality (A.36), we obtain from (A.62) that there exists a constant

$$(A.63) \quad C_{wp}(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 1$$

independent of τ_{max} , such that

$$(A.64) \quad E_{k+1}[\psi](\tau') \leq C_{wp} E_{k+1}[\psi](\tau_1),$$

for all $\tau' \in [\tau_1, \tau_1 + \tau_{max}]$. Moreover, by using (A.64) and the smallness $E_{k+1}[\psi](\tau_1) \leq \epsilon$, we also prove that

$$(A.65) \quad E_{k+2}[\psi](\tau') \leq C_{wp} E_{k+2}[\psi](\tau_1),$$

by repeating the arguments of the above proof, and after redefining $C_{wp}(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) > 1$ appropriately, provided that $E_{k+2}[\psi](\tau_1) < \infty$.

Finally, for a sufficiently large $C_b(k, M, \Lambda, A_{[k+1]}, B_{[k+1]}) \gg C_{wp}$ and a sufficiently small $\epsilon > 0$, we improve the bootstrap assumption (A.38) by Sobolev inequalities, see Lemma 2.2, on the left hand side of inequality (A.64).

We conclude the energy estimates (3.9), (3.10) and therefore the Proposition. \square

Note also the following Remark

Remark A.1. *A posteriori, one can remove the dependence of δ on the τ_{step} parameter. To do so, note that we dropped the terms on the hypersurfaces $\mathcal{H}_\delta^+, \bar{\mathcal{H}}_\delta^+$ from (A.61). By using the N redshift vector field in the spacetime regions*

$$(A.66) \quad \{r_+ - \tilde{\delta} \leq r \leq r_+ - \delta\}, \quad \{\bar{r}_+ + \delta \leq r \leq \bar{r}_+ + \tilde{\delta}\}$$

for $\delta > 0$ as in Theorem 1, and for a sufficiently small $\tilde{\delta}$ independent of τ_{step} , we can absorb the contributions at \mathcal{H}_δ^+ , $\tilde{\mathcal{H}}_\delta^+$ by (A.61), and conclude the Cauchy stability result of Proposition 3.1 for a $\tilde{\delta} > 0$ in the place of δ .

REFERENCES

- [1] Lars Andersson, Thomas Bäckdahl, Pieter Blue, and Siyuan Ma. Stability for linearized gravity on the Kerr spacetime, 2019. arXiv:1903.03859.
- [2] Jean-François Bony and Dietrich Häfner. Decay and non-decay of the local energy for the wave equation on the de Sitter-Schwarzschild metric. *Comm. Math. Phys.*, 282(3):697–719, 2008.
- [3] N. Burq. Smoothing effect for Schrödinger boundary value problems. *Duke Math. J.*, 123(2):403–427, 2004.
- [4] Nicolas Burq, Colin Guillarmou, and Andrew Hassell. Strichartz estimates without loss on manifolds with hyperbolic trapped geodesics. *Geom. Funct. Anal.*, 20(3):627–656, 2010.
- [5] Mihalis Dafermos, Gustav Holzegel, and Igor Rodnianski. Boundedness and decay for the Teukolsky equation on Kerr spacetimes I: The case $|a| \ll M$. *Ann. PDE*, 5(1):Paper No. 2, 118, 2019.
- [6] Mihalis Dafermos, Gustav Holzegel, and Igor Rodnianski. The linear stability of the Schwarzschild solution to gravitational perturbations. *Acta Math.*, 222(1):1–214, 2019.
- [7] Mihalis Dafermos, Gustav Holzegel, Igor Rodnianski, and Martin Taylor. The non-linear stability of the Schwarzschild family of black holes, 2021. arXiv:2104.08222.
- [8] Mihalis Dafermos and Igor Rodnianski. A note on energy currents and decay for the wave equation on a Schwarzschild background. September 2007. arXiv:0710.0171.
- [9] Mihalis Dafermos and Igor Rodnianski. The wave equation on Schwarzschild-de Sitter spacetimes, September 2007. arXiv:0709.2766.
- [10] Mihalis Dafermos and Igor Rodnianski. The red-shift effect and radiation decay on black hole spacetimes. *Comm. Pure Appl. Math.*, 62(7):859–919, 2009.
- [11] Mihalis Dafermos and Igor Rodnianski. A new physical-space approach to decay for the wave equation with applications to black hole spacetimes. In *XVIth International Congress on Mathematical Physics*, pages 421–432. World Sci. Publ., Hackensack, NJ, 2010.
- [12] Mihalis Dafermos and Igor Rodnianski. Lectures on black holes and linear waves. In *Evolution equations*, volume 17 of *Clay Math. Proc.*, pages 97–205. Amer. Math. Soc., Providence, RI, 2013.
- [13] Mihalis Dafermos, Igor Rodnianski, and Yakov Shlapentokh-Rothman. Decay for solutions of the wave equation on Kerr exterior spacetimes III: The full subextremal case $|a| < M$. *Ann. of Math. (2)*, 183(3):787–913, 2016.
- [14] Semyon Dyatlov. Exponential energy decay for Kerr–de Sitter black holes beyond event horizons. *Math. Res. Lett.*, 18(5):1023–1035, 2011.
- [15] Semyon Dyatlov. Quasi-normal modes and exponential energy decay for the Kerr-de Sitter black hole. *Comm. Math. Phys.*, 306(1):119–163, 2011.
- [16] Semyon Dyatlov. Spectral gaps for normally hyperbolic trapping. *Ann. Inst. Fourier (Grenoble)*, 66(1):55–82, 2016.
- [17] Helmut Friedrich. On the existence of n -geodesically complete or future complete solutions of Einstein’s field equations with smooth asymptotic structure. *Comm. Math. Phys.*, 107(4):587–609, 1986.
- [18] Dietrich Häfner, Peter Hintz, and András Vasy. Linear stability of slowly rotating Kerr black holes. *Invent. Math.*, 223(3):1227–1406, 2021.
- [19] Peter Hintz and András Vasy. Non-trapping estimates near normally hyperbolic trapping. *Math. Res. Lett.*, 21(6):1277–1304, 2014.
- [20] Peter Hintz and András Vasy. Semilinear wave equations on asymptotically de Sitter, Kerr–de Sitter and Minkowski spacetimes. *Anal. PDE*, 8(8):1807–1890, 2015.
- [21] Peter Hintz and András Vasy. Global analysis of quasilinear wave equations on asymptotically de Sitter spaces. *Annales de l’Institut Fourier*, 66(4):1285–1408, 2016.
- [22] Peter Hintz and András Vasy. Global analysis of quasilinear wave equations on asymptotically Kerr–de Sitter spaces. *Int. Math. Res. Not. IMRN*, (17):5355–5426, 2016.
- [23] Peter Hintz and András Vasy. The global non-linear stability of the Kerr–de Sitter family of black holes. *Acta Math.*, 220(1):1–206, 2018.
- [24] Gustav Holzegel and Christopher Kauffman. A note on the wave equation on black hole spacetimes with small non-decaying first order terms, May 2020. arXiv:2005.13644.
- [25] Mitsuru Ikawa. Decay of solutions of the wave equation in the exterior of several convex bodies. *Ann. Inst. Fourier (Grenoble)*, 38(2):113–146, 1988.
- [26] S. Klainerman. The null condition and global existence to nonlinear wave equations. In *Nonlinear systems of partial differential equations in applied mathematics, Part 1 (Santa Fe, N.M., 1984)*, volume 23 of *Lectures in Appl. Math.*, pages 293–326. Amer. Math. Soc., Providence, RI, 1986.
- [27] Sergiu Klainerman and Jérémie Szeftel. *Global nonlinear stability of Schwarzschild spacetime under polarized perturbations*, volume 210 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2020.

- [28] Hans Lindblad and Mihai Tohaneanu. Global existence for quasilinear wave equations close to Schwarzschild. *Comm. Partial Differential Equations*, 43(6):893–944, 2018.
- [29] Jonathan Luk. The null condition and global existence for nonlinear wave equations on slowly rotating Kerr spacetimes. *J. Eur. Math. Soc. (JEMS)*, 15(5):1629–1700, 2013.
- [30] Georgios Mavrogiannis. Morawetz estimates without relative degeneration and exponential decay on Schwarzschild de Sitter spacetimes. preprint.
- [31] Georgios Mavrogiannis. Relatively non degenerate estimates on Kerr de Sitter and applications. in preparation.
- [32] Richard Melrose, Antônio Sá Barreto, and Andrés Vasy. Asymptotics of solutions of the wave equation on de Sitter-Schwarzschild space. *Comm. Partial Differential Equations*, 39(3):512–529, 2014.
- [33] Stéphane Nonnenmacher and Maciej Zworski. Quantum decay rates in chaotic scattering. *Acta Math.*, 203(2):149–233, 2009.
- [34] Antônio Sá Barreto and Maciej Zworski. Distribution of resonances for spherical black holes. *Math. Res. Lett.*, 4(1):103–121, 1997.
- [35] Volker Schlue. Global results for linear waves on expanding Kerr and Schwarzschild de Sitter cosmologies. *Comm. Math. Phys.*, 334(2):977–1023, 2015.
- [36] Volker Schlue. Decay of the Weyl curvature in expanding black hole cosmologies, 2021. arXiv:1610.04172.
- [37] Yakov Shlapentokh-Rothman and Rita Teixeira da Costa. Boundedness and decay for the Teukolsky equation on Kerr in the full subextremal range $|a| < M$: frequency space analysis, 2020. arXiv:2005.13644.
- [38] Daniel Tataru and Mihai Tohaneanu. A local energy estimate on Kerr black hole backgrounds. *Int. Math. Res. Not. IMRN*, (2):248–292, 2011.
- [39] Andrés Vasy. The wave equation on asymptotically de Sitter-like spaces. *Adv. Math.*, 223(1):49–97, 2010.
- [40] Andrés Vasy. Microlocal analysis of asymptotically hyperbolic and Kerr-de Sitter spaces (with an appendix by Semyon Dyatlov). *Invent. Math.*, 194(2):381–513, 2013.