

Spin Entropy

Davi Geiger and Zvi M. Kedem

Courant Institute of Mathematical Sciences

New York University, New York, New York 10012

Abstract

In classical physics, entropy quantifies the randomness of large systems, where the complete specification of the state, though possible in theory, is not possible in practice. In quantum physics, despite its inherently probabilistic nature, the concept of entropy has been elusive. The von Neumann entropy, currently adopted in quantum information and computing, models only the randomness associated with unknown specifications of a state and is zero for pure quantum states, and thus cannot quantify the inherent randomness of its observables. Our goal is to provide such quantification.

This paper focuses on the quantification of the observed spin values associated with a pure quantum state, given an axis z . To this end, we define a spin entropy, which is not zero for pure states, and its minimum is $\ln 2\pi$, reflecting the uncertainty principle for the spin observables. We study the spin entropy for single massive particles with spin $1/2$ and spin 1 , photons, and two fermions in entangled and disentangled states. The spin entropy attains local minima for Bell states, which are pure entangled states of two fermions, and local maxima for disentangled states. The spin entropy may be useful for developing robust quantum computational processes.

INTRODUCTION

The transition of concepts developed in classical physics to their counterparts in quantum physics is still a subject of ongoing interest.

One area of interest is the quantification of randomness associated with a state. In classical physics, entropy was introduced to handle large ensembles of particles, where the complete specification of the state is not practical. However, it has been an elusive concept in quantum physics. Von Neumann entropy [20] quantifies the lack of knowledge an observer has about the quantum state, i.e., it quantifies the randomness in specifying the degrees of freedom (DOFs) of a quantum state. However, it does not quantify the intrinsic randomness of the observables associated with a quantum state whose DOFs have been specified. Thus, all single particles and entangled particles have zero von Neumann entropy even though only probabilities can be known about the set of observables. The Stern–Gerlach experiment [10] illustrates the quantum scenario for spin, where spin $1/2$ particle state with a z up direction are prepared and, despite having zero von Neuman entropy, randomness on the spin values along any direction perpendicular to z is observed. Clearly, this uncertainty is due to the intrinsic randomness of the observables given a quantum state, the z up state.

We propose a definition of quantum entropy in order to quantify the randomness of the observables once the DOFs of a quantum state have been specified. Following on the earlier example, our focus is spin systems. We adopt the geometric quantization method of the spin to create a spin phase space, in which the randomness of the state can be quantified. We show that this entropy has the minimum value of $\ln 2\pi$, due to the uncertainty principle for spin.

In this new quantification of randomness, we explore entanglement and disentanglement of two particles. Entanglement states tend to disentangle during time evolution in nature, posing a challenge to build quantum computer processes. Entan-

glement is a topic of much interest in quantum information and quantum computing. A better understanding of temporal evolution of such states will influence the choice of quantum processes for constructing robust quantum algorithms and machines. For the case of two particles, the behavior of von Neumann entropy of entanglement has been studied by tracing out the states of one particle to obtain a mixture of states for the remaining particle. While our proposed entropy decreases the more entangled the states are, von Neumann entropy increases the more the states are entangled. Thus, our entropy will be increasing during the evolution of an entangled state, up to when disentanglement occurs.

The paper is organized as follows: First we describe our entropy. We prove this entropy's minimum possible value. We then evaluate the entropy for massive single particles of spin $1/2$ and 1 . We extend the study to photons and to two fermions with spin $1/2$ in entangled and disentangled states. Finally, we conclude the paper.

Spin Entropy

The spin matrix associated with a particle can be specified, e.g., [7], as

$$\begin{aligned} \vec{S} &= S_x \hat{x} + S_y \hat{y} + S_z \hat{z} \quad \text{and} \quad S^2 = S_x^2 + S_y^2 + S_z^2, \\ [S_a, S_b] &= i\hbar S_c, \quad \text{where } a, b, c \text{ is a cyclic permutation of } x, y, z, \text{ and} \\ [S^2, S_a] &= 0, \quad \text{for } a = x, y, z. \end{aligned}$$

The spin value of a particle is a Casimir invariant but it is not possible to know the spin projections in all directions in three dimensions. Knowing the value of the spin along the z direction implies that the x or the y direction spin values cannot be known. This uncertainty reflects the close relation between spin matrices, their unitary transformations, and the rotation group $SO(3)$. For spin $1/2$ particles, any spin state is reachable from any other spin state via a 2×2 unitary transformation,

a local isomorphism (and a global homomorphism) to the $SO(3)$ group. For spin 1, the matrices are unitarily similar to $SO(3)$, and one can transform them into generators of $SO(3)$ via unitary transformations.

The Geometric Quantization Approach to Spin Phase Space

The DOFs of a spin specify the state along a selected z axis. Our goal is to quantify the randomness associated with the spin values given a state. The commutation property $[S_x, S_y] = i\hbar S_z$ shows that an eigentate of S_z yields randomness in both x and y spin directions. Our approach to capture all the randomness without double counting is to describe the spin operators with the least number of variables. Then, we exploit the phase space of a spin, which is derived from a quantized sphere as developed by the geometric quantization (GQ) method, e.g., see [4, 15, 23].

We start by summarizing the creation of the spin phase space. On the sphere, the surface of the ball with a radius given by the spin magnitude $s\hbar$, the z values are described by the polar representation $s\hbar \cos \theta$, while the values on the intersection with planes perpendicular to the z axis are described by the azimuth angle ϕ . Treating the sphere as the phase space, one assigns a rescaled symplectic 2-form in spherical polar coordinates $d\omega = dp \wedge dq = s\hbar \sin \theta d\theta \wedge d\phi$, and so the Lagrangian is $\mathcal{L} = p\dot{q} = s\hbar \cos \theta \dot{\phi}$, and the action is $S_{\mathcal{L}} = \int \mathcal{L} dt = s\hbar \int \cos \theta d\phi$ [?]. From this Lagrangian, the GQ method derives the uncertainty commutation relation

$$[\phi, s\hbar \cos \theta] = i\hbar,$$

where the conjugate pair of eigenvalues $(\phi, s\hbar \cos \theta) \in [0, 2\pi) \times \{-s\hbar, \dots, s\hbar\}$ forms a spin phase space with a finite Hilbert space volume $2(2s + 1)\pi\hbar$.

At the northern pole ($\cos \theta = 1$) and the southern pole ($\cos \theta = -1$) the angle ϕ is not defined. Thus, in the basis $|\phi\rangle$ that diagonalizes ϕ we can have two operators,

namely $s\hbar \cos \theta = s\hbar - i\hbar \frac{\partial}{\partial \phi}$ for the northern hemisphere and $s\hbar \cos \theta = -s\hbar - i\hbar \frac{\partial}{\partial \phi}$ for the southern hemisphere.

In the basis $|\phi\rangle$, an eigenstate of the operator $S_z = s\hbar \cos \theta$ is then

$$|\xi_{s,m}\rangle = \int |\phi\rangle \langle \phi | \xi_{s,m}\rangle d\phi = \int \psi_{s,m}(\phi) |\phi\rangle d\phi,$$

where $m = -s, \dots, s$, and

$$\psi_{s,m}(\phi) = \begin{cases} \frac{1}{\sqrt{2\pi}} e^{i(s+m)\phi}, & m \geq 0 \quad (\text{northern hemisphere}) \\ \frac{1}{\sqrt{2\pi}} e^{i(-s+m)\phi}, & m < 0 \quad (\text{southern hemisphere}) \end{cases} \quad (1)$$

The two solutions in (1) differ by a phase (gauge) transformation of $e^{-i2s\phi}$.

Consider a particle of spin magnitude $s\hbar$ in a state $|\xi_s\rangle$. This state in the basis of the eigenvectors of $S_z = s\hbar \cos \theta$ and S^2 is

$$|\xi_s\rangle = \sum_{m=-s}^s \alpha_{s,m} |\xi_{s,m}\rangle,$$

with $1 = \sum_{m=-s}^s |\alpha_{s,m}|^2$. In the basis of the conjugate variable ϕ , the state $|\xi_s\rangle$ is

$$\begin{aligned} |\xi_s\rangle &= \int_0^{2\pi} |\phi\rangle \langle \phi | \xi_s\rangle d\phi = \int_0^{2\pi} |\phi\rangle \sum_{m=-s}^s \alpha_{s,m} \langle \phi | \xi_{s,m}\rangle d\phi \\ &= \int_0^{2\pi} \sum_{m=-s}^s \alpha_{s,m} \psi_{s,m}(\phi) |\phi\rangle d\phi = \int_0^{2\pi} \lambda_s(\phi) |\phi\rangle d\phi, \end{aligned} \quad (2)$$

where the wave function in the $|\phi\rangle$ basis is

$$\lambda_s(\phi) = \langle \phi | \xi_s\rangle = \sum_{m=-s}^s \alpha_{s,m} \psi_{s,m}(\phi). \quad (3)$$

The spin phase state with wave functions $\{\alpha_{s,m}\}$, $\{\lambda_s(\phi)\}$ has associated probabilities $\{P_{s,m} = |\alpha_{s,m}|^2\}$ and probability densities $\{\rho(\phi) = |\lambda_s(\phi)|^2\}$, respectively.

The Entropy in Spin Phase Space

We define the entropy of a state with spin s in spin phase space to be

$$\mathbf{S}_s = \mathbf{S}_{s,\theta}^z + \mathbf{S}_{s,\phi}^{z^\perp} = - \sum_{m=-s}^s |\alpha_{s,m}|^2 \ln |\alpha_{s,m}|^2 - \int |\lambda_s(\phi)|^2 \ln |\lambda_s(\phi)|^2 d\phi. \quad (4)$$

The first term is the Shannon entropy capturing the randomness of the spin value along the z axis. The second term is differential entropy capturing the randomness of the spin value in the plane perpendicular to the z axis. We define *intrinsic-information*, represented by Γ , as the inverse of the entropy, i.e.,

$$\Gamma_s = \frac{-1}{\sum_{m=-s}^s |\alpha_{s,m}|^2 \ln |\alpha_{s,m}|^2 + \int |\lambda_s(\phi)|^2 \ln |\lambda_s(\phi)|^2 d\phi}.$$

Γ_s is non-negative and quantifies how much it is known about the observables of a state, where infinite value would imply infinite precision, or full information, on all observable values simultaneously. To illustrate, consider the case of $\alpha_{s,m} = \sqrt{\frac{2\pi}{2s+1}} \psi_{s,m}^*(\phi_0)$, which yields the uniform distribution $|\alpha_{s,m}|^2 = \frac{1}{2s+1}$, i.e., it is the a state with the lowest intrinsic-information about the spin value along the z axis.

Then

$$\begin{aligned} \lambda_s(\phi) &= \frac{1}{\sqrt{2\pi} 2s+1} \sum_{m=-s}^s e^{i((2\theta(m)-1)s+m)(\phi-\phi_0)} \\ &= \frac{1}{\sqrt{2\pi} (2s+1)} \left(1 - \text{mod}(2s, 2) + 2 \sum_{j=j_{\min}}^{2s} \cos(j(\phi - \phi_0)) \right), \end{aligned}$$

where $j_{\min} = s + 1 - \frac{1}{2} \text{mod}(2s, 2)$, and mod is the modulus function. Note that the distribution $\lambda_s(\phi)$ becomes more concentrated around $\phi = \phi_0$ as s increases. The intrinsic-information about the spin on the plane z^\perp increases as s increases. However, the total intrinsic-information reduces due to the increase in entropy along

z. For a fermion of spin $1/2$ the entropy of (4) becomes

$$\begin{aligned} S_s &= \ln 2 + \ln \pi - \frac{1}{\pi} \int_0^{2\pi} \cos^2(\phi - \phi_0) \ln \cos^2(\phi - \phi_0) d\phi \\ &= \ln 2 + \ln \pi + \ln \frac{4}{e} \approx 1.531 + \ln 2, \end{aligned}$$

and $\Gamma_s \approx 0.450$. For spin 1 we get $S_s \approx 1.270 + \ln 3$ with $\Gamma_s \approx 0.422$.

We conjecture that the spin entropy (4) depends only on the variables that define the spin entropy component along the z axis. This is further elaborated and utilized in Conjecture 1. We also conjecture that the ordering of the states according to the spin entropy can be obtained solely from the spin entropy component along the z axis. This conjecture implies that states with maximum and minimum spin entropy values are the ones with maximum and minimum spin entropy component along the z axis.

Minimum Entropy

The third law of thermodynamics establishes zero as the minimum of thermodynamics entropy. The use of differential entropy for the term $S_{s,\phi}^{z\perp}$ may suggest that the proposed entropy could be negative, but we will establish a positive minimum for it.

In quantum mechanics, ignoring the spin, the entropic uncertainty principle [1, 3, 11], establishes that $S_x + S_p \geq 3 \ln(e \hbar \pi)$, with equality for the normal distribution. This is a stronger bound than the Robertson's uncertainty principle, though equality is attained for the same normal distribution. A similar bound can now be derived for the spin entropy.

We will need the following lemma

Lemma 1 (Hausdorff-Young Inequality Holds). *The Hausdorff-Young inequality*

holds for the spin states phase space, i.e., for a number p in the interval $(1, 2]$ and for every conjugate exponent q satisfying $1 = \frac{1}{p} + \frac{1}{q}$, with $1 \leq p \leq 2$,

$$\left(\sum_{m=-s}^s |\alpha_{s,m}|^q \right)^{1/q} \leq \left((2\pi)^{\frac{p-2}{2}} \int_0^{2\pi} |\lambda_s(\phi)|^p d\phi \right)^{1/p}.$$

Proof. We first expand the set of coefficients $\alpha_{s,m}$ to include the ones with $|m| > s$, with increments by 1, and set $\alpha_{s,m} = 0$ for $|m| > s$. So we can write (3) as

$$\lambda_s(\phi) = \sum_{m=-\infty}^{\infty} \alpha_{s,m} \psi_{s,m}(\phi). \quad (5)$$

Note that the sum on m is either on all even or all odd multiples of $\frac{1}{2}$, depending on whether $2s$ is even or odd. Then, we rewrite (5) as

$$\lambda_s(\phi) = \frac{1}{\sqrt{2\pi}} \sum_{m=-\infty}^{\infty} \alpha_{s,m} e^{i[(2\theta(m)-1)s+m]\phi} = \frac{1}{\sqrt{2\pi}} \sum_{j=-\infty}^{\infty} \beta_j e^{ij\phi}, \quad (6)$$

where $j \in \mathbb{Z}$, $\beta_j = \alpha_{s,m}$ for $j = (2\theta(m) - 1)s + m$ and $\beta_j = 0$ otherwise. Clearly, for any value of r

$$\sum_{j=-\infty}^{\infty} |\beta_j|^r = \sum_{m=-s}^s |\alpha_{s,m}|^r.$$

Then (6) describes a Fourier series expansion of the periodic complex-valued function $\lambda_s(\phi)$. The Fourier inverse is

$$\frac{1}{\sqrt{2\pi}} \int_0^{2\pi} \lambda_s(\phi) e^{-ij\phi} d\phi = \beta_j \quad \text{or} \quad \int_0^1 \sqrt{2\pi} \lambda_s(2\pi\mu) e^{-2\pi i j \mu} d\mu = \beta_j.$$

Thus, the Hausdorff-Young inequality

$$\left(\sum_{j=-\infty}^{\infty} |\beta_j|^q \right)^{1/q} \leq \left(\int_0^1 \left| \sqrt{2\pi} \lambda_s(2\pi\mu) \right|^p d\mu \right)^{1/p} \quad (7)$$

holds. We can then replace back into (7) the variables $\alpha_{s,m}$'s and $\phi = 2\pi\mu$ to complete the proof. \square

Theorem 1. *The entropy (4) satisfies the inequality*

$$S_s \geq \ln 2\pi, \quad (8)$$

with equality attained for the eigenstates of the spin operator $S_z = s\hbar \cos \theta$.

Proof. For the proof of the inequality we adapt the work of [1, 3, 11] to the specific spin phase space.

First, due to Lemma 1 we can apply Beckner's theorem [1]

$$\left(\frac{1}{q}\right)^{\frac{n}{2q}} \left(\frac{1}{p}\right)^{-\frac{n}{2p}} \geq \frac{\left(\sum_{m=-s}^s |\alpha_{s,m}|^q\right)^{1/q}}{\left((2\pi)^{\frac{p-2}{2}} \int_0^{2\pi} |\lambda_s(\phi)|^p d\phi\right)^{1/p}},$$

where $\frac{1}{p} + \frac{1}{q} = 1$, $1 \leq p \leq 2$, and n is the dimensionality of the space where the functions are defined. For the sphere $n = 2$. We apply the logarithm on both sides of the inequality, noting that all the quantities are non-negative, so

$$\ln \left[\left(\frac{1}{q}\right)^{\frac{1}{q}} \left(\frac{1}{p}\right)^{-\frac{1}{p}} \right] + \ln \left((2\pi)^{\frac{p-2}{2}} \int_0^{2\pi} |\lambda_s(\phi)|^p d\phi \right)^{1/p} \geq \ln \left(\sum_{m=-s}^s |\alpha_{s,m}|^q \right)^{1/q}. \quad (9)$$

Substituting variables $p = 2\alpha$ and $q = 2\beta$, we get the constraint $\frac{\alpha}{1-\alpha} = -\frac{\beta}{1-\beta} \in [1, \infty)$ from $1 = \frac{1}{p} + \frac{1}{q}$. Multiplying the terms on both sides of (9) by either one of these equal and positive expressions $\frac{\alpha}{1-\alpha}$ or $-\frac{\beta}{1-\beta}$, and rearranging terms, we get

$$\begin{aligned} & \frac{1}{1-\alpha} \ln \left(\int_0^{2\pi} |\lambda_s(\phi)|^{2\alpha} d\phi \right) + \frac{1}{1-\beta} \ln \left(\sum_{m=-s}^s |\alpha_{s,m}|^{2\beta} \right) \\ & \geq \frac{1}{1-\alpha} \ln \left(\frac{1}{2\alpha} \right) + \frac{1}{1-\beta} \ln \left(\frac{1}{2\beta} \right) + \ln 2\pi, \end{aligned}$$

where the two first terms are the Rényi's differential entropy and the Rényi's entropy, forming an entropic inequality. Accordingly, taking the limits $\alpha \rightarrow 1^-$ and $\beta \rightarrow 1^+$, we obtain (8).

The minimum entropy is reached for the eigenstates of S_z , $|\xi_{s,z}\rangle = |\xi_{s,m_0}\rangle$, for any eigenvalue $\hbar m_0$ where $-s \leq m_0 \leq s$. Then $\alpha_{s,m} = \delta_{m,m_0}$, and so $\lambda_s(\phi) = \psi_{s,m_0}$. The probabilities are then $|\alpha_{s,m}|^2 = \delta_{m,m_0}$, and so the Shannon entropy is zero. The probability density is then $|\lambda_s(\phi)|^2 = \frac{1}{2\pi}$, with differential entropy $\ln 2\pi$. \square

Thus, the intrinsic-information is bounded as follows: $0 \leq I_s \leq \frac{1}{\ln 2\pi}$.

The entropic principle inequality for unbounded variables reaches equality, a minimum entropy, for a state that balances the two entropy terms in phase space, namely states that yield normal distributions on both variables [1, 3]. However, a bounded variable and the integer variable resulting from the Fourier series transformation creates a phase space with a continuous probability distribution for the bounded variable and discrete probabilities associated with the integer variable. Then, the minimum of the entropy occurs for quantum states that minimize the entropy term associated with the discrete probabilities, and for spin those are the eigenstates of the S_z, S^2 operators.

Preparing a system to align a spin state with a particular direction, an eigenstate $|\xi_{s,m=s}\rangle$ of a z axis, as it is done in many experiments with spin up along z , provides the knowledge of the spin along that direction. Thus it is expected that the intrinsic-information is gained by such a preparation.

System with Multiple Particles and Quantum Field Theory

We now consider the spin of a system with N particles of the same species, each with spin s . The maximum spin value s_{\max} occurs when all the N particles are aligned yielding $s_{\max} = Ns$. For a state with spin value s_{\max} , the possible z values

are $m = -s_{\max}, \dots, s_{\max}$. However, for the system with N particles all different values $s' \in [s_{\min}, s_{\max}]$ can occur, where $s_{\min} = s \bmod(N, 2)$ for fermions and $s_{\min} = 0$ for bosons. For each s' , all possible z components must be considered, so $m_{s'} = -s', \dots, s'$. The eigenstate basis of the S_z, S^2 operators associated with the N particles is then

$$\{|\xi_{N,s}\rangle\} = \bigcup_{s'=s_{\min}}^{s_{\max}} \bigcup_{m'=-s'}^{s'} |\xi_{s',m'}\rangle .$$

For example, two fermions with $s = \frac{1}{2}$ will produce four z states, namely three states with $s' = 1$ and $m' = -1, 0, 1$, and the fourth state with $s' = 0$ and $m' = 0$. Thus, the eigenstate basis of S_z, S^2 operators for two fermions is $\{|\xi_{2,1/2}\rangle\} = \{|\xi_{1,1}\rangle, |\xi_{1,0}\rangle, |\xi_{1,-1}\rangle, |\xi_{0,0}\rangle\}$. Expanding the proposed entropy (4) to N particles of the same species yields

$$\begin{aligned} S_{N,s,\phi,\theta} &= S_{N,s,\theta}^z + S_{N,s,\phi}^{z^\perp} = \sum_{s=s_{\min}}^{s_{\max}} \left(S_{s,\theta}^z + S_{s,\phi}^{z^\perp} \right) \\ &= - \sum_{s=s_{\min}}^{s_{\max}} \left(\sum_{m_s=-s}^s |\alpha_{s,m_s}|^2 \ln |\alpha_{s,m_s}|^2 - \int |\lambda_s(\phi)|^2 \ln |\lambda_s(\phi)|^2 d\phi \right) . \end{aligned} \quad (10)$$

where $1 = \sum_{m_s=-s}^s |\alpha_{s,m_s}|^2$. Thus, the phase space to describe a system of N particles of spin s consists of all the spheres with radiuses that are integer multiples of $\hbar s$, from $\hbar s_{\min}$ up to $\hbar s_{\max}$.

In quantum field theory (QFT), superpositions of states with any number of particles are also considered. One can write a state as

$$|\xi_{\text{QFT}}\rangle = \sum_{N=0}^{\infty} \gamma_N \sum_{s'=s_{\min}}^{s_{\max}} \sum_{m=-s'}^{s'} \alpha_{s',m} |\xi_{s',m}\rangle ,$$

where $\sum_{N=0}^{\infty} |\gamma_N|^2 = 1$. Due to the orthogonality of states with different number of

particles, it is straightforward to expand (10) to

$$S_{\text{QFT},s,\phi,\theta} = \sum_{N=0}^{\infty} |\gamma_N|^2 \left(S_{N,s,\theta}^z + S_{N,s,\phi}^{z^\perp} \right) = \sum_{N=0}^{\infty} |\gamma_N|^2 \left(\sum_{s'=s_{\min}}^{s_{\max}} \left(S_{s',\theta}^z + S_{s',\phi}^{z^\perp} \right) \right),$$

where both, s_{\min} and s_{\max} depend on N and s . Thus, in QFT the entropy is the weighted average of the entropy terms associated with the exact number of particles.

SPIN ENTROPY FOR ONE PARTICLE

We first analyze the spin entropy for a fermion with spin value $1/2$, then for a massive boson with spin 1, and we conclude this section by analyzing the photon entropy.

One fermion with spin $1/2$

A spin state of a particle with spin $1/2$, represented by a set of two orthonormal eigenstates $|+\rangle = |\xi_{1/2,1/2}\rangle = (1, 0)^\top$ and $|-\rangle = |\xi_{1/2,-1/2}\rangle = (0, 1)^\top$ of the operators (S^2, S_z) with associated eigenvalues $(s = 1/2, m = \pm 1/2)$, is

$$|\xi_{1/2,z}\rangle = e^{i\varphi} \left(e^{i\nu} \cos \theta_\alpha |+\rangle + \sin \theta_\alpha |-\rangle \right) = e^{i\varphi} \left(e^{i\nu} \cos \theta_\alpha, \sin \theta_\alpha \right)^\top \quad (11)$$

with DOFs $\theta_\alpha \in [0, \frac{\pi}{2}]$ and $\varphi, \nu \in [0, \pi)$.

Proposition 1 (spin entropy for $s = 1/2$). *The spin entropy (4) of a spin state with $s = 1/2$, described by (11), is*

$$S_{\frac{1}{2}}(\theta_\alpha) = -\cos^2 \theta_\alpha \ln \cos^2 \theta_\alpha - \sin^2 \theta_\alpha \ln \sin^2 \theta_\alpha + \ln 2\pi - \frac{1}{2\pi} \int_0^{2\pi} (1 + \sin 2\theta_\alpha \cos 2\phi) \ln (1 + \sin 2\theta_\alpha \cos 2\phi) d\phi, \quad (12)$$




Figure 1. A plot of spin entropy (12) vs $\theta_\alpha \in [0, \frac{\pi}{2}]$, for $s = 1/2$.

which, of course, is invariant under changes of the phase factors φ and ν .

Proof. A state $|\xi_{1/2}\rangle$ assigns the probability distribution along the z axis eigenstates $P_z = (|\langle +|\xi_{1/2}\rangle|^2, |\langle -|\xi_{1/2}\rangle|^2)^\top = (\cos^2 \theta_\alpha, \sin^2 \theta_\alpha)^\top$. Thus, from (2) and (3) we get $|\xi_{1/2,\phi}\rangle = \int e^{i\varphi} \frac{1}{\sqrt{2\pi}} (e^{i\nu} e^{i\phi} \cos \theta_\alpha + e^{-i\phi} \sin \theta_\alpha) |\phi\rangle d\phi$ and so $\rho_{\theta_\alpha,\nu}(\phi) = \frac{1}{2\pi} (1 + \sin 2\theta_\alpha \cos(2\phi + \nu))$. \square

For a visualization of the entropy, see Figure 1.

Spin 1

For massive particles with spin $s = 1$, the spin matrices are

$$S_x = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad S_y = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -i & 0 \\ i & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad S_z = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad S^2 = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix},$$

yielding a basis representation formed with the eigenvectors of S_z and S^2 : $|\uparrow\rangle_z = |\xi_{1,1}\rangle = (1, 0, 0)^\top$, $|\rightarrow\rangle_z = |\xi_{1,0}\rangle = (0, 1, 0)^\top$, $|\downarrow\rangle_z = |\xi_{1,-1}\rangle = (0, 0, 1)^\top$. A

general state of spin $s = 1$ in the basis aligned with the z axis is

$$|\xi_z\rangle = e^{i\varphi_y} \left(\sin \theta_\alpha \cos \theta_\beta e^{i\varphi_x} |\uparrow\rangle_z + \cos \theta_\alpha |\rightarrow\rangle_z + \sin \theta_\alpha \sin \theta_\beta e^{i\varphi_z} |\downarrow\rangle_z \right), \quad (13)$$

with the DOFs $\theta_\alpha, \theta_\beta \in [0, \frac{\pi}{2}]$, and $\varphi_x, \varphi_z, \varphi_y \in [0, 2\pi)$. Following (2), we write the state in the $|\phi\rangle$ basis as

$$|\xi_{z^\perp}\rangle = \frac{e^{i\varphi_y}}{\sqrt{2\pi}} \int \left(\sin \theta_\alpha \cos \theta_\beta e^{i(\varphi_x+2\phi)} + \cos \theta_\alpha e^{i\phi} + \sin \theta_\alpha \sin \theta_\beta e^{i(\varphi_z-2\phi)} \right) |\phi\rangle d\phi. \quad (14)$$

Proposition 2. *The spin entropy (4) of a particle with spin $s = 1$ in a state given by (13) is*

$$\begin{aligned} S_1 = & - \int_0^{2\pi} \rho_1(\phi, \varphi_x, \varphi_z, \theta_\alpha, \theta_\beta) \ln \rho_1(\phi, \varphi_x, \varphi_z, \theta_\alpha, \theta_\beta) d\phi \\ & + S_c(\cos^2 \theta_\alpha) + \sin^2 \theta_\alpha S_c(\cos^2 \theta_\beta), \end{aligned} \quad (15)$$

where

$$\begin{aligned} S_c(\cos^2 \theta) &= -\cos^2 \theta \ln(\cos^2 \theta) - (1 - \cos^2 \theta) \ln(1 - \cos^2 \theta) \\ \rho_1(\phi) &= \frac{1}{2\pi} \left| \sin \theta_\alpha \cos \theta_\beta e^{i(\varphi_x+2\phi)} + \cos \theta_\alpha e^{i\phi} + \sin \theta_\alpha \sin \theta_\beta e^{i(\varphi_z-2\phi)} \right|^2 \\ &= \frac{1}{2\pi} \left[1 + \sin^2 \theta_\alpha \sin(2\theta_\beta) \cos(4\phi + \varphi_x - \varphi_z) \right. \\ &\quad \left. + \sin(2\theta_\alpha) (\cos \theta_\beta \cos(\phi + \varphi_x) + \sin \theta_\beta \cos(3\phi - \varphi_z)) \right]. \end{aligned}$$

Proof. Computing the three probabilities associated with state $|\xi_z\rangle$ and deriving its entropy term $S_{1,\theta}^z$ yields

$$S_{1,\theta}^z = -\cos^2 \theta_\alpha \ln \cos^2 \theta_\alpha - \sin^2 \theta_\alpha \left[\ln \sin^2 \theta_\alpha - S_c(\cos^2 \theta_\beta) \right].$$

The entropy term $S_{1,\phi}^{z^\perp}$ is derived from the probability density associated with the state $|\xi_{z^\perp}\rangle$ from (14), and so

$$\rho_1(\phi) = \frac{1}{2\pi} \left| \sin \theta_\alpha \cos \theta_\beta e^{i(\varphi_x+2\phi)} + \cos \theta_\alpha e^{i\phi} + \sin \theta_\alpha \sin \theta_\beta e^{i(\varphi_z-2\phi)} \right|^2$$

□

Partially observed in simulations and in (12), we conjecture that

Conjecture 1. *The spin entropy (4) depends only on the variables that define the spin entropy component along the z axis. Other variables present in the state can be set to any convenient values as the spin entropy is invariant under changes of these values. In particular, the spin entropy (15) of a state given by (13) can be written more simply as*

$$S_1(\theta_\alpha, \theta_\beta) = - \int_0^{2\pi} \rho(\phi, \theta_\alpha, \theta_\beta) \ln \rho(\phi, \theta_\alpha, \theta_\beta) d\phi \quad (16)$$

$$+ S_c(\cos^2 \theta_\alpha) + \sin^2 \theta_\alpha S_c(\cos^2 \theta_\beta),$$

where

$$\rho(\phi, \theta_\alpha, \theta_\beta) = \frac{1}{2\pi} \left| \sin \theta_\alpha \cos \theta_\beta e^{i2\phi} + \cos \theta_\alpha e^{i\phi} + \sin \theta_\alpha \sin \theta_\beta e^{-i2\phi} \right|^2$$

which, of course, is invariant under changes of the phase factors φ_x and φ_z .

By Theorem 1, the minimum entropy does occur at $\theta_\alpha = 0$, and at $(\theta_\alpha = \frac{\pi}{2}, \theta_\beta = 0)$, and at $(\theta_\alpha = \frac{\pi}{2}, \theta_\beta = \frac{\pi}{2})$, the three eigenstates along the z axis. Thus, preparing a spin state orientation, by aligning it with an axis, reduces the entropy. The entropy given by (16) is visualized in Figure 2.




Figure 2. Spin entropy for $s = 1$ (16) vs $(\theta_\alpha, \theta_\beta)$. Note that for $\theta_\alpha = 0$ and for $\theta_\alpha = \frac{\pi}{2}$, $\theta_\beta = 0, \frac{\pi}{2}$, describing the three eigenstates along the z axis, the spin entropy reaches its minimum value 1.838, approximating $\ln 2\pi$.

Photon Entropy

Photon is a massless spin 1 particle. Its group representation is $ISO(2)$ or $E(2)$, with the gauge transformation accounting for the transverse direction. Then, this results in a particle with helicity ± 1 and a two dimensional polarization field in a plane propagating along or against the helicity direction. Formally one creates two circular polarized states, a right hand side, $|h_+\rangle$, and a left hand side, $|h_-\rangle$, which are the quantum states of the polarization field. Given an x, y, z coordinate system, we write $|h_+\rangle = \frac{1}{\sqrt{2}}(1, i)^T$ and $|h_-\rangle = \frac{1}{\sqrt{2}}(1, -i)^T$ in the basis $|x\rangle = (1, 0)^T$, $|y\rangle = (0, 1)^T$ in the plane perpendicular to z . The photon-spin operator along z is then

$$S_z = \hbar (|h_+\rangle \langle h_+| - |h_-\rangle \langle h_-|) = \hbar \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix},$$

with the two eigenstates, $|h_+\rangle, |h_-\rangle$ and eigenvalues ± 1 , which are the helicity values. The general state of the photon is then $|\psi\rangle = \alpha_+ |h_+\rangle + \alpha_- |h_-\rangle$ where $1 = |\alpha_+|^2 + |\alpha_-|^2$. The DOFs of the photon spin are captured by these two coefficients that produce the probabilities $|\alpha_+|^2, |\alpha_-|^2$ for the photon to be in each of the helicities eigenvectors.

The rotation operator $e^{-i\frac{S_z}{\hbar}\phi}$ of an angle ϕ around the z axis describes the polarization angle in the x - y plane. There is a clear parallel with the GC, where the polarization angle is the azimuth angle of the GC, i.e., the GQ of the spin is already used in physics to describe the photon. Robertson's uncertainty between the helicity and the angle ϕ has the physical intuition of being the uncertainty between the observed quantities angular momentum (helicity) and the polarization angle (ϕ). It is clear that the formalism for the photon is the same as the one obtained from the GQ of the spin $1/2$ particles state, where there are only two eigenvalues/eigenstates along the z axis. The spin entropy of the photon, quantified by the Shannon entropy along the z axis and the relative entropy on the polarization plane, captures this intrinsic randomness and it is given by (12).

TWO FERMIONS AND ENTANGLEMENT

We now examine the spin entropy of a system with two fermions, say A and B , with spin $s = 1/2$. We focus on entanglement. Consider the spin basis, the product of individual spin eigenstates along the z axis,

$$\begin{aligned} |++\rangle &= |+\rangle^A |+\rangle^B, & |+-\rangle &= |+\rangle^A |-\rangle^B, \\ |-+\rangle &= |-\rangle^A |+\rangle^B, & |--\rangle &= |-\rangle^A |-\rangle^B. \end{aligned}$$

Theorem 2 (Two spin $s = 1/2$ entanglement). *Consider a system with two identical particles, each with $s = 1/2$, in a subspace of spin 0 along z . The Ising-like spin-spin*

interaction is described by a Hamiltonian $H = \begin{pmatrix} E & 0 \\ 0 & E \end{pmatrix}$. A spinless environment state $|e\rangle$ with energy $\hbar\gamma$ produces an orthogonal basis of the system $|+-\rangle \otimes |e\rangle$ and $| -+\rangle \otimes |e\rangle$. Assume that due to the presence of the environment the two system states interact via a Hamiltonian $H^I = \hbar \begin{pmatrix} 0 & -i\omega \\ i\omega & 0 \end{pmatrix}$ with eigenvalues $\pm\omega$. Let the initial state be entangled $|\Psi^+, e\rangle = \frac{1}{\sqrt{2}}(|+-\rangle + |-+\rangle) \otimes |e\rangle$. Then, spin entropy of the evolution of this state $|\Psi^+(t), e(t)\rangle = U(t)|\Psi^+, e\rangle = e^{-i\frac{(E+\hbar\gamma)I+H^I}{\hbar}t} |\Psi^+, e\rangle$ is

$$S(t) = -\sin^2(\omega t) \ln \sin^2(\omega t) - \cos^2(\omega t) \ln \cos^2(\omega t) \quad (17)$$

$$- \int_0^{2\pi} \frac{(1 + \sin(2\omega t) \cos \phi)}{2\pi} \ln \frac{(1 + \sin(2\omega t) \cos \phi)}{2\pi} d\phi,$$

which is invariant under E and γ , and reaches its maximum at time $T = \frac{\pi}{4\omega}$, when the state is disentangled with probability 1.

Proof. The Hamiltonian can be diagonalized as

$$H + \hbar\gamma I + H^I = \frac{\hbar}{2} \begin{pmatrix} 1 & -i \\ i & -1 \end{pmatrix} \begin{pmatrix} \frac{E}{\hbar} + \gamma + \omega & 0 \\ 0 & \frac{E}{\hbar} + \gamma - \omega \end{pmatrix} \begin{pmatrix} 1 & -i \\ i & -1 \end{pmatrix}.$$

Thus, the unitary evolution operator $U(t)$ is

$$U(t) = e^{-i(\frac{E}{\hbar} + \gamma)t} \begin{pmatrix} \cos \omega t & \sin \omega t \\ -\sin \omega t & \cos \omega t \end{pmatrix}.$$

To evaluate the spin entropy, we develop the spin phase space for the two fermions case. Spin matrices associated with the two fermions can be written in terms of Pauli matrices $\sigma_x, \sigma_y, \sigma_z$ of the individual fermions as

$$S_{x,y,z} = \frac{1}{2} \sigma_{x,y,z} \otimes I + I \otimes \frac{1}{2} \sigma_{x,y,z} \quad \text{and} \quad S^2 = S_x^2 + S_y^2 + S_z^2,$$

and so

$$S_x = \frac{1}{2} \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix}, \quad S_y = \frac{1}{2} \begin{pmatrix} 0 & -i & -i & 0 \\ i & 0 & 0 & -i \\ i & 0 & 0 & -i \\ 0 & i & i & 0 \end{pmatrix}, \quad S_z = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}.$$

A basis constructed from the eigenvectors of S_z and S^2 is $\{|++\rangle, \frac{1}{\sqrt{2}}(|+-\rangle + |-+\rangle), \frac{1}{\sqrt{2}}(|+-\rangle - |-+\rangle), |--\rangle\}$. In particular, we are exploring the subspace with spin zero along the z axis ($m = 0$), i.e., the subspace generated by the two Bell states

$$|\Psi^+\rangle = |\xi_{2,1,0}\rangle = \frac{1}{\sqrt{2}}(|+-\rangle + |-+\rangle), \quad |\Psi^-\rangle = |\xi_{2,0,0}\rangle = \frac{1}{\sqrt{2}}(|+-\rangle - |-+\rangle),$$

which are entangled states and eigenstates of S_z with $m = 0$. The basis transformation from $|+\rangle|-\rangle, |+\rangle|-\rangle$ to $|\Psi^+\rangle, |\Psi^-\rangle$ is $B = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix}$ and so, in the basis $|\Psi^+, e\rangle, |\Psi^-, e\rangle$ the unitary evolution operator is

$$U_z(t) = BU(t)B^{-1} = e^{-i(\frac{E}{\hbar} + \gamma)t} \begin{pmatrix} \cos \omega t & -\sin \omega t \\ \sin \omega t & \cos \omega t \end{pmatrix},$$

where the subscript in U_z indicates that this representation is associated with eigenstates of the spin operators S_z, S^2 . Thus, $|\Psi^+(t), e(t)\rangle = U_z(t) |\Psi^+\rangle = e^{-i(\frac{E}{\hbar} + \gamma)t} \begin{pmatrix} \cos \omega t \\ \sin \omega t \end{pmatrix}$ with probabilities $P(t) = \begin{pmatrix} \cos^2 \omega t \\ \sin^2 \omega t \end{pmatrix}$. We now examine the entropy term of the state $|\Psi^+(t), e(t)\rangle$ in the conjugate basis $|\phi, e\rangle$. From (2)

$$|\Psi_{z^\perp}^+(t), e(t)\rangle = \frac{e^{-i(\frac{E}{\hbar} + \gamma)t}}{\sqrt{2\pi}} \int \left(e^{i\phi} \cos \omega t + \sin \omega t \right) |\phi, e\rangle d\phi,$$

implying that $\rho_{z^\perp}(\phi, t) = \frac{1}{2\pi} (1 + \sin(2\omega t) \cos \phi)$ and the entropy (10) can be readily




Figure 3. a. spin entropy (17) and its components $S^z(t)$, $S^{z^+}(t)$. b. spin entropy vs von Neumann entropy. The magnitudes are significantly different and their maxima and minima occur in opposition, at values of ωt that are multiples of $\frac{\pi}{4}$.

evaluated. There is an oscillation of period $\frac{\pi}{2\omega}$. The entropy is maximized at $T = \frac{\pi}{4\omega}$, where $|\Psi^+(T), e(T)\rangle = e^{-i(\frac{E}{\hbar}+\gamma)\frac{\pi}{4\omega}} \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = e^{-i(\frac{E}{\hbar}+\gamma)\frac{\pi}{4\omega}} |+-, e\rangle$. Thus, with probability 1, at $T = \frac{\pi}{4\omega}$ the spin state is in the disentangled state $|+-, e\rangle$. \square

For a visualization of this entropy $S(t)$ and of its two components, $S^z(t)$ and $S^{z^+}(t)$, see Figure 3a.

Work exists exploring the information content of entangled physical systems considering von Neumann entropy, e.g., [2, 5, 18, 22]. In that approach the required choice of basis functions is the product of one-particle eigenstates $|+\rangle |-\rangle$, $|+\rangle |-\rangle$ so that one can trace out over one-particle spin eigenstates. Thus, the evolution of the initial state $|\Psi^+, e\rangle = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ in this basis is described by

$$|\Psi^+(t), e(t)\rangle = U(t) |\Psi^+, e\rangle = e^{-i(\frac{E}{\hbar}+\gamma)t} \begin{pmatrix} \sin(\omega t + \frac{\pi}{4}) \\ \cos(\omega t + \frac{\pi}{4}) \end{pmatrix}.$$

After tracing out any of the two particles states, the probabilities of the resulting mixture state are $P_{A_+}(t) = \sin^2(\omega t + \frac{\pi}{4})$ and $P_{A_-}(t) = \cos^2(\omega t + \frac{\pi}{4})$ with von

Neumann entropy

$$S_{\text{vN}}(t) = -\cos^2\left(\omega t + \frac{\pi}{4}\right) \log \cos^2\left(\omega t + \frac{\pi}{4}\right) - \sin^2\left(\omega t + \frac{\pi}{4}\right) \log \sin^2\left(\omega t + \frac{\pi}{4}\right),$$

which is also plotted in Figure 3b for a visual comparison to the spin entropy.

Disentanglement occurs at time steps $t = \frac{\pi}{4\omega} + n\frac{\pi}{\omega}$; $n = 0, 1, \dots$. The spin entropy predicts highest entropy at these times, while the tracing out procedure predicts the lowest possible entropy, 0. This 0 value for the von Neumann entropy is to be expected since the mixture of states becomes a pure state when disentanglement occurs. Also, the Bell states (the eigenstates of S_z, S^2 for $m = 0$), are the ones that have the lowest spin entropy while yielding the highest von Neumann entropy. These min/max entropy differences is, technically, due to the spin basis describing the z axis probabilities. For the spin entropy, the spin basis needs to be the eigenfunctions of the two-particles spin operators S_z, S^2 , while for the von Neumann entropy, one must trace out the eigenstates of the individual particle eigenstates. The large entropy magnitude difference everywhere is due to the spin entropy's contribution from the ϕ variable, the conjugate variable to $z = \cos \theta$.

We point out that the environment was only modeled as one state, i.e., we did not consider the pioneer work [12, 24] of decoherence. The main reason is that our focus here is to demonstrate that for the entangled and the disentangled states, spin entropy behaves quite differently than von Neumann entropy. It would be interesting in future research to study how the two entropies can complement each other in the study of the impact of the environment on quantum states.

CONCLUSIONS

The concept of spin entropy in a spin phase space was proposed. The spin phase space of a particle is defined via the quantization of a sphere surface. The operators

associated with the spherical polar representation do not commute, yielding the uncertainty principle for the spin values in phase space. The proposed entropy captures the randomness present in the spin phase space, for any specified z axis and specified spin state for such axis. In the case of a photon, the z axis is the one of the propagation. The formulation is general for a system of many spin particles and extends to quantum fields. We studied spin entropy for a single particle with spin $1/2$, spin 1, photons, and for two entangled fermions of spin $1/2$.

The proposed entropy differs from von Neumann entropy, which is the one currently adopted in quantum information and quantum computing. The von Neumann entropy models the randomness associated with the unknown specifications of a state, and it is non zero only for a classical probabilistic mixture of quantum states. In contrast, our proposed entropy works with pure quantum states and models the intrinsic randomness of the observables: the possible observed spin values along any direction in three dimensions.

We have examined entangled states and their time evolution. Bell's entangled states that are eigenstates of the spin S_z, S^2 operators with total zero spin value along the z axis have lower spin entropy than that of the product of one particle states (disentangled states). In contrast, von Neumann entropy is highest for Bell's entangled states and lowest for disentangled states. We studied the dynamics of entangled states with an Ising-like Hamiltonian model of the interaction between spins, and a model of the environment as one state and its impact on a Hamiltonian mediating the interaction between Bell's entangled states. In a simulation, the time evolution of such Bell entangled states the entropy increases up to its maximum value when disentanglement occurs.

ACKNOWLEDGEMENT

This paper is partially based upon work supported by both the National Science Foundation under Grant No. DMS-1439786 and the Simons Foundation Institute Grant Award ID 507536 while the first author was in residence at the Institute for Computational and Experimental Research in Mathematics in Providence, RI, during the spring 2019 semester “Computer Vision” program.

-
- [1] W. Beckner. *Inequalities in Fourier analysis*. PhD thesis, Princeton., 1975.
 - [2] C. H. Bennett, H. J. Bernstein, S. Popescu, and B. Schumacher. Concentrating partial entanglement by local operations. *Phys. Rev. A*, 53:2046–2052, Apr 1996.
 - [3] I. Białyński-Birula and J. Mycielski. Uncertainty relations for information entropy in wave mechanics. *Commun. Math. Phys.*, 44(2):129–132, 1975.
 - [4] M. Blau. Symplectic geometry and geometric quantization. *Lecture Notes*, <http://www.ictp.trieste.it/~mblau/lecturesGQ.ps.gz>, 1992.
 - [5] P. Calabrese and J. Cardy. Entanglement entropy and quantum field theory. *J. Stat. Mech.*, 2004(06):P06002, 2004.
 - [6] R. Clausius. *The mechanical theory of heat: With its applications to the steam-engine and to the physical properties of bodies*. J. van Voorst, 1867.
 - [7] C. Cohen-Tannoudji, B. Diu, and F. Laloe. *Quantum Mechanics: Volume 1, Basic concepts, tool, and applications*. Wiley-VCH, 2 edition, 2019.
 - [8] P. A. M. Dirac. A theory of electrons and protons. *Proceedings of the Royal Society of London. Series A, Containing papers of a mathematical and physical character*, 126(801):360–365, 1930.
 - [9] D. Geiger and Z. M. Kedem. spin entropy, 2021. <https://arxiv.org/abs/2111.11605>.

- [10] W. Gerlach and O. Stern. Der experimentelle Nachweis des magnetischen Moments des Silberatoms. *Z. Phys.*, 8(1):110–111, 1922.
- [11] I. I. Hirschman. A note on entropy. *Am. J. Math.*, 79(1):152–156, 1957.
- [12] E. Joos and H. D. Zeh. The emergence of classical properties through interaction with the environment. *Zeitschrift für Physik B Condensed Matter*, 59(2):223–243, 1985.
- [13] L. D. Landau. The damping problem in wave mechanics. *Z. Phys*, 45(63):430–441, 1927.
- [14] M. Le Bellac. *Quantum physics*. Cambridge University Press, 2011.
- [15] V. P. Nair. Elements of geometric quantization and applications to fields and fluids. *arXiv preprint arXiv:1606.06407*, 2016.
- [16] M. A. Nielsen and I. L. Chuang. Quantum computation and quantum information. Cambridge University Press, 2011.
- [17] H. P. Robertson. The uncertainty principle. *Physical Review*, 34:163–164, Jul 1929.
- [18] M. Srednicki. Entropy and area. *Phys. Rev. Lett.*, 71:666–669, Aug 1993.
- [19] C. E. Shannon. A mathematical theory of communication. *The Bell system technical journal*, 27,3:379–423, 1948.
- [20] J. von Neumann. Mathematical foundations of quantum mechanics: New edition Princeton University Press, 2018.
- [21] S. Weinberg. *The quantum theory of fields: Volume 1, (Foundations)*. Cambridge University Press, 1995.
- [22] E. Witten. A mini-introduction to information theory. preprint arXiv:1805.11965, 2018.
- [23] N.M.J Woodhouse. *Geometric quantization*. Oxford University Press, 1997.
- [24] H. D. Zeh. On the interpretation of measurement in quantum theory. *Foundations of Physics*, 1(1):69–76, 1970.
- [25] W. H. Zurek. Pointer basis of quantum apparatus: Into what mixture does the wave packet collapse? *Physical review D*, 24(6):1516, 1981.

- [26] W. H. Zurek. Environment-induced superselection rules. *Physical review D*, 26(8):1862, 1982.