

HR-LENGTH OF A FREE GROUP VIA POLYNOMIAL FUNCTORS

SERGEI O. IVANOV AND ROMAN MIKHAILOV

ABSTRACT. We prove that for a subring $R \subseteq \mathbb{Q}$ and a free group F of rank at least 2 the length of the Bousfield's HR -localization tower for F is at least $\omega + \omega$. The key ingredient of the proof is the theory of polynomial functors over \mathbb{Q} .

1. Introduction

In [2], Bousfield developed a theory of localization of spaces with respect to a homology theory. In particular, for a ring R , which is either a subring of \mathbb{Q} or $R = \mathbb{Z}/n$, he considers the localization with respect to $H_*(-, R)$. He proved that on the level of fundamental groups this localization corresponds to the so called HR -localization of groups that we denote by

$$(1.1) \quad \eta : G \longrightarrow L^R G.$$

Later [1] he found a construction of the group $L^R G$ as a limit of a natural transfinite tower $L_\alpha^R G$, where α is an ordinal, $L_\alpha^R G$ are constructed inductively, and $L^R G = L_\alpha^R G$ for a large enough α . In fact $L_\alpha^R G = L^R G / \gamma_\alpha^R(L^R G)$, where γ_α^R denotes the transfinite lower R -central series. The HR -localization of a group G is closely related to its R -completion \widehat{G}_R . Namely, if $H_1(G, R)$ is finitely generated over R , then $L_\omega^R G = \widehat{G}_R$.

The HR -length of a group G is defined as the minimal ordinal α such that $L^R G = L_\alpha^R G$. This invariant of a group is interesting because if two spaces are R -homologically equivalent, then the HR -lengths of their fundamental groups are the same. For the case $R = \mathbb{Z}$ this notion studied by authors in [11]. In particular, we found a characterisation of matrices $a \in \mathrm{GL}_n(\mathbb{Z})$ such that the $H\mathbb{Z}$ -length of the group $\mathbb{Z} \rtimes_a \mathbb{Z}^n$ is ω or $\omega + 1$ and constructed a matrix $a \in \mathrm{GL}_2(\mathbb{Z})$ such that the $H\mathbb{Z}$ -length of $\mathbb{Z} \rtimes_a \mathbb{Z}^2$ is at least $\omega + 2$. The characterization is given on the language of eigenvalues of the matrices.

In this paper we study the HR -localization of a free group F of rank at least 2. Bousfield in [1] proved that the integral homology of the pronilpotent completion $H_2(\widehat{F}_{\mathbb{Z}}, \mathbb{Z})$ is nontrivial. This is equivalent to the fact that $H\mathbb{Z}\text{-length}(F) \geq \omega + 1$. The authors proved in [8] that $H_2(\widehat{F}_{\mathbb{Q}}, \mathbb{Q}) \neq 0$ and in [10] that $H_2(\widehat{F}_{\mathbb{Z}/p}, \mathbb{Z}/p) \neq 0$ which is equivalent to $H\mathbb{Q}\text{-length}(F) \geq \omega + 1$ and $H\mathbb{Z}/p\text{-length}(F) \geq \omega + 1$. In [11] we proved that $H\mathbb{Z}\text{-length}(F) \geq \omega + 2$. This paper is devoted to the following theorem.

Theorem. *For a subring $R \subseteq \mathbb{Q}$ and a free group F of rank at least 2 we have*

$$HR\text{-length}(F) \geq \omega + \omega.$$

The key ingredient of the proof is the theory of polynomial functors over \mathbb{Q} . We construct a group $\mathcal{G}(V)$ that depends naturally on a finite dimensional \mathbb{Q} -vector space V and reduce the question about the free group to the question about the group $\mathcal{G}(V)$, that we solve using the theory of polynomial functors. We use the fact that any polynomial functor $\mathbf{Vect}^{\text{fin}}(\mathbb{Q}) \rightarrow \mathbf{Vect}(\mathbb{Q})$ can be decomposed as a direct sum of homogeneous functors [3, Prop.2.8] and that there are no natural transformations between homogeneous functors of different degrees. We use this fact in our analysis of the differentials of the spectral sequences of the extensions $L_{\omega+n+1}^{\mathbb{Q}}\mathcal{G}(V) \rightarrow L_{\omega+n}^{\mathbb{Q}}\mathcal{G}(V)$ decomposing the terms of the spectral sequences into homogeneous polynomial functors.

2. Polynomial functors over \mathbb{Q}

In this section we recall the definition of polynomial functors in the sense of Eilenber-MacLane and state a result about decomposition of polynomial functors over \mathbb{Q} into direct sum of homogeneous components. The main references for us are [6, §9] and [3] (see also [4]).

2.1. Polynomial functors. Further in this section \mathcal{A} denotes an additive category and \mathcal{V} denotes an abelian category. For a functor $\Phi : \mathcal{A} \rightarrow \mathcal{V}$ and $n \geq 0$ the n -th crossed effect of Φ is a functor $\text{cr}_n(\Phi) : \mathcal{A}^n \rightarrow \mathcal{V}$ given by

$$(2.1) \quad \text{cr}_n(\Phi)(A_1, \dots, A_n) = \text{Ker} \left(\Phi \left(\bigoplus_{i=1}^n A_i \right) \longrightarrow \bigoplus_{i=1}^n \Phi \left(\bigoplus_{j \neq i} A_j \right) \right),$$

where the homomorphisms are induced by the canonical projections. In particular, $\text{cr}_0(\Phi) = \Phi(0)$ and $\text{cr}_1(\Phi)(A) = \text{Ker}(\Phi(A) \rightarrow \Phi(0))$. In fact, the cross effect is a direct summand and there is a decomposition

$$(2.2) \quad \Phi \left(\bigoplus_{i=1}^n A_i \right) = \bigoplus_{s=0}^n \bigoplus_{1 \leq j_1 < \dots < j_s \leq n} \text{cr}_s(\Phi)(A_{j_1}, \dots, A_{j_s}).$$

A functor Φ is called polynomial (in the sense of Eilenbrg-MacLane) of degree $\leq d$ if $\text{cr}_{d+1}(\Phi) = 0$. It is easy to see that the functor $\text{cr}_{d+1} : \mathbf{Fun}(\mathcal{A}, \mathcal{V}) \rightarrow \mathbf{Fun}(\mathcal{A}^{d+1}, \mathcal{V})$ is exact, and hence, the class of functors of degree $\leq d$ is closed under extensions and subquotients.

2.2. Homogeneous components of polynomial functors over \mathbb{Q} . A polynomial functor $\Phi : \mathcal{A} \rightarrow \mathcal{V}$ is called homogeneous of degree d if $\Phi(\lambda \cdot \text{id}_A) = \lambda^d \cdot \text{id}_{\Phi(A)}$ for any $\lambda \in \mathbb{Z}$ and any object A of \mathcal{A} .

Further in this section we assume that K is an algebra over \mathbb{Q} and $\mathcal{V} = \mathbf{Mod}(K)$.

Lemma 2.1. *If $\Phi, \Psi : \mathcal{A} \rightarrow \mathbf{Mod}(K)$ are two homogeneous functors of different degrees, then $\mathbf{Hom}(\Phi, \Psi) = 0$.*

Proof. Denote by d and d' the degrees of Φ and Ψ . For any natural transformation $f : \Phi \rightarrow \Psi$ and any object A of \mathcal{A} we have $f_A \circ \Phi(\lambda \cdot \text{id}_A) = \Psi(\lambda \cdot \text{id}_A) \circ f_A$. Hence $\lambda^d f_A = \lambda^{d'} f_A$ for any $\lambda \in \mathbb{Z}$. Since $\mathbf{Hom}(\Phi(A), \Psi(A))$ is a \mathbb{Q} -vector space, this implies $f_A = 0$. \square

Theorem 2.2 ([3, Prop.2.8]). *Let K be an algebra over \mathbb{Q} . Then for any polynomial functor $\Phi : \mathcal{A} \rightarrow \text{Mod}(K)$ there is an isomorphism*

$$(2.3) \quad \Phi \cong \Phi_0 \oplus \Phi_1 \oplus \cdots \oplus \Phi_d,$$

where Φ_i is a homogeneous functor of degree i .

Lemma 2.1 implies that for two polynomial functors $\Phi, \Psi : \mathcal{A} \rightarrow \text{Mod}(K)$ we have

$$(2.4) \quad \text{Hom}(\Phi, \Psi) \cong \bigoplus_i \text{Hom}(\Phi_i, \Psi_i),$$

where $\Phi \cong \bigoplus \Phi_i$ and $\Psi \cong \bigoplus \Psi_i$ are decompositions as in Theorem 2.2. In particular, it follows that the components Φ_i of Φ are uniquely defined up to isomorphism. It also follows that the class of homogeneous functors of degree d is closed under subquotients and extensions.

Further we will use the following notation for a polynomial functor $\Phi : \mathcal{A} \rightarrow \text{Mod}(K)$

$$(2.5) \quad \min.\text{deg}(\Phi) = \min\{i \geq 0 \mid \Phi_i \neq 0\}.$$

If $\Phi = 0$, we set $\min.\text{deg}(0) = \infty$.

3. R -stem extensions

Eckmann, Hilton and Stambach in [5] proved that for a central extension

$$(3.1) \quad 1 \longrightarrow A \longrightarrow E \longrightarrow G \longrightarrow 1$$

there is an exact sequence

$$(3.2) \quad H_3(E) \rightarrow H_3(G) \rightarrow (E_{ab} \otimes A)/U \xrightarrow{\kappa} H_2(E) \rightarrow H_2(G) \rightarrow A \rightarrow E_{ab} \rightarrow G_{ab},$$

where $H_*(-) = H_*(-, \mathbb{Z})$ and U is defined as follows. We consider the map $\mu : A \otimes A \rightarrow E_{ab} \otimes A$ induced by the map $A \rightarrow E$ and set

$$(3.3) \quad U = \mu(\tilde{\Gamma}^2 A),$$

where $\tilde{\Gamma}^2 A = \text{Ker}(A \otimes A \rightarrow \Lambda^2 A)$. Moreover, the sequence is natural with respect to the central extension.

Eckmann-Hilton-Stambach say that a central extension is stem if the map $E_{ab} \rightarrow G_{ab}$ is an isomorphism and prove that in this case $U = 0$. Further we give a slight generalisation of this result. If R is a subring of \mathbb{Q} , we say that a central extension is R -stem if the map $H_1(E, R) \rightarrow H_1(G, R)$ is an isomorphism.

Proposition 3.1. *Let R be a subring of \mathbb{Q} and (3.1) be an R -stem extension. Then there is an exact sequence*

$$(3.4) \quad H_3(E) \rightarrow H_3(G) \rightarrow H_1(G) \otimes H_1(A) \xrightarrow{\kappa} H_2(E) \rightarrow H_2(G) \rightarrow H_1(A) \rightarrow 0,$$

where $H_*(-) = H_*(-, R)$. Moreover, this sequence is natural in two following senses:

- a morphism of R -stem extensions induces a morphism of the exact sequences;
- if $R \subseteq S \subseteq \mathbb{Q}$ then any R -stem extension is also an S -stem extension and the exact sequence for S is the exact sequence for R tensored by S .

Proof. Since R is torsion free, the functor of tensor product $- \otimes R$ is exact. Consider the sequence (3.2) tensored by R . Since $A \rightarrow E_{ab} \rightarrow G_{ab} \rightarrow 0$ is exact and $E_{ab} \otimes R \rightarrow G_{ab} \otimes R$ is an isomorphism, we obtain that the map $A \otimes R \rightarrow E_{ab} \otimes R$ is trivial. Hence, the map $A \otimes A \otimes R \rightarrow E_{ab} \otimes A \otimes R$ is also trivial and $((E_{ab} \otimes A)/U) \otimes R = E_{ab} \otimes A \otimes R = H_1(E) \otimes H_1(A)$. The naturalness are obvious from the construction. \square

4. Universal relative R -central extensions

In this section for a subring $R \subseteq \mathbb{Q}$ we give a definition of a relative R -central extension and R -perfect homomorphism. For any R -perfect homomorphism f we construct a relative R -central extension $E^R(f)$ that we call the universal relative R -central extension (but we do not prove any universal property). Universal properties of this construction are not important for us. For us it is important that this construction is natural in all parameters, including the subring $R \subseteq \mathbb{Q}$, and that this construction can be used to construct the HR -localization of a group [1, §3.4]. For the case $R = \mathbb{Z}$ a detailed theory of relative central extensions was developed in [7] and [11].

An R -central extension is a central extension, where the kernel is an R -module. A relative R -central extension (under a group Γ) is a commutative diagram of groups

$$(4.1) \quad \begin{array}{ccccccc} & & & \Gamma & & & \\ & & & \downarrow & \searrow f & & \\ 0 & \longrightarrow & A & \longrightarrow & E & \longrightarrow & G \longrightarrow 1 \end{array}$$

where the bottom line is an R -central extension.

For a group homomorphism $f : \Gamma \rightarrow G$ we consider the space

$$(4.2) \quad C(f) = \text{Cone}(K(\Gamma, 1) \rightarrow K(G, 1))$$

and denote by $H_*(f, R)$ (resp. $H^*(f, R)$) the homology (resp. cohomology) of the space $C(f)$ with coefficients in R . Then there is an exact sequence

$$(4.3) \quad H_2(\Gamma, R) \rightarrow H_2(G, R) \rightarrow H_2(f, R) \rightarrow H_1(\Gamma, R) \rightarrow H_1(G, R) \rightarrow H_1(f, R) \rightarrow 0.$$

We say that f is R -perfect if $H_1(f, R) = 0$ (i.e. $H_1(\Gamma, R) \rightarrow H_1(G, R)$ is surjective). For an R -perfect homomorphism f the universal coefficient theorem says that we have an isomorphism

$$(4.4) \quad H^2(f, H_2(f, R)) \cong \text{Hom}(H_2(f, R), H_2(f, R)).$$

The cohomology class corresponding to $\text{id}_{H_2(f, R)}$ is denoted by t_f^R . It corresponds to a map $C(f) \rightarrow K(H_2(f, R), 2)$ whose composition with $K(G, 1) \rightarrow C(f)$ is denoted by $\tilde{t}_f^R : K(G, 1) \rightarrow K(H_2(f, R), 2)$. Consider the principal fibration

$$(4.5) \quad \begin{array}{ccccc} K(H_2(f, R), 1) & \longrightarrow & X & \longrightarrow & K(G, 1) \\ \downarrow & & \downarrow & & \downarrow \tilde{t}_f^R \\ K(H_2(f, R), 1) & \longrightarrow & P(K(H_2(f, R), 2)) & \longrightarrow & K(H_2(f, R), 2) \end{array}$$

and set

$$(4.6) \quad E^R(f) = \pi_1(X).$$

Applying π_1 , we obtain an R -central extension $H_2(f, R) \twoheadrightarrow E^R(f) \twoheadrightarrow G$. Since the composition map $K(\Gamma, 1) \rightarrow K(G, 1) \rightarrow K(H_2(f, R), 2)$ is null-homotopic, we obtain a map to the path space such that the square

$$(4.7) \quad \begin{array}{ccc} K(\Gamma, 1) & \longrightarrow & K(G, 1) \\ \downarrow & & \downarrow \\ P(K(H_2(f, R), 2)) & \longrightarrow & K(H_2(f, R), 2). \end{array}$$

is commutative. This defines a map $K(\Gamma, 1) \rightarrow X$ such that its composition with $X \rightarrow K(G, 1)$ is the map $K(\Gamma, 1) \rightarrow K(G, 1)$. The induced map on fundamental groups is denoted by $e^R(f) : \Gamma \rightarrow E^R(f)$. Therefore, any R -perfect homomorphism $f : \Gamma \rightarrow G$ defines a relative R -central extension

$$(4.8) \quad \begin{array}{ccccccc} & & & \Gamma & & & \\ & & & \downarrow e^R(f) & \searrow f & & \\ 0 & \longrightarrow & H_2(f, R) & \longrightarrow & E^R(f) & \longrightarrow & G \longrightarrow 1. \end{array}$$

It is easy to check that $E^R(f)$ is natural in R and f .

5. HR -localization

5.1. HR -localization. Further in this section R denotes a fixed subring of \mathbb{Q} and we set

$$(5.1) \quad H_*(-) = H_*(-, R).$$

For a group G we denote by LG the HR -localization of G . If we want to emphasize the ring R , we use the notation $L^R G$. We denote by $\gamma_\alpha G = \gamma_\alpha^R G$ the transfinite R -lower central series of G , which is defined by the equations $\gamma_1 G = G$,

$$(5.2) \quad \gamma_{\alpha+1} G = \text{Ker} \left(\gamma_\alpha G \rightarrow \frac{\gamma_\alpha G}{[\gamma_\alpha G, G]} \otimes R \right), \quad \gamma_\beta G = \bigcap_{\alpha < \beta} \gamma_\alpha G,$$

for a limit ordinal β . Finally we set

$$(5.3) \quad L_\alpha G = LG / \gamma_\alpha(LG)$$

and denote by $\eta_\alpha : G \rightarrow L_\alpha G$ the composition homomorphism. We also set

$$(5.4) \quad h_\alpha G = H_2(\eta_\alpha) = \text{Coker}(H_2(G) \rightarrow H_2(L_\alpha G)).$$

Again, if we want to emphasize the ring R , we use the notation $h_\alpha^R G$.

Bousfield proved [1, §3.4] that for any α we have

$$(5.5) \quad L_{\alpha+1} G = E^R(\eta_\alpha), \quad \eta_{\alpha+1} = e^R(\eta_\alpha).$$

Hence we obtain a universal R -central extension (see (4.8)):

$$(5.6) \quad \begin{array}{ccccccc} & & & G & & & \\ & & & \downarrow \eta_{\alpha+1} & \searrow \eta_\alpha & & \\ 0 & \longrightarrow & h_\alpha G & \longrightarrow & L_{\alpha+1} G & \longrightarrow & L_\alpha G \longrightarrow 1. \end{array}$$

The HR -length of a group G can be defined as

$$(5.7) \quad HR\text{-length}(G) = \sup\{\alpha \mid h_\alpha G \neq 0\}$$

If $R \subseteq S \subseteq \mathbb{Q}$ then any S -local group is R -local, and we obtain homomorphisms under G :

$$(5.8) \quad \varphi : L^R G \longrightarrow L^S G, \quad \varphi_\alpha : L_\alpha^R G \longrightarrow L_\alpha^S G.$$

Using that the construction of universal relative R -central extension is natural, we obtain that the diagram

$$(5.9) \quad \begin{array}{ccccccc} 0 & \longrightarrow & h_\alpha^R G & \longrightarrow & L_{\alpha+1}^R G & \longrightarrow & L_\alpha^R G \longrightarrow 1 \\ & & \downarrow \psi_\alpha & & \downarrow \varphi_{\alpha+1} & & \downarrow \varphi_\alpha \\ 0 & \longrightarrow & h_\alpha^S G & \longrightarrow & L_{\alpha+1}^S G & \longrightarrow & L_\alpha^S G \longrightarrow 1, \end{array}$$

is commutative, where $\psi_\alpha : h_\alpha^R G \rightarrow h_\alpha^S G$ is induced by the map $H_2(L_\alpha^R G, R) \rightarrow H_2(L_\alpha^S G, S)$.

Proposition 5.1. *Let G be a group and α be an ordinal number. Then there exists an exact sequence*

$$(5.10) \quad H_3(L_{\alpha+1}) \rightarrow H_3(L_\alpha) \xrightarrow{\delta} H_1(G) \otimes h_\alpha \xrightarrow{\kappa} H_2(L_{\alpha+1}) \rightarrow H_2(L_\alpha) \rightarrow h_\alpha,$$

where $L_\beta = L_\beta^R G$ and $h_\alpha = h_\alpha^R G$ and $H_*(-) = H_*(-, R)$. Moreover, if we denote this sequence by $\mathcal{A}(G, \alpha, R)$, it is natural in the following two senses:

- a group homomorphism $G \rightarrow G'$ induces a morphism of the sequences $\mathcal{A}(G, \alpha, R) \rightarrow \mathcal{A}(G', \alpha, R)$;
- for any $R \subseteq S \subseteq \mathbb{Q}$ the morphism of central extensions (5.9) induces a morphism $\mathcal{A}(G, \alpha, R) \rightarrow \mathcal{A}(G, \alpha, S)$.

Proof. It follows from Proposition 3.1 and using that (5.6) is an R -stem extension. The naturalness in R and G follows from the fact that E_f^R is natural in f and R , for $R \subseteq S \subseteq \mathbb{Q}$ any R -stem extension is an S -stem extension and the fact that the sequence (3.4) is natural in the R -stem extensions. \square

5.2. Natural maps connecting $h_{\alpha+n}$. In Proposition 5.1 we have a map

$$(5.11) \quad \kappa : H_1(G) \otimes h_\alpha G \longrightarrow H_2(L_{\alpha+1} G).$$

We can compose it with the obvious projection and obtain a map

$$(5.12) \quad \bar{\kappa} : H_1(G) \otimes h_\alpha G \longrightarrow h_{\alpha+1} G.$$

Then inductively for $n \geq 0$ we can define a natural map

$$(5.13) \quad \bar{\kappa}_n : H_1(G)^{\otimes n} \otimes h_\alpha G \longrightarrow h_{\alpha+n} G$$

such that $\bar{\kappa}_n = \bar{\kappa} \circ (\text{id} \otimes \bar{\kappa}_{n-1})$ and $\bar{\kappa}_0 = \text{id} : h_\alpha G \rightarrow h_\alpha G$. Any element $g \in G$ defines an element $g' \in H_1(G)$ and we consider the map

$$(5.14) \quad \bar{\kappa}_{n,g} : h_\alpha G \longrightarrow h_{\alpha+n} G,$$

defined by $\bar{\kappa}_{n,g}(x) = \bar{\kappa}_n(g' \otimes \cdots \otimes g' \otimes x)$.

Note that Proposition 5.1 implies that the map $\bar{\kappa}_{n,g}^R$ is natural in the following sense. Let $f : G \rightarrow G'$ be a homomorphism and $g \in G, g' \in G'$ be elements such

that $f(g) = g'$ and $R \subseteq S \subseteq \mathbb{Q}$. Then the diagram

$$(5.15) \quad \begin{array}{ccc} h_\alpha^R G & \xrightarrow{\bar{r}_{n,g}^R} & h_{\alpha+n}^R G \\ \downarrow & & \downarrow \\ h_\alpha^S G' & \xrightarrow{\bar{r}_{n,g'}^S} & h_{\alpha+n}^S G' \end{array}$$

is commutative.

6. The $H\mathbb{Q}$ -localization of the V -lamplighter group

In this section everything is rational, $R = \mathbb{Q}$,

$$(6.1) \quad h_\alpha = h_\alpha^\mathbb{Q}, \quad H_*(-) = H_*(-, \mathbb{Q}).$$

We denote by $C = \langle t \rangle$ the infinite cyclic group, by $\mathbb{Q}[C] = \mathbb{Q}[t, t^{-1}]$ the group algebra and by $I \triangleleft \mathbb{Q}[C]$ the augmentation ideal, which is generated by one element $t - 1$.

6.1. $\mathbb{Q}[C]$ -modules. For a $\mathbb{Q}[C]$ -module M we denote by \widehat{M}_I the I -adic completion $\widehat{M}_I = \varprojlim M/MI^n$. It is easy to check that the \mathbb{Q} -adic completion of $\mathbb{Q}[C]$ is the ring of power series $\widehat{\mathbb{Q}[C]}_I = \mathbb{Q}[[x]]$, where the map $\mathbb{Q}[C] \rightarrow \mathbb{Q}[[x]]$ is induced by $t \mapsto 1 + x$. Therefore, there is a natural structure of $\mathbb{Q}[[x]]$ -module on the I -adic completion \widehat{M}_I .

Consider the group $C \otimes \mathbb{Q}$, which is isomorphic to \mathbb{Q} but written multiplicatively $C \otimes \mathbb{Q} = \{t^\alpha \mid \alpha \in \mathbb{Q}\}$. Then there is a multiplicative homomorphism

$$(6.2) \quad C \otimes \mathbb{Q} \longrightarrow \mathbb{Q}[[x]], \quad t^\alpha \mapsto (1+x)^\alpha = \sum_{n \geq 0} \binom{\alpha}{n} x^n,$$

where $\binom{\alpha}{n} = \alpha(\alpha-1)\dots(\alpha-n+1)/n!$. Therefore, for any $\mathbb{Q}[C]$ -module M there is a natural structure of $\mathbb{Q}[C \otimes \mathbb{Q}]$ -module on \widehat{M}_I .

6.2. Free $\mathbb{Q}[C]$ -modules. For a finite dimensional vector space V we consider the free $\mathbb{Q}[C]$ -module

$$(6.3) \quad \mathcal{M} = \mathcal{M}(V) := \mathbb{Q}[C] \otimes V.$$

We will think about it as about a functor from the category of finite dimensional vector spaces over \mathbb{Q}

$$(6.4) \quad \mathcal{M} : \mathbf{Vect}^{\text{fin}}(\mathbb{Q}) \rightarrow \mathbf{Mod}(\mathbb{Q}[C]).$$

It is easy to check that the I -completion of this module can be computed as

$$(6.5) \quad \widehat{\mathcal{M}}_I = \mathbb{Q}[[x]] \otimes V,$$

where the map $\mathcal{M} \rightarrow \widehat{\mathcal{M}}_I$ is induced by the map $\mathbb{Q}[C] \rightarrow \mathbb{Q}[[x]]$, $t \mapsto 1 + x$.

Lemma 6.1. *For any $n \geq 1$ we have*

$$H_1(C, \Lambda^n(\mathcal{M})) = 0, \quad H_1(C \otimes \mathbb{Q}, \Lambda^n(\widehat{\mathcal{M}}_I)) = 0.$$

Proof. First we prove that $H_1(C, \Lambda^n(\mathcal{M})) = 0$. For any C -module M there is an isomorphism $H_1(C, M) \cong H^0(C, M) = M^C$. So in order to prove the first equation we need to prove that the module $\Lambda^n(\mathcal{M})$ has no invariant elements. For any torsion free group A there is a standard embedding $\Lambda^n(A) \rightarrow A^{\otimes n}$, given by $a_1 \wedge \cdots \wedge a_n \mapsto \sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{\sigma(1)} \otimes \cdots \otimes a_{\sigma(n)}$. If A is a C -module, this embedding respects the C -action. Then it is enough to prove that $\mathcal{M}^{\otimes n}$ has no non-trivial invariant elements. Note that $\mathcal{M}^{\otimes n} = \mathbb{Q}[t_1^{\pm 1}, \dots, t_n^{\pm 1}] \otimes V^{\otimes n}$ with the action of t induced by multiplication on the product $t_1 \cdots t_n$. Since the functor of invariants is additive and $V \cong \mathbb{Q}^k$ for some k , it is enough to prove that $\mathbb{Q}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ has no non-trivial invariants. Finally $\mathbb{Q}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ has no non-trivial invariants of the action by $t_1 \cdots t_n$ because it is an integral domain.

Now we prove that $H_1(C \otimes \mathbb{Q}, \Lambda^n(\widehat{\mathcal{M}}_I)) = 0$. The group \mathbb{Q} is the union of the groups $\frac{1}{m}\mathbb{Z}$. Since homology commute with filtered colimits, it is enough to prove that

$$(6.6) \quad H_1(C \otimes \frac{1}{m}\mathbb{Z}, \Lambda^n(\widehat{\mathcal{M}}_I)) = 0.$$

The group $C \otimes \frac{1}{m}\mathbb{Z}$ is cyclic, so it is enough to prove there are no non-trivial invariants of the multiplication of $t^{\frac{1}{m}}$ on $\Lambda^n(\widehat{\mathcal{M}}_I)$. As in the previous case, we can embed the exterior power into the tensor power $(\widehat{\mathcal{M}}_I)^{\otimes n} = \mathbb{Q}[[x]]^{\otimes n} \otimes V^{\otimes n}$ and reduce the question to the question about $t^{\frac{1}{m}}$ -invariant elements of $\mathbb{Q}[[x]]^{\otimes n}$. Further, note that there is an embedding of $C \otimes \mathbb{Q}$ -modules $\mathbb{Q}[[x]]^{\otimes n} \hookrightarrow \mathbb{Q}[[x_1, x_2, \dots, x_n]]$ where $t^{\frac{1}{m}}$ acts on the algebra of power series by the multiplication on the power series $(1+x_1)^{\frac{1}{m}} \cdots (1+x_n)^{\frac{1}{m}}$. Since $\mathbb{Q}[[x_1, \dots, x_n]]$ is an integral domain, we obtain that it has no non-trivial $t^{\frac{1}{m}}$ -invariant elements. \square

6.3. Semidirect products $C \rtimes M$. Let M be an $\mathbb{Q}[C]$ -module. The \mathbb{Q} -completion of the semidirect product $C \rtimes M$ can be computed by the formula

$$(6.7) \quad (C \rtimes M)_{\mathbb{Q}}^{\wedge} = (C \otimes \mathbb{Q}) \rtimes \widehat{M}_I$$

(see [9, Prop.4.7]).

For any abelian group A of homological dimension 1 and any $\mathbb{Q}[A]$ -module M the spectral sequence of the extension $M \twoheadrightarrow A \rtimes M \twoheadrightarrow A$ for any $n \geq 1$ gives short exact sequences

$$(6.8) \quad 0 \rightarrow H_0(A, H_n(M)) \rightarrow H_n(A \rtimes M) \rightarrow H_1(A, H_{n-1}(M)) \rightarrow 0.$$

The first homology can be computed directly:

$$(6.9) \quad H_1(A \rtimes M) = (A \otimes \mathbb{Q}) \oplus M_A.$$

Since $H_*(M) = \Lambda^* M$, we obtain

$$(6.10) \quad 0 \rightarrow (\Lambda^n M)_C \rightarrow H_n(C \rtimes M) \rightarrow H_1(C, \Lambda^{n-1} M) \rightarrow 0$$

and

$$(6.11) \quad 0 \rightarrow (\Lambda^n \widehat{M}_I)_{C \otimes \mathbb{Q}} \rightarrow H_n((C \rtimes M)_{\mathbb{Q}}^{\wedge}) \rightarrow H_1(C \otimes \mathbb{Q}, \Lambda^{n-1} \widehat{M}_I) \rightarrow 0.$$

6.4. V -lamplighter group. For a finite dimensional \mathbb{Q} -vector space V we consider the semidirect product

$$(6.12) \quad \mathcal{G} = \mathcal{G}(V) = C \ltimes \mathcal{M}(V) = C \ltimes (\mathbb{Q}[C] \otimes V).$$

We consider the group \mathcal{G} as a functor from the category of finite dimensional \mathbb{Q} -vector spaces to the category of groups

$$(6.13) \quad \mathcal{G} : \mathbf{Vect}^{\text{fin}}(\mathbb{Q}) \longrightarrow \mathbf{Gr}.$$

By (6.7) the \mathbb{Q} -completion of this group is

$$(6.14) \quad \widehat{\mathcal{G}}_{\mathbb{Q}} = (C \otimes \mathbb{Q}) \ltimes \widehat{\mathcal{M}}_I.$$

Proposition 6.2. *There are isomorphisms*

$$(6.15) \quad H_1(\mathcal{G}) \cong \mathbb{Q} \oplus V, \quad H_n(\mathcal{G}) \cong (\Lambda^n \mathcal{M})_C,$$

$$(6.16) \quad H_1(\widehat{\mathcal{G}}_{\mathbb{Q}}) \cong \mathbb{Q} \oplus V, \quad H_n(\widehat{\mathcal{G}}_{\mathbb{Q}}) \cong (\Lambda^n \widehat{\mathcal{M}})_{C \otimes \mathbb{Q}},$$

where $n \geq 2$.

Proof. It follows from equation (6.9), short exact sequences (6.10), (6.11) and Lemma 6.1. \square

6.5. $H\mathbb{Q}$ -localization of the V -lamplighter group. Since $H_1(\mathcal{G}) = \mathbb{Q} \oplus V$ is finite dimensional, we have $L_{\omega}\mathcal{G} = \widehat{G}_{\mathbb{Q}}$. Hence Proposition 6.2 implies that

$$(6.17) \quad h_{\omega}\mathcal{G} = \text{Coker}((\Lambda^2 \mathcal{M})_C \longrightarrow (\Lambda^2 \widehat{\mathcal{M}}_I)_{C \otimes \mathbb{Q}}).$$

Proposition 6.3. *The functors*

$$(6.18) \quad h_{\omega}\mathcal{G}, H_2(\widehat{\mathcal{G}}_{\mathbb{Q}}) : \mathbf{Vect}^{\text{fin}}(\mathbb{Q}) \longrightarrow \mathbf{Vect}(\mathbb{Q})$$

are homogeneous quadratic functors such that $h_{\omega}\mathcal{G}(V)$ and $H_2(\widehat{\mathcal{G}}(V)_{\mathbb{Q}})$ are uncountable for any $V \neq 0$.

Proof. The functors are homogeneous quadratic because the functor $\Lambda^2(\mathbb{Q}[x] \otimes V)$ is obviously homogeneous quadratic and a quotient of homogeneous quadratic functor is homogeneous quadratic. The fact that the functors applied to $V = \mathbb{Q}$ are uncountable follows from [8, Prop.2.1]. Since \mathbb{Q} is a retract of any non-trivial \mathbb{Q} -vector space, we obtain this result for any $V \neq 0$. \square

Recall that the cyclic group C is generated by an element t . It defines an element $(t, 0) \in \mathcal{G}$. The map $\mathcal{G} \rightarrow H_1(\mathcal{G}) \cong \mathbb{Q} \oplus V$ sends this element to the element $(1, 0)$. Consider the map (5.14) for this element and $\alpha = \omega$

$$(6.19) \quad \bar{\kappa}_{(t,0)}^n : h_{\omega}\mathcal{G} \rightarrow h_{\omega+n}\mathcal{G}.$$

Further we say that a set X is *countable* if $|X| \leq \aleph_0$ (in particular finite sets are countable). If X is not countable, we call it *uncountable*.

Theorem 6.4. *For any $n \geq 0$ and $i \geq 0$ the functors*

$$(6.20) \quad h_{\omega+n}\mathcal{G}, H_i(L_{\omega+n}\mathcal{G}) : \mathbf{Vect}^{\text{fin}}(\mathbb{Q}) \longrightarrow \mathbf{Vect}(\mathbb{Q})$$

are polynomial and the following holds.

- (1) *The map $\bar{\kappa}_{(t,0)}^n : h_{\omega}\mathcal{G} \rightarrow h_{\omega+n}\mathcal{G}$ has a countable kernel for any V . In particular, $h_{\omega+n}\mathcal{G}(V)$ is uncountable for any $V \neq 0$.*
- (2) $\min.\text{deg}(h_{\omega+n}\mathcal{G}) = \min.\text{deg}(H_2(L_{\omega+n}\mathcal{G})) = 2$;

$$(3) \min.\deg(H_3(L_{\omega+n}\mathcal{G})) \geq 3.$$

Proof. For this proof we set $L_{\omega+n} := L_{\omega+n}\mathcal{G}$, $h_{\omega+n} := h_{\omega+n}\mathcal{G}$. and $m = \min.\deg$. The proof is by induction. First we check it for $n = 0$. The fact that the functors are polynomial and the statements (2),(3) follow from the explicit formulas from Proposition 6.2: $H_i(\widehat{\mathcal{G}}_{\mathbb{Q}}) = (\Lambda^i(\mathbb{Q}[[x]] \otimes V))_{C \otimes \mathbb{Q}}$; and the fact that a quotient of a homogeneous functor of degree d is a homogeneous functor of degree d . The statement (1) is obvious.

Now assume that we proved the statement for some n and let's prove it for $n + 1$. The extension $h_{\omega+n} \twoheadrightarrow L_{\omega+n+1} \twoheadrightarrow L_{\omega+n}$ gives a spectral sequence

$$(6.21) \quad H_i(L_{\omega+n}) \otimes \Lambda^j h_{\omega+n} \Rightarrow H_{i+j}(L_{\omega+n+1}).$$

Since polynomial functors are closed under tensor products, subquotients and extensions, the spectral sequence implies that $H_i(L_{\omega+n+1})$ is a polynomial functor. The functor $h_{\omega+n+1}$ is a quotient of $H_2(L_{\omega+n+1})$, so, it is also polynomial.

Prove (3). The third diagonal of the second page of the spectral sequence consists of the four functors

$$(6.22) \quad H_3(L_{\omega+n}), H_2(L_{\omega+n}) \otimes h_{\omega+n}, H_1(L_{\omega+n}) \otimes \Lambda^2 h_{\omega+n}, \Lambda^3 h_{\omega+n}.$$

Note that for any two functors Φ, Ψ we have

$$(6.23) \quad m(\Phi \otimes \Psi) \geq m(\Phi) + m(\Psi), \quad m(\Phi \circ \Psi) \geq m(\Phi) \cdot m(\Psi).$$

Using these formulas and induction hypothesis, it easy to see that minimal degrees of these four functors are at least 3. It follows that $m(H_3(L_{\omega+n+1})) \geq 3$.

Prove (2). Proposition 5.1 implies that there is an exact sequence

$$(6.24) \quad H_3(L_{\omega+n}) \xrightarrow{\delta} (\mathbb{Q} \oplus V) \otimes h_{\omega+n} \xrightarrow{\kappa} H_2(L_{\omega+n+1}) \rightarrow H_2(L_{\omega+n}).$$

First note that $m((\mathbb{Q} \oplus V) \otimes h_{\omega+n}) = m(h_{\omega+n}) = 2$ and $m(H_3(L_{\omega+n})) \geq 3$. It follows that $m(\text{Im}(\kappa)) = 2$. By the induction hypothesis we have $m(H_2(L_{\omega+n})) = 2$. Therefore $m(H_2(L_{\omega+n+1})) = 2$.

Prove (1). Since h_{ω} is homogeneous of degree 2 and $m(H_3(L_{\omega+n})) \geq 3$ we obtain that the image of the composition map

$$(6.25) \quad h_{\omega} \xrightarrow{\bar{\kappa}_{(t,0)}^n} h_{\omega+n} \xrightarrow{t \otimes \text{id}} (\mathbb{Q} \oplus V) \otimes h_{\omega+n}$$

does not intersect the image of δ . It follows that the composition of this map with κ

$$(6.26) \quad \kappa(t \otimes \text{id})\bar{\kappa}_{(t,0)}^n : h_{\omega} \rightarrow H_2(L_{\omega+n+1})$$

has the same kernel as $\bar{\kappa}_{(t,0)}^n$, which is countable. The map $\bar{\kappa}_{(t,0)}^{n+1} : h_{\omega} \rightarrow h_{\omega+n+1}$ is the composition of the map (6.26) and the map $H_2(L_{\omega+n+1}) \twoheadrightarrow h_{\omega+n+1}$. Both these maps have countable kernels, hence, $\bar{\kappa}_{(t,0)}^{n+1} : h_{\omega} \rightarrow h_{\omega+n+1}$ has also a countable kernel. \square

Corollary 6.5. *For any finite dimensional non-trivial \mathbb{Q} -vector space V we have*

$$(6.27) \quad H\mathbb{Q}\text{-length}(\mathcal{G}) \geq \omega + \omega.$$

Proof. It follows from the fact that $h_{\omega+n}\mathcal{G}$ is non-trivial for any n and any $V \neq 0$, and (5.7). \square

7. *HR*-localization of the free group

Let $F = F(a, b)$ be the free group of rank 2, $C = \langle t \rangle$ be the cyclic group and G be the \mathbb{Q} -lamplighter group $G = \mathcal{G}(\mathbb{Q}) = C \rtimes \mathbb{Q}[C]$. Then $\widehat{G}_{\mathbb{Q}} = (C \otimes \mathbb{Q}) \rtimes \mathbb{Q}[[x]]$. Consider the homomorphism

$$(7.1) \quad \varphi : F \longrightarrow G, \quad a \mapsto (t, 0), b \mapsto (1, 1).$$

The following lemma was proved in [8] (Section 5 “Proof of Theorems 1 and 2”, (5-1)).

Lemma 7.1 ([8, (5-1)]). *The image of the map*

$$(7.2) \quad H_2(\widehat{F}_{\mathbb{Z}}, \mathbb{Z}) \longrightarrow H_2(\widehat{G}_{\mathbb{Q}}, \mathbb{Q})$$

is uncountable.

Corollary 7.2. *The image of the map*

$$(7.3) \quad h_{\omega}^{\mathbb{Z}}F \longrightarrow h_{\omega}^{\mathbb{Q}}G$$

is uncountable.

Proposition 7.3. *For any natural $n \geq 0$ the image of the map*

$$(7.4) \quad h_{\omega}^{\mathbb{Z}}F \xrightarrow{\widehat{\kappa}_{n,a}^{\mathbb{Z}}} h_{\omega+n}^{\mathbb{Z}}F \longrightarrow h_{\omega+n}^{\mathbb{Q}}G$$

is uncountable.

Proof. By (5.15) the diagram

$$(7.5) \quad \begin{array}{ccc} h_{\omega}^{\mathbb{Z}}F & \xrightarrow{\widehat{\kappa}_{n,a}^{\mathbb{Z}}} & h_{\omega+n}^{\mathbb{Z}}F \\ \downarrow & & \downarrow \\ h_{\omega}^{\mathbb{Q}}G & \xrightarrow{\widehat{\kappa}_{n,(t,0)}^{\mathbb{Q}}} & h_{\omega+n}^{\mathbb{Q}}G \end{array}$$

is commutative. Corollary 7.2 implies that the image of the left-hand vertical map is uncountable. Theorem 6.4 implies that the kernel of the bottom arrow is countable. Hence the image of the composition map is uncountable. \square

Theorem 7.4. *For a subring $R \subseteq \mathbb{Q}$ and a free group F of rank at least 2 we have*

$$(7.6) \quad HR\text{-length}(F) \geq \omega + \omega.$$

Proof. Since a free group of rank 2 is a retract of any free group of rank ≥ 2 , we can assume that F is of rank 2. By Proposition 7.3 the image of the map $h_{\omega+n}^{\mathbb{Z}}F \rightarrow h_{\omega+n}^{\mathbb{Q}}G$ is uncountable. Since it factors through $h_{\omega+n}^R F$, the group $h_{\omega+n}^R F$ is also uncountable. Using that $HR\text{-length}(F) = \sup\{\alpha \mid h_{\alpha}^R F \neq 0\}$, we obtain the assertion. \square

References

- [1] Aldridge Knight Bousfield. *Homological Localization Towers for Groups and Pi-Modules*. Vol. 186. American Mathematical Soc., 1977.
- [2] Aldridge Knight Bousfield. “The localization of spaces with respect to homology”. In: *Topology* 14.2 (1975), pp. 133–150.

- [3] Aurélien Djament, Antoine Touzé, and Christine Vespa. “Décompositions à la Steinberg sur une catégorie additive”. In: *arXiv preprint arXiv:1904.09190* (2019).
- [4] Yuriy A Drozd. “On cubic functors”. In: *Communications in Algebra* 31.3 (2003), pp. 1147–1173.
- [5] Beno Eckmann, Peter Hilton, and Urs Stammbach. “On the homology theory of central group extensions II. The exact sequence in the general case”. In: *Commentarii Mathematici Helvetici* 47.1 (1972), pp. 171–178.
- [6] Samuel Eilenberg and Saunders MacLane. “On the groups $H(\Pi, n)$, II: methods of computation”. In: *Annals of Mathematics* (1954), pp. 49–139.
- [7] Emmanuel D Farjoun and Yoav Segev. “Relative Schur multipliers and universal extensions of group homomorphisms”. In: *Manifolds and K-Theory* 682 (2017), p. 65.
- [8] Sergei Ivanov and Roman Mikhailov. “A finite \mathbb{Q} -bad space”. In: *Geometry & Topology* 23.3 (2019), pp. 1237–1249.
- [9] Sergei O Ivanov and Roman Mikhailov. “On a problem of Bousfield for metabelian groups”. In: *Advances in Mathematics* 290 (2016), pp. 552–589.
- [10] Sergei O Ivanov and Roman Mikhailov. “On discrete homology of a free pro- p -group”. In: *Compositio Mathematica* 154.10 (2018), pp. 2195–2204.
- [11] Sergei O Ivanov and Roman Mikhailov. “On lengths of $\mathbb{H}\mathbb{Z}$ -localization towers”. In: *Israel Journal of Mathematics* 226.2 (2018), pp. 635–683.

LABORATORY OF MODERN ALGEBRA AND APPLICATIONS, ST. PETERSBURG STATE UNIVERSITY,
14TH LINE, 29B, SAINT PETERSBURG, 199178 RUSSIA
Email address: ivanov.s.o.1986@gmail.com

LABORATORY OF MODERN ALGEBRA AND APPLICATIONS, ST. PETERSBURG STATE UNIVERSITY,
14TH LINE, 29B, SAINT PETERSBURG, 199178 RUSSIA, AND ST. PETERSBURG DEPARTMENT OF
STEKLOV MATHEMATICAL INSTITUTE
Email address: rmikhailov@mail.ru