

## AN IDENTITY FOR $(-q^5; q^{10})_\infty$

SUMIT KUMAR JHA

ABSTRACT. We prove that

$$\prod_{n=0}^{\infty} (1 + q^{10n+5}) = \frac{\sum_{n=-\infty}^{\infty} q^{n^2} \sum_{n=-\infty}^{\infty} (-1)^n (-q)^{n(3n-1)/2}}{4 \sum_{n \geq 0} (-q)^{\frac{n(n+1)}{2}} \sin \left\{ \frac{(2n+1)3\pi}{10} \right\} \sum_{n \geq 0} (-q)^{\frac{n(n+1)}{2}} \sin \left\{ \frac{(2n+1)\pi}{10} \right\}} \quad |q| < 1$$

using identities due to Ramanujan.

### 1. NOTATIONS AND DEFINITIONS

We assume that  $|q| < 1$  hereon. The functions  $\varphi(q)$ ,  $\psi(q)$ ,  $f(-q)$ , and  $\chi(q)$  are defined as

$$\begin{aligned} \varphi(q) &:= \sum_{n=-\infty}^{\infty} q^{n^2}, \\ \psi(q) &:= \sum_{n=0}^{\infty} q^{n(n+1)/2}, \\ f(-q) &:= \sum_{n=-\infty}^{\infty} (-1)^n q^{n(3n-1)/2} = \prod_{n \geq 1} (1 - q^n), \\ \chi(q) &:= \prod_{n \geq 0} (1 + q^{2n+1}). \end{aligned}$$

The four *Jacobi theta functions* [1, p. 464] are defined by:

$$\begin{aligned} \theta_1(z, q) &= 2 \sum_{n=0}^{\infty} (-1)^n q^{(n+\frac{1}{2})^2} \sin((2n+1)z), \\ \theta_2(z, q) &= 2 \sum_{n=0}^{\infty} q^{(n+\frac{1}{2})^2} \cos((2n+1)z), \\ \theta_3(z, q) &= 1 + 2 \sum_{n=1}^{\infty} q^{n^2} \cos(2nz), \\ \theta_4(z, q) &= 1 + 2 \sum_{n=1}^{\infty} (-1)^n q^{n^2} \cos(2nz). \end{aligned}$$

We wish to prove that

$$(1) \quad \chi(q^5) = \frac{\varphi(q) f(q)}{4 \sum_{n \geq 0} (-q)^{\frac{n(n+1)}{2}} \sin \left\{ \frac{(2n+1)3\pi}{10} \right\} \sum_{n \geq 0} (-q)^{\frac{n(n+1)}{2}} \sin \left\{ \frac{(2n+1)\pi}{10} \right\}}.$$

### 2. PROOF OF (1)

*Proof.* We begin with an identity of Ramanujan's in his lost notebook [3, p. 27, Theorem 1.6.1 (ii)]

$$\varphi^2(q) - 5 \varphi^2(q^5) = -4 f^2(-q^2) \frac{\chi(q^5)}{\chi(q)}.$$

---

2010 *Mathematics Subject Classification.* 33D99.

*Key words and phrases.* Theta functions, Ramanujan's theta function, Jacobi's theta function.

The factorizations of the above are given by [3, p. 29, Entry 1.7.2 (i)]

$$(2) \quad \varphi(q) + \sqrt{5} \varphi(q^5) = \frac{(1 + \sqrt{5}) f(-q^2)}{\prod_{n \text{ odd}} (1 + \alpha q^n + q^{2n}) \prod_{n \text{ even}} (1 - \beta q^n + q^{2n})},$$

and [3, p. 30, Entry 1.7.2 (ii)]

$$(3) \quad \varphi(q) - \sqrt{5} \varphi(q^5) = \frac{(1 - \sqrt{5}) f(-q^2)}{\prod_{n \text{ odd}} (1 + \beta q^n + q^{2n}) \prod_{n \text{ even}} (1 - \alpha q^n + q^{2n})},$$

where

$$\alpha = \frac{1 - \sqrt{5}}{2}, \quad \beta = \frac{1 + \sqrt{5}}{2}.$$

Multiplying (2) and (3) and using the following Fourier expansions [1, p. 469] for  $\theta_1(z, q)$  and  $\theta_3(z, q)$ :

$$\theta_1(z, q) = 2q^{1/4} \sin z \prod_{n=1}^{\infty} (1 - q^{2n}) (1 - 2q^{2n} \cos(2z) + q^{4n}),$$

$$\theta_3(z, q) = \prod_{n=1}^{\infty} (1 - q^{2n}) (1 + 2q^{2n-1} \cos(2z) + q^{4n-2})$$

we have

$$\frac{\chi(q^5)}{\chi(q)} = \frac{q^{1/2} f^4(-q^2)}{\theta_1(\pi/10, q) \theta_3(\pi/10, q) \theta_1(3\pi/10, q) \theta_3(3\pi/10, q)}.$$

Using the identity [4]

$$\frac{\theta_1(z, q^2) \theta_3(z, q^2)}{\theta_1(z, iq)} = i^{-1/4} \sqrt{\frac{\theta_2(0, q^2) \theta_4(0, q^2)}{2}}$$

we now have

$$\begin{aligned} \frac{\chi(q^5)}{\chi(q)} &= \frac{2 q^{1/2} i^{1/2} f^4(-q^2)}{\theta_1(\pi/10, i\sqrt{q}) \theta_1(3\pi/10, i\sqrt{q}) \theta_2(0, q) \theta_4(0, q)} \\ &= \frac{q^{1/4} i^{1/2} f^4(-q^2)}{\psi(q^2) \varphi(-q) \theta_1(\pi/10, i\sqrt{q}) \theta_1(3\pi/10, i\sqrt{q})} \\ &= \frac{q^{1/4} i^{1/2} f(-q^2) \varphi^2(-q^2)}{\varphi(-q) \theta_1(\pi/10, i\sqrt{q}) \theta_1(3\pi/10, i\sqrt{q})} \\ &= \frac{q^{1/4} i^{1/2} f(-q^2) \varphi(q)}{\theta_1(\pi/10, i\sqrt{q}) \theta_1(3\pi/10, i\sqrt{q})} \\ &= \frac{\sum_{n=-\infty}^{\infty} q^{n^2} \sum_{m=-\infty}^{\infty} (-1)^n q^{n(3n-1)}}{4 \sum_{n \geq 0} (-q)^{\frac{n(n+1)}{2}} \sin \left\{ \frac{(2n+1)3\pi}{10} \right\} \sum_{n \geq 0} (-q)^{\frac{n(n+1)}{2}} \sin \left\{ \frac{(2n+1)\pi}{10} \right\}}, \end{aligned}$$

where we have used [2, p. 15, (1.3.34)]

$$f^3(-q^2) = \psi(q^2) \varphi^2(-q^2)$$

and [2, p. 15, (1.3.32)]

$$\varphi^2(-q^2) = \varphi(q) \varphi(-q).$$

Now we finally use [3, p.15, (1.3.31)]

$$f(q) = \chi(q) f(-q^2)$$

to conclude our main assertion.  $\square$

## REFERENCES

- [1] E. T. Whittaker and G. N. Watson, *A course of modern analysis*, Cambridge University Press, 1996.
- [2] B. C. Berndt, *Number theory in the spirit of Ramanujan*, American Mathematical Society, 2012.
- [3] G. E. Andrews and B. C. Berndt, *Ramanujan's lost notebook: Part 1*, Springer, 2012.
- [4] P. L. Walker, Reduction formulae for products of theta functions, *Journal of Research of the National Institute of Standards and Technology* **117** (2012), 297–303. Available at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4553875/>.

G-2, C-185, RAMPRASTHA, GHAZIABAD, U.P.-201011, INDIA  
Email address: sumitkumarjha.iiit@gmail.com