

Some Generalized Multi-sum Chu-Vandermonde Identities

M.J. Kronenburg

Abstract

Some generalized multi-sum Chu-Vandermonde identities are presented and proved, generalizing some known multi-sum Chu-Vandermonde identities from literature and adding some quadratic and cubic examples of these identities. Some other closely related identities are provided. The identities are proved with the integral representation method using complex residues, a method also used in some earlier papers.

Keywords: binomial coefficient, combinatorial identities.

MSC 2010: 05A10, 05A19

1 Generalized Multi-sum Chu-Vandermonde Identities

Let $\{a_i\}$ and $\{c_i\}$ be two sets each of m nonnegative integers, and let $\{x_i\}$ and $\{y_i\}$ be two sets each of m complex numbers, and let \bar{w} be the complex conjugate of w . Let the following definitions be given.

$$A_{p,q} = \sum_{i=1}^m x_i^p a_i^q \quad (1.1)$$

$$A_p = A_{p,1} = \sum_{i=1}^m x_i^p a_i \quad (1.2)$$

$$A_{p,q}^* = \sum_{i=1}^m x_i^p y_i^q a_i \quad (1.3)$$

$$A_p^* = A_{0,p}^* = \sum_{i=1}^m y_i^p a_i \quad (1.4)$$

$$A_{\text{abs}} = \sum_{i=1}^m |x_i|^2 a_i \quad (1.5)$$

$$C_{p,q} = \sum_{i=1}^m x_i^p c_i^q \quad (1.6)$$

$$C_p = C_{p,1} = \sum_{i=1}^m x_i^p c_i \quad (1.7)$$

$$C_{p,q}^* = \sum_{i=1}^m x_i^p y_i^q c_i \quad (1.8)$$

$$C_p^* = C_{0,p}^* = \sum_{i=1}^m y_i^p c_i \quad (1.9)$$

$$C_{\text{abs}} = \sum_{i=1}^m |x_i|^2 c_i \quad (1.10)$$

$$S_{p,q} = \sum_{i=1}^m x_i a_i^p c_i^q \quad (1.11)$$

The following generalized multi-sum Chu-Vandermonde identities are proved.
For integer $m \geq 1$ and $n \geq 0$:

$$\sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}} \prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} = \binom{A_0 - C_0}{n - C_0} \prod_{i=1}^m \binom{a_i}{c_i} \quad (1.12)$$

$$\begin{aligned} & \sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i=1}^m x_i k_i \\ &= \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{(n - C_0)A_1 - (n - A_0)C_1}{A_0 - C_0} \end{aligned} \quad (1.13)$$

$$\begin{aligned} & \sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left(\sum_{i=1}^m x_i k_i \right)^2 = \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \\ & \cdot \frac{(n - C_0)[(n - C_0 - 1)A_1^2 + (n - A_0)(C_2 - A_2 - 2A_1C_1)] + (n - A_0)(n - A_0 + 1)C_1^2}{(A_0 - C_0)(A_0 - C_0 - 1)} \end{aligned} \quad (1.14)$$

$$\begin{aligned} & \sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left| \sum_{i=1}^m x_i k_i \right|^2 \\ &= \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)} \\ & \cdot \{ (n - C_0)[(n - C_0 - 1)|A_1|^2 + (n - A_0)(C_{\text{abs}} - A_{\text{abs}} - A_1\overline{C_1} - \overline{A_1}C_1)] \\ & \quad + (n - A_0)(n - A_0 + 1)|C_1|^2 \} \end{aligned} \quad (1.15)$$

$$\begin{aligned}
& \sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}}^{a_1 \cdots a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left(\sum_{i=1}^m x_i k_i \right) \left(\sum_{i=1}^m y_i k_i \right) \\
&= \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)} \\
&\quad \cdot \{ (n - C_0)[(n - C_0 - 1)A_1 A_1^* + (n - A_0)(C_{1,1}^* - A_{1,1}^* - A_1 C_1^* - A_1^* C_1)] \\
&\quad + (n - A_0)(n - A_0 + 1)C_1 C_1^* \}
\end{aligned} \tag{1.16}$$

$$\begin{aligned}
& \sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}}^{a_1 \cdots a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i=1}^m x_i k_i^2 \\
&= \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)} \\
&\quad \cdot \{ (n - C_0)[(n - C_0 - 1)A_{1,2} + (n - A_0)(C_1 - A_1 - 2S_{1,1})] \\
&\quad + (n - A_0)(n - A_0 + 1)C_{1,2} \}
\end{aligned} \tag{1.17}$$

$$\begin{aligned}
& \sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}}^{a_1 \cdots a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left(\sum_{i=1}^m x_i k_i \right)^3 \\
&= \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)(A_0 - C_0 - 2)} \\
&\quad \cdot \{ (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)A_1^3 \\
&\quad + (n - A_0)(A_3 - C_3 + 3A_1(C_2 - A_2 - A_1 C_1))] \\
&\quad - (n - A_0)(n - A_0 + 1)[(n - A_0 + 2)C_1^3 \\
&\quad + (n - C_0)(C_3 - A_3 + 3C_1(C_2 - A_2 - A_1 C_1))] \}
\end{aligned} \tag{1.18}$$

$$\begin{aligned}
& \sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}}^{a_1 \cdots a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i=1}^m x_i k_i^3 \\
&= \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)(A_0 - C_0 - 2)} \\
&\quad \cdot \{ (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)A_{1,3} \\
&\quad + (n - A_0)(A_1 - C_1 - 3(A_{1,2} + S_{2,1} - S_{1,1}))] \\
&\quad - (n - A_0)(n - A_0 + 1)[(n - A_0 + 2)C_{1,3} \\
&\quad + (n - C_0)(C_1 - A_1 + 3(C_{1,2} - S_{1,2} - S_{1,1}))] \}
\end{aligned} \tag{1.19}$$

Taking all $c_i = 0$ and therefore all $C_{p,q} = C_p = C_{p,q}^* = C_p^* = C_{\text{abs}} = 0$, identities (1.12), (1.13) and (1.15) can be found in [9], and taking $m = 2$, $x_1 = 1$ and $x_2 = 0$, and therefore $A_0 = a + b$, $C_0 = c + d$, $A_1 = A_2 = A_3 = a$ and $C_1 = C_2 = C_3 = c$, identities (1.12),

(1.13), (1.14) and (1.18) can be found in [6].

2 Unrestricted Multi-sum Identities

When the multiple sum is without the restriction on the k_i -indices, other closely related identities result which are given below.

For integer $m \geq 1$:

$$\sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} = 2^{A_0-C_0} \prod_{i=1}^m \binom{a_i}{c_i} \quad (2.1)$$

$$\sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i=1}^m x_i k_i = 2^{A_0-C_0-1} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (A_1 + C_1) \quad (2.2)$$

$$\begin{aligned} & \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left(\sum_{i=1}^m x_i k_i \right)^2 \\ &= 2^{A_0-C_0-2} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] [A_2 - C_2 + (A_1 + C_1)^2] \end{aligned} \quad (2.3)$$

$$\begin{aligned} & \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left| \sum_{i=1}^m x_i k_i \right|^2 \\ &= 2^{A_0-C_0-2} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (A_{\text{abs}} - C_{\text{abs}} + |A_1 + C_1|^2) \end{aligned} \quad (2.4)$$

$$\begin{aligned} & \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left(\sum_{i=1}^m x_i k_i \right) \left(\sum_{i=1}^m y_i k_i \right) \\ &= 2^{A_0-C_0-2} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] [A_{1,1}^* - C_{1,1}^* + (A_1 + C_1)(A_1^* + C_1^*)] \end{aligned} \quad (2.5)$$

$$\begin{aligned} & \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i=1}^m x_i k_i^2 \\ &= 2^{A_0-C_0-2} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (A_1 - C_1 + A_{1,2} + C_{1,2} + 2S_{1,1}) \end{aligned} \quad (2.6)$$

$$\begin{aligned} & \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left(\sum_{i=1}^m x_i k_i \right)^3 \\ &= 2^{A_0-C_0-3} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (A_1 + C_1)[(A_1 + C_1)^2 + 3(A_2 - C_2)] \end{aligned} \quad (2.7)$$

$$\begin{aligned}
& \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i=1}^m x_i k_i^3 \\
& = 2^{A_0 - C_0 - 3} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] [A_{1,3} + C_{1,3} + 3(A_{1,2} - C_{1,2} + S_{1,2} + S_{2,1})]
\end{aligned} \tag{2.8}$$

3 Proof of the Multi-sum Chu-Vandermonde Identities

For the proof of the multi-sum identities, the order of summation is changed:

$$\begin{aligned}
& \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i_1=0}^m \cdots \sum_{i_s=0}^m \prod_{j=1}^s w_{i_j} k_{i_j}^{p_j} \\
& \quad \sum_{i=0}^m k_i = n \\
& = \sum_{i_1=0}^m \cdots \sum_{i_s=0}^m \left[\prod_{j=1}^s w_{i_j} \right] \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \prod_{j=1}^s k_{i_j}^{p_j} \\
& \quad \sum_{i=0}^m k_i = n
\end{aligned} \tag{3.1}$$

The powers p_j in the right side must be constant in the second multiple summation over the k_i , which means that the indices i_j must be mutually unequal. A product of sums in the summand is therefore splitted into multiple sums over mutually unequal indices:

$$\left(\sum_{i=1}^m x_i k_i \right) \left(\sum_{i=1}^m y_i k_i \right) = \sum_{p=1}^m \sum_{\substack{q=1 \\ p \neq q}}^m x_p y_q k_p k_q + \sum_{p=1}^m x_p y_p k_p^2 \tag{3.2}$$

$$\begin{aligned}
& \left(\sum_{i=1}^m x_i k_i \right) \left(\sum_{i=1}^m y_i k_i \right) \left(\sum_{i=1}^m z_i k_i \right) = \sum_{p=1}^m \sum_{\substack{q=1 \\ p \neq q}}^m \sum_{\substack{r=1 \\ p \neq q \neq r, \\ p \neq r}}^m x_p y_q z_r k_p k_q k_r \\
& + \sum_{p=1}^m \sum_{\substack{q=1 \\ p \neq q}}^m (x_p y_q z_q + y_p x_q z_q + z_p x_q y_q) k_p k_q^2 + \sum_{p=1}^m x_p y_p z_p k_p^3
\end{aligned} \tag{3.3}$$

For the multi-sum identities up to cubic order, in the right side of (3.1) the power combinations 1, k_p , $k_p k_q$, k_p^2 , $k_p k_q k_r$, $k_p k_q^2$ and k_p^3 are needed, where as mentioned $p \neq q \neq r$ and $p \neq r$, and the second multiple summation over the k_i is evaluated below with the integral representation method using complex residues as in earlier papers [6, 7]:

$$\binom{n}{k} = \text{Res}_w \frac{(1+w)^n}{w^{k+1}} \tag{3.4}$$

$$\text{Res}_{w=w_p} \frac{f(w)}{(w-w_p)^k} = \frac{1}{(k-1)!} D_w^{k-1} f(w) |_{w=w_p} \tag{3.5}$$

where $D_w^n f(w) |_{w=w_p}$ is the n -th derivative of $f(w)$ at $w = w_p$.

Theorem 3.1. For integer $m \geq 1$:

$$\sum_{\substack{a_1 \cdots a_m \\ k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}} \prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} = \binom{A_0 - C_0}{n - C_0} \prod_{i=1}^m \binom{a_i}{c_i} \quad (3.6)$$

Proof. The trinomial revision identity [5, 8] is applied:

$$\binom{a_i}{k_i} \binom{k_i}{c_i} = \binom{a_i - c_i}{k_i - c_i} \binom{a_i}{c_i} \quad (3.7)$$

Because this expression is zero when $c_i > a_i$, in the multiple summation it can be assumed that $0 \leq c_i \leq a_i$ and that $c_i \leq k_i \leq a_i$. The restriction on the k_i -indices in the multiple sum can therefore be realized by setting k_m equal to:

$$k_m = n - \sum_{i=1}^{m-1} k_i \quad (3.8)$$

reducing an m -fold sum to an $(m-1)$ -fold sum. The infinite geometric series [7] is used:

$$\sum_{k=0}^{\infty} w^k = \frac{1}{1-w} \quad (3.9)$$

The upper bounds a_i can be replaced by ∞ because the summand is zero when any $k_i > a_i$.

$$\begin{aligned} & \sum_{\substack{a_1 \cdots a_m \\ k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}} \prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \\ &= \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \sum_{k_1=0 \cdots k_{m-1}=0} \binom{a_m - c_m}{n - c_m - \sum_{i=1}^{m-1} k_i} \prod_{i=1}^{m-1} \binom{a_i - c_i}{k_i - c_i} \\ &= \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \sum_{k_1=0 \cdots k_{m-1}=0} \left[\prod_{i=1}^m \text{Res}_{w_i} \right] \frac{(1+w_m)^{a_m - c_m}}{w_m^{n - c_m - \sum_{i=1}^{m-1} k_i + 1}} \prod_{i=1}^{m-1} \frac{(1+w_i)^{a_i - c_i}}{w_i^{k_i - c_i + 1}} \\ &= \left[\prod_{i=1}^m \binom{a_i}{c_i} \text{Res}_{w_i} \right] \frac{1}{w_m^n} \left[\prod_{i=1}^m (1+w_i)^{a_i - c_i} w_i^{c_i - 1} \right] \sum_{k_1=0 \cdots k_{m-1}=0} \prod_{i=1}^{m-1} \left(\frac{w_m}{w_i} \right)^{k_i} \quad (3.10) \\ &= \left[\prod_{i=1}^m \binom{a_i}{c_i} \text{Res}_{w_i} \right] \frac{1}{w_m^n} \left[\prod_{i=1}^m (1+w_i)^{a_i - c_i} w_i^{c_i - 1} \right] \prod_{i=1}^{m-1} \sum_{k_i=0}^{\infty} \left(\frac{w_m}{w_i} \right)^{k_i} \\ &= \left[\prod_{i=1}^m \binom{a_i}{c_i} \text{Res}_{w_i} \right] \frac{1}{w_m^n} \left[\prod_{i=1}^m (1+w_i)^{a_i - c_i} w_i^{c_i - 1} \right] \prod_{i=1}^{m-1} \frac{1}{1 - \frac{w_m}{w_i}} \\ &= \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \text{Res}_{w_m} \frac{(1+w_m)^{a_m - c_m}}{w_m^{n - c_m + 1}} \prod_{i=1}^{m-1} \text{Res}_{w_i} \frac{(1+w_i)^{a_i - c_i} w_i^{c_i}}{w_i - w_m} \end{aligned}$$

The residue of w_i is evaluated with (3.5), using that $c_i \geq 0$ and $a_i - c_i \geq 0$:

$$\begin{aligned} &= \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \operatorname{Res}_{w_m} \frac{(1+w_m)^{a_m-c_m}}{w_m^{n-c_m+1}} \prod_{i=1}^{m-1} (1+w_m)^{a_i-c_i} w_m^{c_i} \\ &= \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \operatorname{Res}_{w_m} \frac{(1+w_m)^{\sum_{i=1}^m (a_i-c_i)}}{w_i^{n-\sum_{i=1}^m c_i+1}} = \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \binom{\sum_{i=1}^m (a_i-c_i)}{n-\sum_{i=1}^m c_i} \end{aligned} \quad (3.11)$$

□

Theorem 3.2. For integer $m \geq 1$ and $1 \leq p \leq m$:

$$\sum_{\substack{k_1=0 \dots k_m=0 \\ \sum_{i=0}^m k_i=n}} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] k_p = \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{(n - C_0)a_p - (n - A_0)c_p}{A_0 - C_0} \quad (3.12)$$

Proof. The following modified infinite geometric series [7] is used:

$$\sum_{k=0}^{\infty} k w^k = \frac{w}{(1-w)^2} \quad (3.13)$$

The theorem is first proved for $1 \leq p < m$:

$$\begin{aligned} &\left[\prod_{i=1}^m \operatorname{Res}_{w_i} \right] \frac{1}{w_m^n} \left[\prod_{i=1}^m (1+w_i)^{a_i-c_i} w_i^{c_i-1} \right] \sum_{k_1=0 \dots k_{m-1}=0}^{\infty \dots \infty} k_p \prod_{i=1}^{m-1} \left(\frac{w_m}{w_i} \right)^{k_i} \\ &= \left[\prod_{i=1}^m \operatorname{Res}_{w_i} \right] \frac{1}{w_m^n} \left[\prod_{i=1}^m (1+w_i)^{a_i-c_i} w_i^{c_i-1} \right] \frac{\frac{w_m}{w_p}}{\left(1 - \frac{w_m}{w_p}\right)^2} \prod_{\substack{i=1 \\ i \neq p}}^{m-1} \frac{1}{1 - \frac{w_m}{w_i}} \\ &= \operatorname{Res}_{w_m} \frac{(1+w_m)^{a_m-c_m}}{w_m^{n-c_m}} \operatorname{Res}_{w_p} \frac{(1+w_p)^{a_p-c_p} w_p^{c_p}}{(w_p-w_m)^2} \prod_{\substack{i=1 \\ i \neq p}}^{m-1} \operatorname{Res}_{w_i} \frac{(1+w_i)^{a_i-c_i} w_i^{c_i}}{w_i-w_m} \\ &= \operatorname{Res}_{w_m} \frac{(1+w_m)^{a_m-c_m}}{w_m^{n-c_m}} \operatorname{Res}_{w_p} \frac{(1+w_p)^{a_p-c_p} w_p^{c_p}}{(w_p-w_m)^2} \prod_{\substack{i=1 \\ i \neq p}}^{m-1} (1+w_m)^{a_i-c_i} w_m^{c_i} \end{aligned} \quad (3.14)$$

The residue of w_p is evaluated with (3.5) using the product rule for the derivative:

$$\operatorname{Res}_{w_p} \frac{(1+w_p)^{a_p-c_p} w_p^{c_p}}{(w_p-w_m)^2} = (a_p-c_p)(1+w_m)^{a_p-c_p-1} w_m^{c_p} + c_p(1+w_m)^{a_p-c_p} w_m^{c_p-1} \quad (3.15)$$

and evaluating the residue of w_m and using the absorption identity [5, 8]:

$$\begin{aligned}
& \text{Res}_{w_m} \left[(a_p - c_p) \frac{(1 + w_m)^{\sum_{i=1}^m (a_i - c_i) - 1}}{w_m^{n - \sum_{i=1}^m c_i}} + c_p \frac{(1 + w_m)^{\sum_{i=1}^m (a_i - c_i)}}{w_m^{n - \sum_{i=1}^m c_i + 1}} \right] \\
&= \binom{A_0 - C_0 - 1}{n - C_0 - 1} (a_p - c_p) + \binom{A_0 - C_0}{n - C_0} c_p \\
&= \binom{A_0 - C_0}{n - C_0} \left[\frac{(a_p - c_p)(n - C_0)}{A_0 - C_0} + c_p \right] \\
&= \binom{A_0 - C_0}{n - C_0} \frac{(n - C_0)a_p - (n - A_0)c_p}{A_0 - C_0}
\end{aligned} \tag{3.16}$$

The theorem is now proved for $1 \leq p < m$, but because the indices of the k_i in the left side of the theorem can be interchanged, the theorem is true for $1 \leq p \leq m$. \square

Theorem 3.3. For integer $m \geq 1$:

$$\begin{aligned}
& \sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i=1}^m x_i k_i \\
&= \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{(n - C_0)A_1 - (n - A_0)C_1}{A_0 - C_0}
\end{aligned} \tag{3.17}$$

Proof. Changing the order of summation as in (3.1) and using theorem 3.2:

$$\begin{aligned}
& \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \sum_{i=1}^m x_i \frac{(n - C_0)a_i - (n - A_0)c_i}{A_0 - C_0} \\
&= \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{(n - C_0)A_1 - (n - A_0)C_1}{A_0 - C_0}
\end{aligned} \tag{3.18}$$

\square

Theorem 3.4. For integer $m \geq 1$ and $1 \leq p \leq m$:

$$\begin{aligned}
& \sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] k_p^2 = \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \\
& \cdot \frac{(n - C_0)[(n - C_0 - 1)a_p^2 + (n - A_0)(c_p - a_p - 2a_p c_p)] + (n - A_0)(n - A_0 + 1)c_p^2}{(A_0 - C_0)(A_0 - C_0 - 1)}
\end{aligned} \tag{3.19}$$

Proof. The following modified infinite geometric series [7] is used:

$$\sum_{k=0}^{\infty} k^2 w^k = \frac{w}{(1 - w)^2} + \frac{2w^2}{(1 - w)^3} \tag{3.20}$$

The first term gives a result identical to theorem 3.2, and for the second term:

$$\begin{aligned} & \left[\prod_{i=1}^m \text{Res}_{w_i} \right] \frac{1}{w_m^n} \left[\prod_{i=1}^m (1+w_i)^{a_i-c_i} w_i^{c_i-1} \right] \frac{2\left(\frac{w_m}{w_p}\right)^2}{\left(1-\frac{w_m}{w_p}\right)^3} \prod_{\substack{i=1 \\ i \neq p}}^{m-1} \frac{1}{1-\frac{w_m}{w_i}} \\ &= \text{Res}_{w_m} \frac{(1+w_m)^{a_m-c_m}}{w_m^{n-c_m-1}} \text{Res}_{w_p} \frac{2(1+w_p)^{a_p-c_p} w_p^{c_p}}{(w_p-w_m)^3} \prod_{\substack{i=1 \\ i \neq p}}^{m-1} (1+w_m)^{a_i-c_i} w_m^{c_i} \end{aligned} \quad (3.21)$$

The residue of w_p is evaluated with (3.5) using the product rule for the second derivative:

$$\begin{aligned} \text{Res}_{w_p} \frac{2(1+w_p)^{a_p-c_p} w_p^{c_p}}{(w_p-w_m)^3} &= (a_p-c_p)(a_p-c_p-1)(1+w_m)^{a_p-c_p-2} w_m^{c_p} \\ &+ 2(a_p-c_p)c_p(1+w_m)^{a_p-c_p-1} w_m^{c_p-1} + c_p(c_p-1)(1+w_m)^{a_p-c_p} w_m^{c_p-2} \end{aligned} \quad (3.22)$$

Using the absorption identity [5, 8]:

$$\begin{aligned} & \binom{A_0-C_0-2}{n-C_0-2} (a_p-c_p)(a_p-c_p-1) + 2 \binom{A_0-C_0-1}{n-C_0-1} (a_p-c_p)c_p \\ &+ \binom{A_0-C_0}{n-C_0} c_p(c_p-1) \\ &= \binom{A_0-C_0}{n-C_0} \left[\frac{(n-C_0)(n-C_0-1)}{(A_0-C_0)(A_0-C_0-1)} (a_p-c_p)(a_p-c_p-1) \right. \\ &\quad \left. + 2 \frac{n-C_0}{A_0-C_0} (a_p-c_p)c_p + c_p(c_p-1) \right] \\ &= \binom{A_0-C_0}{n-C_0} \frac{1}{(A_0-C_0)(A_0-C_0-1)} \\ &\quad \cdot \{ (n-C_0)(a_p-c_p)[(n-C_0-1)(a_p-c_p-1) + 2(A_0-C_0-1)c_p] \\ &\quad + (A_0-C_0)(A_0-C_0-1)c_p(c_p-1) \} \end{aligned} \quad (3.23)$$

As mentioned the result of theorem 3.2 must be added, which gives:

$$\begin{aligned} & \binom{A_0-C_0}{n-C_0} \frac{1}{(A_0-C_0)(A_0-C_0-1)} \\ &\quad \cdot \{ (n-C_0)(a_p-c_p)[(n-C_0-1)(a_p-c_p-1) + 2(A_0-C_0-1)c_p] \\ &\quad + (A_0-C_0-1)[(A_0-C_0)c_p(c_p-1) + (n-C_0)a_p - (n-A_0)c_p] \} \end{aligned} \quad (3.24)$$

This result simplifies to the theorem. \square

Theorem 3.5. For integer $m \geq 2$, $1 \leq p \leq m$, $1 \leq q \leq m$ and $p \neq q$:

$$\begin{aligned} & \sum_{\substack{k_1=0 \dots k_m=0 \\ \sum_{i=0}^n k_i=n}} \dots \sum_{a_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] k_p k_q = \binom{A_0-C_0}{n-C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \\ & \cdot \frac{(n-C_0)[(n-C_0-1)a_p a_q - (n-A_0)(a_p c_q + c_p a_q)] + (n-A_0)(n-A_0+1)c_p c_q}{(A_0-C_0)(A_0-C_0-1)} \end{aligned} \quad (3.25)$$

Proof. The proof is similar to theorem 3.2 but with two quadratic residues instead of one.

$$\begin{aligned} & \text{Res}_{w_m} \frac{(1+w_m)^{a_m-c_m}}{w_m^{n-c_m-1}} \text{Res}_{w_p} \frac{(1+w_p)^{a_p-c_p} w_p^{c_p}}{(w_p-w_m)^2} \text{Res}_{w_q} \frac{(1+w_q)^{a_p-c_p} w_q^{c_p}}{(w_q-w_m)^2} \\ & \cdot \prod_{\substack{i=1 \\ i \neq p, i \neq q}}^{m-1} (1+w_m)^{a_i-c_i} w_m^{c_i} \end{aligned} \quad (3.26)$$

The two last residues are given by (3.15), and multiplying these out gives:

$$\begin{aligned} & [(a_p-c_p)(1+w_m)^{a_p-c_p-1} w_m^{c_p} + c_p(1+w_m)^{a_p-c_p} w_m^{c_p-1}] \\ & \cdot [(a_q-c_q)(1+w_m)^{a_q-c_q-1} w_m^{c_q} + c_q(1+w_m)^{a_q-c_q} w_m^{c_q-1}] \\ & = (a_p-c_p)(a_q-c_q)(1+w_m)^{a_p-c_p+a_q-c_q-2} w_m^{c_p+c_q} \\ & \quad + [(a_p-c_p)c_q + (a_q-c_q)c_p](1+w_m)^{a_p-c_p+a_q-c_q-1} w_m^{c_p+c_q-1} \\ & \quad + c_p c_q (1+w_m)^{a_p-c_p+a_q-c_q} w_m^{c_p+c_q-2} \end{aligned} \quad (3.27)$$

This yields the following:

$$\begin{aligned} & \binom{A_0-C_0-2}{n-C_0-2} (a_p-c_p)(a_q-c_q) + \binom{A_0-C_0-1}{n-C_0-1} [(a_p-c_p)c_q + (a_q-c_q)c_p] \\ & + \binom{A_0-C_0}{n-C_0} c_p c_q \end{aligned} \quad (3.28)$$

and using again the absorption identity:

$$\begin{aligned} & \binom{A_0-C_0}{n-C_0} \frac{1}{(A_0-C_0)(A_0-C_0-1)} \\ & \cdot \{(n-C_0)[(n-C_0-1)(a_p-c_p)(a_q-c_q) + (A_0-C_0-1)((a_p-c_p)c_q + (a_q-c_q)c_p)] \\ & \quad + (A_0-C_0)(A_0-C_0-1)c_p c_q\} \end{aligned} \quad (3.29)$$

This result simplifies to the theorem. \square

Theorem 3.6. For integer $m \geq 1$:

$$\begin{aligned} & \sum_{\substack{k_1=0 \dots k_m=0 \\ \sum_{i=0}^m k_i=n}}^{a_1 \dots a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left(\sum_{i=1}^m x_i k_i \right) \left(\sum_{i=1}^m y_i k_i \right) \\ & = \binom{A_0-C_0}{n-C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0-C_0)(A_0-C_0-1)} \\ & \quad \cdot \{(n-C_0)[(n-C_0-1)A_1 A_1^* + (n-A_0)(C_{1,1}^* - A_{1,1}^* - A_1 C_1^* - A_1^* C_1)] \\ & \quad + (n-A_0)(n-A_0+1)C_1 C_1^*\} \end{aligned} \quad (3.30)$$

Proof. The product in the summand becomes:

$$\left(\sum_{i=1}^m x_i k_i\right)\left(\sum_{i=1}^m y_i k_i\right) = \sum_{i=1}^m \sum_{\substack{j=1 \\ i \neq j}}^m x_i y_j k_i k_j + \sum_{i=1}^m x_i y_i k_i^2 \quad (3.31)$$

Changing the order of summation as in (3.1) and using theorem 3.4 and 3.5.

$$\begin{aligned} & \sum_{i=1}^m \sum_{\substack{j=1 \\ i \neq j}}^m x_i y_j \{(n - C_0)[(n - C_0 - 1)a_i a_j - (n - A_0)(a_i c_j + c_i a_j)] \\ & \quad + (n - A_0)(n - A_0 + 1)c_i c_j\} \\ & \quad + \sum_{i=1}^m x_i y_i \{(n - C_0)[(n - C_0 - 1)a_i^2 + (n - A_0)(c_i - a_i - 2a_i c_i)] \\ & \quad + (n - A_0)(n - A_0 + 1)c_i^2\} \\ & = \sum_{i=1}^m \sum_{j=1}^m x_i y_j \{(n - C_0)[(n - C_0 - 1)a_i a_j - (n - A_0)(a_i c_j + c_i a_j)] \\ & \quad + (n - A_0)(n - A_0 + 1)c_i c_j\} \\ & \quad + \sum_{i=1}^m x_i y_i (n - C_0)(n - A_0)(c_i - a_i) \\ & = (n - C_0)[(n - C_0 - 1)A_1 A_1^* + (n - A_0)(C_{1,1}^* - A_{1,1}^* - A_1 C_1^* - A_1^* C_1)] \\ & \quad + (n - A_0)(n - A_0 + 1)C_1 C_1^* \end{aligned} \quad (3.32)$$

□

Theorem 3.7. For integer $m \geq 1$:

$$\begin{aligned} & \sum_{\substack{k_1=0 \dots k_m=0 \\ \sum_{i=0}^m k_i=n}}^{a_1 \dots a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left(\sum_{i=1}^m x_i k_i \right)^2 = \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \\ & \cdot \frac{(n - C_0)[(n - C_0 - 1)A_1^2 + (n - A_0)(C_2 - A_2 - 2A_1 C_1)] + (n - A_0)(n - A_0 + 1)C_1^2}{(A_0 - C_0)(A_0 - C_0 - 1)} \end{aligned} \quad (3.33)$$

Proof. In theorem 3.6 taking $y_i = x_i$, then $A_1^* = A_1$, $C_1^* = C_1$, $A_{1,1}^* = A_2$ and $C_{1,1}^* = C_2$, giving this theorem. □

Theorem 3.8.

$$\begin{aligned}
& \sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}} \cdots \sum_{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left| \sum_{i=1}^m x_i k_i \right|^2 \\
&= \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)} \\
&\quad \cdot \{ (n - C_0)[(n - C_0 - 1)|A_1|^2 + (n - A_0)(C_{\text{abs}} - A_{\text{abs}} - A_1 \overline{C_1} - \overline{A_1} C_1)] \\
&\quad + (n - A_0)(n - A_0 + 1)|C_1|^2 \}
\end{aligned} \tag{3.34}$$

Proof. For this theorem the following is used.

$$\begin{aligned}
\left| \sum_{i=1}^m x_i k_i \right|^2 &= [\text{Re}(\sum_{i=1}^m x_i k_i)]^2 + [\text{Im}(\sum_{i=1}^m x_i k_i)]^2 \\
&= \left(\sum_{i=0}^m \text{Re}(x_i) k_i \right)^2 + \left(\sum_{i=0}^m \text{Im}(x_i) k_i \right)^2
\end{aligned} \tag{3.35}$$

Both of these terms are evaluated by theorem 3.7, replacing x_i by $\text{Re}(x_i)$ and then by $\text{Im}(x_i)$, and then the results are added, using the following.

$$\left(\sum_{i=1}^m \text{Re}(x_i) a_i \right)^2 + \left(\sum_{i=1}^m \text{Im}(x_i) a_i \right)^2 = [\text{Re}(\sum_{i=1}^m x_i a_i)]^2 + [\text{Im}(\sum_{i=1}^m x_i a_i)]^2 = |A_1|^2 \tag{3.36}$$

$$\left(\sum_{i=1}^m \text{Re}(x_i) c_i \right)^2 + \left(\sum_{i=1}^m \text{Im}(x_i) c_i \right)^2 = [\text{Re}(\sum_{i=1}^m x_i c_i)]^2 + [\text{Im}(\sum_{i=1}^m x_i c_i)]^2 = |C_1|^2 \tag{3.37}$$

$$\sum_{i=1}^m \text{Re}^2(x_i) a_i + \sum_{i=1}^m \text{Im}^2(x_i) a_i = \sum_{i=1}^m [\text{Re}^2(x_i) + \text{Im}^2(x_i)] a_i = \sum_{i=1}^m |x_i|^2 a_i = A_{\text{abs}} \tag{3.38}$$

$$\sum_{i=1}^m \text{Re}^2(x_i) c_i + \sum_{i=1}^m \text{Im}^2(x_i) c_i = \sum_{i=1}^m [\text{Re}^2(x_i) + \text{Im}^2(x_i)] c_i = \sum_{i=1}^m |x_i|^2 c_i = C_{\text{abs}} \tag{3.39}$$

and using for complex a, c : $2[\text{Re}(a)\text{Re}(c) + \text{Im}(a)\text{Im}(c)] = a\overline{c} + \overline{a}c$ [1]:

$$\begin{aligned}
& 2 \left[\left(\sum_{i=1}^m \text{Re}(x_i) a_i \right) \left(\sum_{i=1}^m \text{Re}(x_i) c_i \right) + \left(\sum_{i=1}^m \text{Im}(x_i) a_i \right) \left(\sum_{i=1}^m \text{Im}(x_i) c_i \right) \right] \\
&= 2 \left[\text{Re} \left(\sum_{i=1}^m x_i a_i \right) \text{Re} \left(\sum_{i=1}^m x_i c_i \right) + \text{Im} \left(\sum_{i=1}^m x_i a_i \right) \text{Im} \left(\sum_{i=1}^m x_i c_i \right) \right] \\
&= A_1 \overline{C_1} + \overline{A_1} C_1
\end{aligned} \tag{3.40}$$

and the theorem is proved. \square

Theorem 3.9.

$$\begin{aligned}
& \sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}}^{a_1 \cdots a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i=1}^m x_i k_i^2 \\
&= \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)} \\
&\quad \cdot \{(n - C_0)[(n - C_0 - 1)A_{1,2} + (n - A_0)(C_1 - A_1 - 2S_{1,1})] \\
&\quad + (n - A_0)(n - A_0 + 1)C_{1,2}\}
\end{aligned} \tag{3.41}$$

Proof. Changing the order of summation as in (3.1) and using theorem 3.4.

$$\begin{aligned}
& \sum_{i=1}^m x_i \{(n - C_0)[(n - C_0 - 1)a_i^2 + (n - A_0)(c_i - a_i - 2a_i c_i)] \\
&\quad + (n - A_0)(n - A_0 + 1)c_i^2\} \\
&= (n - C_0)[(n - C_0 - 1)A_{1,2} + (n - A_0)(C_1 - A_1 - 2S_{1,1})] \\
&\quad + (n - A_0)(n - A_0 + 1)C_{1,2}
\end{aligned} \tag{3.42}$$

□

Theorem 3.10. For integer $m \geq 1$ and $1 \leq p \leq m$:

$$\begin{aligned}
& \sum_{\substack{k_1=0 \cdots k_m=0 \\ \sum_{i=0}^m k_i=n}}^{a_1 \cdots a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] k_p^3 \\
&= \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)(A_0 - C_0 - 2)} \\
&\quad \cdot \{(n - C_0)(n - C_0 - 1)[(n - C_0 - 2)a_p^3 - (n - A_0)(3a_p^2 c_p + 3a_p^2 - 3a_p c_p - a_p + c_p)] \\
&\quad - (n - A_0)(n - A_0 + 1)[(n - A_0 + 2)c_p^3 - (n - C_0)(3a_p c_p^2 - 3c_p^2 + 3a_p c_p + a_p - c_p)]\}
\end{aligned} \tag{3.43}$$

Proof. The following modified infinite geometric series [7] is used:

$$\sum_{k=0}^{\infty} k^3 w^k = \frac{w}{(1-w)^2} + \frac{6w^2}{(1-w)^3} + \frac{6w^3}{(1-w)^4} \tag{3.44}$$

The first two terms gives a result identical to theorem 3.2 and 3.4, and for the third term:

$$\begin{aligned}
& \left[\prod_{i=1}^m \text{Res}_{w_i} \right] \frac{1}{w_m^n} \left[\prod_{i=1}^m (1 + w_i)^{a_i - c_i} w_i^{c_i - 1} \right] \frac{6(\frac{w_m}{w_p})^3}{(1 - \frac{w_m}{w_p})^4} \prod_{\substack{i=1 \\ i \neq p}}^{m-1} \frac{1}{1 - \frac{w_m}{w_i}} \\
&= \text{Res}_{w_m} \frac{(1 + w_m)^{a_m - c_m}}{w_m^{n - c_m - 2}} \text{Res}_{w_p} \frac{6(1 + w_p)^{a_p - c_p} w_p^{c_p}}{(w_p - w_m)^4} \prod_{\substack{i=1 \\ i \neq p}}^{m-1} (1 + w_m)^{a_i - c_i} w_m^{c_i}
\end{aligned} \tag{3.45}$$

The residue of w_p is evaluated with (3.5) using the product rule for the third derivative:

$$\begin{aligned}
& \operatorname{Res}_{w_p} \frac{6(1+w_p)^{a_p-c_p} w_p^{c_p}}{(w_p-w_m)^4} \\
&= (a_p-c_p)(a_p-c_p-1)(a_p-c_p-2)(1+w_m)^{a_p-c_p-3} w_m^{c_p} \\
&\quad + 3(a_p-c_p)(a_p-c_p-1)c_p(1+w_m)^{a_p-c_p-2} w_m^{c_p-1} \\
&\quad + 3(a_p-c_p)c_p(c_p-1)(1+w_m)^{a_p-c_p-1} w_m^{c_p-2} \\
&\quad + c_p(c_p-1)(c_p-2)(1+w_m)^{a_p-c_p} w_m^{c_p-3}
\end{aligned} \tag{3.46}$$

Using the absorption identity [5, 8] as in theorem 3.4, and adding as mentioned the result of theorem 3.2 and 3.4:

$$\begin{aligned}
& \binom{A_0-C_0}{n-C_0} \frac{1}{(A_0-C_0)(A_0-C_0-1)(A_0-C_0-2)} \\
& \cdot \{ (n-C_0)(n-C_0-1)(a_p-c_p)(a_p-c_p-1) \\
& \quad \cdot [(n-C_0-2)(a_p-c_p-2) + 3(A_0-C_0-2)c_p] \\
& \quad + (A_0-C_0-1)(A_0-C_0-2)c_p(c_p-1) \\
& \quad \cdot [(A_0-C_0)(c_p-2) + 3(n-C_0)(a_p-c_p)] \\
& \quad + (A_0-C_0-2)[3(n-C_0)(a_p-c_p)((n-C_0-1)(a_p-c_p-1) + 2(A_0-C_0-1)c_p) \\
& \quad + (A_0-C_0-1)(3(A_0-C_0)c_p(c_p-1) + (n-C_0)a_p - (n-A_0)c_p) \}
\end{aligned} \tag{3.47}$$

This result simplifies to the theorem. \square

Theorem 3.11. For integer $m \geq 2$, $1 \leq p \leq m$, $1 \leq q \leq m$ and $p \neq q$:

$$\begin{aligned}
& \sum_{\substack{k_1=0 \dots k_m=0 \\ \sum_{i=0}^m k_i=n}} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] k_p k_q^2 \\
&= \binom{A_0-C_0}{n-C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0-C_0)(A_0-C_0-1)(A_0-C_0-2)} \\
& \cdot \{ (n-C_0)(n-C_0-1)[(n-C_0-2)a_p a_q^2 - (n-A_0)(c_p a_q^2 + a_p a_q - a_p c_q + 2a_p a_q c_q)] \\
& \quad - (n-A_0)(n-A_0+1)[(n-A_0+2)c_p c_q^2 - (n-C_0)(a_p c_q^2 - c_p c_q + c_p a_q + 2c_p a_q c_q)] \}
\end{aligned} \tag{3.48}$$

Proof. The proof is similar to theorem 3.5 but with two quadratic and one cubic residues instead of two quadratic residues.

$$\begin{aligned}
& \operatorname{Res}_{w_m} \frac{(1+w_m)^{a_m-c_m}}{w_m^{n-c_m-2}} \operatorname{Res}_{w_p} \frac{(1+w_p)^{a_p-c_p} w_p^{c_p}}{(w_p-w_m)^2} \\
& \cdot \left[\operatorname{Res}_{w_q} \frac{(1+w_q)^{a_p-c_p} w_q^{c_p}}{(w_q-w_m)^2} + \operatorname{Res}_{w_q} \frac{2(1+w_q)^{a_p-c_p} w_q^{c_p}}{(w_q-w_m)^3} \right] \prod_{\substack{i=1 \\ i \neq p, i \neq q}}^{m-1} (1+w_m)^{a_i-c_i} w_m^{c_i}
\end{aligned} \tag{3.49}$$

The first product gives a result identical to theorem 3.5, and for the second product the residues are given by (3.15), and multiplying these out gives:

$$\begin{aligned}
& [(a_p - c_p)(1 + w_m)^{a_p - c_p - 1} w_m^{c_p} + c_p(1 + w_m)^{a_p - c_p} w_m^{c_p - 1}] \\
& \cdot [(a_q - c_q)(a_q - c_q - 1)(1 + w_m)^{a_q - c_q - 2} w_m^{c_q} + 2(a_q - c_q)c_q(1 + w_m)^{a_q - c_q - 1} w_m^{c_q - 1} \\
& \quad + c_q(c_q - 1)(1 + w_m)^{a_q - c_q} w_m^{c_q - 2}] \\
& = (a_p - c_p)(a_q - c_q)(a_q - c_q - 1)(1 + w_m)^{a_p - c_p + a_q - c_q - 3} w_m^{c_p + c_q} \\
& + [c_p(a_q - c_q)(a_q - c_q - 1) + 2(a_p - c_p)(a_q - c_q)c_q](1 + w_m)^{a_p - c_p + a_q - c_q - 2} w_m^{c_p + c_q - 1} \\
& + [(a_p - c_p)c_q(c_q - 1) + 2c_p(a_q - c_q)c_q](1 + w_m)^{a_p - c_p + a_q - c_q - 1} w_m^{c_p + c_q - 2} \\
& + c_p c_q (c_q - 1)(1 + w_m)^{a_p - c_p + a_q - c_q} w_m^{c_p + c_q - 3}
\end{aligned} \tag{3.50}$$

Using the absorption identity [5, 8] as in theorem 3.10, and adding as mentioned the result of theorem 3.5:

$$\begin{aligned}
& \binom{A_0 - C_0}{n - C_0} \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)(A_0 - C_0 - 2)} \\
& \cdot \{ (n - C_0)(n - C_0 - 1)(a_q - c_q)[(n - C_0 - 2)(a_p - c_p)(a_q - c_q - 1) \\
& \quad + (A_0 - C_0 - 2)(c_p(a_q - c_q - 1) + 2(a_p - c_p)c_q)] \\
& + (A_0 - C_0 - 1)(A_0 - C_0 - 2)c_q[(A_0 - C_0)c_p(c_q - 1) \\
& \quad + (n - C_0)((a_p - c_p)(c_q - 1) + 2c_p(a_q - c_q))] \\
& + (A_0 - C_0 - 2)[(n - A_0)(n - A_0 + 1)c_p c_q \\
& \quad + (n - C_0)((n - C_0 - 1)a_p a_q - (n - A_0)(a_p c_q + c_p a_q)] \}
\end{aligned} \tag{3.51}$$

This result simplifies to the theorem. \square

Theorem 3.12. For integer $m \geq 3$, $1 \leq p \leq m$, $1 \leq q \leq m$, $1 \leq r \leq m$, $p \neq q \neq r$ and $p \neq r$:

$$\begin{aligned}
& \sum_{\substack{k_1=0 \dots k_m=0 \\ \sum_{i=0}^m k_i = n}} \dots \sum_{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] k_p k_q k_r \\
& = \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)(A_0 - C_0 - 2)} \\
& \cdot \{ (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)a_p a_q a_r - (n - A_0)(a_p a_q c_r + a_p c_q a_r + c_p a_q a_r)] \\
& - (n - A_0)(n - A_0 + 1)[(n - A_0 + 2)c_p c_q c_r - (n - C_0)(c_p c_q a_r + c_p a_q c_r + a_p c_q c_r)] \}
\end{aligned} \tag{3.52}$$

Proof. The proof is similar to theorem 3.5 but with three quadratic residues instead of

two.

$$\begin{aligned}
& \text{Res}_{w_m} \frac{(1+w_m)^{a_m-c_m}}{w_m^{n-c_m-2}} \text{Res}_{w_p} \frac{(1+w_p)^{a_p-c_p} w_p^{c_p}}{(w_p-w_m)^2} \text{Res}_{w_q} \frac{(1+w_q)^{a_p-c_p} w_q^{c_p}}{(w_q-w_m)^2} \\
& \cdot \text{Res}_{w_r} \frac{(1+w_r)^{a_p-c_p} w_r^{c_p}}{(w_r-w_m)^2} \prod_{\substack{i=1 \\ i \neq p, i \neq q, i \neq r}}^{m-1} (1+w_m)^{a_i-c_i} w_m^{c_i}
\end{aligned} \tag{3.53}$$

The three last residues are given by (3.15), and multiplying these out gives:

$$\begin{aligned}
& [(a_p - c_p)(1+w_m)^{a_p-c_p-1} w_m^{c_p} + c_p(1+w_m)^{a_p-c_p} w_m^{c_p-1}] \\
& \cdot [(a_q - c_q)(1+w_m)^{a_q-c_q-1} w_m^{c_q} + c_q(1+w_m)^{a_q-c_q} w_m^{c_q-1}] \\
& \cdot [(a_r - c_r)(1+w_m)^{a_r-c_r-1} w_m^{c_r} + c_r(1+w_m)^{a_r-c_r} w_m^{c_r-1}] \\
& = (a_p - c_p)(a_q - c_q)(a_r - c_r)(1+w_m)^{a_p-c_p+a_q-c_q+a_r-c_r-3} w_m^{c_p+c_q+c_r} \\
& + [(a_p - c_p)(a_q - c_q)c_r + (a_p - c_p)c_q(a_r - c_r) + c_p(a_q - c_q)(a_r - c_r)] \\
& \cdot (1+w_m)^{a_p-c_p+a_q-c_q+a_r-c_r-2} w_m^{c_p+c_q+c_r-1} \\
& + [(a_p - c_p)c_q c_r + c_p(a_q - c_q)c_r + c_p c_q(a_r - c_r)] \\
& \cdot (1+w_m)^{a_p-c_p+a_q-c_q+a_r-c_r-1} w_m^{c_p+c_q+c_r-2} \\
& + c_p c_q c_r (1+w_m)^{a_p-c_p+a_q-c_q+a_r-c_r} w_m^{c_p+c_q+c_r-3}
\end{aligned} \tag{3.54}$$

Using the absorption identity [5, 8] as in theorem 3.11:

$$\begin{aligned}
& \left(\frac{A_0 - C_0}{n - C_0} \right) \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)(A_0 - C_0 - 2)} \\
& \cdot \{ (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)(a_p - c_p)(a_q - c_q)(a_r - c_r) + (A_0 - C_0 - 2) \\
& \quad \cdot ((a_p - c_p)(a_q - c_q)c_r + (a_p - c_p)c_q(a_r - c_r) + c_p(a_q - c_q)(a_r - c_r))] \\
& + (A_0 - C_0 - 1)(A_0 - C_0 - 2)[(A_0 - C_0)c_p c_q c_r \\
& \quad + (n - C_0)((a_p - c_p)c_q c_r + c_p(a_q - c_q)c_r + c_p c_q(a_r - c_r))] \}
\end{aligned} \tag{3.55}$$

This result simplifies to the theorem. \square

Theorem 3.13. For integer $m \geq 1$:

$$\begin{aligned}
& \sum_{\substack{k_1=0 \dots k_m=0 \\ \sum_{i=0}^m k_i=n}}^{a_1 \dots a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left(\sum_{i=1}^m x_i k_i \right)^3 \\
& = \left(\frac{A_0 - C_0}{n - C_0} \right) \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)(A_0 - C_0 - 2)} \\
& \cdot \{ (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)A_1^3 \\
& \quad + (n - A_0)(A_3 - C_3 + 3A_1(C_2 - A_2 - A_1 C_1))] \\
& - (n - A_0)(n - A_0 + 1)[(n - A_0 + 2)C_1^3 \\
& \quad + (n - C_0)(C_3 - A_3 + 3C_1(C_2 - A_2 - A_1 C_1))] \}
\end{aligned} \tag{3.56}$$

Proof. The product of the sum in the summand is expressed in sums with unequal indices:

$$\left(\sum_{i=1}^m x_i k_i\right)^3 = \sum_{\substack{p=1 \\ p \neq q \neq r, \\ p \neq r}}^m \sum_{q=1}^m \sum_{r=1}^m x_p x_q x_r k_p k_q k_r + 3 \sum_{p=1}^m \sum_{\substack{q=1 \\ p \neq q}}^m x_p x_q^2 k_p k_q^2 + \sum_{p=1}^m x_p^3 k_p^3 \quad (3.57)$$

Changing the order of summation, the multiple sum over the products of k_p , k_q and k_r is substituted from theorems 3.12, 3.11 and 3.10:

$$\begin{aligned} & \sum_{i=1}^m \sum_{\substack{j=1 \\ i \neq j \neq k, \\ i \neq k}}^m \sum_{k=1}^m x_i x_j x_k \{ (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)a_i a_j a_k - 3(n - A_0)a_i a_j c_k] \\ & \quad - (n - A_0)(n - A_0 - 1)[(n - A_0 - 2)c_i c_j c_k - 3(n - C_0)c_i c_j a_k] \} \\ & + 3 \sum_{i=1}^m \sum_{\substack{j=1 \\ i \neq j}}^m x_i x_j^2 \{ (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)a_i a_j^2 \\ & \quad - (n - A_0)(c_i a_j^2 + a_i a_j - a_i c_j + 2a_i a_j c_j)] \\ & \quad - (n - A_0)(n - A_0 + 1)[(n - A_0 + 2)c_i c_j^2 \\ & \quad - (n - C_0)(a_i c_j^2 - c_i c_j + c_i a_j + 2c_i a_j c_j) \} \\ & + \sum_{i=1}^m x_i^3 \{ (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)a_i^3 - \\ & \quad (n - A_0)(3a_i^2 c_i + 3a_i^2 - 3a_i c_i - a_i + c_i)] \\ & \quad - (n - A_0)(n - A_0 + 1)[(n - A_0 + 2)c_i^3 - \\ & \quad (n - C_0)(3a_i c_i^2 - 3c_i^2 + 3a_i c_i + a_i - c_i) \} \end{aligned} \quad (3.58)$$

The expressions of A_1^3 , C_1^3 , $A_1^2 C_1$ and $A_1 C_1^2$ as sums over unequal indices are given by equation (3.3):

$$\begin{aligned} & (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)A_1^3 - 3(n - A_0)A_1^2 C_1] \\ & - (n - A_0)(n - A_0 + 1)[(n - A_0 + 2)C_1^3 - 3(n - A_0)A_1 C_1^2] \\ & - 3 \sum_{i=1}^m \sum_{\substack{j=1 \\ i \neq j}}^m x_i x_j^2 [(n - C_0)(n - C_0 - 1)(n - A_0)(a_i a_j - a_i c_j) \\ & \quad + (n - A_0)(n - A_0 + 1)(n - C_0)(c_i c_j - c_i a_j)] \\ & - \sum_{i=1}^m x_i^3 [(n - C_0)(n - C_0 - 1)(n - A_0)(3a_i^2 - 3a_i c_i - a_i + c_i) \\ & \quad + (n - A_0)(n - A_0 - 1)(n - C_0)(3c_i^2 - 3a_i c_i - a_i + c_i)] \end{aligned} \quad (3.59)$$

The remaining sums can be expressed in A_1, C_1, A_2, C_2, A_3 and C_3 :

$$\begin{aligned}
& (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)A_1^3 - 3(n - A_0)A_1^2C_1] \\
& - (n - A_0)(n - A_0 + 1)[(n - A_0 + 2)C_1^3 - 3(n - A_0)A_1C_1^2] \\
& + 3(n - C_0)(n - A_0)[(n - C_0 - 1)(A_1C_2 - A_1A_2) \\
& \quad + (n - A_0 + 1)(C_1A_2 - C_1C_2)] \\
& + (n - C_0)(n - A_0)[(n - C_0 - 1) + (n - A_0 + 1)](A_3 - C_3)
\end{aligned} \tag{3.60}$$

This result simplifies to the theorem. \square

Theorem 3.14. For integer $m \geq 1$:

$$\begin{aligned}
& \sum_{\substack{k_1=0 \dots k_m=0 \\ \sum_{i=0}^m k_i=n}} \dots \sum_{i=1}^m \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i=1}^m x_i k_i^3 \\
& = \binom{A_0 - C_0}{n - C_0} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] \frac{1}{(A_0 - C_0)(A_0 - C_0 - 1)(A_0 - C_0 - 2)} \\
& \quad \cdot \{ (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)A_{1,3} \\
& \quad \quad + (n - A_0)(A_1 - C_1 - 3(A_{1,2} + S_{2,1} - S_{1,1}))] \\
& \quad \quad - (n - A_0)(n - A_0 + 1)[(n - A_0 + 2)C_{1,3} \\
& \quad \quad + (n - C_0)(C_1 - A_1 + 3(C_{1,2} - S_{1,2} - S_{1,1}))] \}
\end{aligned} \tag{3.61}$$

Proof. Changing the order of summation as in (3.1) and using theorem 3.10.

$$\begin{aligned}
& \sum_{i=1}^m x_i \{ (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)a_i^3 \\
& \quad - (n - A_0)(3a_i^2c_i + 3a_i^2 - 3a_ic_i - a_i + c_i)] \\
& \quad - (n - A_0)(n - A_0 + 1)[(n - A_0 + 2)c_i^3 \\
& \quad - (n - C_0)(3a_ic_i^2 - 3c_i^2 + 3a_ic_i + a_i - c_i)] \} \\
& = (n - C_0)(n - C_0 - 1)[(n - C_0 - 2)A_{1,3} \\
& \quad - (n - A_0)(3S_{2,1} + 3A_{1,2} - 3S_{1,1} - A_1 + C_1)] \\
& \quad - (n - A_0)(n - A_0 + 1)[(n - A_0 + 2)C_{1,3} \\
& \quad - (n - C_0)(3S_{1,2} - 3C_{1,2} + 3S_{1,1} + A_1 - C_1)] \}
\end{aligned} \tag{3.62}$$

which results in the theorem. \square

4 Proof of the Unrestricted Multi-sum Identities

Theorem 4.1. For integer $m \geq 1$:

$$\sum_{k_1=0}^{a_1} \dots \sum_{k_m=0}^{a_m} \prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} = 2^{A_0 - C_0} \prod_{i=1}^m \binom{a_i}{c_i} \tag{4.1}$$

Proof. The multiple sum of the product on the left side is a product of sums:

$$\sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} = \prod_{i=1}^m \sum_{k_i=0}^{a_i} \binom{a_i}{k_i} \binom{k_i}{c_i} \quad (4.2)$$

The trinomial revision identity [5, 8] is applied as in theorem 3.1:

$$\binom{a_i}{k_i} \binom{k_i}{c_i} = \binom{a_i - c_i}{k_i - c_i} \binom{a_i}{c_i} \quad (4.3)$$

Because this expression is zero when $c_i > a_i$, in the individual sums it can be assumed that $0 \leq c_i \leq a_i$. Then each sum can be evaluated as follows, where the upper bound a can be replaced by ∞ because the summand is zero when $k > a$:

$$\begin{aligned} \sum_{k=0}^a \binom{a-c}{k-c} &= \sum_{k=0}^{\infty} \operatorname{Res}_w \frac{(1+w)^{a-c}}{w^{k-c+1}} = \operatorname{Res}_w (1+w)^{a-c} w^{c-1} \sum_{k=0}^{\infty} \left(\frac{1}{w}\right)^k \\ &= \operatorname{Res}_w \frac{(1+w)^{a-c} w^{c-1}}{1-1/w} = \operatorname{Res}_w \frac{(1+w)^{a-c} w^c}{w-1} = 2^{a-c} \end{aligned} \quad (4.4)$$

The product of these individual sums gives the theorem. \square

Theorem 4.2. For integer $m \geq 1$ and $1 \leq p \leq m$:

$$\sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] k_p = 2^{A_0 - C_0 - 1} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (a_p + c_p) \quad (4.5)$$

Proof. The following modified infinite geometric series [7] is used:

$$\sum_{k=0}^{\infty} k w^k = \frac{w}{(1-w)^2} \quad (4.6)$$

One of the individual sums of the product becomes:

$$\begin{aligned} \sum_{k=0}^a \binom{a-c}{k-c} k &= \sum_{k=0}^a k \operatorname{Res}_w \frac{(1+w)^{a-c}}{w^{k-c+1}} = \operatorname{Res}_w (1+w)^{a-c} w^{c-1} \sum_{k=0}^{\infty} k \left(\frac{1}{w}\right)^k \\ &= \operatorname{Res}_w \frac{(1+w)^{a-c} w^{c-1}}{w(1-1/w)^2} = \operatorname{Res}_w \frac{(1+w)^{a-c} w^c}{(w-1)^2} = (a-c)2^{a-c-1} + c2^{a-c} \\ &= 2^{a-c-1}(a-c+2c) = 2^{a-c-1}(a+c) \end{aligned} \quad (4.7)$$

The other individual sums of the product remain identical, so the product becomes the theorem. \square

Theorem 4.3. For integer $m \geq 1$:

$$\sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i=1}^m x_i k_i = 2^{A_0 - C_0 - 1} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (A_1 + C_1) \quad (4.8)$$

Proof. Changing the order of summation and applying theorem 4.2:

$$\sum_{i=1}^m x_i(a_i + c_i) = A_1 + C_1 \quad (4.9)$$

□

Theorem 4.4. For integer $m \geq 1$ and $1 \leq p \leq m$:

$$\sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] k_p^2 = 2^{A_0 - C_0 - 2} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] [(a_p + c_p)^2 + a_p - c_p] \quad (4.10)$$

Proof. The following modified infinite geometric series [7] is used:

$$\sum_{k=0}^{\infty} k^2 w^k = \frac{w}{(1-w)^2} + \frac{2w^2}{(1-w)^3} \quad (4.11)$$

One of the individual sums of the product becomes:

$$\begin{aligned} \sum_{k=0}^a \binom{a-c}{k-c} k^2 &= \text{Res}_w (1+w)^{a-c} w^{c-1} \sum_{k=0}^{\infty} k^2 \left(\frac{1}{w}\right)^k \\ &= \text{Res}_w (1+w)^{a-c} w^{c-1} \left[\frac{1}{w(1-1/w)^2} + \frac{2}{w^2(1-1/w)^3} \right] \\ &= \text{Res}_w (1+w)^{a-c} w^c \left[\frac{1}{(w-1)^2} + \frac{2}{(w-1)^3} \right] \\ &= 2^{a-c-1} (a+c) + (a-c)(a-c-1)2^{a-c-2} + 2(a-c)c2^{a-c-1} + c(c-1)2^{a-c} \\ &= 2^{a-c-2} [2(a+c) + (a-c)(a-c-1) + 4(a-c)c + 4c(c-1)] \\ &= 2^{a-c-2} [(a+c)^2 + a-c] \end{aligned} \quad (4.12)$$

The other individual sums of the product remain identical, so the product becomes the theorem. □

Theorem 4.5. For integer $m \geq 1$, $1 \leq p \leq m$, $1 \leq q \leq m$ and $p \neq q$:

$$\sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] k_p k_q = 2^{A_0 - C_0 - 2} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (a_p + c_p)(a_q + c_q) \quad (4.13)$$

Proof. This theorem follows from applying theorem 4.2 twice. □

Theorem 4.6. For integer $m \geq 1$:

$$\begin{aligned} \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left(\sum_{i=1}^m x_i k_i \right) \left(\sum_{i=1}^m y_i k_i \right) \\ = 2^{A_0 - C_0 - 2} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] [A_{1,1}^* - C_{1,1}^* + (A_1 + C_1)(A_1^* + C_1^*)] \end{aligned} \quad (4.14)$$

Proof. The product in the summand becomes:

$$\left(\sum_{i=1}^m x_i k_i\right)\left(\sum_{i=1}^m y_i k_i\right) = \sum_{i=1}^m \sum_{\substack{j=1 \\ i \neq j}}^m x_i y_j k_i k_j + \sum_{i=1}^m x_i y_i k_i^2 \quad (4.15)$$

Changing the order of summation and using theorem 4.4 and 4.5.

$$\begin{aligned} & \sum_{i=1}^m \sum_{\substack{j=1 \\ i \neq j}}^m x_i y_j (a_i + c_i)(a_j + c_j) + \sum_{i=1}^m x_i y_i [(a_i + c_i)^2 + a_i - c_i] \\ &= \sum_{i=1}^m \sum_{j=1}^m x_i y_j (a_i + c_i)(a_j + c_j) + \sum_{i=1}^m x_i y_i (a_i - c_i) \\ &= A_{1,1}^* - C_{1,1}^* + (A_1 + C_1)(A_1^* + C_1^*) \end{aligned} \quad (4.16)$$

□

Theorem 4.7. For integer $m \geq 1$:

$$\begin{aligned} & \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left(\sum_{i=1}^m x_i k_i \right)^2 \\ &= 2^{A_0 - C_0 - 2} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] [A_2 - C_2 + (A_1 + C_1)^2] \end{aligned} \quad (4.17)$$

Proof. In the previous theorem taking $y_i = x_i$, then $A_1^* = A_1$, $C_1^* = C_1$, $A_{1,1}^* = A_2$ and $C_{1,1}^* = C_2$ gives this theorem. □

Theorem 4.8. For integer $m \geq 1$:

$$\begin{aligned} & \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left| \sum_{i=1}^m x_i k_i \right|^2 \\ &= 2^{A_0 - C_0 - 2} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (A_{\text{abs}} - C_{\text{abs}} + |A_1 + C_1|^2) \end{aligned} \quad (4.18)$$

Proof. Using the previous theorem and using the same method as in theorem 3.8 and using:

$$[\text{Re}(A_1) + \text{Re}(C_1)]^2 + [\text{Im}(A_1) + \text{Im}(C_1)]^2 = \text{Re}^2(A_1 + C_1) + \text{Im}^2(A_1 + C_1) = |A_1 + C_1|^2 \quad (4.19)$$

gives the theorem. □

Theorem 4.9. For integer $m \geq 1$:

$$\begin{aligned} & \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i=1}^m x_i k_i^2 \\ &= 2^{A_0 - C_0 - 2} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (A_1 - C_1 + A_{1,2} + C_{1,2} + 2S_{1,1}) \end{aligned} \quad (4.20)$$

Proof. Changing the order of summation and using theorem 4.4.

$$\begin{aligned} \sum_{i=1}^m x_i [(a_i + c_i)^2 + a_i - c_i] &= \sum_{i=1}^m x_i (a_i^2 + c_i^2 + 2a_i c_i + a_i - c_i) \\ &= A_{1,2} + C_{1,2} + 2S_{1,1} + A_1 - C_1 \end{aligned} \quad (4.21)$$

□

Theorem 4.10. For integer $m \geq 1$ and $1 \leq p \leq m$:

$$\sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] k_p^3 = 2^{A_0 - C_0 - 3} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (a_p + c_p) [(a_p + c_p)^2 + 3(a_p - c_p)] \quad (4.22)$$

Proof. The following modified infinite geometric series [7] is used:

$$\sum_{k=0}^{\infty} k^3 w^k = \frac{w}{(1-w)^2} + \frac{6w^2}{(1-w)^3} + \frac{6w^3}{(1-w)^4} \quad (4.23)$$

One of the individual sums of the product becomes:

$$\begin{aligned} \sum_{k=0}^a \binom{a-c}{k-c} k^3 &= \text{Res}_w (1+w)^{a-c} w^{c-1} \sum_{k=0}^{\infty} k^3 \left(\frac{1}{w}\right)^k \\ &= \text{Res}_w (1+w)^{a-c} w^{c-1} \left[\frac{1}{w(1-1/w)^2} + \frac{6}{w^2(1-1/w)^3} + \frac{6}{w^3(1-1/w)^4} \right] \\ &= \text{Res}_w (1+w)^{a-c} w^c \left[\frac{1}{(w-1)^2} + \frac{6}{(w-1)^3} + \frac{6}{(w-1)^4} \right] \\ &= 2^{a-c-1} (a+c) + 3[(a-c)(a-c-1)2^{a-c-2} + 2(a-c)c2^{a-c-1} + c(c-1)2^{a-c}] \\ &\quad + (a-c)(a-c-1)(a-c-2)2^{a-c-3} + 3(a-c)(a-c-1)c2^{a-c-2} \\ &\quad + 3(a-c)c(c-1)2^{a-c-1} + c(c-1)(c-2)2^{a-c} \\ &= 2^{a-c-3} (a+c) [(a+c)^2 + 3(a-c)] \end{aligned} \quad (4.24)$$

The other individual sums of the product remain identical, so the product becomes the theorem. □

Theorem 4.11. For integer $m \geq 1$, $1 \leq p \leq m$, $1 \leq q \leq m$ and $p \neq q$:

$$\sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] k_p k_q^2 = 2^{A_0 - C_0 - 3} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (a_p + c_p) [(a_q + c_q)^2 + a_q - c_q] \quad (4.25)$$

Proof. This theorem follows from applying theorem 4.2 and 4.4. □

Theorem 4.12. For integer $m \geq 3$, $1 \leq p \leq m$, $1 \leq q \leq m$, $1 \leq r \leq m$, $p \neq q \neq r$ and $p \neq r$:

$$\sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] k_p k_q k_r = 2^{A_0 - C_0 - 3} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (a_p + c_p)(a_q + c_q)(a_r + c_r) \quad (4.26)$$

Proof. This theorem follows from applying theorem 4.2 three times. \square

Theorem 4.13. For integer $m \geq 1$:

$$\begin{aligned} & \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \left(\sum_{i=1}^m x_i k_i \right)^3 \\ &= 2^{A_0 - C_0 - 3} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] (A_1 + C_1) [(A_1 + C_1)^2 + 3(A_2 - C_2)] \end{aligned} \quad (4.27)$$

Proof. Changing the order of summation and using theorem 4.10, 4.11 and 4.12.

$$\begin{aligned} & \sum_{\substack{i=1 \\ i \neq j \neq k}}^m \sum_{\substack{j=1 \\ i \neq j}}^m \sum_{\substack{k=1 \\ i \neq j \neq k}}^m x_i x_j x_k (a_i + c_i)(a_j + c_j)(a_k + c_k) \\ &+ 3 \sum_{\substack{i=1 \\ i \neq j}}^m \sum_{j=1}^m x_i x_j^2 (a_i + c_i)((a_j + c_j)^2 + a_j - c_j) \\ &+ \sum_{i=1}^m x_i^3 (a_i + c_i)[(a_i + c_i)^2 + 3(a_i - c_i)] \\ &= (A_1 + C_1)^3 + 3 \sum_{\substack{i=1 \\ i \neq j}}^m \sum_{j=1}^m x_i x_j^2 (a_i + c_i)(a_j - c_j) + 3 \sum_{i=1}^m x_i^3 (a_i + c_i)(a_i - c_i) \\ &= (A_1 + C_1)^3 + 3(A_1 + C_1)(A_2 - C_2) = (A_1 + C_1)[(A_1 + C_1)^2 + 3(A_2 - C_2)] \end{aligned} \quad (4.28)$$

\square

Theorem 4.14. For integer $m \geq 1$:

$$\begin{aligned} & \sum_{k_1=0}^{a_1} \cdots \sum_{k_m=0}^{a_m} \left[\prod_{i=1}^m \binom{a_i}{k_i} \binom{k_i}{c_i} \right] \sum_{i=1}^m x_i k_i^3 \\ &= 2^{A_0 - C_0 - 3} \left[\prod_{i=1}^m \binom{a_i}{c_i} \right] [A_{1,3} + C_{1,3} + 3(A_{1,2} - C_{1,2} + S_{1,2} + S_{2,1})] \end{aligned} \quad (4.29)$$

Proof. Changing the order of summation and using theorem 4.10.

$$\begin{aligned}
& \sum_{i=1}^m x_i(a_i + c_i)[(a_i + c_i)^2 + 3(a_i - c_i)] \\
&= \sum_{i=1}^m x_i(a_i^3 + c_i^3 + 3a_i^2 - 3c_i^2 + 3a_i c_i^2 + 3a_i^2 c_i) \\
&= A_{1,3} + C_{1,3} + 3(A_{1,2} - C_{1,2} + S_{1,2} + S_{2,1})
\end{aligned} \tag{4.30}$$

□

References

- [1] L.V. Ahlfors, *Complex Analysis*, McGraw-Hill, 1979.
- [2] G.P. Egorychev, *Integral Representation and the Computation of Combinatorial Sums*, Translations of Mathematical Monographs, 59, Amer. Math. Soc., 1984.
- [3] H.W. Gould, *Combinatorial Identities*, rev. ed., Morgantown, 1972.
- [4] H.W. Gould, H.M. Srivastava, Some Combinatorial Identities Associated with the Vandermonde Convolution, *Appl. Math. Comput.* 84 (1997) 97-102.
- [5] R.L. Graham, D.E. Knuth, O. Patashnik, *Concrete Mathematics, A Foundation for Computer Science*, 2nd ed., Addison-Wesley, 1994.
- [6] M.J. Kronenburg, A Generalization of the Chu-Vandermonde Convolution and some Harmonic Number Identities, [arXiv:1701.02768 \[math.CO\]](#)
- [7] M.J. Kronenburg, Some Weighted Generalized Fibonacci Number Summation Identities, Part 1, [arXiv:1903.01407 \[math.NT\]](#)
- [8] M.J. Kronenburg, The Binomial Coefficient for Negative Arguments, [arXiv:1105.3689 \[math.CO\]](#)
- [9] R. Meštrović, Several Generalizations and Variations of Chu-Vandermonde Identity, [arXiv:1807.10604 \[math.CO\]](#)