

Non-asymptotic estimation of risk measures using stochastic gradient Langevin dynamics

Jiarui Chu*

Ludovic Tangpi†

November 25, 2021

Abstract

In this paper we will study the approximation of arbitrary law invariant risk measures. As a starting point, we approximate the average value at risk using stochastic gradient Langevin dynamics, which can be seen as a variant of the stochastic gradient descent algorithm. Further, the Kusuoka’s spectral representation allows us to bootstrap the estimation of the average value at risk to extend the algorithm to general law invariant risk measures. We will present both theoretical, non-asymptotic convergence rates of the approximation algorithm and numerical simulations.

Keywords Convex risk measure · Stochastic Optimization · Risk minimization · Average value at risk · Stochastic gradient Langevin

Mathematics Subject Classification 91G70 · 90C90

Statements and Declarations The authors gratefully acknowledge support from the NSF grant DMS-2005832. The authors have no competing interests to declare.

1 Introduction

Every financial decision involves some degree of risk. Quantifying risk associated with a future random outcome allows organizations to compare financial decisions and develop risk management plans to prepare for potential loss and uncertainty. By the seminal work of Artzner, Delbaen, Eber, and Heath [4], the canonical way to quantify the riskiness of a *random* financial position X is to compute the *number* $\rho(X)$ for a convex risk measure ρ , whose definition we recall.

Definition 1.1. A mapping $\rho : \mathbb{L}^\infty \rightarrow \mathbb{R}$ is a convex risk measure if it satisfies the following conditions for all $X, Y \in \mathcal{X}$:

- *translation invariance:* $\rho(X + m) = \rho(X) - m$ for all $m \in \mathbb{R}$
- *monotonicity:* $\rho(X) \geq \rho(Y)$ if $X \leq Y$
- *convexity:* $\rho(\lambda X + (1 - \lambda)Y) \leq \lambda\rho(X) + (1 - \lambda)\rho(Y)$ for $\lambda \in [0, 1]$.

Intuitively¹, $\rho(X)$ measures the minimum amount of capital that should be added to the current financial position X to make it acceptable. Due to its fundamental importance in quantitative finance, the theory of risk measures (sometimes called quantitative risk management) has been extensively developed. We refer for instance to [20; 30; 44; 18; 47; 17; 35; 48] for a few milestones and the influential textbooks of Alexander J. McNeil and Frey [2] and Föllmer and Schied [31] for overviews.

An important problem for risk managers in practice is to efficiently simulate the number $\rho(X)$ for a financial position X and a risk measure ρ . The difficulty here stems from the fact that, unless ρ is a “simple enough” risk measure and the law of X belongs to a tractable family of distributions, there are no closed form formula allowing to compute $\rho(X)$. The goal of this paper is to develop a method allowing to *numerically simulate* the riskiness $\rho(X)$ for general convex risk measures, and when the law of X is not necessarily known (as it is the case in practical applications).

One commonly used measure of the riskiness of a financial position is the value at risk (VaR). For a given risk intolerance $u \in (0, 1)$, the value at risk $\text{VaR}_u(X)$ of X is the $(1 - u)$ -quantile of the distribution of X . Despite the various shortcomings

*Princeton University, ORFE, jiaruic@princeton.edu

†Princeton University, ORFE, ludovic.tangpi@princeton.edu

¹In the rest of the paper, for notation simplicity, we will assume risk measures to be increasing. That is, we work with $\rho(-X)$. This does not restrict the generality.

of this measure of risk documented by the academic community [2], VaR remains the standard in the banking industry, and due to its widespread use, the computation of VaR has been extensively studied. We refer interested readers for instance to [37; 40; 12; 27], and references therein for various simulation techniques. Recommendations [53] from the Basel Committee on Banking Supervision which oversees risk management for financial institutions have revived the development of convex risk measures such as the average value at risk (AVaR), also called conditional value at risk or expected shortfall. This risk measure is the expected loss given that losses are greater than or equal to the VaR. That is, AVaR_u is given by:

$$\text{AVaR}_u(X) := \mathbb{E}[X|X > \text{VaR}_u(X)]. \quad (1.1)$$

For general distributions of X , $\text{AVaR}_u(X)$ usually does not have closed form expressions. Therefore, in practice, numerical estimations are often required. As a result, the estimation of AVaR has received considerable attention. We refer for instance to works by Eckstein and Kupper [26] and Bühler, Gonon, Teichmann, and Wood [11] in which (among other things) the simulation of optimized certainty equivalents (of which AVaR is a particular case) are considered using deep learning techniques. One approximation technique for the AVaR is based on Monte-Carlo type algorithm. In this direction, let us refer for instance to Hong and Liu [39] and Chen [15] on Monte Carlo estimation of VaR and AVaR, Zhu and Zhou [67] on nested Monte Carlo estimation. More recently, motivated by developments of gradient descent methods in stochastic optimization, (in particular the stochastic Langevin gradient descent (SGLD) technique), Sabanis and Zhang [59] provide non-asymptotic error bounds for the estimation of AVaR. Other works developing such gradient descent techniques in the context of risk management include Iyengar and Ma [43], Tamar, Glassner, and Mannor [62] and Soma and Yoshida [61]. Essentially, these papers take advantage of new developments in machine learning and optimization, see e.g. Allen-Zhu [3], Gelfand and Mitter [36], Nesterov [52] and Raginsky, Rakhlin, and Telgarsky [56]. Let us also mention the recent work of Reppen and Soner [57] who develop a data-driven approach based on ideas from learning theory.

In this work we go beyond the numerical simulation of AVaR by extending stochastic gradient descent type techniques to compute a large family of risk measures, including the AVaR. We are interested in this work in deriving explicit (non-asymptotic) error estimates for the approximation, as well as deviation inequalities. We will restrict our attention to law-invariant convex risk measures (whose definition we recall below), since in practice, only the law of a financial position can be (approximately) observed. In fact, the requirement for a risk measure to be law-invariant is natural and is satisfied by most risk measures².

Definition 1.2. [34] *A risk measure ρ is law-invariant if for all X, Y with the same distribution, we have $\rho(X) = \rho(Y)$.*

To the best of our knowledge, the papers considering (non-parametric) estimation of general convex risk measures are Weber [64], Belomestny and Krättschmer [7] and Bartl and Tangp [5]. These papers consider a (data-driven) Monte-Carlo estimation method by proposing a plug-in estimator based on the empirical measure of the historical observations of the underlying distribution of the random outcome. Weber [64] proves a large deviation theorem, and Belomestny and Krättschmer [7] provide a central limit theorem. Note that both of these papers give asymptotic estimation results. Bartl and Tangpi [5] provide sharp non-asymptotic convergence rates for the estimation.

To estimate general law-invariant convex risk measures, we rely on the Kusuoka’s spectral representation [48]. Intuitively, this representation says any law invariant risk measure can be constructed as an integral of the AVaR risk measure. Therefore, the first step of our approximation of general law-invariant risk measures is to estimate AVaR. Since we would like to analyze approximation algorithms for the risk of claims with possibly non-convex payoffs, we employ the idea of Raginsky, Rakhlin, and Telgarsky [56] and use the stochastic gradient Langevin dynamic which, essentially, adds a Gaussian noise to the unbiased estimate of the gradient in stochastic gradient descent. To quantify the distance between the estimator and the true value of the risk measure, we present non-asymptotic rates on the mean squared estimation error and a matching deviation inequality in the case of a AVaR, and more importantly provide rate on the mean squared estimation error of general law-invariant risk measures. The proof of the mean squared error of estimating the AVaR makes use of the observation that the SGLD algorithm is a variant of the Euler-Maruyama discretization of the solution of the Langevin stochastic differential equation (SDE). This observation allows us to use results on the convergence rate of the Euler-Maruyama scheme and classical techniques of deriving the convergence rate of the solution of the Langevin SDE to the invariant measure. For the proof of the deviation inequality in the case of AVaR, we prove the existence of an exponential moment of the discretized solution of the Langevin SDE in addition to using a concentration inequality for the Euler-Maruyama scheme of SDEs [33]. For the rate on the mean squared estimation error of the general case, our proof relies heavily on Kusuoka’s representation which allows to build general law-invariant risk measures from AVaR.

Beyond our theoretical guarantees for the convergence of approximation algorithms for general convex risk measures, the present work also contributes to the non-convex optimization literature in that we propose a new proof for the

²All risk measures considered in this work will be implicitly assumed to be convex law-invariant risk measures.

convergence of SGLD algorithms for some non-convex objective functions. The idea is essentially to reduce the problem into the analysis of contractivity properties of the semi-group originating from a Langevin diffusion with non-convex potential. This problem was notably investigated by Eberle [25].

The paper is organized as follows: We start by describing formally the approximation techniques and presenting the main results in Section 2. In the same section, we also present numerical results on the estimation of AVaR. In Section 3, we prove the rates on the mean squared error and the deviation inequality for the estimation of AVaR. The derivation of the mean squared error for the estimation of a general law-invariant risk measure is done in Section 4.

Notations: Let $\mathbb{N}^* := \mathbb{N} \setminus \{0\}$ and let \mathbb{R}_+^* be the set of real positive numbers. Fix an arbitrary Polish space E endowed with a metric d_E . Throughout this paper, for every p -dimensional E -valued vector e with $p \in \mathbb{N}^*$, we denote by e^1, \dots, e^p its coordinates. For $(\alpha, \beta) \in \mathbb{R}^p \times \mathbb{R}^p$, we also denote by $\alpha \cdot \beta$ the usual inner product, with associated norm $\|\cdot\|$, which we simplify to $|\cdot|$ when p is equal to 1. For any $(\ell, c) \in \mathbb{N}^* \times \mathbb{N}^*$, $E^{\ell \times c}$ will denote the space of $\ell \times c$ matrices with E -valued entries.

Let $\mathcal{B}(E)$ be the Borel σ -algebra on E (for the topology generated by the metric d_E on E). For any $p \geq 1$, for any two probability measures μ and ν on $(E, \mathcal{B}(E))$ with finite p -moments, we denote by $\mathcal{W}_p(\mu, \nu)$ the p -Wasserstein distance between μ and ν , that is

$$\mathcal{W}_p(\mu, \nu) := \left(\inf_{\alpha \in \Gamma(\mu, \nu)} \int_{E \times E} d(x, y)^p \alpha(dx, dy) \right)^{1/p},$$

where the infimum is taken over the set $\Gamma(\mu, \nu)$ of all couplings π of μ and ν , that is, probability measures on $(E^2, \mathcal{B}(E)^{\otimes 2})$ with marginals μ and ν on the first and second factors respectively.

2 Approximation technique and main results

In this section we rigorously describe the approximation method developed in this article as well as our main results. Throughout, we fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ on which all random variables will be defined, unless otherwise stated. Let us denote by \mathbb{L}^∞ the space of essentially bounded random variables on this probability space. The starting point of our method is based on the following spectral representation of law-invariant risk measures due to Kusuoka [48]:

Theorem 2.1. *A mapping $\rho : \mathbb{L}^\infty \rightarrow \mathbb{R}$ is a law-invariant risk measure if and only if it satisfies*

$$\rho(X) = \sup_{\gamma \in \mathcal{M}} \left(\int_{[0,1]} AVaR_u(X) \gamma(du) - \beta(\gamma) \right), \text{ for all } X \in \mathbb{L}^\infty \quad (2.1)$$

for some functional $\beta : \mathcal{M} \rightarrow [0, \infty)$, where \mathcal{M} is the set of all Borel probability measures on $[0, 1]$.

In fact, this spectral representation suggests that the risk measure AVaR is the ‘‘basic building block’’ allowing to construct all convex law-invariant risk measures. Thus, the idea will be to propose an approximation algorithm for AVaR that will be later bootstrapped to derive an algorithm for general law invariant risk measures. This approach is also used in [5] for a very different approximation method.

2.1 Approximation of average value at risk

Let us first focus on estimating AVaR. For this purpose, recall (see e.g. [31, Proposition 4.51]) that for every $X \in \mathbb{L}^\infty$ and $u \in (0, 1)$, $AVaR_u(X)$ takes the form

$$AVaR_u(X) = \inf_{q \in \mathbb{R}} \left(\frac{1}{1-u} \mathbb{E}[(X - q)^+] + q \right).$$

In other words, $AVaR_u(X)$ is nothing but the value of a stochastic optimization problem. In most financial applications, the contingent claim X whose risk is assessed is of the form $X = f(r, S)$ where $S = (S^1, \dots, S^d)$ is a d -dimensional random vector of risk factors, and $r \in \mathcal{X}$, see e.g. [2, Section 2.1] for details. Note that here the space \mathcal{X} can be infinite dimensional. A standard practice to approach the infinite dimensional case is to use neural networks for approximation, which leads to non-convex objective functions. Therefore, we allow f to be non-convex with certain regularity conditions. A standard example arises when X is the profit and loss (P&L) of an investment strategy. In this case, r is the portfolio and the random vector S represents (increments) of the stock prices. That is, $f(r, S) := \sum_{i=1}^d r_i (S_1^i - S_0^i)$ where S_1^i and

S_0^i are the values of the stock i at time 1 and 0, respectively. Hence, we let r be in a finite dimensional space, and our goal will be to estimate the value of the (multi-dimensional) risk minimization problem

$$\begin{aligned}\overline{\text{AVaR}}_u(f) &:= \inf_{r \in \mathbb{R}^d} \text{AVaR}_u(f(r, S)) \\ &= \inf_{r \in \mathbb{R}^d, q \in \mathbb{R}} \left(\frac{1}{1-u} \mathbb{E} \left[(f(r, S) - q)^+ \right] + q \right).\end{aligned}\tag{2.2}$$

A natural way to numerically solve such problems is by gradient descent. However, when the dataset is large, gradient descent usually does not perform well, since computing the gradient on the full dataset at each iteration is computationally expensive.

Among others, one method that has been proposed to get around the high computational cost of gradient descent is the stochastic gradient descent (SGD) algorithm, which replaces the true gradient with an unbiased estimate calculated from a random subset of the data. A more recent approach, called the Stochastic Langevin Gradient Descent, injects a random noise to an unbiased estimate of the gradient at each iteration of the SGD algorithm. Originally introduced by [Welling and Teh](#). [65] as a tool for Bayesian posterior sampling on large scale and high dimensional datasets, SGLD maintains the scalability property of SGD, and has a few advantages over the SGD. By adding a noise to SGD, SGLD navigates out of saddle points and local minima more easily [9], outperforms SGD in terms of accuracy [21], and overcomes the curse of dimensionality [16]. Moreover, SGLD also applies to cases where the objective function is non-convex but sufficiently regular [36] [56].

We will apply the SGLD in the present context of estimation of AVaR. Recall that our goal is to solve the optimization problem given in equation (2.2). Consider the (objective) function

$$\tilde{L}(r, q) := \frac{1}{1-u} \mathbb{E} \left[(f(r, S) - q)^+ \right] + q$$

and, given a strictly positive constant $\gamma > 0$, let

$$L(r, q) := \tilde{L}(r, q) + \frac{\gamma}{2} q^2\tag{2.3}$$

be the usual penalized objective function. Since for γ small we have

$$\inf_{(m, r) \in \mathbb{R}^{d+1}} L(r, q) \approx \overline{\text{AVaR}}_u(f),$$

we will approximate the left hand side above using the SGLD algorithm, which consists in approximating its minimizer by the (support of the) invariant measure of the Markov chain $(Z_{m,h}^\lambda)$ given by

$$Z_{m+1,h}^\lambda = Z_{m,h}^\lambda - \nabla L(Z_{m,h}^\lambda)h + \sqrt{2\lambda^{-1}}\xi_m,\tag{2.4}$$

where $(\xi_m)_{m \geq 1}$ are independent Gaussian random variables. In the practice of financial risk management, the distribution of S is typically unknown. This is a well-studied issue in quantitative finance, refer for instance to [6; 19; 46] and the references therein. In particular, ∇L cannot be directly computed. It will be replaced by an unbiased estimator. Following Monte-Carlo simulation ideas, we let (S^1, \dots, S^P) be independent copies of S and $(\widetilde{W}^1, \dots, \widetilde{W}^N)$ be independent Brownian motions, and we thus let

$$\tilde{\ell}(z) := \frac{1}{P} \sum_{p=1}^P \frac{1}{1-u} (f(r, S^p) - m)^+ + m \quad , \quad \text{and } \ell(z) = \tilde{\ell}(z) + \frac{\gamma}{2} m^2 \quad \text{with } z = (m, r).$$

In the following, we will take $P = N$ for simplicity. Put

$$\tilde{Z}_{m+1,h}^{\lambda,n} = \tilde{Z}_{m,h}^{\lambda,n} - \nabla \ell(\tilde{Z}_{m,h}^{\lambda,n})h + \sqrt{2\lambda^{-1}}\Delta \widetilde{W}_h^n,\tag{2.5}$$

with $\Delta \widetilde{W}_h^n := \widetilde{W}_{m+1}^n - \widetilde{W}_m^n$. Hence we will show that

$$\widetilde{\text{AVaR}}_u(f) := \frac{1}{N} \sum_{n=1}^N \ell(\tilde{Z}_{M,h}^{\lambda,n})\tag{2.6}$$

approximates $\overline{\text{AVaR}}_u(f)$. Note that the optimal portfolio r can be easily recovered. It is simply the last d coordinates of $\tilde{Z}_{M,h}^{\lambda,n}$. Similarly, the value-at-risk can be obtained from the Markov chain $\tilde{Z}_{M,h}^{\lambda,n}$. See Remark 3.1 for details. Let us now formulate the assumptions we make on f and the random vector S .

Assumption 2.2. The random variable S takes values in \mathbb{R}^d and the function $f : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ is Borel measurable, and they satisfy

- (i) S has finite fourth moment.
- (ii) The function $r \mapsto f(r, s)$ Lipschitz-continuous and continuously differentiable for all $s \in \mathbb{R}^d$.
- (iii) $\inf_s f(r, s)/\|r\| \rightarrow \infty$ when $\|r\| \rightarrow \infty$ and $\inf_r \mathbb{E}[f(r, S)] > 0$.
- (iv) The random variable $\nabla_r f(r, S)$ is bounded, uniformly in r , and $\nabla_s f(r, \cdot)$ is Lipschitz, uniformly in r
- (v) Consider the function κ defined as

$$\kappa(u) := \inf \left\{ -\sqrt{2\lambda} \frac{(z - z') \cdot (\nabla L(z') - \nabla L(z))}{\|z - z'\|}, z, z' \in \mathbb{R}^{d+1} : \|z - z'\| = u \right\}.$$

It holds

$$\liminf_{u \rightarrow \infty} \kappa(u) > 0 \quad \text{and} \quad \int_0^1 u \kappa(u)^- du < \infty.$$

Let us briefly comment on these conditions before stating the result. The integrability, regularity, coercivity and lower boundedness conditions (i) – (iii) allow to ensure that the problem is well-posed. The boundedness condition (iv) is assumed mostly to simplify the exposition. Most of our statements will remain true if it is replaced by a suitable integrability condition. We introduce the more involved condition (v) to make for the possible lack of convexity of the objective function L . This condition is by now standard when employing coupling by reflection techniques to prove contractivity of diffusion semigroups. We refer for instance to Eberle [25; 24] or the earlier work of Chen and Li [14]. Note, for instance, that this condition is automatically satisfied if f is convex (since in this case L is strongly convex) or when L is strictly convex outside a given ball (see [25, Example 1]).

The following is the first main result of this work:

Theorem 2.3. Let Assumptions 2.2 hold. Let $t, M > 0$ be such that $h = \frac{t}{M^2}$. For all $t, \gamma, \lambda > 0$ and $M, N \in \mathbb{N}^*$, if $h < \frac{1}{\frac{2}{1-u} + \gamma} \wedge 1$ and $\frac{1}{\lambda} \leq \gamma \leq 1$, we have

$$\mathbb{E} \left[\left| \frac{1}{N} \sum_{n=1}^N \ell(\tilde{Z}_{M,h}^{\lambda,n}) - \overline{\text{AVaR}}_u(f) \right|^2 \right] \leq C_{(M,\lambda,t,u)} \gamma^2 + C_{(u,S,\lambda,t)} \frac{1}{N} + \frac{C_{(u,t)}}{M^4} + C_{(d,\lambda,u)} e^{-tC_{(d,\lambda)}} + \frac{C_{(u)}}{\lambda^2} \quad (2.7)$$

for some positive constants $C_{(M,\lambda,t,u)}, C_{(u,S,\lambda,t)}, C_{(u,t)}, C_{(d,\lambda,u)}, C_{(d,\lambda)}$ and $C_{(u)}$.

Theorem 2.3 provides a non-asymptotic rate for the convergence of the estimator $\frac{1}{N} \sum_{n=1}^N \ell(\tilde{Z}_{M,h}^{\lambda,n})$ to the (optimized) average value at risk. Such a rate is crucial in applications since it gives a precise order of magnitude for the choice of the parameters M, N, γ and λ needed to achieve a desired order of accuracy. Moreover, the rate is independent of the dimension d , implying in particular that the rate is not made worst when increasing the size of the portfolio $S = (S^1, \dots, S^d)$ (or in general the number of risk factors). Furthermore, observe that this estimator $\overline{\text{AVaR}}_u(f)$ is rather easy to simulate: one only needs to simulate N independent Gaussian random variables, for each of them simulate the iterative scheme (2.5) and compute the empirical average of the outcomes. We provide numerical results on the estimation of AVaR in Section 2.3 below.

Our second main result complements the above convergence rate with a matching deviation inequality.

Theorem 2.4. Let the conditions of Theorem 2.3 be satisfied. Let $h = \frac{t}{M^2}$, and $h, \lambda > 0, 0 < \gamma < 1$ and $M, N \in \mathbb{N}^*$ be given such that $h < \frac{1}{\frac{2}{1-u} + \gamma}$. If γ is small enough and λ, M large enough, then for every $\epsilon > 0$, we have

$$\mathbb{P} \left(\left| \frac{1}{N} \sum_{n=1}^N \ell \left(\tilde{Z}_{M,h}^{\lambda,n} \right) - \overline{\text{AVaR}}(f) \right| \geq \epsilon \right) \leq 2 \exp \left(- \frac{\epsilon^2 N}{\Psi(M, t, u)} \right) + C_{(M,\lambda,\gamma)} e^{-\epsilon^2/64}$$

with $\Psi(M, h, u)$ given by

$$\Psi(M, h, u) := 1152 \frac{t}{M} (1-u)^{-1} \lambda^{-1} \exp \left(2 \left(\frac{1}{1-u} + 1 \right) \frac{t}{M} \right) \quad (2.8)$$

and for some constants $C_{(m,h,\lambda)}, C_{(\lambda,\gamma,h,M,u)} > 0$.

Remark 2.5. Observe that the method developed here can also allow (with minor changes) to simulate the value function of utility maximization problems of the form

$$\sup_r \mathbb{E}^\mu [U(f(r, S))]$$

where U is a concave utility function and \mathbb{E}^μ the expectation when $S \sim \mu$, or even of the robust utility maximization problem

$$\sup_r \inf_{\mu \in \mathcal{P}} \mathbb{E}^\mu[U(f(r, S))]$$

where \mathcal{P} is the set of possible distributions of S . In the latter case, one will need to compute (or find an appropriate unbiased estimator of) ∇L with

$$L(r) := \inf_{\mu \in \mathcal{P}} \mathbb{E}^\mu[U(f(r, S))].$$

This is easily done for instance when \mathcal{P} is a ball with respect to the Wasserstein metric around a given distribution μ_0 , see e.g. [6].

2.2 Approximation of general convex risk measures

Let us return to the problem of approximating general law-invariant convex risk measures. In this context (as in the case of AVaR) our goal is to simulate the optimized risk measure

$$\bar{\rho}(f) := \inf_{r \in \mathbb{R}^d} \rho(f(r, S)).$$

To that end, let us recall a notion of regularity of risk measures introduced in [5] that will be needed to derive an explicit non-asymptotic convergence rate. Recall that a random variable X^* is said to follow the Pareto distribution with scale parameter $x > 0$ and shape parameter $q > 0$ if

$$P(X \geq t) = \begin{cases} (x/t)^q & \text{if } t \geq x \\ 1 & \text{if } t < x. \end{cases}$$

Definition 2.6. [5] Let $q \in (1, \infty)$, and let X^* follow Pareto distribution with scale parameter 1 and shape parameter q . A convex risk measure $\rho : \mathbb{L}^\infty \rightarrow \mathbb{R}$ is said to be q -regular if it satisfies

$$\sup_{n \in \mathbb{N}} \rho(X^* \wedge n) < \infty.$$

We refer to [5] for a discussion on this notion of regularity, but note for instance that AVaR is q -regular for all $q > 1$ and that this notion of regularity is slightly stronger than the well-known Fatou property and the Lebesgue property often assumed for risk measures, see e.g. Föllmer and Schied [31]. Moreover, one consequence of q -regularity is the following slight refinement of Kusuoka's representation: The risk measure ρ satisfies

$$\rho(f(r, S)) = \sup_{\gamma \in \mathcal{M}: s.t. \beta(\gamma) \leq b} \left(\int_{[0,1]} \text{AVaR}_u(f(r, S)) \gamma(du) - \beta(\gamma) \right). \quad (2.9)$$

for some $b > 0$, see [5, Lemma 4.4] for details. Thus, the estimator we consider for $\bar{\rho}$ is given by

$$\tilde{\rho}^\delta(f) := \sup_{\gamma \in \mathcal{M}: \beta(\gamma) \leq b} \left(\int_{[0,\delta]} \widetilde{\text{AVaR}}_u(f) \gamma(du) - \beta(\gamma) \right) \quad (2.10)$$

for some $\delta \in (0, 1)$, and where $\widetilde{\text{AVaR}}_u(f)$ is the estimator of $\overline{\text{AVaR}}_u(f)$ given by (2.6), which implicitly depends on u through the objective function ℓ . Note that by the lower-semicontinuity of β and the fact that \mathcal{P} is separable, it follows by standard arguments, using e.g. [8, Proposition 7.50], that $\tilde{\rho}^\delta$ is measurable. The following theorem gives a convergence rate for the approximation of the general law-invariant convex risk measure $\bar{\rho}(f)$ by $\tilde{\rho}^\delta(f)$.

Theorem 2.7. Let ρ be a q -regular convex risk measure with $q > 1$. Let f be bounded and satisfy the assumptions of Theorem 2.3. Let $h = \frac{t}{M^2}$. For all $h, \lambda > 0$, $0 < \gamma < 1$, $\delta \in (0, 1)$ and $M, N \in \mathbb{N}^*$, we have

$$\mathbb{E}[|\bar{\rho}(f) - \tilde{\rho}^\delta(f)|^2] \leq C(1 - \delta)^{1/2q} + C\delta^{1/q} \left(\frac{C_{(\delta,t,\lambda)}}{M^2} + \frac{C_{(\delta,t,\lambda)}}{N} + C_{(t,\delta,\lambda,M)}\gamma^2 + C_{(\delta,d,\lambda)}e^{-C_{(d,\lambda)}t} + \frac{C_{(u)}}{\lambda^2} \right),$$

for some constants $C_{(\delta,t,\lambda)}, C_{(\delta,t,\lambda)}, C_{(t,\delta,\lambda,M)}, C_{(\delta,d,\lambda)}, C_{(d,\lambda)}, C_{(u)} > 0$, and a universal constant $C > 0$.

2.3 Numerical results on AVaR

Let us complement the above theoretical guarantees with empirical experiments³. We first focus on the approximation of the average value at risk and the value at risk with respect to the time evolution of the Markov chain in the SGLD algorithm. Thus, for the numerical computations, we set

$$\lambda = 10^8, \quad \gamma = 10^{-8}, \quad h = 10^{-4}, \quad \text{and} \quad u = 0.95.$$

We will consider two cases in our experiments. In the first case we assume the underlying distribution to be known and use Monte–Carlo simulation, and in the second case we use real historical stock price data.

2.3.1 Monte Carlo simulation

For the Monte Carlo experiments, we set $N = 5000$. Figure 1a shows the convergence of AVaR in the 1 dimensional case with $f(r, S) = S$, where S is sampled from Gaussian distributions. Figure 1b shows the estimation error, $\widehat{\text{AVaR}}_u - \overline{\text{AVaR}}_u$, where $\overline{\text{AVaR}}_u$ is the theoretical average value at risk for 1-dimensional Gaussian distributions given by

$$\overline{\text{AVaR}}_u = \mu + \sigma \frac{\phi(\Phi^{-1}(u))}{1 - u}, \quad (2.11)$$

where ϕ and Φ are respectively, the PDF and the CDF of a standard Gaussian distribution. For the multi-dimensional




Figure 1: Numerical Results for 1-d Gaussian random variables

case, we take the function


$$f(r, S) = \sum_{i=1}^d \frac{e^{r_i}}{\sum_{j=1}^d e^{r_j}} S_i, \quad \text{for } i = 1, \dots, d.$$

Figure 2a and Figure 2b show the convergence of VaR and AVaR in the 2-dimensional case, where S^1 is sampled from $\mathcal{N}(1, 4)$ and S^2 is sampled from $\mathcal{N}(0, 1)$.


2.3.2 Numerical results with real data

In this subsection, we compute AVaR for a portfolio of 106 stocks using real aggregated stock prices over 15-minute time intervals from January 2, 2015 to August 31, 2015. Among 128 NASDAQ stocks that are "sufficiently liquid", we remove the ones with missing values, and use the remaining 106 stocks. For a detailed description of the data used and for a definition of "sufficiently liquid", please refer to Section 3.2 of Pohl, Ristig, Schachermayer, and Tangpi [55]. We use changes in stock prices instead of stock prices themselves, because stock prices are highly dependent. We present paths of the estimated optimized VaR and AVaR of the portfolio of 106 stocks in Figures 3a and 3b respectively.

³Code available at https://github.com/jiaruic/sfld_risk_measures




(a) Path of VaR




(b) Path of AVaR

Figure 2: Portfolio of two Gaussian random variables



(a) Paths of VaR



(b) Paths of AVaR

Figure 3: Computations on real price increments of portfolio of 106 stocks.

2.4 Numerical results on general risk measures

In order to simulate general risk measures one needs to specify the penalty function β , or alternatively the precise form of ρ since β is given by [32]

$$\beta(\gamma) = \sup_{X \in \mathcal{A}_\rho} \int_{[0,1]} \text{AVaR}_u(X) \gamma(du), \quad \text{with } \mathcal{A}_\rho := \{X \in \mathbb{L}^\infty : \rho(X + m) \leq 0\}.$$

In general, the simulation of $\tilde{\rho}^\delta(f)$ as given in (2.10) will probably require introducing neural networks since it is the value of an infinite dimensional optimization problem. This will be addressed in future research. We will focus here on a case where the problem can be simplified.

In fact, denote by $\partial_\mu \beta$ the so-called linear functional derivative of β . It is defined as the function $\partial_\mu \beta : \mathcal{M}([0, 1]) \times [0, 1] \rightarrow \mathbb{R}$ such that

$$\beta(\mu') - \beta(\mu) = \int_0^1 \int_0^1 \partial_\mu \beta((1 - \lambda)\mu + \lambda\mu', x) (\mu' - \mu)(dx) d\lambda.$$

Up to an additive constant, there exists a unique such derivative $\partial_\mu \beta$, see e.g. [13]. We have the following:

Proposition 2.8. *Let the assumptions of Theorem 2.7 hold and assume that β admits a second order linear functional*

derivative that is jointly continuous and such that

$$\sup_{\eta, \eta \in \mathbb{L}^2} \mathbb{E} \left[\sup_{\mu \in \mathcal{M}([0,1])} |\partial_\mu \beta(\mu, \eta_1)| + \sup_{\mu \in \mathcal{M}([0,1])} |\partial_\mu^2 \beta(\mu, \eta_1, \eta_2)| \right] < K.$$

Then, it holds

$$\mathbb{E} \left[\left| \inf_{(x_i)_{i=1, \dots, J} \subset [0,1]} F \left(\frac{1}{J} \sum_{i=1}^J \delta_{x_i} \right) - \rho(f) \right|^2 \right] \leq \frac{4K^2}{J^2} + C(1-\delta)^{1/2q} + C_{(\delta, \epsilon)} \frac{1}{M^7} + \frac{C_{(\delta, t, \gamma, \lambda)}}{N} + C_{(t, \delta, \lambda)} \gamma^2 + C_{(\delta, \gamma, \lambda)} e^{-C_{(\gamma, \lambda)} t} + \frac{C_{(\gamma)}}{\lambda^2},$$

with $F(\mu) := \int_0^1 \widetilde{\text{AVaR}}_u(f) \mu(du) - \beta(\mu)$ (recall Equation (2.6)).

This is a direct consequence of Theorem 2.7 and [41, Theorem 2.4]. The proof is omitted.

As an illustrative example, we consider the so-called entropic value-at-risk introduced by Ahmadi-Javid [1] and studied e.g. by Pichler and Schlotter [54] and Föllmer and Knispel [29] in connection to large portfolio asymptotics. This is a risk measure based on the Rényi entropy given by

$$\rho_u(X) = \sup \left\{ \mathbb{E}[ZX] : Z \geq 0, \mathbb{E}[Z] = 1, H_q(Z) \leq \log \frac{1}{1-u} \right\}$$


with $H_q(Z) := \frac{1}{q-1} \log \mathbb{E} Z^q$ for $q \in \mathbb{R}_+^* \setminus \{1\}$. The associated penalty function takes the form

$$\beta(\gamma) := \begin{cases} 0 & \text{if } \int_0^1 \sigma_\gamma(x)^q dx \leq \left(\frac{1}{1-u}\right)^{q-1} \\ +\infty & \text{else} \end{cases}, \quad \text{with } \sigma_\gamma(x) := \int_0^x \frac{1}{1-v} \gamma(dv).$$


To numerically compute entropic value-at-risk, for a large k , we simulate

$$\sup_{(x_i)_{i=1, \dots, J} \subset [0,1]} F \left(\frac{1}{J} \sum_{i=1}^J \delta_{x_j} \right) := \sup_{(x_i)_{i=1, \dots, J} \subset [0,1]} \left(\frac{1}{J} \sum_{i=1}^J \widetilde{\text{AVaR}}_{x_i}(f) - k \left\{ \int_0^1 \left(\frac{1}{J} \sum_{i=1}^N \frac{1}{1-x_i} 1_{x_j \leq x} \right)^q dx - \left(\frac{1}{1-u} \right)^{q-1} \right\}^+ \right).$$

For Monte Carlo simulation, we set $J = 5000$, $k = 10^{18}$, $q = 1.00001$, and estimate the supremum over $(x_i)_{i=1, \dots, J} \subset [0, 1]$ by the maximum of 5000 random partitions, each consisting of J points, of the interval $[0, 1]$. Figure 4a shows the convergence of the entropic value at risk in the 1 dimensional case with $f(r, S) = S$, where S is sampled from $\mathcal{N}(1, 2)$. Figure 4b shows the estimation error compared to the theoretical entropic value-at-risk for $\mathcal{N}(1, 2)$ given by $\rho_u(X) = 1 + \sqrt{-2 \log((1-u)2)}$.



(a) Paths of Entropic Value-at-risk



(b) Estimation Error of Entropic Value-at-risk

Figure 4: Numerical Results Entropic Value-at-risk

3 Rates for the average value at risk

This section is dedicated to the proofs of Theorems 2.3 and 2.4. We will start by some preliminary considerations allowing us to introduce ideas used in the proofs. The details of the proofs will be given in the subsections 3.2 and 3.3.

3.1 Preliminaries

The starting point of our method is to recognize (2.5) as the m -th step of the Euler-Maruyama scheme that discretizes the stochastic differential equation

$$dZ_t^\lambda = -\nabla L(Z_t^\lambda) dt + \sqrt{2\lambda^{-1}} dW_t, \quad Z_0^\lambda = z \in \mathbb{R}^{d+1} \quad (3.1)$$

where W is a $d + 1$ -dimensional Brownian motion. This SDE is the Langevin SDE, with inverse temperature parameter λ . The Langevin SDE is widely studied in physics [60] and for the sampling of Gibbs distribution via Markov chain Monte-Carlo methods [23]. Equipping the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with the \mathbb{P} -completion of the filtration of W , the equation (3.1) admits a unique strong solution. It is well-known that this solution has a unique invariant measure (that we denote by μ_∞^λ) and whose density reads⁴

$$\mu_\infty^\lambda(x) = \frac{e^{-\lambda L(x)}}{\int_{\mathbb{R}^{d+1}} e^{-\lambda L(z)} dz}, \quad (3.2)$$

see e.g. [49, Lemma 2.1]. In this work, the interest of the Langevin equation (aside from its analytical tractability) stems from the fact that the limiting measure μ_∞ of μ_∞^λ as $\lambda \rightarrow \infty$ concentrates on the minimizers of L , which we will show exist. This follows from results of Hwang [42]. Intuitively, this means that if (r^*, m^*) is the minimizer of L , then for $\lambda \rightarrow \infty$

$$\int_{\mathbb{R}^{d+1}} L(z) \mu_\infty^\lambda(dz) \approx L(r^*, m^*). \quad (3.3)$$

Moreover, the Langevin equation allows us to exploit classical techniques in order to derive explicit convergence rates to the invariant measure in the present non-convex potential case.

Remark 3.1. *One interesting byproduct of our method is that, the simulation of AVaR directly allows to compute the value at risk and the optimal portfolios, as well as deriving non-asymptotic rates. Let us illustrate this on the problem of simulation of optimal portfolios r^* in Equation (2.2). As observed above, μ_∞^λ converges to a measure μ supported on the optimal portfolios. Now, let $G : \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ be a strictly convex function such that the gradient ∇G is invertible. Then, by Taylor's expansion we have*

$$G(\widetilde{Z}_{M,h}^{\lambda,n}) - G(m^*, r^*) \geq \nabla G(K)(\widetilde{Z}_{M,h}^{\lambda,n} - (m^*, r^*))$$

for some random variable K , showing that

$$\|\widetilde{Z}_{M,h}^{\lambda,n} - (m^*, r^*)\| \leq \|\nabla G(K)^{-1}\| |G(\widetilde{Z}_{M,h}^{\lambda,n}) - G(m^*, r^*)|.$$

Therefore, provided that the inverse of ∇G does not grow too fast, the argument we give below to prove Theorem 2.3 would allow to derive theoretical guarantees for the optimal portfolio as well, replacing L by G .

3.2 Proof of Theorem 2.3

Throughout this section we assume that the assumptions of Theorem 2.3 are satisfied. We split the proof into several intermediate lemmas. The first one is probably well known, it asserts that the optimization problem defining $\overline{\text{AVaR}}(f)$ admits a solution.

Lemma 3.2. *The function \widetilde{L} defined in Equation (2.3) admits a minimum.*

Proof. By [42, Propositions 2.1 and 2.3], in order to show that \widetilde{L} has a minimum, it is sufficient to check that there exists $\varepsilon > 0$ such that $\{z \in \mathbb{R}^{d+1} : \widetilde{L}(z) \leq \varepsilon\}$ is compact.

Since \widetilde{L} is continuous, the set $A = \{z \in \mathbb{R}^{d+1} : \widetilde{L}(z) \leq \varepsilon\}$ is closed as the pre-image of the closed set $(-\infty, \varepsilon]$. In addition, A is bounded if $\varepsilon < \inf_r \mathbb{E}[f(r, S)]$. In fact, if there exists a sequence $\{z_n\}$ in A such that $\|z_n\| \rightarrow \infty$, then by Assumption 2.2. (iii), it holds $\widetilde{L}(z)/\|z\| \rightarrow \infty$ as $\|z\| \rightarrow \infty$. This contradicts $z_n \in A$ for all n . Thus, the set A is bounded, and is therefore compact. \square

⁴As usual, in this article, we use the same notation for a probability measure on \mathbb{R}^n for any $n \in \mathbb{N}$ and its density function.

To derive the claimed convergence rate, we decompose the expected error into terms that will be handled independently. First, we will exploit the approximation (3.3). Next, using the N independent Brownian motions \widetilde{W}^n introduced just before (2.5), we construct N i.i.d. copies $\widetilde{Z}_t^{\lambda,n}$ of the solution of the Langevin equation as solutions of the SDEs

$$d\widehat{Z}_t^{\lambda,n} = -\nabla L(\widehat{Z}_t^{\lambda,n}) dt + \sqrt{2\lambda^{-1}} d\widetilde{W}^n, \quad n = 1, \dots, N. \quad (3.4)$$

This allows to decompose the error as

$$\begin{aligned} \mathbb{E} \left[\left| \frac{1}{N} \sum_{n=1}^N \ell(\widetilde{Z}'_{M,h}{}^{\lambda,n}) - \overline{\text{AVAR}}_u(f) \right|^2 \right] &\leq 2^5 \mathbb{E} \left[\left| \frac{1}{N} \sum_{n=1}^N \ell(\widetilde{Z}'_{M,h}{}^{\lambda,n}) - \frac{1}{N} \sum_{n=1}^N L(\widetilde{Z}_{M,h}{}^{\lambda,n}) \right|^2 \right] \\ &+ 2^5 \mathbb{E} \left[\left| \frac{1}{N} \sum_{n=1}^N L(\widetilde{Z}_{M,h}{}^{\lambda,n}) - \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda,n}) \right|^2 \right] + 2^5 \mathbb{E} \left[\left| \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda,n}) - \mathbb{E}[L(Z_t^\lambda)] \right|^2 \right] + 2^5 \left| \mathbb{E}[L(Z_t^\lambda)] - \int_{\mathbb{R}^{d+1}} L d\mu_\infty^\lambda \right|^2 \\ &+ 2^5 \left| \int_{\mathbb{R}^{d+1}} L d\mu_\infty^\lambda - \int_{\mathbb{R}^{d+1}} \widetilde{L} d\mu_\infty^\lambda \right|^2 + 2^5 \left| \int_{\mathbb{R}^{d+1}} \widetilde{L} d\mu_\infty^\lambda - \overline{\text{AVAR}}_u(f) \right|^2. \end{aligned} \quad (3.5)$$

The rest of the proof consists in controlling each term above separately.

Lemma 3.3. *Under the conditions of Theorem 2.1, for all $t, \gamma > 0, u \in (0, 1), N, M \in \mathbb{N}^*$ and $\lambda > 1$ if $h < \frac{1}{(\frac{2}{1-u} + \gamma)}$ then we have*

$$\mathbb{E} \left[\left| \frac{1}{N} \sum_{n=1}^N \ell(\widetilde{Z}'_{M,h}{}^{\lambda,n}) - \frac{1}{N} \sum_{n=1}^N L(\widetilde{Z}_{M,h}{}^{\lambda,n}) \right|^2 \right] \leq C_{(M,\lambda,t,u)} \gamma^2 + C_{(u,S)} \frac{1}{N}$$

for some constants $C_{(M,\lambda,t,u)}, C_{(u,S)} > 0$.

Proof. By the definition of L and Jensen's inequality, we have

$$\begin{aligned} &\mathbb{E} \left[\left| \frac{1}{N} \sum_{n=1}^N \ell(\widetilde{Z}'_{M,h}{}^{\lambda,n}) - \frac{1}{N} \sum_{n=1}^N L(\widetilde{Z}_{M,h}{}^{\lambda,n}) \right|^2 \right] \\ &= \mathbb{E} \left[\left| \frac{1}{N} \sum_{n=1}^N \left\{ \widetilde{L}(\widetilde{Z}_{M,h}{}^{\lambda,n}) + \frac{\gamma}{2} \|\widetilde{Z}_{M,h}{}^{\lambda,n}(1)\|^2 - \widetilde{\ell}(\widetilde{Z}'_{M,h}{}^{\lambda,n}) - \frac{\gamma}{2} \|\widetilde{Z}'_{M,h}{}^{\lambda,n}(1)\|^2 \right\} \right|^2 \right] \\ &\leq C \left(\frac{1}{N} \sum_{n=1}^N \mathbb{E} \left[(\widetilde{L}(\widetilde{Z}_{M,h}{}^{\lambda,n}) - \widetilde{\ell}(\widetilde{Z}'_{M,h}{}^{\lambda,n}))^2 \right] + \frac{\gamma^2}{N} \sum_{n=1}^N \mathbb{E} \left[(\|\widetilde{Z}_{M,h}{}^{\lambda,n}(1)\|^2 - \|\widetilde{Z}'_{M,h}{}^{\lambda,n}(1)\|^2)^2 \right] \right) \\ &\leq \frac{C}{N} \sum_{n=1}^N \mathbb{E} \left[(\widetilde{L}(\widetilde{Z}_{M,h}{}^{\lambda,n}) - \widetilde{\ell}(\widetilde{Z}'_{M,h}{}^{\lambda,n}))^2 \right] + C \frac{\gamma^2}{N} \sum_{n=1}^N \left(\mathbb{E} \left[\|\widetilde{Z}_{M,h}{}^{\lambda,n}\|^4 \right] + \mathbb{E} \left[\|\widetilde{Z}'_{M,h}{}^{\lambda,n}\|^4 \right] \right)^{\frac{1}{2}} \mathbb{E} \left[\|\widetilde{Z}_{M,h}{}^{\lambda,n} - \widetilde{Z}'_{M,h}{}^{\lambda,n}\|^4 \right]^{\frac{1}{2}}, \end{aligned} \quad (3.6)$$

with $\widetilde{Z}'_{M,h}{}^{\lambda,n}(1)$ and $\widetilde{Z}_{M,h}{}^{\lambda,n}(1)$ the first components of $\widetilde{Z}'_{M,h}{}^{\lambda,n}$ and $\widetilde{Z}_{M,h}{}^{\lambda,n}$, respectively, and where the last inequality follows by Cauchy-Schwarz. We will show in Lemma 3.9 below that $\|\widetilde{Z}_{M,h}{}^{\lambda,n}\|$ have exponential moments bounded by a constant $C_{M,h,\lambda}$, and similar arguments allow to show that $\|\widetilde{Z}'_{M,h}{}^{\lambda,n}\|^4$ also has exponential moment bounded by the same constant. This shows in particular that

$$\mathbb{E} \left[\|\widetilde{Z}_{M,h}{}^{\lambda,n}\|^4 \right] + \mathbb{E} \left[\|\widetilde{Z}'_{M,h}{}^{\lambda,n}\|^4 \right] \leq 2C_{(M,t,\lambda)}. \quad (3.7)$$

Next, we will bound the fourth moment of the difference between $\widetilde{Z}_{M,h}{}^{\lambda,n}$ and $\widetilde{Z}'_{M,h}{}^{\lambda,n}$. Using the definitions of $\widetilde{Z}_{M,h}{}^{\lambda,n}$ and $\widetilde{Z}'_{M,h}{}^{\lambda,n}$, given in Equation (2.4) and Equation (2.5) respectively, and setting $h = \frac{t}{M^2}$, we have

$$\begin{aligned} \mathbb{E} \left[\|\widetilde{Z}_{M,h}{}^{\lambda,n} - \widetilde{Z}'_{M,h}{}^{\lambda,n}\|^4 \right] &= \mathbb{E} \left[\left\| \widetilde{Z}_{M-1,h}{}^{\lambda,n} - \frac{t}{M^2} \nabla L(\widetilde{Z}_{M-1,h}{}^{\lambda,n}) - \widetilde{Z}'_{M-1,h}{}^{\lambda,n} + \frac{t}{M^2} \nabla \ell(\widetilde{Z}'_{M-1,h}{}^{\lambda,n}) \right\|^4 \right] \\ &\leq C \mathbb{E} \left[\|\widetilde{Z}_{M-1,h}{}^{\lambda,n} - \widetilde{Z}'_{M-1,h}{}^{\lambda,n}\|^4 \right] + C \frac{t^4}{M^8} \mathbb{E} \left[\|\nabla L(\widetilde{Z}_{M-1,h}{}^{\lambda,n}) - \nabla \ell(\widetilde{Z}'_{M-1,h}{}^{\lambda,n})\|^4 \right]. \end{aligned}$$

Using this recursive relationship, it can be checked by induction that we have

$$\mathbb{E} \left[\left\| \tilde{Z}_{M,h}^{\lambda,n} - \tilde{Z}'_{M,h}{}^{\lambda,n} \right\|^4 \right] \leq C \sum_{m=0}^{M-1} \frac{t^4}{M^8} \mathbb{E} \left[\left\| \nabla L(\tilde{Z}_{m,h}^{\lambda,n}) - \nabla \ell(\tilde{Z}'_{m,h}{}^{\lambda,n}) \right\|^4 \right]. \quad (3.8)$$

Let $\tilde{\mathbb{E}}$ be the expectation that acts only on S , but not on $\hat{Z}_r^{\lambda,n}$. Finally, let R_m (resp. R'_m) denote the last d coordinates of $\tilde{Z}_{M,h}^{\lambda,n}$ (resp. $\tilde{Z}'_{M,h}{}^{\lambda,n}$). Define the vector $\mathbf{S} := (-1, S^1, \dots, S^d)$, and let $e_1 \in \mathbb{R}^{d+1}$ be a vector with $1 + \gamma m$ in the first entry and 0 everywhere else. Adding and subtracting $\nabla L(\tilde{Z}'_{M,h}{}^{\lambda,n})$, we get

$$\begin{aligned} & \mathbb{E} \left[\left\| \nabla L(\tilde{Z}_{M,h}^{\lambda,n}) - \nabla \ell(\tilde{Z}'_{M,h}{}^{\lambda,n}) \right\|^4 \right] \\ & \leq C \mathbb{E} \left[\left\| \nabla L(\tilde{Z}_{M,h}^{\lambda,n}) - \nabla L(\tilde{Z}'_{M,h}{}^{\lambda,n}) \right\|^4 \right] + C \mathbb{E} \left[\left\| \nabla L(\tilde{Z}'_{M,h}{}^{\lambda,n}) - \nabla \ell(\tilde{Z}'_{M,h}{}^{\lambda,n}) \right\|^4 \right] \\ & \leq C \left(\frac{1}{1-u} + \gamma \right)^4 \mathbb{E} \left[\left\| \tilde{Z}_{M,h}^{\lambda,n} - \tilde{Z}'_{M,h}{}^{\lambda,n} \right\|^4 \right] \\ & + C \mathbb{E} \left[\left| \left(e_1 + \frac{1}{1-u} \tilde{\mathbb{E}}[\nabla f(R'_m, \mathbf{S}) 1_{\{f(R'_m, \mathbf{S}) \geq \tilde{Z}'_{M,h}{}^{\lambda,n}(1)\}}] \right) - \left(e_1 + \frac{1}{1-u} \frac{1}{N} \sum_{i=1}^N (\nabla f(R'_M, S^i) 1_{\{f(R'_M, S^i) \geq \tilde{Z}'_{M,h}{}^{\lambda,n}(1)\}}) \right) \right|^4 \right] \\ & \leq C \left(\frac{1}{1-u} + \gamma \right)^4 \mathbb{E} \left[\left\| \tilde{Z}_{M,h}^{\lambda,n} - \tilde{Z}'_{M,h}{}^{\lambda,n} \right\|^4 \right] + C \left(\frac{1}{1-u} \right)^4, \end{aligned} \quad (3.9)$$

where the last inequality follows by Assumption 2.2 (iii). Putting equations (3.8), (3.9) together, we have

$$\mathbb{E} \left[\left\| \tilde{Z}_{M,h}^{\lambda,n} - \tilde{Z}'_{M,h}{}^{\lambda,n} \right\|^4 \right] \leq C \frac{t^2}{M^8} \left(\frac{1}{1-u} + \gamma \right)^4 \sum_{m=0}^{M-1} \mathbb{E} \left[\left\| \tilde{Z}_{m,h}^{\lambda,n} - \tilde{Z}'_{m,h}{}^{\lambda,n} \right\|^4 \right] + C \frac{t^4}{M^7} \left(\frac{1}{1-u} \right)^4.$$

Using the discrete version of the Grönwall's inequality [38, Proposition 5], we have

$$\mathbb{E} \left[\left\| \tilde{Z}_{M,h}^{\lambda,n} - \tilde{Z}'_{M,h}{}^{\lambda,n} \right\|^4 \right] \leq \frac{t^4}{M^7} \left(\frac{1}{1-u} \right)^4 \exp \left(\frac{t^4}{M^7} \left(\frac{1}{1-u} + \gamma \right)^4 \right). \quad (3.10)$$

Now, we put equations (3.7) and (3.10) together to have

$$\frac{\gamma^2}{N} \sum_{n=1}^N \left(\mathbb{E} \left[\left\| \tilde{Z}_{M,h}^{\lambda,n} \right\|^4 \right] + \mathbb{E} \left[\left\| \tilde{Z}'_{M,h}{}^{\lambda,n} \right\|^4 \right] \right)^{\frac{1}{2}} \left(\mathbb{E} \left[\left\| \tilde{Z}_{M,h}^{\lambda,n} - \tilde{Z}'_{M,h}{}^{\lambda,n} \right\|^4 \right] \right)^{\frac{1}{2}} \quad (3.11)$$

$$\leq C_{(M,h,\lambda)} \gamma^2 \frac{t^2}{M^{7/2}} \left(\frac{1}{1-u} \right)^2 \exp \left(\frac{t^4}{2M^7} \left(\frac{1}{1-u} + \gamma \right)^4 \right) \leq C_{(M,t,\lambda)} \gamma^2. \quad (3.12)$$

Let us now turn to the first term on the right hand side of (3.6). Adding and subtracting $\tilde{L}(\tilde{Z}'_{M,h}{}^{\lambda,n})$, we have

$$\mathbb{E} \left[\left(\tilde{L}(\tilde{Z}_{M,h}^{\lambda,n}) - \tilde{\ell}(\tilde{Z}'_{M,h}{}^{\lambda,n}) \right)^2 \right] \leq 2 \mathbb{E} \left[\left(\tilde{L}(\tilde{Z}_{M,h}^{\lambda,n}) - \tilde{L}(\tilde{Z}'_{M,h}{}^{\lambda,n}) \right)^2 \right] + 2 \mathbb{E} \left[\left(\tilde{L}(\tilde{Z}'_{M,h}{}^{\lambda,n}) - \tilde{\ell}(\tilde{Z}'_{M,h}{}^{\lambda,n}) \right)^2 \right]. \quad (3.13)$$

Using that \tilde{L} is $\frac{2}{1-u}$ -Lipschitz, it follows that

$$\begin{aligned} \mathbb{E} \left[\left(\tilde{L}(\tilde{Z}_{M,h}^{\lambda,n}) - \tilde{L}(\tilde{Z}'_{M,h}{}^{\lambda,n}) \right)^2 \right] & \leq \frac{4}{(1-u)^2} \mathbb{E} \left[\left\| \tilde{Z}_{M,h}^{\lambda,n} - \tilde{Z}'_{M,h}{}^{\lambda,n} \right\|^2 \right] \\ & \leq \frac{4}{(1-u)^4} \frac{t^2}{M^{7/2}} \exp \left(\frac{t^4}{2M^7} \left(\frac{1}{1-u} + \gamma \right)^4 \right) \end{aligned}$$

where the second inequality follows in view of (3.10). For the second term on the right hand side of (3.13), we use a law

of large number type argument. In fact, we have

$$\begin{aligned}
& \mathbb{E} \left[\left(\widetilde{L}(\widetilde{Z}'_{M,h}{}^{\lambda,n}) - \widetilde{\ell}(\widetilde{Z}'_{M,h}{}^{\lambda,n}) \right)^2 \right] = \mathbb{E} \left[\mathbb{E} \left[\left(\widetilde{L}(\widetilde{Z}'_{M,h}{}^{\lambda,n}) - \widetilde{\ell}(\widetilde{Z}'_{M,h}{}^{\lambda,n}) \right)^2 \middle| \widetilde{Z}'_{M,h}{}^{\lambda,n} \right] \right] \\
& = \mathbb{E} \left[\mathbb{E} \left[\left(\frac{1}{1-u} \widetilde{\mathbb{E}}[(f(R'_M, S) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] - \frac{1}{1-u} \frac{1}{N} \sum_{i=1}^N (f(R'_M, S^i) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ \right)^2 \middle| \widetilde{Z}'_{M,h}{}^{\lambda,n} \right] \right] \\
& = \left(\frac{1}{1-u} \right)^2 \frac{1}{N^2} \mathbb{E} \left[\mathbb{E} \left[\sum_{i,j=1}^N \left((f(R'_M, S^i) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ - \widetilde{\mathbb{E}}[(f(R'_M, S) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] \right) \right. \right. \\
& \quad \left. \left. \cdot \left((f(R'_M, S^j) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ - \widetilde{\mathbb{E}}[(f(R'_M, S) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] \right) \middle| \widetilde{Z}'_{M,h}{}^{\lambda,n} \right] \right] \tag{3.14}
\end{aligned}$$

For $i \neq j$, using that S^i and S^j are independent, we have

$$\begin{aligned}
& \mathbb{E} \left[\mathbb{E} \left[\left((f(R'_M, S^i) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ - \widetilde{\mathbb{E}}[(f(R'_M, S) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] \right) \right. \right. \\
& \quad \left. \left. \cdot \left((f(R'_M, S^j) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ - \widetilde{\mathbb{E}}[(f(R'_M, S) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] \right) \middle| \widetilde{Z}'_{M,h}{}^{\lambda,n} \right] \right] \\
& = \mathbb{E} \left[\mathbb{E} \left[(f(R'_M, S^i) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ - \widetilde{\mathbb{E}}[(f(R'_M, S) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] \middle| \widetilde{Z}'_{M,h}{}^{\lambda,n} \right] \right] \\
& \quad \cdot \mathbb{E} \left[\mathbb{E} \left[(f(R'_M, S^j) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ - \widetilde{\mathbb{E}}[(f(R'_M, S) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] \middle| \widetilde{Z}'_{M,h}{}^{\lambda,n} \right] \right] = 0.
\end{aligned}$$

Therefore, we can estimate the desired term as

$$\begin{aligned}
& \mathbb{E} \left[\left(\widetilde{L}(\widetilde{Z}'_{M,h}{}^{\lambda,n}) - \widetilde{\ell}(\widetilde{Z}'_{M,h}{}^{\lambda,n}) \right)^2 \right] \tag{3.15} \\
& = \left(\frac{1}{1-u} \right)^2 \frac{1}{N^2} \sum_{i=1}^N \mathbb{E} \left[\mathbb{E} \left[\left((f(R'_M, S^i) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ - \widetilde{\mathbb{E}}[(f(R'_M, S) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] \right)^2 \middle| \widetilde{Z}'_{M,h}{}^{\lambda,n} \right] \right] \\
& = \left(\frac{1}{1-u} \right)^2 \frac{1}{N^2} \sum_{i=1}^N \mathbb{E} \left[\mathbb{E} \left[\left((f(R'_M, S^i) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ - \mathbb{E}[(f(R'_M, S) - \widetilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ | \widetilde{Z}'_{M,h}{}^{\lambda,n}] \right)^2 \middle| \widetilde{Z}'_{M,h}{}^{\lambda,n} \right] \right] \\
& = \left(\frac{1}{1-u} \right)^2 \frac{1}{N} \mathbb{E} \left[\text{Var} \left(f(R'_M, S) \middle| \widetilde{Z}'_{M,h}{}^{\lambda,n} \right) \right] \\
& = \left(\frac{1}{1-u} \right)^2 \frac{1}{N} \text{Var}(f(R'_M, S)) \leq \left(\frac{1}{1-u} \right)^2 \frac{1}{N} \text{Var}(C(R_M + S^2)) \leq \left(\frac{1}{1-u} \right)^2 \frac{C_{(u,S)}}{N}, \tag{3.16}
\end{aligned}$$

where the third to last step follows by Jensen's inequality and tower property. The penultimate step follows by the boundedness of $\nabla_r f(r, S)$ and Lipschitzness of $\nabla_s f(r, S)$. In the last step, we used that S has finite fourth moment, and equation (3.7). Finally, putting (3.6), (3.12) and (3.16) together yields the lemma. \square

We now investigate the second term on the right hand side of (3.5).

Lemma 3.4. *Under the conditions of Theorem 2.3, for all $t, \gamma > 0, u \in (0, 1), M, N \in \mathbb{N}^*$ and $\lambda > 1$, if $h < \frac{1}{(\frac{2}{1-u} + \gamma)}$ we have*

$$\mathbb{E} \left[\left| \frac{1}{N} \sum_{n=1}^N L(\widetilde{Z}'_{M,h}{}^{\lambda,n}) - \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}'_t{}^{\lambda,n}) \right|^2 \right] \leq \frac{C_{(u,t)}}{M^4} + C_{(M,t,\lambda)} \gamma^2,$$

for $C, C_{(u,t)}, C_{(M,t,\lambda)} > 0$.

Proof. Following the same argument as in the proof of Lemma 3.2, we have

$$\mathbb{E} \left[\left| \frac{1}{N} \sum_{n=1}^N L(\tilde{Z}_{M,h}^{\lambda,n}) - \frac{1}{N} \sum_{n=1}^N L(\hat{Z}_t^{\lambda,n}) \right|^2 \right] \leq C \left(\frac{1}{N} \sum_{n=1}^N \mathbb{E} \left[\left(\tilde{L}(\tilde{Z}_{M,h}^{\lambda,n}) - \tilde{L}(\hat{Z}_t^{\lambda,n}) \right)^2 \right] + \frac{\gamma^2}{N} \sum_{n=1}^N \mathbb{E} \left[\left(\|\tilde{Z}_{M,h}^{\lambda,n}\|^2 - \|\hat{Z}_t^{\lambda,n}\|^2 \right)^2 \right] \right). \quad (3.17)$$

Let $Z^\lambda(s)$ be a continuous time approximation of the Euler-Maruyama scheme in (2.5). One way to define such an approximation is by setting

$$Z^\lambda(s) := z - \int_0^{n_s} \nabla L(Z^\lambda(r)) dr + \int_0^{n_s} \sqrt{2\lambda^{-1}} dW_r,$$

with $n_s := \max\{\frac{t}{M^2}n : \frac{t}{M^2}n \leq s, n \in \mathbb{Z}\}$. Note that for each $1 \leq i \leq N$, $1 \leq m \leq M$, and $t > 0$, we have $Z^\lambda(s_m) = \tilde{Z}_{m,h}^{\lambda,i}$. In other words, $Z^\lambda(s)$ coincides with $\tilde{Z}_{m,h}^{\lambda,i}$ at the time discretization points. For the first term in (3.17), we have

$$\begin{aligned} \frac{1}{N} \sum_{n=1}^N \mathbb{E} \left[\left| \tilde{L}(\tilde{Z}_{M,h}^{\lambda,n}) - \tilde{L}(\hat{Z}_t^{\lambda,n}) \right|^2 \right] &\leq \frac{4}{(1-u)^2 N} \sum_{n=1}^N \mathbb{E} \left[\|\tilde{Z}_{M,\frac{t}{M^2}}^{\lambda,n} - \hat{Z}_t^{\lambda,n}\|^2 \right] \\ &\leq \frac{4}{(1-u)^2 N} \sum_{n=1}^N \mathbb{E} \left[\sup_{0 \leq s \leq t} \|Z^\lambda(s) - \hat{Z}_s^{\lambda,n}\|^2 \right], \end{aligned}$$

where we used that \tilde{L} is $\frac{2}{1-u}$ -Lipschitz in the first inequality. By standard results on the error estimation for SDE approximations, see e.g. [45, Theorems 10.3.5 and 10.6.3], we have

$$\mathbb{E} \left[\sup_{0 \leq s \leq t} |Z^\lambda(s) - \hat{Z}_s^{\lambda,n}|^4 \right] \leq C \frac{t^4}{M^8} \quad (3.18)$$

for some constant $C > 0$. Thus, we have

$$\frac{1}{N} \sum_{n=1}^N \mathbb{E} \left[\left| \tilde{L}(\tilde{Z}_{M,h}^{\lambda,n}) - \tilde{L}(\hat{Z}_t^{\lambda,n}) \right|^2 \right] \leq \frac{Ct^2}{(1-u)^2 M^4}. \quad (3.19)$$

For the second term on the right hand side of (3.17), by Cauchy-Schwartz inequality, we have

$$\begin{aligned} \mathbb{E} \left[\left| (\tilde{Z}_{M,h}^{\lambda,n})^2 - (\hat{Z}_t^{\lambda,n})^2 \right|^2 \right] &\leq \mathbb{E} \left[\|\tilde{Z}_{M,h}^{\lambda,n}\|^4 + \|\hat{Z}_t^{\lambda,n}\|^4 \right]^{\frac{1}{2}} \mathbb{E} \left[\|\tilde{Z}_{M,h}^{\lambda,n} - \hat{Z}_t^{\lambda,n}\|^4 \right]^{\frac{1}{2}} \\ &\leq \mathbb{E} \left[\|\tilde{Z}_{M,h}^{\lambda,n}\|^4 + \|\hat{Z}_t^{\lambda,n}\|^4 \right]^{\frac{1}{2}} \mathbb{E} \left[\sup_{0 \leq s \leq t} \|Z^\lambda(s) - \hat{Z}_s^{\lambda,n}\|^4 \right]^{\frac{1}{2}} \\ &\leq C \mathbb{E} \left[\|\tilde{Z}_{M,h}^{\lambda,n}\|^4 + \|\hat{Z}_t^{\lambda,n}\|^4 \right]^{\frac{1}{2}} \frac{t^2}{M^4} \end{aligned} \quad (3.20)$$

where we used equation (3.18) in the last step.

It remains to control the fourth moment of $\hat{Z}_t^{\lambda,n}$, since that of $\tilde{Z}_{M,h}^{\lambda,n}$ was bounded by a constant $C_{M,h,\lambda}$ in Lemma 3.3 using the exponential moment given in Lemma 3.9. Since $\hat{Z}_t^{\lambda,n}$ solves the SDE

$$d\hat{Z}_t^{\lambda,n} = -\nabla L(\hat{Z}_t^{\lambda,n}) dt + \sqrt{2\lambda^{-1}} d\tilde{W}^n,$$

with a linearly growing drift, the following bound on the fourth moment of the solution follows by standard arguments:

$$\mathbb{E}[\|\hat{Z}_t^{\lambda,n}\|^4] \leq C \left(1 + \frac{t}{(1-u)^4} + \frac{1}{\lambda^2} t^2 \right) e^{\gamma^4 C t} \leq C_{(\lambda,t)} e^{C\gamma^4 t}. \quad (3.21)$$

We omit the proof. Putting equations (3.19), (3.20), (3.21) and (3.7) together yields the lemma. \square

Now we move on to analyzing the third term in (3.5).

Lemma 3.5. *Under the conditions of Theorem 2.3, for all $h, t > 0, u \in (0, 1), N \in \mathbb{N}^*, 0 < \gamma < 1$, and $\lambda > 1$, we have*

$$\mathbb{E} \left[\left| \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda, n}) - \mathbb{E}[L(Z_t^\lambda)] \right|^2 \right] \leq \frac{C_{(\lambda, u, t)}}{N}$$

for $C_{(\lambda, t)} > 0$.

Proof. Since $(\widehat{Z}_t^{\lambda, n})_{n \geq 1}$ are i.i.d. copies of Z_t^λ , a standard law of large numbers argument gives

$$\mathbb{E} \left[\left| \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda, n}) - \mathbb{E}[L(Z_t^\lambda)] \right|^2 \right] \leq \frac{1}{N} \text{Var}(L(Z_t^\lambda))$$

where $\text{Var}(L(Z_t^\lambda))$ is the variance of $L(Z_t^\lambda)$. Since ∇L is $(\frac{1}{1-u} + \gamma)$ -Lipschitz, it follows by [22, Corollary 5.11], that the law of Z_t^λ satisfies the Poincaré inequality. That is,

$$\text{Var}(L(Z_t^\lambda)) \leq \frac{2}{\lambda(\frac{1}{1-u} + \gamma)^2} \mathbb{E} \left[\|\nabla L(Z_t^\lambda)\|^2 \right].$$

Now using that $\nabla \tilde{L}$ is $\frac{C}{1-u}$ -Lipschitz, we have $\|\nabla L(x)\| \leq C + (\frac{C}{1-u} + \gamma)\|x\|$ for some constant $C > 0$. Thus, it follows that

$$\begin{aligned} \text{Var}(L(Z_t^\lambda)) &\leq \frac{C}{\lambda(\frac{C}{1-u} + \gamma)^2} \mathbb{E} \left[\left| C + \left(\frac{C}{1-u} + \gamma \right) \|Z_t^\lambda\| \right|^2 \right] \\ &\leq C \left(\frac{1}{\lambda(\frac{1}{1-u} + \gamma)^2} + \frac{1}{\lambda} (\mathbb{E}[\|Z_t^\lambda\|^4])^{\frac{1}{2}} \right) \\ &\leq C \left(\frac{1}{\lambda(\frac{1}{1-u} + \gamma)^2} + \frac{1}{\lambda} \left(1 + \frac{\sqrt{t}}{(1-u)^2} + \frac{t}{\lambda} \right) e^{\gamma^4 C t} \right), \end{aligned}$$

where the second to last inequality follows from (3.21). Taking $0 < \gamma < 1$ yields the result of the lemma. \square

In the next lemma we analyze the fourth term in (3.5). Essentially, this concerns the rate of convergence of the law μ_t^λ of the solution of the Langevin equation to its invariant measure μ_∞^λ .

Lemma 3.6. *For all $t > 0, u, \gamma \in (0, 1), \lambda > 1$ and an initial position $Z_0^\lambda = z \in \mathbb{R}^{d+1}$, we have*

$$\left| \mathbb{E}L(Z_t^\lambda) - \int L(x) d\mu_\infty^\lambda \right|^2 \leq C_{(d, \lambda)} \frac{1}{(1-u)^2} e^{-2tC_{(d, \lambda)}} \mathcal{W}_1^2(\delta_z, \mu_\infty^\lambda) + \gamma^2 C_{(\lambda, t)}.$$

Proof. The investigation of convergence rates to the invariance measure is an active research area, see e.g. [10]. In the present case with non-convex potential functions ∇L , this follows from the so-called coupling by reflection arguments of Eberle [24]. In fact, by Assumption 2.2. (iv), it follows from [24, Corollary 2.1], (see also [25, Corollary 2]) that there is a constant $C_{(d, \lambda)} > 0$ depending only on d, γ and λ such that

$$\mathcal{W}_1(\mu_t^\lambda, \mu_\infty^\lambda) \leq C_{(d, \lambda)} e^{-tC_{(d, \lambda)}} \mathcal{W}_1(\delta_z, \mu_\infty^\lambda) \quad \text{for all } t > 0. \quad (3.22)$$

It now remains to bound $|\mathbb{E}L(Z_t^\lambda) - \int L(x) d\mu_\infty^\lambda|$ by $\mathcal{W}_1(\mu_t^\lambda, \mu_\infty^\lambda)$. Let $\hat{\alpha} \in \Gamma(\mu_\infty^\lambda, \mu_t^\lambda)$ be an optimal coupling of μ_∞^λ and μ_t^λ , i.e. such that

$$\mathbb{E}_{\hat{\alpha}} \|X - Y\| = \inf_{\alpha \in \Gamma(\mu_\infty^\lambda, \mu_t^\lambda)} \mathbb{E}_\alpha \|X - Y\|,$$

see e.g. [63] for existence of $\hat{\alpha}$. Above, we denote by $\mathbb{E}_\alpha[\|X - Y\|]$ the expectation of when $(X, Y) \sim \alpha$. By definition of L and Lipschitz-continuity of \tilde{L} , it holds

$$|L(x) - L(y)| \leq \frac{2}{1-u} \|x - y\| + \frac{\gamma}{2} (\|x\|^2 + \|y\|^2).$$

Let \mathbb{E}_∞ and \mathbb{E}_t be the expectations with respect to μ_∞^λ and μ_t^λ respectively. Taking the expectation with respect to $\hat{\alpha} \in \Gamma(\mu_\infty^\lambda, \mu_t^\lambda)$ we have

$$\begin{aligned} \mathbb{E}_{\hat{\alpha}} \left[|L(X) - L(Y)| \right] &\leq \frac{2}{1-u} \mathbb{E}_{\hat{\alpha}} [\|X - Y\|] + \frac{\gamma}{2} \mathbb{E}_{\hat{\alpha}} [\|X\|^2 + \|Y\|^2] \\ &\leq \frac{2}{1-u} \mathcal{W}_1(\mu_\infty^\lambda, \mu_t^\lambda) + \frac{\gamma}{2} \left(\mathbb{E}_\infty[\|X\|^2] + \mathbb{E}[\|Z_t^\lambda\|^2] \right) \end{aligned} \quad (3.23)$$

As in the proof of Lemma 3.4, see e.g. Equation (3.21), Z_t^λ has second moment bounded by a constant $C_{(\lambda,t)} > 0$. Concerning the term $\mathbb{E}_\infty[\|X\|^2]$, note that it holds

$$\mathbb{E}_\infty[\|X\|^2] = \int \|x\|^2 d\mu_\infty^\lambda = \int \|x\|^2 \frac{e^{-\lambda L(x)}}{\int e^{-\lambda L(a)} da} dx.$$

Since \tilde{L} is $\frac{2}{1-u}$ -Lipschitz, $|\tilde{L}(x)| \leq C + \frac{2}{1-u}\|x\|$ for some constant C . We thus have

$$\int \|x\|^2 e^{-\lambda L(x)} dx = \int \|x\|^2 e^{-\lambda(\tilde{L}(x) + \gamma\|x\|^2/2)} dx \leq \int \|x\|^2 e^{C\lambda + 2\lambda\|x\|/(1-u) - \|x\|^2/2} dx < C_{(\lambda)} \quad (3.24)$$

for a constant $C_{(\lambda)} > 0$ depending only on λ . Using the same argument, $0 < \int e^{-\lambda L(a)} da < \infty$. Therefore, we have $\mathbb{E}_\infty[\|X\|^2] < \infty$. Combining this with (3.22) and (3.23) yields the desired result. \square

Remark 3.7. If the function f is convex, it follows that L is strongly convex in the sense that $\nabla^2 L \geq \gamma I_{d+1}$, where I_{d+1} is the $(d+1) \times (d+1)$ identity matrix. In this case, the exponential convergence to equilibrium follows by standard arguments, see e.g. [10]. In fact, we have the following bound is second order Wasserstein distance:

$$\mathcal{W}_2^2(\mu_t^\lambda, \mu_\infty^\lambda) \leq e^{-2\gamma t} \mathcal{W}_2^2(\delta_z, \mu_\infty^\lambda). \quad (3.25)$$

In this case, a slight modification of the above arguments allow to get the bound

$$\left| \mathbb{E}L(Z_t^\lambda) - \int L(x) d\mu_\infty^\lambda \right|^2 \leq C \left(\frac{1}{(1-u)^2} + 1 \right) e^{-\gamma t} \mathcal{W}_2^2(\delta_z, \mu_\infty^\lambda).$$

The estimation of the fifth term in the decomposition (3.5) is an immediate consequence of the existence of second moment of the invariant measure μ_∞^λ obtained in the proof of the preceding lemma. In fact, by definition of L and \tilde{L} we have

$$\left| \int L(x) d\mu_\infty^\lambda - \int \tilde{L}(x) d\mu_\infty^\lambda \right|^2 = \left(\frac{\gamma}{2} \right)^2 \int \|x\|^2 d\mu_\infty^\lambda \leq C\gamma^2, \quad (3.26)$$

for a constant $C > 0$. We conclude the proof of the theorem with the following lemma estimating the last term on the right hand side of (3.5).

Lemma 3.8. Under the conditions of Theorem 2.3, we have

$$\left| \int \tilde{L}(x) d\mu_\infty^\lambda - \overline{\text{AVaR}}(f) \right|^2 \leq C_{(u)} \frac{\gamma^2}{(1+\gamma)^2} + \frac{C_{(u)}}{\lambda^2}. \quad (3.27)$$

Proof. First recall, see Equation (2.2) and Lemma 3.2 that $\overline{\text{AVaR}}(f) = \tilde{L}(z^*)$ where $z^* = (r^*, m^*)$ is the optimizer in (2.2). Now, consider the differential entropy

$$\begin{aligned} - \int \mu_\infty^\lambda \log \mu_\infty^\lambda dx &= - \int \frac{e^{-\lambda L(x)}}{\int e^{-\lambda L(u)} du} \log \frac{e^{-\lambda L(x)}}{\int e^{-\lambda L(u)} du} dx \\ &= - \int \frac{e^{-\lambda L(x)}}{\int e^{-\lambda L(u)} du} \left(-\lambda L(x) - \log \int e^{-\lambda L(u)} du \right) dx \\ &= \lambda \int L(x) d\mu_\infty^\lambda + \log \int e^{-\lambda L(x)} dx \\ &= \lambda \int \left(\tilde{L}(x) + \frac{\gamma}{2} \|x\|^2 \right) d\mu_\infty^\lambda + \log \int e^{-\lambda \tilde{L}(x) - \frac{\lambda\gamma}{2} \|x\|^2} dx. \end{aligned}$$

Rearranging the terms gives the following expression for the integral of \tilde{L} :

$$\int \tilde{L}(x) d\mu_\infty^\lambda = -\frac{\gamma}{2} \int \|x\|^2 d\mu_\infty^\lambda - \frac{1}{\lambda} \int \mu_\infty^\lambda \log \mu_\infty^\lambda dx - \frac{1}{\lambda} \log \int e^{-\lambda \tilde{L}(x) - \frac{\lambda\gamma}{2} \|x\|^2} dx. \quad (3.28)$$

Since for any continuous random variable, a Gaussian distribution with the same second moment maximizes the differential entropy ([66, Theorem 10.48]), it holds

$$-\int \mu_\infty^\lambda \log \mu_\infty^\lambda dx \leq \frac{1}{2} \log \left((2\pi e)^{d+1} \int \|x\|^2 d\mu_\infty^\lambda \right), \quad (3.29)$$

and using that $\int \|x\|^2 d\mu_\infty^\lambda < \infty$ (see equation (3.24)) and subtracting $\tilde{L}(z^*)$ from both sides of (3.28), we have

$$\begin{aligned} \int \tilde{L}(x) d\mu_\infty^\lambda - \tilde{L}(z^*) &\leq C \left(-\gamma + \frac{1}{\lambda} - \frac{1}{\lambda} \log \int e^{-\lambda \tilde{L}(x) - \frac{\lambda\gamma}{2} \|x\|^2} dx - \tilde{L}(z^*) \right) \\ &\leq C \left(\gamma + \frac{1}{\lambda} - \frac{1}{\lambda} \log \left(e^{-\lambda \tilde{L}(z^*)} \int e^{-\lambda(\tilde{L}(x) - \tilde{L}(z^*)) - \frac{\lambda\gamma}{2} \|x\|^2} dx \right) - \tilde{L}(z^*) \right) \\ &= C \left(\gamma + \frac{1}{\lambda} - \frac{1}{\lambda} \log \int e^{-\lambda(\tilde{L}(x) - \tilde{L}(z^*)) - \frac{\lambda\gamma}{2} \|x\|^2} dx \right). \end{aligned}$$

Hence, using the fact that $\nabla \tilde{L}$ is $\frac{1}{1-u}$ -Lipschitz, we can estimate the exponent in the integral above as

$$|\tilde{L}(x) - \tilde{L}(z^*) - \nabla \tilde{L}(z^*) \cdot (x - z^*)| \leq \frac{1}{2(1-u)} \|x - z^*\|^2.$$

Thanks to the above inequality, using $\nabla \tilde{L}(z^*) = 0$ we obtain

$$\begin{aligned} \int \tilde{L}(x) d\mu_\infty^\lambda - L(z^*) &\leq C \left(\gamma + \frac{1}{\lambda} - \frac{1}{\lambda} \log \int e^{-\frac{\lambda}{2(1-u)} \|x - z^*\|^2 - \frac{\lambda\gamma}{2} \|x\|^2} dx \right) \\ &= C \left(\gamma + \frac{1}{\lambda} - \frac{1}{\lambda} \log \left(\sqrt{\frac{2\pi}{\lambda(\gamma + \frac{1}{1-u})}} e^{-\frac{\gamma\lambda \|z^*\|^2}{2(1-u)(\frac{1}{1-u} + \gamma)}} \right) \right) \\ &\leq C \left(\gamma + \frac{1}{\lambda} - \frac{1}{\lambda} \log \left(\sqrt{\frac{2\pi}{\lambda(\gamma + \frac{1}{1-u})}} \right) + \frac{\gamma}{(1-u)(\frac{1}{1-u} + \gamma)} \right). \end{aligned}$$

For the other side of the inequality, since z^* is a minimizer of \tilde{L} , we have

$$\tilde{L}(x^*) - \int \tilde{L}(x) d\mu_\infty^\lambda \leq \tilde{L}(z^*) - \tilde{L}(z^*) \int d\mu_\infty^\lambda = 0.$$

Using $0 < \gamma < 1$ concludes the proof. \square

3.3 Proof of Theorem 2.4

In this section we present the proof of concentration inequality given in Theorem 2.4. Due to the identification of the SGLD scheme as the discretization of a Langevin equation, the main argument for the proof is based on concentration inequalities for the Euler-Maruyama scheme of stochastic differential equations, see e.g. Menozzi and Lemaire [51], Frikha and Menozzi. [33], Malrieu and Talay [50] and the references therein for results in this direction. Moreover, since the objective function L is not Lipschitz-continuous (it is quadratic) we further need to prove an integrability property (see Lemma 3.9) which makes up the bulk of the proof.

Proof of Theorem 2.4. We start by decomposing the deviation by four terms that will be dealt with independently: Let z^* be an optimizer. Recall that $\text{AVaR}(f) = L(z^*)$. Thus, we have

$$\begin{aligned} \mathbb{P} \left(\left| \frac{1}{N} \sum_{n=1}^N \ell \left(\tilde{Z}_{M,h}^{\lambda,n} \right) - \tilde{L}(x^*) \right| \geq \epsilon \right) &\leq \mathbb{P} \left(\left| \frac{1}{N} \sum_{n=1}^N \ell \left(\tilde{Z}'_{M,h} \right) - \frac{1}{N} \sum_{n=1}^N L \left(\tilde{Z}_{M,h}^{\lambda,n} \right) \right| \geq \frac{\epsilon}{4} \right) \\ &+ \mathbb{P} \left(\left| \frac{1}{N} \sum_{n=1}^N L \left(\tilde{Z}_{M,h}^{\lambda,n} \right) - \mathbb{E} \left[L \left(\tilde{Z}_{M,h}^{\lambda,n} \right) \right] \right| \geq \frac{\epsilon}{4} \right) + \mathbb{P} \left(\left| \mathbb{E} \left[L \left(\tilde{Z}_t^\lambda \right) \right] - \tilde{L}(x^*) \right| \geq \frac{\epsilon}{4} \right). \end{aligned} \quad (3.30)$$

By lemmas 3.6, 3.8 and Equation (3.26), the last term is zero for λ and t large enough and γ small enough. Similarly, the third term is zero for h small enough. Thus we can focus on the first two terms.

For the first term, by Jensen's inequality we have

$$\begin{aligned} \mathbb{E} \left[\exp \left(\left| \frac{1}{N} \sum_{n=1}^N \ell \left(\widetilde{Z}'_{M,h}{}^{\lambda,n} \right) - \frac{1}{N} \sum_{n=1}^N L \left(\widetilde{Z}_{M,h}{}^{\lambda,n} \right) \right|^2 \right) \right] &\leq \mathbb{E} \left[\frac{1}{N} \sum_{n=1}^N \exp \left(\left| \ell \left(\widetilde{Z}'_{M,h}{}^{\lambda,n} \right) - L \left(\widetilde{Z}_{M,h}{}^{\lambda,n} \right) \right|^2 \right) \right] \\ &\leq \mathbb{E} \left[\exp \left(4 \left(\ell \left(\widetilde{Z}'_{M,h}{}^{\lambda,n} \right) \right)^2 \right) \right]^{1/2} \mathbb{E} \left[\exp \left(4 \left(L \left(\widetilde{Z}_{M,h}{}^{\lambda,n} \right) \right)^2 \right) \right]^{1/2}, \end{aligned}$$

where the last inequality follows by the Cauchy-Schwarz inequality. Since \widetilde{L} is $\frac{2}{1-u}$ -Lipschitz, and we have $L(x) = \widetilde{L}(x) + \frac{\gamma}{2}x^2$, we can continue the estimation as

$$\begin{aligned} &\mathbb{E} \left[\exp \left(\left| \frac{1}{N} \sum_{n=1}^N \ell \left(\widetilde{Z}'_{M,h}{}^{\lambda,n} \right) - \frac{1}{N} \sum_{n=1}^N L \left(\widetilde{Z}_{M,h}{}^{\lambda,n} \right) \right|^2 \right) \right] \\ &\leq \left(\mathbb{E} \left[\exp \left(C \left(\frac{1}{1-u} \vee \gamma \right) \left\| \widetilde{Z}'_{M,h}{}^{\lambda,n} \right\|^4 \right) \right] \right)^{1/2} \left(\mathbb{E} \left[\exp \left(C \left(\frac{1}{1-u} \vee \gamma \right) \left\| \widetilde{Z}_{M,h}{}^{\lambda,n} \right\|^4 \right) \right] \right)^{1/2}. \end{aligned}$$

Then it follows by Lemma 3.9 below that

$$\mathbb{E} \left[\exp \left(\left| \frac{1}{N} \sum_{n=1}^N \ell \left(\widetilde{Z}'_{M,h}{}^{\lambda,n} \right) - \frac{1}{N} \sum_{n=1}^N L \left(\widetilde{Z}_{M,h}{}^{\lambda,n} \right) \right| \right) \right] \leq C_{\lambda,\gamma,t}.$$

Thus, by Markov's inequality, we have

$$\mathbb{P} \left(\left| \frac{1}{N} \sum_{n=1}^N \ell \left(\widetilde{Z}'_{M,h}{}^{\lambda,n} \right) - \frac{1}{N} \sum_{n=1}^N L \left(\widetilde{Z}_{M,h}{}^{\lambda,n} \right) \right| \geq \frac{\varepsilon}{4} \right) \leq C_{\lambda,\gamma,t} e^{-\varepsilon^2/16}.$$

Regarding the second term in (3.30), using the definition of L , we have

$$\begin{aligned} \mathbb{P} \left(\left| \frac{1}{N} \sum_{n=1}^N L \left(\widetilde{Z}_{M,h}{}^{\lambda,n} \right) - \mathbb{E} \left[L \left(\widetilde{Z}_{M,h}{}^{\lambda,n} \right) \right] \right| \geq \frac{\varepsilon}{4} \right) &\leq \mathbb{P} \left(\left| \frac{1}{N} \sum_{n=1}^N \widetilde{L} \left(\widetilde{Z}_{M,h}{}^{\lambda,n} \right) - \mathbb{E} \left[\widetilde{L} \left(\widetilde{Z}_{M,h}{}^{\lambda,n} \right) \right] \right| \geq \frac{\varepsilon}{8} \right) \\ &\quad + \mathbb{P} \left(\left| \frac{1}{N} \sum_{n=1}^N \frac{\gamma}{2} \left\| \widetilde{Z}_{M,h}{}^{\lambda,n} \right\|^2 - \mathbb{E} \left[\frac{\gamma}{2} \left\| \widetilde{Z}_{M,h}{}^{\lambda,n} \right\|^2 \right] \right| \geq \frac{\varepsilon}{8} \right). \end{aligned} \quad (3.31)$$

Since \widetilde{L} is Lipschitz-continuous, the first term on the right hand side above is estimated using the concentration inequality of [33, Theorem 2.1]. Note that assumption $GC(\alpha)$ therein is satisfied with $\alpha = 2$, and assumption (A) therein is satisfied since the drift coefficient ∇L is uniformly Lipschitz, and the diffusion coefficient $\sqrt{2\lambda^{-1}}$ is a constant. Thus, we have

$$\mathbb{P} \left(\left| \frac{1}{N} \sum_{n=1}^N \widetilde{L} \left(\widetilde{Z}_{M,h}{}^{\lambda,n} \right) - \mathbb{E} \left[\widetilde{L} \left(\widetilde{Z}_{M,h}{}^{\lambda,n} \right) \right] \right| \geq \frac{\varepsilon}{8} \right) \leq 2 \exp \left(- \frac{\varepsilon^2 N}{\Psi(M, h, u)} \right),$$

where the constant $\Psi(M, h, u)$ is given in equation (2.8). The estimation of the second term on the right hand side of (3.31) follows again by exponential integrability and Markov's inequality:

$$\begin{aligned} \mathbb{P} \left(\left| \frac{1}{N} \sum_{n=1}^N \frac{\gamma}{2} \left\| \widetilde{Z}_{M,h}{}^{\lambda,n} \right\|^2 - \mathbb{E} \left[\frac{\gamma}{2} \left\| \widetilde{Z}_{M,h}{}^{\lambda,n} \right\|^2 \right] \right| \geq \frac{\varepsilon}{8} \right) &\leq e^{-\varepsilon^2/64} \mathbb{E} \left[\exp \left(\left| \frac{1}{N} \sum_{n=1}^N \gamma \left\| \widetilde{Z}_{M,h}{}^{\lambda,n} \right\|^2 - \mathbb{E} \left[\gamma \left\| \widetilde{Z}_{M,h}{}^{\lambda,n} \right\|^2 \right] \right| \right) \right] \\ &\leq e^{-\varepsilon^4/64} \mathbb{E} \left[\exp(\gamma \left\| \widetilde{Z}_{M,h}{}^{\lambda,n} \right\|^2) \right] \\ &\leq C_{m,h,\lambda} e^{-\varepsilon^2/64}, \end{aligned}$$

where the second inequality follows by Jensen's inequality and the fact that $(\widetilde{Z}_{M,h}{}^{\lambda,n})_n$ are i.i.d., and the last inequality follows by Lemma 3.9 given below. \square

We conclude this section with the following exponential moment property used in the previous proof.

Lemma 3.9. *For any $m \in \{1, \dots, M\}$, $t > 0$, $0 < \gamma < 1$, and $\lambda > 0$ with $h < \frac{1}{(\frac{2}{1-u} + \gamma)}$, we have $\mathbb{E} \left[\exp(C \|\tilde{Z}_{M,h}^{\lambda,n}\|^4) \right] \leq C_{(M,h,\lambda,u)}$ for some constant $C_{(M,h,\lambda,u)} > 0$ depending only on M, h, λ and u .*

Proof. We will first compute $E \left[\|\tilde{Z}_{M,h}^{\lambda,n}\|^{4k} \right]$, for any $k \in \mathbb{N}$ and then use the series expansion $e^{x^4} = \sum_{k=0}^{\infty} \frac{x^{4k}}{k!}$ to conclude. Since ∇L is $\frac{2}{1-u}$ -Lipschitz, we have $\nabla L(x) = \nabla \tilde{L}(x) + \gamma x \leq \frac{2}{1-u} + \gamma x \leq C_{u,\gamma}(1+x)$, with $C_{u,\gamma} = (\frac{2}{1-u} + \gamma)$. Using the definition of $\tilde{Z}_{M,h}^{\lambda,n}$, we have

$$\begin{aligned} \|\tilde{Z}_{m+1,h}^{\lambda,n}\|^{4k} &= \|\tilde{Z}_{M,h}^{\lambda,n} - \nabla L(\tilde{Z}_{M,h}^{\lambda,n}) \frac{t}{M^2} + \Delta \tilde{W}_m \sqrt{2\lambda^{-1}}\|^{4k} \\ &\leq 3^{4k} \left(\|\tilde{Z}_{M,h}^{\lambda,n}\|^{4k} + \left\| \nabla L(\tilde{Z}_{M,h}^{\lambda,n}) \frac{t}{M^2} \right\|^{4k} + \|\Delta \tilde{W}_m \sqrt{2\lambda^{-1}}\|^{4k} \right) \\ &\leq 3^{4k} \left(\|\tilde{Z}_{M,h}^{\lambda,n}\|^{4k} + \left(C_{u,\gamma} \frac{t}{M^2} \right)^{4k} (1 + \|\tilde{Z}_{M,h}^{\lambda,n}\|)^{4k} + \left(\frac{2}{\lambda} \right)^{2k} \|\Delta \tilde{W}_m\|^{4k} \right) \\ &\leq 3^{4k} \left(\|\tilde{Z}_{M,h}^{\lambda,n}\|^{4k} + 2^{4k} + 2^{4k} \|\tilde{Z}_{M,h}^{\lambda,n}\|^{4k} + \left(\frac{2}{\lambda} \right)^{2k} \|\Delta \tilde{W}_m\|^{4k} \right) \\ &\leq 3^{4k} \left((1 + 2^{4k}) \|\tilde{Z}_{M,h}^{\lambda,n}\|^{4k} + 2^{4k} + \left(\frac{2}{\lambda} \right)^{2k} \|\Delta \tilde{W}_m\|^{4k} \right). \end{aligned}$$

Taking expectation of both sides, we have

$$\mathbb{E} \left[\|\tilde{Z}_{m+1,h}^{\lambda,n}\|^{4k} \right] \leq 3^{4k} \left((1 + 2^{4k}) \mathbb{E} \left[\|\tilde{Z}_{M,h}^{\lambda,n}\|^{4k} \right] + 2^{4k} + \left(\frac{2}{\lambda} \right)^{2k} \mathbb{E} \left[\|\Delta \tilde{W}_m\|^{4k} \right] \right). \quad (3.32)$$

We will now show by induction that for any $m \in \{1, \dots, M\}$, it holds that

$$\mathbb{E} \left[\|\tilde{Z}_{m,h}^{\lambda,n}\|^{4k} \right] \leq (3^{4k} (1 + 2^{4k}))^m \|z\|^{4k} + \left(2^{4k} + \left(\frac{2}{\lambda} \right)^{2k} \mathbb{E} \left[\|\Delta W_{m-1}\|^{4k} \right] \right) \sum_{p=0}^{m-1} (3^{4k})^{p+1} (1 + 2^{4k})^p. \quad (3.33)$$

For the base case, from equation (3.32), we have

$$\mathbb{E} \left[\|\tilde{Z}_{1,h}^{\lambda,n}\|^{4k} \right] \leq 3^{4k} \left((1 + 2^{4k}) \|z\|^{4k} + 2^{4k} + \left(\frac{2}{\lambda} \right)^{2k} \mathbb{E} \left[\|\Delta W_0\|^{4k} \right] \right).$$

Now suppose Equation (3.33) holds for $\mathbb{E} \left[\|\tilde{Z}_{M,h}^{\lambda,n}\|^{4k} \right]$. Using the inductive assumption in Equation (3.33), we have

$$\begin{aligned} \mathbb{E} \left[\|\tilde{Z}_{m+1,h}^{\lambda,n}\|^{4k} \right] &\leq 3^{4k} \left\{ (1 + 2^{4k}) \left[(3^{4k} (1 + 2^{4k}))^m \|z\|^{4k} \right. \right. \\ &\quad \left. \left. + \left(2^{4k} + \left(\frac{2}{\lambda} \right)^{2k} \mathbb{E} \left[\|\Delta W_m\|^{4k} \right] \right) \sum_{p=0}^{m-1} (3^{4k})^{p+1} (1 + 2^{4k})^p \right] + 2^{4k} + \left(\frac{2}{\lambda} \right)^{2k} \mathbb{E} \left[\|\Delta W_m\|^{4k} \right] \right\} \\ &\leq (3^{4k} (1 + 2^{4k}))^{m+1} \|z\|^{4k} + \left(2^{4k} + \left(\frac{2}{\lambda} \right)^{2k} \mathbb{E} \left[\|\Delta W_m\|^{4k} \right] \right) \sum_{p=0}^m (3^{4k})^{p+1} (1 + 2^{4k})^p. \end{aligned} \quad (3.34)$$

Thus, the induction is established. Since $\Delta W_m \sim N(0, \frac{t}{M^2} I_{d+1})$, where I_{d+1} is a $(d+1) \times (d+1)$ identity matrix, we have $\mathbb{E} \left[\|\Delta W_m\|^{4k} \right] = C \frac{t^{2k} (4k-1)!}{2^{2k} M^{4k} (2k-1)!}$. Then, simplifying (3.33), yields

$$\mathbb{E} \left[\|\tilde{Z}_{M,h}^{\lambda,n}\|^{4k} \right] \leq (3^{4k} (1 + 2^{4k}))^m \|z\|^{4k} + \frac{3^{4k} (3^{4k} (1 + 2^{4k}))^m}{6^{4k} + 3^{4k} - 1} \left(2^{4k} + \left(\frac{2}{\lambda} \right)^{2k} C \frac{t^{2k} (4k-1)!}{2^{4k} M^{4k} (2k-1)!} \right). \quad (3.35)$$

Therefore, using the series expansion of exponential gives

$$\begin{aligned}
\mathbb{E} \left[\exp \left(C \left\| \tilde{Z}_{m,h}^{\lambda,n} \right\|^4 \right) \right] &= \sum_{k=0}^{\infty} \frac{C^k}{k!} \mathbb{E} \left[\left\| \tilde{Z}_{m,h}^{\lambda,n} \right\|^{4k} \right] \\
&\leq \sum_{k=0}^{\infty} \frac{C^k (3^{4k} (1 + 2^{4k}))^m \|z\|^{4k}}{k!} + \sum_{k=0}^{\infty} \frac{C^k 3^{4k} (3^{4k} (1 + 2^{4k}))^m}{6^{4k} + 3^{4k} - 1} \cdot \frac{2^{4k}}{k!} + \sum_{k=0}^{\infty} \frac{C^k 3^{4k} (3^{4k} (1 + 2^{4k}))^m}{6^{4k} + 3^{4k} - 1} \left(\frac{2}{\lambda} \right)^{2k} \frac{t^{2k} (4k - 1)!}{2^{2k} M^{4k} (2k - 1)!} \\
&=: \sum_{k=0}^{\infty} a_k + \sum_{k=0}^{\infty} b_k + \sum_{k=0}^{\infty} c_k.
\end{aligned}$$

We will show convergence of each term using the ratio test. In fact,

$$\frac{a_{k+1}}{a_k} = 3^{4m} \|z\|^{4k} C \frac{(1 + 2^{4k+4})^m}{(1 + 2^{4k})^m} \frac{k!}{(k+4)!},$$

which converges to 0 as $k \rightarrow \infty$. Thus, $\sum_{k=0}^{\infty} a_k < \infty$. Similarly,

$$\lim_{k \rightarrow \infty} \frac{b_{k+1}}{b_k} = \lim_{k \rightarrow \infty} C 6^4 \left(\frac{3^4 (1 + 2^{4k+4})}{(1 + 2^{4k})} \right)^m \frac{6^{4k} + 3^{4k} - 1}{6^{4k+4} + 3^{4k+4} - 1} \frac{k!}{(k+4)!} = 0,$$

and

$$\lim_{k \rightarrow \infty} \frac{c_{k+1}}{c_k} = \frac{C k!}{(k+4)!} \left(\frac{3^4 (1 + 2^{4k+4})}{(1 + 2^{4k})} \right)^m \frac{6^{4k} + 3^{4k} - 1}{6^{4k+4} + 3^{4k+4} - 1} \frac{16h^2 (4k+3)!(2k+1)!}{\lambda^2 (4k-1)!(2k-1)!} = 0.$$

Therefore, we have $\mathbb{E} \left[\exp \left(C \left\| \tilde{Z}_{m,h}^{\lambda,n} \right\|^4 \right) \right] < C_{(m,h,\lambda,u)}$ for a constant $C_{(m,h,\lambda,u)} > 0$ depending only on m, t, λ , and u . \square

4 Rate for general convex risk measures

We now focus on the estimation of the approximation error of general convex risk measures. As explained in Section 2, the main argument for the derivation of the rate is the representation of the (law invariant) convex risk measure with respect to AVaR. One technical difficulty is that this representation involves an integral with respect to the risk aversion level u of AVaR. Notice that both the functions L and \bar{L} as well as the Markov chain $\tilde{Z}_{m,h}^{\lambda,n}$ in the approximation scheme depend on u . We will make this dependence explicit in this section by writing L^u , $\hat{Z}_t^{\lambda,n,u}$, and $\tilde{Z}_{M,h}^{\lambda,n,u}$ for the function and processes defined in (2.3), (3.4) and (2.5) respectively.

4.1 Proof of Theorem 2.7

This subsection covers the proof of Theorem 2.7. Some auxiliary lemmas needed in the proof are postponed to the final subsection of the article. Recall that we approximate a general law invariant convex risk measure by

$$\tilde{\rho}^\delta(f) := \sup_{\gamma \in \mathcal{M}: \beta(\gamma) \leq b} \left(\int_0^\delta \widetilde{AVaR}_u(f) \gamma(du) - \beta(\gamma) \right),$$

with $\widetilde{AVaR}_u(f) = \frac{1}{N} \sum_{n=1}^N \ell^u(\tilde{Z}_{M,h}^{\lambda,n,u})$. Further define

$$\rho^\delta(f) := \inf_{r \in \mathbb{R}^d} \sup_{\gamma \in \mathcal{M}: \beta(\gamma) \leq b} \left(\int_0^\delta \text{AVaR}_u(f(r, S)) \gamma(du) - \beta(\gamma) \right), \quad \delta \in (0, 1). \quad (4.1)$$

To begin, we decompose the approximation error into two parts:

$$\mathbb{E} \left[|\rho(f) - \tilde{\rho}^\delta(f)|^2 \right] \leq 2|\rho(f) - \rho^\delta(f)|^2 + 2\mathbb{E} \left[|\rho^\delta(f) - \tilde{\rho}^\delta(f)|^2 \right]. \quad (4.2)$$

Let us estimate the first term. First, we will show that for all $\delta \in (0, 1)$, ρ^δ satisfies

$$\begin{aligned}\rho^\delta(f) &= \sup_{\gamma \in \mathcal{M}: \beta(\gamma) \leq b} \left(\int_0^\delta \inf_{r \in \mathbb{R}} \text{AVaR}_u(f(r, S)) \gamma(du) - \beta(\gamma) \right) \\ &= \sup_{\gamma \in \mathcal{M}: \beta(\gamma) \leq b} \left(\int_0^\delta \overline{\text{AVaR}}_u(f) \gamma(du) - \beta(\gamma) \right).\end{aligned}\quad (4.3)$$

Note that going from the definition in (4.1) to the expression in (4.3) requires interchanging the infimum and the supremum and then the infimum and the integral in the definition given in 4.1. Since the supremum is taken over a compact set and the function $(r, \gamma) \mapsto \int_0^\delta \text{AVaR}_u(f(r, S)) \gamma(du) - \beta(\gamma)$ is continuous in r and upper semi-continuous and concave in γ , by Fan's minimax theorem, see [28, Theorem 2], we can interchange the supremum and the infimum. To interchange the infimum and the integral, we apply Rockafellar's interchange theorem [58]. For the first term, using Equation (2.9) and Equation (4.3), it holds that

$$|\rho(f) - \rho^\delta(f)|^2 \leq \sup_{\gamma \in \mathcal{M}: \beta(\gamma) \leq b} \int_\delta^1 |\overline{\text{AVaR}}(f)|^2 \gamma(du).$$

Let us partition the interval $[\delta, 1)$ into $\bigcup_{n \geq 1} I_n$, where $I_n = [\delta + (1 - 2^{-n+1})(1 - \delta), \delta + (1 - 2^{-n})(1 - \delta))$ for every $n \geq 1$. Defining

$$\Gamma_b(I_n) := \sup_{\gamma \in \mathcal{M} \text{ s.t. } \beta(\gamma) \leq b} \gamma(I_n)$$

we obtain the estimation,

$$|\bar{\rho}(f) - \rho^\delta(f)|^2 \leq \sum_{n \geq 1} \Gamma_b(I_n) \sup_{u \in I_n} |\overline{\text{AVaR}}_u(X)|^2. \quad (4.4)$$

Now by [5, Lemma 4.5 and Lemma 4.3], we have

$$\Gamma_b(I_n) \leq C((1 - \delta)2^{-n})^{1/q} \quad \text{and} \quad |\overline{\text{AVaR}}_u(f)| \leq \frac{C}{(1 - u)^{1/p}}, \quad (4.5)$$

for every $p \in (1, \infty)$. Therefore, for any given $p \in (1, \infty)$ and $n \in \mathbb{N}$, it holds

$$\sup_{u \in I_n} |\overline{\text{AVaR}}_u(f)|^2 \leq \frac{C}{((1 - \delta)2^{-n})^{2/p}}. \quad (4.6)$$

Choosing $p = 4q$ and using (4.4), (4.5) and (4.6), we have

$$\begin{aligned}|\bar{\rho}(f) - \rho^\delta(f)|^2 &\leq \sum_{n \geq 1} ((1 - \delta)2^{-n})^{1/q} \frac{C}{((1 - \delta)2^{-n})^{2/p}} \\ &\leq C(1 - \delta)^{1/q - 2/p} \sum_{n \geq 1} 2^{n(2/p - 1/q)} \\ &\leq C(1 - \delta)^{1/2q} \sum_{n \geq 1} 2^{-n/2q}.\end{aligned}$$

Since $q \in (1, \infty)$, it holds $\sum_{n \geq 1} 2^{-n/2q} \leq C$ for some universal constant C , and thus

$$|\bar{\rho}(f) - \rho^\delta(f)|^2 \leq C(1 - \delta)^{1/2q}. \quad (4.7)$$

The estimation of the second term is more involved, and it constitutes the rest of the section. Inspiring ourselves from the proof of Theorem 2.3, it will be adequate to decompose this into terms that can be analyzed separately. First, observe that due to the bound (4.5) on Γ_b we have

$$\begin{aligned}\mathbb{E} \left[|\rho^\delta(f) - \tilde{\rho}^\delta(f)|^2 \right] &= \mathbb{E} \left[\sup_{\gamma \in \mathcal{M}: \beta(\gamma) \leq b} \int_{[0, \delta)} \left| \overline{\text{AVaR}}_u(f) - \widetilde{\text{AVaR}}_u(f) \right|^2 \gamma(du) \right] \\ &\leq \Gamma_b([0, \delta)) \mathbb{E} \left[\sup_{u \in [0, \delta)} \left| \overline{\text{AVaR}}_u(X) - \widetilde{\text{AVaR}}_u(X) \right|^2 \right] \\ &\leq C\delta^{1/q} \mathbb{E} \left[\sup_{u \in [0, \delta)} \left| \overline{\text{AVaR}}_u(f) - \widetilde{\text{AVaR}}_u(f) \right|^2 \right].\end{aligned}\quad (4.8)$$

Thus, adapting the decomposition (3.5) to the present situation allows to obtain

$$\begin{aligned}
& \mathbb{E} \left[\sup_{u \in [0, \delta]} \left| \overline{\text{AVaR}}_u(f) - \widetilde{\text{AVaR}}_u(f) \right|^2 \right] \leq 2^5 \mathbb{E} \left[\sup_{u \in [0, \delta]} \left| \frac{1}{N} \sum_{n=1}^N \ell(\widetilde{Z}'_{M,h}{}^{\lambda, n, u}) - \frac{1}{N} \sum_{n=1}^N L(\widetilde{Z}_{M,h}{}^{\lambda, n, u}) \right|^2 \right] \\
& + 2^5 \mathbb{E} \left[\sup_{u \in [0, \delta]} \left| \frac{1}{N} \sum_{n=1}^N L(\widetilde{Z}_{M,h}{}^{\lambda, n, u}) - \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda, n, u}) \right|^2 \right] + 2^5 \mathbb{E} \left[\sup_{u \in [0, \delta]} \left| \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda, n, u}) - \mathbb{E} \left[L(Z_t^{\lambda, u}) \right] \right|^2 \right] \\
& + 2^5 \sup_{u \in [0, \delta]} \left| \mathbb{E} \left[L(Z_t^{\lambda, u}) \right] - \int L^u d\mu_\infty^\lambda \right|^2 + 2^5 \sup_{u \in [0, \delta]} \left| \int L^u d\mu_\infty^\lambda - \int \widetilde{L}^u d\mu_\infty^\lambda \right|^2 + 2^5 \sup_{u \in [0, \delta]} \left| \int \widetilde{L}^u d\mu_\infty^\lambda - \overline{\text{AVaR}}_u(f) \right|^2.
\end{aligned} \tag{4.9}$$

The estimation of the last three terms follows directly from Equation (3.26) and the lemmas 3.6, 3.8. We omit the details. Unfortunately the estimation of the first three terms does not follow directly from the results obtained in Section 3.2. This is due to the optimization over different risk aversion coefficients $u \in [0, \delta]$ *inside* the expectation.

The estimation of the first term in (4.9) is very similar to Lemma 3.3, except for the fact that we need uniform moment bounds on the processes. The detailed derivation of these uniform moment bounds is deferred to the next subsection. As in the proof of Lemma 3.3, we have

$$\begin{aligned}
& \mathbb{E} \left[\sup_{u \in [0, \delta]} \left| \frac{1}{N} \sum_{n=1}^N l(\widetilde{Z}'_{M,h}{}^{\lambda, n, u}) - \frac{1}{N} \sum_{n=1}^N L(\widetilde{Z}_{M,h}{}^{\lambda, n, u}) \right|^2 \right] \\
& \leq \frac{C}{N} \sum_{n=1}^N \mathbb{E} \left[\sup_{u \in [0, \delta]} \left(\widetilde{L}(\widetilde{Z}_{M,h}{}^{\lambda, n, u}) - \widetilde{L}(\widetilde{Z}'_{M,h}{}^{\lambda, n, u}) \right)^2 \right] + \frac{C}{N} \sum_{n=1}^N \mathbb{E} \left[\sup_{u \in [0, \delta]} \left(\widetilde{L}(\widetilde{Z}'_{M,h}{}^{\lambda, n, u}) - \widetilde{\ell}(\widetilde{Z}'_{M,h}{}^{\lambda, n, u}) \right)^2 \right] \\
& + C \frac{\gamma^2}{N} \sum_{n=1}^N \left(\mathbb{E} \left[\sup_{u \in [0, \delta]} \|\widetilde{Z}_{M,h}{}^{\lambda, n, u}\|^4 \right] + \mathbb{E} \left[\sup_{u \in [0, \delta]} \|\widetilde{Z}'_{M,h}{}^{\lambda, n, u}\|^4 \right] \right)^{\frac{1}{2}} \mathbb{E} \left[\sup_{u \in [0, \delta]} \|\widetilde{Z}_{M,h}{}^{\lambda, n, u} - \widetilde{Z}'_{M,h}{}^{\lambda, n, u}\|^4 \right]^{\frac{1}{2}}.
\end{aligned} \tag{4.10}$$

By $\frac{1}{1-u}$ -Lipschitz-continuity of \widetilde{L}^u , for the first term in (4.10), we have

$$\begin{aligned}
\mathbb{E} \left[\sup_{u \in [0, \delta]} \left(\widetilde{L}(\widetilde{Z}_{M,h}{}^{\lambda, n, u}) - \widetilde{L}(\widetilde{Z}'_{M,h}{}^{\lambda, n, u}) \right)^2 \right] & \leq \left(\frac{1}{1-\delta} \right)^2 \mathbb{E} \left[\sup_{u \in [0, \delta]} \|\widetilde{Z}_{M,h}{}^{\lambda, n, u} - \widetilde{Z}'_{M,h}{}^{\lambda, n, u}\|^2 \right] \\
& \leq C_{(\delta, t)} \frac{1}{M^7}
\end{aligned} \tag{4.11}$$

where the second inequality follows from Lemma 4.3.

The estimation of the second term on the right hand side of (4.10) builds on a law of large numbers type argument. To this end, we will need to estimate the supremum by a countable sum. Thus, let I_c be a countable dense subset of the

interval $[0, \delta)$. The first equality below is Lemma 4.1, and the last one follows since the random variables S^i are i.i.d.

$$\begin{aligned}
& \mathbb{E} \left[\sup_{u \in [0, \delta)} \left(\tilde{L}(\tilde{Z}'_{M,h}{}^{\lambda,n,u}) - \tilde{\ell}(\tilde{Z}'_{M,h}{}^{\lambda,n,u}) \right)^2 \right] \\
&= \left(\frac{1}{1-\delta} \right)^2 \mathbb{E} \left[\sup_{u \in I_c} \left| \tilde{\mathbb{E}}[(f(r'_M, S) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] - \frac{1}{N} \sum_{i=1}^N (f(r'_M, S^i) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ \right|^2 \right] \\
&= \left(\frac{1}{1-\delta} \right)^2 \frac{1}{N^2} \mathbb{E} \left[\sup_{u \in I_c} \sum_{i \neq j} \left(\tilde{\mathbb{E}}[(f(r'_M, S) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] - (f(r'_M, S^i) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ \right) \right. \\
&\quad \cdot \left. \left(\tilde{\mathbb{E}}[(f(r'_M, S) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] - (f(r'_M, S^j) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ \right) \right] \\
&\quad + \left(\frac{1}{1-\delta} \right)^2 \frac{1}{N^2} \mathbb{E} \left[\sup_{u \in I_c} \sum_i \left(\tilde{\mathbb{E}}[(f(r'_M, S) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] - (f(r'_M, S^i) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ \right)^2 \right] \\
&\leq \left(\frac{1}{1-\delta} \right)^2 \frac{1}{N^2} \mathbb{E} \left[\sum_{u \in I_c} \sum_{i \neq j} \left(\tilde{\mathbb{E}}[(f(r'_M, S) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] - (f(r'_M, S^i) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ \right) \right. \\
&\quad \cdot \left. \left(\tilde{\mathbb{E}}[(f(r'_M, S) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] - (f(r'_M, S^j) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ \right) \right] \\
&\quad + \left(\frac{1}{1-\delta} \right)^2 \frac{1}{N^2} \sum_{i=1}^N \mathbb{E} \left[\sup_{u \in I_c} \left(\tilde{\mathbb{E}}[(f(r'_M, S) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] - (f(r'_M, S^i) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ \right)^2 \right] \\
&= \left(\frac{1}{1-\delta} \right)^2 \frac{1}{N^2} \sum_{i=1}^N \mathbb{E} \left[\sup_{u \in I_c} \left(\tilde{\mathbb{E}}[(f(r'_M, S) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+] - (f(r'_M, S^i) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+ \right)^2 \right] \\
&\leq \left(\frac{1}{1-\delta} \right)^2 \frac{C}{N^2} \sum_{i=1}^N \left(\mathbb{E} \left[\sup_{u \in I_c} |\tilde{\mathbb{E}}[(f(r'_M, S) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+]|^2 \right] + \mathbb{E} \left[\sup_{u \in I_c} |(f(r'_M, S^i) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+|^2 \right] \right).
\end{aligned}$$

Due to equations (4.18) and (4.19), and taking $0 < \gamma < 1$, we have

$$\left(\frac{1}{1-\delta} \right)^2 \mathbb{E} \left[\sup_{u \in I_c} |\tilde{\mathbb{E}}[(f(r'_M, S) - \tilde{Z}'_{M,h}{}^{\lambda,n}(1))^+]|^2 \right] \leq C_{(\delta,t,\lambda)}. \quad (4.12)$$

Combining (4.10), (4.11), (4.12), (4.18), and Lemma 4.3, we have

$$\mathbb{E} \left[\sup_{u \in [0, \delta)} \left| \frac{1}{N} \sum_{n=1}^N l(\tilde{Z}'_{M,h}{}^{\lambda,n,u}) - \frac{1}{N} \sum_{n=1}^N L(\tilde{Z}_{M,h}{}^{\lambda,n,u}) \right|^2 \right] \leq C_{(\delta,t)} \frac{1}{M^7} + \frac{1}{N} C_{(\delta,t,\gamma,\lambda)} + C_{(h,M,\delta,\lambda)} \gamma^2. \quad (4.13)$$

Regarding the second term in (4.9), applying subsequently Jensen's inequality and Cauchy-Schwartz inequality we have

$$\begin{aligned}
& \mathbb{E} \left[\sup_{u \in [0, \delta)} \left| \frac{1}{N} \sum_{n=1}^N L^u(\tilde{Z}_{M,h}{}^{\lambda,n,u}) - \frac{1}{N} \sum_{n=1}^N L^u(\hat{Z}_t^{\lambda,n,u}) \right|^2 \right] \leq \frac{C}{N} \sum_{n=1}^N \mathbb{E} \left[\sup_{u \in [0, \delta)} |\tilde{L}^u(\tilde{Z}_{M,h}{}^{\lambda,n,u}) - \tilde{L}^u(\hat{Z}_t^{\lambda,n,u})|^2 \right] \\
&\quad + \frac{C\gamma}{N} \sum_{n=1}^N \left(\mathbb{E} \left[\sup_{u \in [0, \delta)} \|\tilde{Z}_{M,h}{}^{\lambda,n,u}\|^4 \right] + \mathbb{E} \left[\sup_{u \in [0, \delta)} \|\hat{Z}_t^{\lambda,n,u}\|^4 \right] \right)^{1/2} \left(\mathbb{E} \left[\sup_{u \in [0, \delta)} \|\tilde{Z}_{M,h}{}^{\lambda,n,u} - \hat{Z}_t^{\lambda,n,u}\|^4 \right] \right)^{1/2} \\
&\leq \frac{C}{N} \sum_{n=1}^N \mathbb{E} \left[\sup_{u \in [0, \delta)} \frac{1}{(1-u)^2} \|\tilde{Z}_{M,h}{}^{\lambda,n,u} - \hat{Z}_t^{\lambda,n,u}\|^2 \right] \\
&\quad + \frac{C\gamma}{N} \sum_{n=1}^N \left(\mathbb{E} \left[\sup_{u \in [0, \delta)} \|\tilde{Z}_{M,h}{}^{\lambda,n,u}\|^4 \right] + \mathbb{E} \left[\sup_{u \in [0, \delta)} \|\hat{Z}_t^{\lambda,n,u}\|^4 \right] \right)^{1/2} \left(\mathbb{E} \left[\sup_{u \in [0, \delta)} \|\tilde{Z}_{M,h}{}^{\lambda,n,u} - \hat{Z}_t^{\lambda,n,u}\|^4 \right] \right)^{1/2} \\
&\leq C_{(\delta,t,\lambda)} \frac{1}{M^2}, \quad (4.14)
\end{aligned}$$

where the last inequality follows by the moment bounds given in Lemma 4.2 and Lemma 4.4 below and by taking $0 < \gamma < 1$.

Finally, following the same idea of estimating the supremum by a countable sum and then applying a law of large numbers argument as in the second term of (4.10), we have

$$\begin{aligned} & \mathbb{E} \left[\sup_{u \in [0, \delta]} \left| \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda, n, u}) - \mathbb{E} [L(Z_t^{\lambda, u})] \right|^2 \right] = \mathbb{E} \left[\sup_{u \in I_c} \left| \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda, n, u}) - \mathbb{E} [L(Z_t^{\lambda, u})] \right|^2 \right] \\ & \leq \frac{C}{N^2} \sum_{i=1}^N \left(\mathbb{E} \left[\sup_{u \in I_c} |L^u(\widehat{Z}_t^{\lambda, n, u})|^2 \right] + \sup_{u \in I_c} \mathbb{E} \left[|L^u(Z_t^{\lambda, u})|^2 \right] \right). \end{aligned}$$

Thus, by Lemma 4.2 it holds

$$\mathbb{E} \left[\sup_{u \in [0, \delta]} \left| \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda, n, u}) - \mathbb{E} [L(Z_t^{\lambda, u})] \right|^2 \right] \leq \frac{C(\delta, t, \lambda)}{N}. \quad (4.15)$$

Now, combine the expressions (4.2), (4.7), (4.8), (4.9), (4.14) and (4.15) to conclude the proof.

4.2 Auxiliary lemmas

This final section of the paper gathers lemmas used in the proof of the error bound given in Theorem 2.7. Throughout this section, we assumed that the conditions of Theorem 2.7 are satisfied. The first lemma allows us to interchange supremum and infimum in the definition of $\bar{\rho}$.

Lemma 4.1. *Let I_c be a countable dense subset of $[0, \delta]$. Then it holds*

$$\mathbb{E} \left[\sup_{u \in [0, \delta]} \left| \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda, n, u}) - \mathbb{E} [L(Z_t^{\lambda, u})] \right|^2 \right] = \mathbb{E} \left[\sup_{u \in I_c} \left| \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda, n, u}) - \mathbb{E} [L(Z_t^{\lambda, u})] \right|^2 \right],$$

and

$$\mathbb{E} \left[\sup_{u \in [0, \delta]} \left(\widetilde{L}(\widetilde{Z}'_{M, h}{}^{\lambda, n, u}) - \widetilde{\ell}(\widetilde{Z}'_{M, h}{}^{\lambda, n, u}) \right)^2 \right] = \mathbb{E} \left[\sup_{u \in I_c} \left(\widetilde{L}(\widetilde{Z}'_{M, h}{}^{\lambda, n, u}) - \widetilde{\ell}(\widetilde{Z}'_{M, h}{}^{\lambda, n, u}) \right)^2 \right] \quad (4.16)$$

Proof. Let $u \in [0, \delta]$ and a sequence (u_k) in I_c converging to u . Let \widehat{R}^u and \widehat{R}^{u_k} denote the last d coordinates of $\widehat{Z}^{\lambda, n, u}$ and $\widehat{Z}^{\lambda, n, u_k}$. Let us show that, $L^{u_k}(\widehat{Z}_t^{\lambda, n, u_k}) \rightarrow \widetilde{L}^u(\widehat{Z}_t^{\lambda, n, u})$ \mathbb{P} -a.s., $\widetilde{L}^{u_k}(\widetilde{Z}'_{M, h}{}^{\lambda, n, u_k}) \rightarrow L^u(\widetilde{Z}'_{M, h}{}^{\lambda, n, u})$ \mathbb{P} -a.s., and $\widetilde{l}^{u_k}(\widetilde{Z}'_{M, h}{}^{\lambda, n, u_k}) \rightarrow \widetilde{l}^u(\widetilde{Z}'_{M, h}{}^{\lambda, n, u})$ \mathbb{P} -a.s. We first show that $\widehat{Z}_t^{\lambda, n, u_k} \rightarrow \widehat{Z}_t^{\lambda, n, u}$ \mathbb{P} -a.s. Using the definition of $\widehat{Z}_r^{\lambda, n, u}$, then adding and subtracting $\nabla L^u(\widehat{Z}_r^{\lambda, n, u})$, we have

$$\begin{aligned} & \|\widehat{Z}_t^{\lambda, n, u_k} - \widehat{Z}_t^{\lambda, n, u}\| \\ & \leq \int_0^t \|\nabla L^u(\widehat{Z}_s^{\lambda, n, u}) - \nabla L^{u_k}(\widehat{Z}_s^{\lambda, n, u_k})\| ds \\ & \leq \int_0^t \|\nabla L^u(\widehat{Z}_s^{\lambda, n, u}) - \nabla L^u(\widehat{Z}_s^{\lambda, n, u_k})\| ds + \int_0^t \|\nabla L^u(\widehat{Z}_s^{\lambda, n, u_k}) - \nabla L^{u_k}(\widehat{Z}_s^{\lambda, n, u_k})\| ds \\ & \leq \int_0^t \left(\frac{1}{1-u} + \gamma \right) \|\widehat{Z}_s^{\lambda, n, u_k} - \widehat{Z}_s^{\lambda, n, u}\| ds \\ & + \int_0^t \left\| e_1 + \frac{1}{1-u} \mathbb{E}[\nabla f(\widehat{R}^{u_k}, S) \mathbf{1}_{\{f(\widehat{R}^{u_k}, S) \geq \widehat{Z}_s^{\lambda, n, u_k}(1)\}}] - \left(e_1 + \frac{1}{1-u_k} \mathbb{E}[\nabla f(\widehat{R}^{u_k}, S) \mathbf{1}_{\{f(\widehat{R}^{u_k}, S) \geq \widehat{Z}_s^{\lambda, n, u_k}(1)\}}] \right) \right\| dr \\ & \leq \int_0^t \left(\frac{1}{1-u} + \gamma \right) \|\widehat{Z}_s^{\lambda, n, u_k} - \widehat{Z}_s^{\lambda, n, u}\| ds + \int_0^t \left| \frac{1}{1-u} - \frac{1}{1-u_k} \right| \mathbb{E}[\|\nabla f(\widehat{R}^{u_k}, S) \mathbf{1}_{\{f(\widehat{R}^{u_k}, S) \geq \widehat{Z}_s^{\lambda, n, u}(1)\}}\|] ds \\ & \leq \int_0^t \left(\frac{1}{1-u} + \gamma \right) \|\widehat{Z}_s^{\lambda, n, u_k} - \widehat{Z}_s^{\lambda, n, u}\| ds + Ct \left| \frac{1}{1-u} - \frac{1}{1-u_k} \right|, \end{aligned}$$

where we used the boundedness of $\|\nabla f\|$ in the last step. Applying Grönwall's inequality leads to

$$\|\widehat{Z}_t^{\lambda, n, u_k} - \widehat{Z}_t^{\lambda, n, u}\| \leq Ct \left| \frac{1}{1-u} - \frac{1}{1-u_k} \right| \exp \left\{ \left(\frac{1}{1-u} + \gamma \right) t \right\}.$$

Thus, we have $\lim_{u_k \rightarrow u} \|\widehat{Z}_t^{\lambda, n, u_k} - \widehat{Z}_t^{\lambda, n, u}\| = 0$. Next, subtracting and adding $L^{u_k}(\widehat{Z}_t^{\lambda, n, u})$, we have

$$\begin{aligned} |L^{u_k}(\widehat{Z}_t^{\lambda, n, u_k}) - L^u(\widehat{Z}_t^{\lambda, n, u})| &\leq |L^{u_k}(\widehat{Z}_t^{\lambda, n, u_k}) - L^{u_k}(\widehat{Z}_t^{\lambda, n, u})| + |L^{u_k}(\widehat{Z}_t^{\lambda, n, u}) - L^u(\widehat{Z}_t^{\lambda, n, u})| \\ &\leq \frac{2}{1-u_k} \|\widehat{Z}_t^{\lambda, n, u_k} - \widehat{Z}_t^{\lambda, n, u}\| + \frac{\gamma}{2} \left(\|\widehat{Z}_t^{\lambda, n, u_k}\|^2 - \|\widehat{Z}_t^{\lambda, n, u}\|^2 \right) \\ &\quad + \left| \frac{1}{1-u_k} - \frac{1}{1-u} \right| \mathbb{E}[(f(\widehat{R}^u, S) - \widehat{Z}_t^{\lambda, n, u}(1))^+]. \end{aligned}$$

Since $\mathbb{E}[\widehat{Z}_t^{\lambda, n, u}(1)] \leq \mathbb{E}[\|\widehat{Z}_t^{\lambda, n, u}\|] \leq C_{(t, \lambda, \gamma)}$, and by the finite fourth moment of S , we have $\mathbb{E}[(f(\widehat{R}^u, S) - \widehat{Z}_t^{\lambda, n, u}(1))^+] \leq C_{(t, \lambda, \gamma)}$. Thus, we have that almost surely,

$$\lim_{u_k \rightarrow u} |L^{u_k}(\widehat{Z}_t^{\lambda, n, u_k}) - L^u(\widehat{Z}_t^{\lambda, n, u})| = 0.$$

Next, we will show $\widetilde{Z}'_{M, h}{}^{\lambda, n, u_k} \rightarrow \widetilde{Z}'_{M, h}{}^{\lambda, n, u}$ \mathbb{P} -a.s. Using the definition of $\widetilde{Z}'_{M, h}{}^{\lambda, n, u}$, and the boundedness of $\|\nabla f\|$, we have

$$\|\widetilde{Z}'_{M, h}{}^{\lambda, n, u_k} - \widetilde{Z}'_{M, h}{}^{\lambda, n, u}\| \leq \left(1 + \frac{t}{(1-u)M^2} + \gamma \frac{t}{M^2}\right) \|\widetilde{Z}'_{M-1, h}{}^{\lambda, n, u_k} - \widetilde{Z}'_{M-1, h}{}^{\lambda, n, u}\| + \left| \frac{1}{1-u} - \frac{1}{1-u_k} \right| C \frac{t}{M^2}.$$

Using the above recursive relationship, we have

$$\|\widetilde{Z}'_{M, h}{}^{\lambda, n, u_k} - \widetilde{Z}'_{M, h}{}^{\lambda, n, u}\| \leq \left| \frac{1}{1-u} - \frac{1}{1-u_k} \right| Ch \sum_{i=1}^M \left(1 + \frac{t}{(1-u)M^2} + \gamma \frac{t}{M^2}\right)^{i-1},$$

where we used $\widetilde{Z}'_{0, h}{}^{\lambda, n, u_k} = \widetilde{Z}'_{0, h}{}^{\lambda, n, u} = z$. Thus, $\lim_{u_k \rightarrow u} \widetilde{Z}'_{M, h}{}^{\lambda, n, u_k} - \widetilde{Z}'_{M, h}{}^{\lambda, n, u} = 0$. Then, following the exact same argument as in the case of L , we have that $\widetilde{L}^{u_k}(\widetilde{Z}'_{M, h}{}^{\lambda, n, u}) \rightarrow L^u(\widetilde{Z}'_{M, h}{}^{\lambda, n, u})$ \mathbb{P} -a.s., and $\widetilde{l}^{u_k}(\widetilde{Z}'_{M, h}{}^{\lambda, n, u}) \rightarrow \widetilde{l}^u(\widetilde{Z}'_{M, h}{}^{\lambda, n, u})$ \mathbb{P} -a.s.. Since $I_c \subset [0, \delta)$ is dense, for any $u \in I$, there exists a sequence $u_k \in I_c$ such that $u_k \rightarrow u$. Then, we have

$$\sup_{u \in I_c} \left| \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda, n, u}) - \mathbb{E}[L(Z_t^{\lambda, u})] \right|^2 \geq \sup_{u \in [0, \delta)} \left| \frac{1}{N} \sum_{n=1}^N L(\widehat{Z}_t^{\lambda, n, u}) - \mathbb{E}[L(Z_t^{\lambda, u})] \right|^2,$$

and

$$\sup_{u \in I_c} \left| \widetilde{L}(\widetilde{Z}'_{M, h}{}^{\lambda, n, u}) - \widetilde{\ell}(\widetilde{Z}'_{M, h}{}^{\lambda, n, u}) \right|^2 \geq \sup_{u \in [0, \delta)} \left| \widetilde{L}(\widetilde{Z}'_{M, h}{}^{\lambda, n, u}) - \widetilde{\ell}(\widetilde{Z}'_{M, h}{}^{\lambda, n, u}) \right|^2$$

Thanks to the first part of the proof, first taking the limit as $u_k \rightarrow u$ and then the supremum over $u \in [0, \delta)$ yields the inequality “ \geq ”. The reverse inequality is clear. \square

Lemma 4.2. *We have the following uniform moment bounds:*

$$\mathbb{E} \left[\sup_{u \in [0, \delta)} \left\| \widehat{Z}_t^{\lambda, n, u} \right\|^4 \right] \leq C \left(1 + \frac{t}{(1-\delta)^4} + \frac{t^2}{\lambda^2} \right) e^{C\gamma^4 t}, \quad (4.17)$$

$$\mathbb{E} \left[\sup_{u \in [0, \delta)} \left\| \widetilde{Z}'_{M, h}{}^{\lambda, n, u} \right\|^4 \right] + \mathbb{E} \left[\sup_{u \in [0, \delta)} \left\| \widetilde{Z}_{M, h}{}^{\lambda, n, u} \right\|^4 \right] \leq C \left(\frac{h^{4M+4}}{(1-\delta)^4} + \frac{h^{4M+2} \gamma^{4M}}{\lambda^2} \right), \quad (4.18)$$

and

$$\mathbb{E} \left[\sup_{u \in I_c} |L^u(\widehat{Z}_t^{\lambda, i, u})|^2 \right] \leq C_{(\delta, t, \lambda)} \quad (4.19)$$

Proof. Recall that $\widehat{Z}_t^{\lambda, n, u}$ solves the SDE

$$d\widehat{Z}_t^{\lambda, n, u} = -\nabla L^u(\widehat{Z}_t^{\lambda, n, u}) dt + \sqrt{2\lambda^{-1}} d\widetilde{W}^n.$$

Thus, we have

$$\begin{aligned}
\mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widehat{Z}_t^{\lambda, n, u} \right\|^4 \right] &\leq C \left(\|z\|^4 + \mathbb{E} \left[\int_0^t \sup_{u \in [0, \delta]} \left\| \nabla L(\widehat{Z}_{s, u}^{\lambda, n}) \right\|^4 ds \right] + \frac{4}{\lambda^2} \mathbb{E}[\|W_t\|^4] \right) \\
&\leq C \left(\|z\|^4 + \mathbb{E} \left[\int_0^t \sup_{u \in [0, \delta]} \left\| \frac{2}{1-u} + \gamma \widehat{Z}_{s, u}^{\lambda, n} \right\|^4 ds \right] + \frac{1}{\lambda^2} t^2 \right) \\
&\leq C \left(\|z\|^4 + \frac{t}{(1-\delta)^4} + \gamma^4 \int_0^t \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widehat{Z}_{s, u}^{\lambda, n} \right\|^4 \right] ds + \frac{t^2}{\lambda^2} \right),
\end{aligned}$$

where in the second line, we used that \widetilde{L} is $\frac{2}{1-u}$ -Lipschitz. Applying Grönwall's inequality, we have the desired result.

For Equation (4.18), recall that $\widetilde{Z}_{M, h}^{\lambda, n, u}$ is the M -th step of the SGLD algorithm with $\Delta \widetilde{W}_M^n$. We have

$$\begin{aligned}
\mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widetilde{Z}_{M, h}^{\lambda, n, u} \right\|^4 \right] &\leq \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widetilde{Z}_{M-1, h}^{\lambda, n, u} - h \nabla L(\widetilde{Z}_{M-1, h}^{\lambda, n, u}) + \Delta \widetilde{W}_M^n \right\|^4 \right] \\
&\leq C \left(\mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widetilde{Z}_{M-1, h}^{\lambda, n, u} \right\|^4 \right] + h^4 \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \nabla L(\widetilde{Z}_{M-1, h}^{\lambda, n, u}) \right\|^4 \right] + \mathbb{E} \left\| \Delta \widetilde{W}_M^n \right\|^4 \right) \\
&\leq C \left(\mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widetilde{Z}_{M-1, h}^{\lambda, n, u} \right\|^4 \right] + h^4 \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \frac{2}{1-u} + \gamma \widetilde{Z}_{M-1, h}^{\lambda, n, u} \right\|^4 \right] + \frac{t^2}{M^2 \lambda^2} \right) \\
&\leq C \left((1 + \gamma^4 h^4) \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widetilde{Z}_{M-1, h}^{\lambda, n, u} \right\|^4 \right] + \frac{t^4}{(1-\delta)^4 M^8} + \frac{t^2}{\lambda^2 M^4} \right).
\end{aligned}$$

Using this recursive relationship, we get

$$\begin{aligned}
\mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widetilde{Z}_{M, h}^{\lambda, n, u} \right\|^4 \right] &\leq C \left((1 + \gamma^4 h^4)^M \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widetilde{Z}_{0, h}^{\lambda, n, u} \right\|^4 \right] + \left(\frac{h^4}{(1-\delta)^4} + \frac{h^2}{\lambda^2} \right) \sum_{i=1}^M (1 + \gamma^4 h^4)^i \right) \\
&\leq C \left(\frac{h^{4M+4}}{(1-\delta)^4} + \frac{h^{4M+2} \gamma^{4M}}{\lambda^2} \right).
\end{aligned}$$

Exactly the same argument yields the bound for $\mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widetilde{Z}'_{M, h}^{\lambda, n, u} \right\|^4 \right]$. Now, using the relation between L^u and \widetilde{L}^u , and that \widetilde{L}^u is $\frac{2}{1-u}$ -Lipschitz, we have

$$\begin{aligned}
&\mathbb{E} \left[\sup_{u \in I_c} |L^u(\widehat{Z}_t^{\lambda, i, u})|^2 \right] \\
&\leq C \left(\frac{1}{(1-\delta)^2} \mathbb{E} \left[\sup_{u \in I_c} \left\| \widehat{Z}_t^{\lambda, i, u} \right\|^2 \right] + \gamma \mathbb{E} \left[\sup_{u \in I_c} \left\| \widehat{Z}_t^{\lambda, i, u} \right\|^4 \right] + 1 \right) \\
&\leq C \left(\frac{1}{(1-\delta)^2} \left(z^2 + \frac{\sqrt{t}}{(1-\delta)^2} + \frac{t}{\lambda} \right) e^{C\gamma^4 t} + \gamma \left(z^4 + \frac{t}{(1-\delta)^2} + \frac{t^2}{\lambda^2} \right) e^{C\gamma^4 t} + 1 \right) \\
&\leq C_{(\delta, t, \lambda)},
\end{aligned}$$

where in the last inequality, we used Lemma 4.2 and $0 < \gamma < 1$. □

Lemma 4.3. For all $t, \gamma, \lambda > 0$, $\delta \in (0, 1)$ and $M, n \in \mathbb{N}^*$, it holds

$$\mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widetilde{Z}_{M, h}^{\lambda, n, u} - \widetilde{Z}'_{M, h}^{\lambda, n} \right\|^2 \right] \leq C_{(\delta, t)} \frac{1}{M^7}.$$

Proof. Using the definitions of $\widetilde{Z}_{M, h}^{\lambda, n, u}$ and $\widetilde{Z}'_{M, h}^{\lambda, n}$, we have

$$\begin{aligned}
\mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widetilde{Z}_{M, h}^{\lambda, n, u} - \widetilde{Z}'_{M, h}^{\lambda, n, u} \right\|^4 \right] &= \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widetilde{Z}_{M-1, h}^{\lambda, n} - h \nabla L(\widetilde{Z}_{M-1, h}^{\lambda, n}) - \widetilde{Z}'_{M-1, h}^{\lambda, n} + h \nabla \ell(\widetilde{Z}'_{M-1, h}^{\lambda, n}) \right\|^4 \right] \\
&\leq C \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \widetilde{Z}_{M-1, h}^{\lambda, n} - \widetilde{Z}'_{M-1, h}^{\lambda, n} \right\|^4 \right] + Ch^4 \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \nabla L(\widetilde{Z}_{M-1, h}^{\lambda, n}) - \nabla \ell(\widetilde{Z}'_{M-1, h}^{\lambda, n}) \right\|^4 \right].
\end{aligned}$$

Using this recursive relationship, it can be checked by induction that we have

$$\mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \tilde{Z}_{M,h}^{\lambda, n, u} - \tilde{Z}'_{M,h}{}^{\lambda, n, u} \right\|^4 \right] \leq C \sum_{m=0}^{M-1} h^4 \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \nabla L(\tilde{Z}_{M,h}^{\lambda, n, u}) - \nabla \ell(\tilde{Z}'_{M,h}{}^{\lambda, n, u}) \right\|^4 \right]. \quad (4.20)$$

Let $m \in \{1, \dots, M-1\}$. Let \hat{R}^u denote the last d coordinates of $\tilde{Z}'_{M,h}{}^{\lambda, n, u}$. Adding and subtracting $\nabla L(\tilde{Z}'_{M,h}{}^{\lambda, n, u})$, we get

$$\begin{aligned} & \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \nabla L(\tilde{Z}_{M,h}^{\lambda, n, u}) - \nabla \ell(\tilde{Z}'_{M,h}{}^{\lambda, n, u}) \right\|^4 \right] \\ & \leq C \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \nabla L(\tilde{Z}_{M,h}^{\lambda, n, u}) - \nabla L(\tilde{Z}'_{M,h}{}^{\lambda, n, u}) \right\|^4 \right] + C \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \nabla L(\tilde{Z}'_{M,h}{}^{\lambda, n, u}) - \nabla \ell(\tilde{Z}'_{M,h}{}^{\lambda, n, u}) \right\|^4 \right] \\ & \leq C \sup_{u \in [0, \delta]} \left(\frac{2}{1-u} \right)^4 \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \tilde{Z}_{M,h}^{\lambda, n, u} - \tilde{Z}'_{M,h}{}^{\lambda, n, u} \right\|^4 \right] \\ & + C \mathbb{E} \left[\sup_{u \in [0, \delta]} \left| \left(e_1 + \frac{1}{1-u} \tilde{\mathbb{E}}[\nabla f(r, S) 1_{\{f(\hat{R}^u, S) \geq (\tilde{Z}'_{M,h}{}^{\lambda, n, u})_1\}}] \right) - \left(e_1 + \frac{1}{1-u} \frac{1}{N} \sum_{i=1}^N (\nabla f(\hat{R}^u, S_i) 1_{\{f(\hat{R}^u, S_i) \geq (\tilde{Z}'_{M,h}{}^{\lambda, n, u})_1\}}) \right) \right|^4 \right] \\ & \leq C \left(\frac{1}{1-\delta} \right)^4 \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \tilde{Z}_{M,h}^{\lambda, n, u} - \tilde{Z}'_{M,h}{}^{\lambda, n, u} \right\|^4 \right] + C \left(\frac{1}{1-\delta} \right)^4, \end{aligned} \quad (4.21)$$

where the last inequality follows since $\nabla f(r, S)$ is bounded. Putting equations (4.20) and (4.21) together, we have

$$\mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \tilde{Z}_{M,h}^{\lambda, n, u} - \tilde{Z}'_{M,h}{}^{\lambda, n, u} \right\|^4 \right] \leq Ch^4 \left(\frac{1}{1-\delta} \right)^4 \sum_{m=0}^{M-1} \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \tilde{Z}_{M,h}^{\lambda, n, u} - \tilde{Z}'_{M,h}{}^{\lambda, n, u} \right\|^4 \right] + Ch^4 M \left(\frac{1}{1-\delta} \right)^4.$$

Now, using the discrete version of the Grönwall's inequality [38, Proposition 5] and $h = \frac{t}{M^2}$ yields

$$\mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| \tilde{Z}_{M,h}^{\lambda, n, u} - \tilde{Z}'_{M,h}{}^{\lambda, n, u} \right\|^2 \right] \leq C \frac{t^4}{M^7} \left(\frac{1}{1-\delta} \right)^4 \exp \left(C \frac{t^4}{M^7} \left(\frac{2}{1-\delta} \right)^4 \right).$$

This gives the result. \square

Lemma 4.4. *For all $t, \gamma, \lambda > 0$, $\delta \in (0, 1)$ and $M, n \in \mathbb{N}^*$, it holds*

$$\mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| Z^{\lambda, u}(t) - \hat{Z}_t^{\lambda, n, u} \right\|^4 \right] \leq C_{(\delta, t, \lambda)} \frac{1}{M^8}. \quad (4.22)$$

Proof. The main idea of this proof comes from the standard argument for the strong convergence of the Euler-Maruyama scheme. Let $n_t := \max\{hn : hn \leq t, n \in \mathbb{Z}\}$, and recall that $Z^{\lambda, u}$ is the continuous time approximation of the Euler-Maruyama scheme given in 2.5. Using the definition of $\hat{Z}_t^{\lambda, n, u}$ and $Z^{\lambda, u}(t)$, we have

$$\begin{aligned} \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| Z^{\lambda, u}(t) - \hat{Z}_t^{\lambda, n, u} \right\|^4 \right] & \leq C \mathbb{E} \left[\sup_{u \in [0, \delta]} \left| \int_0^{n_t} \left\| \nabla L(Z^{\lambda, u}(s)) - \nabla L(\hat{Z}_s^{\lambda, n, u}) \right\| ds \right|^4 \right] + C \mathbb{E} \left[\sup_{u \in [0, \delta]} \left| \int_{n_t}^t \left\| \nabla L(\hat{Z}_s^{\lambda, n, u}) \right\| ds \right|^4 \right] \\ & + C \mathbb{E} \left[\left\| \int_{n_t}^t \sqrt{\frac{2}{\lambda}} dW_s \right\|^4 \right]. \end{aligned}$$

Using Jensen's inequality gives

$$\begin{aligned} \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| Z^{\lambda, u}(t) - \hat{Z}_t^{\lambda, n, u} \right\|^4 \right] & \leq Ct^3 \mathbb{E} \left[\sup_{u \in [0, \delta]} \int_0^{n_t} \left\| \nabla L(Z^{\lambda, u}(s)) - \nabla L(\hat{Z}_s^{\lambda, n, u}) \right\|^4 dr \right] \\ & + C \frac{t^3}{M^6} \mathbb{E} \left[\sup_{u \in [0, \delta]} \int_{n_t}^t \left\| \nabla L(\hat{Z}_s^{\lambda, n, u}) \right\|^4 ds \right] + C \frac{4}{\lambda^2} \mathbb{E}[\|W_t - W_{n_t}\|^4]. \end{aligned}$$

Using that $\nabla \tilde{L}$ is $\frac{1}{1-u}$ -Lipschitz and that \tilde{L} is $\frac{2}{1-u}$ -Lipschitz, we have

$$\begin{aligned} \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| Z^{\lambda, u}(t) - \hat{Z}_t^{\lambda, n, u} \right\|^4 \right] &\leq Ct^3 \left(\frac{1}{1-\delta} + \gamma \right)^4 \int_0^{n_t} \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| Z^{\lambda, u}(s) - \hat{Z}_s^{\lambda, n, u} \right\|^4 ds \right] \\ &\quad + Ch^3 \mathbb{E} \int_{n_t}^t \sup_{u \in [0, \delta]} \left| \frac{2}{1-u} \right|^4 dr + Ch^3 \gamma^4 \mathbb{E} \left[\sup_{u \in [0, \delta]} \int_{n_t}^t \left\| \hat{Z}_s^{\lambda, n, u} \right\|^4 ds \right] + C \frac{h^2}{\lambda^2} \\ &\leq Ct^3 \left(\frac{1}{1-\delta} + \gamma \right)^4 \int_0^{n_t} \mathbb{E} \left[\sup_{u \in [0, \delta]} \left\| Z^{\lambda, u}(r) - \hat{Z}_r^{\lambda, n, u} \right\|^4 ds \right] + \frac{Ch^4}{(1-\delta)^4} \\ &\quad + Ch^3 \gamma^4 \left(1 + \frac{t}{(1-\delta)^4} + \frac{t^2}{\lambda^2} \right) e^{C\gamma^4 t} + \frac{Ch^2}{\lambda^2}, \end{aligned}$$

where we used (4.17) in the last step. Finally, applying Grönwall's inequality and taking $0 < \gamma < 1$, we have the result of the lemma. \square

References

- [1] A. Ahmadi-Javid. Entropic value-at-risk: A new coherent risk measure. *J Optimiz Theory App*, 155(3):1105–1123, 2012.
- [2] Paul Embrechts Alexander J. McNeil and Rüdiger Frey. *Quantitative Risk Management*. Princeton University Press, 2015.
- [3] Zeyuan Allen-Zhu. Natasha 2: Faster non-convex optimization than sgd. *arXiv preprint arXiv:1708.08694*, 2017.
- [4] Philippe Artzner, Freddy Delbaen, Jean Marc Eber, and David Heath. Coherent measures of risk. *Math. Finance*, 9:203–228, 1999.
- [5] Daniel Bartl and Ludovic Tangp. Non-asymptotic rates for the estimation of risk measures. *arXiv:2003.10479*, 17, 2020.
- [6] Daniel Bartl, Samuel Drapeau, and Ludovic Tangpi. Computational aspects of robust optimized certainty equivalents and option pricing. *Math. Finance*, 30:287–309, 2020.
- [7] Denis Belomestny and Volker Krätschmer. Central limit theorems for law-invariant coherent risk measures. *J Appl Probab*, 49(1):1–21, 2012.
- [8] D. P. Bertsekas and S. E Shreve. *Stochastic Optimal Control. The discrete time case*, volume 139 of *Mathematics in Science and Engineering*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers, New York, London, 1978.
- [9] Chandrasekaran Anirudh Bhardwaj. Adaptively preconditioned stochastic gradient langevin dynamics. *arXiv preprint arXiv:1906.04324*, 2019.
- [10] François Bolley, Ivan Gentil, and Armand Guillin. Convergence to equilibrium in Wasserstein distance for Fokker-Planck equations. *J. Funct. Anal.*, 263(8):2430–2457, 2012.
- [11] Hans Bühler, Lukas Gonon, Josef Teichmann, and Ben Wood. Deep hedging. *Quant Financ*, 19(8):1271–1291, 2019.
- [12] JS Butler and Barry Schachter. Estimating value-at-risk with a precision measure by combining kernel estimation with historical simulation. *Rev. Deriv. Res.*, 1:371–390, 1997.
- [13] René Carmona and François Delarue. *Probabilistic theory of mean field games with applications. I*, volume 83 of *Probability Theory and Stochastic Modelling*. Springer, Cham, 2018. ISBN 978-3-319-56437-1; 978-3-319-58920-6. Mean field FBSDEs, control, and games.
- [14] Mu-Fa Chen and Shao-Fu Li. Coupling methods for multidimensional diffusion processes. *The Ann. Probab.*, pages 151–177, 1989.
- [15] Song Xi Chen. Nonparametric estimation of expected shortfall. *J Financ Economet*, 6(1):87–107, 2008.

- [16] Tianqi Chen, Emily Fox, and Carlos Guestrin. Stochastic gradient hamiltonian monte carlo. In *ICML*, pages 1683–1691. PMLR, 2014.
- [17] Patrick Cheridito and Tianhui Li. Risk measures on Orlicz hearts. *Math. Finance*, 19(2):189–214, 2009.
- [18] Patrick Cheridito, Freddy Delbaen, and Michael Kupper. Dynamic monetary risk measures for bounded discrete-time processes. *Electron. J. Probab.*, 11(3):57–106, 2006.
- [19] Rama Cont, Romain Deguest, and Giacomo Scandolo. Robustness and sensitivity analysis of risk measurement procedures. *Quantitative finance*, 10(6):593–606, 2010.
- [20] Freddy Delbaen. *Monetary Utility Functions*. Osaka University Press, 2012.
- [21] Wei Deng, Guang Lin, and Faming Liang. A contour stochastic gradient langevin dynamics algorithm for simulations of multi-modal distributions. *arXiv preprint arXiv:2010.09800*, 2020.
- [22] H. Djellout, A. Guillin, and L. Wu. Transportation cost-information inequalities and applications to random dynamical systems and diffusions. *Ann. Probab.*, 32(3B):2702–2732, 2004.
- [23] Alain "Durmus, Szymon Majewski, and Blażej" Miasojedow. Analysis of langevin monte carlo via convex optimization. *J. Mach. Learn. Res.*, 20(1):2666–2711, 2019.
- [24] Andreas Eberle. Reflection coupling and wasserstein contractivity without convexity. *C. R. Math*, 349(19-20): 1101–1104, 2011.
- [25] Andreas Eberle. Reflection couplings and contraction rates for diffusions. *Probab. Theory Relat. Fields*, 166(3): 851–886, 2016.
- [26] Stephan Eckstein and Michael Kupper. Computation of optimal transport and related hedging problems via penalization and neural networks. *Forthcoming in Appl. Math. Optim.*, 2019.
- [27] Jianqing Fan and Juan Gu. Semiparametric estimation of value at risk. *Econom. J.*, 6(2):261–290, 2003.
- [28] Ky Fan. Minimax theorems. *PNAS USA*, 39(1):42, 1953.
- [29] Hans Föllmer and T. Knispel. Entropic risk measures: Coherence vs. convexity, model ambiguity and robust large deviations. *Stoch. Dyn.*, 11(02n03):333–351, 2011.
- [30] Hans Föllmer and Alexander Schied. Convex measures of risk and trading constraint. *Financ Stoch*, 6(4):429–447, 2002.
- [31] Hans Föllmer and Alexander Schied. *Stochastic Finance. An Introduction in Discrete Time*. de Gruyter Studies in Mathematics. Walter de Gruyter, Berlin, New York, 2 edition, 2004.
- [32] Hans Föllmer and Alexander Schied. *Stochastic finance*. de Gruyter, 2016.
- [33] Noufel Frikha and Stéphane Menozzi. Concentration bounds for stochastic approximations. *Electron. Commun. Probab.*, 17, 2012.
- [34] Marco Frittelli and Emanuela Rosazza Gianin. Law invariant convex risk measures. *Adv. Math. Econ.*, pages 33–46, 2005.
- [35] Marco Frittelli and Emanuela Rosazza Gianin. Putting order in risk measures. *Journal of Banking & Finance*, 26(7):1473–1486, July 2002.
- [36] Saul B Gelfand and Sanjoy K Mitter. Recursive stochastic algorithms for global optimization in \mathbb{R}^d . *SIAM J Control Optim*, 29(5):999–1018, 1991.
- [37] Paul Glasserman, Philip Heidelberger, and Perwez Shahabuddin. Variance reduction techniques for estimating value-at-risk. *Manag. Sci*, 46(10):1349–1364, 2000.
- [38] John M Holte. Discrete gronwall lemma and applications. In *MAA-NCS meeting at the University of North Dakota*, volume 24, pages 1–7, 2009.

- [39] L Jeff Hong and Guangwu Liu. Monte carlo estimation of value-at-risk, conditional value-at-risk and their sensitivities. In *Proc. 2011 WSC*, pages 95–107. IEEE, 2011.
- [40] Lennart Hoogerheide and Herman K van Dijk. Bayesian forecasting of value at risk and expected shortfall using adaptive importance sampling. *Int. J. Forecast.*, 26(2):231–247, 2010.
- [41] Kaitong Hu, Shenjie Ren, David Siska, and Lukasz Szpruch. Mean-field Langevin dynamics and energy landscape of neural networks. *Annales de l’Institut Henri Poincaré (B) Probabilités and Statistiques*, to appear, 2020.
- [42] C. R. Hwang. Laplace’s method revisited: weak convergence of probability measures. *Ann. Probab.*, pages 2189–2211, 1980.
- [43] Garud Iyengar and Alfred Ka Chun Ma. Fast gradient descent method for mean-cvar optimization. *Ann. Oper. Res.*, 205(1):203–212, 2013.
- [44] Elyès Jouini, Walter Schachermayer, and Nizar Touzi. Law invariant risk measures have the fatou property. In Shigeo Kusuoka and Akira Yamazaki, editors, *Advances in Mathematical Economics*, volume 9 of *Advances in Mathematical Economics*, pages 49–71. Springer Japan, 2006.
- [45] P. E. Kloeden and E. Platen. *Numerical solution of stochastic differential equations*. Springer Science and Business Media, 2013.
- [46] Volker Krättschmer, Alexander Schied, and Henryk Zähle. Comparative and quantitative robustness for law-invariant risk measures. *Finance Stoch.*, 2014.
- [47] Michael Kupper and Walter Schachermayer. Representation results for law invariant time consistent functions. *Math. Financ. Econ.*, 2(3):189–210, 2009.
- [48] Shigeo Kusuoka. On law invariant coherent risk measures. In *Advances in mathematical economics*, pages 83–95. Springer, 2001.
- [49] D. Lacker, M. Shkolnikov, and J. Zhang. Inverting the markovian projection, with an application to local stochastic volatility models. *Ann. Probab.*, 48(5):2189–2211, 2020.
- [50] Florent Malrieu and Denis Talay. Concentration inequalities for Euler schemes. *Monte Carlo and quasi-Monte Carlo methods*, 2004(355-371), 2006.
- [51] Stéphane Menozzi and Vincent Lemaire. On some non asymptotic bounds for the euler scheme. *Electron. J. Probab.*, 15:1645–1681, 2010.
- [52] Yu Nesterov. Smooth minimization of non-smooth functions. *Math. Program.*, 103(1):127–152, 2005.
- [53] Basel Committee on Banking Supervision. Fundamental review of the trading book: A revised market risk framework. Technical report, Bank of international settlements, 2014.
- [54] Alois Pichler and R. Schlotter. Entropic based risk measures. *European Journal on Operational Research*, 285(1): 223–236, 2020.
- [55] Mathias Pohl, Alexander Ristig, Walter Schachermayer, and Ludovic Tangpi. Theoretical and empirical analysis of trading activity. *Math. Program.*, 181(2):405–434, 2020.
- [56] Maxim Raginsky, Alexander Rakhlin, and Matus Telgarsky. Non-convex learning via stochastic gradient langevin dynamics: a nonasymptotic analysis. In *COLT*, pages 1674–1703. PMLR, 2017.
- [57] A Max Reppen and H Mete Soner. Bias-variance trade-off and overlearning in dynamic decision problems. *arXiv preprint arXiv:2011.09349*, 2020.
- [58] Ralph Rockafellar. Integrals which are convex functionals. *Pacific journal of mathematics*, 24(3):525–539, 1968.
- [59] Sotirios Sabanis and Ying Zhang. A fully data-driven approach to minimizing cvar for portfolio of assets via sgld with discontinuous updating. *arXiv preprint arXiv:2007.01672*, 2020.

- [60] Ken Sekimoto. *Stochastic energetics*, volume 799. Springer, 2010.
- [61] Tasuku Soma and Yuichi Yoshida. Statistical learning with conditional value at risk. *arXiv preprint arXiv:2002.05826*, 2020.
- [62] Aviv Tamar, Yonatan Glassner, and Shie Mannor. Optimizing the cvar via sampling. In *AAAI-15*, 2015.
- [63] C. Villani. *Optimal Transport. Old and New*, volume 338 of *Grundlehren der mathematischen Wissenschaften*. Springer, 2009.
- [64] Stefan Weber. Distribution-invariant risk measures, Entropy, and large deviations. *J. Appl. Prob.*, 44:16–40, 2007.
- [65] Max Welling and Yee W. Teh. Bayesian learning via stochastic gradient langevin dynamics. *Proc. ICML-11*, pages 681–688, 2011.
- [66] R. W. Yeung. *Information theory and network coding*. Springer Science and Business Media, 2008.
- [67] Helin Zhu and Enlu Zhou. Estimation of conditional value-at-risk for input uncertainty with budget allocation. In *Proc. 2015 WSC*, pages 655–666. IEEE, 2015.

,