

Tight bounds on the expected number of holes in random point sets

Martin Balko^{1*} Manfred Scheucher^{2†} Pavel Valtr^{1‡}

¹ Department of Applied Mathematics,
Faculty of Mathematics and Physics, Charles University, Czech Republic
balko@kam.mff.cuni.cz

² Institut für Mathematik, Technische Universität Berlin, Germany
scheucher@math.tu-berlin.de

Abstract

For integers $d \geq 2$ and $k \geq d + 1$, a k -hole in a set S of points in general position in \mathbb{R}^d is a k -tuple of points from S in convex position such that the interior of their convex hull does not contain any point from S . For a convex body $K \subseteq \mathbb{R}^d$ of unit d -dimensional volume, we study the expected number $EH_{d,k}^K(n)$ of k -holes in a set of n points drawn uniformly and independently at random from K .

We prove an asymptotically tight lower bound on $EH_{d,k}^K(n)$ by showing that, for all fixed integers $d \geq 2$ and $k \geq d + 1$, the number $EH_{d,k}^K(n)$ is at least $\Omega(n^d)$. For some small holes, we even determine the leading constant $\lim_{n \rightarrow \infty} n^{-d} EH_{d,k}^K(n)$ exactly. We improve the currently best known lower bound on $\lim_{n \rightarrow \infty} n^{-d} EH_{d,d+1}^K(n)$ by Reitzner and Temesvari (2019). In the plane, we show that the constant $\lim_{n \rightarrow \infty} n^{-2} EH_{2,k}^K(n)$ is independent of K for every fixed $k \geq 3$ and we compute it exactly for $k = 4$, improving earlier estimates by Fabila-Monroy, Huemer, and Mitsche (2015) and by the authors (2020).

1 Introduction

For a positive integer d , let S be a set of points from \mathbb{R}^d in *general position*. That is, for every $k \in \{1, \dots, d - 1\}$, no $k + 2$ points from S lie on a k -dimensional affine subspace

*M. Balko was supported by the grant no. 21-32817S of the Czech Science Foundation (GAČR), by the Center for Foundations of Modern Computer Science (Charles University project UNCE/SCI/004), and by the PRIMUS/17/SCI/3 project of Charles University. This article is part of a project that has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 810115).

†M. Scheucher was supported by the DFG Grants SCHE 2214/1-1 and FE 340/12-1, and by the internal research funding “Post-Doc-Funding” from Technische Universität Berlin. We also acknowledge support by the internal research program IFFP 2016–2020 of the FernUniversität in Hagen.

‡P. Valtr was supported by the grant no. 21-32817S of the Czech Science Foundation (GAČR) and by the PRIMUS/17/SCI/3 project of Charles University.

of \mathbb{R}^d . Throughout the whole paper we only consider point sets in \mathbb{R}^d that are finite and in general position.

A point set P is in *convex position* if no point from P is contained in the convex hull of the remaining points from P . For an integer $k \geq d + 1$, a k -hole H in S is a set of k points from S in convex position such that the convex hull $\text{conv}(H)$ of H does not contain any point of S in its interior.

The study of k -holes in point sets was initiated by Erdős [Erd78], who asked whether, for each $k \in \mathbb{N}$, every sufficiently large point set in the plane contains a k -hole. This was known to be true for $k \leq 5$, but, in the 1980s, Horton [Hor83] constructed arbitrarily large point sets without 7-holes. The question about the existence of 6-holes was a longstanding open problem until 2007, when Gerken [Ger08] and Nicolas [Nic07] showed that every sufficiently large set of points in the plane contains a 6-hole.

The existence of k -holes was considered also in higher dimensions. Valtr [Val92] showed that, for $k \leq 2d + 1$, every sufficiently large set of points in \mathbb{R}^d contains a k -hole. He also constructed arbitrarily large sets of points in \mathbb{R}^d that do not contain any k -hole with $k > 2^{d-1}(P(d-1) + 1)$, where $P(d-1)$ denotes the product of the first $d-1$ prime numbers. Very recently Bukh, Chao, and Holzman [BCH20] improved this construction and Colon and Li [CL21] constructed new sets in \mathbb{R}^d with no large holes.

Estimating the number of k -holes in point sets in \mathbb{R}^d attracted a lot of attention; see [ABH⁺20]. In particular, it is well-known that the minimum number of $(d+1)$ -holes (also called *empty simplices*) in sets of n points in \mathbb{R}^d is of order $O(n^d)$. This is tight, as every set of n points in \mathbb{R}^d contains at least $\binom{n-1}{d}$ $(d+1)$ -holes [BF87, KM88].

The tight upper bound $O(n^d)$ can be obtained by considering random point sets drawn from a convex body. More formally, a *convex body* in \mathbb{R}^d is a compact convex subset of \mathbb{R}^d with a nonempty interior. We use λ_d to denote the d -dimensional Lebesgue measure on \mathbb{R}^d and \mathcal{K}_d to denote the set of all convex bodies in \mathbb{R}^d of volume $\lambda_d(K) = 1$. For an integer $k \geq d + 1$ and a convex body $K \in \mathcal{K}_d$, let $EH_{d,k}^K(n)$ be the expected number of k -holes in a set S of n points chosen uniformly and independently at random from K . Note that S is in general position with probability 1.

Bárány and Füredi [BF87] proved the upper bound $EH_{d,d+1}^K(n) \leq (2d)^{2d^2} \cdot \binom{n}{d}$ for every $K \in \mathcal{K}_d$. Valtr [Val95] improved this bound in the plane by showing $EH_{2,3}^K(n) \leq 4 \binom{n}{2}$ for any $K \in \mathcal{K}_2$. Very recently, Reitzner and Temesvári [RT19, Theorem 1.4] showed that this bound on $EH_{2,3}^K(n)$ is asymptotically tight for every $K \in \mathcal{K}_2$. This follows from their more general bounds

$$\lim_{n \rightarrow \infty} n^{-2} EH_{2,3}^K(n) = 2 \tag{1}$$

and

$$\frac{2}{d!} \leq \lim_{n \rightarrow \infty} n^{-d} EH_{d,d+1}^K(n) \leq \frac{d}{(d+1)} \frac{\kappa_{d-1}^{d+1} \kappa_d^2}{\kappa_d^{d-1} \kappa_{(d-1)(d+1)}} \tag{2}$$

for $d \geq 2$, where $\kappa_d = \pi^{\frac{d}{2}} \Gamma(\frac{d}{2} + 1)^{-1}$ is the volume of the d -dimensional Euclidean unit ball. Moreover, the upper bound in (2) holds with equality in the case $d = 2$, and if K is a d -dimensional ellipsoid with $d \geq 3$. Note that, by (2), there are absolute positive

constants c_1, c_2 such that

$$d^{-c_1 d} \leq \lim_{n \rightarrow \infty} n^{-d} EH_{d,d+1}^K(n) \leq d^{-c_2 d}$$

for every $d \geq 2$ and $K \in \mathcal{K}_d$.

Considering general k -holes in random point sets in \mathbb{R}^d , the authors [BSV21] recently proved that $EH_{d,k}^K(n) \leq O(n^d)$ for all fixed integers $d \geq 2$ and $k \geq d+1$ and every $K \in \mathcal{K}_d$. More precisely, we showed

$$EH_{d,k}^K(n) \leq 2^{d-1} \cdot \left(2d^{2d-1} \binom{k}{\lfloor d/2 \rfloor} \right)^{k-d-1} \cdot \frac{n(n-1) \cdots (n-k+2)}{(k-d-1)! \cdot (n-k+1)^{k-d-1}}. \quad (3)$$

In this paper, we also study the expected number $EH_{d,k}^K(n)$ of k -holes in random sets of n points in K . In particular, we derive a lower bound that asymptotically matches the upper bound (3) for all fixed values of k . Moreover, for some small holes, we even determine the leading constants $\lim_{n \rightarrow \infty} n^{-d} EH_{d,k}^K(n)$.

2 Our Results

First, we show that for all fixed integers $d \geq 2$ and $k \geq d+1$ the number $EH_{d,k}^K(n)$ is in $\Omega(n^d)$, which matches the upper bound (3) by the authors [BSV21] up to the leading constant.

Theorem 1. *For all integers $d \geq 2$ and $k \geq d+1$, there are constants $C = C(d, k) > 0$ and $n_0 = n_0(d, k)$ such that, for every integer $n \geq n_0$ and every convex body $K \subseteq \mathbb{R}^d$ of unit volume, we have $EH_{d,k}^K(n) \geq C \cdot n^d$.*

In particular, we see that random point sets typically contain many k -holes no matter how large k is, as long as it is fixed. This contrasts with the fact that, for every $d \geq 2$, there is a number $t = t(d)$ and arbitrarily large sets of points in \mathbb{R}^d without any t -holes [Hor83, Val92].

Theorem 1 together with (3) shows that $EH_{d,k}^K(n) = \Theta(n^d)$ for all fixed integers d and k and every $K \in \mathcal{K}_d$, which determines the asymptotic growth rate of $EH_{d,k}^K(n)$. We thus focus on determining the leading constants $\lim_{n \rightarrow \infty} n^{-d} EH_{d,k}^K(n)$ for small holes.

For a convex body $K \subseteq \mathbb{R}^d$ (of a not necessarily unit volume), we use p_d^K to denote the probability that the convex hull of $d+2$ points chosen uniformly and independently at random from K is a d -simplex. That is, the probability that one of the $d+2$ points falls in the convex hull of the remaining $d+1$ points. The problem of computing p_d^K is known as the d -dimensional *Sylvester's convex hull problem* for K and it has been studied extensively. Let $p_d = \max_K p_d^K$, where the maximum is taken over all convex bodies $K \subseteq \mathbb{R}^d$. We note that the maximum is achieved, since it is well-known that every affine-invariant continuous functional on the space of convex bodies attains a maximum.

First, we prove the following lower bound on the expected number $EH_{d,d+1}^K(n)$ of empty simplices in random sets of n points in K , which improves the lower bound from (2) by Reitzner and Temesvari [RT19] by a factor of d/p_{d-1} .

Theorem 2. For every integer $d \geq 2$ and every convex body $K \subseteq \mathbb{R}^d$ of unit volume, we have

$$\lim_{n \rightarrow \infty} n^{-d} EH_{d,d+1}^K(n) \geq \frac{2}{(d-1)! p_{d-1}}.$$

Using the trivial fact $p_1 = 1$ with the inequality $EH_{2,3}^K(n) \leq 2(1 + o(1))n^2$ proved by Valtr [Val95], we see that the leading constant in our estimate is asymptotically tight in the planar case. Results of Blaschke [Bla17, Bla23] (see Theorem 15) and Groemer [Gro73] (see Theorem 16) imply the following bounds showing that, in any dimension $d \geq 3$, the leading constant in $EH_{d,d+1}^K(n)$ depends on the convex body.

Corollary 3. For every $d \geq 3$, every d -dimensional simplex S of unit volume fulfills

$$\lim_{n \rightarrow \infty} n^{-d} EH_{d,d+1}^S(n) < \lim_{n \rightarrow \infty} n^{-d} EH_{d,d+1}^B(n),$$

where B is a d -dimensional ball of unit volume. Moreover, for every convex body $K \subseteq \mathbb{R}^3$ of unit volume, we have

$$3 \leq \lim_{n \rightarrow \infty} n^{-3} EH_{3,4}^K(n) \leq \frac{12\pi^2}{35} \approx 3.38386.$$

We performed computer experiments to obtain estimates on the constants $c_{3,4}^K = \lim_{n \rightarrow \infty} n^{-3} EH_{3,4}^K(n)$ when K is a Platonic solid or a ball. The idea was to repeatedly sample 3 points p_1, p_2, p_3 uniformly at random from K and to compute the ratio of the so-called *Fischer triangle* spanned by p_1, p_2, p_3 which is contained inside K . This ratio coincides with $c_{3,4}^K$; see Section 3 in [BSV21]. Since this ratio is a realization of an independent and identically distributed random variable, the sample average is normally distributed by the central limit theorem. After evaluating 16 million samples by our program, we now have a statistical evidence that

- $3.095 \leq c_{3,4}^T \leq 3.097$ for a regular tetrahedron T ,
- $3.264 \leq c_{3,4}^C \leq 3.266$ for a cube C ,
- $3.289 \leq c_{3,4}^O \leq 3.291$ for a regular octahedron O ,
- $3.354 \leq c_{3,4}^D \leq 3.356$ for a regular dodecahedron D , and
- $3.362 \leq c_{3,4}^I \leq 3.364$ for a regular icosahedron I .

Moreover, our program gave a statistical evidence that $3.383 \leq c_{3,4}^B \leq 3.385$ for a ball B , which for dimension 2 confirms the constant by Reitzner and Temesvari [RT19]; see the upper bound in (2). All the bounds are with confidence level 99.99%. The source code of our program and more information is available on the supplemental website [Sch].

By Theorem 2, better upper bounds on p_{d-1} imply stronger lower bounds on $EH_{d,d+1}^K(n)$. The problem of estimating p_d is equivalent to the problem of estimating the expected d -dimensional volume EV_d^K of the convex hull of $d+1$ points drawn from

a convex body $K \subseteq \mathbb{R}^d$ uniformly and independently at random, since $p_d^K = \frac{(d+2)EV_d^K}{\lambda_d(K)}$; see [Kle69, Sch88]. In the plane, Blaschke [Bla17, Bla23] showed that EV_2^K is maximized if K is a triangle (see Theorem 15), which we use in Section 4 to derive the lower bound in Corollary 3. For $d \geq 3$, it is one of the major problems in convex geometry to decide whether EV_d^K is maximized if K is a simplex [SW08]. Meckes [Mec04] proved that there is a constant $c > 0$ such that every convex body $K \in \mathcal{K}_d$ with $d \geq 2$ satisfies

$$EV_d^K \leq (cd^{-1/4} \log d)^d,$$

which by Theorem 2 implies the following improved lower bound on $EH_{d,d+1}^K(n)$.

Corollary 4. *There is a constant $c > 0$ such that for every integer $d \geq 2$ and every convex body $K \subseteq \mathbb{R}^d$ of unit volume, we have*

$$\lim_{n \rightarrow \infty} n^{-2} EH_{d,d+1}^K(n) \geq \frac{2}{(d+2)(d-1)!} \left(\frac{d^{1/4}}{c \log d} \right)^d.$$

We also note that there are connections between the probabilities p_d , the famous *Slicing conjecture* (also known as *Hyperplane conjecture*), and the *isotropic constant*; see for example [Mec04].

For convex bodies K with small diameter, we can obtain the following better estimate on p_d^K .

Proposition 5. *Let $\varepsilon > 0$ be a real number and let $d \geq 1$ be an integer. Let $K \subseteq \mathbb{R}^d$ be a convex body. If there is a volume-preserving affine transformation $f: \mathbb{R}^d \rightarrow \mathbb{R}^d$ such that $f(K)$ has diameter D , then $p_d^K \leq \frac{(d+2)D^d}{d! \lambda_d(K)}$. In particular, if $D \leq d^{1-\varepsilon} \lambda_d(K)^{1/d}$, then $p_d^K \leq \frac{(d+2)d^{(1-\varepsilon)d}}{d!}$.*

We note that there are convex bodies that do not satisfy the assumption from Proposition 5, for example the regular d -dimensional simplex.

Besides empty simplices, we also consider larger k -holes. The expected number $EH_{2,4}^K(n)$ of 4-holes in random planar sets of n points was considered by Fabila-Monroy, Huemer, and Mitsche [FMHM15], who showed

$$EH_{2,4}^K(n) \leq 18\pi D^2 n^2 + o(n^2)$$

for any $K \in \mathcal{K}_2$, where $D = D(K)$ is the diameter of K . Since we have $D \geq 2/\sqrt{\pi}$, by the Isodiametric inequality [EG15], the leading constant in their bound is at least 72 for any $K \in \mathcal{K}_2$. This result was strengthened by the authors [BSV21] to $EH_{2,4}^K(n) \leq 12n^2 + o(n^2)$ for every $K \in \mathcal{K}_2$. Here, we determine the leading constant in $EH_{2,4}^K(n)$ exactly.

Theorem 6. *For every convex body $K \subseteq \mathbb{R}^2$ of unit area, we have*

$$\lim_{n \rightarrow \infty} n^{-2} EH_{2,4}^K(n) = 10 - \frac{2\pi^2}{3} \approx 3.42026.$$

Our computer experiments [Sch] support this result and also equation (1). We sampled random sets of $n = 25000$ points from a triangle, a square, and a disk, and for each shape the average number of 3-holes and 4-holes was around $2n^2$ and $3.42n^2$, respectively. Moreover, the average number of 5-holes and 6-holes was around $3.4n^2$ and $3n^2$, respectively. It may turn out that the expected numbers of 4-holes and 5-holes coincide asymptotically. Note that all these bounds for random point sets are worse than the ones obtained by the squared Horton sets of size n , which asymptotically contain $c_k n^2$ k -holes for $k \in \{3, \dots, 6\}$, where $c_3 \approx 1.6195$, $c_4 \approx 1.9396$, $c_5 \approx 1.0206$, and $c_6 \approx 0.2005$ [BV04].

For larger k -holes in the plane, we do not determine the value $\lim_{n \rightarrow \infty} n^{-2} EH_{2,k}^K(n)$ exactly, but we can show that it exists and does not depend on the convex body K . We recall that this is not true in larger dimensions already for empty simplices by Corollary 3.

Theorem 7. *For every integer $k \geq 3$, there is a constant $C = C(k)$ such that, for every convex body $K \subseteq \mathbb{R}^2$ of unit area, we have*

$$\lim_{n \rightarrow \infty} n^{-2} EH_{2,k}^K(n) = C.$$

The idea of the proof of Theorem 1 is novel and the proof is presented in Section 3. The proof of Theorem 2 is a combination of ideas used by Reitzner and Temesvari [RT19] and of a result by Blaschke [Bla17, Bla23]. The proof can be found in Section 4, where we also prove Proposition 5 using simple geometric arguments. Finally, Theorems 6 and 7 are proved in Sections 5 and 6, respectively. Both these proofs combine arguments used by Reitzner and Temesvari [RT19] and by Valtr [Val95].

Open problems We determined the leading constants $\lim_{n \rightarrow \infty} n^{-d} EH_{d,k}^K(n)$ exactly for small holes and, in particular, we showed that these limits exist in such cases. However, we do not have any argument that would yield the existence of these limits for all values of d and k . It is thus an interesting open problem to determine whether $\lim_{n \rightarrow \infty} n^{-d} EH_{d,k}^K(n)$ exists for all positive integers d and k with $k \geq d + 1$. It follows from a result by Reitzner and Temesvari [RT19, Theorem 1.4] and from Theorem 7 that this limit exists if $k = d + 1$ or if $k \geq 3$ and $d = 2$, respectively.

We also pose the following problem asking for the value of the expected number of empty simplices in a tetrahedron.

Problem 1. *Let K be a 3-dimensional simplex of unit volume. Determine the leading constant $\lim_{n \rightarrow \infty} n^{-d} EH_{3,4}^K(n)$.*

It might also be interesting to determine $\lim_{n \rightarrow \infty} n^{-2} EH_{2,k}^K(n)$ exactly for as many values $k > 4$ as possible. Recall that, by Theorem 7, the number $\lim_{n \rightarrow \infty} n^{-2} EH_{2,k}^K(n)$ is the same for all convex bodies $K \in \mathcal{K}_2$.

3 Proof of Theorem 1

We show that, for all integers $d \geq 2$ and $k \geq d + 1$, there are constants $C = C(d, k) > 0$ and $n_0 = n_0(d, k)$ such that, for every $n \geq n_0$ and every convex body $K \in \mathcal{K}_d$, we have

$EH_{d,k}^K(n) \geq C \cdot n^d$. In our proof, we make no serious attempt to optimize the constants. We start by stating some auxiliary results and definitions.

We use B^d to denote the d -dimensional ball of radius 1 centered at the origin and S^{d-1} to denote the sphere that is the boundary of B^d . For $h \in [0, 1]$, let $C(h) = \{(x_1, \dots, x_d) \in B^d : x_1 \geq 1 - h\}$ be the *spherical cap* of S^{d-1} of height h . For $x \in S^{d-1}$, we use $C(h, x)$ to denote the image of $C(h)$ under a rotation r of S^{d-1} around the origin such that $r(1, 0, \dots, 0) = x$. We call the set $C(h, x)$ the *spherical cap centered at x* . Note that $C(h, x)$ and $C(h)$ have the same height, volume, and surface area for every $x \in S^{d-1}$.

We first give a high-level overview of the proof. For a given convex body $K \in \mathcal{K}_d$, we choose a sufficiently large $(d-1)$ -dimensional ellipsoid S'' contained in K . We then select d points q_1, \dots, q_d on S'' and we consider an ε -neighborhood N_i of q_i for every $i = 1, \dots, d$ for a sufficiently small constant $\varepsilon = \varepsilon(d) > 0$. When we draw a set S of n points uniformly and independently at random from K , some d -tuple p_1, \dots, p_d of the points from S satisfies $p_i \in N_i$ with positive probability. It then remains to show that the points p_1, \dots, p_d are contained in a k -hole in S with positive probability. To do that, we use the choice of S'' , which enables us to choose a $(d-1)$ -dimensional ellipsoid $F \subseteq K$ containing the points p_1, \dots, p_d . Moreover, since each p_i is close to S'' , the choice of S'' also guarantees the existence of a d -dimensional ellipsoid $G \subseteq K$ of volume $1/(n-d)$ such that $F \subseteq G$. We then choose $k-d$ pairwise disjoint caps C_1, \dots, C_{k-d} of G , each with volume at least c for some constant $c = c(d, k) > 0$, and we show that, with positive probability, there is a point p_{i+d} of S inside each cap C_i and there is no point of P inside $G \setminus (C_1 \cup \dots \cup C_{k-d})$. The points p_1, \dots, p_k then determine a k -hole in S with positive probability.

Although the main idea of the proof of Theorem 1 is relatively simple, there are many technical steps along the way and we need to prove several auxiliary results. As the first such result, we use the following lower bound on the volume of spherical caps.

Lemma 8 ([MV10]). *For every integer $d \geq 2$ and $h \in (0, 1]$, we have*

$$\lambda_d(C(h)) > \left(\sqrt{2h - h^2}\right)^{d-1} \cdot \frac{h}{2d} \cdot \lambda_d(B^d).$$

The following lemma is used to estimate the surface area of $C(h)$.

Lemma 9. *For every $h \in (0, 1/4]$, we have*

$$\lambda_{d-1}(C(h) \cap S^{d-1}) \leq (4\sqrt{h})^{d-1} \lambda_{d-1}(S^{d-1}).$$

Proof. Let \overline{C} be the convex hull of $C(h) \cup \{0\}$, where 0 denotes the origin. Then $\frac{\lambda_{d-1}(C(h) \cap S^{d-1})}{\lambda_{d-1}(S^{d-1})} = \frac{\lambda_d(\overline{C})}{\lambda_d(B^d)}$ and it thus suffices to find an upper bound on $\lambda_d(\overline{C})$. The basis of $C(h)$ on S^{d-1} is a $(d-1)$ -dimensional ball of radius $\sqrt{2h - h^2}$. Let B be the $(d-1)$ -dimensional ball with radius $\sqrt{2h - h^2}/(1-h)$ centered at $(1, 0, \dots, 0)$ lying in the hyperplane $x_1 = 1$. The set \overline{C} is then contained in the convex hull \overline{B} of $B \cup \{0\}$. Thus,

$$\lambda_d(\overline{C}) \leq \lambda_d(\overline{B}) = \frac{1}{d} \cdot \left(\frac{\sqrt{h(2-h)}}{1-h}\right)^{d-1} \cdot \lambda_{d-1}(B^{d-1}) \leq (4\sqrt{h})^{d-1} \lambda_d(B^d),$$

where we used $\sqrt{2-h}/(1-h) \leq 2$ for $h \leq 1/4$ together with a crude estimate $\lambda_{d-1}(B^{d-1}) \leq d2^{d-1}\lambda_d(B^d)$. We thus obtain $\lambda_{d-1}(C(h) \cap S^{d-1}) \leq (4\sqrt{h})^{d-1}\lambda_{d-1}(S^{d-1})$. \square

Using this lemma, we show that, for every $k \in \mathbb{N}$, we can place k pairwise disjoint spherical caps on S^{d-1} so that each cap has height that depends only on d and k .

Lemma 10. *For integers $d \geq 2$, $k \geq d$, and $h \in (0, 1/(64k^{2/(d-1)}))$, let x_1, \dots, x_d be points from S^{d-1} . Then, there are pairwise disjoint spherical caps $C(h, x_{d+1}), \dots, C(h, x_k)$ on B^d such that $x_i \notin C(h, x_j)$ for all $i = 1, \dots, d$ and $j = d+1, \dots, k$.*

Proof. We proceed by induction on k . The statement is trivial for $k = d$, so we assume that there are pairwise disjoint spherical caps $C(h, x_{d+1}), \dots, C(h, x_{k-1})$ for some $k \geq d+1$ and $h \in (0, 1/(64k^{2/(d-1)}]) \subseteq (0, 1/(64(k-1)^{2/(d-1)}])$ such that $x_i \notin C(h, x_j)$ for all $i = 1, \dots, d$ and $j = d+1, \dots, k-1$. Let $U = S^{d-1} \cap (\bigcup_{i=1}^{k-1} C(4h, x_i))$.

By Lemma 9, we have

$$\lambda_{d-1}(U) \leq (k-1)(8\sqrt{h})^{d-1}\lambda_{d-1}(S^{d-1}).$$

Since $h \leq 1/(64k^{2/(d-1)})$, we obtain $\lambda_{d-1}(U) < \lambda_{d-1}(S^{d-1})$ and thus there is a point $x_k \in S^{d-1} \setminus U$.

For every $h' \in [0, 1]$ and $x \in S^{d-1}$, the basis of each spherical cap $C(h', x)$ on S^{d-1} is a $(d-1)$ -dimensional ball of radius $\sqrt{2h' - (h')^2}$. Thus, the distance between x and any point of $C(h', x)$ is at most $\sqrt{2h' - (h')^2 + (h')^2} = \sqrt{2h'}$. On the other hand, every point of S^{d-1} at distance at most $\sqrt{2h'}$ from x lies in $C(h', x)$. Since $x_k \notin U$, the distance between x_k and x_i is larger than $\sqrt{2 \cdot 4h} = 2\sqrt{2h}$ for every $i = 1, \dots, k-1$. Also, any point $y \in C(h, x_k)$ is at distance larger than $2\sqrt{2h} - \sqrt{2h} = \sqrt{2h}$ from x_i and thus $y \notin C(h, x_i)$. Therefore, by the induction hypothesis, the spherical caps $C(h, x_{d+1}), \dots, C(h, x_k)$ are pairwise disjoint and $x_i \notin C(h, x_j)$ for all $i = 1, \dots, d$ and $j = d+1, \dots, k$. \square

For an integer $n > d$, a d -dimensional ellipsoid $D \subseteq K$ with $\lambda_d(\text{conv}(D)) = 1/(n-d)$, and points q_1, \dots, q_d on the boundary of D , we use $E(D, q_1, \dots, q_d)$ to denote the event that, for the set S of $n-d$ points drawn uniformly independently at random from K , there is a k -hole $\{p_1, \dots, p_k\}$ in $S \cup \{q_1, \dots, q_d\}$ satisfying $p_i = q_i$ for every $i = 1, \dots, d$.

Lemma 11. *There is $n_0 = n_0(d, k) \in \mathbb{N}$ such that for every integer $n \geq n_0$, every d -dimensional ellipsoid $D \subseteq K$ with $\lambda_d(\text{conv}(D)) = 1/(n-d)$, and any points q_1, \dots, q_d on the boundary of D , there is a constant $C = C(d, k) > 0$ such that the probability of the event $E(D, q_1, \dots, q_d)$ is at least C .*

Proof. Let $f: \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a one-to-one affine mapping such that $f(D) = B^d$. We set $h = 1/(64k^{2/(d-1)})$. By Lemma 10, there are pairwise disjoint spherical caps $C(h, f(q_{d+1})), \dots, C(h, f(q_k))$ on B^d such that $f(q_i) \notin C(h, f(q_j))$ for all $i = 1, \dots, d$ and $j = d+1, \dots, k$. We let $C'_i = f^{-1}(C(h, f(q_i)))$ for every $i = d+1, \dots, k$. We then fix the set $D' = f^{-1}(B^d \setminus (\bigcup_{i=d+1}^k C(h, f(q_i)))) = D \setminus (\bigcup_{i=d+1}^k C'_i)$.

Let S be the set of $n - d$ points drawn at random from K . Let $E(D', 0)$ be the event that no point from S lies in the set D' . By Lemma 8, the volume of each spherical cap $C(h, f(q_i))$ is at least $c \cdot \lambda_d(B^d)$ for some constant $c = c(d, k) > 0$. Since f preserves ratios of volumes, the volume of each set C'_i is at least $c/(n - d)$. It then follows from the fact that the spherical caps $C(h, f(q_{d+1})), \dots, C(h, f(q_k))$ are pairwise disjoint that the volume of D' is at most $\frac{1}{n-d} - (k-d)\frac{c}{n-d} = \frac{1-c(k-d)}{n-d}$. We may assume that $\frac{1-c(k-d)}{n-d} > 0$, as otherwise $D' = \emptyset$ and $\Pr[E(D', 0)] = 1$.

Since the volume of the set D' is at most $(1 - c(k - d))/(n - d)$, we have

$$\Pr[E(D', 0)] \geq \left(1 - \frac{1 - c(k - d)}{n - d}\right)^{n-d} \geq e^{-2+2c(k-d)} \geq e^{-2},$$

where we used the inequality $1 - x \geq e^{-2x}$ for $x \in [0, 1/2]$. Note that indeed $(1 - c(k - d))/(n - d) \in [0, 1/2]$ if n is sufficiently large with respect to d and k .

We partition the set S into $k - d$ sets B_{d+1}, \dots, B_k , each of size at least $\lfloor (n - d)/(k - d) \rfloor \geq (n - d)/(k - d) - 1 \geq 1$. For each $i \in \{d + 1, \dots, k\}$, let $E(D', i)$ be the event that no point from B_i appears in C'_i . Then

$$\begin{aligned} \Pr[E(D', i) \mid E(D', 0)] &= \left(\frac{1 - \lambda_d(D') - \lambda_d(C'_i)}{1 - \lambda_d(D')}\right)^{|B_i|} = \left(1 - \frac{\lambda_d(C'_i)}{1 - \lambda_d(D')}\right)^{|B_i|} \\ &\leq (1 - \lambda_d(C'_i))^{|B_i|} \leq \left(1 - \frac{c}{n - d}\right)^{\frac{n-d}{k-d}-1} \leq 2e^{-c/(k-d)}, \end{aligned}$$

since $\lambda_d(C'_i) \geq c/(n - d)$ and $1/(1 - \frac{c}{n-d}) \leq 2$ for n sufficiently large. Thus the event $\overline{E(D', i)}$ that there is a point from B_i lying in C'_i satisfies $\Pr[\overline{E(D', i)} \mid E(D', 0)] \geq 1 - 2e^{-c/(k-d)}$. Since the events $\overline{E(D', d+1)}, \dots, \overline{E(D', k)}$ are mutually independent, the conditioned probability that there is a point from B_i in C'_i for every $i \in \{d + 1, \dots, k\}$ satisfies

$$\Pr\left[\bigcap_{i=d+1}^k \overline{E(D', i)} \mid E(D', 0)\right] \geq \left(1 - 2e^{-c/(k-d)}\right)^{k-d}.$$

We now show that if the event $E(D', 0) \cap \left(\bigcap_{i=d+1}^k \overline{E(D', i)}\right)$ holds, then there is a k -hole $\{p_1, \dots, p_k\}$ in $S \cup \{q_1, \dots, q_d\}$ satisfying $p_i = q_i$ for every $i = 1, \dots, d$ and $p_j \in C'_j$ for every $j = d + 1, \dots, k$. That is, we show $E(D, q_1, \dots, q_d) \supseteq E(D', 0) \cap \left(\bigcap_{i=d+1}^k \overline{E(D', i)}\right)$. Since there is a point from B_i in C'_i for every $i = d + 1, \dots, k - d$, we can choose a point $p_i \in C'_i \cap S$ that is closest to D' among all points from $C'_i \cap S$. We also let $p_i = q_i$ for every $i = 1, \dots, d$. By the choice of the spherical caps $C(h, q_{d+1}), \dots, C(h, q_k)$, the points p_1, \dots, p_k are in convex position. Since each point p_i is the closest one to D' , there is no point from $S \cap C'_i$ in the interior of $\text{conv}(\{p_1, \dots, p_k\})$. Since there is no point from S in the set D' , we see that the points p_1, \dots, p_k form a desired k -hole in $S \cup \{q_1, \dots, q_d\}$.

It thus remains to estimate the probability of the event $E(D', 0) \cap \left(\bigcap_{i=d+1}^k \overline{E(D', i)} \right)$ from below. We have

$$\begin{aligned} \Pr \left[E(D', 0) \cap \left(\bigcap_{i=d+1}^k \overline{E(D', i)} \right) \right] &= \Pr \left[\bigcap_{i=d+1}^k \overline{E(D', i)} \mid E(D', 0) \right] \cdot \Pr [E(D', 0)] \\ &\geq \left(1 - 2e^{-c/(k-d)} \right)^{k-d} \cdot \frac{1}{e^2} > 0, \end{aligned}$$

which finishes the proof of Lemma 11. \square

For $\varepsilon > 0$, a d -dimensional simplex with vertices r_1, \dots, r_{d+1} is ε -regular, if there exists a regular d -dimensional simplex with vertices r'_1, \dots, r'_{d+1} such that r_i is at distance at most ε from r'_i for every $i = 1, \dots, d+1$.

The following auxiliary result states that, for every ε -regular simplex R with a sufficiently small $\varepsilon > 0$, there is a circumscribed ellipsoid of R with roughly the same diameter.

Lemma 12. *For an integer $d \geq 1$ and a real number $D > 0$, let $\varepsilon \in (0, D/(4\sqrt{2d(d+1)}))$. If R is a d -dimensional ε -regular simplex in \mathbb{R}^d with diameter D , then there is a number $c = c(d)$ and a circumscribed ellipsoid of R with diameter at most cD .*

Proof. Let r_1, \dots, r_{d+1} be the vertices of R . Since R is ε -regular, there is a regular simplex R' with vertices r'_1, \dots, r'_{d+1} such that r_i is at distance at most ε from r'_i for every $i = 1, \dots, d+1$. By the choice of R , the diameter of R' is at least $D - 2\varepsilon$ and most $D + 2\varepsilon$. In particular, the side length a of R' satisfies $a \in [D - 2\varepsilon, D + 2\varepsilon] \subseteq [D/2, 2D]$, as $\varepsilon \leq D/4$.

The distance of the barycenter of R' from its facets is $a\sqrt{\frac{d+1}{2d}} \cdot \frac{1}{d+1} = \frac{a}{\sqrt{2d(d+1)}}$. Thus R' contains an inscribed d -dimensional ball B of radius $\frac{a}{\sqrt{2d(d+1)}}$. Since $\varepsilon < \frac{D}{4\sqrt{2d(d+1)}}$ and $a \geq D/2$, we have $\varepsilon < \frac{a}{2\sqrt{2d(d+1)}}$. On the other hand, since every vertex of R' lies at distance $a\sqrt{\frac{d+1}{2d}} \cdot \frac{d}{d+1} = a\sqrt{\frac{d}{2(d+1)}}$ from the barycenter of R' , there is a circumscribed ball B' of R' with radius $a\sqrt{\frac{d}{2(d+1)}}$. Note that $B' = dB$.

Since R is an affine image of R' , there is a circumscribed ellipsoid E of R such that R contains an inscribed copy of $\frac{1}{d}E$. If $a_1 \leq \dots \leq a_d$ are the semi axes of E , then the volume V of E equals $\kappa_d(a_1 \cdots a_d)$, where κ_d is the volume of the d -dimensional unit ball. Clearly, the volume of R is at most $D^d/d!$. Since $\frac{1}{d}E \subseteq R \subseteq E$, we have $Vd^{-d} \leq \lambda_d(R) \leq V$ and, in particular, $V \leq d^d \lambda_d(R) \leq d^d D^d/d!$.

Since $B \subseteq R'$ and $\varepsilon < \frac{a}{2\sqrt{2d(d+1)}}$, it follows that R contains a ball of radius $r = \frac{a}{\sqrt{2d(d+1)}} - \varepsilon \geq \frac{a}{2\sqrt{2d(d+1)}}$ and thus $a_1, \dots, a_d \geq r$. The radius r is at least $\frac{D}{4\sqrt{2d(d+1)}}$, since $a \geq D/2$. Since $\kappa_d(a_1 \cdots a_d) = V \leq d^d D^d/d!$, we obtain

$$a_d \leq \frac{d^d D^d}{d! \cdot \kappa_d \cdot a_1 \cdots a_{d-1}} \leq \frac{d^d D^d}{d! \cdot \kappa_d \cdot r^{d-1}} \leq \frac{d^d (4\sqrt{2d(d+1)})^{d-1}}{d! \cdot \kappa_d} \cdot D,$$

where the second inequality follows from $a_1, \dots, a_{d-1} \geq r$ and the third one from $r \geq \frac{D}{4\sqrt{2d(d+1)}}$. Since $a_1 \leq \dots \leq a_d$, the diameter of E equals $2a_d \leq cD$, where $c = \frac{2d^d(4\sqrt{2d(d+1)})^{d-1}}{d! \cdot \kappa_d}$. \square

We will also need the following classical result, known as *John's lemma* [Joh48].

Lemma 13 ([Mat02, Theorem 13.4.1]). *For every d -dimensional convex body $K \subseteq \mathbb{R}^d$, there is a d -dimensional ellipsoid E with the center s satisfying*

$$E \subseteq K \subseteq s + d(E - s).$$

Finally, we proceed with the proof of Theorem 1. Let K be a d -dimensional convex body of unit volume and let $k \geq d+1$ be an integer. We set $\varepsilon > 0$ to be sufficiently small with respect to d .

By Lemma 13, there is a d -dimensional ellipsoid E with the center s that satisfies $E \subseteq K \subseteq s + d(E - s)$. In particular, $\lambda_d(E) \geq \lambda_d(K)/d^d = d^{-d}$. By applying a suitable affine volume preserving transformation, we can assume without loss of generality that E is a d -dimensional ball B of radius r with the center at the origin. Since $d^{-d} \leq \lambda_d(B) = \kappa_d r^d \leq \pi^{d/2} r^d$, we have $r \geq 1/(d\sqrt{\pi})$.

We let B' be the ball of radius $r' = r/2$ centered at the origin and B'' be the ball of radius $r'' = (r' - 3\varepsilon)/(2c + 1)$ centered at the origin, where $c = c(d)$ is the constant from Lemma 12. Since $3\varepsilon \leq 1/(4d\sqrt{\pi}) \leq r'/2 = r/4$ for $\varepsilon = \varepsilon(d)$ sufficiently small, we have $r'' \geq r/(8c + 4)$ and the volume of the ball B'' is at least $(4d(2c + 1))^{-d}$. Moreover, since $r' = r'' + 3\varepsilon + 2cr''$, all points lying at distance at most $2cr'' + 3\varepsilon$ from an arbitrary point of B'' lie in B' .

Let S'' be a $(d-1)$ -dimensional sphere obtained as the intersection of the boundary of B'' with the hyperplane $x_d = 0$ and let q_1, \dots, q_d be points of a regular $(d-1)$ -dimensional simplex R' inscribed to S'' . We let N_i be an ε -neighborhood of q_i for each $i = 1, \dots, p$. Note that $\lambda_d(N_i) = \varepsilon^d \lambda_d(B^d)$ and $N_i \subseteq B'$ for every $i = 1, \dots, d$.

Let p_1, \dots, p_n be the points drawn independently and uniformly at random from K . Assume that some distinct points p_{i_1}, \dots, p_{i_d} satisfy $p_{i_j} \in N_j$ for every $j = 1, \dots, d$. We will show that with positive probability that depends only on d and k , there is a k -hole in S containing the points p_{i_1}, \dots, p_{i_d} .

Let R be the $(d-1)$ -dimensional simplex with vertices p_{i_1}, \dots, p_{i_d} . Since $p_{i_j} \in N_j$ for every $j = 1, \dots, d$, the simplex R is ε -regular. Since $R' \subseteq B''$, the diameter of R is at most $2r'' + 2\varepsilon$ and at least $2r'' - 2\varepsilon \geq r/(8c + 4) - 2\varepsilon \geq 1/(d\sqrt{\pi}(8c + 4)) - 2\varepsilon > 4(\sqrt{2(d-1)d})\varepsilon$ for $\varepsilon = \varepsilon(d)$ sufficiently small. Thus, by Lemma 12, there is a circumscribed $(d-1)$ -dimensional ellipsoid F to R with diameter at most $2c(r'' + \varepsilon)$. Since the points p_{i_1}, \dots, p_{i_d} lie at distance at most ε from B'' and all points lying at distance at most $2cr'' + 3\varepsilon$ from an arbitrary point of B'' lie in B' , the triangle inequality gives $F \subseteq B'$ and, in particular, the diameter of F is at most $2r'$. Since $B' \subseteq K$, we have $F \subseteq K$ as well.

Since F is circumscribed to the ε -regular simplex R , F contains a $(d-1)$ -dimensional ball of radius at least t for some constant $t = t(d) > 0$ and thus the volume of F is at least t^{d-1} . Let G be a d -dimensional ellipsoid of d -dimensional volume $1/(n-d)$ such

that F is the intersection of G with a hyperplane. All points of G lie at distance at most $d/((n-d) \cdot \lambda_{d-1}(F))$ from F , since otherwise G contains the cone with basis F and height larger than $d/((n-d) \cdot \lambda_{d-1}(F))$, which has d -dimensional volume larger than $\frac{1}{d}d\lambda_{d-1}(F)/((n-d) \cdot \lambda_{d-1}(F)) = 1/(n-d)$. Since $\lambda_{d-1}(F) \geq t^{d-1}$, by choosing n sufficiently large with respect to d , we have $d/((n-d) \cdot \lambda_{d-1}(F)) \leq r/2$. Since all points at distance at most $r/2$ from B' lie in B and since $F \subseteq B'$, we see that the set of all points lying at distance at most $r/2$ from F lie in B . It follows that $G \subseteq B \subseteq K$.

By Lemma 11 applied to G and points p_{i_1}, \dots, p_{i_d} , the probability that there is a k -hole $\{p'_1, \dots, p'_k\}$ in S with $p'_j = p_{i_j}$ for $j = 1, \dots, d$ is at least C for some positive constant $C = C(d, k)$.

Let $E(i_1, \dots, i_d)$ be the event that $p_{i_j} \in N_j$ for every $j = 1, \dots, d$. The probability that $p_{i_j} \in N_j$ equals $\lambda_d(N_j) = (\varepsilon^d \lambda_d(B^d))$ and thus $\Pr[E(i_1, \dots, i_d)] = (\varepsilon^d \lambda_d(B^d))^d$. If $H(i_1, \dots, i_d)$ denotes the event that there is a k -hole in S containing the points p_{i_1}, \dots, p_{i_d} , then we have $\Pr[H(i_1, \dots, i_d) \mid E(i_1, \dots, i_d)] \geq C$. Altogether, we obtain $\Pr[H(i_1, \dots, i_d)] \geq C(\varepsilon^d \lambda_d(B^d))^d$.

It follows that the expected number of d -tuples of points from S that are contained in a k -hole in S is at least $C(\varepsilon^d \lambda_d(B^d))^d \binom{n}{d}$. Since every k -hole contains $\binom{k}{d}$ d -tuples of points, the expected number of k -holes in S is at least

$$C(\varepsilon^d \lambda_d(B^d))^d \binom{k}{d}^{-1} \binom{n}{d}.$$

This finishes the proof of Theorem 1.

4 Proof of Theorem 2

Here we prove the lower bound $\lim_{n \rightarrow \infty} n^{-d} EH_{d,d+1}^K(n) \geq \frac{2}{(d-1)! p_{d-1}}$ for every integer $d \geq 2$ and every convex body $K \in \mathcal{K}_d$, where p_{d-1} is the maximum probability that the convex polytope formed by the convex hull of $d+1$ points chosen uniformly and independently at random from a convex body in \mathbb{R}^{d-1} has only d vertices. We also prove Corollary 3 and Proposition 5 at the end of this section.

First, we state some definitions. Recall that λ_d denotes the Lebesgue measure on \mathbb{R}^d . For a subspace E of \mathbb{R}^d , we use λ_E to denote the Lebesgue measure on E and, for a compact convex set F with affine hull E , we use λ_F to denote the restriction of λ_E to F .

Recall that the volume of the d -dimensional unit ball B^d in \mathbb{R}^d is denoted by

$$\kappa_d = \lambda_d(B^d) = \frac{\pi^{d/2}}{\Gamma(1 + d/2)}.$$

Similarly, we denote the surface area of the d -dimensional unit sphere S^{d-1} by

$$\omega_d = \lambda_{d-1}(S^{d-1}) = d \cdot \kappa_d = \frac{2\pi^{d/2}}{\Gamma(d/2)}.$$

We also let $\Delta_q(x_0, \dots, x_q)$ be the q -dimensional volume of the simplex $\text{conv}(\{x_0, \dots, x_q\})$.

For integers d and k with $1 \leq k \leq d$, we let $A(d, k)$ be the affine Grassmannian of k -dimensional affine subspaces of \mathbb{R}^d equipped with its standard topology. The Grassmannian $A(d, k)$ is also equipped with the unique rigid motion invariant Haar measure μ_k , normalized by $\mu_k(\{E \in A(d, k) : E \cap B^d \neq \emptyset\}) = \kappa_{d-k}$. For more background, see Sections 1.3 and 5.1 in [SW08].

The following result is a special case of the affine Blaschke–Petkantschin formula; see Theorem 7.2.7 in [SW08].

Lemma 14. *For every integer $d \geq 2$, if K' is a d -dimensional convex body, then*

$$\lambda_d(K')^d = (d-1)! \frac{\omega_d}{\omega_1} \int_{A(d, d-1)} \int_{(K' \cap E)^d} \Delta_{d-1} \, d\lambda_E^d \mu_{d-1}(dE).$$

Let $EV_d^{K'}$ denote the expected d -dimensional volume of the convex hull of $d+1$ points drawn uniformly and independently at random from a convex body $K' \subseteq \mathbb{R}^d$. That is,

$$EV_d^{K'} = \frac{1}{\lambda(K')^{d+1}} \int_{K'} \cdots \int_{K'} \Delta_d(x_0, \dots, x_d) \lambda(dx_0) \cdots \lambda(dx_d).$$

Let K be a convex body in \mathbb{R}^d of unit d -dimensional volume. Since $\lambda_d(K) = 1$, Lemma 14 gives

$$1 = (d-1)! \frac{\omega_d}{\omega_1} \int_{A(d, d-1)} \int_{(K \cap E)^d} \Delta_{d-1} \, d\lambda_E^d \mu_{d-1}(dE), \quad (4)$$

where the right side equals

$$(d-1)! \frac{\omega_d}{\omega_1} \int_{A(d, d-1)} \lambda_{d-1}(K \cap E)^d EV_{d-1}^{K \cap E} \mu_{d-1}(dE) \quad (5)$$

by the definition of $EV_{d-1}^{K \cap E}$. Since

$$\frac{(d+1)EV_{d-1}^{K \cap E}}{\lambda_{d-1}(K \cap E)} = p_{d-1}^{K \cap E} \leq p_{d-1} \quad (6)$$

(see [Kle69, Sch88]), we have $EV_{d-1}^{K \cap E} \leq \frac{\lambda_{d-1}(K \cap E) p_{d-1}}{d+1}$. Using this estimate, we bound (5) from above by

$$\frac{p_{d-1}}{d+1} (d-1)! \frac{\omega_d}{\omega_1} \int_{A(d, d-1)} \lambda_{d-1}(K \cap E)^{(d+1)} \mu_{d-1}(dE).$$

By combining this estimate with (4), we obtain

$$\frac{\omega_1(d+1)}{\omega_d(d-1)! p_{d-1}} \leq \int_{A(d, d-1)} \lambda_{d-1}(K \cap E)^{d+1} \mu_{d-1}(dE). \quad (7)$$

Now, we apply a result of Reitzner and Temesvari [RT19, Theorem 1.4], which states

$$\lim_{n \rightarrow \infty} n^{-d} EH_{d, d+1}^K(n) = \frac{d\kappa_d}{(d+1)} \lambda_d(K)^{-d} \int_{A(d, d-1)} \lambda_{d-1}(K \cap E)^{d+1} \mu_{d-1}(dE).$$

Together with (7), $\lambda_d(K) = 1$, $\omega_d = d\kappa_d$, and $\omega_1 = 2$, this yields the desired estimate

$$\lim_{n \rightarrow \infty} n^{-d} EH_{d,d+1}^K(n) \geq \frac{d\kappa_d}{(d+1)} \cdot \frac{\omega_1(d+1)}{\omega_d(d-1)!p_{d-1}} = \frac{2}{(d-1)!p_{d-1}},$$

which completes the proof of Theorem 2. \square

We now prove Corollary 3 using the following bounds on p_2^K by Blaschke [Bla17, Bla23] and a result of Groemer [Gro73].

Theorem 15 ([Bla17, Bla23]). *For every 2-dimensional convex body $K \subseteq \mathbb{R}^2$, we have*

$$\frac{35}{12\pi^2} \leq p_2^K \leq \frac{1}{3}.$$

Moreover, the left inequality is tight if and only if K is an ellipse, and the right inequality is tight if and only if K is a triangle.

Theorem 16 ([Gro73]). *For every integer $d \geq 2$ and every d -dimensional convex body $K \subseteq \mathbb{R}^d$, we have*

$$EV_d^B \leq EV_d^K,$$

where B is the d -dimensional ball of the same volume as K . Moreover, the inequality is tight if and only if K is an ellipsoid.

Proof of Corollary 3. Let \mathcal{C}_K be the space of all $(d-1)$ -dimensional convex bodies that can be obtained as an intersection of a d -dimensional convex body K with a hyperplane. The probability p_{d-1}^K is affine invariant and thus attains its minimum and maximum on \mathcal{C}_K for every K ; we refer the reader to [Mac51] and Chapter 1 in [Sch14].

Note that \mathcal{C}_B contains only ellipsoids. Since \mathcal{C}_S does not contain ellipsoids, Theorem 16 gives

$$\min_{K \in \mathcal{C}_S} p_{d-1}^K > p_{d-1}^{B^{d-1}} = \max_{K \in \mathcal{C}_B} p_{d-1}^K.$$

By estimating $EV_{d-1}^{K \cap E}$ in (5) from below with $(d+1)\lambda_{d-1}(K \cap E) \min_{K \in \mathcal{C}_S} p_{d-1}^K$ for S and from above with $(d+1)\lambda_{d-1}(K \cap E) p_{d-1}^{B^{d-1}}$ for B , it then follows from the proof of Theorem 2 that

$$\lim_{n \rightarrow \infty} n^{-d} EH_{d,d+1}^S(n) < \lim_{n \rightarrow \infty} n^{-d} EH_{d,d+1}^B(n).$$

For the bounds in \mathbb{R}^3 , the upper bound was proved by Reitzner and Temesvari [RT19]. Thus, we only need to show the lower bound. Theorem 15 implies $p_2 = \frac{1}{3}$ and by plugging this identity into Theorem 2, we obtain

$$\lim_{n \rightarrow \infty} n^{-3} EH_{3,4}^K \geq \frac{2}{2! \cdot \frac{1}{3}} = 3.$$

\square

Finally, we show an upper bound on p_d^K for every $d \geq 2$ and every convex body $K \subseteq \mathbb{R}^d$ with small diameter. That is, we prove Proposition 5.

Proof of Proposition 5. We can assume without loss of generality that $f(K) = K$. We prove the upper bound on p_d^K . Since $\frac{(d+2)EV_d^K}{\lambda_d(K)} = p_d^K$ by (6), it is sufficient to estimate the term $\frac{EV_d^K}{\lambda_d(K)}$, which is the expected d -dimensional volume EV_d^K of the convex hull $\text{conv}(\{p_1, \dots, p_{d+1}\})$ of $d+1$ points drawn uniformly from K divided by the volume $\lambda_d(K)$ of K . The volume of $\text{conv}(\{p_1, \dots, p_{d+1}\})$ can be expressed as $\frac{1}{d!} \prod_{i=2}^{d+1} h_i$, where h_i is the distance of p_i from the affine hull of the set $\{p_1, \dots, p_{i-1}\}$. Since K is convex, the distance h_i is at most the diameter D of K for every $i \in \{2, \dots, d+1\}$ and thus

$$\frac{EV_d^K}{\lambda_d(K)} \leq \frac{1}{d! \lambda_d(K)} \prod_{i=2}^{d+1} h_i \leq \frac{D^d}{d! \lambda_d(K)}.$$

Altogether, we have $p_d \leq \frac{(d+2)D^d}{d! \lambda_d(K)}$. □

5 Proof of Theorem 6

Let K be a planar convex body of unit area. Let S be a set of n points chosen uniformly and independently at random from K . We prove Theorem 6 by showing that the expected number $E_{2,4}^K(n)$ of 4-holes in S satisfies

$$\lim_{n \rightarrow \infty} n^{-2} E_{2,4}^K(n) = 10 - \frac{2\pi^2}{3}.$$

We distinguish two possible *types* of 4-holes and estimate their expected numbers separately. In the first type, the pair of vertices of a 4-hole H in S at the largest distance between any two vertices of H forms a diagonal of H . In the other type, this pair of vertices forms an edge of H ; see Figure 1.



Figure 1: Two types of 4-holes.

We now estimate the expected number of the 4-holes of the first type. For points $p_1, p_2, p_3, p_4 \in S$, we estimate the probability of the event $E_1(p_3, p_4 \mid p_1, p_2)$ that there is a 4-hole $H = \{p_1, \dots, p_4\}$ in S , where the points p_1 and p_2 form a diagonal and determine the largest distance between any two points from H , conditioned on the fact that p_1 and p_2 are already placed in K . That is, the probability that, for fixed p_1 and p_2 , the points p_3 and p_4 determine a 4-hole of the first type with diameter $|p_1 p_2|$.

Without loss of generality we can assume that $p_1 = (0, 0)$ and $p_2 = (\ell, 0)$ for some $\ell > 0$, as otherwise we apply a suitable isometry to S . Let R be the set of points from

$K \cap ([0, \ell] \times [-\frac{2}{\ell}, \frac{2}{\ell}])$ that are at distance at most ℓ from p_1 and also from p_2 . Note that the set R is convex. For every point $p \neq p_1, p_2$ of the 4-hole H , we have $x(p_1) < x(p) < x(p_2)$, as otherwise $|p_1 p_2|$ is not the diameter of H . If $|y(p)| > \frac{2}{\ell}$, then the triangle T spanned by p_1, p_2 , and p has area larger than 1, which is impossible, since we have $T \subseteq K$ and $\lambda_2(K) = 1$ by the convexity of K . Consequently, p lies in R . For a real number $y \in [-\frac{2}{\ell}, \frac{2}{\ell}]$, let I_y be the line segment formed by points $r \in R$ with $y(r) = y$. Note that $|I_y| \leq \ell$ for every y and that $|I_0| = \ell$.

The event $E_1(p_3, p_4 \mid p_1, p_2)$ is partitioned into two events $E_1^\uparrow(p_3, p_4 \mid p_1, p_2)$ and $E_1^\downarrow(p_3, p_4 \mid p_1, p_2)$ distinguished by the position of p_3 and p_4 (either p_3 lies above $\overline{p_1 p_2}$ and p_4 below or vice versa). By symmetry, it suffices to estimate the probability of $E_1^\uparrow(p_3, p_4 \mid p_1, p_2)$.

The event $E_1^\uparrow(p_3, p_4 \mid p_1, p_2)$ happens if and only if p_3, p_4 lie in R , both triangles $p_1 p_2 p_3$ and $p_1 p_2 p_4$ are 3-holes in S , and $|p_3 p_4| \leq |p_1 p_2|$. Note that $\lambda_2(p_1 p_2 p_3) + \lambda_2(p_1 p_2 p_4) = \frac{\ell}{2}(y(p_3) - y(p_4)) \leq 1$ by the convexity of K , so we get $-y(p_4) \leq 2/\ell - y(p_3)$. Therefore, the probability $\Pr[E_1^\uparrow(p_3, p_4 \mid p_1, p_2)]$ is at most

$$\int_{y=0}^{2/\ell} \int_{z=0}^{2/\ell-y} |I_y| \cdot |I_{-z}| \cdot \Pr[p_1 p_2 p_3 p_4 \text{ is empty in } S \mid p_3 \in I_y, p_4 \in I_{-z}] \, dz dy. \quad (8)$$

Let I_z^y be the set of points from I_z that are at distance at most ℓ from any point of I_y and observe that $I_0^0 = I_0$. Then, the probability $\Pr[E_1^\uparrow(p_3, p_4 \mid p_1, p_2)]$ is at least

$$\int_{y=0}^{2/\ell} \int_{z=0}^{2/\ell-y} |I_y| \cdot |I_{-z}^y| \cdot \Pr[p_1 p_2 p_3 p_4 \text{ is empty in } S \mid p_3 \in I_y, p_4 \in I_{-z}^y] \, dz dy. \quad (9)$$

Note that

$$\Pr[p_1 p_2 p_3 p_4 \text{ is empty in } S \mid p_3 \in I_y, p_4 \in I_{-z}] = \left(1 - \frac{\ell \cdot (y + z)}{2}\right)^{n-4},$$

as all the remaining $n - 4$ points from S must lie outside of $\text{conv}(\{p_1, p_2, p_3, p_4\})$, which has area $\lambda_2(p_1 p_2 p_3) + \lambda_2(p_1 p_2 p_4) = \frac{\ell}{2}(y(p_3) - y(p_4))$. The same expression holds for $\Pr[p_1 p_2 p_3 p_4 \text{ is empty in } S \mid p_3 \in I_y, p_4 \in I_{-z}^y]$. We now plug in this expression to (8) and (9) and substitute $Y = yn$ and $Z = zn$. Then, (8) becomes

$$\frac{1}{n^2} \int_{Y=0}^{2n/\ell} \int_{Z=0}^{2n/\ell-Y} |I_{Y/n}| \cdot |I_{-Z/n}| \cdot \left(1 - \frac{\ell \cdot (Y + Z)}{2n}\right)^{n-4} \, dZ dY$$

while (9) becomes

$$\frac{1}{n^2} \int_{Y=0}^{2n/\ell} \int_{Z=0}^{2n/\ell-Y} |I_{Y/n}| \cdot |I_{-Z/n}^Y| \cdot \left(1 - \frac{\ell \cdot (Y + Z)}{2n}\right)^{n-4} \, dZ dY.$$

Note that we integrate over the set $\{(Y, Z) \in \mathbb{R}^2 : Y + Z \leq 2n/\ell, Y, Z \geq 0\}$, which becomes $\{(Y, Z) \in \mathbb{R}^2 : Y, Z \geq 0\}$ for n going to infinity. Also note all sets $I_{Y/n}, I_{-Z/n}$,

and $I_{-Z/n}^{Y/n}$ become I_0 as n goes to infinity. Thus, by the monotone convergence theorem, we have

$$\lim_{n \rightarrow \infty} n^2 \Pr[E_1^\uparrow(p_3, p_4 | p_1, p_2)] = \int_{Y=0}^{\infty} \int_{Z=0}^{\infty} |I_0| \cdot |I_0| \cdot e^{-\ell \cdot (Y+Z)/2} dZ dY.$$

Since $|I_0| = \ell$, we can simplify this expression as

$$\begin{aligned} \lim_{n \rightarrow \infty} n^2 \Pr[E_1^\uparrow(p_3, p_4 | p_1, p_2)] &= \int_{Y=0}^{\infty} \int_{Z=0}^{\infty} \ell^2 e^{-\ell \cdot (Y+Z)/2} dZ dY \\ &= \int_{Y=0}^{\infty} \ell e^{-\ell Y/2} dY \cdot \int_{Z=0}^{\infty} \ell e^{-\ell Z/2} dZ = 2 \cdot 2 = 4. \end{aligned}$$

Thus, the expected number of 4-holes of the first type determined by p_1 and p_2 is $\binom{n}{2} \cdot 2 \cdot \frac{4}{n^2} = 4(1 + o(1))$ for n going to infinity. Thus, the overall expected number of 4-holes of the first type is $4(1 + o(1))\binom{n}{2} = 2(1 + o(1))n^2$.

It remains to estimate the expected number of the 4-holes of the second type. For points $p_1, p_2, p_3, p_4 \in S$, we estimate the probability of the event $E_2(p_3, p_4 | p_1, p_2)$ that, for fixed p_1 and p_2 , there is a 4-hole $H = \{p_1, p_2, p_3, p_4\}$ in S , where p_1 and p_2 form an edge of $\text{conv}(H)$ and determine the largest distance between any two points from H and p_4 is closer to the line $\overline{p_1 p_2}$ than p_3 . In particular, p_1 and p_2 determine a 4-hole of the second type of diameter $|p_1 p_2|$.

Again, we can assume without loss of generality that $p_1 = (0, 0)$ and $p_2 = (\ell, 0)$ for some $\ell > 0$. We consider the sets R and I_y for every $y \in [-\frac{2}{\ell}, \frac{2}{\ell}]$ defined exactly as before. Again, every point $p \neq p_1, p_2$ of the 4-hole H lies in R .

The event $E_2(p_3, p_4 | p_1, p_2)$ is partitioned into four pairwise disjoint events $E_2^{\nwarrow}(p_3, p_4 | p_1, p_2)$, $E_2^{\nearrow}(p_3, p_4 | p_1, p_2)$, $E_2^{\searrow}(p_3, p_4 | p_1, p_2)$, and $E_2^{\swarrow}(p_3, p_4 | p_1, p_2)$ distinguished by the position of p_3 and p_4 with respect to the line $\overline{p_1 p_2}$ (either both lie above the line or below) and by the position of p_4 with respect to p_3 (p_4 is either to the left or to the right of p_3). First, we estimate the probability $\Pr[E_2^{\nwarrow}(p_3, p_4 | p_1, p_2)]$ of the event $E_2(p_3, p_4 | p_1, p_2)$ conditioned on the fact that p_3 and p_4 both lie above $\overline{p_1 p_2}$ and p_4 lies to the left of p_3 ; see Figure 2.

The event $E_2^{\nwarrow}(p_3, p_4 | p_1, p_2)$ happens if and only if p_3, p_4 lie in R and both triangles $p_1 p_2 p_3$ and $p_1 p_3 p_4$ are 3-holes in S . For $y \in [0, 2/\ell]$, let $(l(y), y)$ and $(r(y), y)$ be the left and the right endpoint of the line segment I_y , respectively. Note that $l(0) = 0$ and $r(0) = \ell$. The point p_3 lies in I_y for some $y \in [0, 2/\ell]$ and the x -coordinate of p_3 then lies in the interval $[l(y), r(y)]$. Since p_4 lies to the left of $\overline{p_1 p_3}$, the x -coordinate of p_4 lies in the interval $[l(y'), xy'/y']$, where $p_3 = (x, y)$ and y' is the y -coordinate of p_4 .

Thus, we can express the probability $\Pr[E_2^{\nwarrow}(p_3, p_4 | p_1, p_2)]$ as

$$\int_0^{2/\ell} \int_{l(y)}^{r(y)} \int_0^y \int_{l(y')}^{xy'/y'} P(p_1, p_2, p_3, p_4) dx' dy' dx dy,$$

where $P(p_1, p_2, p_3, p_4)$ is the probability that $\text{conv}(\{p_1, p_2, p_3, p_4\})$ is empty in S , conditioned on the event $p_3 = (x, y)$ and $p_4 = (x', y')$.




Figure 2: The case when the points $p_3 = (x, y)$ and $p_4 = (x', y')$ are both above $\overline{p_1 p_2}$ and p_4 lies to the left of p_3 .




Figure 3: The partitioning of $\text{conv}(\{p_1, p_2, p_3, p_4\})$.

By partitioning the set $\text{conv}(H) = \text{conv}(\{p_1, p_2, p_3, p_4\})$ into four regions A_1 , A_2 , A_3 , and A_4 as in Figure 3, we can express the area $\lambda_2(\text{conv}(H))$ as $\lambda_2(A_1) + \dots + \lambda_2(A_4)$, which equals

$$\frac{x'y'}{2} + (x - x')y' + \frac{(\ell - x)y}{2} + \frac{(x - x')(y - y')}{2}.$$

Rewriting this expression, we obtain $\lambda_2(\text{conv}(H)) = \frac{(\ell - x')y + xy'}{2}$. Thus, we have

$$P(p_1, p_2, p_3, p_4) = (1 - \lambda_2(\text{conv}(H)))^{n-4} = \left(1 - \frac{(\ell - x')y + xy'}{2}\right)^{n-4},$$

as $\lambda_2(K) = 1$ and all the remaining $n - 4$ points from S must lie in $K \setminus \text{conv}(H)$.

Therefore, the probability $\Pr[E_2^K(p_3, p_4 \mid p_1, p_2)]$ equals

$$\int_0^{2/\ell} \int_{l(y)}^{r(y)} \int_0^y \int_{l(y')}^{xy'/y} \left(1 - \frac{(\ell - x')y + xy'}{2}\right)^{n-4} dx' dy' dx dy.$$

By Fubini's theorem, we can exchange the order of the integrals as

$$\int_0^{2/\ell} \int_0^y \int_{l(y)}^{r(y)} \int_{l(y')}^{xy'/y} \left(1 - \frac{(\ell - x')y + xy'}{2}\right)^{n-4} dx' dx dy' dy.$$

We now substitute $Y = yn$ and $Y' = y'n$ and express $\Pr[E_2^\wedge(p_3, p_4 | p_1, p_2)]$ as

$$\frac{1}{n^2} \int_0^{\frac{2n}{\ell}} \int_0^Y \int_{l(\frac{Y}{n})}^{r(\frac{Y}{n})} \int_{l(\frac{Y'}{n})}^{\frac{xY'}{Y}} \left(1 - \frac{(\ell - x')Y + xY'}{2n}\right)^{n-4} dx' dx dY' dY.$$

Since $l(0) = 0$ and $r(0) = \ell$, the monotone convergence theorem implies that

$$\lim_{n \rightarrow \infty} n^2 \Pr[E_2^\wedge(p_3, p_4 | p_1, p_2)]$$

equals

$$\int_0^\infty \int_0^Y \int_0^\ell \int_0^{xY'/Y} e^{-((\ell-x')Y+xY')/2} dx' dx dY' dY,$$

which can be rewritten as

$$\int_0^\infty e^{-\ell Y/2} \int_0^Y \int_0^\ell e^{-xY'/2} \int_0^{xY'/Y} e^{x'Y/2} dx' dx dY' dY.$$

Computing the innermost integral, we obtain

$$\lim_{n \rightarrow \infty} n^2 \Pr[E_2^\wedge(p_3, p_4 | p_1, p_2)] = 2 \int_0^\infty \frac{e^{-\ell Y/2}}{Y} \int_0^Y \int_0^\ell (1 - e^{-xY'/2}) dx dY' dY,$$

which can be further rewritten as

$$2 \int_0^\infty \frac{e^{-\ell Y/2}}{Y} \int_0^Y \left(\ell - \frac{1 - e^{-\ell Y'/2}}{Y'/2} \right) dY' dY = 4 \int_0^\infty \frac{e^{-z}}{z} \int_0^z \left(1 - \frac{1 - e^{-t}}{t} \right) dt dz$$

by substituting $\frac{\ell}{2}Y = z$ and $\frac{\ell}{2}Y' = t$. We split the integral into two parts. The first part equals

$$4 \int_0^\infty \frac{e^{-z}}{z} \int_0^z 1 dt dz = 4 \int_0^\infty e^{-z} dz = 4.$$

Using the modified exponential integral function (see Chapter 5.1 in [AS65])

$$\int_0^z \frac{1 - e^{-t}}{t} dt = \sum_{k=1}^{\infty} \frac{(-1)^{k+1} \cdot z^k}{k \cdot k!},$$

we can write the second part as

$$4 \int_0^\infty \frac{e^{-z}}{z} \int_0^z \frac{1 - e^{-t}}{t} dt dz = 4 \int_0^\infty \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k \cdot k!} \cdot e^{-z} z^{k-1} dz.$$

The integral of the absolute value is finite, because

$$\begin{aligned} 4 \int_0^\infty \sum_{k=1}^{\infty} \frac{1}{k \cdot k!} \cdot e^{-z} z^{k-1} dz &= 4 \sum_{k=1}^{\infty} \int_0^\infty \frac{1}{k \cdot k!} \cdot e^{-z} z^{k-1} dz \\ &= 4 \sum_{k=1}^{\infty} \frac{1}{k \cdot k!} \int_0^\infty e^{-z} z^{k-1} dz = 4 \sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{2\pi^2}{3}, \end{aligned}$$

where the first equality follows from Tonelli's theorem, the second equality from the linearity of the integral, and the third equality uses $\int_0^\infty e^{-z} z^{k-1} dz = \Gamma(k) = (k-1)!$ for $k \in \mathbb{N}$. Therefore, we can apply Fubini's theorem and obtain

$$4 \int_0^\infty \sum_{k=1}^\infty \frac{(-1)^{k+1}}{k \cdot k!} \cdot e^{-z} z^{k-1} dz = 4 \sum_{k=1}^\infty \int_0^\infty \frac{(-1)^{k+1}}{k \cdot k!} e^{-z} z^{k-1} dz.$$

Again, since $\int_0^\infty e^{-z} z^{k-1} dz = \Gamma(k) = (k-1)!$ for $k \in \mathbb{N}$, we have

$$4 \sum_{k=1}^\infty \frac{(-1)^{k+1}}{k \cdot k!} \int_0^\infty e^{-z} z^{k-1} dz = 4 \sum_{k=1}^\infty \frac{(-1)^{k+1}}{k^2} = \frac{\pi^2}{3}.$$

Altogether, we have

$$\lim_{n \rightarrow \infty} n^2 \Pr[E_2^\wedge(p_3, p_4 \mid p_1, p_2)] = 4 - \frac{\pi^2}{3}.$$

The probability of each of the remaining three pairwise disjoint events $E_2^\nearrow(p_3, p_4 \mid p_1, p_2)$, $E_2^\searrow(p_3, p_4 \mid p_1, p_2)$, and $E_2^\times(p_3, p_4 \mid p_1, p_2)$ can be computed in an analogous way and all of them are equal. We then obtain

$$\begin{aligned} \Pr[E_2(p_3, p_4 \mid p_1, p_2)] &= \Pr[E_2^\wedge(p_3, p_4 \mid p_1, p_2)] + \Pr[E_2^\nearrow(p_3, p_4 \mid p_1, p_2)] \\ &\quad + \Pr[E_2^\searrow(p_3, p_4 \mid p_1, p_2)] + \Pr[E_2^\times(p_1, p_2)] \\ &= 4 \Pr[E_2^\wedge(p_3, p_4 \mid p_1, p_2)]. \end{aligned}$$

Thus, the expected number of 4-holes of the second type determined by p_1 and p_2 is $n^2 \left(\frac{16}{n^2} - \frac{4\pi^2}{3n^2} \right) = 16 - \frac{4\pi^2}{3}$ for n going to infinity. Thus, the overall expected number of 4-holes of the second type is $(1 + o(1)) \left(16 - \frac{4\pi^2}{3} \right) \binom{n}{2} = (1 + o(1)) \left(8 - \frac{2\pi^2}{3} \right) n^2$. Adding the expected number of the 4-holes of the first type, we see that

$$\lim_{n \rightarrow \infty} n^{-2} EH_{2,4}^K(n) = 2 + \left(8 - \frac{2\pi^2}{3} \right) = 10 - \frac{2\pi^2}{3},$$

which completes the proof of Theorem 6.

6 Proof of Theorem 7

We extend the methods used in Section 5 to larger holes in random planar point sets and we show that, for every fixed integer $k \geq 3$, the limit $\lim_{n \rightarrow \infty} n^{-2} EH_{2,k}^K(n)$ exists and does not depend on the convex body $K \in \mathcal{K}_2$.

Let K be a convex body of unit area in the plane and let $k \geq 3$ be an integer. We use S to denote a set of n points chosen uniformly and independently at random from K .

We distinguish several types of k -holes in S and we compute the expected number for each type separately. Let H be a k -hole in S and let p_1, \dots, p_k be the points of H such that $|p_1 p_k|$ is the diameter of $\text{conv}(H)$. We can assume without loss of generality that $p_1 = (0, 0)$ and $p_k = (\ell, 0)$ for some $\ell > 0$. In particular, all the remaining points p_2, p_3, \dots, p_{k-1} lie between p_1 and p_k , since $|p_1 p_k|$ is the diameter of H . We then label the points p_1, \dots, p_k so that they are ordered by their increasing x -coordinates. For $T \subseteq \{2, 3, \dots, k-1\}$, we use \bar{T} to denote the set $\{2, 3, \dots, k-1\} \setminus T$. For $a \in \bar{T}$ and $b \in T$, we say that H is of *type* (T, a, b) if all points p_i with $i \in T$ lie above the line $\overline{p_1 p_k}$, all points from \bar{T} lie below the line $\overline{p_1 p_k}$, the point p_b has the largest y -coordinate among the points in T and p_a has the smallest y -coordinate among the points in \bar{T} ; see Figure 4. If T or \bar{T} is empty, then we let a or b be equal to \emptyset , respectively. Observe that there are exactly

$$\begin{aligned} 2(k-2) + \sum_{\substack{T \subseteq \{2, 3, \dots, k-1\} \\ T, \bar{T} \neq \emptyset}} |T| \cdot |\bar{T}| &= 2(k-2) + \sum_{i=1}^{k-3} \binom{k-2}{i} i(k-2-i) \\ &= 2k-4 + 2^{k-4}(k^2 - 5k + 6) \end{aligned}$$

types of k -holes in S .




Figure 4: An example of an 8-hole $\{p_1, \dots, p_8\}$ in S of the type $(\{2, 3, 5, 7\}, 6, 7)$ with the diameter $\ell = |p_1 p_8|$.

Let H be a k -hole of a type (T, a, b) and let T^\wedge be the set of points p_i with $i \in T$ and $i < b$. Then, since the points p_1, \dots, p_k are in convex position, the points p_i with $i \in T^\wedge \cup \{1, b\}$ have increasing y -coordinates. Similarly, we define T^\nearrow to be the set of points p_i with $i \in T$ and $i > b$ and T^\swarrow and T^\searrow to be the sets of points p_i with $i \in \bar{T}$ and with $i < a$ and $i > a$, respectively. The convex position of the points p_1, \dots, p_k then implies that the points p_i with $i \in T^\nearrow \cup \{b, k\}$ and $T^\swarrow \cup \{1, a\}$ have decreasing y -coordinates while the points p_i with $T^\searrow \cup \{a, k\}$ have increasing y -coordinates.

Let (T, a, b) be a fixed type of k -holes in S . For any two points $p_1, p_k \in K$, we estimate the probability of the event $E_{T,a,b}(p_1, \dots, p_k \mid p_1, p_k)$ that, for fixed p_1 and p_k , there is a k -hole $H = \{p_1, \dots, p_k\}$ in S of the type (T, a, b) with the diameter $|p_1 p_k|$.

We use R to denote the set of points from $K \cap ([0, \ell] \times [-\frac{2}{\ell}, \frac{2}{\ell}])$ that are at distance at most ℓ from p_1 and also from p_k . Similarly as in Section 5, note that R is convex and that every point $p \neq p_1, p_k$ of the k -hole H lies in R . For $y \in [-\frac{2}{\ell}, \frac{2}{\ell}]$, let I_y be the line segment formed by points $r \in R$ with $y(r) = y$. We use $(l(y), y)$ and $(r(y), y)$ to denote the left and the right endpoint of I_y , respectively. Note that $l(0) = 0$ and $r(0) = \ell$.

The point p_a lies in I_y for some $y \in [-\frac{2}{\ell}, 0]$ and p_b in $I_{y'}$ for some $y' \in [0, \frac{2}{\ell}]$. The x -coordinates of p_a and p_b then lie in the intervals $[l(y), r(y)]$ and $[l(y'), r(y')]$, respectively.

Let p_{i_1}, \dots, p_{i_m} be the points from $T^\times \cup \{p_1, p_b\}$ ordered by increasing x -coordinates. We assume $p_{i_j} = (x_{i_j}, y_{i_j})$ for every $j = 1, \dots, m$. Since the points from $T^\times \cup \{p_1, p_b\}$ are in convex position with increasing x - and y -coordinates, we have, if $m \geq 3$,

$$0 \leq y_{i_{m-1}} \leq y_{i_m} \quad \text{and} \quad l(y_{i_{m-1}}) \leq x_{i_{m-1}} \leq x_{i_m} \frac{y_{i_{m-1}}}{y_{i_m}} \quad (10)$$

as $p_{i_{m-1}} \in I_{y_{i_{m-1}}}$ has to lie above the line $\overline{p_1 p_{i_m}}$. Moreover, for every $j = m-2, \dots, 3, 2$, we have

$$0 \leq y_{i_j} \leq y_{i_{j+1}} \quad (11)$$

and

$$\max \left\{ l(y_{i_j}), x_{i_{j+1}} + \frac{(x_{i_{j+2}} - x_{i_{j+1}})(y_{i_j} - y_{i_{j+1}})}{y_{i_{j+2}} - y_{i_{j+1}}} \right\} \leq x_{i_j} \leq x_{i_{j+1}} \frac{y_{i_j}}{y_{i_{j+1}}} \quad (12)$$

as $p_{i_j} \in I_{y_j}$ has to lie below the line $\overline{p_{i_{j+1}} p_{i_{j+2}}}$ and above the line $\overline{p_1 p_{i_{j+1}}}$. Let $J_{T,a,b}^\times$ be the set of vectors $(x_2, y_2, x_3, y_3, \dots, x_{k-1}, y_{k-1})$ such that $0 \leq x_2 \leq \dots \leq x_{k-1} \leq \ell$ and the points $(x_{i_1}, y_{i_1}), \dots, (x_{i_{m-1}}, y_{i_{m-1}})$ satisfy the conditions (10), (11) and (12). We analogously define the sets $J_{T,a,b}^\nearrow$, $J_{T,a,b}^\searrow$, and $J_{T,a,b}^\swarrow$ for the sets T^\nearrow , T^\searrow , and T^\swarrow , respectively.

We let $A_{T,a,b}^\times$ be the area of the convex hull of the points from $T^\times \cup \{p_1, p_b, (x(p_b), 0)\}$; see Figure 5. By considering the partitioning into triangles and rectangles as in Figure 5, the area $A_{T,a,b}^\times$ can be expressed as

$$\sum_{j=2}^m \frac{(x_{i_j} - x_{i_{j-1}})(y_{i_j} - y_{i_{j-1}})}{2} + \sum_{j=2}^{m-1} (x_{i_m} - x_{i_j})(y_{i_j} - y_{i_{j-1}}). \quad (13)$$

We also let $A_{T,a,b}^\nearrow$, $A_{T,a,b}^\searrow$, and $A_{T,a,b}^\swarrow$ be the corresponding areas and we note that they can be evaluated using an analogous approach. Observe that $\lambda_2(\text{conv}(H)) = A_{T,a,b}^\times + A_{T,a,b}^\nearrow + A_{T,a,b}^\searrow + A_{T,a,b}^\swarrow$.

The event $E_{T,a,b}(p_1, \dots, p_k \mid p_1, p_k)$ happens if all points from $S \setminus \{p_1, \dots, p_k\}$ lie in $K \setminus \text{conv}(H)$. Since $\lambda_2(K) = 1$, we can express the probability $\Pr[E_{T,a,b}(p_1, \dots, p_k \mid p_1, p_k)]$ in the case $a < b$ as

$$\int_0^{2/\ell} \int_{l(y_b)}^{r(y_b)} \int_{-2/\ell+y_b}^0 \int_{l(y_a)}^{\min\{x_b, r(y_a)\}} \int_{J_{T,a,b}} (1 - \lambda_2(\text{conv}(H)))^{n-k} \, d\mathbf{z} \, dx_a \, dy_a \, dx_b \, dy_b,$$




Figure 5: The area $A_{T,a,b}^{\wedge}$ of the convex hull of the points from $\{p_2, p_3, p_5\} \cup \{p_1, p_b, (x(p_b), 0)\}$ is denoted by four grey triangles and three grey rectangles.

where $p_i = (x_i, y_i)$ for every $i = 1, \dots, k$ and $J_{T,a,b}$ is the set of vectors $(x_2, y_2, \dots, x_{k-1}, y_{k-1})$ from $J_{T,a,b}^{\nwarrow} \cap J_{T,a,b}^{\nearrow} \cap J_{T,a,b}^{\searrow} \cap J_{T,a,b}^{\swarrow}$ where any two points (x_i, y_i) and (x_j, y_j) are at distance at most ℓ . In the case $a > b$ we can write $\Pr[E_{T,a,b}(p_1, \dots, p_k \mid p_1, p_k)]$ as

$$\int_0^{2/\ell} \int_{l(y_b)}^{r(y_b)} \int_{-2/\ell+y_b}^0 \int_{\max\{x_b, l(y_a)\}}^{r(y_a)} \int_{J_{T,a,b}} (1 - \lambda_2(\text{conv}(H))^{n-k} \, dz \, dx_a \, dy_a \, dx_b \, dy_b.$$

In the case $a = \emptyset$, $\Pr[E_{T,a,b}(p_1, \dots, p_k \mid p_1, p_k)]$ equals

$$\int_0^{2/\ell} \int_{l(y_b)}^{r(y_b)} \int_{J_{T,\emptyset,b}} (1 - \lambda_2(\text{conv}(H))^{n-k} \, dz \, dx_b \, dy_b$$

and, similarly, if $b = \emptyset$, we have

$$\int_{-2/\ell}^0 \int_{l(y_a)}^{r(y_a)} \int_{J_{T,a,\emptyset}} (1 - \lambda_2(\text{conv}(H))^{n-k} \, dz \, dx_a \, dy_a.$$

If we substitute $Y_i = ny_i$ for every $i = 2, 3, \dots, k-1$ and we let n go to infinity, then conditions (10), (11), and (12) become

$$0 \leq Y_{i_{m-1}} \leq Y_{i_m} \quad \text{and} \quad 0 \leq x_{i_{m-1}} \leq x_{i_m} \frac{Y_{i_{m-1}}}{Y_{i_m}},$$

$$0 \leq Y_{i_j} \leq Y_{i_{j+1}},$$

and

$$\max \left\{ 0, x_{i_{j+1}} + \frac{(x_{i_{j+2}} - x_{i_{j+1}})(Y_{i_j} - Y_{i_{j+1}})}{Y_{i_{j+2}} - Y_{i_{j+1}}} \right\} \leq x_{i_j} \leq x_{i_{j+1}} \frac{Y_{i_j}}{Y_{i_{j+1}}},$$

respectively, since $l(0) = 0$. In particular, none of the new conditions depends on K . Let $G_{T,a,b}^{\wedge}$ be the set of vectors $(x_2, Y_2, x_3, Y_3, \dots, x_{k-1}, Y_{k-1})$, where $0 \leq x_2 \leq \dots \leq x_{k-1} \leq \ell$

and the points $(x_{i_1}, Y_{i_1}), \dots, (x_{i_{m-1}}, Y_{i_{m-1}})$ satisfy these new conditions. We analogously define the sets $G_{T,a,b}^{\nearrow}$, $G_{T,a,b}^{\searrow}$, and $G_{T,a,b}^{\swarrow}$.

By substituting $Y_i = ny_i$ for every $i = 2, 3, \dots, k-1$ and letting n go to infinity, the expression (13) for the area $A_{T,a,b}^{\nwarrow}$ becomes

$$\sum_{j=2}^m \frac{(x_{i_j} - x_{i_{j-1}})(Y_{i_j} - Y_{i_{j-1}})}{2n} + \sum_{j=2}^{m-1} \frac{(x_{i_m} - x_{i_j})(Y_{i_j} - Y_{i_{j-1}})}{n} = \frac{B_{T,a,b}^{\nearrow}}{n},$$

where $B_{T,a,b}^{\nearrow}$ is independent of K . Similarly, the expressions for the areas $A_{T,a,b}^{\nearrow}$, $A_{T,a,b}^{\searrow}$, $A_{T,a,b}^{\swarrow}$ become $\frac{B_{T,a,b}^{\nearrow}}{n}$, $\frac{B_{T,a,b}^{\searrow}}{n}$, $\frac{B_{T,a,b}^{\swarrow}}{n}$, respectively, where $B_{T,a,b}^{\nearrow}$, $B_{T,a,b}^{\searrow}$, $B_{T,a,b}^{\swarrow}$ are independent of K . Since $\lambda_2(\text{conv}(H)) = A_{T,a,b}^{\nwarrow} + A_{T,a,b}^{\nearrow} + A_{T,a,b}^{\searrow} + A_{T,a,b}^{\swarrow}$, we have $\lambda_2(\text{conv}(H)) = (B_{T,a,b}^{\nwarrow} + B_{T,a,b}^{\nearrow} + B_{T,a,b}^{\searrow} + B_{T,a,b}^{\swarrow})/n$.

Thus by substituting $Y_i = ny_i$ for every $i = 2, 3, \dots, k-1$ and using the monotone convergence theorem with the facts $l(0) = 0$ and $r(0) = \ell$, we can express $\lim_{n \rightarrow \infty} n^{k-2} \Pr[E_{T,a,b}(p_1, \dots, p_k \mid p_1, p_k)]$ in the case $a < b$ as

$$\int_0^\infty \int_0^\ell \int_{-\infty}^0 \int_0^{x_b} \int_{G_{T,a,b}} e^{-B_{T,a,b}^{\nwarrow} - B_{T,a,b}^{\nearrow} - B_{T,a,b}^{\searrow} - B_{T,a,b}^{\swarrow}} d\mathbf{Z} dx_a dY_a dx_b dY_b,$$

where $p_i = (x_i, Y_i/n)$ for every $i = 1, \dots, k$ and $G_{T,a,b}$ is the set of vectors $(x_2, Y_2, \dots, x_{k-1}, Y_{k-1})$ from $G_{T,a,b}^{\nwarrow} \cap G_{T,a,b}^{\nearrow} \cap G_{T,a,b}^{\searrow} \cap G_{T,a,b}^{\swarrow}$ such that the distance between any two points p_i and p_j is at most ℓ . Also note that this expression does not depend on K and is finite [BSV21]. This is true also in the remaining three cases $a > b$, $a = \emptyset$, and $b = \emptyset$.

Let $E(p_1, \dots, p_k \mid p_1, p_k)$ be the event that, for fixed p_1 and p_k , there is a k -hole $\{p_1, \dots, p_k\}$ in S with the diameter $|p_1 p_k|$. Then the limit $\lim_{n \rightarrow \infty} n^{k-2} \Pr[E(p_1, \dots, p_k \mid p_1, p_k)]$ equals

$$\begin{aligned} & \sum_{a \in \{2, \dots, k-1\}} \Pr[E_{\emptyset, a, \emptyset}(p_1, \dots, p_k \mid p_1, p_k)] + \sum_{b \in \{2, \dots, k-1\}} \Pr[E_{\{2, \dots, k-1\}, \emptyset, b}(p_1, \dots, p_k \mid p_1, p_k)] \\ & + \sum_{\substack{T \subseteq \{2, \dots, k-1\} \\ T, \bar{T} \neq \emptyset}} \sum_{a \in \bar{T}} \sum_{b \in T} \Pr[E_{T, a, b}(p_1, \dots, p_k \mid p_1, p_k)], \end{aligned}$$

which also does not depend on the convex body K .

The expected number of k -holes in S determined by p_1 and p_k with the diameter $|p_1 p_k|$ converges to $n^{k-2} \frac{\Pr[E(p_1, \dots, p_k \mid p_1, p_k)]}{n^{k-2}} = \Pr[E(p_1, \dots, p_k \mid p_1, p_k)]$. Considering all possible pairs $\{p_1, p_k\}$, we see that $\lim_{n \rightarrow \infty} n^{-2} EH_{2,k}^K(n) = \frac{1}{2} \Pr[E(p_1, \dots, p_k \mid p_1, p_k)]$, which does not depend on K . This finishes the proof of Theorem 7.

Acknowledgements We thank Sophia and Jonathan Rau for helping with the computation in the proof of Theorem 6. We would also like to thank to the anonymous reviewers for several helpful comments and for raising the question about the existence of the limits $\lim_{n \rightarrow \infty} EH_{d,k}^K(n)$ for general values of d and k and pointing out the connections to the Hyperplane conjecture.

References

- [ABH⁺20] O. Aichholzer, M. Balko, T. Hackl, J. Kynčl, I. Parada, M. Scheucher, P. Valtr, and B. Vogtenhuber. A superlinear lower bound on the number of 5-holes. *Journal of Combinatorial Theory. Series A*, 173:105236, 2020.
- [AS65] M. Abramowitz and I. Stegun. *Handbook of Mathematical Functions: With Formulas, Graphs, and Mathematical Tables*. Applied mathematics series. Dover Publications, 1965.
- [BCH20] B. Bukh, T. Chao, and R. Holzman. On convex holes in d -dimensional point sets. <http://arXiv.org/abs/2007.08972>, 2020.
- [BF87] I. Bárány and Z. Füredi. Empty simplices in Euclidean space. *Canadian Mathematical Bulletin*, 30(4):436–445, 1987.
- [Bla17] W. Blaschke. Lösung des 'Vierpunktproblems' von Sylvester aus der Theorie der geometrischen Wahrscheinlichkeiten. *Berichte über die Verhandlungen der Sächsischen Akademie der Wissenschaften zu Leipzig. Mathematisch-Physische Klasse*, 69:436–453, 1917.
- [Bla23] W. Blaschke. *Vorlesungen über Differentialgeometrie und geometrische Grundlagen von Einsteins Relativitätstheorie II*. Die Grundlehren der mathematischen Wissenschaften. Springer, 1923.
- [BSV21] M. Balko, M. Scheucher, and P. Valtr. Holes and Islands in Random Point Sets. *Random Structures & Algorithms*, 2021.
- [BV04] I. Bárány and P. Valtr. Planar point sets with a small number of empty convex polygons. *Studia Scientiarum Mathematicarum Hungarica*, 41(2):243–266, 2004.
- [CL21] D. Conlon and J. Lim. Fixing a hole. <http://arXiv.org/abs/2108.07087>, 2021.
- [EG15] L. C. Evans and R. F. Gariepy. *Measure theory and fine properties of functions*. Textbooks in Mathematics. CRC Press, revised edition, 2015.
- [Erd78] P. Erdős. Some more problems on elementary geometry. *Australian Mathematical Society Gazette*, 5:52–54, 1978.
- [FMHM15] R. Fabila-Monroy, C. Huemer, and D. Mitsche. Empty non-convex and convex four-gons in random point sets. *Studia Scientiarum Mathematicarum Hungarica*, 52(1):52–64, 2015.
- [Ger08] T. Gerken. Empty Convex Hexagons in Planar Point Sets. *Discrete & Computational Geometry*, 39(1):239–272, 2008.

- [Gro73] H. Groemer. On some mean values associated with a randomly selected simplex in a convex set. *Pacific J. Math.*, 45:525–533, 1973.
- [Hor83] J. Horton. Sets with no empty convex 7-gons. *Canadian Mathematical Bulletin*, 26:482–484, 1983.
- [Joh48] F. John. Extremum problems with inequalities as subsidiary conditions. In *Studies and Essays, presented to R. Courant on his 60th birthday, January 8, 1948*, pages 187–204. Interscience Publ., New York, 1948.
- [Kle69] V. Klee. Research Problems: What is the Expected Volume of a Simplex Whose Vertices are Chosen at Random from a Given Convex Body? *American Mathematical Monthly*, 76(3):286–288, 1969.
- [KM88] M. Katchalski and A. Meir. On empty triangles determined by points in the plane. *Acta Mathematica Hungarica*, 51(3-4):323–328, 1988.
- [Mac51] A. M. Macbeath. A compactness theorem for affine equivalence-classes of convex regions. *Canad. J. Math.*, 3:54–61, 1951.
- [Mat02] J. Matoušek. *Lectures on Discrete Geometry*, volume 212 of *Graduate Texts in Mathematics*. Springer, New York, 2002.
- [Mec04] M. W. Meckes. Volumes of symmetric random polytopes. *Archiv der Mathematik*, 82(1):85–96, 2004.
- [MV10] D. Micciancio and P. Voulgaris. Faster exponential time algorithms for the shortest vector problem. In *Proceedings of the Twenty-First Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1468–1480. SIAM, Philadelphia, PA, 2010.
- [Nic07] M. C. Nicolas. The Empty Hexagon Theorem. *Discrete & Computational Geometry*, 38(2):389–397, 2007.
- [RT19] M. Reitzner and D. Temesvari. Stars of empty simplices. <http://arxiv.org/abs/1808.08734>, 2019.
- [Sch] M. Scheucher. Supplemental program for computer experiments. http://page.math.tu-berlin.de/~scheuch/suppl/holes_in_random_sets/.
- [Sch88] R. Schneider. Random approximation of convex sets. *Journal of Microscopy*, 151(3):211–227, 1988.
- [Sch14] R. Schneider. *Convex bodies: the Brunn-Minkowski theory*, volume 151 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, expanded edition, 2014.
- [SW08] R. Schneider and W. Weil. *Stochastic and integral geometry*. Probability and its Applications. Springer, 2008.

- [Val92] P. Valtr. Sets in \mathbb{R}^d with no large empty convex subsets. *Discrete Mathematics*, 108(1):115–124, 1992.
- [Val95] P. Valtr. On the minimum number of empty polygons in planar point sets. *Studia Scientiarum Mathematicarum Hungarica*, pages 155–163, 1995.