

# FLOWS WITH TIME-REVERSAL SYMMETRIC LIMIT SETS ON SURFACES

TOMOO YOKOYAMA

ABSTRACT. The Long-time behavior of orbits is one of the most fundamental properties in dynamical systems. Poincaré studied the Poisson stability, which satisfies a time-reversal symmetric condition, to capture a property whether points return arbitrarily near the initial positions after a sufficiently long time. Birkhoff introduced and studied the concept of non-wandering points, which is one of time-reversal symmetric conditions. Moreover, minimality and pointwise periodicity satisfy the time-reversal symmetric condition for limit sets. Athanassopoulos characterized a flow that is either irrational or Denjoy by using a time-reversal symmetric condition for limit sets. In this paper, we refine the characterization of irrational or Denjoy flows into one of the flows with the time-reversal symmetric condition for limit sets. Moreover, we demonstrate that limit quasi-circuits can become like the boundary of the lakes of Wada.

## 1. INTRODUCTION

The Long-time behavior of orbits is one of the most fundamental properties in dynamical systems. In [6, 7], Poincaré studied the Poisson stability, which satisfies a time-reversal symmetric condition, to capture a property whether points return arbitrarily near the initial positions after a sufficiently long time. In [3], Birkhoff introduced and studied the concept of non-wandering points, which is one of time-reversal symmetric conditions, by introducing the concepts of  $\omega$ -limit set and  $\alpha$ -limit set of a point. Moreover, minimality and pointwise periodicity satisfy the time-reversal symmetric condition for limit sets. Athanassopoulos characterized a flow that is either irrational or Denjoy on a closed surface by using a time-reversal symmetric condition for non-closed Poisson stable orbits [2]. In this paper, we refine the characterization of irrational or Denjoy flows into one of the flows with time-reversal symmetric limit sets. To describe the statement, we define some concepts. A point  $x$  has the time-reversal symmetric limit set if  $\alpha(x) = \omega(x)$ . A subset (resp. flow) has the time-reversal symmetric limit set if any points in it (resp. in the whole space) have the time-reversal symmetric limit sets. A non-recurrent orbit is connecting quasi-separatrix if each of the  $\omega$ -limit and the  $\alpha$ -limit sets is contained in a boundary component of the singular point set. A closed connected invariant subset is a non-trivial quasi-circuit if it is a boundary component of an open annulus, contains a non-recurrent point, and consists of non-recurrent points and singular

---

*Date:* November 29, 2021.

*2020 Mathematics Subject Classification.* Primary 37E35; Secondary 70H33.

*Key words and phrases.* Surface flow, Time-reversal symmetric conditions, Lakes of Wada. The author was partially supported by JSPS Grant Number 20K03583.

points. A non-trivial quasi-circuit is a limit quasi-circuit if it is the  $\omega$ -limit or  $\alpha$ -limit set of a point outside of it. Then we have the following characterization of flows with time-reversal symmetric limit sets.

**Theorem A.** *A flow on a compact connected surface has time-reversal symmetric limit sets if and only if one of the following statements holds exclusively:*

- (1) *The flow is an irrational flow.*
- (2) *The flow is a Denjoy flow.*
- (3) *The surface is either a torus or a Klein bottle, and there is a limit cycle whose complement is a transverse annulus.*
- (4) *Each recurrent orbit is closed, and each non-recurrent orbit is either a connecting quasi-separatrix with time-reversal symmetric limit sets or is contained in an open invariant transverse annulus with time-reversal symmetric limit sets which are limit quasi-circuits.*

*In case (4), the periodic point set is open, and the  $\omega$ -limit and  $\alpha$ -limit sets of any non-recurrent point are either contained in the singular point set or a limit quasi-circuit.*

Notice that limit quasi-circuits for a flow with time-reversal symmetric limit sets on a compact connected surface can become like the boundary of the lakes of Wada (see Example 3 for details).

The present paper consists of four sections. In the next section, as preliminaries, we introduce fundamental concepts. In §3, we topologically characterize flows with time-reversal symmetric limit sets. In the final section, some examples are illustrated to describe the time-reversal symmetric condition for limit sets.

## 2. PRELIMINARIES

**2.1. Topological notion.** A surface is a two dimensional paracompact manifold with or without boundary.

Recall that a one-dimensional cell complex  $\gamma$  is essential in a compact surface  $S$  if and only if  $\gamma$  is not null homotopic in  $S^*$ , where  $S^*$  is the resulting surface from  $S$  by collapsing all boundary components into singletons. We call that a subset of a surface  $S$  is essential if it is not null homotopic in the resulting surface  $S^*$  from  $S$  by collapsing all boundary components into singletons. Here a singleton is a set consisting of a point.

**2.1.1. End completions.** Consider the direct system  $\{K_\lambda\}$  of compact subsets of a topological space  $X$  and inclusion maps such that the interiors of  $K_\lambda$  cover  $X$ . There is a corresponding inverse system  $\{\pi_0(X - K_\lambda)\}$ , where  $\pi_0(Y)$  denotes the set of connected components of a space  $Y$ . Then the set of ends of  $X$  is defined to be the inverse limit of this inverse system. Notice that  $X$  has one end  $x_\mathcal{U}$  for each sequence  $\mathcal{U} := (U_i)_{i \in \mathbb{Z}_{>0}}$  with  $U_i \supseteq U_{i+1}$  such that  $U_i$  is a connected component of  $X - K_{\lambda_i}$  for some  $\lambda_i$ . Considering the disjoint union  $X_{\text{end}}$  of  $X$  and  $\{\pi_0(X - K_\lambda)\}$  as set, a subset  $V$  of the union  $X_{\text{end}}$  is an open neighborhood of an end  $x_\mathcal{U}$  if there is some  $i \in \mathbb{Z}_{>0}$  such that  $U_i \subseteq V$ . Then the resulting topological space  $X_{\text{end}}$  is called the end completion (or end compactification) of  $X$ . Note that the end completion is not compact in general.

**2.2. Notion of dynamical systems.** By a flow, we mean a continuous  $\mathbb{R}$ -action on a surface. Let  $v: \mathbb{R} \times S \rightarrow S$  be a flow on a compact surface  $S$ . Then  $v_t := v(t, \cdot)$




FIGURE 1. Left, flow box; middle, periodic annulus; right, transverse annulus.

is a homeomorphism on  $S$ . A subset of  $S$  is said to be invariant (or saturated) if it is a union of orbits. An invariant closed subset is minimal if it has no non-empty proper invariant closed subsets. For a closed invariant set  $\gamma$ , define the stable manifold  $W^s(\gamma) := \{y \in X \mid \omega(y) \subseteq \gamma\}$  and the unstable manifold  $W^u(\gamma) := \{y \in X \mid \alpha(y) \subseteq \gamma\}$ . A point  $x$  of  $S$  is singular if  $x = v_t(x)$  for any  $t \in \mathbb{R}$ , is periodic if there is a positive number  $T > 0$  such that  $x = v_T(x)$  and  $x \neq v_t(x)$  for any  $t \in (0, T)$ , and if closed if it is either singular or periodic. Denote by  $\text{Sing}(v)$  (resp.  $\text{Per}(v)$ ,  $\text{Cl}(v)$ ) the set of singular (resp. periodic, closed) points. For a point  $x \in S$ , define the  $\omega$ -limit set  $\omega(x)$  and the  $\alpha$ -limit set  $\alpha(x)$  of  $x$  as follows:  $\omega(x) := \bigcap_{n \in \mathbb{R}} \overline{\{v_t(x) \mid t > n\}}$ ,  $\alpha(x) := \bigcap_{n \in \mathbb{R}} \overline{\{v_t(x) \mid t < -n\}}$ . A point  $x$  of  $S$  is Poisson stable (or strongly recurrent) if  $x \in \omega(x) \cap \alpha(x)$ . A point  $x$  of  $S$  is recurrent if  $x \in \omega(x) \cup \alpha(x)$ . Denote by  $\text{P}(v)$  (resp.  $\text{R}(v)$ ) the set of non-recurrent (resp. non-closed recurrent) points. Then  $S = \text{Cl}(v) \sqcup \text{P}(v) \sqcup \text{R}(v)$ . The closure of a non-closed recurrent orbit is called a Q-set (or quasi-minimal set). An orbit is singular (resp. periodic, closed, recurrent, Poisson stable) if it consists of singular (resp. periodic, closed, recurrent, Poisson stable) points. A periodic orbit is a limit cycle if it is the  $\omega$ -limit or  $\alpha$ -limit set of a point outside of it. A flow is Poisson stable (resp. recurrent) if each point is Poisson stable (resp. recurrent).

An invariant subset is a quasi-Q-set if it is an  $\omega$ -limit or  $\alpha$ -limit set of a point and intersects an essential closed transversal infinitely many times.

**2.2.1. Flow boxes, periodic annuli, and transverse annuli.** A closed trivial flow box is homeomorphic to  $[0, 1]^2$  each of whose orbit arcs correspond to  $[0, 1] \times \{t\}$  for some  $t \in [0, 1]$  as on the left of Figure 1. An open trivial flow box is homeomorphic to  $(0, 1)^2$  each of whose orbit arcs correspond to  $(0, 1) \times \{t\}$  for some  $t \in (0, 1)$ . A subset  $U$  is an open (resp. closed) periodic annulus if there are an interval  $I := (0, 1)$  (resp.  $I := [0, 1]$ ) and a homeomorphism  $h: U \rightarrow I \times \mathbb{S}^1$  whose images of any orbits in  $U$  are maximal orbits  $\{t\} \times \mathbb{S}^1$  for some  $t \in I$  as on the middle of Figure 1. A subset  $U$  is an open (resp. closed) transverse annulus if there are an interval  $I := (0, 1)$  (resp.  $I := [0, 1]$ ) and a homeomorphism  $h: U \rightarrow I \times \mathbb{S}^1$  whose images of any maximal orbit arcs in  $U$  are maximal orbit arcs  $I \times \{\theta\}$  for some  $\theta \in \mathbb{S}^1$  as on the right of Figure 1.

**2.2.2. Time-reversal symmetric conditions for subsets.** A point  $x$  has the time-reversal symmetric limit set if  $\alpha(x) = \omega(x)$ . A subset has the time-reversal symmetric limit set if any points in it have the time-reversal symmetric limit sets. A flow has time-reversal symmetric limit sets if each point has the time-reversal symmetric limit set.

3. CHARACTERIZATION OF FLOWS WITH TIME-REVERSAL SYMMETRIC LIMIT SETS

We have the following observation.

**Lemma 3.1.** *The following statements are equivalent for a flow on a compact connected surface:*

- (1) *The flow is either an irrational flow or a Denjoy flow.*
- (2) *There are a Poisson stable orbit and its open invariant neighborhood with time-reversal symmetric limit sets.*
- (3) *The flow has a non-closed recurrent point and time-reversal symmetric limit sets.*

*Proof.* [2, Theorem 1.1] implies that assertions (1) and (2) are equivalent. Assertion (1) implies that assertion (3). Let  $v$  be a flow on a compact connected surface  $S$ . Suppose that  $v$  has a non-closed recurrent point  $x$  and time-reversal symmetric limit sets. By [4, Theorem VI], the closure of the non-closed recurrent orbit  $O(x)$  contains uncountably many Poisson orbits whose closures are  $\overline{O}$ . Therefore, assertion (2) holds.  $\square$

We obtain the following statement.

**Lemma 3.2.** *The following statements hold for a flow without non-closed recurrent points on a compact connected surface  $S$ :*

- (1) *If there are a limit cycle and its open invariant neighborhood with time-reversal symmetric limit sets, then the surface is either a torus or a Klein bottle and its complement is a transverse annulus.*
- (2) *If there are a non-recurrent point  $x$  whose  $\omega(x)$ -limit set is a limit quasi-circuit and the open invariant neighborhood of  $\omega(x)$  with time-reversal symmetric limit sets, then the connected component of  $W^u(\omega(x)) - \omega(x) = W^s(\omega(x)) - \omega(x)$  containing  $x$  is an open invariant transverse annulus.*

*Proof.* Let  $\gamma$  be a limit cycle or limit quasi-circuit, an open invariant neighborhood  $U$  of  $\gamma$ , and  $x$  a point with  $\gamma = \omega(x)$ . Then  $x \in U$  and  $\gamma = \alpha(x) = \omega(x)$ . Fix any non-recurrent point  $y$  in  $\gamma$  and a small open transverse arc  $I$  intersecting  $\{y\} = I \cap \gamma$ . [9, Theorem A] implies that if  $\gamma$  is not periodic then  $\alpha(y) = \omega(y) \subseteq \text{Sing}(v)$ . Let  $I_-$  and  $I_+$  be the connected components of  $I - \{y\}$  with  $O^+(x) \cap I \subset I_+$  and  $O^-(x) \cap I \subset I_-$ . Taking  $I$  small if necessary, we may assume that  $O^+(x) \cap I_+$  converges to  $y$  monotonically in  $I$  and that  $O^-(x) \cap I_-$  converges to  $y$  monotonically in  $I$ . For any  $i \in \mathbb{Z}_{>0}$ , denote by  $x_i$  the  $i$ -th intersection  $O^+(x) \cap I_+$  for  $v$  and by  $x_{-i}$  the  $i$ -th intersection  $O^-(x) \cap I_-$  for  $-v$ . Let  $f_v : I \rightarrow I$  be the first return map on  $I$  induced by  $v$ ,  $C_{a,b} \subset O(x)$  the orbit arc from  $a$  to  $b$ , and  $I_{a,b} \subset I$  the subinterval between  $a$  and  $b$  of  $I$ . Then  $x_{i+1} = f_v(x_i)$  and  $x_{-i-1} = f_v(x_{-i})$  for any  $i \in \mathbb{Z}_{>0}$ . For any  $i \in \mathbb{Z}_{>0}$ , put  $C_i := C_{x_i, x_{i+1}}$ ,  $J_i := I_{x_i, x_{i+1}}$ ,  $C_{-i} := C_{x_{-i}, x_{-i-1}}$ ,  $J_{-i} := I_{x_{-i}, x_{-i-1}}$ . Then the unions  $\mu_i := C_i \cup J_i$  and  $\mu_{-i} := C_{-i} \cup J_{-i}$  for any  $i \in \mathbb{Z}_{>0}$  are pairwise disjoint loops. By compactness of  $S$ , by taking a subarc of  $I$ , we may assume that the connected component of  $S - (\mu_i \sqcup \mu_{i+2})$  (resp.  $S - (\mu_{-i} \sqcup \mu_{-i-2})$ ) containing  $\mu_{i+1}$  (resp.  $\mu_{-i-1}$ ) for any  $i \in \mathbb{Z}_{>0}$  is an open annulus. Applying [9, Lemma 3.1] to the loops  $\mu_i := C_i \cup J_i$  and  $\mu_{-i} := C_{-i} \cup J_{-i}$  for any  $i \in \mathbb{Z}_{>0}$ , there are closed transversal  $\gamma_i$  and  $\gamma_{-i}$  near  $\mu_i$  and  $\mu_{-i}$  respectively. By construction, the connected component of  $S - (\gamma_i \sqcup \gamma_{i+2})$  (resp.  $S - (\gamma_{-i} \sqcup \gamma_{-i-2})$ ) containing  $\gamma_{i+1}$  (resp.  $\gamma_{-i-1}$ ) for any  $i \in \mathbb{Z}_{>0}$  is an open transverse annulus  $A_i$  (resp.  $A_{-i}$ ). Then the union  $A_+ := \bigsqcup_{i=1}^{\infty} A_i \sqcup \gamma_{i+1}$  is a positive invariant open transverse annulus

and the union  $A_- := \bigsqcup_{i=1}^{\infty} A_{-i} \sqcup \gamma_{-i-1}$  is a negative invariant open transverse annulus. By construction, we have that  $\gamma = \omega(x) = \omega(z_+)$  for any  $z_+ \in A_+$  and  $\gamma = \alpha(x) = \alpha(z_-)$  for any  $z_- \in A_-$ . Moreover, the union  $A_- \sqcup \gamma \sqcup A_+$  is a neighborhood of  $\gamma \setminus \text{Sing}(v) = \gamma \cap (\text{Per}(v) \sqcup \text{P}(v))$  and is contained in any open invariant neighborhood of  $\gamma$ . Then  $A_- \sqcup \gamma \sqcup A_+$  has time-reversal symmetric limit sets. Since any  $\omega$ -limit and  $\alpha$ -limit sets of points in  $A_- \sqcup \gamma \sqcup A_+$  are time-reversal symmetric, the intersection  $O^-(z_+) \cap A_-$  for any  $z_+ \in \gamma_1$  is negative invariant and the intersection  $O^+(z_-) \cap A_+$  for any  $z_- \in \gamma_{-1}$  is positive invariant. For any  $z_+ \in \gamma_1$ , denote by  $C_{z_+}$  the closed orbit arc from  $z_+$  to the intersection  $\gamma_{-1} \cap O^-(z_+)$ . By the flow box theorem for a continuous flow on a compact surface (cf. Theorem 1.1, p.45[1]), for any  $z_+ \in \gamma_1$ , there is a closed flow box  $B_{z_+}$  containing  $C_{z_+}$  such that  $\bigsqcup_{z'_+ \in \gamma_1 \cap B_{z_+}} C_{z'_+} = B_{z_+}$ . From the compactness of  $\gamma_1$ , there are finitely many closed flow boxes  $B_{z_{+,i}}$  whose union is a cover of  $\gamma_1$ . Then the union  $A_0 := \bigcup_i B_{z_{+,i}}$  is a closed transverse annulus whose boundary is  $\gamma_1 \sqcup \gamma_{-1}$ . Therefore the disjoint union  $A := A_- \sqcup A_0 \sqcup A_+$  is an invariant open annulus with  $x \in A$ . Since the union  $A \sqcup \gamma$  is a neighborhood of  $\gamma \setminus \text{Sing}(v) = \gamma \cap (\text{Per}(v) \sqcup \text{P}(v))$ , the connected component of  $W^u(\gamma) - \gamma = W^s(\gamma) - \gamma$  containing  $x$  is the open invariant transverse annulus  $A$ .

Suppose that  $\gamma$  is a limit cycle. Then the invariant open subset  $A \sqcup \gamma$  is a neighborhood of  $\gamma$  with  $\gamma = \partial A$ . Therefore  $A \sqcup \gamma$  is closed and open. Since  $S$  is connected, we have that  $S = U = A \sqcup \gamma$  is the resulting surface from a closed annulus gluing the boundary and is either a torus or a Klein bottle.  $\square$

Notice that, for any positive integer  $k$ , there is a limit quasi-circuit  $\gamma$  for a flow with time-reversal symmetric limit sets but without non-closed recurrent points on a compact connected surface such that the complement  $W^u(\gamma) - \gamma = W^s(\gamma) - \gamma$  consists of  $k$  connected components. Indeed, by a Wada-lakes-like construction, we can construct such a limit quasi-circuit whose complement consists of invariant subsets in  $\text{Cl}(v)$  and open invariant transverse annuli (see Example 3 for details).

The previous lemma implies the following statement.

**Corollary 3.3.** *The following statements are equivalent for a flow on a compact connected surface:*

- (1) *The surface is either a torus or a Klein bottle, and there is a limit cycle whose complement is a transverse annulus.*
- (2) *There are a limit cycle and its open invariant neighborhood with time-reversal symmetric limit sets.*
- (3) *The flow has a limit cycle and time-reversal symmetric limit sets.*

*Proof.* Lemma 3.2 implies that assertions (1) and (2) are equivalent. Assertion (1) implies assertion (3). Assertion (3) implies assertion (2).  $\square$

The non-existence of quasi-Q-sets consisting of singular points and non-recurrent points is followed from the time-reversal symmetric condition for limit sets.

**Lemma 3.4.** *A flow with time-reversal symmetric limit sets on a compact connected surface has no quasi-Q-sets that consist of singular points and non-recurrent points.*

*Proof.* Let  $v$  be a flow with time-reversal symmetric limit sets on a compact connected surface  $S$ . Assume that there is a quasi-Q-set  $Q$  that consists of singular

points and non-recurrent points. Fix a point  $x$  whose  $\omega$ -limit and  $\alpha$ -limit sets are  $Q$ . Denote by  $T$  the connected component of the complement  $S - (\text{Sing}(v) \cap Q)$  containing  $x$ . Then  $T$  is an invariant open subset of  $S$  such that  $\partial T \subseteq \text{Sing}(v) \cap Q$ . By [8, Theorem 3], the open surface  $T$  is homeomorphic to the resulting surface from a compact surface by removing a closed totally disconnected subset and so the end completion  $T_{\text{end}}$  of  $T$  is a compact surface. Let  $v_{\text{end}}$  be the resulting flow on  $T_{\text{end}}$  from  $v$  by adding singular points. Collapsing any connected component of  $\text{Sing}(v) \cap Q$  into singletons, we obtain the resulting space  $T'$  from  $T \sqcup (\text{Sing}(v) \cap Q)$ . Let  $p_0: T \sqcup (\text{Sing}(v) \cap Q) \rightarrow T'$  be the quotient map. Then  $T$  is a proper subset of both  $T_{\text{end}}$  and  $T'$ . By construction, the compact metrizable space  $T'$  can be obtained from  $T_{\text{end}}$  by identifying ends. Let  $p: T_{\text{end}} \rightarrow T'$  be the quotient map. Note that the restriction  $p|_T$  is identical. Put  $U := S - (T \sqcup (\text{Sing}(v) \cap Q))$ . Then  $U$  is an open invariant subset of  $S$  with  $S = T \sqcup (\text{Sing}(v) \cap Q) \sqcup U$ .

We claim that the set of points on  $T'$  whose inverse images by  $p$  are disconnected are finite. Indeed, assume that there are infinitely many singletons  $e_i$  on  $T'$  whose images are disconnected. Then there are infinitely many loops  $\gamma_i \subset T$  which bound open disks intersecting  $p^{-1}(e_i)$  such that  $p(\gamma_i) = e_i$  and that the complement  $T' - \bigsqcup_i \gamma_i$  is connected. Since the resulting space from a disconnected topological space by collapsing connected subsets into singletons is disconnected, the set difference  $p_0^{-1}(T' - \bigsqcup_i \gamma_i) = (T - \bigsqcup_i \gamma_i) \sqcup (\text{Sing}(v) \cap Q)$  is connected. Since the closures of any connected components of  $U$  intersects  $\text{Sing}(v) \cap Q$ , the union  $(T - \bigsqcup_i \gamma_i) \sqcup (\text{Sing}(v) \cap Q) \sqcup U = S - \bigsqcup_i \gamma_i$  is connected, which contradicts the compactness of  $S$ .

We claim that any inverse images of singletons on  $T'$  by  $p$  have at most finitely many connected components. Indeed, assume that there is a point  $e$  on  $T'$  whose inverse image by  $p$  has infinitely many connected components. Then there are infinitely many loops  $\gamma_i \subset T$  which bound open disks intersecting  $p^{-1}(e)$  such that  $p(\gamma_i) = e$  and that the complement  $T' - \bigsqcup_i \gamma_i$  is connected. As above, the complement  $S - \bigsqcup_i \gamma_i$  is connected, which contradicts the compactness of  $S$ .

Thus the compact metrizable space  $T'$  can be obtained from  $T_{\text{end}}$  by identifying finitely many ends finitely many times. Let  $E \subset \text{Sing}(v_{\text{end}})$  be the finite union of ends in  $T_{\text{end}}$  each of whose inverse images by  $p$  is not a singleton. By definition of quasi-Q-set, there is a closed transversal  $\mu \subset T$  intersecting  $Q$  infinitely many times.

We claim that there are infinitely many pairwise disjoint non-recurrent orbits in  $Q$  intersecting  $\mu$ . Indeed, by [9, Theorem A], the closures in  $T_{\text{end}}$  of any non-recurrent orbits in  $Q$  is a closed interval and so intersects  $\mu$  at most finitely many times. Therefore any non-recurrent orbits in  $Q$  intersect  $\mu$  at most finitely many times. This means that there are infinitely many pairwise disjoint non-recurrent orbits in  $Q$  intersecting  $\mu$ .

From [9, Theorem A], the  $\omega$ -limit and  $\alpha$ -limit sets of non-recurrent orbits in  $Q$  are contained in  $\text{Sing}(v) \cap Q$ . Then there are infinitely many pairwise disjoint non-recurrent orbits  $O_i$  in  $Q \subset T$  intersecting  $\mu$ . Then the inverse images  $p^{-1}(p_0(\omega(O_i))) \subset T_{\text{end}}$  of ends are singletons which singular points for any  $i$ . Since  $T'$  is obtained from  $T_{\text{end}}$  by identifying finitely many ends finitely many times, there are infinitely many loops  $\nu_i \subset T_{\text{end}}$  such that either  $\nu_i = \{\omega_{v_{\text{end}}}(O_{i_1})\} \sqcup O_{i_1}$  or  $\{\omega_{v_{\text{end}}}(O_{i_1}), \alpha_{v_{\text{end}}}(O_{i_1})\} \sqcup O_{i_1} \sqcup O_{i_2}$  for some  $i_1, i_2$ . Then any pair of distinct loops  $\nu_i$  and  $\nu_j$  intersects at most two points. Since any loops  $\nu_i$  intersect the closed

transversal  $\mu$ , the loops  $p(\nu_i)$  are essential in  $T'$ . Therefore any loops  $\nu_i$  either is essential or bound open disks containing ends in  $E$ .

We claim that there are no infinitely many pairwise distinct loops in  $\bigcup_i \nu_i$ . Indeed, assume that there are infinitely many pairwise disjoint loops in  $\bigcup_i \nu_i$ . By renumbering of  $O_i$ , we may assume that loops  $\nu_i$  are pairwise disjoint. From the finite existence of genus and boundary components, there are infinitely many loops such that any pairs of distinct such loops bound invariant open annuli. Therefore there are a closed annulus  $A$  whose boundary consists of two distinct loops  $\nu'$  and  $\nu''$  in  $\bigcup_i \nu_i$  and a loop  $\nu'''$  in  $\bigcup_i \nu_i$  such that  $A - \nu'''$  is a disjoint union of two invariant annuli  $A' \sqcup A''$ . By  $\nu' \sqcup \nu'' \sqcup \nu''' \subset p^{-1}(p(Q)) = \omega_{v_{\text{end}}}(x)$ , we have  $x \in A' \cap A'' = \emptyset$ , which contradict  $x \notin \emptyset$ .

Thus we may assume that  $\nu_i \cap \nu_j \neq \emptyset$  for any  $i \neq j$ . We claim that there are no infinitely many loops  $\nu_i$  such that  $\nu_i = \{\omega_{v_{\text{end}}}(O_{i_1})\} \sqcup O_{i_1} \subset T_{\text{end}}$  for some  $i_1$ . Indeed, assume that there are infinitely many loops  $\nu_i$  such that  $\nu_i = \{\omega_{v_{\text{end}}}(O_{i_1})\} \sqcup O_{i_1}$  for some  $i_1$ . Since  $v$  has the time-reversal symmetric limit set, by renumbering of  $O_i$ , we may assume that there is a singular points  $s$  such that  $\nu_i = \{s\} \sqcup O_i$  for any  $i$ . Then  $\bigcup_i \nu_i = \{s\} \sqcup \bigcup_i O_i$ . From the finite existence of genus and boundary components, there are infinitely many loops such that any pairs of distinct orbits  $O_i$  and  $O_j$  equipped with  $s$  bound invariant open disks. Therefore there are a closed disk  $B$  whose boundary is  $\nu' \sqcup \{s\} \sqcup \nu''$  in  $\bigcup_i \nu_i$  and an orbit  $\nu'''$  in  $\bigcup_i \nu_i$  such that  $B - \nu'''$  is a disjoint union of two invariant disks  $B' \sqcup B''$ . By  $\nu' \sqcup \nu'' \sqcup \nu''' \subset p^{-1}(p(Q)) = \omega_{v_{\text{end}}}(x)$ , we have  $x \in B' \cap B'' = \emptyset$ , which contradict  $x \notin \emptyset$ .

Thus there are infinitely many loops  $\nu_i$  such that  $\nu_i = \{\omega_{v_{\text{end}}}(O_{i_1}), \alpha_{v_{\text{end}}}(O_{i_1})\} \sqcup O_{i_1} \sqcup O_{i_2}$  for some  $i_1, i_2$ . Since  $T'$  is obtained from  $T_{\text{end}}$  by identifying finitely many ends finitely many times, we have that  $p(\omega_{v_{\text{end}}}(O_{i_1})) = p(\alpha_{v_{\text{end}}}(O_{i_1})) \in p(E)$  for any  $i$ , and that  $\bigcup_i \nu_i \cap \text{Sing}(v_{\text{end}}) = \bigcup_i \{\omega_{v_{\text{end}}}(O_{i_1}), \alpha_{v_{\text{end}}}(O_{i_1})\} \subseteq E$  are finite. By renumbering of  $O_i$ , there are singular points  $\omega$  and  $\alpha$  such that  $\nu_i = \{\omega, \alpha\} \sqcup O_{2i} \sqcup O_{2i-1} \subset T_{\text{end}}$  for any  $i$ . Since any loops  $\nu_i$  either is essential or bound open disks containing ends in  $E$ , by renumbering of  $O_i$ , we may assume that any loops  $\nu_i$  are homotopic relative to  $\{\omega, \alpha\}$  in  $T_{\text{end}}$  and so that any orbits  $O_i$  are homotopic relative to  $\{\omega, \alpha\}$  in  $T_{\text{end}}$ . Therefore there are pairwise distinct integers  $i, j, k$  and a closed disk  $B$  whose boundary is  $O_i \sqcup \{\omega, \alpha\} \sqcup O_j$  such that  $B - O_k$  is a disjoint union of two invariant disks  $B' \sqcup B''$ . By  $O_i \sqcup O_j \sqcup O_k \subset p_0^{-1}(p(Q)) \setminus \text{Sing}(v_{\text{end}}) \subset \omega_{v_{\text{end}}}(x) \cup \alpha_{v_{\text{end}}}(x)$ , we have  $x \in B' \cap B'' = \emptyset$ , which contradict  $x \notin \emptyset$ .

Thus there are no quasi-Q-sets that consist of singular points and non-recurrent points.  $\square$

**3.1. Proof of Theorem A.** Let  $v$  be a flow on a compact connected surface  $S$ . Suppose that  $v$  has time-reversal symmetric limit sets. Lemma 3.2 implies that the existence of non-closed recurrent points implies that the flow  $v$  is either an irrational flow or a Denjoy flow. Thus we may assume that any recurrent orbits are closed. Then  $S = \text{Cl}(v) \sqcup \text{P}(v)$ . If there is a limit cycle, then Lemma 3.2 implies that the surface is either a torus or a Klein bottle and its complement is a transverse annulus. Thus we may assume that there are no limit cycles. By [10, Lemma 3.4], the non-existence of limit cycles implies that  $\text{Per}(v)$  is open. Then  $\overline{\text{P}(v)} \subseteq \text{Sing}(v) \sqcup \text{P}(v)$ . Lemma 3.4 implies the non-existence of quasi-Q-sets that consist of singular points and non-recurrent points. [9, Theorem A] implies that any  $\omega$ -limit and  $\alpha$ -limit sets of non-recurrent points are either nowhere dense subsets in  $\partial \text{Sing}(v)$  or limit

quasi-circuits. For any point  $x$  whose  $\omega$ -limit set is a limit quasi-circuit, Lemma 3.2 implies that the connected component of  $W^u(\gamma) - \gamma = W^s(\gamma) - \gamma$  containing  $x$  is an open invariant transverse annulus. Each non-recurrent orbit is either a connecting quasi-separatrix with time-reversal symmetric limit sets or is contained in an open invariant transverse annulus with time-reversal symmetric limit sets which are limit quasi-circuits.

Conversely, if  $v$  is either an irrational flow or a Denjoy flow, then  $v$  has time-reversal symmetric limit sets. If the surface is either a torus or a Klein bottle and there is a limit cycle whose complement is a transverse annulus, then  $v$  has time-reversal symmetric limit sets. Suppose that each recurrent orbit is closed and each non-recurrent orbit is either a connecting quasi-separatrix with time-reversal symmetric limit sets or is contained in an open invariant transverse annulus with time-reversal symmetric limit sets which are limit quasi-circuits. Then  $S = \text{Cl}(v) \sqcup P(v)$  and each non-recurrent point has time-reversal symmetric limit sets. This means that  $v$  has time-reversal symmetric limit sets.

#### 4. EXAMPLES

By definition of time-reversal symmetric limit sets, we have the following observation.

**Example 1.** *Any minimal flows and pointwise periodic flows have time-reversal symmetric limit sets.*

Any generic Hamiltonian flows on closed surfaces have time-reversal symmetric limit sets. More precisely, we have the following examples.

**Example 2.** *Structurally stable Hamiltonian  $C^r$ -vector fields ( $r \in \mathbb{Z}_{>0}$ ) on closed surfaces form an open dense subset in the set of Hamiltonian  $C^r$ -vector fields and have time-reversal symmetric limit sets [5, Theorem 2.3.8, p. 74].*

We have the following example of a flow with Wada-lakes-like structures.

**Example 3.** *There is a flow with a limit quasi-circuit whose complement consists of an invariant open periodic center disk and an open invariant transverse annulus on a torus as follows. Indeed, let  $v_0$  be a flow with one limit cycle and the complement is an open invariant transverse annulus as on the upper left figure of Figure 2. Replace a point in the limit cycle by a singular point, we obtain the resulting flow  $v'_0$  as on the upper middle figure of Figure 2. Replacing the singular point into a rectangle  $D$  consisting singular points, we obtain the resulting flow  $v''_0$  as on the upper right figure of Figure 2. Dividing the rectangle  $D$  into 9 equal small rectangles and replacing the small middle rectangle  $D_{22}$  into a center disk as on the middle left in Figure 3, we obtain the resulting flow  $v_1$  as on the lower left figure of Figure 2. As the construction of lakes of Wada, dig a lake as on the middle in Figure 3, we obtain the resulting flow  $v_1$  as on the lower middle figure of Figure 2. As the construction of lakes of Wada, dig another lake as on the middle right in Figure 3, we obtain the resulting flow  $v_1$  as on the lower right figure of Figure 2. As the construction of lakes of Wada, extending the center disk, and two lakes infinitely many times, the resulting flow is desired.*

*In addition, increasing center disks and lakes, we can realize arbitrarily finite numbers of connected components of the complement of the quasi-circuit. Moreover, replacing center disks into periodic Möbius bands, and cutting pairs of center disks*






FIGURE 2. A construction of a flow with a limit quasi-circuit whose complement consists of an invariant open periodic center disk and an open invariant transverse annulus




FIGURE 3. A construction of a limit quasi-circuit whose complement consists of an invariant open periodic center disk and an open invariant transverse annulus by modifying the rectangle  $D$

and gluing the boundaries, the connected components can be not only center disks and open transverse annuli but also periodic Möbius bands and periodic annuli.

REFERENCES

[1] S. K. Aranson, G. R. Beliskii, and E. Zhuzhoma. *Introduction to the qualitative theory of dynamical systems on surfaces*. American Mathematical Society, 1996.  
 [2] K. Athanassopoulos. A characterization of denjoy flows. *Bulletin of the London Mathematical Society*, 24(1):83–86, 1992.

- [3] G. D. Birkhoff. *Dynamical systems*, volume 9. American Mathematical Soc., 1927.
- [4] T. Cherry. Topological properties of the solutions of ordinary differential equations. *American Journal of Mathematics*, 59(4):957–982, 1937.
- [5] T. Ma and S. Wang. *Geometric theory of incompressible flows with applications to fluid dynamics*. Number 119. American Mathematical Soc., 2005.
- [6] H. Poincaré. Sur le problème des trois corps et les équations de la dynamique. *Acta Mathematica*, 13(1–2):1–270, 1890.
- [7] H. Poincaré. *Les méthodes nouvelles de la mécanique céleste, Tome III*. Gauthier-Villars (Paris), 1899.
- [8] I. Richards. On the classification of noncompact surfaces. *Transactions of the American Mathematical Society*, 106(2):259–269, 1963.
- [9] T. Yokoyama. A Poincaré-Bendixson theorem for flows with arbitrarily many singular points. *arXiv preprint arXiv:2109.12478*, 2017.
- [10] T. Yokoyama. Decompositions of surface flows. *arXiv preprint arXiv:1703.05501*, 2017.

APPLIED MATHEMATICS AND PHYSICS DIVISION, GIFU UNIVERSITY, YANAGIDO 1-1, GIFU, 501-1193, JAPAN,

*Email address:* tomoo@gifu-u.ac.jp