

# Flexible forward improvement iteration for infinite time horizon Markovian optimal stopping problems

Sören Christensen\*, Albrecht Irle† and Julian Peter Lemburg

November 29, 2021

In this paper, we propose an extension of the forward improvement iteration algorithm, originally introduced in Irle [2006] and recently re-considered in Miclo and Villeneuve [2021]. The main new ingredient is a flexible window parameter describing the look-ahead distance in the improvement step. We consider the framework of a Markovian optimal stopping problem in discrete time with random discounting and infinite time horizon. We prove convergence and show that the additional flexibility may significantly reduce the runtime.

**Keywords:** Discrete time Markov chain, Optimal stopping, forward improvement iteration,  $k$ -step look ahead, infinite time horizon

**AMS MSC2010:** 60G40; 60J05; 91B28; 62L15.

## 1. Introduction

We consider a Markovian optimal stopping problem with random discounting of the form

$$\sup_{\tau} \mathbb{E} \left( \left( \prod_{i=0}^{\tau-1} \alpha(Z_i) \right) g(Z_{\tau}) \middle| Z_0 = z \right).$$

As is well-known, an optimal stopping rule is (if it exists) given as a first entrance time of the underlying Markov chain  $Z$  into the stopping set, see Shiryaev [2008]. The forward forward improvement iteration algorithm, see Irle [2006], provides an upper approximation of this stopping set. More precisely, it consists of computing a decreasing sequence  $B^0 \supseteq B^1 \supseteq B^2 \supseteq \dots$  of subsets of the state space in the

---

\*Christian-Albrechts-Universität, Mathematisches Seminar, Heinrich-Hecht-Platz 6, 24118 Kiel, Germany, E-mail: christensen@math.uni-kiel.de

†Deceased on July 21, 2021

following way:  $B^{n+1}$  is constructed from  $B^n$  by taking out those states  $z$  where using the first entrance time into  $B^n$ , that is  $\inf\{n \geq 1 : Z_n \in B^n\}$ , gives more expected reward than stopping immediately. In this sense, 1-step look ahead rules are used in each iteration step to identify points which are not in the stopping set. The main finding reported in the literature are the convergence of this algorithm to the optimal value. For finite states spaces, the algorithm terminates in a finite number of steps. In the numerical examples it turns out that this number is often surprisingly low.

The aim of this paper is to study the following modification of the forward improvement iteration algorithm. Instead of considering 1-step look ahead rules,  $\kappa$ -steps look ahead rules are used in each iteration step, where  $\kappa$  denotes a window parameter  $\geq 1$  which may vary over time. This makes the analysis considerably more involved. The purpose of this paper is twofold. First, we prove convergence of this algorithm. Second, we will show in an example that using a flexible window parameter notably speeds up the algorithm.

### 1.1. Some related literature

Pricing of American and Bermudan style derivatives on a high dimensional system of underlyings is considered an enduring problem for the last decades and many different methods have been proposed. Prices for such high dimensional options are difficult, if not impossible, to compute by standard partial differential equation methods. Therefore, various Monte Carlo algorithms for pricing Bermudan options have been developed and described in the literature. For a detailed and general overview we refer to Glasserman [2003].

However, a major part of these methods is based on a backward structure in time, which requires a finite time horizon. For infinite horizon problems, the number of suitable algorithms is more limited. Here, we just mention methods being based on stochastic approximation, see Tsitsiklis and Van Roy [1999], and linear programming, see Cho and Stockbridge [2002], Christensen [2014].

Another stream of literature, which we follow here, is based in the classical method of policy iteration as proposed by Howard in Howard [1960]. Adapting this idea to problems of optimal stopping yields a method first used in Irle [1980] for tackling best choice problems with a random population size, and later called forward improvement iteration, see Irle [2006, 2009] for a detailed analysis. The method has recently been reconsidered in Miclo and Villeneuve [2021] and, e.g., applied to restricted problems. See also Presman [2011] for a related approach.

Let us mention some further related literature. Bender, Kolodko and Schoenmakers have developed a similar approach in Kolodko and Schoenmakers [2006] and Bender and Schoenmakers [2006] with some interesting new features, including a window parameter, for problems with a finite set of exercise dates. Their analysis is based on backward induction. In a continuous time framework, a related approach for one-dimensional diffusions is presented in Sonin [1999]. Let us also mention that a related form of forward algorithm to find equilibrium stopping times in

time-inconsistent problems is studied in [Christensen and Lindensjö, 2018, Section 3]. Further references are given below.

## 1.2. Structure of the paper

We will introduce the framework and basic notions in Section 2 and then state the main result (Theorem 2.2). As a key ingredient for the proof of this result, we first analyze the one-step improvement of stopping rules in Section 3, where two lemmata are moved to an appendix to enhance readability. These results are then used to prove Theorem 2.2 in Section 4. For the implementation it is interesting to note that for finite state spaces the procedure can be described in terms of linear equations as detailed in Section 5. This is then used in Section 6 where an example is given.

## 2. Setting and main result

Let  $(Z_n)_{n \in \mathbb{N}_0}$  be a time-homogeneous Markov chain on a state space  $S$  on some probability space with filtration  $\mathcal{A}_n$ ,  $n \in \mathbb{N}_0$ . To avoid technicalities, we assume  $S$  to be countable (but not necessarily finite). Let furthermore  $g : S \rightarrow \mathbb{R}$  and  $\alpha : S \rightarrow [0, 1]$ . We write  $\mathcal{T}$  for the set of all stopping rules and look at the problem of optimal stopping for the pay-off

$$X_n := \left( \prod_{i=0}^{n-1} \alpha(Z_i) \right) g(Z_n) \text{ for all } n \in \mathbb{N}_0, \quad X_\infty := \limsup_n X_n.$$

We write for simpler notation formally

$$\left( \prod_{i=0}^{\infty} \alpha(Z_i) \right) g(Z_\infty) = X_\infty$$

so that we have

$$X_\tau = \left( \prod_{i=0}^{\tau-1} \alpha(Z_i) \right) g(Z_\tau) \text{ for all stopping rules } \tau.$$

Assume as usual

$$\mathbb{E}(X_\tau | Z_0 = z) \text{ exists for all } \tau \in \mathcal{T} \text{ and } z \in S.$$

Furthermore, we assume that the integrability condition

$$\mathbb{E} \left( X_{\lim_{n \rightarrow \infty} \sigma_n} \middle| Z_0 = z \right) = \lim_{n \rightarrow \infty} \mathbb{E}(X_{\sigma_n} | Z_0 = z)$$

holds true for all  $z \in S$  and all non-decreasing sequences  $(\sigma_n)_{n \in \mathbb{N}_0}$  of stopping rules. It is easily seen that the previous assumptions are fulfilled under the standard conditions

$$\mathbb{E} \left( \sup_{n \in \mathbb{N}} |X_n| \middle| Z_0 = z \right) < \infty \text{ and } X_\infty = \lim_{n \rightarrow \infty} X_n.$$

In this framework, we consider the problem of optimal stopping for  $X$ , i.e., the problem of maximizing, for all  $z \in S$ , the expectation

$$\mathbb{E} (X_\tau | Z_0 = z)$$

over all stopping rules  $\tau \in \mathcal{T}$ . Here, we even consider a more general constraint problem of stopping, namely the maximization over  $\tau$  in

$$\mathcal{T}(B) := \{ \tau \in \mathcal{T} ; Z_\tau \in B \text{ on } \{ \tau < \infty \} \}$$

for some given set  $B \subseteq S$ . We remark that this class of problems naturally arises in problems of optimal control that are based on stopping, see – in a continuous time framework – e.g. Christensen and Sohr [2020].

**Remark 2.1.** *In the recent article Miclo and Villeneuve [2021], a (time-homogeneous) Markov chain in continuous time  $(\widehat{Z}_t)_{t \in [0, \infty[}$  operating on the discrete state space  $S$  is considered. Assuming no instantaneous jumps (i. e. between two jump times there is always a positive amount of time) and no absorption, the above Markov chain has a pure jump structure, it is a special Markov jump process and we can describe it up to explosion time completely by a sequence of jump times and a sequence of states visited with adequate parameters, see [Asmussen, 2003, p. 39-40]. When focusing on the problem of optimal stopping of the above continuous time Markov chain we can reduce this problem to one studied in the present paper.*

For all  $B \subseteq S$  and  $\sigma \in \mathcal{T}$  define

$$\tau_\sigma(B) := \inf \{ k \geq \sigma ; Z_k \in B \},$$

the time of first entrance in  $B$  at or after time  $\sigma$ . Furthermore, write for all  $B \subseteq S$  and  $k \in \mathbb{N}$

$$B^{*k} := \{ z \in B ; g(z) \geq \mathbb{E} (X_{\tau_l(B)} | Z_0 = z) \text{ for all } l = 1, \dots, k \}$$

for the set of all states such that looking  $l = 1, \dots, k$  steps ahead and then waiting until reaching  $B$  does not yield more expected reward than immediate stopping. Note that for  $B = S$ , the first entrance time into  $S^{*k}$  is called  $k$ -step look ahead rule and had been studied extensively in the early days of optimal stopping, see [Ferguson, 2008, Chapter 5] for an overview with many examples. The case  $k = 1$  corresponds to the myopic rule, which is optimal for problems of monotone type under natural assumptions, see Chow et al. [1971] for an early treatment and Christensen and Irle [2020] for an overview on more recent developments.

For each iteration in the algorithm, we now want to choose some window parameter  $k$ .  $k$  may be chosen differently in each step, so we therefore fix a window parameter function

$$\kappa : \mathbb{N} \rightarrow \mathbb{N}$$

To make the procedure precise, for a given  $B^0 \subseteq S$ , we define recursively

$$B^k := (B^{k-1})^{*\kappa(k)} \text{ for all } k \in \mathbb{N}, \quad F := \bigcap_{k \in \mathbb{N}_0} B^k.$$

Our main result is the following:

**Theorem 2.2.** *Let  $B^0 \subseteq S$ . Then  $\tau_0(F)$  is optimal in  $\mathcal{T}(B^0)$  and moreover*

$$\tau_0(B^k) \leq \tau_0(B^{k+1}) \text{ for all } k \in \mathbb{N}_0, \quad (1)$$

$$\lim_{k \rightarrow \infty} \tau_0(B^k) = \tau_0(F), \quad (2)$$

$$E\left(X_{\tau_0(B^k)} \middle| Z_0 = z\right) \leq E\left(X_{\tau_0(B^{k+1})} \middle| Z_0 = z\right) \text{ for all } k \in \mathbb{N}_0 \text{ for all } z \in S. \quad (3)$$

We will give the proof in Section 4 below.

### 3. One-step improvement

Before proving the main result, we first study a one-step policy improvement of a given stopping rule inspired by the algorithm described before.

While running numerical examples (see Section 6 below) it appears that using more general  $D \subseteq \mathbb{N}$  (usually with  $1 \in D$ ) for the improvement step instead of an initial sequence of  $\mathbb{N}$  can be of interest. Hence let us for all  $B \subseteq S$  and  $D \subseteq \mathbb{N}$  generalize the notion from above by introducing the notation

$$B^{*D} := \{z \in B ; g(z) \geq E(X_{\tau_k(B)} \middle| Z_0 = z) \text{ for all } k \in D\}.$$

For all  $B \subseteq S$ ,  $D \subseteq \mathbb{N}$ ,  $\sigma \in \mathcal{T}(B)$  and  $\rho \in \mathcal{T}(B)$  with  $\sigma \leq \rho \leq \tau_\sigma(B^{*D})$  define the *improved* stopping rule  $\hat{\rho}$  via

$$\hat{\rho} := \begin{cases} \rho, & \text{on } \{\rho = \tau_\sigma(B^{*D})\}, \\ \tau_{n+j}(B) \wedge \tau_\sigma(B^{*D}), & \text{on } \{\rho = n, \inf\{i \in D ; Z_n \notin B^{*D \leq i}\} = j\}, \end{cases} \quad (4)$$

where we use the notation  $D_{\leq i} = \{l \in D : l \leq i\}$ . We now show that  $\hat{\rho}$  is indeed an improvement of  $\rho$  in the following sense:

**Proposition 3.1.** *Let  $B \subseteq S$ ,  $D \subseteq \mathbb{N}$ ,  $\sigma \in \mathcal{T}(B)$  and*

$$\rho \in \mathcal{T}(B) \text{ with } \sigma \leq \rho \leq \tau_\sigma(B^{*D}).$$

1. It holds that

$$\hat{\rho} \in \mathcal{T}(B) \text{ with } \sigma \leq \hat{\rho} \leq \tau_\sigma(B^{*D}), \quad (5)$$

$$\rho \leq \hat{\rho}, \quad (6)$$

$$\rho + 1 \leq \hat{\rho} \text{ on } \{\rho < \tau_\sigma(B^{*D})\}. \quad (7)$$

2. If  $D = \{l \in \mathbb{N} : l \leq k\}$  for some  $k \in \mathbb{N} \cup \{\infty\}$ , then

$$E(X_\rho | Z_0 = z) \leq E(X_{\hat{\rho}} | Z_0 = z) \text{ for all } z \in S, \quad (8)$$

and

$$E(X_\sigma | Z_0 = z) \leq E(X_{\tau_\sigma(B^{*D})} | Z_0 = z) \text{ for all } z \in S. \quad (9)$$

*Proof.* We start by proving (5), (6), (7). We have

$$\rho = \hat{\rho} = \tau_\sigma(B^{*D}) \text{ on } \{\rho = \tau_\sigma(B^{*D})\} = \{\rho = \infty\} \cup \{Z_\rho \in B^{*D}\}.$$

Furthermore,  $B^{*\emptyset} = B$  and

$$B^{*D} = \bigcap_{j \in D} B^{*D \leq j}.$$

Hence

$$\{Z_n \notin B^{*D}\} = \bigcup_{j \in D} \{j = \inf \{i \in D ; Z_n \notin B^{*D \leq i}\}\} \text{ for all } n \in \mathbb{N}_0.$$

So for all  $n \in \mathbb{N}_0$  and  $j \in D$  we have that on the event  $\{\rho = n\} \cap \{j = \inf \{i \in D ; Z_n \notin B^{*D \leq i}\}\}$  it holds that

$$\rho < \rho + 1 = n + 1 \leq \overbrace{(\tau_{n+j}(B) \wedge \tau_\sigma(B^{*D}))}^{=\hat{\rho}} \leq \tau_\sigma(B^{*D}).$$

Hence

$$\{\rho < \tau_\sigma(B^{*D})\} = \{\rho < \infty\} \cap \{Z_\rho \notin B^{*D}\} = \bigcup_{n=0}^{\infty} \{\rho = n\} \cap \{Z_n \notin B^{*D}\}.$$

Now we come to the proof of (8): Let  $z \in S$  and define

$$A := \{\rho < \tau_\sigma(B^{*D})\} = \{\rho < \infty\} \cap \{Z_\rho \notin B^{*D}\}.$$

We have

$$A^c = \{\rho = \infty\} \cup \{Z_\rho \in B^{*D}\} = \{\rho = \tau_\sigma(B^{*D})\} \subseteq \{\rho = \hat{\rho}\}$$

and thus obviously

$$\mathbb{E}(\mathbf{1}_{A^c} X_\rho | Z_0 = z) \leq \mathbb{E}(\mathbf{1}_{A^c} X_{\hat{\rho}} | Z_0 = z).$$

Define for all  $n \in \mathbb{N}_0$  and  $j \in D$

$$A^{n,j} := \{\rho = n\} \cap \{j = \inf\{i \in D; Z_n \notin B^{*D \leq i}\}\}.$$

Hence

$$A = \bigsqcup_{n=0}^{\infty} \bigsqcup_{j \in D} A^{n,j}.$$

For all  $n \in \mathbb{N}_0$  and  $j \in D$  we have by Lemma A.1

$$\mathbb{E}(\mathbf{1}_{A^{n,j}} X_\rho | Z_0 = z) = \mathbb{E}(\mathbf{1}_{A^{n,j}} X_n | Z_0 = z) \leq \mathbb{E}(\mathbf{1}_{A^{n,j}} X_{\tau_{n+j}(B)} | Z_0 = z).$$

Due to dominated convergence it suffices to show for all  $n \in \mathbb{N}_0$  and  $j \in D$  the inequality

$$\mathbb{E}(\mathbf{1}_{A^{n,j}} X_{\tau_{n+j}(B)} | Z_0 = z) \leq \mathbb{E}(\mathbf{1}_{A^{n,j}} X_{\hat{\rho}} | Z_0 = z).$$

To this end, define

$$E := \{\tau_{n+j}(B) \leq \tau_\sigma(B^{*D})\}.$$

We have

$$\tau_{n+j}(B) = \tau_{n+j}(B) \wedge \tau_\sigma(B^{*D}) = \hat{\rho} \text{ on } A^{n,j} \cap E,$$

hence

$$\mathbb{E}(\mathbf{1}_{A^{n,j} \cap E} X_{\tau_{n+j}(B)} | Z_0 = z) \leq \mathbb{E}(\mathbf{1}_{A^{n,j} \cap E} X_{\hat{\rho}} | Z_0 = z).$$

For all  $m \in \mathbb{N}$  with  $m < j$  define

$$A_m := A^{n,j} \cap \{Z_{n+m} \in B^{*D}\} \cap \bigcap_{l=1}^{m-1} \{Z_{n+l} \notin B^{*D}\}.$$

For all  $m \in \mathbb{N}$  with  $m < j$  and  $j - m \in D$  it holds by Lemma A.2

$$\int_{A_m} X_{\tau_{n+j}(B)} dP_z \leq \int_{A_m} X_{n+m} dP_z = \int_{A_m} X_{\tau_\sigma(B^{*D})} dP_z = \int_{A_m} X_{\hat{\rho}} dP_z.$$

We will prove next

$$(A^{n,j} \cap E^c) = \bigsqcup_{m=1}^{j-1} A_m.$$

Due to linearity of expectation this finishes the proof.

For all  $m, l \in \mathbb{N}$  with  $m, l < j$ ,  $m \neq l$  we have

$$A_m \cap A_l \subseteq \{Z_{n+m} \in B^{*D}\} \cap \{Z_{n+m} \notin B^{*D}\} = \emptyset, \text{ hence } A_m \cap A_l = \emptyset.$$

For all  $m \in \mathbb{N}$  with  $m < j$  we have due to  $n + m < n + j \leq \tau_{n+j}(B)$  and the definition of  $A_m$

$$A_m \subseteq \{\tau_\sigma(B^{*D}) = n + m\} \subseteq \{\tau_\sigma(B^{*D}) < \tau_{n+j}(B)\} = E^c.$$

Let  $\omega \in A^{n,j} \cap E^c$ .

For all  $l \in \mathbb{N}_0$  with  $n + j \leq l < \tau_{n+j}(B)(\omega) = \inf \{l \geq n + j; Z_l(\omega) \in B\}$  we obviously have  $Z_l(\omega) \notin B$  and due to  $B^{*D} \subseteq B$  we have  $Z_l(\omega) \notin B^{*D}$ .

Since we consider

$$\omega \in A^{n,j} \cap E^c \subseteq \{\tau_\sigma(B^{*D}) < \tau_{n+j}(B)\} \cap \{\rho = n\},$$

hence

$$\tau_{n+j}(B)(\omega) > \tau_\sigma(B^{*D})(\omega) = \inf \{l \geq \sigma(\omega); Z_l(\omega) \in B^{*D}\},$$

we obtain  $n + j > \tau_\sigma(B^{*D})(\omega)$  and  $\tau_\sigma(B^{*D})(\omega) \geq \rho(\omega) = n$ . By the definition of  $A^{n,j}$  we have  $Z_n(\omega) \notin B^{*D \leq j} \supseteq B^{*D}$ , hence  $\tau_\sigma(B^{*D})(\omega) \neq n$ . So we have  $n < \tau_\sigma(B^{*D})(\omega) < n + j$ , thus it follows  $\omega \in A_{[\tau_\sigma(B^{*D})(\omega) - n]}$  by definition.

It remains to prove (9). Define  $\sigma_0 := \sigma$  and inductively

$$\sigma_{k+1} := \hat{\sigma}_k \text{ for all } k \in \mathbb{N}_0.$$

A simple induction yields

$$\sigma_k \leq \sigma_{k+1} \leq \tau_\sigma(B^{*D}) \text{ for all } k \in \mathbb{N}_0.$$

Therefore,

$$\lim_{k \rightarrow \infty} \sigma_k \leq \tau_\sigma(B^{*D})$$

and by (7)

$$\sigma_k + 1 \leq \sigma_{k+1} \text{ on } \{\sigma_k < \tau_\sigma(B^{*D})\} \text{ for all } k \in \mathbb{N},$$

hence we have

$$\sigma_0 \leq \sigma_1 \leq \sigma_2 \leq \dots \leq \lim_{k \rightarrow \infty} \sigma_k = \tau_\sigma(B^{*D}).$$

Let  $z \in S$ . We have for all  $k \in \mathbb{N}_0$  by (8)

$$\mathbb{E}(X_{\sigma_k} | Z_0 = z) \leq \mathbb{E}(X_{\sigma_{k+1}} | Z_0 = z).$$

We infer using our standing assumption

$$\begin{aligned} \mathbb{E}(X_\sigma | Z_0 = z) &= \mathbb{E}(X_{\sigma_0} | Z_0 = z) \leq \lim_{k \rightarrow \infty} \mathbb{E}(X_{\sigma_k} | Z_0 = z) \\ &= \mathbb{E}\left(X_{\lim_{k \rightarrow \infty} \sigma_k} \middle| Z_0 = z\right) = \mathbb{E}\left(X_{\tau_\sigma(B^{*D})} \middle| Z_0 = z\right) \end{aligned}$$


□



**Example 3.2.** Using the stopping rule

$$\tilde{\rho} := \begin{cases} \rho, & \text{on } \{\rho = \tau_\sigma(B^{*D})\}, \\ \tau_{n+j}(B), & \text{on } \{\rho = n, \inf \{i \in D ; Z_n \notin B^{*D \leq i}\} = j\}, \end{cases}$$

instead of  $\hat{\rho}$  in (4) looks natural at first glance. However, then the properties (5) and (9) may not hold. We can see this by the following example: Consider a Markov chain given by



and the rewards, i.e. the function  $g$

state	payment
a	3
b	4
c	1.5
d	2.5
e	2

and we will use  $D = \{1, 2\}$  with  $\alpha \equiv 1$ . We initialize the algorithm with

$$B = \{a, b, c, d, e\}.$$

Hence we have

$$B^{*\{1\}} = \{a, b, d, e\}$$

and

$$B^{*\{1,2\}} = \{b, d, e\}.$$

Now we determine  $\tilde{\rho}$  for  $\rho := \tau_\sigma(B^{*\{1\}})$ . On  $\{Z_0 \neq a\}$  we have  $\tilde{\rho} = \tau_\sigma(B^{*\{1,2\}})$ . If  $Z_0 = a$ , then  $\rho = n = 0$ , hence

$$j := \inf \{i \in D ; Z_n \notin B^{*D \leq i}\} = \inf \{i \in D ; a \notin B^{*D \leq i}\} = 2$$

and thus on  $\{Z_0 = a\}$

$$\begin{aligned} \tilde{\rho} &= \tau_{n+j}(B) = \tau_{0+2}(B) = \tau_2(B) = \inf \{n \geq 2 ; Z_n \in \{b, e\}\} \\ &> \inf \{n \geq 0 ; Z_n \in \{b, d, e\}\} = \tau_\sigma(B^{*\{1,2\}}). \end{aligned}$$

Hence it is necessary to take the minimum with  $\sigma^{*D}$  for a sufficient improvement, in the sense of having  $\tilde{\rho}$  always smaller than  $\sigma^{*D}$  and not giving a smaller expected revenue.

## 4. Proof of Theorem 2.2

*Proof.* Let  $k \in \mathbb{N}_0$ . We have

$$B^{k+1} = (B^k)^{* \kappa(k)} \supseteq B^k$$

and

$$\tau_0(B^k) \leq \tau_{\tau_0(B^k)}((B^k)^{* \kappa(k)}) = \tau_{\tau_0(B^k)}(B^{k+1}) = \tau_0(B^{k+1}).$$

Hence we can infer (1) and (2), see Proposition 3.1. (3) also immediately follows using Proposition 3.1.

Now, we come to the proof of the optimality.

(1) At first we will show that for any  $\rho \in \mathcal{T}(B^0)$  there exists

$$\rho^\infty \in \mathcal{T}(F) \text{ with } \rho^\infty \geq \rho \text{ and } \mathbb{E}(X_\rho | Z_0 = z) \leq \mathbb{E}(X_{\rho^\infty} | Z_0 = z) \text{ for all } z \in S.$$

Let  $\rho \in \mathcal{T}(B^0)$ . Define

$$\begin{aligned} \rho^k &:= \inf \left\{ n \geq \rho; Z_n \in B^k \right\} \text{ for all } k \in \mathbb{N}_0, \\ \rho^\infty &:= \inf \left\{ n \geq \rho; Z_n \in F \right\}. \end{aligned}$$

Since  $(B^k)_{k \in \mathbb{N}_0}$  is a non-increasing sequence of sets with limit  $F$ ,  $(\rho^k)_{k \in \mathbb{N}_0}$  is a non-decreasing sequence of stopping rules with limit  $\rho^\infty$ . Since  $\rho \in \mathcal{T}(B^0)$ , it furthermore holds that  $\rho = \rho^0$ . Using Proposition 3.1 and our standing assumption, it follows

$$\begin{aligned} \mathbb{E}(X_\rho | Z_0 = z) &\leq \lim_{k \rightarrow \infty} \mathbb{E}(X_{\rho^k} | Z_0 = z) \\ &\leq \mathbb{E} \left( X_{\lim_{k \rightarrow \infty} \rho^k} \middle| Z_0 = z \right) = \mathbb{E}(X_{\rho^\infty} | Z_0 = z) \text{ for all } z \in S. \end{aligned}$$

(2) Let  $\rho, \tau \in \mathcal{T}(F)$  such that  $\rho \leq \tau$ . We will now show

$$\mathbb{E}(X_\tau | Z_0 = z) \leq \mathbb{E}(X_\rho | Z_0 = z) \text{ for all } z \in S.$$

Define

$$\begin{aligned} \rho_k &:= \rho \mathbf{1}_{\{\rho = \tau\}} + \mathbf{1}_{\{\rho < \tau\}} \sum_{n=0}^{\infty} \mathbf{1}_{\{\rho = n\}} \tau_{n+1}(B^k) \text{ for all } k \in \mathbb{N}_0, \\ \rho_\infty &:= \rho \mathbf{1}_{\{\rho = \tau\}} + \mathbf{1}_{\{\rho < \tau\}} \sum_{n=0}^{\infty} \mathbf{1}_{\{\rho = n\}} \tau_{n+1}(F). \end{aligned}$$

Then we have  $\rho \leq \rho_k \leq \rho_{k+1} \leq \rho_\infty \leq \tau$  for all  $k \in \mathbb{N}_0$  and  $\lim_{k \rightarrow \infty} \rho_k = \rho_\infty$ . Furthermore we have

$$\rho + 1 = n + 1 \leq \tau_{n+1}(F) = \rho_\infty \text{ on } \{\rho < \tau\} \cap \{\rho = n\} \text{ for all } n \in \mathbb{N}_0,$$

hence

$$\rho + 1 \leq \rho_\infty \text{ on } \bigcup_{n=0}^{\infty} \{\rho = n\} \cup \{\rho < \tau\} = \{\rho \neq \tau\}. \quad (10)$$

Let  $(Z'_n)_{n \in \mathbb{N}_0}$  be an independent copy of  $(Z_n)_{n \in \mathbb{N}_0}$  and for all  $p \in \mathbb{N}_0$  and  $B \subseteq S$  let  $\tau'_p(B) = \inf \{q \geq p; Z'_q \in B\}$ . Let  $n \in \mathbb{N}_0$  and  $k \in \mathbb{N}_0$ . Since  $\rho \in \mathcal{T}(F)$  we have  $Z_\rho \in F$  on  $\{\rho < \infty\}$  and thus

$$\begin{aligned} \{\rho = n\} \cap \{\rho < \tau\} &\subseteq \{Z_n \in F\} \\ &\subseteq \{Z_n \in (B^k)^{* \kappa(k)}\} \\ &\subseteq \left\{ g(Z_n) \geq \mathbb{E} \left( \left( \prod_{i=0}^{\tau'_1(B^k)-1} \alpha(Z'_i) \right) g(Z'_{\tau'_{n+1}(B^k)}) \middle| Z'_0 = Z_n \right) \right\}, \end{aligned}$$

hence we have for all  $z \in S$

$$\begin{aligned} \int_{\{\rho=n<\tau\}} X_{\rho^k} dP_z &= \int_{\{\rho=n<\tau\}} \left( \prod_{i=0}^{\tau_{n+1}(B^k)-1} \alpha(Z_i) \right) g(Z_{\tau_{n+1}(B^k)}) dP_z \\ &= \int_{\{\rho=n<\tau\}} \mathbb{E} \left( \left( \prod_{i=0}^{\tau'_1(B^k)-1} \alpha(Z'_i) \right) g(Z'_{\tau'_{n+1}(B^k)}) \middle| Z'_0 = Z_n \right) \prod_{i=0}^{n-1} \alpha(Z_i) dP_z \\ &\leq \int_{\{\rho=n<\tau\}} g(Z_n) \prod_{i=0}^{n-1} \alpha(Z_i) dP_z \\ &= \int_{\{\rho=n<\tau\}} X_\rho dP_z. \end{aligned}$$

By dominated convergence follows

$$\mathbb{E}(X_{\rho^k} | Z_0 = z) \leq \mathbb{E}(X_\rho | Z_0 = z) \text{ for all } z \in S.$$

Letting  $k \rightarrow \infty$  we obtain

$$\mathbb{E}(X_{\rho_\infty} | Z_0 = z) \leq \mathbb{E}(X_\rho | Z_0 = z) \text{ for all } z \in S.$$

Define  $\rho^0 = \rho$  and  $\rho^k = (\rho^{k-1})_\infty$  for all  $k \in \mathbb{N}$ . Then obviously  $\rho \leq \rho^k \leq \rho^{k+1} \leq \tau$ , and  $\rho^k \in \mathcal{T}(F)$  for all  $k \in \mathbb{N}$ . Hence we have by (10)  $\lim_{k \rightarrow \infty} \rho^k = \tau$ .

It follows for all  $z \in S$  using our standing assumption

$$\mathbb{E}(X_\tau | Z_0 = z) = \mathbb{E} \left( X_{\lim_{k \rightarrow \infty} \rho^k} \middle| Z_0 = z \right) = \lim_{k \rightarrow \infty} \mathbb{E}(X_{\rho^k} | Z_0 = z) \leq \mathbb{E}(X_\rho | Z_0 = z).$$

(3) Now, we prove that  $\tau(F)$  is optimal. Let  $\sigma \in \mathcal{T}(B^0)$ . By step 1, there exists  $\tau \in \mathcal{T}(F)$  with  $\tau \geq \sigma$  such that

$$\mathbb{E}(X_\sigma | Z_0 = z) \leq \mathbb{E}(X_\tau | Z_0 = z) \text{ for all } z \in S.$$

Obviously we have  $\tau_0(F) \leq \tau$  due to  $\tau \in \mathcal{T}(F)$ . Hence it follows by step 2 that

$$\mathbb{E}(X_\sigma | Z_0 = z) \leq \mathbb{E}(X_\tau | Z_0 = z) \leq \mathbb{E}(X_{\tau_0(F)} | Z_0 = z) \text{ for all } z \in S.$$

□

## 5. Finite state space

We now assume additionally that we have a finite state space. The algorithm requires the calculation of many conditional expectations. We show in this section that these expectations can be found by solving linear equations.

Let  $\Pi$  be the transition matrix and define  $\Psi$  by  $\Psi_{z,y} := \alpha(z) \cdot \Pi_{z,y}$  for all  $z, y \in S$ .

Write for all  $p \in \mathbb{N}$  and all  $B \subseteq S$

$$h'_{B,p} : S \rightarrow \mathbb{R}, z \mapsto \mathbb{E}(X_{\tau_p(B)} | Z_0 = z) = \mathbb{E}\left(g(Z_{\tau_p(B)}) \prod_{j=0}^{\tau_p(B)-1} \alpha(Z_j) \middle| Z_0 = z\right).$$

**Proposition 5.1.** *For all  $B \subseteq S$ ,  $p \in \mathbb{N}$  and  $z \in S$  we have*

$$h'_{B,p}(z) = \sum_{y \in S} (\Psi^p)_{z,y} h'_{B,0}(y).$$

*Proof.* Let  $B \subseteq S$ .

We have for all  $p \in \mathbb{N}$  and  $z \in S$  using the Markov property

$$\begin{aligned} h'_{B,p}(z) &= \mathbb{E}\left(g(Z_{\tau_p(B)}) \prod_{j=0}^{\tau_p(B)-1} \alpha(Z_j) \middle| Z_0 = z\right) \\ &= \sum_{y \in S} \alpha(z) P(Z_1 = y | Z_0 = z) \mathbb{E}\left(g(Z_{\tau_{p-1}(B)}) \prod_{j=0}^{\tau_{p-1}(B)-1} \alpha(Z_j) \middle| Z_0 = y\right) \\ &= \sum_{y \in S} \Psi_{z,y} \mathbb{E}\left(g(Z_{\tau_{p-1}(B)}) \prod_{j=0}^{\tau_{p-1}(B)-1} \alpha(Z_j) \middle| Z_0 = y\right) \\ &= \sum_{y \in S} \Psi_{z,y} h'_{B,p-1}(y). \end{aligned}$$

Hence we have

$$h'_{B,p} = \Psi \cdot h'_{B,p-1}.$$

By induction we have for all  $p \in \mathbb{N}$

$$h'_{B,p} = \Psi^p \cdot h'_{B,0}.$$

□

We immediately obtain:

**Corollary 5.2.** *We have for all  $B \subseteq S$  and  $D \subseteq \mathbb{N}$*

$$B^{*D} = \left\{ z \in B ; g(z) \geq \sup_{p \in D} \sum_{y \in S} (\Psi^p)_{z,y} h'_{B,0}(y) \right\}.$$

We may now rewrite the approach using matrices: Define the vector  $b$  by

$$b_i := \begin{cases} 1 & i \in B, \\ 0 & i \notin B, \end{cases} \quad \text{for all } i \in S,$$

the vector  $d$  by

$$d_i := b_i \cdot g(i) \quad \text{for all } i \in S$$

and the matrix  $A$  by

$$A_{i,j} := \delta_{i,j} - (1 - b_i) \Psi_{i,j} \quad \text{for all } i, j \in S.$$

**Corollary 5.3.** *Let  $B \subseteq S$  and assume that for all  $z \in S$*

$$\alpha(z) < 1 \text{ or } P(\tau_0(B) < \infty \mid Z_0 = z) = 1. \quad (11)$$

*Then  $h'_{B,0}$  is the unique solution of the system of linear equations  $Ah = d$ .*

*Proof.* For all  $h \in \mathbb{R}^S$  we have by definition

$$d = Ah \iff \forall z \in B \quad h(z) = g(z) \wedge \forall z \notin B \quad h(z) = \alpha(z) \sum_{y \in S} P(Z_1 = y \mid Z_0 = z) h(y).$$

For all  $z \in B$  we have  $Z_0 = z \in B$  and thus  $\tau_0(B) = 0$  and  $\prod_{j=0}^{\tau_0(B)-1} \alpha(Z_j) = 1$ , hence

$$h'_{B,0}(z) = \mathbb{E} \left( g(Z_{\tau_0(B)}) \prod_{j=0}^{\tau_0(B)-1} \alpha(Z_j) \middle| Z_0 = z \right) = \mathbb{E} (g(Z_0) \cdot 1 \mid Z_0 = z) = g(z)$$

and for all  $z \notin B$  we have using the Markov property

$$\begin{aligned} h'_{B,0}(z) &= \alpha(z) \sum_{y \in S} P(Z_1 = y \mid Z_0 = z) \mathbb{E} \left( g(Z_{\tau_0(B)}) \prod_{j=0}^{\tau_0(B)-1} \alpha(Z_j) \middle| Z_0 = y \right) \\ &= \alpha(z) \sum_{y \in S} P(Z_1 = y \mid Z_0 = z) h'(y). \end{aligned}$$

So we have  $d = Ah'_{B,0}$ .

We now prove uniqueness by doing so for the corresponding homogeneous system of linear equations: Hence consider some  $h \in \mathbb{R}^S$  with  $0 = Ah$ . It is obvious that for all  $y \in B$  we have  $h(y) = 0$ . For all  $z \notin B$

$$\begin{aligned} h(z) &= \alpha(z) \sum_{y \in S} P(Z_1 = y \mid Z_0 = z) h(y) \\ &= \sum_{y \notin B} \alpha(z) P(Z_1 = y \mid Z_0 = z) h(y) \\ &= \sum_{y \notin B} \Psi_{z,y} h(y). \end{aligned}$$

Moreover,

$$\begin{aligned} |h(z)| &= \left| \sum_{y \notin B} \Psi_{z,y} h(y) \right| \leq \sum_{y \notin B} \Psi_{z,y} |h(y)| \\ &\leq \max \{ |h(x)| \ ; \ x \notin B \} \sum_{y \notin B} \Psi_{z,y}. \end{aligned}$$

Hence

$$\max \{ |h(x)| \ ; \ x \notin B \} \leq \max \{ |h(x)| \ ; \ x \notin B \} \sum_{y \notin B} \Psi_{z,y}.$$

By (11) we on the other hand have

$$\sum_{y \notin B} \Psi_{z,y} = \alpha(z) \sum_{y \notin B} \Pi_{z,y} < 1 \text{ for all } z \notin B. \quad (12)$$

We obtain

$$0 = \max \{ |h(x)| \ ; \ x \notin B \} \sum_{y \notin B} \Psi_{z,y}.$$

Hence  $h \equiv 0$ , proving the claim.  $\square$

## 6. A Numerical Experiment

To start with a toy example, consider  $\{0, \dots, 20\} \times \{0, \dots, 20\}$  as the state space for the chain and a payoff function  $g$  with  $g(z, n) = \alpha^n f(z)$  with  $f((5, 5)) = 10$ ,  $f((5, 15)) = f((15, 15)) = 0$ ,  $f((x, y)) = 5$  otherwise, see Figure 1. The transition probabilities are given in the form

$$\begin{aligned} p(x+1, y \mid x, y) &= 0.5 \cdot p_x, \quad p(x-1, y \mid x, y) = 0.5 \cdot (1 - p_x), \\ p(x, y+1 \mid x, y) &= 0.5 \cdot p_y, \quad p(x, y-1 \mid x, y) = 0.5 \cdot (1 - p_y), \end{aligned}$$

with reflecting boundaries.

With a discount-factor  $\alpha = 0.98^{1/20}$  our program recommends a stopping rule as given in Figure 2 with corresponding optimal value shown in Figure 3. For constant




Figure 1: Undiscounted Pay-Offs in the toy example




Figure 2: Stopping-Points (big squares) of the optimal stopping rule in the toy example




Figure 3: Expected Values in the toy example

$\kappa$  equivalent to  $\{1\}$  (the traditional forward improvement iteration algorithm) the program is solving the problem above within 19 iterations. The speed of solving the toy example above is so high that the effect of changing  $\kappa$  cannot be examined therein.

Thus we will increase the resolution by factor 10 in each dimension to study the effect of  $\kappa$ , hence we study a state space of  $\{0, \dots, 200\} \times \{0, \dots, 200\}$  with pay-off 10 at  $(50, 50)$ , pay-off 0 at  $(50, 150)$  and  $(150, 150)$  and 5 otherwise. We set  $\alpha = 0.9999$ , leading to an optimal stopping rule of stopping at the four surrounding points of  $(50, 150)$  and  $(150, 150)$  as well as at  $(50, 50)$  and continuing at all other points.

At first we will examine  $\kappa \equiv k$  for different  $k$ . We have to take into account two effects: With increasing  $k$ , more powers of the transition matrix have to be calculated while less iterations are needed. In this case, a choice of  $k = 5$  seems to be a good balance, see Figure 4.

## A. Some Lemmata

**Lemma A.1.** *Assume  $B \subseteq S$ ,  $D \subseteq \mathbb{N}$ ,  $n \in \mathbb{N}_0$ ,  $j \in D$ ,  $A \in \mathcal{A}_n$  with  $A \subseteq \{Z_n \in B^{*D < j} \setminus B^{*D \leq j}\}$ ,  $z \in S$ .*

*Then we have*

$$\int_A X_n dP_z \leq \int_A X_{\tau_{n+j}(B)} dP_z.$$






Figure 4: Runtime of the algorithm (in log-scale) plotted against  $\kappa \equiv k$ .

*Proof.* For all  $E \subseteq D$ ,  $j \in E$  we have

$$\begin{aligned}
 B^{*E} &\subseteq B^{*E \setminus \{j\}}, \\
 B^{*E} &= \left\{ z \in B^{*E \setminus \{j\}} ; g(z) \geq \mathbf{E} \left( X_{\tau_j(B)} \mid Z_0 = z \right) \right\}, \\
 B^{*E \setminus \{j\}} \setminus B^{*E} &= \left\{ z \in B^{*E \setminus \{j\}} ; g(z) < \mathbf{E} \left( X_{\tau_j(B)} \mid Z_0 = z \right) \right\}.
 \end{aligned}$$

Assume  $(Z'_m)_{m \in \mathbb{N}_0}$  being an independent copy of  $(Z_m)_{m \in \mathbb{N}_0}$  and

$$\tau'_p(B) = \inf \{ q \geq p ; Z'_q \in B \} \text{ for all } p \in \mathbb{N}_0.$$

Then we have

$$\begin{aligned}
& \int_A X_n dP_z \\
&= \int_A \left( \prod_{i=0}^{n-1} \alpha(Z_i) \right) g(Z_n) dP_z \\
&\leq \int_A \left( \prod_{i=0}^{n-1} \alpha(Z_i) \right) \mathbb{E} \left( g \left( Z'_{\tau'_j(B)} \right) \prod_{i=0}^{\tau'_j(B)-1} \alpha(Z'_i) \middle| Z'_0 = Z_n \right) dP_z \\
&= \int_A \left( \prod_{i=0}^{n-1} \alpha(Z_i) \right) \mathbb{E} \left( g \left( Z_{\tau_{n+j}(B)} \right) \prod_{i=n}^{\tau_{n+j}(B)-1} \alpha(Z_i) \middle| \mathcal{A}_n \right) dP_z \\
&= \int_A \left( \prod_{i=0}^{n-1} \alpha(Z_i) \right) g \left( Z_{\tau_{n+j}(B)} \right) \prod_{i=n}^{\tau_{n+j}(B)-1} \alpha(Z_i) dP_z \\
&= \int_A \left( \prod_{i=0}^{\tau_{n+j}(B)-1} \alpha(Z_i) \right) g \left( Z_{\tau_{n+j}(B)} \right) dP_z \\
&= \int_A X_{\tau_{n+j}(B)} dP_z.
\end{aligned}$$

□

**Lemma A.2.** Assume  $B \subseteq S$ ,  $D \subseteq \mathbb{N}$ ,  $s, t \in \mathbb{N}$  with  $s \leq t$  and  $t - s \in D$ ,  $A \in \mathcal{A}_s$  with  $A \subseteq \{Z_s \in B^{*D}\}$ . Then we have

$$\int_A X_{\tau_t(B)} dP_z \leq \int_A X_s dP_z.$$

*Proof.* Assume  $(Z'_n)_{n \in \mathbb{N}_0}$  being an independent copy of  $(Z_n)_{n \in \mathbb{N}_0}$  and

$$\tau'_p(B) = \inf \{q \geq p ; Z'_q \in B\} \text{ for all } p \in \mathbb{N}_0.$$

Then we have

$$\begin{aligned}
& \int_A X_{\tau_t(B)} dP_z \\
&= \int_A \left( \prod_{i=0}^{\tau_t(B)-1} \alpha(Z_i) \right) g(Z_{\tau_t(B)}) dP_z \\
&= \int_A \left( \prod_{i=0}^{s-1} \alpha(Z_i) \right) g(Z_{\tau_t(B)}) \prod_{i=s}^{\tau_t(B)-1} \alpha(Z_i) dP_z \\
&= \int_A \left( \prod_{i=0}^{s-1} \alpha(Z_i) \right) \mathbb{E} \left( g(Z_{\tau_t(B)}) \prod_{i=s}^{\tau_t(B)-1} \alpha(Z_i) \middle| \mathcal{A}_s \right) dP_z \\
&= \int_A \left( \prod_{i=0}^{s-1} \alpha(Z_i) \right) \mathbb{E} \left( g(Z'_{\tau'_{t-s}(B)}) \prod_{i=0}^{\tau'_{t-s}(B)-1} \alpha(Z'_i) \middle| Z'_0 = Z_s \right) dP_z \\
&\stackrel{Z_s \in B^{*D}}{\leq} \int_A \left( \prod_{i=0}^{s-1} \alpha(Z_i) \right) g(Z_s) dP_z \\
&\stackrel{t-s \in D}{=} \int_A X_s dP_z.
\end{aligned}$$

□

## References

- Søren Asmussen. *Applied Probability and Queues*. Applications of mathematics 51. Springer, 2003. ISBN 0387002111.
- Christian Bender and John Schoenmakers. An iterative method for multiple stopping: convergence and stability. *Advances in Applied Probability*, 38(3):729–749, 2006. ISSN 0001-8678.
- Moon Jung Cho and Richard H Stockbridge. Linear programming formulation for optimal stopping problems. *SIAM Journal on Control and Optimization*, 40(6):1965–1982, 2002.
- Yuan Shih Chow, Herbert Robbins, and David Siegmund. *Great expectations: the theory of optimal stopping*. Houghton Mifflin Company Boston, 1971. ISBN 0395053145, 9780395053140.
- Sören Christensen. A Method for Pricing American Options Using Semi-Infinite Linear Programming. *Mathematical Finance*, 24(1):156–172, 2014.

- Sören Christensen and Albrecht Irlé. The monotone case approach for the solution of certain multidimensional optimal stopping problems. *Stochastic Processes and their Applications*, 130(4):1972–1993, 2020.
- Sören Christensen and Kristoffer Lindensjö. On finding equilibrium stopping times for time-inconsistent markovian problems. *SIAM Journal on Control and Optimization*, 56(6):4228–4255, 2018.
- Sören Christensen and Tobias Sohr. A solution technique for lévy driven long term average impulse control problems. *Stochastic Processes and their Applications*, 130(12):7303–7337, 2020.
- Thomas S. Ferguson. Optimal stopping and applications. <http://www.math.ucla.edu/~tom/Stopping/Contents.html>, 2008.
- Paul Glasserman. *Monte Carlo Methods in Financial Engineering (Stochastic Modelling and Applied Probability)*. Springer, 2003. ISBN 0387004513.
- Ronald A. Howard. *Dynamic Programming and Markov Processes*. MIT Press, Cambridge, MA, 1960.
- Albrecht Irlé. On the best choice problem with random population size. *Zeitschrift für Operations Research. Serie A. Serie B*, 24(5):177–190, 1980. ISSN 0340-9422.
- Albrecht Irlé. A forward algorithm for solving optimal stopping problems. *Journal of Applied Probability*, 43(1):102–113, 2006. ISSN 0021-9002.
- Albrecht Irlé. On forward improvement iteration for stopping problems. In *Proceeding of the International Workshop of Sequential Methodologies*, Troyes, 2009.
- Anastasia Kolodko and John Schoenmakers. Iterative construction of the optimal Bermudan stopping time. *Finance and Stochastics*, 10(1):27–49, 2006. ISSN 0949-2984.
- Laurent Miclo and Stéphane Villeneuve. On the forward algorithm for stopping problems on continuous-time markov chains. *Journal of Applied Probability*, 58(4):1043–1063, 2021. doi: 10.1017/jpr.2021.11.
- Ernst Presman. A new approach to the solution of optimal stopping problem in a discrete time. *Stochastics An International Journal of Probability and Stochastic Processes*, 83(4-6):467–475, 2011.
- Albert N. Shiryaev. *Optimal stopping rules*, volume 8 of *Stochastic Modelling and Applied Probability*. Springer-Verlag, Berlin, 2008. ISBN 978-3-540-74010-0. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=2374974>. Translated from the 1976 Russian second edition by A. B. Aries, Reprint of the 1978 translation.

Isaac Sonin. The elimination algorithm for the problem of optimal stopping. *Mathematical methods of operations research*, 49(1):111–123, 1999.

John N Tsitsiklis and Benjamin Van Roy. Optimal stopping of markov processes: Hilbert space theory, approximation algorithms, and an application to pricing high-dimensional financial derivatives. *IEEE Transactions on Automatic Control*, 44(10):1840–1851, 1999.