

A product forcing model in which the Russell-nontypical sets satisfy ZFC strictly between HOD and the universe*

Vladimir Kanovei[†] Vassily Lyubetsky[‡]

Abstract

A set is nontypical in the Russell sense, if it belongs to a countable ordinal definable set. The class **HNT** of all hereditarily nontypical sets satisfies all axioms of **ZF** and the double inclusion $\mathbf{HOD} \subseteq \mathbf{HNT} \subseteq \mathbf{V}$ holds. Solving a problem recently proposed by Tzouvaras, a generic extension $\mathbf{L}[a, x]$ of \mathbf{L} , by two reals a, x , is presented in which $\mathbf{L} = \mathbf{HOD} \subsetneq \mathbf{L}[a] = \mathbf{HNT} \subsetneq \mathbf{V} = \mathbf{L}[a, x]$, so that **HNT** is a model of **ZFC** strictly between **HOD** and the universe.

1 Introduction

A set x is *nontypical with a cardinal parameter* κ , for short $x \in \mathbf{NT}_\kappa$, if it belongs to an **OD** (ordinal definable) set X of cardinality $\text{card} X < \kappa$. A set x is *hereditarily nontypical with a cardinal parameter* κ , for short $x \in \mathbf{HNT}_\kappa$, if it itself, all its elements, elements of elements, and so on, are all nontypical, in other words the *transitive closure* $\text{TC}(x)$ satisfies $\text{TC}(x) \subseteq \mathbf{NT}_\kappa$. These notions Tzouvaras [22, 21] connected with some philosophical and mathematical ideas of Bertrand Russell and works of van Lambalgen [19] *etc.* on the concept of randomness. They contribute to the ongoing study of important classes of sets in the set theoretic universe \mathbf{V} which themselves satisfy the axioms of set theory, similarly to the Gödel class \mathbf{L} of all *constructible* sets and the class **HOD** of all *hereditarily ordinal definable* sets [7].

It is clear that $\mathbf{NT}_2 = \mathbf{OD}$ and $\mathbf{HNT}_2 = \mathbf{HOD}$, thus the case $\kappa = 2$ corresponds to the ordinal definability. The classes \mathbf{NT}_ω (elements of finite ordinal definable sets) and \mathbf{HNT}_ω correspond to *algebraically definability* recently studied in [4, 5, 6]. The following classes correspond to the next cardinality level $\kappa = \omega_1$:

$$\mathbf{NT} := \mathbf{NT}_{\omega_1} \quad \text{and} \quad \mathbf{HNT} := \mathbf{HNT}_{\omega_1}.$$

Thus $x \in \mathbf{NT}$ iff x belongs to a countable **OD** set, and $x \in \mathbf{HNT}$ iff $\text{TC}(x) \subseteq \mathbf{NT}$.

The class **HNT** is transitive and, as shown in [21], satisfies all axioms of **ZF** (the axiom of choice **AC** not included), and also satisfies the relation $\mathbf{HOD} \subseteq \mathbf{HNT} \subseteq \mathbf{V}$. Tzouvaras [21, 2.15] asks whether the double strict inequality $\mathbf{HOD} \subsetneq \mathbf{HNT} \subsetneq \mathbf{V}$

*Partial support of RFBR grant 20-01-00670 acknowledged.

[†]IITP RAS, Moscow, Russia, kanovei@iitp.ru.

[‡]IITP RAS, Moscow, Russia, lyubetsk@iitp.ru

can be realized in an appropriate model of **ZFC**. The following theorem, the main result of this paper, answers this question in the affirmative.

Theorem 1.1. *Let $\mathbb{C} = \omega^{<\omega}$ be the Cohen forcing for adding a generic real $x \in \omega^\omega$ to \mathbf{L} . There is a forcing notion $\mathbb{P} \in \mathbf{L}$, which consists of Silver trees, and such that if a pair of reals $\langle a, x \rangle$ is $(\mathbb{P} \times \mathbb{C})$ -generic over \mathbf{L} then it is true in $\mathbf{L}[a, x]$ that*

$$\mathbf{L} = \mathbf{HOD} \subsetneq \mathbf{L}[a] = \mathbf{HNT} \subsetneq \mathbf{V} = \mathbf{L}[a, x].$$

Note that **HNT** satisfies **ZFC**, not merely **ZF**, in the model $\mathbf{L}[a, x]$ of the theorem.

Remark 1.2. This result is an essential strengthening of [17, Theorem 9.1]. Comparably to the latter, the claims that $\mathbf{L} = \mathbf{HOD}$ (instead of simply $a \notin \mathbf{HOD}$) and especially $\mathbf{L}[a] = \mathbf{HNT}$ (instead of just $x \notin \mathbf{HNT}$) are added here, w.r.t. basically the same model, which makes the research more accomplished.

To make the text of this preprint more self-contained, we decided to near-copy-paste some definitions and auxiliary results from [17], instead of briefly citing them as it would be more accustomed in a journal paper. \square

To prove the theorem, we make use of a forcing notion \mathbb{P} introduced in [10] in order to define a generic real $a \in 2^\omega$ whose \mathbf{E}_0 -equivalence class $[a]_{\mathbf{E}_0}$ is a lightface Π_2^1 (hence **OD**) set of reals with no **OD** element. This property of \mathbb{P} is responsible for a \mathbb{P} -generic real a to belong to **HNT**, and ultimately to $\mathbf{L}[a] \subseteq \mathbf{HNT}$, in $\mathbf{L}[a, x]$. This will be based on some results on Silver trees and Borel functions in Sections 2,3,4. The construction of \mathbb{P} in \mathbf{L} is given in Sections 5,6. The proof that $\mathbf{L}[a] \subseteq \mathbf{HNT}$ in $\mathbf{L}[a, x]$ follows in Section 8.

The inverse inclusion $\mathbf{HNT} \subseteq \mathbf{L}[a]$ in $\mathbf{L}[a, x]$ will be proved in Section 9 on the basis of our earlier result [11] on countable **OD** sets in Cohen-generic extensions.

2 Perfect trees and Silver trees

Our results will involve forcing notions that consist of perfect trees and Silver trees. Here we introduce the relevant terminology from our earlier works [10, 12, 13].

By $2^{<\omega}$ we denote the set of all *tuples* (finite sequences) of terms 0, 1, including the empty tuple Λ . The length of a tuple s is denoted by $\text{lh } s$, and $2^n = \{s \in 2^{<\omega} : \text{lh } s = n\}$ (all tuples of length n). A tree $\emptyset \neq T \subseteq 2^{<\omega}$ is *perfect*, symbolically $T \in \mathbf{PT}$, if it has no endpoints and isolated branches. In this case, the set

$$[T] = \{a \in 2^\omega : \forall n (a \upharpoonright n \in T)\}$$

of all *branches* of T is a perfect set in 2^ω . Note that $[S] \cap [T] = \emptyset$ iff $S \cap T$ is finite.

- If $u \in T \in \mathbf{PT}$, then a *portion* (or a *pruned tree*) $T \upharpoonright_u \in \mathbf{PT}$ is defined by $T \upharpoonright_u = \{s \in T : u \subset s \vee s \subseteq u\}$.
- A tree $S \subseteq T$ is *clopen* in T iff it is equal to the union of a finite number of portions of T . This is equivalent to $[S]$ being clopen in $[T]$.

A tree $T \subseteq 2^{<\omega}$ is a *Silver tree*, symbolically $T \in \mathbf{ST}$, if there is an infinite sequence of tuples $u_k = u_k(T) \in 2^{<\omega}$, such that T consists of all tuples of the form

$$s = u_0 \hat{\ } i_0 \hat{\ } u_1 \hat{\ } i_1 \hat{\ } u_2 \hat{\ } i_2 \hat{\ } \dots \hat{\ } u_n \hat{\ } i_n$$

and their sub-tuples, where $n < \omega$ and $i_k = 0, 1$. Then the *stem* $\mathbf{stem}(T) = u_0(T)$ is equal to the largest tuple $s \in T$ with $T = T \upharpoonright_s$, and $[T]$ consists of all infinite sequences $a = u_0 \hat{\ } i_0 \hat{\ } u_1 \hat{\ } i_1 \hat{\ } u_2 \hat{\ } i_2 \hat{\ } \dots \in 2^\omega$, where $i_k = 0, 1, \forall k$. Put

$$\mathbf{spl}_n(T) = \mathbf{lh} u_0 + 1 + \mathbf{lh} u_1 + 1 + \dots + \mathbf{lh} u_{n-1} + 1 + \mathbf{lh} u_n.$$

In particular, $\mathbf{spl}_0(T) = \mathbf{lh} u_0$. Thus $\mathbf{spl}(T) = \{\mathbf{spl}_n(T) : n < \omega\} \subseteq \omega$ is the set of all *splitting levels* of the Silver tree T .

Action. Let $\sigma \in 2^{<\omega}$. If $v \in 2^{<\omega}$ is another tuple of length $\mathbf{lh} v \geq \mathbf{lh} \sigma$, then the tuple $v' = \sigma \cdot v$ of the same length $\mathbf{lh} v' = \mathbf{lh} v$ is defined by $v'(i) = v(i) +_2 \sigma(i)$ (addition modulo 2) for all $i < \mathbf{lh} \sigma$, but $v'(i) = v(i)$ whenever $\mathbf{lh} \sigma \leq i < \mathbf{lh} v$. If $\mathbf{lh} v < \mathbf{lh} \sigma$, then we just define $\sigma \cdot v = (\sigma \upharpoonright \mathbf{lh} v) \cdot v$.

If $a \in 2^\omega$, then similarly $a' = \sigma \cdot a \in 2^\omega$, $a'(i) = a(i) +_2 \sigma(i)$ for $i < \mathbf{lh} \sigma$, but $a'(i) = a(i)$ for $i \geq \mathbf{lh} \sigma$. If $T \subseteq 2^{<\omega}$, $X \subseteq 2^\omega$, then the sets

$$\sigma \cdot T = \{\sigma \cdot v : v \in T\} \quad \text{and} \quad \sigma \cdot X = \{\sigma \cdot a : a \in X\}$$

are *shifts* of the tree T and the set X accordingly.

Lemma 2.1 ([13], 3.4). *If $n < \omega$ and $u, v \in T \cap 2^n$, then $T \upharpoonright_u = v \cdot u \cdot (T \upharpoonright_v)$.*

If $t \in T \in \mathbf{ST}$ and $\sigma \in 2^{<\omega}$, then $\sigma \cdot T \in \mathbf{ST}$ and $T \upharpoonright_s \in \mathbf{ST}$. □

Definition 2.2 (refinements). Assume that $T, S \in \mathbf{ST}$, $S \subseteq T$, $n < \omega$. We define $S \subseteq_n T$ (the tree S *n-refines* T) if $S \subseteq T$ and $\mathbf{spl}_k(T) = \mathbf{spl}_k(S)$ for all $k < n$. This is equivalent to $(S \subseteq T \text{ and } u_k(S) = u_k(T) \text{ for all } k < n)$, of course. □

Then $S \subseteq_0 T$ is equivalent to $S \subseteq T$, and $S \subseteq_{n+1} T$ implies $S \subseteq_n T$ (and $S \subseteq T$), but if $n \geq 1$ then $S \subseteq_n T$ is equivalent to $\mathbf{spl}_{n-1}(T) = \mathbf{spl}_{n-1}(S)$.

Lemma 2.3. *Assume that $T, U \in \mathbf{ST}$, $n < \omega$, $h > \mathbf{spl}_{n-1}(T)$, $s_0 \in 2^h \cap T$, and $U \subseteq T \upharpoonright_{s_0}$. Then there is a unique tree $S \in \mathbf{ST}$ such that $S \subseteq_n T$ and $S \upharpoonright_{s_0} = U$.*

If in addition U is clopen in T then S is clopen in T as well.

Proof (sketch). Define a tree S so that $S \cap 2^h = T \cap 2^h$, and if $t \in T \cap 2^h$ then, by Lemma 2.1, $S \upharpoonright_t = (t \cdot s_0) \cdot U$; then $S \upharpoonright_{s_0} = U$. To check that $S \in \mathbf{ST}$, we can easily compute the tuples $u_k(S)$. Namely, as $U \subseteq T \upharpoonright_{s_0}$, we have $s_0 \subseteq u_0(U) = \mathbf{stem}(U)$, hence $\ell = \mathbf{lh}(u_0(U)) \geq h > m = \mathbf{spl}_{n-1}(T)$. Then $u_k(S) = u_k(T)$ for all $k < n$, $u_n(S) = u_0(U) \upharpoonright [m, \ell]$ (thus $u_n(S) \in 2^{\ell-m}$), and $u_k(S) = u_k(U)$ for all $k > n$. □

Lemma 2.4 ([13], Lemma 4.4). *Let $\dots \subseteq_4 T_3 \subseteq_3 T_2 \subseteq_2 T_1 \subseteq_1 T_0$ be a sequence of trees in \mathbf{ST} . Then $T = \bigcap_n t_n \in \mathbf{ST}$.* □

Proof (sketch). By definition we have $u_k(T_n) = u_k(T_{n+1})$ for all $k \leq n$. Then one easily computes that $u_n(T) = u_n(T_n)$ for all n . □

3 Reduction of Borel maps to continuous ones

A classical theorem claims that in Polish spaces every Borel function is continuous on a suitable dense \mathbf{G}_δ set (Theorem 8.38 in Kechris [18]). It is also known that a Borel map defined on 2^ω is continuous on a suitable Silver tree. The next lemma combines these two results. Our interest in functions defined on $2^\omega \times \omega^\omega$ is motivated by further applications to reals in generic extensions of the form $\mathbf{L}[a, x]$, where $a \in 2^\omega$ is \mathbb{P} -generic real for some $\mathbb{P} \subseteq \mathbf{ST}$ while $x \in \omega^\omega$ is just Cohen generic.

In the remainder, if $v \in \omega^{<\omega}$ (a tuple of natural numbers), then we define $\mathcal{N}_v = \{x \in \omega^\omega : v \subset x\}$, a *Baire interval* or *portion* in the Baire space ω^ω .

Lemma 3.1. *Let $T \in \mathbf{ST}$ and $f : 2^\omega \times \omega^\omega \rightarrow 2^\omega$ be a Borel map. There is a Silver tree $S \subseteq T$ and a dense \mathbf{G}_δ set $D \subseteq \omega^\omega$ such that f is continuous on $[S] \times D$.*

Proof. By the abovementioned classical theorem, f is already continuous on some dense \mathbf{G}_δ set $Z \subseteq [T] \times \omega^\omega$. It remains to define a Silver tree $S \subseteq T$ and a dense \mathbf{G}_δ set $D \subseteq \omega^\omega$ such that $[S] \times D \subseteq Z$. This will be our goal.

We have $Z = \bigcap_n Z_n$, where each $Z_n \subseteq [T] \times \omega^\omega$ is open dense.

We fix a recursive enumeration $\omega \times \omega^{<\omega} = \{\langle N_k, v_k \rangle : k < \omega\}$. We will define a sequence of Silver trees S_k and tuples $w_k \in \omega^{<\omega}$ satisfying the following:

- (1) $\dots \subseteq_4 S_3 \subseteq_3 S_2 \subseteq_2 s_1 \subseteq_1 S_0 = T$, as in Lemma 2.4;
- (2) if $k < \omega$ then S_{k+1} is **clopen** in S_k (see Section 2);
- (3) $v_k \subseteq w_k$ and $[S_{k+1}] \times \mathcal{N}_{w_k} \subseteq Z_{N_k}$, for all k .

At step 0 we already have $S_0 = T$.

Assume that $S_k \in \mathbf{ST}$ has already been defined. Let $h = \mathbf{spl}_{k+1}(S_k)$.

Consider any tuple $t \in 2^h \cap S_k$. As Z_{N_k} is open dense, there is a tuple $u_1 \in \omega^{<\omega}$ and a Silver tree $A_1 \subseteq S_k \upharpoonright_t$, clopen in S_k (for example, a portion in S_k) such that $v_k \subseteq u_1$ and $[A_1] \times \mathcal{N}_{u_1} \subseteq Z_{N_k}$. According to Lemma 2.3, there exists a Silver tree $U_1 \subseteq_{k+1} S_k$, clopen in S_k along with A , such that $U_1 \upharpoonright_t = A_1$, so $[U_1 \upharpoonright_t] \times \mathcal{N}_{u_1} \subseteq Z_{N_k}$ by construction.

Now take another tuple $t' \in 2^h \cap S_k$, and similarly find $u_2 \in \omega^{<\omega}$ and a Silver tree $A_2 \subseteq U_1 \upharpoonright_{t'}$, clopen in U_1 , such that $u_1 \subseteq u_2$ and $[A_2] \times \mathcal{N}_{u_2} \subseteq Z_{N_k}$. Once again there is a Silver tree $U_2 \subseteq_{k+1} U_1$, clopen in S_k and such that $[U_2 \upharpoonright_{t'}] \times \mathcal{N}_{u_2} \subseteq Z_{N_k}$.

We iterate this construction over all tuples $t \in 2^h \cap S_k$, \subseteq_{k+1} -shrinking trees and extending tuples in $\omega^{<\omega}$. We get a Silver tree $U \subseteq_{k+1} S_k$, clopen in S_k , and a tuple $w \in \omega^{<\omega}$, that $v_k \subseteq w$ and $[U] \times \mathcal{N}_w \subseteq Z_{N_k}$. Take $w_k = w$, $S_{k+1} = U$. This completes the inductive step.

As a result we get a sequence $\dots \subseteq_4 S_3 \subseteq_3 S_2 \subseteq_2 S_1 \subseteq_1 S_0 = T$ of Silver trees S_k , and tuples $w_k \in \omega^{<\omega}$ ($k < \omega$), which satisfy (1),(2),(3).

We put $S = \bigcap_k S_k$; then $S \in \mathbf{ST}$ by (1) and Lemma 2.4, and $S \subseteq T$.

If $n < \omega$ then let $W_n = \{w_k : N_k = n\}$. We claim that $D_n = \bigcup_{w \in W_n} \mathcal{N}_w$ is an open dense set in ω^ω . Indeed, let $v \in \omega^{<\omega}$. Consider any k such that that $v_k = v$

and $N_k = n$. By construction, we have $v \subseteq w_k \in W_n$, as required. We conclude that the set $D = \bigcap_n D_n$ is dense and \mathbf{G}_δ .

To check $[S] \times D \subseteq Z$, let $n < \omega$; we show that $[S] \times D \subseteq Z_n$. Let $a \in [S]$ and $x \in D$, in particular $x \in D_n$, so $x \in \mathcal{N}_{w_k}$ for some k with $N_k = n$. However, $[S_{k+1}] \times \mathcal{N}_{w_k} \subseteq Z_n$ by (3), and at the same time obviously $a \in [S_{k+1}]$. We conclude that in fact $\langle a, x \rangle \in Z_n$, as required. \square (Lemma 3.1)

Corollary 3.2. *Let $T \in \mathbf{ST}$ and $f : 2^\omega \rightarrow 2^\omega$ be a Borel map. There is a Silver tree $S \subseteq T$ such that f is continuous on $[S]$.* \square

We add the following result that belongs to the folklore of the Silver forcing. See Corollary 5.4 in [12] for a proof.

Lemma 3.3. *Assume that $T \in \mathbf{ST}$ and $f : 2^\omega \rightarrow 2^\omega$ is a continuous map. Then there is a Silver tree $S \subseteq T$ such that f is either a bijection or a constant on $[S]$.* \square

4 Normalization of Borel maps

Definition 4.1. A map $f : 2^\omega \times \omega^\omega \rightarrow 2^\omega$ is *normalized on $T \in \mathbf{ST}$* for $\mathcal{U} \subseteq \mathbf{ST}$ if there exists a dense \mathbf{G}_δ set $X \subseteq \omega^\omega$ such that f is continuous on $[T] \times X$ and:

- either (a) there are tuples $v \in \omega^{<\omega}$, $\sigma \in 2^{<\omega}$ such that $f(a, x) = \sigma \cdot a$ for all $a \in [T]$ and $x \in \mathcal{N}_v \cap X$, where, we remind, $\mathcal{N}_v = \{x \in \omega^\omega : v \subset x\}$;
- or (b) $f(a, x) \notin \bigcup_{\sigma \in 2^{<\omega} \wedge S \in \mathcal{U}} \sigma \cdot [S]$ for all $a \in [T]$ and $x \in X$. \square

Theorem 4.2. *Let $\mathcal{U} = \{T_0, T_1, T_2, \dots\} \subseteq \mathbf{ST}$ and $f : 2^\omega \times \omega^\omega \rightarrow 2^\omega$ be a Borel map. There is a set $\mathcal{U}' = \{S_0, S_1, S_2, \dots\} \subseteq \mathbf{ST}$, such that $S_n \subseteq T_n$ for all n and f is normalized on S_0 for \mathcal{U}' .*

Proof. First of all, according to Lemma 3.1, there is a Silver tree $T' \subseteq T_0$ and a dense \mathbf{G}_δ set $W \subseteq \omega^\omega$ such that f is continuous on $[T'] \times W$. And since any dense \mathbf{G}_δ set $X \subseteq \omega^\omega$ is homeomorphic to ω^ω , we can w.l.o.g. assume that $W = \omega^\omega$ and $T' = T_0$. Thus, we simply suppose that f is already *continuous on $[T_0] \times \omega^\omega$* .

Assume that option (a) of the definition of 4.1 does not take place, i.e.

- (*) if $X \subseteq \omega^\omega$ is dense \mathbf{G}_δ , and $v \in \omega^{<\omega}$, $\sigma \in 2^{<\omega}$, $S \in \mathbf{ST}$, $S \subseteq T_0$, then there are reals $a \in [S]$ and $x \in \mathcal{N}_v \cap X$ such that $f(a, x) \neq \sigma \cdot a$.

We'll construct Silver trees $S_n \subseteq T_n$ and a dense \mathbf{G}_δ set $X \subseteq \omega^\omega$ satisfying (b) of Definition 4.1, that is, in our case, the relation $f(a, x) \notin \bigcup_{\sigma \in 2^{<\omega} \wedge n < \omega} \sigma \cdot [S_n]$ will be fulfilled for all $a \in [S_0]$ and $x \in X$.

To maintain the construction, we fix any enumeration $\omega \times 2^{<\omega} \times \omega^{<\omega} = \{\langle N_k, \sigma_k, v_k \rangle : k < \omega\}$. Auxiliary Silver trees S_k^n ($n, k < \omega$) and tuples $w_k \in \omega^{<\omega}$ ($k < \omega$), satisfying the following conditions, will be defined.

- (1) $\dots \subseteq_4 S_3^n \subseteq_3 S_2^n \subseteq_2 S_1^n \subseteq_1 S_0^n = T_n$ as in Lemma 2.4, for each $n < \omega$;

- (2) $S_{k+1}^n = S_k^n$ for all $n > 0$, $n \neq N_k$;
- (3) $S_{k+1}^0 \subseteq_{k+1} S_k^0$, $S_{k+1}^{N_k} \subseteq_{k+1} S_k^{N_k}$, $v_k \subseteq w_k$, and $f(a, x) \notin \sigma_k \cdot [S_{k+1}^N]$ for all reals $a \in [S_{k+1}^0]$ and $x \in \mathcal{N}_{w_k}$.

At step 0 of the construction, we put $S_0^n = T_n$ for all n , by (1).

Assume that $k < \omega$ and all Silver trees S_k^n , $n < \omega$ are already defined. We put $S_{k+1}^n = S_k^n$ for all $n > 0$, $n \neq N_k$, by (2).

To define the trees S_{k+1}^0 and $S_{k+1}^{N_k}$, we put $h = \text{spl}_{k+1}(S_k^0)$, $m = \text{spl}_{k+1}(S_k^{N_k})$.

Case 1: $N_k > 0$. Take any pair of tuples $s \in 2^h \cap S_k^0$, $t \in 2^m \cap S_k^{N_k}$ and any reals $a_0 \in [S_k^0 \upharpoonright_s]$ and $x_0 \in \omega^\omega$. Consider any real $b_0 \in [S_k^{N_k} \upharpoonright_t]$ not equal to $\sigma_k \cdot f(a_0, x_0)$. Let's say $b_0(\ell) = i \neq j = (\sigma_k \cdot f(a_0, x_0))(\ell)$, where $i, j \leq 1$, $\ell < \omega$. By the continuity of f , there is a tuple $u_1 \in \omega^{<\omega}$ and Silver tree $A \subseteq S_k^0 \upharpoonright_s$ such that $v_k \subseteq u_1 \subset x_0$, $a_0 \in [A]$, and $(\sigma_k \cdot f(a, x))(\ell) = j$ for all $x \in \mathcal{N}_{u_1}$ and $a \in [A]$. It is also clear that $B = \{\tau \in S_k^{N_k} \upharpoonright_t : \text{lh } \tau \leq \ell \vee \tau(\ell) = i\}$ is a Silver tree containing b_0 , and $b(\ell) = i$ for all $b \in [B]$. According to Lemma 2.3, there are Silver trees $U_1 \subseteq_{k+1} S_k^0$ and $V_1 \subseteq_{k+1} S_k^{N_k}$, such that $U_1 \upharpoonright_s = A$ and $V_1 \upharpoonright_t = B$, hence by construction we have $\sigma_k \cdot f(a, x) \notin [V_1 \upharpoonright_t]$ for all $a \in [U_1 \upharpoonright_s]$ and $x \in \mathcal{N}_{u_1}$.

Now consider another pair of tuples $s \in 2^h \cap S_k^0$, $t \in 2^m \cap S_k^{N_k}$. We similarly get Silver trees $U_2 \subseteq_{k+1} U_1$ and $V_2 \subseteq_{k+1} V_1$, and a tuple $u_2 \in \omega^{<\omega}$, such that $u_1 \subseteq u_2$ and $\sigma_k \cdot f(a, x) \notin [V_2 \upharpoonright_{s'}]$ for all $a \in [U_2 \upharpoonright_{s'}]$ and $x \in \mathcal{N}_{u_2}$. In this case, we have $V_2 \upharpoonright_t \subseteq V_1 \upharpoonright_t$ and $U_2 \upharpoonright_s \subseteq U_1 \upharpoonright_s$, so that what has already been achieved at the previous step is preserved.

We iterate through all pairs of $s \in 2^h \cap S_k^0$, $t \in 2^m \cap S_k^{N_k}$, \subseteq_{k+1} -shrinking trees and extending tuples in $\omega^{<\omega}$ at each step. This results in a pair of Silver trees $U \subseteq_{k+1} S_k^0$, $V \subseteq_{k+1} S_k^{N_k}$ and a tuple $w \in \omega^{<\omega}$ such that $v_k \subseteq w$ and $\sigma_k \cdot f(a, x) \notin [V]$ for all reals $a \in [U]$ and $x \in \mathcal{N}_w$. Now to fulfill (3), take $w_k = w$, $S_{k+1}^0 = U$, and $S_{k+1}^{N_k} = V$. Recall that here $N_k > 0$.

Case 2: $N_k = 0$. Here the construction somewhat changes, and hypothesis (*) will be used. We claim that there exist:

- (4) a tuple $w_k \in \omega^{<\omega}$ and a Silver tree $S_{k+1}^0 \subseteq_{k+1} S_k^0$ such that $v_k \subseteq w_k$ and $f(a, x) \notin \sigma_k \cdot [S_{k+1}^0]$ for all $a \in [S_{k+1}^0]$, $x \in \mathcal{N}_{w_k}$. (**Equivalent to (3) as $N_k = 0$.**)

Take any pair of tuples $s, t \in 2^h \cap S_k^0$, where $h = \text{spl}_{k+1}(S_k^0)$ as above. Thus $S_k^0 \upharpoonright_t = t \cdot s \cdot (S_k^0 \upharpoonright_s)$, by Lemma 2.1. According to (*), there are reals $x_0 \in \mathcal{N}_v$ and $a_0 \in [S_k^0 \upharpoonright_s]$ satisfying $f(a_0, x_0) \neq \sigma_k \cdot s \cdot t \cdot a_0$, or equivalently, $\sigma_k \cdot f(a_0, x_0) \neq s \cdot t \cdot a_0$.

Similarly to Case 1, we have $(\sigma_k \cdot f(a_0, x_0))(\ell) = i \neq j = (s \cdot t \cdot a_0)(\ell)$ for some $\ell < \omega$ and $i, j \leq 1$. By the continuity of f , there is a tuple $u_1 \in \omega^{<\omega}$ and a Silver tree $A \subseteq S_k^0 \upharpoonright_s$, clopen in S_k^0 , such that $v_k \subseteq u_1 \subset x_0$, $a_0 \in [A]$, and $(\sigma_k \cdot f(a, x))(\ell) = j$ but $(s \cdot t \cdot a)(\ell) = j$ for all $x \in \mathcal{N}_{u_1}$ and $a \in [A]$. Lemma 2.3 gives us a Silver tree $U_1 \subseteq_{k+1} S_k^0$, clopen in S_k^0 as well, such that $U_1 \upharpoonright_s = A$ — and then $U_1 \upharpoonright_t = s \cdot t \cdot A$. Therefore $\sigma_k \cdot f(a, x) \notin [U_1 \upharpoonright_t]$ holds for all $a \in [U_1 \upharpoonright_s]$ and $x \in \mathcal{N}_{u_1}$ by construction.

Having worked out all pairs of tuples $s, t \in 2^h \cap S_k^0$, we obtain a Silver tree $U \subseteq_{k+1} S_k^0$ and a tuple $w \in \omega^{<\omega}$, such that $v_k \subseteq w$ and $\sigma_k \cdot f(a, x) \notin [U]$ for all $a \in [U]$ and $x \in \mathcal{N}_w$. Now to fulfill (4), take $w_k = w$ and $S_{k+1}^0 = U$.

To conclude, we have for each n a sequence $\dots \subseteq_4 S_3^n \subseteq_3 S_2^n \subseteq_2 S_1^n \subseteq_1 S_0^n = T_n$ of Silver trees S_k^n , along with tuples $w_k \in \omega^{<\omega}$ ($k < \omega$), and these sequences satisfy the requirements (1),(2),(3) (equivalent to (4) in case $N_k = 0$).

We put $S_n = \bigcap_k S_k^n$. Then $S_n \in \mathbf{ST}$ by Lemma 2.4, and $S_n \subseteq T_n$.

If $n < \omega$ and $\sigma \in 2^{<\omega}$ then let $W_{n\sigma} = \{w_k : N_k = n \wedge \sigma_k = \sigma\}$. The set $X_{n\sigma} = \bigcup_{w \in W_{n\sigma}} \mathcal{N}_w$ is then open dense in ω^ω . Indeed, if $v \in \omega^\omega$ then we take k such that $v_k = v$, $N_k = n$, $\sigma_k = \sigma$; then $v \subseteq w_k \in W_{n\sigma}$ by construction. Therefore, $X = \bigcap_{n < \omega, \sigma \in 2^{<\omega}} X_{n\sigma}$ is a dense \mathbf{G}_δ set. Now to check property (b) of Definition 4.1, consider any $n < \omega$, $\sigma \in 2^{<\omega}$, $a \in [S_0]$, $x \in X$; we claim that $f(a, x) \notin \sigma \cdot [S_n]$.

By construction, we have $x \in X_{n\sigma}$, i.e. $x \in \mathcal{N}_{w_k}$, where $k \in W_{n\sigma}$, so that $N_k = n$, $\sigma_k = \sigma$. Now $f(a, x) \notin \sigma \cdot [S_n]$ directly follows from (3) for this k , since $S_0 \subseteq S_{k+1}^0$ and $S_n \subseteq S_{k+1}^n$. \square (Theorem 4.2)

5 The forcing notion for Theorem 1.1

Using the standard encoding of Borel sets, as e.g. in [20] or [9, §1D], we fix a coding of Borel functions $f : 2^\omega \rightarrow 2^\omega$. As usual, it includes a Π_1^1 -set¹ of codes $\mathbf{BC} \subseteq \omega^\omega$, and for each code $r \in \mathbf{BC}$ a certain Borel function $F_r : 2^\omega \rightarrow 2^\omega$ coded by r . We assume that each Borel function has some code, and there is a Σ_1^1 relation $\mathfrak{S}(\cdot, \cdot, \cdot)$ and a Π_1^1 relation $\mathfrak{P}(\cdot, \cdot, \cdot)$ such that for all $r \in \mathbf{BC}$ and $a, b \in 2^\omega$ it holds $F_r(a) = b \iff \mathfrak{S}(r, a, b) \iff \mathfrak{P}(r, a, b)$.

Similarly, we fix a coding of Borel functions $f : 2^\omega \times \omega^\omega \rightarrow 2^\omega$, that includes a Π_1^1 -set of codes $\mathbf{BC}_2 \subseteq \omega^\omega$, and for each code $r \in \mathbf{BC}_2$ a Borel function $F_r^2 : 2^\omega \times \omega^\omega \rightarrow 2^\omega$ coded by r , such that each Borel function has some code, and there is a Σ_1^1 relation $\mathfrak{S}^2(\cdot, \cdot, \cdot, \cdot)$ and a Π_1^1 relation $\mathfrak{P}^2(\cdot, \cdot, \cdot, \cdot)$ such that for all $r \in \mathbf{BC}$, $x \in \omega^\omega$, and $a, b \in 2^\omega$ it holds $F_r^2(a, x) = b \iff \mathfrak{S}^2(r, a, x, b) \iff \mathfrak{P}^2(r, a, x, b)$.

If $\mathbb{U} \subseteq \mathbf{ST}$, then $\mathbf{Clos}(\mathbb{U})$ denotes the set of all trees of the form $\sigma \cdot (T \upharpoonright_s)$, where $\sigma \in 2^{<\omega}$ and $s \in T \in \mathbb{U}$, i.e. the closure of \mathbb{U} w.r.t. both shifts and portions.

The following construction is maintained in L. We define a sequence of countable sets $\mathbb{U}_\alpha \subseteq \mathbf{ST}$, $\alpha < \omega_1$ satisfying the following conditions 1°–6°.

1°. Each $\mathbb{U}_\alpha \subseteq \mathbf{ST}$ is countable, \mathbb{U}_0 consists of a single tree $2^{<\omega}$.

We then define $\mathbb{P}_\alpha = \mathbf{Clos}(\mathbb{U}_\alpha)$, $\mathbb{P}_{<\alpha} = \bigcup_{\xi < \alpha} \mathbb{P}_\xi$. These sets are obviously closed with respect to shifts and portions, that is $\mathbf{Clos}(\mathbb{P}_\alpha) = \mathbb{P}_\alpha$ and $\mathbf{Clos}(\mathbb{P}_{<\alpha}) = \mathbb{P}_{<\alpha}$.

2°. For every $T \in \mathbb{P}_{<\alpha}$ there is a tree $S \in \mathbb{U}_\alpha$, $S \subseteq T$.

Let \mathbf{ZFC}^- be the subtheory of the theory \mathbf{ZFC} , containing all axioms except the power set axiom, and additionally containing an axiom asserting the existence of

¹The letters Σ and Π denote effective (lightface) projective classes.

the power set $\mathcal{P}(\omega)$. This implies the existence of $\mathcal{P}(X)$ for any countable X , the existence of ω_1 and 2^ω , as well as the existence of continual sets like 2^ω or **ST**.

By \mathfrak{M}_α we denote the smallest model of **ZFC**⁻ of the form \mathbf{L}_λ containing the sequence $\langle \mathbb{U}_\xi \rangle_{\xi < \alpha}$, in which α and all sets \mathbb{U}_ξ , $\xi < \alpha$, are countable.

3°. If a set $D \in \mathfrak{M}_\alpha$, $D \subseteq \mathbb{P}_{<\alpha}$ is dense in $\mathbb{P}_{<\alpha}$, and $U \in \mathbb{U}_\alpha$, then $U \subseteq^{\text{fin}} \bigcup D$, meaning that there is a finite set $D' \subseteq D$ such that $U \subseteq \bigcup D'$.

4°. If a set $D \in \mathfrak{M}_\alpha$, $D \subseteq \mathbb{P}_{<\alpha} \times \mathbb{P}_{<\alpha}$ is dense in $\mathbb{P}_{<\alpha} \times \mathbb{P}_{<\alpha}$, and $U \neq V$ belong to \mathbb{U}_α , then $U \times V \subseteq^{\text{fin}} \bigcup D$, meaning that there is a finite set $D' \subseteq D$ such that $[U] \times [V] \subseteq \bigcup_{\langle U', V' \rangle \in D'} [U'] \times [V']$.

Given that $\text{Clos}(\mathbb{P}_{<\alpha}) = \mathbb{P}_{<\alpha}$, this is automatically transferred to all trees $U \in \mathbb{P}_\alpha$ as well. It follows that D remains predense in $\mathbb{P}_{<\alpha} \cup \mathbb{P}_\alpha$.

To formulate the next property, we fix an enumeration

$$\mathbf{ST} \times \mathbf{BC} \times \mathbf{BC}_2 = \{ \langle T_\xi, b_\xi, c_\xi \rangle : \xi < \omega_1 \}$$

in \mathbf{L} , which 1) is definable in \mathbf{L}_{ω_1} , and 2) each value in $\mathbf{ST} \times \mathbf{BC} \times \mathbf{BC}_2$ is taken uncountably many times.

5°. If $T_\alpha \in \mathbb{P}_{<\alpha}$ then there is a tree $S \in \mathbb{U}_\alpha$ such that $S \subseteq T$ and:

- (a) $F_{b_\alpha}^2$ is normalized for \mathbb{U}_α on $[S]$ in the sense of Definition 4.1, and
- (b) F_{c_α} is continuous and either a bijection or a constant on $[S]$.

6°. The sequence $\langle \mathbb{U}_\alpha \rangle_{\alpha < \omega_1}$ is \in -definable in \mathbf{L}_{ω_1} .

The construction goes on as follows. **Arguing in \mathbf{L}** , suppose that

- (†) $\alpha < \omega_1$, the subsequence $\langle \mathbb{U}_\xi \rangle_{\xi < \alpha}$ has been defined and satisfies 1°, 2° below α , and the sets $\mathbb{P}_\xi = \text{Clos}(\mathbb{U}_\xi)$ (for $\xi < \alpha$), $\mathbb{P}_{<\alpha}$, \mathfrak{M}_α are defined as above.

See the proof of the next lemma in Section 6 below.

Lemma 5.1 (\mathbb{U} -extension lemma, in \mathbf{L}). *Under the assumptions of (†), there is a countable set $\mathbb{U}_\alpha \subseteq \mathbf{ST}$ satisfying 2°, 3°, 4°, 5°.*

To accomplish the construction, we take \mathbb{U}_α to be the smallest, in the sense of the Gödel wellordering of \mathbf{L} , of those sets that exist by Lemma 5.1. Since the whole construction is relativized to \mathbf{L}_{ω_1} , the requirement 6° is also met.

We put $\mathbb{P}_\alpha = \text{Clos}(\mathbb{U}_\alpha)$ for all $\alpha < \omega_1$, and $\mathbb{P} = \bigcup_{\alpha < \omega_1} \mathbb{P}_\alpha$.

The following result, in part related to CCC, is a fairly standard consequence of 3° and 4°, see for example [10, 6.5], [12, 12.4], or [8, Lemma 6]; we will skip the proof.

Lemma 5.2 (in \mathbf{L}). *The forcing notion \mathbb{P} belongs to \mathbf{L} , satisfies $\mathbb{P} = \text{Clos}(\mathbb{P})$ and satisfies CCC in \mathbf{L} . The product $\mathbb{P} \times \mathbb{P}$ satisfies CCC in \mathbf{L} as well. \square*

Lemma 5.3 (in **L**). *Assume that $T \in \mathbb{P}$. If $g : 2^\omega \rightarrow 2^\omega$ is a Borel map then there is a tree $S \in \mathbb{U}_\alpha$, $S \subseteq T$, such that g is either a bijection or a constant on $[S]$.*

If $f : 2^\omega \times \omega^\omega \rightarrow 2^\omega$ is a Borel map then there is an ordinal $\alpha < \omega_1$ and a tree $S \in \mathbb{U}_\alpha$, $S \subseteq T$, such that f is normalized for \mathbb{U}_α on $[S]$.

Proof. By the choice of the enumeration of triples in $\mathbf{ST} \times \mathbf{BC} \times \mathbf{BC}_2$, there is an ordinal $\alpha < \omega_1$ such that $T \in \mathbb{P}_{<\alpha}$ and $T = T_\alpha$, $f = F_{b_\alpha}^2$, $g = F_{b_\alpha}$. It remains to refer to 5°. \square

6 Proof of the extension lemma

This section is entirely devoted to the **proof of Lemma 5.1**.

We work in **L under the assumptions of (†) above.**

We first define a set $\mathbb{U} = \{U_n : n < \omega\}$ of Silver trees $U_n \subseteq 2^\omega$ satisfying 2°, 3° 4°; then further narrowing of the trees will be made to also satisfy 5°. This involves a splitting/fusion construction known from our earlier papers, see [10, §4], [13, §9–10], [12, §10], [16, §7], and to some extent from the proof of Theorem 4.2 above.

We fix enumerations

$$\mathcal{D} = \{D(j) : j < \omega\} \quad \text{and} \quad \mathcal{D}_2 = \{D_2(j) : j < \omega\}$$

of the set \mathcal{D} of all sets $D \in \mathfrak{M}_\alpha$, $D \subseteq \mathbb{P}_{<\alpha}$ open-dense in $\mathbb{P}_{<\alpha}$, and the set \mathcal{D}_2 of all sets $D \in \mathfrak{M}_\alpha$, $D \subseteq \mathbb{P}_{<\alpha} \times \mathbb{P}_{<\alpha}$ open-dense in $\mathbb{P}_{<\alpha} \times \mathbb{P}_{<\alpha}$. We also fix a bijection $\beta : \omega \xrightarrow{\text{onto}} \omega^4$ which assumes each value $\langle j, j', M, N \rangle \in \omega^4$ infinitely many times.

The construction of the trees U_n is organized in the form $U_n = \bigcup_k U_k^n$, where the Silver trees U_k^n satisfy the following requirements:

- (1) $\dots \subseteq_4 U_3^n \subseteq_3 U_2^n \subseteq_2 U_1^n \subseteq_1 U_0^n$ as in Lemma 2.4 for each $n < \omega$;
- (2) if $T \in \mathbb{P}_{<\alpha}$ then $T = U_0^n$ for some n ;
- (3) each U_k^n is a k -collage over $\mathbb{P}_{<\alpha}$.

A Silver tree T is a k -collage over $\mathbb{P}_{<\alpha}$ [13, 12] when $T \upharpoonright_s \in \mathbb{P}_{<\alpha}$ for each tuple $s \in T \cap 2^h$, where $h = \mathbf{spl}_k(T)$. Then 0-collages are just trees in $\mathbb{P}_{<\alpha}$, and every k -collage is a $k+1$ -collage as well since $\mathbf{Clos}(\mathbb{P}_{<\alpha}) = \mathbb{P}_{<\alpha}$.

- (4) if $k \geq 1$, $\beta(k) = \langle j, j', M, N \rangle$, $\mu = \mathbf{spl}_k(U_k^M)$, $\nu = \mathbf{spl}_k(U_k^N)$ (integers), $s \in U_k^M \cap 2^\mu$, $t \in U_k^N \cap 2^\nu$ (tuples of length resp. μ, ν), $M \neq N$, then the tree $U_k^M \upharpoonright_s$ belongs to $D(j)$ and the pair $\langle U_k^M \upharpoonright_s, U_k^N \upharpoonright_t \rangle$ belongs to $D_2(j')$.

It follows that $U_k^M \subseteq^{\text{fin}} \bigcup D(j)$ and $\langle U_k^M, U_k^N \rangle \subseteq^{\text{fin}} \bigcup D_2(j')$ in the sense of 3° and 4° of Section 5.

To begin the inductive construction, we assign $U_0^n \in \mathbb{P}_{<\alpha}$ so that $\{U_0^n : n < \omega\} = \mathbb{P}_{<\alpha}$, to get (2). Now let's maintain the step $k \rightarrow k+1$. Thus suppose that $k < \omega$, and all Silver trees U_k^n , $n < \omega$ are defined and are k -collages over $\mathbb{P}_{<\alpha}$.

Let $\beta(k) = \langle j, j', M, N \rangle$. If $N = M$ then put $U_{k+1}^n = U_k^n$ for all n .

Now assume that $M \neq N$. Put $U_{k+1}^n = U_k^n$ for all $n \notin \{M, N\}$.

It takes more effort to define U_{k+1}^M and U_{k+1}^N . Let $\mu = \mathbf{spl}_{k+1}(U_k^M)$, $\nu = \mathbf{spl}_{k+1}(U_k^N)$. To begin with we put $U_{k+1}^M := U_k^M$ and $U_{k+1}^N := U_k^N$. These $k+1$ -collages are the initial values for the trees U_{k+1}^M and U_{k+1}^N , to be \subseteq_{k+1} -shrunk in a finite number of substeps (within the step $k \rightarrow k+1$), each corresponding to a pair of tuples $s \in U_k^M \cap 2^\mu$ and $t \in U_k^N \cap 2^\nu$.

Namely let $s \in U_{k+1}^M \cap 2^\mu$, $t \in U_{k+1}^N \cap 2^\nu$ be the first such pair. The trees $U_{k+1}^M \upharpoonright_s$, $U_{k+1}^N \upharpoonright_t$ belong to $\mathbb{P}_{<\alpha}$ as U_{k+1}^M , U_{k+1}^N are $k+1$ -collages over $\mathbb{P}_{<\alpha}$. Therefore by the open density there exist trees $A, B \in D(j)$ such that the pair $\langle U_{k+1}^M \upharpoonright_s, U_{k+1}^N \upharpoonright_t \rangle$ belongs to $D_2(j')$ and $A \subseteq U_{k+1}^M \upharpoonright_s$, $B \subseteq U_{k+1}^N \upharpoonright_t$. Now Lemma 2.3 gives us Silver trees $S \subseteq_{k+1} U_k^M$ and $T \subseteq_{k+1} U_k^N$ satisfying $S \upharpoonright_s \subseteq A$, $T \upharpoonright_t \subseteq B$. Moreover, by Lemma 2.1, S and T still are $k+1$ -collages over $\mathbb{P}_{<\alpha}$ since $\mathbb{P}_{<\alpha}$ is closed under shifts by construction. To conclude, we have defined $k+1$ -collages $S \subseteq_{k+1} U_{k+1}^M$ and $T \subseteq_{k+1} U_{k+1}^N$ over $\mathbb{P}_{<\alpha}$, satisfying $S \upharpoonright_s \in D(j)$, $T \upharpoonright_t \in D(j)$, and $\langle S \upharpoonright_s, T \upharpoonright_t \rangle \in D_2(j')$. We re-assign the “new” U_{k+1}^M and U_{k+1}^N to be equal to resp. S, T .

Applying this \subseteq_{k+1} -shrinking procedure consecutively for all pairs of tuples $s \in U_k^M \cap 2^\mu$ and $t \in U_k^N \cap 2^\nu$, we eventually (after finitely many substeps according to the number of all such pairs), we get a pair of $k+1$ -collages $U_{k+1}^M \subseteq_{k+1} U_k^M$ and $U_{k+1}^N \subseteq_{k+1} U_k^N$ over $\mathbb{P}_{<\alpha}$, such that for every pair of tuples $s \in U_k^M \cap 2^\mu$ and $t \in U_k^N \cap 2^\nu$, we have $U_{k+1}^M \upharpoonright_s \in D(j)$ and $\langle U_{k+1}^M \upharpoonright_s, U_{k+1}^N \upharpoonright_t \rangle \in D_2(j')$, so conditions (3) and (4) are satisfied.

Having defined, in \mathbf{L} , a system of Silver trees U_k^n satisfying (1),(2),(3),(4), we then put $U_n = \bigcap_k U_k^N$ for all n . Those are Silver trees by Lemma 2.4. The collection $\mathcal{U}_\alpha := \{U_n : n < \omega\}$ satisfies 2° of Section 5 by (2).

To check condition 3° of Section 5, let $D \in \mathfrak{M}_\alpha$, $D \subseteq \mathbb{P}_{<\alpha}$ be dense in $\mathbb{P}_{<\alpha}$, and $U \in \mathcal{U}_\alpha$. We can w.l.o.g. assume that D is open-dense, for if not then replace T by $D' = \{S \in \mathbb{P}_{<\alpha} : \exists T \in D (S \subseteq T)\}$. Then $D = D(j)$ for some j , and $U = U_M$ for some M by construction. Now consider any index k such that $\beta(k) = \langle M, N, j, j' \rangle$ for M, j as above and any N, j' . Then we have $U = U_M \subseteq U_k^M$ by construction, and $U_k^M \subseteq^{\mathbf{fin}} \bigcup D$ by (4), thus $U \subseteq^{\mathbf{fin}} \bigcup D$, as required.

Condition 4° is verified similarly.

It remains to somewhat shrink all trees U_n to also fulfill 5° . We still work in \mathbf{L} .

Recall that an enumeration $\mathbf{ST} \times \mathbf{BC} \times \mathbf{BC}_2 = \{\langle T_\xi, b_\xi, c_\xi \rangle : \xi < \omega_1\}$, parameter-free definable in \mathbf{L}_{ω_1} , is fixed in Section 5. We suppose that the tree T_α belongs to $\mathbb{P}_{<\alpha}$. (If not then we don't worry about 5° .) Consider, according to 2° , a tree $U = U_M \in \mathcal{U}_\alpha$ satisfying $T \subseteq T_\alpha$. Using Corollary 3.2, Lemma 3.3, and Theorem 4.2, we shrink each tree $U_n \in \mathcal{U}_\alpha$ to a tree $U'_n \in \mathbf{ST}$, $U' \subseteq U$, so that the function $F_{b_\alpha}^2$ is normalized on U'_M for $\mathcal{U}' = \{U'_n : n < \omega\}$ and F_{c_α} is continuous and either a bijection or a constant on $[U'_M]$. Take \mathcal{U}' as the final \mathcal{U}_α and T' as U'_M to fulfill 5° .

□ (Lemma 5.1)

7 The model, part I

We use the product $\mathbb{P} \times \mathbb{C}$ of the forcing notion \mathbb{P} defined in \mathbf{L} in Section 5 and satisfying conditions 1°–6° as above, and the Cohen forcing, here in the form of $\mathbb{C} = \omega^{<\omega}$, to prove the following more detailed form of Theorem 1.1. The proof of this theorem in the next three sections is based on a combination of different ideas.

Theorem 7.1. *Let a pair of reals $\langle a_0, x_0 \rangle$ be $\mathbb{P} \times \mathbb{C}$ -generic over \mathbf{L} . Then*

- (I) a_0 is not **OD**, and moreover, $\mathbf{HOD} = \mathbf{L}$ in $\mathbf{L}[a_0, x_0]$;
- (II) a_0 belongs to **HNT**, and moreover, $\mathbf{L}[a_0] \subseteq \mathbf{HNT}$ in $\mathbf{L}[a_0, x_0]$;
- (III) x_0 does not belong to **HNT**, and moreover, $\mathbf{HNT} \subseteq \mathbf{L}[a_0]$ in $\mathbf{L}[a_0, x_0]$.

We prove Claim (I) of the theorem in this section. The proof is based on several lemmas. According to the next lemma, it suffices to prove that $\mathbf{HOD} = \mathbf{L}$ in $\mathbf{L}[a_0]$.

Lemma 7.2. $(\mathbf{HOD})^{\mathbf{L}[a_0, x_0]} \subseteq (\mathbf{HOD})^{\mathbf{L}[a_0]}$.

Proof. By the forcing product theorem, x_0 is a Cohen generic real over $\mathbf{L}[a_0]$. It follows by a standard argument based on the full homogeneity of the Cohen forcing \mathbb{C} that if $H \subseteq \mathbf{Ord}$ is **OD** in $\mathbf{L}[a_0, x_0]$ then $H \in \mathbf{L}[a_0]$ and H is **OD** in $\mathbf{L}[a_0]$.

Now prove the implication $Y \in (\mathbf{HOD})^{\mathbf{L}[a_0, x_0]} \implies Y \in \mathbf{L} \wedge Y \in (\mathbf{HOD})^{\mathbf{L}[a_0]}$ by induction on the set-theoretic rank $\mathbf{rk} x$ of $x \in \mathbf{L}[a_0, x_0]$. Since each set consists only of sets of strictly lower rank, it is sufficient to check that if a set $H \in \mathbf{L}[a_0, x_0]$ satisfies $H \subseteq (\mathbf{HOD})^{\mathbf{L}[a_0]}$ and $H \in \mathbf{HOD}$ in $\mathbf{L}[a_0, x_0]$ then $H \in \mathbf{L}[a_0]$ and $H \in (\mathbf{OD})^{\mathbf{L}[a_0]}$. Here we can assume that in fact $H \subseteq \mathbf{Ord}$, since **HOD** allows an **OD** wellordering and hence an **OD** bijection onto **Ord**. But in this case $H \in \mathbf{L}[a_0]$ and H is **OD** in $\mathbf{L}[a_0]$ by the above, as required. \square

Lemma 7.3 (Lemma 7.5 in [10]). a_0 is not **OD** in $\mathbf{L}[a_0]$.

Proof. Suppose towards the contrary that a_0 is **OD** in $\mathbf{L}[a_0]$. But a_0 is a \mathbb{P} -generic real over \mathbf{L} , so the contrary assumption is forced. In other words, there is a tree $T \in \mathbb{P}$ with $a_0 \in [T]$ and a formula $\vartheta(x)$ with ordinal parameters, such that if $a \in [T]$ is \mathbb{P} -generic over \mathbf{L} then a is the only real in $\mathbf{L}[a]$ satisfying $\vartheta(a)$. Let $s = \mathbf{stem}(\langle T \rangle)$. Then both $s \smallfrown 0$ and $s \smallfrown 1$ belong to T , and either $s \smallfrown 0 \subset a_0$ or $s \smallfrown 1 \subset a_0$. Let, say, $s \smallfrown 0 \subset a_0$. Let $n = \mathbf{lh}(s)$ and $\sigma = 0^n \smallfrown 1$, so that all three strings $s \smallfrown 0$, $s \smallfrown 1$, σ belong to 2^{n+1} , and $s \smallfrown 0 = \sigma \cdot (s \smallfrown 1)$. As the forcing \mathbb{P} is invariant under the action of σ , the real $a_1 = \sigma \cdot a_0$ is \mathbb{P} -generic over \mathbf{L} , and $\sigma \cdot T = T$. We conclude that it is true in $\mathbf{L}[a_1] = \mathbf{L}[a_0]$ that a_1 is still the only real in $\mathbf{L}[a_1]$ satisfying $\vartheta(a_1)$. However obviously $a_1 \neq a_0$! \square

Lemma 7.4. *If $b \in \mathbf{L}[a_0] \setminus \mathbf{L}$ is a real then b is not **OD** in $\mathbf{L}[a_0]$.*

Proof. It follows from Lemma 5.2 (and the countability of \mathbb{C}) that the forcing $\mathbb{P} \times \mathbb{C}$ preserves cardinals. We conclude that $b = g(a_0)$ for some Borel function $g = F_r : 2^\omega \rightarrow 2^\omega$ with a code $r \in \mathbf{BC} \cap \mathbf{L}$. Now by Lemma 5.3 there is a tree $S \in \mathbb{P}$

such that $a_0 \in [S]$ and $h = g \upharpoonright [S]$ is a bijection of a constant. If h is a bijection then $b \notin \mathbf{OD}$ in $\mathbf{L}[a_0]$ since otherwise $a_0 = h^{-1}(b) \in \mathbf{OD}$, contrary to Lemma 7.3. If h is a constant, so that there is a real $b_0 \in \mathbf{L} \cap 2^\omega$ such that $h(a) = b_0$ for all $a \in [S]$, then $b = h(a_0) = c \in \mathbf{L}$, contrary to the choice of b . \square

Lemma 7.5. *If $X \subseteq \mathbf{Ord}$, $X \in \mathbf{L}[a_0] \setminus \mathbf{L}$, then X is not \mathbf{OD} in $\mathbf{L}[a_0]$.*

Proof. Suppose to the contrary that $X \subseteq \mathbf{Ord}$, $X \in \mathbf{L}[a_0] \setminus \mathbf{L}$, and X is \mathbf{OD} in $\mathbf{L}[a_0]$. Let t be a \mathbb{P} -name for X . Then a condition $T_0 \in \mathbb{P}$ (a Silver tree) \mathbb{P} -forces

$$t \in \mathbf{L}[a_0] \setminus \mathbf{L} \wedge t \in \mathbf{OD}$$

over \mathbf{L} . Say that t *splits* conditions $S, T \in \mathbb{P}$ if there is an ordinal γ such that S forces $\gamma \in t$ but T forces $\gamma \notin t$ or vice versa; let γ_{ST} be the least such an ordinal γ .

We claim that the set

$$D = \{\langle S, T \rangle : S, T \in \mathbb{P} \wedge S \cup T \subseteq T_0 \wedge t \text{ splits } S, T\} \in \mathbf{L}$$

is dense in $\mathbb{P} \times \mathbb{P}$ above $\langle T_0, T_0 \rangle$. Indeed let $S, T \in \mathbb{P}$ be subtrees of T_0 . If t splits no stronger pair of trees $S' \subseteq S$, $T' \subseteq T$ in \mathbb{P} then easily both S and T decide $\gamma \in t$ for every ordinal γ , a contradiction with the choice of T_0 . Thus D is indeed dense.

Let, in \mathbf{L} , $A \subseteq D$ be a maximal antichain; A is countable in \mathbf{L} by Lemma 5.2, and hence the set $W = \{\gamma_{ST} : \langle S, T \rangle \in A\} \in \mathbf{L}$ is countable in \mathbf{L} . We claim that

(\ddagger) the intersection $b = X \cap W$ does not belong to \mathbf{L} .

Indeed otherwise there is a tree $T_1 \in \mathbb{P}$, $T_1 \subseteq T_0$, which \mathbb{P} -forces that $t \cap W = b$. (The sets $W, b \in \mathbf{L}$ are identified with their names.)

By the countability of A, W there is an ordinal $\alpha < \omega_1^{\mathbf{L}}$ such that $A \subseteq \mathbb{P}_{<\alpha} \times \mathbb{P}_{<\alpha}$, $T_1 \in \mathbb{P}_{<\alpha}$, and $W \subseteq \alpha$. We can w.l.o.g. assume that $A \in \mathfrak{M}_\alpha$, for if not then further increase α below $\omega_1^{\mathbf{L}}$ accordingly. Let $u = \text{stem}(T_1)$. The trees $T_{10} = T_1 \upharpoonright_{u \frown 0}$ and $T_{11} = T_1 \upharpoonright_{u \frown 1}$ belong to $\mathbb{P}_{<\alpha}$ along with T_1 , and hence there are trees $U, V \in \mathcal{U}_\alpha$ with $U \subseteq T_{10}$ and $V \subseteq T_{11}$. Clearly $U \neq V$, so that we have $[U] \times [V] \subseteq \bigcup_{\langle U', V' \rangle \in A'} [U'] \times [V']$ for a finite set $A' \subseteq A$ by 4° of Section 5. Now take reals $a' \in [U]$ and $a'' \in [V]$ both \mathbb{P} -generic over \mathbf{L} . Then there is a pair of trees $\langle U', V' \rangle \in A'$ such that $a' \in [U']$ and $a'' \in [V']$. The interpretations $X' = t[a']$ and $X'' = t[a'']$ are then different on the ordinal $\gamma = \gamma_{U'V''} \in W$ since $A' \subseteq A \subseteq D$. Thus the restricted sets $b' = X' \upharpoonright W$ and $b'' = X'' \upharpoonright W$ differ from each other. In particular at least one of b', b'' is not equal to b . But $a', a'' \in [T_1]$ by construction, hence this contradicts the choice of T_1 and completes the proof of (\ddagger).

Recall that $b \subseteq W$, and $W \in \mathbf{L}$ is countable in \mathbf{L} . It follows that b can be considered as a real, so we conclude that b is not \mathbf{OD} in $\mathbf{L}[a_0]$ by Lemma 7.4 and (\ddagger).

However $b = X \cap W$, where X is \mathbf{OD} and $W \in \mathbf{L}$, hence W is \mathbf{OD} in $\mathbf{L}[a_0]$ and b is \mathbf{OD} in $\mathbf{L}[a_0]$. The contradiction obtained ends the proof of the lemma. \square (Lemma)

Now Theorem 7.1(I) immediately follows from Lemma 7.2 and Lemma 7.5.

\square (Claim (II) of Theorem 7.1)

8 The model, part II

Here we establish Claim (II) of Theorem 7.1. To prove $\mathbf{L}[a_0] \subseteq \mathbf{HNT}$ it suffices to show that a_0 itself belongs to \mathbf{HNT} , and then make use of the fact that by Gödel every set $z \in \mathbf{L}[a_0]$ has the form $x = F(a_0)$, where F is an \mathbf{OD} function.

Further, to prove $a_0 \in \mathbf{HNT}$ it suffices to check that the \mathbf{E}_0 -equivalence class² $[a_0]_{\mathbf{E}_0} = \{b \in 2^\omega : a_0 \mathbf{E}_0 b\}$ (which is a countable set) of our generic real a_0 is an \mathbf{OD} set in $\mathbf{L}[a_0, x_0]$. According to 6°, it suffices to establish the equality

$$[a_0]_{\mathbf{E}_0} = \bigcap_{\xi < \omega_1} \bigcup_{T \in \mathbb{P}_\xi} [T]. \quad (*)$$

Note that every set \mathbb{P}_ξ is pre-dense in \mathbb{P} ; this follows from 3° and 5°, see, for example, Lemma 6.3 in [10]. This immediately implies $a_0 \in \bigcup_{T \in \mathbb{P}_\xi} [T]$ for each ξ . Yet all sets \mathbb{P}_ξ are invariant w.r.t. shifts by construction. Thus we have \subseteq in (*).

To prove the inverse inclusion, assume that a real $b \in 2^\omega$ belongs to the right-hand side of (*) in $\mathbf{L}[a_0, x_0]$. It follows from Lemma 5.2 (and the countability of \mathbb{C}) that the forcing $\mathbb{P} \times \mathbb{C}$ preserves cardinals. We conclude that $b = g(a_0, x_0)$ for some Borel function $g = F_q : 2^\omega \times \omega^\omega \rightarrow 2^\omega$ with a code $q \in \mathbf{BC} \cap \mathbf{L}$.

Assume to the contrary that $b = g(a_0, x_0) \notin [a_0]_{\mathbf{E}_0}$.

Since $x_0 \in \omega^\omega$ is a \mathbb{C} -generic real over $\mathbf{L}[a_0]$ by the forcing product theorem, this assumption is forced, so that there is a tuple $u \in \mathbb{C} = \omega^{<\omega}$ such that

$$f(a_0, x) \in \bigcap_{\xi < \omega_1} \bigcup_{T \in \mathbb{P}_\xi} [T] \setminus [a_0]_{\mathbf{E}_0},$$

whenever a real $x \in \mathcal{N}_u$ is \mathbb{C} -generic over $\mathbf{L}[a_0]$. (Recall that $\mathcal{N}_u = \{y \in \omega^\omega : u \subset y\}$.) Let H be the canonical homomorphism of ω^ω onto \mathcal{N}_u . We put $f(a, x) = g(a, H(x))$ for $a \in 2^\omega$, $x \in \omega^\omega$. Then H preserves the \mathbb{C} -genericity, and hence

$$f(a_0, x) \in \bigcap_{\xi < \omega_1} \bigcup_{T \in \mathbb{P}_\xi} [T] \setminus [a]_{\mathbf{E}_0}, \quad (**)$$

whenever $x \in \omega^\omega$ is \mathbb{C} -generic over $\mathbf{L}[a_0]$. Note that f also has a Borel code $r \in \mathbf{BC}$ in \mathbf{L} , so that $f = F_r$.

It follows from Lemma 5.3 that there is an ordinal $\alpha < \omega_1$ and a tree $S \in \mathcal{U}_\alpha$, on which f is normalized for \mathcal{U}_α , and which satisfies $a_0 \in [S]$. Normalization means that, in \mathbf{L} , there is a dense \mathbf{G}_δ set $X \subseteq \omega^\omega$ satisfying one of the two options of Definition 4.1. Consider a real $z \in \omega^\omega \cap \mathbf{L}$ (a \mathbf{G}_δ -code for X in \mathbf{L}) such that $X = X_z = \bigcap_k \bigcup_{z(2^k \cdot 3^j) = 1} \mathcal{N}_{w_j}$, where $2^{<\omega} = \{w_j : j < \omega\}$ is a fixed recursive enumeration of tuples.

Case 1: there are tuples $v \in \omega^{<\omega}$, $\sigma \in 2^{<\omega}$, such that $f(a, x) = \sigma \cdot a$ for all points $a \in [S]$ and $x \in \mathcal{N}_v \cap X$. In other words, it is true in \mathbf{L} that

$$\forall a \in [S] \forall x \in \mathcal{N}_v \cap X_z (f(a, x) = \sigma \cdot a).$$

²Recall that the equivalence relation \mathbf{E}_0 is defined on 2^ω so that $a \mathbf{E}_0 b$ iff the set $a \Delta b = \{k : a(k) \neq b(k)\}$ is finite. Equivalently, $a \mathbf{E}_0 b$ iff $a = \sigma \cdot b$ for some tuple $\sigma \in 2^{<\omega}$. Then $[a]_{\mathbf{E}_0} = \{b \in 2^\omega : a \mathbf{E}_0 b\} = \{\sigma \cdot a : \sigma \in 2^{<\omega}\}$ is the \mathbf{E}_0 -equivalence class of a .

But this formula is absolute by Shoenfield, so it is also true in $\mathbf{L}[a_0, x_0]$. Take $a = a_0$ (recall: $a_0 \in [S]$) and any real $x \in \mathcal{N}_v$, \mathbb{C} -generic over $\mathbf{L}[a_0]$. Then $x \in X_z$, because X_z is a dense \mathbf{G}_δ with a code even from \mathbf{L} . Thus $f(a_0, x) = \sigma \cdot a_0 \in [a_0]_{E_0}$, which contradicts (**).

Case 2: $f(a, x) \notin \bigcup_{\sigma \in 2^{<\omega} \wedge U \in \mathcal{U}_\alpha} \sigma \cdot [U]$ for all $a \in [S]$ and $x \in X$. By the definition of \mathbb{P}_α , this implies $f(a, x) \notin \bigcup_{T \in \mathbb{P}_\alpha} [T]$ for all $a \in [S]$ and $x \in X$, and this again contradicts (**) for $a = a_0$.

The resulting contradiction in both cases refutes the contrary assumption above and completes the proof.

□ (Claim (II) of Theorem 7.1)

9 The model, part III

Here we prove Claim (III) of Theorem 7.1. We make use of the following result here.

Lemma 9.1. *Let $x \in \omega^\omega$ be Cohen-generic over a set universe \mathbf{V} . Then it holds in $\mathbf{V}[x]$ that if $Z \subseteq 2^\omega$ is a countable \mathbf{OD} set then $Z \in \mathbf{V}$. More generally if $q \in 2^\omega \cap \mathbf{V}$ then it holds in $\mathbf{V}[x]$ that if $Z \subseteq 2^\omega$ is a countable $\mathbf{OD}(q)$ set then $Z \in \mathbf{V}$. □*

Proof (sketch). The pure \mathbf{OD} case is Theorem 1.1 in [11].³ The proof of the general case does not differ, q is present in the flow of arguments as a passive parameter. □

This result admits the following extension for the case $\mathbf{V} = \mathbf{L}$. Here $\mathbf{OD}(a)$ naturally means sets definable by a formula containing a_0 and ordinals as parameters

Corollary 9.2. *Assume that $a \in 2^\omega$ and $x \in \omega^\omega$ is Cohen-generic over $\mathbf{L}[a]$. Then it holds in $\mathbf{L}[a, x]$ that if $X \in \mathbf{L}[a]$ and $A \subseteq 2^X$ is a countable $\mathbf{OD}(a)$ set then $A \subseteq \mathbf{L}$.*

Proof. As the Cohen forcing is countable, there is a set $Y \subseteq X$, $Y \in \mathbf{L}[a]$, countable in $\mathbf{L}[a]$ and such that if $f \neq g$ belong to 2^X then $f(x) \neq g(x)$ for some $x \in Y$. Then Y is countable and $\mathbf{OD}(a)$ in $\mathbf{L}[a, x]$, so the *projection* $B = \{f \upharpoonright Y : f \in A\}$ of the set A will also be countable and $\mathbf{OD}(a)$ in $\mathbf{L}[a, x]$. We have $B \in \mathbf{L}[a]$ by Lemma 9.1. (The set Y here can be identified with ω .) Hence, each $f \in B$ is $\mathbf{OD}(a)$ in $\mathbf{L}[a, x]$. However, if $f \in A$ and $w = f \upharpoonright Y$, then by the choice of Y it holds in $\mathbf{L}[a, x]$ that f is the only element in A satisfying $f \upharpoonright Y = w$. Therefore $f \in \mathbf{OD}(a)$ in $\mathbf{L}[a, x]$. We conclude that $f \in \mathbf{L}[a]$. □

Proof (Claim (III) of Theorem 7.1). We prove an even stronger claim

$$x \in \mathbf{HNT}(a_0) \implies x \in \mathbf{L}[a_0]$$

in $\mathbf{L}[a_0, x_0]$ by induction on the set-theoretic rank $\mathbf{rk} x$ of sets $x \in \mathbf{L}[a_0, x_0]$. Here $\mathbf{HNT}(a_0)$ naturally means all sets hereditarily $\mathbf{NT}(a_0)$, the latter means all elements of countable sets in $\mathbf{OD}(a_0)$.

³See our papers [11, 15, 14] for more on countable and Borel \mathbf{OD} sets in Cohen and some other generic extensions.

Since each set consists only of sets of strictly lower rank, it is sufficient to check that if a set $H \in \mathbf{L}[a_0, x_0]$ satisfies $H \subseteq \mathbf{L}[a_0]$ and $H \in \mathbf{HNT}(a_0)$ in $\mathbf{L}[a_0, x_0]$ then $H \in \mathbf{L}[a_0]$. Here we can assume that in fact $H \subseteq \mathbf{Ord}$, since $\mathbf{L}[a_0]$ allows an $\mathbf{OD}(a_0)$ wellordering. Thus, let $H \subseteq \lambda \in \mathbf{Ord}$. Additionally, since $H \in \mathbf{HNT}(a_0)$, we have, in $\mathbf{L}[a_0, x_0]$, a countable $\mathbf{OD}(a_0)$ set $A \subseteq \mathcal{P}(\lambda)$ containing H . However, $A \in \mathbf{L}[a_0]$ by Corollary 9.2. This implies $H \in \mathbf{L}[a_0]$ as required. \square

\square (Claim (III) and Theorem 7.1 as a whole)

\square (Theorem 1.1)

10 Comments and questions

1. Recall that if x is a Cohen real over \mathbf{L} then $\mathbf{HNT} = \mathbf{L}$ in $\mathbf{L}[x]$ by Lemma 9.1.

Problem 10.1. Is it true in generic extensions of \mathbf{L} by a single Cohen generic real that a countable \mathbf{OD} set of any kind necessarily consists only of \mathbf{OD} elements?

We cannot solve this even for *finite* \mathbf{OD} sets.

By the way it is not that obvious to expect the *positive* answer. Indeed, the problem solves in the *negative* for Sacks and some other generic extensions even for *pairs*, see [1, 2]. For instance, if x is a Sacks-generic real over \mathbf{L} then it is true in $\mathbf{L}[x]$ that there is an \mathbf{OD} unordered pair $\{X, Y\}$ of sets of reals $X, Y \subseteq \mathcal{P}(2^\omega)$ such that X, Y themselves are non- \mathbf{OD} sets. See [1] for a proof of this rather surprising result originally by Solovay.

2. See Fuchs [3] (unpublished) for some other research lines related to Russell-nontypical sets with various cardinal parameters.

References

- [1] Ali Enayat and Vladimir Kanovei, *An unpublished theorem of Solovay on OD partitions of reals into two non-OD parts, revisited*, J. Math. Log. **21** (2021), no. 3, 1–22, Article No. 2150014.
- [2] Ali Enayat, Vladimir Kanovei, and Vassily Lyubetsky, *On effectively indiscernible projective sets and the Leibniz-Mycielski axiom*, Mathematics **9** (2021), no. 14, 1–19 (English), Article No 1670.
- [3] Gunter Fuchs, *Blurry definability*, Preprint, 2021.
- [4] Gunter Fuchs, Victoria Gitman, and Joel David Hamkins, *Ehrenfeucht’s lemma in set theory*, Notre Dame J. Formal Logic **59** (2018), no. 3, 355–370 (English).
- [5] Marcia J. Groszek and Joel David Hamkins, *The implicitly constructible universe*, J. Symb. Log. **84** (2019), no. 4, 1403–1421 (English).
- [6] Joel David Hamkins and Cole Leahy, *Algebraicity and implicit definability in set theory*, Notre Dame J. Formal Logic **57** (2016), no. 3, 431–439 (English).
- [7] Thomas Jech, *Set theory*, The third millennium revised and expanded ed., Springer-Verlag, Berlin-Heidelberg-New York, 2003 (English), Pages xiii + 769.

- [8] Ronald Jensen, *Definable sets of minimal degree*, Math. Logic Found. Set Theory, Proc. Int. Colloqu., Jerusalem 1968 (Yehoshua Bar-Hillel, ed.), Studies in logic and the foundations of mathematics, vol. 59, North-Holland, Amsterdam-London, 1970, pp. 122–128.
- [9] Vladimir Kanovei and Vassily Lyubetsky, *On some classical problems in descriptive set theory*, Russ. Math. Surv. **58** (2003), no. 5, 839–927 (Russian, English).
- [10] ———, *A definable E_0 class containing no definable elements*, Arch. Math. Logic **54** (2015), no. 5-6, 711–723 (English).
- [11] ———, *Countable OD sets of reals belong to the ground model*, Arch. Math. Logic **57** (2018), no. 3-4, 285–298 (English).
- [12] ———, *Definable E_0 classes at arbitrary projective levels*, Ann. Pure Appl. Logic **169** (2018), no. 9, 851–871 (English).
- [13] ———, *Non-uniformizable sets of second projective level with countable cross-sections in the form of Vitali classes*, Izvestiya: Mathematics **82** (2018), no. 1, 61–90.
- [14] ———, *Borel OD sets of reals are OD-Borel in some simple models*, Proc. Am. Math. Soc. **147** (2019), no. 3, 1277–1282 (English).
- [15] ———, *Definable elements of definable Borel sets*, Math. Notes **105** (2019), no. 5, 684–693 (English).
- [16] ———, *Models of set theory in which separation theorem fails*, Izvestiya: Mathematics **85** (2021), no. 6, to appear.
- [17] ———, *On Russell typicality in Set Theory*, arXiv e-prints (2021), arXiv:2111.07654.
- [18] Alexander S. Kechris, *Classical descriptive set theory*, Springer-Verlag, New York, 1995. MR 96e:03057
- [19] Michiel van Lambalgen, *The axiomatization of randomness*, J. Symb. Log. **55** (1990), no. 3, 1143–1167 (English).
- [20] Robert M. Solovay, *A model of set-theory in which every set of reals is Lebesgue measurable*, Ann. Math. (2) **92** (1970), 1–56.
- [21] Athanassios Tzouvaras, *Typicality á la Russell in set theory*, ResearchGate Preprint, May 2021, 14 pp., ResearchGate Link.
- [22] ———, *Russell’s typicality as another randomness notion*, Mathematical Logic Quarterly **66** (2020), no. 3, 355–365.