

PERMUTATIONS ON WEIERSTRASS PRYM EIGENFORMS

RODOLFO GUTIÉRREZ-ROMO AND ANGEL PARDO

ABSTRACT. Let $X \in \mathcal{H}(2)$ be a Veech surface of discriminant D and let $G(X)$ be the permutation group induced by the affine group of X on the set of Weierstrass points of X . We show that $G(X) \cong \text{Dih}_4$ if $D \equiv_4 0$, $G(X) \cong \text{Dih}_5$ if $D \equiv_8 5$, and $G(X) \cong \text{Dih}_6$ if $D \equiv_8 1$, where Dih_n is the dihedral group of order $2n$. Thus, $G(X)$ is a weak invariant, as it can distinguish the residue class of $D \pmod 8$, but it cannot tell different spin invariants apart when $D \equiv_8 1$. Moreover, we show that the same groups arise when we only consider the action of the *parabolic* elements of the Veech group of X on the Weierstrass points.

We prove a similar result in the Weierstrass Prym locus in genus three. Indeed, we show that the permutation group induced by the affine group on the set of fixed points for the Prym involution of a Weierstrass Prym eigenform of discriminant D is isomorphic to Sym_2 when D is an even quadratic residue modulo 16, and it is isomorphic to Sym_3 otherwise. In genus three, the same group is also obtained when we consider only *parabolic* elements. Since the permutation action of the affine group on Prym fixed points is trivial in genus four and no Weierstrass Prym eigenforms exist for genus five or larger, we completely classify this action for all Weierstrass Prym eigenforms.

CONTENTS

1. Introduction	2
2. Teichmüller curves in $\mathcal{H}(2)$	6
3. Action of parabolic elements in genus two	9
4. Restrictions for permutations in genus two	14
5. Weierstrass eigenforms in genus two of small discriminant	20
6. Classification of permutations in genus two	23
7. Weierstrass Prym eigenforms	24
8. Action of parabolic elements in genus three	27
9. Restrictions for permutations in genus three	29
10. Weierstrass Prym eigenforms in genus three of small discriminant	31
11. Classification of permutations in genus three	33
References	34

2020 *Mathematics Subject Classification.* 37C85 (primary), and 14H55, 37D40 (secondary).

Key words and phrases. Veech group, translation surface, Veech surface, lattice surface, Weierstrass point, Prym involution, Weierstrass Prym eigenform, permutation group, dihedral group.

This work was supported by the Center for Mathematical Modeling (CMM), ACE210010 and FB210005, BASAL funds for centers of excellence from ANID-Chile, and the MATH-AmSud 21-MATH-07 grant. The first named author was also supported by ANID-Chile through the FONDECYT Iniciación 11190034 grant, while the second named author was also supported by ANID-Chile through the FONDECYT 3190257 grant.

1. INTRODUCTION

A *translation surface* is a pair (X, ω) , where X is a Riemann surface and ω is a nonzero Abelian differential on X . If $\omega(p) = 0$ for $p \in X$, we say that p is a *zero* or *singularity*. The *order* of p is the order of p viewed as a zero of ω . On the other hand, if $\omega(p) \neq 0$, we say that p is a *regular point*. By a slight abuse of notation, we will often omit the differential and say that X is a translation surface. For an introduction and more details on this subject, we refer the reader to Zorich's survey [Zor06].

The moduli space of translation surfaces is stratified by the list of orders of zeros of the Abelian differential: the *stratum* $\mathcal{H}(\kappa_1, \dots, \kappa_n)$ consists of translation surfaces whose singularities have prescribed orders $\kappa_1, \dots, \kappa_n$ (with multiplicities). The genus $g \geq 1$ of X can be deduced from the combinatorial data by the Gauss–Bonnet-like formula $\kappa_1 + \dots + \kappa_n = 2g - 2$. Strata are endowed with the natural topology coming from period coordinates [Zor06, Section 3].

A translation surface may also be defined as an equivalence class of collections of polygons in \mathbb{C} in which every side is identified with another parallel, equal-length, opposite side by translation. The equivalence relation is the one induced by *cut-and-paste operations*: two such collections represent the same translation surface if it is possible to cut pieces of the former along straight lines and glue them using the side identifications in order to obtain the latter. This polygon definition induces a *flat metric* on the translation surface which becomes singular at the zeros. Indeed, an order- k zero of the Abelian differential corresponds to a singular point for the flat metric, with cone angle $2\pi(1+k)$.

There exists a natural action of $\mathrm{SL}(2, \mathbb{R})$ on a stratum given by linearly deforming the collection of polygons (by using $\mathbb{C} \cong \mathbb{R}^2$). With the first definition of translation surface, this action is exactly post-composition with coordinate charts (again using $\mathbb{C} \cong \mathbb{R}^2$). The $\mathrm{SL}(2, \mathbb{R})$ -action preserves both the stratum in which a translation surface X lies and the total area of X . Hence, it induces an action on the set $\mathcal{H}^{(r)}(\kappa) \subseteq \mathcal{H}(\kappa)$ of area- r translation surfaces inside a given stratum $\mathcal{H}(\kappa)$, for some fixed $r > 0$.

Let $X \in \mathcal{H}^{(r)}(\kappa)$ be an area- r translation surface. Regarding its orbit by the $\mathrm{SL}(2, \mathbb{R})$ -action, two extreme cases can arise:

- X is a *Veech surface* (or simply *Veech*), that is, its $\mathrm{SL}(2, \mathbb{R})$ -orbit is closed inside $\mathcal{H}^{(r)}(\kappa)$; or
- X is *generic*, that is, its $\mathrm{SL}(2, \mathbb{R})$ -orbit is dense in \mathcal{C} , where \mathcal{C} is the connected component of $\mathcal{H}^{(r)}(\kappa)$ containing X .

The stabiliser $\mathrm{SL}(X)$ of X for the $\mathrm{SL}(2, \mathbb{R})$ -action is called the *Veech group* of X . We have that X is a Veech surface if and only if $\mathrm{SL}(X)$ is a lattice inside $\mathrm{SL}(2, \mathbb{R})$, that is, $\mathrm{SL}(2, \mathbb{R})/\mathrm{SL}(X)$ has finite measure¹. For this reason, Veech surfaces are also often called *lattice surfaces*.

The projection of the $\mathrm{SL}(2, \mathbb{R})$ -orbit of a Veech surface into the moduli space of Riemann surfaces is called a *Teichmüller curve*. They come in two different flavours:

- the *arithmetic* ones, which arise as the projection of the $\mathrm{SL}(2, \mathbb{R})$ -orbit of a *square-tiled surface*, that is, a translation surface that admits a covering of the unit torus branched over a single point; and
- *non-arithmetic* ones, which cannot be defined in terms of a square-tiled surface.

¹Veech surfaces were originally defined as those whose Veech group is a lattice. The fact that this is indeed equivalent to having a closed $\mathrm{SL}(2, \mathbb{R})$ -orbit is known as *Smillie's theorem*, since it was originally proved, though not published, by John Smillie; it was first referenced in Veech's work [Vee95].

The distinction between arithmetic and non-arithmetic Teichmüller curves is important, as different tools are usually used for their study. Indeed, square-tiled surfaces admit purely combinatorial descriptions, unlike those belonging to non-arithmetic Teichmüller curves.

The $\mathrm{SL}(2, \mathbb{R})$ -action can be thought of as part of a larger action by $\mathrm{GL}^+(2, \mathbb{R})$ on a stratum; the main difference being that the latter does not necessarily preserve area. Nevertheless, scaling a translation surface will not change the aforementioned properties (such as being Veech or generic). More precisely, for any $r, r' > 0$, there exists a homeomorphism between the spaces $\mathcal{H}^{(r)}(\kappa)$ and $\mathcal{H}^{(r')}(\kappa)$ simply given by the action of a constant multiple of the identity matrix in $\mathrm{GL}^+(2, \mathbb{R})$. As these matrices commute with any other matrix, the homeomorphism is $\mathrm{SL}(2, \mathbb{R})$ -equivariant. We will sometimes make use of such scalings to simplify some computations.

Closed orbits in $\mathcal{H}(2)$. We are interested in translation surfaces X in the minimal stratum $\mathcal{H}(2)$ in genus two. Every such surface is *hyperelliptic*, that is, it admits an involution $\iota: X \rightarrow X$ satisfying $\iota^*\omega = -\omega$ and such that X/ι is (topologically) a sphere. This involution, known as the *hyperelliptic involution*, has exactly six fixed points: the order-2 zero of ω and five regular points w_1, \dots, w_5 . Collectively, these six points are known as the *Weierstrass points* of X .

By the work of McMullen [McM07], every surface in $\mathcal{H}(2)$ is either Veech or generic. Moreover, McMullen [McM05] also showed that Teichmüller curves coming from closed $\mathrm{SL}(2, \mathbb{R})$ -orbits in $\mathcal{H}(2)$ are classified by two invariants: a positive integer $D > 4$ satisfying $D \equiv_4 0$ or $D \equiv_4 1$, known as the *discriminant*, and, when $D \equiv_8 1$ and $D \neq 9$, the *spin invariant*, which can be either even or odd. The *discriminant* of a Veech surface is the discriminant of the Teichmüller curve that it induces.

The action of the Veech group on Weierstrass points in $\mathcal{H}(2)$. Let X be a translation surface in $\mathcal{H}(2)$. We say that a bijective, orientation-preserving map $f: X \rightarrow X$ is an *affine transformation* if its local expressions (on the charts given by the flat metric) are affine maps of \mathbb{R}^2 . We denote the group of affine transformations on X by $\mathrm{Aff}(X)$.

The linear part of an affine transformation does not depend on the choice of flat charts. This observation gives rise to a group homomorphism $D: \mathrm{Aff}(X) \rightarrow \mathrm{SL}(2, \mathbb{R})$, known as the *derivative map*. The *Veech group* $\mathrm{SL}(X)$ of X can be defined as the image of the derivative map, that is, $\mathrm{SL}(X) = D\mathrm{Aff}(X)$. Veech showed that the derivative map is finite-to-one; the kernel of this map is hence a finite group, which is known as the automorphism group $\mathrm{Aut}(X)$ of X . In other words, we have a short exact sequence

$$1 \rightarrow \mathrm{Aut}(X) \rightarrow \mathrm{Aff}(X) \rightarrow \mathrm{SL}(X) \rightarrow 1.$$

In the case of minimal strata, the group $\mathrm{Aut}(X)$ is trivial, so D is one-to-one [HL06, Proposition 4.4]. Thus, in what follows, we will implicitly make use of the identification $\mathrm{SL}(X) \cong \mathrm{Aff}(X)$ for surfaces in $\mathcal{H}(2)$.

Now, we turn our attention to the action of $\mathrm{SL}(X)$ on regular Weierstrass points. More precisely, if $X \in \mathcal{H}(2)$, the group $\mathrm{SL}(X)$ acts by permutations on the set $\{w_1, \dots, w_5\}$. The permutation group $G(X) \leq \mathrm{Sym}(\{w_1, \dots, w_5\})$ is the group obtained by all such permutations. Observe that this group is only well-defined up to conjugation by elements of the Veech group. In other words, we do not keep track the “names” of the regular Weierstrass points.

The HLK-invariant. In the arithmetic case, a regular Weierstrass point on X projects down to a 2-torsion point on the unit torus $\mathbb{C}/\mathbb{Z}[i]$. Moreover, any element of $\mathrm{SL}(2, \mathbb{R})$

preserves both the sets of 1-torsion points and of 2-torsion points. These observations give rise to an invariant for the $\mathrm{SL}(2, \mathbb{R})$ -orbit of X , defined as the distribution of these projections on the unit torus; it is known as *HLK-invariant* since it was first considered by Kani [Kan06] and Hubert–Lelièvre [HL06] (see also the work by Matheus–Möller–Yoccoz [MMY15]). This invariant naturally restricts the action of $\mathrm{Aff}(X)$ on the set of regular Weierstrass points. We prove that this is the only restriction.

Moreover, in the non-arithmetic case, two distinct situations can arise. If the discriminant D is a quadratic residue modulo 8, a suitable reduction allows us to apply the same criterion even if the periods are no longer integral (they belong to a real quadratic field). This is no longer true when D is a quadratic nonresidue modulo 8 and thus, we are not able to extend this argument for $D \equiv_8 5$. However, we are still able to adapt these ideas to constrain the action of the affine group on regular Weierstrass points and show that these restrictions are, as well, the only ones.

Main results for genus two. Our main result is thus the following:

Theorem 1.1. *Let $X \in \mathcal{H}(2)$ be a Veech surface of discriminant $D \geq 9$ and let w_1, \dots, w_5 be its regular Weierstrass points. Let $G(X) \leq \mathrm{Sym}(\{w_1, \dots, w_5\})$ be the permutation group induced by the action of $\mathrm{SL}(X)$ on $\{w_1, \dots, w_5\}$. Then, $G(X)$ is isomorphic to:*

- Dih_4 if $D \equiv_4 0$;
- Dih_5 if $D \equiv_8 5$; and
- Dih_6 if $D \equiv_8 1$, regardless of the spin;

where Dih_n is the dihedral group of order $2n$.

Note that the previous theorem implies that the isomorphism class of $G(X)$ is a “weak” invariant: it allows us to deduce the residue class of the discriminant D , but not the parity of the spin invariant. Furthermore, the classification does not depend on whether the Teichmüller curve induced by X is arithmetic or not.

A natural follow-up question is which elements of the Veech group are required to generate $G(X)$. As a corollary of our proof of **Theorem 1.1** we obtain the following:

Corollary 1.2. *For any discriminant $D > 4$ with $D \notin \{9, 33\}$, the group $G(X)$ is generated by the action of two parabolic elements in $\mathrm{SL}(X)$. If $D \in \{9, 33\}$, nevertheless, the minimal number of parabolic elements that are needed to generate the group $G(X)$ is three.*

Weierstrass Prym eigenforms in genus three. The classification of Veech surfaces in $\mathcal{H}(2)$ relies on two fundamental features of this stratum:

- (1) every translation surface (X, ω) admits an involution ι whose (topological) quotient reduces the genus by two; and
- (2) such involution satisfies $\iota^*\omega = -\omega$.

It is possible to find “moral analogues” of $\mathcal{H}(2)$ in higher genera sharing these features. For genus $g > 2$, these spaces are proper subvarieties of strata and are known as *Prym loci*. More precisely, the Prym locus $\mathrm{Prym}(\kappa)$ is the subset of $\mathcal{H}(\kappa)$ of translation surfaces (X, ω) that admit a holomorphic involution (known as a *Prym involution*) $\iota: X \rightarrow X$, whose quotient surface X/ι has genus $g - 2$, and such that $\iota^*\omega = -\omega$. By the Riemann–Hurwitz formula, the Prym involution has exactly $10 - 2g$ fixed points. Thus, Prym loci can only arise up to genus five.

Prym loci contain smaller loci known as *Prym eigenform loci*, introduced by McMullen [McM06], which can be studied in a similar fashion as the classification of Veech surfaces in $\mathcal{H}(2)$. Prym eigenform loci inside $\mathrm{Prym}(2g - 2)$, also known as *Weierstrass*

Prym eigenforms, were completely classified by Lanneau–Nguyen [LN14; LN20]. Indeed, they are distinguished by a *discriminant* D and, for $g = 3$ and some values of D , a moral analogue of the spin invariant. As the classification of Prym eigenform loci in non-minimal strata is not completely understood, we restrict our study to Weierstrass Prym eigenform loci, that is, those inside $\text{Prym}(2g - 2)$. Observe that these loci are empty for $g \geq 5$; the order- $(2g - 2)$ zero is necessarily a fixed point of the Prym involution.

By McMullen’s work [McM07], every Veech surface in a Prym locus is a Prym eigenform. In particular, Veech surfaces in $\text{Prym}(2g - 2)$ are completely classified. By a dimension count, non-Veech surfaces in $\text{Prym}(2g - 2)$ are generic, so a complete classification of all orbit closures inside $\text{Prym}(2g - 2)$ is known.

Main results for genus three. As in the case of Veech surfaces in $\mathcal{H}(2)$, we are interested in the permutations of the regular fixed points for the Prym involution induced by the action of the Veech group. In genus four there is only one such regular fixed point, so the only interesting remaining case is $\text{Prym}(4)$ in genus three. We prove the following result, which is analogous to [Theorem 1.1](#) and [Corollary 1.2](#).

Theorem 1.3. *Let $X \in \text{Prym}(4)$ be a Veech surface of discriminant $D \geq 8$. Let $G(X)$ be the permutation group induced by the action of $\text{SL}(X)$ on the three regular fixed points for the Prym involution on X . Then $G(X)$ is isomorphic to:*

- Sym_2 if D is even and a quadratic residue modulo 16; and
- Sym_3 otherwise.

Moreover, $G(X)$ is generated by the action of one parabolic element in $\text{SL}(X)$ in the former case, and by the action of two parabolic elements in $\text{SL}(X)$ in the latter case.

1.1. Strategy of the proof. McMullen’s classification [McM05] of Teichmüller curves in $\mathcal{H}(2)$ relies on a combinatorial description of two-cylinder cusps, called *splitting prototypes*. This classification gives rise to *prototypical surfaces* on each orbit of a Veech surface. Since, for any Veech surface X , the groups $G(X)$ and $G(g \cdot X)$ are conjugate for every $g \in \text{SL}(2, \mathbb{R})$, it is enough to compute $G(X)$ for prototypical surfaces.

Given a prototypical surface S , we start by computing the action of some parabolic elements of $\text{SL}(S)$ on its regular Weierstrass points. This shows that a specific dihedral group H is contained inside $G(S)$. The cases where S has discriminant 9 or 33 have to be treated separately, as the general argument is insufficient.

Then, we use the geometric restrictions imposed by the locations of the Weierstrass points, and the algebraic restrictions imposed by the coefficients of $\text{SL}(S)$, to show that the inclusion $H \leq G(S)$ is actually an equality. In other words, we show that the reverse containment also holds. In the arithmetic case, these restrictions are those given by the HLK-invariant. In the non-arithmetic case, we are able to extend the HLK-invariant when the discriminant D is a quadratic residue modulo 8 and obtain the desired reverse inclusion in this way. For the remaining case of $D \equiv_8 5$, we show the reverse containment by analysing the same geometric and algebraic restrictions. However, we do *not* find an extension of the HKL-invariant for this latter case.

By combining both containments, we conclude the desired results and prove [Theorem 1.1](#) and [Corollary 1.2](#).

In the case of Weierstrass Prym eigenforms in genus 3, we use the classification by Lanneau–Nguyen [LN14] and follow the same strategy as for genus 2. Several computations and details differ from the genus 2 case. However, the strategy of the proof of [Theorem 1.3](#) is essentially the same.

1.2. Structure of the paper. In [Section 2](#), we recall McMullen’s classification of Teichmüller curves in $\mathcal{H}(2)$, and introduce splitting prototypes and prototypical surfaces. [Section 3](#) is devoted to the study of the action of parabolic elements on regular Weierstrass points: we compute their action on prototypical surfaces, showing that the permutation group given by action of the Veech group on regular Weierstrass points contains a subgroup isomorphic to an appropriate dihedral group. In [Section 4](#), we compute the corresponding reverse inclusions; this section is divided into several subsections. Indeed, in [Section 4.1](#) we study the arithmetic case and compute the distribution of regular Weierstrass points over 2-torsion in the unit torus, giving the corresponding restrictions for the action on regular Weierstrass points. In [Section 4.2](#), we study the non-arithmetic case; the case of quadratic residues modulo 8 is treated in [Section 4.2.1](#) and, in [Section 4.2.2](#), we study the case of $D \equiv_8 5$.

For some small discriminants the general arguments are insufficient; we treat these exceptional cases separately in [Section 5](#). Finally, in [Section 6](#), we summarise our results and give a proof of [Theorem 1.1](#) and [Corollary 1.2](#).

The case of Weierstrass Prym eigenforms in genus 3 is studied in [Section 7](#) following the same structure. In [Section 7.1](#), we recall Lanneau–Nguyen’s classification and introduce splitting prototypes and prototypical surfaces. In [Section 8](#) we study the action of parabolic elements on prototypical surfaces. In [Section 9](#), we compute the reverse inclusion; the arithmetic case is treated in [Section 9.1](#) and the non-arithmetic case, in [Section 9.2](#). Some small discriminants are treated separately in [Section 10](#). Finally, in [Section 11](#), we collect our results giving a proof of [Theorem 1.3](#).

Acknowledgements. We are grateful to Vincent Delecroix and Carlos Matheus for their interesting comments, which motivated some parts of this work.

2. TEICHMÜLLER CURVES IN $\mathcal{H}(2)$

In this background section we recall McMullen’s classification of Teichmüller curves in $\mathcal{H}(2)$ and *splitting prototypes*, which describe two-cylinder cusps.

2.1. Classification of Teichmüller curves in $\mathcal{H}(2)$. Fix a Riemann surface X . Recall that $H^{1,0}(X)$ is the complex vector space of Abelian differentials on X . As a real vector space, $H^{1,0}(X)$ is canonically isomorphic to $H^1(X; \mathbb{R})$ via the isomorphism $\omega \mapsto \operatorname{Re}(\omega)$. This isomorphism also endows $H^{1,0}(X)$ with a symplectic form by pulling back the one on $H^1(X; \mathbb{R})$.

The *Jacobian* of X , denoted by $\operatorname{Jac}(X)$, is the complex torus $H^{1,0}(X)^*/H_1(X; \mathbb{Z})$ endowed with the symplectic form on $T_0 \operatorname{Jac}(X) = H^{1,0}(X)^*$ induced by the symplectic form on $H^{1,0}(X)$. An *endomorphism* of $\operatorname{Jac}(X)$ is an endomorphism of the complex torus. By the isomorphism $H^{1,0}(X) \cong H^1(X; \mathbb{R})$, we can think of an endomorphism of $\operatorname{Jac}(X)$ as an endomorphism of $H_1(X; \mathbb{Z})$ whose real-linear extension to $H^{1,0}(X)^*$ is complex-linear. An endomorphism of $\operatorname{Jac}(X)$ is said to be *self-adjoint* if it is self-adjoint with respect to the symplectic form on $T_0 \operatorname{Jac}(X)$.

Fix an integer $D > 0$ satisfying $D \equiv_4 0$ or $D \equiv_4 1$. Such D is known as a *real quadratic discriminant*. Let \mathcal{O}_D be the *real quadratic order* of discriminant D , that is,

$$\mathcal{O}_D = \mathbb{Z}[\lambda] \cong \mathbb{Z}[T]/(T^2 - eT - bc),$$

where b, c and e are integers satisfying $D = e^2 + 4bc$ and $\lambda = \frac{e + \sqrt{D}}{2}$. Note that if D is odd and square-free, or if D is even and $D/4$ is square-free, then \mathcal{O}_D is the *maximal order* inside $\mathbb{Q}(\sqrt{D})$ and coincides with its ring of integers.

We say that an Abelian variety A admits real multiplication by \mathbb{O}_D if $\dim_{\mathbb{C}} A = 2$ and \mathbb{O}_D occurs as an indivisible, self-adjoint subring of $\text{End}(A)$.

With these definitions in mind, we can now define the loci in which we are interested:

Definition 2.1. The *Weierstrass curve* W_D of discriminant D is the locus of Riemann surfaces $X \in \mathcal{M}_2$ such that:

- $\text{Jac}(X)$ admits real multiplication by \mathbb{O}_D ; and
- there exists an Abelian differential ω on X such that:
 - $(X, \omega) \in \mathcal{H}(2)$; and
 - ω is an eigenform, that is, $\mathbb{O}_D \cdot \omega \subseteq \mathbb{C} \cdot \omega$.

By the work of McMullen [McM05] (see also the articles by Calta [Cal04] and Hubert-Lelièvre [HL06]), we have the following:

Theorem 2.2. *Every irreducible component of a Weierstrass curve is a Teichmüller curve. Conversely, any Teichmüller curve coming from a translation surface in $\mathcal{H}(2)$ is an irreducible component of some Weierstrass curve.*

Let D be a real quadratic discriminant. If $D \leq 4$, then the Weierstrass curve W_D is empty. On the other hand, if $D \geq 5$, then W_D is irreducible if $D \not\equiv_8 1$ or if $D = 9$ and, otherwise, W_D can be decomposed as $W_D^0 \sqcup W_D^1$, where W_D^0 and W_D^1 are irreducible and distinguished by the spin invariant.

For the definition of the spin invariant, we refer the reader to McMullen's original work [McM05, Section 5].

Given a real quadratic discriminant D , the Weierstrass curve W_D is a locus in the moduli space of Riemann surfaces. As we will work with *translation* surfaces, we define ΩW_D as the locus of translation surfaces in $\mathcal{H}(2)$ projecting down onto W_D . We have that each ΩW_D consists entirely on Veech surfaces. We similarly define ΩW_D^ε for $\varepsilon \in \{0, 1\}$ when $D > 9$ and $D \equiv_8 1$.

2.2. Prototypes. Following McMullen's classification [McM05], we say that a quadruple of integers (a, b, c, e) is a *splitting prototype* of discriminant D , if

$$\begin{aligned} D &= e^2 + 4bc, & 0 \leq a < \gcd(b, c), & \quad c + e < b, \\ & & 0 < b, c & \quad \text{and} \quad \gcd(a, b, c, e) = 1. \end{aligned}$$

With each splitting prototype (a, b, c, e) , we associate a translation surface as in [Figure 1](#), where $\lambda = \frac{e + \sqrt{D}}{2}$. We call such surface the *prototypical surface* associated with the prototype (a, b, c, e) and denote it by $P(a, b, c, e) \in \Omega W_D$.

Splitting prototypes are in a one-to-one correspondence with two-cylinder cusps in W_D [McM05, Theorem 4.1] (see [Section 3.1](#) for more details on cylinder decompositions). In particular, any two cylinder decomposition of a translation surface in ΩW_D corresponds to a splitting prototype of discriminant D and can be turned into a surface as in [Figure 1](#), up to a rotation (that maps the two-cylinder direction to the horizontal direction) and the action of a diagonal matrix in $\text{GL}^+(2, \mathbb{R})$ (in order to match the side lengths). We use this fact in [Section 3](#) to understand how parabolic elements induced by two-cylinder decompositions act on Weierstrass points.

2.2.1. The spin invariant. The spin invariant can be used to distinguish the two components of ΩW_D when $D > 9$ and $D \equiv_8 1$. McMullen showed [McM05, Theorem 5.3] that the spin invariant of a splitting prototype (a, b, c, e) of discriminant $D \equiv_8 1$ is given




FIGURE 1. Prototypical surface $P(a, b, c, e)$, with $\lambda = \frac{e+\sqrt{D}}{2}$. This example corresponds to $P(2, 4, 4, -3)$ for $D = 73$.

by the parity of

$$(1) \quad \frac{e-f}{2} + (c+1)(a+b+ab),$$

where $f > 0$ is the conductor of \mathcal{O}_D , that is, the index of \mathcal{O}_D inside the maximal order of $\mathcal{O}_D \otimes \mathbb{Q} \cong \mathbb{Q}(\sqrt{D})$. In particular, when D is odd, we have $D = Ef^2$ with E square-free.

2.2.2. Reduced prototypes. A splitting prototype that has the form

$$(a, b, c, e) = \left(0, \frac{D-e^2}{4}, 1, e\right),$$

that is, such that $c = 1$, is called a *reduced prototype* [McM05, Sections 8 and 10]. Given a real quadratic discriminant $D > 4$, reduced prototypes are parametrised only by e , where $e \in \mathbb{Z}$ satisfies

$$e \equiv_2 D, \quad e^2 < D \quad \text{and} \quad (e+2)^2 < D.$$

Observe that the last two inequalities are equivalent to $0 < \lambda < b$, where $\lambda = \frac{e+\sqrt{D}}{2}$, and can also be restated as $-\sqrt{D} < e < \sqrt{D} - 2$. Moreover, when $D \equiv_8 1$ and $D > 9$, Equation (1) shows that the spin of the prototypical surface $P\left(0, \frac{D-e^2}{4}, 1, e\right)$ is given by the parity of $\frac{e-f}{2}$, where f is the conductor of \mathcal{O}_D . Thus, it is equal to 0 if $e \equiv_4 f$, and it is equal to 1 if $e \not\equiv_4 f$.

We denote the prototypical surface associated with the reduced prototype e by $P_D(e)$, that is, $P_D(e) = P\left(0, \frac{D-e^2}{4}, 1, e\right) \in \Omega W_D$, as in the left of Figure 2.

Let R_D be the set of e -parameters of reduced prototypes of discriminant D and, for D satisfying $D \equiv_8 1$ and $D > 9$, let R_D^ε be the set of e -parameters of reduced prototypes corresponding to surfaces in ΩW_D^ε , for $\varepsilon \in \mathbb{Z}/2\mathbb{Z}$. That is, for any discriminant $D > 4$, let

$$R_D = \{e \in \mathbb{Z} \mid e \equiv_2 D, e^2 < D, (e+2)^2 < D\},$$

and, for $D \equiv_8 1$ and $D > 9$, let

$$R_D^0 = \{e \in R_D \mid e \equiv_4 f\} \quad \text{and} \quad R_D^1 = \{e \in R_D \mid e \not\equiv_4 f\},$$




FIGURE 2. Prototypical surface $P_D(e)$ (left) and L-shaped surface $L_D(e)$ (right), for a reduced prototype $e \in R_D$. This example shows $P_{73}(1)$ and $L_{73}(1)$.

where f is the conductor of \mathcal{O}_D , that is, the largest positive integer whose square divides D . Since, for fixed D , there is a one-to-one correspondence between reduced prototypes and their e -parameters, we will refer to $e \in R_D$ as a reduced prototype.

2.2.3. L-shaped representatives. We also use, in Section 4, an L-shaped version of the surface associated with a reduced prototype. This surface is obtained by the action of the diagonal matrix $\begin{pmatrix} \lambda^{-1} & 0 \\ 0 & 1 \end{pmatrix} \in \mathrm{GL}^+(2, \mathbb{R})$ on $P_D(e)$, with $e \in R_D$, as in Figure 2. We denote such surface by $L_D(e) \in \Omega\mathcal{W}_D$.

3. ACTION OF PARABOLIC ELEMENTS IN GENUS TWO

In this section, we show that the permutation group induced by the action of *parabolic* elements of the Veech group contains a dihedral group. More precisely, using the reduced prototypes $P_D(e)$, we compute a subgroup of $G_D(e) = G(P_D(e))$, generated by the action of certain parabolic elements in the Veech group of $P_D(e)$.

3.1. Parabolic elements and cylinder decompositions. In what follows, we assume that a translation surface (X, ω) is endowed with the flat metric coming from ω . If (X, ω) is viewed as a set of polygons in \mathbb{C} with side identifications, then a geodesic that does not contain any singularities in its interior is simply a straight line that respects said side identifications.

A *cylinder of circumference* $w > 0$ and *height* $h > 0$ on X is a maximal open annulus filled by isotopic simple closed regular geodesics, isometric to $\mathbb{R}/w\mathbb{Z} \times (0, h)$. Its *modulus* is the ratio between its circumference w and its height h , that is, $m = \frac{w}{h} > 0$. The *core curve* of a cylinder is the simple closed geodesic identified with $\mathbb{R}/w\mathbb{Z} \times \{h/2\}$. A *saddle connection* is a geodesic joining two different singularities or a singularity to itself, with no singularities in its interior. Cylinders are bounded by parallel (concatenations of) saddle connections.

A *cylinder decomposition* of X is a collection of parallel cylinders whose closures cover X . Given $\theta \in [0, 2\pi)$, we say that a cylinder decomposition is *in direction* θ if the angle between the horizontal direction and the core curve of any cylinder (or, equivalently, all cylinders) is θ .

By the work of Veech [Vee89], cylinder decompositions and parabolic elements are closely related. Indeed, fix an angle $\theta \in [0, 2\pi)$ and let R_θ be the matrix rotating the plane (counterclockwise) by θ . Given $t \in \mathbb{R}$, let $T_t = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}$ be a matrix twisting the horizontal direction by t . Then, a matrix of the form $P_\theta^t = R_\theta T_t R_\theta^{-1}$ belongs to $\mathrm{SL}(X)$ if and only if there exists a cylinder decomposition $\{C_i\}_{i \in I}$ of X in direction θ such that

the moduli m_i of the cylinders C_i are pairwise rationally related, that is, if $r_{ij} = \frac{m_i}{m_j} \in \mathbb{Q}$ for every $i, j \in I$. Furthermore, if $t \in \mathbb{R}$ is an integer multiple of the modulus of each cylinder, that is, for each $i \in I$ there exist $k_i \in \mathbb{Z}$ such that $t = k_i m_i$, then $P_\theta^t \in \text{SL}(X)$. The corresponding affine transformation preserves the cylinder decomposition, as it twists each cylinder C_i exactly k_i times along itself. In particular, P_θ^t acts on each core curve γ_i by a rotation by angle $k_i \pi$ (that is, k_i half-turns). If the k_i are minimal and positive, we call this action “the” *twist* in direction θ and we say that θ is a *parabolic direction*.

In the case of Veech surfaces, parabolic directions are easily characterised by the so-called *Veech dichotomy* [Vee89]. Indeed, a direction $\theta \in [0, 2\pi)$ is parabolic if and only if there exists a saddle connection or a closed geodesic in direction θ , meaning that the angle between the horizontal and the curve is θ .

3.2. Cylinder decompositions in $\mathcal{H}(2)$. Let $X \in \mathcal{H}(2)$. It is known that there are at most two cylinders in a cylinder decomposition of X . Moreover, the core curve of each cylinder contains exactly two Weierstrass points [Par20, Remark 5]. In particular, a parabolic element $P_\theta^t \in \text{SL}(X)$ exchanges two (different) regular Weierstrass points $x, y \in X$ if and only if

- (1) X admits a cylinder decomposition $\{C_i\}_i$ in direction θ ;
- (2) x and y lie on the same core curve γ_i of one of the cylinders C_i ; and
- (3) the corresponding integer k_i of half-turns along γ_i is odd (recall that $t = k_i m_i$).

Let $\theta \in [0, 2\pi)$ and assume that $\{C_1, C_2\}$ is a two-cylinder decomposition of X in direction θ . We can write

$$r = \frac{m_1}{m_2} = \frac{p}{q} \in \mathbb{Q},$$

where p and q are positive integers with $\gcd(p, q) = 1$. Then, the twist in direction θ is described by taking $k_1 = q$ and $k_2 = p$. In other words, if $t = qm_1 = pm_2$, then the parabolic element P_θ^t belongs to $\text{SL}(X)$, and $t > 0$ is minimal for this property. Since t will be always chosen in this way, we can omit it from the notation and define $r_\theta = \frac{m_1}{m_2} = \frac{p}{q}$.

By the previous discussion, we can compute the action of P_θ^t at the level of (regular) Weierstrass points, which we denote by $\tau_\theta \in \text{Sym}(\{w_1, \dots, w_5\})$. Indeed, τ_θ exchanges the two regular Weierstrass points in γ_i , the core curve of C_i , if and only if k_i is odd. Since $\gcd(k_1, k_2) = \gcd(q, p) = 1$, τ_θ is either a transposition or a product of two disjoint transpositions.

Notation. In what follows we denote

$$\begin{aligned} \frac{\text{odd}}{\text{odd}} &= \left\{ \frac{p}{q} \in \mathbb{Q} \mid p, q \in \mathbb{N}, \gcd(p, q) = 1, p \equiv_2 q \equiv_2 1 \right\}, \\ \frac{\text{even}}{\text{odd}} &= \left\{ \frac{p}{q} \in \mathbb{Q} \mid p, q \in \mathbb{N}, \gcd(p, q) = 1, p \equiv_2 0, q \equiv_2 1 \right\} \quad \text{and} \\ \frac{\text{odd}}{\text{even}} &= \left\{ \frac{p}{q} \in \mathbb{Q} \mid p, q \in \mathbb{N}, \gcd(p, q) = 1, p \equiv_2 1, q \equiv_2 0 \right\}. \end{aligned}$$

Moreover, for a two-cylinder decomposition of a surface in $\mathcal{H}(2)$, we always choose C_1 to be the cylinder with the largest circumference. We will refer to C_1 as the *long cylinder* and, accordingly, to C_2 as the *short cylinder*.

We can summarise the previous discussions in the form of the following lemma which we will repeatedly use throughout the article.

Lemma 3.1. *Let X be a translation surface in $\mathcal{H}(2)$. Let $\theta \in [0, 2\pi)$ be a parabolic direction of X and assume that there exists a two-cylinder decomposition $\{C_1, C_2\}$ of X in direction $\theta \in [0, 2\pi)$. Recall that $r_\theta = \frac{m_1}{m_2}$, where m_i is the modulus of the cylinder C_i , and that $\tau_\theta \in \text{Sym}(\{w_1, \dots, w_5\})$ is the action of the twist in direction θ on the regular Weierstrass points w_1, \dots, w_5 of X . We have that*

- (1) *if $r_\theta \in \frac{\text{odd}}{\text{odd}}$, then τ_θ is a product of two disjoint transpositions, exchanging the regular Weierstrass points in the core curves of C_1 and C_2 ;*
- (2) *if $r_\theta \in \frac{\text{even}}{\text{odd}}$, then τ_θ only exchanges the two regular Weierstrass points in the core curve of the long cylinder C_1 ; and*
- (3) *if $r_\theta \in \frac{\text{odd}}{\text{even}}$, then τ_θ only exchanges the two regular Weierstrass points in the core curve of the short cylinder C_2 .*

□

3.3. Horizontal and vertical twists on reduced prototypes. Let $D > 0$ be a real quadratic discriminant. Given a splitting prototype (a, b, c, e) of discriminant D , the horizontal direction induces a two-cylinder decomposition $\{C_1, C_2\}$ of $P(a, b, c, e)$, where C_1 is the long cylinder. Referring to [Figure 1](#) we see that the moduli m_i of the cylinders C_i are given by

$$m_1 = \frac{b}{c} \quad \text{and} \quad m_2 = \frac{\lambda}{\lambda} = 1.$$

The ratio of the moduli is then

$$r_h = r_0 = \frac{m_1}{m_2} = \frac{b}{c} = \frac{D - e^2}{4c^2},$$

where the label “h” stands for *horizontal*.

Consider now the case of a reduced prototype $e \in R_D$. In this case, we have that $r_h = b = \frac{D - e^2}{4}$. Moreover, the vertical direction induces a two-cylinder decomposition $\{C_1, C_2\}$ of $P_D(e)$, where C_1 is the long cylinder. From [Figure 2](#) (left) we see that the corresponding moduli are

$$m_1 = \frac{\lambda + 1}{\lambda} \quad \text{and} \quad m_2 = \frac{1}{b - \lambda} = \frac{1}{\frac{D - e^2}{4} - \lambda}.$$

Since $\lambda = \frac{e + \sqrt{D}}{2}$, it follows from an elementary computation that the ratio of the moduli is

$$r_v = r_{\pi/2} = \frac{m_1}{m_2} = \frac{D - (e + 2)^2}{4} = b - e - 1,$$

where the label “v” stands for *vertical*. In particular, we obtain that $r_v = r_h - e - 1$.

We can now summarise this discussion for reduced prototypes:

Lemma 3.2. *Let $D > 4$ be a real quadratic discriminant. Let $e \in R_D$ be a reduced prototype and let $b = \frac{D - e^2}{4} \in \mathbb{N}$. Let $X = P_D(e)$ be the prototypical surface associated with the reduced prototype $(0, b, 1, e)$ and let w_1, \dots, w_5 be its regular Weierstrass points. We set $\tau_h = \tau_0$ and $\tau_v = \tau_{\pi/2}$, that is, τ_h and τ_v are the permutations in $\text{Sym}(\{w_1, \dots, w_5\})$ associated with the horizontal and vertical twists, respectively. Then, we have that the permutations τ_h and τ_v always exchange the regular Weierstrass points in the long horizontal and long vertical cylinders, respectively. Moreover, τ_h exchanges the regular Weierstrass points in the short horizontal cylinder if and only if b is odd. Similarly, τ_v exchanges the regular Weierstrass points in the short vertical cylinder if and only if $b - e - 1$ is odd.*

Proof. From the previous discussion, we have that $r_h = b$ and $r_v = b - e - 1$. Thus, r_h and r_v are integers. In particular, they belong to $\frac{\text{odd}}{\text{odd}} \sqcup \frac{\text{even}}{\text{odd}}$. By [Lemma 3.1](#), the horizontal and vertical twists always exchange the regular Weierstrass points in the corresponding long cylinders. Finally, [Lemma 3.1](#) also implies that whether they exchange the regular Weierstrass points in the corresponding short cylinders only depends on the parities of b and $b - e - 1$. \square

Remark 3.3. We have refrained from introducing the notion of *butterfly moves* to simplify the exposition. In fact, several cylinder decompositions on a prototypical surface can be thought of as a butterfly move on splitting prototypes (see McMullen's work [[McM05](#), Section 7]). For the case of reduced prototypes, the vertical cylinder decomposition corresponds to the butterfly move B_∞ , which maps the prototypical surface $P_D(e)$ into the prototypical surface $P_D(-e - 4)$ [[McM05](#), Theorem 7.5]. Thus, the fact that the action induced by the vertical twist depends on the parity of $\frac{D-(e+4)^2}{4}$ can also be explained in this way.

In what follows, given a prototypical surface, we fix w_1, \dots, w_5 as in [Figure 1](#) and identify $\text{Sym}(\{w_1, \dots, w_5\})$ with Sym_5 in the natural way. Thus, for example, the transposition $(i j) \in \text{Sym}_5$ corresponds to $(w_i w_j) \in \text{Sym}(\{w_1, \dots, w_5\})$.

Proposition 3.4. *Let $D > 4$ be a real quadratic discriminant with $D \equiv_4 0$ and $X \in \Omega W_D$. Then, $G(X)$ has a subgroup conjugate to Dih_4 generated by the action of two parabolic elements in $\text{SL}(X)$.*

Proof. Let $e \in R_D$. Since W_D is irreducible, there is an element in $\text{GL}^+(2, \mathbb{R})$ that maps X to $P_D(e)$. In particular $G(X)$ is conjugate to $G_D(e) = G(P_D(e))$, and the parabolic elements of $\text{SL}(X)$ are conjugate to the parabolic elements of $\text{SL}(P_D(e))$. Thus, it is enough to prove the result for $G_D(e)$.

By definition of R_D , we have that $e \equiv_2 D$, so e is even. Therefore, $r_h \not\equiv_2 r_h - e - 1 = r_v$. In the case that $D \equiv_8 e^2$, $r_h = b = \frac{D-e^2}{4}$ is even and r_v is odd. Thus, by [Lemma 3.2](#), the horizontal twist induces the permutation $\tau_h = (1\ 2)$, and the vertical twist induces the permutation $\tau_v = (1\ 3)(2\ 4)$. It follows that

$$\langle \tau_h, \tau_v \rangle = \text{Dih} \left(\begin{array}{ccc} & 1 & \\ 4 & \swarrow & \searrow & 3 \\ & 2 & \end{array} \right) \leq G_D(e).$$

On the other hand, if $D \not\equiv_8 e^2$, $r_h = b = \frac{D-e^2}{4}$ is odd and r_v is even. Thus, by [Lemma 3.2](#), the horizontal twist induces the permutation $\tau_h = (1\ 2)(3\ 5)$, and the vertical twist induces the permutation $\tau_v = (1\ 3)$. It follows that

$$\langle \tau_h, \tau_v \rangle = \text{Dih} \left(\begin{array}{ccc} & 1 & \\ 5 & \swarrow & \searrow & 2 \\ & 3 & \end{array} \right) \leq G_D(e).$$

\square

Proposition 3.5. *Let $D > 4$ be a real quadratic discriminant with $D \equiv_8 5$ and $X \in \Omega W_D$. Then, $G(X)$ has a subgroup conjugate to Dih_5 generated by the action of two parabolic elements in $\text{SL}(X)$.*

Proof. Let $e \in R_D$. Since W_D is irreducible, there is an element in $\text{GL}^+(2, \mathbb{R})$ that maps X to $P_D(e)$. In particular $G(X)$ is conjugate to $G_D(e) = G(P_D(e))$, and the parabolic

elements of $\mathrm{SL}(X)$ are conjugate to the parabolic elements of $\mathrm{SL}(\mathbb{P}_D(e))$. Thus, it is enough to prove the result for $G_D(e)$.

By definition of R_D , we have that $e \equiv_2 D$, so e is odd. Therefore, $r_h \equiv_2 r_h - e - 1 = r_v$ and $e^2 \equiv_8 1$. Hence, $r_h = b = \frac{D-e^2}{4}$ is odd. Thus, by [Lemma 3.2](#), the horizontal twist induces the permutation $\tau_h = (1\ 2)(3\ 5)$, and the vertical twist induces the permutation $\tau_v = (1\ 3)(2\ 4)$. It follows that

$$\langle \tau_h, \tau_v \rangle = \mathrm{Dih} \left(\begin{array}{ccc} & 1 & \\ 3 & & 2 \\ & 4 & 5 \end{array} \right) \leq G_D(e).$$

□

Proposition 3.6. *Let $D > 4$ be a real quadratic discriminant with $D \equiv_8 1$ and $X \in \Omega W_D$. Then, $G(X)$ has a subgroup conjugate to Sym_3 generated by the action of two parabolic elements in $\mathrm{SL}(X)$.*

Proof. In this case $W_D = W_D^0 \sqcup W_D^1$. Let $\varepsilon \in \mathbb{Z}/2\mathbb{Z}$ be the corresponding spin, that is, such that $X \in \Omega W_D^\varepsilon$. Let $e \in R_D^\varepsilon$. Since W_D^ε is irreducible, there is an element in $\mathrm{GL}^+(2, \mathbb{R})$ that maps X to $\mathbb{P}_D(e)$. In particular $G(X)$ is conjugate to $G_D(e) = G(\mathbb{P}_D(e))$, and, similarly, the parabolic elements of $\mathrm{SL}(X)$ are conjugate to the parabolic elements of $\mathrm{SL}(\mathbb{P}_D(e))$. Thus, it is enough to prove the result for $G_D(e)$.

By definition of R_D , we have that $e \equiv_2 D$, so e is odd. Therefore, $r_h \equiv_2 r_h - e - 1 = r_v$ and $e^2 \equiv_8 1$. Hence, $r_h = b = \frac{D-e^2}{4}$ is even. Thus, by [Lemma 3.2](#), the horizontal twist induces the permutation $\tau_h = (1\ 2)$, and the vertical twist induces the permutation $\tau_v = (1\ 3)$, regardless of the spin. It follows that

$$\langle \tau_h, \tau_v \rangle = \mathrm{Sym}(\{1, 2, 3\}) \leq G_D(e).$$

□

Proposition 3.7. *Let $D > 4$ be a real quadratic discriminant with $D \equiv_8 1$ and $D \notin \{9, 33\}$. Let $X \in \Omega W_D$. Then, $\mathrm{SL}(X)$ contains a parabolic element that induces a product of two disjoint transpositions inside $G(X)$.*

Proof. Let D be as in the statement of the proposition. Observe that, since $D \neq 9$ and $D \equiv_8 1$, W_D is reducible and $W_D = W_D^0 \sqcup W_D^1$. It is enough to find a prototypical surface for each spin such that its horizontal twist induces a product of two disjoint transpositions. Consider the choices $b = 4k + 2$, $c = 2$ and $e \in \{-1, -3, -5, -7\}$. Observe that, with these choices, we have that

$$D = e^2 + 4bc = e^2 + 16 + 32k = \begin{cases} 17 + 32k & \text{if } e = -1 \\ 25 + 32k & \text{if } e = -3 \\ 41 + 32k & \text{if } e = -5 \\ 65 + 32k & \text{if } e = -7, \end{cases}$$

where $k \geq 0$. Then, both $(0, b, c, e)$ and $(1, b, c, e)$ are splitting prototypes. Indeed, $a < \gcd(b, c) = 2$ if $a \in \{0, 1\}$, $c + e \leq 1 < 2 \leq b$, and $\gcd(a, b, c, e) = 1$ since $c = 2$ and e is odd. Moreover, since b and c are even, [Equation \(1\)](#) shows that these two prototypes correspond to different spins.

Finally, since $r_0 = \frac{b}{c} = 2k + 1$, we have $r_0 \in \frac{\text{odd}}{\text{odd}}$. Thus, by [Lemma 3.1](#), $G(X)$ contains an element that is the product of two disjoint transpositions. □

Remark 3.8. If $D \in \{9, 33\}$, an exhaustive search shows that no prototype satisfies that the ratio of moduli r_0 belongs to $\frac{\text{odd}}{\text{odd}}$ (for both spins in the case $D = 33$). In particular, the proof of [Proposition 3.7](#) does not work in these cases. Since any two parabolic directions can be mapped to the horizontal and vertical directions on some prototypical surface, we conclude that, if we only restrict to only two parabolic elements, the largest group that can be generated by the action of these elements on the regular Weierstrass points of a surface in $\Omega\mathcal{W}_D$ is $\text{Sym}_3 \cong \text{Dih}_3$. However, it is still possible to exhibit a *third* explicit parabolic direction on particular representatives that allows us to conclude that $G(X) \cong \text{Sym}_3 \times \text{Sym}_2 \cong \text{Dih}_6$. See [Sections 5.1](#) and [5.2](#) below.

4. RESTRICTIONS FOR PERMUTATIONS IN GENUS TWO

In this section, we constrain the group of permutations of Weierstrass points. To this end, we use geometric restrictions on the location of the Weierstrass points, and algebraic restrictions on the coefficients of the elements of the Veech group.

Since these restrictions differ in the arithmetic case (when D is a perfect square) and the non-arithmetic case, we separate the analysis in different subsections. We start with the arithmetic case, which is more straightforward.

4.1. Arithmetic case. In this section, we analyse the case where $D \geq 9$ is a perfect square, say $D = d^2$ for $d \geq 3$ an integer. Observe first that if D is odd (or, equivalently, if d is odd), then $D \equiv_8 1$. Thus, the case $D \equiv_8 5$ does not arise.

The Weierstrass locus $\Omega\mathcal{W}_{d^2}$ consists of $\text{GL}^+(2, \mathbb{R})$ orbits of square-tiled surfaces, which we will properly define in what follows. A (primitive) *square-tiled surface* is a translation surface (X, ω) such that the set of periods of ω is the lattice $\mathbb{Z}[i]$ in \mathbb{C} . That is, the integration of ω along any saddle connection produces an element of $\mathbb{Z}[i]$. For such a surface, we obtain again by integrating ω a holomorphic map

$$p: X \rightarrow E = \mathbb{C}/\mathbb{Z}[i],$$

which can be normalised so it is branched only over $z = 0$.

If we assume that $(X, \omega) \in \mathcal{H}(2)$ is a square-tiled surface, we have, by McMullen's work [[McM05](#), Section 6], that $(X, \omega) \in \Omega\mathcal{W}_{d^2}$ for some d . Moreover, the degree of the covering map p is equal to d , and the d preimages of the unit-square fundamental domain for E provide the square tiling of X .

A Weierstrass point $w \in X$ can only project down to a 2-torsion point in E . Clearly, modulo the lattice $\mathbb{Z}[i]$, the 2-torsion points in E are $0, h = \frac{1}{2}, v = \frac{i}{2}$ and $c = \frac{1}{2} + \frac{i}{2}$. We say that the Weierstrass point $w \in X$ is of type $0, h, v$ or c , accordingly. In addition, a Weierstrass point $w \in X$ is called *integral* if $p(w) = 0 \pmod{\mathbb{Z}[i]}$, that is, if w is of type 0 . To conclude the proof of [Theorem 1.1](#) for the arithmetic case, we use these geometric constraints imposed by the projection of the regular Weierstrass points of X . In fact, if an affine transformation on X maps a point of a certain type t_0 to another point of type t_1 , then it has to map *every* point of type t_0 to a point of type t_1 , since it induces a well-defined quotient map on E . Moreover, the set of integral points must be preserved. Recall that the HLK-invariant is defined as the number of integral points together with the (unordered) list of points of type h, v and c . We will use square brackets to denote unordered lists. The restrictions of the previous paragraph are precisely equivalent to the fact that the HLK-invariant is preserved.

For $e \in R_D$, it is not difficult to see that the L-shaped representatives $L_D(e)$ in [Figure 2](#) are in fact square-tiled surfaces when $D = d^2$ is a perfect square. Indeed, we have that $\lambda = \frac{e+d}{2} \in \mathbb{N}$, so these surfaces consist on $\lambda + 1$ unit squares stacked upwards, and

$\lambda - e$ unit squares stacked rightwards. See [Figures 3 and 4](#). In particular, the type of $w_1, \dots, w_5 \in L_D(e)$ depends only on the parity of λ and $\lambda - e - 1$. More precisely, we have that

- w_1 is always of type c ;
- w_2 is of type c whenever $\lambda - e - 1$ is odd, and of type v otherwise;
- w_3 is of type c whenever λ is odd, and of type h otherwise;
- w_4 is of type h whenever $\lambda - e - 1$ is odd, and integral otherwise; and
- w_5 is of type v whenever λ is odd, and integral otherwise.

Furthermore, recall that these L-shaped surfaces are essentially prototypical surfaces for a reduced prototype (up to a scaling). This allows us to obtain the following simple lemma:

Lemma 4.1. *Let $d > 2$ be an odd integer and let $e \in R_D$. Set $\lambda = \frac{e+d}{2}$. Then, $L_{d^2}(e) \in \Omega W_{d^2}^0$ if λ is odd, and $L_{d^2}(e) \in \Omega W_{d^2}^1$ if λ is even.*

Proof. Fix an odd integer $d > 2$ and let $e \in R_{d^2}$. Recall that $L_{d^2}(e)$ is a scaling of the reduced prototype $\left(0, \frac{d^2-e^2}{4}, 1, e\right)$. Thus, its spin coincides with the spin of the prototypical surface $P\left(0, \frac{d^2-e^2}{4}, 1, e\right)$.

Since $D = d^2$, the conductor f of D is equal to d . By [Equation \(1\)](#), we see that the spin of this prototypical surface is equal to the parity of $\frac{e-d}{2}$. Since d is odd, the spin is also equal to the parity of $\lambda + 1 = \frac{e+d}{2} + 1$, which gives the desired conclusion.

This lemma also follows directly from McMullen's work [[McM05](#), Theorem 6.1] by counting the number of integral points in each case. \square

Proposition 4.2. *Let $d > 2$ be an even integer and let $X \in \Omega W_{d^2}$. Then, $G(X)$ is contained in a group conjugate to Dih_4 .*

Proof. Fix an even integer $d > 2$. Let $e \in R_D$. When $d^2 \equiv_8 e^2$, we have that λ is even and $\lambda - e - 1$ is odd. Thus, w_1, w_2 are of type c , w_3, w_4 are of type h and w_5 is integral. Thus, the HLK-invariant is $(1, [2, 2, 0])$. See [Figure 3](#) (left). It follows that

$$G_{d^2}(e) \leq \text{Dih} \left(\begin{array}{ccc} & 1 & \\ 4 & \diagdown & \diagup \\ & 2 & 3 \end{array} \right).$$

When $d^2 \not\equiv_8 e^2$, we have that λ is even and $\lambda - e - 1$ is odd. Thus, w_1, w_3 are of type c , w_2, w_5 are of type v and w_4 is integral. Thus, the HLK-invariant is $(1, [2, 2, 0])$. See [Figure 3](#) (right). It follows that

$$G_{d^2}(e) \leq \text{Dih} \left(\begin{array}{ccc} & 1 & \\ 5 & \diagdown & \diagup \\ & 3 & 2 \end{array} \right).$$

\square

Proposition 4.3. *Let $d > 2$ be an odd integer and let $X \in \Omega W_{d^2}$. Then, $G(X)$ is contained in a group isomorphic to $\text{Sym}_3 \times \text{Sym}_2 \cong \text{Dih}_6$.*

Proof. Fix an odd integer $d > 2$. Let $e \in R_D$. When λ is odd, [Lemma 4.1](#) shows that $L_{d^2}(e) \in \Omega W_{d^2}^0$. In this case, we have no integral regular Weierstrass points, as in




FIGURE 3. The L-shaped square-tiled surface $L_{d^2}(e) \in \Omega W_{d^2}$, when d is even. The case $d^2 \equiv_8 e^2$ (left) and $d^2 \not\equiv_8 e^2$ (right). These examples show $L_{12^2}(0)$ and $L_{10^2}(0)$, respectively.




FIGURE 4. The L-shaped square-tiled surface $L_{d^2}(e) \in \Omega W_{d^2}$, when d is odd. The case $W_{d^2}^0$ (left) and $W_{d^2}^1$ (right). These examples show $L_{13^2}(1)$ and $L_{13^2}(-1)$, respectively.

Figure 4 (left). Moreover, $\lambda - e - 1$ is even and therefore, w_1, w_2, w_3 are of type c , w_4 is of type h and w_5 , of type v . Thus, the HLK-invariant is $(0, [3, 1, 1])$. It follows that

$$G_{d^2}(e) \leq \text{Sym}(\{1, 2, 3\}) \times \text{Sym}(\{4, 5\}).$$

Observe that the previous situation is the only one that can arise for the particular case of $d = 3$, as $R_{3^2} = \{-1\}$ (and, hence, $\lambda = 1$).

When λ is even, Lemma 4.1 shows that $L_{d^2}(e) \in \Omega W_{d^2}^1$. In this case, we have two integral regular Weierstrass points, namely w_4 and w_5 , as in Figure 4 (right). Moreover, $\lambda - e - 1$ is odd and therefore, w_1, w_2 and w_3 are of type c, v and h , respectively. Thus, the HLK-invariant is $(2, [1, 1, 1])$. It follows that

$$G_{d^2}(e) \leq \text{Sym}(\{1, 2, 3\}) \times \text{Sym}(\{4, 5\}).$$

□

4.2. Non-arithmetic case. In this section, we analyse the case where $D > 9$ is a real quadratic discriminant that is not a perfect square. We first show that if D is a quadratic residue modulo 8, then the problem can be treated completely analogously to the arithmetic case. In the case where D is a quadratic nonresidue modulo 8, that is,

when $D \equiv_8 5$, we show that the action of the affine group on the set of regular Weierstrass points can be restricted by using similar ideas, although it cannot be reduced to the arithmetic case.

Lemma 4.4. *Let $D > 4$ be a real quadratic discriminant. Let $e \in R_D$ be any reduced prototype. Then, $\mathrm{SL}(L_D(e)) \leq \mathrm{SL}(2, \mathcal{O}_D)$.*

Proof. Fix $A \in \mathrm{SL}(L_D(e))$. We have that A must map every saddle connection into a saddle connection, and, therefore, must preserve the set of holonomy vectors of saddle connections.

In particular, consider the two saddle connections obtained by tracing a ray from the lower-left corner of **Figure 2** (right) and moving rightwards or upwards until the singularity is again met. The holonomy vectors of these saddle connections are $(1, 0)^\top$ and $(0, 1)^\top$, respectively. We conclude that both columns of A are holonomy vectors of saddle connections.

Finally, observe that the holonomy vector of any saddle connection must belong to $\mathcal{O}_D \times \mathcal{O}_D$, since every side length and translation vector of $L_D(e)$ belongs to \mathcal{O}_D . \square

In what follows, we are interested in the relative position of the regular Weierstrass points in $L_D(e)$, with respect to the singularity. In particular, we restrict our attention to elements in $\frac{1}{2}(\mathcal{O}_D \times \mathcal{O}_D)$.

Fix $\rho \in \mathbb{Q}(\sqrt{D})$ such that $\mathcal{O}_D = \mathbb{Z}[\rho]$; the choice of ρ will be made explicit later as needed. Each element x of the field $\mathbb{Q}(\sqrt{D})$ can be written uniquely as $x = p + q\rho$ with $p, q \in \mathbb{Q}$. We say that p is the *rational part* of x (note that this depends on the choice of ρ). Moreover, we define $\mathrm{fr}(x)$ as the fractional part of the rational part of x . That is, $\mathrm{fr}(x) = \{p\} = (p \bmod 1)$.

Notation.

- For $A = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{SL}(2, \mathcal{O}_D)$, we write $\eta = p_\eta + q_\eta \rho \in \mathcal{O}_D$, where $p_\eta, q_\eta \in \mathbb{Z}$, for each $\eta \in \{\alpha, \beta, \gamma, \delta\}$. We also write $P_A = \begin{pmatrix} p_\alpha & p_\beta \\ p_\gamma & p_\delta \end{pmatrix}$ and $Q_A = \begin{pmatrix} q_\alpha & q_\beta \\ q_\gamma & q_\delta \end{pmatrix}$, so $A = P_A + Q_A \rho$.
- For $v \in \frac{1}{2}(\mathcal{O}_D \times \mathcal{O}_D)$, we write $v = \frac{1}{2}(p_1 + q_1 \rho, p_2 + q_2 \rho)$, where $p_i, q_i \in \mathbb{Z}$ for each $i \in \{1, 2\}$. We also write $p_v = (p_1, p_2)$ and $q_v = (q_1, q_2)$, so $v = \frac{1}{2}(p_v + q_v \rho)$.

Then,

$$(2) \quad Av = \frac{1}{2}(P_A + Q_A \rho)(p_v + q_v \rho) = \frac{1}{2}(P_A p_v + Q_A q_v \rho^2 + (P_A q_v + Q_A p_v) \rho)$$

4.2.1. Quadratic residues modulo 8. We will now show that, if the discriminant is a quadratic residue modulo 8, then the constraints on the group of permutations on the regular Weierstrass points are exactly the same as in the arithmetic case. We do this by showing that if we only keep track of rational parts, then there is an analogue of the HKL-invariant that has to be preserved. The next lemma provides the first step in this direction:

Lemma 4.5. *Let $D > 0$ be a real quadratic discriminant and assume that it is a quadratic residue modulo 8, say $D \equiv_8 d^2$ for some $d \in \mathbb{Z}$, and let $A \in \mathrm{SL}(2, \mathcal{O}_D)$. Take $\rho = \frac{\sqrt{D}-d}{2}$. Then, we have $\mathrm{fr}(Av) = \frac{1}{2}P_A p_v \bmod 1$ for any $v \in \frac{1}{2}(\mathcal{O}_D \times \mathcal{O}_D)$. Moreover, P_A is non-singular modulo 2.*

Proof. We have that

$$\rho^2 = \frac{D + d^2 - 2d\sqrt{D}}{4} = \frac{D - d^2}{4} - d\rho$$

By Equation (2), we obtain that the rational part of Av is

$$\frac{1}{2}P_A p_v + \frac{D-d^2}{8}Q_A q_v$$

Since $\frac{D-d^2}{8}$ is an integer, this implies $\text{fr}(Av) = \frac{1}{2}P_A p_v \pmod{1}$.
Now, a direct computation shows that

$$1 = \det(A) = \det(P_A + Q_A \rho) = \det(P_A) + \det(Q_A) \rho^2 + \begin{vmatrix} p_\alpha & p_\beta \\ q_\gamma & q_\delta \end{vmatrix} \rho + \begin{vmatrix} q_\alpha & q_\beta \\ p_\gamma & p_\delta \end{vmatrix} \rho.$$

Taking rational parts, we obtain $\det(P_A) + \frac{D-d^2}{4} \det(Q_A) = 1$, so $\det(P_A) \equiv_2 1$. \square

The previous lemmas show that the cases $D \equiv_8 d^2$ can be treated as the arithmetic case. That is, if we set $\rho = \frac{\sqrt{D-d}}{2}$ and we only keep track of the fractional part of the rational part of the vectors $v_i \in \frac{1}{2}(\mathbb{O}_D \times \mathbb{O}_D)$ associated with the regular Weierstrass points w_i , for $i \in \{1, \dots, 5\}$, on a surface $X \in \mathcal{H}(2)$, then the action of $\text{SL}(X)$ behaves as a linear action on points over the 2-torsion points on a torus. This is exactly the case for the regular Weierstrass points in the arithmetic case (see Section 4.1). As in that case, we say that a regular Weierstrass point w_i is of type 0, h , v or c depending on where $\text{fr}(v_i)$ lies on the unit square. Since P_A is non-singular modulo 2, we conclude that it acts bijectively on the sets of 1-torsion and 2-torsion points of the unit torus, so we may extend the definition of the HLK-invariant to this case. More precisely, we define the HLK-invariant as the following data:

- the number of points of type 0; and
- the (unordered) list of number of points of type h , v and c .

Given $d \in \mathbb{Z}$, a real quadratic discriminant $D > 4$ with $D \equiv_8 d^2$ and $e \in R_D$, it is straightforward to compute $\text{fr}(v_i)$ for each $i \in \{1, \dots, 5\}$ in the surface $L_D(e)$. The following lists summarise these computations.

(1) If $D \equiv_8 0$ or $D \equiv_8 4$, we have two cases:

(a) If $e^2 \equiv_8 D$, the HLK-invariant is $(1, [2, 2, 0])$ since

$$\begin{aligned} \text{fr}(v_1) &= (1/2, 1/2), & \text{fr}(v_2) &= (1/2, 1/2), & \text{fr}(v_3) &= (1/2, 0), \\ \text{fr}(v_4) &= (1/2, 0) & \text{and} & & \text{fr}(v_5) &= (0, 0) \end{aligned}$$

(b) Otherwise, if $e^2 \not\equiv_8 D$, the HLK-invariant is also $(1, [2, 2, 0])$ since

$$\begin{aligned} \text{fr}(v_1) &= (1/2, 1/2), & \text{fr}(v_2) &= (0, 1/2), & \text{fr}(v_3) &= (1/2, 1/2), \\ \text{fr}(v_4) &= (0, 0) & \text{and} & & \text{fr}(v_5) &= (0, 1/2). \end{aligned}$$

(2) If $D \equiv_8 1$, we have two cases:

(a) If $e \equiv_4 1$, the HLK-invariant is $(2, [1, 1, 1])$ since

$$\begin{aligned} \text{fr}(v_1) &= (1/2, 1/2), & \text{fr}(v_2) &= (0, 1/2), & \text{fr}(v_3) &= (1/2, 0), \\ \text{fr}(v_4) &= (0, 0) & \text{and} & & \text{fr}(v_5) &= (0, 0). \end{aligned}$$

(b) If $e \equiv_4 3$, the HLK-invariant is $(0, [3, 1, 1])$ since

$$\begin{aligned} \text{fr}(v_1) &= (1/2, 1/2), & \text{fr}(v_2) &= (1/2, 1/2), & \text{fr}(v_3) &= (1/2, 1/2), \\ \text{fr}(v_4) &= (1/2, 0) & \text{and} & & \text{fr}(v_5) &= (0, 1/2). \end{aligned}$$

Observe that the HLK-invariants that we obtain are precisely the same as the ones found for the arithmetic case.

The linear action on these points restricts the possibilities of the action of $\text{Aff}(L_D(e))$ on regular Weierstrass points. We get the following results which are completely analogous to the arithmetic case (that is, to [Propositions 4.2](#) and [4.3](#)):

- (1) If D is even, that is, $D \equiv_8 0$ or $D \equiv_8 4$, we have two cases:
 (a) If $e^2 \equiv_8 D$, then

$$G_D(e) \leq \text{Dih} \left(\begin{array}{ccc} & 1 & \\ 4 & \swarrow \quad \searrow & 3 \\ & 2 & \end{array} \right),$$

- (b) Otherwise, if $e^2 \not\equiv_8 D$, then

$$G_D(e) \leq \text{Dih} \left(\begin{array}{ccc} & 1 & \\ 5 & \swarrow \quad \searrow & 2 \\ & 3 & \end{array} \right),$$

- (2) If D is odd, that is $D \equiv_8 1$, then

$$G_D(e) \leq \text{Sym}(\{1, 2, 3\}) \times \text{Sym}(\{4, 5\}).$$

Thus, we get the following result:

Corollary 4.6. *Let $D > 4$ be a real quadratic discriminant which is a quadratic residue modulo 8. We have the following:*

- (a) *If $D \equiv_4 0$ and $X \in \Omega W_D$, then $G(X)$ is contained in a group conjugate to Dih_4 .*
 (b) *If $D \equiv_8 1$ and $X \in \Omega W_D$ (regardless of the spin), then $G(X)$ is contained in a group isomorphic to $\text{Sym}_3 \times \text{Sym}_2 \cong \text{Dih}_6$. \square*

4.2.2. Quadratic nonresidues modulo 8. In the remaining case of $D \equiv_8 5$, let $e \in R_D$ and consider $\rho = \lambda = \frac{\sqrt{D+e}}{2}$. With this choice of ρ , we get that the location of the regular Weierstrass points (relative to the singularity) in $L_D(e)$, up to addition by elements in \mathcal{O}_D , are

$$\begin{aligned} v_1 &= \left(\frac{1}{2}, \frac{1}{2} \right), & v_2 &= \left(\frac{1}{2} + \frac{1}{2}\rho, \frac{1}{2} \right), & v_3 &= \left(\frac{1}{2}, \frac{1}{2} + \frac{1}{2}\rho \right), \\ v_4 &= \left(\frac{1}{2} + \frac{1}{2}\rho, 0 \right) & \text{and} & & v_5 &= \left(0, \frac{1}{2} + \frac{1}{2}\rho \right). \end{aligned}$$

Lemma 4.7. *Let $D > 4$ be a real quadratic discriminant with $D \equiv_8 5$. Let $e \in R_D$, and observe that e is odd. Fix $\rho = \frac{\sqrt{D+e}}{2}$. If $A \in \text{SL}(L_D(e))$ fixes $v_1 \pmod{\mathcal{O}_D}$, then $\text{fr}(Av_2) = \text{fr}(Av_3)$.*

Proof. Note first that

$$\frac{1}{2}\rho^2 = \frac{1}{2}\rho + \frac{D-e^2}{8} \equiv_{\mathcal{O}_D} \frac{1}{2}\rho + \frac{1}{2}.$$

Then, by [Equation \(2\)](#), we have that

$$\begin{aligned} Av &\equiv_{\mathcal{O}_D} \frac{1}{2} \begin{pmatrix} p_\alpha p_1 + p_\beta p_2 + q_\alpha q_1 + q_\beta q_2 \\ p_\gamma p_1 + p_\delta p_2 + q_\gamma q_1 + q_\delta q_2 \end{pmatrix} \\ &+ \frac{\rho}{2} \begin{pmatrix} q_\alpha q_1 + q_\beta q_2 + p_\alpha q_1 + p_\beta q_2 + q_\alpha p_1 + q_\beta p_2 \\ q_\gamma q_1 + q_\delta q_2 + p_\gamma q_1 + p_\delta q_2 + q_\gamma p_1 + q_\delta p_2 \end{pmatrix}. \end{aligned}$$

for any $v \in \frac{1}{2}(\mathcal{O}_D \times \mathcal{O}_D)$. In particular, we have that

$$\text{fr}(Av) \equiv_1 \frac{1}{2} \begin{pmatrix} p_\alpha p_1 + p_\beta p_2 + q_\alpha q_1 + q_\beta q_2 \\ p_\gamma p_1 + p_\delta p_2 + q_\gamma q_1 + q_\delta q_2 \end{pmatrix}.$$

Since A fixes $v_1 \bmod \mathcal{O}_D$, we obtain that

$$v_1 = \frac{1}{2} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \equiv_{\mathcal{O}_D} \frac{1}{2} \begin{pmatrix} p_\alpha + p_\beta \\ p_\gamma + p_\delta \end{pmatrix} + \frac{\rho}{2} \begin{pmatrix} q_\alpha + q_\beta \\ q_\gamma + q_\delta \end{pmatrix}.$$

It follows that

$$\begin{aligned} p_\alpha + p_\beta &\equiv_2 1, & q_\alpha + q_\beta &\equiv_2 0, \\ p_\gamma + p_\delta &\equiv_2 1, & q_\gamma + q_\delta &\equiv_2 0. \end{aligned}$$

Applying this to v_2 , we get

$$\text{fr}(Av_2) \equiv_1 \frac{1}{2} \begin{pmatrix} p_\alpha + p_\beta + q_\alpha \\ p_\gamma + p_\delta + q_\gamma \end{pmatrix} \equiv_1 \frac{1}{2} \begin{pmatrix} 1 + q_\alpha \\ 1 + q_\gamma \end{pmatrix}.$$

Similarly,

$$\text{fr}(Av_3) \equiv_1 \frac{1}{2} \begin{pmatrix} p_\alpha + p_\beta + q_\beta \\ p_\gamma + p_\delta + q_\delta \end{pmatrix} \equiv_1 \frac{1}{2} \begin{pmatrix} 1 + q_\alpha \\ 1 + q_\gamma \end{pmatrix}.$$

□

Proposition 4.8. *Let $D > 4$ be a real quadratic discriminant with $D \equiv_8 5$. Then, for any $e \in R_D$, the group $G_D(e) = G(L_D(e))$ of permutations of the regular Weierstrass points of $L_D(e)$ is such that*

$$G \leq \text{Dih} \left(\begin{array}{c} 1 \\ 3 \quad \diagdown \quad \diagup \quad 2 \\ 4 \quad \text{---} \quad 5 \end{array} \right).$$

Proof. Using $\rho = \frac{\sqrt{D+e}}{2}$ as in the previous lemma, we have that

$$\begin{aligned} \text{fr}(v_1) &= (1/2, 1/2), & \text{fr}(v_2) &= (1/2, 1/2), & \text{fr}(v_3) &= (1/2, 1/2), \\ \text{fr}(v_4) &= (1/2, 0) & \text{and} & & \text{fr}(v_5) &= (0, 1/2). \end{aligned}$$

Let $A \in \text{SL}(L_D(e))$ that fixes w_1 . Then, A fixes $v_1 \bmod \mathcal{O}_D$ and, by the previous lemma, $\text{fr}(Av_2) = \text{fr}(Av_3)$. But $\{v_2, v_3\}$ is the only possible such pair, so A fixes $\{w_2, w_3\}$. Since A fixes $\{w_1\}$ and $\{w_2, w_3\}$, it also fixes $\{w_4, w_5\}$.

In particular, every element in $G_D(e)$ that fixes 1 has order at most 2. On the other hand, by [Proposition 3.5](#), $G_D(e)$ has a subgroup of order 10. But, up to automorphisms, the subgroups of Sym_5 of order divisible by 10 are Dih_5 , $\text{GA}(1, 5)$, A_5 and Sym_5 . All of them, but Dih_5 , have elements of order greater than 2 that stabilise 1 (in particular, such elements can be taken to be cycles). □

We obtain the following straightforward corollary:

Corollary 4.9. *Let $D > 4$ be a real quadratic discriminant with $D \equiv_8 5$ and $X \in \Omega W_D$. Then, $G(X)$ is contained in a group conjugate to Dih_5 .* □

5. WEIERSTRASS EIGENFORMS IN GENUS TWO OF SMALL DISCRIMINANT

In this section we treat the remaining cases are not covered by our previous arguments. That is, the cases where the discriminant D is either 9 or 33.




FIGURE 5. In the case of $D = 9$, $L_9(-1) \in \Omega\mathcal{W}_9$ is a square-tiled surface and the diagonal direction induces the permutation $(4, 5) \in G_9(-1)$.




FIGURE 6. In the case $D = 33$, we exhibit explicit directions that induce the permutation $(4, 5) \in G_{33}(e)$, for $e = -1$ (left) and $e = 1$ (right). The cylinder C_1 is shown in a darker shade, while C_2 is depicted in a lighter shade.

5.1. **Discriminant 9.** We will start by fixing $D = 9$, so $R_9 = \{-1\}$ and that the surface $S = L_{25}(-1)$ is the primitive square-tiled surface depicted in Figure 5.

The action of the horizontal and vertical twists on S generates $\text{Sym}(\{1, 2, 3\}) \leq G(S)$, as was seen in Proposition 3.6. On the other hand, the diagonal direction, (that is, taking $\theta = \arctan(1) = \pi/4$) on S gives a one-cylinder decomposition whose core curve contains w_4 and w_5 , as shown in Figure 5. It follows that $(4\ 5) \in G(S)$ and, by Corollary 4.6, we conclude that $G(S) = \text{Sym}(\{1, 2, 3\}) \times \text{Sym}(\{4, 5\}) \cong \text{Dih}_6$.

Finally, since $\Omega\mathcal{W}_9$ is connected, we obtain that $G(X) \cong \text{Sym}_3 \times \text{Sym}_2 \cong \text{Dih}_6$ for every $X \in \Omega\mathcal{W}_9$. Combining this with Proposition 3.6, we obtain that $G(X)$ can be generated by the action of three parabolic elements. \square

5.2. **Discriminant 33.** We now focus our attention to the case where $D = 33$. We have that $R_{33} = \{-5, -3, -1, 1\}$ and we will consider the surfaces $L_{33}(-1)$ and $L_{33}(1)$, depicted in Figure 6. By Equation (1), these surfaces belong to distinct components of $\Omega\mathcal{W}_{33}$.

By Proposition 3.6, the action of the horizontal and vertical twists on both surfaces generates $\text{Sym}_3 \leq G(S)$. In both cases, we will exhibit an explicit direction that induces the remaining transposition $(4\ 5)$. After this is done, we will obtain that $G(X)$

is isomorphic to $\text{Sym}_3 \times \text{Sym}_2 \cong \text{Dih}_6$ for every $X \in \Omega W_{33}$. Combining this with [Proposition 3.6](#), we obtain that $G(X)$ can be generated by the action of three parabolic elements.

To simplify the computations, we define the *horizontal height* of a cylinder to be the length of a maximal horizontal segment contained in the cylinder. Indeed, when computing the ratio of the moduli of two cylinders in a cylinder decomposition, we may replace the height by the horizontal height and obtain the same result. In fact, the horizontal height is equal to the height up to a linear factor that is cancelled out when taking the ratio. In other words, the horizontal height corresponds to the actual height of the same cylinder when considering a sheared version of the surface; the shear is performed in the direction of the cylinder decomposition and, thus, does not change any cylinder width. By a slight abuse of notation, we will denote the horizontal heights by h as we previously did for the actual heights.

Similarly, we will also use the *horizontal width* of a cylinder, which is the length of the projection of the cylinder to the horizontal axis. An argument analogous to the paragraph above shows that, when computing ratio of moduli, using the actual width of the horizontal width makes no difference. We denote the horizontal width by w .

First consider the case $L_{33}(-1)$ as in [Figure 6a](#). We take the cylinder decomposition with slope $\frac{1}{\lambda}$, that is, in direction $\theta = \arctan\left(\frac{1}{\lambda}\right)$.

Let C_1 be the cylinder containing the points w_4 and w_5 , and let C_2 be the other cylinder of the decomposition. We need to show that the ratio of the moduli of these cylinders belongs to $\frac{\text{odd}}{\text{odd}}$ or $\frac{\text{even}}{\text{odd}}$ to show that the transposition $(4\ 5)$ belongs to $G(L_{33}(-1))$.

As already discussed, we will use horizontal heights instead of heights to compute the moduli. We denote these heights by h_1 and h_2 , respectively. We have that these numbers satisfy the equations

$$\begin{aligned} h_1 + h_2 &= 1 \\ 3h_1 + 2h_2 &= \lambda. \end{aligned}$$

so $h_1 = \lambda - 2$ and $h_2 = 3 - \lambda$.

The modulus of C_1 is then

$$m_1 = \frac{\lambda + 1 + 1 + 1 + 1 + 1 + h_2 + \lambda + h_1 + (\lambda - h_1) + (\lambda - 1) + 1}{h_1} = \frac{3\lambda + 8}{\lambda - 2},$$

and that the modulus of C_2 is then

$$m_2 = \frac{\lambda + (h_1 + h_2) + (\lambda - (h_1 + h_2)) + 2(h_1 + h_2) + 1 + 1 + 1 + 1 + 1 + 1}{h_2} = \frac{2\lambda + 8}{\lambda - 3}.$$

We obtain that $r_\theta = \frac{m_1}{m_2} = \frac{-3\lambda^2 + \lambda + 24}{2(\lambda^2 + 2\lambda - 8)}$. Since $\lambda = \frac{\sqrt{33}-1}{2}$, an elementary computation shows that $r_\theta = 2$. Thus, $\tau_\theta = (4\ 5) \in G(L_{33}(-1))$. Combining this with [Proposition 3.6](#), we obtain that $G(L_{33}(-1)) \cong \text{Sym}_3 \times \text{Sym}_2 \cong \text{Dih}_6$.

We now focus our attention on $L_{33}(1)$ and consider the cylinder decomposition with slope $\frac{3}{\lambda-2}$, that is, with direction $\theta = \arctan\left(\frac{3}{\lambda-2}\right)$.

Let C_1 be the cylinder containing the points w_4 and w_5 , and let C_2 be the other cylinder of the decomposition. As in the previous case, we need to show that the ratio of the moduli of these cylinders has odd denominator (when written as an irreducible fraction) to obtain that $(4\ 5) \in G(L_{33}(1))$. Let h_1 and h_2 be the horizontal heights of C_1 and C_2 , respectively. We have that

$$h_1 + h_2 = 1$$

$$3h_1 = \lambda - 2.$$

so $h_1 = \frac{\lambda-2}{3}$ and $h_2 = \frac{5-\lambda}{3}$.

The modulus of C_1 is then

$$m_1 = \frac{h_1 + h_1 + h_1 + 1 + 1 - h_1 + h_1}{h_1} = \frac{3\lambda}{\lambda - 2},$$

and that the modulus of C_2 is then

$$m_2 = \frac{1 + 1}{h_2} = \frac{6}{5 - \lambda}.$$

We obtain that $r_\theta = \frac{m_1}{m_2} = \frac{\lambda(5-\lambda)}{2(\lambda-2)}$. Since $\lambda = \frac{\sqrt{33}+1}{2}$, an elementary computation shows that $r_\theta = 2$. Thus, $\tau_\theta = (4\ 5) \in G(\mathbb{L}_{33}(-1))$. Combining this with [Proposition 3.6](#), we obtain that $G(\mathbb{L}_{33}(-1)) \cong \text{Sym}_3 \times \text{Sym}_2 \cong \text{Dih}_6$. \square

6. CLASSIFICATION OF PERMUTATIONS IN GENUS TWO

In this section we collect our results to give a complete proof of [Theorem 1.1](#) and [Corollary 1.2](#).

Proof of [Theorem 1.1](#). In the case $D \equiv_4 0$, [Proposition 3.4](#) shows a first containment, namely that $G(X)$ contains a subgroup conjugate to Dih_4 for any $X \in \Omega\mathcal{W}_D$. Moreover, [Proposition 4.2](#) gives the reverse inclusion in the arithmetic case, and [Corollary 4.6\(a\)](#) provides the corresponding reverse inclusion in the non-arithmetic case.

In the case $D \equiv_8 1$, [Proposition 4.3](#) and [Corollary 4.6\(b\)](#) show a first containment, that is, that $G(X)$ is contained in a group conjugate to $\text{Sym}_3 \times \text{Sym}_2 \cong \text{Dih}_6$, for the arithmetic and non-arithmetic cases, respectively. The reverse inclusion can be deduced *a posteriori* from [Propositions 3.6](#) and [3.7](#), for $D \notin \{9, 33\}$. Indeed, the former proposition shows that $G(X)$ contains a subgroup conjugate to Sym_3 , while the latter shows that it contains a product of two disjoint transpositions. Since no product of disjoint two transpositions belongs to Sym_3 , we conclude that $G(X)$ contains a *proper* subgroup conjugate to Sym_3 . Thus, $G(X)$ must be isomorphic to $\text{Sym}_3 \times \text{Sym}_2 \cong \text{Dih}_6$.

In the case $D \equiv_8 5$, [Proposition 3.5](#) shows the first containment, namely that $G(X)$ contains a subgroup conjugate to Dih_5 for any $X \in \Omega\mathcal{W}_D$. Finally, [Corollary 4.9](#) gives the corresponding reverse inclusion.

The only remaining cases are the exceptional discriminants $D = 9$ and $D = 33$. These cases are treated in [Section 5.1](#) and [Section 5.2](#), respectively. \square

Proof of [Corollary 1.2](#). For $D \equiv_4 0$ and $D \equiv_8 5$, we have shown this result in [Propositions 3.4](#) and [3.5](#).

In the case of $D \equiv_8 1$ and $D \notin \{9, 33\}$, we have by [Proposition 3.7](#) that $G(X)$ contains a product of two disjoint transpositions induced by the action of a *single* parabolic element. Moreover, [Proposition 3.6](#) shows that $G(X)$ contains a subgroup conjugate to Sym_3 , which is, in turn, generated by the action of two parabolic elements. We know that these three parabolic elements generate $G(X)$, but one of these elements is redundant. Indeed, assume that we label the regular Weierstrass points in such a way that $G(X)$ is contained in $\text{Sym}(\{1, 2, 3\}) \times \text{Sym}(\{4, 5\})$. Then, the product of two disjoint transpositions must be of the form $\tau = (r\ s)(4\ 5)$, where $r < s$ and $r, s \in \{1, 2, 3\}$. Since there exist two parabolic elements in $\text{SL}(X)$ whose actions generate a subgroup of $G(X)$ conjugate to $\text{Sym}(\{1, 2, 3\})$, some parabolic element gives a transposition

$\sigma \neq (r\ s)$. We have that $\langle \sigma, \tau \rangle = \text{Sym}(\{1, 2, 3\}) \times \text{Sym}(\{4, 5\})$, so we obtain the desired result.

For $D \in \{9, 33\}$, no parabolic element can produce a product of disjoint transpositions (see [Remark 3.8](#)). In particular, since no pair of transpositions can generate a group conjugate to $\text{Sym}(\{1, 2, 3\}) \times \text{Sym}(\{4, 5\})$, there is no pair of parabolic elements that can generate the corresponding permutation group. On the other hand, we show in [Section 5.1](#) that $G(X)$ can be generated by the action of three parabolic elements for any $X \in \Omega\mathcal{W}_9$ and, similarly, we show in [Section 5.2](#) that $G(X)$ can be generated by the action of three parabolic elements for any $X \in \Omega\mathcal{W}_{33}$. \square

7. WEIERSTRASS PRYM EIGENFORMS

In this section we treat the case of Weierstrass Prym eigenforms. For more detailed discussion and definitions we refer to McMullen's seminal work [\[McM06\]](#) and the classification by Lanneau–Nguyen [\[LN14\]](#).

7.1. Prym varieties and eigenforms. Fix a Riemann surface X . Recall that $H^{1,0}(X)$ is the complex vector space of Abelian differentials on X . As a real vector space, $H^{1,0}(X)$ is canonically isomorphic to $H^1(X; \mathbb{R})$ via the isomorphism $\omega \mapsto \text{Re}(\omega)$. This isomorphism also endows $H^{1,0}(X)$ with a symplectic form by pulling back the intersection form on $H^1(X; \mathbb{R})$.

Suppose that $\iota: X \rightarrow X$ is a holomorphic involution with $\iota^*\omega = -\omega$. Then, the *Prym variety* of (X, ι) is the subtorus of the Jacobian of X determined by the anti-invariant eigenspace of ι :

$$\text{Prym}(X, \iota) = (H^{1,0}(X)^-)^* / (H_1(X; \mathbb{Z})^-).$$

Recall that an Abelian variety A admits real multiplication by \mathbb{O}_D (the real quadratic order of discriminant $D > 0$) if $\dim_{\mathbb{C}} A = 2$ and \mathbb{O}_D occurs as an indivisible, self-adjoint subring of $\text{End}(A)$.

Now suppose that $P = \text{Prym}(X, \iota)$ is a Prym variety with real multiplication by \mathbb{O}_D . Then \mathbb{O}_D also acts on $H^{1,0}(X)^-$ and we say that $\omega \in H^{1,0}(X)^-$ is a *Prym eigenform* (for real multiplication by \mathbb{O}_D) if $\mathbb{O}_D \cdot \omega \subseteq \mathbb{C} \cdot \omega$ (and $\omega \neq 0$). We say that ι is the *Prym involution*.

Let $E_D^g \subseteq \Omega\mathcal{M}_g$ denote the space of all genus- g Prym eigenforms for real multiplication by \mathbb{O}_D . By the seminal work of McMullen [\[McM06\]](#), E_D is a closed, $\text{GL}^+(2, \mathbb{R})$ -invariant subset of $\Omega\mathcal{M}_g$.

Note that $2 = \dim_{\mathbb{C}} \text{Prym}(X, \iota) \leq g$ and, by Riemann–Hurwitz formula, ι has exactly $10 - 2g$ fixed points. Thus, we have that $2 \leq g \leq 5$.

In this work, we focus our attention on *Weierstrass Prym eigenform loci*, that is, loci of Prym eigenforms with a single zero. We define $E_D(2g - 2) = E_D^g \cap \mathcal{H}(2g - 2)$. By a dimension count, $E_D(2g - 2)$ consists entirely of Veech surfaces (see McMullen's work [\[McM06, Theorem 1.2\]](#) and [Theorem 7.1](#) below).

It is not difficult to see that $E_D(2) = \Omega\mathcal{W}_D \subseteq \mathcal{H}(2)$. Thus, Weierstrass Prym eigenforms in genus two correspond exactly to Veech surfaces in $\mathcal{H}(2)$, and the Prym involution corresponds to the hyperelliptic involution.

As in the case of Veech surfaces in $\mathcal{H}(2)$, we are interested in the action of $\text{Aff}(X, \omega)$ on the set of fixed points for the Prym involution.

For a Prym eigenform in $E_D(2g - 2)$, the only zero of order $2g - 2$ is necessarily fixed by the corresponding Prym involution. Thus:




FIGURE 7. Prototypical surfaces $A^+(a, b, c, e)$ (left) and $A^-(a, b, c, e)$ (right), with $\lambda = \frac{e+\sqrt{D}}{2}$. These examples depict $A^+(1, 4, 2, -3)$ for $D = 73$, and $A^-(1, 7, 3, -3)$ for $D = 177$, respectively.

- in genus three, Prym eigenforms in $E_D(4)$ have exactly three fixed regular points for the Prym involution;
- in genus four, Prym eigenforms in $E_D(6)$ have only one fixed regular point for the Prym involution; and
- in genus five, $E_D(8) = \emptyset$.

We conclude that the action of the Veech group on the regular fixed points of the Prym involution can be nontrivial only in genus two and three. The genus-two case was already established in [Theorem 1.1](#). Thus, we focus our attention to Weierstrass Prym eigenform loci in genus three.

By the work of Lanneau–Nguyen [\[LN14\]](#), we have the following classification:

Theorem 7.1 ([\[LN14, Theorem 1.1\]](#)). *For $D \geq 17$, the locus $E_D(4)$ is non-empty if and only if $D \equiv_8 1$ or $D \equiv_4 0$. Moreover, $E_D(4)$ has exactly two connected components for $D \equiv_8 1$ and is connected for $D \equiv_4 0$.*

For $D < 17$, the locus $E_D(4)$ is non-empty if and only if $D \in \{8, 12\}$ and, in these cases, it is connected.

Finally, each component of $E_D(4)$ is a closed $\mathrm{GL}^+(2, \mathbb{R})$ -orbit. Moreover, the loci $E_D(4)$ are pairwise disjoint for distinct D .

7.2. Prototypes in $E_D(4)$. As in genus two, and following Lanneau–Nguyen [\[LN14\]](#), we say that a quadruple of integers (a, b, c, e) is a *splitting prototype* of discriminant D if:

$$\begin{aligned} D &= e^2 + 8bc, & 0 \leq a < \gcd(b, c), & \quad 2c + e < b, \\ & & 0 < b, c, & \quad \text{and} \quad \gcd(a, b, c, e) = 1. \end{aligned}$$

With each splitting prototype (a, b, c, e) , we associate two translation surfaces as in [Figure 7](#), where $\lambda = \frac{e+\sqrt{D}}{2}$. We call such surfaces *prototypical surfaces* associated with the splitting prototype (a, b, c, e) and denote them by $A^+(a, b, c, d)$ and $A^-(a, b, c, e)$; we refer to them as *model A^+* and *model A^-* , respectively. The work of Lanneau–Nguyen [\[LN14\]](#) shows that these surfaces belong to $E_D(4)$.

Unlike in genus two, these prototypical surfaces do not account for all cusps. However, for each splitting prototype (a, b, c, e) of discriminant D , the models A^+ and A^-




FIGURE 8. Re-scaled prototypical surface $Z_D(e)$ obtained by applying $\begin{pmatrix} 2\lambda^{-1} & 0 \\ 0 & 1 \end{pmatrix}$ to $A_D^-(e)$. This example shows $Z_{116}(-6)$.

can be identified with distinct cusp of $E_D(4)$. In most cases this is enough to prove [Theorem 1.3](#). In fact, we have the following.

Theorem 7.2 ([LN14, Proposition 4.7]). *Let $X \in E_D(4)$. If X admits no cylinder decompositions in model A^+ or model A^- , then $D = 8$.*

7.2.1. Reduced prototypes. We can define *reduced* prototypes in a similar fashion as for genus two: those satisfying $c = 1$. Unlike genus two, however, in genus three we will work exclusively with reduced prototypes. These splitting prototypes necessarily have the form

$$(a, b, c, e) = \left(0, \frac{D - e^2}{8}, 1, e \right)$$

and are therefore parametrised by

$$S_D = \{e \in \mathbb{Z} \mid e^2 \equiv_8 D, e^2 < D, (e + 4)^2 < D\}.$$

We denote the prototypical surface associated with the reduced prototype $e \in S_D$ by $A_D^+(e)$ and $A_D^-(e)$. That is, $A_D^\pm(e) = A^\pm\left(0, \frac{D - e^2}{8}, 1, e\right) \in E_D(4)$.

7.2.2. Disconnected loci. When $E_D(4)$ is disconnected, there is a parity invariant analogous to the spin invariant in the case of Veech surfaces in $\mathcal{H}(2)$. Even if it is possible to obtain a closed expression for this invariant (as [Equation \(1\)](#) in genus two), this is outside of the scope of our work (and there is no direct reference as in the case of genus two). Indeed, we only need to tell some particular cases apart for our purposes. More precisely, we use the following:

Lemma 7.3. *Let D be an odd discriminant such that $E_D(4)$ is non-empty (and, thus, disconnected). Then,*

- If $e \in S_D$, then $A_D^+(e)$ and $A_D^-(e)$ lie at different components of $E_D(4)$.
- If $e, e' \in S_D$, then $A_D^+(e)$ and $A_D^+(e')$ (resp. $A_D^-(e)$ and $A_D^-(e')$) lie at the same component of $E_D(4)$ if and only if $e \equiv_4 e'$.

That is, $A_D^+(e)$ and $A_D^-(e')$ lie at the same component of $E_D(4)$ if and only if $e \equiv_4 -e'$.

Proof. The first claim follows directly from Lanneau–Nguyen [LN14, Theorem 6.1]. The proof of the second claim is analogous to McMullen’s work in genus two [McM05, Theorem 5.3] (cf. Lanneau–Nguyen [LN14, Lemma 6.2]). \square

7.2.3. Re-scaled prototypical surfaces. We also use, in [Section 9](#), a re-scaled version of the surface associated with a reduced prototype. This surface is obtained by the action of the diagonal matrix $\begin{pmatrix} 2\lambda^{-1} & 0 \\ 0 & 1 \end{pmatrix} \in \text{GL}^+(2, \mathbb{R})$ on $A_D^-(e)$, with $e \in S_D$, as in [Figure 8](#). We denote such surface by $Z_D(e) \in E_D(4)$.

8. ACTION OF PARABOLIC ELEMENTS IN GENUS THREE

Given a reduced prototype $e \in S_D$ of discriminant D , a quick examination of [Figure 7](#) shows that the horizontal and vertical twists on the associated prototypical surfaces $A^\pm(e)$ induce either trivial elements or transpositions in $G(A^\pm(e))$. The following lemma explicitly states the conditions so that Prym fixed points are exchanged or not.

Lemma 8.1. *Let $D > 4$ be a real quadratic discriminant such that $E_D(4)$ is non-empty and let $e \in S_D$. Let $b = \frac{D-e^2}{8}$. We have that:*

- *the horizontal twist acts on the Prym fixed points of $A_D^\pm(e)$ either trivially or by the transposition (1 2). In the case of $A_D^+(e)$, the action is non-trivial if and only if b is odd, while it is always non-trivial for $A_D^-(e)$; and*
- *the vertical twist acts on the Prym fixed points of $A_D^\pm(e)$ either trivially or by the transposition (1 3). In the case of $A_D^+(e)$ the action is always non-trivial, while it is non-trivial for $A_D^-(e)$ if and only if $b - e - 2$ is odd.*

Proof. The strategy of the proof is simply to analyse the cylinder decompositions and the ratio of the moduli of the resulting cylinders for the horizontal and vertical directions, as we did in [Section 3.2](#).

For the horizontal direction, the prototypical surface $A_D^+(e)$ has two long cylinders of moduli $m_1 = b$, and one short cylinder of modulus $m_2 = 1$. The Prym fixed points w_1 and w_2 lie in the interior of the short cylinder. See [Figure 7](#) (left). Therefore, the corresponding ratio is $r_h = \frac{m_1}{m_2} = b$. Thus, the points are exchanged if and only if $r_h \in \frac{\text{odd}}{\text{odd}} \sqcup \frac{\text{odd}}{\text{even}}$, which clearly happens if and only if b is odd, as it is an integer.

Moreover, the prototypical surface $A_D^-(e)$ has one long cylinder of modulus $m_1 = b$, and two short cylinders of moduli $m_2 = 1$. The Prym fixed points w_1 and w_2 lie in the interior of the long cylinder. See [Figure 7](#) (right). We obtain that the corresponding ratio is $r_h = \frac{m_1}{m_2} = b$ as in the previous case. Thus, the points are exchanged if and only if $r_h \in \frac{\text{odd}}{\text{odd}} \sqcup \frac{\text{even}}{\text{odd}}$, which happens regardless of r as it is an integer.

The vertical direction can be treated similarly. Indeed, observe that the prototypical surface $A_D^+(e)$ has a long cylinder of modulus $m_1 = \frac{\lambda+2}{\lambda}$, and two short cylinders of moduli $m_2 = \frac{1}{b-\lambda}$. The points w_1 and w_3 lie in the interior of the long cylinder. The corresponding ratio is

$$r_v = \frac{m_1}{m_2} = \frac{\frac{\lambda+2}{\lambda}}{\frac{1}{b-\lambda}} = \frac{D - (e+4)^2}{8} = b - e - 2$$

which is also an integer so the points are exchanged unconditionally.

Finally, the prototypical surface $A_D^-(e)$ has two long cylinders of modulus $m_1 = \frac{\lambda/2+1}{\lambda/2}$ and a short cylinder of modulus $m_2 = \frac{1}{b-\lambda}$. The points w_1 and w_3 lie in the interior of the short cylinder. Thus, $r_v = \frac{m_1}{m_2} = b - e - 2$, so these points are exchanged by the vertical twist if and only if $b - e - 2$ is odd. \square

Remark 8.2. We have refrained from introducing the notion of *butterfly moves* to simplify the exposition, but, as in the genus-two case (see [Remark 3.3](#)), the cylinder decomposition of the vertical direction on $A_D^\pm(e)$ can be thought of as the butterfly move B_∞ , which transforms it into the prototypical surface $A_D^\mp(-e-4)$ [[LN14](#), Proposition 7.5(2)]. Thus, again, the fact that the action induced by the vertical twist depends on the parity of $\frac{D-(e+4)^2}{8}$ can be also explained in this way.

The previous lemma already allows us to show that $G(X)$ is non-trivial in most cases.

Proposition 8.3. *Let $D \neq 8$ be a real quadratic discriminant such that $E_D(4)$ is non empty and let $X \in E_D(4)$. Then, $G(X)$ contains a transposition, which is the action of a parabolic element in $\mathrm{SL}(X)$.*

Proof. By [Theorem 7.2](#), for any D as in the statement there exists a model- A^\pm cylinder decomposition. In particular, $S_D \neq \emptyset$.

Now, given a reduced prototype $e \in S_D$, by [Lemma 8.1](#), the horizontal twist on $A_D^-(e)$ gives $(1\ 2) \in G(A_D^-(e))$ and the vertical twist on $A^+(e)$ gives $(1\ 3) \in G(A_D^+(e))$.

If D is even, $E_D(4)$ is connected, so this already proves the desired result for every $X \in E_D(4)$. On the other hand, if D is odd and $E_D(4)$ is disconnected, [Theorem 7.1](#) shows that $A_D^-(e)$ and $A^+(e)$ lie on different component, so the result also follows for any $X \in E_D(4)$. \square

Remark 8.4. By [Lemma 8.1](#), in order to get the entire symmetric group Sym_3 from the horizontal and vertical twists on $A_D^+(e)$, we need that $r_h = b = \frac{D-e^2}{8}$ is odd, or, equivalently, that $D \equiv_{16} 8+e^2$. Similarly, on $A_D^-(e)$, we need $r_v = \frac{D-(e+4)^2}{8}$ to be odd, that is, we need that $D \equiv_{16} 8+(e+4)^2$. Since [Lemma 7.3](#) shows that $A^+(e)$ and $A^-(e+4)$ lie on the same component, verifying these conditions for either case is actually equivalent.

The next series of lemmas establish [Theorem 1.3](#) for $D \geq 12$ satisfying $D \equiv_4 0$ or $D \equiv_8 1$, with $D \notin \{25, 33\}$, and such that D is either odd or an even quadratic nonresidue modulo 16. More precisely, we will show that $G(X) = \mathrm{Sym}_3$ for any $X \in E_D(4)$ for such D . Observe that if D is even, it is a quadratic nonresidue if and only if $D \equiv_{16} 8$ or $D \equiv_{16} 12$. We will analyse these cases separately.

Lemma 8.5. *Let $D > 0$ be a real quadratic discriminant with $D \equiv_{16} 8$ and $X \in E_D(4)$. If $D \neq 8$, then $G(X) = \mathrm{Sym}_3$ and is generated by the action of two parabolic elements in $\mathrm{SL}(X)$.*

Proof. Clearly, $D \equiv_8 0$. By hypothesis, $D \geq 24$, so we have that $0 \in S_D$. Since $E_D(4)$ is connected, it is enough to prove the result for the prototypical surface $A_D^+(0)$. As discussed in [Remark 8.4](#), it suffices to show that $D \equiv_{16} 8 + 0^2$, which also holds by hypothesis. \square

Remark 8.6. Observe that the previous proof fails for $D = 8$, as $0 \notin S_8 = \emptyset$. We postpone the proof for the case $D = 8$ to [Section 10.1](#).

Lemma 8.7. *Let $D > 0$ be a real quadratic discriminant with $D \equiv_{16} 12$ and let $X \in E_D(4)$. Then, $G(X) = \mathrm{Sym}_3$ and is generated by the action of two parabolic elements in $\mathrm{SL}(X)$.*

Proof. We have that $D \equiv_8 4$. By hypothesis, $D \geq 12$, so we have that $-2 \in S_D$. Since $E_D(4)$ is connected, it is enough to prove the result for the prototypical surface $A_D^+(-2)$. As discussed in [Remark 8.4](#), it suffices to show that $D \equiv_{16} 8 + (-2)^2$, which also holds by hypothesis. \square

Lemma 8.8. *Let $D > 0$ be a real quadratic discriminant with $D \equiv_8 1$ and let $X \in E_D(4)$. If $D \geq 33$, then $G(X) = \mathrm{Sym}_3$ and is generated by the action of two parabolic elements in $\mathrm{SL}(X)$.*

Proof. Assume first that $D \equiv_{16} 1$. By hypothesis, $D \geq 33$, so we have that $\{-5, -3\} \subseteq S_D$. Observe that $D \equiv_{16} 8 + e^2$ for $e \in \{-5, -3\}$, so, as discussed in [Remark 8.4](#), we obtain directly that $G(A_D^+(-5)) = \mathrm{Sym}_3$ and that $G(A_D^+(-3)) = \mathrm{Sym}_3$. Since, by [Lemma 7.3](#), these surfaces lie on different components, we obtain the result for any $X \in E_D(4)$.




FIGURE 9. The square-tiled surface $Z_{d^2}(e) \in E_{d^2}(4)$, for even $d > 4$. This example shows $Z_{100}(-2)$.

Now, assume that $D \equiv_{16} 9$. By hypothesis, $D \geq 41$, so we have that $\{-1, 1\} \subseteq S_D$. In this case, $D \equiv_{16} 8 + e^2$ for $e \in \{-1, 1\}$. The result follows analogously. \square

Remark 8.9. Observe that the previous proof fails for $D \in \{17, 25\}$. We postpone the proof for these cases to [Sections 10.2](#) and [10.3](#).

9. RESTRICTIONS FOR PERMUTATIONS IN GENUS THREE

In the cases where $D \not\equiv_{16} 8 + e^2$ for every $e \in S_D$, the arguments in the previous section do not allow us to obtain the full symmetric group Sym_3 and, in fact, generating this group is not possible. More precisely, we shall see that, in these cases, the group of permutations of Prym fixed points consists only on the identity and a transposition. We follow a similar strategy to [Section 4](#) by imposing restrictions for a matrix to act on the Prym fixed points in terms of their relative position on a given surface. Note that $D \equiv_{16} e^2$ for every $e \in S_D$ if and only if $D \equiv_{16} 0$ or 4 , so the remaining cases are precisely those where D is a quadratic residue modulo 16.

9.1. Arithmetic case. Let $D = d^2$ for $d > 4$ an even integer. The Prym locus $E_{d^2}(4)$ consists of $\text{GL}^+(2, \mathbb{R})$ -orbits of square-tiled surfaces (see [Section 4.1](#) for more details on square-tiled surfaces). As in the genus 2 case, a Prym fixed point $w \in X$ can only project down to a 2-torsion point on the unit torus.

Let $e \in S_{d^2}$ and consider the re-scaled prototypical surface $X = Z_D(e)$. This surface is a primitive square-tiled surface as in [Figure 9](#). Moreover, since $d^2 \equiv_{16} e^2$, we have that $d \equiv_4 e$. It follows that $\lambda - e = \frac{d-e}{2}$ is even and, therefore, that w_1 and w_2 project down to $v = i/2 \in \mathbb{C}/\mathbb{Z}[i]$, and that w_3 projects down to $0 \in \mathbb{C}/\mathbb{Z}[i]$. Thus, it is not possible to exchange w_3 with w_1 or w_2 . By [Proposition 8.3](#), we get that $G(X) \cong \text{Sym}_2$.

Since for even D the locus $E_D(4)$ is connected, we get the following.

Proposition 9.1. *Let $d > 4$ be an even integer and let $X \in E_{d^2}(4)$. Then, $G(X) \cong \text{Sym}_2$. Moreover, $G(X)$ is generated by the action of a single parabolic element in $\text{SL}(X)$. \square*

9.2. Non-arithmetic case. Following [Section 4.2](#), we reduce the non-arithmetic case to the arithmetic case. The computations in the genus-three case are much more subtle, so we need some simple results before we are able to state the reduction.

Lemma 9.2. *Let $D > 16$ be an even real quadratic discriminant such that $E_D(4)$ is non-empty. Then, $\frac{1}{2}S_D = R_{D/4}$.*

Proof. Recall first that $D \equiv_4 0$, so $\frac{D}{4}$ is indeed an integer. Likewise, every element of S_D is even, so $\frac{1}{2}S_D$ consists entirely of integers.

Observe that each condition in the definition of $\frac{1}{2}S_D$ translates to a condition in the definition of $R_{D/4}$. More precisely, given $e \in S_D$, we have that $e^2 \equiv_8 D$ if and only if $(\frac{e}{2})^2 \equiv_2 \frac{e}{2} \equiv_2 \frac{D}{4}$. Moreover, the condition $e^2 < D$ is equivalent to $(\frac{e}{2})^2 < \frac{D}{4}$. Finally, $(e+4)^2 < D$ holds if and only if $(\frac{e}{2} + 2)^2 < \frac{D}{4}$. \square

If $D > 16$ is an even quadratic discriminant and $e \in S_D$, then we know that $\mathbb{Z}[\lambda] = \mathbb{O}_D$. The previous lemma shows that, moreover, $\mathbb{Z}[\lambda/2] = \mathbb{O}_{D/4}$. Indeed, we have that

$$\frac{\lambda}{2} = \frac{e + \sqrt{D}}{4} = \frac{e/2 + \sqrt{D/4}}{2}$$

and $\frac{e}{2} \in R_D$. This can be extended to other generators of \mathbb{O}_D as shown by the next lemma:

Lemma 9.3. *Let $D > 0$ be an even real quadratic discriminant and assume that it is a quadratic residue modulo 16, say $D \equiv_{16} d^2$ for some $d \in \mathbb{Z}$. Take $\rho = \frac{\sqrt{D}-d}{2}$. Then, we have that $\mathbb{Z}[\rho/2] = \mathbb{O}_{D/4}$.*

Proof. If $e \in S_D$ and $\lambda = \frac{e+\sqrt{D}}{2}$, we have that $\rho = \lambda - \frac{e+d}{2}$. Note that $e+d \equiv_4 0$. Indeed, we have that $D \equiv_{16} d^2$ and $D \equiv_8 e^2$, which implies that $e^2 \equiv_8 d^2$. Since both e and d are even, this implies that $e \equiv_4 d$, and, again using that they are both even, we obtain that $e+d \equiv_4 0$. Thus,

$$\frac{\rho}{2} = \frac{\lambda}{2} - \frac{e+d}{4} = \frac{e/2 + \sqrt{D/4}}{2} - \frac{e+d}{4} = \frac{\lambda}{2} - \frac{e+d}{4}$$

Since $\frac{e+d}{4}$ an integer, we conclude that $\mathbb{Z}[\rho/2] = \mathbb{O}_{D/4}$. \square

Lemma 9.4. *Let $D > 4$ be even and such that $E_D(4)$ is non-empty. Let $e \in S_D$ and let $A \in \text{SL}(\mathbb{Z}_D(e))$. Then, $A = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ for some $\alpha, \beta \in \mathbb{O}_D$ and $\gamma, \delta \in \mathbb{O}_{D/4}$.*

Proof. Fix $A \in \text{SL}(\mathbb{Z}_D(e))$. We proceed exactly as in [Lemma 4.4](#), which allows us to show that each column of A is a holonomy vector of a saddle connection. As the horizontal sides and the horizontal coordinates of translation vectors of $\mathbb{Z}_D(e)$ belong to \mathbb{O}_D , while the vertical sides and vertical coordinates of translation vectors of $\mathbb{Z}_D(e)$ belong to $\mathbb{O}_{D/4}$, we get the desired result. \square

In what follows, we are interested in the relative position of the regular Prym fixed points in $\mathbb{Z}_D(e)$, with respect to the singularity. In particular, we restrict our attention to elements in $\mathbb{O}_{D/4} \times \frac{1}{2}\mathbb{O}_D$.

For the rest of this section, we assume that D is a quadratic residue modulo 16, with $D \equiv_{16} d^2$. We also fix $\rho = \frac{\sqrt{D}-d}{2}$.

Notation.

- For $A = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$, with $\alpha, \beta \in \mathbb{O}_D$ and $\gamma, \delta \in \mathbb{O}_{D/4}$, we write
 - $\eta = p_\eta + q_\eta \rho \in \mathbb{O}_D$, where $p_\eta, q_\eta \in \mathbb{Z}$ for each $\eta \in \{\alpha, \beta\}$; and
 - $\eta = p_\eta + q_\eta \frac{\rho}{2} \in \mathbb{O}_{D/4}$, where $p_\eta, q_\eta \in \mathbb{Z}$ for each $\eta \in \{\gamma, \delta\}$.

We also write $P_A = \begin{pmatrix} p_\alpha & p_\beta \\ p_\gamma & p_\delta \end{pmatrix}$ and $Q_A = \begin{pmatrix} 2q_\alpha & 2q_\beta \\ q_\gamma & q_\delta \end{pmatrix}$, so $A = P_A + Q_A \frac{\rho}{2}$. Observe that both P_A and Q_A have integer coefficients and, moreover, that $\det(Q_A) \equiv_2 0$.

- For $v \in \mathcal{O}_{D/4} \times \left(\frac{1}{2}\mathcal{O}_D\right)$, we write $v = \left(p_1 + q_1 \frac{\rho}{2}, \frac{p_2}{2} + \frac{q_2}{2} \rho\right)$, where $p_i, q_i \in \mathbb{Z}$ for each $i \in \{1, 2\}$. We also write $p_v = (2p_1, p_2)$ and $q_v = (q_1, q_2)$, so we have $v = \frac{1}{2}(p_v + q_v \rho)$.

Then,

$$\begin{aligned}
 Av &= \frac{1}{2} \left(P_A + Q_A \frac{\rho}{2} \right) (p_v + q_v \rho) = \frac{1}{2} \left(P_A p_v + Q_A q_v \frac{\rho^2}{2} + (P_A q_v + Q_A p_v) \frac{\rho}{2} \right) \\
 &= \frac{1}{2} P_A p_v + Q_A q_v \left(\frac{D - d^2}{16} - \frac{d\rho}{4} \right) + (P_A q_v + Q_A p_v) \frac{\rho}{4} \\
 (3) \quad &= \frac{1}{2} P_A p_v + \frac{D - d^2}{16} Q_A q_v + (P_A q_v + Q_A p_v - d Q_A q_v) \frac{\rho}{4}
 \end{aligned}$$

Lemma 9.5. *Let $D > 0$ be a real quadratic discriminant and assume that it is a quadratic residue modulo 16, say $D \equiv_{16} d^2$ for some $d \in \mathbb{Z}$, and let $e \in S_D$ and $A \in \mathrm{SL}(Z_D(e))$. Take $\rho = \frac{\sqrt{D}-d}{2}$. Then, we have $\mathrm{fr}(Av) = \frac{1}{2} P_A p_v \pmod{1}$ for any $v \in \mathcal{O}_{D/4} \times \left(\frac{1}{2}\mathcal{O}_D\right)$. Moreover, P_A is non-singular modulo 2.*

Proof. By Equation (3), we obtain that the rational part of Av is

$$\frac{1}{2} P_A p_v + \frac{D - d^2}{16} Q_A q_v$$

Since $\frac{D-d^2}{16}$ is an integer, this implies $\mathrm{fr}(Av) = \frac{1}{2} P_A p_v \pmod{1}$. Now, a direct computation shows that

$$1 = \det(A) = \det \left(P_A + Q_A \frac{\rho}{2} \right) = \det(P_A) + \det(Q_A) \frac{\rho^2}{4} + \begin{vmatrix} p_\alpha & p_\beta \\ q_\gamma & q_\delta \end{vmatrix} \frac{\rho}{2} + \begin{vmatrix} q_\alpha & q_\beta \\ p_\gamma & p_\delta \end{vmatrix} \rho.$$

Taking rational parts, we obtain $\det(P_A) + \frac{D-d^2}{16} \det(Q_A) = 1$, so $\det(P_A) \equiv_2 1$ since $\det(Q_A) \equiv_2 0$. \square

In an analogous way to Section 4.2, the previous lemmas show that, when $D \equiv_{16} d^2$, the action of $\mathrm{SL}(Z_D(e))$ on Prym fixed points behaves as in the arithmetic case treated in Section 9.1.

A straightforward computation shows that the fractional part of the rational part of the vectors $v_i \in \mathcal{O}_{D/4} \times \frac{1}{2}\mathcal{O}_D$ on the surface $Z_D(e)$, for $e \in S_D$, are

$$\mathrm{fr}(v_1) = (0, 1/2), \quad \mathrm{fr}(v_2) = (0, 1/2) \quad \text{and} \quad \mathrm{fr}(v_3) = (0, 0),$$

when D is an even quadratic residue modulo 16. It follows that

$$G(Z_D(e)) \leq \mathrm{Sym}(\{1, 2\}).$$

Thus, we get the following result, analogous to Proposition 9.1 in the arithmetic case.

Corollary 9.6. *Let $D \geq 20$ be an even quadratic residue modulo 16 and $X \in E_D(4)$. Then $G(X)$ is conjugate to Sym_2 . Moreover, $G(X)$ is generated by the action of a single parabolic element in $\mathrm{SL}(X)$. \square*

10. WEIERSTRASS PRYM EIGENFORMS IN GENUS THREE OF SMALL DISCRIMINANT

In this section we treat the remaining cases that are not covered by our previous arguments. That is, we focus on the specific cases where $D \in \{8, 17, 25\}$.




FIGURE 10. The surface $B_8(0) \in E_8(4)$ and its Prym fixed points.




FIGURE 11. Re-scaled prototypical surface $Z_{17}(-3)$ and the cylinder decomposition induced by the direction $\arctan(1/2)$.

10.1. **Discriminant 8.** By [Theorem 7.2](#), there are no model A cusps on $E_8(4)$. However, we have the representative $S = B_8(0) \in E_8(4)$, with the corresponding Prym fixed points as shown in [Figure 10](#), as is shown explicitly by Lanneau–Nguyen [[LN14](#), Proposition 4.7 and Figure 6].

Since the moduli of all the horizontal cylinders in S are 1, we get that $(1 \ 2) \in G(S)$. Similarly, the moduli of all the vertical cylinders are 1 as well, and we get $(1 \ 3) \in G(S)$. Thus, $G(S) = \text{Sym}_3$ and, since $E_8(4)$ is connected, we get that $G(X) = \text{Sym}_3$ for every $X \in E_8(4)$. \square

10.2. **Discriminant 17.** We now assume that $D = 17$, so $S_{17} = \{-3, -1\}$. If we take $e = -3$, we get that $r = \frac{D-e^2}{8} = 1$ is odd and, by [Remark 8.4](#), we conclude that $G(A_{17}^+(-3)) = \text{Sym}_3$.

On the other hand, consider the re-scaled prototypical surface $Z_{17}(-3)$ in [Figure 11](#). Recall that this surface lies on the same component as $A_{17}^-(-3)$ (since it is obtained by re-scaling this latter surface). It is then enough to exhibit an explicit parabolic element exchanging w_2 and w_3 , since, by [Lemma 8.1](#), the horizontal twist exchanges w_1 and w_2 .

The cylinder decomposition along $\theta = \arctan(1/2)$ (or along the direction $(2, 1)$) of $Z_{17}(-3)$ is shown in [Figure 11](#). As was done in [Section 5.2](#), we will compute the ratio between the *horizontal* circumference w of a cylinder and its *horizontal* h . Let C_1 be the




FIGURE 12. For $D = 25$, $S = A_{25}^-(-1) \in E_{25}(4)$ is a square-tiled surface and the diagonal direction gives the permutation $(1\ 3) \in G(S)$.

“narrow” cylinder containing w_2 and w_3 in its core curve, and let C_2 be any of the two other cylinders (since they have equal moduli).

We have that the modulus of the cylinder C_1 is

$$m_1 = \frac{(\lambda + 1) + 1 + 2 + \lambda + (2 - \lambda) + \lambda}{1 - \lambda} = \frac{2(\lambda + 3)}{1 - \lambda}$$

and that the modulus of the cylinder C_2 is

$$m_2 = \frac{(\lambda + 1) + 1 + 2}{\lambda} = \frac{\lambda + 4}{\lambda}.$$

Thus, the ratio of the moduli is

$$\frac{m_1}{m_2} = \frac{2\lambda(\lambda + 3)}{(1 - \lambda)(\lambda + 4)}.$$

Since $\lambda = \frac{\sqrt{17}-3}{2}$, a straightforward computation then shows that $\frac{m_1}{m_2} = 2$, so we conclude that $G(Z_{17}(-3)) = \text{Sym}_3$. \square

10.3. Discriminant 25. We start by fixing $D = 25$, so $S_{25} = \{-3, -1\}$. Taking $e = -3$, we get that $r = \frac{D-(e+4)^2}{8} = 3$ is odd. Thus, by [Remark 8.4](#), $G(A_{25}^-(-3)) = \text{Sym}_3$.

On the other hand, $S = A_{25}^-(-1)$ is the primitive square-tiled surface in [Figure 12](#).

The vertical direction on S corresponds to the model A^- reduced prototype $e' = -1$.

Since $r' = \frac{D-e'^2}{8} = 3$ is odd and the core curve of the central vertical cylinder passes through w_2 and w_3 , we get $(2\ 3) \in G(S)$ (cf. [Proposition 8.3](#)).

On the other hand, the diagonal direction (with angle $\pi/4$) on S gives a one-cylinder decomposition whose core curve passes through w_1 and w_3 . It follows that $(1\ 3) \in G(S)$. Thus, $G(S) = \text{Sym}_3$.

Finally, by [Lemma 7.3](#), we know that $A_{25}^-(-1)$ and $A_{25}^-(-3)$ lie on different components, showing that $G(X) = \text{Sym}_3$ for every $X \in E_{25}(4)$. \square

11. CLASSIFICATION OF PERMUTATIONS IN GENUS THREE

We now summarise the results that give the proof of [Theorem 1.3](#).

Proof of Theorem 1.3. In the case where $D > 8$ is odd or a quadratic nonresidue modulo 16, [Lemmas 8.5](#), [8.7](#) and [8.8](#) show that $G(X) = \text{Sym}_3$ for every $X \in E_D(4)$.

In the case where D is an even square, [Proposition 9.1](#) shows that $G(X) \cong \text{Sym}_2$, for every $X \in E_D(4)$. In the non-arithmetic case, when D is an even quadratic residue modulo 16, [Corollary 9.6](#) shows that $G(X) \cong \text{Sym}_2$, for every $X \in E_D(4)$.

The exceptional cases are treated separately: $D = 8$ is treated in [Section 10.1](#), $D = 17$ is treated in [Section 10.2](#), and $D = 25$ is treated in [Section 10.3](#). Thus, $G(X) = \text{Sym}_3$ for every $X \in E_D(4)$, for $D \in \{8, 17, 25\}$. \square

REFERENCES

- [Cal04] K. Calta. “Veech surfaces and complete periodicity in genus two”. In: *J. Amer. Math. Soc.* 17.4 (2004), pp. 871–908. doi: [10.1090/S0894-0347-04-00461-8](https://doi.org/10.1090/S0894-0347-04-00461-8).
- [HL06] P. Hubert and S. Lelièvre. “Prime arithmetic Teichmüller discs in $\mathcal{H}(2)$ ”. In: *Israel J. Math.* 151 (2006), pp. 281–321. doi: [10.1007/BF02777365](https://doi.org/10.1007/BF02777365).
- [Kan06] E. Kani. “The number of genus 2 covers of an elliptic curve”. In: *Manuscripta Math.* 121.1 (2006), pp. 51–80. doi: [10.1007/s00229-006-0012-z](https://doi.org/10.1007/s00229-006-0012-z).
- [LN14] E. Lanneau and D.-M. Nguyen. “Teichmüller curves generated by Weierstrass Prym eigenforms in genus 3 and genus 4”. In: *J. Topol.* 7.2 (2014), pp. 475–522. doi: [10.1112/jtopol/jtt036](https://doi.org/10.1112/jtopol/jtt036).
- [LN20] E. Lanneau and D.-M. Nguyen. “Weierstrass Prym eigenforms in genus four”. In: *J. Inst. Math. Jussieu* 19.6 (2020), pp. 2045–2085. doi: [10.1017/s1474748019000057](https://doi.org/10.1017/s1474748019000057).
- [McM05] C. T. McMullen. “Teichmüller curves in genus two: discriminant and spin”. In: *Math. Ann.* 333.1 (2005), pp. 87–130. doi: [10.1007/s00208-005-0666-y](https://doi.org/10.1007/s00208-005-0666-y).
- [McM06] C. T. McMullen. “Prym varieties and Teichmüller curves”. In: *Duke Math. J.* 133.3 (2006), pp. 569–590. doi: [10.1215/S0012-7094-06-13335-5](https://doi.org/10.1215/S0012-7094-06-13335-5).
- [McM07] C. T. McMullen. “Dynamics of $SL_2(\mathbb{R})$ over moduli space in genus two”. In: *Ann. of Math. (2)* 165.2 (2007), pp. 397–456. doi: [10.4007/annals.2007.165.397](https://doi.org/10.4007/annals.2007.165.397).
- [MMY15] C. Matheus, M. Möller and J.-C. Yoccoz. “A criterion for the simplicity of the Lyapunov spectrum of square-tiled surfaces”. In: *Invent. Math.* 202.1 (2015), pp. 333–425. doi: [10.1007/s00222-014-0565-5](https://doi.org/10.1007/s00222-014-0565-5).
- [Par20] A. Pardo. “A non-varying phenomenon with an application to the wind-tree model”. In: *Int. Math. Res. Not. IMRN* 18 (2020), pp. 5642–5660. doi: [10.1093/imrn/rny188](https://doi.org/10.1093/imrn/rny188).
- [Vee89] W. A. Veech. “Teichmüller curves in moduli space, Eisenstein series and an application to triangular billiards”. In: *Invent. Math.* 97.3 (1989), pp. 553–583. doi: [10.1007/BF01388890](https://doi.org/10.1007/BF01388890).
- [Vee95] W. A. Veech. “Geometric realizations of hyperelliptic curves”. In: *Algorithms, fractals, and dynamics (Okayama/Kyoto, 1992)*. Plenum, New York, 1995, pp. 217–226. doi: [10.1007/978-1-4613-0321-3_19](https://doi.org/10.1007/978-1-4613-0321-3_19).
- [Zor06] A. Zorich. “Flat surfaces”. In: *Frontiers in number theory, physics, and geometry. I*. Springer, Berlin, 2006, pp. 437–583. doi: [10.1007/978-3-540-31347-2_13](https://doi.org/10.1007/978-3-540-31347-2_13).

(Rodolfo Gutiérrez-Romo) CENTRO DE MODELAMIENTO MATEMÁTICO, CNRS-IRL 2807, UNIVERSIDAD DE CHILE, BEAUCHEF 851, SANTIAGO, CHILE.

Email address: g-r@rodol.fo

URL: <http://rodol.fo>

(Angel Pardo) DEPARTAMENTO DE INGENIERÍA MATEMÁTICA & CENTRO DE MODELAMIENTO MATEMÁTICO, CNRS-IRL 2807, UNIVERSIDAD DE CHILE, BEAUCHEF 851, SANTIAGO, CHILE.

Email address: aapardo@dim.uchile.com