

Punctures and p -spin curves from matrix models III : logarithmic potential

Shinobu Hikami

Okinawa Institute of Science and Technology Graduate University, 1919-1
Tancha, Okinawa 904-0495, Japan. e-mail: hikami@oist.jp

Abstract

The intersection numbers for general integer p spin curves of the moduli space $\overline{\mathcal{M}}_{g,n}$ are evaluated from the n point functions in matrix models in Laurent expansions. The results coincide with the previous values derived from the recursive p -th KdV equation. The extension to the half-integers $p = \frac{1}{2}$, $p = -\frac{1}{2}$ and to the negative integer $p = -2, -3$ cases are investigated in this formulation. The strong coupling expansions with a logarithmic potential are examined by Laplace-Borel transformation for the character expansions. The intersection numbers of D types of the ADE singularities are derived by this matrix model.

Contents

1. Introduction.
2. One point function for the A_l type.
3. One point function for the D_l type.
4. One point function for the A_l type of non-positive integer p -spin curves.
5. n -points function for the A_l type of positive integer p -spin curves.
6. String equation, dilaton equation and WDVV equation.
7. n -point functions of $p = \frac{1}{2}, -\frac{1}{2}, -\frac{3}{2}$ and $-2, -3$.
- 7-1: n -point function for $p = \frac{1}{2}$.
- 7-2: n -point function for $p = -\frac{1}{2}$.
- 7-3: Strong coupling of $p = -2$.
- 7-4: Strong coupling of $p = -\frac{3}{2}$.
- 7-5: Strong coupling of $p = -3$.
8. General p and recursion equation.
9. Characteristic polynomials and p -spin curves.
10. Summary and discussions.

1 Introduction

It is well known that the generating function of the intersection numbers on moduli space of p spin stable curves becomes the τ function of the generalized KdV hierarchies (p -reduced KP hierarchies), which are related to a two dimensional gravity [1, 2]. There are now many studies of the intersection numbers including Gelfand-Dikii pseudo differential equation [3, 4]. We have proposed a method of the calculation for this intersection numbers by a generalized Kontsevich matrix model, which was derived by the duality and replica method [5, 6, 7]. For one point function, an expression for single intersection numbers has been derived explicitly [5, 7].

For the purpose of the extension to half integer spin p , we reformulated this integral expressions by a new change of variables, and we have investigated the intersection numbers of Ramond punctures in previous articles I, II [8, 9].

This reformulation enables us to obtain easily the intersection numbers for integer p (Neveu-Schwarz punctures) for n point functions, which should be consistent with the results obtained by the recursive method [10, 11, 13] due to Gelfand-Dikii equation. The evaluation of several marked points in general p and for genus g was obtained in the recursive calculations [11], and we show in this article that our method of the Laurent expansion agrees with them for the lower orders, especially for three point function.

The ADE singularities characterized by Dynkin diagrams are well known. The D_l singularity is represented by $y^{l-1} + yx^2 + z^2 = 0$ ($l \geq 4$). The Coxeter number defined by $p = 2l - 2$ in D_l singularity can be interpreted as a spin p . The weight system of D_l is known as $(a, b, c; h) = (2, l - 2, l - 1; 2(l - 1))$. The simply laced Lie algebra, A_l, B_l, C_l, D_l , are related to classical Lie groups $SU(l+1), SO(2l+1), SP(l), SO(2l)$, respectively. We have discussed the generalized Airy matrix model with an external source by the use of HarishChandra theorem for the cases A_l, B_l, C_l, D_l cases, where HarishChandra theorem can be applied for the matrix models with external sources [7, 14]. These matrix models were applied for the non-orientable surfaces or Klein surfaces. The D_l singularity for integrable hierarchy has been studied and the intersection numbers are deduced

from the integrable hierarchy [17].

Recently open intersection numbers were discussed based on the logarithmic matrix models [18, 19, 21, 25, 22, 23, 26, 27]. The open intersection theory means the extension of the intersection theory of Riemann surface to open Riemann surface, which has boundaries [21, 25]. The boundaries are represented by the logarithmic terms, which is analogous to the D brane. The matrix model with a logarithmic potential for the open intersection theory is written for general p (generalized Airy matrix model with a logarithmic potential) [18],

$$Z = \int dB e^{-\frac{c}{p+1}\text{tr}B^{p+1} + k\text{tr}\log B + \text{tr}B\Lambda} \quad (1.1)$$

where B is a Hermitian matrix and k is a parameter. The matrix Λ is an external source. This model is called as Kontsevich-Penner model for $p = 2$. When $k = 0$ and $p = 2$, it reduces to Kontsevich Airy matrix model. When $c = 0$, it reduces to Penner model, which provides Euler characteristics for orbifold. The spin p is related to l as $p = 2l - 2$ for D_l type singularity. We investigated before the case of Lie algebra $so(N)$, $sp(N)$ and found that the correspondent $u(s)$ has a logarithmic terms [14], which makes a difference from A_l type. These cases were discussed as a manifestation of non-orientable surface (Klein surface) further in [19, 7]. In this paper, we explicitly show that the intersection numbers of D_l type is derived from the one point function $u(s)$, which has a logarithmic term. By the Gaussian integral of x for $y^{l-1} + x^2y + z^2$, we have $1/\sqrt{y}$ which turns to be a logarithmic potential by the exponentiation. This may give simple explanation of the appearance of the logarithmic term in the matrix model of D_l singularity. In general, an oscillating integrals of n -variables have asymptotic expansions with logarithmic terms, related to Newton polygon [20] and it is well known that such expansion is related to the resolution of the singularities. In D_l types, the simple logarithmic term appears. The D_l type is related to a real algebraic curves and the Euler characteristics χ of a real algebraic curve is obtained in the case of $p = -1$ in D_l type ($p = 2l - 2$) [34, 14].

It is known that A_5, A_4, A_3 ($p=6,5,4$) singularities correspond to Ashkin-Teller model, 3-state Potts model and Ising model respectively [28, 29]. The A_{p-1} singularity has a central charge $C = 2 - 6/p$. The central charge $C = \frac{1}{2}$ and critical exponent of the energy ν are consistent with the values of Ising model. The anomalous conformal dimensions are $\Delta_\epsilon = \frac{4}{p}$, $\Delta_\phi = \frac{p-3}{p(p-2)}$. These dimensions and central charges agree with the well-known values of Ising ($p = 4$) and 3-state Potts model ($p = 5$).

For D_l type singularities, the central charge is also given by $C = 2 - \frac{6}{p}$, where p is Coxeter number $p = 2l - 2$. The interesting applications are found in the condensed matter physics for the topological excitation of electron at the boundaries, as Majorana fermions [25], and the edge excitation on the boundary as Quantum Hall effect, for instance. Furthermore, in D_4 singularity, there are intriguing Ramond sector [33], which is related to the vanishing relation for one point function of $g = 2 + 3m$, $m \in \mathbb{Z}$.

The spin p takes a value of positive integer according to the singularity theory and the central charge $C = 2 - \frac{6}{p}$. However, as discussed for $p = -1$, Euler characteristic is obtained from the continuation from positive integer to $p = -1$ in the expression of the intersection numbers. The cases of $p = \frac{1}{2}$ and $p = \frac{3}{2}$, which are fractional spins, have been discussed in our previous papers [8, 9]. These cases correspond to "fermion", and the genus expansion of one point function agree with the selection rule due to Riemann-Roch theorem. As

a conformal field theory (CFT) in two dimensions, $p = \frac{3}{2}$ case exists as $\beta - \gamma$ system in the supersymmetric non-linear sigma model, which also corresponds to spanning forrest model with a central charge $C = -2$. There is an interesting observation that for $p = \frac{1}{2}$ case, the tautological relations become simple [35]. In this paper, in addition to the results of I,II [8, 9], we investigate the m point correlation functions of $p = \frac{1}{2}$, and find that the punctures of Ramond type appear in a pairwise as same as $p = -2$ case.

The case $p = -2$ corresponds to the unitary matrix model of the lattice gauge theory. The strong coupling region of this case has a character expansion. We consider this character expansion [31] for the negative p case, $p = -2, -3, \dots$ by the n -point function of $U(s_1, \dots, s_n) = \langle \prod_i \text{tr} e^{s_i M} \rangle$, which can be interpreted as Wilson loops. This unitary matrix model is related to Chern-Simons theory, which describes torus links. Therefore, p spin curves for the negative cases are related to the knot theory, and the ADE type singularities. In this paper, we consider the factorization of the characters through Laplace-Borel transformations of $U(s_1, \dots, s_n)$ with a logarithmic potential. The Laplace transform of $U(s_1, \dots, s_n)$ with a logarithmic term is related to Hilbert-Poincare characteristic polynomials. The Laurent expansion about y variable is similar to knot polynomials, particularly Alexander polynomials in the knot theory.

The organization of this article is following: We investigate one point intersection numbers of p spin curves for general genus g in section 2 for A_l . In section 3, one point functions of D_l type is evaluated. In section 4, one point function for non-integer p case is discussed for $p = \frac{3}{2}, \frac{1}{2}, -\frac{1}{2}, -1, -2$ both for A type and D type. In section 5, the intersection numbers for the positive integers are explicitly derived up to four marked points in many examples. We confirm the intersection numbers, which are obtained by the integral formula, agree with the values derived through recursion relations. In section 6, we discuss the relations of string equation, dilaton equation and WDVV equation. We prove the string equation and the dilaton equation for the positive integer p by the n -point correlation function of the matrix models. The section 7 is devoted to the evaluations of $p = \frac{1}{2}, -\frac{1}{2}$ and $p = -2$ cases for multi-correlation functions. In section 8, general p spin curves for the negative integer case is discussed for the strong coupling regions and the relation to the characters of $U(N)$ is discussed. The relation of the Laurent polynomials of y of p spin curves to knot polynomials is further discussed in the section 9. In the section 10, we give summary and discussions the future problems.

2 One point function for A_l type

Since the p spin curves of the moduli space has a correspondence to A_{p-1} singularity by mirror symmetry, we use the terminology of A_l type for the $l = p - 1$ spin curves, which distinguishes the case D type.

For A_l case, the generating function of the intersection number for one marked point is expressed as [5, 6]

$$\begin{aligned} u(s) &= \langle \text{tr} e^{sB} \rangle \\ &= \frac{1}{s} \int \frac{du}{2i\pi} e^{-\frac{c}{p+1}[(u+\frac{s}{2})^{p+1} - (u-\frac{s}{2})^{p+1}]} \end{aligned} \quad (2.1)$$

The generating function for the intersection numbers of A_l ($p = l + 1$) is evalu-

ated for small s by the replacement $u = (\frac{t}{cs})^{1/p}$,

$$u(s) = \frac{1}{sp\pi} \cdot \frac{1}{(cs)^{1/p}} \int_0^\infty dt t^{\frac{1}{p}-1} e^{-t} \\ \times e^{-\frac{p(p-1)}{3!4} s^{2+\frac{2}{p}} c^{\frac{2}{p}} t^{1-\frac{2}{p}} - \frac{p(p-1)(p-2)(p-3)}{5!4^2} s^{4+\frac{4}{p}} c^{\frac{4}{p}} t^{1-\frac{4}{p}} + \dots} \quad (2.2)$$

This leads to

$$u(s) = \frac{1}{c^{\frac{1}{p}} s^{1+\frac{1}{p}} \pi} [\Gamma(1 + \frac{1}{p}) - \frac{p-1}{24} z \Gamma(1 - \frac{1}{p}) \\ + \frac{(p-1)(2p+1)(p-3)}{5760} z^2 \Gamma(1 - \frac{3}{p}) + \dots] \quad (2.3)$$

where $z = c^{\frac{2}{p}} s^{2+\frac{2}{p}}$. Writing this expansion with the intersection numbers $\langle \tau_{n,j} \rangle_g$ (n is integer, and spin component $j = 0, 1, 2, \dots, p-2$) as

$$u(s) = \sum_g \langle \tau_{n,j} \rangle_g \frac{1}{\pi} \Gamma(1 - \frac{j+1}{p}) c^{\frac{2g-1}{p}} p^{g-1} s^{(2g-1)(1+\frac{1}{p})}, \quad (2.4)$$

one obtain the intersection numbers as a polynomial of p . We have a relation of n and j for non-vanishing intersection numbers,

$$(2g-1)(1 + \frac{1}{p}) = n + \frac{j+1}{p} \quad (2.5)$$

which comes from the Riemann-Roch relation (RR) for \tilde{s} marked points by $\tilde{s} = 1$

$$3g - 3 + \tilde{s} = \sum_{i=1}^{\tilde{s}} n_i + (g-1)(1 - \frac{2}{p}) + \frac{1}{p} \sum_{i=1}^{\tilde{s}} j_i \quad (2.6)$$

Thus the s dependence appears for one point case ($\tilde{s} = 1$) as a power $s^{(2g-1)(1+\frac{1}{p})} = s^{n+\frac{j+1}{p}}$ in (2.4). The intersection numbers (in the case of $p = 2$), are given by the first Chern class c_1 or ψ , as

$$\langle \tau_{n_1} \dots \tau_{n_s} \rangle_g = \int_{\bar{\mathcal{M}}_{g,s}} \psi^{n_1} \dots \psi^{n_s} \quad (2.7)$$

where $\bar{\mathcal{M}}_{g,s}$ is a compactified moduli space with s marked points on genus g Riemann surface. The intersection numbers of one point $\tau_{A_i}(g)$ (Coxeter

number $p= l+1$) is thus given as [8, 9]

$$\begin{aligned}
\langle \tau_{n,j} \rangle_{g=1} &= \frac{p-1}{24} \\
\langle \tau_{n,j} \rangle_{g=2} &= \frac{(p-1)(2p+1)(p-3)}{p \cdot 5! \cdot 4^2 \cdot 3} \frac{\Gamma(1 - \frac{3}{p})}{\Gamma(1 - \frac{1+j}{p})} \\
\langle \tau_{n,j} \rangle_{g=3} &= \frac{(p-1)(2p+1)(p-5)(8p^2 - 13p - 13)}{p^2 \cdot 7! \cdot 4^3 \cdot 3^2} \frac{\Gamma(1 - \frac{5}{p})}{\Gamma(1 - \frac{1+j}{p})} \\
\langle \tau_{n,j} \rangle_{g=4} &= \frac{(p-1)(2p+1)(p-7)(72p^4 - 298p^3 - 17p^2 + 562p + 281)}{p^3 \cdot 9!4^4 \cdot 15} \\
&\quad \times \frac{\Gamma(1 - \frac{7}{p})}{\Gamma(1 - \frac{1+j}{p})} \\
\langle \tau_{n,j} \rangle_{g=5} &= (p-1)(2p+1)(p-3)(p-9)(4p+3)(32p^4 - 162p^3 + p^2 \\
&\quad + 326p + 163) \frac{1}{p^4 11!4^5 3} \frac{\Gamma(1 - \frac{9}{p})}{\Gamma(1 - \frac{1+j}{p})} \\
\langle \tau_{n,j} \rangle_{g=6} &= (p-1)(2p+1)(p-11)(530688p^8 - 5830544p^7 + 16589332p^6 \\
&\quad + 8955300p^5 - 65056373p^4 - 26944928p^3 + 85178190p^2 \\
&\quad + 80708428p + 20177107) \frac{1}{p^5 \cdot 13!7 \cdot 5 \cdot 4^6 3^3} \frac{\Gamma(1 - \frac{11}{p})}{\Gamma(1 - \frac{1+j}{p})} \quad (2.8)
\end{aligned}$$

where $\langle \tau_{n,j} \rangle_{g=1} = \langle \tau_{1,0} \rangle_{g=1}$ [6]. In [7], $\langle \tau_{n,j} \rangle$ up to $g = 9$ is evaluated.

When $p = 2$ (Kontsevich model), it leads to a simple expression,

$$\langle \tau_{3g-2} \rangle = \frac{1}{g!(24)^g} \quad (2.9)$$

There appear interesting vanishing relations for $\langle \tau_{A_l} \rangle$. For instance, at $p = 3$ case, $\langle \tau_{A_l} \rangle$ are vanishing at $g = 2 + 3m$ ($m \in \mathbb{Z}$). For $p = 5$, $\langle \tau_{A_l} \rangle$ are vanishing at $g = 3 + 5m$ ($m \in \mathbb{Z}$). In general odd integer p , $\langle \tau_{A_l} \rangle$ are vanishing at $g = (p+1)/2 + pm$ ($m \in \mathbb{Z}$). Some of these relations can be seen in the expressions of $\langle \tau_{A_l} \rangle$ up to genus $g = 9$ in [7]. We will see later that D_4 ($p = 6$) type has this periodicity of the vanishing relation at $g = 2 + 3m$ ($m \in \mathbb{Z}$) for D_4 .

For the case A_2 ($p = 3$), one point function $u(s)$ is given by the Airy function as [6]

$$\begin{aligned}
u(s) &= \frac{1}{Ns(3Ns)^{1/3}} A_i(\zeta) \\
&= \frac{1}{Ns(3Ns)^{1/3}} \left[A_i(0) \left(1 + \frac{1}{3!} \zeta^3 + \frac{1 \cdot 4}{6!} \zeta^6 + \frac{1 \cdot 4 \cdot 7}{9!} \zeta^9 + \dots \right) \right. \\
&\quad \left. + A_i'(0) \left(\zeta + \frac{2}{4!} \zeta^4 + \frac{2 \cdot 5}{7!} \zeta^7 + \frac{2 \cdot 5 \cdot 8}{10!} \zeta^{10} + \dots \right) \right] \quad (2.10)
\end{aligned}$$

where $\zeta = -N^{2/3}(4 \cdot 3^{1/3})^{-1} s^{8/3}$, $A_i(0) = 3^{-2/3}/\Gamma(\frac{2}{3})$ and $A_i'(0) = -3^{-1/3}/\Gamma(\frac{2}{3})$. This Airy function leads to the intersection numbers of

$$\langle \tau_{\frac{8g-5-j}{3}, j} \rangle_g = \frac{1}{(12)^g g!} \frac{\Gamma(\frac{g+1}{3})}{\Gamma(\frac{2-j}{3})} \quad (2.11)$$

which shows the vanishing relations for $g = 2, 5, 8, \dots$ ($g = 2 + 3m$, $m \in \mathbb{Z}$), for such case the value of spin j takes 2. The absence of $g = 2, 5, 8, \dots$ is due to Stokes phenomena.

For $p = 4$, $u(s)$ is written by the Bessel function, [7]

$$\begin{aligned} u(s) &= \frac{1}{2\sqrt{8}} e^{\frac{3}{160}s^5} \frac{1}{2\sin(\frac{\pi}{4})} \left[I_{-\frac{1}{4}}\left(\frac{1}{32}s^5\right) + I_{\frac{1}{4}}\left(\frac{1}{32}s^5\right) \right] \\ &= \frac{1}{8} \sum_{m,n=0}^{\infty} \frac{1}{m!n!\Gamma(n + \frac{5}{4})} \left(\frac{3}{160}\right)^m \left(\frac{1}{64}\right)^{2n+\frac{1}{4}} s^{5m+10n+\frac{1}{4}} \quad (2.12) \end{aligned}$$

We have Riemann-Roch relation (RR) of (2.6) for the s-point intersection numbers $\langle \tau_{n_1, j_1} \tau_{n_2, j_2} \cdots \tau_{n_s, j_s} \rangle_g$ [1]. The factor $(1 - \frac{2}{p})$ is a central charge $\hat{c} = \frac{p-2}{p}$. The last term is also charge for p spin curves. Note this central charge \hat{c} is also valid for D_l singularity, since the weight system of D_l singularity $W = y^{l-1} + xy^2 + z^2 = 0$ ($l \geq 4$). The weight system is $(a, b, c; h) = (2, l-2, l-1; 2(l-1))$, where h is called as Coxeter. For D_l case, we use the spin p value for Coxeter number h , which is related to l as $p = 2l - 2$. For D_4 , we have spin curve of $p = 6$.

3 One point function for D_l type

The singularity theory of D_l type is described as a two dimensional normal singularity by the equation of $x^{l-1} + xy^2 + z^2 = 0$ ($l \geq 4$). The weight system is $(a, b, c; h) = (2, l-2, l-1; 2(l-1))$, where h is called as Coxeter. For D_l case, we use the spin p value for Coxeter number h , which is related to l as $p = 2l - 2$. For D_4 , we have spin curve of $p = 6$.

The intersection numbers are extended from (2.7) to the one includes the boundary. The tangent bundle is trivially on the boundary of the moduli space. We need analogous term for the first Chern class on the boundary and introduce the correspondent quantity σ for the boundary [25].

$$\langle \tau_{n_1} \cdots \tau_{n_s} \sigma^m \rangle = \int_{\mathcal{M}} \psi^{n_1} \cdots \psi^{n_s} \quad (3.1)$$

where m punctures on the boundary are added to (2.7). This is generalization to open intersection numbers, and related to D_l singularity as we will see. Instead of working of the geometrical moduli space \mathcal{M} , we study the equivalent partition function of a matrix model as same as A_l case. The partition function of a matrix model is expressed by the n point correlation function $u(s_1, s_2, \dots, s_n)$ with a logarithmic potential [7].

We have the following one point function $\hat{u}(s)$ in the integral form for D_l ,

Theorem 2 (D_l)

The generating function $\hat{u}(s)$ of the intersection numbers $\langle \tau \rangle$ for D_l ($p = 2l - 2$) is given by

$$\hat{u}(s) = \frac{1}{s} \int_0^\infty du e^{-\frac{c}{(p+1)}[(u+\frac{s}{2})^{p+1} - (u-\frac{s}{2})^{p+1}]} \frac{1}{\sqrt{1 - \frac{s^2}{4u^2}}} \quad (3.2)$$

where $c = \frac{N}{p-1} \sum \frac{1}{a_\alpha^{p+1}}$ as shown in [6]. a_α is eigenvalues of the external source.

The last factor is absent for A_l case. It is written as

$$\begin{aligned} \frac{1}{\sqrt{1 - \frac{s^2}{4u^2}}} &= \frac{1}{2} \left(\sqrt{\frac{u + \frac{s}{2}}{u - \frac{s}{2}}} + \sqrt{\frac{u - \frac{s}{2}}{u + \frac{s}{2}}} \right) \\ &= \frac{1}{2} \left(e^{\frac{1}{2} \log(u - \frac{s}{2}) - \frac{1}{2} \log(u + \frac{s}{2})} + e^{\frac{1}{2} \log(u + \frac{s}{2}) - \frac{1}{2} \log(u - \frac{s}{2})} \right) \end{aligned} \quad (3.3)$$

Note that if we change $u \rightarrow -u$, above term is invariant. If we write the coefficient of the logarithm as k instead of $\frac{1}{2}$, two terms are k and $-k$ coefficient, and it leads to the polynomial of even power of k . This characterizes the D_l type as we will discuss later.

The small s expansion of $u(s)$ with $u = t^{\frac{1}{p}}$, becomes

$$\begin{aligned} \hat{u}(s) &= \frac{1}{N\pi s} \frac{1}{(cs)^{1/p}} \int_0^\infty dt t^{\frac{1}{p}-1} e^{-t} \left[1 - \frac{p(p-1)}{24} s^2 (cs)^{2/p} t^{1-\frac{2}{p}} \right. \\ &\quad - \frac{p(p-1)(p-2)(p-3)}{5! \cdot 16} s^4 (cs)^{4/p} t^{1-\frac{4}{p}} + \frac{p^2(p-1)^2}{2 \cdot (24)^2} s^4 (cs)^{4/p} t^{2-\frac{4}{p}} \\ &\quad - \frac{p^3(p-1)^3}{3!(24)^3} s^6 (cs)^{6/p} t^{3-\frac{6}{p}} + \frac{p^2(p-1)^2(p-2)(p-3)}{5! \cdot 14 \cdot 16} s^6 (cs)^{6/p} t^{2-\frac{6}{p}} \\ &\quad - \frac{p^3(p-1)^3}{3!(24)^3} s^6 (cs)^{6/p} t^{3-\frac{6}{p}} \\ &\quad \left. - \frac{p(p-1)(p-2)(p-3)(p-4)(p-5)}{7!4^3} s^6 (cs)^{6/p} t^{1-\frac{6}{p}} + \dots \right] \\ &\times \left[1 + \frac{1}{8} s^2 (cs)^{2/p} t^{-\frac{2}{p}} + \frac{3}{128} s^4 (cs)^{4/p} t^{-\frac{4}{p}} + \frac{15}{3072} s^6 (cs)^{6/p} t^{-\frac{6}{p}} + \dots \right] \end{aligned} \quad (3.4)$$

The last factor is the expansion of (3.3). The integer power of s , denoted as m , shows the relation to genus g as $2g - 1 = m$.

The small s expansion of $\hat{u}(s)$, with the normalization factors $\frac{1}{p^{g-1}}$ and gamma factor $1/\Gamma(1 - \frac{1+j}{p})$, gives the intersection numbers of one point case,

$$\begin{aligned} \langle \tau_{1,0} \rangle_{g=1} &= \frac{p+2}{24} \\ \langle \tau_{n,j} \rangle_{g=2} &= \frac{(p+2)(2p+1)(p-6)}{5760p} \frac{\Gamma(1 - \frac{3}{p})}{\Gamma(1 - \frac{1+j}{p})} \\ \langle \tau_{n,j} \rangle_{g=3} &= \frac{(p+2)(2p+1)(8p^3 - 77p^2 + 196p + 188)}{2903040p^2} \frac{\Gamma(1 - \frac{5}{p})}{\Gamma(1 - \frac{1+j}{p})} \\ \langle \tau_{n,j} \rangle_{g=4} &= \frac{(p+2)(2p+1)(4p^2 - 27p - 22)(18p^3 - 133p^2 + 308p + 332)}{1393459200p^3} \\ &\quad \times \frac{\Gamma(1 - \frac{7}{p})}{\Gamma(1 - \frac{1+j}{p})} \\ \langle \tau_{n,j} \rangle_{g=5} &= \frac{(p+2)(2p+1)(4p+3)(p-6)}{122624409600p^4} (32p^5 - 450p^4 + 1741p^3 \\ &\quad - 1642p^2 - 6788p - 3288) \frac{\Gamma(1 - \frac{9}{p})}{\Gamma(1 - \frac{1+j}{p})} \end{aligned} \quad (3.5)$$

As a check, we find for $p = 2$, the explicit expression for one point function as ($j = 0$ for $p = 2$)

$$\langle \tau_{3g-2} \rangle = \frac{1}{g!6^g} \quad (3.6)$$

which leads to $\frac{1}{6}$, $\frac{1}{72}$, and $\frac{1}{1296} = \frac{1}{316^3}$, for $g = 1, 2, 3$, respectively. The derivation of this result will be provided later in (4.7).

There is a relation to the open intersection numbers. The logarithmic potential with a coefficient k has been studied so called as Kontsevich-Penner model or Airy matrix model with a logarithmic potential [18, 19, 22]. Putting $p = 2$ and $k = \frac{1}{2}$ in the expressions of (1.1) [18, 22], the results agree with (3.5). For instance, $\langle \tau_{1,0} \rangle_{g=1} = \frac{p+2}{24}$ becomes $\frac{1}{6}$ for $p = 2$, which agrees with $\langle \tau_1 \rangle = \frac{1+12k^2}{24} = \frac{1}{6}$ with $k = \frac{1}{2}$ in [18, 22]. $\langle \tau_{n,j} \rangle_{g=2} = \frac{1}{72}$ for $p = 2$ agrees with $\langle \tau_4 \rangle = \frac{1+56k^2+16k^4}{1152} = \frac{1}{72}$ for $k = \frac{1}{2}$. For non orientable surface, taking account of projective plane ($g = \frac{1}{2}$), Klein surface ($g = 1$), and crosscap ($g = \frac{3}{2}$), genus g is considered as a fractional (double genus) [14]. Thus for the comparison with the expressions of $\langle \tau_{\frac{3g'-1}{2}} \rangle$ in [19, 22], we need a relation $g' = 2g - 1$ i.e. the one point function of (3.5) corresponds to $\langle \tau_{\frac{3g'-1}{2}} \rangle$ in [19, 22], and the result completely agrees for the case of integer $\frac{3g'-1}{2}$. The case $p = 2$ in D_l type means D_2 , so D_2 case has a meaning of the Kontsevich-Penner model as an open intersection theory, although $l \geq 4$ case is discussed in the singularity theory. The terms of $\langle \tau_{\frac{3g'-1}{2}} \rangle$, which appear with half-integer $\frac{3g'-1}{2}$, have expressions of odd k polynomial, and they do not appear in D_l type in (3.5).

For the open intersection numbers, there appear intersection numbers such that $\langle \tau_{\frac{5}{2}} \rangle = \frac{1}{12}(k + k^3)$. These are odd power of k . As we observed in (4.7), we have sum of k and $-k$ for the logarithmic correction, therefore adding these two contributions, they are cancelled for such $\langle \tau_{\frac{5}{2}} \rangle$ intersection numbers. This is a reason that we have no half integer indexed intersection numbers in D_l type.

As noted in [18], such odd k intersection numbers with half integer indexed τ are evaluated by the contour integral, and therefore can be regarded as Ramond sectors.

Thus the intersection numbers belong to the orientable surfaces, which do not include the case of half integer $\frac{3g'-1}{2}$. For general p , if we put $k = \frac{1}{2}$ in [18], we find also the agreement with (3.5).

For $p = 6$, which corresponds to D_4 , the intersection numbers become $\tau_{D_4} = 1, \frac{1}{2}, 0, \frac{1}{40824}, \frac{13}{122472}, 0$ for $g = 0, 1, 2, 3, 4, 5$, respectively. These values are obtained from (3.5) as the coefficients of Gamma function factors. They agree with [17]. It is remarkable that τ_{D_4} is vanishing at $g = 2, 5$, since a factor $(p-6)$ appears in (3.5). This suggests τ_{D_4} ($p = 6$ case) is vanishing periodically at $g = 2 + 3k$, $k \in \mathbb{Z}$.

This periodic vanishing reallation of the intersection numbers is due to the selection rule for spin p . We have from Riemann-Roch formula the relation between the spin p and the genus g ,

$$(p+1)(2g-1) = pn + j + 1 \quad (3.7)$$

where $j = 0, 1, 2, \dots, p-2$. For the singularity theory, D_l has a relation

$$2g-1 = pn' + m_\alpha \quad (3.8)$$

where m_α is defined by the characteristic polynomial $\chi(t)$ for the weight system $(a, b, c; h)$ as

$$\chi(t) = \frac{1}{t^h} \frac{(t^h - t^a)(t^h - t^b)(t^h - t^c)}{(t^a - 1)(t^b - 1)(t^c - 1)} = \sum_{\alpha=1}^l t^{m_\alpha} \quad (3.9)$$

with l is Milnor number. j is $m_\alpha - 1$, which is the spin components of spin p .

For D_4 , the weight system becomes $(2, 2, 3; 6)$, and $m_\alpha = 1, 3, 5$ ($m_2 = m_3 = 3$, double). The relation of (3.7) is written by $n' = n + 1$, $j = m_\alpha - 1$ in (3.8). For $p = 6$ in D_4 , $g = 1$ corresponds to $m_\alpha = 1$ ($j=0, n=1$). $g = 2$ corresponds to $m_\alpha = 3$ ($j = 2, n = 3$). $g = 5, g = 8$ correspond to $m_\alpha = 3$. Thus for $g = 2 + 3k$ ($k \in \mathbb{Z}$), the exponent becomes $m_\alpha = 3$, which is doubled. This correspondence is related to the vanishing relation of the intersection numbers of D_4 for $g = 2 + 3k$.

From (3.5), It is easy to check the cases; (i) large p behavior is same as (2.8), (ii) the intersection numbers of $p = -1/2$ are vanishing for $g > 1$ for all order of g , (iii) the intersection numbers of $p = -2$ are vanishing for all genus $g > 0$.

Since we have derived exact one point function of D_l type, it may be interesting to consider the negative values of p and half-integer p as discussed in A_l type [7, 8, 9]. In the next section, we examine these non positive integer values of p in details. Interesting applications of non-positive integer cases were discussed in [9] for A_l type.

4 One point function of D_l type for the non-positive integer cases

We have discussed the non-positive integer spin p for one marked point of the A type in the previous articles [8, 9]. Here we extend these results of A type to D type for one marked point.

For D_l singularity, the relation of $p = 2l - 2$ gives the constraint that spin p is even integer. Since the intersection numbers are expressed by the polynomial of p as (3.5), the analytic continuation of p to the general values including the non-integer case may be possible. Our formulation of a matrix model allows the non-positive integer value of p .

• $p = 1$ case in D_l

The case $p = 1$ in D_l singularity corresponds to $D_{\frac{3}{2}}$ by the relation $p = 2l - 2$. The non positive integer l of D_l is out of the definition in the standard singularity theory. Here we consider a matrix model with general real power exponent p with a logarithmic term, and we notate this case as $D_{\frac{3}{2}}$ for convenience.

As discussed in [8, 33], there is no Ramond contribution for A_l case in the positive integer p . However, in D_l , there are Ramond punctures [9, 33]. Here we identify the Ramond contribution by the residue of $y = 0$ in the following integral representation. We use a change of variable from u to y following the discussion of [8, 9] as

$$u = \frac{i}{2} \left(y^2 - \frac{1}{y^2} \right) \quad (4.1)$$

The factor of D_l logarithmic potential becomes by the change of variable of (4.1),

$$\frac{u}{\sqrt{u^2-1}} = \frac{y - \frac{1}{y^3}}{y + \frac{1}{y^3}} \quad (4.2)$$

The measure is $du = i(y + 1/y^3)dy$, which is cancelled by the denominator of (4.2).

For instance $p = 1$ case of A_l ,

$$\begin{aligned} u(s) &= \frac{i}{2} \oint \frac{dy}{2i\pi} (y + \frac{1}{y^3}) e^{-\frac{i}{4}s(y^2 - \frac{1}{y^2})} \\ &= \frac{i}{2} \oint \frac{dy}{2i\pi} (-\frac{2}{is}) \frac{d}{dy} e^{-\frac{i}{4}s(y^2 - \frac{1}{y^2})} \\ &= 0 \end{aligned} \quad (4.3)$$

which becomes vanishing by the total derivative. This is consistent with a factor $(p-1)$ in all order in (2.8). For D_l case, instead, we take a factor $(y - \frac{1}{y^3})$ instead of $(y + \frac{1}{y^3})$,

$$\begin{aligned} u(s) &= \frac{i}{2} \oint \frac{dy}{2i\pi} [2y - (y + \frac{1}{y^3})] e^{-\frac{i}{4}s(y^2 - \frac{1}{y^2})} \\ &= \frac{i}{2} \oint \frac{dt}{2i\pi} t e^{-\frac{i}{4}s(t - \frac{1}{t})} \\ &= - \sum_{n=0}^{\infty} \frac{1}{2^{4n+3} n! (n+1)!} s^{2n+1} \end{aligned} \quad (4.4)$$

This is consistent with the expression of (3.5) for $p = 1$. The genus g is equal to $n + 1$, and we have taken $cs = 1$ in (2.4) for simplicity.

• $p=2$ case and open intersection numbers

In the previous works of open intersection numbers, the logarithmic potential has been considered. For $p = 2$ case, with the logarithmic potential (Eq.(116) of [8]),

$$u(s) = \frac{1}{s} \oint \frac{du}{2i\pi} e^{-\frac{c}{3}[(u+\frac{c}{2})^3 - (u-\frac{c}{2})^3] + k \log(u+\frac{c}{2}) - k \log(u-\frac{c}{2})} \quad (4.5)$$

The one point function $\hat{u}(s)$ of (3.2) is equal to $u(s)|_{m=\frac{1}{2}} + u(s)|_{m=-\frac{1}{2}}$. Indeed we have intersection number of $p = 2$ as $\langle \tau_D \rangle = \frac{p+2}{24} = \frac{1}{6}$. The same number is obtained from the generalized open intersection number $\langle \tau_1 \rangle_{open} = (1 + 12k^2)/24 = \frac{1}{6}$ by putting $k = \pm \frac{1}{2}$. Note the open intersection number is described by the logarithmic potential with $k = 1$ [18, 19, 6, 21, 26, 27]. When $p = 2$, the generalized open intersection numbers are evaluated in higher orders [22], in which a parameter N is same as our k . The odd power terms are cancelled by adding $\pm k$ contributions, and the results agree with D_l type intersection numbers with $k = \frac{1}{2}$.

For $p = 2$, using the representation of y , one point function $u(s)$ is expressed as

$$u(s) = \frac{i}{2} e^{-\frac{c}{12}s^3 - \frac{c}{8}cs^3} \oint \frac{dy}{2i\pi} (y \pm \frac{1}{y^3}) e^{\frac{cs^3}{16}(y^4 + \frac{1}{y^4})} \quad (4.6)$$

where \pm means A type and D type, respectively. With the change of variable $y = t^{\frac{1}{4}}$, this is written by the modified Bessel function $I_\nu(z)$,

$$\begin{aligned} u(s) &= \frac{i}{8} e^{-\frac{c}{12}s^3 - \frac{c}{8}s^3} \oint \frac{dt}{2i\pi} (t^{-\frac{1}{2}} \pm t^{-\frac{3}{2}}) e^{\frac{cs^3}{16}(t+\frac{1}{t})} \\ &= \frac{i}{8} e^{-\frac{c}{12}s^3 - \frac{c}{8}s^3} [I_{-\frac{1}{2}}(\frac{cs^3}{8}) \pm I_{\frac{1}{2}}(\frac{cs^3}{8})] \\ &= \frac{i}{2} e^{-\frac{c}{12}s^3 - \frac{c}{8}s^3} \sqrt{\frac{1}{\pi(cs^3)}} e^{\pm \frac{cs^3}{8}} \end{aligned} \quad (4.7)$$

Taking $c = -\frac{1}{2}$, we find for A type, the result of (2.9) $\langle \tau_{3g-2} \rangle^{(A)} = \frac{1}{g!(24)^g}$, and for D type, the result of (3.6) $\langle \tau_{3g-2} \rangle^{(D)} = \frac{1}{g!6^g}$.

As another method, we may use the integral representation of the factor of D type

$$\frac{1}{\sqrt{1 - \frac{s^2}{4u^2}}} = \frac{2}{\sqrt{\pi}} \int_0^\infty dt e^{-(1 - \frac{s^2}{4u^2})t^2} \quad (4.8)$$

Then from (4.5),

$$U(s) = \frac{2}{s\sqrt{\pi}} e^{-\frac{c}{12}s^3} \int_0^\infty dt e^{-t^2} \int \frac{du}{2i\pi} e^{-csu^2 - \frac{s^2}{4u^2}t^2} \quad (4.9)$$

$$I = \int_0^\infty dt e^{-t^2} \int_0^\infty du e^{-csu^2} \sum \frac{(-1)^n}{n!} \left(\frac{s^2 t^2}{4u^2}\right)^n \quad (4.10)$$

The integral part is evaluated as $e^{-\frac{c}{4}s^3}$, then we obtain (4.7) and $\langle \tau_{3g-2} \rangle = \frac{1}{g!6^g}$.

- **general p case of open intersection numbers**

Extension of Airy with logarithmic potential (Kontsevich-Penner model) to general p spin case has been investigated with the logarithmic potential with coefficient k in [19, 18].

For genus $g = 1$, the intersection number is

$$\langle \tau \rangle_{g=1} = \frac{p-1+12k^2}{24} \quad (4.11)$$

With $k = \pm\frac{1}{2}$, it becomes

$$\langle \tau \rangle = \frac{p+2}{24} \quad (4.12)$$

which agrees with τ_{D_l} . For genus $g = 2$, the generalized Kontsevich-Penner model gives (Eq.(5.28) of [19]),

$$\langle \tau \rangle_{g=2} = \frac{1}{p(12)^2} \left[\frac{(p-1)(p-3)(2p+1)}{40} - (3p+1)k^2 - 2k^4 \right] \quad (4.13)$$

By putting $k = \pm\frac{1}{2}$, we find exactly $\tau_{D_l}(2) = \frac{1}{5760p}(p+2)(2p+1)(p-6)$ in (3.5).

Thus we find that the one point intersection number of D_l type ($p = 2l - 2$) is same as the intersection numbers of generalized Kontsevich-Penner matrix model with $k = \frac{1}{2}$.

- $p = -1$

When $p = -1$, one point function $u(s)$ provides Euler characteristic $\chi(\bar{M}_{g,1})$ for A_l type singularity [6] as

$$\begin{aligned} u(s) &= \frac{1}{N} \int \frac{du}{2i\pi} \left(\frac{u - \frac{1}{2}}{u + \frac{1}{2}} \right) \\ &= -\frac{1}{N} \int_0^\infty \frac{dz}{2\pi} \frac{e^{-z}}{(1 - e^{-z})^2} e^{-Nz} \\ &= \int_0^\infty \frac{dz}{2\pi} \left(\sum B_n \frac{z^{n-1}}{n!} \right) e^{-Nz} \end{aligned} \quad (4.14)$$

where a change of variable $(u - \frac{1}{2})/(u + \frac{1}{2}) = e^{-z}$ is used and a factor N is putted for the genus g expansion. This $u(s)$ gives Euler characteristic $\chi(\bar{M}_{g,1}) = \zeta(1 - 2g) = -\frac{1}{2g} B_g$, where B_n is a Bernoulli number. $B_1 = \frac{1}{6}, B_2 = \frac{1}{30}, B_3 = \frac{1}{42}, \dots$

For D_l type, we put $p = -1$ in (3.5) with $j = 0$ (gamma function of the denominator becomes one for $p = -1, j = 0$, and numerator gamma function gives $(2g - 1)!$), and find that $\langle \tau_D(g) \rangle$ as $\frac{1}{24}, -\frac{7}{960}, \frac{31}{8064}, \dots$ for $g = 1, 2, 3, \dots$. These numbers are equal to Euler characteristics

$$\chi = (1 - 2^{1-2g}) \frac{B_g}{2g}. \quad (4.15)$$

In the limit $p \rightarrow -1$, (3.2) becomes

$$\begin{aligned} \hat{u}(s) &= \frac{1}{2} \int du e^{-c \log(\frac{u+1}{u-1})} \frac{u}{\sqrt{u^2 - 1}} \\ &= \frac{1}{2} \int du \left(\frac{u-1}{u+1} \right)^c \frac{u}{\sqrt{u^2 - 1}} \end{aligned} \quad (4.16)$$

By the change of variable $(u - 1)/(u + 1) = e^{-z}$, $du = -2e^{-z}/(1 - e^{-z})^2 dz = (-2)(e^{z/2} - e^{-z/2})^{-2} dz$, $u/\sqrt{u^2 - 1} = \frac{1}{2}(e^{z/2} + e^{-z/2})$, it becomes after partial integration,

$$\hat{u}(s) = \int dz \frac{1}{e^{z/2} - e^{-z/2}} e^{-Nz} \quad (4.17)$$

where c is replaced by N to make clear of genus dependence, as $\frac{1}{N^{2g}}$. Above integral reduces to (4.15), with $1/24, 7/960, \dots$ for $g = 1, 2, \dots$. This χ is same as virtual Euler characteristics, obtained for $o(2N)$ matrix model as a non-orientable surface [7]. For $p = -1$, there is no ψ class, and only Euler class (Witten class) exists. Since D_l type is related to $o(2N)$ Lie algebra, it is reasonable to obtain the result of (4.15), which is same as a virtual Euler characteristics of real algebraic curves [34]. Indeed, we have obtained this virtual Euler characteristic of real algebraic curves for $so(2n + 1), so(2n), sp(n)$ cases as antisymmetric matrix models [14, 7].

- $p = -\frac{1}{2}$

It is remarkable that the intersection number has a factor of $(2p + 1)$ for D_l type. Note that this factor appears also in A_l case, where $p = l + 1$ as shown in (2.8). The reason of this factor exists for all genus $g > 1$ is same as A_l case, and it can be explained by the change of variable of (4.1). The exponential factor in (3.2) becomes simply as $e^{-c \frac{z^{1/2}}{y}}$ for $p = -1/2$. Therefore we have for D_l case

with $p = -\frac{1}{2}$ from (4.2),

$$\hat{u}(s) = \oint_c \frac{dy}{2i\pi} \left(y - \frac{1}{y^3}\right) e^{-c \frac{s^{1/2}}{y}} \quad (4.18)$$

where the integral is evaluated by a contour around $y = 0$. This gives non-vanishing term of order s , which means $g = 1$, but the remaining terms of all higher genus ($g > 1$) are vanishing. This gives a factor $(2p + 1)$, which should appear for $g > 1$ in (3.5).

• **p = - 2**

In A_{p-1} matrix model, the case $p = -2$ becomes equivalent to the unitary matrix model (BGW) model [37, 38], and the intersection numbers agree with BGW matrix model as shown in [31].

For D_l ($p = 2l - 2$) case, there is a factor $(p + 2)$ for all genus case as in (3.5). This may be explained as follows [31]. We have in the $p = -2$ case for $D_l = D_0$,

$$\begin{aligned} \hat{u}(s) &= \frac{1}{s} \oint \frac{du}{2i\pi} e^{-\frac{1}{u+s} + \frac{1}{u}} \left(\frac{u+s}{u}\right)^k \\ &= -\frac{1}{4\pi\sqrt{s}} \int dx \frac{1}{x^2} e^{-\frac{4x^2}{1+sx^2}} \left(\frac{1-ix\sqrt{s}}{1+ix\sqrt{s}}\right)^k \\ &= -\frac{1}{2\sqrt{\pi s}} \left(-s \frac{1}{8}(4k^2 - 1) + s^2 \frac{1}{3!2^7}(4k^2 - 1)(4k^2 - 9)\right. \\ &\quad \left.- s^3 \frac{1}{5!2^9}(4k^2 - 1)(4k^2 - 9)(4k^2 - 25) + \dots\right) \end{aligned} \quad (4.19)$$

The second line is derived by the change of variable $u = \frac{1}{2}(-1 + \frac{i}{x\sqrt{s}})$. Since the coefficient of the logarithmic factor of D_l is $k = \frac{1}{2}$ as (4.5), we find the vanishing results of $\hat{u}(s)$ from above equation. This gives a proof for a factor $p + 2$ for the intersection numbers in all orders of (3.5).

BGW model is considered for the unitary matrix, and its extension to O(N) group was studied in [40]. There appears a phase transition between weak coupling (small s) and a strong coupling regions (large s).

• **Large s expansion for p= -2**

The large s expansion was investigated in [31] based on (4.19) for $p = -2$ of A_l type. Due to two terms of (1.1), the extension of the large s expansion for $p = -2$ case of D_l type is straightforward. We obtain by the shift $u \rightarrow (u - 1)/2$ following [31],

$$\begin{aligned} \hat{u}(s) &= \frac{1}{2} \oint \frac{du}{2i\pi} e^{\frac{4}{2(u^2-1)}} \left[\left(\frac{u+1}{u-1}\right)^k + \left(\frac{u+1}{u-1}\right)^{-k}\right] \\ &= \frac{1}{2} \sum_1^\infty \frac{4^m}{m!s^m} \oint \frac{du}{2i\pi} \frac{1}{(u^2-1)^m} \left[\left(\frac{u+1}{u-1}\right)^k + \left(\frac{u+1}{u-1}\right)^{-k}\right] \end{aligned} \quad (4.20)$$

with $k = \frac{1}{2}$. Noting that

$$\begin{aligned} \oint \frac{du}{2i\pi} \frac{1}{(u^2-1)^m} \left(\frac{u+1}{u-1}\right)^k &= -\frac{2}{\pi} \sin\pi k \int_1^\infty dx \frac{(x+1)^{k-m}}{(x-1)^{k+m}} \\ &= -\frac{2}{\pi} (\sin\pi k) 2^{1-2m} (2m-2)! \frac{\Gamma(-k-m+1)}{\Gamma(-k+m)} \end{aligned} \quad (4.21)$$

we obtain $\hat{u}(s)$ in the large s expansion. In the unitary matrix model, we put $k = -N$ and the result agree with the strong coupling expansion (character expansion) [31].

• **Half-integer $p = \frac{1}{2}$ and $\frac{3}{2}$**

For D_l type, with the change of variables $u \rightarrow y$ in (4.1), one point function $\hat{u}(s)$ is

$$\hat{u}(s) = \frac{1}{s} \oint \frac{dy}{2i\pi} \left(y - \frac{1}{y^3}\right) g(y) \quad (4.22)$$

Note that the difference between A_l and D_l is a factor of $(y \pm \frac{1}{y^3})$, where (+) is for A_l and (-) is for D_l . The factor $g(y)$ is same as A_l case. We have for A_l ,

$$u(s) = \frac{1}{s} \int \frac{du}{2i\pi} e^{-\frac{1}{(p+1)}[(u+\frac{s}{2})^{p+1} - (u-\frac{s}{2})^{p+1}]} \quad (4.23)$$

By the change of variable of $u = \frac{i}{2}(y^2 - y^{-2})$, it becomes after the replace $u \rightarrow su/2$, we have for A_l case (+) and D_l case (-)

$$u(s) = \frac{i}{2} \int \frac{dy}{2i\pi} \left(y \pm \frac{1}{y^3}\right) e^{-\frac{sp+1}{(p+1)}(\frac{i}{4})^{p+1} \frac{1}{y^{2(p+1)}} [(y^2-i)^{2p+2} - (y^2+i)^{2p+2}]} \quad (4.24)$$

where the exponential factor $g(y)$ is expressed as

$$g(y) = e^{c' s^{3/2} (3y - \frac{1}{y^3})} \quad (p = \frac{1}{2}, c' = -\frac{1}{6} i^{1/2}) \quad (4.25)$$

The half-integer p case is related to Ramond sector [8, 9]. We assume Riemann-Roch relation for $p = \frac{1}{2}$ is valid as

$$3g - 3 + 1 = n + (g-1)\left(1 - \frac{2}{p}\right) + \frac{j}{p} \quad (4.26)$$

with $p = \frac{1}{2}$ and $j = -1$. $j = -1$ means Ramond puncture [9]. This selection rule indeed valid for the small s expansion of $u(s)$. We have from (4.26)

$$n = 6g - 3 \quad (4.27)$$

which means that $u(s)$ is a power series of s^{6g-3} and the intersection number is given by the coefficient, $\langle \tau_{n,j} \rangle_g = \langle \tau_{6g-3,-1} \rangle_g$.

The one point function $u(s)$ for $p = \frac{1}{2}$ for A_l is evaluated with the integral by part,

$$\begin{aligned} u(s) &= \frac{i}{2} \oint \frac{dy}{2i\pi} \left(y + \frac{1}{y^3}\right) e^{c' s^{3/2} (3y - \frac{1}{y^3})} = \frac{i}{6c' s^{3/2}} \oint \frac{dy}{2i\pi} y \frac{d}{dy} e^{c' s^{3/2} (3y - \frac{1}{y^3})} \\ &= -\frac{i}{6c' s^{3/2}} \oint \frac{dy}{2i\pi} e^{c' s^{3/2} (3y - \frac{1}{y^3})} \\ &= \frac{i}{6} \sum_{m=1}^{\infty} \frac{(-1)^m 3^{3m-1} c'^{4m-2}}{m!(3m-1)!} s^{6m-3} \\ &= 2 \sum_{m=1}^{\infty} \frac{(-1)^m}{m!(3m-1)!} \left(\frac{1}{48}\right)^m s^{6m-3} \end{aligned} \quad (4.28)$$

with $c' = -i^{1/2}/6$. The power m is equal to genus g from (4.27). We find the precise agreement with (2.8). Note that $\Gamma(1 - \frac{2g-1}{p})$ gives the additional

factor for the intersection numbers in (2.8) by the normalization of $\Gamma(-1)$, and it agrees with (4.28). The case of $p = \frac{1}{2}$ is interpreted as a manifestation of Ramond puncture since spin component is $j = -1$.

For D_l type ($p = 1/2, l = 5/4$), it becomes from (4.2)

$$\begin{aligned}\hat{u}(s) &= \frac{i}{2} \oint \frac{dy}{2i\pi} \left(y - \frac{1}{y^3}\right) e^{cs^{3/2}(3y - \frac{1}{y^3})} \\ &= \frac{i}{2s} \oint \frac{dy}{2i\pi} \left(y - \frac{s^2}{y^3}\right) e^{cs(3y - \frac{s^2}{y^3})}\end{aligned}\quad (4.29)$$

This integral is written as the derivative of the exponent by s . Using the same integral as (4.28), we obtain

$$\begin{aligned}\hat{u}(s) &= \frac{i}{6cs} \frac{d}{ds} \left(\oint \frac{dy}{2i\pi} e^{3csy - \frac{cs^3}{y^3}} \right) \\ &= 2 \sum_{m=1}^{\infty} \frac{(-1)^m (6m-1)}{m!(3m-1)!} \left(\frac{1}{48}\right)^m s^{6m-3}\end{aligned}\quad (4.30)$$

where $c = -\frac{1}{6}i^{\frac{1}{2}}$. This result is consistent for $p = \frac{1}{2}$ in (3.5) as $\langle \tau \rangle_{g=1} = \frac{p+2}{24} = \frac{5}{48}$. The expression of $p = \frac{1}{2}$ for D_l ($l = \frac{1}{2}(p+2) = \frac{5}{2}$) is obtained solely from the residue calculation, which means that this case is Ramond puncture with $j = -1$ [9].

For $p = \frac{3}{2}$ of A_l type, it corresponds to $\beta-\gamma$ system as discussed in [8]. There are two different punctures belongs to Neveu-Schwarz and Ramond sectors. From (4.24), D_l case is expressed as

$$g(y) = e^{cs^{3/2}(5y^3 - \frac{10}{y} + \frac{1}{y^5})} \quad (p = \frac{3}{2}, c = \frac{4i}{5} \left(\frac{i}{4}\right)^{5/2}) \quad (4.31)$$

$$u(s) = \oint \frac{dy}{2i\pi} \left(y - \frac{1}{y^3}\right) e^{cs^{\frac{3}{2}}[5y^3 - \frac{10}{y} + \frac{1}{y^5}]} \quad (4.32)$$

The small s expansion gives $\langle \tau_{n,j} \rangle_g s^{n+\frac{2}{3}(j+1)}$, where $j = -1, -\frac{1}{2}, 0$. The spin $j = -1$ corresponds to Ramond sector, and Neveu-Schwarz sector is $j = -\frac{1}{2}, 0$. The Ramond sector (R) is evaluated by the contour integral for A_l of $p = \frac{3}{2}$ in [8],

$$u_R(s) = \frac{5}{2^5} c^2 s^5 - \frac{7}{3} \cdot \frac{5^4}{2^{16}} c^6 s^{15} + \frac{79 \cdot 11}{10!} \cdot \frac{3^{25} 5^7}{2^{24}} c^{10} s^{25} + \dots \quad (4.33)$$

where $s^{(2g-1)(1+\frac{1}{p})} = s^{n+\frac{1}{p}(j+1)}$. If we take $p = \frac{3}{2}$ and $j = -1$, then we have $s^{\frac{10}{3}g - \frac{5}{3}}$. The first term of above equation is for $g = 2$, and the second term is for $g = 5$. For D_l , $p = \frac{3}{2}$, the contribution of Ramond sector becomes

$$u_R(s) = \frac{105}{2^5} c^2 s^5 - \frac{19 \cdot 17 \cdot 5^4}{2^{19}} c^6 s^{15} + \frac{29 \cdot 1663 \cdot 5^5}{3^2 2^{32}} c^{10} s^{25} + \dots \quad (4.34)$$

5 n- point functions for the A_l type of the positive integer p

The punctures are classified into Neveu-Schwarz (NS) and Ramond (R) types. They appear as the boundary condition for anti-periodic and periodic conditions

in the super conformal field theory. For the correlation functions of m -punctures, the path of integrals becomes different. We have used the contour integral for the correlation functions in the variable of u . We then introduced a new variable y , which gives Laurent series for the correlation functions. If we further make a change of variable $y^2 \rightarrow z$, we have two different Laurent series with \sqrt{z} or without \sqrt{z} , which leads to two different types of integrals for $p = 2$, i. e. Neveu-Schwarz and Ramond types, which are familiar to the super conformal field.

For both types, the integral of the variable y gives Laurent series in the genus g expansions. In the variable of y , with a change of variable as $u = \frac{i}{2}(y^2 - \frac{1}{y^2})$, the correlation function $u(s)$ is given by

$$\begin{aligned} u(s) &= \frac{1}{s} \oint \frac{du}{2i\pi} e^{-\frac{c'}{p+1}[(u+\frac{s}{2})^{p+1} - (u-\frac{s}{2})^{p+1}]} \\ &= \frac{1}{4\pi} \oint dy (y + \frac{1}{y^3}) e^{-\frac{c'}{p+1}(\frac{i}{4})^{p+1} s^{p+1} [(y-\frac{i}{y})^{2p+2} - (y+\frac{i}{y})^{2p+2}]} \\ &= \frac{1}{4\pi} \oint dy (y + \frac{1}{y^3}) e^{-c'(\frac{i}{4})^p s^{p+1} (y^{2p} + \dots)} \end{aligned} \quad (5.1)$$

By the scale as $y \rightarrow s^{-\frac{p+1}{2p}} y$, with $c = c'(\frac{i}{4})^p$, it becomes

$$\begin{aligned} u(s) &= \frac{1}{4\pi} \frac{1}{s^{1+\frac{1}{p}}} \int dy (y \pm \frac{s^{2+\frac{2}{p}}}{y^3}) \\ &\quad e^{-c(y^{2p} - \frac{p(2p+1)}{3} s^{2+\frac{2}{p}} y^{2p-4} + \frac{(2p+1)p(2p-1)(p-1)}{30} s^{4+\frac{4}{p}} y^{2p-8} + \dots)} \end{aligned} \quad (5.2)$$

This is a power series of $s^{(2n-1)(1+\frac{1}{p})}$, and by the change of variable $t = y^{2p}$, we obtain the same results of A type (+) and D type (-) as (2.2) and (3.4) for Neveu-Schwarz type.

The genus $g = 1$ term is obtained as

$$u(s) = \frac{1}{4\pi} \frac{1}{s^{1+\frac{1}{p}}} \int dy (y \pm \frac{s^{2+\frac{2}{p}}}{y^3}) e^{-cy^{2p}} [1 + \frac{cp(2p+1)}{3} s^{2+\frac{2}{p}} y^{2p-4}] \quad (5.3)$$

With $y^{2p} = \frac{1}{c}t$, it becomes

$$\begin{aligned} u(s) &= \frac{1}{4\pi} \frac{1}{s^{1+\frac{1}{p}}} \frac{1}{2pc^{\frac{1}{p}}} \int_0^\infty dt e^{-t} t^{\frac{1}{p}-1} [1 + c^{\frac{2}{p}} s^{2+\frac{2}{p}} (\pm \frac{1}{t^{\frac{2}{p}}} + \frac{t}{t^{\frac{2}{p}}} \frac{p(2p+1)}{3})] \\ &= \frac{1}{8\pi pc^{\frac{1}{p}} s^{1+\frac{1}{p}}} [\Gamma(\frac{1}{p}) + c^{\frac{2}{p}} s^{2+\frac{2}{p}} p \Gamma(1 - \frac{1}{p}) (\mp 1 + \frac{2p+1}{3})] \end{aligned} \quad (5.4)$$

This leads to the intersection number $\langle \tau_{1,0} \rangle_{g=1} = \frac{p-1}{24}$ for A type (+) and $\frac{p+2}{24}$ for D type (-), which are consistent with (2.8) and (3.5). Higher genus intersection numbers are obtained similarly. For instance, $g = 2$ of A type, from the terms of order $s^{3+\frac{3}{p}}$, it becomes $\langle \tau_{3,2} \rangle = (p-1)(2p+1)(p-3)/(45 \cdot 2^7 p)$, which is consistent with (2.8).

- two point functions for integer p

The two point function $u(s_1, s_2)$ is written as, (for $p = \frac{1}{2}$, see (7.11)),

$$\begin{aligned}
u(s_1, s_2) &= -4 \left(\frac{s_1}{s_2}\right)^{\frac{1}{p}} \oint \frac{dy_1 dy_2}{(2i\pi)^2} \left(y_1 + \frac{s_1}{y_1^3}\right) \left(y_2 + \frac{s_2}{y_2^3}\right) \\
&\times \frac{1}{y_1^4} \frac{\exp[\sum_{i=1}^2 (-c) (y_i^{2p} - \frac{p(2p+1)}{3} s_i^{2+\frac{2}{p}} y_i^{2p-4} + \frac{(2p+1)p(2p-1)(p-1)}{30} s_i^{4+\frac{4}{p}} y_i^{2p-8} + \dots)]}{\left[1 - \frac{s_1}{y_1^4} - \left(\frac{s_1}{s_2}\right)^{\frac{1}{p}} \frac{1}{y_1} \left(y_2^2 - \frac{s_2}{y_2^2}\right)\right]^2 + \frac{4s_1^{\frac{2}{p}}}{y_1^4} (s_1 + s_2)^2}
\end{aligned} \tag{5.5}$$

For $p = \frac{1}{2}$, it becomes as (7.11) with a slight difference of notation of c (a factor 3 difference).

The selection rule of (2.6) is

$$2(g-1)\left(1 + \frac{1}{p}\right) + 2 = n_1 + n_2 + \frac{1}{p}(j_1 + j_2) \tag{5.6}$$

The two point function is expressed as

$$u(s_1, s_2) = C \sum_{n_i, j_i} \langle \tau_{n_1, j_1} \tau_{n_2, j_2} \rangle s_1^{n_1 + \frac{1}{p}(1+j_1)} s_2^{n_2 + \frac{1}{p}(1+j_2)} \tag{5.7}$$

where C is a constant., which involves factors of gamma-function. This two point function and the intersection numbers $\langle \tau_{n_1, j_1} \tau_{n_2, j_2} \rangle$ for Neveu-Schwarz punctures are evaluated in general p in [32]. Here we reconsider two point functions under the formula by the integral y in (5.5), which may be easily obtained in more systematic ways for both Neveu-Schwarz and Ramond punctures. The numerator and denominator of (5.5) are expanded in the small s_1 and s_2 .

The string equation is easily obtained with the term of $s_1^{\frac{1}{p}}$ in the y_1 integral, where other terms of $s_1^{2+\frac{2}{p}}$ are neglected. The remaining integral of y_2 is just the term of $u(s_2)$ multiplied by s_2 for (5.2).

$$u(s_1, s_2) = 2s_1^{\frac{1}{p}} c^{\frac{1}{p}} s_2 u(s_2) \tag{5.8}$$

Since $s_1^{\frac{1}{p}}$ corresponds to $\langle \tau_{0,0} \rangle$ and $u(s_2)$ corresponds to $\langle \tau_{n,j} \rangle$, we have a string equation such that

$$\langle \tau_{0,0} \tau_{n,j} \rangle_g = \langle \tau_{n-1,j} \rangle_g \tag{5.9}$$

For the genus one case, we have $\langle \tau_{0,2} \tau_{1,p-2} \rangle_{g=1}$ if $p \geq 4$, and $\langle \tau_{0,3} \tau_{1,p-3} \rangle_{g=1}$ if $g \geq 5$. These intersection numbers were evaluated as $\langle \tau_{0,2} \tau_{1,2} \rangle_{g=1} = \frac{1}{96}$ for $p = 4$ and $\langle \tau_{0,2} \tau_{1,3} \rangle_{g=1} = \langle \tau_{0,3} \tau_{1,2} \rangle_{g=1} = \frac{1}{60}$ for $p = 5$ [10, 11]. The result of $\langle \tau_{0,2} \tau_{1,p-2} \rangle_{g=1} = (p-3)/24p$ has been obtained for $p \geq 4$ in [32].

Here we evaluate these cases for general p based on (5.5).

For $\langle \tau_{0,2} \tau_{1,p-2} \rangle_{g=1}$, we have $s_1^{\frac{3}{2}} s_2^{2-\frac{1}{p}}$ term from (5.5) as

$$\begin{aligned}
U(s_1, s_2) &= -4 \left(s_1^{\frac{3}{2}} s_2^{2-\frac{1}{p}}\right) \int_0^\infty dy_1 dy_2 \frac{1}{y_1^7} (-7y_2 + p(2p+1)y_2^{2p+1}) e^{-c(y_1^{2p} + y_2^{2p})} \\
&= \frac{p-3}{24p} s_1^{\frac{3}{2}} s_2^{2-\frac{1}{p}} \Gamma\left(1 - \frac{3}{p}\right) \Gamma\left(\frac{1}{p}\right)
\end{aligned} \tag{5.10}$$

where $c = (i/4)^p$, and the coefficient is the intersection number $\langle \tau_{0,2} \tau_{1,p-2} \rangle_{g=1} = (p-3)/24p$.

For the intersection $\langle \tau_{0,3}\tau_{1,p-3} \rangle$, we evaluate the term of $s_1^{\frac{4}{p}} s_2^{2-\frac{2}{p}}$ as

$$\begin{aligned} u(s_1, s_2) &= s_1^{\frac{4}{p}} s_2^{2-\frac{2}{p}} \int_0^\infty dy_1 dy_2 e^{-cy_1^{2p}-cy_2^{2p}} \frac{1}{y_1^9} (96y_2^2 - \frac{16}{3}p(1+2p))y_2^{3+2p} \\ &= \frac{1}{12p} (p-4) s_1^{\frac{4}{p}} s_2^{2-\frac{2}{p}} \Gamma(1-\frac{4}{p}) \Gamma(1-\frac{p-2}{p}) \end{aligned} \quad (5.11)$$

which gives the intersection number $\langle \tau_{0,3}\tau_{1,p-3} \rangle = \frac{p-3}{24p}$, which gives $\frac{1}{60}$ for $p=5$ and it is consistent with [11].

In the case $p \geq 6$, we obtain $\langle \tau_{0,4}\tau_{1,p-4} \rangle = \frac{p-10}{16p}$, since it is evaluated as

$$u(s_1, s_2) = \frac{1}{16} \frac{p-10}{p} s_1^{\frac{4}{p}} s_2^{2-\frac{2}{p}} \Gamma(1-\frac{5}{p}) \Gamma(\frac{3}{p}) \quad (5.12)$$

For the genus two case, the intersection numbers are $\langle \tau_{0,1}\tau_{n,j} \rangle$, which are obtained from the term of $s_1^{\frac{2}{p}} s_2^{n+\frac{1+j}{p}}$. Previously such intersection numbers as $\langle \tau_{0,1}\tau_{4,1} \rangle_{g=2} = \frac{1}{320}$ for $p=4$ [10] and $\langle \tau_{0,1}\tau_{4,1} \rangle_{g=2} = \frac{7}{1200}$ for $p=5$ [11], are reported. The intersection number $\langle \tau_{0,1}\tau_{4,1} \rangle_{g=2}$ for general spin p is obtained from the integral of order of $s_1^{\frac{2}{p}} s_2^{4+\frac{2}{p}}$ by the expansion of the denominator and the exponents as

$$\begin{aligned} u(s_1, s_2) &= c^{\frac{4}{p}} s_1^{\frac{2}{p}} s_2^{4+\frac{2}{p}} \int_0^\infty \frac{1}{y_1^5} e^{-y_1^{2p}} \int_0^\infty dy_2 e^{-y_2^{2p}} (\frac{8}{y_2^5} \\ &\quad + \frac{4}{15}(p-1)p(2p-1)(2p+1)y_2^{2p-5} - \frac{4}{9}p^2(2p+1)^2] \\ &= \frac{1}{2880} (p-1)(p+2)(p-2) \{ \Gamma(1-\frac{2}{p}) \}^2 s_1^{\frac{2}{p}} s_2^{4+\frac{2}{p}} \end{aligned} \quad (5.13)$$

By the normalization of the intersection number with $p^{-(g-1)}$, we have $\langle \tau_{0,1}\tau_{4,1} \rangle_{g=2} = \frac{(p-1)(p+2)(p-2)}{2880p}$, which agrees with the previous result of Eq. (5.24) of [32], and with results of [10, 11], i.e. $\langle \tau_{0,1}\tau_{4,1} \rangle_{g=2} = \frac{1}{320}$ for $p=4$ and $\langle \tau_{0,1}\tau_{4,1} \rangle_{g=2} = \frac{7}{1200}$.

For the case of $\langle \tau_{0,2}\tau_{4,0} \rangle_{g=2}$, which is proportional to $s_1^{\frac{3}{p}} s_2^{4+\frac{1}{p}}$, is obtained by the expansion of (5.5) in this order,

$$\begin{aligned} u(s_1, s_2) &= c^{\frac{4}{p}} s_1^{\frac{3}{p}} s_2^{4+\frac{1}{p}} \int_0^\infty dy_1 \frac{1}{y_1^7} e^{-y_1^{2p}} \int_0^\infty dy_2 e^{-y_2^{2p}} [\frac{28}{y_2^3} \\ &\quad + (\frac{146p}{15} + \frac{274p^2}{15} - \frac{8p^3}{5} + \frac{8p^4}{5})y_2^{2p-3} - (\frac{2p^2}{3} + \frac{8p^3}{3} + \frac{8p^4}{3})y_2^{4p-3}] \\ &= \frac{1}{5760} (p-1)(p-3)(2p+11) \Gamma(1-\frac{3}{p}) \Gamma(1-\frac{1}{p}) \end{aligned} \quad (5.14)$$

which is consistent with the previous result of $\langle \tau_{0,2}\tau_{4,0} \rangle_{g=2} = \frac{(p-1)(p-3)(2p+11)}{5760p}$ with the normalization constant $\frac{1}{p}$, i.e. $\langle \tau_{0,2}\tau_{4,0} \rangle = \frac{19}{7680}$ for $p=4$ and $\langle \tau_{0,2}\tau_{4,0} \rangle = \frac{7}{1200}$ for $p=5$ in [32, 10, 11].

In the genus $g=2$, two point function is evaluated from (5.5). We represent examples of (ii.1) - (ii.5), which are consistent with the results in [10].

For the case of $\langle \tau_{0,2}\tau_{4,0} \rangle_{g=2}$, which is proportional to $s_1^{\frac{3}{p}} s_2^{4+\frac{1}{p}}$, is obtained

by the expansion of (5.5) in this order,

$$\begin{aligned}
(ii.1) \quad u(s_1, s_2) &= c^{\frac{4}{p}} s_1^{\frac{3}{p}} s_2^{4+\frac{1}{p}} \int_0^\infty dy_1 \frac{1}{y_1} e^{-y_1^{2p}} \int_0^\infty dy_2 e^{-y_2^{2p}} \left[\frac{28}{y_2^3} \right. \\
&\quad \left. + \left(\frac{146p}{15} + \frac{274p^2}{15} - \frac{8p^3}{5} + \frac{8p^4}{5} \right) y_2^{2p-3} - \left(\frac{2p^2}{3} + \frac{8p^3}{3} + \frac{8p^4}{3} \right) y_2^{4p-3} \right] \\
&= \frac{1}{5760} (p-1)(p-3)(2p+11) \Gamma\left(1 - \frac{3}{p}\right) \Gamma\left(1 - \frac{1}{p}\right)
\end{aligned} \tag{5.15}$$

which is consistent with the previous result of $\langle \tau_{0,2}\tau_{4,0} \rangle_{g=2} = (p-1)(p-3)(2p+11)/5760p$ with the normalization constant $\frac{1}{p}$, i.e. $\langle \tau_{0,2}\tau_{4,0} \rangle = \frac{19}{7680}$ for $p=4$ and $\langle \tau_{0,2}\tau_{4,0} \rangle = \frac{7}{1200}$ for $p=5$ in [32, 10, 11].

In the case of $\langle \tau_{1,1}\tau_{3,1} \rangle_{g=2}$ for $p=3$, from the expansion of order $s_1^{5/3} s_2^{11/3} = t_{1,1}t_{3,1}$, we have from (5.5),

$$\begin{aligned}
(ii.2a) \quad &(-4)c^{\frac{4}{3}} \int_0^\infty dy_1 dy_2 e^{-y_1^6 - y_2^6} y_1^{-11} (66y_2 - 420y_2^7 + 245/2y_2^{13}) \\
&= \frac{11}{1440} \left(\frac{1}{4}\right)^4 [\Gamma(1 - \frac{2}{3})]^2.
\end{aligned} \tag{5.16}$$

By multiplying a normalization constant $1/p^{g-1} = \frac{1}{3}$, it leads to $\langle \tau_{1,1}\tau_{3,1} \rangle_{g=2} = \frac{11}{4320}$ for $p=3$, which agrees with the value of [10, 12].

For $p=4$, $\langle \tau_{1,1}\tau_{3,1} \rangle$, it becomes

$$\begin{aligned}
(ii.2b) \quad u(s_1, s_2) &= s_1^{\frac{3}{4}} s_2^{\frac{7}{4}} c \int dy_1 dy_2 \frac{1}{y_1^{13}} (-1144y_2^3 + \frac{27984}{5}y_2^{11} - 1728y_2^{19}) \\
&\quad \times e^{-y_1^8 - y_2^8} = \frac{7}{240} s_1^{\frac{3}{4}} s_2^{\frac{7}{4}} [\Gamma(\frac{1}{2})]^2
\end{aligned} \tag{5.17}$$

It leads to $\langle \tau_{1,1}\tau_{3,1} \rangle_{g=2} = \frac{7}{960}$ for $p=4$ by multiplying a normalization factor $\frac{1}{p^{g-1}} = \frac{1}{4}$ by the definition of the intersection number, which agree with [10, 11]. For $p=5$, $\langle \tau_{1,1}\tau_{3,1} \rangle$, it becomes

$$\begin{aligned}
(ii.2c) \quad u(s_1, s_2) &= s_1^{\frac{7}{5}} s_2^{\frac{17}{5}} c^{\frac{4}{5}} \int_0^\infty dy_1 dy_2 \frac{1}{y_1^{15}} (-3276y_2^5 + \frac{44044}{3}y_2^{15} \\
&\quad - \frac{42350}{9}y_2^{25}) e^{-y_1^{10} - y_2^{10}} = \frac{17}{240} s_1^{\frac{7}{5}} s_2^{\frac{17}{5}} [\Gamma(\frac{3}{5})]^2
\end{aligned} \tag{5.18}$$

with $c = (\frac{1}{4})^p$. It leads to $\langle \tau_{1,1}\tau_{3,1} \rangle = \frac{17}{1200}$ for $p=5$ with a factor $\frac{1}{5}$, which agrees with [11].

For $\langle \tau_{1,2}\tau_{3,0} \rangle$ of $p=4$, we have

$$\begin{aligned}
(ii.3) \quad u(s_1, s_2) &= s_1^{\frac{7}{4}} s_2^{\frac{13}{4}} \int_0^\infty dy_1 dy_2 \left(\frac{32}{y_1^7 y_2^3} - \frac{3276y_2^5}{y_1^{15}} + \frac{384y_2^5}{y_1^7} + \frac{45528y_2^{13}}{5y_1^{15}} \right. \\
&\quad \left. - \frac{2016y_2^{21}}{y_1^{15}} \right) e^{-c(y_1^8 + y_2^8)} = \frac{41}{1920} s_1^{\frac{7}{4}} s_2^{\frac{13}{4}} \Gamma\left(\frac{1}{4}\right) \Gamma\left(\frac{3}{4}\right)
\end{aligned} \tag{5.19}$$

This leads to $\langle \tau_{1,2}\tau_{3,0} \rangle = \frac{41}{7680}$ for $p=4$ with a normalization factor $\frac{1}{p}$, which agrees with [10].

For $\langle \tau_{2,0}\tau_{2,2} \rangle$ of $p=4$, we get

$$\begin{aligned}
(ii.4) \quad u(s_1, s_2) &= s_1^{\frac{9}{4}} s_2^{\frac{11}{4}} \int_0^\infty dy_1 dy_2 \left(-\frac{576y_2}{y_1^{11}} + \frac{3840y_2^9}{y_1^{11}} - \frac{15504y_2^9}{y_1^{19}} \right. \\
&\quad \left. + \frac{100296y_2^{17}}{5y_1^{19}} - \frac{2592y_2^{25}}{y_1^{19}} \right) e^{-c(y_1^8 + y_2^8)} = \frac{49}{1920} s_1^{\frac{9}{4}} s_2^{\frac{11}{4}} \Gamma\left(\frac{3}{4}\right) \Gamma\left(\frac{1}{4}\right).
\end{aligned} \tag{5.20}$$

which leads to $\langle \tau_{2,0}\tau_{2,2} \rangle = \frac{49}{7680}$ with a factor $\frac{1}{p} = \frac{1}{4}$. This result also agrees with [10].

For $\langle \tau_{2,1}\tau_{2,1} \rangle$ of $p = 4$, we find

$$(ii.5) \quad u(s_1, s_2) = s_1^{\frac{5}{2}} s_2^{\frac{5}{2}} \int_0^\infty dy_1 dy_2 e^{-c(y_1^8 + y_2^8)} \left(\frac{1}{y_1^{13}} (-2816y_2^3 + 7680y_2^{11}) \right. \\ \left. + \frac{1}{y_1^{21}} (-28728y_2^{11} + 27888y_2^{19} - 2880y_2^{27}) \right) = \frac{11}{240} s_1^{\frac{5}{2}} s_2^{\frac{5}{2}} \left[\Gamma\left(\frac{1}{2}\right) \right]^2 \quad (5.21)$$

By multiplying a factor $\frac{1}{p} = \frac{1}{4}$, it leads to $\langle \tau_{2,1}\tau_{2,1} \rangle = \frac{11}{960}$, which agrees with [10].

In the genus $g = 3$, $\langle \tau_{7,0}\tau_{0,1} \rangle_{g=3} = \frac{1}{15552}$ for $p = 3$ is reported in [46, 48, 10, 32]. This case of $p = 3$ is obtained by the expansion of the exponent of

$$e^{-y_i^6 + s_1^{\frac{8}{3}} y_i^2 - 7s_i^{\frac{16}{3}} y_i^{-2} + s_i^8 y_i^{-6}} \quad (5.22)$$

up to order s_i^8 . We obtain $u(s_1, s_2) = \frac{1}{1728} s_1^{\frac{22}{3}} s_2^{\frac{2}{3}} \Gamma\left(\frac{2}{3}\right) \Gamma\left(\frac{1}{3}\right)$. By the normalization of $\frac{1}{9}$ for genus two, we obtain the result of $\langle \tau_{7,0}\tau_{0,1} \rangle_{g=3} = \frac{1}{15552}$ for $p = 3$.

Thus we verified the integral of (5.5) is a generating function for the intersection numbers of two point function. The evaluation is straightforward for arbitrary genus g in the variables of y_1 and y_2 , which is reduced to the gamma-function of $\Gamma\left(1 - \frac{1+j}{p}\right)$ for $s^{n + \frac{1+j}{p}}$. Each terms satisfy the selection rule of (2.6) for genus g .

• three point functions

Three point function $u(s_1, s_2, s_3)$ is given for A type by

$$u(s_1, s_2, s_3) = -8 \left(\frac{s_1}{s_3} \right)^{\frac{1}{p}} \int \frac{1}{(2i\pi)^3} \prod_{i=1}^3 dy_i \prod_{i=1}^3 \left(y_i + \frac{s_i}{y_i^3} \right) e^{-c \sum (y_i^{2p} + \dots)} \\ \times \frac{1}{y_1^4} \frac{1}{1 - \frac{s_1}{y_1^4} - \left(\frac{s_2}{s_3} \right)^{\frac{1}{p}} \frac{1}{y_1^2} \left(y_2^2 - \frac{s_2}{y_2^2} \right) - \frac{2is_1^{\frac{1}{p}}}{y_1^2} (s_1 + s_2)} \\ \times \frac{1}{y_2^2} \frac{1}{1 - \frac{s_2}{y_2^2} - \left(\frac{s_3}{s_1} \right)^{\frac{1}{p}} \frac{1}{y_2^2} \left(y_3^2 - \frac{s_3}{y_3^2} \right) - \frac{2is_2^{\frac{1}{p}}}{y_2^2} (s_2 + s_3)} \\ \times \frac{1}{1 - \frac{s_1}{y_1^4} - \left(\frac{s_3}{s_1} \right)^{\frac{1}{p}} \frac{1}{y_1^2} \left(y_3^2 - \frac{s_3}{y_3^2} \right) + \frac{2is_1^{\frac{1}{p}}}{y_1^2} (s_1 + s_3)} \quad (5.23)$$

The exponential term is expressed after the scaling of s as (5.5),

$$e^{-y_i^{2p} + \frac{p(2p+1)}{3} s_i^{\frac{2}{p}} y_i^{2p-2} - \frac{(2p+1)p(2p-1)(p-1)}{30} s_i^{\frac{4}{p}} y_i^{2p-8} + \dots} \quad (5.24)$$

where the factor c is absorbed in y_i , which follows the scaling to s_i as cs^{p+1} , and it is expressed as an over all factor in the front of the integral.

For the term of $s_1^{\frac{1}{p}}$, as a simplest case, we obtain in the genus $g = 0$ case by

neglecting all terms of $s_i^{2+\frac{2}{p}}$,

$$\begin{aligned}
u(s_1, s_2, s_3) &= -8\left(\frac{s_1}{s_3}\right)^{\frac{1}{p}} \int \frac{dy_1}{2i\pi} \frac{1}{y_1^3} e^{-cy_1^{2p}} \int \frac{dy_2}{2i\pi} \frac{dy_3}{2i\pi} \frac{1}{y_2} y_3 e^{-cy_2^{2p}-cy_3^{2p}} \\
&\times \frac{1}{1 - \frac{s_2^{\frac{p}{2}}}{y_2^{\frac{1}{2}} \left(\frac{y_3^2}{s_3^{\frac{1}{p}}} + 2is_3\right)}}
\end{aligned} \tag{5.25}$$

The term of order $s_2^{\frac{n}{p}}$ is obtained in the geometrical series, and from $s_2^{\frac{n}{p}} \left(\frac{y_3^2}{s_3^{\frac{1}{p}}} + 2is_3\right)^n \sim 2ins_2^{\frac{n}{p}} s_3^{1-\frac{n-1}{p}} y_3^{2(n-1)}$, and by the integrations of y_i , we obtain

$$u(s_1, s_2, s_3) = -8s_1^{\frac{1}{p}} s_2^{\frac{n}{p}} s_3^{1-\frac{n}{p}} (2in) \frac{1}{(2p)^3} \Gamma\left(-\frac{1}{p}\right) \Gamma\left(-\frac{n}{p}\right) \Gamma\left(\frac{n}{p}\right) \tag{5.26}$$

Noting that $\Gamma\left(\frac{n}{p}\right) = \Gamma\left(1 - \frac{p-n}{p}\right)$, we obtain the intersection number of $\langle \tau_{0,0}\tau_{0,n-1}\tau_{p-n-1} \rangle_{g=0} = 1$, which satisfies

$$\langle \tau_{0,j_1}\tau_{0,j_2}\tau_{0,j_3} \rangle_{g=0} = \delta_{j_1+j_2+j_3,p-2} \tag{5.27}$$

for $j_1 = 0, j_2 = n-1, j_3 = p-n-1$.

Above formula is general. For instance, for $p = 8$, we have the case, which satisfies the selection rule of (2.6) as $\langle \tau_{0,1}\tau_{0,2}\tau_{0,3} \rangle_{g=0}$. We have a correspondent term of $s_1^{\frac{2}{p}} s_2^{\frac{3}{p}} s_3^{1-\frac{4}{p}}$ with $p = 8$. This term is easily obtained by the expansion of the denominator in (5.23), which includes a term of $s_2^{\frac{1}{2}} s_3$ same as (5.26). Other sets of j_i satisfy the formula of (5.27), and it is proved from (5.23).

For $p = 3$, we have $g = 0$ contribution from (5.23), as an example $\langle \tau_{0,0}\tau_{0,0}\tau_{0,1} \rangle$,

$$\begin{aligned}
u(s_1, s_2, s_3) &= (-8)(2i)s_1^{\frac{1}{3}} s_2^{\frac{1}{3}} s_3^{\frac{2}{3}} \int_0^\infty \prod dy_i \left(\frac{y_3}{y_1^2 y_2}\right) e^{-c \sum y_i^6} \\
&= -\frac{2i}{3} \left[\Gamma\left(\frac{2}{3}\right)\right]^2 \Gamma\left(\frac{1}{3}\right) c^{\frac{1}{3}}
\end{aligned} \tag{5.28}$$

Noting $c = c' \left(\frac{1}{4}\right)^p$, ($c' = 1$), and with p^{g-1} factor, we obtain the result of (5.27).

The genus $g = 1$ is evaluated from (5.23). For instance, $\langle \tau_{0,1}\tau_{0,1}\tau_{2,1} \rangle_{g=1} = \frac{1}{36}$, which is evaluated from (5.23) as

$$\begin{aligned}
\text{(iii.0)} \quad u(s_1, s_2, s_3) &= (-8)s_1^{\frac{2}{3}} s_2^{\frac{2}{3}} s_3^{\frac{8}{3}} \int_0^\infty \prod dy_i \left[-\frac{8iy_3}{y_1^5 y_2^5} + \frac{56iy_3^7}{y_1^5 y_2^5}\right] e^{-c \sum y_i^6} \\
&= \frac{1}{72} s_1^{\frac{2}{3}} s_2^{\frac{2}{3}} s_3^{\frac{8}{3}} \left[\Gamma\left(\frac{1}{3}\right)\right]^3
\end{aligned} \tag{5.29}$$

with $c = \left(\frac{1}{4}\right)^3$. The factor 2 is needed to compare with the result of [10]. The factor 2 is due to the symmetric factor of the connected longest cycles in the determinant of 3×3 matrix of the three point function [6]. The intersection number $\langle \tau_{0,1}\tau_{1,1}\tau_{1,1} \rangle_{g=1} = \frac{1}{36}$ is obtained as a coefficient of $s_1^{\frac{2}{3}} s_2^{\frac{2}{3}} s_3^{\frac{8}{3}}$ with a normalization factor 2.

There are results of the intersection numbers of three-point for $p = 3$ up to $g = 2$ for different 6 intersection numbers[10]. We evaluate these 6 cases ((3.1)-(3.6)) here to verify the validity the integral representation of (5.23) based on

the random matrix theory. The genus $g = 2$ calculation, we need the expansion of the exponent in (5.24) up to $s_i^{4+\frac{4}{p}}$. The genus $g = 2$ intersection number (1) $\langle \tau_{0,1}\tau_{0,1}\tau_{5,0} \rangle_{g=2} = \frac{1}{432}$, and (2) $\langle \tau_{0,1}\tau_{1,1}\tau_{4,0} \rangle_{g=2} = \frac{13}{2160}$ are obtained from (5.23) with a normalization factor $\frac{2}{3}$ (3 is due to p^{g-1}). For $\langle \tau_{0,1}\tau_{0,1}\tau_{5,0} \rangle_{g=2}$ of $p = 3$, (5.23) becomes as a term of $s_1^{\frac{2}{3}}s_2^{\frac{2}{3}}s_3^{\frac{16}{3}}$,

$$(iii.1) \quad u(s_1, s_2, s_3) = 8\frac{1}{4^5} \int_0^\infty dy_1 dy_2 dy_3 e^{-\sum y_i^6} \left[-\frac{8}{y_1^5 y_2^5 y_3^3} - \frac{112y_3^3}{y_1^5 y_2^5} + \frac{196y_3^9}{y_1^5 y_2^5} \right] \\ = \frac{1}{288} s_1^{\frac{2}{3}} s_2^{\frac{2}{3}} s_3^{\frac{16}{3}} \left[\Gamma\left(\frac{1}{3}\right) \right]^2 \Gamma\left(\frac{2}{3}\right) \quad (5.30)$$

which leads to $\langle \tau_{0,1}\tau_{0,1}\tau_{5,0} \rangle_{g=2} = \frac{1}{432}$ with a normalization factor of $\frac{2}{3}$. For $\langle \tau_{0,1}\tau_{1,1}\tau_{4,0} \rangle_{g=2}$ of $p = 3$, we have

$$(iii.2) \quad u(s_1, s_2, s_3) = 8\frac{1}{4^5} \int dy_1 dy_2 dy_3 e^{-\sum y_i^6} \left[-\frac{28}{y_1^5 y_2^5 y_3^3} + \frac{484y_3^3}{y_1^5 y_2^{11}} - \frac{280y_3^9}{y_1^5 y_2^5} \right. \\ \left. - \frac{1820y_3^9}{y_1^5 y_2^{11}} + \frac{294y_3^9}{y_1^5 y_2^5} + \frac{490y_3^{15}}{y_1^5 y_2^{11}} \right] = \frac{13}{1440} s_1^{\frac{2}{3}} s_2^{\frac{5}{3}} s_3^{\frac{13}{3}} \left[\Gamma\left(\frac{1}{3}\right) \right]^2 \Gamma\left(\frac{2}{3}\right) \quad (5.31)$$

With a factor $\frac{2}{3}$, it leads to $\langle \tau_{0,1}\tau_{1,1}\tau_{4,0} \rangle_{g=2} = \frac{13}{2160}$.

In the case of $\langle \tau_{0,1}\tau_{2,1}\tau_{3,0} \rangle_{g=2}$,

$$(iii.3) \quad u(s_1, s_2, s_3) = -8is_1^{\frac{2}{3}}s_2^{\frac{8}{3}}s_3^{\frac{10}{3}}c^{\frac{5}{3}} \int \prod dy_i e^{-\sum y_i^6} \left(-\frac{32}{y_1^5 y_2^5 y_3^3} + \frac{2040y_3^3}{y_1^5 y_2^{11}} \right. \\ \left. - \frac{224y_3^3}{y_1^5 y_2^5} + \frac{8704y_3^9}{y_1^5 y_2^{11}} - \frac{5320y_3^9}{y_1^5 y_2^{11}} - \frac{8064y_3^{15}}{y_1^5 y_2^{17}} + \frac{588y_3^{15}}{y_1^5 y_2^{11}} + \frac{784y_3^{21}}{y_1^5 y_2^{17}} \right) \\ = \frac{23}{1440} \left(\Gamma\left(\frac{1}{3}\right) \right)^2 \Gamma\left(\frac{2}{3}\right) s_1^{\frac{2}{3}} s_2^{\frac{8}{3}} s_3^{\frac{10}{3}} \quad (5.32)$$

With a normalization factor $\frac{2}{3}$, it becomes $\frac{23}{2160}$, which agrees with the result of [10].

In the case of $\langle \tau_{0,1}\tau_{2,0}\tau_{3,1} \rangle_{g=2}$,

$$(iii.4) \quad u(s_1, s_2, s_3) = -8is_1^{\frac{2}{3}}s_2^{\frac{7}{3}}s_3^{\frac{11}{3}}c^{\frac{5}{3}} \int \prod dy_i \left(\frac{1}{y_1^5 y_2^9} (392y_3 - 2576y_3^7 \right. \\ \left. + 490y_3^{13}) + \frac{1}{y_1^5 y_2^{15}} (3976y_3^7 - 5292y_3^{13} + 686y_3^{19}) \right) e^{-\sum y_i^6} \\ = \frac{1}{72} s_1^{\frac{2}{3}} s_2^{\frac{7}{3}} s_3^{\frac{11}{3}} \left(\Gamma\left(\frac{1}{3}\right) \right)^2 \Gamma\left(\frac{2}{3}\right) \quad (5.33)$$

With a normalization factor $\frac{2}{3}$, it becomes $\langle \tau_{0,1}\tau_{2,0}\tau_{3,1} \rangle_{g=2} = \frac{1}{108}$.

In the case of $\langle \tau_{1,1}\tau_{1,1}\tau_{3,0} \rangle_{g=2}$,

$$(iii.5) \quad u(s_1, s_2, s_3) = -8c^{\frac{5}{3}}s_1^{\frac{5}{3}}s_2^{\frac{5}{3}}s_3^{\frac{10}{3}} \int \prod dy_i e^{-\sum y_i^6} \left(-560\frac{y_2 y_3^3}{y_1^{11}} - 80\frac{y_2}{y_1^{11} y_3^3} \right. \\ \left. + \frac{1}{y_1^{11} y_2^5} (2860y_3^3 - 7700y_3^9 + 1470y_3^{15}) + \frac{1}{y_1^{11} y_2^{11}} (8520y_3^9 - 8400y_3^{15} \right. \\ \left. + 1225y_3^{21}) \right) \\ = \frac{29}{1440} s_1^{\frac{5}{3}} s_2^{\frac{5}{3}} s_3^{\frac{10}{3}} \left(\Gamma\left(\frac{1}{3}\right) \right)^2 \Gamma\left(\frac{2}{3}\right) \quad (5.34)$$

This leads to $\langle \tau_{1,1}\tau_{1,1}\tau_{3,0} \rangle_{g=2} = \frac{29}{2160}$ for $p = 3$ by the normalization of $\frac{2}{3}$.

In the case of $\langle \tau_{1,1}\tau_{2,0}\tau_{2,1} \rangle_{g=2}$,

$$\begin{aligned}
\text{(iii.6)} \quad u(s_1, s_2, s_3) &= -8c^{\frac{5}{3}}s_1^{\frac{5}{3}}s_2^{\frac{7}{3}}s_3^{\frac{8}{3}} \int \prod dy_i e^{-\sum y_i^6} \left(\frac{1}{y_1^{11}y_2^3} (1440y_3 \right. \\
&\quad \left. - 5600y_3^7) + \frac{1}{y_1^{11}y_2^9} (22880y_3^7 - 21840y_3^{13} + 1960y_3^{19}) \right. \\
&\quad \left. + \frac{1}{y_1^{11}y_2^{155}} (30954y_3^{13} - 18620y_3^{19} + 1715y_3^{25}) \right) \\
&= \frac{19}{720} s_1^{\frac{5}{3}} s_2^{\frac{7}{3}} s_3^{\frac{8}{3}} \left(\Gamma\left(\frac{1}{3}\right) \right)^2 \Gamma\left(\frac{1}{3}\right)
\end{aligned} \tag{5.35}$$

which leads to $\langle \tau_{1,1}\tau_{2,0}\tau_{2,1} \rangle_{g=2} = \frac{19}{1080}$ by the multiplying a factor $\frac{2}{3}$.

We have correctly derived the 6 intersection numbers (3.1)-(3.6) of three punctures of $p = 3$ in genus $g = 2$ with a normalization constant $\frac{2}{3}$, which agrees the values of [10]. The 3 point function of (3.1)-(3.6), is expressed in the case of $p = 3$ for genus $g = 2$ as,

$$\begin{aligned}
U(s_1, s_2, s_3)_{g=2, p=3} &= s_1^{\frac{2}{3}} s_2^{\frac{2}{3}} s_3^{\frac{16}{3}} \langle \tau_{0,1}\tau_{0,1}\tau_{5,0} \rangle + s_1^{\frac{2}{3}} s_2^{\frac{5}{3}} s_3^{\frac{13}{3}} \langle \tau_{0,1}\tau_{1,1}\tau_{4,0} \rangle \\
&\quad + s_1^{\frac{2}{3}} s_2^{\frac{8}{3}} s_3^{\frac{10}{3}} \langle \tau_{0,1}\tau_{2,1}\tau_{3,0} \rangle + s_1^{\frac{2}{3}} s_2^{\frac{5}{3}} s_3^{\frac{10}{3}} \langle \tau_{1,1}\tau_{1,1}\tau_{3,0} \rangle \\
&\quad + s_1^{\frac{2}{3}} s_2^{\frac{7}{3}} s_3^{\frac{11}{3}} \langle \tau_{0,1}\tau_{2,0}\tau_{3,1} \rangle + s_1^{\frac{5}{3}} s_2^{\frac{7}{3}} s_3^{\frac{8}{3}} \langle \tau_{1,1}\tau_{2,0}\tau_{2,1} \rangle
\end{aligned} \tag{5.36}$$

where the interaction numbers are given above. Thus we confirm that the three point function of (5.23) gives a correct generating function of the intersection numbers for three marked point for $p = 3$. In general p , it also gives the intersection numbers, which are consistent with the results of $p = 4, 5$ derived through recursion relations [11, 13].

• four point functions

Four point function $u(s_1, s_2, s_3, s_4)$ is given for A type (for D type, the (+) sign of the first integrand is replaced by (-)),

$$\begin{aligned}
u(s_1, s_2, s_3, s_4) &= -16 \left(\frac{s_1}{s_4} \right)^{\frac{1}{p}} \int \prod_{i=1}^4 \frac{dy_i}{2i\pi} \prod_{i=1}^4 \left(y_i + \frac{s_i}{y_i^{\frac{2+p}{p}}} \right) \\
&\quad \times e^{c \frac{i}{4p+4} \sum s_i^{-1-\frac{1}{p}} \left(\left(y_i + \frac{s_i}{y_i} \right)^{2p+2} - \left(y_i - \frac{s_i}{y_i} \right)^{2p+2} \right)} \\
&\quad \times \frac{1}{y_1^4} \frac{1}{1 - \frac{s_1}{y_1^{\frac{2+p}{p}}} - \left(\frac{s_1}{s_2} \right)^{\frac{1}{p}} \frac{1}{y_1^{\frac{2}{p}}} \left(y_2^2 - \frac{s_2}{y_2^{\frac{2+p}{p}}} \right) - \frac{2is_1^{\frac{1}{p}}}{y_1^{\frac{2}{p}}} (s_1 + s_2)} \\
&\quad \times \frac{1}{y_2^2} \frac{1}{1 - \frac{s_2}{y_2^{\frac{2+p}{p}}} - \left(\frac{s_2}{s_3} \right)^{\frac{1}{p}} \frac{1}{y_2^{\frac{2}{p}}} \left(y_3^2 - \frac{s_3}{y_3^{\frac{2+p}{p}}} \right) - \frac{2is_2^{\frac{1}{p}}}{y_2^{\frac{2}{p}}} (s_2 + s_3)} \\
&\quad \times \frac{1}{y_3^2} \frac{1}{1 - \frac{s_3}{y_3^{\frac{2+p}{p}}} - \left(\frac{s_3}{s_4} \right)^{\frac{1}{p}} \frac{1}{y_3^{\frac{2}{p}}} \left(y_4^2 - \frac{s_4}{y_4^{\frac{2+p}{p}}} \right) - \frac{2is_3^{\frac{1}{p}}}{y_3^{\frac{2}{p}}} (s_3 + s_4)} \\
&\quad \times \frac{1}{1 - \frac{s_1}{y_1^{\frac{2+p}{p}}} - \left(\frac{s_1}{s_4} \right)^{\frac{1}{p}} \frac{1}{y_1^{\frac{2}{p}}} \left(y_4^2 - \frac{s_4}{y_4^{\frac{2+p}{p}}} \right) + \frac{2is_1^{\frac{1}{p}}}{y_1^{\frac{2}{p}}} (s_1 + s_4)}
\end{aligned} \tag{5.37}$$

Or in a symmetric form between s_1 and s_3 , it can be written as

$$\begin{aligned}
u(s_1, s_2, s_3, s_4) &= -16 \left(\frac{s_1 s_3}{s_2 s_4} \right)^{\frac{1}{p}} \int \prod_{i=1}^4 \frac{dy_i}{2i\pi} \prod_{i=1}^4 \left(y_i + \frac{s_i}{y_i^3} \right)^{2+\frac{2}{p}} \\
&\times \frac{1}{y_1^4 y_3^4} e^{c \frac{i}{4p+4} \sum s_i^{-1-\frac{1}{p}} \left((y_i + \frac{s_i}{y_i^3})^{2p+2} - (y_i - \frac{s_i}{y_i^3})^{2p+2} \right)} \\
&\times \frac{1}{1 - \frac{s_1}{y_1^4} - \left(\frac{s_1}{s_2} \right)^{\frac{1}{p}} \frac{1}{y_1^2} \left(y_2^2 - \frac{s_2}{y_2^2} \right) - \frac{2is_1^{\frac{1}{p}}}{y_1^2} (s_1 + s_2)} \\
&\times \frac{1}{1 - \frac{s_1}{y_1^4} - \left(\frac{s_1}{s_4} \right)^{\frac{1}{p}} \frac{1}{y_1^2} \left(y_4^2 - \frac{s_4}{y_4^2} \right) + \frac{2is_1^{\frac{1}{p}}}{y_1^2} (s_1 + s_4)} \\
&\times \frac{1}{1 - \frac{s_3}{y_3^4} - \left(\frac{s_3}{s_2} \right)^{\frac{1}{p}} \frac{1}{y_3^2} \left(y_2^2 - \frac{s_2}{y_2^2} \right) + \frac{2is_3^{\frac{1}{p}}}{y_3^2} (s_2 + s_3)} \\
&\times \frac{1}{1 - \frac{s_3}{y_3^4} - \left(\frac{s_3}{s_4} \right)^{\frac{1}{p}} \frac{1}{y_3^2} \left(y_4^2 - \frac{s_4}{y_4^2} \right) - \frac{2is_3^{\frac{1}{p}}}{y_3^2} (s_3 + s_4)} \tag{5.38}
\end{aligned}$$

The genus $g = 0$ case for four point function is obtained from (5.38). The four point function in genus zero has $\langle \tau_{0,1} \tau_{0,1} \tau_{0,p-2} \tau_{0,p-2} \rangle$, for $p \geq 4$, which is obtained from the coefficient of $s_1^{2/p} s_3^{2/p} s_2^{1-1/p} s_4^{1-1/p}$. We use the expression of (5.38). The expansion of the denominator of (5.38), we take a term of $2is_1^{\frac{1}{p}}(s_2 - s_4) \times 2is_3^{\frac{1}{p}}(s_4 - s_2) \sim 8s_1^{\frac{1}{p}} s_3^{\frac{1}{p}} s_2 s_4$. It becomes with $c = (\frac{1}{4})^p$,

$$\begin{aligned}
(4.1) \quad U(s_1, \dots, s_4) &= c^{\frac{2}{p}} s_1^{2/p} s_2^{1-1/p} s_3^{2/p} s_4^{1-1/p} \int_0^\infty \prod dy_i e^{-\sum y_i^{2p}} \left(\frac{128 y_2 y_4}{y_1^5 y_3^5} \right) \\
&= \frac{1}{8p^2} s_1^{2/p} s_2^{2/p} s_3^{1-1/p} s_4^{1-1/p} (\Gamma(1 - \frac{2}{p}))^2 (\Gamma(\frac{1}{p}))^2 \tag{5.39}
\end{aligned}$$

With a normalization factor $8p$, it leads to $\langle \tau_{0,1} \tau_{0,1} \tau_{0,p-2} \tau_{0,p-2} \rangle_{g=0} = \frac{1}{p}$.

For each $p = 3, 4, 5, 6, 7$, it is confirmed from (5.38) that $u(s_1, \dots, s_4)$ is described by (5.39) for $\langle \tau_{0,1} \tau_{0,1} \tau_{0,p-2} \tau_{0,p-2} \rangle$, namely

$$u(s_1, \dots, s_4) = c^{\frac{2}{p}} s_1^{2/p} s_2^{1-1/p} s_3^{2/p} s_4^{1-1/p} \int_0^\infty \prod dy_i e^{-\sum y_i^{2p}} \left(\frac{128 y_2 y_4 h^2}{y_1^5 y_3^5} \right) \tag{5.40}$$

where we put some parameter h (which should be 1) in the front of the term $s_i^{\frac{1}{p}}(s_i + s_{i+1})$ in the denominators of (5.38). Thus putting $h = 1$, leads to the intersection number $\frac{1}{p}$. This shows that the intersection number comes through the term of h . In these calculations, where the correction terms in the measure $s_i^{2+2/p}/y_i^3$ and the correction terms in the exponent of order of $s_i^{2+2/p}$ are neglected, since we are considering $g = 0$ contribution. In these four point functions for the intersection numbers, the representation of (5.38) is more useful than (5.37).

For $p = 7$, we find by the similar calculations for genus $g = 0$ with a normalization factor of $8p$ as

$$\langle \tau_{0,1} \tau_{0,2} \tau_{0,4} \tau_{0,5} \rangle = \langle \tau_{0,2} \tau_{0,2} \tau_{0,3} \tau_{0,5} \rangle = \langle \tau_{0,1} \tau_{0,1} \tau_{0,5} \tau_{0,5} \rangle = \frac{1}{7} \tag{5.41}$$

which are consistent with [13].

From the selection rule, the genus 0 case corresponds to $2p - 2 = \sum_{i=1}^4 j_i$ for $\langle \tau_{0,j_1} \tau_{0,j_2} \tau_{j_3} \tau_{j_4} \rangle$. Above result is consistent with

$$\langle \tau_{0,j_1} \tau_{0,j_2} \tau_{0,j_3} \tau_{0,j_4} \rangle_{g=0} = \frac{1}{p} \min(j_i, p - 1 + j_i) \quad (5.42)$$

This result is analyzed from the cohomological consideration in algebraic geometry [4].

We here evaluate $g = 0$ and $p = 3$ case, which will be used the result for the discussion of WDVV equation in the next section. We derive the three terms for four point function in (5.37) as

$$u(s_1, s_2, s_3, s_4) = d_1 s_1^{\frac{4}{3}} s_2^{\frac{1}{3}} s_3^{\frac{2}{3}} s_4^{\frac{1}{3}} + d_2 s_1^{\frac{5}{3}} s_2^{\frac{1}{3}} s_3^{\frac{1}{3}} s_4^{\frac{1}{3}} + d_3 s_1^{\frac{2}{3}} s_2^{\frac{2}{3}} s_3^{\frac{2}{3}} s_4^{\frac{2}{3}} \quad (5.43)$$

The third term of d_3 has been evaluated in (5.40). The integral of y_i becomes $d_3 = \frac{1}{72}$ with $c = (1/4)^3$. The first and the second term of d_1 and d_2 correspond to a dilation equation and string equation since they contain $\tau_{0,0}$ or $\tau_{1,0}$.

6 String, dilaton and WDVV equations

For integer value of p , the intersection numbers satisfy the following string equation and the dilaton equation.

$$\begin{aligned} \langle \tau_{0,0} \prod_{i=1}^s \tau_{n_i, m_i} \rangle_{g=0} &= \sum_{j=1}^s \langle \tau_{n_j-1, m_j} \prod_{i \neq j} \tau_{n_i, m_i} \rangle_{g=0} \\ \langle \tau_{1,0} \prod_{i=1}^s \tau_{n_j, m_j} \rangle_{g=0} &= (2g - 2 + s) \langle \prod_{i=1}^s \tau_{n_i, m_i} \rangle_{g=0} \end{aligned} \quad (6.1)$$

where $m_i = 0, 1, 2, \dots, p - 2$. These equation is for Neveu-Schwarz sector, and for $m_j = p - 1$, these correlations are vanishing (Ramond sector).

The string and dilaton equation can be derived from a matrix model of generalized Kotsevich model by Virasoro constraints, which are obtained by the equation of motion [42]

$$\int dB \frac{\partial}{\partial B_{ab}} e^{\frac{1}{p+1} \text{tr} B^{p+1} + \text{tr} B \Lambda} = 0 \quad (6.2)$$

or by using Gelfand-Dikii pseudo differential operator [41], which will be discussed in the next section. The string equation of (6.1) is equivalent to

$$\frac{\partial F}{\partial t_{0,0}} = \frac{1}{2} \sum_{i,j=0}^{p-2} \delta_{i+j,p-2} t_{0,i} t_{0,j} + \sum_{n=0}^{\infty} \sum_{m=0}^{p-2} t_{n+1,m} \frac{\partial F}{\partial t_{n,m}} \quad (6.3)$$

where the free energy, which is a generating function of the intersection numbers, is

$$F(t_{0,0}, t_{0,1}, \dots) = \sum_{d_{n,m}} \langle \prod_{n,m} t_{n,m}^{d_{n,m}} \rangle \prod_{n,m} \frac{t_{n,m}^{d_{n,m}}}{d_{n,m}!} \quad (6.4)$$

We prove the string and dilaton equations in the integral representation of $u(s_1, \dots, s_n)$, which is derived by a duality to above matrix model [43]. For two

point function $u(s_1, s_2)$, we have proved a string equation in (5.8), which is expressed as

$$\langle \tau_{0,0} \tau_{n,j} \rangle = \langle \tau_{n-1,j} \rangle \quad (6.5)$$

Since the left hand side is described by (5.5) and the right hand side is one point function of (5.2). The term of $s_1^{\frac{1}{p}}$ is present as a coefficient of the integral, the denominator can be put one. the difference to one point function is just a factor s_2 , which shifts n to $n-1$. as (5.8). Therefore, the string equation of (6.5) holds. This also holds for D type singularity, since the difference between A type and D type is the measure part of the integrals.

The selection rule of 2 point ($s=2$) is written as

$$3g-3+s = \sum_{i=1}^s n_i + (g-1)\left(1-\frac{2}{p}\right) + \frac{1}{p} \sum_{i=1}^s j_i \quad (6.6)$$

If we eliminate $\tau_{0,0}$ term ($n_1=0, j_1=0$) from above equation, s becomes $s-1$, and for the compensation to this change, n_2 should be changed to n_2-1 to maintain this equation of the selection rule when $s=2$.

For three marked point, the string equation becomes

$$\langle \tau_{0,0} \tau_{n_1, j_1} \tau_{n_2, j_2} \rangle = \langle \tau_{n_1-1, j_1} \tau_{n_2, j_2} \rangle + \langle \tau_{n_1, j_1} \tau_{n_2-1, j_2} \rangle \quad (6.7)$$

This equation satisfies the selection rule of (6.6). From (5.23) and (5.5), we are able to prove the string equation. We have an expression of $u(s_1, s_2, s_3)$ as (5.23). Taking the term of order $s_1^{\frac{1}{p}}$ in (5.23), it is expressed as

$$u(s_1, s_2, s_3) = \frac{(-8)s_1^{\frac{1}{p}} \int dy_1 \frac{1}{y_1^3} e^{-cy_1^{2p}} \int dy_2 dy_3 (y_2 + \frac{s_2}{y_2^2})(y_3 + \frac{s_3}{y_3^2})}{e^{-cy_2^{2p} - cy_3^{2p} + \dots}} \\ \frac{1}{s_3^{\frac{1}{p}} (y_2^2 - \frac{s_2}{y_2^2}) - s_2^{\frac{1}{p}} (y_3^2 - \frac{s_3}{y_3^2}) - 2i(s_2 s_3)^{\frac{2}{p}} (s_2 + s_3)} \quad (6.8)$$

By the exchange of $s_2 \leftrightarrow s_3, y_2 \leftrightarrow y_3$, we obtain $u(s_1, s_3, s_2)$. By adding these two terms, and using the expression of $u(s_2, s_3)$ in (5.5), we find

$$\frac{1}{2} \left(u(s_1, s_2, s_3) + u(s_1, s_3, s_2) \right) = i s_1^{\frac{1}{p}} \left(\int dy_1 \frac{1}{y_1^3} e^{-cy_1^{2p}} \right) u(s_2, s_3) (s_2 + s_3) \quad (6.9)$$

Noting that $u(s_1, s_2, s_3) = \sum \langle \tau_{0,0} \tau_{n_2, m_2} \tau_{n_3, m_3} \rangle = s_1^{\frac{1}{p}} s_2^{\frac{n_2 + \frac{1+m_2}{p}}{p}} s_3^{\frac{n_3 + \frac{1+m_3}{p}}{p}}$ and $u(s_2, s_3) = \sum \langle \tau_{n_2, m_2} \tau_{n_3, m_3} \rangle = s_2^{\frac{n_2 + \frac{1+m_2}{p}}{p}} s_3^{\frac{n_3 + \frac{1+m_3}{p}}{p}}$, (6.9) becomes a string equation of

$$\langle \tau_{0,0} \tau_{n_2, m_2} \tau_{n_3, m_3} \rangle = \langle \tau_{n_2-1, m_2} \tau_{n_3, m_3} \rangle + \langle \tau_{n_2, m_2} \tau_{n_3-1, m_3} \rangle \quad (6.10)$$

This is a proof of the string equation for three marked points. For higher marked points, we are able to prove similarly from the expressions of $u(s_1, \dots, s_n)$. The derivation of four point function comes from the determinant [6]. For a 4×4

determinant, the components of the connected longest cycles are 6 terms as

$$\text{Det} \begin{vmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{vmatrix} = -(a_{12}a_{23}a_{34}a_{41} + a_{13}a_{32}a_{24}a_{41} + a_{14}a_{23}a_{31}a_{42} \\ + a_{13}a_{21}a_{34}a_{42} + a_{12}a_{24}a_{31}a_{43} + a_{14}a_{21}a_{32}a_{43}) \quad (6.11)$$

where

$$a_{ij} = \frac{1}{u_i - u_j + \frac{1}{2}(s_i + s_j)} \quad (6.12)$$

By adding these terms, with the variables y_i , we obtain as same as three point function,

$$u(s_1, s_2, s_3, s_4) = C s_1^{\frac{1}{p}} (s_2 + s_3 + s_4) u(s_2, s_3, s_4) \quad (6.13)$$

which leads to the string equation,

$$\langle \tau_{0,0} \tau_{n_2, m_2} \tau_{n_3, m_3} \tau_{n_4, m_4} \rangle = \sum_{j=2}^4 \langle \tau_{n_j-1, m_j} \prod_{i \neq j} \tau_{n_i, m_i} \rangle \quad (6.14)$$

The equation of elimination of $\tau_{0,1}$ is related to W-constraints [12]. For two marked point in genus $g = 2$, we find for p , the following relation which is suggested by the selection rule of (6.6),

$$\langle \tau_{0,1} \tau_{n,j} \rangle_{g=2} = C \langle \tau_{n-1, j+1} \rangle_g \quad (6.15)$$

From the general p analysis, we obtain [32]

$$\langle \tau_{0,1} \tau_{4,1} \rangle_{g=2} = \frac{(p-1)(p-2)(p+2)}{2880p}, \quad \langle \tau_{3,2} \rangle_{g=2} = \frac{(p-1)(2p+1)(p-3)}{5760p} \quad (6.16)$$

Then the ratio C is given for this case by

$$C = \frac{2(p-2)(p+2)}{(2p+1)(p-3)} \quad (6.17)$$

This result is consistent with the case of $\langle \tau_{0,1} \tau_{4,1} \rangle_{g=2} = \frac{8}{3} \langle \tau_{3,2} \rangle_{g=2} = \frac{1}{320}$ for $p = 4$, and $\langle \tau_{0,1} \tau_{4,1} \rangle_{g=2} = \frac{21}{11} \langle \tau_{3,2} \rangle_{g=2} = \frac{7}{1200}$ for $p = 5$.

• dilaton equation

The dilaton equation is written as (6.1) for $\tau_{1,0}$, which comes from $s^{1+\frac{1}{p}}$ dependence.

$$\langle \tau_{1,0} \prod_{i=1}^s \tau_{n_i, j_i} \rangle_{g=2} = (2g-2+s) \langle \prod_{i=1}^s \tau_{n_i, j_i} \rangle_g \quad (6.18)$$

For two point function in (5.5), the relevant term of $s_1^{1+\frac{1}{p}}$ is obtained by the expansion of the denominator in (5.5). The denominator is expressed by the neglecting $s_1^{2+\frac{2}{p}}$ term as

$$\left(\frac{1}{-4i \frac{s_1^{\frac{1}{p}} s_2}{y_1^2}} \right) \left[\frac{1}{1 - \left(\frac{s_1}{s_2} \right)^{\frac{1}{p}} \left(\frac{y_2}{y_1} \right)^2 \left(1 - \frac{s_2^{\frac{2}{p}}}{y_2^4} \right) + 2i \frac{s_1^{\frac{1}{p}} s_2}{y_1^2}} - c.c \right] \quad (6.19)$$

where *c.c.* means a complex conjugate. Since the first factor $s_1^{\frac{1}{p}}$ in (6.19) is cancelled with a factor of $s_1^{\frac{1}{p}}$ in the front of the integral (5.5), we need to derive the terms of order of $s_1^{1+\frac{1}{p}}$ from (6.19). From the expansion of the denominator, we have $s_1^{1+\frac{1}{p}} \frac{1}{y_1^{2p+2}} Z^{p+1}$ term, where Z is

$$\begin{aligned} Z &= s_2^{-\frac{1}{p}} y_2^2 \left(1 - \frac{s_2^{\frac{2+\frac{2}{p}}{p}}}{y_2^4}\right) - 2i s_2 \\ &= s_2^{-\frac{1}{p}} \left(y_2 - i \frac{s_2^{1+\frac{1}{p}}}{y_2}\right)^2 \end{aligned} \quad (6.20)$$

By taking the power of $(p+1)$, it becomes

$$Z^{p+1} = s_2^{-1-\frac{1}{p}} \left(y_2 - i \frac{s_2^{1+\frac{1}{p}}}{y_2}\right)^{2p+2} \quad (6.21)$$

By subtraction of the complex conjugate term $Z^{*(p+1)}$, it is remarkably same as the exponent of (5.5); $e^{-c[(y_2 - \frac{i s_2^{1+\frac{1}{p}}}{y_2})^{2p+2} - (y_2 - \frac{i s_2^{1+\frac{1}{p}}}{y_2})^{2p+2}]}$. Thus the factor $Z^{p+1} - Z^{*(p+1)}$ is expressed as the derivative by the coefficient c in the exponent,

$$\frac{d}{dc} u(s_2) = \frac{d}{dc} \left\{ \sum_{n,j} \langle \tau_{n,j} \rangle_g s_2^{n+\frac{1+j}{p}} \right\} \quad (6.22)$$

From the scaling cs^{1+p} , and the selection rule $2g-1 = n + \frac{1}{p}(j-2(g-1))$, the term of $\langle \tau_{n,j} \rangle$ is order of $c^{\frac{p}{p+1}(n+\frac{1+j}{p})}$. The derivative of c gives a factor $\frac{p}{p+1}(n+\frac{1+j}{p})$. This factor is written by the selection rule of (2.6) as

$$2g-1 = \frac{p}{p+1} \left(n + \frac{1+j}{p}\right) \quad (6.23)$$

The left hand side coincides with a coefficient of dilaton equation of (6.18) for one-point ($s=1$) case. The integral of y_1 gives a gamma function, which is multiplied to (6.22). Thus we prove the dilaton equation of two point function, which satisfies (6.18) for the positive integer p . (This holds also for D type).

For three point function, the proof of the dilaton equation is similar to two point function of (6.22). The integral formula of three point function $u(s_1, s_2, s_3)$ provides the term of $s_1^{1+\frac{1}{p}}$ by the expansion of the denominator of (5.23). We denote three denominators as g_1, g_2, g_3 in (5.23) in orders. We neglect the terms of $s_1^{\frac{2+\frac{2}{p}}{p}}$ in them for a dilaton equation. Since g_1 and g_3 involve the factor $s_1^{\frac{1}{p}}$, we write as

$$\frac{1}{1-g_1} \frac{1}{1-g_3} = \left(\frac{1}{1-g_1} - \frac{1}{1-g_3}\right) \frac{1}{g_1-g_3} \quad (6.24)$$

The term $\frac{1}{g_1-g_3}$ is given by the neglecting a factor $s_1^{\frac{2+\frac{2}{p}}{p}}$,

$$\frac{1}{g_1-g_3} = \left(\frac{y_1^2}{s_1^{\frac{1}{p}}}\right) \frac{s_2^{\frac{1}{p}}}{y_2^2 - \frac{s_2^{\frac{2+\frac{2}{p}}{p}}}{y_2^2} - \left(\frac{s_2}{s_3}\right)^{\frac{1}{p}} \left(y_3^2 - \frac{s_3^{\frac{2+\frac{2}{p}}{p}}}{y_3^2}\right) + 2i s_2^{\frac{1}{p}} (s_2 + s_3)} \quad (6.25)$$

The factor $\frac{1}{y_2^2(1-g_2)}$ is written,

$$\frac{1}{y_2^2(1-g_2)} = \frac{1}{y_2^2 - \frac{s_2^{2+\frac{2}{p}}}{y_2^2} - (\frac{s_2}{s_3})^{\frac{1}{p}}(y_3^2 - \frac{s_3^{2+\frac{2}{p}}}{y_3^2}) - 2is_2^{\frac{1}{p}}(s_2 + s_3)}. \quad (6.26)$$

We obtain the term of product of these factors as

$$\frac{1}{y_2^2(1-g_2)} \frac{1}{g_1 - g_3} = \frac{y_1^2 s_2^{\frac{1}{p}}}{s_1^{\frac{1}{p}}} \frac{1}{(y_2^2 - \frac{s_2^{2+\frac{2}{p}}}{y_2^2} - (\frac{s_2}{s_3})^{\frac{1}{p}}(y_3^2 - \frac{s_3^{2+\frac{2}{p}}}{y_3^2}))^2 + 4s_2^{\frac{2}{p}}(s_2 + s_3)^2} \quad (6.27)$$

which is proportional to $u(s_2, s_3)$ in (5.5). From the the expansion of the factor $\frac{1}{1-g_1}$,

$$\frac{1}{1-g_1} = \frac{1}{1 - (\frac{s_1}{s_2})^{\frac{1}{p}} \frac{1}{y_1} (y_2 - i \frac{s_2^{1+\frac{1}{p}}}{y_2})^2} \quad (6.28)$$

we have a term g_1^{p+1} , which is order of $s_1^{1+\frac{1}{p}}$ as

$$\frac{s_1^{1+\frac{1}{p}}}{y_1^{2(p+1)}} \frac{1}{s_2^{1+\frac{1}{p}}} (y_2 - i \frac{s_2^{1+\frac{1}{p}}}{y_2})^{2p+2} \quad (6.29)$$

Noting that the exponential term of $u(s_2, s_3)$ in (5.5) has $(y_2 - i \frac{s_2^{1+\frac{1}{p}}}{y_2})^{2p+2}$ term, the dilaton term of $s_1^{1+\frac{1}{p}}$ is obtained by the derivative of the exponential term (Hamiltonian) by the coefficient c . Two point function $u(s_2, s_3)$ is

$$u(s_2, s_3) = \sum \langle \tau_{n_2, m_2} \tau_{n_3, m_3} \rangle s_2^{n_2 + \frac{1+m_2}{p}} s_3^{n_3 + \frac{1+m_3}{p}} \quad (6.30)$$

Thus we have

$$\frac{d}{dc} u(s_2, s_3) = \sum_{n_2, m_2, n_3, m_3} \frac{d}{dc} c^{\frac{p}{p+1}(n_2+n_3+\frac{1}{p}(m_2+m_3+2))} s_2^{n_2 + \frac{1+m_2}{p}} s_3^{n_3 + \frac{1+m_3}{p}} \quad (6.31)$$

From the selection rule of (2.6), we have

$$2(g-1) + (s-1) = \frac{p}{p+1} \left(\sum_{j=2}^s n_j + \frac{1}{p} \sum_{j=2}^s (1+m_j) \right) \quad (6.32)$$

where we put $s = 3$ for three point function. Thus the constant of (6.31) is $2g - 2 + (s - 1)$, which provides the dilaton equation.

For half integer p , above argument does not hold since the power $p + 1$ of Z^{p+1} becomes not integer, and the geometrical series of the integer p does not work. Indeed we found that there is no dilaton equation for $p = \frac{1}{2}$. The term of the dilaton $\tau_{1,0}$ corresponds to $s^{1+\frac{1}{p}}$, which becomes s^3 for $p = \frac{1}{2}$. The term of s^3 does not exist in the contour integral of $u(s_1, s_2)$ for $p = \frac{1}{2}$. We took a spin component j as -1 for $p = \frac{1}{2}$.

The absence of the dilaton equation for $p = \frac{1}{2}$ may be related to the simplicity of Faber conjecture [35]. We now discuss the case of $p = \frac{1}{2}$.

• **Gelfand-Dikii and WDVV equation**

The computational algorithm of the intersection numbers of p -spin curves has been found by the use of Gelfand-Dikii equation [3, 1, 2]. In this section, we briefly discuss the relation of our integral formula to Gelfand-Dikii residual calculations [4, 10, 11].

The differential operator Q of Gelfand-Dikii equation is defined as

$$Q = D^p + \sum_{i=0}^{p-2} \gamma_i(x) D^i \quad (6.33)$$

where $D = \frac{i}{\sqrt{p}} \frac{\partial}{\partial x}$. The pseudo differential operator with a fractional power $Q^{\frac{1}{p}}$ is defined as

$$Q^{\frac{1}{p}} = D + \sum_{i>0} \alpha_i D^{-i} \quad (6.34)$$

Gelfand-Dikii equation is

$$i \frac{\partial Q}{\partial t_{n,m}} = \frac{c_{n,m}}{\sqrt{p}} [Q_+^{n+\frac{m}{p}}, Q] \quad (6.35)$$

where

$$c_{n,m} = \frac{(-1)^n p^{n+1}}{(m+1)(p+m+1) \cdots (np+m+1)} \quad (6.36)$$

We have a relation of the residue,

$$\langle\langle \tau_{0,0} \tau_{n,m} \rangle\rangle = \frac{\partial^2}{\partial t_{0,0} \partial t_{n,m}} F = -c_{n,m} \text{Res}(Q^{n+\frac{1+m}{p}}) \quad (6.37)$$

Since the evaluation of Gelfand-Dikii equation is a bit complicated, here we restrict ourself to the case $p = 3$, for instance [35]. More details for general p are shown in [4, 11]. The differential operator Q becomes

$$Q = D^3 + \gamma_1 D + \gamma_2 \quad (6.38)$$

which is written as

$$Q = D^3 + 3 \text{Res}(Q^{\frac{1}{3}}) D + \frac{3}{2} \text{Res}(Q^{\frac{2}{3}}) + D \text{Res}(Q^{\frac{1}{3}}) \quad (6.39)$$

Since x is same as $t_{0,0}$, the last term is $-\frac{i}{2\sqrt{3}} \langle\langle \tau_{0,0}^3 \rangle\rangle$. Thus we have

$$\gamma_1 = -\langle\langle \tau_{0,0}^2 \rangle\rangle, \quad \gamma_2 = -\langle\langle \tau_{0,0} \tau_{0,1} \rangle\rangle - \frac{i}{2\sqrt{3}} \langle\langle \tau_{0,0}^3 \rangle\rangle \quad (6.40)$$

The pseudo differential operator $Q^{n-1+\frac{m}{p}}$ is

$$Q^{n-1+\frac{m}{3}} = \alpha_1 D^{-1} + \alpha_2 D^{-2} + \alpha_3 D^{-3} + \cdots \quad (6.41)$$

We obtain

$$\text{Res}(Q^{n+\frac{m}{3}}) = \alpha_2 \gamma_1 + \alpha_1 \gamma_2 + \alpha_4 - \alpha_1 D \gamma_1 \quad (6.42)$$

From Gelfand-Dikii equation of (6.35), we have

$$\begin{aligned}
\frac{i\sqrt{3}}{c_{n-1,m}} \frac{\partial \gamma_1}{\partial t_{n-1,m}} &= 3D\alpha_1, \\
\frac{i\sqrt{3}}{c_{n-1,m}} \frac{\partial \gamma_2}{\partial t_{n-1,m}} &= 3D^2\alpha_1 + 3D\alpha_2, \\
\frac{i\sqrt{3}}{c_{n,m}} \frac{\partial \gamma_1}{\partial t_{n,m}} &= 3DRes(Q^{n+\frac{m}{3}})
\end{aligned} \tag{6.43}$$

Then, we obtain the following equation [35]

$$\begin{aligned}
(3n+m+1) \langle\langle \tau_{n,m} \tau_{0,0}^2 \rangle\rangle &= \langle\langle \tau_{n-1,m} \tau_{0,1} \rangle\rangle \langle\langle \tau_{0,0}^3 \rangle\rangle \\
+ 2 \langle\langle \tau_{n-1,m} \tau_{0,0} \rangle\rangle \langle\langle \tau_{0,1} \tau_{0,0}^2 \rangle\rangle & \\
+ 2 \langle\langle \tau_{n-1,m} \tau_{0,1} \tau_{0,0} \rangle\rangle \langle\langle \tau_{0,0}^2 \rangle\rangle & \\
+ \frac{2}{3} \langle\langle \tau_{n-1,m} \tau_{0,1} \tau_{0,0}^3 \rangle\rangle + 3 \langle\langle \tau_{n-1,m} \tau_{0,0}^2 \rangle\rangle \langle\langle \tau_{0,1} \tau_{0,0} \rangle\rangle & \tag{6.44}
\end{aligned}$$

This equation provides the intersection numbers for $p = 3$ case.

WDVV equation [1, 2] is written as for $p = 2$, which is directly related to KdV equations,

$$\langle\langle \tau_{1,0} \tau_{0,0} \rangle\rangle_g = \frac{1}{2} \langle\langle \tau_{0,0}^2 \rangle\rangle_{g'} \langle\langle \tau_{0,0}^2 \rangle\rangle_{g-g'} + \frac{1}{12} \langle\langle \tau_{0,0}^4 \rangle\rangle_{g-1} \tag{6.45}$$

where the notation of $\langle\langle \tau_{1,0} \tau_{0,0} \rangle\rangle$ is $\frac{\partial^2 F}{\partial t_{1,0} \partial t_{0,0}}$. The free energy F is expressed as

$$\begin{aligned}
F &= \sum_g \sum_{d_{n,m}} \langle \prod_{n,m} \tau_{n,m}^{d_{n,m}} \rangle_g \prod_{n,m} \frac{t_{n,m}^{d_{n,m}}}{d_{n,m}!} \\
&= \sum_g F_g
\end{aligned} \tag{6.46}$$

The free energy F of genus zero ($g = 0$) is expressed for $p = 2$ as

$$F_{g=0} = \frac{t_0^3}{3!} + \frac{t_0^3 t_1}{3!} + 2 \cdot \frac{t_0^3 t_1^2}{3!2!} + \frac{t_0^4 t_2}{4!} + 6 \frac{t_0^3 t_1^3}{3!3!} + \dots \tag{6.47}$$

where t_n means $t_{n,0}$. For $p = 3$, genus 0 free energy becomes

$$\begin{aligned}
F_{g=0} &= \frac{1}{2} t_{0,0}^2 t_{0,1} + \frac{1}{72} t_{0,1}^4 + \frac{1}{2} t_{1,0} t_{0,1} t_{0,0}^2 + \frac{1}{6} t_{1,1} t_{0,0}^3 \\
&+ \frac{1}{6} t_{2,0} t_{0,1} t_{0,0}^3 + \frac{1}{3} t_{1,1} t_{1,0} t_{0,0}^3 + \frac{1}{4} t_{1,0}^2 t_{0,1} t_{0,0}^2 + O(t^6)
\end{aligned} \tag{6.48}$$

These terms are obtained from the selection rule and the string equation. The last three terms are derived from (5.43). From above expressions, with the notation of $\langle\langle \dots \rangle\rangle_g$ as the derivatives of F_g , we find for $p = 2$ case,

$$\begin{aligned}
\langle\langle \tau_0 \tau_1 \rangle\rangle_{g=0} &= \frac{1}{2} t_0^2 + t_0^2 t_1 + \frac{3}{2} t_0^2 t_1^2 + \dots \\
\langle\langle \tau_0^2 \rangle\rangle_{g=0} &= t_0 + t_0 t_1 + t_1^2 t_0 + \frac{1}{2} t_0^2 t_2 + \dots
\end{aligned} \tag{6.49}$$

which satisfies $\langle\langle \tau_0 \tau_1 \rangle\rangle_{g=0} = \frac{1}{2} [\langle\langle \tau_0^2 \rangle\rangle_{g=0}]^2$ of (6.45).

For $p = 2$, the free energy F is a τ -function for the KdV hierarchy. The quantity $U = \frac{\partial^2 F}{\partial t_{0,0}^2}$ follows all equations in the KdV equation. The first equation of KdV hierarchy is

$$\frac{\partial U}{\partial t_{1,0}} = \frac{\partial^3 F}{\partial t_{1,0} \partial t_{0,0}^2} = \left(\frac{\partial^2 F}{\partial t_{0,0}^2} \right) \left(\frac{\partial^3 F}{\partial t_{0,0}^3} \right) + \frac{1}{12} \frac{\partial^5 F}{\partial t_{0,0}^5} \quad (6.50)$$

This equation is same as (6.45) with the integration of $t_{0,0}$.

The n -point function of $g = 0$ is obtained from previous n point functions for $p = 2$ as

$$\begin{aligned} u(s_1, s_2, s_3) &= \frac{1}{4} \left(\Gamma\left(\frac{1}{2}\right) \right)^3 (s_1 s_2 s_3)^{\frac{1}{2}} \\ u(s_1, s_2, s_3, s_4) &= \frac{1}{16} \left(\Gamma\left(\frac{1}{2}\right) \right)^4 (s_1 s_2 s_3)^{\frac{1}{2}} s_4^{\frac{3}{2}} \\ u(s_1, s_2, s_3, s_4, s_5) &= \frac{1}{16} \left(\Gamma\left(\frac{1}{2}\right) \right)^5 (s_1 s_2 s_3)^{\frac{1}{2}} (s_4 s_5)^{\frac{3}{2}} + \frac{1}{16} \left(\Gamma\left(\frac{1}{2}\right) \right)^5 (s_1 s_2 s_3 s_4)^{\frac{1}{2}} s_5^{\frac{5}{2}} \end{aligned} \quad (6.51)$$

These terms of $g = 0$ correspond to F in (6.47). We assign $s_1^{n+\frac{1+j}{p}}$ to $t_{n,j}$.

By repeating the application of the string equation and from $\langle \tau_0^3 \rangle = 1$, we find $\langle \tau_0^3 \tau_1 \rangle = 1$, $\langle \tau_0^3 \tau_1^2 \rangle = 2$, $\langle \tau_0^4 \tau_2 \rangle = 1$, $\langle \tau_0^3 \tau_1^3 \rangle = 6$, $\langle \tau_0^4 \tau_2 \tau_1 \rangle = 3$, $\langle \tau_0^5 \tau_3 \rangle = 1$ for $g = 0, p = 2$. From these intersection numbers, we obtain the expression of the free energy F for $p = 2$ in genus $g = 0$ as (6.47).

For genus one in the case $p = 2$, the free energy F becomes

$$\begin{aligned} F_{g=1} &= \frac{1}{24} t_1 + \frac{1}{48} t_1^2 + \frac{1}{24} t_0 t_2 + \frac{1}{72} t_1^3 + \frac{1}{12} t_0 t_1 t_2 + \frac{1}{48} t_0^2 t_3 \\ &+ \frac{1}{8} t_0 t_1^2 t_2 + \frac{1}{24} t_0^2 t_2^2 + \frac{1}{16} t_0^2 t_1 t_3 + \frac{1}{96} t_1^4 + \frac{1}{24} t_0^4 t_2 + \frac{1}{8} t_0^4 t_1 t_2 \\ &+ \frac{1}{120} t_0^5 t_3 + \dots \end{aligned} \quad (6.52)$$

From this, by taking the derivatives, we find

$$\langle\langle \tau_{1,0} \tau_{0,0} \rangle\rangle_{g=1} = \frac{1}{12} t_2 + \frac{1}{4} t_1 t_2 + \frac{1}{8} t_0 t_3 + \dots \quad (6.53)$$

which satisfies WDVV equation of (6.45).

$$\left(\frac{\partial^2 F}{\partial t_{1,0} \partial t_{0,0}} \right)_{g=1} = \left(\frac{\partial^2 F}{\partial t_{0,0}^2} \right)_{g=1} \left(\frac{\partial^2 F}{\partial t_{0,0}^2} \right)_{g=0} + \frac{1}{12} \left(\frac{\partial^4 F}{\partial t_{0,0}^4} \right)_{g=0} \quad (6.54)$$

The genus zero case of WDVV equation is equivalent to the string equation and the dilaton equation. The string equation for p -spin curves is

$$\frac{\partial F}{\partial t_{0,0}} = \frac{1}{2} \sum_{i,j=0}^{p-2} \delta_{i+j,p-2} t_{0,i} t_{0,j} + \sum_{n=0}^{\infty} \sum_{m=0}^{p-2} t_{n+1,m} \frac{\partial F}{\partial t_{n,m}} \quad (6.55)$$

or equivalently, it is written as

$$\langle \tau_{0,0} \prod_{i=1}^s \tau_{n_i, m_i} \rangle_g = \sum_{j=1}^s \langle \tau_{n_j-1, m_j} \prod_{i=1, i \neq j}^s \tau_{n_i, m_i} \rangle_g \quad (6.56)$$

For $p = 3$, it becomes [10, 11]

$$\langle\langle \tau_{1,0}\tau_{0,0} \rangle\rangle_g = \langle\langle \tau_{0,0}\tau_{0,1} \rangle\rangle_{g'} \langle\langle \tau_{0,0}^2 \rangle\rangle_{g-g'} + \frac{1}{6} \langle\langle \tau_{0,0}^3\tau_{0,1} \rangle\rangle_{g-1} \quad (6.57)$$

It is related to Boussinesq equation with variables $t_{0,0}$, $t_{0,1}$ and $t_{1,0}$. From the explicit expression of free energy of (6.48), we verify above equation for $g = 0$.

Our formula for n point function provides the coefficients of $\prod_{i=1}^n s_i^{n_i + \frac{1+m_i}{p}}$, which are the intersection numbers as shown in the previous sections. The free energy is consist of the sum of the n point functions as

$$F = \sum_{n=1}^{\infty} u(s_1, \dots, s_n) \quad (6.58)$$

So $s^{n+\frac{1+m}{p}}$ is regarded as $t_{n,m}$ in the free energy of (6.46). The derivative of the free energy $\langle\langle \tau_{0,1}\tau_{0,0} \rangle\rangle$ is a coefficient of $s_1^{\frac{2}{p}} s_2^{\frac{1}{p}}$ in the n -point functions.

$$\langle\langle \tau_{0,1}\tau_{0,0} \rangle\rangle_g = \sum_k \langle \tau_{0,1}\tau_{0,0} \prod_{i=1}^k \tau_{n_i, m_i} \rangle_g \prod_{i=1}^k s_i^{n_i + \frac{1+m_i}{p}} \quad (6.59)$$

where genus g follows the selection rule of (2.6).

WDVV equation is written as [11] for genus zero,

$$\begin{aligned} & \sum_{m', m''}^{p-2} \prod_{\{5, \dots, s\} = I \cup J} \langle \tau_{n_1, m_1} \tau_{n_2, m_2} (\prod_{i \in I} \tau_{n_i, m_i}) \tau_{0, m'} \rangle_{g=0} \\ & \eta^{m', m''} \langle \tau_{0, m''} \tau_{n_3, m_3} \tau_{n_4, m_4} (\prod_{i \in J} \tau_{n_i, m_i}) \rangle_{g=0} \\ & = \sum_{m', m''}^{p-2} \prod_{\{5, \dots, s\} = I \cup J} \langle \tau_{n_1, m_1} \tau_{n_3, m_3} (\prod_{i \in I} \tau_{n_i, m_i}) \tau_{0, m'} \rangle_{g=0} \\ & \eta^{m', m''} \langle \tau_{0, m''} \tau_{n_2, m_2} \tau_{n_4, m_4} (\prod_{i \in J} \tau_{n_i, m_i}) \rangle_{g=0} \end{aligned} \quad (6.60)$$

More explicitly, for four point intersection numbers, we have [4]

$$\begin{aligned} & \langle \tau_{n_1+1, m_1} \tau_{n_2, m_2} \tau_{n_3, m_3} \tau_{n_4, m_4} \rangle \\ & = \sum_{m', m''} \langle \tau_{n_1, m_1} \tau_{n_2, m_2} \tau_{0, m'} \rangle \eta^{m', m''} \langle \tau_{0, m''} \tau_{n_3, m_3} \tau_{n_4, m_4} \rangle \end{aligned} \quad (6.61)$$

where $\eta^{m', m''} = \delta_{m'+m'', p-2}$. Above equation is generalized to

$$\begin{aligned} & \langle\langle \tau_{n_1, m_1} \tau_{n_2, m_2} \tau_{0, m'} \rangle\rangle \eta^{m', m''} \langle\langle \tau_{0, m''} \tau_{n_3, m_3} \tau_{n_4, m_4} \rangle\rangle \\ & = \langle\langle \tau_{n_1, m_1} \tau_{n_3, m_3} \tau_{0, m'} \rangle\rangle \eta^{m', m''} \langle\langle \tau_{0, m''} \tau_{n_2, m_2} \tau_{n_4, m_4} \rangle\rangle \end{aligned} \quad (6.62)$$

In the next section, we will investigate the case of $p = \frac{1}{2}, -\frac{1}{2}$, for which Gelfand-Dikii pseudo differential method does not work since $Q^{n+\frac{1+m}{p}}$ commutes with Q .

7 n-point functions of $p = \frac{1}{2}, -\frac{1}{2}, -\frac{3}{2}$ and $-2, -3$

In this section, we consider the non-positive integer $p = \frac{1}{2}, -\frac{1}{2}, -\frac{3}{2}$ and $p = -2, -3$ cases. Some of these one point functions are derived in the previous articles [8, 9].

7-1: n-point function for $p = \frac{1}{2}$

Since $\frac{1}{p} \sum j_i$ becomes integer in this case, we are able to include them in the integer part of $\sum n_i$. Thus the spin component j_i can be chosen as arbitrary value as mod p . Here we take the spin component j_i as -1 . Other choice may be $j = 0$ or $j = -\frac{1}{2}$. All these cases provide the shift of the integer n . From the continuation of Ramond spin component $j = p - 1$, the choice of $j = -\frac{1}{2}$ may be considered a natural value. However, the difference between $j = -1$ and $j = -\frac{1}{2}$ leads the shift of the integer n , this choice may do not cause a serious conclusion. For the integer p , the algebraic geometry has been studied [4]. For $p = \frac{1}{2}$ case, the study of algebraic geometry has not been investigated.

One point function $u(s)$ is given by (4.28), which becomes after a partial integral,

$$u(s) = -\frac{i}{6c's^{3/2}} \oint \frac{dy}{2i\pi} e^{c's^{3/2}(3y - \frac{1}{y^3})} \quad (7.1)$$

and it becomes a series of $\sum_g a_g s^{6g-3}$.

• string equation for $p = \frac{1}{2}$

We discuss the string equation for $p = \frac{1}{2}$. Although the usual string equation may be not applied for this case, analogous equation about a forgetting of a marked point can be written as (6.1).

The selection rule for $p = \frac{1}{2}$ becomes from (2.6) for s marked points,

$$6g - 6 + s = \sum_{i=1}^s n_i + 2 \sum_{i=1}^s j_i \quad (7.2)$$

Possible values of j_i may be $0, -\frac{1}{2}, -1$, for which the last term becomes integers. The term $s_i^{n_i + \frac{1+m_i}{p}}$ becomes $s_i^{n_i+2}, s_i^{n_i+1}$ and $s_i^{n_i}$, respectively.

By taking $j_i = -1$, it becomes

$$6g - 6 + 3s = \sum_{i=1}^s n_i \quad (7.3)$$

The term of order $s_1^2 s_2^{6g-2}$ are derived from (5.5), which is considered as a string equation, since $s^{\frac{1}{p}} = s^2$ for $p = \frac{1}{2}$.

$$\begin{aligned} u(s_1, s_2) &= -\frac{4s_1^2}{s_2^2} \oint \frac{dy_1 dy_2}{(2i\pi)^2} \frac{1}{y_1^3} (y_2 \pm \frac{s_2^6}{y_2^3}) e^{c_1(3y_1) + c_2(3y_2 - \frac{s_2^6}{y_2^3})} \\ &= \sum_g a_g s_1^2 s_2^{6g-2} c_1^2 c_2^{4g-2} \end{aligned} \quad (7.4)$$

The coefficient a_g is denoted by the intersection numbers as $\langle \tau_{2,-1} \tau_{6g-2,-1} \rangle$, where we defined n of $\tau_{n,-1}$ as the power of $s^{n + \frac{1+j}{p}}$. $\langle \tau_{n,-1} \rangle$ is a coefficient

of s^n . This term of order s_2^{6g-2} is one point function in (4.28). Thus we have a string equation,

$$\langle \tau_{2,-1} \tau_{6g-2,-1} \rangle = \langle \tau_{6g-3,-1} \rangle \quad (7.5)$$

in which $\tau_{2,-1}$ operates on $\tau_{6g-2,-1}$ for the change to $\tau_{6g-3,-1}$. Note that the usual notation of $\tau_{n,j}$ is different from above as a shift $n \rightarrow n-2$, since we took $j_i = -1$.

Three points function of $p = \frac{1}{2}$ is studied for A_l case in [9]. From the selection rule of $j_i = -1$ ($i = 1, 2, 3$), we have from (7.3) ; $6g+3 = n_1 + n_2 + n_3$ for $\langle \tau_{n_1,-1} \tau_{n_2,-1} \tau_{n_3,-1} \rangle$.

The term of order s_1^2 is obtained from three point function as

$$u(s_1, s_2, s_3) = -\frac{s_1^2}{(s_2 s_3)^2} \oint \frac{dy_1}{2i\pi} \frac{1}{y_1^3} e^{3c_1 y_1} \oint \frac{dy_2 dy_3}{(2i\pi)^2} (y_2 \pm \frac{s_2^6}{y_2^6})(y_3 \pm \frac{s_3^6}{y_3^6}) e^{c_2(3y_2 - \frac{s_2^6}{y_2^6}) + c_3(3y_3 - \frac{s_3^6}{y_3^6})} \frac{1}{2i(s_2 + s_3) + \frac{y_2^2}{s_2^2} - \frac{s_2^4}{y_2^2} - \frac{y_3^2}{s_3^2} + \frac{s_3^4}{y_3^2}} \quad (7.6)$$

Since two point function $u(s_2, s_3)$ is written as

$$u(s_2, s_3) = \frac{i}{s_2^2 s_3^2 (s_2 + s_3)} \oint \frac{dy_2 dy_3}{(2i\pi)^2} (y_2 \pm \frac{s_2^6}{y_2^6})(y_3 \pm \frac{s_3^6}{y_3^6}) e^{c_2(3y_2 - \frac{s_2^6}{y_2^6}) + c_3(3y_3 - \frac{s_3^6}{y_3^6})} \left(\frac{1}{\frac{y_2^2}{s_2^2} - \frac{s_2^4}{y_2^2} - \frac{y_3^2}{s_3^2} + \frac{s_3^4}{y_3^2} - 2i(s_2 + s_3)} - c.c. \right) \quad (7.7)$$

Thus we have a string equation

$$\langle \tau_{2,-1} \tau_{n_2,-1} \tau_{n_3,-1} \rangle_g = \langle \tau_{n_2-1,-1} \tau_{n_3,-1} \rangle_g + \langle \tau_{n_2,-1} \tau_{n_3-1,-1} \rangle_g \quad (7.8)$$

The difference between A_l and D_l is due to the sign (± 1) in the representation of y variable in (4.24), and the string equation of three points function for A_l is valid also for D_l case.

• **two point function and absence of dilaton equation for $p = \frac{1}{2}$**

For integer spin p , a dilaton field corresponds to $\tau_{1,0}$, which corresponds to $s^{1+\frac{1}{p}}$. When $p = \frac{1}{2}$ is inserted into this term, it becomes s^3 term.

We have found that there is no dilaton equation for $p = \frac{1}{2}$ [9]. The one marked point intersection number $\langle \tau_{1,0} \rangle_{g=1}$ corresponds to $\langle \tau_{3,-1} \rangle_{g=1}$ in our notation. However, we do not find two point function

$$\langle \tau_3 \tau_{6g-3} \rangle_g = 0 \times \langle \tau_{6g-3} \rangle_g \quad (7.9)$$

which means a vanishing coefficient. Above equation plays a role of a dilaton equation, but the coefficient becomes zero.

The two point function of $p = \frac{1}{2}$ is obtained by the expansions of small s_1 and s_2 in a genus expansion. For this purpose, we write the integral representation of (5.5) as

$$u(s_1, s_2) = -\frac{4}{s_1^2 s_2^2} \oint \frac{dy_1 dy_2}{(2i\pi)^2} (y_1 + \frac{s_1^6}{y_1^6})(y_2 + \frac{s_2^6}{y_2^6}) e^{c \sum (y_i - \frac{s_i^6}{3y_i^3})} \times \frac{1}{(\frac{1}{s_1^2}(y_1^2 - \frac{s_1^6}{y_1^6}) - \frac{1}{s_2^2}(y_2^2 - \frac{s_2^6}{y_2^6}))^2 + 4(s_1 + s_2)^2} \quad (7.10)$$

If we expand in the inverse of y_1 , we obtain

$$\begin{aligned}
u(s_1, s_2) &= -4 \frac{s_1^2}{s_2^2} \oint \frac{dy_1 dy_2}{(2i\pi)^2} (y_1 + \frac{s_1^6}{y_1^3})(y_2 + \frac{s_2^6}{y_2^3}) e^{c(y_1 - \frac{s_1^6}{3y_1^3}) + c(y_2 - \frac{s_2^6}{3y_2^3})} \\
&\times \frac{1}{y_1^4} [1 + g + g^2 + g^3 + \dots] \tag{7.11}
\end{aligned}$$

where

$$\begin{aligned}
g &= 2 \left(\frac{s_1^6}{y_1^4} + \frac{s_1^2 y_2^2}{s_2^2 y_1^2} - \frac{s_1^2 s_2^4}{y_1^2 y_2^2} \right) - \left(\frac{s_1^6}{y_1^4} + \frac{s_1^2 y_2^2}{s_2^2 y_1^2} - \frac{s_1^2 s_2^4}{y_1^2 y_2^2} \right)^2 \\
&- \frac{4s_1^4 (s_1 + s_2)^2}{y_1^4} \tag{7.12}
\end{aligned}$$

By the contour integral at $y_1 = 0, y_2 = 0$, we have expanding the exponential term proportional to c ,

$$\begin{aligned}
u(s_1, s_2) &= -\left[\frac{1}{3} c^4 s_1^2 s_2^4 \right. \\
&+ c^8 \left(\frac{1}{360} s_1^8 s_2^4 + \frac{1}{135} s_1^7 s_2^5 + \frac{1}{180} s_1^6 s_2^6 + \frac{1}{216} s_1^4 s_2^8 + \frac{1}{1080} s_1^2 s_2^{10} \right) + O(c^{12}) \\
&\left. + [s_1 \leftrightarrow s_2] \right] \tag{7.13}
\end{aligned}$$

The expression should be symmetric by adding the terms of $s_1 \leftrightarrow s_2$. The terms of order $g = 2$ becomes by this symmetrization,

$$c^8 \frac{1}{3^3 \cdot 5} \left((s_1^8 s_2^4 + s_1^4 s_2^8 + s_1^7 s_2^5 + s_1^5 s_2^7) + \frac{3}{2} s_1^6 s_2^6 + \frac{1}{8} (s_1^2 s_2^{10} + s_1^{10} s_2^2) \right) \tag{7.14}$$

The first term shows the string equation; $s_1^2 s_2^4 \rightarrow s_2^3$. Note that there is no s^3 term in two point function, which is supposed to be present for the term for a dilaton equation. This is related to a fact that the power $p+1$ of Z^{p+1} becomes not integer in (6.20), and the argument of the proof of a dilaton equation does not hold. The term $s_1^7 s_2^5$ comes from the last term of g .

From (2.6) for $p = \frac{1}{2}$, we have $6g = n_1 + n_2$ for $s_1^{n_1} s_2^{n_2}$. The first term of (7.13) is genus one, and the term of c^8 is genus two, since $n_1 + n_2 = 12$. It is interesting to note that n_1 and n_2 are both even, or both odd, since $6g$ is an even number. The terms of $s_1^2 s_2^{10}$ is consistent with a string equation of (7.4).

It may be easy to obtain the higher order terms. The parameter c is $c = \frac{i}{4}$. Each term of two point function is order of $c^{4g} s_1^{n_1} s_2^{6g-n_1}$ for genus g , which agrees with the selection rule of (6.6). The scaling relation between c and s can be seen in (2.4). After the rescaling of $u \rightarrow \frac{s}{2} u$, the exponent becomes cs^{p+1} . This leads to the scaling $cs^{\frac{3}{2}}$, and $u(s_1, s_2) \sim \sum_{n=2}^{\infty} \sum_{g=1}^{\infty} c^{4g} s_1^n s_2^{6g-n}$. (7.13) is consistent with this behavior.

There may be another way for the evaluation of the contour integral. For genus one case, the last factor of (7.10) is approximated by

$$\frac{1}{\left(\frac{y_1^2}{s_1^2} - \frac{y_2^2}{s_2^2}\right)^2} = \frac{1}{\left(\frac{y_1}{s_1} - \frac{y_2}{s_2}\right)^2 \left(\frac{y_1}{s_1} + \frac{y_2}{s_2}\right)^2} \tag{7.15}$$

We take residues at $y_1 = \frac{s_1}{s_2} y_2$ and at $y_1 = -\frac{s_1}{s_2} y_2$. Expanding the exponent in order c^4 , we find the residue of $y_1 = \frac{s_1}{s_2} y_2$ and $y_2 = 0$ as

$$U(s_1, s_2) = \frac{c^4}{6s_2^3} (s_1^9 + 2s_1^8 s_2 + s_1^7 s_2^2 - s_1^3 s_2^6 - 2s_1^2 s_2^7 - s_1 s_2^8) \tag{7.16}$$

and from the pole at $y_1 = -\frac{s_1}{s_2}y_2$ and $y_2 = 0$ as

$$u(s_1, s_2) = \frac{c^4}{6s_2^3}(-s_1^9 + 2s_1^8s_2 - s_1^7s_2^2 + s_1^3s_2^6 - 2s_1^2s_2^7 + s_1s_2^8) \quad (7.17)$$

Adding these two contributions, by noting the cancellation of terms, we have

$$u(s_1, s_2) = \frac{2c^4s_1^8}{3s_2^2} - \frac{2}{3}c^4s_1^2s_2^4 \quad (7.18)$$

The term of $s_1^2s_2^4$ agrees with (7.13) in genus one.

One may evaluate the residue of (7.10) by taking 8 poles of y_1 . There are 8 poles, except a pole of numerator $y_1 = 0$,

$$y_1 = \pm \frac{is_1s_2^2}{2y_2} \pm \frac{s_1y_2}{2s_2} \pm \frac{s_1\sqrt{-s_2^6 \pm 4is_1s_2^2y_2^2 \pm 2is_2^3y_2^2 + y_2^4}}{2s_2y_2} \quad (7.19)$$

These residues for 8 poles are simply expressed as

$$\begin{aligned} & \oint \frac{dy_1}{2i\pi} \frac{(y_1 + \frac{s_1^6}{y_1})(y_2 + \frac{s_2^6}{y_2})}{(\frac{1}{s_1^2}(y_1^2 - \frac{s_1^6}{y_1}) - \frac{1}{s_2^2}(y_2^2 - \frac{s_2^6}{y_2}))^2 + 4(s_1 + s_2)^2} \\ &= \pm i \frac{s_1^2(s_2^6 + y_2^4)}{8(s_1 + s_2)y_2^3} \end{aligned} \quad (7.20)$$

Adding the contribution of 8 poles, there is a cancellation. Thus the contribution comes only from the pole of $y_1 = y_2 = 0$.

• **three-point function for $p = \frac{1}{2}$**

We have three point function of $p = \frac{1}{2}$ by putting $p = \frac{1}{2}$ in (5.23),

$$\begin{aligned} u(s_1, s_2, s_3) &= -8 \frac{s_1^2}{s_3^2} \oint \prod_{i=1}^3 \frac{dy_i}{2i\pi} \prod_{i=1}^3 (y_i + \frac{s_i^6}{y_i}) e^{c \sum (3y_i - \frac{s_i^6}{y_i})} \\ &\times \left(\frac{1}{y_1^4 y_2^2} \frac{1}{1 - \frac{s_1^6}{y_1^4} - \frac{s_1^2}{s_2^2 y_1^2} (y_2^2 - \frac{s_2^6}{y_2}) - \frac{2is_1^2}{y_1^2} (s_1 + s_2)} \right) \\ &\times \frac{1}{1 - \frac{s_2^6}{y_2^4} - \frac{s_2^2}{s_3^2 y_2^2} (y_3^2 - \frac{s_3^6}{y_3}) - \frac{2is_2^2}{y_2^2} (s_2 + s_3)} \\ &\times \frac{1}{1 - \frac{s_3^6}{y_3^4} - \frac{s_3^2}{s_1^2 y_3^2} (y_1^2 - \frac{s_1^6}{y_1}) + \frac{2is_3^2}{y_3^2} (s_1 + s_3)} \end{aligned} \quad (7.21)$$

where $c = -\frac{1}{6}i^{1/2}$. Expanding the denominators in the power series of s_i , we obtain by the evaluation of residues,

$$u(s_1, s_2, s_3) = -8 \frac{s_1^2}{s_3^2} \left(\frac{27}{4} c^4 s_3^6 + i \frac{81}{4} c^6 (s_1^2 s_2 s_3^6 + 3s_2^3 s_3^6 + 3s_2^2 s_3^7) + O(c^8) \right) \quad (7.22)$$

These terms are order of $c^{4g+2}s^{6g+3}$ according to the scaling relation between c and s as $cs^{3/2}$. After the contour integration of y_1 , the remaining term is odd

for the exchange of $(s_2, y_2) \leftrightarrow (s_3, y_3)$. For this order of c^4 , the contour integral of y_1 is factorized as

$$\begin{aligned}
U(s_1, s_2, s_3) &= -8s_1^2 \oint \frac{dy_1 y_1}{2i\pi y_1^4} e^{c(3y_1 - \frac{s_1^6}{y_1^4})} \oint \frac{dy_2 dy_3}{(2i\pi)^2} (y_2 + \frac{s_2^6}{y_2^3}) (y_3 + \frac{s_3^6}{y_3^3}) \\
&\times \frac{e^{c(3y_2 - \frac{s_2^6}{y_2^3}) + c(3y_3 - \frac{s_3^6}{y_3^3})}}{s_3^2 y_2^2 - \frac{s_2^6 s_3^2}{y_2^2} - s_2^2 y_3^2 + \frac{s_2^2 s_3^6}{y_3^2} - 2is_2^2 s_3^2 (s_2 + s_3)} \quad (7.23)
\end{aligned}$$

The denominator becomes $(s_3^2 y_2^2 - s_2^2 y_3^2)^{-1}$ in this order, and it is odd for the exchange $(s_2, y_2) \leftrightarrow (s_3, y_3)$. Therefore the first term of order c^4 is vanishing by adding the symmetric terms of $s_2 \leftrightarrow s_3$. Absence of order c^4 term agrees with (6.6), which implies the order of $c^{4g+2} s^{6g+3}$ for a genus g term. The term of order c^6 in (7.22) contains $i = (-1)^{1/2}$, which comes from the term of $i(s_i + s_j)$ in the denominator of (7.21). Note that $c = -\frac{1}{6}i^{3/2}$, ic^6 is real number. The term of order c^6 is also vanishing by the followin reason. We have expanded the large $y_1 > y_2 > y_3$. We have to consider also the case of $y_3 > y_2$. For this case, we exchange $y_2 \leftrightarrow y_3$ as well as $s_2 \leftrightarrow s_3$ with the complex conjugate of the term $-2i(s_2 + s_3)$, and this leads to the opposite sign of the term of c^6 with the exchange $s_2 \leftrightarrow s_3$. Thus after adding this symmetric term, we obtain the cancellation of (7.22) and vanishing result of c^6 term. The fact that there is no three point function suggests the punctures are pairwise, and the vanishing three point function gives the simple structure of the spin $p = \frac{1}{2}$ case, which may be related to the simplification for Faber conjecture [54] as discussed in [9, 55].

- **four-point function for $p = \frac{1}{2}$**

The selection rule for four point function is $6g+6 = \sum_{i=1}^4 n_i$. The connected four point function $u(s_1, s_2, s_3, s_4)$ is expressed as in [9] with the same change of variables $u_i \rightarrow \frac{i}{2}(y_i^2 - \frac{1}{y_i^2})$, by assuming order of smallness $s_1 < s_2 < s_3 < s_4$,

$$\begin{aligned}
u(s_1, s_2, s_3, s_4) &= -16 \frac{s_2^2}{s_4^2} \oint \prod_{i=1}^4 (y_i + \frac{s_i^6}{y_i^3}) e^{c \sum_{i=1}^4 (3y_i - \frac{s_i^6}{y_i^3})} \\
&\times \frac{1}{y_1^2 - \frac{s_1^6}{y_1^2} - \frac{s_2^2}{s_2^2} (y_2^2 - \frac{s_2^6}{y_2^2}) - 2is_1^2 (s_1 + s_2)} \frac{1}{y_2^2 - \frac{s_2^6}{y_2^2} - \frac{s_2^2}{s_3^2} (y_3^2 - \frac{s_3^6}{y_3^2}) - 2is_2^2 (s_2 + s_3)} \\
&\times \frac{1}{y_3^2 - \frac{s_3^6}{y_3^2} - \frac{s_2^2}{s_4^2} (y_4^2 - \frac{s_4^6}{y_4^2}) - 2is_3^2 (s_3 + s_4)} \frac{1}{y_1^2 - \frac{s_1^6}{y_1^2} - \frac{s_1^2}{s_4^2} (y_4^2 - \frac{s_4^6}{y_4^2}) + 2is_1^2 (s_1 + s_4)} \quad (7.24)
\end{aligned}$$

Since the four point function is cyclic about y_i ($i = 1, \dots, 4$), we take y_1 and y_3 are large, and expansion of $1/y_1$ and $1/y_3$ in (7.24). Similar to the three point function, we find that the contour integrals of (7.24) are vanishing for the terms of order c^i ($i = 1, \dots, 7$). The non-vanishing term appears at the order c^8 with a result of

$$u(s_1, s_2, s_3, s_4) = -729c^8 s_1^2 s_3^2 s_2^4 s_4^4 + O(c^{12} s^{18}) \quad (7.25)$$

where $c = -\frac{1}{6}i^{1/2}$. This is consistent with the selection rule of $6g+6 = \sum_{i=1}^4 n_i$, where n_i is a power of s_i . In this order, the genus g is one, and it is consistent with the scaling of $c^{4g+4} s^{6g+6} = c^8 s^{12}$. The result of (7.25) is obtained for $y_1 >$

$\{y_2, y_4\}$ and $y_3 > \{y_2, y_4\}$. The total expression is obtained by the permutation of the variables $\{s_1, \dots, s_4\}$.

In this four point function, the numbers of punctures are even numbers (i.e. it is four). The punctures have the spin indices $j = -1$ as a Ramond puncture, and according to this spin indices, the selection rule $6g + 6 = \sum_{i=1}^4 n_i$ is obtained. Thus our explicit result of the four point function is consistent with the conjecture that Ramond punctures are pairwise.

The term of (7.25) is consistent with the successive string equations, which give the reduction of $s_1^2 s_3^2 s_2^4 s_4^4 \rightarrow s_1^2 s_2^3 s_4^4 \rightarrow s_3^2 s_4^4 \rightarrow s_4^3$. Note that $\langle \tau_3 \rangle$ is non-vanishing one point intersection number.

For n -point function ($n > 1$), this Ramond puncture pairing conjecture states that if n is odd, the n -point function is vanishing, and if n is even, it is non-vanishing. For n -point function ($n \geq 6$), the contour integral becomes similar to $n = 4$. Assuming y_j (j is odd) is large, and expand in the inverse power of y_j , we obtain the n -point function $u(s_1, \dots, s_n)$ from the evaluation of the residues of y_i at $y_i = 0$. Through these residual calculations, we find the selection rule of (2.6) for $p = \frac{1}{2}$ is valid with the Ramond values of spin component $j = -1$, and Ramond punctures ($j = -1$) appear as pair-wise form.

• **continuation from integer p to $p = \frac{1}{2}$**

For the continuation to $p = \frac{1}{2}$, the spin component j is assumed to be (-1) [8, 9]. From the selection rule of (2.6), we have for $p = \frac{1}{2}$

$$6g - 6 + 3 = \sum_{i=1}^3 n_i + 2 \sum_{i=1}^3 j_i \quad (7.26)$$

For general p , the three point function $\langle \tau_{0,0} \tau_{0,0} \tau_{0,p-2} \rangle > t_{0,0} t_{0,0} t_{0,p-2}$ is genus zero term, which corresponds to $s_1^{\frac{1}{p}} s_2^{\frac{1}{p}} s_3^{\frac{p-1}{p}}$ term. Putting $p = \frac{1}{2}$, this term becomes $s_1^2 s_2^2 s_3^{-1}$. This term is obtained from the contour integral in (7.21) by expanding the second denominator about $2is_2^2 s_3 / y_2^2$ as

$$u(s_1, s_2, s_3)_{g=0} = -8s_1^2 s_2^2 s_3^{-1} \oint \prod_{i=1}^3 \frac{dy_i}{2i\pi} \frac{y_3^3}{y_1^3 y_2^2} e^{c \sum y_i} \quad (7.27)$$

However the contour integral of y_3 gives zero around $y_3 = 0$. Thus we have no 3-point function of $p = \frac{1}{2}$ at genus zero. This result is consistent with the conjecture of the punctures of Ramond type, which should be paired.

The string equation of $p = \frac{1}{2}$ has been found in [9]. It is transform from $t_{0,0} \rightarrow t_2$, ($t_2 = s^2$).

For $u(s)$ is given by (4.28).

$$u(s) = \frac{c^2}{48} s^3 - \frac{1}{5!} \left(\frac{1}{48}\right)^2 c^6 s^9 + \dots \quad (7.28)$$

This expression is continuation of $u(s)$ for the integer p , with $p = \frac{1}{2}$, since for general p , it is given by [6] as

$$u(s) = \frac{1}{s^{1+\frac{1}{p}}} \left[\Gamma\left(1 + \frac{1}{p}\right) - \frac{p-1}{24} y \Gamma\left(1 - \frac{1}{p}\right) + \frac{(p-1)(p-3)(1+2p)}{5!4^2 3} y^2 \Gamma\left(1 - \frac{3}{p}\right) + \dots \right] \quad (7.29)$$

with $y = s^{2+\frac{2}{p}}$. There appears a divergence in the limit of $p = \frac{1}{2}$ for Γ -function, and with the normalization of this factor, (7.29) becomes equivalent to (7.28) as noted in Eq.(18) of [9].

From the results of (7.13), $u(s_1, s_2)$ is expressed as

$$\begin{aligned} u(s_1, s_2) = & -\frac{1}{3}c^4 s_1^2 s_2^4 + c^8 \left(\frac{1}{1080} s_1^2 s_2^{10} + \frac{1}{216} s_1^4 s_2^8 + \frac{1}{180} s_1^6 s_2^6 + \frac{1}{135} s_1^7 s_2^5 \right. \\ & + \frac{1}{360} s_1^8 s_2^4 \left. \right) \\ & - c^{12} \left(\frac{s_1^2 s_2^{16}}{3265920} + \frac{s_1^4 s_2^{14}}{204120} + \frac{7s_1^6 s_2^{12}}{291600} + \frac{s_1^7 s_2^{11}}{48600} + \frac{313s_1^8 s_2^{10}}{8164800} + \frac{s_1^9 s_2^9}{22680} \right. \\ & + \frac{443s_1^{10} s_2^8}{16329600} + \frac{11s_1^{11} s_2^7}{510300} + \frac{169s_1^{12} s_2^6}{8164800} + \frac{19s_1^{13} s_2^5}{2041200} + \left. \frac{s_1^{14} s_2^4}{583200} \right) + O(c^{16}) \end{aligned} \quad (7.30)$$

This expression is interpreted as a continuation of $p \rightarrow \frac{1}{2}$. s_2^4 corresponds to $s_2^{\frac{2}{p}}$ with $p = \frac{1}{2}$. Since $s_2^{\frac{2}{p}}$ is $t_{0,1}$, s^4 corresponds to $t_{0,1}$. The term s^5 corresponds to $t_{1,1}$.

The intersection number for $s_1^{\frac{2}{p}} s_2^{4+\frac{2}{p}}$ is given

$$\langle \tau_{0,1} \tau_{4,1} \rangle_{g=2} = \frac{(p-1)(p+2)(p-2)}{2880p} \rightarrow \frac{1}{768} \quad (7.31)$$

This expression agrees for the value of [10, 11], $\frac{1}{320}$ and $\frac{7}{1200}$ for $p = 4$ and $p = 5$. (7.31) corresponds to $s_1^{\frac{2}{p}} s_2^{4+\frac{2}{p}} \rightarrow s_1^4 s_2^8$ for $p = \frac{1}{2}$. The term, which gives $s_1^4 s_2^8$ in the limit $p \rightarrow \frac{1}{2}$, may exist also from another term $\langle \tau_{2,0} \tau_{2,2} \rangle$ with $s_1^{2+\frac{1}{p}} s_2^{2+\frac{3}{p}}$. Therefore, $s_1^4 s_2^8$ is obtained from the limit of different terms of general p . The term $s_1^{2+\frac{1}{p}} s_2^{2+\frac{3}{p}}$ is as

$$u(s_1, s_2) = -\frac{1}{576}(p+1)(p-1)\Gamma\left(1-\frac{1}{p}\right)\Gamma\left(1-\frac{3}{p}\right)s_1^{2+\frac{2}{p}}s_2^{2+\frac{1}{p}} \quad (7.32)$$

which by the change $s_1 \leftrightarrow s_2$, it becomes $s_1^4 s_2^8$ for $p \rightarrow \frac{1}{2}$.

For $s_1^6 s_2^6$ term in (7.30), we have

$$\langle \tau_{0,2} \tau_{4,0} \rangle_{g=2} = \frac{(p-1)(p-3)(2p+11)}{5760p} \rightarrow \frac{1}{192} \quad (7.33)$$

which corresponds to $s_1^{\frac{3}{p}} s_2^{4+\frac{1}{p}} \rightarrow s_1^6 s_2^6$ for $p = \frac{1}{2}$. The values of these intersection numbers agree with the intersection numbers from (7.30).

For $\langle \tau_{2,1} \tau_{2,1} \rangle_{g=2}$ corresponds to $s_1^{2+\frac{2}{p}} s_2^{2+\frac{2}{p}} \rightarrow s_1^6 s_2^6$, which degenerates to (7.33). Thus the continuation to $p = \frac{1}{2}$ is not unique.

The term $s^{\frac{2}{p}}$ and $s^{2+\frac{1}{p}}$ become same s^4 in the limit $p \rightarrow \frac{1}{2}$. The terms $s^{\frac{2}{p}}$, $s^{2+\frac{2}{p}}$ and $s^{4+\frac{1}{p}}$ become s^6 in the limit of $p \rightarrow \frac{1}{2}$.

7-2: n-point function of $p = -\frac{1}{2}$

For $p = -\frac{1}{2}$, the selection rule becomes for k -marked points

$$2g - 2 + k = -\sum_{i=1}^k n_i \quad (7.34)$$

where we take $j_i = -1$. This requires the negative power n_i of s for higher genus g , which leads to the inverse power of s_i in m -point function $u(s_1, \dots, s_m)$. As discussed in [8, 9], one point function $u(s)$ is a contour integral,

$$\begin{aligned} u(s) &= s \oint \frac{dy}{2i\pi} (y + \frac{1}{s^2 y^3}) e^{-\frac{c}{y}} \\ &= \frac{1}{2} c^2 s \end{aligned} \quad (7.35)$$

This term belongs to genus $g = 0$ order, which leads to $n_1 = 1$ from (7.34) as s^{n_1} . The coefficient c^2 is given by the scaling $(cs^{p+1})^2 = c^2 s$. From the expansion of the intersection numbers in (2.8), the higher order terms are vanishing for $p = -\frac{1}{2}$ by the existent factor $(2p+1)$ for $g > 1$. In the genus $g = 1$ order, the intersection number $\langle \tau_{1,0} \rangle = \frac{p-1}{24}$ in (2.3) for the integer p suggests the nonvanishing value for $p = -\frac{1}{2}$. This order is given by the second term of the integrand. By the change of $y \rightarrow \frac{1}{t}$, (7.35) is written as

$$u(s) = s \int \frac{dt}{2i\pi} (-\frac{1}{t^2}) (\frac{1}{t} + \frac{t^3}{s^2}) e^{-ct} \quad (7.36)$$

The second term of the integrand becomes by the integration as $-\frac{1}{c^2 s}$ ($c = (\frac{1}{4})^p = 2$), which is a continuation of $\frac{p-1}{24} = -\frac{1}{16}$. There is a factor $\Gamma(3) = 2$ in (2.3). Thus genus $g = 1$ term is consistent with (2.3).

Two point function of (5.5) is written for $p = -\frac{1}{2}$ as

$$\begin{aligned} u(s_1, s_2) &= -4 (\frac{s_2}{s_1})^2 \oint \frac{dy_1 dy_2}{(2i\pi)^2} \prod_{i=1}^2 (y_i + \frac{1}{s_i^2 y_i^3}) e^{-c \sum \frac{1}{y_i}} \\ &= \frac{1}{y_1^4 (1 - \frac{1}{s_1^2 y_1^4} - (\frac{s_2}{s_1})^2 \frac{1}{y_1^2} (y_2^2 - \frac{1}{s_2^2 y_2^2}))^2 + \frac{4}{s_1^4 y_1^4} (s_1 + s_2)^2} \end{aligned} \quad (7.37)$$

The contour integral is evaluated for the poles of $y_1 = y_2 = 0$. By the expansion of the exponential term as $\sum (-c(\frac{1}{y_1} + \frac{1}{y_2}))^k / k!$, we obtain

$$\begin{aligned} u(s_1, s_2) &= -4s_2^2 (\frac{1}{4}c^4 - \frac{c^8}{2 \cdot 6!} (s_1^2 + 8s_1s_2 + 12s_2^2) \\ &+ \frac{c^{12}}{2 \cdot 10!} (s_1^4 + 24s_1^3s_2 + 161s_1^2s_2^2 + 384s_1s_2^3 + 288s_2^4) \\ &- \frac{c^{16}}{2 \cdot 14!} (s_1^6 + 48s_1^5s_2 + 715s_1^4s_2^2 + 4528s_1^3s_2^3 + 13441s_1^2s_2^4 + 18304s_1s_2^5 \\ &+ 9152s_2^6) + O(c^{20})) + (s_1 \leftrightarrow s_2) \end{aligned} \quad (7.38)$$

Each term is written as $c^{4g} s_1^{|n_1|} s_2^{|n_2|}$, $n_1 + n_2 = 2g$. For $p = -\frac{1}{2}$, the selection rule becomes for two point function as ($m_i = -1$)

$$-2g = n_1 + n_2 \quad (7.39)$$

Thus n_1, n_2 are negative integers for $s_1^{-n_1} s_2^{-n_2}$. The expansion of (7.38) is similar to the weak coupling expansion of $p = -2$ case.

By the change of variable $y_i = \frac{1}{t_i}$, ($i = 1, 2$), it becomes by $t_i \rightarrow t_i/c$,

$$\begin{aligned} u(s_1, s_2) &= -4 (\frac{s_2}{s_1})^2 \oint \frac{dt_1 dt_2}{(2i\pi)^2} (t_1 + \frac{t_1^5}{c^4 s_1^2}) e^{-t_1} (\frac{1}{t_2^3} + \frac{t_2}{c^4 s_2^2}) e^{-t_2} \\ &= \frac{1}{(1 - \frac{t_1^4}{c^4 s_1^2} - (\frac{s_2}{s_1})^2 t_1^2 (\frac{1}{t_2^2} - \frac{t_2^2}{c^4 s_2^2}))^2 + \frac{4t_1^4}{c^4 s_1^4} (s_1 + s_2)^2} \end{aligned} \quad (7.40)$$

From the selection rule of (7.34), in the case two point function $n = 2$, we have

$$2g = -(n_1 + n_2) \quad (7.41)$$

for $u(s_1, s_2) = \sum c_{n_1, n_2} s_1^{n_1} s_2^{n_2}$. The values of n_1 and n_2 can be both positive and negative integers. The expansion of (7.40) is

$$\begin{aligned} u(s_1, s_2) = & -4 \left(\frac{s_2}{s_1} \right)^2 \oint \frac{dt_1 dt_2}{(2i\pi)^2} e^{-t_1 - t_2} \frac{t_1 t_2}{(t_2^2 - \frac{s_2^2}{s_1^2} t_1^2)^2} \left(1 + \frac{t_1^4}{c^4 s_1^2} + \frac{t_2^4}{c^4 s_2^2} \right. \\ & \left. + 2 \frac{t_1^2}{c^4 s_1^2} \frac{t_1^2 - t_2^2}{1 - \frac{s_2^2}{s_1^2} \frac{t_1^2}{t_2^2}} - 4 \frac{(s_1 + s_2)^2}{c^4 s_1^4} \frac{t_1^4}{(1 - \frac{s_2^2}{s_1^2} \frac{t_1^2}{t_2^2})^2} + O\left(\frac{1}{c^8}\right) \right) \end{aligned} \quad (7.42)$$

where the dependence of c is obtained from the scaling to s as a combination of $c^2 s$.

7-3: Strong coupling of $p = -2$

• $p = -2$ and unitary matrix model

It was pointed out that for $p = -2$ corresponds to unitary matrix model [39, 31]. The unitary matrix model has two phases of the weak coupling and string coupling phases [37, 38].

For one point function, we have for large s ,

$$\begin{aligned} u(s) &= \frac{i}{2} \oint \frac{dy}{2i\pi} \left(y + \frac{1}{y^3} \right) e^{\frac{c}{s} \frac{y^4}{(y^4+1)^2}} \\ &= \frac{2}{s} + \frac{2}{s^2} + \frac{2}{s^3} + \frac{5}{3s^4} + \frac{7}{6s^5} + \frac{7}{10s^6} + \frac{11}{30s^7} + \frac{11 \cdot 13}{840s^8} + \frac{11 \cdot 13}{2016s^9} \\ &+ \frac{11 \cdot 13 \cdot 17}{2^5 3^4 5 \cdot 7 s^{10}} + \frac{13 \cdot 17 \cdot 19}{2^5 3^4 5^2 \cdot 7 s^{11}} + \frac{13 \cdot 17 \cdot 19}{2^6 3^4 5^2 \cdot 11 s^{12}} + \frac{17 \cdot 19 \cdot 23}{2^7 3^5 5^2 \cdot 11 s^{13}} + \dots \end{aligned} \quad (7.43)$$

where $c = (\frac{1}{4})^p = 2^4$. The contour integral reduces to the calculation of the residue at $y = (-1)^{1/4}$. This expansion is written by the modified Bessel function as

$$\begin{aligned} u(s) &= \frac{1}{s} \sum_{k=0}^{\infty} \frac{2^{2k+1} \Gamma(k + \frac{1}{2})}{\Gamma(\frac{1}{2}) \Gamma(k+2) k!} \frac{1}{s^k} \\ &= \frac{2}{s} F\left(\frac{1}{2}, 2; \frac{4}{s}\right) = \frac{2}{s} e^{\frac{2}{s}} [I_0\left(\frac{2}{s}\right) - I_1\left(\frac{2}{s}\right)] \end{aligned} \quad (7.44)$$

where $F(\alpha, \gamma; z)$ is a confluent hypergeometric function, and $I_0(z)$ is a modified Bessel function. The crossover to the weak coupling of small s is at $s = 2$, which corresponds to $s = \frac{1}{\Lambda} = 2$. The external source is Λ . The small s expansion is given by (2.8) through the intersection numbers of $p = -2$ [31]. We have

$$\begin{aligned} u(s) &= \frac{1}{s} \oint \frac{du}{2i\pi} e^{-\frac{1}{u+s} + \frac{1}{u}} \\ &= -\frac{1}{4\pi\sqrt{s}} \int \frac{dx}{x^2} e^{-\frac{4x^2}{1+x^2s}} \\ &= -\frac{1}{4\pi\sqrt{s}} \sum_{n=0}^{\infty} \frac{(2n-1)!!}{2^n n!} \Gamma\left(n - \frac{1}{2}\right) \left(\frac{s}{4}\right)^n \end{aligned} \quad (7.45)$$

By noting that $(2n-1)!! = 2^n \Gamma(n + \frac{1}{2}) / \sqrt{\pi}$, the last sum is written as

$$u(s) = -\frac{1}{4\pi^{\frac{3}{2}}\sqrt{s}} \sum_{n=0}^{\infty} \frac{\Gamma(n + \frac{1}{2})\Gamma(n - \frac{1}{2})}{n!} \left(\frac{s}{4}\right)^n \quad (7.46)$$

This weak coupling expansion of one point function has been derived and the comparison with the unitary matrix model has been discussed in [31].

The model $p = -2$ has naturally a logarithmic potential, which becomes equivalent to unitary $U(N)$ matrix model. The strong coupling expansion in (7.44) is evaluated by the character expansion of $U(N)$ group, which character χ satisfies

$$\int dU \chi_r(AU) \chi_{r'}(U^\dagger A^\dagger) = \delta_{rr'} \frac{\chi_r(AA^\dagger)}{d_r} \quad (7.47)$$

The character of $U(N)$ is expressed by

$$\chi_{n_1, n_2, \dots} = \frac{\det[t_i^{n_j + N - j}]}{\det[t_i^{N - j}]} \quad (7.48)$$

where t_i is eigenvalue of U . The dimension d_r is equal to $d_r = \chi_r(1)$. For the eigenvalue 1, ($t_i = 1$),

$$d_{[2,0]} = \chi_{[2,0]}(1) = \frac{1}{2}(N^2 + N), \quad d_{[1,1]} = \chi_{[1,1]}(1) = \frac{1}{2}(N^2 - N) \quad (7.49)$$

The integration of unitary matrix U is written as

$$\int dU \operatorname{tr}(AU) \operatorname{tr}(BU^\dagger) = C_1 \operatorname{tr}(AB) \quad (7.50)$$

where A and B are arbitrary matrices,

$$\begin{aligned} \int DU \operatorname{tr}(A_1 U) \operatorname{tr}(A_2 U) \operatorname{tr}(B_1 U^\dagger) \operatorname{tr}(B_2 U^\dagger) &= C_{12} [\operatorname{tr}(A_1 B_1) \operatorname{tr}(A_2 B_2) \\ &+ \operatorname{tr}(A_1 B_2) \operatorname{tr}(A_2 B_1)] + C_2 [\operatorname{tr}(A_1 B_1 A_2 B_2) + \operatorname{tr}(A_1 B_2 A_2 B_1)] \end{aligned} \quad (7.51)$$

Generalization of this equation for n -times trace is described by C_{l_1, \dots, l_n} .

The coefficient of the single trace is $C_n = C_{n, 0, \dots, 0}$, which are

$$\begin{aligned} C_1 &= 1, \quad C_2 = -\frac{1}{N^2 - 1}, \quad C_3 = \frac{4}{(N^2 - 1)(N^2 - 4)}, \\ C_4 &= -\frac{30}{(N^2 - 1)(N^2 - 4)(N^2 - 9)}, \\ C_5 &= \frac{336}{(N^2 - 1)(N^2 - 4)(N^2 - 9)(N^2 - 16)} \end{aligned} \quad (7.52)$$

The general formula of C_n is [49]

$$\begin{aligned} C_n &= (-1)^n \frac{[(n-1)!]^3}{n} \frac{1}{(N^2 - 1)(N^2 - 4)(N^2 - 9) \dots (N^2 - (n-1)^2)} \\ &\times \sum_{q=0}^{n-1} \frac{1}{q! q! (n-q-1)! (n-q-1)!} \end{aligned} \quad (7.53)$$

The coefficient of $1/s^n$ in (7.43) is $2C_n$, and agrees with $N = 0$ [31, 49].

It is interesting to note that when we change the measure $(y + \frac{1}{y^3})$ to $(y - \frac{1}{y^3})$, i.e. D-type, one point function is vanishing i.e. $u(s) = 0$. The replacement of this measure $y + 1/y^3$ by y , the values of $u(s)$ becomes half of (7.43).

For the derivation of the N dependence, we make a change of variable u to z as

$$\frac{u-1}{u+1} = e^{-z}, \quad u = \frac{1+e^{-z}}{1-e^{-z}} \quad (7.54)$$

and

$$du = -2 \frac{e^{-z}}{(1-e^{-z})^2} \quad (7.55)$$

Thus we obtain

$$\begin{aligned} U(s) &= \frac{N}{2} \int du e^{\frac{4c}{s(u^2-1)}} \left(\frac{u-1}{u+1}\right)^N \\ &= -N \int_0^\infty dz \sum_{n=0}^\infty \frac{1}{n!} e^{-Nz} \left[\frac{c}{s} e^z (1-e^{-z})^2\right]^n \frac{e^{-z}}{(1-e^{-z})^2} \end{aligned} \quad (7.56)$$

The leading term ($c = 0$) gives the Euler characteristics $\zeta(1-2g)$ [6], the next order is $\frac{c}{s}$, and the second order becomes

$$N \frac{c^2}{2s^2} \int_0^\infty dz e^{2z} (1-e^{-z})^2 e^{-Nz} = \frac{c^2}{s^2} \frac{1}{N^2-1} \quad (7.57)$$

The next order is

$$\begin{aligned} & -\frac{N}{6} \left(\frac{c}{s}\right)^3 \int dz e^{2z} (1-e^{-z})^4 e^{-Nz} \\ &= -\frac{N}{6} \left(\frac{c}{s}\right)^3 \left[\frac{1}{N-2} - \frac{4}{N-1} + \frac{6}{N} - \frac{4}{N+1} + \frac{1}{N+2} \right] \\ &= -4N \left(\frac{c}{s}\right)^3 \frac{1}{N(N^2-1)(N^2-4)} \end{aligned} \quad (7.58)$$

Extending these evaluations, we find from (7.56)

$$U(s) = \sum_{n=1}^\infty \frac{(2n-2)!}{n! s^n} \frac{1}{\prod_{l=1}^{n-1} (N^2 - l^2)} \quad (7.59)$$

where $c = 1$. The number $(2n-2)!/n!$ coincides with the numerator of C_n in (7.52). Thus we find that one point function $U(s)$, which is a single trace operator, is a generating function of the coefficient C_n in (7.53).

For two point function $U(s_1, s_2)$ with a logarithmic term is written as

$$\begin{aligned} U(s_1, s_2) &= \int du_1 du_2 e^{\frac{4c}{s_1(u_1^2-1)} + \frac{4c}{s_2(u_2^2-1)}} \left(\frac{u_1+1}{u_1-1}\right)^N \left(\frac{u_2+1}{u_2-1}\right)^N \\ &\times \frac{1}{(s_2 u_2 - s_1 u_1 + (s_1 + s_2))(s_2 u_2 - s_1 u_1 - (s_1 + s_2))} \end{aligned} \quad (7.60)$$

Using the change of variables in (7.54), one obtain

$$\begin{aligned}
U(s_1, s_2) &= s_1 s_2 \int dz_1 dz_2 e^{-Nz_1 - Nz_2} \frac{e^{-z_1}}{(1 - e^{-z_1})^2} \frac{e^{-z_2}}{(1 - e^{-z_2})^2} \\
&\times \frac{1}{s_2(1 + \frac{2e^{-z_2}}{1 - e^{-z_2}}) - s_1(1 + \frac{2e^{-z_1}}{1 - e^{-z_1}}) + (s_1 + s_2)} \\
&\times \frac{1}{s_2(1 + \frac{2e^{-z_2}}{1 - e^{-z_2}}) - s_1(1 + \frac{2e^{-z_1}}{1 - e^{-z_1}}) - (s_1 + s_2)} \\
&\times e^{\frac{c}{s_1} e^{z_1} (1 - e^{-z_1})^2 + \frac{c}{s_2} e^{z_2} (1 - e^{-z_2})^2}
\end{aligned} \tag{7.61}$$

Expanding the two denominators for $s_2 > s_1$,

$$\frac{1}{4s_2^2} \sum_{m=0}^{\infty} \sum_{l=0}^{\infty} \left(\frac{1 - e^{-z_2}}{1 - e^{-z_1}} e^{-z_1} \right)^m \left(\frac{1 - e^{-z_2}}{1 - e^{-z_1}} e^{z_2} \right)^l \left(\frac{s_1}{s_2} \right)^{m+l} e^{z_2} (1 - e^{-z_2})^2 \tag{7.62}$$

with a measure of $du_i = e^{-z_i} / (1 - e^{-z_i})^2 dz_i$, we obtain the strong coupling expansion of $1/s_1^{n_1} s_2^{n_2}$,

$$\begin{aligned}
U(s_1, s_2) &= (s_1 s_2) \int dz_1 dz_2 e^{-Nz_1 - Nz_2} \frac{e^{-z_1}}{(1 - e^{-z_1})^2} \frac{e^{-z_2}}{(1 - e^{-z_2})^2} \\
&\times \frac{1}{4s_2^2} \sum_{m=0}^{\infty} \sum_{l=0}^{\infty} \left(\frac{1 - e^{-z_2}}{1 - e^{-z_1}} e^{-z_1} \right)^m \left(\frac{1 - e^{-z_2}}{1 - e^{-z_1}} e^{z_2} \right)^l \left(\frac{s_1}{s_2} \right)^{m+l} e^{z_2} (1 - e^{-z_2})^2 \\
&\times e^{\frac{c}{s_1} e^{z_1} (1 - e^{-z_1})^2 + \frac{c}{s_2} e^{z_2} (1 - e^{-z_2})^2}
\end{aligned} \tag{7.63}$$

The leading term is given by Euler characteristics $\zeta(1 - 2g)$,

$$\begin{aligned}
U(s_1, s_2) &= \frac{1}{4s_2^2} \int dz_1 dz_2 e^{-Nz_1 - Nz_2} \frac{e^{-z_1}}{(1 - e^{-z_1})^2} \\
&= \frac{1}{4s_2^2 N^2} \sum_{n=0}^{\infty} \frac{1}{N^{2n}} (-1)^{n-1} \frac{B_n}{2n} \\
&= \frac{1}{4s_2^2 N^2} \sum_n \frac{1}{N^{2n}} \zeta(1 - 2n)
\end{aligned} \tag{7.64}$$

where we have to count $s_1 \leftrightarrow s_2$, and B_n is Bernoulli number $B_1 = \frac{1}{6}, B_2 = \frac{1}{30}, B_3 = \frac{1}{42}, \dots$. Note this term has no s_1 dependence.

The next term in the strong coupling is the term of order $1/s_1 s_2$ as we will see later for $N = 0$ in (7.68). In the strong coupling region, $U(s_1, s_2)$ is expanded in the inverse power of s_1 and s_2 . The pole of $(1 - e^{-z_1})$ in (7.63) should be cancelled with the expansion of the exponent in the last term. Then, similar to one point function $U(s)$, two point function is evaluated in the form of characters of $U(N)$ as shown in [31], which used a different method as shown here.

We have from (7.63) two point function for $p = -2$,

$$\begin{aligned}
U(s_1, s_2) &= \frac{c}{4s_2 N^2} + \frac{c^2}{4s_1 s_2} \frac{1}{N^2(N^2 - 1)} + \frac{c^3}{s_1^2 s_2} \frac{1}{N^2(N^2 - 1)(N^2 - 4)} \\
&+ \frac{30c^4}{4s_1^3 s_2} \frac{1}{N^2(N^2 - 1)(N^2 - 4)(N^2 - 9)} + O\left(\frac{1}{s_1^n s_2^n}\right)
\end{aligned} \tag{7.65}$$

The coefficient $C_{1,n}$ is [31]

$$\begin{aligned} C_{1,1} &= \frac{1}{N^2 - 1}, & C_{1,2} &= -\frac{12}{(N^2 - 1)(N^2 - 4)} \\ C_{1,3} &= \frac{120}{(N^2 - 1)(N^2 - 4)(N^2 - 9)} \\ C_{1,4} &= -\frac{1680}{(N^2 - 1)(N^2 - 4)(N^2 - 9)(N^2 - 16)} \end{aligned} \quad (7.66)$$

The expression of $U(s_1, s_2)$ of (7.63) in a strong coupling agrees with the coefficients of C_{n_1, n_2} .

This leads to the following conjecture such that the n point function $U(s_1, \dots, s_n)$ of $p = -2$ case in the inverse s expansion is a generating function of the coefficient C_{l_1, \dots, l_n} of unitary integral of the multi-trace products.

We now turn to the change of variable y . We here use the expression of (5.5) for $p = -2$. It becomes for two point function,

$$\begin{aligned} u(s_1, s_2) &= -4s_1s_2 \oint \frac{dy_1 dy_2}{(2i\pi)^2} \left(\frac{1+y_1^4}{y_1^3} \right) \left(\frac{1+y_2^4}{y_2^3} \right) e^{-\frac{c}{s_1} \frac{y_1^4}{(1+y_1^4)^2} - \frac{c}{s_2} \frac{y_2^4}{(1+y_2^4)^2}} \\ &\times \frac{1}{(s_1(y_1^2 - \frac{1}{y_1^2}) - s_2(y_2^2 - \frac{1}{y_2^2}))^2 + 4(s_1 + s_2)^2} \end{aligned} \quad (7.67)$$

For the large s expansion, the expansion parameter is coupled to c as $\frac{c}{s}$ by a scaling. The exponent of the integrand has a pole at $1 + y_i^4 = 0$, which leads to $y_i = \pm(-1)^{\frac{1}{4}}$, $y_i = \pm(-1)^{\frac{3}{4}}$. This pole appears for $p \leq -\frac{3}{2}$, since the term of exponent is $(y^2 + \frac{1}{y^2})^{2p+2}$. Taking the pair of poles $y_1 = \pm(-1)^{\frac{1}{4}}$, $y_2 = \pm(-1)^{\frac{1}{4}}$ or $y_1 = \pm(-1)^{\frac{3}{4}}$, $y_2 = \pm(-1)^{\frac{3}{4}}$, two point function in the large s is evaluated with a contour integral as,

$$\begin{aligned} u(s_1, s_2) &= \frac{c^2}{2^6} \frac{1}{s_1 s_2} - \frac{c^3}{2^{10}} \left(\frac{1}{s_1 s_2^2} + \frac{1}{s_1^2 s_2} \right) + \frac{c^4}{3 \cdot 2^{15}} \left(\frac{5}{s_1 s_2^3} + \frac{7}{s_1^2 s_2^2} + \frac{5}{s_1^3 s_2} \right) \\ &- \frac{c^5}{2^{20} \cdot 3} \left(\frac{7}{s_1 s_2^4} + \frac{13}{s_1^2 s_2^3} + \frac{13}{s_1^3 s_2^2} + \frac{7}{s_1^4 s_2} \right) \\ &+ \frac{c^6}{2^{24} \cdot 15} \left(\frac{21}{s_1 s_2^5} + \frac{49}{s_1^2 s_2^4} + \frac{66}{s_1^3 s_2^3} + \frac{49}{s_1^4 s_2^2} + \frac{21}{s_1^5 s_2} \right) \\ &- \frac{c^7}{2^{28} \cdot 15} \left(\frac{11}{s_1 s_2^6} + \frac{31}{s_1^2 s_2^5} + \frac{53}{s_1^3 s_2^4} + \frac{53}{s_1^4 s_2^3} + \frac{31}{s_1^5 s_2^2} + \frac{11}{s_1^6 s_2} \right) + O(c^8) \end{aligned} \quad (7.68)$$

where $c = (\frac{1}{4})^p = 2^4$. This result is consistent with [31], where there is a factor N , which comes from the logarithmic potential. Above expansion coincides with the result of $N = 0$ in [31], since we do not take the logarithmic potential in (7.67). From this derivation, it can be found that the large s region is related to the pole at the solution of $y^4 + 1 = 0$. This pole exists for $p = -2$, while this pole does not exist for the positive integer p . It may be interesting to discuss $p = -3/2$ and $p = -3$ cases, which seem to be related to the interesting phase of the gravity. The relation to the black hole for $p \rightarrow -\infty$ has been discussed in [7].

For two point function in the weak coupling region, we use the expression of (5.5) with the value of $p = -2$. Since the leading term in the exponent

is $-c\frac{1}{y_i^4}$, we make a change of variable $\frac{1}{y_i^4} = t_i$. Then the exponent becomes $-ct_i/(1+s_it_i)^2$. The two point function $u(s_1, s_2)$ is written as

$$u(s_1, s_2) = -\frac{1}{4}\left(\frac{s_2}{s_1}\right)^{\frac{1}{2}} \oint \frac{dt_1 dt_2}{(2i\pi)^2} t_1^{-\frac{1}{2}} t_2^{-\frac{3}{2}} (1+s_1 t_1)(1+s_2 t_2) e^{-c \sum \frac{t_i}{(1+s_i t_i)^2}} \times \frac{1}{1-g} \quad (7.69)$$

where g is

$$g = 2s_1 t_1 + 2\left(\frac{s_2 t_1}{s_1 t_2}\right)^{\frac{1}{2}} (1-s_2 t_2) - [s_1 t_1 + \left(\frac{s_2 t_1}{s_1 t_2}\right)^{\frac{1}{2}} (1-s_2 t_2)]^2 - \frac{4t_1}{s_1} (s_1 + s_2)^2 \quad (7.70)$$

By the expansion of $1/(1-g) = \sum g^n$, we evaluate the terms of order $s_2^{n_2} s_1^{n_1}$. The selection rule in (2.6) becomes for two marked points of $p = -2$,

$$g + 1 + \frac{1}{2}(j_1 + j_2) = n_1 + n_2 \quad (7.71)$$

From (7.69), we find the terms in the lower orders similar to (2.1)-(2.5) in the previous section. For $g = 1$, $j_1 = j_2 = 0$, we obtain the term $s_1^{\frac{1}{2}} s_2^{\frac{1}{2}}$ as

$$u(s_1, s_2) = c^{-1} s_1^{\frac{1}{2}} s_2^{\frac{1}{2}} \int \frac{dt_1 s t_2}{(2i\pi)^2} \frac{1}{2t_2^{\frac{3}{2}}} (t_1^{\frac{1}{2}} - t_1^{\frac{3}{2}}) e^{-t_1 - t_2} = -\frac{1}{16\pi} s_1^{\frac{1}{2}} s_2^{\frac{1}{2}} \quad (7.72)$$

For the term of order $s_1 s_2$, which is $g = 2$, $j_1 = j_2 = -1$,

$$u(s_1, s_2) = c^{-2} s_1 s_2 \int \frac{dt_1 dt_2}{(2i\pi)^2} e^{-t_1 - t_2} \frac{1}{t_2^2} (-7t_1^2 + 13t_1^3 - t_1^4) = -\frac{5}{64\pi} s_1 s_2 \quad (7.73)$$

where the integral of t_2 is a residue evaluation, and the integral t_1 is the exponential integral for $0 < t_1 < \infty$. The coefficient c is $c = 4^{-p} = 4^2$.

The terms $s_1^{n_1} s_2^{n_2}$ (n_1, n_2 : integers) belong to Ramond sector ($j_1 = j_2 = -1$), and they appear in a pair. The terms of $s_1^{n_1 + \frac{1}{2}} s_2^{n_2 + \frac{1}{2}}$ belong to Neveu-Schwarz sectors with the spin components $j_1 = j_2 = 0$ in $u(s_1, s_2) \sim s_1^{n_1 - \frac{1+j_1}{2}} s_2^{n_2 - \frac{1+j_2}{2}}$.

7-4: Strong coupling of $p = -\frac{3}{2}$

For $p = -\frac{3}{2}$, one point function $u(s)$ is given by

$$u(s) = \frac{i}{2} \oint \frac{dy}{2i\pi} \left(y + \frac{1}{y^3}\right) e^{\frac{c}{\sqrt{s}} \frac{y}{y^4+1}} \quad (7.74)$$

Expanding the exponential, the $u(s)$ is reduced to the contour integral. The term of order c , the integral becomes

$$\oint \frac{dy}{2i\pi} \left(y + \frac{1}{y^3}\right) \frac{c}{\sqrt{s}} \frac{y}{y^4+1} = \oint \frac{dy}{2i\pi} \frac{c}{\sqrt{s}} \frac{1}{y^2} = 0 \quad (7.75)$$

The order of c^2 becomes

$$I = \frac{c^2}{s} \oint \frac{dy}{2i\pi} \frac{1}{y(y^4+1)} = 0 \quad (7.76)$$

The poles are at $y = 0$ and at $y = \pm(-1)^{1/4}, \pm(-1)^{3/4}$. The residues of these poles are 1 and $-\frac{1}{4}, -\frac{1}{4}, -\frac{1}{4}, -\frac{1}{4}$, respectively. Adding these contributions, one finds the integral is zero. For the term of order c^3 , with adding four poles, it becomes

$$I = \frac{c^3}{s^{3/2}} \oint \frac{dy}{2i\pi} \frac{1}{(y^4 + 1)^2} = 0 \quad (7.77)$$

since the contributions from four poles are $\pm\frac{3}{16}(-1)^{1/4}, \pm\frac{3}{16}(-1)^{3/4}$. Thus the contour integral of $u(s)$ is expanded in the series of $\frac{c^{4n}}{s^{2n}}$.

$$\begin{aligned} u(s) = & 2i \left(-\frac{ic^4}{2^8 s^2} - \frac{1}{2^{19} \cdot 3 \cdot 5} \frac{ic^8}{s^4} - \frac{13}{2^{30} \cdot 3^4 \cdot 5^2 \cdot 7} \frac{ic^{12}}{s^6} \right. \\ & - \frac{17 \cdot 19}{2^{42} \cdot 3^5 \cdot 5^2 \cdot 7^2 \cdot 11 \cdot 13} \frac{ic^{16}}{s^8} - \frac{23ic^{20}}{2^{54} \cdot 3^7 \cdot 5^2 \cdot 7 \cdot 11 \cdot 13 \cdot 17 \cdot s^{10}} \\ & - \frac{29 \cdot 31ic^{24}}{2^{65} \cdot 3^7 \cdot 5^3 \cdot 7^3 \cdot 11^2 \cdot 13 \cdot 17 \cdot 19s^{12}} - \frac{29 \cdot 31 \cdot 37ic^{28}}{2^{76} \cdot 3^{10} \cdot 5^6 \cdot 7^3 \cdot 11 \cdot 13^2 \cdot 17 \cdot 19 \cdot 23s^{14}} \\ & \left. + O(c^{32}) \right) \end{aligned} \quad (7.78)$$

This series for $p = -\frac{3}{2}$ can be expressed as

$$\begin{aligned} u(s) &= 2 \sum_{n=1}^{\infty} \frac{(6n-3)!!}{2^{10n-2} (4n-1)!! (4n-3)!! (2n-1)!! (n-1)! n! (2n-1)} \frac{c^{4n}}{s^{2n}} \\ &= 2 \sum_{n=1}^{\infty} \frac{(6n-3)!(2n-2)!}{(3n-2)!(4n-3)!(4n-1)!(2n-1)! n! 2^{8n}} \frac{c^{4n}}{s^{2n}} \end{aligned} \quad (7.79)$$

When a logarithmic term is added, the term $((u+1)/(u-1))^N$ is multiplied to (7.75). This term due to a logarithmic potential is written in the variable $u+1 = 2/(1-e^{-z})$ and $u-1 = 2e^{-z}/(1-e^{-z})$ as

$$\left(\frac{u-1}{u+1} \right)^N = e^{-Nz} \quad (7.80)$$

one obtain

$$\begin{aligned} U(s) &= \int du e^{cs^{-1/2}((u+1)^{-1/2} - (u-1)^{-1/2})} \left(\frac{u-1}{u+1} \right)^N \\ &= -2 \int dz \frac{e^{-z}}{(1-e^{-z})^2} e^{-Nz} e^{cs^{-1/2} 2^{-1/2} (1-e^{-z})^{1/2} (1-e^{z/2})} \end{aligned} \quad (7.81)$$

For the terms of $(c/\sqrt{s})^4$, it leads to

$$\begin{aligned} & -2 \int dz e^{-Nz} e^{-z} \frac{1}{4!} \frac{c^4}{s^2} 2^{-2} (1-e^{\frac{z}{2}})^4 \\ &= (-2) \frac{1}{4!} \frac{1}{4} \frac{c^4}{s^2} \frac{6}{N(N^2-1)(4N^2-1)} \end{aligned} \quad (7.82)$$

The term of $(c/\sqrt{s})^8$ is

$$-2 \left(\frac{1}{8! 2^4} \right) \left(\frac{c^8}{s^4} \right) \frac{45 \cdot 7! (4N^2 - 3)}{N(N^2 - 1)(N^2 - 4)(N^2 - 9)(4N^2 - 1)(4N^2 - 9)(4N^2 - 25)} \quad (7.83)$$

For the sector of $\frac{c^{4n}}{s^{2n}}$, one point function $U(s)$ of $p = -\frac{3}{2}$ has a form of characteristic polynomials, which are factorized. In the limit $N \rightarrow 0$, the expression becomes consistent with (7.79).

7-5: Strong coupling of $p = -3$

For $p = -3$ case, the strong coupling expansion of the inverse of s is

$$u(s) = \frac{i}{2} \oint \frac{dy}{2i\pi} \left(y + \frac{1}{y^3}\right) e^{\frac{c}{s^2} \frac{y^6(y^4-1)}{(y^4+1)^4}} \quad (7.84)$$

This large s expansion is given by the residue at $y = (-1)^{1/4}$,

$$\begin{aligned} u(s) &= \frac{1}{s^4} + \frac{3}{8s^8} + \frac{13}{2^6 \cdot 3s^{12}} + \frac{17 \cdot 19}{2^{10} \cdot 3^2 \cdot 5s^{16}} + \frac{23}{2^{14} \cdot 3s^{20}} \\ &+ \frac{29 \cdot 31}{2^{17} \cdot 3^2 \cdot 5 \cdot 7s^{24}} + \frac{29 \cdot 31 \cdot 37}{2^{20} \cdot 3^5 \cdot 5^2 \cdot 7s^{28}} + \frac{37 \cdot 41 \cdot 43}{2^{25} \cdot 3^4 \cdot 5^2 \cdot 7^2 s^{32}} \\ &+ \frac{37 \cdot 41 \cdot 43 \cdot 47}{2^{30} \cdot 3^6 \cdot 5^3 \cdot 7 \cdot 11s^{36}} + \frac{41 \cdot 43 \cdot 47 \cdot 53}{2^{33} 3^8 5^3 \cdot 7 \cdot 13s^{40}} + \frac{47 \cdot 53 \cdot 59 \cdot 61}{2^{36} 3^7 5^4 \cdot 7 \cdot 11 \cdot 13s^{44}} \\ &+ \dots \end{aligned} \quad (7.85)$$

with $c = 2^6$. These terms of order of $\frac{1}{s^{2k}}$ is expressed as

$$\begin{aligned} u(s) &= \sum_{n=1}^{\infty} \frac{(6n-5)!!}{(4n-1)!!(2n-3)!!(n-1)!(n-1)!} \frac{3}{n} \frac{1}{2^{2n-2} s^{4n}} \\ &= \sum_{n=0}^{\infty} \frac{(6n+1)!(2n+1)}{(4n+4)!(3n)!n!} \frac{8}{4^n s^{4n+4}} \end{aligned} \quad (7.86)$$

With a logarithmic potential, $p = -3$ one point function $U(s)$ becomes

$$U(s) = \oint \frac{du}{2i\pi} e^{\frac{c}{s^2} \left(\frac{1}{(u+1)^2} - \frac{1}{(u-1)^2}\right)} \left(\frac{u+1}{u-1}\right)^N \quad (7.87)$$

Similar to $p = -2$ case, we make a change of variables of (7.54). Then we have

$$\begin{aligned} U(s) &= -2 \int dz e^{-Nz} \frac{e^{-z}}{(1-e^{-z})^2} e^{-\frac{c}{4s^2} e^{2z} (1-e^{-z})^3 (1+e^{-z})} \\ &= -2 \int dz \frac{e^{-z}}{(1-e^{-z})^2} e^{\frac{c}{4s^2} [-2e^{-z} + e^{-2z} - e^{2z} + 2e^z]} e^{-Nz} \end{aligned} \quad (7.88)$$

This gives a strong coupling expansion. The term of order $\frac{c}{s^2}$ is

$$-\left(\frac{c}{s^2}\right) \frac{1}{N^2 - 1}$$

The term of order $\frac{c^2}{s^4}$ is

$$\begin{aligned} &-\left(\frac{c}{4s^2}\right)^2 \int dz e^{-Nz} e^{3z} (1-e^{-z})^4 (1+e^{-z})^2 \\ &= -\left(\frac{c}{4s^2}\right)^2 \frac{48(2N^2 - 3)}{N(N^2 - 1)(N^2 - 4)(N^2 - 9)} \end{aligned} \quad (7.89)$$

The order of $\frac{c^3}{s^6}$ in $U(s)$ for $p = -3$ becomes

$$\frac{8965}{4} \left(\frac{c^3}{s^6}\right) \frac{N^2 - 7}{(N^2 - 1)(N^2 - 4)(N^2 - 9)(N^2 - 16)(N^2 - 25)}$$

The order of $\frac{c^4}{s^8}$ becomes

$$-9450 \frac{c^4}{s^8} \frac{2N^4 - 38N^2 + 63}{N(N^2 - 1)(N^2 - 4)(N^2 - 9)(N^2 - 16)(N^2 - 25)(N^2 - 36)(N^2 - 49)}$$

The denominator is same as $p = -2$ case, but the numerator is different from $p = -2$ case. There is no term of order $\frac{1}{s^{2n+1}}$. The terms of order $\frac{1}{s^{4n}}$ have a factor $\frac{1}{N}$ and there is no such factor for other terms. When $N \rightarrow 0$, $U(s)$ becomes in the power series of $\frac{1}{s^{4n}}$. Indeed the expansion of (7.85) agrees with the expansion of (7.88), and the coefficient is consistent with $c = 2$.

8 General p and recursion equation

For general p ($p < 0$), One point function with a logarithmic term in a strong coupling region is expressed with $r = -p - 1$, ($p < 0$),

$$U(s) = -2 \int_0^\infty dz e^{-Nz} \frac{e^{-z}}{(1 - e^{-z})^2} e^{\frac{c}{(2s)^r} (1 - e^{-z})^r (1 - e^{rz})} \quad (8.1)$$

where the logarithmic term is expressed as $[(u - 1)/(u + 1)]^N = e^{-Nz}$ in (7.54). The strong coupling expansion becomes

$$\begin{aligned} U(s) &= -2 \sum_{l=0}^{\infty} \frac{c^l}{(2s)^{rl} l!} \int_0^\infty dz e^{-Nz} e^{-z} (1 - e^{-z})^{rl-2} (1 - e^{rz})^l \\ &= -2 \sum_{l=0}^{\infty} \frac{c^l}{(2s)^{rl} l!} I_l[r] \end{aligned} \quad (8.2)$$

where $I_l[r]$ becomes for $l = 1$ as

$$\begin{aligned} I_1[1] &= \frac{1}{N}, \quad I_1[2] = -\frac{2}{N^2 - 1}, \quad I_1[3] = -\frac{6N}{(N^2 - 1)(N^2 - 4)} \\ I_1[4] &= -\frac{24(N^2 + 1)}{(N^2 - 1)(N^2 - 4)(N^2 - 9)} \\ I_1[5] &= -\frac{120N(N^2 + 5)}{(N^2 - 1)(N^2 - 4)(N^2 - 9)(N^2 - 16)} \\ I_1[6] &= -\frac{720(N^4 + 15N^2 + 8)}{(N^2 - 1)(N^2 - 4)(N^2 - 9)(N^2 - 16)(N^2 - 25)} \\ I_1[7] &= -\frac{5040N(N^4 + 35N^2 + 84)}{(N^2 - 1)(N^2 - 4)(N^2 - 9)(N^2 - 25)(N^2 - 36)} \end{aligned} \quad (8.3)$$

The numerator of $I_{l=1}[r]$ is $r!(r - 2)!$ for $N = 1$. We have investigated $I_l[1]$ in (7.59) for arbitrary l . For $l = 2$, it becomes

$$\begin{aligned} I_2[1] &= \frac{2}{N(N^2 - 1)} \\ I_2[2] &= \frac{48(2N^2 - 3)}{N(N^2 - 1)(N^2 - 4)(N^2 - 9)} \\ I_2[3] &= \frac{2160(3N^4 - 11N^2 + 80)}{N(N^2 - 1)(N^2 - 4)(N^2 - 9)(N^2 - 16)(N^2 - 25)} \end{aligned} \quad (8.4)$$

For $l = 3$, it leads to

$$\begin{aligned}
I_3[1] &= -\frac{24}{N(N^2-1)(N^2-4)} \\
I_3[2] &= \frac{40320(N^2-7)}{(N^2-1)(N^2-4)(N^2-9)(N^2-16)(N^2-25)} \\
I_3[3] &= -3259200(3N^6-51N^4+888N^2+6272) \prod_{m=1}^8 \frac{1}{N^2-m^2}
\end{aligned} \tag{8.5}$$

For $l = 4$, it becomes

$$\begin{aligned}
I_4[1] &= \frac{720}{N(N^2-1)(N^2-4)(N^2-9)} \\
I_4[2] &= (10)! \cdot 8(2N^4-38N^2+63) \frac{1}{N} \prod_{m=1}^7 \frac{1}{N^2-m^2} \\
I_4[3] &= (14)! \cdot 81(N^8-46N^6+1649N^4+14396N^2+422400) \\
&\times \frac{1}{N} \prod_{m=1}^{11} \frac{1}{N^2-m^2}
\end{aligned} \tag{8.6}$$

For two point function $U(s_1, s_2)$ is evaluated for arbitrary r , similar to (7.65) for $p = -2$ ($r = 1$) case.

These integrands of $I_l[r]$ in (8.2) is polynomials of q , if we replace $q = e^{-z}$. For instance, for a few r - cases, the polynomial $P_{I_l[r]}(q)$ are

$$\begin{aligned}
P_{I_l[1]}(q) &= (-1)^l q^{N-l+1} (1-q)^{2l-2} = (-1)^l q^N (q^{-1/2} - q^{1/2})^{2l-2} \\
P_{I_l[2]}(q) &= (-1)^l q^{N-2l+1} (1-q)^{3l-2} (q+1)^l \\
&= (-1)^l q^N (q^{-1/2} - q^{1/2})^{3l-2} (q^{-1/2} + q^{1/2})^l \\
P_{I_l[3]}(q) &= (-1)^l q^{N-3} ((1-q)^{4l-2} (q^2+q+1)^l \\
&= (-1)^l q^N (q^{-1/2} - q^{1/2})^{4l-2} (q+1+q^{-1})^l \\
P_{I_l[4]}(q) &= (-1)^l q^N (q^{-1/2} - q^{1/2})^{5l-2} (q^{-1/2} + q^{1/2})^l (q+q^{-1})^l
\end{aligned} \tag{8.7}$$

These polynomials leads to the expressions of $I_l[r]$. From the above polynomial $P_{I_l[r]}(q)$, we obtain the recursion relation. For $r = 1$, dividing by $(-1)^l q^N$, obtained polynomial $\hat{P}_{I_l[1]}(q)$ has a relation,

$$\hat{P}_{I_{l+1}[1]}(q) = \hat{P}_{I_l[1]}(q)(-2+q^{-1}+q) \tag{8.8}$$

The last factor make shifts of the exponent N to $N-1$ and $N+1$ for q^{-1} and q . Thus $I_2[1]$ is obtained from $I_1[1]$ as

$$I_2[1] = -\frac{2}{N} + \frac{1}{N-1} + \frac{1}{N+1} = \frac{2}{N(N^2-1)} \tag{8.9}$$

$I_3[1]$ is obtained from a recursion relation of (8.8), as

$$\begin{aligned}
I_3[1] &= -2 \frac{2}{N(N^2-1)} + \frac{2}{(N-1)((N-1)^2-1)} + \frac{2}{(N+1)((N+1)^2-1)} \\
&= \frac{24}{N(N^2-1)(N^2-4)}
\end{aligned} \tag{8.10}$$

This leads to (7.59) for general l . The factorized polynomial of $I_l[1]$ corresponds to a row of Young diagram of the length l , which is obtained for a single trace $U(s)$. For other value of r , the recursion equations for $I_l[r]$ are obtained from (8.7).

For $r = 2$, the recursion relation becomes

$$P_{I_{l+1}[2]} = P_{I_l[2]} \times (q^{-2} - 2q^{-1} + 2q - q^2) \quad (8.11)$$

and from $I_1[2] = -2/(N^2 - 1)$, it leads to

$$\begin{aligned} I_2[2] &= \left(-\frac{2}{N^2-1}\right)|_{N \rightarrow N+2} - 2\left(-\frac{2}{N^2-1}\right)|_{N \rightarrow N+1} + 2\left(-\frac{2}{N^2-1}\right)|_{N \rightarrow N-1} \\ &\quad - \left(-\frac{2}{N^2-1}\right)|_{N \rightarrow N-2} \\ &= \frac{48(2N^2-3)}{N(N^2-1)(N^2-4)(N^2-9)} \end{aligned} \quad (8.12)$$

By replacing N by $N+2, N+1, N-1$, and $N-2$ according to the recursion relation of (8.11), we obtain the next order,

$$I_3[2] = -\frac{40320(N^2-7)}{N(N^2-1)(N^2-4)(N^2-9)(N^2-16)(N^2-25)} \quad (8.13)$$

$$I_4[2] = \frac{29030400(2N^4-38N^2+63)}{N \prod_{m=1}^7 (N^2-m^2)} \quad (8.14)$$

This successive recursion by the replacement $[(N \rightarrow N+2) - 2(N \rightarrow N+1) + 2(N \rightarrow N-1) - (N \rightarrow N-2)]$ to $I_l[2]$ gives a convenient method of $I_{l+1}[2]$, which agrees with (8.5). Thus we find a recursive method of $I_l[r]$ for arbitrary l and r .

The strong coupling expansion of two point function $U(s_1, s_2)$ of $r = -p-1$ ($p < 0$) becomes

$$\begin{aligned} U(s_1, s_2) &= \frac{s_1}{s_2} \int dz_1 dz_2 \frac{q_1^{N_1} q_2^{N_2} q_1}{(1-q_1)^2} \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \left(\frac{s_1}{s_2}\right)^{m+n} \left(\frac{1-q_2}{1-q_1}\right)^{m+n} q_1^m q_2^n \\ &\quad \times e^{\frac{c}{(2s_1)^r} (1-q_1)^r (1-q_1^{-r}) + \frac{c}{(2s_2)^r} (1-q_2)^r (1-q_2^{-r})} \end{aligned} \quad (8.15)$$

where $q_1 = e^{-z_1}$, $q_2 = e^{-z_2}$. The distinction of N_1 and N_2 is for the convenience of the recursion relation, and we put $N_1 = N_2 = N$ after all. The strong coupling expansion of $\frac{1}{s_1^{l_1} s_2^{l_2}}$ is obtained for $r = 1$ ($p=-2$). The two variable polynomial $P_{l_1, l_2}(q_1, q_2)$ is obtained by the expansion of the exponential term. Writing $U(s_1, s_2)$ as

$$U(s_1, s_2) = \sum_{l_1=0}^{\infty} \sum_{l_2=0}^{\infty} \frac{1}{s_1^{l_1} s_2^{l_2}} I_{l_1, l_2}[r] \quad (8.16)$$

We have for $r = 1$, the following recursion relation is obtained,

$$I_{1,1}[1] = \frac{1}{N_2} \frac{1}{N_1(N_1^2-1)} \quad (8.17)$$

$$I_{1,n}[1] = -(n+1)I_{n+1}[1] \quad (8.18)$$

This reads to

$$\begin{aligned} I_{1,2}[1] &= -\frac{12}{(N^2-1)(N^2-4)}, \quad I_{1,3}[1] = \frac{120}{(N^2-1)(N^2-4)}, \\ I_{1,4}[1] &= -\frac{1680}{(N^2-1)(N^2-4)(N^2-9)(N^2-16)} \end{aligned} \quad (8.19)$$

and they agree with the character expansion of the unitary matrix model ($C_{1,2}^c(N)$, $C_{1,3}^c(N)$, $C_{1,4}^c(N)$, which are the coefficients of $(\frac{\text{tr}AB}{N})(\frac{\text{tr}(AB)^m}{N})$ in the unitary matrix model [49]).

The coefficients C_α^c appears in the integration of the unitary matrix U for the unitary matrix model

$$\begin{aligned} J(AB) &= \int dU e^{g(\text{tr}AU + \text{tr}BU^\dagger)} \\ &= \text{Exp}[\sum_n \frac{g^{2n}}{n!} \sum_\alpha C_\alpha^c (\text{tr}AB)^{\alpha_1} (\text{tr}(AB)^2)^{\alpha_2} \dots (\text{tr}(AB)^n)^{\alpha_n}] \end{aligned} \quad (8.20)$$

From (8.15), $I_{2,n}[1]$ is evaluated,

$$I_{2,2}[1] = \frac{18(3N^2-7)}{(N^2-1)^2(N^2-4)(N^2-9)} \quad (8.21)$$

$$I_{2,3}[1] = \frac{-480(3N^2-13)}{(N^2-1)^2(N^2-4)(N^2-9)(N^2-16)} \quad (8.22)$$

which are again consistent with $C_{2,2}^c(N)$, $C_{2,3}^c(N)$ in [49, 31]. Thus $U(s_1, s_2)$ in (8.15) becomes a generated function for the coefficients $C_{2,n}^c$ in the unitary matrix model.

For r ($p = -r - 1$), the unitary matrix model of (8.20) is modified as

$$\begin{aligned} J(AB)[r] &= \int dU e^{g(\text{tr}AU^r + \text{tr}BU^\dagger)} \\ &= \text{Exp}[\sum_n \frac{g^{2n}}{n!} \sum_\alpha C_\alpha^c[r] (\text{tr}AB^r)^{\alpha_1} (\text{tr}(AB^r)^2)^{\alpha_2} \dots (\text{tr}(AB^r)^n)^{\alpha_n}] \end{aligned} \quad (8.23)$$

n point function $U(s_1, \dots, s_n) = \langle \text{tr}e^{s_1 M} \dots \text{tr}e^{s_n M} \rangle$ is expressed for the external source $A = \{a_\alpha\}$ [7],

$$U(s_1, \dots, s_n) = e^{\sum_{i=1}^n s_i^2} \oint \prod_{i=1}^n \frac{du_i}{2i\pi} e^{\sum u_i s_i} \prod_{\alpha=1}^N \prod_{i=1}^n \left(\frac{a_\alpha - (u_i + s_i)}{a_\alpha - u_i} \right) \det \frac{1}{u_i - u_j + s_i} \quad (8.24)$$

With the tuning of a_α , it leads to n -point function for the p spin curve,

$$U(s_1, \dots, s_n) = \int \prod_{i=1}^n \frac{du_i}{2i\pi} e^{\sum c 2^{p+1} [(u_i + \frac{s_i}{2})^{p+1} - (u_i - \frac{s_i}{2})^{p+1}]} \det \frac{1}{u_i - u_j + \frac{1}{2}(s_i + s_j)} \quad (8.25)$$

In the strong coupling expansion, by taking the longest cycles for the connected parts,

$$U(s_1, s_2, \dots, s_n) = \sum_{l_i} \frac{1}{s_1^{l_1} s_2^{l_2} \dots s_n^{l_n}} I_{l_1, l_2, \dots, l_n}[r] \quad (8.26)$$

From this expression, three point function for r ($r = -p-1$), reads by adding the logarithmic potential,

$$\begin{aligned}
U(s_1, s_2, s_3) &= -\frac{s_3}{s_1} \int \prod_{i=1}^3 du_i e^{\sum_{i=1}^3 \frac{c}{(2s_i)^r} [(u_i+1)^{-r} - (u_i-1)^{-r}]} \prod_{i=1}^3 \left(\frac{u_i-1}{u_i+1} \right)^N \\
&\times \sum_{l_1=0}^{\infty} \sum_{l_2=0}^{\infty} \sum_{l_3=0}^{\infty} \frac{s_2^{l_1-l_2} s_3^{l_2+l_3}}{s_1^{l_1+l_3}} \frac{(u_2-1)^{l_1} (u_2+1)^{-l_2-1} (u_3+1)^{l_3} (u_3-1)^{l_2}}{(u_1+1)^{l_1+1} (u_1-1)^{l_3+1}} \\
&= -\frac{s_3}{s_1} \int \prod_{i=1}^3 dz_i \prod_{i=1}^3 \frac{q_i}{(1-q_i)^2} e^{\sum_{i=1}^3 \frac{c}{(2s_i)^r} [(1-q_i)^r (1-q_i^{-r})]} q_1^N q_2^N q_3^N \\
&\sum_{l_1, l_2, l_3=0}^{\infty} \frac{s_2^{l_1-l_2} s_3^{l_2+l_3}}{s_1^{l_1+l_3}} q_1^{-l_3-1} q_2^{l_1} q_3^{l_2} (1-q_1)^{l_1+l_3+2} (1-q_2)^{l_2-l_1+1} (1-q_3)^{-l_2-l_3}
\end{aligned} \tag{8.27}$$

where $u_i = \frac{1+e^{-z_i}}{1-e^{-z_i}} = \frac{1+q_i}{1-q_i}$, $q_i = e^{-z_i}$. This gives the polynomial $P(q_1, q_2, q_3)$ and $I_{n_1, n_2, n_3}[r]$. The term of order $\frac{1}{s_1}$ corresponds to $l_1 = l_3 = 0$. Thus expanding in the power series of $\frac{1}{s_1}$, we obtain the function $U(s_1, s_2, s_3) = \sum_n f_n(s_2, s_3) \frac{1}{s_1^n}$. From the determinat, there are two terms for the longest cyclic terms.

$$\begin{aligned}
f_1(s_2, s_3) &= -\frac{1}{N} s_2 s_3 \int \prod_{i=2}^3 dz_i (q_2 q_3)^N \frac{q_2 q_3}{(1-q_2)(1-q_3)} \\
&\left(\frac{1}{s_2(1-q_3) - s_3 q_3(1-q_2)} + \frac{1}{s_3(1-q_2) - s_2 q_2(1-q_3)} \right) \\
&\times e^{\sum_{i=2}^3 \frac{c}{(2s_i)^r} [(1-q_i)^r (1-q_i^{-r})]}
\end{aligned} \tag{8.28}$$

When $r = 1$, we obtain $U(s_1, s_2, s_3) = \frac{1}{s_1 s_2 s_3} \frac{8}{N^2(N^2-1)(N^2-4)}$, which is consistent with a quantity $C_{(1,1,1)}$ in [49, 31].

In the next section, we discuss the relation to characteristic polynomials and to Chern-Simons theory, which is realized by the matrix model of Stieltjes-Wigert polynomial [58].

9 Characteristic polynomials and p spin curves

As discussed in I, II [8, 9] and in the previous section 5, the correlation function $u(s_1, \dots, s_n)$ is given by the integral of the Laurent expansion of y_i . The integral simply becomes an exponential integral of y_i (Laplace transformation) and the intersection numbers are essentially obtained from the coefficients of the combinatorial terms of the Laurent expansion of y_i . The polynomial of y is introduced for the investigation of half integer p in [8, 9], but as shown in this article, the Laurent expansion of y is powerful for the evaluation of the intersection numbers of n -point function. This Laurent expansion is similar to the Laurent expansions in knot theories, Alexander, Jones and HOMFLYPT polynomials in terms of t . In this section, we discuss the relation to characters and knot polynomials.

The knot matrix model is discussed through Harer-Zagier (HZ) formula defined below [50, 51], which becomes a generating function of Gaussian mean.

The n point function $U(s_1, \dots, s_n)$ is interpreted as the Wilson loops, which are generating functions of the following Gaussian means

$$\langle \text{tr}M^{i_1} \dots \text{tr}M^{i_n} \rangle = \frac{\int \text{tr}M^{i_1} \dots \text{tr}M^{i_n} e^{-\frac{1}{2}\text{tr}M^2} dM}{\int e^{-\frac{1}{2}\text{tr}M^2} dM} \quad (9.1)$$

where M is $N \times N$ Hermitian matrix. HZ function is defined by [51, 52]

$$\Phi(z, N) = \sum_{i=0}^{\infty} \frac{z^{2i}}{(2i-1)!!} \langle \text{tr}M^{2i} \rangle = \frac{1}{2z^2} \left[\left(\frac{1+z^2}{1-z^2} \right)^N - 1 \right] \quad (9.2)$$

from which one can obtain, for instance,

$$\langle \text{tr}M^4 \rangle = 2N^3 + N \quad (9.3)$$

By the sum of N , and by the use of HZ transform, one can write a Laplace transform of above quantity,

$$Z(\lambda) = \sum_{N=0}^{\infty} \lambda^N \langle \text{tr}M^4 \rangle = \sum_{N=0}^{\infty} (2N^3 + N) \lambda^N = \frac{3\lambda(1+\lambda)}{(1-\lambda)^4}. \quad (9.4)$$

The knot polynomial of HOMFLYPT for a torus knot is realized as an average of a character [51]. For instance, the trefoil $T(2, 3)$, which is a torus knot of $(2, 3)$ type, and the normalized HOMFLYPT polynomial \bar{P} , by the multiplication of $(v - v^{-1})/z$ with the replacement of $v \rightarrow q^N$ and $z \rightarrow q - q^{-1}$, is expressed as

$$\bar{P}(T(2, 3) : v, z) = \frac{(q^2 + q^{-2})q^{5N} - (q^2 + 1 + q^{-2})q^{3N} + q^N}{q - q^{-1}} \quad (9.5)$$

By the HZ transformation of $\bar{P}(T(2, 3) : v, z)$, one obtains

$$Z(q, \lambda) = \sum_{N=0}^{\infty} \lambda^N \bar{P}(T(2, 3) : v, z) = \frac{\lambda(1 - \lambda q^9)}{(1 - q\lambda)(1 - q^3\lambda)(1 - q^5\lambda)} \quad (9.6)$$

which is Hilbert-Poincare polynomial with ideals. This factorized form may be interpreted as BPS spectrum.

The variable q , which appears as $z = q - \frac{1}{q}$, provides the Laurent expansion of q . There is a similarity between p -spin curve and torus knot $T(2, 2a + 1)$. The HOMFLYPT polynomial $P(T(2, 2a + 1) : v, z)$ for torus knot $T(2, 2a + 1)$ is obtained from skein relation as

$$\begin{aligned} P(T(2, 2a + 1) : v, z) &= \frac{v^{2a}}{2^{2a+1}\sqrt{z^2 + 4}} [(z + \sqrt{z^2 + 4})^{2a+1} - (z - \sqrt{z^2 + 4})^{2a+1}] \\ &+ 2(1 - v^2) \frac{1}{z} [(z + \sqrt{z^2 + 4})^{2a} - (z - \sqrt{z^2 + 4})^{2a}] \end{aligned} \quad (9.7)$$

It reduces to Jones polynomial when $v = t$, $z = t^{\frac{1}{2}} - t^{-\frac{1}{2}}$ for torus knot; $P = t + t^3 - t^4$.

For $v = 1$, $P(T(2, 2a + 1) : v, z)$ becomes Alexander polynomial $\Delta(T(2, 2a + 1) : t)$,

$$\begin{aligned} &P(T(2, 2a + 1) : v = 1, z) \\ &= \frac{1}{2^{2a+1}\sqrt{z^2 + 4}} [(z + \sqrt{z^2 + 4})^{2a+1} - (z - \sqrt{z^2 + 4})^{2a+1}] \end{aligned} \quad (9.8)$$

Putting $z = t - \frac{1}{t}$, $\sqrt{z^2 + 4}$ becomes $t + \frac{1}{t}$. Then, above polynomial becomes

$$P(T(2, 2a + 1) : v = 1, t - \frac{1}{t}) = \frac{1}{t + \frac{1}{t}} (t^{2a+1} - \frac{(-1)^{2a+1}}{t^{2a+1}}) \quad (9.9)$$

Alexander polynomial for a torus link $T(n, m)$ is denoted by $\Delta(T(n, m) : t)$, and it is expressed as

$$\Delta(T(n, m) : t) = t^{-\frac{(n-1)(m-1)}{2}} \frac{(t^{\frac{nm}{d}} - 1)^d (t - 1)}{(t^n - 1)(t^m - 1)} \quad (9.10)$$

where d is a greatest common divisor of n and m ; $d = gcd(n, m)$. Alexander polynomial $\Delta(T(2, n) : t^2)$ of torus link $T(2, n)$, when n is odd it becomes a torus knot, which is defined as is expressed as

$$\begin{aligned} \Delta(T(2, 2) : t^2) &= t - \frac{1}{t} \\ \Delta(T(2, 3) : t^2) &= t^2 + \frac{1}{t^2} - 1 \\ \Delta(T(2, 2n) : t^2) &= \sum_{l=1}^n (t^{2l-1} - \frac{1}{t^{2l-1}}) (-1)^{n+l} \\ \Delta(T(2, 2n+1) : t^2) &= \sum_{l=1}^n (t^{2l} + \frac{1}{t^{2l}}) (-1)^{n+l} + (-1)^n \end{aligned} \quad (9.11)$$

Noting the expression of the exponent I_p of $U(s)$ is written as

$$\begin{aligned} I_p(y) &= cs^{p+1} [(u+1)^{p+1} - (u-1)^{p+1}] \\ &= c \left(\frac{is}{2}\right)^{p+1} \left[\left(y - \frac{i}{y}\right)^{2p+2} - \left(y + \frac{i}{y}\right)^{2p+2} \right], \end{aligned} \quad (9.12)$$

where $u = \frac{i}{2}(y^2 - \frac{1}{y^2})$. This polynomial becomes when p is odd,

$$I_p(y) = 2ic \left(\frac{is}{2}\right)^{p+1} \sum_{l=1}^{(p+1)/2} \frac{(2p+2)! (-1)^{l+1}}{(2l-1)! (2p+3-2l)!} \left[y^{2p+4-4l} - \frac{1}{y^{2p+4-4l}} \right] \quad (9.13)$$

and for p is even number,

$$\begin{aligned} I_p(y) &= 2ic \left(\frac{is}{2}\right)^{p+1} \left(\sum_{l=1}^{p/2} \frac{(2p+2)! (-1)^l}{(2l-1)! (2p+3-2l)!} \left[y^{2p+4-4l} + \frac{1}{y^{2p+4-4l}} \right] \right. \\ &\quad \left. + \frac{(2p+2)!}{(p+1)! (p+1)!} (-1)^{\frac{p}{2}+1} \right) \end{aligned} \quad (9.14)$$

Thus the exponent of p -spin curve $I_p(y)$ is written by a linear combination of Alexander polynomial Δ in (9.11), by putting $y^2 = t$

For $p = 3$, the term of (9.12) becomes in terms of t in (9.9) with $y^2 = t$,

$$I_p(y) = -ics^4 \left[\left(y^6 - \frac{1}{y^6}\right) - 7 \left(y^2 - \frac{1}{y^2}\right) \right] \quad (9.15)$$

which is written as a combination of $\Delta(T(2, 4) : t^2)$ and $\Delta(T(2, 2) : t^2)$ in (9.11).

For $p = 2$, the term of (9.12) becomes

$$I_p(y) = -cs^3 \left(\frac{3}{2} \left(y^4 + \frac{1}{y^4} \right) - 5 \right) \quad (9.16)$$

which is written by Alexander knot polynomial $\Delta(T(2, 3) : t^2)$ of (9.11). When p is odd, I is expressed as linear combination of Alexander polynomial of torus link $\Delta(T(2, 2n) : t^2)$. For even p , it is written as a linear combination of Alexander polynomial of torus knot $\Delta(T(2, 2n + 1) : t^2)$. The even and odd types correspond to even p and odd integer p . The odd integer p case represents the closure of the gap [61].

For half integer spin p , the polynomial I_p of (9.12) is described by the mixture of $\Delta(T(2, 2n) : t^2)$ and $\Delta(T(2, 2n + 1) : t^2)$. For $p = \frac{2n+1}{2}$ ($n=0,1,2$), I_p is given by

$$\begin{aligned} I_{p=\frac{1}{2}}(y) &= 2ic \left(\frac{is}{2} \right)^{\frac{3}{2}} \frac{1}{y} \left[\left(y^2 + \frac{1}{y^2} \right) + 2 \left(y^2 - \frac{1}{y^2} \right) \right] \\ I_{p=\frac{3}{2}}(y) &= 2ic \left(\frac{is}{2} \right)^{\frac{5}{2}} \frac{1}{y} \left[3 \left(y^4 + \frac{1}{y^4} \right) + 2 \left(y^4 - \frac{1}{y^4} \right) - 10 \right] \\ I_{p=\frac{5}{2}}(y) &= 2ic \left(\frac{is}{2} \right)^{\frac{7}{2}} \frac{1}{y} \left[3 \left(y^6 + \frac{1}{y^6} \right) + 4 \left(y^6 - \frac{1}{y^6} \right) - 7 \left(y^2 + \frac{1}{y^2} \right) - 28 \left(y^2 - \frac{1}{y^2} \right) \right] \end{aligned} \quad (9.17)$$

In this case, the combination of odd and even polynomial, divided by y , appears.

The relation of knots invariants to Chern-Simons theory is known. The Wilson loop average over Chern-Simons action is written by Homflypt polynomials with $q = e^{\frac{2\pi i}{k+N}}$, where k is a level of $SU(N)$. In our notation, k is given by $p = k + 2$ for p spin curve. The variable y is replaced by $y^2 = q = e^{\frac{2\pi i}{p}}$. Putting this replacement in $I_p(y)$, we have a polynomial of $e^{\frac{2i\pi m}{p}}$, where $m = 0, 1, \dots, p-1$. Thus there appears interesting relation between polynomial $I_p(y)$ and Wilson loop average in the Chern-Simons theory. The matrix model for the torus knot $T(n, m)$ is known as [60]. The average W_R of Wilson loop in the representation of R is expressed by a matrix model,

$$W_R = \int \int dv e^{-\sum_i \frac{v_i^2}{2g_s}} \prod_{i < j} \sinh\left(\frac{v_i - v_j}{n}\right) \sinh\left(\frac{v_i - v_j}{m}\right) ch_R(e^v) \quad (9.18)$$

where ch_R is a character. The sinh measure in the Chern-Simons matrix model has been considered for the lattice gauge matrix models for $U(N)$ in the large N limit (Table I in [40]). The exponent v_i^2 is obtained from the lattice gauge action in the limit $g_s \rightarrow 0$ by the replacement of $\cos v_i \sim 1 - \frac{1}{2}v_i^2$. This matrix model for torus knot $T(n, m)$ may be similar to two unitary matrix model,

$$Z = \int dU dV e^{\text{tr}(A_1 U^n + A_2 V^m + B_1 U^\dagger + B_2 V^\dagger + UV)} \quad (9.19)$$

With the condition of the equal eigenvalues of U and V , $\theta_j = \phi_j$, ($e^{i\theta_j}$ and $e^{i\phi_j}$ are eigenvalues of U and V), and with HarishChandra integral, it reduces the matrix model of torus knot. Thus the p spin curves, which is represented by (8.23) is related to torus knot $T(n, m)$. The Chern-Simons matrix model for the torus knot of (9.18) is derived by the extension of the lattice gauge theory of $U(N)$ by a hyperbolic gamma function [59, 53].

We now consider HZ transform for p -spin curve. The matrix model for the p spin intersection numbers has a logarithmic potential in the formulation of the super matrix model [24]. And also two matrix model and a model for the open-intersection numbers have a logarithmic potential [18, 19]. By applying HZ transformation of (9.4) for the logarithmic term, we obtain characters or Hilbert-Poincare polynomials. For p -spin curves, one point function $u(s)$ with a logarithmic potential is given as

$$U(s) = \frac{1}{2} \oint \frac{du}{2i\pi} e^{cs^{p+1}[(u+1)^{p+1} - (u-1)^{p+1}]} \left(\frac{u+1}{u-1}\right)^N \quad (9.20)$$

The last factor becomes a logarithmic term by the exponentiation. By HZ transform, we have

$$U(s|\lambda) = \frac{1}{2(1-\lambda)} \oint \frac{du}{2i\pi} e^{cs^{p+1}[(u+1)^{p+1} - (u-1)^{p+1}]} \frac{u-1}{u - \left(\frac{1+\lambda}{1-\lambda}\right)} \quad (9.21)$$

• **example of $p = -2$ case**

In the case of $p = -2$, strong coupling expansion becomes

$$\begin{aligned} U(s) &= \frac{1}{2} \oint \frac{du}{2i\pi} e^{\frac{s}{c}(u^2-1)} \left(\frac{u+1}{u-1}\right)^N \\ &= \sum_{n=1}^{\infty} \frac{(2n-2)!}{n!s^n} \frac{1}{\prod_{l=1}^{n-1} (N^2 - l^2)} \end{aligned} \quad (9.22)$$

The coefficients of $\frac{1}{s^n}$ is related to the characters of $U(N)$ as discussed in (7.47). HZ transformation for $p = -2$ becomes ($c \rightarrow \frac{1}{2}c$ is taken here),

$$U(s|\lambda) = \frac{1}{2(1-\lambda)} \oint \frac{du}{2i\pi} \sum_n \frac{1}{n!} \left(-\frac{2c}{s} \frac{1}{u^2-1}\right)^n \frac{u-1}{u - \left(\frac{1+\lambda}{1-\lambda}\right)} \quad (9.23)$$

The Borel transform, which eliminates $n!$ factor of exponential function, becomes

$$\hat{U} = \int_0^{\infty} dz e^{-z} u\left(\frac{s}{z}|\lambda\right) = \frac{1}{2(1-\lambda)} \oint \frac{du}{2i\pi} \frac{(u+1)(u-1)^2}{(u^2-1 + \frac{2c}{s})(u - \left(\frac{1+\lambda}{1-\lambda}\right))} \quad (9.24)$$

Three poles are at $u = \pm\sqrt{1-2c/s}$, $u = (1+\lambda)/(1-\lambda)$. The sum of the residues at these poles becomes

$$\hat{U} = \frac{\lambda}{(\lambda-1)^2} \quad (9.25)$$

which is independent of s (zero mode). The contributions from three residues are cancelled for finite s . If one takes only the pole at $u = (1+\lambda)/(1-\lambda)$, it leads to the residue as

$$\hat{U} = \frac{2\lambda^2 s}{c(\lambda-1)^2[(\lambda-1)^2 + 2\lambda s/c]} \quad (9.26)$$

The strong coupling expansion for small $\frac{2c}{s}$ in (9.24) is evaluated by the

residues at $u = \pm 1$,

$$\begin{aligned}
\hat{U} &= \frac{1}{2(1-\lambda)} \oint \frac{du}{2i\pi} \frac{(u+1)(u-1)^2}{(u^2-1)} \frac{1}{u - \frac{1+\lambda}{1-\lambda}} \sum_{n=0}^{\infty} \left(-\frac{2c}{s(u^2-1)}\right)^n \\
&= \frac{c}{2s} - \frac{(\lambda-1)^2}{\lambda} \left(\frac{c}{2s}\right)^2 + \frac{(\lambda-1)^4}{\lambda^2} \left(\frac{c}{2s}\right)^3 - \dots \\
&= \frac{\lambda}{(\lambda-1)^2 + \frac{2\lambda s}{c}}
\end{aligned} \tag{9.27}$$

This result is same as the contribution of the pole at $u = \pm\sqrt{1-2c/s}$, since these poles can be expanded in the power of $\frac{2c}{s}$. Adding the contribution from HZ for the logarithmic term in (9.26) to the strong coupling expansion in (9.27), one finds

$$\hat{U} = \frac{\lambda}{(\lambda-1)^2} \tag{9.28}$$

which is same again as (9.25). Therefore to obtain the strong coupling expansion, the residue at the pole of $u = \pm\sqrt{1-2c/s}$ should be taken. The transformation of HZ and Borel transform for $U(s)$ are useful to find the relation to the characters.

It may be interesting to compare with Hilbert-Poincare polynomials of the orbifold C^2/Z_N , P_{C^2/Z_N} . Since it is related to $y_1 y_2 = y_3^N$ (A_{N-1} type singularity), and the degree of y_1, y_2 is N , and the degree of y_3 is 2. These ideals give the Hilbert-Poincare polynomials. Thus we have

$$P_{C^2/Z_N} = \frac{1-t^{2N}}{(1-t^N)^2(1-t^2)} = \frac{1+t^N}{(1-t^N)(1-t^2)} \tag{9.29}$$

which is equal to

$$P_{C^2/Z_N} = \frac{1}{N} \sum_{l=0}^{N-1} \frac{1}{(1-q^l t)(1-q^{-l} t)} = \frac{1}{N} \sum_{l=0}^{N-1} \frac{1}{1-(q^l + q^{-l})t + t^2} \tag{9.30}$$

where $q = e^{2i\pi/N}$ (root of unity).

It may be also interesting to compare with the character of $\beta - \gamma$ system. The character $\chi(q, y)$ for $\beta - \gamma$ system of $su(2)_{-\frac{1}{2}}$ ($p = \frac{3}{2}, k = -\frac{1}{2}$) is written as [56]

$$\chi(q, t) = q^{\frac{1}{24}} q^{-\frac{1}{8}(1-(1-j)^2)} \prod_n \frac{1}{1+tq^{n+\frac{1}{2}(1-j)}} \frac{1}{1+t^{-1}q^{n+\frac{1}{2}(1+j)}} \tag{9.31}$$

where $j = 1$ for Ramond sector and $j = 0$ for Neveu-Schwarz sector. For Ramond sector ($j = 1$), $P_\gamma P_\beta$ becomes

$$P_\gamma P_\beta = \prod_{n=1} \frac{1}{1-tq^n} \frac{1}{1-t^{-1}q^{n+1}} \tag{9.32}$$

where P are generating functions for the partition of positive integer n in k pairs, $p(n, k)$; $P_\gamma = \sum p(n, k) q^n t^k = \prod (1-tq^n)^{-1}$, and $P_\beta = \sum p(n, k) q^n (q/t)^k = \prod (1-q^{n+1}/t)^{-1}$.

The analysis of HZ transform and Borel transform may be applied for any p . The exponent becomes in the variable of y , $u = \frac{i}{2}(y^2 - \frac{1}{y^2})$, $\exp[\frac{ics^{p+1}}{2y^{2p+2}}((y^2 - i)^{2p+2} - (y^2 + i)^{2p+2})]$.

For $p = -\frac{1}{2}$, the HZ transform for the logarithmic term is modified as

$$\sum_{N=0}^{\infty} \lambda^{2N} \frac{(y^2 - i)^{2N}}{(y^2 + i)^{2N}} = \frac{1}{1 - \lambda^2 \left(\frac{y^2 - i}{y^2 + i}\right)^2} \quad (9.33)$$

$$U(s|\lambda) = \frac{i}{2} \oint \frac{dy}{2i\pi} \left(y + \frac{1}{y^3}\right) e^{cs^{1/2} \frac{1}{y}} \frac{1}{1 - \lambda^2 \left(\frac{y^2 - i}{y^2 + i}\right)^2} \quad (9.34)$$

The Borel transform $u(st^2|\lambda)$ becomes

$$\hat{U} = \frac{i}{2(1 - \lambda^2)} \oint \frac{dy}{2i\pi} \left(y^2 + \frac{1}{y^2}\right) \left(\frac{1}{y - cs^{1/2}}\right) \frac{(y^2 + i)^2}{(y^2 + i(\frac{1-\lambda}{1+\lambda})) (y^2 + i(\frac{1+\lambda}{1-\lambda}))} \quad (9.35)$$

Above contour integral has poles at (i) $y = cs^{1/2}$, (ii) $y = 0$, (iii) $y = \pm i^{1/2}(\lambda + 1)^{1/2}/(\lambda - 1)^{1/2}$, $y = \pm i^{1/2}(\lambda - 1)^{1/2}/(\lambda + 1)^{1/2}$. From the residue at (i) $y = cs^{1/2}$,

$$\hat{U} = -\frac{1}{2} \frac{(ic^2s + 1)(ic^2s - 1)^3}{(ic^2s)(ic^2s(\lambda - 1) + (1 + \lambda))(ic^2s(1 + \lambda) - (1 - \lambda))} \quad (9.36)$$

and from the residue at (ii) $y = 0$,

$$\hat{U} = \frac{1}{2(1 - \lambda^2)(ic^2s)} \quad (9.37)$$

From adding the (iii) residues of the poles of $y = \pm \frac{i^{1/2}(\lambda \mp 1)^{1/2}}{(\lambda \pm 1)^{1/2}}$, one obtains

$$\hat{U} = -\frac{\lambda^2}{(\lambda^2 - 1)^2} \frac{1 - \lambda^2 - (1 + \lambda^2)(ic^2s)}{[1 + \lambda + ic^2s(\lambda - 1)][\lambda - 1 + ic^2s(1 + \lambda)]} \quad (9.38)$$

The total contribution is the sum of (9.36), (9.37) and (9.38). When $ic^2s = 0$, (i) and (ii) are cancelled, and total contribution becomes $\hat{U} = \lambda^2/(\lambda^2 - 1)^2$. Note that we used λ^2 for HZ transform instead of λ , thus above result becomes same as (9.25).

For $p = -3$ with HZ transform, one obtains

$$U(s|\lambda) = \frac{1}{2(1 - \lambda)} \oint \frac{du}{2i\pi} \sum_{n=0}^{\infty} \frac{1}{n!} \left(-\frac{4cu}{s^2(u^2 - 1)^2}\right)^n \frac{u - 1}{u - (\frac{1+\lambda}{1-\lambda})} \quad (9.39)$$

With Borel transform (replacing $\frac{1}{n!}$ by one in above formula), it becomes

$$\hat{U} = \int_0^{\infty} u \left(\frac{s}{\sqrt{t}}|\lambda\right) e^{-t} = \frac{1}{2(1 - \lambda)} \oint \frac{du}{2i\pi} \left(\frac{s^2(u^2 - 1)^2}{s^2(u^2 - 1)^2 + 4cu}\right) \frac{u - 1}{u - (\frac{1+\lambda}{1-\lambda})} \quad (9.40)$$

By taking the residues at $u = \pm 1$ in this strong coupling, it becomes

$$\begin{aligned} \hat{U} &= -\frac{c}{s^2} \frac{\lambda^2 - 1}{4\lambda} \sum_{n=0}^{\infty} \left(\frac{c(\lambda^2 - 1)(\lambda - 1)^2}{4s^2\lambda^2}\right)^n \\ &= -\left(\frac{c}{s^2}\right) \frac{\lambda^2 - 1}{4\lambda} \frac{1}{1 - \frac{c}{s^2} \frac{(\lambda^2 - 1)(\lambda - 1)^2}{4\lambda^2}} \end{aligned} \quad (9.41)$$

The Borel transform \hat{u} for p spin curve is the replacement of the exponential by the geometric series. This is equivalent to the evaluation of the intersection numbers. We have

$$U(s) = \frac{1}{s} \oint \frac{du}{2i\pi} e^{cs^{p+1} \frac{1}{2(p+1)} ((u+1)^{p+1} - (u-1)^{p+1})} \quad (9.42)$$

$$\begin{aligned} \hat{U}(s) &= \oint \frac{du}{2i\pi} \frac{1}{1 - \frac{cs^{p+1}}{2(p+1)} [(u+1)^{p+1} - (u-1)^{p+1}]} \\ &= -\frac{1}{cs^{p+1}} \oint \frac{du}{2i\pi} \frac{1}{z - \frac{1}{cs^{p+1}}} \end{aligned} \quad (9.43)$$

where

$$z = \frac{1}{2(p+1)} [(u+1)^{p+1} - (u-1)^{p+1}] \quad (9.44)$$

This equation is solved recursively as

$$\begin{aligned} z &= u^p + \frac{1}{6}p(p-1)u^{p-2} + \frac{1}{5!}p(p-1)(p-2)(p-3)u^{p-4} \\ &+ \frac{1}{7!}p(p-1)(p-2)(p-3)(p-4)(p-5)u^{p-6} + O(u^{p-8}) \end{aligned} \quad (9.45)$$

$$\begin{aligned} u &= z^{\frac{1}{p}} - \frac{1}{6}(p-1)z^{-\frac{1}{p}} + \frac{1}{360}(p-1)(2p+1)(p-3)z^{-\frac{3}{p}} \\ &- \frac{(p-1)(2p+1)(p-5)(8p^2 - 16p - 13)}{7! \cdot 3^2} z^{-\frac{5}{p}} + O(z^{-\frac{7}{p}}) \end{aligned} \quad (9.46)$$

and $du = J(z)dz$, where Jacobian $J(z)$ is obtained by the derivative of above equation. The residue at $z = (cs^{p+1})^{-1}$ gives the contour integral of $\hat{u}(s)$. Thus one finds the intersection numbers, which are equal to (2.3) and (2.8). This method may be easier than the exponential integral, which is derived in the section 2. The residue is the replacement of z by $(cs^{p+1})^{-1}$; $\hat{U} = (cs^{p+1})^{-1} J(\frac{1}{cs^{p+1}})$.

$$\begin{aligned} \hat{U} &= \frac{1}{p}(cs^{p+1})^{-\frac{1}{p}} + \frac{p-1}{6p}(cs^{p+1})^{\frac{1}{p}} - \frac{1}{120p}(p-1)(2p+1)(p-3)(cs^{p+1})^{\frac{3}{p}} \\ &+ \frac{5(p-1)(2p+1)(p-5)(8p^2 - 13p - 13)}{7! \cdot 3^2 \cdot p}(cs^{p+1})^{\frac{5}{p}} + O((cs^{p+1})^{\frac{7}{p}}) \end{aligned} \quad (9.47)$$

This expression is compared to $U(s)$ in (2.3) [6], which has an expansion as

$$\begin{aligned} U(s) &= c^{-\frac{1}{p}} s^{-1-\frac{1}{p}} \Gamma(1 + \frac{1}{p}) - \frac{p-1}{6 \cdot 4} c^{\frac{1}{p}} s^{1+\frac{1}{p}} \Gamma(1 - \frac{1}{p}) \\ &+ \frac{(p-1)(2p+1)(p-3)}{5! \cdot 4^2 \cdot 3} c^{\frac{3}{p}} s^{3+\frac{3}{p}} \Gamma(1 - \frac{3}{p}) \\ &- \frac{(p-1)(2p+1)(p-5)(8p^2 - 13p - 13)}{7! \cdot 4^3 \cdot 3^2} c^{\frac{5}{p}} s^{5+\frac{5}{p}} \Gamma(1 - \frac{5}{p}) + \dots \end{aligned} \quad (9.48)$$

When one replace c by c/t , and integrate \hat{U} by multiplying e^{-t} , one obtains $U(s)$ from \hat{U} from the relations of (9.42) and (9.43).

For instance, $p = 3$ case,

$$\begin{aligned} U(s) &= \frac{1}{2s} \oint \frac{du}{2i\pi} e^{-cs^{p+1}[(u+1)^{p+1} - (u-1)^{p+1}]} \\ &= \frac{1}{s(3s)(1/3)} Ai(\zeta) \end{aligned} \quad (9.49)$$

where $Ai(\zeta)$ is Airy function and $\zeta = -(4 \cdot 3^{1/3})^{-1} s^{8/3}$. By the replacement of the exponential as

$$\begin{aligned} \hat{U} &= \int_0^\infty dt e^{-t} \oint \frac{du}{2i\pi} e^{s^4 t (u^3 + u)} \\ &= \oint \frac{du}{2i\pi} \frac{1}{1 - cs^4(u^3 + u)} \\ &= \oint \frac{dz}{2i\pi} \frac{J(z)}{1 - cs^4 z} \end{aligned} \quad (9.50)$$

where $u^3 + u = z$, which is expanded in the series of $u = z^{1/3} - \frac{1}{3}z^{-1/3} + O(z^{-5/3})$. The measure is $J(z) = \frac{1}{3}z^{-2/3} + \frac{1}{9}z^{-4/3} + O(z^{-8/3})$. By the residue, one obtain

$$\hat{U} = -\frac{1}{3}(cs^4)^{-1/3} - \frac{1}{9}(cs^4)^{1/3} + \dots \quad (9.51)$$

which is consistent with Airy function of (9.49) and leads to the intersection numbers.

With a logarithmic potential, it is written as by HZ transform,

$$\begin{aligned} \hat{U} &= \frac{1}{1-\lambda} \oint \frac{du}{2i\pi} \frac{1}{1 - cs^4(u^3 + u)} \frac{u-1}{u - (\frac{\lambda+1}{\lambda-1})} \\ &= \frac{2\lambda(1-\lambda)}{(1-\lambda)^3 - 2cs^4(1+\lambda)(1+\lambda^2)} \end{aligned} \quad (9.52)$$

where the pole at $u = (1+\lambda)/(1-\lambda)$ is considered.

Since the representation of Borel transform \hat{U} seems easier than the previous evaluations in [6] as shown in (9.46) of this section, it is interesting to apply this method for the multi-point correlation functions and to more general situations of the singularities.

Two point function of p spin curve, $U(s_1, s_2)$, is defined as

$$U(s_1, s_2) = - \oint \frac{du_1 du_2}{(2i\pi)^2} \frac{1}{(u_1 - u_2)^2 - \frac{1}{4}(s_1 + s_2)^2} e^{\sum_{i=1}^2 \frac{cs_i^{p+1}}{2(p+1)} [(u_i+1)^{p+1} - (u_i-1)^{p+1}]} \quad (9.53)$$

Applying the Borel transformation, it becomes

$$\hat{U}(s_1, s_2) = - \oint \frac{du_1 du_2}{(2i\pi)^2} \frac{1}{(u_1 - u_2)^2 - \frac{1}{4}(s_1 + s_2)^2} \frac{1}{(cs_1^{p+1} z_1 - 1)(cs_2^{p+1} z_2 - 1)} \quad (9.54)$$

where u_i is expressed by z_i as (9.46). From the residue at $z_i = (cs_i^{p+1})^{-1}$, $\hat{U}(s_1, s_2)$ is evaluated.

For a simple case, $p = -\frac{1}{2}$, one finds $u_i = 1/z_i^2 + z_i^2/4$ from (9.46). One point function $U(s)$ is

$$U(s) = \oint \frac{du}{2i\pi} (y + \frac{1}{y^3}) e^{\frac{cs}{y}} \quad (9.55)$$

Borel transform \hat{U} is

$$\begin{aligned}\hat{U} &= \oint \frac{du}{2i\pi} \left(y + \frac{1}{y^3}\right) \frac{y}{y - cs^{\frac{1}{2}}} \\ &= cs^{\frac{1}{2}}\end{aligned}\tag{9.56}$$

where the residues of $y = 0$ and $y = cs^{\frac{1}{2}}$ are considered. The residue at $y = 0$ gives $-\frac{1}{c^2s}$, and the residue at $y = cs^{1/2}$ gives $\frac{1}{c^2s} + c^2s$. However, the factor $\frac{y}{y - cs^{\frac{1}{2}}}$ should be expanded as $1 + \frac{cs^{\frac{1}{2}}}{y} + (\frac{cs^{\frac{1}{2}}}{y})^2 + \dots$, and take the residue at $y = 0$. Then we have the result of (9.56).

The Borel transform of two point function for $p = -\frac{1}{2}$ becomes from (5.5)

$$\begin{aligned}\hat{U}(s_1, s_2) &= -16s_1s_2 \oint \frac{dy_1 dy_2}{(2i\pi)^2} \left(y_1 + \frac{1}{y_1^3}\right) \left(y_2 + \frac{1}{y_2^3}\right) \frac{1}{\left(1 - \left(\frac{cs_1^{\frac{1}{2}}}{y_1} + \frac{cs_2^{\frac{1}{2}}}{y_2}\right)\right)} \\ &\times \frac{1}{\left(s_1\left(y_1^2 - \frac{1}{y_1^2}\right) - s_2\left(y_2^2 - \frac{1}{y_2^2}\right)\right)^2 + 4(s_1 + s_2)^2}\end{aligned}\tag{9.57}$$

The term of $\left(1 - \left(\frac{cs_1^{\frac{1}{2}}}{y_1} + \frac{cs_2^{\frac{1}{2}}}{y_2}\right)\right)^{-1}$ should be expanded for small c . Then taking the residues at $y_1 = 0$ and y_2 . The order of the taking the residues gives a different results. So the first y_1 and the second y_2 , and the reverse order should be summed. (or simply add the term of $s_1 \leftrightarrow s_2$). One then finds

$$\begin{aligned}\hat{U} &= -24c^4(s_1^2 + s_2^2) - 112c^8(2s_1^2s_2^2 + 8(s_1s_2^3 + s_1^3s_2) + 12(s_1^4 + s_2^4)) \\ &- 264c^{12}(s_1^4s_2^2 + s_1^2s_2^4 + 48s_1^3s_2^3 + 161(s_1^2s_2^4 + s_1^4s_2^2) + 384(s_1s_2^5 + s_5s_2) \\ &+ 288(s_1^6 + s_2^6)) + O(c^{16})\end{aligned}\tag{9.58}$$

We have evaluated the two point function $U(s_1, s_2)$ in (7.38). \hat{U} and $U(s_1, s_2)$ are different only by a factor $n!$ for the term of c^n due to the expansion of the exponential term. For the two point function, the pole at $1 - cs_1^{1/2}/y_1 - cs_2^{1/2}/y_2 = 0$, which appears in Borel transformation, does not give the expression of (9.58).

10 Summary and discussions

In this article, we generalized the results of the correlation functions of p spin curve in [8, 9]. The multiple correlation functions are studied as examples for various values of p . For the positive integer p , we derived the intersection for such examples, which agree with the previous reported values based on the Gelfand Dikii equations [10, 11]. We use the explicit integral representation of the n -point correlation function for a fixed number of the marked points. This integral representation enables us to continue the integer p to the non-positive integer p , like $p = \frac{1}{2}$, $p = -\frac{1}{2}$ and $p = -2$ for which we evaluated n point functions explicitly as discussed in I and II [8, 9].

Since the central charge $C = 2 - \frac{6}{p}$, the case $p = -\frac{1}{2}$ gives $C = 14$ for instance, which is larger than one. Such extension is interesting from the conformal field theory, since it goes over a barrier at $C = 1$. Recently the CFT for $C > 1$ has attracted interest for the case $1 < C < 26$, which is related to 2d gravity coupled to matter field of the central charge C [44], and in this region it is expected the

behavior like a branched polymer, which is related to quantum Liouville field [45]. The area of $1 < C$ for $p < 0$ is interesting in the relation to gauge theory as we discussed in chapter 9. The p spin curve is related to the singularity theory of $f = x^p + y^2$, noting that the torus knot $T_{p,q}$ is given by $f = x^p + y^q$. The weight ω is $\{\omega_i\} = \{\frac{1}{p}, \frac{1}{q}\}$. The central charge $\hat{c} = \sum_{i=1}^n (1 - 2\omega_i)$ in the elliptic genus, which gives a central charge $1 - \frac{2}{p}$. This factor of central charge appears in Riemann-Roch equation in (2.6). For the strong coupling expansion of $p < 0$, the relation to knot polynomials and Laplace-Borel transform is discussed in this article. The non-torus knots has no factorization [51] and its representation by a matrix model will be an interesting subject in a future.

The D_l type singularity is investigated, in the new representation of the contour integrals by the variables y_i for m -point correlation functions of a matrix model. The difference between A_l and D_l types is characterized by the factor, which provides the different measure of the contour integrals. We have shown that D_l type is obtained by the logarithmic term, which is based on the supersymmetric random matrices [24]. The other type singularity like E_6 type [17] will be in a future work.

Acknowledgement

The author thank Edouard Brézin for the encouragement and joint works, which appeared in I, II in the previous articles for the p spin curves [8, 9]. He also thank Andreani Petrou for the discussions of knot polynomials in OIST. This work is supported by JSPS Kakenhi 19H01813.

References

- [1] E. Witten, Two-dimensional gravity and intersection theory on moduli space, Surveys in differential geometry 1 (1991) 243.
- [2] R. Dijkgraaf, H. Verlinde and E. Verlinde, Topological strings in $d < 1$, Nucl. Phys. B352 (1991) 59.
- [3] I. Gelfand and L. Dikii, Fractional powers of operators and hamiltonian systems, Functional Anal. Appl.10 (1977), 259.
- [4] E. Witten, Algebraic geometry associated with matrix models of two dimensional gravity, in Topological Methods in Modern Mathematics, Publish or Perish, Houston, 1993, 235.
- [5] E. Brézin and S. Hikami, Intersection numbers of Riemann surfaces from Gaussian matrix models, JHEP10 (2007) 096.
- [6] E. Brézin and S. Hikami, Computing topological invariants with one and two-matrix models, JHEP 04 (2009)110.
- [7] E. Brézin and S. Hikami, Random matrix theory with an external source, SpringerBriefs in Mathematical Physics Vol. 19 (2016), Springer.
- [8] E. Brézin and S. Hikami, Punctures and p-spin curves from matrix models, J. Stat. Phys. 180 (2020) 1031
- [9] S. Hikami and E. Brézin, Punctures and p-spin curves from matrix models II, J. Stat. Phys. 183 (2021) 36.

- [10] K. Liu and H. Xu, Decendent integrals and tautological rings of moduli spaces of curves, *Geometry and Analysis Vol. 2, Adv. Lect. Math.* 18 (2010) p.137. arXiv: 0912.0584.
- [11] K. Liu, R. Vakil and H. Xu, Formal pseudodifferential operators and Witten's r -spin numbers, *J. Reine Angew. Math.* 728 (2017), 1.
- [12] J. Zhou, Solution of W-constraints for r -spin intersection numbers, arXiv:1305.6991.
- [13] X-M. Ding, Y. Li and L. Meng, From r -spin intersection numbers to Hodge integrals, *JHEP01* (2016), 015.
- [14] E. Brézin and S. Hikami, Intersection numbers from the antisymmetric Gaussian matrix model, *JHEP07(2008)050*.
- [15] M. Bertola, B. Dubrovin and D. Yang, Correlation functions of the KdV hierarchy and applications to intersection numbers over $\bar{M}_{g,n}$. *Physica D: Nonlinear Phenomena*, 327 (2016) 30, arXiv: 1504.06452.
- [16] M. Bertola, B. Dubrovin and D. Yang, Simple Lie algebras and topological ODEs, *IMRN* 2018 (5) (2018) 1368. arXiv:1508.03750.
- [17] B. Dubrovin, D. Yang and D. Zagier, Geometry and arithmetic of integrable hierarchies of KdV type. I. Integrality, arXiv: 2101.10924.
- [18] E. Brézin and S. Hikami, On an Airy matrix model with a logarithmic potential, *J. Phys. A: Math. Theor.* 45 (2012) 045203.
- [19] E. Brézin and S. Hikami, Random matrix, singularities and open/close intersection numbers, *J. Phys. A: Mathematical and Theoretical*, 48 (2015) 475201.
- [20] A. N. Varchenko, Newton polyhedra and estimation of oscillating integrals, *Functional Analysis and its applications.* 10 (3) (1976) 175.
- [21] R. Pandharipande, J. R. Solomon, R.J. Tessler, Intersection theory on moduli of desks, open KdV and Virasoro. arXiv:1409.2191.
- [22] M. Bertola and G. Ruzza, The Kontsevich-Penner matrix integral, isomonodromic tau functions and open intersection numbers, *Ann. Henri Poincaré* 20 (2019) 393.
- [23] B. Safnuk, Combinatorial models for moduli spaces of open Riemann surfaces (2016). arXiv:1609.07226v2
- [24] E. Brézin and S. Hikami, Random supermatrices with an external source, *JHEP* 08 (2018) 086.
- [25] R. Dijkgraaf and E. Witten, Developments in topological gravity, arXiv: 1804.03275.
- [26] A. Buryak, E. Clader and R. J. Tessler, Open r -spin theory and the Gelfand-Dickey wave function, arXiv: 1809.02536.
- [27] A. Alexandrov, A. Buryak R.J.Tessler, Refined open intersection numbers and Kontsevich-Penner matrix model, *JHEP* 03 (2017) 123.

- [28] A.B. Zamolodchikov and V.A. Fateev, Nonlocal (parafermion) currents in two-dimensional conformal quantum field theory and self-dual critical points in Z_N -symmetric statistical systems, *Sov. Phys. JETP* 62 (1985) 215.
- [29] A.I.B. Zamolodchikov, Conformal scalar field on the hyperelliptic curve and critical Ashkin-Teller multipoint correlation functions, *Nucl. Phys. B* 285 [FS19] (1987) 481.
- [30] S.A. Apikyan and A. B. Zamolodchikov, Conformal blocks, related to conformally invariant Ramond states of a free scalar field. *Sov. Phys. JETP* 65 (1987) 19. *Zh. Eksp. Teor. Fiz.* 92 (1987) 34
- [31] E. Brézin and S. Hikami, Duality and replicas for a unitary matrix model, *JHEP*07 (2010) 067.
- [32] E. Brézin and S. Hikami, The intersection numbers of the p -spin curves from random matrix theory, *JHEP*02(2013), 035.
- [33] H. Fan, A. Francis, T. Jarvis and E. Merrell and Y. Ruan, Witten's D_4 integrable hierarchies conjectures, arXiv: 1008.0927.
- [34] I. P. Goulden, J. L. Harer and D. M. Jackson, A geometric parametrization for the virtual Euler characteristics of the moduli spaces of real and complex algebraic curves, *Commun. Math. Phys.* 353 (2001) 4405.
- [35] E. Garcia-Failde, R. Kramer, D. Lewanski and S. Shadrin, Half-spin tautological relation via r -spin structures, *SIGMA* 15 (2019) 080. arXiv: 1902.02742.
- [36] P. Di Francesco, F. Lesage and J.-B. Zuber, Graph rings and integrable perturbations of $N=2$ superconformal theories, *Nucl.Phys. B* 408(1993)[FS]600.
- [37] E. Brézin and D. J. Gross, The external field problem in the large N limit of QCD, *Phys. Lett. B* 97 (1980) 120.
- [38] D.J. Gross and E. Witten, Possible third-order phase transition in the large- N lattice gauge theory, *Phys. Rev. D* 21 (1980) 446.
- [39] A. Mironov, A. Morozov and G.W.Semenoff, Unitary matrix integrals in the framework of the generalized Kontsevich model, *Int. Journal Mod. Phys. A* 11 (1996) 5031.
- [40] S. Hikami and T. Maskawa, Nonlinear σ models on symmetric spaces and large N limit, *Prog. Theor. Phys.* 67 (1982) 1038.
- [41] M. Adler and P. van Moerbeke A matrix integral solution to two-dimensional W_p -gravity, *Commun. Math. Phys.* 147 (1992) 25.
- [42] E. Brézin and S. Hikami, On an Airy matrix model with a logarithmic potential, *J. Phys. A: Math. Theor.* 45 (2012) 045203.
- [43] E. Brézin and S. Hikami, Intersection theory from duality and replica, *Commun. Math. Phys.* 283 (2008) 507.

- [44] E. Brézin and S. Hikami, A naive matrix model approach to 2D quantum gravity coupled to matter of arbitrary central charge, *Phys. Lett. B.* 283 (1992) 203.
- [45] E. Gwynne, N. Holden, S. Pfeffer and G. Remy, Liouville quantum gravity with matter central charge in $(1, 25)$: a probabilistic approach, *Commun. Math. Phys.* 376 (2020) 1573.
- [46] S. Shadrin, Geometry of meromorphic functions and intersections on moduli spaces of curves. *Int. Math. Res. Not.* no 38 (2003) 2051.
- [47] T. Jarvis, T. Kimura and Vaintrob, Moduli spaces of higher spin curves and integrable hierarchies, *Compositio Math.* 126 (2001) 157.
- [48] T. Kimura and X. Liu, A genus 3 topological recursion relation, *Commun. Math. Phys.* 262 (2006), 645.
- [49] S. Samuel, $U(N)$ integrals, $1/N$, and the De Wit-'t Hooft anomalies, *J. Math. Phys.* 21 (1980) 2695.
- [50] Harer and D. Zagier, The Euler characteristic of the moduli space of curves, *Inv. Math.* 85 (1986) 457.
- [51] A. Morozov, A. Popolitov and Sh. Shakirov, Harer-Zagier formulas for knot matrix models. *Phys. Lett. B* 818 (2021) 136370. arXiv: 2102.11187.
- [52] A. Morozov and Sh. Shakirov, From Brezin-Hikami to Harer-Zagier formulas for Gaussian correlators. arXiv: 1007.4100.
- [53] I.G. Macdonald, Symmetric functions and Hall polynomials, Oxford Science Publication, (1995).
- [54] C. Faber, A Conjectural description of the tautological ring of the moduli space of curves, C. Faber and E. Looijenga (Eds.) , *Moduli of Curves and Abelian Varieties*, Vieweg Braunschweig (1999).
- [55] E. Garcia-Failde, R. Kramer, D. Lewanski and S. Shadrin, Half-spin tautological relations and Faber's proportionalities of kappa classes. *SIGMA* 15 (2019) 080.
- [56] F. Lesage, P. Methieu, J. Rasmussen, H. Saleur, The $\hat{su}(2)_{-1/2}$ WZW model and the $\beta\gamma$ system, *Nucl. Phys. B*647 [FS] (2002) 363.
- [57] M. Gisonni, T. Grava and G. Ruzza, Jacobi ensemble, Hurwitz numbers and Wilson polynomials. *Lett. Math. Phys.* 111 (2020) 67.
- [58] Y. Dolivet and M. Tierz, Chern-Simons matrix models and Stieltjes-Wigert polynomials. *J. Math. Phys.* 48 (2007) 023507.
- [59] M. Romo and M. Tierz, Unitary Chern-Simons matrix model and the Villain lattice action, *Phys. Rev. D*86 (2012) 045027.
- [60] A. Brini, M. Marino and B. Eynard, Torus knots and mirror symmetry, *Annales Henri Poincaré* 13 (2012) 1873.
- [61] E. Brézin and S. Hikami, Universal singularities at the closure of a gap in a random matrix theory. *Phys. Rev.* E57 (1998) 4140.