

# OPTIMAL CONVERGENCE RATE IN THE QUANTUM ZENO EFFECT FOR OPEN QUANTUM SYSTEMS IN INFINITE DIMENSIONS

TIM MÖBUS<sup>1,2</sup>, CAMBYSE ROUZÉ<sup>1,2</sup>

ABSTRACT. In open quantum systems, the quantum Zeno effect consists in frequent applications of a given quantum operation, e.g. a measurement, used to restrict the time evolution (due e.g. to decoherence) to states that are invariant under the quantum operation. In an abstract setting, the Zeno sequence is an alternating concatenation of a contraction operator (quantum operation) and a  $C_0$ -contraction semigroup (time evolution) on a Banach space. In this paper, we prove the optimal convergence rate of the Zeno sequence under weak boundedness assumptions, which induce a Zeno dynamics generated by an unbounded generator. For that, we derive a new Chernoff-type  $\sqrt{n}$ -lemma, which improves the convergence rate for vector-valued function converging linearly to the identity and which we believe to be of independent interest. We achieve the optimal Zeno convergence rate of order  $\frac{1}{n}$ .

## TABLE OF CONTENTS

1. Introduction	2
2. Preliminaries	3
3. Main results	5
4. Chernoff $\sqrt{n}$ -Lemma and Trotter-Kato's Product Formula	7
5. Strongly Continuous Zeno Dynamics	9
5.1. Proof of Theorem 3.1	9
5.2. Generalized Power Convergence	14
6. Uniform Power Convergence with finitely many Eigenvalues	16
6.1. Proof of Theorem 3.3	16
7. Examples	21
8. Discussion and Open Questions	22
9. References	23
Appendix A. Spectral Gap Assumption	26
Appendix B. Holomorphic Functional Calculus and Semicontinuous Spectra	26

## 1. INTRODUCTION

In its original form, the quantum Zeno effect is defined for closed finite quantum systems. Misra and Sudarshan predicted that "an unstable particle which is continuously observed to see whether it decays will never be found to decay!" [24, Abst.]. Experimentally, the Zeno effect is verified for instance in [19, 12]. In addition to its theoretical value, the quantum Zeno effect is used in error correction schemes to suppress decoherence in open quantum systems [18, 13, 3, 1, 23]. The idea is to frequently measure the quantum state and thereby force the evolution to remain within the code space. With an appropriate measurement, one can even decouple the system from its environment [10, 4] and show that appropriately encoded states can be protected from decoherence with arbitrary accuracy [6, 8].

Introduced by Misra and Sudarshan, the Zeno effect in its simplest form can be stated as follows: given a projective measurement  $P$  and a unitary time-evolution generated by a Hamiltonian  $H$  acting on a finite dimensional Hilbert space  $\mathcal{H}$  [24]:

$$(Pe^{\frac{it}{n}H})^n \longrightarrow e^{itPHP} \quad \text{as} \quad n \rightarrow \infty.$$

Since the seminal work of [24], the result was extended to the infinite dimensional setting with generalized measurements and unbounded Hamiltonians [28, 11, 19, 9]. A first approach to extend the Zeno effect to unbounded Hamiltonians was to assume that  $(\mathbb{1} - P)e^{itH}P$  is Lipschitz continuous at  $t = 0$ , a condition referred to as the *uniform asymptotic Zeno condition* [28]. This assumption is natural because it is related to the boundedness of the first moment of the Hamiltonian in the initial state and is efficiently verifiable in practice. With the works [4, 25, 1], the quantum Zeno effect was generalized to open and infinite dimensional quantum systems with general instruments and uniformly continuous time evolutions. Note that in open quantum systems, we are dealing with operators acting on the Banach space  $\mathcal{T}_1(\mathcal{H})$  of trace-class operators. More recently, Becker, Datta and Salzmann proved a quantitative bound on the convergence rate for the alternating concatenation of a  $C_0$ -contraction semigroup  $e^{\frac{t}{n}\mathcal{L}}$  (the quantum evolution) and a contraction operator  $M$  (the quantum instrument) both defined on an abstract Banach space  $\mathcal{X}$  [2]: under a condition of *uniform strong power convergence* of the power sequence  $\{M^k\}_{k \in \mathbb{N}}$  towards some projection  $P$  and boundedness of  $M\mathcal{L}$  and  $\mathcal{L}M$ ,

$$(Me^{\frac{t}{n}\mathcal{L}})^n - e^{tP\mathcal{L}P} = \mathcal{O}(n^{-\frac{2}{3}}), \quad (1)$$

where the convergence is in the strong topology. However, the optimality of (1) was left open.

*Main contributions:* In this paper, we achieve the optimal convergence rate  $\mathcal{O}(n^{-1})$  of the *Zeno sequence*. Moreover, we generalize the results of [2] in two complementary directions: Firstly, we weaken the assumption on the semigroup to the *uniform asymptotic Zeno condition* inherited from the unitary setting of [28], that is

$$\left\| (\mathbb{1} - P)e^{t\mathcal{L}}P \right\|_{\infty} = \mathcal{O}(t) \quad \text{and} \quad \left\| Pe^{t\mathcal{L}}(\mathbb{1} - P) \right\|_{\infty} = \mathcal{O}(t)$$

for  $t \rightarrow 0$  in the uniform topology. Here  $P$  is defined by a special case of the uniform power convergence:  $\|M^n - P\| \leq \delta^n$ . This leads to a possibly unbounded generator describing the Zeno dynamics since we only assume that  $P\mathcal{L}P$  is the generator of a  $C_0$ -semigroup. Second, under slightly weaker conditions as in [2], we prove that the convergence to the Zeno sequence holds in the uniform topology in addition to providing the optimal convergence rate.

In order to achieve these results, we prove a modified Chernoff  $\sqrt{n}$ -lemma in Lemma 4.2, find tight assumptions for the upper semicontinuity of parts of the spectrum of  $Me^{t\mathcal{L}}$ , and prove a quantitative convergence rate for  $\exp(nP(e^{\frac{1}{n}t\mathcal{L}} - \mathbb{1})P)P - e^{tP\mathcal{L}P}$  as  $n \rightarrow \infty$ , where  $P\mathcal{L}P$  is possibly unbounded.

*Organization of the paper:* In Section 2, we provide a short recap on bounded and unbounded operator theory. We expose our main results in Section 3. Section 4 deals with the modified Chernoff  $\sqrt{n}$ -Lemma and some of its implications as regards to Trotter-Kato's product formula. Then, we prove our main theorems under the weakest possible assumptions on the  $C_0$ -semigroup in Section 5, and under the weakest possible assumptions on  $M$  in Section 6. Our results are illustrated by two large classes of examples, namely finite and infinite dimensional quantum systems, in Section 7. Finally, we discuss some remaining open questions in Section 8.

## 2. PRELIMINARIES

Let  $(\mathcal{X}, \|\cdot\|)$  be a Banach space and  $(\mathcal{B}(\mathcal{X}), \|\cdot\|_\infty)$  be the associated space of bounded linear operators over  $\mathbb{C}$  equipped with the operator norm and the identity  $\mathbb{1} \in \mathcal{B}(\mathcal{X})$ . By a slight abuse of notation, we extend all densely defined and bounded operators by the *bounded linear extension theorem* to bounded operators on  $\mathcal{X}$  [21, Thm. 2.7-11]. A sequence  $(C_k)_{k \in \mathbb{N}} \subset \mathcal{B}(\mathcal{X})$  converges uniformly to  $C \in \mathcal{B}(\mathcal{X})$  if  $\lim_{k \rightarrow \infty} \|C_k - C\|_\infty = 0$  and strongly if  $\lim_{k \rightarrow \infty} \|C_k x - Cx\| = 0$  for all  $x \in \mathcal{X}$ . The integral over bounded vector-valued maps, e.g.  $[a, b] \rightarrow \mathcal{X}$  or  $[a, b] \rightarrow \mathcal{B}(\mathcal{X})$  with  $a < b$ , is defined by the *Bochner integral*, which satisfies the triangle inequality, is invariant under bounded linear operators, and satisfies the *fundamental theorem of calculus* if the map is continuously differentiable [17, Sec. 3.7-8, 22, Sec. A.1-2].

We define the *resolvent set* of  $C \in \mathcal{B}(\mathcal{X})$  by  $\rho(C) := \{z \in \mathbb{C} \mid (z - C) \text{ bijective}\}$  and its *spectrum* by  $\sigma(C) := \mathbb{C} \setminus \rho(C)$ . Then, we define the *resolvent* of  $C$  by  $R(z, C) := (z - C)^{-1}$  for all  $z \in \rho(C)$ . A point  $\lambda \in \sigma(C)$  is called *isolated* if there is a neighbourhood  $V \subset \mathbb{C}$  of  $\lambda$  such that  $V \cap \sigma(C) = \{\lambda\}$ . Moreover, the spectrum is separated by a curve  $\Gamma : I \rightarrow \rho(C)$  if  $\sigma(C) = \sigma_1 \cup \sigma_2$ ,  $\sigma_1, \sigma_2$  have disjoint neighbourhoods  $V_1, V_2$ , and the curve is simple, closed, rectifiable<sup>1</sup>, and encloses one of the neighbourhoods without intersecting  $V_1$  and  $V_2$  [20, Sec. III§6.4]. This definition can be generalized to a finite sum of subsets separated by curves. An example for a simple, closed, and rectifiable curve is the parametrization of the boundary of a complex ball around the origin with radius  $r$ . We denote the open ball by  $\mathbb{D}_r$  and its boundary by  $\partial\mathbb{D}_r$ .

We define the *spectral projection* with respect to a separated subset of  $\sigma(C)$  enclosed by a curve  $\Gamma$  via the *holomorphic functional calculus* [29, Thm. 2.3.1-3]

$$P = \frac{1}{2\pi i} \oint_{\Gamma} R(z, C) dz. \quad (2)$$

As the name suggests,  $P$  satisfies the projection property,  $P^2 = P$ . We denote the perpendicular projection  $\mathbb{1} - P$  by  $P^\perp$ . If  $\Gamma$  encloses the isolated one-point subset  $\{\lambda\} \subset \sigma(C)$ , then  $CP = \lambda P + N$  with the *quasinilpotent operator* [29, Thm. 2.3.5]

$$N := \frac{1}{2\pi i} \oint_{\Gamma} (z - \lambda) R(z, C) dz \in \mathcal{B}(\mathcal{X}). \quad (3)$$

A quasinilpotent operator  $N \in \mathcal{B}(\mathcal{X})$  is characterized by  $\lim_{n \rightarrow \infty} \|N^n\|_\infty^{\frac{1}{n}} = 0$ .

The time evolution in the Zeno sequence is described by a  $C_0$ -contraction semigroup [7, Def. 5.1]. A family of operators  $(T_t)_{t \geq 0} \subset \mathcal{B}(\mathcal{X})$  is called a  $C_0$ -semigroup if the family satisfies the semigroup properties, namely: (i)  $T_t T_s = T_{t+s}$  for all  $t, s \geq 0$ , (ii)  $T_0 = \mathbb{1}$ , and (iii)  $t \mapsto T_t$  is strongly continuous in 0. The generator of a  $C_0$ -semigroup is an unbounded operator defined by taking the strong derivative in 0, that is

$$\mathcal{L}x := \lim_{h \downarrow 0} \frac{T(h)x - x}{h}$$

for all  $x \in \mathcal{D}(\mathcal{L}) := \{x \in \mathcal{X} \mid t \mapsto T_t x \text{ differentiable}\}$  and is denoted by  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  (cf. [20, p. 152]). For ease of notations, we denote a  $C_0$ -semigroup by  $(e^{t\mathcal{L}})_{t \geq 0}$ , although the series representation of the exponential is not well-defined (cf. [20, Sec. IX§1]). A semigroup is called (uniformly) bounded if there exists  $c \in \mathbb{R}_{\geq 0}$  such that  $\sup_{t \geq 0} \|e^{t\mathcal{L}}\|_\infty \leq c$  and contractive if  $c = 1$ . Note that

<sup>1</sup>A simple, closed, and rectifiable curve is continuous, has finite length, joins up, and does not cross itself.

the linear combination and concatenation of two unbounded operators  $(\mathcal{K}, \mathcal{D}(\mathcal{K}))$  and  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  is defined on  $\mathcal{D}(\mathcal{K} + \mathcal{L}) = \mathcal{D}(\mathcal{K}) \cap \mathcal{D}(\mathcal{L})$  and  $\mathcal{D}(\mathcal{K}\mathcal{L}) = \mathcal{L}^{-1}(\mathcal{D}(\mathcal{K}))$  (cf. [20, Sec. III§5.1]). The following lemma summarizes some properties of  $C_0$ -semigroups which are used in this work:

**Lemma 2.1** ([7, Lem. 1.3]) *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of the  $C_0$ -semigroup  $(e^{t\mathcal{L}})_{t \geq 0}$  defined on  $\mathcal{B}(\mathcal{X})$ . Then the following properties hold:*

(1) *If  $x \in \mathcal{D}(\mathcal{L})$ , then  $e^{t\mathcal{L}}x \in \mathcal{D}(\mathcal{L})$  and*

$$\frac{\partial}{\partial t} e^{t\mathcal{L}}x = e^{t\mathcal{L}}\mathcal{L}x = \mathcal{L}e^{t\mathcal{L}}x \quad \text{for all } t \geq 0.$$

(2) *For every  $t \geq 0$  and  $x \in \mathcal{X}$ , one has*

$$\int_0^t e^{\tau\mathcal{L}}x d\tau \in \mathcal{D}(\mathcal{L}).$$

(3) *For every  $t \geq 0$ , one has*

$$\begin{aligned} e^{t\mathcal{L}}x - x &= \mathcal{L} \int_0^t e^{\tau\mathcal{L}}x d\tau \quad \text{for all } x \in \mathcal{X} \\ &= \int_0^t e^{\tau\mathcal{L}}\mathcal{L}x d\tau \quad \text{if } x \in \mathcal{D}(\mathcal{L}). \end{aligned}$$

Additionally, a general *integral formulation* for the difference of two semigroups is discussed in the following lemma (cf. [7, Cor. III.1.7]<sup>2</sup>).

**Lemma 2.2** (Integral equation for semigroups) *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  and  $(\mathcal{K}, \mathcal{D}(\mathcal{K}))$  be the generators of two  $C_0$ -semigroups on  $\mathcal{B}(\mathcal{X})$  and  $t \geq 0$ . Assume  $\mathcal{D}(\mathcal{L}) \subset \mathcal{D}(\mathcal{K})$  and  $[0, t] \ni s \mapsto e^{s\mathcal{K}}(\mathcal{K} - \mathcal{L})e^{(t-s)\mathcal{L}}x$  is continuous for all  $x \in \mathcal{D}(\mathcal{L})$ . Then,*

$$e^{t\mathcal{K}}x - e^{t\mathcal{L}}x = \int_0^t e^{s\mathcal{K}}(\mathcal{K} - \mathcal{L})e^{(t-s)\mathcal{L}}x ds$$

for all  $x \in \mathcal{D}(\mathcal{L})$ . If the semigroup  $(e^{t\mathcal{L}})_{t \geq 0}$  is a quasi-contraction, i.e. there exists a  $w \geq 0$  so that  $\|e^{t\mathcal{L}}\|_\infty \leq e^{tw}$ , and  $\mathcal{K} = \mathcal{L} + A$  with  $A \in \mathcal{B}(\mathcal{X})$ , then  $(\mathcal{L} + A, \mathcal{D}(\mathcal{L}))$  is a generator of a  $C_0$ -semigroup and

$$\|e^{t(\mathcal{L}+A)} - e^{t\mathcal{L}}\|_\infty \leq e^{tw}(e^{t\|A\|_\infty} - 1).$$

*Proof.* In the first part, we follow the proof of Lemma B.16 and Corollary III.1.7 in [7]. By definition, the vector-valued maps  $[0, t] \ni s \mapsto e^{s\mathcal{K}}x$  and  $[0, t] \ni s \mapsto e^{(t-s)\mathcal{L}}x$  are continuously differentiable for all  $x \in \mathcal{D}(\mathcal{L})$  so that

$$\begin{aligned} \frac{1}{h} \left( e^{(s_0+h)\mathcal{K}}e^{(t-s_0-h)\mathcal{L}}x - e^{s_0\mathcal{K}}e^{(t-s_0)\mathcal{L}}x \right) &= e^{(s_0+h)\mathcal{K}} \frac{e^{(t-s_0-h)\mathcal{L}} - e^{(t-s_0)\mathcal{L}}}{h} x \\ &\quad + \frac{e^{(s_0+h)\mathcal{K}} - e^{s_0\mathcal{K}}}{h} e^{(t-s_0)\mathcal{L}}x \end{aligned}$$

for all  $h > 0$  and  $x \in \mathcal{D}(\mathcal{L})$ . The second term converges to  $\mathcal{K}e^{s_0\mathcal{K}}e^{(t-s_0)\mathcal{L}}x$  for  $h \downarrow 0$ . Moreover,

$$\left\{ \frac{e^{(t-s_0-h)\mathcal{L}} - e^{(t-s_0)\mathcal{L}}}{h} x \mid h \in [0, 1] \right\}$$

is relatively compact in  $\mathcal{X}$  by the continuous differentiability of  $(e^{(t-s)\mathcal{L}})_{s \geq 0}$ , and the *uniform boundedness principle* shows that  $\{e^{(s_0+h)\mathcal{L}} \mid h \in [0, 1]\}$  is bounded [7, Lem. 5.2]. Then, the topologies of pointwise convergence and uniform convergence on relatively compact sets coincide [7, Prop. A.3] so that

$$\lim_{h \downarrow 0} e^{(s_0+h)\mathcal{K}} \frac{e^{(t-s_0-h)\mathcal{L}} - e^{(t-s_0)\mathcal{L}}}{h} x = e^{s_0\mathcal{K}}\mathcal{L}e^{(t-s_0)\mathcal{L}}x$$

<sup>2</sup>The abbreviation Cor. III.1.7 refers to Corollary 1.7 in Chapter III

and

$$\frac{\partial}{\partial s} e^{s\mathcal{K}} e^{(t-s)\mathcal{L}} x = e^{s\mathcal{K}} (\mathcal{K} - \mathcal{L}) e^{(t-s)\mathcal{L}} x$$

for all  $x \in \mathcal{D}(\mathcal{L})$ . Since  $[0, t] \ni s \mapsto e^{s\mathcal{K}} e^{(t-s)\mathcal{L}} x$  is continuously differentiable, the fundamental theorem of calculus shows for all  $x \in \mathcal{D}(\mathcal{L})$

$$e^{t\mathcal{K}} x - e^{t\mathcal{L}} x = \int_0^t e^{s\mathcal{K}} (\mathcal{K} - \mathcal{L}) e^{(t-s)\mathcal{L}} x ds.$$

In the second part,  $\mathcal{K} = \mathcal{L} + A$  where  $A \in \mathcal{B}(\mathcal{X})$ . By [17, Thm. 13.2.1], the unbounded operator  $(\mathcal{L} + A, \mathcal{D}(\mathcal{L}))$  is a generator of a  $C_0$ -semigroup and  $[0, t] \ni s \mapsto e^{s\mathcal{K}} A e^{(t-s)\mathcal{L}} x$  is continuous by [7, Lem. B.15]. Then

$$e^{t(\mathcal{L}+A)} x - e^{t\mathcal{L}} x = \int_0^t e^{s(\mathcal{L}+A)} A e^{(t-s)\mathcal{L}} x ds$$

for all  $x \in \mathcal{D}(\mathcal{L})$ . The equality can be extended to  $x \in \mathcal{X}$  since all operators are bounded and  $\mathcal{D}(\mathcal{L})$  is dense in  $\mathcal{X}$ . If  $(e^{t\mathcal{L}})_{t \geq 0}$  is a quasi-contraction, then

$$\|e^{t(\mathcal{L}+A)}\|_\infty \leq e^{t(w+\|A\|_\infty)}$$

by [17, p. 390], which finishes the proof because of

$$\begin{aligned} \|e^{t(\mathcal{L}+A)} - e^{t\mathcal{L}}\|_\infty &\leq \int_0^t \|e^{s(\mathcal{L}+A)} A e^{(t-s)\mathcal{L}}\|_\infty ds \\ &\leq e^{tw} \|A\|_\infty \int_0^t e^{s\|A\|_\infty} ds \\ &= e^{tw} (e^{t\|A\|_\infty} - 1). \end{aligned}$$

□

### 3. MAIN RESULTS

In this section, we list the main results achieved in the paper. Since the quantum Zeno effect consists of a contraction operator  $M \in \mathcal{B}(\mathcal{X})$  (e.g. measurement) and a  $C_0$ -semigroup (e.g. quantum time evolution), the assumptions on the contraction operator and on the  $C_0$ -semigroup influence each other. We first start with weak assumptions on the semigroup and stronger assumption on the contraction operator:

**Theorem 3.1** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$ ,  $M \in \mathcal{B}(\mathcal{X})$  a contraction, and  $P$  projection satisfying*

$$\|M^n - P\|_\infty \leq \delta^n \tag{4}$$

for a  $\delta \in (0, 1)$  and all  $n \in \mathbb{N}$ . Moreover, assume there is  $b \geq 0$  so that for all  $t \geq 0$

$$\|P e^{t\mathcal{L}} P^\perp\|_\infty \leq tb \quad \text{and} \quad \|P^\perp e^{t\mathcal{L}} P\|_\infty \leq tb. \tag{5}$$

If  $(P\mathcal{L}P, \mathcal{D}(\mathcal{L}P))$  is the generator of a  $C_0$ -semigroup, then for any  $t \geq 0$

$$\left\| \left( M e^{\frac{t}{n}\mathcal{L}} \right)^n x - e^{tP\mathcal{L}P} P x \right\| \leq \frac{c}{n} \left( \|x\| + \|\mathcal{L}P x\| + \|(\mathcal{L}P)^2 x\| \right) + \delta^n \|x\|$$

for a constant  $c > 0$  and all  $x \in \mathcal{D}((\mathcal{L}P)^2)$ .

Note that the assumption on  $M$  is a special case of the so-called *uniform power convergence* assumption (q.v. Eq. (9)). Moreover, the assumption on the  $C_0$ -semigroup is a generalization of the uniform *asymptotic Zeno condition* which implies the convergent for unitary evolution and a projective measurement [28, Sec. 3.1]. The following example confirms the optimality of the achieved convergence rate.

*Example 1.* Let  $\mathcal{X}$  be a Banach space,  $\mathcal{L}$  be a bounded generator of a contraction semigroup on  $\mathcal{B}(\mathcal{X})$  such that  $\mathcal{L}^2 = 0$ , and  $M \in \mathcal{B}(\mathcal{X})$  be a contraction which satisfies (4) for some projection  $P$  and  $\delta \in (0, 1)$  (q.v. Sec. 3). Moreover, assume that  $M^\perp := P^\perp M$  commutes with  $e^{t\mathcal{L}}$  and  $\mathcal{L}P = 0$ . Then,

$$\begin{aligned} \left(Me^{\frac{1}{n}\mathcal{L}}\right)^n &= \left(P^\perp Me^{\frac{1}{n}\mathcal{L}} + Pe^{\frac{1}{n}\mathcal{L}}\right)^n \\ &= \left(M^\perp\right)^n e^\mathcal{L} + \left(P + \frac{1}{n}P\mathcal{L}\right)^n \\ &= \left(M^\perp\right)^n e^\mathcal{L} + Pe^{P\mathcal{L}P} + \frac{1}{n}P\mathcal{L}. \end{aligned} \quad (6)$$

Let  $\{|1\rangle, |2\rangle, |3\rangle\}$  be an orthonormal basis of  $\mathbb{R}^3$  and  $\delta \in (0, 1)$ . An explicit non-trivial example with the property that  $P$  does not commute with  $\mathcal{L}$  can be constructed on a qutrit system:

$$\mathcal{L} := |1\rangle\langle 2| \quad \text{and} \quad M := |1\rangle\langle 1| + \delta|3\rangle\langle 3|.$$

Then,  $\mathcal{L}$ ,  $P = |1\rangle\langle 1|$ , and  $M^\perp = \delta|3\rangle\langle 3|$  satisfies the assumptions. Equation (6) shows

$$\left(Me^{\frac{1}{n}\mathcal{L}}\right)^n = \underbrace{\delta^n|3\rangle\langle 3|}_{(P^\perp M)^n e^\mathcal{L}} + \underbrace{|1\rangle\langle 1|}_{Pe^{P\mathcal{L}P}} + \frac{1}{n}\underbrace{|1\rangle\langle 2|}_{P\mathcal{L}}$$

so that

$$\left\| \left(Me^{\frac{1}{n}\mathcal{L}}\right)^n - Pe^{P\mathcal{L}P} \right\|_\infty = \max\left\{\frac{1}{n}, \delta^n\right\},$$

which shows the optimality of our convergence rate in Theorem 3.1.

Our next goal is to weaken the assumption on the contraction  $M$ .

**Proposition 3.2** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$  and  $M \in \mathcal{B}(\mathcal{X})$  a contraction such that*

$$\|M^n - P\|_\infty \leq \tilde{c}\delta^n \quad (7)$$

for some projection  $P$ ,  $\delta \in (0, 1)$  and  $\tilde{c} \geq 0$ . Moreover, we assume that there is  $b \geq 0$  so that

$$\|Pe^{t\mathcal{L}}P^\perp\|_\infty \leq tb, \quad \|M^\perp e^{t\mathcal{L}} - M^\perp\|_\infty \leq tb, \quad \text{and} \quad \|P^\perp e^{t\mathcal{L}}P\|_\infty \leq tb. \quad (8)$$

If  $(P\mathcal{L}P, \mathcal{D}(\mathcal{L}P))$  generates a  $C_0$ -semigroup, then there is  $\epsilon > 0$  such that for all  $t \geq 0$ ,  $n \in \mathbb{N}$  satisfying  $t \in [0, n\epsilon]$ ,  $\tilde{\delta} \in (\delta, 1)$ , and  $x \in \mathcal{D}((\mathcal{L}P)^2)$

$$\left\| \left(Me^{\frac{t}{n}\mathcal{L}}\right)^n x - e^{tP\mathcal{L}P}Px \right\| \leq \frac{c_1}{n} \left( \|x\| + \|\mathcal{L}Px\| + \|(\mathcal{L}P)^2x\| \right) + c_2\tilde{\delta}^n\|x\|,$$

for some constants  $c_1, c_2 \geq 0$ .

Finally, we extend the assumption on  $M$  (q.v. Eq. (4), (7)) to the *uniform power convergence* introduced in [2]. Let  $\{P_j, \lambda_j\}_{j=1}^J$  be a set of orthogonal projections and associated eigenvalues on the unit circle  $\partial\mathbb{D}_1$ . Then,  $M$  is called uniformly power convergent with rate  $\delta \in (0, 1)$  if  $M^n - \sum_{j=1}^J \lambda_j^n P_j = \mathcal{O}(\delta^n)$  uniformly for  $n \rightarrow \infty$ . To prove Theorem 3.3, we assume that  $M\mathcal{L}$  and  $\mathcal{L}P_\Sigma$  with  $P_\Sigma := \sum_{j=1}^J P_j$  are densely defined and bounded (cf. [2]):

**Theorem 3.3** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$  and  $M \in \mathcal{B}(\mathcal{X})$  a contraction satisfying the following uniform power convergence: there is  $\tilde{c} > 0$  so that*

$$\left\| M^n - \sum_{j=1}^J \lambda_j^n P_j \right\|_\infty \leq \tilde{c}\delta^n \quad (9)$$

for a set of orthogonal projections  $\{P_j\}_{j=1}^J$ , eigenvalues  $\{\lambda_j\}_{j=1}^J \subset \partial\mathbb{D}_1$ , and a rate  $\delta \in (0, 1)$ . For  $P_\Sigma := \sum_{j=1}^J P_j$ , we assume that  $M\mathcal{L}$  and  $\mathcal{L}P_\Sigma$  are densely defined and bounded by  $b \geq 0$ .

Then, there is an  $\epsilon > 0$  such that for all  $n \in \mathbb{N}$ ,  $t \geq 0$  satisfying  $t \in [0, n\epsilon]$

$$\left\| \left( M e^{\frac{t}{n} \mathcal{L}} \right)^n - \sum_{j=1}^J \lambda_j^n e^{t P_j \mathcal{L} P_j} P_j \right\|_{\infty} \leq \frac{c_1 J}{n} + c_2 \delta^n,$$

for some constants  $c_1, c_2 \geq 0$ .

In comparison to Theorem 3 in [2], Theorem 3.3 achieves the optimal convergence rate and is formulated in the uniform topology under slightly weaker assumptions on the generator.

*Remark.* A natural way to weaken the above assumption is to assume that the power converges in the strong topology (cf. [2, Thm. 2]).

#### 4. CHERNOFF $\sqrt{n}$ -LEMMA AND TROTTER-KATO'S PRODUCT FORMULA

In previous works [25, 2], Chernoff's  $\sqrt{n}$ -Lemma [5, Lem. 2], which we restate here, is used as a proof technique to approximate the Zeno product by a semigroup (q.v. Eq. (11)).

**Lemma 4.1** (Chernoff  $\sqrt{n}$  Lemma) *Let  $C \in \mathcal{B}(\mathcal{X})$  be a contraction. Then,  $(e^{t(C-1)})_{t \geq 0}$  is a uniformly continuous contraction semigroup and for all  $x \in \mathcal{X}$*

$$\|C^n x - e^{n(C-1)} x\| \leq \sqrt{n} \|(C - \mathbb{1})x\|.$$

*Remark.* In Lemma 2.1 in [31], the dependence on  $n$  is improved to  $n^{\frac{1}{3}}$ . This is crucial in the proof of the convergence rate in [2, Lem. 5.4-5.5]. Unfortunately, we found an inconsistency in the proof of [31, Lem. 2.1].

In the case of the quantum Zeno effect, the contraction  $C$  is a vector-valued map satisfying  $\|(C(\frac{1}{n}) - \mathbb{1})x\| = \mathcal{O}(n^{-1} \|\mathcal{L}x\|)$  for all  $x \in \mathcal{D}(\mathcal{L})$ . Then, Chernoff's  $\sqrt{n}$ -Lemma sets a convergence rate  $n^{-\frac{1}{2}}$ . Here, we prove an alternative bound, which allows us to get the optimal rate for the quantum Zeno effect

**Lemma 4.2** (Modified Chernoff Lemma) *Let  $C \in \mathcal{B}(\mathcal{X})$  be a contraction and  $n \in \mathbb{N}$ . Then,  $(e^{t(C-1)})_{t \geq 0}$  is a contraction semigroup and for all  $x \in \mathcal{X}$*

$$\|(C^n - e^{n(C-1)})x\| \leq \frac{n}{2} \|(C - \mathbb{1})^2 x\|.$$

*Remark.* At first glance, this seems to be worse than the original and improved Chernoff  $\sqrt{n}$ -Lemmas in [5, 31]. However, if  $C$  is a vector-valued map satisfying  $\|C(\frac{1}{n}) - \mathbb{1}\|_{\infty} = \mathcal{O}(n^{-1})$ , then the modified Chernoff lemma gives

$$\left\| C\left(\frac{1}{n}\right)^n - e^{n(C\left(\frac{1}{n}\right) - \mathbb{1})} \right\|_{\infty} \leq \frac{n}{2} \left\| C\left(\frac{1}{n}\right) - \mathbb{1} \right\|_{\infty}^2 = \mathcal{O}(n^{-1})$$

which proves the optimal convergence rate of the quantum Zeno effect under the assumptions that the contraction  $M$  satisfies the uniform power convergence and that the semigroup  $(e^{t\mathcal{L}})_{t \geq 0}$  is uniformly continuous (q.v. Lem. 5.5).

*Proof of Lemma 4.2.* Similarly to Chernoff's proof [5, Lem. 2],  $(e^{t(C-1)})_{t \geq 0}$  is a contraction semigroup. We define  $C_t := (1-t)\mathbb{1} + tC = \mathbb{1} + t(C - \mathbb{1})$ , which itself is a contraction, and we use the fundamental theorem of calculus so that

$$\begin{aligned} \|(C^n - e^{n(C-1)})x\| &\leq \int_0^1 \left\| \frac{\partial}{\partial t} (C_t^n e^{(1-t)n(C-1)})x \right\| dt \\ &\leq n \int_0^1 \left\| C_t^{n-1} e^{(1-t)n(C-1)} \right\|_{\infty} \|(1-C)(\mathbb{1} - C_t)x\| dt \\ &\leq n \|(1-C)^2 x\| \int_0^1 t dt \\ &= \frac{n}{2} \|(C - \mathbb{1})^2 x\|, \end{aligned}$$

which proves the lemma.  $\square$

In [5, p. 241], Chernoff proves the convergence of Trotter's product formula by approximating the product using the Chernoff  $\sqrt{n}$ -Lemma. For bounded generators, Chernoff's proof gives a convergence rate of order  $n^{-\frac{1}{2}}$ . Following his proof and using our modified Chernoff Lemma, we achieve the well-known optimal convergence rate of order  $n^{-1}$  [16, Thm. 2.11, 26, p. 1-2]:

**Proposition 4.3** *Let  $F : \mathbb{R}_{\geq 0} \rightarrow \mathcal{B}(\mathcal{X})$  be a continuously differentiable function (in the uniform topology) satisfying  $\sup_{t \in \mathbb{R}_{\geq 0}} \|F(t)\|_{\infty} \leq 1$ . Assume that  $F(0) = \mathbb{1}$  and denote the derivative in  $t = 0$  by  $\mathcal{L} \in \mathcal{B}(\mathcal{X})$ . Then, for all  $t \geq 0$*

$$\left\| F\left(\frac{t}{n}\right)^n - e^{t\mathcal{L}} \right\|_{\infty} \leq \|n(F\left(\frac{t}{n}\right) - \mathbb{1}) - t\mathcal{L}\|_{\infty} + \frac{n}{2} \left\| F\left(\frac{t}{n}\right) - \mathbb{1} \right\|_{\infty}^2.$$

*Proof.* The case  $t = 0$  is clear. For  $t > 0$ , applying Lemma 4.2, we get

$$\begin{aligned} \left\| F\left(\frac{t}{n}\right)^n - e^{t\mathcal{L}} \right\|_{\infty} &\leq \left\| F\left(\frac{t}{n}\right)^n - e^{n(F\left(\frac{t}{n}\right) - \mathbb{1})} \right\|_{\infty} + \left\| e^{n(F\left(\frac{t}{n}\right) - \mathbb{1})} - e^{t\mathcal{L}} \right\|_{\infty} \\ &\leq \frac{n}{2} \left\| F\left(\frac{t}{n}\right) - \mathbb{1} \right\|_{\infty}^2 + \left\| e^{n(F\left(\frac{t}{n}\right) - \mathbb{1})} - e^{t\mathcal{L}} \right\|_{\infty}. \end{aligned}$$

For the second term above, we apply Lemma 2.2:

$$\begin{aligned} \left\| e^{n(F\left(\frac{t}{n}\right) - \mathbb{1})} - e^{t\mathcal{L}} \right\|_{\infty} &= \int_0^1 \left\| e^{sn(F\left(\frac{t}{n}\right) - \mathbb{1})} (n(F\left(\frac{t}{n}\right) - \mathbb{1}) - t\mathcal{L}) e^{(1-s)t\mathcal{L}} \right\|_{\infty} ds \\ &\leq \|n(F\left(\frac{t}{n}\right) - \mathbb{1}) - t\mathcal{L}\|_{\infty}, \end{aligned}$$

where in the last inequality we also used that  $\|F\left(\frac{t}{n}\right)\|_{\infty} \leq 1$  and that  $e^{sn(F\left(\frac{t}{n}\right) - \mathbb{1})}$  is a uniformly continuous contraction semigroup.  $\square$

Applying the proposition to the case of Trotter's product formula, we achieve the optimal convergence rate for bounded generators on Banach spaces [16, Thm. 2.11, 26, p. 1-2]:

**Corollary 4.4** *Let  $\mathcal{L}_1$  and  $\mathcal{L}_2$  be two bounded generators of uniformly continuous contraction semigroups. Then, for  $n \rightarrow \infty$*

$$\left\| \left( e^{\frac{1}{n}\mathcal{L}_1} e^{\frac{1}{n}\mathcal{L}_2} \right)^n - e^{\mathcal{L}_1 + \mathcal{L}_2} \right\|_{\infty} = \mathcal{O}\left(\frac{1}{n}\right).$$

*Proof.* We define  $F\left(\frac{1}{n}\right) := e^{\frac{1}{n}\mathcal{L}_1} e^{\frac{1}{n}\mathcal{L}_2}$  for which

$$\begin{aligned} F\left(\frac{1}{n}\right) - \mathbb{1} &= e^{\frac{1}{n}\mathcal{L}_1} \left( e^{\frac{1}{n}\mathcal{L}_2} - \mathbb{1} \right) + e^{\frac{1}{n}\mathcal{L}_1} - \mathbb{1} \\ &= \left( e^{\frac{1}{n}\mathcal{L}_1} - \mathbb{1} \right) \left( e^{\frac{1}{n}\mathcal{L}_2} - \mathbb{1} \right) + e^{\frac{1}{n}\mathcal{L}_2} - \mathbb{1} + e^{\frac{1}{n}\mathcal{L}_1} - \mathbb{1} \end{aligned}$$

holds. Moreover,

$$\left\| e^{\frac{1}{n}\mathcal{L}_1} - \mathbb{1} \right\|_{\infty} = \frac{1}{n} \left\| \int_0^1 e^{\frac{\tau_1}{n}\mathcal{L}_1} \mathcal{L}_1 d\tau_1 \right\|_{\infty} \leq \frac{1}{n} \|\mathcal{L}_1\|_{\infty}$$

and

$$\left\| n \left( e^{\frac{1}{n}\mathcal{L}_1} - \mathbb{1} \right) - \mathcal{L}_1 \right\|_{\infty} = \frac{1}{n} \left\| \int_0^1 \int_0^1 \tau_1 e^{\frac{\tau_1\tau_2}{n}\mathcal{L}_1} \mathcal{L}_1^2 d\tau_2 d\tau_1 \right\|_{\infty} \leq \frac{1}{2n} \|\mathcal{L}_1\|_{\infty}^2.$$

Then,

$$\left\| n \left( F\left(\frac{1}{n}\right) - \mathbb{1} \right) - \mathcal{L}_1 - \mathcal{L}_2 \right\|_{\infty} \leq \frac{1}{n} \left( \|\mathcal{L}_1\|_{\infty} \|\mathcal{L}_2\|_{\infty} + 2\|\mathcal{L}_1\|_{\infty}^2 + 2\|\mathcal{L}_2\|_{\infty}^2 \right)$$

and the statement follows from Proposition 4.3.  $\square$



## 5. STRONGLY CONTINUOUS ZENO DYNAMICS

We proceed with proving our first main result, namely Theorem 3.1, which we restate here for sake of clarity of exposition:

**Theorem 3.1** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$ ,  $M \in \mathcal{B}(\mathcal{X})$  a contraction, and  $P$  projection satisfying*

$$\|M^n - P\|_\infty \leq \delta^n \quad (4)$$

for a  $\delta \in (0, 1)$  and all  $n \in \mathbb{N}$ . Moreover, assume there is  $b \geq 0$  so that for all  $t \geq 0$

$$\|Pe^{t\mathcal{L}}P^\perp\|_\infty \leq tb \quad \text{and} \quad \|P^\perp e^{t\mathcal{L}}P\|_\infty \leq tb. \quad (5)$$

If  $(P\mathcal{L}P, \mathcal{D}(P\mathcal{L}P))$  is the generator of a  $C_0$ -semigroup, then for any  $t \geq 0$

$$\left\| \left( Me^{\frac{t}{n}\mathcal{L}} \right)^n x - e^{tP\mathcal{L}P} Px \right\| \leq \frac{c}{n} \left( \|x\| + \|\mathcal{L}Px\| + \|(\mathcal{L}P)^2 x\| \right) + \delta^n \|x\|$$

for a constant  $c > 0$  and all  $x \in \mathcal{D}((\mathcal{L}P)^2)$ .

Before proving Theorem 3.1, we consider a slight weakening of the condition on the map  $M$ :

**Corollary 5.1** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$  and  $M \in \mathcal{B}(\mathcal{X})$  a contraction satisfying*

$$\|M^n - P\|_\infty \leq \tilde{c}\delta^n$$

for some projection  $P$ ,  $\delta \in (0, 1)$ ,  $\tilde{c} \geq 0$  and all  $n \in \mathbb{N}$ . Moreover, assume there is  $b \geq 0$  so that

$$\|Pe^{t\mathcal{L}}P^\perp\|_\infty \leq tb \quad \text{and} \quad \|P^\perp e^{t\mathcal{L}}P\|_\infty \leq tb.$$

If  $(P\mathcal{L}P, \mathcal{D}(P\mathcal{L}P))$  is the generator of a  $C_0$ -semigroup, then there exists a constant  $c > 0$  and  $n_0 \in \mathbb{N}$  so that  $\tilde{c}\delta^{n_0} =: \tilde{\delta} < 1$  and for all  $x \in \mathcal{D}((\mathcal{L}P)^2)$

$$\left\| \left( M^{n_0} e^{\frac{t}{n}\mathcal{L}} \right)^n x - e^{tP\mathcal{L}P} Px \right\| \leq \frac{c}{n} \left( \|x\| + \|\mathcal{L}Px\| + \|(\mathcal{L}P)^2 x\| \right) + \tilde{\delta}^n \|x\|.$$

A more physically motivated result treating the same generalization as the corollary above is provided in Section 5.2. Before, Section 5.1 below is devoted to the proof of Theorem 3.1.

### 5.1. Proof of Theorem 3.1.

We assume for sake of clarity that  $t = 1$ , and split our proof in three parts:

$$\left\| \left( Me^{\frac{1}{n}\mathcal{L}} \right)^n x - e^{P\mathcal{L}P} Px \right\| \leq \left\| \left( Me^{\frac{1}{n}\mathcal{L}} \right)^n x - \left( Pe^{\frac{1}{n}\mathcal{L}}P \right)^n x \right\| \quad (10)$$

$$+ \left\| \left( Pe^{\frac{1}{n}\mathcal{L}}P \right)^n x - e^{nP(e^{\frac{1}{n}\mathcal{L}}-1)P} Px \right\| \quad (11)$$

$$+ \left\| e^{nP(e^{\frac{1}{n}\mathcal{L}}-1)P} Px - e^{P\mathcal{L}P} Px \right\| \quad (12)$$

for all  $x \in \mathcal{D}((\mathcal{L}P)^2)$ .

Upper bound on Equation (10):

**Lemma 5.2** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$  and  $M \in \mathcal{B}(\mathcal{X})$  a contraction. Assume that both satisfy the same assumption as in Theorem 3.1. Then, for all  $x \in \mathcal{X}$*

$$\left\| \left( Me^{\frac{1}{n}\mathcal{L}} \right)^n x - \left( Pe^{\frac{1}{n}\mathcal{L}}P \right)^n x \right\| \leq \left( \delta^n + \frac{b}{n} + \frac{1}{n} \frac{\delta}{1-\delta} e^{4b} \right) \|x\|.$$

The proof of the above lemma relies on a counting method: more precisely, we need to count the number transitions in a binary sequence. This is related to the *urn problem*, where  $k$  indistinguishable balls are placed in  $l$  distinguishable urns [30, Chap. 1.9]. Then, there are

$$\binom{k-1}{l-1} \quad (13)$$

possibilities to distribute the balls so that each urn contains at least one ball.

**Definition 5.3** Let  $S = \{A, B\}$ ,  $j, n, k \in \mathbb{N}$ , and  $n \geq 1$ . We define

$$\begin{aligned} S_{n,k} &:= \{s \in S^n \mid A \text{ appears } k \text{ times in } s\} \\ N(j, n, k) &:= \#\{s \in S_{n,k} \mid s \text{ includes } j \text{ transitions } AB \text{ or } BA\}. \end{aligned}$$

In words,  $N(j, n, k)$  counts the number of sequences consisting of  $k$   $A$ 's and  $n - k$   $B$ 's with the restriction that  $A$  changes to  $B$  or vice versa  $j$  times.

*Example 2.* Let  $S = \{A, B\}$ ,  $n = 4$ , and  $k = 2$ . Then,

$$\begin{aligned} N(0, n, k) &= \#\emptyset &&= 0 \\ N(1, n, k) &= \#\{AABB, BBAA\} &&= 2 \\ N(2, n, k) &= \#\{ABBA, BAAB\} &&= 2 \\ N(3, n, k) &= \#\{ABAB, BABA\} &&= 2. \end{aligned}$$

**Lemma 5.4** Let  $S = \{A, B\}$  and  $n, k, j \in \mathbb{N}$  with  $k \leq n$ . Then,

$$N(j, n, k) = \begin{cases} 2 \binom{n-k-1}{l-1} \binom{k-1}{l-1} & \text{if } j = 2l - 1 \\ \frac{n-2l}{l} \binom{n-k-1}{l-1} \binom{k-1}{l-1} & \text{if } j = 2l \\ \delta_{k \in \{0, n\}} & \text{if } j = 0 \end{cases}$$

for  $j \in \{0, \dots, 2 \min\{k, n - k\} - 1_{2k=n}\}$ . Otherwise  $N(j, n, k) = 0$ .

*Proof.* For  $j \leq 2 \min\{n - k, k\} - 1_{2k=n}$ , then  $N(j, n, k) \neq 0$  and we let  $s \in S_{n,k}$  so that  $s$  includes  $j$  transitions  $AB$  or  $BA$ . Next we assume that  $j = 0$ , the only possible sequences are  $s = A^n$  ( $k = n$ ) and  $s = B^n$  ( $k = 0$ ) so that  $N(0, n, k) = 1_{k \in \{0, n\}}$ . Assume now,  $k < n$ .

In the odd case  $j = 2l - 1$  for  $l \in \{1, \dots, \min\{k, n - k\}\}$ , the element  $s$  is constructed by  $l$  blocks of  $A$ 's and  $l$  blocks of  $B$ 's:

$$\begin{aligned} s &= \underbrace{A \dots A}_1 \underbrace{B \dots B}_1 \underbrace{A \dots A}_2 B \quad \dots \quad A \underbrace{B \dots B}_{l-1} \underbrace{A \dots A}_l \underbrace{B \dots B}_l, \\ s &= \underbrace{B \dots B}_1 \underbrace{A \dots A}_1 \underbrace{B \dots B}_2 A \quad \dots \quad B \underbrace{A \dots A}_{l-1} \underbrace{B \dots B}_l \underbrace{A \dots A}_l. \end{aligned}$$

Identifying these blocks with distinguishable urns and the elements  $A$  and  $B$  with indistinguishable balls (q.v. Eq. (13)), the task is to count the possibilities of placing  $k$   $A$ 's in  $l$  urns and vice versa  $n - k$   $B$ 's in  $l$  urns with the additional assumption that each urn must contain at least one  $A$  or one  $B$ . By changing the roles of  $A$  and  $B$ , we get twice the number of possible combinations. Therefore, one of the *Twelvefold Ways* [30, Chap. 1.9] shows

$$N(j, n, k) = 2 \binom{n-k-1}{l-1} \binom{k-1}{l-1}.$$

In the even case  $j = 2l$  for  $l \in \{1, \dots, \min\{k, n - k\} - 1_{2k=n}\}$ , we can argue similarly to the odd case. The only difference is that  $s$  is constructed by  $l + 1$  blocks of  $A$ 's and  $l$  blocks of  $B$ 's or vice versa:

$$\begin{aligned} s &= \underbrace{A \dots A}_1 \underbrace{B \dots B}_1 \underbrace{A \dots A}_2 B \quad \dots \quad A \underbrace{B \dots B}_{l-1} \underbrace{A \dots A}_l \underbrace{B \dots B}_l \underbrace{A \dots A}_{l+1}, \\ s &= \underbrace{B \dots B}_1 \underbrace{A \dots A}_1 \underbrace{B \dots B}_2 B \quad \dots \quad A \underbrace{A \dots A}_{l-1} \underbrace{B \dots B}_l \underbrace{A \dots A}_l \underbrace{B \dots B}_{l+1}. \end{aligned}$$

Then the *Twelvetfold Ways* [30, Chap. 1.9] gives:

$$\begin{aligned} N(j, n, k) &= \binom{n-k-1}{l-1} \binom{k-1}{l} + \binom{n-k-1}{l} \binom{k-1}{l-1} \\ &= \frac{n-2l}{l} \binom{n-k-1}{l-1} \binom{k-1}{l-1} \end{aligned}$$

which proves the statement.  $\square$

With the help of this counting method, we are ready to prove Lemma 5.2. In what follows, we identify the couple  $(A, B)$  with the product  $AB$  by slight abuse of notations.

*Proof of Lemma 5.2.* By the assumption on  $M$ ,  $MP = P$  because for all  $n \in \mathbb{N}$

$$\|PM - P\|_\infty \leq \|(P - M^n)M\|_\infty + \|M^{n+1} - P\|_\infty \leq \delta^n + \delta^{n+1}. \quad (14)$$

The main idea is to split  $M = P + M^\perp$  with  $M^\perp := P^\perp M$  and order the terms after expanding the following polynomial appropriately. Let  $A := M^\perp e^{\frac{1}{n}\mathcal{L}}$  and  $B := P e^{\frac{1}{n}\mathcal{L}}$  so that

$$\left( (P + M^\perp) e^{\frac{1}{n}\mathcal{L}} \right)^n = B^n + \sum_{k=1}^{n-1} \sum_{s \in S_{n,k}} s + A^n \quad (15)$$

where elements in  $S_{n,k}$  are identified with sequences of concatenated operators and denoted by  $s$ . Then, we partition summands by the number of transitions from  $A$  to  $B$  or vice versa and use

$$\|A\|_\infty \leq \delta \quad \text{and} \quad \|AB\|_\infty = \left\| M^\perp P^\perp e^{\frac{1}{n}\mathcal{L}} P M e^{\frac{1}{n}\mathcal{L}} \right\|_\infty \leq \delta \frac{b}{n}.$$

The number of summands with  $j$  transitions is equal to  $N(j, n, k)$  given by Lemma 5.4 for  $j \in \{1, \dots, m\}$  and  $m := 2 \min\{k, n-k\} - 1_{2k=n}$ . Then, the inequality above shows

$$\begin{aligned} \left\| \left( M e^{\frac{1}{n}\mathcal{L}} \right)^n - \left( P e^{\frac{1}{n}\mathcal{L}} \right)^n \right\|_\infty &\leq \delta^n + \sum_{k=1}^{n-1} \sum_{j=1}^m \delta^k N(j, n, k) \left( \frac{b}{n} \right)^j \\ &= \delta^n + \sum_{k=1}^{n-1} \sum_{l=1}^{\lfloor \frac{m}{2} \rfloor} \delta^k 2 \binom{n-k-1}{l-1} \binom{k-1}{l-1} \left( \frac{b}{n} \right)^{2l-1} \\ &\quad + \sum_{k=1}^{n-1} \sum_{l=1}^{\lfloor \frac{m}{2} \rfloor} \delta^k \frac{b(n-2l)}{nl} \binom{n-k-1}{l-1} \binom{k-1}{l-1} \left( \frac{b}{n} \right)^{2l-1} \\ &\leq \delta^n + (2+b) \sum_{l=1}^{n-1} \sum_{k=1}^{n-1} \delta^k \frac{n^{2l-2}}{(l-1)!^2} \left( \frac{b}{n} \right)^{2l-1} \\ &\leq \delta^n + \frac{b(2+b)}{n} \frac{\delta + \delta^n}{1-\delta} \left( \sum_{l=0}^{n-2} \frac{b^l}{l!} \right)^2 \\ &\leq \delta^n + \frac{1}{n} \frac{\delta}{1-\delta} e^{4b}. \end{aligned}$$

In the third inequality, we used the upper bound  $\binom{n}{k} \leq \frac{n^k}{k!}$  to shows

$$\binom{n-k-1}{l-1} \binom{k-1}{l-1} \leq \frac{n^{2l-2}}{(l-1)!^2}.$$

Additionally, we increase the upper index to  $n-1$ , and upper bound

$$2 + \frac{b(n-2l)}{nl} \leq 2 + b.$$

Applying the assumptions again to

$$\left(Pe^{\frac{1}{n}\mathcal{L}}\right)^n x - \left(Pe^{\frac{1}{n}\mathcal{L}}P\right)^n x = \left(Pe^{\frac{1}{n}\mathcal{L}}P\right)^{n-1} Pe^{\frac{1}{n}\mathcal{L}}P^\perp x$$

proves the lemma:

$$\left\| \left(Me^{\frac{1}{n}\mathcal{L}}\right)^n x - \left(Pe^{\frac{1}{n}\mathcal{L}}P\right)^n x \right\| \leq \left( \delta^n + \frac{b}{n} + \frac{1}{n} \frac{\delta}{1-\delta} e^{4b} \right) \|x\|. \quad \square$$

*Remark.* By the introduced counting method, we can approximate  $(Me^{\frac{1}{n}\mathcal{L}})^n$  by  $(Pe^{\frac{1}{n}\mathcal{L}}P)^n$ , which is independent of  $M^\perp$ . In previous works [25, 2], the operators considered in similar proof steps as Equation (11) and (12) depend on  $M^\perp$ .

*Upper bound on Equation (11):* In the next step, we apply our modified Chernoff Lemma 4.2:

**Lemma 5.5** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$  and  $P \in \mathcal{B}(\mathcal{X})$  be a projection. Assume that both operators satisfy the same assumption as in Theorem 3.1. Then, for all  $x \in \mathcal{D}((\mathcal{L}P)^2)$*

$$\left\| \left(Pe^{\frac{1}{n}\mathcal{L}}P\right)^n x - e^{nP(e^{\frac{1}{n}\mathcal{L}}-1)P} Px \right\| \leq \frac{1}{2n} \left( b^2 \|x\| + b \|\mathcal{L}Px\| + \|(\mathcal{L}P)^2 x\| \right).$$

*Proof.* The proof relies on the modified Chernoff Lemma (q.v. Lem. 4.2) applied to the contraction  $C(\frac{1}{n}) = Pe^{\frac{1}{n}\mathcal{L}}P$  on  $\mathcal{B}(Px)$ . Then, for all  $x \in \mathcal{X}$

$$\left\| \left(Pe^{\frac{1}{n}\mathcal{L}}P\right)^n x - e^{nP(e^{\frac{1}{n}\mathcal{L}}-1)P} Px \right\| \leq \frac{n}{2} \left\| \left(P \left(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1}\right) P\right)^2 x \right\|.$$

Moreover, the asymptotic Zeno condition (5) implies

$$\left\| P^\perp \mathcal{L}Px \right\| = \lim_{h \rightarrow 0} \frac{1}{h} \left\| P^\perp e^{h\mathcal{L}} Px \right\| \leq b \|x\|$$

for all  $x \in \mathcal{D}(\mathcal{L}P)$ , and hence  $P^\perp \mathcal{L}P$  is a bounded operator with  $\|P^\perp \mathcal{L}P\|_\infty \leq b$ . Next, given  $x \in \mathcal{D}(\mathcal{L}P)$ , the  $C_0$ -semigroup properties (q.v. Lem. 2.1) imply

$$\begin{aligned} n \left\| \left(P \left(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1}\right) P\right)^2 x \right\| &= \left\| P \left(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1}\right) P \int_0^1 e^{\frac{\tau_1}{n}\mathcal{L}} (\mathbb{1} - P + P) \mathcal{L}Px d\tau_1 \right\| \\ &\leq \left\| P \left(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1}\right) P \int_0^1 e^{\frac{\tau_1}{n}\mathcal{L}} P \mathcal{L}Px d\tau_1 \right\| \\ &\quad + 2 \int_0^1 \left\| P e^{\frac{\tau_1}{n}\mathcal{L}} P^\perp \right\|_\infty d\tau_1 \left\| P^\perp \mathcal{L}P \right\|_\infty \|x\|. \end{aligned}$$

Note that  $\int_0^1 e^{\frac{\tau_1}{n}\mathcal{L}} P \mathcal{L}Px$  belongs to  $\mathcal{D}(\mathcal{L})$  by Lemma 2.1, but not necessarily to  $\mathcal{D}(\mathcal{L}P)$ . However, for all  $x \in \mathcal{D}((\mathcal{L}P)^2)$

$$\begin{aligned} n \left\| \left(P \left(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1}\right) P\right)^2 x \right\| &\leq \left\| P \left(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1}\right) P \int_0^1 e^{\frac{\tau_1}{n}\mathcal{L}} P \mathcal{L}Px d\tau_1 \right\| + \frac{b^2}{n} \|x\| \\ &\leq \left\| P \left(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1}\right) \int_0^1 e^{\frac{\tau_1}{n}\mathcal{L}} P \mathcal{L}Px d\tau_1 \right\| \\ &\quad + \frac{b^2}{n} \|x\| + \left\| P e^{\frac{1}{n}\mathcal{L}} P^\perp \right\|_\infty \|\mathcal{L}Px\| \\ &\leq \frac{1}{n} \left\| P \int_0^1 e^{\frac{\tau_2}{n}\mathcal{L}} \mathcal{L} \int_0^1 e^{\frac{\tau_1}{n}\mathcal{L}} P \mathcal{L}Px d\tau_1 d\tau_2 \right\| \\ &\quad + \frac{b^2}{n} \|x\| + \frac{b}{n} \|\mathcal{L}Px\| \\ &\leq \frac{1}{n} \left( b^2 \|x\| + b \|\mathcal{L}Px\| + \|(\mathcal{L}P)^2 x\| \right), \end{aligned}$$

which proves Lemma 5.5. □

*Remark.* As regards to the convergence rate of the quantum Zeno effect, Lemma 5.5 constitutes our main improvement compared to the work [2]. The modified Chernoff lemma allows to improve the convergence rate to  $n^{-1}$ .

*Upper bound on Equation (12):* Finally, we prove an upper bound on Equation (12), which can be interpreted as a modified *Dunford-Segal approximation* [15].

**Lemma 5.6** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$  and  $P \in \mathcal{B}(\mathcal{X})$  be a projection. Assume that both operators satisfy the assumptions of Theorem 3.1. Then, for all  $x \in \mathcal{D}((\mathcal{L}P)^2)$*

$$\left\| e^{nP(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1})P} Px - e^{P\mathcal{L}P} Px \right\| \leq \frac{e^{\bar{b}}}{2n} \left( b^2 \|x\| + b \|\mathcal{L}Px\| + \|(\mathcal{L}P)^2 x\| \right)$$

with  $e^{\bar{b}} := \sup_{s \in [0,1]} \|e^{sP\mathcal{L}P}\|_\infty < \infty$ .

*Proof.* The proof relies on the integral equation for semigroups from Lemma 2.2. We start by proving the continuity of

$$[0, 1] \ni s \mapsto e^{snP(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1})P} \left( nP(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1})P - P\mathcal{L}P \right) e^{(1-s)P\mathcal{L}P} Px. \quad (16)$$

Let  $x_s := e^{(1-s)P\mathcal{L}P} Px$ . By Lemma 2.1  $x_s \in \mathcal{D}((\mathcal{L}P)^2)$  if  $x \in \mathcal{D}((\mathcal{L}P)^2)$  and

$$\begin{aligned} nP(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1})Px_s - P\mathcal{L}Px_s &= P \int_0^1 e^{\frac{\tau_1}{n}\mathcal{L}} \mathcal{L}Px_s d\tau_1 - P\mathcal{L}Px_s \\ &= \frac{1}{n} P \int_0^1 \int_0^1 \tau_1 e^{\frac{\tau_1\tau_2}{n}\mathcal{L}} (\mathcal{L}P)^2 x_s d\tau_2 d\tau_1 \\ &\quad + P \int_0^1 e^{\frac{\tau_1}{n}\mathcal{L}} P^\perp \mathcal{L}Px_s d\tau_1 \\ &= P \int_0^1 \int_0^1 \frac{\tau_1}{n} e^{\frac{\tau_1\tau_2}{n}\mathcal{L}} P^\perp \mathcal{L}P\mathcal{L}Px_s d\tau_2 d\tau_1 \\ &\quad + P \int_0^1 \int_0^1 \frac{\tau_1}{n} e^{\frac{\tau_1\tau_2}{n}\mathcal{L}} (\mathcal{L}P)^2 x_s d\tau_2 d\tau_1 \\ &\quad + P \int_0^1 e^{\frac{\tau_1}{n}\mathcal{L}} P^\perp \mathcal{L}Px_s d\tau_1 \end{aligned} \quad (17)$$

Moreover,  $P\mathcal{L}Pe^{(1-s)P\mathcal{L}P}x = e^{(1-s)P\mathcal{L}P}P\mathcal{L}Px$  for all  $x \in \mathcal{D}(\mathcal{L}P)$  so that for all  $x \in \mathcal{D}((\mathcal{L}P)^2)$

$$s \mapsto P\mathcal{L}Px_s = e^{(1-s)P\mathcal{L}P}P\mathcal{L}Px \quad \text{and} \quad s \mapsto (P\mathcal{L}P)^2 x_s = e^{(1-s)P\mathcal{L}P}(P\mathcal{L}P)^2 x,$$

which are both continuous vector-valued maps. Since  $[0, 1] \ni s \mapsto \exp(snP(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1})P)$  is uniformly continuous and  $P^\perp \mathcal{L}P$  bounded, the vector-valued map in Equation (16) is a sum of continuous maps and thereby continuous. Therefore, Lemma 2.2 gives for all  $x \in \mathcal{D}((\mathcal{L}P)^2)$

$$\begin{aligned} e^{nP(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1})P} Px - e^{P\mathcal{L}P} Px &= \int_0^1 e^{snP(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1})P} \left( nP(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1})P - P\mathcal{L}P \right) e^{(1-s)P\mathcal{L}P} Px ds. \end{aligned}$$

In the following, we apply the upper bound  $\|P^\perp \mathcal{L}P\|_\infty \leq b$ , the assumption  $\|Pe^{s\mathcal{L}}P^\perp\|_\infty \leq sb$ , and the property that  $(e^{snP(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1})P})_{s \geq 0}$  is a contraction to Equation (16) and (17):

$$\begin{aligned} \left\| e^{nP(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1})P} Px - e^{P\mathcal{L}P} Px \right\| &\leq \int_0^1 \left\| \left( nP(e^{\frac{1}{n}\mathcal{L}} - \mathbb{1})P - P\mathcal{L}P \right) x_\tau \right\| d\tau \\ &\leq \frac{e^{\bar{b}}}{2n} \left( b^2 \|x\| + b \|\mathcal{L}Px\| + \|(\mathcal{L}P)^2 x\| \right) \end{aligned}$$

where  $e^{\tilde{b}} = \sup_{s \in [0,1]} \|e^{sP\mathcal{L}P}\|_\infty$ . □

The above approximation of  $e^{P\mathcal{L}P}$  by  $e^{nP(e^{\frac{1}{n}\mathcal{L}}-1)P}$  is similar to the Dunford-Segal approximation, which would be given by  $\exp(n(\exp(\frac{1}{n}P\mathcal{L}P) - \mathbb{1}))$ : for a generator  $(\mathcal{K}, \mathcal{D}(\mathcal{K}))$  of a bounded  $C_0$ -semigroup, Gomilko and Tomilov proved [15, Cor. 1.4]

$$\left\| e^{nt(e^{\frac{1}{n}\mathcal{K}}-1)}x - e^{t\mathcal{K}}x \right\| \leq 8\tilde{b}\frac{t}{n}\|\mathcal{K}^2x\|$$

for all  $x \in \mathcal{D}(\mathcal{K}^2)$  and  $\tilde{b} := \sup_{t \geq 0} \|e^{t\mathcal{K}}\|_\infty$ . In our case, it is not clear whether  $(e^{sP\mathcal{L}P})_{s \geq 0}$  is uniformly bounded.

*Remark.* The specificity of the last step stems from the fact that  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  is unbounded. In the previous works [25, 2] a similar step exists but in both papers  $\mathcal{L}$  was assumed to be bounded.

*End of the proof of Theorem 3.1:* We combine Lemma 5.2, 5.5, and 5.6 to prove Theorem 3.1.

*Proof of Theorem 3.1.* Let  $x \in \mathcal{D}((\mathcal{L}P)^2)$ . Then,

$$\begin{aligned} \left\| \left( Me^{\frac{1}{n}\mathcal{L}} \right)^n x - e^{P\mathcal{L}P}Px \right\| &\leq \left( \delta^n + \frac{b}{n} + \frac{1}{n} \frac{\delta}{1-\delta} e^{4b} \right) \|x\| && \text{(Lem. 5.2)} \\ &+ \frac{1}{2n} \left( b^2 \|x\| + b \|\mathcal{L}Px\| + \|(\mathcal{L}P)^2x\| \right) && \text{(Lem. 5.5)} \\ &+ \frac{e^{\tilde{b}}}{2n} \left( b^2 \|x\| + b \|\mathcal{L}Px\| + \|(\mathcal{L}P)^2x\| \right) && \text{(Lem. 5.6)}. \end{aligned}$$

Redefining  $\mathcal{L}$  by  $t\mathcal{L}$  and  $b$  by  $tb$ , we get

$$\left\| \left( Me^{\frac{t}{n}\mathcal{L}} \right)^n x - e^{tP\mathcal{L}P}Px \right\| \leq \frac{c}{n} \left( \|x\| + \|\mathcal{L}Px\| + \|(\mathcal{L}P)^2x\| \right) + \delta^n \|x\|$$

with the constant  $c := \frac{1}{1-\delta} e^{4tb} + t^2 \left( 1 + \frac{\tilde{b}}{2} \right) \max\{b^2, b, 1\}$  and  $e^{\tilde{b}} = \sup_{s \in [0,t]} \|e^{sP\mathcal{L}P}\|_\infty$ . □

*Remark.* The upper bound in Theorem 3.1 can be formulated for all  $x \in \mathcal{D}(\mathcal{L}P)$ . For that one must stop at an earlier stage of the proof and express the error terms by appropriate integrals.

## 5.2. Generalized Power Convergence.

In this subsection, we weaken the assumptions on the contraction  $M$  at the cost of stronger assumptions on the  $C_0$ -semigroup. For that, we combine the holomorphic functional calculus with the semicontinuity of the spectrum of  $M$  perturbed by the semigroup under certain conditions. We refer to Appendix B for details on the tools needed to prove the main result of this section. With the help of these results, we can weaken the assumption on  $M$  in Equation (4).

**Proposition 3.2** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$  and  $M \in \mathcal{B}(\mathcal{X})$  a contraction such that*

$$\|M^n - P\|_\infty \leq \tilde{c}\delta^n \tag{7}$$

for some projection  $P$ ,  $\delta \in (0, 1)$  and  $\tilde{c} \geq 0$ . Moreover, we assume that there is  $b \geq 0$  so that

$$\left\| Pe^{t\mathcal{L}}P^\perp \right\|_\infty \leq tb, \quad \left\| M^\perp e^{t\mathcal{L}} - M^\perp \right\|_\infty \leq tb, \quad \text{and} \quad \left\| P^\perp e^{t\mathcal{L}}P \right\|_\infty \leq tb. \tag{8}$$

If  $(P\mathcal{L}P, \mathcal{D}(\mathcal{L}P))$  generates a  $C_0$ -semigroup, then there is  $\epsilon > 0$  such that for all  $t \geq 0$ ,  $n \in \mathbb{N}$  satisfying  $t \in [0, n\epsilon]$ ,  $\tilde{\delta} \in (\delta, 1)$ , and  $x \in \mathcal{D}((\mathcal{L}P)^2)$

$$\left\| \left( Me^{\frac{t}{n}\mathcal{L}} \right)^n x - e^{tP\mathcal{L}P}Px \right\| \leq \frac{c_1}{n} \left( \|x\| + \|\mathcal{L}Px\| + \|(\mathcal{L}P)^2x\| \right) + c_2 \tilde{\delta}^n \|x\|,$$

for some constants  $c_1, c_2 \geq 0$ .

The only difference to the proof of Theorem 3.1 is summarized in the question: How can we upper bound  $\|(M^\perp e^{\frac{t}{n}\mathcal{L}})^k\|_\infty$  for all  $k \in \{1, \dots, n\}$  with the weaker assumption (7) on  $M$ ? For that, we need to replace the argument in the proof of Lemma 5.2, which only worked for the special case  $\tilde{c} = 1$ .

*Proof.* Since the bounds found in Lemma 5.5 and 5.6 are independent of the value of  $\tilde{c}$ , it is enough to improve Lemma 5.2:


$$\begin{aligned} \left\| \left( M e^{\frac{t}{n}\mathcal{L}} \right)^n x - e^{tP\mathcal{L}P} P x \right\| &\leq \left\| \left( M e^{\frac{t}{n}\mathcal{L}} \right)^n x - \left( P e^{\frac{t}{n}\mathcal{L}P} \right)^n x \right\| \\ &+ \frac{t^2}{n} \left( 2b^2 \|x\| + b \|\mathcal{L}P x\| + \|(\mathcal{L}P)^2 x\| \right) \end{aligned} \quad (\text{Lem. 5.5})$$

$$+ \frac{e^{\tilde{b}t^2}}{2n} \left( b^2 \|x\| + b \|\mathcal{L}P x\| + \|(\mathcal{L}P)^2 x\| \right) \quad (\text{Lem. 5.6})$$

with  $e^{\tilde{b}} = \sup_{s \in [0, t]} \|e^{sP\mathcal{L}P}\|_\infty < \infty$ . Since the assumption (7) on  $M$  is a special case of the uniform power convergence (q.v. Eq. (9)), Proposition A.1 shows the equivalence of the assumption to the spectral gap assumption, that is

$$\sigma(M) \subset \mathbb{D}_\delta \cup \{1\},$$

where the quasinilpotent operator corresponding to the eigenvalue 1 is zero. Therefore, the eigenprojection  $P$  w.r.t. 1 satisfies  $MP = PM = P$  and the curve  $\gamma : [0, 2\pi] \rightarrow \mathbb{C}$ ,  $\varphi \mapsto \tilde{\delta} e^{i\varphi}$  encloses the spectrum of  $M^\perp := MP^\perp$ . Together with the second bound in (8), Lemma B.2 shows that there exists  $\epsilon > 0$  such that the spectrum of  $M^\perp e^{s\mathcal{L}}$  can be separated by  $\gamma$  for all  $s \in [0, \epsilon]$ . Therefore, the holomorphic functional calculus (q.v. Prop. B.1) shows for all  $t \in [0, n\epsilon]$  and  $k \in \mathbb{N}$



$$\left( M^\perp e^{\frac{t}{n}\mathcal{L}} \right)^k = \frac{1}{2\pi i} \oint_\gamma z^k R(z, M^\perp e^{\frac{t}{n}\mathcal{L}}) dz. \quad (18)$$

Let  $t \geq 0$ ,  $n \in \mathbb{N}$  so that  $t \in [0, n\epsilon]$ . By the *principle of stability of bounded invertibility* [20, Thm. IV.2.21],  $R(z, M^\perp e^{s\mathcal{L}})$  is well-defined and bounded for all  $z \in \gamma$  and  $s \in [0, \epsilon]$ . More explicitly, using the *second Neumann series* for the resolvent [20, p. 67], we have

$$\begin{aligned} \left\| R(z, M^\perp e^{s\mathcal{L}}) \right\|_\infty &= \left\| R(z, M^\perp) \sum_{p=0}^{\infty} \left( (M^\perp e^{s\mathcal{L}} - M^\perp) R(z, M^\perp) \right)^p \right\|_\infty \\ &\leq \left\| R(z, M^\perp) \right\|_\infty \sum_{p=0}^{\infty} \left( sb \left\| R(z, M^\perp) \right\|_\infty \right)^p \\ &\leq \sup_{z \in \gamma} \left\| R(z, M^\perp) \right\|_\infty \frac{2 + 2\tilde{\delta}^2}{1 + 2\tilde{\delta}^2}, \end{aligned} \quad (19)$$

where we applied the assumption (8), the following upper bound on  $s$  (q.v. Eq. (38)):

$$s \leq \epsilon < \frac{1}{2b} (1 + \tilde{\delta}^2)^{-1} \left( 1 + \sup_{z \in \gamma} \left\| R(z, M^\perp) \right\|_\infty^2 \right)^{-\frac{1}{2}} \leq \frac{1}{2b} (1 + \tilde{\delta}^2)^{-1} \sup_{z \in \gamma} \left\| R(z, M^\perp) \right\|_\infty^{-1},$$

and finally computed the geometric series. Combining Equation (18) and (19) shows for all  $k \geq 1$

$$\left\| \left( M^\perp e^{\frac{t}{n}\mathcal{L}} \right)^{k-1} M^\perp \right\|_\infty \leq \frac{1}{2\pi} \oint_\gamma |z|^{k-1} \left\| R(z, M^\perp e^{\frac{t}{n}\mathcal{L}}) \right\|_\infty dz \leq \tilde{\delta}^k \sup_{z \in \gamma} \left\| R(z, M^\perp) \right\|_\infty \frac{2 + 2\tilde{\delta}^2}{1 + 2\tilde{\delta}^2}.$$

Defining  $c_2 := \sup_{z \in \gamma} \left\| R(z, M^\perp) \right\|_\infty \frac{2 + 2\tilde{\delta}^2}{1 + 2\tilde{\delta}^2}$ , we achieve

$$\left\| \left( M^\perp e^{\frac{t}{n}\mathcal{L}} \right)^{k-1} M^\perp \right\|_\infty \leq c_2 \tilde{\delta}^k \quad (20)$$

for all  $k \in \{1, \dots, n\}$ . Next, we expand

$$\left(Me^{\frac{t}{n}\mathcal{L}}\right)^n = \left(PMe^{\frac{t}{n}\mathcal{L}} + M^\perp e^{\frac{t}{n}\mathcal{L}}\right)^n.$$

Denoting  $A := M^\perp e^{\frac{t}{n}\mathcal{L}}$  and  $B := Pe^{\frac{t}{n}\mathcal{L}}$ , the above  $n^{\text{th}}$ -power can be expanded in terms of sequences of the form

$$A\dots AB\dots BA\dots AB\dots \quad \text{or} \quad B\dots BA\dots AB\dots BA\dots$$

Similarly to Lemma 5.2, we can upper bound every sequence w.r.t. the number of transitions  $AB$  or  $BA$  using the assumptions (8) on the  $C_0$ -semigroup as well as the inequality (20). The only difference to the proof of Lemma 5.2 is the constant  $c_2$  in the inequality so that

$$\left\| \left(Me^{\frac{t}{n}\mathcal{L}}\right)^n - \left(Pe^{\frac{t}{n}\mathcal{L}}\right)^n \right\|_\infty \leq c_2 \tilde{\delta}^n + \sum_{k=1}^{n-1} \sum_{j=1}^m \tilde{\delta}^k c_2^j N(j, n, k) \left(\frac{tb}{n}\right)^j \leq c_2 \tilde{\delta}^n + \frac{1}{n} \frac{\tilde{\delta}}{1 - \tilde{\delta}} e^{4tbc_2},$$

where  $m := 2 \min\{k, n - k\} - 1_{2k=n}$ . Then, for all  $x \in \mathcal{D}((\mathcal{L}P)^2)$

$$\left\| \left(Me^{\frac{t}{n}\mathcal{L}}\right)^n x - e^{tP\mathcal{L}P} Px \right\| \leq \frac{c_1}{n} \left( \|x\| + \|\mathcal{L}Px\| + \|(\mathcal{L}P)^2 x\| \right) + c_2 \tilde{\delta}^n \|x\|$$

with  $c_1 := \frac{\tilde{\delta}}{1 - \tilde{\delta}} e^{4tbc_2} + t^2 \left(2 + \frac{e^{\tilde{b}}}{2}\right) \max\{b^2, b, 1\}$  and  $e^{\tilde{b}} = \sup_{s \in [0, t]} \|e^{sP\mathcal{L}P}\|_\infty$ .  $\square$

## 6. UNIFORM POWER CONVERGENCE WITH FINITELY MANY EIGENVALUES

In this section, we weaken the assumption on  $M$  to the uniform power convergence assumption (q.v. Eq. (9)), that is we allow for finitely many eigenvalues  $\{\lambda_j\}_{j=1}^J$  and associated projections  $\{P_j\}_{j=1}^J$ . Similarly to Theorem 3 in [2], we strengthen the assumptions on the  $C_0$ -semigroup to  $M\mathcal{L}$  and  $\mathcal{L}P_\Sigma$  being densely defined and bounded by  $b \geq 0$ , where  $P_\Sigma := \sum_{j=1}^J P_j$ . Under those assumptions, we can prove the Zeno convergence in the uniform topology:

**Theorem 3.3** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$  and  $M \in \mathcal{B}(\mathcal{X})$  a contraction satisfying the following uniform power convergence: there is  $\tilde{c} > 0$  so that*

$$\left\| M^n - \sum_{j=1}^J \lambda_j^n P_j \right\|_\infty \leq \tilde{c} \delta^n \quad (9)$$

for a set of orthogonal projections  $\{P_j\}_{j=1}^J$ , eigenvalues  $\{\lambda_j\}_{j=1}^J \subset \partial\mathbb{D}_1$ , and a rate  $\delta \in (0, 1)$ . For  $P_\Sigma := \sum_{j=1}^J P_j$ , we assume that  $M\mathcal{L}$  and  $\mathcal{L}P_\Sigma$  are densely defined and bounded by  $b \geq 0$ . Then, there is an  $\epsilon > 0$  such that for all  $n \in \mathbb{N}$ ,  $t \geq 0$  satisfying  $t \in [0, n\epsilon]$

$$\left\| \left(Me^{\frac{t}{n}\mathcal{L}}\right)^n - \sum_{j=1}^J \lambda_j^n e^{tP_j\mathcal{L}P_j} P_j \right\|_\infty \leq \frac{c_1 J}{n} + c_2 \delta^n,$$

for some constants  $c_1, c_2 \geq 0$ .

### 6.1. Proof of Theorem 3.3.

Similarly to the papers [25] and [2], we use the holomorphic functional calculus to separate the spectrum of the perturbed contraction  $Me^{t\mathcal{L}}$  appearing in the Zeno sequence. However, and in contrast to [2] where the  $C_0$ -semigroup is approximated by a sequence of uniformly continuous semigroups, we instead crucially rely upon the uniform continuity of the perturbed contraction



to recover the optimal convergence rate. We upper bound the following terms:

$$\left\| \left( M e^{\frac{t}{n} \mathcal{L}} \right)^n - \sum_{j=1}^J \lambda_j e^{t P_j \mathcal{L} P_j} P_j \right\|_{\infty} \leq \left\| \left( M e^{\frac{t}{n} \mathcal{L}} \right)^n - \left( P_{\Sigma} M e^{\frac{t}{n} \mathcal{L}} P_{\Sigma} \right)^n \right\|_{\infty} \quad (21)$$

$$+ \left\| \left( P_{\Sigma} M e^{\frac{t}{n} \mathcal{L}} P_{\Sigma} \right)^n - \sum_{j=1}^J \lambda_j^n e^{n(C_j(\frac{t}{n}) - P_j(\frac{t}{n}))} P_j(\frac{t}{n}) \right\|_{\infty} \quad (22)$$

$$+ \left\| \sum_{j=1}^J \lambda_j^n e^{n(C_j(\frac{t}{n}) - P_j(\frac{t}{n}))} P_j(\frac{t}{n}) - \sum_{j=1}^J \lambda_j^n e^{t P_j \mathcal{L} P_j} P_j \right\|_{\infty} \quad (23)$$

where the definitions of the perturbed spectral projection  $P_j(\frac{t}{n})$  and  $C_j(\frac{t}{n})$  are postponed to Lemma 6.3.

*Approximation of the Perturbed Spectral Projection:* In the following result, we construct a perturbed spectral projection for which we obtain an approximation bound (cf. [2, Lem. 5.3]). The key tools are the holomorphic functional calculus and the semicontinuity of the spectrum under uniform perturbations. The statements are summarized in Proposition B.1 and Lemma B.2.

**Lemma 6.1** *Let  $A \in \mathcal{B}(\mathcal{X})$ ,  $t \mapsto B(t)$  be a vector-valued map on  $\mathcal{B}(\mathcal{X})$  which is uniformly continuous at  $t = 0$  with  $\sup_{t \geq 0} \|B(t)\|_{\infty} \leq b$ , and  $\Gamma : [0, 2\pi] \rightarrow \rho(A)$  be a curve separating  $\sigma(A)$ . Then, there exists  $\epsilon > 0$  so that for all  $t \in [0, \epsilon]$*

$$P(t) := \frac{1}{2\pi i} \oint_{\Gamma} R(z, A + tB(t)) dz$$

defines a projection. The perturbation  $P(t) - P(0)$  is controlled by

$$\|P(t) - P(0)\|_{\infty} \leq t \sup_{z \in \Gamma} \|R(z, A)\|_{\infty}^2 b \inf_{z \in \Gamma} \frac{2 + 2|z|^2}{1 + 2|z|^2} \frac{|\Gamma|}{2\pi}, \quad (24)$$

where  $|\Gamma|$  denotes the length of the curve  $\Gamma$ . The derivative at  $t = 0$  corresponding to the vector-valued map  $t \mapsto P(t)$  is

$$P' := \lim_{t \rightarrow 0} \frac{P(t) - P(0)}{t} = \frac{1}{2\pi i} \oint_{\Gamma} R(z, A) B(0) R(z, A) dz. \quad (25)$$

Finally, there exist  $d_1 > 0$  and  $d_2 > 0$  so that for all  $t \in [0, \epsilon]$

$$\|P(t) - P(0) - tP'\|_{\infty} \leq d_1 t^2 + d_2 t \|B(t) - B(0)\|_{\infty}.$$

*Proof.* Since  $t \mapsto B(t)$  is uniformly continuous at  $t = 0$ , the vector-valued map  $t \mapsto A + tB(t)$  is as well. Then, Lemma B.2 states that there exists  $\epsilon > 0$  such that  $\sigma(A + tB(t))$  is separated by  $\Gamma$  for all  $t \in [0, \epsilon]$  and Proposition B.1 shows that

$$P(t) = \frac{1}{2\pi i} \oint_{\Gamma} R(z, A + tB(t)) dz$$

defines a projection on  $\mathcal{B}(\mathcal{X})$  for all  $t \in [0, \epsilon]$ . Let  $R := \sup_{z \in \Gamma} \|R(z, A)\|_{\infty} < \infty$ , then using the same steps as in Equation (19), we have that for all  $\eta \in \Gamma$

$$\|R(\eta, A + tB(t))\|_{\infty} \leq R \inf_{z \in \Gamma} \frac{2 + 2|z|^2}{1 + 2|z|^2}. \quad (26)$$

Therefore, the perturbed resolvent is uniformly bounded. To prove the explicit representation of the derivative and the quantitative approximation, we follow the ideas of [2, Lem. 5.2-5.3]:

$$\begin{aligned} \frac{P(t) - P(0)}{t} &= \frac{1}{t 2\pi i} \left( \oint_{\Gamma} R(z, A + tB(t)) dz - \oint_{\Gamma} R(z, A) dz \right) \\ &= \frac{1}{2\pi i} \oint_{\Gamma} R(z, A + tB(t)) B(t) R(z, A) dz, \end{aligned}$$

proves Equation (25) by Lebesgue's dominated convergence theorem [17, Thm. 3.7.9]. Moreover,

$$\begin{aligned}
\|P(t) - P(0) - tP'\|_\infty &\leq \frac{t}{2\pi} \left\| \oint_\Gamma (R(z, A + tB(t)) - R(z, A))B(t)R(z, A)dz \right\|_\infty \\
&\quad + \frac{t}{2\pi} \left\| \oint_\Gamma R(z, A) (B(t) - B(0)) R(z, A)dz \right\|_\infty \\
&\leq \frac{t}{2\pi} \left\| \oint_\Gamma R(z, A + tB(t))tB(t)R(z, A)B(t)R(z, A)dz \right\|_\infty \\
&\quad + \frac{t}{2\pi} \left\| \oint_\Gamma R(z, A) (B(t) - B(0)) R(z, A)dz \right\|_\infty \\
&\leq \frac{tR^2|\Gamma|}{2\pi} \left( tb^2R \inf_{z \in \Gamma} \frac{2 + 2|z|^2}{1 + 2|z|^2} + \|B(t) - B(0)\|_\infty \right) \\
&\leq c_1t^2 + tc_2\|B(t) - B(0)\|_\infty
\end{aligned} \tag{27}$$

with  $d_1 := \frac{R^3b^2|\Gamma|}{2\pi} \inf_{z \in \Gamma} \frac{2+2|z|^2}{1+2|z|^2}$ , where  $|\Gamma|$  stands for the length of the curve  $\Gamma$ , and  $d_2 := \frac{R^2|\Gamma|}{2\pi}$ . In the second line, we used the *second resolvent identity*, i.e.  $R(z, A + tB(t))tB(t)R(z, A) = R(z, A) - R(z, A + tB(t))$  for all  $z \in \Gamma$  and  $t \in [0, \epsilon]$ . The third inequality comes from another use of Equation (26). The bound (24) follows from the estimates above.  $\square$

Now, we are ready to prove Theorem 3.3.

*Upper bound on Equation (21):*

**Lemma 6.2** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$  and  $M \in \mathcal{B}(\mathcal{X})$  a contraction with the same assumption as in Theorem 3.3. Then, there is  $\epsilon_1 > 0$  and a  $c_2 > 0$  so that for all  $t \geq 0$  and  $n \in \mathbb{N}$  satisfying  $t \in [0, n\epsilon_1]$*

$$\left\| \left( M e^{\frac{t}{n}\mathcal{L}} \right)^n - \left( P_\Sigma M e^{\frac{t}{n}\mathcal{L}} P_\Sigma \right)^n \right\| \leq c_2 \tilde{\delta}^n + \frac{tb}{n} + \frac{1}{n} \frac{\tilde{\delta}}{1 - \tilde{\delta}} e^{4tbc_2}.$$

*Proof.* As in the proof of Proposition 3.2, Proposition A.1 shows that the uniform power convergence assumption (9), that is  $\|M^n - \sum_{j=1}^J \lambda_j^n P_j\|_\infty \leq \tilde{c} \delta^n$  for all  $n \in \mathbb{N}$ , is equivalent to the spectral gap assumption (q.v. Appx. A),

$$\sigma(M) \subset \mathbb{D}_\delta \cup \{\lambda_1, \dots, \lambda_J\},$$

with corresponding quasinilpotent operators being zero. Therefore, the curve  $\gamma : [0, 2\pi] \rightarrow \mathbb{C}, \varphi \mapsto \tilde{\delta} e^{i\varphi}$ , with  $\tilde{\delta} \in (\delta, 1)$ , encloses the spectrum of  $M^\perp := M P_\Sigma^\perp = P_\Sigma^\perp M$ . By Lemma 2.1


$$\left\| M^\perp e^{s\mathcal{L}} - M^\perp \right\|_\infty = s \left\| M^\perp \mathcal{L} \int_0^1 e^{\tau s \mathcal{L}} d\tau \right\|_\infty \leq s \|P_\Sigma^\perp M \mathcal{L}\|_\infty \leq sb,$$

so that  $M^\perp e^{s\mathcal{L}}$  converges uniformly to  $M^\perp$  for  $s \downarrow 0$ . Hence, Lemma B.2 shows that there exists  $\epsilon_1 > 0$  such that the spectrum of  $M^\perp e^{s\mathcal{L}}$  can be separated by  $\gamma$  for all  $s \in [0, \epsilon_1]$ . Therefore, we can apply the holomorphic functional calculus (Prop. B.1) to conclude that for all  $t \geq 0$  and  $n \in \mathbb{N}$  satisfying  $t \in [0, n\epsilon_1]$

$$\left( M^\perp e^{\frac{t}{n}\mathcal{L}} \right)^k = \frac{1}{2\pi i} \oint_\gamma z^k R(z, M^\perp e^{\frac{t}{n}\mathcal{L}}) dz,$$

where  $k \in \{1, \dots, n\}$ . By Equation (19) and the definition  $c_2 := \sup_{z \in \gamma} \|R(z, M^\perp)\|_\infty \frac{2+2\tilde{\delta}^2}{1+2\tilde{\delta}^2}$ , we achieve for all  $k \in \{1, \dots, n\}$

$$\left\| \left( M^\perp e^{\frac{t}{n}\mathcal{L}} \right)^{k-1} M^\perp \right\|_\infty \leq \frac{1}{2\pi} \oint_\gamma |z|^{k-1} \left\| R(z, M^\perp e^{\frac{t}{n}\mathcal{L}}) \right\|_\infty dz \leq c_2 \tilde{\delta}^k.$$



By the same expansion of  $(Me^{\frac{t}{n}\mathcal{L}})^n = (P_\Sigma Me^{\frac{t}{n}\mathcal{L}} + M^\perp e^{\frac{t}{n}\mathcal{L}})^n$  as in the proof of Proposition 3.2,

$$\left\| \left( Me^{\frac{t}{n}\mathcal{L}} \right)^n - \left( P_\Sigma Me^{\frac{t}{n}\mathcal{L}} \right)^n \right\|_\infty \leq c_2 \tilde{\delta}^n + \frac{1}{n} \frac{\tilde{\delta}}{1 - \tilde{\delta}} e^{4tbc_2}.$$

Finally, Lemma 2.1 shows

$$\left\| P_\Sigma Me^{\frac{t}{n}\mathcal{L}} (\mathbb{1} - P_\Sigma) \right\|_\infty = \frac{t}{n} \left\| P_\Sigma M \mathcal{L} \int_0^1 e^{\tau \frac{t}{n} \mathcal{L}} (\mathbb{1} - P_\Sigma) d\tau \right\|_\infty \leq \frac{tb}{n}$$

so that

$$\left\| \left( Me^{\frac{t}{n}\mathcal{L}} \right)^n - \left( P_\Sigma Me^{\frac{t}{n}\mathcal{L}} P_\Sigma \right)^n \right\| \leq c_2 \tilde{\delta}^n + \frac{tb}{n} + \frac{1}{n} \frac{\tilde{\delta}}{1 - \tilde{\delta}} e^{4tbc_2},$$

which finishes the proof.  $\square$

*Upper bound on Equation (22).*

As in Lemma 5.5, we apply the modified Chernoff Lemma (Lem. 4.2) to upper bound the second term (22). However, our proof strategy includes two crucial improvements compared to Theorem 3 in [2]. Firstly, we show that the spectrum of the perturbed contraction is upper semicontinuous under certain assumptions on  $M$  and the  $C_0$ -semigroup. Therefore, we can use the holomorphic functional calculus and apply the modified Chernoff Lemma with respect to each eigenvalue separately, which allows us to achieve the optimal convergence as in Lemma 5.5.

**Lemma 6.3** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$  and  $M \in \mathcal{B}(\mathcal{X})$  a contraction satisfying the same assumption as in Theorem 3.3. Then,*

$$\left\| \left( P_\Sigma Me^{\frac{t}{n}\mathcal{L}} P_\Sigma \right)^n - \sum_{j=1}^J \lambda_j^n e^{n(C_j(\frac{t}{n}) - P_j(\frac{t}{n}))} P_j(\frac{t}{n}) \right\|_\infty \leq \frac{Jd}{n}$$

where  $C_j(\frac{1}{n}) := \bar{\lambda}_j P_j(\frac{1}{n}) P_\Sigma M e^{\frac{1}{n}\mathcal{L}} P_\Sigma P_j(\frac{1}{n})$  is defined on  $P_j(\frac{1}{n})\mathcal{X}$ , and for some  $d > 0$ .

*Proof.* By Proposition A.1,  $P_\Sigma M = P_\Sigma M P_\Sigma = \sum_{j=1}^J \lambda_j P_j$  so that  $\sigma(P_\Sigma M)$  consists of  $J$  isolated eigenvalues on the unit circle separated by  $\Gamma_j : [0, 2\pi] \rightarrow \mathbb{C}, \phi \mapsto \lambda_j + re^{i\phi}$  with

$$r := \min_{i \neq j} \left\{ \frac{|\lambda_i - \lambda_j|}{3} \right\}. \quad (28)$$

Note that we use the curve interchangeably with its image and denote the formal sum of all curves around the eigenvalues  $\{\lambda_j\}_{j=1}^J$  by  $\Gamma$ . In the following, we define the vector-valued function:


$$s \mapsto P_\Sigma M + sB(s) := P_\Sigma M + s P_\Sigma M \mathcal{L} \int_0^1 e^{s\tau \mathcal{L}} P_\Sigma d\tau = P_\Sigma M e^{s\mathcal{L}} P_\Sigma. \quad (29)$$

By the boundedness of  $M\mathcal{L}$ , the defined vector-valued map converges in the uniform topology to  $P_\Sigma M$ . Moreover, Lemma 2.1 shows that  $s \mapsto B(s)$  is uniformly bounded and continuous in  $s = 0$  because

$$\|B(s) - B(0)\|_\infty = s \left\| P_\Sigma M \mathcal{L} \int_0^1 \int_0^1 \tau_1 e^{\tau_1 \tau_2 s \mathcal{L}} \mathcal{L} P_\Sigma d\tau_2 d\tau_1 \right\|_\infty \leq s \frac{b^2}{2}. \quad (30)$$

Then, we can apply Lemma 6.1 which shows that there exists  $\epsilon_2 > 0$  such that for all  $s \in [0, \epsilon_2]$

$$P_j(s) := \frac{1}{2\pi i} \oint_{\Gamma_j} (z - (P_\Sigma M + sB(s)))^{-1} dz$$



defines the perturbed spectral projection w.r.t.  $\lambda_j$ . Assume  $t \geq 0$  and  $n \in \mathbb{N}$  such that  $t \in [0, n\epsilon_2]$  and redefine  $t\mathcal{L}$  by  $\mathcal{L}$  and  $tb$  by  $b$ . Then, by spectral decomposition

$$\left(P_\Sigma M e^{\frac{1}{n}\mathcal{L}} P_\Sigma\right)^n = \sum_{j=1}^J \left(P_j\left(\frac{1}{n}\right) P_\Sigma M e^{\frac{1}{n}\mathcal{L}} P_\Sigma P_j\left(\frac{1}{n}\right)\right)^n.$$

Thereby, we fix  $j \in \{1, \dots, J\}$  and apply the modified Chernoff lemma (q.v. Lem. 4.2) separately to  $C_j(\frac{1}{n}) := \bar{\lambda}_j P_j(\frac{1}{n}) P_\Sigma M e^{\frac{1}{n}\mathcal{L}} P_\Sigma P_j(\frac{1}{n})$  defined on  $P_j(\frac{1}{n})\mathcal{X}$ . This gives

$$\left\| \left(C_j\left(\frac{1}{n}\right)\right)^n - e^{n(C_j(\frac{1}{n}) - P_j(\frac{1}{n}))} P_j\left(\frac{1}{n}\right) \right\|_\infty \leq \frac{n}{2} \left\| C_j\left(\frac{1}{n}\right) - P_j\left(\frac{1}{n}\right) \right\|_\infty^2.$$

Then, we approximate  $s \mapsto P_j(s)$  via Lemma 6.1 combined with Equation (30):

$$\left\| P_j\left(\frac{1}{n}\right) - P_j - \frac{1}{n} P_j' \right\|_\infty \leq \frac{d_1 + \frac{b^2}{2} d_2}{n^2} =: \frac{c_j}{n^2}. \quad (31)$$

Then, by Lemma 2.1

$$\begin{aligned} \left\| C_j\left(\frac{1}{n}\right) - P_j\left(\frac{1}{n}\right) \right\|_\infty &\leq \frac{1}{n} \left\| \bar{\lambda}_j P_j\left(\frac{1}{n}\right) P_\Sigma M \mathcal{L} \int_0^1 e^{\frac{s}{n}\mathcal{L}} P_\Sigma P_j\left(\frac{1}{n}\right) ds \right\|_\infty \\ &\quad + \left\| \bar{\lambda}_j P_j\left(\frac{1}{n}\right) P_\Sigma M P_j\left(\frac{1}{n}\right) - P_j\left(\frac{1}{n}\right) \right\|_\infty \\ &\leq \frac{b}{n} + \left\| \bar{\lambda}_j \left(P_j\left(\frac{1}{n}\right) - P_j - \frac{1}{n} P_j'\right) P_\Sigma M P_j\left(\frac{1}{n}\right) \right\|_\infty \\ &\quad + \left\| \bar{\lambda}_j \left(P_j + \frac{1}{n} P_j'\right) P_\Sigma M \left(P_j\left(\frac{1}{n}\right) - P_j - \frac{1}{n} P_j'\right) \right\|_\infty \\ &\quad + \left\| \bar{\lambda}_j \left(P_j + \frac{1}{n} P_j'\right) P_\Sigma M \left(P_j + \frac{1}{n} P_j'\right) - P_j - \frac{1}{n} P_j' \right\|_\infty \\ &\quad + \left\| P_j\left(\frac{1}{n}\right) - P_j - \frac{1}{n} P_j' \right\|_\infty \\ &\leq \frac{b}{n} + \frac{c_j}{n^2} (3 + R_j^2 b |\Gamma_j|) + \frac{1}{n^2} (R_j^2 b |\Gamma_j|)^2, \end{aligned} \quad (32)$$

where we used the expression in Lemma 6.1 for  $P_j'$  with  $R_j := \sup_{z \in \Gamma_j} \|R(z, P_\Sigma M)\|_\infty$ . In the inequalities above, we used that the product rule for derivatives applied to  $P_j(s)^2 = P_j(s)$  implies  $P_j' = P_j P_j' + P_j' P_j$  (cf. [25, Lem. 3]). Finally, we square the expression, maximize over all  $j \in \{1, \dots, n\}$ , multiply by  $\frac{n}{2}$ , and redefined  $b$  by  $tb$  and  $\mathcal{L}$  by  $t\mathcal{L}$ .  $\square$

*Upper bound on Equation (23):*

**Lemma 6.4** *Let  $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$  be the generator of a  $C_0$ -contraction semigroup on  $\mathcal{B}(\mathcal{X})$  and  $M \in \mathcal{B}(\mathcal{X})$  a contraction with the same assumption as in Theorem 3.3. Then, there exists a constant  $c_3 \geq 0$  such that*

$$\left\| \sum_{j=1}^J \lambda_j^n e^{n(C_j(\frac{t}{n}) - P_j(\frac{t}{n}))} P_j\left(\frac{t}{n}\right) - \sum_{j=1}^J \lambda_j^n e^{tP_j\mathcal{L}} P_j \right\|_\infty \leq \frac{J}{n} e^{tb} c_3.$$

*Proof.* For ease of notation, we absorb the time parameter  $t$  into the generator  $\mathcal{L}$  and  $b$ . In order to prove the convergence of the generator, we slightly adapt the calculation in Equation (32):

$$\begin{aligned} \left\| n \left(C_j\left(\frac{1}{n}\right) - P_j\left(\frac{1}{n}\right)\right) - P_j \mathcal{L} P_j \right\|_\infty &\leq \left\| \bar{\lambda}_j P_j\left(\frac{1}{n}\right) P_\Sigma M \mathcal{L} \int_0^1 e^{\frac{s}{n}\mathcal{L}} P_\Sigma P_j\left(\frac{1}{n}\right) ds - P_j \mathcal{L} P_j \right\|_\infty \\ &\quad + \frac{c_j}{n} (3 + R_j^2 b |\Gamma_j|) + \frac{1}{n} (R_j^2 b |\Gamma_j|)^2, \end{aligned}$$

Then, we apply Lemma 2.1 and Lemma 6.1 on the first term:

$$\begin{aligned}
 & \left\| \bar{\lambda}_j P_j(\tfrac{1}{n}) P_\Sigma M \mathcal{L} \int_0^1 e^{\frac{\tau_1}{n} \mathcal{L}} P_\Sigma P_j(\tfrac{1}{n}) d\tau_1 - P_j \mathcal{L} P_j \right\|_\infty \\
 & \leq \frac{1}{n} \left\| \bar{\lambda}_j P_j(\tfrac{1}{n}) P_\Sigma M \mathcal{L} \int_0^1 \int_0^1 \tau_1 e^{\frac{\tau_1 \tau_2}{n} \mathcal{L}} \mathcal{L} P_\Sigma P_j(\tfrac{1}{n}) d\tau_2 d\tau_1 \right\|_\infty \\
 & \quad + \left\| \bar{\lambda}_j P_j(\tfrac{1}{n}) P_\Sigma M \mathcal{L} P_\Sigma P_j(\tfrac{1}{n}) - P_j \mathcal{L} P_j \right\|_\infty \\
 & \leq \frac{b^2}{2n} + \left\| \bar{\lambda}_j P_j(\tfrac{1}{n}) P_\Sigma M \mathcal{L} P_\Sigma P_j(\tfrac{1}{n}) - P_j \mathcal{L} P_j \right\|_\infty \\
 & \leq \frac{b^2}{2n} + \frac{1}{n} R_j^2 b^2 \frac{2 + 2(1+r)^2}{1 + 2(1+r)^2} \frac{|\Gamma_j|}{\pi},
 \end{aligned}$$

where we used Equation (24) in the last inequality with  $\|B(t)\|_\infty \leq b$ ,  $r$  is defined in (28) and  $t \mapsto B(t)$  is defined in Equation (29) and satisfies  $\sup_t \|B(t)\|_\infty \leq b$ . In combination with the calculation above, we have found a constant  $d_j > 0$  such that

$$\left\| n(C_j(\tfrac{1}{n}) - P_j(\tfrac{1}{n})) - P_j \mathcal{L} P_j \right\|_\infty \leq \frac{d_j}{n}.$$

Moreover, we have from Lemma 2.2 and Lemma 6.1 that

$$\begin{aligned}
 & \left\| \sum_{j=1}^J \lambda_j^n e^{n(C_j(\frac{1}{n}) - P_j(\frac{1}{n}))} P_j(\tfrac{1}{n}) - \sum_{j=1}^J \lambda_j^n e^{P_j \mathcal{L} P_j} P_j \right\|_\infty \\
 & \leq \left\| \sum_{j=1}^J \lambda_j^n e^{n(C_j(\frac{1}{n}) - P_j(\frac{1}{n}))} P_j(\tfrac{1}{n}) - \sum_{j=1}^J \lambda_j^n e^{P_j \mathcal{L} P_j} P_j(\tfrac{1}{n}) \right\|_\infty + \left\| \sum_{j=1}^J \lambda_j^n e^{P_j \mathcal{L} P_j} (P_j(\tfrac{1}{n}) - P_j) \right\|_\infty \\
 & \leq \frac{J}{n} e^b c_3
 \end{aligned}$$

for some constant  $c_3 \geq 0$ . This proves the lemma after scaling of  $b$  and  $\mathcal{L}$  by  $t$ .  $\square$

*End of the Proof of Theorem 3.3:* Finally, we combine the lemmas in order to finish the proof of Theorem 3.3.

*Proof of Theorem 3.3.* To combine Lemma 6.2, 6.3, and 6.4, we define  $\epsilon := \min\{\epsilon_1, \epsilon_2\}$  where  $\epsilon_1$  is defined in Lemma 6.2 and  $\epsilon_2$  in Lemma 6.3. The result follows after choosing

$$c_1 := tb + \frac{\tilde{\delta}}{1 - \tilde{\delta}} e^{4tbc_2} + Jd + J e^{tb} c_3. \quad (33)$$

$\square$

## 7. EXAMPLES

In this section, we present two classes of examples, which illustrate the range of applicability of our results. In the examples, we denote by  $\rho, \sigma$  quantum states.

*Example 3* (Finite dimensional quantum systems). We choose  $\mathcal{X} = \mathcal{B}(\mathcal{H})$  to be the algebra of linear operators over a finite dimensional Hilbert space  $\mathcal{H}$  endowed with the trace norm  $\|x\|_1 = \text{tr}|x|$ ,  $M : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  a quantum channel, i.e. a completely positive, trace preserving linear map, and  $\mathcal{L}$  the generator of a semigroup of quantum channels over  $\mathcal{B}(\mathcal{H})$ . Here, we further assume the existence of a projection  $P : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{N}$  onto a subalgebra  $\mathcal{N} \subset \mathcal{B}(\mathcal{H})$  with  $MP = PM$  and such that the following *strong data processing inequality* holds for some  $\tilde{\delta} \in (0, 1)$ : for all states  $\rho$  on  $\mathcal{H}$ ,

$$D(M(\rho) \| M \circ P(\rho)) \leq \tilde{\delta} D(\rho \| P(\rho)), \quad (34)$$

where we recall that the relative entropy between two quantum states, i.e. positive, trace-one operators on  $\mathcal{H}$ , is defined as  $D(\rho\|\sigma) := \text{tr}[\rho \log \rho - \rho \log \sigma]$ , whenever  $\text{supp}(\rho) \subseteq \text{supp}(\sigma)$ . Equation (34) was recently shown to hold under a certain detailed balance condition for  $M$  in [14]: there exists a full-rank state  $\sigma$  such that for any two  $x, y \in \mathcal{B}(\mathcal{H})$ ,

$$\text{tr}[\sigma x^* M^*(z)] = \text{tr}[\sigma M^*(x^*)y].$$

Here  $x^*$ , resp.  $M^*$ , denotes the adjoint of  $x$  w.r.t. the inner product on  $\mathcal{H}$ , resp. the adjoint of  $M$  w.r.t. the Hilbert-Schmidt inner product on  $\mathcal{B}(\mathcal{H})$ . In finite dimensions, the quantity  $\sup_{\rho} D(\rho\|P(\rho)) < \infty$  is called the Pimsner-Popa index of  $P$  [27]. Using Pinsker's inequality, we see that the assumption of Theorem 3.3 is satisfied: for all  $x = x^* \in \mathcal{T}_1(\mathcal{H})$  with  $\|x\|_1 \leq 1$  and decomposition  $x = x_+ - x_-$  into positive and negative parts and corresponding states  $\rho_{\pm} = x_{\pm}/\text{tr}[x_{\pm}]$ ,

$$\begin{aligned} \|(M^n - M^n \circ P)(x)\|_1 &\leq \text{tr}[x_+] \|(M^n - M^n \circ P)(\rho_+)\|_1 + \text{tr}[x_-] \|(M^n - M^n \circ P)(\rho_-)\|_1 \\ &= \|x\|_1 \sqrt{2} \max_{\rho \in \{\rho_+, \rho_-\}} D(M^n(\rho)\|M^n \circ P(\rho))^{\frac{1}{2}} \\ &\leq \sqrt{2} \sup_{\rho} D(\rho\|P(\rho))^{\frac{1}{2}} \tilde{\delta}^{\frac{n}{2}} \equiv c \delta^n. \end{aligned}$$

*Example 4* (Infinite dimensional quantum systems and unbounded generators). Here, we pick  $\mathcal{H} = L^2(\mathbb{R})$ , denote by  $I$  the identity operator on  $\mathcal{H}$ , let  $\sigma$  be a quantum state on  $\mathcal{H}$  and  $M$  a generalized depolarizing channel of depolarizing parameter  $p \in (\frac{1}{2}, 1)$  and fixed point  $\sigma$ :

$$M(\rho) := (1 - p)\rho + p \text{tr}(\rho)\sigma. \quad (35)$$

It is clear by convexity that  $M$  satisfies the uniform strong power convergence with projection  $P(\rho) = \text{tr}(\rho)\sigma$  and parameter  $\delta = 2(1 - p) < 1$ . We also assume that  $\sigma \in \mathcal{D}(\mathcal{L})$ . Then, we have that for all  $t \geq 0$ :

$$\|(\mathbb{1} - P)e^{t\mathcal{L}}P\|_{1 \rightarrow 1} = \sup_{\|x\|_1 \leq 1} |\text{tr}(x)| \|e^{t\mathcal{L}}(\sigma) - \sigma\|_1 \leq t \|\mathcal{L}(\sigma)\|_1. \quad (36)$$

Moreover, by duality and the unitality of the maps  $e^{t\mathcal{L}^*}$  we have that

$$\|Pe^{t\mathcal{L}}(\mathbb{1} - P)\|_{1 \rightarrow 1} = \|(\mathbb{1} - P)^*e^{t\mathcal{L}^*}P^*\|_{\infty \rightarrow \infty} = \sup_{\|y\| \leq 1} |\text{tr}(\sigma y)| \|(\mathbb{1} - P)(I)\| = 0, \quad (37)$$

Therefore, the assumptions of Theorem 3.1 are satisfied and we find the convergence rate  $\mathcal{O}(n^{-1})$ . Interestingly, this answers a conjecture of [2, Ex. 3, 5] for the Hamiltonian evolution generated by the one-dimensional harmonic oscillator. There, the authors had numerically guessed the optimal rate which we prove here. However their analytic bounds could only provide a decay of order  $\mathcal{O}(n^{-\frac{2}{3}})$  and for a restriction of  $H$  to a finite dimensional stable subspace, which effectively assumed the boundedness of the generator. As illustrated here, our asymptotic Zeno condition is easily verifiable and provides a much richer class of examples.

## 8. DISCUSSION AND OPEN QUESTIONS

In this paper, we proved the optimal convergence rate of the quantum Zeno effect in two results: Theorem 3.1 focuses on weakening the assumptions of the  $C_0$ -semigroup to the so-called asymptotic Zeno condition. Hence, Theorem 3.1 allows strongly continuous Zeno dynamics which is novel for open systems. In Theorem 3.3 instead, we weaken the assumption on  $M$  to the uniform power convergence as in [2, Thm. 3]. Additionally, we presented an example which shows the optimality of the achieved convergence rate. This brings up the question whether our assumptions are optimal and how the assumption on the contraction correlates with the assumption on the  $C_0$ -semigroup. For example, is it possible to weaken the uniform power convergence in Theorem 3.1 or Proposition 3.2 to finitely many eigenvalues on the unit circle without assuming stronger assumption on the semigroup? Following our proof strategy (q.v. Lem. 5.2), this is related to

the conjecture of a generalized version of *Trotter's product formula for finitely many projections* under certain assumptions on the generator,

$$\left\| \left( \sum_{j=1}^J \lambda_j P_j e^{\frac{1}{n} \mathcal{L}} \right)^n x - \sum_{j=1}^J \left( \lambda_j P_j e^{\frac{1}{n} \mathcal{L}} \right)^n x \right\| \stackrel{?}{=} \mathcal{O} \left( \frac{1}{n} \|x\| \right),$$

On a related note, it would be interesting to see if Lemma 4.2 can be used to improve the convergence rate of *Trotter's product formula* in the strong topology.

Another line of generalization, which is not discussed in this paper, would be to weaken the assumption on  $M$  to the strong topology as in Theorem 2 in [2]. There, the authors assume that  $M^n$  converges to  $P$  in the strong topology and that the semigroup is uniformly continuous. It would be interesting to know whether an extension to  $C_0$ -semigroups is possible. Finally, another important line of generalization would be to extend our results to time-dependent semigroups as in [25].

*Acknowledgments:* We would like to thank Michael Wolf and Markus Hasenöhrl for their support on this project. Moreover, we would like to thank Valentin A. Zagrebnoy for his helping out with questions on the Chernoff Lemma. T.M. and C.R. acknowledge the support of the Munich Center for Quantum Sciences and Technology, and C.R. that of the Humboldt Foundation.

## 9. REFERENCES

- [1] Norbert Barankai and Zoltán Zimborás. *Generalized quantum Zeno dynamics and ergodic means*. 2018. URL: <https://arxiv.org/pdf/1811.02509>.
- [2] Simon Becker, Nilanjana Datta and Robert Salzmänn. ‘Quantum Zeno Effect in Open Quantum Systems’. In: *Annales Henri Poincaré* 22.11 (2021), pp. 3795–3840. ISSN: 1424-0661. DOI: [10.1007/s00023-021-01075-8](https://doi.org/10.1007/s00023-021-01075-8).
- [3] Beige, Braun, Tregenna and Knight. ‘Quantum computing using dissipation to remain in a decoherence-free subspace’. In: *Physical review letters* 85.8 (2000), pp. 1762–1765. ISSN: 0031-9007. DOI: [10.1103/PhysRevLett.85.1762](https://doi.org/10.1103/PhysRevLett.85.1762).
- [4] Daniel Burgarth, Paolo Facchi, Hiromichi Nakazato, Saverio Pascazio and Kazuya Yuasa. ‘Quantum Zeno Dynamics from General Quantum Operations’. In: *Quantum* 4 (2020), p. 289. DOI: [10.22331/q-2020-07-06-289](https://doi.org/10.22331/q-2020-07-06-289).
- [5] Paul R. Chernoff. ‘Note on product formulas for operator semigroups’. In: *Journal of Functional Analysis* 2 (1968), pp. 238–242. ISSN: 00221236. DOI: [10.1016/0022-1236\(68\)90020-7](https://doi.org/10.1016/0022-1236(68)90020-7).
- [6] Jason M. Dominy, Gerardo A. Paz-Silva, A. T. Rezakhani and D. A. Lidar. ‘Analysis of the quantum Zeno effect for quantum control and computation’. In: *Journal of Physics A: Mathematical and Theoretical* 46.7 (2013). ISSN: 1751-8121. DOI: [10.1088/1751-8113/46/7/075306](https://doi.org/10.1088/1751-8113/46/7/075306).
- [7] Klaus-Jochen Engel and R. Nagel. *One-parameter semigroups for linear evolution equations*. Vol. 194. Graduate texts in mathematics. New York and London: Springer, 2000. DOI: [10.1007/b97696](https://doi.org/10.1007/b97696).
- [8] Noam Erez, Yakir Aharonov, Benni Reznik and Lev Vaidman. ‘Correcting quantum errors with the Zeno effect’. In: *Physical review. A, Atomic, molecular, and optical physics* 69.6 (2004). ISSN: 1050-2947. DOI: [10.1103/PhysRevA.69.062315](https://doi.org/10.1103/PhysRevA.69.062315).
- [9] Pavel Exner and Takashi Ichinose. ‘Note on a Product Formula Related to Quantum Zeno Dynamics’. In: *Annales Henri Poincaré* 22.5 (2021), pp. 1669–1697. ISSN: 1424-0661. DOI: [10.1007/s00023-020-01014-z](https://doi.org/10.1007/s00023-020-01014-z).
- [10] Paolo Facchi, Daniel A. Lidar and Saverio Pascazio. ‘Unification of dynamical decoupling and the quantum Zeno effect’. In: *Physical review. A, Atomic, molecular, and optical physics* 69.3 (2004). ISSN: 1050-2947. DOI: [10.1103/PhysRevA.69.032314](https://doi.org/10.1103/PhysRevA.69.032314).

- [11] Paolo Facchi and Saverio Pascazio. ‘Quantum Zeno dynamics: mathematical and physical aspects’. In: *Journal of Physics A: Mathematical and Theoretical* 41.49 (2008). ISSN: 1751-8121. DOI: [10.1088/1751-8113/41/49/493001](https://doi.org/10.1088/1751-8113/41/49/493001).
- [12] Martin C. Fischer, Braulio Gutiérrez-Medina and Mark G. Raizen. ‘Observation of the quantum zeno and anti-zeno effects in an unstable system’. In: *Physical review letters* 87.4 (2001). ISSN: 0031-9007. DOI: [10.1103/PhysRevLett.87.040402](https://doi.org/10.1103/PhysRevLett.87.040402).
- [13] James D. Franson, Bart C. Jacobs and Todd B. Pittman. ‘Quantum computing using single photons and the Zeno effect’. In: *Physical review. A, Atomic, molecular, and optical physics* 70.6 (2004). ISSN: 1050-2947. DOI: [10.1103/PhysRevA.70.062302](https://doi.org/10.1103/PhysRevA.70.062302).
- [14] Li Gao and Cambyse Rouzé. ‘Complete entropic inequalities for quantum Markov chains’. In: *arXiv preprint arXiv:2102.04146* (2021).
- [15] Alexander Gomilko and Yuri Tomilov. ‘On convergence rates in approximation theory for operator semigroups’. In: *Journal of Functional Analysis* 266.5 (2014), pp. 3040–3082. ISSN: 00221236. DOI: [10.1016/j.jfa.2013.11.012](https://doi.org/10.1016/j.jfa.2013.11.012).
- [16] Brian C. Hall. *Lie groups, Lie algebras, and Representations: An Elementary Introduction*. Second edition. Vol. 222. Graduate texts in mathematics. Springer, 2015. ISBN: 978-3-319-13466-6. DOI: [10.1007/978-3-319-13467-3](https://doi.org/10.1007/978-3-319-13467-3).
- [17] Einar Hille and Ralph S. Phillips. *Functional analysis and Semi-Groups*. Rev. and expanded ed. Vol. vol. 31. Colloquium publications (American Mathematical Society). Providence: American Mathematical Society, 2000. ISBN: 9780821810316. DOI: [10.1090/coll/031](https://doi.org/10.1090/coll/031).
- [18] Onur Hosten, Matthew T. Rakher, Julio T. Barreiro, Nicholas A. Peters and Paul G. Kwiat. ‘Counterfactual quantum computation through quantum interrogation’. In: *Nature* 439.7079 (2006), pp. 949–952. ISSN: 1476-4687. DOI: [10.1038/nature04523](https://doi.org/10.1038/nature04523).
- [19] Wayne M. Itano, Daniel J. Heinzen, John J. Bollinger and David J. Wineland. ‘Quantum Zeno effect’. In: *Physical review. A, Atomic, molecular, and optical physics* 41.5 (1990), pp. 2295–2300. ISSN: 1050-2947. DOI: [10.1103/physreva.41.2295](https://doi.org/10.1103/physreva.41.2295).
- [20] Toshio Katō. *Perturbation Theory for Linear Operators*. 2nd ed. Vol. 132. A Series of Comprehensive Studies in Mathematics. Springer International Publishing, 1995. ISBN: 9783642662829. DOI: [10.1007/978-3-642-66282-9](https://doi.org/10.1007/978-3-642-66282-9).
- [21] Erwin Kreyszig. *Introductory Functional Analysis with Applications*. New York: Wiley classics library, 1989. ISBN: 0471504599.
- [22] Wei Liu and Michael Röckner. *Stochastic Partial Differential Equations: An Introduction*. Springer, 2015. ISBN: 9783319223544.
- [23] Ilia A. Luchnikov and Sergey N. Filippov. ‘Quantum evolution in the stroboscopic limit of repeated measurements’. In: *Physical Review A* 95.2 (2017). DOI: [10.1103/PhysRevA.95.022113](https://doi.org/10.1103/PhysRevA.95.022113).
- [24] Baidyanath Misra and George Sudarshan. ‘The Zeno’s paradox in quantum theory’. In: *Journal of Mathematical Physics* 18.4 (1977), pp. 756–763. ISSN: 0022-2488. DOI: [10.1063/1.523304](https://doi.org/10.1063/1.523304).
- [25] Tim Möbus and Michael M. Wolf. ‘Quantum Zeno effect generalized’. In: *Journal of Mathematical Physics* 60.5 (2019). ISSN: 0022-2488. DOI: [10.1063/1.5090912](https://doi.org/10.1063/1.5090912).
- [26] Hagen Neidhardt, Artur Stephan and Valentin A. Zagrebnov. ‘Operator-Norm Convergence of the Trotter Product Formula on Hilbert and Banach Spaces: A Short Survey’. In: *Current research in nonlinear analysis*. Ed. by Themistocles M. Rassias, H. Brézis and L. Nirenberg. Vol. 135. Springer, 2018, pp. 229–247. DOI: [10.1007/978-3-319-89800-1\\_{\text{underscore}}9](https://doi.org/10.1007/978-3-319-89800-1_{\text{underscore}}9).
- [27] Mihai Pimsner and Sorin Popa. ‘Entropy and index for subfactors’. In: *Annales scientifiques de l’Ecole normale supérieure*. Vol. 19. 1. 1986, pp. 57–106.



- [28] Andreas U. Schmidt. ‘Mathematics of the Quantum Zeno Effect’. In: *Mathematical physics research on the leading edge*. Ed. by Charles V. Benton. Nova Science Publishers, 2004, pp. 113–143. ISBN: 1590339053.
- [29] Barry Simon. *Operator theory*. Vol. 4. A comprehensive course in analysis. Providence, Rhode Island: American Mathematical Society, 2015. ISBN: 9781470411039.
- [30] Richard P. Stanley. *Enumerative combinatorics*. Boston, MA: Springer, Boston, MA, 1986. ISBN: 9781461597636. DOI: [10.1007/978-1-4615-9763-6](https://doi.org/10.1007/978-1-4615-9763-6).
- [31] Valentin Zagrebnov. ‘Comments on the Chernoff  $\sqrt{n}$ -Lemma’. In: *Functional analysis and operator theory for quantum physics*. Ed. by J. Dittrich, Hynek Kovařík and Ari Laptev. EMS series of congress reports. Zurich Switzerland: European Mathematical Society, 2017. ISBN: 978-3-03719-175-0.

*Email address:* [tim.moebus@tum.de](mailto:tim.moebus@tum.de), [rouzecambyse@gmail.com](mailto:rouzecambyse@gmail.com)

<sup>1</sup> DEPARTMENT OF MATHEMATICS, TECHNICAL UNIVERSITY OF MUNICH

<sup>2</sup> MUNICH CENTER FOR QUANTUM SCIENCE AND TECHNOLOGY (MCQST), MÜNCHEN, GERMANY

## APPENDIX A. SPECTRAL GAP ASSUMPTION

In the proofs of Proposition 3.2 and Theorem 3.3, we use the equivalence [2, Prop. 3.1] of the uniform power convergence and the spectral gap assumption with corresponding quasinilpotent operators being zero, that is there are eigenvalues  $\{\lambda_j\}_{j=1}^J \subset \partial\mathbb{D}_1$  and a gap  $\delta \in (0, 1)$  so that

$$\sigma(M) \subset \mathbb{D}_\delta \cup \{\lambda_j\}_{j=1}^J.$$

Moreover, the quasinilpotent operators are assumed to be zero, that is

$$N_j = \frac{1}{2\pi i} \oint_{\Gamma_j} (z - \lambda_j) R(z, M) dz = 0$$

for  $\Gamma_j : [0, 2\pi) \ni \varphi \mapsto \lambda_j + \frac{1}{2} \min_{k,l \in \{1, \dots, J\}} \{|\lambda_k - \lambda_l|, 1 - \delta\} e^{i\varphi}$  (q.v. Eq. (3)). Additionally, the projections defined in the uniform power convergence are equal to the spectral projections

$$P_j = \frac{1}{2\pi i} \oint_{\Gamma_{\lambda_j}} R(z, M) dz$$

for all  $j \in \{1, \dots, J\}$  (q.v. Eq. (2)). Since  $N_j = 0$ ,  $P_j M = M P_j = \lambda_j P_j$  [29, Thm. 2.3.5]. The equivalence between the uniform power convergence and the spectral gap assumption with corresponding quasinilpotent operators being zero is given in [2, Prop. 3.1].

**Proposition A.1** ([2, Prop. 3.1]) *Let  $M \in \mathcal{B}(\mathcal{X})$  be a contraction,  $J \in \mathbb{N}$ ,  $\delta \in (0, 1)$ , and  $\{\lambda_j\}_{j=1}^J \subset \partial\mathbb{D}_1$ . Then, the following statements are equivalent:*

- (1) *The contraction  $M$  satisfies the spectral gap assumption w.r.t.  $\{\lambda_j\}_{j=1}^J$ , the gap  $\delta$ , and the corresponding quasinilpotent operators being zero.*
- (2) *There are projections  $\{P_j\}_{j=1}^J$ , a  $\delta \in (0, 1)$  and  $\tilde{c} \geq 0$  such that  $P_j M = P_j$  for all  $j = \{1, \dots, J\}$  and*

$$\left\| M^n - \sum_{j=1}^J \lambda_j^n P_j \right\|_\infty \leq \tilde{c} \delta^n.$$

- (3) *There are contractions  $\{C_j\}_{j=1}^J$  so that  $\lim_{n \rightarrow \infty} \|M^n - \sum_{j=1}^J \lambda_j^n C_j\|_\infty = 0$ .*

## APPENDIX B. HOLOMORPHIC FUNCTIONAL CALCULUS AND SEMICONTINUOUS SPECTRA

In this section, we introduce the holomorphic functional calculus, consider some continuity properties of the spectrum of perturbed operators, and finally prove an approximation result for perturbed spectral projections. Let's start with the holomorphic functional: To be precise, the holomorphic functional calculus defines  $f(A)$  for  $A \in \mathcal{B}(\mathcal{X})$  and a function  $f : \mathbb{C} \rightarrow \mathbb{C}$  holomorphic on a neighbourhood of  $\sigma(A)$ . One advantage, for example, is that if the spectrum is separated into two parts, then the theorem shows that the functions reduced to the separated subsets can be treated "independently". This is used to define the spectral projection. More details on this topic can be found in Section 2.3 in [29]. We use the holomorphic functional calculus to introduce the integral representation of  $f(A)$  for a holomorphic function  $f$ , and the spectral projection:

**Proposition B.1** (Holomorphic Functional Calculus [29, Thm. 2.3.1-3]) *Let  $A \in \mathcal{B}(\mathcal{X})$ ,  $\Gamma : [0, 2\pi] \rightarrow \mathbb{C}$  be a curve around  $\sigma(A)$ , and  $f$  be a function holomorphic in the neighbourhood of  $\sigma(A)$ . Then,*

$$f(A) = \frac{1}{2\pi i} \oint_{\Gamma} f(z) R(z, A) dz$$

*and independent of  $\Gamma$ . If a subset  $\tilde{\sigma}$  of the spectrum is separated by a curve  $\Gamma$ , the spectral projection w.r.t. the enclosed subset is*

$$P := \frac{1}{2\pi i} \oint_{\Gamma} R(z, A) dz,$$

and satisfies  $P^2 = P$  and  $PA = AP$ . If  $\tilde{\sigma} = \{\lambda_0\}$  is an isolated eigenvalue, the associated nilpotent operator is

$$N := \frac{1}{2\pi i} \oint_{\Gamma} (z - \lambda_0) R(z, A) dz,$$

so that  $NP = PN = N$  and  $AP = \lambda_0 P + N$ .

In our case,  $M$  has finitely many isolated eigenvalues on the unit circle. However, if the semigroup perturbs those eigenvalues, the following lemma states a conditions under which the separation of the spectrum maintain.

**Lemma B.2** (Semicontinuous Spectra [20, Sec. IV.3.1]) *Let  $b \geq 0$  and  $M : \mathbb{R}_{\geq 0} \rightarrow \mathcal{B}(\mathcal{X})$  be a vector-valued function which satisfies*

$$\|M(t) - M(0)\|_{\infty} \leq tb.$$

*Moreover, assume that  $\sigma(M(0)) = \sigma_1 \cup \sigma_2$  is a disjoint union separated by a curve  $\gamma : [0, 2\pi] \rightarrow \rho(M(0))$ . For  $\epsilon > 0$  satisfying*

$$\epsilon < \frac{1}{2b} \inf_{z \in \gamma} (1 + |z|^2)^{-1} \left( 1 + \sup_{z \in \gamma} \|R(z, M(0))\|_{\infty}^2 \right)^{-\frac{1}{2}}, \quad (38)$$

*the spectrum of  $\sigma(M(t))$  is separated by  $\gamma$  for all  $t \in [0, \epsilon]$ .*

*Proof.* Firstly, we show that Equation (38) is well-defined — the value  $\epsilon$  exists. The compactness of the image of  $\gamma$  and the continuity of the resolvent [20, Thm. II.1.5] show that  $\sup_{z \in \gamma} \|R(z, M(0))\|_{\infty} < \infty$  and  $\epsilon > 0$  exists. Then, we can apply Theorem 3.16 from [20, Sec. IV], which proves the lemma.  $\square$

*Remark.* It is possible, to generalize the lemma above to uniformly continuous vector-valued maps. However, we would loose the explicit choice of  $\epsilon$ .