

# A propagation of chaos result for a class of mean-field reflected BSDEs with jumps

Boualem Djehiche <sup>\*</sup>      Roxana Dumitrescu <sup>†</sup>      Jia Zeng<sup>‡</sup>

November 30, 2021

## Abstract

In this article, we establish the propagation of chaos property for weakly interacting nonlinear Snell envelopes which converge to a class of mean-field reflected backward stochastic differential equations (BSDEs) with jumps, where the mean-field interaction in terms of the distribution of the  $Y$ -component of the solution enters both the driver and the lower obstacle. Under mild Lipschitz and integrability conditions on the coefficients, we prove existence and uniqueness of the solution to both the mean-field reflected BSDEs with jumps and the corresponding system of weakly interacting particles and provide the propagation of chaos result for the whole solution  $(Y, Z, U, K)$ .

*Keywords:* Mean-field, Backward SDEs with jumps, Snell envelope, Interacting particle system, Propagation of chaos

*2010 Mathematics Subject Classification:* 60H10, 60H07, 49N90

## 1 Introduction

The notion of propagation of chaos can be traced back at least to the work by Kac [31] whose initial motivation was to investigate the particle system approximation of some nonlocal partial differential equations (PDEs) arising in thermodynamics. He studied the evolution of a large number  $n$  of random particles starting from  $n$  given independent and identically distributed random variables, i.e. the initial configuration is chaotic, and whose respective dynamics weakly interact. Kac's insight is that the initial chaotic configuration propagates over time. Precise mathematical treatments of this theory can be found in McKean [33], Sznitman [40] and Gärtner [19]. For further development and applications of propagation of chaos theory, see e.g. Jabin [23], Lacker [27], Shkolnikov [39] and references therein. More recently, due to the very active development within mean-field type control and games (see Lasry

---

<sup>\*</sup>Department of Mathematics, KTH Royal Institute of Technology, Stockholm, Sweden, email: [boualem@kth.se](mailto:boualem@kth.se)

<sup>†</sup>Department of Mathematics, King's College London, United Kingdom, email: [roxana.dumitrescu@kcl.ac.uk](mailto:roxana.dumitrescu@kcl.ac.uk)

<sup>‡</sup>Department of Mathematics, King's College London, United Kingdom and The University of Hong Kong, Hong Kong, email: [jia.zeng@kcl.ac.uk](mailto:jia.zeng@kcl.ac.uk)

and Lions [29], Andersson and Djehiche [1]), there is a renewed interest to study propagation of chaos for stochastic systems with a view towards optimal control and games.

A question is to ask whether propagation of chaos carries over to systems of particles with chaotic terminal configuration. To the best of our knowledge, only the papers of Buckdahn et al. [8], Hu et al. [22], Laurière and Tangpi [28] and Briand et al. [5] studied limit results for interacting backward particles. All these papers about backward propagation of chaos only considered the case of non-reflected BSDE driven by Brownian motion. Similar results were obtained in Briand and Hibon [6] for a specific class of mean-reflected BSDEs where the constraint acts on the law of the  $Y$ -component rather than on its path.

The objective of the present paper is to establish the propagation of chaos result for a class of mean-field reflected backward stochastic differential equations (BSDEs) with jumps having the particularity that the obstacle process might depend on the solution too. BSDEs have been widely studied since the pioneering work by Pardoux and Peng [34]. In particular, they appear as a powerful tool to solve stochastic optimization problems in many fields such as engineering, investment science including mathematical finance, game theory and insurance, among many other areas. In the Markovian case, BSDEs provide a probabilistic representation of a class of semilinear PDEs. Later Tang and Li [41] considered the more involved case when the noise is driven not only by a Brownian motion but also by an independent Poisson random measure. They proved existence and uniqueness of the solution. Barles et al. [3] studied the link of those BSDEs with viscosity solutions of integral-partial differential equations. The notion of reflected BSDEs was first introduced in El Karoui et al. [17]. In the setting of those BSDEs, one of the components of the solution is forced to stay above a given process. The extension to the case of reflected BSDEs with jumps can be found in e.g. [13], [14], [15], [16], [18], [20], [21].

The interaction of particles is through the empirical distribution of the system. In the main contributions of the paper we derive convergence results of the  $n$ -particle system to solutions of mean-field BSDEs under mild Lipschitz and integrability conditions on the coefficients, as well as related propagation of chaos results. Mean-field BSDEs have been introduced in Buckdahn et al. [7] and motivated by their connection to control of McKean-Vlasov equation or mean-field games, see Carmona et al. [9] and Acciaio et al. [2] for the discussions of their connections. Later, the addition of a reflection to these BSDEs has been considered in Li [30] and Djehiche et al. [12]. In particular, motivated by the setting of dynamical solvency level in life insurance and pension, Djehiche et al. [12] studied a new class of mean-field type reflected BSDEs, where the mean-field interaction in terms of the distribution of the  $Y$ -component of the solution enters both the driver and the lower obstacle. They obtained the well-posedness of such a class of equations under mild Lipschitz, integrability conditions on the coefficients, and monotonicity assumptions on the driver and the obstacle, using the classical penalization scheme.

To the best of our knowledge, this is the first paper which establishes the well-posedness and the propagation of chaos property of the mean-field reflected BSDEs with jumps by constructing a system of weakly interacting nonlinear Snell envelopes in a general setting. In order to deal with the possible dependence of the obstacle on the solution, one needs to develop specific techniques. In particular,

using a new method based on the characterization of the solution of the mean-field reflected BSDE as a nonlinear optimal stopping problem of mean-field type, the existence and uniqueness are proved without further assumptions of monotone driver and obstacle with respect to the measure component, which represents a major improvement compared to Djehiche et al. [12]. We also establish a sufficient condition on the type of dependence of the obstacle on the solution which ensures the continuity of the increasing predictable process, which pushes the solution above the obstacle in a minimal way. Furthermore, we provide a particle system approximation in terms of a system of weakly interacting Snell envelopes and show its well-posedness. In order to show the convergence of the weakly interacting system of particles, we provide a law of large numbers type result for independent copies of the  $Y$ -component of the solution to the mean-field reflected BSDE with jumps, which seems new in the literature and relies on a careful study of the Skorohod space of right continuous functions with left limits.

The paper is organized as follows. In Section 2, we prove the existence and uniqueness of the mean-field reflected BSDEs with jumps, where the mean-field interaction in terms of the distribution of the  $Y$ -component of the solution enters both the driver and the lower obstacle. The well-posedness of the mean-field reflected BSDE with jumps is obtained by extending the notion of Peng's nonlinear expectation (e.g. Peng [36]). In Section 3, we discuss the interpretation of the mean-field reflected BSDEs with jumps at the particle level and show the existence and uniqueness of the system of weakly interacting particles. In Section 4, we give a law of large numbers result for the empirical measure associated with  $\mathbb{D} := D([0, T], \mathbb{R})$ -valued independent copies of the  $Y$ -component of the solution to the mean-field reflected BSDE with jumps w.r.t. the  $p$ -Wasserstein distance. With this law of large numbers, we prove the convergence of the particle system to solutions of mean-field reflected BSDEs with jumps, and obtain a related propagation of chaos result for the solution  $(Y, Z, U, K)$ .

**Notation.** Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a complete probability space.  $B = (B_t)_{0 \leq t \leq T}$  is a standard one dimensional Brownian motion and  $N(dt, de)$  is a Poisson random measure, independent of  $B$ , with compensator  $\nu(de)dt$  such that  $\nu$  is a  $\sigma$ -finite measure on  $\mathbb{R}^*$  equipped with its Borel  $\sigma$ -field  $\mathcal{B}(\mathbb{R}^*)$ . Let  $\tilde{N}(dt, du)$  be its compensated process. We denote by  $\mathbb{F} = \{\mathcal{F}_t\}$  the (completed) natural filtration associated with  $B$  and  $N$ . Let  $\mathcal{P}$  be the  $\sigma$ -algebra on  $\Omega \times [0, T]$  of  $\mathcal{F}_t$ -progressively measurable sets. Next, we introduce the following spaces, for  $p \geq 1$  and  $\beta \geq 0$ :

- $\mathcal{T}_t$  is the set of  $\mathbb{F}$ -stopping times  $\tau$  such that  $\tau \in [t, T]$  a.s.
- $L^p(\mathcal{F}_T)$  is the set of random variables  $\xi$  which are  $\mathcal{F}_T$ -measurable and square integrable.
- $\mathcal{S}_\beta^p$  is the set of real-valued càdlàg adapted processes  $y$  such that  $\|y\|_{\mathcal{S}_\beta^p}^p := \mathbb{E}[\sup_{0 \leq u \leq T} e^{\beta ps} |y_u|^p] < \infty$ . We set  $\mathcal{S}^p = \mathcal{S}_0^p$ .
- $\mathcal{S}_{\beta, i}^p$  is the subset of  $\mathcal{S}_\beta^p$  such that the process  $k$  is non-decreasing and  $k_0 = 0$ . We set  $\mathcal{S}_i^p = \mathcal{S}_{0, i}^p$ .
- $\mathbb{L}_\beta^p$  is the set of real-valued càdlàg adapted processes  $y$  such that  $\|y\|_{\mathbb{L}_\beta^p}^p := \sup_{\tau \in \mathcal{T}_0} \mathbb{E}[e^{\beta p \tau} |y_\tau|^p] < \infty$ .  $\mathbb{L}_\beta^p$  is a Banach space (see Theorem 22 in [11], pp. 60 when  $p = 1$ ). We set  $\mathbb{L}^p = \mathbb{L}_0^p$ .

- $\mathcal{H}^{p,d}$  is the set of  $\mathcal{P}$ -measurable,  $\mathbb{R}^d$ -valued processes such that  $\mathbb{E}[(\int_0^T |v_s|^2 ds)^{p/2}] < \infty$ .
- $L_\nu^p$  is the set of measurable functions  $l : \mathbb{R}^* \rightarrow \mathbb{R}$  such that  $\int_{\mathbb{R}^*} |l(u)|^p \nu(du) < +\infty$ . The set  $L_\nu^2$  is a Hilbert space equipped with the scalar product  $\langle \delta, l \rangle_\nu := \int_{\mathbb{R}^*} \delta(u) l(u) \nu(du)$  for all  $(\delta, l) \in L_\nu^2 \times L_\nu^2$ , and the norm  $\|l\|_{\nu,2} := (\int_{\mathbb{R}^*} |l(u)|^2 \nu(du))^{1/2}$ . If there is no risk for confusion, we sometimes denote  $\|l\|_{\nu,2} := \|l\|_\nu$ .
- $\mathcal{B}(\mathbb{R}^d)$  (resp.  $\mathcal{B}(L_\nu^p)$ ) is the Borel  $\sigma$ -algebra on  $\mathbb{R}^d$  (resp. on  $L_\nu^p$ ).
- $\mathcal{H}_\nu^{p,d}$  is the set of predictable processes  $l$ , i.e. measurable

$$l : ([0, T] \times \Omega \times \mathbb{R}^*, \mathcal{P} \otimes \mathcal{B}(\mathbb{R}^*)) \rightarrow (\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d)); \quad (\omega, t, u) \mapsto l_t(\omega, u)$$

such that  $\|l\|_{\mathcal{H}_\nu^{p,d}}^p := \mathbb{E} \left[ \left( \int_0^T \sum_{j=1}^d |l_t^j|^2 dt \right)^{\frac{p}{2}} \right] < \infty$ . For  $d = 1$ , we denote  $\mathcal{H}_\nu^{p,1} := \mathcal{H}_\nu^p$ .

- $\mathcal{P}_p(\mathbb{R})$  is the set of probability measures on  $\mathbb{R}$  with finite  $p$ th moment. We equip the space  $\mathcal{P}_p(\mathbb{R})$  with the  $p$ -Wasserstein distance denoted by  $\mathcal{W}_p$  and defined as

$$\mathcal{W}_p(\mu, \nu) := \inf \left\{ \int_{\mathbb{R} \times \mathbb{R}} |x - y|^p \pi(dx, dy) \right\}^{1/p},$$

where the infimum is over probability measures  $\pi \in \mathcal{P}_p(\mathbb{R} \times \mathbb{R})$  with first and second marginals  $\mu$  and  $\nu$ , respectively.

## 2 Existence and uniqueness of the solution of mean-field reflected BSDE with jumps

In this section, we study the well-posedness of a new class of reflected BSDE of mean-field type with jumps, which have the additional feature that the obstacle might depend on the solution too. We start with the following definition.

**Definition 2.1.** *We say that the quadruple of progressively measurable processes  $(Y_t, Z_t, U_t, K_t)_{t \leq T}$  is a solution of the mean-field reflected BSDE associated with  $(f, \xi, h)$  if*

$$\begin{cases} Y \in \mathcal{S}^p, Z \in \mathcal{H}^{p,1}, U \in \mathcal{H}_\nu^p \text{ and } K \in \mathcal{S}_i^p \text{ and predictable,} \\ Y_t = \xi + \int_t^T f(s, Y_s, Z_s, U_s, \mathbb{P}_{Y_s}) ds + K_T - K_t - \int_t^T Z_s dB_s - \int_t^T \int_{\mathbb{R}^*} U_s(e) \tilde{N}(ds, de), \quad t \in [0, T], \\ Y_t \geq h(t, Y_t, \mathbb{P}_{Y_t}), \quad t \in [0, T], \\ \int_0^T (Y_{t-} - h(t^-, Y_{t-}, \mathbb{P}_{Y_{t-}})) dK_t = 0. \quad (\text{Skorohod's flatness condition}) \end{cases} \quad (2.1)$$

**Remark 2.1.** *The flatness condition is equivalent to the following condition: if  $K = K^c + K^d$ , where  $K^c$  (resp.  $K^d$ ) represents the continuous (resp. the discontinuous) part of  $K$ , then*

$$\begin{aligned} & \int_0^T (Y_t - h(t, Y_t, \mathbb{P}_{Y_t})) dK_t^c = 0, \text{ and } \forall \tau \in \mathcal{T}_T \text{ predictable,} \\ & \Delta Y_\tau = Y_\tau - Y_{\tau-} = -\Delta K_\tau^d = -(h(\tau^-, Y_{\tau-}, \mathbb{P}_{Y_{\tau-}|t=\tau}) - Y_\tau)^+ \mathbf{1}_{\{Y_{\tau-} = h(\tau^-, Y_{\tau-}, \mathbb{P}_{Y_{\tau-}|t=\tau})\}}. \end{aligned}$$

Throughout the paper, we assume  $p \geq 2$  (see Remark 4.3 below about a possible extension to  $p \in (1, 2)$ ) and make the following assumption on  $(f, h, \xi)$ .

**Assumption 2.1.** *The coefficients  $f, h$  and  $\xi$  satisfy*

(i)  *$f$  is a mapping from  $[0, T] \times \Omega \times \mathbb{R} \times \mathbb{R} \times L_\nu^p \times \mathcal{P}_p(\mathbb{R})$  into  $\mathbb{R}$  such that*

(a) *the process  $(f(t, 0, 0, 0, \delta_0))_{t \leq T}$  is  $\mathcal{P}$ -measurable and belongs to  $\mathcal{H}^{p,1}$ ;*

(b)  *$f$  is Lipschitz w.r.t.  $(y, z, u, \mu)$  uniformly in  $(t, \omega)$ , i.e. there exists a positive constant  $C_f$  such that  $\mathbb{P}$ -a.s. for all  $t \in [0, T]$ ,*

$$|f(t, y_1, z_1, u_1, \mu_1) - f(t, y_2, z_2, u_2, \mu_2)| \leq C_f(|y_1 - y_2| + |z_1 - z_2| + |u_1 - u_2|_\nu + \mathcal{W}_p(\mu_1, \mu_2))$$

*for any  $y_1, y_2 \in \mathbb{R}$ ,  $z_1, z_2 \in \mathbb{R}$ ,  $u_1, u_2 \in L_\nu^p$  and  $\mu_1, \mu_2 \in \mathcal{P}_p(\mathbb{R})$ .*

(c) *Assume that  $d\mathbb{P} \otimes dt$  a.e. for each  $(y, z, u_1, u_2, \mu) \in \mathbb{R}^2 \times (L_\nu^2)^2 \times \mathcal{P}_p(\mathbb{R})$ ,*

$$f(t, y, z, u_1, \mu) - f(t, y, z, u_2, \mu) \geq \langle \gamma_t^{y, z, u_1, u_2, \mu}, l_1 - l_2 \rangle_\nu, \quad (2.2)$$

*with*

$$\begin{aligned} \gamma : [0, T] \times \Omega \times \mathbb{R}^2 \times (L_\nu^2)^2 \times \mathcal{P}_p(\mathbb{R}) &\mapsto L_\nu^2; \\ (\omega, t, y, z, u_1, u_2, \mu) &\mapsto \gamma_t^{y, z, u_1, u_2, \mu}(\omega, \cdot) \end{aligned}$$

*$\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^2) \otimes \mathcal{B}((L_\nu^2)^2) \otimes \mathcal{B}(\mathcal{P}_p(\mathbb{R}))$  measurable satisfying  $\|\gamma_t^{y, z, u_1, u_2, \mu}(\cdot)\|_\nu \leq C$  for all  $(y, z, u_1, u_2, \mu) \in \mathbb{R}^2 \times (L_\nu^2)^2 \times \mathcal{P}_p(\mathbb{R})$ ,  $d\mathbb{P} \otimes dt$ -a.e., where  $C$  is a positive constant, and such that  $\gamma_t^{y, z, u_1, u_2, \mu}(e) \geq -1$ , for all  $(y, z, u_1, u_2, \mu) \in \mathbb{R}^2 \times (L_\nu^2)^2 \times \mathcal{P}_p(\mathbb{R})$ ,  $d\mathbb{P} \otimes dt \otimes d\nu(e)$ -a.e.*

(ii)  *$h$  is a mapping from  $[0, T] \times \Omega \times \mathbb{R} \times \mathcal{P}_p(\mathbb{R})$  into  $\mathbb{R}$  such that*

(a) *For all  $(y, \mu) \in \mathbb{R} \times \mathcal{P}_p(\mathbb{R})$ ,  $(h(t, y, \mu))_t$  is a right-continuous left-limited process;*

(b) *the process  $\left(\sup_{(y, \mu) \in \mathbb{R} \times \mathcal{P}_p(\mathbb{R})} |h(t, y, \mu)|\right)_{0 \leq t \leq T}$  belongs to  $\mathcal{S}^p$ ;*

(c)  *$h$  is Lipschitz w.r.t.  $(y, \mu)$  uniformly in  $(t, \omega)$ , i.e. there exists two positive constants  $\gamma_1$  and  $\gamma_2$  such that  $\mathbb{P}$ -a.s. for all  $t \in [0, T]$ ,*

$$|h(t, y_1, \mu_1) - h(t, y_2, \mu_2)| \leq \gamma_1 |y_1 - y_2| + \gamma_2 \mathcal{W}_p(\mu_1, \mu_2)$$

*for any  $y_1, y_2 \in \mathbb{R}$  and  $\mu_1, \mu_2 \in \mathcal{P}_p(\mathbb{R})$ .*

(iii)  *$\xi \in L^p(\mathcal{F}_T)$  and satisfies  $\xi \geq h(T, \xi, \mathbb{P}_\xi)$ .*

We now prove the existence and the uniqueness of the solution for the mean-field reflected BSDE with jumps (2.1), for which the law of the  $Y$ -component of the solution enters both the driver and the obstacle. Our method relies on Peng's nonlinear expectation (see e.g. Peng [36]) and a characterization of the solution of a mean-field reflected BSDE with jumps as a specific *mean-field nonlinear optimal*

*stopping problem.* This approach is completely different from the one used in [12] (where the penalization technique under a domination condition has been used in the case of a Brownian filtration) and allows to weaken the assumptions on the coefficients  $f$  and  $h$ , in particular we get rid of the monotonicity assumptions with respect to the measure component  $\mu$ .

Let  $\widehat{\Phi} : \mathbb{L}_\beta^p \rightarrow \mathbb{L}_\beta^p$  be the mapping that associates to a process  $Y$  another process  $\widehat{\Phi}(Y)$  defined by:

$$\widehat{\Phi}(Y)_t := \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathcal{E}_{t,\tau}^{f \circ Y} [h(\tau, Y_\tau, \mathbb{P}_{Y_s|s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau = T\}}], \quad (2.3)$$

where  $(f \circ Y)(t, y, z, u) = f(t, y, z, u, \mathbb{P}_{Y_t})$ .

**Theorem 2.1.** *Suppose that Assumption 2.1 is in force for some  $p \geq 2$ . Assume that  $\gamma_1$  and  $\gamma_2$  satisfy*

$$\gamma_1^p + \gamma_2^p < 2^{1-p}. \quad (2.4)$$

*Then the system (2.1) has a unique solution in  $\mathcal{S}^p \times \mathcal{H}^{p,1} \times \mathcal{H}_\nu^p \times \mathcal{S}_i^p$ .*

**Proof.** The proof is organized in three steps.

Step 1. We first show that  $\widehat{\Phi}$  is a well-defined map from  $\mathbb{L}_\beta^p$  to itself. Indeed, let  $\bar{Y} \in \mathbb{L}_\beta^p$ . Since  $h$  satisfies Assumption 2.1 (ii), it follows that  $h(t, \bar{Y}_t, \mathbb{P}_{\bar{Y}_t}) \in \mathcal{S}^p$ . Therefore, there exists a unique solution  $(\hat{Y}, \hat{Z}, \hat{U}, \hat{K}) \in \mathcal{S}^p \times \mathcal{H}^{p,1} \times \mathcal{H}_\nu^p \times \mathcal{S}_i^p$  of the reflected BSDE associated with obstacle  $h(t, \bar{Y}_t, \mathbb{P}_{\bar{Y}_t})$ , terminal condition  $\xi$  and driver  $(f \circ \bar{Y})(t, \omega, y, z, u)$ . Since  $f$  satisfies Assumption 2.1 (i.c), by classical results on the link between the  $Y$ -component of the solution of a reflected BSDE and optimal stopping with nonlinear expectations (see e.g. [15], [16]), we get that  $\widehat{\Phi}(\bar{Y}) = \hat{Y} \in \mathcal{S}^p \subset \mathbb{L}_\beta^p$ .

Step 2. We show that  $\widehat{\Phi} : \mathbb{L}_\beta^p \rightarrow \mathbb{L}_\beta^p$  is a contraction on the time interval  $[T - \delta, T]$ .

First, note that by the Lipschitz continuity of  $f$  and  $h$ , for  $Y, \bar{Y} \in \mathcal{S}_\beta^p$ ,  $Z, \bar{Z} \in \mathcal{H}^{p,1}$ , and  $U, \bar{U} \in \mathcal{H}_\nu^p$ ,

$$|f(s, Y_s, Z_s, U_s, \mathbb{P}_{Y_s}) - f(s, \bar{Y}_s, \bar{Z}_s, \bar{U}_s, \mathbb{P}_{\bar{Y}_s})| \leq C_f(|Y_s - \bar{Y}_s| + |Z_s - \bar{Z}_s| + |U_s - \bar{U}_s|_\nu + \mathcal{W}_p(\mathbb{P}_{Y_s}, \mathbb{P}_{\bar{Y}_s})),$$

$$|h(s, Y_s, \mathbb{P}_{Y_s}) - h(s, \bar{Y}_s, \mathbb{P}_{\bar{Y}_s})| \leq \gamma_1 |Y_s - \bar{Y}_s| + \gamma_2 \mathcal{W}_p(\mathbb{P}_{Y_s}, \mathbb{P}_{\bar{Y}_s}). \quad (2.5)$$

For the  $p$ -Wasserstein distance, we have the following inequality: for  $0 \leq s \leq u \leq t \leq T$ ,

$$\sup_{u \in [s, t]} \mathcal{W}_p(\mathbb{P}_{Y_u}, \mathbb{P}_{\bar{Y}_u}) \leq \sup_{u \in [s, t]} (\mathbb{E}[|Y_u - \bar{Y}_u|^p])^{1/p}, \quad (2.6)$$

from which we derive the following useful inequality

$$\sup_{u \in [s, t]} \mathcal{W}_p(\mathbb{P}_{Y_u}, \delta_0) \leq \sup_{u \in [s, t]} (\mathbb{E}[|Y_u|^p])^{1/p}. \quad (2.7)$$

Fix  $Y, \hat{Y} \in \mathbb{L}_\beta^p$ . For any  $t \leq T$ , by the estimates (A.1) on BSDEs, we have

$$\begin{aligned}
& |\widehat{\Phi}(Y)_t - \widehat{\Phi}(\bar{Y})_t|^p \\
&= \left| \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathcal{E}_{t,\tau}^{f \circ Y} [h(\tau, Y_\tau, \mathbb{P}_{Y_s|s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau=T\}}] - \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathcal{E}_{t,\tau}^{f \circ \bar{Y}} [h(\tau, \bar{Y}_\tau, \mathbb{P}_{\bar{Y}_s|s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau=T\}}] \right|^p \\
&\leq \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \left| \mathcal{E}_{t,\tau}^{f \circ Y} [h(\tau, Y_\tau, \mathbb{P}_{Y_s|s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau=T\}}] - \mathcal{E}_{t,\tau}^{f \circ \bar{Y}} [h(\tau, \bar{Y}_\tau, \mathbb{P}_{\bar{Y}_s|s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau=T\}}] \right|^p \\
&\leq \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[ \int_t^\tau e^{p\beta(s-t)} \eta^p |(f \circ Y)(s, \widehat{Y}_s^\tau, \widehat{Z}_s^\tau, \widehat{U}_s^\tau) - (f \circ \bar{Y})(s, \widehat{Y}_s^\tau, \widehat{Z}_s^\tau, \widehat{U}_s^\tau)|^p ds \right. \\
&\quad \left. + e^{p\beta(\tau-t)} |h(\tau, Y_\tau, \mathbb{P}_{Y_s|s=\tau}) - h(\tau, \bar{Y}_\tau, \mathbb{P}_{\bar{Y}_s|s=\tau})|^p | \mathcal{F}_t \right] \\
&= \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[ \int_t^\tau e^{p\beta(s-t)} \eta^p |f(s, \widehat{Y}_s^\tau, \widehat{Z}_s^\tau, \widehat{U}_s^\tau, \mathbb{P}_{Y_s}) - f(s, \widehat{Y}_s^\tau, \widehat{Z}_s^\tau, \widehat{U}_s^\tau, \mathbb{P}_{\bar{Y}_s})|^p ds \right. \\
&\quad \left. + e^{p\beta(\tau-t)} |h(\tau, Y_\tau, \mathbb{P}_{Y_s|s=\tau}) - h(\tau, \bar{Y}_\tau, \mathbb{P}_{\bar{Y}_s|s=\tau})|^p | \mathcal{F}_t \right],
\end{aligned}$$

with  $\eta, \beta > 0$  such that  $\eta \leq \frac{2c(p)}{pC_f^p}$  and  $\beta \geq C_f + \frac{1}{\eta} + \frac{p-1}{p} \left( \frac{1}{\eta} \right)^{\frac{p}{p-1}}$ , where  $(\widehat{Y}^\tau, \widehat{Z}^\tau, \widehat{U}^\tau)$  is the solution of the BSDE associated with driver  $f \circ \bar{Y}$  and terminal condition  $h(\tau, \bar{Y}_\tau, \mathbb{P}_{\bar{Y}_s|s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau=T\}}$ . Therefore, using (2.5) and the fact that

$$\mathcal{W}_p^p(\mathbb{P}_{Y_s|s=\tau}, \mathbb{P}_{\bar{Y}_s|s=\tau}) \leq \mathbb{E}[|Y_s - \bar{Y}_s|^p]_{s=\tau} \leq \sup_{\tau \in \mathcal{T}_t} \mathbb{E}[|Y_\tau - \bar{Y}_\tau|^p], \quad (2.8)$$

we have, for any  $t \in [T - \delta, T]$ ,

$$\begin{aligned}
e^{p\beta t} |\widehat{\Phi}(Y)_t - \widehat{\Phi}(\bar{Y})_t|^p &\leq \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[ \int_t^\tau e^{p\beta(s-t)} \eta^p C_f^p \mathbb{E}[|Y_s - \bar{Y}_s|^p] ds \right. \\
&\quad \left. + e^{p\beta(\tau-t)} \{ \gamma_1 |Y_\tau - \bar{Y}_\tau| + \gamma_2 (\mathbb{E}[|Y_s - \bar{Y}_s|^p]_{s=\tau})^{1/p} \}^p | \mathcal{F}_t \right], \\
&\leq \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[ \int_t^\tau e^{p\beta s} \eta^p C_f^p \mathbb{E}[|Y_s - \bar{Y}_s|^p] ds \right. \\
&\quad \left. + e^{p\beta \tau} \{ 2^{p-1} \gamma_1^p |Y_\tau - \bar{Y}_\tau|^p + 2^{p-1} \gamma_2^p \mathbb{E}[|Y_s - \bar{Y}_s|^p]_{s=\tau} \} | \mathcal{F}_t \right].
\end{aligned} \quad (2.9)$$

Therefore,

$$e^{p\beta t} |\widehat{\Phi}(Y)_t - \widehat{\Phi}(\bar{Y})_t|^p \leq \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E}[G(\tau) | \mathcal{F}_t] := V_t,$$

where

$$G(\tau) := \int_{T-\delta}^\tau e^{p\beta s} \eta^p C_f^p \mathbb{E}[|Y_s - \bar{Y}_s|^p] ds + e^{p\beta \tau} (2^{p-1} \gamma_1^p |Y_\tau - \bar{Y}_\tau|^p + 2^{p-1} \gamma_2^p \mathbb{E}[|Y_s - \bar{Y}_s|^p]_{s=\tau}).$$

which yields

$$\sup_{\tau \in \mathcal{T}_{T-\delta}} \mathbb{E}[e^{p\beta \tau} |\widehat{\Phi}(Y)_\tau - \widehat{\Phi}(\bar{Y})_\tau|^p] \leq \sup_{\tau \in \mathcal{T}_{T-\delta}} \mathbb{E}[V_\tau]. \quad (2.10)$$

We have

$$\sup_{\tau \in \mathcal{T}_{T-\delta}} \mathbb{E}[V_\tau] \leq \alpha \sup_{\tau \in \mathcal{T}_{T-\delta}} \mathbb{E}[e^{p\beta \tau} |Y_\tau - \bar{Y}_\tau|^p]. \quad (2.11)$$

Indeed, by Lemma D.1 in [25], for any  $\sigma \in \mathcal{T}_{T-\delta}$ , there exists a sequence  $(\tau_n)_n$  of stopping times in  $\mathcal{T}_\sigma$  such that

$$V_\sigma = \lim_{n \rightarrow \infty} \mathbb{E}[G(\tau_n) | \mathcal{F}_\sigma]$$

and so, by Fatou's Lemma, we have

$$\mathbb{E}[V_\sigma] \leq \liminf_{n \rightarrow \infty} \mathbb{E}[G(\tau_n)] \leq \sup_{\tau \in \mathcal{T}_{T-\delta}} \mathbb{E}[G(\tau)].$$

Therefore,

$$\sup_{\tau \in \mathcal{T}_{T-\delta}} \mathbb{E}[e^{p\beta t} |\widehat{\Phi}(Y)_t - \widehat{\Phi}(\bar{Y})_t|^p] \leq \sup_{\tau \in \mathcal{T}_{T-\delta}} \mathbb{E}[G(\tau)].$$

Using (2.8) and noting that  $e^{p\beta\tau} \mathbb{E}[|Y_s - \bar{Y}_s|^p]_{s=\tau} = \mathbb{E}[e^{p\beta s} |Y_s - \bar{Y}_s|^p]_{s=\tau}$ , we obtain

$$\sup_{\tau \in \mathcal{T}_{T-\delta}} \mathbb{E}[G(\tau)] \leq \alpha \sup_{\tau \in \mathcal{T}_{T-\delta}} \mathbb{E}[e^{p\beta\tau} |Y_\tau - \bar{Y}_\tau|^p]$$

where  $\alpha := \delta \eta^p C_f^p + 2^{p-1}(\gamma_1^p + \gamma_2^p)$ . This in turn yields (2.11).

Assuming  $(\gamma_1, \gamma_2)$  satisfies

$$\gamma_1^p + \gamma_2^p < 2^{1-p}$$

we can choose

$$0 < \delta < \frac{1}{\eta^p C_f^p} (1 - 2^{p-1}(\gamma_1^p + \gamma_2^p))$$

to make  $\widehat{\Phi}$  a contraction on  $\mathbb{L}_\beta^p$  over the time interval  $[T - \delta, T]$ , i.e.  $\widehat{\Phi}$  admits a unique fixed point over  $[T - \delta, T]$ .

Step 3. We finally show that the mean-field RBSDE with jumps (2.1) has a unique solution.

Given  $Y \in \mathbb{L}_\beta^p$  the solution over  $[T - \delta, T]$  obtained in *Step 2*, let  $(\hat{Y}, \hat{Z}, \hat{U}, \hat{K}) \in \mathcal{S}^p \times \mathcal{H}^{p,1} \times \mathcal{H}_v^p \times \mathcal{S}_i^p$  be the unique solution of the standard reflected BSDE, with barrier  $h(s, Y_s, \mathbb{P}_{Y_s})$  (which belongs to  $\mathcal{S}^p$  by Assumption 2.1 (ii)(a)) and driver  $g(s, y, z, u) := f(s, y, z, u, \mathbb{P}_{Y_s})$ . Then,

$$\hat{Y}_t = \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathcal{E}_{t,\tau}^g [h(\tau, Y_\tau, \mathbb{P}_{Y_s}|_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau = T\}}].$$

By the fixed point argument from Step 2, we have  $\hat{Y} = Y$  a.s., which implies that  $Y \in \mathcal{S}^p$  and

$$Y_t = \xi + \int_t^T f(s, Y_s, \hat{Z}_s, \hat{U}_s, \mathbb{P}_{Y_s}) ds + \hat{K}_T - \hat{K}_t - \int_t^T \hat{Z}_s dB_s - \int_t^T \int_{R^*} \hat{U}_s(e) \tilde{N}(ds, de), \quad T - \delta \leq t \leq T,$$

which yields existence of a solution. Furthermore, the process  $Y_t$  is a *nonlinear* g-supermartingale. Hence, by the uniqueness of the *nonlinear* Doob-Meyer decomposition, we obtain uniqueness of the associated processes  $(\hat{Z}, \hat{U}, \hat{K})$ . This yields existence and uniqueness of (2.1) in this general case.

Applying the same reasoning on each time interval  $[T - (i+1)\delta, T - i\delta]$  with a similar dynamics but terminal condition  $Y_{T-i\delta}$  at time  $T - i\delta$ , we build recursively for  $i = 1$  to any  $n$  a solution  $(Y, Z, U, K) \in \mathcal{S}^p \times \mathcal{H}^{p,1} \times \mathcal{H}_v^p \times \mathcal{S}_i^p$  on each time interval  $[T - (i+1)\delta, T - i\delta]$ . Pasting these processes, we derive a unique solution  $(Y, Z, U, K)$  satisfying (2.1) on the full time interval  $[0, T]$ .  $\square$

**Remark 2.2** (On the different smallness conditions). *We note that the following smallness condition, suggested in [12], imposed on  $(\gamma_1, \gamma_2)$ ,*

$$(\gamma_1 + \gamma_2)^{\frac{p-1}{p}} \left\{ \left( \frac{p}{p-1} \right)^p \gamma_1 + \gamma_2 \right\}^{\frac{1}{p}} < 1,$$

*implies (2.4). This follows from the inequality*

$$(\gamma_1 + \gamma_2)^{p-1} \left\{ \left( \frac{p}{p-1} \right)^p \gamma_1 + \gamma_2 \right\} \geq (\gamma_1 + \gamma_2)^p \geq 2^p \max(\gamma_1^p, \gamma_2^p) \geq 2^{p-1}(\gamma_1^p + \gamma_2^p). \quad (2.12)$$



Suppose that the assumptions of Theorem 2.1 hold. We now give a sufficient condition on the barrier  $h$  under which the component  $(Y_t)$  of the solution  $(Y, Z, U, K)$  of the mean-field reflected BSDE (2.1) has only totally inaccessible jumps, i.e. the process  $(K_t)$  is continuous. We first recall the following definition (see e.g. [15], [16]).

**Definition 2.2.** *A progressive process  $(\phi_t)$  is called left-upper semicontinuous (in short l.u.s.c) along stopping times if for all  $\tau \in \mathcal{T}_0$  and for each non-decreasing sequence of stopping times  $(\tau_n)$  such that  $\tau_n \uparrow \tau$  a.s.,  $\phi_\tau \geq \limsup_{n \rightarrow \infty} \phi_{\tau_n}$  a.s.*

Note that no condition is required at totally inaccessible jumps. When the process  $(\phi_t)$  is left limited,  $(\phi_t)$  is l.u.s.c. along stopping times if and only if for each predictable stopping time  $\tau \in \mathcal{T}_0$ ,  $\phi_\tau \geq \phi_{\tau-}$  a.s.

**Theorem 2.2.** *Suppose that  $\gamma_1$  and  $\gamma_2$  satisfy the condition (2.4). Assume that  $h$  takes the form  $h(t, \omega, y, \mu) := \xi_t(\omega) + \kappa(y, \mu)$ , where  $\xi$  belongs to  $\mathcal{S}^p$  and is a left upper semicontinuous process along stopping times, and  $\kappa$  is a bounded and Lipschitz function with respect to  $(y, \mu)$ . Let  $(Y, Z, U, K)$  be the unique solution of the mean-field reflected BSDE (2.1). Then  $(Y_t)$  has jumps only at totally inaccessible stopping times (i.e. the predictable process  $(K_t)$  is continuous).*

**Proof.** Let  $Y^0 \equiv 0$ . Define  $Y_t^{n+1} := \widehat{\Phi}(Y^n)_t$ , where the operator  $\widehat{\Phi}$  is defined by (2.3). Since the function  $\kappa$  is continuous with respect to  $(y, \mu)$  and  $\xi_t$  is left upper semicontinuous along stopping times, it follows that the process  $h(t, Y_t^0, \mathbb{P}_{Y_t^0})$  is also left upper semicontinuous along stopping times. Then  $Y^1$  corresponds to the first component of the reflected BSDE with driver  $f \circ Y^0$ , terminal condition  $\xi$  and obstacle process  $h(t, Y_t^0, \mathbb{P}_{Y_t^0})$ . Since  $h(t, Y_t^0, \mathbb{P}_{Y_t^0})$  is left upper semicontinuous along stopping times, it follows that  $Y^1$  has jumps only at totally inaccessible jumps and thus the map  $t \mapsto \mathbb{P}_{Y_t^0}$  is continuous. We deduce that  $h(t, Y_t^1, \mathbb{P}_{Y_t^1})$  is left upper semicontinuous along stopping times, and by the same arguments as above,  $Y^2$  has jumps only at totally inaccessible jumps too. By induction, one can show that  $Y_t^n$  has jumps only at totally inaccessible jumps for all  $n \geq 1$ .

Since the condition (2.4) is satisfied, by Theorem 2.1, the sequence  $Y_t^n$  is Cauchy for the norm  $\mathbb{L}^p$  and thus converges in  $\mathbb{L}^p$  to the component  $Y$  of the unique solution of the mean-field reflected BSDE (2.1). Let  $\tau \in \mathcal{T}_0$  be a predictable stopping time. Since  $\Delta Y_\tau^n = 0$  a.s. for all  $n$ , we obtain

$$\mathbb{E}[|\Delta Y_\tau|^p] = \mathbb{E}[|\Delta Y_\tau - \Delta Y_\tau^n|^p] \leq 2^p \sup_{\tau \in \mathcal{T}_0} \mathbb{E}[|Y_\tau - Y_\tau^n|^p], \quad (2.13)$$

which implies that  $\Delta Y_\tau = 0$  a.s. By the arbitrariness of the predictable stopping time  $\tau \in \mathcal{T}_0$ , the result follows.

### 3 Weakly interacting nonlinear Snell envelopes

We will now discuss the interpretation of the mean-field reflected BSDE with jumps studied in the previous section at the particle level and study the well-posedness of the associated particle system. To

this end, for a given vector  $\mathbf{x} := (x^1, \dots, x^n) \in \mathbb{R}^n$ , denote the empirical measure associated to  $\mathbf{x}$  by

$$L_n[\mathbf{x}] := \frac{1}{n} \sum_{k=1}^n \delta_{x^k}.$$

Let  $\{B^i\}_{1 \leq i \leq n}$ ,  $\{\tilde{N}^i\}_{1 \leq i \leq n}$  be independent copies of  $B$  and  $\tilde{N}$  and denote by  $\mathbb{F}^n := \{\mathcal{F}_t^n\}_{t \in [0, T]}$  the completion of the filtration generated by  $\{B^i\}_{1 \leq i \leq n}$  and  $\{\tilde{N}^i\}_{1 \leq i \leq n}$ . Thus, for each  $1 \leq i \leq n$ , the filtration generated by  $(B^i, \tilde{N}^i)$  is a sub-filtration of  $\mathbb{F}^n$ , and we denote by  $\tilde{\mathbb{F}}^i := \{\tilde{\mathcal{F}}_t^i\}$  its completion. Let  $\mathcal{T}_t^n$  be the set of  $\mathbb{F}^n$  stopping times with values in  $[0, T]$ .

Consider a family of weakly interacting processes  $\mathbf{Y}^n := (Y^{1,n}, \dots, Y^{n,n})$  evolving backward in time as follows: for  $i = 1, \dots, n$ ,

$$\begin{cases} Y_t^{i,n} = \xi^{i,n} + \int_t^T f(s, Y_s^{i,n}, Z_s^{i,i,n}, U_s^{i,i,n}, L_n[\mathbf{Y}_s^n]) ds + K_T^{i,n} - K_t^{i,n} \\ \quad - \int_t^T \sum_{j=1}^n Z_s^{i,j,n} dB_s^j - \int_t^T \int_{R^*} \sum_{j=1}^n U_s^{i,j,n}(e) \tilde{N}^j(ds, de), \quad t \in [0, T], \\ Y_t^{i,n} \geq h(t, Y_t^{i,n}, L_n[\mathbf{Y}_t^n]), \quad \forall t \in [0, T], \\ \int_0^T (Y_{t^-}^{i,n} - h(t^-, Y_{t^-}^{i,n}, L_n[\mathbf{Y}_{t^-}^n])) dK_t^{i,n} = 0. \end{cases} \quad (3.1)$$

**Assumption 3.1.** *We assume that*

- *$f$  and  $h$  are deterministic functions of  $(t, y, z, u, \mu)$  and  $(t, y, \mu)$ , respectively.*
- *$\xi^{i,n} \in L^p(\mathcal{F}_T^n)$ ,  $i = 1, \dots, n$ .*
- *$\xi^{i,n} \geq h(T, \xi^{i,n}, L_n[\xi^n])$  a.s.  $i = 1, \dots, n$ .*

For a more general form of the driver  $f$  and the obstacle  $h$  see Remark 4.2, below.

**Remark 3.1.**

- (1) *Note that in the system (3.1) we need to use all the noise processes  $((B^i, \tilde{N}^i), i = 1, \dots, n)$  to describe the dynamics  $Y^{i,n}$  of each particle  $i$ . This is due to the fact that the construction of each individual backward SDE is based on the martingale representation theorem which should involve the whole noise processes  $((B^i, \tilde{N}^i), i = 1, \dots, n)$  and not only individual noise, as is the case for forward SDEs.*
- (2) *The system (3.1) is a 'true' system of weakly interacting processes, and is different from the construction in [8] where the MF-BSDE is obtained a limit of one BSDE indexed by  $n$ , driven by one fixed Brownian motion and whose driver is an average of the Wiener measure preserving shift operator applied to measurable functions of the Wiener path, for which the strong law of large numbers for i.i.d. sequences holds.*

Note that we have the inequality

$$\mathcal{W}_p^p(L_n[\mathbf{x}], L_n[\mathbf{y}]) \leq \frac{1}{n} \sum_{j=1}^n |x_j - y_j|^p, \quad (3.2)$$

and, in particular,

$$\mathcal{W}_p^p(L_n[\mathbf{x}], L_n[\mathbf{0}]) \leq \frac{1}{n} \sum_{j=1}^n |x_j|^p, \quad (3.3)$$

where we note that  $L_n[\mathbf{0}] = \frac{1}{n} \sum_{j=1}^n \delta_0 = \delta_0$ .

Endow the product spaces  $\mathcal{S}_\beta^{p,\otimes n} := \mathcal{S}_\beta^p \times \mathcal{S}_\beta^p \times \cdots \times \mathcal{S}_\beta^p$  and  $\mathbb{L}_\beta^{p,\otimes n} := \mathbb{L}_\beta^p \times \mathbb{L}_\beta^p \times \cdots \times \mathbb{L}_\beta^p$  with the respective norms

$$\|h\|_{\mathcal{S}_\beta^{p,\otimes n}}^p := \sum_{1 \leq i \leq n} \|h^i\|_{\mathcal{S}_\beta^p}^p, \quad \|h\|_{\mathbb{L}_\beta^{p,\otimes n}}^p := \sum_{1 \leq i \leq n} \|h^i\|_{\mathbb{L}_\beta^p}^p. \quad (3.4)$$

Note that  $\mathcal{S}_\beta^{p,\otimes n}$  and  $\mathbb{L}_\beta^{p,\otimes n}$  are complete metric spaces. We denote by  $\mathcal{S}^{p,\otimes n} := \mathcal{S}_0^{p,\otimes n}$ ,  $\mathbb{L}^{p,\otimes n} := \mathbb{L}_0^{p,\otimes n}$ . Let  $\tilde{\Phi} : \mathbb{L}_\beta^{p,\otimes n} \rightarrow \mathbb{L}_\beta^{p,\otimes n}$  to be the mapping that associates to a process  $\mathbf{Y}^n := (Y^{1,n}, Y^{2,n}, \dots, Y^{n,n})$  the process  $\tilde{\Phi}(\mathbf{Y}^n) = (\tilde{\Phi}^1(\mathbf{Y}^n), \tilde{\Phi}^2(\mathbf{Y}^n), \dots, \tilde{\Phi}^n(\mathbf{Y}^n))$  defined by the following system: for every  $i = 1, \dots, n$  and  $t \leq T$ ,

$$\tilde{\Phi}^i(\mathbf{Y}^n)_t := \text{ess sup}_{\tau \in \mathcal{T}_t^n} \mathcal{E}_{t,\tau}^{\mathbf{f}^i \circ \mathbf{Y}^n} [h(\tau, Y_\tau^{i,n}, L_n[\mathbf{Y}_s^n])_{s=\tau} \mathbf{1}_{\{\tau < T\}} + \xi^{i,n} \mathbf{1}_{\{\tau = T\}}], \quad (3.5)$$

where

$$\begin{aligned} \mathbf{f}^i \circ \mathbf{Y}^n : [0, T] \times \Omega \times \mathbb{R} \times \mathbb{R}^n \times (\mathbb{R}^*)^n &\mapsto \mathbb{R} \\ (\mathbf{f}^i \circ \mathbf{Y}^n)(t, \omega, y, z, u) &= f(t, \omega, y, z^i, u^i, L_n[\mathbf{Y}_t^n](\omega)), \quad i = 1, \dots, n. \end{aligned}$$

**Theorem 3.1.** *Suppose that Assumptions 2.1 and 3.1 are in force for some  $p \geq 2$ . Suppose further that  $\gamma_1$  and  $\gamma_2$  satisfy (2.4) i.e.*

$$\gamma_1^p + \gamma_2^p < 2^{1-p}.$$

*Then the system (3.1) has a unique solution in  $\mathcal{S}^{p,\otimes n} \otimes \mathcal{H}^{p,n \otimes n} \otimes \mathcal{H}_V^{p,n \otimes n} \otimes \mathcal{S}_i^{p,\otimes n}$ .*

**Proof.** Step 1. We first show that  $\tilde{\Phi}$  is a well-defined map from  $\mathbb{L}_\beta^{p,\otimes n}$  to itself. To this end, we linearize the mapping  $h$  as follows: for  $i = 1, \dots, n$  and  $0 \leq s \leq T$ ,

$$h(s, Y_s^{i,n}, L_n[\mathbf{Y}_s^n]) = h(s, 0, L_n[\mathbf{0}]) + a_h^i(s) Y_s^{i,n} + b_h^i(s) \mathcal{W}_p(L_n[\mathbf{Y}_s^n], L_n[\mathbf{0}]),$$

where  $a_h^i(\cdot)$  and  $b_h^i(\cdot)$  are adapted processes given by

$$\begin{cases} a_h^i(s) := \frac{h(s, Y_s^{i,n}, L_n[\mathbf{Y}_s^n]) - h(s, 0, L_n[\mathbf{Y}_s^n])}{Y_s^{i,n}} \mathbf{1}_{\{Y_s^{i,n} \neq 0\}}, \\ b_h^i(s) := \frac{h(s, 0, L_n[\mathbf{Y}_s^n]) - h(s, 0, L_n[\mathbf{0}])}{\mathcal{W}_p(L_n[\mathbf{Y}_s^n], L_n[\mathbf{0}])} \mathbf{1}_{\{\mathcal{W}_p(L_n[\mathbf{Y}_s^n], L_n[\mathbf{0}]) \neq 0\}}. \end{cases} \quad (3.6)$$

and which, by the Lipschitz continuity of  $h$ , satisfy  $|a_h^i(\cdot)| \leq \gamma_1$ ,  $|b_h^i(\cdot)| \leq \gamma_2$ .

By the estimates (A.1) on BSDEs established in Proposition A.1, below, we obtain, for any stopping time  $\tau \in \mathcal{T}_t^n$  the following:

$$\begin{aligned} &\left| \mathcal{E}_{t,\tau}^{\mathbf{f}^i \circ \mathbf{Y}^n} [\xi^{i,n} \mathbf{1}_{\{\tau = T\}} + h(\tau, Y_\tau^{i,n}, L_n[\mathbf{Y}_s^n])_{s=\tau} \mathbf{1}_{\{\tau < T\}}] \right|^p \\ &\leq \mathbb{E} \left[ e^{p\beta(\tau-t)} \left| \xi^{i,n} \mathbf{1}_{\{\tau = T\}} + h(\tau, Y_\tau^{i,n}, L_n[\mathbf{Y}_s^n])_{s=\tau} \mathbf{1}_{\{\tau < T\}} \right|^p \middle| \mathcal{F}_t \right] \\ &\quad + \eta^p \mathbb{E} \left[ \int_t^\tau e^{p\beta(s-t)} |f|^p(s, 0, 0, 0, L_n[\mathbf{Y}_s^n]) ds \middle| \mathcal{F}_t \right], \end{aligned} \quad (3.7)$$

with  $\eta, \beta > 0$  such that  $\eta \leq \frac{2c(p)}{pC_f^2}$  and  $\beta \geq C_f + \frac{1}{\eta} + \frac{p-1}{p} \left(\frac{1}{\eta}\right)^{\frac{p}{p-1}}$ . Moreover, since  $f(t, 0, 0, 0, \delta_0) \in \mathcal{H}^{p,1}$ ,  $\xi^{i,n} \in L^p(\mathcal{F}_T)$ ,  $(h(t, 0, \delta_0))_{0 \leq t \leq T} \in \mathcal{S}_\beta^p$  and  $(Y^{i,n})_{1 \leq i \leq n} \in \mathbb{L}_\beta^{p,\otimes n}$ , the non-negative càdlàg process  $(M_t^{i,\beta})_{0 \leq t \leq T}$  defined by

$$\begin{aligned} M_t^{i,\beta} &:= \left( e^{\beta t} |h(t, 0, L_n[\mathbf{0}])| + \gamma_1 e^{\beta t} |Y_t^{i,n}| + \gamma_2 e^{\beta t} \mathcal{W}_p(L_n[\mathbf{Y}_t^n], L_n[\mathbf{0}]) + e^{\beta T} |\xi^{i,n}| \mathbf{1}_{\{t=T\}} \right)^p \\ &\quad + \int_0^t \eta^p (e^{\beta s} |f(s, 0, 0, 0, L_n[\mathbf{0}])| + C_f e^{\beta s} \mathcal{W}_p(L_n[\mathbf{Y}_s^n], L_n[\mathbf{0}]))^p ds \end{aligned}$$

belongs to  $\mathbb{L}^1$ . Therefore, using the same arguments as Step 1 in the proof of Theorem 2.1, for  $i = 1, \dots, n$ , it follows that  $\tilde{\Phi}(\mathbf{Y}^n) \in \mathbb{L}_\beta^{p,\otimes n}$ .

Step 2. We now show that  $\tilde{\Phi}$  is a contraction on the time interval  $[T - \delta, T]$ .

Fix  $\mathbf{Y}^n = (Y^{1,n}, \dots, Y^{n,n})$ ,  $\bar{\mathbf{Y}}^n = (\bar{Y}^{1,n}, \dots, \bar{Y}^{n,n}) \in \mathbb{L}_\beta^{p,\otimes n}$ ,  $(\hat{Y}, \tilde{Y}) \in (\mathcal{S}_\beta^p)^2$ ,  $(\hat{Z}, \tilde{Z}) \in (\mathcal{H}^{p,1})^2$ ,  $(\hat{U}, \tilde{U}) \in (\mathcal{H}_\nu^p)^2$ . By the Lipschitz continuity of  $f$  and  $h$ , we get

$$\begin{aligned} |f(s, \hat{Y}_s, \hat{Z}_s, \hat{U}_s, L_n[\mathbf{Y}_s^n]) - f(s, \tilde{Y}_s, \tilde{Z}_s, \tilde{U}_s, L_n[\bar{\mathbf{Y}}_s^n])| &\leq C_f (|\hat{Y}_s - \tilde{Y}_s| + |\hat{Z}_s - \tilde{Z}_s| \\ &\quad + |\hat{U}_s - \tilde{U}_s|_\nu + \mathcal{W}_p(L_n[\mathbf{Y}_s^n], L_n[\bar{\mathbf{Y}}_s^n])), \\ |h(s, Y_s^{i,n}, L_n[\mathbf{Y}_s^n]) - h(s, \bar{Y}_s^{i,n}, L_n[\bar{\mathbf{Y}}_s^n])| &\leq \gamma_1 |Y_s^{i,n} - \bar{Y}_s^{i,n}| + \gamma_2 \mathcal{W}_p(L_n[\mathbf{Y}_s^n], L_n[\bar{\mathbf{Y}}_s^n]). \end{aligned} \quad (3.8)$$

By (3.2), we have

$$\mathcal{W}_p^p(L_n[\mathbf{Y}_s^n], L_n[\bar{\mathbf{Y}}_s^n]) \leq \frac{1}{n} \sum_{j=1}^n |Y_s^{j,n} - \bar{Y}_s^{j,n}|^p. \quad (3.9)$$

Then, using the estimates (A.1) again, for any  $t \leq T$  and  $i = 1, \dots, n$ , we have

$$\begin{aligned} &|\tilde{\Phi}^i(\mathbf{Y}^n)_t - \tilde{\Phi}^i(\bar{\mathbf{Y}}^n)_t|^p \\ &= |\text{ess sup}_{\tau \in \mathcal{T}_t^n} \mathcal{E}_{t,\tau}^{\mathbf{f}^i \circ \mathbf{Y}^n} [h(\tau, Y_\tau^{i,n}, L_n[\mathbf{Y}_s^n]_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi^{i,n} \mathbf{1}_{\{\tau=T\}}] \\ &\quad - \text{ess sup}_{\tau \in \mathcal{T}_t^n} \mathcal{E}_{t,\tau}^{\mathbf{f}^i \circ \bar{\mathbf{Y}}^n} [h(\tau, \bar{Y}_\tau^{i,n}, L_n[\bar{\mathbf{Y}}_s^n]_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi^{i,n} \mathbf{1}_{\{\tau=T\}}]|^p \\ &\leq \text{ess sup}_{\tau \in \mathcal{T}_t^n} \left| \mathcal{E}_{t,\tau}^{\mathbf{f}^i \circ \mathbf{Y}^n} [h(\tau, Y_\tau^{i,n}, L_n[\mathbf{Y}_s^n]_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi^{i,n} \mathbf{1}_{\{\tau=T\}}] \right. \\ &\quad \left. - \mathcal{E}_{t,\tau}^{\mathbf{f}^i \circ \bar{\mathbf{Y}}^n} [h(\tau, \bar{Y}_\tau^{i,n}, L_n[\bar{\mathbf{Y}}_s^n]_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi^{i,n} \mathbf{1}_{\{\tau=T\}}] \right|^p \\ &\leq \text{ess sup}_{\tau \in \mathcal{T}_t^n} \mathbb{E}[\int_t^\tau e^{p\beta(s-t)} \eta^p |(\mathbf{f}^i \circ \mathbf{Y}^n)(s, \hat{Y}_s^{i,\tau}, \hat{Z}_s^{i,\tau}, \hat{U}_s^{i,\tau}) - (\mathbf{f}^i \circ \bar{\mathbf{Y}}^n)(s, \hat{Y}_s^{i,\tau}, \hat{Z}_s^{i,\tau}, \hat{U}_s^{i,\tau})|^p ds \\ &\quad + e^{p\beta(\tau-t)} |h(\tau, Y_\tau^{i,n}, L_n[\mathbf{Y}_s^n]_{s=\tau}) - h(\tau, \bar{Y}_\tau^{i,n}, L_n[\bar{\mathbf{Y}}_s^n]_{s=\tau})|^p | \mathcal{F}_t] \\ &\leq \text{ess sup}_{\tau \in \mathcal{T}_t^n} \mathbb{E}[\int_t^\tau e^{p\beta(s-t)} \eta^p |f(s, \hat{Y}_s^{i,\tau}, \hat{Z}_s^{i,\tau}, \hat{U}_s^{i,\tau}, L_n[\mathbf{Y}_s^n]) - f(s, \hat{Y}_s^{i,\tau}, \hat{Z}_s^{i,\tau}, \hat{U}_s^{i,\tau}, L_n[\bar{\mathbf{Y}}_s^n])|^p ds \\ &\quad + e^{p\beta(\tau-t)} |h(\tau, Y_\tau^{i,n}, L_n[\mathbf{Y}_s^n]_{s=\tau}) - h(\tau, \bar{Y}_\tau^{i,n}, L_n[\bar{\mathbf{Y}}_s^n]_{s=\tau})|^p | \mathcal{F}_t], \end{aligned}$$

where  $(\hat{Y}^{i,\tau}, \hat{Z}^{i,\tau}, \hat{U}^{i,\tau})$  is the solution of the BSDE associated with driver  $\mathbf{f}^i \circ \bar{\mathbf{Y}}^n$  and terminal condition  $h(\tau, \bar{Y}_\tau^{i,n}, L_n[\bar{\mathbf{Y}}_s^n]_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi^{i,n} \mathbf{1}_{\{\tau=T\}}$ .

Therefore, using (3.8) and (3.9), we have, for any  $t \leq T$  and  $i = 1, \dots, n$ ,

$$\begin{aligned} e^{p\beta t} |\tilde{\Phi}^i(\mathbf{Y}^n)_t - \tilde{\Phi}^i(\bar{\mathbf{Y}}^n)_t|^p &\leq \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t^n} \mathbb{E} \left[ \int_t^T \eta^p C_f^p \left( \frac{1}{n} \sum_{j=1}^n e^{p\beta s} |Y_s^{j,n} - \bar{Y}_s^{j,n}|^p \right) ds \right. \\ &\quad \left. + \left( \gamma_1 e^{p\beta \tau} |Y_\tau^{i,n} - \bar{Y}_\tau^{i,n}| + \gamma_2 \left( \frac{1}{n} \sum_{j=1}^n e^{p\beta \tau} |Y_\tau^{j,n} - \bar{Y}_\tau^{j,n}|^p \right)^{\frac{1}{p}} \right)^p \middle| \mathcal{F}_t \right]. \end{aligned} \quad (3.10)$$

Next, let  $\delta > 0$ ,  $t \in [T - \delta, T]$  and with  $\alpha := \max(\delta \eta^p C_f^p + 2^{p-1} \gamma_1^p, \delta \eta^p C_f^p + 2^{p-1} \gamma_2^p)$ . By applying the same arguments used in Step 2 from the proof of Theorem 2.1, and the definition (3.4), we have

$$\|\tilde{\Phi}(\mathbf{Y}^n) - \tilde{\Phi}(\bar{\mathbf{Y}}^n)\|_{\mathbb{L}_\beta^{p,\otimes n}[T-\delta, T]}^p \leq \alpha \|\mathbf{Y}^n - \bar{\mathbf{Y}}^n\|_{\mathbb{L}_\beta^{p,\otimes n}[T-\delta, T]}^p.$$

Hence, if  $\gamma_1$  and  $\gamma_2$  satisfy

$$\gamma_1^p + \gamma_2^p < 2^{1-p},$$

we can choose

$$0 < \delta < \frac{1}{\eta^p C_f^p} (1 - 2^{p-1}(\gamma_1^p + \gamma_2^p))$$

to make  $\tilde{\Phi}$  a contraction on  $\mathbb{L}_\beta^{p,\otimes n}([T - \delta, T])$ , i.e.  $\tilde{\Phi}$  admits a unique fixed point over  $[T - \delta, T]$ . Moreover, in view of Assumption 2.1 (ii)(a), we get that  $\mathbf{Y}^n \in \mathcal{S}^{p,\otimes n}([T - \delta, T])$ .

Step 3. The system (3.1) has a unique solution. We first denote

$$\begin{aligned} \boldsymbol{\xi}^n &:= (\xi^{1,n}, \dots, \xi^{n,n}); & \mathbf{Z}^{i,n} &:= (Z^{i,1,n}, \dots, Z^{i,n,n})_{i=1, \dots, n}; \\ \mathbf{U}^{i,n} &:= (U^{i,1,n}, \dots, U^{i,n,n})_{i=1, \dots, n}; & \mathbf{K}^n &:= (K^{1,n}, \dots, K^{n,n}). \end{aligned}$$

Given  $\mathbf{Y}^n = (Y^{1,n}, \dots, Y^{n,n}) \in \mathbb{L}_\beta^{p,\otimes n}$  the solution obtained in Step 2, let  $(\hat{\mathbf{Y}}^n, \hat{\mathbf{Z}}^n, \hat{\mathbf{U}}^n, \hat{\mathbf{K}}^n) \in \mathcal{S}^{p,\otimes n} \otimes \mathcal{H}^{p,n,\otimes n} \otimes \mathcal{H}_\nu^{p,n,\otimes n} \otimes \mathcal{S}_i^{p,\otimes n}$  be the unique solution of the RBSDE system over  $[T - \delta, T]$ , with barrier  $(h(t, Y^{i,n}, L_n(\mathbf{Y}^n)))_{i=1, \dots, n}$  (which belongs to  $\mathcal{S}^{p,\otimes n}$ ) and driver  $(\mathbf{f}^i \circ \mathbf{Y}^n)_{i=1, \dots, n}$ . Then, for each  $i = 1, \dots, n$ , we get

$$\hat{Y}_t^{i,n} = \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t^n} \mathcal{E}_{t,\tau}^{\mathbf{f}^i \circ \mathbf{Y}^n} [h(\tau, \mathbf{Y}_\tau^n) \mathbf{1}_{\{\tau < T\}} + \xi^{i,n} \mathbf{1}_{\{\tau = T\}}].$$

By the fixed point argument from Step 2, we have  $\hat{\mathbf{Y}}^n = \mathbf{Y}^n$  a.s., where the equality is component-wise. This implies that  $\mathbf{Y}^n \in \mathcal{S}^{p,\otimes n}$  and for all  $1 \leq i \leq n$  and  $T - \delta \leq t \leq T$ ,

$$Y_t^{i,n} = \xi^{i,n} + \int_t^T f(s, Y_s^{i,n}, \hat{Z}_s^{i,i,n}, \hat{U}_s^{i,i,n}) ds + \hat{K}_T^{i,n} - \hat{K}_t^{i,n} - \int_t^T \sum_{j=1}^n \hat{Z}_s^{i,j,n} dB_s^j - \int_t^T \int_{R^*} \sum_{j=1}^n \hat{U}_s^{i,j,n}(e) \tilde{N}^j(ds, de).$$

Therefore, we obtain existence of a solution. Note that, for each  $i$ , the process  $Y_t^i$  is a *nonlinear*  $\mathbf{f}^i \circ \mathbf{Y}$ -supermartingale. Hence, by the uniqueness of the *nonlinear* Doob-Meyer decomposition, we get uniqueness of the associated processes  $(\hat{\mathbf{Z}}^{i,n}, \hat{\mathbf{U}}^{i,n}, \hat{K}^{i,n})$ . This yields existence and uniqueness of (3.1) on the time interval  $[T - \delta, T]$  in this general case.

Applying the same reasoning on each time interval  $[T - (j+1)\delta, T - j\delta]$ ,  $1 \leq j \leq m$ , with a similar dynamics but terminal condition  $\mathbf{Y}_{T-j\delta}^i$  at time  $T - j\delta$ , we build recursively, for  $j = 1$  to  $n$ , a solution  $(\mathbf{Y}^i, \mathbf{Z}^i, \mathbf{U}^i, \mathbf{K}^i)$  on each time interval  $[T - (j+1)\delta, T - j\delta]$ . Pasting properly these processes, we naturally derive a unique solution  $(\mathbf{Y}, \mathbf{Z}, \mathbf{U}, \mathbf{K})$  satisfying (3.1) on the full time interval  $[0, T]$ .  $\square$

## 4 Convergence of the particle system and propagation of chaos

This section is concerned with the convergence of the sequence of processes  $(Y^{i,n}, Z^{i,n}, U^{i,n}, K^{i,n})$ , solutions of the particle system (3.1), to the independent copies  $(Y^i, Z^i, U^i, K^i)$  of the solution of the mean-field Reflected BSDEs (2.1), as well as a related propagation of chaos result.

### 4.1 Preliminaries

We first introduce the Wasserstein metric on  $\mathbb{D} := D([0, T], \mathbb{R})$  and recall some of its properties along with a law of large numbers result for  $\mathbb{D}$ -valued i.i.d. random variables to be used in our proofs towards the propagation of chaos property.

**Wasserstein metric on  $\mathbb{D} := D([0, T], \mathbb{R})$**

Let  $\mathbb{D} = D([0, T], \mathbb{R})$  be the set of càdlàg functions i.e. the space of functions from  $[0, T]$  to  $\mathbb{R}$  which are right continuous with left limits at each  $t \in [0, T)$  and are left continuous at time  $T$ . Let  $\Lambda$  denote the class of strictly increasing, continuous mappings of  $[0, T]$  onto itself. If  $\lambda \in \Lambda$ , then  $\lambda(0) = 0$  and  $\lambda(T) = T$ . Let  $d^\circ(x, y)$  be the infimum of those positive  $\epsilon$  for which  $\Lambda$  contains some  $\lambda$  such that  $\|\lambda\|^\circ := \sup_{0 \leq s < t \leq T} |\log \frac{\lambda(t) - \lambda(s)}{t - s}| \leq \epsilon$  and for  $x, y \in \mathbb{D}$ ,

$$\sup_{0 \leq t \leq T} |x(t) - y(\lambda(t))| = \sup_{0 \leq t \leq T} |x(\lambda^{-1}(t)) - y(t)| < \epsilon.$$

In other words, let

$$d^\circ(x, y) = \inf_{\lambda \in \Lambda} \{ \|\lambda\|^\circ \vee \sup_{0 \leq t \leq T} |x(t) - y(\lambda(t))| \}. \quad (4.1)$$

The space  $\mathbb{D}$  endowed with the Skorohod metric  $d^\circ$  is a Polish space. By taking  $\lambda$  to be identity function, we obtain  $\|\lambda\|^\circ = 0$ , and then

$$d^\circ(x, y) \leq |x - y|_T,$$

where  $|z|_t := \sup_{0 \leq s \leq t} |z(s)|$ , i.e.  $|\cdot|_T$  induces a stronger topology than  $d^\circ$ .

To derive (4.5) below, we need the metric  $d$  which is equivalent to  $d^\circ$  and for which  $(\mathbb{D}, d)$  is separable but not complete, defined as (see (14.12)-(14.13) in Billingsley [4])

$$d(x, y) = \inf_{\lambda \in \Lambda} \{ \sup_{0 \leq t \leq T} |\lambda(t) - t| \vee \sup_{0 \leq t \leq T} |x(t) - y(\lambda(t))| \}. \quad (4.2)$$

In view of (4.1) and (4.2), we have, for any  $(x, y) \in \mathbb{D} \times \mathbb{D}$ ,

$$d(x, y) \leq |x - y|_T, \quad d(x, y) \leq T[e^{d^\circ(x, y)} - 1], \quad (4.3)$$

where the last inequality follows from the fact that, by using the continuity of the map  $\lambda$ , we get

$$\begin{aligned} \sup_{0 \leq t \leq T} |\lambda(t) - t| &= \sup_{0 < t \leq T} |\lambda(t) - t| = \sup_{0 < t \leq T} t \left| \frac{\lambda(t) - 0}{t - 0} - 1 \right| \\ &\leq T \sup_{0 < t \leq T} \left| \frac{\lambda(t) - 0}{t - 0} - 1 \right| \leq T[e^{\|\lambda\|^\circ} - 1], \end{aligned}$$

where we have used the inequality  $|a - 1| \leq e^{|\log a|} - 1$  for  $a > 0$ .

Following Billingsley [4], (14.6) and (14.7), for  $x \in \mathbb{D}$ ,  $\delta > 0$  and  $0 \leq u \leq v \leq T$ , put

$$\begin{aligned} w_x([u, v]) &= \sup\{|x(t) - x(s)| : s, t \in [u, v]\}, \\ w'_x(\delta) &= \inf_{\{t_i\}} \max_{1 \leq i \leq r} w_x([t_{i-1}, t_i]), \end{aligned}$$

where the infimum extends over the finite sets  $\{t_i\}$  of points satisfying

$$\begin{cases} 0 = t_0 < t_1 < \dots < t_r = T, \\ t_i - t_{i-1} > \delta, \quad i = 1, \dots, r. \end{cases}$$

From this definition, it follows that

$$w_x([u, v]) \leq 2w'_x(v - u) + \sup_{u < t \leq v} |x(t) - x(t^-)|, \quad \text{for all } 0 \leq u \leq v \leq T. \quad (4.4)$$

Furthermore, for any  $x, y \in \mathbb{D}$  and any  $\lambda \in \Lambda$ , it holds that

$$|x(t) - y(t)| \leq |x(t) - y(\lambda(t))| + |y(\lambda(t)) - y(t)|.$$

The definition (4.1), which implies that for any  $\delta > 0$ , for any  $x, y \in \mathbb{D}$  such that  $d^o(x, y) \leq \ln(1 + \delta/T)$  we get  $d(x, y) \leq \delta$  by (4.3), which together with the above relation, yields

$$|x(t) - y(t)| \leq \delta + w_y([t - \delta, t + \delta]).$$

Thus, in view of (4.4), if  $d^o(x, y) \leq \ln(1 + \delta/T)$  then

$$|x(t) - y(t)| \leq \delta + 2w'_y(2\delta) + \sup_{t - \delta < s \leq t + \delta} |y(s) - y(s^-)|. \quad (4.5)$$

The space of probability measures on the space  $\mathbb{D}$  is denoted by  $\mathcal{P}(\mathbb{D})$ . We define the  $p$ -Wasserstein metric between two probability measures  $P$  and  $Q$  on  $(\mathbb{D}, |\cdot|_T)$  as

$$D_t(P, Q) := \inf \left\{ \left( \int_{\mathbb{D} \times \mathbb{D}} |x - y|_t^p R(dx, dy) \right)^{1/p} \right\}, \quad 0 \leq t \leq T, \quad (4.6)$$

over  $R \in \mathcal{P}(\mathbb{D} \times \mathbb{D})$  with marginals  $P$  and  $Q$ .

Let  $\mathcal{P}_p(\mathbb{D})$  denote the space of probability measures  $Q$  on  $(\mathbb{D}, |\cdot|_T)$  with finite  $p$ th-moments:

$$\|Q\|_p^p := \int_{\mathbb{D}} |w|_T^p Q(dw) < +\infty.$$

For  $P, Q \in \mathcal{P}_p(\mathbb{D})$ , we have

$$D_s(P, Q) \leq D_t(P, Q), \quad 0 \leq s \leq t. \quad (4.7)$$

Moreover, for  $P, Q \in \mathcal{P}_p(\mathbb{D})$  with time marginals  $P_t := P \circ \omega^{-1}(t)$  and  $Q_t := Q \circ \omega^{-1}(t)$  ( $\omega$  being the coordinate process in  $\mathbb{D}$ ), the  $p$ -Wasserstein distance between  $P_t$  and  $Q_t$  satisfies

$$\mathcal{W}_p(P_t, Q_t) \leq D_t(P, Q), \quad 0 \leq t \leq T. \quad (4.8)$$

In particular,

$$\sup_{0 \leq t \leq T} \mathcal{W}_p(P_t, Q_t) \leq D_T(P, Q). \quad (4.9)$$

Endowed with the  $p$ -Wasserstein metric  $D_T$ ,  $\mathcal{P}_p(\mathbb{D})$  is a complete metric space.

The  $p$ -Wasserstein metric for probability measures on  $(\mathbb{D}, d^o)$  is defined by

$$D_T^o(P, Q) := \inf \left\{ \left( \int_{\mathbb{D} \times \mathbb{D}} d^o(x, y)^p R(dx, dy) \right)^{1/p} \right\}$$

over  $R \in \mathcal{P}(\mathbb{D} \times \mathbb{D})$  with marginals  $P$  and  $Q$ . Endowed with the  $p$ -Wasserstein metric  $D_T^o$ ,  $\mathcal{P}(\mathbb{D})$  is a Polish space.

Due to the relation  $d^o(x, y) \leq |x - y|_T$ , we have

$$D_T^o(P, Q) \leq D_T(P, Q), \quad P, Q \in \mathcal{P}_p(\mathbb{D}),$$

which means that the metric  $D_T$  is stronger than  $D_T^o$ .

We end these preliminaries with the following law of large numbers.

**Lemma 4.1.** *Let  $P \in \mathcal{P}_p((\mathbb{D}, d^o))$  and  $L^n(\zeta) := \frac{1}{n} \sum_{i=1}^n \delta_{\zeta_i}$  denote the empirical measure associated with the sequence of i.i.d.  $\mathbb{D}$ -valued random variables  $\zeta := (\zeta^i)_{i \geq 1}$  with probability law  $P$ . Then,*

$$\text{for all } n \geq 1, \quad \mathbb{E}[D_T^o(L^n(\zeta), P)^p] \leq 2^p D_T^o(P, \delta_0)^p \quad \text{and} \quad \lim_{n \rightarrow \infty} \mathbb{E}[D_T^o(L^n(\zeta), P)^p] = 0. \quad (4.10)$$

**Proof.** The random variables  $x_i := \delta_{\zeta_i}$  being i.i.d. with values in the Polish space  $(\mathcal{P}(\mathbb{D}), D_T^o)$ , by the strong law of large numbers (SLLN) (Varadarajan's theorem), with probability one,  $L^n(\zeta) := \frac{1}{n} \sum_{i=1}^n x_i$  converges weakly to  $P$ , as  $n$  tends to infinity. Moreover, since  $P \in \mathcal{P}_p((\mathbb{D}, d^o))$ , by the SLLN,  $\frac{1}{n} \sum_{i=1}^n d^0(\zeta_i, 0)^p$  converges to  $D_T^o(P, \delta_0)^p$  a.s.. Therefore, in view of Corollary 12.2.2 in [38],  $D_T^o(L^n(\zeta), P)$  converges to 0 a.s., as  $n \rightarrow \infty$ . We also have

$$D_T^o(L^n(\zeta), P)^p \leq 2^{p-1} \left( \frac{1}{n} \sum_{i=1}^n d^0(\zeta_i, 0)^p + D_T^o(P, \delta_0)^p \right).$$

Thus,

$$\mathbb{E}[D_T^o(L^n(\zeta), P)^p] \leq 2^{p-1} \left( \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n d^0(\zeta_i, 0)^p \right] + D_T^o(P, \delta_0)^p \right) = 2^p D_T^o(P, \delta_0)^p.$$

Therefore,  $D_T^o(L^n(\zeta), P)^p$  is uniformly integrable. Hence it converges in  $L^1$ , i.e.

$$\lim_{n \rightarrow \infty} \mathbb{E}[D_T^o(L^n(\zeta), P)^p] = 0.$$

□

The propagation of chaos is a property of systems of exchangeable (symmetric in law) processes. We first recall the definition of exchangeable (or symmetric in law) random variables and then of that of propagation of chaos. We refer to Sznitman [40] for further details.



**Definition 4.1** (Exchangeable r.v.). *The random variables  $X^1, X^2, \dots, X^n$  are said to be exchangeable if the law of the random vector  $(X^1, X^2, \dots, X^n)$  is the same as that of  $(X^{\sigma(1)}, X^{\sigma(2)}, \dots, X^{\sigma(n)})$  for every permutation  $\sigma$  of the set  $\{1, 2, \dots, n\}$ . We write*

$$\text{Law}(X^1, X^2, \dots, X^n) = \text{Law}(X^{\sigma(1)}, X^{\sigma(2)}, \dots, X^{\sigma(n)}).$$

Let  $E$  be a separable metric space and let  $P$  be a probability measure on  $E$ .

**Definition 4.2** (Propagation of chaos property). *The sequence of exchangeable  $E$ -valued random variables  $X^{1,n}, X^{2,n}, \dots, X^{n,n}$ ,  $n \geq 1$ , is said to be  $P$ -chaotic provided that, for any fixed positive integer  $k$ ,*

$$\lim_{n \rightarrow \infty} \text{Law}(X^{1,n}, X^{2,n}, \dots, X^{k,n}) = P^{\otimes k}.$$

## 4.2 Propagation of chaos

To establish the propagation of chaos property of the particle system (3.1), we shall further impose the following conditions.

**Assumption 4.1.** (i) *The sequence  $\xi^n = (\xi^{1,n}, \xi^{2,n}, \dots, \xi^{n,n})$  is exchangeable i.e. the sequence of probability laws  $\mu^n$  of  $\xi^n$  on  $\mathbb{R}^n$  is symmetric.*

(ii)  *$L_n[\xi^n] = \frac{1}{n} \sum_{i=1}^n \delta_{\xi^{i,n}}$  converges in  $\mathcal{W}_p$  to  $\mu$ . In particular,  $\mu^n$  is  $\mu$ -chaotic.*

(iii) *For each  $i \geq 1$ ,  $\xi^{i,n}$  converges in  $L^p$  to  $\xi^i$ , i.e.*

$$\lim_{n \rightarrow \infty} \mathbb{E}[|\xi^{i,n} - \xi^i|^p] = 0,$$

where the random variables  $\xi^i \in L^p(\mathcal{F}_T^i)$  are independent and equally distributed (iid) with probability law  $\mu$ .

(iv) *The component  $(Y_t)$  of the unique solution of the mean-field reflected BSDE (2.1) has jumps only at totally inaccessible stopping times.*

Sufficient conditions on the barrier  $h$  under which (iv) holds are given in Theorem 2.2.

For  $m \geq 1$ , introduce the Polish spaces  $\mathbb{H}^{2,m} := L^2([0, T]; \mathbb{R}^m)$  and  $\mathbb{H}_\nu^{2,m}$ , the space of measurable functions  $\ell : [0, T] \times \mathbb{R}^* \rightarrow \mathbb{R}^n$  such that  $\int_0^T \int_{\mathbb{R}^*} \sum_{j=1}^m |\ell^j(t, u)|^2 \nu(du) dt < \infty$ .

In the following proposition, we show that the *exchangeability property* transfers from the terminal conditions to the associated solution processes.

**Proposition 4.1** (Exchangeability property). *Assume the sequence  $\xi^n = (\xi^{1,n}, \xi^{2,n}, \dots, \xi^{n,n})$  is exchangeable i.e. the sequence of probability laws  $\mu^n$  of  $\xi^n$  on  $\mathbb{R}^n$  is symmetric. Then the processes  $\Theta^{i,n} := (Y^{i,n}, Z^{i,n}, U^{i,n}, K^{i,n})$ ,  $i = 1, \dots, n$ , solutions of the systems (3.1) are exchangeable.*

**Proof.** Since for any permutation  $\sigma$  of the set  $\{1, 2, \dots, n\}$ , we have

$$\frac{1}{n} \sum_{i=1}^n \delta_{Y^{i,n}} = \frac{1}{n} \sum_{i=1}^n \delta_{Y^{\sigma(i),n}},$$

by the pathwise existence and uniqueness results obtained in Theorem 3.1, we have

$$(\Theta^{1,n}, \Theta^{2,n}, \dots, \Theta^{n,n}) = (\Theta^{\sigma(1),n}, \Theta^{\sigma(2),n}, \dots, \Theta^{\sigma(n),n}), \quad \text{a.s.}$$

whenever  $(\xi^{1,n}, \dots, \xi^{n,n}) = (\xi^{\sigma(1),n}, \dots, \xi^{\sigma(n),n})$  a.s.. Now,  $\xi^n$  being exchangeable, by a careful use of the Skorohod representation theorem applied to the Polish space

$$\mathbb{R}^n \times (\mathbb{D}, d^0)^{\otimes n} \times C([0, T]; \mathbb{R}^n) \times (\mathbb{D}, d^0)^{\otimes n} \times \mathbb{H}^{2,n \otimes n} \times \mathbb{H}_V^{2,n \otimes n} \times (\mathbb{D}, d^0)^{\otimes n}$$

where the process  $(\xi^n, \{B^i\}_{i=1}^n, \{\tilde{N}^i\}_{i=1}^n, \{\Theta^{i,n}\}_{i=1}^n)$  take their values, as used by Yamada and Watanabe [42] to prove uniqueness of solutions to SDEs and extended in Delarue [10] (see the proof after Remark 1.6, pp. 224-227) to prove uniqueness in law for forward-backward SDEs, it follows that each of the systems is exchangeable i.e. for every permutation  $\sigma$  of  $\{1, 2, \dots, n\}$ ,

$$\text{Law}(\Theta^{1,n}, \Theta^{2,n}, \dots, \Theta^{n,n}) = \text{Law}(\Theta^{\sigma(1),n}, \Theta^{\sigma(2),n}, \dots, \Theta^{\sigma(n),n}).$$

□

Consider the product space

$$G := \mathbb{D} \times \mathbb{H}^{2,n} \times \mathbb{H}_V^{2,n} \times \mathbb{D}$$

endowed with the product metric

$$\delta((y, z, u, k), (y', z', u', k')) := \left( d^o(y, y')^p + \|z - z'\|_{\mathbb{H}^{2,n}}^p + \|u - u'\|_{\mathbb{H}_V^{2,n}}^p + d^o(k, k')^p \right)^{\frac{1}{p}}.$$

We define the Wasserstein metric on  $\mathcal{P}_p(G)$  by

$$D_G(P, Q) = \inf \left\{ \left( \int_{G \times G} \delta((y, z, u, k), (y', z', u', k'))^p R(dx, dy) \right)^{1/p} \right\}, \quad (4.11)$$

over  $R \in \mathcal{P}(G \times G)$  with marginals  $P$  and  $Q$ . Since  $(G, \delta)$  is a Polish space,  $(\mathcal{P}_p(G), D_G)$  is a Polish space and induces the topology of weak convergence (see Theorem 6.9 in [43]).

Let  $\Theta^i := (Y^i, Z^i, U^i, K^i)$ ,  $i = 1, \dots, n$ , with independent terminal values  $Y_T^i = \xi^i$ ,  $i = 1, \dots, n$ , be independent copies of  $\Theta := (Y, Z, U, K)$ , the solution of (2.1). More precisely, for each  $i = 1, \dots, n$ ,  $\Theta^i$  is the unique solution of the reflected MF-BSDE

$$\begin{cases} Y_t^i = \xi^i + \int_t^T f(s, Y_s^i, Z_s^i, U_s^i, \mathbb{P}_{Y_s^i}) ds + K_T^i - K_t^i - \int_t^T Z_s^i dB_s^i - \int_t^T \int_{R^*} U_s^i(e) \tilde{N}^i(ds, de), \\ Y_t^i \geq h(t, Y_t^i, \mathbb{P}_{Y_t^i}), \quad \forall t \in [0, T], \\ \int_0^T (Y_{t^-}^i - h(t^-, Y_{t^-}^i, \mathbb{P}_{Y_{t^-}^i})) dK_t^i = 0. \end{cases} \quad (4.12)$$

In the sequel, we denote  $(f \circ Y^i)(t, y, z) := f(t, y, z, \mathbb{P}_{Y^i})$ .

For any fixed  $1 \leq k \leq n$ , let

$$\mathbb{P}^{k,n} := \text{Law}(\Theta^{1,n}, \Theta^{2,n}, \dots, \Theta^{k,n}), \quad \mathbb{P}_{\Theta}^{\otimes k} := \text{Law}(\Theta^1, \Theta^2, \dots, \Theta^k).$$

be the joint probability laws of the processes  $(\Theta^{1,n}, \Theta^{2,n}, \dots, \Theta^{k,n})$  and  $(\Theta^1, \Theta^2, \dots, \Theta^k)$ , respectively.

From the definition of the distance  $D_G$ , we obtain the following inequality.

$$D_G^p(\mathbb{P}^{k,n}, \mathbb{P}_{\Theta}^{\otimes k}) \leq k \sup_{i \leq k} \left( \|Y^{i,n} - Y^i\|_{\mathcal{S}^p}^p + \|Z^{i,n} - Z^i \mathbf{e}_i\|_{\mathcal{H}^{p,n}}^p + \|U^{i,n} - U^i \mathbf{e}_i\|_{\mathcal{H}^{p,n}}^p + \|K^{i,n} - K^i\|_{\mathcal{S}^p}^p \right), \quad (4.13)$$

where for each  $i = 1, \dots, n$ ,  $\mathbf{e}_i := (0, \dots, 0, \underbrace{1}_i, 0, \dots, 0)$ .

Before we state and prove the propagation of chaos result, we derive the following convergence result of the empirical laws of i.i.d. copies  $Y^1, Y^2, \dots, Y^n$  solutions of (4.12) and convergence of the particle system (3.1) to the solution to (2.1).

**Theorem 4.1** (Law of Large Numbers). *Let  $Y^1, Y^2, \dots, Y^n$  with terminal values  $Y_T^i = \xi^i$  be independent copies of the solution  $Y$  of (2.1). Then, we have*

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[ \sup_{0 \leq t \leq T} \mathcal{W}_p^p(L_n[\mathbf{Y}_t], \mathbb{P}_{Y_t}) \right] = 0. \quad (4.14)$$

**Proof.** For any joint law  $R^n(dx, dy)$  with marginals  $L_n[\mathbf{Y}]$  and  $\mathbb{P}_Y$  and any  $\delta > 0$ , we have

$$\begin{aligned} \mathcal{W}_p^p(L_n[\mathbf{Y}_t], \mathbb{P}_{Y_t}) &\leq \int_{\mathbb{D} \times \mathbb{D}} |x(t) - y(t)|^p R^n(dx, dy) \\ &= \int_{\mathbb{D} \times \mathbb{D}} |x(t) - y(t)|^p \mathbf{1}_{\{d^\circ(x,y) \leq \ln(1+\delta/T)\}} R^n(dx, dy) \\ &\quad + \int_{\mathbb{D} \times \mathbb{D}} |x(t) - y(t)|^p \mathbf{1}_{\{d^\circ(x,y) > \ln(1+\delta/T)\}} R^n(dx, dy). \end{aligned}$$

Integrating both sides of (4.5) against the joint law  $R^n(dx, dy)$ , we obtain

$$\begin{aligned} &3^{1-p} \int_{\mathbb{D} \times \mathbb{D}} |x(t) - y(t)|^p \mathbf{1}_{\{d^\circ(x,y) \leq \ln(1+\delta/T)\}} R^n(dx, dy) \\ &\leq \delta^p + 2^p \int_{\mathbb{D}} w'_y(2\delta)^p \mathbb{P}_Y(dy) + \int_{\mathbb{D}} \sup_{(t-\delta) \vee 0 < s \leq t+\delta} |y(s) - y(s^-)|^p \mathbb{P}_Y(dy). \end{aligned}$$

Noting that

$$\int_{\mathbb{D}} \sup_{(t-\delta) \vee 0 < s \leq t+\delta} |y(s) - y(s^-)|^p \mathbb{P}_Y(dy) = \mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < s \leq t+\delta} |\Delta Y(s)|^p \right]$$

where  $\Delta Y(s) := Y(s) - Y(s^-)$  denotes the jump size of the process  $Y$  solution to (2.1), we have

$$\begin{aligned} \mathcal{W}_p^p(L_n[\mathbf{Y}_t], \mathbb{P}_{Y_t}) &\leq \int_{\mathbb{D} \times \mathbb{D}} |x(t) - y(t)|^p \mathbf{1}_{\{d^\circ(x,y) > \ln(1+\delta/T)\}} R^n(dx, dy) + 3^{p-1} \delta^p \\ &\quad + 6^p \int_{\mathbb{D}} w'_y(2\delta)^p \mathbb{P}_Y(dy) + 3^{p-1} \mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < s \leq t+\delta} |\Delta Y(s)|^p \right]. \end{aligned}$$

Since, for any  $M > 0$ , it holds that

$$\max(|x - y|_T^p - M^p, 0) \leq 2^p |x|_T^p \mathbf{1}_{\{|x|_T \geq M/2\}} + 2^p |y|_T^p \mathbf{1}_{\{|y|_T \geq M/2\}},$$

we have

$$\begin{aligned}
& \int_{\mathbb{D} \times \mathbb{D}} |x - y|_T^p \mathbf{1}_{\{d^\circ(x,y) > \ln(1+\delta/T)\}} R^n(dx, dy) \\
& \leq \int_{\mathbb{D} \times \mathbb{D}} \min(|x - y|_T, M)^p \mathbf{1}_{\{d^\circ(x,y) > \ln(1+\delta/T)\}} R^n(dx, dy) \\
& \quad + 2^p \int_{\mathbb{D} \times \mathbb{D}} |x|_T^p \mathbf{1}_{\{|x|_T \geq M/2, d^\circ(x,y) > \ln(1+\delta/T)\}} R^n(dx, dy) \\
& \quad + 2^p \int_{\mathbb{D} \times \mathbb{D}} |y|_T^p \mathbf{1}_{\{|y|_T \geq M/2, d^\circ(x,y) > \ln(1+\delta/T)\}} R^n(dx, dy) \\
& \leq M^p \int_{\mathbb{D} \times \mathbb{D}} \mathbf{1}_{\{d^\circ(x,y) > \ln(1+\delta/T)\}} R^n(dx, dy) + 2^p \int_{\mathbb{D}} |x|_T^p \mathbf{1}_{\{|x|_T \geq M/2\}} L_n[\mathbf{Y}](dx) \\
& \quad + 2^p \int_{\mathbb{D}} |y|_T^p \mathbf{1}_{\{|y|_T \geq M/2\}} \mathbb{P}_Y(dy).
\end{aligned}$$

Therefore, for any  $M > 0, \delta > 0$ ,

$$\begin{aligned}
\sup_{0 \leq t \leq T} \mathcal{W}_p^p(L_n[\mathbf{Y}_t], \mathbb{P}_{Y_t}) & \leq M^p \int_{\mathbb{D} \times \mathbb{D}} \mathbf{1}_{\{d^\circ(x,y) > \ln(1+\delta/T)\}} R^n(dx, dy) \\
& \quad + 2^p \int_{\mathbb{D}} |x|_T^p \mathbf{1}_{\{|x|_T \geq M/2\}} L_n[\mathbf{Y}](dx) + 2^p \int_{\mathbb{D}} |y|_T^p \mathbf{1}_{\{|y|_T \geq M/2\}} \mathbb{P}_Y(dy) \\
& \quad + 3^{p-1} \delta^p + 6^p \int_{\mathbb{D}} w'_y(2\delta)^p \mathbb{P}_Y(dy) + 3^{p-1} \sup_{0 \leq t \leq T} \mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < s \leq t+\delta} |\Delta Y(s)|^p \right].
\end{aligned} \tag{4.15}$$

By (4.8) in Billingsley [4],  $\lim_{\delta \rightarrow 0} w'_y(2\delta) = 0$ , and since

$$\int_{\mathbb{D}} w'_y(2\delta)^p \mathbb{P}_Y(dy) \leq 2\mathbb{E}[|Y|_T^p] < \infty,$$

by Lebesgue's dominated convergence theorem, we obtain

$$\lim_{\delta \rightarrow \infty} \int_{\mathbb{D}} w'_y(2\delta)^p \mathbb{P}_Y(dy) = 0.$$

Moreover, for any  $\delta > 0$ ,

$$\lim_{n \rightarrow \infty} \int_{\mathbb{D} \times \mathbb{D}} |x - y|_T^p \mathbf{1}_{\{d^\circ(x,y) > \ln(1+\delta/T)\}} R^n(dx, dy) = 0, \quad \text{a.s.}$$

Indeed, in view of (4.10), for any  $\delta > 0$ , we have

$$\lim_{n \rightarrow \infty} \int_{\mathbb{D} \times \mathbb{D}} \mathbf{1}_{\{d^\circ(x,y) > \ln(1+\delta/T)\}} R^n(dx, dy) \leq (\ln(1 + \delta/T))^{-1} \lim_{n \rightarrow \infty} D_T^\circ(L_n[\mathbf{Y}], \mathbb{P}_Y)^p = 0, \quad \text{a.s.}$$

Moreover, by the SLLN,

$$\lim_{n \rightarrow \infty} \int_{\mathbb{D}} |x|_T^p \mathbf{1}_{\{|x|_T \geq M/2\}} L_n[\mathbf{Y}](dx) = \int_{\mathbb{D}} |y|_T^p \mathbf{1}_{\{|y|_T \geq M/2\}} \mathbb{P}_Y(dy)$$

and since  $\mathbb{P}_Y \in \mathcal{P}_p(\mathbb{D})$ , it is uniformly integrable. In particular, we have

$$\lim_{M \rightarrow \infty} \int_{\mathbb{D}} |y|_T^p \mathbf{1}_{\{|y|_T \geq M/2\}} \mathbb{P}_Y(dy) = 0.$$

Therefore, for any  $\delta > 0$

$$\lim_{n \rightarrow \infty} \int_{\mathbb{D} \times \mathbb{D}} |x - y|_T \mathbf{1}_{\{d^\circ(x,y) > \ln(1+\delta/T)\}} R^n(dx, dy) \leq 2^{p+1} \lim_{M \rightarrow \infty} \int_{\mathbb{D}} |y|_T^p \mathbf{1}_{\{|y|_T \geq M/2\}} \mathbb{P}_Y(dy) = 0 \quad \text{a.s.}$$

Finally, by taking the limits  $n \rightarrow \infty$ , then  $M \rightarrow \infty$  and lastly  $\delta \rightarrow 0$ , we obtain

$$\lim_{n \rightarrow \infty} \sup_{0 \leq t \leq T} \mathcal{W}_p^p(L_n[\mathbf{Y}_t], \mathbb{P}_{Y_t}) \leq 3^{p-1} \lim_{\delta \rightarrow 0} \sup_{0 \leq t \leq T} \mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < s \leq t+\delta} |\Delta Y(s)|^p \right], \quad \text{a.s.} \tag{4.16}$$

Once we can show that

$$\lim_{\delta \rightarrow 0} \sup_{0 \leq t \leq T} \mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < s \leq t+\delta} |\Delta Y(s)|^p \right] = 0, \quad (4.17)$$

we obtain

$$\lim_{n \rightarrow \infty} \sup_{0 \leq t \leq T} \mathcal{W}_p^p(L_n[\mathbf{Y}_t], \mathbb{P}_{Y_t}) = 0 \quad \text{a.s.}$$

But, since

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} \mathcal{W}_p^p(L_n[\mathbf{Y}_t], \mathbb{P}_{Y_t}) \right] \leq \mathbb{E}[D_T^p(L_n[\mathbf{Y}], \mathbb{P}_Y)] \leq 2^{p-1} \left( \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n |Y^i|_T^p \right] + \|\mathbb{P}_Y\|_p^p \right) = 2^p \|\mathbb{P}_Y\|_p^p,$$

the random variable  $\sup_{0 \leq t \leq T} \mathcal{W}_p^p(L_n[\mathbf{Y}_t], \mathbb{P}_{Y_t})$  is uniformly integrable. Thus, it converges to 0 in  $L^1$ , i.e.

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[ \sup_{0 \leq t \leq T} \mathcal{W}_p^p(L_n[\mathbf{Y}_t], \mathbb{P}_{Y_t}) \right] = 0.$$

It remains to show (4.17). By Assumption (4.1)(iv), it follows that the process  $K_t$  is continuous, which implies that  $\Delta Y_t = \Delta J_t$ , where  $J_t := \int_{(0,t]} \int_{\mathbb{R}^*} U_s(e) N(ds, de)$ .

- Estimation of  $\Delta J_s$  (for the case  $p = 2$ ). Clearly,

$$\Delta J_t = U_t(\Delta \eta_t) \mathbf{1}_{\mathbb{R}^*}(\Delta \eta_t),$$

where  $(\eta_s)_s$  denotes the Lévy process for which

$$N(t, A) := \#\{s \in (0, t], \Delta \eta_t \in A\}, \quad A \subset \mathbb{R}^*.$$

We have

$$\sup_{(t-\delta) \vee 0 < s \leq t+\delta} |\Delta J_s|^2 = \sup_{(t-\delta) \vee 0 < s \leq t+\delta} |U_s(\Delta \eta_s)|^2 \mathbf{1}_{\mathbb{R}^*}(\Delta \eta_s) \leq \sum_{(t-\delta) \vee 0 < s \leq t+\delta} |U_s(\Delta \eta_s)|^2 \mathbf{1}_{\mathbb{R}^*}(\Delta \eta_s)$$

that is

$$\sup_{(t-\delta) \vee 0 < t \leq t+\delta} |\Delta J_s|^2 \leq \int_{(0,T] \times \mathbb{R}^*} \mathbf{1}_{((t-\delta) \vee 0, t+\delta]}(s) |U_s(e)|^2 N(ds, de).$$

Since  $U \in \mathcal{H}_\nu^2$ , we have

$$\mathbb{E} \left[ \int_{(0,T] \times \mathbb{R}^*} \mathbf{1}_{((t-\delta) \vee 0, t+\delta]}(s) |U_s(e)|^2 N(ds, de) \right] = \mathbb{E} \left[ \int_{(t-\delta) \vee 0}^{t+\delta} |U_s|_\nu^2 ds \right].$$

Therefore, for any  $R > 0$ ,

$$\mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < t \leq t+\delta} |\Delta J_s|^2 \right] \leq R^2 \delta + \mathbb{E} \left[ \int_0^T |U_s|_\nu^2 \mathbf{1}_{\{|U_s|_\nu > R\}} ds \right],$$

so

$$\sup_{0 \leq t \leq T} \mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < t \leq t+\delta} |\Delta J_s|^2 \right] \leq R^2 \delta + \mathbb{E} \left[ \int_0^T |U_s|_\nu^2 \mathbf{1}_{\{|U_s|_\nu > R\}} ds \right]. \quad (4.18)$$

Thus,

$$\lim_{\delta \rightarrow 0} \sup_{0 \leq t \leq T} \mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < t \leq t+\delta} |\Delta J_s|^2 \right] \leq \mathbb{E} \left[ \int_0^T |U_s|_\nu^2 \mathbf{1}_{\{|U_s|_\nu > R\}} ds \right].$$

Again, since  $U \in \mathcal{H}_\nu^2$ ,  $\lim_{R \rightarrow \infty} \mathbb{E} \left[ \int_0^T |U_s|_\nu^2 \mathbf{1}_{\{|U_s|_\nu > R\}} ds \right] = 0$ . Hence,

$$\lim_{\delta \rightarrow 0} \sup_{0 \leq t \leq T} \mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < t \leq t+\delta} |\Delta J_s|^2 \right] = 0.$$

• Estimation of  $\Delta J_s$  (for the case  $p > 2$ )

Since the process  $(K_t)$  is continuous, we obtain, for any  $L > 0$ ,

$$\begin{aligned} \sup_{(t-\delta) \vee 0 < s \leq t+\delta} |\Delta J_s|^p &\leq 2^{p-2} \sup_{(t-\delta) \vee 0 < s \leq t+\delta} |\Delta J_s|^2 \sup_{0 \leq s \leq T} |Y_s|^{p-2} \mathbf{1}_{\{\sup_{0 \leq s \leq T} |Y_s| \leq L\}} + 2^p \sup_{0 \leq s \leq T} |Y_s|^p \mathbf{1}_{\{\sup_{0 \leq s \leq T} |Y_s| > L\}} \\ &\leq 2^{p-2} L^{p-2} \sup_{(t-\delta) \vee 0 < s \leq t+\delta} |\Delta J_s|^2 + 2^p \sup_{0 \leq s \leq T} |Y_s|^p \mathbf{1}_{\{\sup_{0 \leq s \leq T} |Y_s| > L\}}. \end{aligned}$$

Therefore, in view of (4.18), for any  $L > 0$  and  $R > 0$ ,

$$\begin{aligned} \mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < t \leq t+\delta} |\Delta J_s|^p \right] &\leq 2^{p-2} L^{p-2} \mathbb{E} \left[ \int_{(t-\delta) \vee 0}^{t+\delta} |U_s|_\nu^2 ds \right] + 2^{p-2} \mathbb{E} \left[ \sup_{0 \leq s \leq T} |Y_s|^p \mathbf{1}_{\{\sup_{0 \leq s \leq T} |Y_s| > L\}} \right] \\ &\leq 2^{p-2} L^{p-2} \left( R^2 \delta + \mathbb{E} \left[ \int_0^T |U_s|_\nu^2 \mathbf{1}_{\{|U_s|_\nu > R\}} ds \right] \right) \\ &\quad + 2^{p-2} \mathbb{E} \left[ \sup_{0 \leq s \leq T} |Y_s|^p \mathbf{1}_{\{\sup_{0 \leq s \leq T} |Y_s| > L\}} \right], \end{aligned}$$

which implies

$$\begin{aligned} \sup_{0 \leq t \leq T} \mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < t \leq t+\delta} |\Delta J_s|^p \right] &\leq 2^{p-2} L^{p-2} \left( R^2 \delta + \mathbb{E} \left[ \int_0^T |U_s|_\nu^2 \mathbf{1}_{\{|U_s|_\nu > R\}} ds \right] \right) \\ &\quad + 2^{p-2} \mathbb{E} \left[ \sup_{0 \leq s \leq T} |Y_s|^p \mathbf{1}_{\{\sup_{0 \leq s \leq T} |Y_s| > L\}} \right]. \end{aligned}$$

Thus,

$$\begin{aligned} \lim_{\delta \rightarrow 0} \sup_{0 \leq t \leq T} \mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < t \leq t+\delta} |\Delta J_s|^p \right] &\leq 2^{p-2} L^{p-2} \mathbb{E} \left[ \int_0^T |U_s|_\nu^2 \mathbf{1}_{\{|U_s|_\nu > R\}} ds \right] \\ &\quad + 2^{p-2} \mathbb{E} \left[ \sup_{0 \leq s \leq T} |Y_s|^p \mathbf{1}_{\{\sup_{0 \leq s \leq T} |Y_s| > L\}} \right]. \end{aligned}$$

Since  $Y \in \mathcal{S}^p$  and  $U \in \mathcal{H}_\nu^p$ ,

$$\lim_{R \rightarrow \infty} \mathbb{E} \left[ \int_0^T |U_s|_\nu^2 \mathbf{1}_{\{|U_s|_\nu > R\}} ds \right] = 0, \quad \lim_{L \rightarrow \infty} \mathbb{E} \left[ \sup_{0 \leq s \leq T} |Y_s|^p \mathbf{1}_{\{\sup_{0 \leq s \leq T} |Y_s| > L\}} \right] = 0.$$

Hence, by taking first the limit in  $R \rightarrow \infty$  and then  $L \rightarrow \infty$ , we obtain

$$\lim_{\delta \rightarrow 0} \sup_{0 \leq t \leq T} \mathbb{E} \left[ \sup_{(t-\delta) \vee 0 < t \leq t+\delta} |\Delta J_s|^p \right] = 0.$$

□

We now provide the following convergence result for the solution  $Y^{i,n}$  of (3.1).

**Proposition 4.2** (Convergence of the  $Y^{i,n}$ 's). *Assume that for some  $p \geq 2$ ,  $\gamma_1$  and  $\gamma_2$  satisfy*

$$2^{2p-1}(\gamma_1^p + \gamma_2^p) < 1. \quad (4.19)$$

Then, under Assumptions 2.1, 3.1 and 4.1, we have

$$\lim_{n \rightarrow \infty} \sup_{0 \leq t \leq T} E[|Y_t^{i,n} - Y_t^i|^p] = 0. \quad (4.20)$$

In particular,

$$\lim_{n \rightarrow \infty} \|Y^{i,n} - Y^i\|_{\mathcal{H}^{p,1}} = 0. \quad (4.21)$$

**Proof.** Let  $\vartheta \in \mathcal{T}_0^n$ . By the estimates on BSDEs in Proposition A.1, we have

$$\begin{aligned} & |Y_\vartheta^{i,n} - Y_\vartheta^i|^p \\ & \leq \text{ess sup}_{\tau \in \mathcal{T}_\vartheta^n} \left| \mathcal{E}_{\vartheta, \tau}^{\mathbf{f}^i \circ \mathbf{Y}^n} [h(\tau, Y_\tau^{i,n}, L_n[\mathbf{Y}_s^n]) \mathbf{1}_{\{\tau < T\}} + \xi^{i,n} \mathbf{1}_{\{\tau = T\}}] - \mathcal{E}_{\vartheta, \tau}^{\mathbf{f}^i \circ Y^i} [h(\tau, Y_\tau^i, \mathbb{P}_{Y_s | s = \tau}) \mathbf{1}_{\{\tau < T\}} + \xi^i \mathbf{1}_{\{\tau = T\}}] \right|^p \\ & \leq \text{ess sup}_{\tau \in \mathcal{T}_\vartheta^n} \mathbb{E} \left[ \int_\vartheta^\tau e^{\beta(s-\vartheta)} \eta^p |(\mathbf{f}^i \circ \mathbf{Y}^n)(s, \widehat{Y}_s^{i,\tau}, \widehat{Z}_s^{i,\tau}, \widehat{U}_s^{i,\tau}) - (\mathbf{f}^i \circ Y^i)(s, \widehat{Y}_s^{i,\tau}, \widehat{Z}_s^{i,\tau}, \widehat{U}_s^{i,\tau})|^p ds \right. \\ & \quad \left. + (e^{\beta(\tau-\vartheta)} |h(\tau, Y_\tau^{i,n}, L_n[\mathbf{Y}_s^n]) - h(\tau, Y_\tau^i, \mathbb{P}_{Y_s | s = \tau})| + e^{\beta(T-\vartheta)} |\xi^{i,n} - \xi^i|)^p | \mathcal{F}_\vartheta \right] \\ & \leq \text{ess sup}_{\tau \in \mathcal{T}_\vartheta^n} \mathbb{E} \left[ \int_\vartheta^\tau e^{\beta(s-\vartheta)} \eta^p C_f^p \mathcal{W}_p^p(L_n[\mathbf{Y}_s^n], \mathbb{P}_{Y_s}) ds + (\gamma_1 e^{\beta(\tau-\vartheta)} |Y_\tau^{i,n} - Y_\tau^i| \right. \\ & \quad \left. + \gamma_2 e^{\beta(\tau-\vartheta)} \mathcal{W}_p(L_n[\mathbf{Y}_\tau^n], \mathbb{P}_{Y_s | s = \tau}) + e^{\beta(T-\vartheta)} |\xi^{i,n} - \xi^i| \mathbf{1}_{\{\tau = T\}})^p | \mathcal{F}_\vartheta \right], \end{aligned} \quad (4.22)$$

where  $\eta, \beta > 0$  such that  $\eta \leq \frac{2c(p)}{pC_f^2}$  and  $\beta \geq C_f + \frac{1}{\eta} + \frac{p-1}{p} \left(\frac{1}{\eta}\right)^{\frac{p}{p-1}}$ ,  $(\widehat{Y}^{i,\tau}, \widehat{Z}^{i,\tau}, \widehat{U}^{i,\tau})$  is the solution of the BSDE associated with driver  $\mathbf{f}^i \circ \mathbf{Y}^n$  and terminal condition  $h(\tau, Y_\tau^{i,n}, L_n[\mathbf{Y}_s^n]_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi^{i,n} \mathbf{1}_{\{\tau = T\}}$ .

Therefore, we have

$$e^{p\beta\vartheta} |Y_\vartheta^{i,n} - Y_\vartheta^i|^p \leq \text{ess sup}_{\tau \in \mathcal{T}_\vartheta^n} \mathbb{E}[G_{\vartheta, \tau}^{i,n} | \mathcal{F}_\vartheta],$$

where

$$\begin{aligned} G_{\vartheta, \tau}^{i,n} & := \int_\vartheta^\tau 2^{p-1} \eta^p C_f^p \left( \frac{1}{n} \sum_{j=1}^n e^{p\beta s} |Y_s^{j,n} - Y_s^j|^p \right) ds \\ & \quad + 2^{2p-2} (\gamma_1^p + \gamma_2^p) \left( e^{p\beta\tau} |Y_\tau^{i,n} - Y_\tau^i|^p + \frac{1}{n} \sum_{j=1}^n e^{p\beta\tau} |Y_\tau^{j,n} - Y_\tau^j|^p \right) \\ & \quad + \left( 2^{2p-2} \gamma_2^p + 2^{p-1} T \eta^p C_f^p \right) \sup_{0 \leq t \leq T} e^{p\beta s} \mathcal{W}_p^p(L_n[\mathbf{Y}_s], \mathbb{P}_{Y_s}) + 2^{2p-2} e^{\beta T} |\xi^{i,n} - \xi^i|^p \mathbf{1}_{\{\tau = T\}}. \end{aligned}$$

Setting  $V_t^{n,p} := \frac{1}{n} \sum_{j=1}^n e^{p\beta t} |Y_t^{j,n} - Y_t^j|^p$  and

$$\Gamma_{n,p} := \left( 2^{2p-2} \gamma_2^p + 2^{p-1} T \eta^p C_f^p \right) \sup_{0 \leq t \leq T} e^{p\beta s} \mathcal{W}_p^p(L_n[\mathbf{Y}_s], \mathbb{P}_{Y_s}) + 2^{2p-2} e^{\beta T} |\xi^{i,n} - \xi^i|^p \mathbf{1}_{\{\tau = T\}}.$$

we obtain

$$V_\vartheta^{n,p} \leq \text{ess sup}_{\tau \in \mathcal{T}_\vartheta^n} \mathbb{E} \left[ \int_\vartheta^\tau 2^{p-1} \eta^p C_f^p V_s^{n,p} ds + 2^{2p-1} (\gamma_1^p + \gamma_2^p) V_\tau^{n,p} + \Gamma_{n,p} | \mathcal{F}_\vartheta \right]. \quad (4.23)$$

In view of Lemma D.1 in [25], we have

$$\mathbb{E}[V_\vartheta^{n,p}] \leq 2^{2p-1}(\gamma_1^p + \gamma_2^p) \sup_{\tau \in \mathcal{T}_\vartheta^n} \mathbb{E}[V_\tau^{n,p}] + \mathbb{E}\left[\int_\vartheta^T 2^{p-1} \eta^p C_f^p V_s^{n,p} ds + \Gamma_{n,p}\right].$$

Since  $\mathcal{T}_\vartheta^n \subset \mathcal{T}_0^n$  and by arbitrariness of  $\vartheta \in \mathcal{T}_0^n$ , we derive

$$\lambda \sup_{0 \leq t \leq T} \mathbb{E}[V_t^{n,p}] \leq \lambda \sup_{\vartheta \in \mathcal{T}_0^n} \mathbb{E}[V_\vartheta^{n,p}] \leq \mathbb{E}\left[\int_0^T 2^{p-1} \eta^p C_f^p V_s^{n,p} ds + \Gamma_{n,p}\right],$$

where  $\lambda := 1 - 2^{2p-1}(\gamma_1^p + \gamma_2^p) > 0$  by the assumption (4.19). By Gronwall inequality we obtain

$$\sup_{0 \leq t \leq T} \mathbb{E}[V_t^{n,p}] \leq e^{K_p} \mathbb{E}[\Gamma_{n,p}]$$

where  $K(p) := \frac{1}{\lambda} 2^{2p-1} \eta^p C_f^p T$ . But, in view of the exchangeability of the processes  $(Y^{i,n}, Y^i)$ ,  $i = 1, \dots, n$  (see Proposition 4.1), we have,  $\mathbb{E}[V_t^{n,p}] = \mathbb{E}[e^{p\beta t} |Y_t^{i,n} - Y_t^i|^p]$ . Thus,

$$\sup_{0 \leq t \leq T} \mathbb{E}[e^{p\beta t} |Y_t^{i,n} - Y_t^i|^p] \leq e^{K_p} \mathbb{E}[\Gamma_{n,p}] \rightarrow 0$$

as  $n \rightarrow \infty$  in view of Theorem 4.1 and Assumption 4.1, as required.  $\square$

**Theorem 4.2** (Propagation of chaos of the  $Y^{i,n}$ 's). *Under the assumptions of Proposition 4.2, the solution  $Y^{i,n}$  of the particle system (3.1) satisfies the propagation of chaos property, i.e. for any fixed positive integer  $k$ ,*

$$\lim_{n \rightarrow \infty} \text{Law}(Y^{1,n}, Y^{2,n}, \dots, Y^{k,n}) = \text{Law}(Y^1, Y^2, \dots, Y^k).$$

**Proof.** Set  $P^{k,n} := \text{law}(Y^{1,n}, Y^{2,n}, \dots, Y^{k,n})$  and  $P^{\otimes k} := \text{Law}(Y^1, Y^2, \dots, Y^k)$ . Consider the Wasserstein metric on  $\mathcal{P}_2(\mathbb{H}^2)$  defined by

$$D_{\mathbb{H}^2}(P, Q) = \inf \left\{ \left( \int_{\mathbb{H}^2 \times \mathbb{H}^2} \|y - y'\|_{\mathbb{H}^2}^2 R(dy, dy') \right)^{1/2} \right\}, \quad (4.24)$$

over  $R \in \mathcal{P}(\mathbb{H}^2 \times \mathbb{H}^2)$  with marginals  $P$  and  $Q$ . Note that, since  $p \geq 2$ , it is enough to show for  $D_{\mathbb{H}^2}(P, Q)$ . Since  $\mathbb{H}^2$  is a Polish space,  $(\mathcal{P}_2(\mathbb{H}^2), D_{\mathbb{H}^2})$  is a Polish space and induces the topology of weak convergence (see Theorem 6.9 in [43]). Thus, we obtain the propagation of chaos property for the  $Y^{i,n}$ 's if we can show that  $\lim_{n \rightarrow \infty} D_{\mathbb{H}^2}(P^{k,n}, P^{\otimes k}) = 0$ . But, this follows from the fact that

$$D_{\mathbb{H}^2}^2(P^{k,n}, P^{\otimes k}) \leq k \sup_{i \leq k} \|Y^{i,n} - Y^i\|_{\mathcal{H}^{2,1}}^2$$

and (4.21) for  $p = 2$ .  $\square$

In the next proposition we provide a convergence result for the whole solution  $(Y^{i,n}, Z^{i,n}, U^{i,n}, K^{i,n})$  of (3.1).

**Proposition 4.3.** *Assume that, for some  $p > 2$ ,  $\gamma_1$  and  $\gamma_2$  satisfy*

$$2^{2p-2}(\gamma_1^p + \gamma_2^p) < \left( \frac{p - \kappa}{2p} \right)^{p/\kappa} \quad (4.25)$$



for some  $\kappa \in [2, p)$ .

Then, under Assumptions 2.1, 3.1 and 4.1, we have

$$\lim_{n \rightarrow \infty} \left( \|Y^{i,n} - Y^i\|_{\mathcal{S}^p} + \|Z^{i,n} - Z^i \mathbf{e}_i\|_{\mathcal{H}^{p,n}} + \|U^{i,n} - U^i \mathbf{e}_i\|_{\mathcal{H}_f^{p,n}} + \|K^{i,n} - K^i\|_{\mathcal{S}^p} \right) = 0, \quad (4.26)$$

where  $\mathbf{e}_1, \dots, \mathbf{e}_n$  are unit vectors in  $\mathbb{R}^n$ .

**Remark 4.1.**

- If the driver  $f$  does not depend of the jump size  $U$ , the smallness condition (4.25) remains valid for any  $p \geq 2$  and any  $\kappa \in (1, p)$  in which case the limit result (4.26) remains valid even for  $p = 2$ .
- If the obstacle  $h$  is independent of the solution  $Y$ , the limit result (4.26) remains valid for  $p \geq 2$  without the condition (4.25).

**Proof.** Step 1. Let us first show that  $\lim_{n \rightarrow \infty} \|Y^{i,n} - Y^i\|_{\mathcal{S}_\beta^p} = 0$ .

In view of (4.22), for any  $\kappa \geq 2$  and any  $t \leq T$ , we have

$$\begin{aligned} |Y_t^{i,n} - Y_t^i|^\kappa &\leq \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t^n} \mathbb{E} \left[ \int_t^\tau e^{\kappa\beta(s-t)} \eta^\kappa C_f^\kappa \mathcal{W}_p^\kappa(L_n[\mathbf{Y}_s^n], \mathbb{P}_{Y_s}) ds + (\gamma_1 e^{\beta(\tau-t)} |Y_\tau^{i,n} - Y_\tau^i| \right. \\ &\quad \left. + \gamma_2 e^{\beta(\tau-t)} \mathcal{W}_p(L_n[\mathbf{Y}_\tau^n], \mathbb{P}_{Y_s|s=\tau}) + e^{\beta(T-t)} |\xi^{i,n} - \xi^i| \mathbf{1}_{\{\tau=T\}})^\kappa \middle| \mathcal{F}_t \right], \end{aligned}$$

where  $\eta, \beta > 0$  are such that  $\eta \leq \frac{2c(\kappa)}{\kappa C_f^2}$  and  $\beta \geq C_f + \frac{1}{\eta} + \frac{\kappa-1}{\kappa} \left(\frac{1}{\eta}\right)^{\kappa-1}$ .

Therefore, for any  $p > \kappa \geq 2$ , we have

$$e^{p\beta t} |Y_t^{i,n} - Y_t^i|^p \leq \mathbb{E}[\mathcal{G}_T^{i,n} | \mathcal{F}_t]^{p/\kappa},$$

where

$$\begin{aligned} \mathcal{G}_T^{i,n} &:= \int_0^T e^{\kappa\beta s} \eta^\kappa C_f^\kappa \mathcal{W}_p^\kappa(L_n[\mathbf{Y}_s^n], \mathbb{P}_{Y_s}) ds \\ &\quad + \left( \gamma_1 \sup_{0 \leq s \leq T} e^{\beta s} |Y_s^{i,n} - Y_s^i| + \gamma_2 \sup_{0 \leq s \leq T} e^{\beta s} \mathcal{W}_p(L_n[\mathbf{Y}_s^n], \mathbb{P}_{Y_s}) + e^{\beta T} |\xi^{i,n} - \xi^i| \right)^\kappa. \end{aligned}$$

Thus, by Doob's inequality, we get

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} e^{p\beta t} |Y_t^{i,n} - Y_t^i|^p \right] \leq \left( \frac{p}{p-\kappa} \right)^{p/\kappa} \mathbb{E} \left[ \left( \mathcal{G}_T^{i,n} \right)^{p/\kappa} \right]. \quad (4.27)$$

Therefore, we have

$$\begin{aligned} 2^{1-\frac{p}{\kappa}} \left( \mathcal{G}_T^{i,n} \right)^{p/\kappa} &\leq C_1 \int_0^T \sup_{0 \leq t \leq s} e^{p\beta t} \mathcal{W}_p^p(L_n[\mathbf{Y}_t^n], L_n[\mathbf{Y}_t]) ds \\ &\quad + 2^{p-1} \left( \gamma_1 \sup_{0 \leq t \leq T} e^{\beta t} |Y_t^{i,n} - Y_t^i| + \gamma_2 \sup_{0 \leq t \leq T} e^{\beta t} \mathcal{W}_p(L_n[\mathbf{Y}_s^n], L_n[\mathbf{Y}_s]) \right)^p + \Lambda_n \end{aligned}$$

where  $C_1 := 2^{p-1} T^{\frac{p}{\kappa}-1} \eta^p C_f^p$  and

$$\Lambda_n := C_1 T \sup_{0 \leq s \leq T} e^{p\beta s} \mathcal{W}_p(L_n[\mathbf{Y}_s], \mathbb{P}_{Y_s}) + 2^{p-1} \left( \gamma_2 \sup_{0 \leq s \leq T} e^{\beta s} \mathcal{W}_p(L_n[\mathbf{Y}_s], \mathbb{P}_{Y_s}) + e^{\beta T} |\xi^{i,n} - \xi^i| \right)^p.$$

But, in view of the exchangeability of the processes  $(Y^{i,n}, Y^i), i = 1, \dots, n$  (see Proposition 4.1), for each  $s \in [0, T]$  we have,

$$\mathbb{E} \left[ \sup_{0 \leq t \leq s} e^{\beta pt} \mathcal{W}_p^p(L_n[\mathbf{Y}_t^n], L_n[\mathbf{Y}_t]) \right] \leq \mathbb{E} \left[ \sup_{0 \leq t \leq s} e^{\beta pt} |Y_t^{i,n} - Y_t^i|^p \right], \quad i = 1, \dots, n,$$

and so

$$2^{1-\frac{p}{\kappa}} \mathbb{E} \left[ \left( \mathcal{G}_T^{i,n} \right)^{p/\kappa} \right] \leq C_1 \mathbb{E} \left[ \int_0^T \sup_{0 \leq t \leq s} e^{\beta pt} \mathcal{W}_p^p(L_n[\mathbf{Y}_t^n], L_n[\mathbf{Y}_t]) ds \right] + \mathbb{E}[\Lambda_n] \\ + 2^{2p-1} (\gamma_1^p + \gamma_2^p) \mathbb{E} \left[ \sup_{0 \leq t \leq T} e^{\beta pt} |Y_t^{i,n} - Y_t^i|^p \right].$$

Therefore, (4.27) becomes

$$\mu \mathbb{E} \left[ \sup_{0 \leq t \leq T} e^{\beta pt} |Y_t^{i,n} - Y_t^i|^p \right] \leq C_1 \mathbb{E} \left[ \int_0^T \sup_{0 \leq t \leq s} e^{\beta pt} |Y_t^{i,n} - Y_t^i|^p ds \right] + \mathbb{E}[\Lambda_n].$$

where  $\mu := 2^{1-\frac{p}{\kappa}} \left( \frac{p}{p-\kappa} \right)^{-p/\kappa} - 2^{2p-1} (\gamma_1^p + \gamma_2^p)$ . Using the condition (4.25), to see that  $\mu > 0$ , and Gronwall's inequality, we finally obtain

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} e^{\beta pt} |Y_t^{i,n} - Y_t^i|^p \right] \leq e^{\frac{C_1 T}{\mu}} \mathbb{E}[\Lambda_n].$$

Next, by (4.14) together with Assumption (4.1) (iii) we have

$$\lim_{n \rightarrow \infty} \mathbb{E}[\Lambda_n] = 0,$$

which yields the desired result.

Step 2. We now show that  $\lim_{n \rightarrow \infty} \|Z^{i,n} - Z^i \mathbf{e}_i\|_{\mathcal{H}^{p,n}} = 0$ ,  $\lim_{n \rightarrow \infty} \|U^{i,n} - U^i \mathbf{e}_i\|_{\mathcal{H}^{p,n}} = 0$  and  $\lim_{n \rightarrow \infty} \|K^{i,n} - K^i\|_{\mathcal{S}^p} = 0$ . We first prove that  $\lim_{n \rightarrow \infty} \|Z^{i,n} - Z^i \mathbf{e}_i\|_{\mathcal{H}^{p,n}} = 0$  and  $\lim_{n \rightarrow \infty} \|U^{i,n} - U^i \mathbf{e}_i\|_{\mathcal{H}^{p,n}} = 0$ . For  $s \in [0, T]$ , denote  $\delta Y_s^{i,n} := Y_s^{i,n} - Y_s^i$ ,  $\delta Z_s^{i,n} := Z_s^{i,n} - Z_s^i \mathbf{e}_i$ ,  $\delta U_s^{i,n} := U_s^{i,n} - U_s^i \mathbf{e}_i$ ,  $\delta K_s^{i,n} := K_s^{i,n} - K_s^i$ ,  $\delta f_s^{i,n} := f(s, Y_s^{i,n}, Z_s^{i,n}, U_s^{i,n}, L_n[\mathbf{Y}_s^n]) - f(s, Y_s^i, Z_s^i, U_s^i, \mathbb{P}_{Y_s^i})$ ,  $\delta \xi^{i,n} := \xi^{i,n} - \xi^i$  and  $\delta h_s^{i,n} := h(s, Y_s^{i,n}, L_n[\mathbf{Y}_s^n]) - h(s, Y_s^i, \mathbb{P}_{Y_s^i})$ . By applying Itô's formula to  $|\delta Y_t^{i,n}|^2$ , we obtain

$$|\delta Y_t^{i,n}|^2 + \int_t^T |\delta Z_s^{i,n}|^2 ds + \int_t^T \int_{\mathbb{R}^*} \sum_{j=1}^n |\delta U_s^{i,j,n}(e)|^2 N^j(ds, de) + \sum_{t < s \leq T} |\Delta K_s^{i,n} - \Delta K_s^i|^2 = |\delta \xi^{i,n}|^2 \\ + 2 \int_t^T \delta Y_s^{i,n} \delta f_s^{i,n} ds - 2 \int_t^T \delta Y_s^{i,n} \sum_{j=1}^n \delta Z_s^{i,j,n} dB_s^j - 2 \int_t^T \int_{\mathbb{R}^*} \delta Y_s^{i,n} \sum_{j=1}^n \delta U_s^{i,j,n}(e) \tilde{N}^j(ds, de) \\ + 2 \int_t^T \delta Y_s^{i,n} d(\delta K_s^{i,n}).$$

By standard estimates, from the assumptions on the driver  $f$ , we get, for all  $\varepsilon > 0$ ,

$$\int_t^T \delta Y_s^{i,n} \delta f_s^{i,n} ds \leq \int_t^T C_f |\delta Y_s^{i,n}|^2 ds + \int_t^T \frac{3}{\varepsilon} C_f^2 |\delta Y_s^{i,n}|^2 ds + \int_t^T \frac{\varepsilon}{4} \{ |\delta Z_s^{i,n}|^2 + |\delta U_s^{i,n}|_\nu^2 \} ds \\ + \int_t^T \frac{\varepsilon}{4} \mathcal{W}_p^2(L_n[\mathbf{Y}_s^n], \mathbb{P}_{Y_s^i}) ds.$$

It follows that, for a given constant  $C_p > 0$ , independent of  $n$ , we have

$$\begin{aligned}
& |\delta Y_0^{i,n}|^p + \left( \int_0^T |\delta Z_s^{i,n}|^2 ds \right)^{\frac{p}{2}} + \left( \int_0^T \int_{\mathbb{R}^*} \sum_{j=1}^n |\delta U_s^{i,j,n}(e)|^2 N^j(ds, de) \right)^{\frac{p}{2}} \leq C_p |\delta \xi^{i,n}|^p \\
& + C_p \left\{ \left( 2C_f + \frac{6}{\varepsilon} C_f^2 \right) T \right\}^{\frac{p}{2}} \sup_{0 \leq s \leq T} |\delta Y_s^{i,n}|^p + C_p \varepsilon^{\frac{p}{2}} \left( \int_0^T |\delta U_s^{i,n}|_\nu^2 ds \right)^{\frac{p}{2}} \\
& + C_p \left( \int_0^T \mathcal{W}_p^2(L_n[\mathbf{Y}_s^n], \mathbb{P}_{Y_s^i}) ds \right)^{\frac{p}{2}} + C_p \left\{ \left| \int_0^T \delta Y_s^{i,n} \sum_{j=1}^n \delta Z_s^{i,j,n} dB_s^j \right|^{\frac{p}{2}} \right. \\
& \left. + \left| \int_0^T \int_{\mathbb{R}^*} \delta Y_s^{i,n} \sum_{j=1}^n \delta U_s^{i,j,n}(e) \tilde{N}^j(ds, de) \right|^{\frac{p}{2}} + \left| \int_0^T \delta Y_s^{i,n} d(\delta K_s^{i,n}) \right|^{\frac{p}{2}} \right\}. \tag{4.28}
\end{aligned}$$

By applying the Burkholder-Davis-Gundy inequality, we derive that there exist some constants  $m_p > 0$  and  $l_p > 0$  such that

$$\begin{aligned}
C_p \mathbb{E} \left[ \left| \int_0^T \delta Y_s^{i,n} \sum_{j=1}^n \delta Z_s^{i,j,n} dB_s^j \right|^{\frac{p}{2}} \right] & \leq m_p \mathbb{E} \left[ \left( \int_0^T (\delta Y_s^{i,n})^2 |\delta Z_s^{i,n}|^2 ds \right)^{\frac{p}{4}} \right] \\
& \leq \frac{m_p^2}{4} \|\delta Y_s^{i,n}\|_{\mathcal{S}^p}^p + \frac{1}{2} \mathbb{E} \left[ \left( \int_0^T |\delta Z_s^{i,n}|^2 ds \right)^{\frac{p}{2}} \right]
\end{aligned}$$

and

$$\begin{aligned}
C_p \mathbb{E} \left[ \left| \int_0^T \int_{\mathbb{R}^*} \delta Y_s^{i,n} \sum_{j=1}^n \delta U_s^{i,j,n}(e) \tilde{N}^j(ds, de) \right|^{\frac{p}{2}} \right] & \leq l_p \mathbb{E} \left[ \left( \int_0^T (\delta Y_s^{i,n})^2 \int_{\mathbb{R}^*} \sum_{j=1}^n (\delta U_s^{i,j,n}(e))^2 N^j(ds, de) \right)^{\frac{p}{4}} \right] \\
& \leq \frac{l_p^2}{4} \|\delta Y_s^{i,n}\|_{\mathcal{S}^p}^p + \frac{1}{2} \mathbb{E} \left[ \left( \int_0^T \int_{\mathbb{R}^*} \sum_{j=1}^n (\delta U_s^{i,j,n}(e))^2 N^j(ds, de) \right)^{\frac{p}{2}} \right].
\end{aligned}$$

Also recall that, for some constant  $e_p > 0$  we have (cf. Eq. (1.3) in [32])

$$\mathbb{E} \left[ \left( \int_0^T \int_{\mathbb{R}^*} |\delta U_s^{i,n}(e)|^2 \nu(de) ds \right)^{\frac{p}{2}} \right] \leq e_p \mathbb{E} \left[ \left( \int_0^T \int_{\mathbb{R}^*} \sum_{j=1}^n |\delta U_s^{i,j,n}(e)|^2 N^j(de, ds) \right)^{\frac{p}{2}} \right].$$

Now, we take the expectation in (4.28), by using the above inequalities and taking  $\varepsilon > 0$  small enough, we obtain

$$\begin{aligned}
& \mathbb{E} \left[ \left( \int_0^T |\delta Z_s^{i,n}|^2 ds \right)^{\frac{p}{2}} + \left( \int_0^T \|\delta U_s^{i,n}\|_\nu^2 ds \right)^{\frac{p}{2}} \right] \leq C_p |\delta \xi^{i,n}|^p + K_{C_f, \varepsilon, T, p} \|\delta Y_s^{i,n}\|_{\mathcal{S}^p}^p \\
& + C_p \mathbb{E} \left[ \sup_{0 \leq s \leq T} \mathcal{W}_p^p(L_n[\mathbf{Y}_s^n], \mathbb{P}_{Y_s^i}) \right] + \mathbb{E} \left[ \left( \sup_{0 \leq s \leq T} |\delta Y_s^{i,n}| (K_T^{i,n} + K_T^i) \right)^{\frac{p}{2}} \right].
\end{aligned}$$

The above inequality, together with the equations

$$\begin{aligned}
K_T^{i,n} & = Y_0^{i,n} - \xi^{i,n} - \int_0^T f(s, Y_s^{i,n}, Z_s^{i,i,n}, U_s^{i,i,n}, L_n[\mathbf{Y}_s^n]) ds \\
& + \sum_{j=1}^n \int_0^T Z_s^{i,j,n} dB_s^j + \int_0^T \int_{\mathbb{R}^*} \sum_{j=1}^n U_s^{i,j,n}(e) \tilde{N}^j(ds, de) \tag{4.29}
\end{aligned}$$

and

$$K_T^i = Y_0^i - \xi^i - \int_0^T f(s, Y_s^i, Z_s^i, U_s^i, \mathbb{P}_{Y_s^i}) ds + \int_0^T Z_s^i dB_s^i + \int_0^T \int_{\mathbb{R}^*} U_s^i(e) \tilde{N}^i(ds, de), \quad (4.30)$$

lead to

$$\begin{aligned} & \mathbb{E} \left[ \left( \int_0^T |\delta Z_s^{i,n}|^2 ds \right)^{\frac{p}{2}} + \left( \int_0^T \int_{\mathbb{R}^*} |\delta U_s^{i,n}(e)|^2 \nu(de) ds \right)^{\frac{p}{2}} \right] \\ & \leq C_p \left( \|\delta Y^{i,n}\|_{\mathcal{S}^p}^p + \mathbb{E} \left[ \sup_{0 \leq s \leq T} \mathcal{W}_p^p(L_n[\mathbf{Y}_s^n], \mathbb{P}_{Y_s^i}) \right] + |\delta \xi^{i,n}|^p + \|\delta Y^{i,n}\|_{\mathcal{S}^p}^p \Psi_p^{i,n} \right), \end{aligned} \quad (4.31)$$

with

$$\Psi_p^{i,n} := \mathbb{E} \left[ \|Y^{i,n}\|_{\mathcal{S}^p}^p + \|Y^i\|_{\mathcal{S}^p}^p + |\xi^{i,n}|^p + |\xi^i|^p + \int_0^T |f(s, 0, 0, 0, \delta_0)|^p ds + \sup_{0 \leq s \leq T} |h(s, 0, \delta_0)|^p \right],$$

for a different constant  $C_p$ .

From Step 1, we have  $\|\delta Y^{i,n}\|_{\mathcal{S}^p}^p \rightarrow 0$ , which also implies the uniform boundedness of the sequence  $(\|Y^{i,n}\|_{\mathcal{S}^p}^p)_{n \geq 0}$ , and in particular the one of  $\Psi_p^{i,n}$ . This result together with Assumption 4.1 and inequality (4.31) gives the desired convergence. In order to show that  $\lim_{n \rightarrow \infty} \|K^{i,n} - K^i\|_{\mathcal{S}^p} = 0$ , we use the equalities (4.29) and (4.30) and the convergence of  $(Y^{i,n}, Z^{i,n}, U^{i,n})$  shown above.  $\square$

**Theorem 4.3** (Propagation of chaos of the whole solution). *Under the assumptions of Proposition 4.3, the particle system (3.1) satisfies the propagation of chaos property, i.e. for any fixed positive integer  $k$ ,*

$$\lim_{n \rightarrow \infty} \text{Law}(\Theta^{1,n}, \Theta^{2,n}, \dots, \Theta^{k,n}) = \text{Law}(\Theta^1, \Theta^2, \dots, \Theta^k).$$

**Proof.** We obtain the propagation of chaos if we can show that  $\lim_{n \rightarrow \infty} D_G^p(\mathbb{P}^{k,n}, \mathbb{P}_\Theta^{\otimes k}) = 0$ . But, in view of the inequality (4.13), it suffices to show that for each  $i = 1, \dots, k$ ,

$$\lim_{n \rightarrow \infty} (\|Y^{i,n} - Y^i\|_{\mathcal{S}^p} + \|Z^{i,n} - Z^i \mathbf{e}_i\|_{\mathcal{H}^{p,n}} + \|U^{i,n} - U^i \mathbf{e}_i\|_{\mathcal{H}^{p,n}} + \|K^{i,n} - K^i\|_{\mathcal{S}^p}) = 0,$$

which is derived in Proposition 4.3.  $\square$

**Remark 4.2.** *All the results of Sections 3 and 4 extend to the case where*

- $f(t, \omega, y, z, \mu) := \hat{f}(t, (B_s(\omega), \tilde{N}_s(\omega))_{0 \leq s \leq t}, y, z, \mu)$ , with  $\hat{f}$  measurable and Lipschitz w.r.t.  $y, z, \mu$ .
- $h(t, \omega, y, \mu) := \hat{h}(t, (B_s(\omega), \tilde{N}_s(\omega))_{0 \leq s \leq t}, y, \mu)$ , with  $\hat{h}$  measurable and Lipschitz w.r.t.  $y, \mu$ .
- $(f^i \circ \mathbf{Y}^n)(t, \omega, y, z, u) = \hat{f}(t, (B_s^i(\omega), \tilde{N}_s^i(\omega))_{0 \leq s \leq t}, y, z^i, u^i, \mathbf{L}_n[\mathbf{Y}_t^n](\omega))$ .
- $(f^i \circ Y)(t, \omega, y, z, u) := \hat{f}(t, (B_s^i(\omega), \tilde{N}_s^i(\omega))_{0 \leq s \leq t}, y, z^i, u^i, \mathbb{P}_{Y_t^i})$ .
- $h(t, \omega, Y_t^{i,n}(\omega), \mathbf{L}_n[\mathbf{Y}_t^n](\omega)) := \hat{h}(t, (B_s^i(\omega), \tilde{N}_s^i(\omega))_{0 \leq s \leq t}, Y_t^{i,n}(\omega), \mathbf{L}_n[\mathbf{Y}_t^n](\omega))$ .
- $h(t, \omega, Y_t^i(\omega), \mathbb{P}_{Y_t^i}) := \hat{h}(t, (B_s^i(\omega), \tilde{N}_s^i(\omega))_{0 \leq s \leq t}, Y_t^i(\omega), \mathbb{P}_{Y_t^i})$ .

**Remark 4.3.** All the results of the paper extend to the case  $1 < p < 2$  following the same steps provided we consider

- either a driver  $f$  which does not depend on the jump size component  $U$ ;
- or we should use for the values of the jump size component the Banach space  $L^p_\nu + L^p_\nu$  endowed with the norm

$$|l|_{\nu,p,2} := \inf_{l^1+l^2=l} (|l^1|_{\nu,p} + |l^2|_{\nu,2}),$$

in which case we use the following Banach space  $\widehat{\mathcal{H}}_{\nu,p,2}^{p,1}$  of predictable processes  $U$  endowed with the norm

$$\|U\|_{\widehat{\mathcal{H}}_{\nu,p,2}^{p,1}} := \inf_{l^1+l^2=l} \left\{ \left( \mathbb{E} \left[ \int_0^T \int_{\mathbb{R}^*} |U_s^1(e)|^p \nu(de) ds \right] \right)^{1/p} + \left( \mathbb{E} \left[ \left( \int_0^T \int_{\mathbb{R}^*} |U_s^2(e)|^2 \nu(de) ds \right)^{p/2} \right] \right)^{1/p} \right\}.$$

These spaces are needed because in the proof of Proposition 4.3 we have to use the following lower bound (see [32]): there is a constant  $c_p$  such that

$$\left( \mathbb{E} \left[ \sup_{0 \leq t \leq T} \left| \int_0^T \int_{\mathbb{R}^*} U_s(e) \tilde{N}(ds, de) \right|^p \right] \right)^{1/p} \geq c_p \|U\|_{\widehat{\mathcal{H}}_{\nu,p,2}^{p,1}}$$

along with the Burkholder-Davis-Gundy inequality.

## A Some technical results

We establish here the following  $L^p$  estimates with *universal constants* for the difference of the solutions  $(Y^i, Z^i, U^i)$ ,  $i = 1, 2$  of BSDEs.

**Proposition A.1** ( *$L^p$  a priori estimates with universal constants*). Let  $T > 0$ . Let  $\tau$  be a  $\mathbb{F}$ -stopping time with values in  $[0, T]$ . Let  $p \geq 2$  and let  $\xi_1$  and  $\xi_2 \in L^p(\mathcal{F}_\tau)$ . Let  $f_1$  be a Lipschitz driver with constant  $C$  and let  $f_2$  be a driver. For  $i = 1, 2$ , let  $(Y^i, Z^i, U^i)$  be a solution of the BSDE associated to terminal time  $\tau$ , driver  $f^i$ , and terminal condition  $\xi^i$ . For  $s \in [0, \tau]$  denote  $\delta Y_s := Y_s^1 - Y_s^2$ ,  $\delta Z_s := Z_s^1 - Z_s^2$ ,  $\delta U_s := U_s^1 - U_s^2$ ,  $\delta f_s := f^1(s, Y_s^2, Z_s^2, U_s^2) - f^2(s, Y_s^2, Z_s^2, U_s^2)$  and  $\delta \xi := \xi^1 - \xi^2$ . Define  $c(p) := \frac{p[(p-1) \wedge 1]}{2}$  and let  $\eta, \beta > 0$  be such that  $\beta \geq C + \frac{1}{\eta} + \frac{p-1}{p} \left(\frac{1}{\eta}\right)^{\frac{p}{p-1}}$ . If  $\eta \leq \frac{2c(p)}{pC^2}$ , then for each  $t \in [0, \tau]$  we have

$$|e^{\beta t} \delta Y_t|^p \leq \mathbb{E} [ |e^{\beta \tau} \delta \xi|^p | \mathcal{F}_t ] + \eta^p \mathbb{E} \left[ \int_t^\tau |e^{\beta s} \delta f_s|^p ds | \mathcal{F}_t \right] \quad a.s. \quad (\text{A.1})$$

Furthermore, when  $p \in (1, 2)$  and the drivers  $f^1$  and  $f^2$  are independent on the jump size component  $U$ , the same estimate holds.

**Proof.** By Corollary (1) in Kruse and Popier [26], it follows that

$$\begin{aligned}
|e^{\beta t} \delta Y_t|^p &\leq |e^{\beta \tau} \delta \xi|^p + p \int_t^\tau |e^{\beta s} \delta Y_s|^{p-1} \phi(e^{\beta s} \delta Y_s) (e^{\beta s} \Delta f_s) ds - p\beta \int_t^\tau |e^{\beta s} \delta Y_s|^p ds \\
&\quad - p \int_t^\tau |e^{\beta s} \delta Y_s|^{p-1} \phi(e^{\beta s} \delta Y_s) (e^{\beta s} \delta Z_s) dW_s - p \int_t^\tau |e^{\beta s} \delta Y_s|^{p-1} \phi(e^{\beta s} \delta Y_s) (e^{\beta s} \int_{\mathbb{R}^*} \delta U_s(e)) \tilde{N}(ds, de) \\
&\quad - c(p) \int_t^\tau |e^{\beta s} \delta Y_s|^{p-2} |e^{\beta s} \delta Z_s|^2 \mathbf{1}_{\delta Y_s \neq 0} ds \\
&\quad - c(p) \int_t^\tau \int_{\mathbb{R}^*} |e^{\beta s} \delta U_s(e)|^2 (|e^{\beta s} \delta Y_{s-}|^2 \vee |e^{\beta s} (\delta Y_{s-} + \delta U_s(e))|^2)^{p/2-1} \mathbf{1}_{|\delta Y_{s-}| \vee |\delta Y_{s-} + \delta U_s(e)| \neq 0} N(ds, de),
\end{aligned} \tag{A.2}$$

where  $\phi(y) := |y|^{-1} y \mathbf{1}_{y \neq 0}$  and  $\Delta f_s := f^1(s, Y_s^1, Z_s^1, U_s^1) - f^2(s, Y_s^2, Z_s^2, U_s^2)$ .

Now, using the Lipschitz property of the driver  $f^1$ , we get

$$\int_t^\tau |e^{\beta s} \delta Y_s|^{p-1} \phi(e^{\beta s} \delta Y_s) |e^{\beta s} \Delta f_s| ds \leq \int_t^\tau |e^{\beta s} \delta Y_s|^{p-1} e^{\beta s} (C|\delta Y_s| + C|\delta Z_s| + C|\delta U_s|_\nu + |\delta f_s|) ds \tag{A.3}$$

By applying Young's inequality, for  $\varepsilon > 0$ , we derive that

$$\int_t^\tau |e^{\beta s} \delta Y_s|^{p-1} e^{\beta s} |\delta f_s| ds \leq \frac{p-1}{p} \left( \frac{1}{\varepsilon^2} \right)^{\frac{p}{p-1}} \int_t^\tau |e^{\beta s} \delta Y_s|^p ds + \frac{\varepsilon^{2p}}{p} \int_t^\tau |e^{\beta s} \delta f_s|^p ds, \tag{A.4}$$

$$C \int_t^\tau |e^{\beta s} \delta Y_s|^{p-1} e^{\beta s} |\delta Z_s| ds \leq \frac{1}{2\varepsilon^2} \int_t^\tau |e^{\beta s} \delta Y_s|^p ds + \frac{C^2 \varepsilon^2}{2} \int_t^\tau |e^{\beta s} \delta Y_s|^{p-2} \mathbf{1}_{\delta Y_s \neq 0} |e^{\beta s} \delta Z_s|^2 ds \tag{A.5}$$

and

$$C \int_t^\tau |e^{\beta s} \delta Y_s|^{p-1} e^{\beta s} |\delta U_s|_\nu ds \leq \frac{1}{2\varepsilon^2} \int_t^\tau |e^{\beta s} \delta Y_s|^p ds + \frac{C^2 \varepsilon^2}{2} \int_t^\tau |e^{\beta s} \delta Y_s|^{p-2} \mathbf{1}_{\delta Y_s \neq 0} |e^{\beta s} \delta U_s|_\nu^2 ds. \tag{A.6}$$

We set  $\eta = \varepsilon^2$  and choose  $\eta \leq \frac{2c(p)}{pC^2}$  and  $\beta \geq C + \frac{1}{\eta} + \frac{p-1}{p} \left( \frac{1}{\eta} \right)^{\frac{p}{p-1}}$ . We take the conditional expectation in (A.2) and since the martingales are true martingales by martingale inequalities, by using the relations (A.3), (A.4), (A.5) and (A.6), we obtain the result.  $\square$

## References

- [1] Andersson, D. and Djehiche, B. (2011). A maximum principle for SDEs of mean-field type. *Applied Mathematics & Optimization*, 63(3), 341–356.
- [2] Acciaio, B., Backhoff-Veraguas, J., and Carmona, R. (2019). Extended mean field control problems: stochastic maximum principle and transport perspective. *SIAM journal on Control and Optimization*, 57(6), 3666-3693.

- [3] Barles, G., Buckdahn, R., and Pardoux, E. (1997). Backward stochastic differential equations and integral-partial differential equations. *Stochastics: An International Journal of Probability and Stochastic Processes*, 60(1-2), 57-83.
- [4] Billingsley, P. (1968). *Convergence of probability measures*. John Wiley & Sons.
- [5] Briand, P., Cardaliaguet, P., de Raynal, P-É C and Hu, Y. (2020). Forward and backward stochastic differential equations with normal constraints in law. *Stochastic Processes and their Applications*, 130, 7021-7097.
- [6] Briand, P., and Hibon, H. (2021). Particles Systems for mean reflected BSDEs. *Stochastic Processes and their Applications*, 131, 253-275.
- [7] Buckdahn, R., Li, J., and Peng, S. (2009). Mean-field backward stochastic differential equations and related partial differential equations. *Stochastic processes and their Applications*, 119(10), 3133-3154.
- [8] Buckdahn, R., Djehiche, B., Li, J., and Peng, S. (2009). Mean-field backward stochastic differential equations: a limit approach. *The Annals of Probability*, 37(4), 1524-1565.
- [9] Carmona, R., Delarue, F., and Lachapelle, A. (2013). Control of McKean–Vlasov dynamics versus mean field games. *Mathematics and Financial Economics*, 7(2), 131-166.
- [10] Delarue, F. (2002). On the existence and uniqueness of solutions to FBSDEs in a non-degenerate case. *Stochastic Processes and their Applications*, 99, 209-286.
- [11] Dellacherie, C., and Meyer, P. A. (1982). *Probabilities and potential B*, Chapter V to VIII. Hermann.
- [12] Djehiche, B., Elie, R., and Hamadène, S. (2019). Mean-field reflected backward stochastic differential equations. *arXiv preprint arXiv:1911.06079*.
- [13] Dumitrescu, R., Labart C. (2016). Reflected scheme for doubly reflected BSDEs with jumps and RCLL obstacles. *Journal of Computational and Applied Mathematics*, 296, 827-839.
- [14] Dumitrescu, R., Labart C. (2016). Numerical approximation of doubly reflected BSDEs with jumps and RCLL obstacles. *Journal of Mathematical Analysis and Applications* 442 (1), 206-243.
- [15] Dumitrescu, R., Quenez, M. C., and Sulem, A. (2015). Optimal stopping for dynamic risk measures with jumps and obstacle problems. *Journal of Optimization Theory and Applications* 167 (1), 219-242.
- [16] Dumitrescu, R., Quenez, M. C., and Sulem, A. (2016). Generalized Dynkin games and doubly reflected BSDEs with jumps. *Electronic Journal of Probability*, 21, 1-32.
- [17] El Karoui, N., Kapoudjian, C., Pardoux, E., Peng, S., and Quenez, M. C. (1997). Reflected solutions of backward SDE's, and related obstacle problems for PDE's. *The Annals of Probability*, 25(2), 702-737.

- [18] Essaky, E. H. (2008). Reflected backward stochastic differential equation with jumps and RCLL obstacle. *Bulletin des sciences mathematiques*, 132(8), 690-710.
- [19] Gärtner, J. (1988). On the McKean-Vlasov limit for interacting diffusions. *Mathematische Nachrichten*, 137(1), 197-248.
- [20] Hamadène, S., and Ouknine, Y. (2003). Reflected backward stochastic differential equation with jumps and random obstacle. *Electronic Journal of Probability*, 8.
- [21] Hamadène, S., and Ouknine, Y. (2016). Reflected backward SDEs with general jumps. *Theory of Probability & Its Applications*, 60(2), 263-280.
- [22] Hu, K., Ren, Z., and Yang, J. (2019). Principal-agent problem with multiple principals. arXiv preprint arXiv:1904.01413.
- [23] Jabin, P. E., and Wang, Z. (2016). Mean field limit and propagation of chaos for Vlasov systems with bounded forces. *Journal of Functional Analysis*, 271(12), 3588-3627.
- [24] Jourdain, B., Méléard, S and Woyczynski, W A. (2008). Nonlinear SDEs driven by Lévy processes and related PDEs, *Alea* 4, 1–29.
- [25] Karatzas, I. and Shreve, S. E. (1998). *Methods of mathematical finance* (Vol. 39, pp. xvi+-407). New York: Springer.
- [26] Kruse, T., and Popier, A. (2016). BSDEs with monotone generator driven by Brownian and Poisson noises in a general filtration. *Stochastics*, 88(4), 491-539.
- [27] Lacker, D. (2018). On a strong form of propagation of chaos for McKean-Vlasov equations. *Electronic Communications in Probability*, 23, 1-11.
- [28] Laurière, M., and Tangpi, L. (2019). Backward propagation of chaos. arXiv preprint arXiv:1911.06835.
- [29] Lasry, J. M., and Lions, P. L. (2007). Mean field games. *Japanese journal of mathematics*, 2(1), 229-260.
- [30] Li, J. (2014). Reflected mean-field backward stochastic differential equations. Approximation and associated nonlinear PDEs. *Journal of Mathematical Analysis and Applications*, 413(1), 47-68.
- [31] Kac, M. (1956). Foundations of Kinetic Theory. In *Third Berkeley Symposium on Mathematical Statistics and Probability* (pp. 171-197).
- [32] Marinelli, C., and Röckner, M. (2014). On maximal inequalities for purely discontinuous martingales in infinite dimensions. *Séminaire de Probabilités XLVI*. Springer, Cham, 2014. 293-315.



- [33] McKean, H. P. (1967). Propagation of chaos for a class of non-linear parabolic equations. *Stochastic Differential Equations (Lecture Series in Differential Equations, Session 7, Catholic Univ., 1967)*, 41-57.
- [34] Pardoux, E., and Peng, S. (1990). Adapted solution of a backward stochastic differential equation. *Systems & Control Letters*, 14(1), 55-61.
- [35] Peng, E. (1999). Monotonic limit theorem of BSDE and nonlinear decomposition theorem of Doob–Meyer’s type. *Probab. Theory Relat. Fields*. 113, 473-499
- [36] Peng, S. (2004). Nonlinear expectations, nonlinear evaluations and risk measures. In *Stochastic methods in finance* (pp. 165-253). Springer, Berlin, Heidelberg.
- [37] Protter, P. E. (2005). *Stochastic integration and differential equations*. Springer, Berlin, Heidelberg.
- [38] Rachev, S. T., Klebanov, L., Stoyanov, S. V., and Fabozzi, F. (2013). *The methods of distances in the theory of probability and statistics*. Springer Science & Business Media.
- [39] Shkolnikov, M. (2012). Large systems of diffusions interacting through their ranks. *Stochastic Processes and their Applications*, 122(4), 1730-1747.
- [40] Sznitman, A. S. (1991). Topics in propagation of chaos. In *Ecole d’été de probabilités de Saint-Flour XIX—1989* (pp. 165-251). Springer, Berlin, Heidelberg.
- [41] Tang, S., and Li, X. (1994). Necessary conditions for optimal control of stochastic systems with random jumps. *SIAM Journal on Control and Optimization*, 32(5), 1447-1475.
- [42] Yamada, T., and Watanabe, S. (1971). On the uniqueness of solutions of stochastic differential equations. *Journal of Mathematics of Kyoto University*, 11(1), 155-167.
- [43] Villani, C (2009). *Optimal transport: old and new*. Vol. 338. Berlin: Springer.