

SUCCESSIVELY ALMOST POSITIVE LINKS

TETSUYA ITO

ABSTRACT. As an extension of positive or almost positive diagrams and links, we introduce a notion of successively almost positive diagrams and links, and good successively almost positive diagrams and links. We review various properties of positive links or almost positive links, and explain how they can be extended to (good) successively almost positive links. Our investigation also leads to an improvement of known results of positive or almost positive links.

1. INTRODUCTION

A link diagram is *positive* if all the crossings are positive, and a *positive link* is a link that can be represented by a positive diagram. Positive links have various nice properties and form an important class of links.

An innocent generalization of a positive diagram is a *k-almost positive diagram*, a diagram such that all but k crossings are positive. A 1-almost positive diagram is usually called an *almost positive diagram* and has been studied in various places.

It is known that almost positive links share various properties with positive links. As is discussed in [PT] there are special properties of 2 or 3-almost positive links. However when k is large, k -almost positive links fail to have nice properties similar to positive links because every knot K is k -almost positive for sufficiently large k .

The aim of this paper is to propose a better and more natural generalization of a (almost) positive diagram which we call a *successively (k-)almost positive diagram*. We also introduce a *good successively (k-)almost positive diagram* which is a successively positive diagram having an additional condition. We show (good) successively almost positive links share various properties with (almost) positive links, for all $k \geq 0$.

The paper has an aspect of survey of (almost) positive links; we review various properties of (almost) positive links which can be found in various places, sometimes in a different context or prospect¹. We try to make it clear how the proof of the properties of (almost) positive links utilizes or reflects the positivity of diagrams, and explain how to extend the proof to a successively almost positive case.

Besides the extension of known properties, our investigation sometimes leads to an improvement of known results or new results of (almost) positive links themselves. See Corollary 1.9, Corollary 1.13, and Corollary 1.15 and Theorem 3.1.

In the following, for a link diagram D we use the following notations.

- $s(D)$ is the number of Seifert circles.
- $c(D)$ is the number of crossings, and $c_{\pm}(D)$ is the number of positive and negative crossings.

2020 *Mathematics Subject Classification*. Primary 57K10.

¹For example, properties of positive links discussed in [Cr] was obtained as a corollary of properties of more general class of links called *homogeneous links*.

- $w(D)(= c_+(D) - c_-(D))$ is the writhe.
- S_D is the *canonical Seifert surface* of D , the Seifert surface obtained from D by applying Seifert's algorithm.
- $\chi(D) = s(D) - c(D) = \chi(S_D)$ is the *euler characteristic* of D . When K is a knot, we often use the *genus* $g(D) = \frac{1-\chi(D)}{2}$ instead of $\chi(D)$.

1.1. Successively k -almost positive diagram.

Definition 1.1 (Successively k -almost positive diagram/link). A knot diagram D is *successively k -almost positive* if all but k crossings of D are positive, and the k negative crossings appear successively along a single overarc (see Figure 1).




FIGURE 1. Successively k -almost positive diagram. * represents the standard base point, which will be used in Section 2.1.

This type of diagram appeared in [IMT, Theorem 5.3] as an extension of positive diagram, designed so that the technique of constructing generalized torsion elements can be applied.

The definition says that a successively 0-almost positive (resp. successively 1-almost positive) diagram is nothing but a positive (resp. almost positive) diagram. On the other hand, a 2-almost positive diagram is not necessarily a successively 2-almost positive diagram; the two-crossing diagram of the negative Hopf link is 2-almost positive but not successively 2-almost positive.

As we will see and discuss later, an investigation of good properties of diagrams themselves leads to the following restricted class of successively k -almost positive diagrams.

Definition 1.2 (Good successively k -almost positive diagram/link). A k -successively almost positive diagram D is *good* if, when two distinct Seifert circles s, s' of D are connected by a negative crossing then there are no other crossings connecting s and s' (see the left figure of Figure 5 for a schematic illustration).

A *successively almost positive link* is a link represented by a successively k -almost positive diagram for some k . Similarly, a *good successively almost positive link* is a link represented by a good successively k -almost positive diagram for some k .

1.2. Properties. We show the following properties, generalizing (almost) positive link cases ([Cr],[PT]).

Theorem 1.3. *If K is successively k -almost positive, then its Levine-Tristram signature² σ_ω satisfies $\sigma_\omega(K) \leq 0$ for all $\omega \in S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$.*

Theorem 1.4. *If K is successively almost positive, the Conway polynomial $\nabla_K(z)$ is non-negative.*

Here we say that a polynomial $f(z) = a_0 + a_1z + \dots$ is *non-negative* if $a_i \geq 0$ for all i .

A positive diagram D has the following nice properties³;

- D represents a split link if and only if the diagram D is split.
- The canonical Seifert surface S_D attains the maximum euler characteristic among its Seifert surfaces; $\chi(D) = \chi(K)$.

It is easy to see these properties fail, even for almost positive diagrams. However, a good successively almost positive diagram has the same properties.

Theorem 1.5. *A good successively k -almost positive diagram D represents a split link if and only if D is non-split.*

Theorem 1.6. *Assume that D is a good successively k -almost positive diagram. Then its canonical Seifert surface S_D attains the maximum euler characteristic.*

Besides these properties, it turns out that a good successively almost positive link have more nice properties.

Let $\chi(K)$ be the maximal euler characteristic of Seifert surfaces of K . For the Conway polynomial $\nabla_K(z)$ of K , $\max \deg_z \nabla_K(z) = 1 - \chi(K)$ holds if K is non-split and positive. Similarly, for the HOMFLY polynomial $P_K(v, z)$ of K ⁴ $\max \deg_z P_K(v, z) = \min \deg_v P_K(v, z) = 1 - \chi(K)$ holds if K is positive [Cr].

We extend these properties for good successively almost positive links.

Theorem 1.7. *$\max \deg_z \nabla_K(z) = 1 - \chi(K)$ if K is a non-split good successively almost positive link K .*

Theorem 1.8. *For a good successively almost positive link K , $\max \deg_z P_K(v, z) = \min \deg_v P_K(v, z) = 1 - \chi(K)$.*

When K is an almost positive link which admits a non-good (successively) 1-almost positive diagram, the equality $\min \deg_v P_K(v, z) = 1 - \chi(K)$ was proven in [St2] (see Corollary 3.2 for a different proof). Therefore Theorem 1.8 positively answers [St2, Question 2];

Corollary 1.9. *$\min \deg_v P_K(v, z) = 1 - \chi(K)$ if K is almost positive.*

A *strongly quasipositive link* is another generalization of positive links. An n -braid is *strongly quasipositive* if it is a product of positive band generators $a_{i,j} = (\sigma_i \sigma_{i+1} \cdots \sigma_{j-2}) \sigma_{j-1} (\sigma_i \sigma_{i+1} \cdots \sigma_{j-2})^{-1}$ ($1 \leq i < j \leq n$), where σ_i is the standard generator of the braid group B_n . We say that a link is *strongly quasipositive* if it is represented as the closure of a strongly quasipositive braid.

²We remark that we adopt the convention that the positive trefoil has negative signature, as opposed to the one adopted in [BDL].

³The first statement can be confirmed by looking at the linking number, and the second statement follows from the Bennequin inequality or [Cr, Corollary 4.1]

⁴Here we use the convention that the skein relation of the HOMFLY polynomial is $v^{-1}P_{K_+}(v, z) - vP_{K_-}(v, z) = zP_{K_0}(v, z)$.

There is an equivalent definition using Seifert surfaces; A Seifert surface is *quasi-positive* if S is realized as an incompressible subsurface of the fiber surface of a positive torus link. A link is strongly quasipositive if and only if it bounds a quasi-positive Seifert surface [Rud1].

In [FLL], Feller-Lewark-Lobb proved that almost positive links are strongly quasipositive. At this moment we do not know whether a k -almost positive link is strongly quasipositive or not. However, for a good successively k -almost positive link, we see that they are strongly quasipositive.

Theorem 1.10. *A good successively almost positive link is strongly quasipositive. Indeed, a canonical Seifert surface S_D of a good successively almost positive diagram D is quasipositive.*

Let $s(K), \tau(K)$ be the Rasmussen invariant [Ra] and Heegard Floer tau-invariant [OS] of a knot⁵ K , and let $g_4(K)$ be the slice genus of K . Since strongly quasipositive knot K satisfies $s(K) = 2\tau(K) = 2g_4(K) = 2g_3(K)$ [Li, Theorem 4], [Sh, Proposition 1.7], we conclude the following.

Corollary 1.11. *If K is a good successively almost positive knot, then $s(K) = 2\tau(K) = 2g_4(K) = 2g(K)$.*

1.3. Signature and concordance finiteness. It is known that the signature of a non-trivial (almost) positive link is always strictly negative; $\sigma(K) < 0$ [PT]. Indeed, for a positive diagram D , a stronger inequality

$$(1.1) \quad \sigma(K) \leq \frac{1}{24}(\chi(D) - 1) \quad \left(= \frac{1}{24}(\chi(K) - 1) \right)$$

was proven in [BDL, Theorem 1.2]. Since positive-to-negative crossing change increases the signature by at most two, (1.1) implies

$$(1.2) \quad \sigma(K) \leq \frac{1}{24}(\chi(D) - 1) + 2c_-(D)$$

for general link diagram D .

In this opportunity, we present an improvement of (1.2).

Theorem 1.12. *Let D be a reduced diagram of a non-trivial link K . Then*

$$\sigma(K) \leq \frac{1}{12}(\chi(D) - 1) + \frac{4}{3}c_-(D) - \frac{1}{2} \leq \frac{1}{12}(\chi(K) - 1) + \frac{4}{3}c_-(D) - \frac{1}{2}$$

This inequality is interesting and useful in its own right. For positive link case, this gives the following improvement of (1.1).

Corollary 1.13. *If K is a non-trivial positive link,*

$$\sigma(K) \leq \frac{1}{12}(\chi(K) - 1) - \frac{1}{2}.$$

In [BDL, Theorem 1.1], as an application of signature estimate they showed that every topological knot concordance class contains finitely many positive knots. Theorem 1.12 leads to a similar finiteness result for successively k -almost positive knots under an additional restriction. Let $g_c^{top}(\mathcal{K}) = \min\{g(K) \mid K \in \mathcal{K}\}$ be the *topological concordance genus* of a topological concordance class \mathcal{K} .

⁵A similar result holds for the link case by using appropriate generalizations of the Rasmussen and the tau invariants for links.

Theorem 1.14. *For any $0 < d < \frac{1}{8}$, every topological knot concordance class \mathcal{K} contains only finitely many successively k -almost positive knots such that $k \leq dg_c^{top}(\mathcal{K})$.*

In particular, this extends the concordance finiteness result for almost positive knots, under the assumption that its topological concordance genus is sufficiently large.

Corollary 1.15. *A topological knot concordance class \mathcal{K} contains only finitely many almost positive knots if $g_c^{top}(\mathcal{K}) \geq 8$.*

ACKNOWLEDGEMENT

The author would like to thank Kimihiko Motegi and Masakazu Teragaito for stimulating conversations and discussions. A notion of successively positive diagram appeared during a joint work [IMT]. The author has been partially supported by JSPS KAKENHI Grant Number 19K03490, 21H04428.

2. PROOF OF PROPERTIES

In this section we present a proof of properties of (good) successively almost positive links.

2.1. Positive skein resolution tree for successively almost positive diagrams.

Definition 2.1 (Skein resolution tree). For a knot diagram D , a *skein resolution tree* is a rooted binary tree T such that

- (i) Each node c is labelled by a knot diagram D_c .
- (ii) The root is labelled by D .
- (iii) At each non-terminal node c with its children c' , c'' , their labellings $(D_c, D_{c'}, D_{c''})$ forms a skein triple (D_+, D_-, D_0) or (D_-, D_+, D_0)
- (iv) At each terminal node c , D_c represents the trivial link.

We say that a skein resolution tree T is *positive* if instead of (iii), the stronger property holds.

- (iii+) At each non-terminal node c with its children c' , c'' , their labellings $D_c, D_{c'}, D_{c''}$ forms a skein triple (D_+, D_-, D_0) .

By the skein relation, one can compute the Conway, the Jones and the HOMFLY polynomial by the skein resolution tree.

There are various ways to construct a skein resolution tree from a diagram. Here we use the one given in [Cr, Theorem 2].

A *based link diagram* D is a link diagram D such that

- The ordering (numbering) of component of D is given; $D = D_1 \cup D_2 \cup \dots \cup D_n$.
- For each component D_i , the base point $*_i \in D_i$ is given.

By suitably assigning the z -coordinates, we view a based link diagram D as a collection of curves $\gamma_i : [0, 1] \rightarrow \mathbb{R}^3$ such that $\gamma_i(0) = \gamma_i(1) = *_i$.

Definition 2.2. A based link diagram D is *descending* if

- (1) The z -coordinate of γ_i is monotone decreasing, except near $t = 1$.

- (2) When $j < i$, the j -th component of D lies entirely above of the i -th component of D .

Obviously, a descending diagram represent the unlink.

For a given based link diagram D , by trying to make D descending, one can construct skein resolution tree as follows (see [Cr, Theorem 2], for details).

We start at the base point $*_1$ of the 1st component D_1 . We walk along the diagram until we encounter the *first non-descending crossing* c , the crossing where the descending condition fails for the first time. Namely, we encounter the crossing c for the first time but we pass the crossing c as the underarc. When we do not encounter such a crossing, then we turn our attention to the 2nd component D_2 , and continue to do the same procedure.

We denote by $d(D)$ the number of crossings that appeared before arriving to the first non-descending crossing. If we cannot find the first non-descending crossing, then D is already descending, and in this case we define $d(D) = c(D)$.

Assume that the first non-descending crossing c is the crossing between the components D_i and D_j ($i \leq j$). We apply the skein resolution at the crossing c . Let D' be the diagram obtained by the crossing change at c , and let D'' be the diagram obtained by the smoothing at c .

The diagram D' is naturally regarded as a based link diagram. We view the diagram D'' as a based diagram as follows:

- When the smoothing merges two different components (i.e. $i \neq j$), we simply forget the base point and merged component D_j .
- When the smoothing splits the component D_i into two components $D_i^{(1)}$ and $D_i^{(2)}$ (i.e. $i = j$), we take the component that contains $*_i$ as the i th component. We take $*_i$ as its base point. We take the other component $D_i^{(2)}$ as $D_{i'}$ where i' is taken so that $i < i'$, and we take a base point near the crossing c .

By induction on $(c(D) - d(D), c(D))$, one can check that this procedure terminates in finitely many steps hence we get a skein resolution tree. We call this *the descending skein resolution tree* of the based diagram D (see Figure 2 for example).

For a successively k -almost positive diagram D , we take a base point $*$ on the overarc of the first negative crossing point. We call $*$ the *standard base point* of D . We view a successively k -almost positive diagram D as a based link diagram by taking the ordering of components so that the component having the standard base point $*$ as D_1 . We take the standard base point as the base point of D_1 , and the rest of the choices (orderings and the base points of other components) are arbitrary.

Proposition 2.3. *The descending skein resolution tree of successively k -almost positive diagram D is positive.*

Proof. By definition of successively k -almost positive diagram D , the diagram D is descending at the first k negative crossings. Thus in the construction of the descending skein resolution tree, all the skein resolutions happen at the positive crossings. \square

Theorem 1.3 and Theorem 1.4 follow from Proposition 2.3 and the following simple observations.

Lemma 2.4. *If a diagram D admits a positive skein resolution tree, then*




FIGURE 2. Descending skein resolution tree of the trefoil knot diagram

- (i) $\nabla_K(z)$ is non-negative.
- (ii) $\sigma_\omega(K) \leq 0$

Proof. (i) follows from the skein relation of the Conway polynomial

$$\nabla_{K_+}(z) = \nabla_{K_-}(z) + z\nabla_{K_0}(z)$$

and the observation that for an n -component unlink U_n , $\nabla_{U_n} = \begin{cases} 1 & (n = 1) \\ 0 & (\text{otherwise}) \end{cases}$

is a non-negative polynomial .

(ii) follows from the property of the Levine-Tristram signature

$$\sigma_\omega(K_-) \geq \sigma_\omega(K_+)$$

and the observation that the existence of positive skein resolution tree implies that one can make K unlink only by crossing changes at positive crossings. \square

2.2. The canonical Seifert surface of a good successively almost positive diagram. Theorem 1.10 and Theorem 1.6 follows from a characterization of quasi-positive canonical Seifert surface [FLL, Theorem B]. Here we give a direct proof based on Murasugi-Prytzcki’s operation reducing the number of Seifert circle [MP]. This tells us how to get a strongly quasipositive braid representative of K .

A crossing c of D is called *independent* if c connects two Seifert circles s, s' then there are no other crossing connecting s and s' .

For an independent crossing c connecting Seifert circles s, s' , let s_1, \dots, s_k be the Seifert circles of D other than s' connected to s by a crossing. Let c_1, \dots, c_n be the crossings that connects s and s_i ($i = 1, \dots, k$).

We move the underarc of the crossing c across one of the Seifert circles s , swallowing all the Seifert circles s_1, \dots, s_n and the crossings c_1, \dots, c_n to get a new diagram D' . We call this operation the *Murasugi-Prytzcki’s move (MP-move, in short)* at c (see Figure 3). The MP-move reduces the number of Seifert circles of D by one.




FIGURE 3. Murasugi-Prytzcki's move at the independent crossing c

We view the MP-move as an isotopy of canonical Seifert surface of D that flips the twisted bands corresponding to c and disk bounded by s to merge the disk bounded by s and s' into a single disk. After the MP-move, the Seifert surface still has a disk-and-twisted-band decomposition structure (see Figure 4).




FIGURE 4. Murasugi-Prytzcki's move viewed as an isotopy of canonical Seifert surface; the Seifert surface inherits a disk-and-twisted-band decomposition.

Proof of Theorem 1.6 and Theorem 1.10. When D is good successively k -almost positive diagram, we can apply the MP-move k times to get a diagram D' (see Figure 5).

Then $s(D') = s(D) - k$ and $w(D') = w(D) + k$. By the Bennequin's inequality $-s(D') + w(D') \leq -\chi(K)$ hence we get

$$\begin{aligned} -s(D) + w(D) + 2k &= -s(D') + w(D') \\ &\leq -\chi(K) \\ &\leq -\chi(S_D) = -s(D) + c(D) = -s(D) + w(D) + 2k. \end{aligned}$$

Therefore the canonical Seifert surface S_D attains the maximal euler characteristic.

Moreover, as in the single MP-move case, the sequence of MP-moves also can be seen as an isotopy of canonical Seifert surface of D ; we see that the canonical Seifert circle S_D can be understood as a surface made of $s(D') = s(D) - k$ disks and

$w(D') = w(D) + k$ positively twisted bands. This disk-and-twisted-band decomposition satisfies a certain nice condition, which we called *quasi-canonical Seifert surface* in [HIK]. As we have discussed in [HIK, Section 6], one can further deform the diagram into a closed braid diagram, preserving the disk and twisted band decomposition structure of S_D . Since every twisted band has twisted in a positive direction, the closed braid obtained from S_D is the closure strongly quasipositive braid (see [HIK, Theorem 6.4]). \square




FIGURE 5. Applying Murasgui-Prytzcki's move k times for good successively k -almost positive diagram ($k = 3$ case illustration).

2.3. Link polynomials and $\chi(K)$. Theorem 1.6 allows us to relate $\chi(K)$ and the knot polynomials.

Proof of Theorem 1.5 and Theorem 1.7. First we note that Theorem 1.5 follows from Theorem 1.7, since $\nabla_K(z) \neq 0$ implies that K is non-split.

We prove the assumption by induction of the number p of positive crossing of D . When $p = 0$, the assertion is clear. Let us consider the skein triple $(D = D_+, D_-, D_0)$ at the root of the descending skein tree (the resolution at the first non-descending crossing). Note that D_0 is a good successively k -almost positive diagram.

When D_0 is non-split, by induction and Theorem 1.6, $\deg_z \nabla_{K_0}(z) = 1 - \chi(K_0) = 1 - s(D) + (c(D) - 1) = -\chi(K)$. On the other hand,

$$\nabla_{K_-}(z) \leq 1 - \chi(K_-) \leq 1 - \chi(D_-) = 1 - \chi(D_+) = 1 - \chi(K).$$

Since $\nabla_K(z) = \nabla_{K_-}(z) + z\nabla_{K_0}(z)$ and both $\nabla_{K_-}(z)$ and $z\nabla_{K_0}(z)$ are non-negative, $\deg \nabla_K(z) = 1 - \chi(K)$ as desired.

When D_0 is split, the crossing c should be a nugatory crossing so K is represented by a non-split good successively k -almost positive diagram having $(p - 1)$ positive crossings. Therefore by induction $\nabla_K(z) = 1 - \chi(K)$. \square

Proof of Theorem 1.8. Assume that K admits a good successively k -almost positive diagram D . We prove the theorem by induction on k .

We consider the skein triple $(D_+, D = D_-, D_0)$ obtained by the first negative crossing D . Then D_+ and D_0 are good successively $(k-1)$ -almost positive diagram hence by Theorem 1.6

$$\chi(K_+) = \chi(K), \chi(K_0) = \chi(K) - 1$$

Therefore by induction

$$\max \deg_z P_{K_+}(v, z) = 1 - \chi(K), \max \deg_z P_{K_0}(v, z) = -\chi(K).$$

By the skein relation $P_K(v, z) = v^{-2}P_{K_+}(v, z) - v^{-1}zP_{K_0}(v, z)$ we get

$$\max \deg_z P_K(v, z) \leq 1 - \chi(K).$$

On the other hand, by Theorem 1.7

$$1 - \chi(K) = \deg_z \nabla_K(z) = \max \deg_z P_K(v, z)|_{v=1} \leq \max \deg_z P_K(v, z)$$

hence we conclude that $\max \deg_z P_K(v, z) = 1 - \chi(K)$.

Next we show $\min \deg_v P_K(v, z) = 1 - \chi(K)$. By the skein relation (characterization) of the HOMFLY polynomial $P_K(v, -v^{-1} + v) = 1$. This implies that $P_K(v, z)$ contains a monomial $v^a z^b$ whose coefficient is non-zero and $a - b \leq 0$. Thus

$$\min \deg_v P_K(v, z) \leq a \leq b \leq \max \deg_z P_K(v, z).$$

On the other hand, by the Morton-Franks-William inequality

$$\overline{sl}(K) + 1 \leq \min \deg_v P_K(v, z)$$

where $\overline{sl}(K)$ is the maximal self-linking number of K . Since we have seen that K is strongly quasipositive, $\overline{sl}(K) = -\chi(K)$. Combining the (in)equalities we get

$$-\chi(K) + 1 = \overline{sl}(K) + 1 \leq \min \deg_v P_K(v, z) \leq \max \deg_z P_K(v, z) = -\chi(K) + 1.$$

□

2.4. Signature estimate. We turn to our attention to Theorem 1.12. In large part, our argument goes the same line as Baader-Dehornoy-Liechti's argument [BDL] adapted so that it can be applied to general link diagrams with slight improvements.

To treat the signature, we use Gordon-Litherland's theorem [GL]. For a (possibly non-orientable) spanning F of a link K , let $\langle \cdot, \cdot \rangle_F : H_1(F) \times H_1(F) \rightarrow \mathbb{Z}$ be the *Gordon-Litherland pairing* of F ; for $a, b \in H_1(F)$, let α, β be curves on S that represent a, b , and let $p_F : \nu(F) \rightarrow F$ be the unit normal bundle of F . The Gordon-Litherland pairing of a and b is defined by $\langle a, b \rangle_F = lk(\alpha, p_S^{-1}(\beta))$, where lk denotes the linking number.

When K is an ℓ -component link $K_1 \cup \cdots \cup K_\ell$, the Gordon-Litherland theorem states

$$(2.1) \quad \sigma(K) = \sigma(F) + \frac{1}{2}e(F, K).$$

Here $\sigma(F)$ is the signature of the Gordon-Litherland pairing of F , and

$$e(F, K) = -\frac{1}{2}\langle [K], [K] \rangle_F - \sum_{1 \leq i < j \leq \ell} lk(K_i, K_j).$$

For a link diagram D , we fix one of its checkerboard coloring, and let B and W be the black and white checkerboard surface of D . We say that a crossing c of D is of *type a* (resp. *of type b*) if, when we put the overarc so that it is an horizontal line, the upper right-hand side and the lower left-handed side (resp. the lower

right-hand side and the upper left-handed side) are colored by black (see Figure 6). Similarly, we say that a crossing c is of *type I* (resp. *of type II*) if the black region is compatible (resp. incompatible) with the orientation of the diagram (see Figure 6). In the definition of type a/b, the orientation of D is irrelevant whereas the definition of type I/II, the over-under information is irrelevant.

We say that a crossing c is of *type Ia*, for example, if c is both of type I and of type a. We put $c_{Ia}, c_{Ib}, c_{IIa}, c_{IIb}$ the number of crossings of type *Ia, Ib, IIa, IIb*, respectively. Note that positive (resp. negative) crossing is either of type Ib or IIa (resp. Ia or IIb) so

$$(2.2) \quad c_+(D) = c_{Ib} + c_{IIa}, \quad c_-(D) = c_{Ia} + c_{IIb}.$$




FIGURE 6. Types of crossings with respect to the checkerboard coloring

By the definition of the Gordon-Litherland pairing,

$$\frac{1}{2}e(B, K) = c_{IIb} - c_{IIa}, \quad \frac{1}{2}e(W, K) = c_{Ia} - c_{Ib}.$$

Thus by (2.1) and (2.2)

$$(2.3) \quad 2\sigma(K) = \sigma(B) + \sigma(W) - c_+(D) + c_-(D)$$

We give an estimation of the signature of Gordon-Litherland pairings. Let $\mathcal{R}(W)$ and $\mathcal{R}(B)$ be the set of white and black regions, respectively. For a white region $R \in \mathcal{R}(W)$, we associate a simple closed curve γ_R on B which is a mild perturbation of the boundary of R (see Figure 7).

We say that the region R is of *type* (α, β) if R contains α type a crossings and β type b crossings as its corners. By definition,

$$\langle [\gamma_R], [\gamma_R] \rangle_B = \alpha - \beta$$

when R is of type (α, β) .




FIGURE 7. Curve γ_R for a white region R

For a black region $R \in \mathcal{R}(B)$, the curve γ_R on W and a notion of type (α, β) are defined similarly and the Gordon-Litherland pairing is given by

$$\langle [\gamma_R], [\gamma_R] \rangle_W = \beta - \alpha$$

when R is of type (α, β) .

Proof of Theorem 1.12. Since D is reduced, there are no regions of type $(1, 0)$ or $(0, 1)$. Moreover, since we assume that K is non-trivial, $\#\mathcal{R}(D), \#\mathcal{B}(D) \geq 2$.

Let Γ be the planar graph whose vertices $\mathcal{V}(\Gamma)$ are white regions of type (α, β) with $\alpha - \beta \leq 0$, and two vertices R, R' are connected by an edge if they share a corner. By Appel-Haken's four-color theorem⁶ [AH] there is a 4-coloring map $col : V(\Gamma) \rightarrow \{1, 2, 3, 4\}$; if two vertices v, v' are connected by an edge, then $col(v) \neq col(v')$. Let $\mathcal{V}' = col^{-1}(\{1, 2\})$. Let Γ' be the subgraph of Γ whose vertices are \mathcal{V}' , and let V_B be the subspace of $H_1(B)$ generated by $[\gamma_R]$ for $R \in \mathcal{V}'$. With no loss of generality, we assume that $\#\mathcal{V}' \geq \frac{1}{2}\#\mathcal{V}(\Gamma)$.

Since Γ' is bipartite, the restriction of Gordon-Litherland pairing $\langle \cdot, \cdot \rangle_F$ on V_B is of the form $\begin{pmatrix} D_1 & X^T \\ X & D_2 \end{pmatrix}$ where D_1, D_2 are diagonal matrices with non-positive diagonals, hence the Gordon-Litherland pairing is non-positive on V_B .

First we assume that \mathcal{V}' is not equal to the whole $\mathcal{R}(W)$. In this case $\{[\gamma_R] \mid R \in \mathcal{V}'\}$ is a basis of V_B so $\dim V_B = \#\mathcal{V}'$. Let $\gamma_{>0}^W$ be the number of white region R such that $\langle [\gamma_R], [\gamma_R] \rangle_B > 0$. Then

$$\dim V_B = \#\mathcal{V}' \geq \frac{1}{2}\#\mathcal{V}(\Gamma) = \frac{1}{2}(\#\mathcal{R}(W) - \gamma_{>0}^W)$$

On the other hand, when \mathcal{V}' coincides with $\mathcal{R}(W)$, then $\dim V_B = \dim H_1(B) = \#\mathcal{R}(W) - 1$. Since $\mathcal{R}(W) \geq 2$, we have the same lower bound of $\dim V_B$;

$$\dim V_B = \#\mathcal{R}(W) - 1 \geq \frac{1}{2}\#\mathcal{R}(W) \geq \frac{1}{2}(\#\mathcal{R}(W) - \gamma_{>0}^W)$$

Therefore we get an upper bound of the signature of the Gordon-Litherland pairing of B^7 .

$$\begin{aligned} \sigma(B) &\leq \dim H_1(B) - \dim V_B = (\#\mathcal{R}(W) - 1) - \dim V_B \\ &\leq \frac{1}{2}\#\mathcal{R}(W) - 1 + \frac{1}{2}\gamma_{>0}^W \end{aligned}$$

By a parallel argument for the white surface W , we get a similar estimate

$$\sigma(W) \leq \frac{1}{2}\#\mathcal{R}(B) - 1 + \frac{1}{2}\gamma_{>0}^B$$

where $\gamma_{>0}^B$ is the number of a black region R such that $\langle \gamma_R, \gamma_R \rangle_W > 0$.

Since $\#\mathcal{R}(W) + \#\mathcal{R}(B) - 2 = c(D)$, by (2.3) we get

$$(2.4) \quad 2\sigma(K) \leq -\frac{1}{2}c(D) + 2c_-(D) + \frac{1}{2}\gamma_{>0}^B + \frac{1}{2}\gamma_{>0}^W - 1$$

It remains to estimate $\gamma_{>0}^B$ and $\gamma_{>0}^W$. Let $\gamma^W(\alpha, \beta)$ be the number of white regions of type (α, β) . By definition of $\gamma_{>0}^W$,

$$\gamma_{>0}^W = \sum_{\substack{\alpha > \beta \geq 0 \\ \alpha \geq 2}} \gamma^W(\alpha, \beta).$$

⁶It is interesting to find an argument that avoids to use the four-color theorem.

⁷This is a point where a minor improvement (constant $-\frac{1}{2}$ in the conclusion) appears.

By counting the number of the crossings of type a that appear as a corner of white regions, we get

$$\begin{aligned}
 2(c_{Ia} + c_{IIa}) &= \sum_{\alpha, \beta \geq 0} \alpha \gamma^W(\alpha, \beta) \\
 &\geq 2\gamma^W(2, 0) + 2\gamma^W(2, 1) + 3 \sum_{\substack{\alpha > \beta \geq 0 \\ \alpha \geq 3}} \gamma^W(\alpha, \beta) \\
 &= -\gamma^W(2, 0) - \gamma^W(2, 1) + 3 \sum_{\substack{\alpha > \beta \geq 0 \\ \alpha \geq 2}} \gamma^W(\alpha, \beta) \\
 &= -\gamma^W(2, 0) - \gamma^W(2, 1) + 3\gamma_{>0}^W.
 \end{aligned}$$

Thus we conclude

$$(2.5) \quad \gamma_{>0}^W \leq \frac{2}{3}(c_{Ia} + c_{IIa}) + \frac{1}{3}\gamma^W(2, 0) + \frac{1}{3}\gamma^W(2, 1)$$

Recall that a positive crossing appears as either of type Ib or of type IIa. If the corner of a white region R is a crossing of type IIa, then the orientation of link and the orientation of the white region switch. Thus if all the corners of R are positive crossings, then the region R must be of type $R(\alpha, 2\beta)$. In particular, if a white region R is of type $(2, 1)$, at least one of its corner is a negative crossing. Similarly, if a white region R is of type $(2, 0)$, its corners are either both positive, or, both negative.

Let $\gamma_{\pm}^W(2, 0)$ be the number of white regions of type $(2, 0)$ whose corners are positive and negative crossings, respectively. By counting the number of negative crossings that appear as a corner of type $(2, 1)$ regions or type $(2, 0)$ regions we get

$$\gamma^W(2, 1) + 2\gamma_{-}^W(2, 0) \leq 2c_{-}.$$

Thus

$$\gamma^W(2, 1) + \gamma_{-}^W(2, 0) \leq 2c_{-}.$$

Let $s_W(D)$ be the number of Seifert circles of D which are the boundary of white bigon. When both corners of a white region R of type $(2, 0)$ are positive, then the boundary of R forms a Seifert circle of D so⁸

$$\gamma_{+}^W(2, 0) \leq s_W(D).$$

By (2.5) we conclude

$$(2.6) \quad \gamma_{>0}^W \leq \frac{2}{3}(c_{Ia} + c_{IIa}) + \frac{2}{3}c_{-} + s_W(D).$$

By the same argument for the white surface, we get a similar inequality

$$(2.7) \quad \gamma_{>0}^B \leq \frac{2}{3}(c_{IIa} + c_{Ib}) + \frac{2}{3}c_{-} + s_B(D).$$

where $s_B(D)$ is the number of Seifert circles which are the boundary of black bigon.

Since a Seifert circle cannot be the boundary of white bigon and black bigon at the same time,

$$s_W(D) + s_B(D) \leq s(D).$$

⁸A substantial improvement appears at this point; In [BDL] they used an upper bound of $\gamma_{+}^W(2, 0)$ in terms of the crossing numbers, as we do for $\gamma_{-}^W(2, 0)$, instead of the number of Seifert circles.

The equality happens only if all the Seifert circles are bigon. Thus the equality occurs only if D is the standard torus $(2, 2n)$ link diagram (with opposite orientation, so that they bound an annulus). In this case, the asserted inequality of signature is obvious, so in the following we can assume that a bit stronger inequality

$$(2.8) \quad s_W(D) + s_B(D) \leq s(D) - 1.$$

By (2.6), (2.7), and (2.8)

$$\begin{aligned} \gamma_{>0}^B + \gamma_{>0}^W &\leq \frac{2}{3}(c_{Ia} + c_{IIa} + c_{Ib} + c_{IIb}) + \frac{1}{3}(s(D) - 1) + \frac{4}{3}c_-(D) \\ &= \frac{2}{3}c(D) + \frac{1}{3}(s(D) - 1) + \frac{4}{3}c_-(D). \end{aligned}$$

Therefore by (2.4) we conclude

$$\begin{aligned} 2\sigma(K) &\leq -\frac{1}{2}c(D) + 2c_-(D) + \frac{1}{2}\gamma_{>0}^B + \frac{1}{2}\gamma_{>0}^W - 1 \\ &\leq -\frac{1}{2}c(D) + 2c_-(D) + \frac{1}{3}c(D) + \frac{1}{6}(s(D) - 1) + \frac{2}{3}c_-(D) - 1 \\ &= \frac{1}{6}(s(D) - c(D) - 1) + \frac{8}{3}c_-(D) - 1. \end{aligned}$$

□

Once a signature estimate from canonical Seifert surface is established, Theorem 1.14 (the concordance finiteness) can be proved by almost the same argument as [BDL, Theorem 1.1].

Recall the Levine-Tristram signature $\sigma_\omega(K)$ of a link K is topological concordance invariant whenever $\Delta_\omega(K) \neq 0$ so if we take ω as non-algebraic number, then σ_ω is always a topological concordance invariant.

Proof of Theorem 1.14. Assume, to the contrary that the topological concordance class \mathcal{K} contains infinitely many successively positive knots $\{K_i\}$.

Let D_i be a successively k_i -almost positive diagram of K_i . By assumption, $k_i \leq dg_c^{top}(K) \leq dg(K_i) \leq dg(D_i)$. Therefore by Theorem 1.12

$$\sigma(\mathcal{K}) = \sigma(K_i) \leq -\frac{1}{6}g(D_i) + \frac{4}{3}k_i - \frac{1}{2} \leq \left(-\frac{1}{6} + \frac{4}{3}d\right)g(D_i).$$

Since $d < \frac{1}{8}$, $(-\frac{1}{6} + \frac{4}{3}d) < 0$. Therefore

$$g(D_i) \leq \frac{6}{8d-1}\sigma(\mathcal{K}).$$

Since the canonical genus of the diagrams $\{D_i\}$ are bounded above, there is a finite set of diagrams \mathcal{D} such that each D_i is obtained from one of a diagram $D'_i \in \mathcal{D}$ by inserting full twists N times (often called the \bar{t}_N -move) for some N , at appropriate crossings of D'_i [St1, Theorem 3.1] (see Figure 8). Since every D_i is successively almost positive, we may assume that the base diagram $D'_i \in \mathcal{D}$ is successively positive, and that D_i is obtained by inserting *positive* twists.

Due to the finiteness of \mathcal{D} , there are only finitely many places to insert full twists. Thus there is a diagram $D_{\text{base}} \in \mathcal{D}$ and a crossing c of D_{base} having the following property; for any $N > 0$, there is a knot K in $\{K_i\}$ such that K is obtained from \mathcal{D} by inserting full twist at least N -times at c and inserting appropriate (positive) twists at other crossings.




FIGURE 8. Inserting N full twists at the crossing c (often called the \bar{t}_N -move)

Let $D(N)$ be the diagram obtained from D_{base} by inserting N -full twists at the crossing c , and let $K(N)$ be the knot represented by $D(N)$. The above observation says that there exists a successively positive knot K in the concordance class \mathcal{K} such that $K(N)$ is obtained from K by the positive-to-negative crossing changes. Thus for every $N > 0$ we have an inequality

$$(2.9) \quad \sigma_\omega(K(N)) \geq \sigma_\omega(K) = \sigma_\omega(\mathcal{K}).$$

Let D_0 be the link diagram obtained by smoothing the crossing c of D_{base} and let L be the link represented by D_0 . Since the canonical Seifert surface S_N of $D(N)$ is obtained from the canonical Seifert surface S_0 of D_0 by adding an N -twisted band, the Seifert matrix $A(N)$ of S_N is of the form

$$A(N) = \begin{pmatrix} -N + a & w \\ v & A(S_0) \end{pmatrix}$$

where a is a constant and $A(S_0)$ is the Seifert matrix of S_0 .

Take a non-algebraic $1 \neq \omega \in \{z \in \mathbb{C} \mid |z| = 1\}$ sufficiently close to 1 so that $\sigma_\omega(\mathcal{K}) = 0$ holds. By definition, $\sigma_\omega(K(N))$ and $\sigma_\omega(L)$ are the signatures of

$$\begin{aligned} A_\omega(N) &= (1 - \omega)A(N) + (1 - \bar{\omega})A(N)^T \\ &= \begin{pmatrix} (2 - 2\text{Re}(\omega))(-N + a) & (1 - \omega)w + (1 - \bar{\omega})v^T \\ (1 - \omega)v + (1 - \bar{\omega})w^T & (1 - \omega)A(S_0) + (1 - \bar{\omega})A(S_0)^T \end{pmatrix} \\ A_\omega(0) &= (1 - \omega)A(S_0) + (1 - \bar{\omega})A(S_0)^T. \end{aligned}$$

Therefore by the cofactor expansion

$$\det A_\omega(N) = (2 - 2\text{Re}(\omega))(-N + a) \det A_\omega(0) + C$$

where C is a constant that does not depend on N . Since we have chosen ω so that it is non-algebraic, $\det A_\omega(S_0) \neq 0$. Thus when N is sufficiently large, the sign of $\det A_\omega(N)$ and $\det A_\omega(S_{D_0})$ are different, which means that the matrix $A_\omega(N)$ has one more negative eigenvalue than $A_\omega(0)$. Thus for sufficiently large N

$$\sigma_\omega(K(N)) = \sigma_\omega(L) - 1.$$

On the other hand, since D_0 is successively positive, by Theorem 1.12 $\sigma_\omega(L) \leq 0$. Therefore by (2.9)

$$\sigma_\omega(\mathcal{K}) \leq \sigma_\omega(K(N)) = \sigma_\omega(L) - 1 \leq -1.$$

Since we have chosen ω so that $\sigma_\omega(\mathcal{K}) = 0$, this is a contradiction. \square

3. DISCUSSIONS AND QUESTIONS

Our argument so far justifies the assertion that a (good) successively k -almost positive diagram is natural and more appropriate generalization of positive links than a k -almost positive diagram. However, since our results also says that the difference between (almost) positive links and (good) successively k -almost positive links are subtle to distinguish.

In fact, at this moment we do not know an example of links which are (good) successively almost positive but is not almost positive, although we believe that there are many such links. This is mainly because we have no useful technique to exclude various candidates are indeed not almost positive.

3.1. Positive v.s. almost positive. Since positive links and almost positive links already share many properties, distinguishing almost positive links with positive link is not an easy task.

The simplest example of almost positive, but not positive knot is 10_{145} ; 10_{145} admits an almost positive diagram of type I. The non-positivity can be detected by Cromwell's theorem [Cr] that $c(K) \leq 4g(K)$ if K is positive and its Conway polynomial $\nabla_K(z)$ is monic, or, the property of the HOMFLY and the Kauffman polynomial of positive knot (3.1) which we discuss in the next section.

3.2. Almost positive of type I v.s. Almost positive of type II. Our argument so far tells that a good successively almost positive diagram has better properties than a usual almost positive diagram.

However, we note that it can happen non-good almost positive links share a property of positive links that fails for good almost positive links.

An almost positive diagram D is

- of type I if D is good successively almost positive; for two Seifert circles s and s' connected by the unique negative crossing c_- , there are no other crossings connecting s and s' .
- of type II otherwise; there are positive crossings connecting s and s' .

An almost positive diagram of type I is nothing but a good successively almost positive diagram. For an almost positive diagram D , S_D attains the maximum euler characteristic if and only if D is of type I [St2]. In particular, when D is an almost positive diagram of type II, we have $\chi(K) = \chi(D) + 2$. The dichotomy plays a fundamental role in a study of almost positive diagrams and links – the proof of various properties of almost positive links often splits into the analysis of two cases [St2, FLL].

Let $D_K(a, z)$ be the Dubrovnik version of the Kauffman polynomial; $D_K(a, z) = a^{-w(D)} \Lambda_D(a, z)$, where $\Lambda_D(a, z)$ is the regular isotopy invariant defined by the skein relations

$$\Lambda_{\nearrow} \searrow (a, z) - \Lambda_{\searrow} \nearrow (a, z) = z(\Lambda_{\smile} - \Lambda_{\frown}) (a, z),$$


$$\Lambda_{\bigcirc} (a, z) = a \Lambda_{\smile} (a, z), \Lambda_{\bigcirc} (a, z) = a^{-1} \Lambda_{\frown} (a, z), \Lambda_{\bigcirc} (a, z) = 1.$$

We express the Dubrovnik polynomial $D_K(a, z) = \sum_i D_K(z; i) a^i$. Similarly, we express the HOMFLY polynomial $P_K(v, z) = \sum_i P_K(z; i) v^i$. In [Yo] Yokota showed that when K is positive,

$$(3.1) \quad P_K(z; 1 - \chi(K)) = D_K(z; 1 - \chi(K)) \neq 0$$

holds. Moreover, this is a non-negative polynomial [Cr, Corollary 4.3].

At first glance, this coincidence sounds mysterious but it turns out Legendrian link point of view brings illuminating explanation.

For a basis of Legendrian links we refer to [Et]. Let \mathcal{D} be the front diagram of a Legendrian link K . The subset ρ of the set of crossings of \mathcal{D} is a *ruling* if the diagram \mathcal{D}_ρ obtained from \mathcal{D} by taking the horizontal smoothing  at each crossing in ρ satisfies the following conditions;

- Each component of \mathcal{D}_ρ is the standard diagram of the Legendrian unknot (i.e it contains two cusps and no crossings).
- For each $c \in \rho$, let P and Q be the component of \mathcal{D}_ρ that contains the smoothed arcs at c . Then P and Q are different component of \mathcal{D}_ρ , and in the vertical slice around c , two components P and Q are not nested – they are aligned in one of the configuration in Figure 9 (ii).




FIGURE 9. (i) Front diagram \mathcal{D} and \mathcal{D}_ρ for $\rho = \{b\}$; ρ is not a ruling. (ii) Normality condition for the ruling.

A ruling is *oriented* if every element of ρ is a positive crossing. Let $\Gamma(\mathcal{D})$ and $O\Gamma(\mathcal{D})$ be the set of rulings and oriented rulings, respectively.

For a ruling ρ we define

$$j(\rho) = \#\rho - l\text{-cusp}(\mathcal{D}) + 1,$$

where $l\text{-cusp}(\mathcal{D})$ denotes the number of the left cusps of \mathcal{D} . In [Rut, Theorem 3.1, Theorem 4.3], Rutherford showed that the ruling polynomials, a graded count of the (oriented) rulings, are equal to the coefficient polynomials of Dubrovinik and HOMFLY polynomials, respectively;

$$(3.2) \quad D_K(z; tb(\mathcal{D}) + 1) = \sum_{\rho \in \Gamma(\mathcal{D})} z^{j(\rho)}, \quad P_K(z; tb(\mathcal{D}) + 1) = \sum_{\rho \in O\Gamma(\mathcal{D})} z^{j(\rho)}.$$

Here $tb(\mathcal{D})$ is the Thurston-Benneuqin invariant, given by

$$tb(\mathcal{D}) = w(\mathcal{D}) - l\text{-cusp}(\mathcal{D}).$$

In particular, by the HOMFLY/Kauffman bound of the maximum Thurston-Bennequin number $\overline{tb}(K)$

$$\overline{tb}(K) + 1 \leq \min \deg_v P_K(v, z), \quad \overline{tb}(K) + 1 \leq \min \deg_a D_K(v, z),$$

if a front diagram \mathcal{D} admits a ruling, then \mathcal{D} attains the maximum Thurston-Bennequin number $\overline{tb}(K)$ among its topological link types, and

$$\overline{tb}(K) + 1 = \min \deg_v P_K(v, z) = \min \deg_a D_K(v, z).$$

One can view a positive diagram D as a front diagram \mathcal{D} [Ta]; each positive crossing is put so that in the form , and each Seifert circle forms a front diagram having exactly two cusps (so $l\text{-cusp}(\mathcal{D}) = s(D)$). Then the set of all crossings forms a (oriented) ruling. Thus \mathcal{D} attains the maximum Thurston-Bennequin number. In particular,

$$\overline{tb}(K) = tb(\mathcal{D}) = c(D) - s(D) = -\chi(D) = -\chi(K).$$

Moreover, since all the crossings of \mathcal{D} are positive, $\Gamma(\mathcal{D}) = O\Gamma(\mathcal{D})$. Thus the ruling polynomial formula (3.2) of the coefficients of the Dubrovnik and the HOMFLY polynomials leads to Yokota's equality (3.1) and its non-negativity.

A mild generalization of this argument shows that an almost positive link of type II shares the same property;

Theorem 3.1. *If K admits an almost positive diagram of type II, then*

$$P_K(z; 1 - \chi(K)) = D_K(z; 1 - \chi(K)) \neq 0$$

and it is a non-negative polynomial.

Proof. We view an almost positive diagram of type II as a front diagram \mathcal{D} as shown in Figure 10, where the box represents a positive diagram part viewed as a front diagram. A ruling of \mathcal{D} cannot contain the negative crossing of \mathcal{D} , so $\Gamma(\mathcal{D}) = O\Gamma(\mathcal{D})$. On the other hand, since D is of type II, the Seifert circles s and s' connected by the negative crossing c_- is also connected by a positive crossing. We take such a crossing c so that there are no such crossing between c and c^- . Let $\rho = C(\mathcal{D}) \setminus \{c_-, c\}$. Then ρ is a ruling so $\Gamma(\mathcal{D}) = O\Gamma(\mathcal{D}) \neq \emptyset$, and

$$\overline{tb}(K) = tb(\mathcal{D}) = (c(D) - 2) - s(D) = -\chi(D) - 2 = -\chi(K).$$

Thus by (3.2) we get the desired identity. \square

Corollary 3.2. *If K is almost positive of type II,*

$$\min \deg_a D_K(z, a) = \max \deg_z P_K(v, z) = 1 - \chi(K).$$

Since there are almost positive knot of type I that fails to have the property stated in Theorem 3.1 (for example 10_{145}), we have the following, which was mentioned in [St2] without proof;

Corollary 3.3. *There exists an almost positive knot of type I which does not admit an almost positive diagram of type II.*

A similar argument can be applied to establish the same equality for certain type of not-good successively almost positive diagrams.





FIGURE 10. (i) Front diagram of almost positive diagram of type II; the box represent a front diagram consisting of positive crossings. (ii) Non-trivial ruling $\rho = C(\mathcal{D}) \setminus \{c_-, c\}$ and its resolution \mathcal{D}_ρ

3.3. (Good) Successively almost positive v.s. strongly quasipositive. A strongly quasipositive knots are not necessarily successively almost positive; for a given link L , there exists a strongly quasipositive link L' such that $\nabla_L(z) = \nabla_{L'}(z)$ [Rud2, 88 Corollary]. See [Si, Section 5] for details and concrete examples. Thus the Conway polynomial of strongly quasipositive link may not be non-negative.

3.4. Questions. We summarize the relations among various positivities discussed in the paper;



Question 1. Are the inclusions (a), (b), (c), (d) strict?

The strictness question of (a) appeared in [St2, Question 3]. We have seen that

$$\{\text{Almost positive of type I}\} \not\subset \{\text{Almost positive of type II}\}$$

We are asking whether the converse inclusion

$$\{\text{Almost positive of type II}\} \subset \{\text{Almost positive of type I}\}$$

holds or not.

As for the strictness of (b), one candidate of the properties of almost positive links which are not extended to good successively almost positive links is [St2, Theorem 5]; it says that $\min \deg V_K(t) = \frac{1}{2}(1 - \chi(K))$ when K is almost positive. Although several simple successively almost positive links still satisfy the same property, the proof of the property utilizes a state-sum argument which we cannot generalize to successively almost positive case in an obvious manner.

Question 2. Is it true that $\min \deg V_K(t) = \frac{1}{2}(1 - \chi(K))$ when K is good successively almost positive?

As for the strictness of (d), it is interesting to ask to what extent a successively almost positive link shares the same properties as good successively almost positive links;

Question 3. *Assume that K is successively almost positive.*

- *Is K strongly quasipositive?*
- *Is $\max \deg_z \nabla_K(z) = 1 - \chi(K)$ when K is non-split?*
- *Is $\max \deg_z P_K(v, z) = \min \deg_v P_K(v, z) = 1 - \chi(K)$?*

In Theorem 1.14 we showed the finiteness of successively almost positive knots in a topological concordance class, under the assumption that the topological concordance genus is sufficiently large. It is natural to investigate whether this additional assumption is necessary or not.

Question 4. *Does every topological concordance class contain at most finitely many successively k -almost positive knots?*

In [Oz] Ozawa showed the visibility of primeness for positive diagrams; a link represented by a positive diagram D is non-prime if and only if the diagram D is non-prime.

Question 5. *If a knot K prime if K is represented by a prime, good successively k -almost positive diagram?*

REFERENCES

- [AH] K. Appel and W. Haken, *Every planar map is four colorable*, Contemporary Mathematics, 98. American Mathematical Society, Providence, RI, 1989. xvi+741 pp.
- [BDL] S. Baader, P. Dehornoy, and L. Liechti, *Signature and concordance of positive knots*. Bull. Lond. Math. Soc. 50 (2018), no. 1, 166–173.
- [Cr] P. Cromwell, *Homogeneous links*, J. London Math. Soc. (2) 39 (1989), no. 3, 535–552.
- [Et] J. Etnyre, *Legendrian and transversal knots*, Handbook of knot theory, 105–185, Elsevier B. V., Amsterdam, 2005.
- [FLL] P. Feller, L. Lewark and A. Lobb, *Almost positive links are strongly quasipositive*, arXiv:1809.06692v2.
- [GL] C. Gordon, C. A. Litherland, *On the signature of a link*, Invent. Math. 47 (1978), no. 1, 53–69.
- [HIK] J. Hamer, T. Ito and K. Kawamuro, *Positivities of knots and links and the defect of Bennequin inequality*, Exp. Math. to appear.
- [IMT] T. Ito, K. Motegi, and M. Teragaito, *Generalized torsion and Dehn filling*, Topol. Appl. 301 (2021) 107515.
- [Li] C. Livingston, *Computations of the Ozsváth-Szabó knot concordance invariant*. Geom. Topol. 8 (2004), 735–742.
- [MP] K. Murasugi and J. Przytycki, *An index of a graph with applications to knot theory*, Mem. Amer. Math. Soc. 106 (1993), no. 508, x+101 pp.
- [Oz] M. Ozawa, *Closed incompressible surfaces in the complements of positive knots*. Comment. Math. Helv. 77 (2002), no. 2, 235–243.
- [OS] P. Ozsváth, and Z. Szabó *Knot Floer homology and the four-ball genus*. Geom. Topol. 7 (2003), 615–639.
- [PT] J. Przytycki and K. Taniyama, *Almost positive links have negative signature*. J. Knot Theory Ramifications 19 (2010), no. 2, 187–289.
- [Ra] J. Rasmussen, *Khovanov homology and the slice genus*, Invent. Math. 182 (2010), no. 2, 419–447.
- [Rud1] L. Rudolph, *Constructions of quasipositive knots and links. III. A characterization of quasipositive Seifert surfaces*, Topology 31 (1992), no. 2, 231–237.
- [Rud2] L. Rudolph, *Knot theory of complex plane curves*, Handbook of knot theory, 349–427, Elsevier B. V., Amsterdam, 2005.

- [Rut] D. Rutherford, *Thurston-Bennequin number, Kauffman polynomial, and ruling invariants of a Legendrian link: the Fuchs conjecture and beyond*. Int. Math. Res. Not. 2006, Art. ID 78591, 15 pp.
- [Sh] A. Shumakovitch, *Rasmussen invariant, slice-Bennequin inequality, and sliceness of knots*. J. Knot Theory Ramifications 16 (2007), no. 10, 1403–1412.
- [Si] M. Silvero, *Strongly quasipositive links with braid index 3 have positive Conway polynomial*, J. Knot Theory Ramifications 25 (2016), no. 12, 1642015, 14 pp.
- [St1] A. Stoimenow, *Knots of genus one or on the number of alternating knots of given genus*. Proc. Amer. Math. Soc. 129 (2001), no. 7, 2141–2156.
- [St2] A. Stoimenow, *On polynomials and surfaces of variously positive links*, J. Eur. Math. Soc. (JEMS) 7 (2005), no. 4, 477–509.
- [Ta] T. Tanaka, *Maximal Bennequin numbers and Kauffman polynomials of positive links*. Proc. Amer. Math. Soc. 127 (1999), no. 11, 3427–3432.
- [Yo] Y. Yokota, *Polynomial invariants of positive links*, Topology 31 (1992), no. 4, 805–811.

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN
Email address: tetitoh@math.kyoto-u.ac.jp