

Geometrodynamical description of two-dimensional electrodynamics

Rodrigo F. Sobreiro*

*UFF - Universidade Federal Fluminense, Instituto de Física,
Av. Litorânea, s/n, 24210-346, Niterói, RJ, Brasil.*

Abstract

Two-dimensional pure electrodynamics is mapped into two-dimensional gravity in the first order formalism at classical and quantum levels. Due to the fact that the degrees of freedom of these two theories do not match, we are enforced to introduce extra fields from the beginning. These fields are introduced through a BRST exact boundary term, so they are harmless to the physical content of the theory. The map between electromagnetism and gravity fields generate a non-trivial Jacobian, which brings extra features (but also harmless to the physical content of the gravity theory) at quantum level.

1 Introduction

Two-dimensional electrodynamics is of great interest in Physics, mainly because of its usage in two-dimensional condensed matter systems, see [1–3] and references therein - It is worth mentioning that two-dimensional classical electrodynamics is an exact solvable system [4]. But also as toy models used to probe more complicated systems. The latter statement is also valid for gravity, specially in the pursuing of a consistent quantization of gravity [5–10]. In the present work we show how we can connect both theories.

Geometrodynamical descriptions of gauge systems appear in the literature quite often, see for instance [11–18] and references therein. Our aim in the present work is to establish a consistent map relating two-dimensional electrodynamics and two-dimensional gravity in the first order formalism (described by the zweibein and the Lorentz connection as fundamental fields). The starting electromagnetic action we work is composed by the Maxwell term and the Chern-Pontryagin density. The main problem we have to face is that two-dimensional electromagnetism is a gauge theory with two degrees of freedom (physical and non-physical) while two-dimensional gravity in the first order formalism carries six degrees of freedom (physical and non-physical). Thence, a naive map between both theories, although possible, is ill defined. To account for the extra degrees of freedom at the gravity side, we introduce, at the electromagnetism side, a set of extra fields via an extra term in the action. The auxiliary term is constructed as a Becchi-Rouet-Stora-Tyutin (BRST) exact quantity in order to not mess with the physical degrees of freedom. Moreover, the auxiliary action is a boundary term. So we can safely say that this term should be “twice harmless”.

*rodrigo_sobreiro@id.uff.br

To actually perform the map we choose a non-trivial way to relate the fields in both theories. Although non-trivial, the map is well defined. At classical level, the gravity theory obtained is composed by the Einstein-Hilbert term which, in two-dimensions, coincides with the Gauss-Bonnet term, a cosmological constant term and another BRST exact boundary term with matter fields. At quantum level, because the map is nontrivial, a non-trivial Jacobian appears in the generating functional. Nevertheless, the Jacobian can be localized with the help of extra fields. The extra fields can also be collected in a BRST exact term, ensuring that the physical sector of the model is untouched.

This work is organized as follows. In Section 2 we construct the two-dimensional electromagnetic theory. In Section 3 we construct the two-dimensional gravity theory. Section 4 is devoted to the map between both theories at classical level. The equivalence is then discussed at quantum level in Section 5. Finally, our conclusions are displayed in Section (6).

2 Electrodynamics

The fundamental ingredient in electrodynamics is the gauge field 1-form $A = A_i dx^i$ where $i, j, k, \dots \in \{0, 1\}$ and dx^i refer to a two-dimensional manifold which we take, for simplicity, as the Euclidean space-time \mathbb{R}^2 . The field A is a connection with respect to the gauge group $U(1)$. The gauge transformation of A is

$$\delta A = d\alpha , \quad (2.1)$$

where $u(x) = \exp[i\alpha(x)] \in U(1)$. The gauge invariant field strength 2-form is given by

$$F = dA = \frac{1}{2} F_{ij} dx^i dx^j . \quad (2.2)$$

The dual field strength $*F$ is obtained from the action of the Hodge dual $*$ on the field strength by means of

$$*F = \frac{1}{2} \epsilon^{ij} F_{ij} , \quad (2.3)$$

where ϵ^{ij} is the Levi-Civita tensor.

At this point, the action we consider is the most general second order action with the properties of being gauge invariant, local and polynomial in the gauge field and its derivatives, and power counting renormalizable, namely

$$S_e = \int \left(\kappa F + \frac{1}{2} F * F \right) , \quad (2.4)$$

where the parameter κ is a mass parameter. Moreover, although we are not considering a specific type of matter interaction, the fundamental electric charge e is at our disposal. The mass dimensions of all (tensorial component) ingredients are given by $[A] = 0$ and $[d] = [e] = [k] = 1$. The second term in action (2.4) is the usual Maxwell term while the first term is known as the topological Chern-Pontryagin density. For a two-dimensional compact region $\Omega \in \mathbb{R}^2$ with boundary $\partial\Omega$, it provides the quantization of the circulation of A along the boundary,

$$\kappa \int_{\Omega} F = \kappa \oint_{\partial\Omega} A = 4\pi\kappa p , \quad (2.5)$$

where $p \in \mathbb{Z}$ is the winding number of the gauge configuration A .

Although we are dealing with electrodynamics at classical level, BRST technology [19] will be evoked along the entire paper. Therefore, defining the Faddeev-Popov ghost field c , and s as the BRST nilpotent

operator, the BRST transformations read

$$\begin{aligned} sA &= dc \\ sc &= 0. \end{aligned} \tag{2.6}$$

For gauge fixing purposes and completeness, one typically introduces the Faddeev-Popov anti-ghost field and the Lautrup-Nakanishi field as a BRST doublet,

$$\begin{aligned} s\bar{c} &= b, \\ sb &= 0. \end{aligned} \tag{2.7}$$

Nevertheless, we will not deal with gauge fixing issues in the present work and fix our attention exclusively to the map.

The electromagnetic action (2.4) is clearly a BRST closed, but not exact, quantity. Thence, it belongs to the nontrivial sector of the BRST cohomology. Consequently, S_{ed} carries physical information. The BRST symmetry is typically used to introduce a gauge fixing in the elimination of spurious degrees of freedom of S_{ed} .

3 Lovelock-Cartan gravity

The second theory we deal with is two-dimensional gravity in the first order formalism [20–26]. It is constructed over a generic two-dimensional differential manifold M which is assumed to be Hausdorff, paracompact and orientable [27]. Over M we define two fundamental fields associated with the geometric properties of M itself, the zweibein 1-form $e^a = e^a_\mu dX^\mu$ and the connection 1-form $\omega^{ab} = \omega^{ab}_\mu dX^\mu$, with X being a generic point in M . The frame indexes $a, b, c, \dots, h \in \{0, 1\}$ are associated with coordinates at the tangent space $T_X(M) \sim T_X$ while world indexes $\mu, \nu, \alpha, \dots \in \{0, 1\}$ are associated with local space-time coordinates. The coordinated point $X \in M$ is introduced in different notation of $x \in \mathbb{R}^2$ in order to avoid confusion with the formulation of electrodynamics in the previous section. The local isometries in tangent space are characterized by the Abelian gauge group $SO(2)$. The $SO(2)$ gauge transformations are typically characterized by

$$\begin{aligned} \delta\omega^a_b &= d\alpha^a_b, \\ \delta e^a &= \alpha^a_b e^b, \end{aligned} \tag{3.1}$$

where $U(X) = \exp(i\alpha^{ab}L_{ab}) \in SO(2)$. The generator of the two-dimensional orthogonal group is denoted by L_{ab} which is anti-symmetric in a and b .

The zweibein is known to have two essential properties. The first one is that it carries the metric properties of spacetime since one can derive the metric tensor from it by means of $g_{\mu\nu} = \eta_{ab}e^a_\mu e^b_\nu$. Second, that it provides the existence of local inertial frames at each point X by means of $dX^a = e^a_\mu dX^\mu$, ensuring the validity of the equivalence principle [28–30]. The coordinate system X^a is identified with the tangent space and all possible local inertial frames are connected by the orthogonal gauge group. Thence, gravity is just a gauge theory with the extra principle that the gauge group is identified with the local isometry group.

Torsion and curvature 2-forms are, respectively, given by

$$\begin{aligned} T^a &= De^a = de^a + \omega^a_b e^b, \\ R^a_b &= d\omega^a_b. \end{aligned} \tag{3.2}$$

In addition, we have at our disposal two invariant objects: the Levi-Civita tensor ϵ_{ab} and the tangent space metric η_{ab} .

With the above ingredients at hand, we are able to construct the most general local and polynomial action, power-counting renormalizable, depending only on the fields and their first order derivatives and not explicitly depending on the metric. The result is the well known Mardones-Zanelli action [24–26, 31] describing Lovelock-Cartan (LC) two-dimensional gravity,

$$S_{LC} = \frac{1}{8\pi G} \int \epsilon_{ab} \left(R^{ab} + \frac{\Lambda^2}{2} e^a e^b \right), \quad (3.3)$$

where G and Λ are Newton and cosmological constants, respectively. The mass dimension of the relevant objects in the action (3.3) are $[d] = [\omega^a_b] = [\Lambda] = 1$ and $[e^a] = [G] = 0$. The last term at the action (3.3) is recognized as the cosmological constant term while the first one is the Einstein-Hilbert term. The Einstein-Hilbert term is topological in two dimensions because it coincides with the Gauss-Bonnet topological invariant. In fact, for a compact and orientable sub-manifold $\Omega \in M$ with boundary $\partial\Omega$, we have

$$\int_{\Omega} \epsilon_{ab} R^{ab} = \oint_{\partial\Omega} \epsilon_{ab} \omega^{ab} = 2\pi\chi_2, \quad (3.4)$$

where χ_2 is the Euler characteristic of Ω .

Again, it is convenient at this point to establish the BRST symmetry of (3.3). Defining the gravitational Faddeev-Popov ghost [31–33] by c^{ab} , the action (3.3) is invariant under

$$\begin{aligned} s\omega^{ab} &= dc^{ab}, \\ se^a &= c^a_b e^b, \\ sc^{ab} &= 0. \end{aligned} \quad (3.5)$$

The action S_{LC} carries physical information since it is a BRST closed but not an exact quantity. Again, for gauge fixing purposes and completeness, we can define

$$\begin{aligned} s\bar{c}^a_b &= b^a_b, \\ sb^a_b &= 0, \end{aligned} \quad (3.6)$$

with \bar{c}^a_b being the gravitational Faddeev-Popov anti-ghost field and b^a_b the Lautrup-Nakanishi field.

4 Electrodynamics-Gravity equivalence

In this section we discuss, at classical level, how we can connect the electromagnetic action (2.4) to the LC action (3.3).

4.1 Naive map

Following [13], we can introduce a 2-form auxiliary field θ , with $[\theta] = 1$, and rewrite action (2.4) as

$$S_e = - \int \left(\kappa F + \theta * F - \frac{1}{2} \theta * \theta \right). \quad (4.1)$$

The auxiliary field is obviously gauge (and BRST) invariant. It is quite intuitive that this action can be mapped into (3.3) due to the homomorphism $U(1) \mapsto SO(2)$. In fact, the following map between fields¹

$$\begin{aligned}\theta(x) &\mapsto \mu \epsilon_{ab} e^a(X) e^b(X), \\ A(x) &\mapsto \frac{1}{2\mu} \epsilon_{ab} \omega^{ab}(X),\end{aligned}\tag{4.2}$$

provides the desired result for $[\mu] = 1$. In (4.2), $x \in \mathbb{R}^2$ and $X \in M$. Along with (4.2), the gauge transformation (2.1) can be mapped to (3.1) in two steps: First, $\alpha(x) \mapsto \epsilon_{ab} \alpha^{ab}(X)$; Second, since θ is gauge invariant, so must be the *rhs* of the first map in (4.2); Therefore, e^a must be an $SO(2)$ invariant quantity.

At the level of the action, the map (4.2) induces a Lovelock-Cartan gravity of the form

$$S_{sLC} = \int \frac{1}{8\pi G} \epsilon_{ab} \left(R^{ab} - \frac{\Lambda^2}{2} e^a e^b \right),\tag{4.3}$$

with²

$$\begin{aligned}G &= \frac{1}{4\pi} \frac{\mu}{(\kappa + 2\mu)}, \\ \Lambda^2 &= 4 \frac{\mu^3}{(\kappa + 2\mu)}.\end{aligned}\tag{4.4}$$

Equations (4.4) can be solved for μ , namely

$$\mu = \frac{1}{\sqrt{\pi G}} \frac{\Lambda}{2}.\tag{4.5}$$

Up to a minus sign in the cosmological constant, the action (4.3) is totally equivalent to the action (3.3). The problem with the present proposal (4.2) is that at one side we have only three degrees of freedom (from $A \equiv (A_0, A_1)$ and $\theta \equiv (\theta_{01})$) and on the other side we have six degrees of freedom ($e^a \equiv (e_0^0, e_1^0, e_0^1, e_1^1)$ and $\omega^{ab} \equiv (\omega_0^{01}, \omega_1^{01})$). Hence, to the map work in both ways, we need extra fields to account for the extra degrees of freedom.

4.2 Improved map

To have the right number of degrees of freedom we introduce the following matter action, to be added to electro-dynamical action (4.1),

$$S_m = \int d \left[(\bar{\varphi} \varphi - \bar{\zeta} \zeta) + (\bar{w} w - \bar{z} z) \right],\tag{4.6}$$

with $(\bar{\varphi}, \varphi)$ and (\bar{w}, w) being real bosonic fields while $(\bar{\zeta}, \zeta)$ and (\bar{z}, z) are hermitian fermionic fields. The fields $(\bar{\varphi}, \bar{\zeta}, \bar{w}, \bar{z})$ are 0-forms while (φ, ζ, w, z) are 1-forms. The mass dimension of all extra fields are

¹The factors depending on μ are required in order to adjust the mass dimensions of the fields. The form of the mapping (4.2) is not unique. In fact, the factor μ is a generic factor with mass dimension 1 constructed from κ and e (e.g., $\mu = e^q \kappa^m f(e^p \kappa^{-p})$ with $q + m = 1$, $p \in \mathbb{R}$ and f any analytical function). The target action (4.3) would be the same but the values for Newton and cosmological constants would be different for each possible μ .

²It is worth mention that, for $\mu = a\kappa$ with $a \in \mathbb{R}$, Newton's constant in (4.4) is fixed to $\frac{1}{4\pi(2a+1)}$. This is consistent with the fact that $[G] = 0$.

$[\bar{\varphi}] = [\bar{\zeta}] = [\bar{w}] = [\bar{z}] = 0$ and $[\varphi] = [\zeta] = [w] = [z] = 1$. The fact that we introduced integer spin fermions must be clarified. The reason is that we are aiming the quantization of the model via BRST framework. Indeed, the auxiliary fields form a BRST octet system and do not affect the physical content of the model [19]. In fact, under BRST symmetry, the new auxiliary fields transform as

$$\begin{aligned} s\bar{\zeta} &= \bar{\varphi}, \\ s\bar{\varphi} &= 0, \\ s\varphi &= \zeta, \\ s\zeta &= 0, \end{aligned} \tag{4.7}$$

and

$$\begin{aligned} s\bar{z} &= \bar{w}, \\ s\bar{w} &= 0, \\ sw &= z, \\ sz &= 0. \end{aligned} \tag{4.8}$$

And the auxiliary action S_m can be written as a BRST exact quantity,

$$S_m = -s \int d(\bar{\zeta}d\varphi + \bar{z}dw). \tag{4.9}$$

Thus, from the BRST point of view [19] we are not introducing any physical degree of freedom. Moreover, action (4.6) is a surface term, only trivially affecting the boundary of \mathbb{R}^2 . For completeness and further use, all quantum numbers of the fields on the electromagnetic side are displayed in Table 1 (including the Faddeev-Popov ghosts and the Lautrup-Nakanishi field).

Fields	A	θ	b	c	\bar{c}	$\bar{\varphi}$	φ	$\bar{\zeta}$	ζ	\bar{w}	w	\bar{z}	z
Dimension	0	1	1	0	0	0	1	0	1	0	1	0	1
Ghost number	0	0	0	1	-1	0	0	-1	1	0	0	-1	1
Form rank	1	2	0	0	0	0	1	0	1	0	1	0	1

Table 1: Quantum numbers of the fields in the electromagnetic theory.

The full action to start with is then

$$S_{em} = S_e + S_m. \tag{4.10}$$

It is easy to see that we have extra degrees of freedom at the side of electrodynamics, a total of fifteen against the six degrees of freedom at the gravity side. The consequence is that we will end up with extra degrees of freedom at the target gravity theory. In fact, a possible map accounting for this reads

$$\begin{aligned} (\bar{\varphi}(x), \varepsilon\bar{\zeta}(x)) &\equiv \bar{\Phi}^a(x) \longmapsto \sigma^a(X), \\ (\varphi(x), -\varepsilon^{-1}\zeta(x)) &\equiv \Phi^a(x) \longmapsto \mu\phi^a_b(X)e^b(X), \end{aligned} \tag{4.11}$$

and

$$\begin{aligned} \bar{w}(x) &\longmapsto \bar{\eta}(X), \\ \bar{z}(x) &\longmapsto \bar{\chi}(X), \\ w(x) &\longmapsto \mu\eta_a(X)e^a(X), \\ z(x) &\longmapsto \mu\chi_a(X)e^a(X), \end{aligned} \tag{4.12}$$

with σ , ϕ^a_b , $\bar{\eta}$, and η^a are real bosonic fields while $\bar{\chi}$ and χ^a are hermitian ghost scalar fields. Moreover, the field ϕ^a_b is assumed to be a Lorentz-algebra-valued field, therefore accounting for only one degree of freedom. Furthermore, all new fields are dimensionless. The maps (4.11) maps 0-forms and 1-forms in the electromagnetic side into algebra-valued 0-forms in the gravity side. Such subtlety is required because the maps must not be $SO(2)$ invariant, such as (4.2) and (4.12). Otherwise, we would still get redundancies and the full set of maps would be not invertible. In fact, the Jacobian of the maps (4.2), (4.11) and (4.12) can be calculated. The result, up to a normalization factor, is simply given by³

$$J = \mu^4 \det^{3/2}(\phi^a_b) \det^3(e^a_\mu) . \quad (4.13)$$

Employing the maps (4.2), (4.11) and (4.12), the action S_m is mapped into

$$\begin{aligned} S_{int} &= \mu \int d \left(\sigma_a \phi^a_b e^b + \bar{\eta} \eta_a e^a - \bar{\chi} \chi_a e^a \right) \\ &= \mu \int \left[D \sigma_a \phi^a_b e^b + \sigma_a (D \phi^a_b e^b + \phi^a_b T^b) + d \bar{\eta} \eta_a e^a + \bar{\eta} (D \eta_a e^a + \eta_a T^a) + \right. \\ &\quad \left. - d \bar{\chi} \chi_a e^a + \bar{\chi} (D \chi_a e^a - \chi_a T^a) \right] . \end{aligned} \quad (4.14)$$

It is quite convenient to redefine the field ϕ^{ab} as

$$\phi_{ab} = \epsilon_{ab} \phi , \quad (4.15)$$

so, action (4.14) reads

$$S_{int} = \mu \int d \left(\epsilon_{ab} \sigma^a e^b \phi + \bar{\eta} \eta_a e^a - \bar{\chi} \chi_a e^a \right) , \quad (4.16)$$

This action is, again, a BRST exact quantity,

$$S_{int} = -\mu s \int d \left(\epsilon_{ab} \xi^a e^b \phi - \bar{\chi} \eta_a e^a \right) , \quad (4.17)$$

if we define,

$$\begin{aligned} s \xi^a &= \sigma^a + c^a_b \xi^b , \\ s \sigma^a &= c^a_b \sigma^b , \\ s \bar{\chi} &= \bar{\eta} , \\ s \bar{\eta} &= 0 , \\ s \eta^a &= \chi^a + c^a_b \eta^b , \\ s \chi^a &= c^a_b \chi^b , \\ s \phi &= 0 . \end{aligned} \quad (4.18)$$

Therefore, the resulting gravity theory is finally given by

$$\begin{aligned} S_{grav} &= S_{sLC} + S_{int} \\ &= \int \frac{1}{8\pi G} \epsilon_{ab} \left(R^{ab} - \frac{\Lambda^2}{2} e^a e^b \right) + \mu \int d \left(\epsilon_{ab} \sigma^a e^b \phi + \bar{\eta} \eta_a e^a - \bar{\chi} \chi_a e^a \right) , \end{aligned} \quad (4.19)$$

which is equivalent to (4.3) up to a BRST trivial boundary term.

For completeness, the quantum numbers of the fields on the gravity theory are collected in Table 2.

³See Appendix A for more details.

Fields	ω^a_b	e^a	b^a_b	c^a_b	\bar{c}^a_b	$\bar{\eta}$	η^a	ϕ^a_b	$\bar{\chi}$	χ^a	σ	$\bar{\eta}$
Dimension	1	0	0	0	0	0	1	0	0	1	0	1
Ghost number	0	0	0	1	-1	0	0	0	-1	1	0	-1
Form rank	1	1	0	0	0	0	0	0	0	0	0	0

Table 2: Quantum numbers of the fields in the gravity theory.

5 Electrodynamics-gravity equivalence at quantum level

The same map can be performed at quantum level. Thence, two-dimensional electrodynamics can be mapped into two-dimensional quantum gravity. Let us start with the generating functional of the electromagnetic theory

$$Z_{qed} = \int [\mathcal{D}\Phi_{ed}] e^{-S_{ed} - S_m}, \quad (5.1)$$

with $[\mathcal{D}\Phi_{ed}] = \mathcal{D}A\mathcal{D}\theta\mathcal{D}\bar{\varphi}\mathcal{D}\phi\mathcal{D}\bar{\zeta}\mathcal{D}\zeta\mathcal{D}\bar{w}\mathcal{D}w\mathcal{D}\bar{z}\mathcal{D}z$ being the functional measure. Applying the maps (4.2), (4.11) and (4.12) and making use of the Jacobian (4.13), one easily achieves

$$Z_{grav} = \int [\mathcal{D}\Phi_{grav}] \mu^4 \det^{3/2}(\phi^a_b) \det^3(e^a_\mu) e^{-S_{grav}}, \quad (5.2)$$

with $[\mathcal{D}\Phi_{grav}] = \mathcal{D}\omega\mathcal{D}e\mathcal{D}\sigma\mathcal{D}\phi\mathcal{D}\bar{\eta}\mathcal{D}\eta\mathcal{D}\bar{\chi}\mathcal{D}\chi$. The determinants can be localized in the usual way. First, we consider

$$\mu^2 \det^3(e^a_\mu) = \mu^2 \det^4(e^a_\mu) \det^{-1}(e^a_\mu) = \int \mathcal{D}\bar{Y}\mathcal{D}Y\mathcal{D}Z \exp \left[-\mu^2 \int d^2X \left(\bar{Y}_a e^a_\mu Y^\mu + \frac{1}{2} Z_a e^a_\mu Z^\mu \right) \right],$$

with \bar{Y}^a and Y^μ being ghost fields while Z_a and Z^μ are bosonic fields. Moreover, the following relations are valid

$$\begin{aligned} Z^a &= e^a_\mu Z^\mu, \\ Y^a &= e^a_\mu Y^\mu, \end{aligned} \quad (5.3)$$

accounting for the correct number of degrees of freedom. Thus, in form notation, (5.3) can be written as

$$\mu^2 \det^3(e^a_\mu) = \exp \left[-\mu^2 \int \left(\bar{Y}_a Y^a + \frac{1}{2} Z_a Z^a \right) \epsilon_{ab} e^a e^b \right]. \quad (5.4)$$

The same trick can be done for the other determinant in (5.2),

$$\mu^2 \det^{3/2}(\phi^a_b) = \mu^2 \det^4(\phi^a_b) \det^{-1/2}(\phi^a_b) = \int \mathcal{D}\bar{W}\mathcal{D}W\mathcal{K} \exp \left[-\mu^2 \int \left(\bar{W}_a \phi^a_b W^b + \tilde{K}_a \phi^a_b K^b \right) \epsilon_{cd} e^c e^d \right], \quad (5.5)$$

with \bar{W}^a and W^a being 0-form ghost fields while \tilde{K}_a and K^a are 0-forms bosonic fields. Moreover, $\tilde{K}_a = \frac{1}{2} \epsilon_{ab} K^b$. We notice that $\tilde{K}_a \phi^a_b K^b = 2\phi K_0 K^0$ with $\phi = \phi^0_1$. Therefore, defining $K_0 = \sigma/2$ and employing (4.15) again, (5.5) becomes

$$\begin{aligned} \mu^2 \det^{3/2}(\phi^a_b) &= \int \mathcal{D}\bar{W}\mathcal{D}W\mathcal{D}\sigma \exp \left[-\mu^2 \int \left(\epsilon_{ab} \bar{W}^a W^b \phi + \frac{1}{2} \sigma^2 \phi \right) \epsilon_{cd} e^c e^d \right], \\ &= \int \mathcal{D}\bar{W}\mathcal{D}W\mathcal{D}\sigma \exp \left[-\mu^2 \int \left(2\bar{W}_a W_b e^a e^b \phi + \frac{1}{2} \sigma^2 \epsilon_{cd} e^c e^d \phi \right) \right], \end{aligned} \quad (5.6)$$

Finally, the gravitational generating functional reads

$$Z_{grav} = \int [\mathcal{D}\Phi_{grav}] e^{-S_{qg}} , \quad (5.7)$$

with

$$S_{qg} = S_{grav} + \mu^2 \int \left(\bar{Y}_a Y^a + \frac{1}{2} Z_a Z^a + \frac{1}{2} \sigma^2 \phi \right) \epsilon_{ab} e^a e^b + 2\mu^2 \int \bar{W}_a W_b e^a e^b \phi , \quad (5.8)$$

and $[\mathcal{D}\Phi_{grav}]$ incorporates all extra fields coming from the determinants.

The extra terms coming from the determinants in (5.8) can also be cast in a BRST exact form. Calling the action coming from the determinants S_{aux} , it reads

$$S_{aux} = \mu^2 s \int \left[\left(\bar{Y}_a Z^a + \frac{1}{2} \bar{\sigma} \sigma \phi \right) \epsilon_{ab} e^a e^b + 2\bar{W}_a M_b e^a e^b \phi \right] , \quad (5.9)$$

with

$$\begin{aligned} s\bar{\sigma} &= \sigma , \\ s\sigma &= 0 , \\ s\bar{Y}^a &= Z^a , \\ sZ^a &= 0 , \\ s\bar{W}^a &= c^a_b \bar{W}^b , \\ sM^a &= W^a + c^a_b M^b , \\ sW^a &= c^a_b W^b . \end{aligned} \quad (5.10)$$

Thence, all physical content are contained in action (4.3), as it should be since we have not introduced any extra physical degree of freedom in the electromagnetic theory.

To end this section, we display the quantum numbers of the new fields in Table 3.

Fields	\bar{Y}^a	Y^a	Z^a	\bar{W}^a	W^a	M^a	$\bar{\sigma}$	σ
Dimension	0	0	0	0	0	0	0	0
Ghost number	-1	1	0	-1	1	0	-1	0
Form rank	0	0	0	0	0	0	0	0

Table 3: Quantum numbers of the localizing fields.

6 Conclusions

In this paper, we have formally developed a map from two-dimensional electrodynamics (action (2.4)) to two-dimensional gravity in the first order formalism (action (4.3)). To account for the different number of degrees of freedom in both theories, we introduced a set of auxiliary matter fields in such a way that the physical content in both theories are not affected - the auxiliary matter action is actually a BRST exact boundary term. Moreover, the respective mapped matter action is also a BRST exact boundary term. The map between theories is, in fact, not trivial. This non-triviality is evident in the computation of the Jacobian the map, which, at quantum level, requires extra localizing fields at the gravity side of the mapping. However, the action containing such fields is also BRST exact. Thus, the Jacobian actually does not affect the physical content of the theory.

We have established the basis to describe two-dimensional electrodynamics as a two-dimensional geometrodynamical theory. Nevertheless, much can be done in the future. For instance, the inclusion of non-trivial charged matter fields such as scalars and spinors would be the natural pathway to describe two-dimensional condensed matter systems⁴. Another approach to attack two-dimensional systems would be considering four-dimensional electrodynamics followed by a suitable dimensional reduction to two dimensions and then, to perform the map. Such approach would be more realistic for actual physical systems. Finally, we can consider the action (5.8) and fix the gauge by considering the full BRST symmetry encompassing the BRST transformations associated to $SO(2)$ and diffeomorphism symmetries [31–33]. The BRST approach can be studied in the pure two-dimensional analysis performed in the present work or in the two-dimensional theory obtained from the dimensional reduction of four-dimensional theories.

A Computation of the Jacobian

To compute the Jacobian of transformations (4.2) and (4.11) we need to construct the Jacobian matrix. For the map (4.2), the non-vanishing components are

$$\begin{aligned}\frac{\delta\theta}{\delta e^a} &= 2\mu\epsilon_{ab}e^b, \\ \frac{\delta A}{\delta\omega^{ab}} &= \frac{1}{2\mu}\epsilon_{ab}.\end{aligned}\tag{A.1}$$

For the map (4.11), the only non-vanishing components are

$$\begin{aligned}\frac{\delta\bar{\Phi}^a}{\delta\sigma^b} &= \delta_b^a, \\ \frac{\delta\Phi^a}{\delta e^a} &= \mu\phi^a{}_b, \\ \frac{\delta\Phi^a}{\delta\phi^b{}_c} &= \delta_b^a e^c.\end{aligned}\tag{A.2}$$

For the map (4.12), the non-vanishing components are

$$\begin{aligned}\frac{\delta\bar{w}}{\delta\bar{\eta}} &= 1, \\ \frac{\delta\bar{z}}{\delta\bar{\chi}} &= 1, \\ \frac{\delta w}{\delta e^a} &= \mu\eta_a, \\ \frac{\delta w}{\delta\eta_a} &= \mu e^a, \\ \frac{\delta z}{\delta e^a} &= -\mu\chi_a, \\ \frac{\delta z}{\delta\chi_a} &= \mu e^a.\end{aligned}\tag{A.3}$$

⁴External matter was not considered here. However, its inclusion would be immediate since the coupling between the electromagnetic potential and an external current is given by jA with j being a 1-form describing the charged source. Clearly, it leads to a gravity coupling of the form $j_{ab}\omega^{ab}$.

The components in (A.1), (A.2) and (A.3), together with all vanishing components, form a 15×15 matrix, namely J_0 . This matrix can be immediately reduced to a 9×9 matrix, named J_1 , due to the trivial relations in (A.1), (A.2) and (A.3). Thus, one can readily show that

$$\det J_0 = \frac{1}{\mu^2} \det J_1, \quad (\text{A.4})$$

where J_1 can be arranged as

$$J_1 = \begin{bmatrix} 0 & 0 & \mu\phi & 0 & \mu\eta_0 & 0 & \mu\chi_0 & 0 & \mu e_1^1 \\ 0 & 0 & 0 & \mu\phi & 0 & \mu\eta_0 & 0 & \mu\chi_0 & -\mu e_0^1 \\ \mu\phi & 0 & 0 & 0 & \mu\eta_1 & 0 & \mu\chi_1 & 0 & -\mu e_1^0 \\ 0 & \mu\phi & 0 & 0 & 0 & \mu\eta_1 & 0 & \mu\chi_1 & \mu e_0^0 \\ \mu e_0^1 & \mu e_1^1 & \mu e_0^0 & \mu e_1^0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \mu e_0^0 & \mu e_1^0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \mu e_0^1 & \mu e_1^1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \mu e_0^0 & \mu e_1^0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \mu e_0^1 & \mu e_1^1 & 0 \end{bmatrix} \quad (\text{A.5})$$

where $\phi = \phi_1^0$. The determinant of J_1 can be systematically computed within Laplace's method. After a long but straightforward computation, the result is given by

$$\det J_1 = \mu^9 \det(\phi^a_b)^{3/2} e^3. \quad (\text{A.6})$$

Thus,

$$\det J_0 = 4\mu^7 \det(\phi^a_b)^{3/2} e^3, \quad (\text{A.7})$$

which, up to a normalization factor, agrees with (4.13).

Acknowledgements

This study was financed in part by The Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

References

- [1] A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, and A. K. Geim, "The electronic properties of graphene". *Rev. Mod. Phys.* **81** (2009) 109–162.
- [2] T. Farajollahpour, Z. Faraei, and S. A. Jafari, "Solid-state platform for space-time engineering: The 8Pmmn borophene sheet". *Phys. Rev. B* **99** no. 23, (2019) 235150.
- [3] S. Cahangirov, M. Topsakal, E. Akturk, H. Sahin, and C. S., "Two- and One-Dimensional Honeycomb Structures of Silicon and Germanium". *Phys. Rev. Lett.* **102** (2009) 236804.
- [4] I. Bialynicki-Birula, "Classical electrodynamics in two dimensions - exact solution". *Phys. Rev. D* **3** (1971) 864–866.

- [5] D. J. Gross and A. A. Migdal, “Nonperturbative Two-Dimensional Quantum Gravity”. *Phys. Rev. Lett.* **64** (1990) 127.
- [6] E. Witten, “On the Structure of the Topological Phase of Two-dimensional Gravity”. *Nucl. Phys. B* **340** (1990) 281–332.
- [7] A. M. Polyakov, “Quantum Gravity in Two-Dimensions”. *Mod. Phys. Lett. A* **2** (1987) 893.
- [8] M. Fukuma, H. Kawai, and R. Nakayama, “Continuum Schwinger-dyson Equations and Universal Structures in Two-dimensional Quantum Gravity”. *Int. J. Mod. Phys. A* **6** (1991) 1385–1406.
- [9] A. H. Chamseddine, “Two-dimensional quantum gravity. A mini review”. *Nucl. Phys. B Proc. Suppl.* **16** (1990) 579–580.
- [10] Y. Kazama and H. Nicolai, “On the exact operator formalism of two-dimensional Liouville quantum gravity in Minkowski space-time: A review”. in *4th Hellenic School on Elementary Particle Physics*, pp. 753–789. 9, 1992.
- [11] A. Achucarro and P. K. Townsend, “A Chern-Simons Action for Three-Dimensional anti-De Sitter Supergravity Theories”. *Phys. Lett.* **B180** (1986) 89.
- [12] E. Witten, “(2+1)-Dimensional Gravity as an Exactly Soluble System”. *Nucl. Phys.* **B311** (1988) 46.
- [13] Yu. N. Obukhov, “Gauge fields and space-time geometry”. *Theor. Math. Phys.* **117** (1998) 1308–1318. [Teor. Mat. Fiz.117,249(1998)].
- [14] R. F. Sobreiro and V. J. Vasquez Otoyá, “Effective gravity from a quantum gauge theory in Euclidean space-time”. *Class. Quant. Grav.* **24** (2007) 4937–4953.
- [15] R. F. Sobreiro, A. A. Tomaz, and V. J. V. Otoyá, “de Sitter gauge theories and induced gravities”. *Eur. Phys. J. C* **72** (2012) 1991.
- [16] T. S. Assimos, A. D. Pereira, T. R. S. Santos, R. F. Sobreiro, A. A. Tomaz, and V. J. Vasquez Otoyá, “From $SL(5, \mathbb{R})$ Yang-Mills theory to induced gravity”. *Int. J. Mod. Phys. D* **26** no. 08, (2017) 1750087.
- [17] T. S. Assimos and R. F. Sobreiro, “Constrained gauge-gravity duality in three and four dimensions”. *Eur. Phys. J. C* **80** no. 1, (2020) 20.
- [18] T. S. Assimos and R. F. Sobreiro, “Topological gauge-gravity equivalence: fiber bundle and homology aspects”.
- [19] O. Piguet and S. P. Sorella, “Algebraic renormalization: Perturbative renormalization, symmetries and anomalies”. *Lect. Notes Phys. Monogr.* **28** (1995) 1–134.
- [20] R. Utiyama, “Invariant theoretical interpretation of interaction”. *Phys. Rev.* **101** (1956) 1597–1607.
- [21] T. W. B. Kibble, “Lorentz invariance and the gravitational field”. *J. Math. Phys.* **2** (1961) 212–221.
- [22] D. W. Sciama, “The Physical structure of general relativity”. *Rev. Mod. Phys.* **36** (1964) 463–469. [Erratum: Rev.Mod.Phys. 36, 1103–1103 (1964)].
- [23] D. Lovelock, “The Einstein tensor and its generalizations”. *J. Math. Phys.* **12** (1971) 498–501.

- [24] A. Mardones and J. Zanelli, “Lovelock-Cartan theory of gravity”. *Class. Quant. Grav.* **8** (1991) 1545–1558.
- [25] J. Zanelli, “Lecture notes on Chern-Simons (super-)gravities. Second edition (February 2008)”. in *Proceedings, 7th Mexican Workshop on Particles and Fields (MWPF 1999): Merida, Mexico, November 10-17, 1999*. 2005.
- [26] J. Zanelli, “Chern-Simons Forms in Gravitation Theories”. *Class. Quant. Grav.* **29** (2012) 133001.
- [27] M. Nakahara, *Geometry, topology and physics*. Boca Raton, USA: Taylor & Francis (2003) 573 p., 2003.
- [28] R. M. Wald, *General Relativity*. Chicago Univ. Pr., Chicago, USA, 1984.
- [29] V. De Sabbata and M. Gasperini, *INTRODUCTION TO GRAVITY*. Singapore, Singapore: World Scientific 346p, 1985.
- [30] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*. W. H. Freeman, San Francisco, 1973.
- [31] O. Moritsch, M. Schweda, and S. P. Sorella, “Algebraic structure of gravity with torsion”. *Class. Quant. Grav.* **11** (1994) 1225–1242.
- [32] L. Baulieu, “Anomalies and Gauge Symmetry”. *Nucl. Phys. B* **241** (1984) 557–588.
- [33] L. Baulieu and J. Thierry-Mieg, “Algebraic Structure of Quantum Gravity and the Classification of the Gravitational Anomalies”. *Phys. Lett. B* **145** (1984) 53–60.