

# THE LATTICE OF NIL-HECKE ALGEBRAS OVER A COXETER GROUP

SUTANAY BHATTACHARYA AND APOORVA KHARE

ABSTRACT. Associated to every Coxeter group  $W$ , we construct a lattice of quotients of its braid group-algebra, which we term nil-Hecke algebras, and which are obtained by killing all braid-words that are “sufficiently long”, as well as some integer power of each generator. These include usual nil-Coxeter algebras, nil-Temperley–Lieb algebras, and their variants, and lead to symmetric semigroup module categories which necessarily cannot be monoidal.

Motivated by classical work of Coxeter (1957), and continuing beyond work of the second author [*Trans. Amer. Math. Soc.* 2018], we obtain a complete classification of the finite-dimensional nil-Hecke algebras for all Coxeter groups  $W$ . These include the usual nil-Coxeter algebras for  $W$  of finite type, their “fully commutative” analogues for  $W$  of FC-finite type, three exceptional algebras (of types  $F_4, H_3, H_4$ ), and three exceptional series (of types  $B_n$  and  $A_n$ ). This classification draws from and brings together results of Stembridge [*J. Alg. Combin.* 1996, 1998], Postnikov via Gowravaram–Khovanova (2015), Hart [*J. Group Th.* 2017], and Khare [*Trans. Amer. Math. Soc.* 2018].

## 1. INTRODUCTION AND MAIN RESULTS

Throughout this paper,  $\mathbb{k}$  will denote a fixed unital commutative ground ring, and by  $\dim M$  we mean the  $\mathbb{k}$ -rank of a free  $\mathbb{k}$ -module  $M$ .

In this paper, we study a class of graded rings over a given Coxeter group  $W$  (finite or infinite). We term such rings *nil-Hecke algebras*. Most of the rings that we study turn out to be the associated graded rings for certain quotients of the corresponding (generic) Iwahori–Hecke algebras – which include, for instance, Temperley–Lieb algebras. Thus, the quotients we study include the corresponding nil-Coxeter as well as nil-Temperley–Lieb algebras.

Our focus in the present work is on exploring the finite-dimensionality of these graded rings. In this, we are motivated by connections to Coxeter group combinatorics, semigroup tensor categories, and flatness of deformations (which in the complex case is related to the Broué–Malle–Rouquier (BMR) freeness conjecture). We elaborate on these connections below.

To introduce nil-Hecke algebras, recall that any Coxeter group  $W$  is accompanied by:

- an associated braid group  $\mathcal{B}_W \twoheadrightarrow W$ ;
- a finite index set  $I$  as well as a set  $\mathbf{S} := \{\mathbf{s}_i : i \in I\}$  that generates  $\mathcal{B}_W$  and yields the simple reflections under the map  $\mathcal{B}_W \twoheadrightarrow W$ ; and
- a finite index set  $J$  (in fact,  $J \subset \binom{I}{2}$ ) as well as relations  $\mathcal{R} := \{\mathbf{v}_j - \mathbf{w}_j : j \in J\}$ , where for each  $j$ , both sides denote words in  $\mathbf{S}$  of equal (and finite) length,

such that  $\langle \mathbf{S} | \mathcal{R} \rangle$  is a presentation of  $\mathcal{B}_W$ . (This presentation was extended by Brieskorn in [9].)

We now define our main object of interest: a lattice of algebras corresponding to  $W$ .

---

*Date:* December 7, 2021.

*2010 Mathematics Subject Classification.* 20F55 (Primary), 20F05, 20C08 (Secondary).

*Key words and phrases.* Generalized Coxeter group, generalized nil-Coxeter algebra, nil-Hecke algebra, Temperley–Lieb algebra.

**Definition 1.1.** Fix a Coxeter group  $W$ , with associated data  $I, J, \mathbf{S}, \mathcal{R}$  (so  $J \subset \binom{I}{2}$ ).

- (1) Given  $J_0 \subset J$  and an integer tuple  $\mathbf{d} = (d_i)_{i \in I}$  with  $1 \leq d_i \leq \infty \forall i$ , define the corresponding *nil-Hecke algebra*  $n\mathcal{H}(W, \mathbf{d}, J_0)$  to be the  $\mathbb{k}$ -algebra generated by  $\mathbf{S}$  with relations:

$$\mathbf{s}_i^{d_i} = 0 \forall i, \quad \mathbf{v}_j = \mathbf{w}_j \forall j \in J_0, \quad \mathbf{v}_j = 0 = \mathbf{w}_j \forall j \in J \setminus J_0.$$

- (2) For  $k \geq 1$ , define  $J_{<k} := \{j \in J : \ell(\mathbf{v}_j) = \ell(\mathbf{w}_j) < k\}$ . Notice that  $J_{<1} = J_{<2} = \emptyset$ , so that  $n\mathcal{H}(W, \mathbf{d}, J_{<1}) = n\mathcal{H}(W, \mathbf{d}, J_{<2})$ . Also define  $J_{<\infty} := J$ .

**Remark 1.2.** Here we omit the relation  $\mathbf{s}_i^{d_i} = 0$  if  $d_i = \infty$ . Also note that all the relations in  $\mathcal{R}$  still hold in  $n\mathcal{H}(W, \mathbf{d}, J_0)$ , but the emphasis is on the additional relations  $\mathbf{v}_j = 0 = \mathbf{w}_j$ ,  $j \in J \setminus J_0$ .

In other words,  $n\mathcal{H}(W, \mathbf{d}, J_0)$  is a quotient of the Artin monoid-algebra of  $W$  by the order relations  $\mathbf{s}_i^{d_i} = 0$  (see [37]). Two special cases of this construction are:

- ( $k = \infty$ .) The *generalized nil-Coxeter algebra* associated to the above data is  $NC_W(\mathbf{d}) := n\mathcal{H}(W, \mathbf{d}, J_{<\infty})$ . These algebras were defined and studied in previous work [27, 28] (see also the extended abstract [29]). In the special case where  $d_i = 2 \forall i$ ,  $NC_W((2, \dots, 2))$  is the “usual” nil-Coxeter algebra associated to  $W$ , with  $J = \binom{I}{2}$ . These algebras are well-studied in connection to flag varieties [7, 34], categorification [30, 31], symmetric function theory [6], and Schubert polynomials [19, 35].
- ( $k = 3$ .) We define the *generalized nil-Temperley–Lieb algebra* associated to the above data to be  $NTL_W(\mathbf{d}) := n\mathcal{H}(W, \mathbf{d}, J_{<3})$ . In the special case where  $d_i = 2 \forall i$ ,  $NTL_W((2, \dots, 2))$  is the “usual” nil-Temperley–Lieb algebra associated to  $W$ . These latter are quotients of nil-Coxeter algebras as well as associated graded algebras of Temperley–Lieb algebras (see e.g. [5, 18]), and have connections to fermionic particle configurations and to quantum Schubert calculus [33, 38]. They were defined and studied in [16, 17, 21]. In the special case of the Coxeter (multi)graph of  $W$  being simply laced, the algebras  $NTL_W((2, \dots, 2))$  were rediscovered by Postnikov as *XYX algebras* [20].

For a given group  $W$  with presentation as above, the set of nil-Hecke  $\mathbb{k}$ -algebras forms a lattice isomorphic to the product of the power set  $2^J$  with  $I$  copies of  $\mathbb{Z}^{\geq 1} \sqcup \{\infty\}$ . Two algebras in this lattice are comparable if and only if one surjects onto the other; the extremal points in the lattice are the source  $\mathcal{B}_W$  (for  $J_0 = J$  and  $d_i = \infty \forall i$ ) and the rank-1 algebra  $\mathbb{k}$  (whenever  $d_i = 1 \forall i$ ). For each  $\mathbf{d} \in (\mathbb{Z}^{\geq 1})^I$ , the corresponding algebras form a sub-lattice with unique sink  $n\mathcal{H}(W, \mathbf{d}, \emptyset)$  and source the generalized nil-Coxeter algebra  $NC_W(\mathbf{d})$ .

We now come to the results in this paper, which classify all nil-Hecke  $\mathbb{k}$ -algebras  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  for  $1 \leq k < \infty$  that are finite-dimensional (for  $\mathbb{k}$  a field, or more precisely, those of finite  $\mathbb{k}$ -rank) of the form  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  with  $1 \leq k \leq \infty$ . In this, we are motivated by classical work of Coxeter [11], in which he considered quotients of the type  $A$  braid group by the relations  $s_i^p = 1 \forall i \geq 1$ , and classified the pairs  $(n, p)$  for which the resulting “generalized Coxeter group” is finite (see also [2]). Our work studies the “Iwahori–Hecke” analogue of Coxeter’s problem.

At the same time, our work also complements (by proving results for all finite  $k$ ) the work of the second author [28], who studied the  $k = \infty$  case. Furthermore, we unify under one umbrella several classical and recent works, including ([28], as well as) results by Stembridge [44, 45, 46], Gowravaram–Khovanova (who attribute their construction to Postnikov) [20], and Hart [23], and show how they fit into this picture.

We begin with the recent work [28], which classified all finite-dimensional objects among the generalized nil-Coxeter algebras (i.e., corresponding to  $k = \infty$ ). Interestingly, this yields a “non-usual” type  $A$  family of such algebras – and it turns out to be the only one:

**Theorem 1.3** ([28]). *Fix a Coxeter group  $W$  with related data  $I, J, \mathbf{S}, \mathcal{R}$ , and integers  $d_i \geq 2 \forall i$ . The corresponding nil-Hecke (or generalized nil-Coxeter) algebra  $n\mathcal{H}(W, \mathbf{d}, J_{<\infty})$  is a finitely generated  $\mathbb{k}$ -module if and only if exactly one of the following occurs:*

- (1)  $W$  is a finite Coxeter group and  $\mathbf{d} = (2, \dots, 2)$  (the “usual” nil-Coxeter algebras); or
- (2)  $W$  is of type  $A$ , and  $\mathbf{d} = (d, 2, \dots, 2)$  or  $(2, \dots, 2, d)$  for some  $d > 2$ . We denote this nil-Hecke algebra by  $NC_A(n, d) := n\mathcal{H}(S_{n+1}, (d, 2, \dots, 2), J_{<\infty})$ .

**Remark 1.4.** Note that setting any  $d_i = 1$  yields degeneracies in the original Definition 1.1, since the generator  $\mathbf{s}_i$  is rendered superfluous. Moreover, if  $d_i = \infty$  then  $n\mathcal{H}(W, \mathbf{d}, J_0)$  contains the  $\mathbb{k}$ -span of  $\mathbf{s}_i^n$  for  $n \geq 0$ . As our goal in this paper is to classify the finite-rank nil-Hecke algebras, throughout this paper we assume  $2 \leq d_i < \infty \forall i \in I$ .

Following Theorem 1.3 from [28], the present work obtains the corresponding classification results for  $J_{<k}$  with  $k < \infty$ . Our first result is for  $k = 1, 2$ :

**Theorem A** ( $k = 1, 2$ ). *Fix a Coxeter group  $W$  with related data  $I, J, \mathbf{S}, \mathcal{R}$ , and integers  $d_i \geq 2 \forall i$ . The corresponding nil-Hecke  $\mathbb{k}$ -algebra  $n\mathcal{H}(W, \mathbf{d}, J_{<1}) = n\mathcal{H}(W, \mathbf{d}, J_{<2})$  is a finitely generated  $\mathbb{k}$ -module if and only if exactly one of the following occurs:*

- (1) The Coxeter (multi)graph of  $W$  is a tree with exactly one multiple edge  $4 \leq m_{i_0 j_0} < \infty$ , and  $d_i = 2 \forall i$ . In this case, if disconnecting  $i_0, j_0$  yields trees  $T, T'$  with  $a, b$  nodes respectively, then

$$\dim n\mathcal{H}(W, \mathbf{d}, J_{<2}) = \begin{cases} \frac{1}{2}m_{i_0 j_0}|I|^2 + 1 - 2ab, & \text{if } m_{i_0 j_0} \text{ is even,} \\ \frac{1}{2}(m_{i_0 j_0} - 1)|I|^2 + 1, & \text{if } m_{i_0 j_0} \text{ is odd.} \end{cases}$$

- (2) The Coxeter (multi)graph of  $W$  is a simply laced tree, and at most one node  $i_0 \in I$  satisfies:  $d_{i_0} \geq 3$ . In this case,

$$\dim n\mathcal{H}(W, \mathbf{d}, J_{<2}) = 1 + |I|^2(d_{i_0} - 1).$$

**Remark 1.5.** Here and below, by the statement “ $\dim n\mathcal{H}(W, \mathbf{d}, J_0) = m$ ” for an integer  $m > 0$ , we mean that  $n\mathcal{H}(W, \mathbf{d}, J_0)$  is a free  $\mathbb{k}$ -module of rank  $m$ .

Thus for  $k = 1, 2$ , there are infinitely many finite-dimensional nil-Hecke algebras corresponding to “non-Coxeter” matrices (i.e., where  $d_{i_0} > 2$ ). In contrast, our next result shows that for  $k \geq 3$ , this already gets reduced to precisely one non-Coxeter family, which for  $k = \infty$  corresponds to  $NC_A(n, d)$  in Theorem 1.3(2).

**Theorem B** ( $k = 3$ ). *Fix a Coxeter group  $W$  with related data  $I, J, \mathbf{S}, \mathcal{R}$ , and integers  $d_i \geq 2 \forall i$ . The corresponding nil-Hecke (or generalized nil-Temperley–Lieb)  $\mathbb{k}$ -algebra  $n\mathcal{H}(W, \mathbf{d}, J_{<3}) = NTL_W(\mathbf{d})$  is a finitely generated  $\mathbb{k}$ -module if and only if exactly one of the following occurs:*

- (1)  $W$  is a finite Coxeter group and  $\mathbf{d} = (2, \dots, 2)$  (these are “usual” nil-Temperley–Lieb algebras);
- (2)  $W$  is a Coxeter group of type  $E_n (n \geq 9)$ ,  $F_n (n \geq 5)$ , or  $H_n (n \geq 5)$  (see Figure 1.1), and  $\mathbf{d} = (2, \dots, 2)$  (these are also “usual” nil-Temperley–Lieb algebras); or
- (3)  $W$  is of type  $A$ , and  $\mathbf{d} = (d, 2, \dots, 2)$  or  $(2, \dots, 2, d)$  for some  $d > 2$ .

In the first two cases,  $\dim NTL_W(\mathbf{d})$  is precisely the number of fully commutative elements in the corresponding Coxeter group  $W$ , i.e., the number of words  $w \in W$  for which any reduced

word can be obtained from any other without using the “non-commutative” braid relations ( $m_{ij} \geq 3$ ). In the last case of  $W = W(A_n)$  and  $d_1 > 2 = d_2 = \dots = d_n$ , we have:

$$\dim NTL_W(\mathbf{d}) = (d-1)C_{n+1} - (d-2)C_n + (d-2) \sum_{j=1}^{n-1} jC_{n-j} \quad (1.6)$$

where  $C_n$  is the  $n^{\text{th}}$  Catalan number.




FIGURE 1.1. Dynkin graphs with  $|W|$  infinite but  $\dim(NTL_W)$  finite

**Remark 1.7.** Following Postnikov’s calling  $NTL_{W(A_n)}((2, \dots, 2))$  an *XYX-algebra* (see [20]), we term  $NTL_{W(A_n)}((d, 2, \dots, 2)) = n\mathcal{H}(W(A_n), (d, 2, \dots, 2), J_{<3})$  a *generalized XYX-algebra*. Its  $\mathbb{k}$ -rank (1.6) involving Catalan numbers, generalizes the  $d = 2$  fact that the type  $A$  XYX-algebra has rank  $C_{n+1}$  – a fact first observed in [3]. We also remark that the final sum in (1.6) is the partial sum of the partial sum of the Catalan numbers, and its first few terms can be found in [43].

In particular, Theorem B together with [28, Theorem B] implies Theorem 1.3 for all generalized Coxeter matrices. This because all Coxeter group algebras  $\mathbb{k}W$  are flat deformations of the corresponding nil-Coxeter algebras  $NC_W((2, \dots, 2))$ .

As the algebras  $NTL_W(\mathbf{d})$  are not our primary focus in the present paper, we will not go beyond the proof of Theorem B; however, we remark that the proof of Theorem B is constructive, and reveals the analogue of fully commutative elements in the “non-usual” case of the generalized XYX-algebras  $NTL_{A_n}((2, \dots, 2, d))$ ,  $d > 2$ . Recall that such elements were introduced and studied by Stembridge in [44, 45], and enumerated in his sequel [46]. In [44], Stembridge also computed the Coxeter groups which contain only finitely many fully commutative elements (see also [15, 21]): these are the finite Coxeter groups as well as the families  $E_n, F_n, H_n$  as above. Theorem B now constructs and enumerates such elements in the “non-usual” generalized nil-Coxeter algebras  $NC_A(n, d)$  (see Theorem 1.3(2)) introduced and studied in [28], adding to our knowledge of them.

Theorem B classified the finite  $\mathbb{k}$ -rank nil-Hecke algebras for  $k = 3$ . Next, we look at the case  $k = 4$ , followed by an immediate corollary for  $k = 5$ .

**Theorem C** ( $k = 4$ ). *Fix a Coxeter group  $W$  with related data  $I, J, \mathbf{S}, \mathcal{R}$ , and integers  $d_i \geq 2 \forall i$ . The corresponding nil-Hecke  $\mathbb{k}$ -algebra  $n\mathcal{H}(W, \mathbf{d}, J_{<4})$  is a finitely generated  $\mathbb{k}$ -module if and only if exactly one of the following occurs:*

- (1)  $W$  is a finite Coxeter group with  $\mathbf{d} = (2, \dots, 2)$ ; or
- (2)  $W$  is of type  $A$ , and  $\mathbf{d} = (d, 2, \dots, 2)$  or  $(2, \dots, 2, d)$  for some  $d > 2$ .

In the first case,  $\dim n\mathcal{H}(W, \mathbf{d}, J_{<4})$  is given by the following table:

$W$	$\dim n\mathcal{H}(W, \mathbf{d}, J_{<4})$
$A_n, D_n, E_6, E_7, E_8$	$ W $
$B_n$	$\sum_{k=0}^n \binom{n}{k}^2 k!$
$F_4$	304
$H_3$	76
$H_4$	1460
$I_2(m)$	$\begin{cases} 2m & \text{if } m < 4 \\ 2m - 1 & \text{if } m \geq 4 \end{cases}$

In the second case, we have

$$\dim n\mathcal{H}(W, \mathbf{d}, J_{<4}) = n!(1 + n(d - 1)).$$

**Corollary 1.8** ( $k = 5$ ). Fix a Coxeter group  $W$  with related data  $I, J, \mathbf{S}, \mathcal{R}$ , and integers  $d_i \geq 2 \forall i$ . The corresponding nil-Hecke  $\mathbb{k}$ -algebra  $n\mathcal{H}(W, \mathbf{d}, J_{<5})$  is a finitely generated  $\mathbb{k}$ -module if and only if exactly one of the following occurs:

- (1)  $W$  is a finite Coxeter group with  $\mathbf{d} = (2, \dots, 2)$ ; or
- (2)  $W$  is of type  $A$ , and  $\mathbf{d} = (d, 2, \dots, 2)$  or  $(2, \dots, 2, d)$  for some  $d > 2$ .

In the first case,  $\dim n\mathcal{H}(W, \mathbf{d}, J_{<5})$  is given by the following table:

$W$	$\dim n\mathcal{H}(W, \mathbf{d}, J_{<5})$
$A_n, B_n, D_n, E_6, E_7, E_8, F_4$	$ W $
$H_3$	76
$H_4$	1460
$I_2(m)$	$\begin{cases} 2m & \text{if } m < 5 \\ 2m - 1 & \text{if } m \geq 5 \end{cases}$

In the second case, we have

$$\dim n\mathcal{H}(W, \mathbf{d}, J_{<5}) = n!(1 + n(d - 1)).$$

**Remark 1.9.** In particular, this work provides some refined combinatorial information that (to the best of our knowledge) was not written down for “usual” nil-Coxeter algebras – nor for their “non-usual” (aka generalized, i.e. with  $\mathbf{d} \neq (2, \dots, 2)$ ) counterparts in type  $A$ .

- For “usual” nil-Coxeter algebras, we prove the above results by computing not just the dimension (or  $\mathbb{k}$ -rank), but by isolating – via Theorem 2.3 – a distinguished subset of words  $S'_k$  inside the “usual” monomial basis, which provides a  $\mathbb{k}$ -basis. For  $k = 1, 2$ , the set  $S'_k$  is simply the full basis (of size  $|W|$ ), and for  $k = 3$  it corresponds to the subset  $W_{\text{fc}}$  of fully commutative elements. However, for  $k = 4, 5$ ,  $S'_k$  was not isolated earlier – even for types  $B_n, F_4, H_3, H_4$  if we restrict to  $S'_k$  finite.

There is also combinatorial information: in all of these cases, the words isolated satisfy the defining condition of avoiding braid relations that are “long enough”. But we show more: e.g. for type  $B_n$ , the words in  $S'_4$  turn out to be precisely the “ $\overline{12}$ -avoiding signed permutations”.

- Moving from combinatorics to algebra: in addition to our identifying a subset of words with some combinatorial properties – our work moreover isolates a multiplication structure on their span. This is akin to the set of fully commutative words inside  $W$  – while their closure under multiplication in  $W$  is unclear, this indeed happens

when looking at their images inside the corresponding nil-Temperley–Lieb algebras. Similarly, the construction of “usual” nil-Hecke algebras in this work reveals a multiplication structure on the  $\mathbb{k}$ -span of the set  $S'_k$  for  $k = 4, 5$ , which is not revealed when considering  $S'_k$  as a subset of the Coxeter group  $W$ . The algebras thus revealed have additional features from the viewpoint of category theory – see Section 2.

- In parallel, in the “non-usual” case, the work [28] that introduced the algebras  $NC_A(n, d)$  (see Theorem 1.3(2)) could not detect the analogue of the fully commutative elements, which we now uncover (via Theorem 2.4) in the proof of Theorem B.

Returning to the classification of finite-rank nil-Hecke algebras, our final result is for  $k \geq 6$ .

**Corollary 1.10** ( $6 \leq k \leq \infty$ ). *Fix a Coxeter group  $W$  with related data  $I, J, \mathbf{S}, \mathcal{R}$ , integers  $d_i \geq 2 \forall i$  and  $6 \leq k \leq \infty$ . The corresponding nil-Hecke  $\mathbb{k}$ -algebra  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  is a finitely generated  $\mathbb{k}$ -module if and only if  $W$  is either a finite Coxeter group, with  $\mathbf{d} = (2, \dots, 2)$  or  $W$  is of type  $A$  with  $\mathbf{d} = (d, 2, \dots, 2)$  or  $(2, \dots, 2, d)$  for some  $d > 2$ . In the first case, the corresponding nil-Hecke algebra has  $\mathbb{k}$ -rank equal to  $|W|$  unless it is of the type  $I_2(m)$ , in which case it is given by*

$$\dim n\mathcal{H}(W, \mathbf{d}, J_{<k}) = \begin{cases} 2m & \text{if } m < k, \\ 2m - 1 & \text{if } m \geq k. \end{cases}$$

In the second case, the  $\mathbb{k}$ -rank is  $n!(1 + n(d - 1))$ .

For ease of reference, we tabulate the above results for  $3 \leq k \leq \infty$  in Table 1, and follow it up with a few comments.

$W$	$J_{<3}$	$J_{<4}$	$J_{<5}$	$J_{<k} (6 \leq k \leq \infty)$
$A_n, D_n, E_6, E_7, E_8$	$ W_{\text{fc}} $	$ W $	$ W $	$ W $
$B_n$	$ W_{\text{fc}} $	$\sum_{k=0}^n \binom{n}{k}^2 k!$	$ W $	$ W $
$F_4$	$ W_{\text{fc}} $	304	$ W $	$ W $
$H_3$	$ W_{\text{fc}} $	76	76	$ W $
$H_4$	$ W_{\text{fc}} $	1460	1460	$ W $
$I_2(m)$	$ W_{\text{fc}} $	$ W $ if $m < 4$ , else $ W  - 1$	$ W $ if $m < 5$ , else $ W  - 1$	$ W $ if $m < k$ else $ W  - 1$
$E_n (n \geq 9),$ $F_n (n \geq 5),$ $H_n (n \geq 5)$	$ W_{\text{fc}} $			
$A_n, \mathbf{d} = (d, 2, \dots, 2),$ $d > 2$	(see (1.6))	$n!(1 + n(d - 1))$	$n!(1 + n(d - 1))$	$n!(1 + n(d - 1))$

TABLE 1. Classification of Coxeter groups with finite-dimensional nil-Hecke algebras  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  for  $3 \leq k \leq \infty$ . Except for the last row,  $\mathbf{d} = (2, \dots, 2)$ .  $W_{\text{fc}}$  denotes the fully commutative elements in  $W$ .

**Remark 1.11.** The fully commutative words in all Coxeter groups were studied and enumerated by Stembridge in [46]. In particular, for  $W$  dihedral (i.e., of type  $I_2(m)$ ),  $W_{\text{fc}} = W \setminus \{w_\circ\}$  comprises all non-longest elements. Thus in Table 1, the  $\mathbb{k}$ -rank of  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  over  $W = W(I_2(m))$  with  $m \geq 3$  is  $|W|$  if  $m < k$ , and  $|W| - 1 = |W_{\text{fc}}|$  otherwise.

This shows that the finite rank nil-Hecke algebras have ranks either (a) equal to  $|W|$  or  $|W_{\text{fc}}|$ , or (b) belonging to the three exceptional cases  $F_4, H_3, H_4$  with  $k = 4$ , or (c) belonging to the three exceptional families:  $B_n$  with  $k = 4$ , or  $A_n$  with  $k = 3, 4$  and  $\mathbf{d} = (2, \dots, 2, d)$  for some  $d \geq 2$ .

Finally, note that as  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  is a quotient of  $n\mathcal{H}(W, \mathbf{d}, J_{<l})$  for  $1 \leq k \leq l \leq \infty$ , the list of finite rank examples for any value of  $k$  – i.e., in any non-initial column in Table 1 – is contained in the rows occupied in the previous column. (This is useful in the proofs below.) In particular, the result in [28] that there is only one family of finite rank generalized nil-Coxeter algebras with  $\mathbf{d} \neq (2, \dots, 2)$ , already follows from the  $J_{<4}$  column in Table 1.

Our final result provides an equivalent condition to the finite-dimensionality of nil-Hecke algebras, motivated by considerations of deformation theory (see the end of Section 2):

**Theorem D.** *Fix a Coxeter group  $W$  with related data  $I, J, \mathbf{S}, \mathcal{R}$ , and integers  $d_i \geq 2 \forall i$ , and let  $k$  be an integer such that  $1 \leq k \leq \infty$ . Then the following are equivalent:*

- (1) *The corresponding nil-Hecke  $\mathbb{k}$ -algebra  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  is a finitely generated  $\mathbb{k}$ -module.*
- (2) *The nil-Hecke algebra is found in the lists in Theorems A, B, C and Corollaries 1.8, 1.10 – or equivalently, in Theorem A and Table 1.*
- (3) *The two-sided augmentation ideal  $\mathfrak{m} \subset n\mathcal{H}(W, \mathbf{d}, J_{<k})$ , which is generated by  $\{\mathbf{s}_i : i \in I\}$ , is nilpotent.*

*If  $\mathbb{k}$  is moreover a field, then the finite-dimensional  $\mathbb{k}$ -algebra  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  is local, with unique maximal ideal  $\mathfrak{m}$ .*

Note that Theorem D generalizes the assertion [28, Theorem D(3)], which is simply the  $k = \infty$  special case.

**Organization.** This paper is organized as follows. We begin by providing several motivations and connections to prior works in the following section, as well as by producing a monomial word  $\mathbb{k}$ -basis for  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  for general  $W$ ,  $\mathbf{d} \in (\mathbb{Z}^{\geq 2})^I$ , and  $1 \leq k \leq \infty$ . The remaining sections are devoted to proving the main theorems above.

**Acknowledgments.** The second author was partially supported by Ramanujan Fellowship grant SB/S2/RJN-121/2017, MATRICS grant MTR/2017/000295, and SwarnaJayanti Fellowship grants SB/SJF/2019-20/14 and DST/SJF/MS/2019/3 from SERB and DST (Govt. of India).

## 2. FLAT DEFORMATIONS, OTHER CONNECTIONS, AND A WORD BASIS

In this section we explain how the problem under consideration, and more broadly, nil-Hecke algebras, are connected to several other areas. We begin with a categorical connection: even before considering finite-dimensionality, the entire lattice of nil-Hecke algebras  $\{n\mathcal{H}(W, \mathbf{d}, J_0) : \mathbf{d} \in (\mathbb{Z}^{\geq 2})^I, J_0 \subset J\}$  yields examples of co-commutative algebras with coproduct, which are not bialgebras.

**Proposition 2.1.** *Let  $A := n\mathcal{H}(W, \mathbf{d}, J_0)$  for some  $\mathbf{d} \in (\mathbb{Z}^{\geq 2})^I$  and  $J_0 \subset J$ . If  $\mathbb{k}$  is a field of characteristic zero, then  $A$  is not a bialgebra.*

*Proof.* We provide a sketch, as the proof is similar to that of [28, Proposition 3.1]. Since each  $d_i < \infty$ , the only possible counit  $\varepsilon : A \rightarrow \mathbb{k}$  sends each  $\mathbf{s}_i$  to 0. Let  $\mathfrak{m}$  denote the two-sided augmentation ideal in  $A$ , generated by  $\{\mathbf{s}_i : i \in I\}$ ; then one shows that any coproduct  $\Delta : A \rightarrow A \otimes A$  must satisfy:

$$\Delta(\mathbf{s}_i) \in 1 \otimes \mathbf{s}_i + \mathbf{s}_i \otimes 1 + \mathfrak{m} \otimes \mathfrak{m}.$$

But now,

$$0 = \Delta(\mathbf{s}_i)^{d_i} = \sum_{k=1}^{d_i-1} \binom{d_i}{k} \mathbf{s}_i^k \otimes \mathbf{s}_i^{d_i-k} + \text{higher degree terms},$$

and this is impossible because  $n\mathcal{H}(W, \mathbf{d}, J_0)$  surjects onto the quotient of the usual nil-Coxeter algebra  $NC_W((2, \dots, 2))$  by some braid words (each of length at least 2), so the  $\mathbf{s}_i$  have nonzero images in this quotient.  $\square$

A consequence of this result is that the module category  $\text{Rep } A$  (with  $A = n\mathcal{H}(W, \mathbf{d}, J_0)$ ) cannot be a tensor category. But since  $\Delta : A \rightarrow A \otimes A$  is co-commutative,  $\text{Rep } A$  is a symmetric semigroup category. We refer the reader to [28, Theorem A] and [14, Proposition 14.2 and Theorem 18.3] for more on the Tannakian formalism behind such categories. Additionally, we mention for completeness how to pass to an “honest” tensor category from  $A$ . Define a central extension of the graded algebra  $A$  by an element  $\mathbf{s}_\infty$ :

$$0 \rightarrow \mathbb{k}\mathbf{s}_\infty \rightarrow \tilde{A} \rightarrow A \rightarrow 0,$$

with the non-graded  $\mathbb{k}$ -algebra  $\tilde{A}$  generated by  $\mathbf{s}'_i, i \in I$  and  $\mathbf{s}_\infty$  via

$$(\mathbf{s}'_i)^{d_i} = \mathbf{s}'_i \mathbf{s}_\infty = \mathbf{s}_\infty \mathbf{s}'_i = \mathbf{s}_\infty^2 = \mathbf{s}_\infty, \quad \forall i \in I,$$

and the analogues for the  $\mathbf{s}'_i$  of the braid-word relations:

$$\mathbf{v}'_j - \mathbf{w}'_j = 0, \quad \forall j \in J_0, \quad \mathbf{v}'_l = \mathbf{w}'_l = \mathbf{s}_\infty, \quad \forall l \in J \setminus J_0.$$

Then  $\tilde{A}$  is indeed a bialgebra, under:

$$\Delta(\mathbf{s}'_i) := \mathbf{s}'_i \otimes \mathbf{s}'_i, \quad \Delta(\mathbf{s}_\infty) := \mathbf{s}_\infty \otimes \mathbf{s}_\infty, \quad \varepsilon(\mathbf{s}_i) = \varepsilon(\mathbf{s}'_\infty) = 1, \quad \forall i \in I,$$

and hence  $\text{Rep } \tilde{A}$  is a monoidal category. We refer the reader to [28, Section 3] for more details in the special case  $J_0 = J$ . Notice also that the algebras  $\tilde{A} = \widetilde{n\mathcal{H}(W, \mathbf{d}, J_0)}$  form a lattice of bialgebras that is isomorphic to the product of  $2^J$  with  $I$  copies of  $\mathbb{Z}^{\geq 2}$ .

A second theme, extensively explored for decades in the Coxeter/Hecke/Lie theory literature, involves classifying finite-dimensional objects in various settings. Such classifications have been of enormous interest in recent and earlier times – including complex simple Lie algebras; real and complex reflection groups [10, 40] and their nil-Coxeter and more general Hecke algebras; finite type quivers, the McKay–Slodowy correspondence, and Kleinian singularities. This evergreen theme has seen recent additions, including for finite-dimensional Nichols algebras [22, 25, 24]; and finite-dimensional pointed Hopf algebras [1], which are intimately connected to small quantum groups. There are also other combinatorial-type results, including by Stembridge and Hart, that are discussed presently; and the second author’s recent work classifying the finite-dimensional generalized nil-Coxeter algebras [28]. The present work is a sequel to this last reference [28], connecting it with the works by Stembridge and Hart, and also going beyond [28] via refined combinatorial phenomena for  $k = 4, 5$  even in the “usual” (i.e.,  $\mathbf{d} = (2, \dots, 2)$ ) Coxeter types  $B_n, F_4, H_3, H_4$ .



We next mention a connection with the question of *flatness of deformations*. In this paper, our goal is to explore the flatness question for e.g. the Temperley–Lieb algebras and their nil-Hecke versions. For instance, in their theses Fan and Graham studied Temperley–Lieb algebras  $TL_W((2, \dots, 2))$  for Coxeter groups  $W$  (see [16, 17, 21]), say with  $\mathbb{k} = \mathbb{Z}[u, u^{-1}]$ . These are quotients of the Iwahori–Hecke algebras  $\mathcal{H}_u(W)$ ; the authors showed in *loc. cit.* that  $TL_W((2, \dots, 2))$  has a  $\mathbb{k}$ -basis in bijection with the fully commutative words in  $W$ . Now Theorem B shows that  $TL_W((2, \dots, 2))$  is indeed a flat deformation of  $N TL_W((2, \dots, 2))$ , meaning that the following diagram is (i) valid, and (ii) a commuting square:

$$\begin{array}{ccc} \mathcal{H}_u(W) & \xrightarrow{\text{gr}} & NC_W((2, \dots, 2)) \\ \downarrow & & \downarrow \\ TL_W((2, \dots, 2)) & \xrightarrow{\text{gr}} & N TL_W((2, \dots, 2)) \end{array}$$

More strongly, we will show in Theorem 2.3 that the above flatness phenomenon holds for nil-Hecke algebras  $n\mathcal{H}(W, (2, \dots, 2), J_{<k})$  for *all*  $k$ . We then extend this to arbitrary  $\mathbf{d}$  in Theorem 2.4.

Our next connection is to the combinatorics of Coxeter groups  $W$ . Stembridge [44] studied the fully commutative elements  $W_{\text{fc}}$  (see Theorem B for the definition) in connection with the Bruhat and weak orderings. In [32] the authors study the 2-sided Kazhdan–Lusztig cell formed by the words  $W_1 \subset W$  that have a unique reduced expression. These words have been counted recently in [23]. In fact all words in the group  $W$  have been studied for the above reasons.

We now introduce a more general notion of word-sets  $W(J_0) \subset W$  for any Coxeter group, of which the above sets  $W_{\text{fc}}, W_1, W$  are special cases for  $J_0 = J_{<3}, J_{<2}, J_{<\infty}$  respectively.

**Definition 2.2.** Given a Coxeter matrix  $M$  with corresponding group  $W$ , define for  $J_0 \subset \binom{I}{2}$  the subset of elements  $W(J_0) \subset W$ , to consist of all  $w \in W$  such that no reduced expression for  $w$  has a substring of positive length that occurs in the set  $\{\mathbf{v}_j, \mathbf{w}_j : j \in J \setminus J_0\}$ .

With this notation in hand, we can state the promised result on flatness of deformations of Temperley–Lieb algebras – and more generally, relate every subset  $W(J_0)$  to the corresponding nil-Hecke algebra  $n\mathcal{H}(W, (2, \dots, 2), J_0)$ .

**Theorem 2.3.** *Suppose  $M$  is a Coxeter matrix, and  $J_0 \subset \binom{I}{2}$  as above. Then the nil-Hecke algebra  $n\mathcal{H}(W, (2, \dots, 2), J_0)$  is a free  $\mathbb{k}$ -module with basis  $\{\overline{\mathbf{s}}_w : w \in W(J_0)\}$ , where  $\{\mathbf{s}_w : w \in W\}$  is the “canonical” basis of the nil-Coxeter algebra  $NC_W((2, \dots, 2))$ .*

In particular, by [16, 17, 21], Temperley–Lieb algebras are flat deformations of their nil-Temperley–Lieb analogues.

Theorem 2.3 thus provides a broader setting into which the families of words  $W(J_0)$  fit: bases of nil-Hecke algebras corresponding to Coxeter matrices. Even more generally, we will consider (bases of) nil-Hecke algebras for generalized Coxeter matrices, i.e. where  $\mathbf{s}_i^{d_i} = 0$  for  $d_i \geq 2$ . Beyond proving Theorem 2.3, in the remaining sections we will characterize when such bases are finite in size.

*Proof of Theorem 2.3.* Begin by observing that the subset  $M$  in  $NC_W(2, \dots, 2)$  consisting of the words

$$M := \{\mathbf{s}_w : w \in W\} \sqcup \{0\}$$

forms a monoid under multiplication; and moreover,  $M \setminus \{0\}$  forms a  $\mathbb{k}$ -basis of  $NC_W((2, \dots, 2))$  (see e.g. [26]). Now let  $I(J_0)$  be the  $\mathbb{k}$ -span in  $NC_W((2, \dots, 2))$  of the words

$$\{\alpha \mathbf{v}_j \beta, \alpha \mathbf{w}_j \beta : j \in J \setminus J_0, \alpha, \beta \in M\}.$$

Then  $I(J_0)$  clearly contains the elements  $\mathbf{s}_w$ ,  $w \notin W(J_0)$ ; moreover, it is a two-sided ideal of  $NC_W((2, \dots, 2))$ . Now suppose a linear combination  $\sum_{w \in W} c_w \mathbf{s}_w \in I(J_0)$ . By definition/ $\mathbb{k}$ -freeness, if  $c_w \neq 0$  then  $w \notin W(J_0)$ , and  $I(J_0)$  intersects the  $\mathbb{k}$ -span of  $\{\mathbf{s}_w : w \in W(J_0)\}$  trivially. This concludes the proof.  $\square$

The above theorem provides a natural basis for nil-Hecke algebras with  $\mathbf{d} = (2, 2, \dots, 2)$ . We now strengthen this result to one which applies to all of our nil-Hecke algebras, and which is key to proving the finite-dimensionality in all of our main results.

**Theorem 2.4.** *Let  $W$  be a Coxeter group generated by  $\{s_i : i \in I\}$ , with corresponding nil-Hecke algebra  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$ , where  $\mathbf{d} \in (\mathbb{Z}^{\geq 2})^I$  and  $1 \leq k \leq \infty$ . Consider equivalence classes of strings in the  $s_i$ , where two strings are equivalent if one can be reached from another by applying the braid relations finitely many times. Let  $S$  denote the set of classes such that no string in the class contains any of the substrings  $s_i^{d_i}$  or the braid words of length  $k$  or more. Then the monomials corresponding to some choice of class representatives from  $S$  form a free  $\mathbb{k}$ -module basis of  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$ .*

*Proof.* Since any monomial not coming from  $S$  can be reduced to zero, these monomials clearly span  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$ . It remains to prove that  $\dim n\mathcal{H}(W, \mathbf{d}, J_{<k}) \geq |S|$ , and we would be done.

Consider a free  $\mathbb{k}$ -module  $\mathcal{M}$  with basis elements indexed by  $S$ . We now define a natural  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$ -module structure on  $\mathcal{M}$  as follows: the element  $\mathbf{s}_j$  acts on the class  $[s_{i_1} \cdots s_{i_n}]$  by sending it to the class  $[s_j s_{i_1} \cdots s_{i_n}]$  if it is in  $S$ , and to zero otherwise. This extends to an action of the free associative algebra  $\mathbb{k}\langle \mathbf{s}_i \mid i \in I \rangle$ .

We now verify this action indeed satisfies the defining relations in  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$ . Indeed, if applying the  $\mathbf{s}_i$ -action  $d_i$  times on a basis element yields a non-zero basis element, then this class contains a string with  $d_i$  consecutive leading  $s_i$ 's and thus cannot be in  $S$ , a contradiction. Similarly, multiplying by braid words of length  $k$  or more gives the zero vector. Finally, the braid relations of lengths less than  $k$  hold: indeed, multiplying a basis element by either side of the braid equations either yields elements in the same equivalence class by definition, or yields elements not in  $S$ , giving the zero vector on either side.

Since  $\mathcal{M}$  is generated as a  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$ -module by the basis vector corresponding to the empty string, this gives a surjection  $n\mathcal{H}(W, \mathbf{d}, J_{<k}) \twoheadrightarrow \mathcal{M}$ . This, combined with the first paragraph, completes our proof.  $\square$

Following Theorem 2.4 on a word basis of  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$ , we describe here a final connection of this work, to the study of PBW bases and deformations – more broadly than the commuting square in this section. The study of PBW (Poincaré–Birkhoff–Witt) type deformations has seen intense activity for several decades now, including foundational works on Drinfeld Hecke algebras [12], graded affine Hecke algebras [36], symplectic reflection algebras [13] (including rational Cherednik algebras), as well as on infinitesimal Hecke algebras, quantum analogues, and their generalizations. The recent program of Shepler and Witherspoon (and their coauthors) has led to a profusion of activity; we mention e.g. [41, 42, 47] here. In all of these works, a bialgebra  $A$  (which is most often a Hopf algebra) acts on a symmetric algebra  $S_V$  of some vector space  $V$ , and one is interested in understanding which deformations of the smash-product algebra  $A \rtimes S_V$  are “PBW”, or flat.

When  $A$  is a nil-Hecke algebra, there are several significant features to note. First, these are not bialgebras by Proposition 2.1 (but possess a coproduct  $\Delta$ ); and yet, a variant of the ‘‘PBW theorem’’ nevertheless holds for the algebras  $A \rtimes S_V$ , which subsumes the PBW theorems in various previous works. (See the previous paper [27] by the second author for the details.) Thus the nil-Hecke algebras  $n\mathcal{H}(W, \mathbf{d}, J_0)$  widen beyond [28] the class of examples of such ‘‘algebras with coproduct’’  $(A, \Delta)$ , which do not possess an antipode or even a counit, yet fit into the framework of the aforementioned works.

Second, there also are technical consequences of finite-dimensionality. We mention two of these; in both, we will assume  $\mathbb{k}$  to be a field. It was shown in [42], [27] that if  $(A, \Delta)$  is finite-dimensional, one can characterize those graded  $\mathbb{k}[t]$ -deformations of  $A \rtimes S_V$ , for which the fiber at  $t = 1$  satisfies the PBW property. This provides a PBW-theoretic motivation to classify the finite-dimensional nil-Hecke algebras.

Even more is true. It was shown in [27] that if  $(A, \Delta)$  is any algebra with coproduct (e.g.  $A = n\mathcal{H}(W, \mathbf{d}, J_{<k})$ ), and if it is local with augmentation ideal  $\mathfrak{m}$  such that  $\Delta(\mathfrak{m}) \subset \mathfrak{m} \otimes \mathfrak{m}$ , then one obtains much additional information regarding the deformations of  $A \rtimes S_V$  – their abelianization, center, and (simple) modules. See e.g. [27, Section 6.1]. Thus, we have a second motivation from the perspective of PBW deformations – to understand which nil-Hecke algebras  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  are local. This is precisely the point of Theorem D; and it shows that this question turns out to be *equivalent* to the main results of the paper:  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  must be finite-dimensional.

### 3. PROOF OF THEOREM A: FINITE RANK NIL-HECKE ALGEBRAS WITH $k = 1, 2$

The remainder of the paper is devoted to proving the above classification theorems on finite-dimensional nil-Hecke algebras. To do so, we will employ the diagrammatic calculus utilized in [28]. We begin by showing Theorem A in this section; the proof is in steps.

**Step 1:** By Theorem 2.3,  $n\mathcal{H}(W, \mathbf{d}, J_{<2})$  has finite  $\mathbb{k}$ -rank if and only if  $W(J_{<2})$  is also finite. By using the results of [23], this proves the result if  $\mathbf{d} = (2, \dots, 2)$ , i.e., when dealing with quotients of usual nil-Coxeter algebras.

**Step 2:** In the remainder of the proof, at least one  $d_i$  is 3 or more. Since  $n\mathcal{H}(W, \mathbf{d}, J_0) \rightarrow n\mathcal{H}(W, (2, \dots, 2), J_0)$ , by the previous step the Coxeter (multi)graph of  $W$  must be a tree with no  $m_{ij} = \infty$  and at most one  $m_{ij} \geq 4$ .

Suppose first that we have a simply laced tree, and  $d_\alpha, d_\gamma \geq 3$  for  $\alpha \neq \gamma \in I$ . As the graph of  $W$  is connected, suppose

$$\alpha \longleftrightarrow \beta_1 \longleftrightarrow \dots \longleftrightarrow \beta_{m-1} \longleftrightarrow \gamma$$


is a path in  $I$  (in the figures below, we write  $m' = m - 1$ ). Now define a free  $\mathbb{k}$ -module  $\mathcal{M}$  with basis vectors

$$\{A_r, B_{1r}, \dots, B_{mr}, C_r, B'_{1r}, \dots, B'_{mr} : r \geq 1\}.$$

Give  $\mathcal{M}$  an  $n\mathcal{H}(W, \mathbf{d}, J_{<2})$ -module structure by letting every  $\mathbf{s}_i$  kill all basis vectors, except for the actions described by the first diagram in Figure 3.1. Explicitly, the basis vectors on which the action is nonzero are:

$$\begin{aligned} \mathbf{s}_\alpha(A_r) &:= B_{1r}, & \mathbf{s}_{\beta_j}(B_{jr}) &:= B_{j+1,r} \quad (1 \leq j \leq m-1), & \mathbf{s}_\gamma(B_{mr}) &:= C_r, \\ \mathbf{s}_\gamma(C_r) &:= B'_{mr}, & \mathbf{s}_{\beta_j}(B_{jr}) &:= B_{j-1,r} \quad (2 \leq j \leq m), & \mathbf{s}_\alpha(B'_{1r}) &:= A_{r+1}. \end{aligned}$$

Thus the ‘‘+’’ at the head of an arrow refers to the index  $r$  increasing by 1. One verifies that the defining relations for  $n\mathcal{H}(W, \mathbf{d}, J_{<2})$  hold on each basis vector, hence in  $\text{End}_{\mathbb{k}}(\mathcal{M})$ . Thus

FIGURE 3.1. The modules  $\mathcal{M}$  for the nil-Hecke algebras in Steps 2 and 3

$\mathcal{M}$  is a cyclic  $n\mathcal{H}(W, \mathbf{d}, J_{<2})$ -module generated by  $A_{11}$ . Now  $\mathcal{M}$ , hence  $n\mathcal{H}(W, \mathbf{d}, J_{<2})$ , is not a finitely generated  $\mathbb{k}$ -module.

**Step 3:** Suppose next that some  $m_{ij} > 3$  and some  $d_{i_0} > 2$ . Without loss of generality there exist nodes  $\beta_1, \dots, \beta_{m-1}$  for some  $m \geq 1$  (in the figure we write  $m' := m - 1$ ), such that

$$\alpha = i_0 \quad \longleftrightarrow \quad \beta_1 \quad \longleftrightarrow \quad \dots \quad \longleftrightarrow \quad \beta_{m-1} = i \quad \longleftrightarrow \quad \gamma = j$$

Now construct a free  $\mathbb{k}$ -module  $\mathcal{M}$  as in Step 2, this time giving it an  $n\mathcal{H}(W, \mathbf{d}, J_{<2})$ -module structure using the second diagram in Figure 3.1. One verifies that the action on  $\mathcal{M}$  satisfies the defining relations. Hence the nil-Hecke algebra is once again of infinite  $\mathbb{k}$ -rank.

**Step 4:** It remains to show the assertion when the Coxeter (multi)graph of  $W$  is a simply laced tree with node set  $I$ , and  $d_{i_0} \geq 3$  for a unique  $i_0 \in I$ . Let us denote the corresponding nil-Hecke algebra by  $n\mathcal{H}(W, i_0, d_{i_0}, J_{<2})$ . For nodes  $s, t \in I$ , let  $[s, t]$  denote the path from source  $s \in I$  to target  $t \in I$ . If  $s = i_1, \dots, i_n = t$  enumerates sequentially the nodes in  $[s, t]$  for some  $n \geq 1$ , define:

$$\mathbf{s}_{[s,t]} := \mathbf{s}_{i_1} \cdots \mathbf{s}_{i_n}. \quad (3.1)$$

We begin by enumerating a spanning set for  $n\mathcal{H}(W, i_0, d_{i_0}, J_{<2})$  of the correct size. Notice that the product  $\mathbf{s}_i \mathbf{s}_j$  (with  $i, j \in I$ ) vanishes unless  $i = j = i_0$  or  $i \neq j$  are adjacent in  $I$ . It follows that the set

$$\{1\} \sqcup \{\mathbf{s}_{[s,t]} : s, t \in I\} \sqcup \{\mathbf{s}_{[s,i_0]} \mathbf{s}_{i_0}^{k-2} \mathbf{s}_{[i_0,t]} : s, t \in I, 2 \leq k \leq d_{i_0} - 1\}$$

spans  $n\mathcal{H}(W, i_0, d_{i_0}, J_{<2})$ . Now by the proof of Theorem 2.4, this is indeed a basis, and we are done.  $\square$

4. PROOF OF THEOREM B: FINITE RANK NIL-HECKE ALGEBRAS WITH  $k = 3$ 

*Proof of Theorem B.* We begin by showing that the only “adjacency graphs”, corresponding to Coxeter groups  $W$  that admit finite  $\mathbb{k}$ -rank nil-Hecke algebras for  $NTL_W(\mathbf{d}) = n\mathcal{H}(W, \mathbf{d}, J_{<3})$  (these algebras are defined following Definition 1.1), are Dynkin diagrams of finite type or type  $E_n, n \geq 9$ . Here, first, by the *adjacency multigraph* of a Coxeter group associated to a Coxeter matrix  $(m_{ij})_{i,j=1}^n$  with  $n = |I|$ , we mean the multigraph on  $n$  nodes with  $m_{ij} - 2$  edges between nodes  $i \neq j \in \{1, \dots, n\}$ . Next, the corresponding *adjacency graph* is the simple graph on  $n$  nodes with  $\mathbf{1}(m_{ij} > 2)$  edges between nodes  $i \neq j$ .

If  $\mathbf{d} = (2, \dots, 2)$ , then  $W$  is a Coxeter group, and by Theorem 2.3, the problem reduces to finding all  $W$  with finitely many fully commutative elements. This was worked out in [15, 21, 44]. In what follows, we therefore assume there exists at least one  $i$  such that  $d_i \geq 3$ . Let this  $i$  correspond to vertex  $\alpha$ .

**Step 1.** Suppose the adjacency multigraph of  $W$  contains a cycle on nodes  $\beta_1, \dots, \beta_m$  with  $m \geq 3$ , i.e.,  $m_{\beta_i, \beta_{i+1}} > 2$  for  $0 < i < m$  and  $m_{\beta_m, \beta_1} > 2$ . Now construct a free  $\mathbb{k}$ -module  $\mathcal{M}$  with basis given by the countable set  $\{A_{1r}, \dots, A_{mr} : r \geq 1\}$ , and define the following  $NTL_W(\mathbf{d})$ -action on it:  $\mathbf{s}_{\beta_i}$  kills  $A_{jr}$  except that  $\mathbf{s}_{\beta_i}(A_{ir}) := A_{i+1,r}$  for  $0 < i < r$  and  $\mathbf{s}_{\beta_m}(A_{mr}) := A_{1,r+1}$ . It is easy to verify that the defining relations of  $NTL_W(\mathbf{d})$  hold in  $\text{End}_{\mathbb{k}}(\mathcal{M})$ , as they hold on each  $A_{ir}$ . Therefore  $\mathcal{M}$  is a cyclic  $NTL_W(\mathbf{d})$ -module generated by  $A_{11}$ , which is not finitely generated as a  $\mathbb{k}$ -module. Hence neither is  $NTL_W(\mathbf{d})$ , as claimed.

The analysis in this step can be conveniently expressed by a diagram of a cycle. We will do so for other cases in the remainder of the proof.

**Step 2.** The remaining adjacency multigraphs are those whose underlying simple graphs are trees. We next claim there is no vertex adjacent to 4 nodes, and at most one vertex adjacent to 3 nodes. Indeed, if  $\beta_1$  is adjacent to nodes  $\alpha_1, \alpha_2, \gamma_1, \gamma_2$ , then we appeal to Figure 4.1 with  $m = 2$ , to generate a  $NTL_W(\mathbf{d})$ -module  $\mathcal{M}$  with basis  $\{A_r, B_r, B_{1r}, B_{2r}, B'_{1r}, B'_{2r}, C_r, D_r : r \geq 1\}$ . The module relations are read off of the diagram. Namely,  $\mathbf{s}_i$  kills all basis vectors for all  $i \in I$ , with the following exceptions:

$$\begin{aligned} \mathbf{s}_{\beta_1}(B_{1r}) &= B_{2r}, & \mathbf{s}_{\gamma_1}(B_{2r}) &= D_r, & \mathbf{s}_{\gamma_2}(B_{2r}) &= C_r, & \mathbf{s}_{\gamma_1}(C_r) &= B'_{2r}, & \mathbf{s}_{\gamma_2}(D_r) &= B'_{2r}, \\ \mathbf{s}_{\beta_1}(B'_{2r}) &= B'_{1r}, & \mathbf{s}_{\alpha_2}(B'_{1r}) &= B_r, & \mathbf{s}_{\alpha_1}(B_r) &= B_{1r}, & \mathbf{s}_{\alpha_2}(A_r) &= B_{1r}, & \mathbf{s}_{\alpha_1}(B'_{1r}) &= A_{r+1}. \end{aligned}$$




FIGURE 4.1. Diagrammatic calculus for the algebras  $NTL(M)$ ; here  $m' = m - 1$ .

(The “+” at the head of an arrow again refers to the index increasing by 1.) It is easy to verify that the defining relations of  $NTL_W(\mathbf{d})$  hold at each node in the diagram (i.e., at the corresponding basis vector) and hence in  $\text{End}_{\mathbb{k}}(\mathcal{M})$ . This yields a cyclic  $NTL_W(\mathbf{d})$ -module  $\mathcal{M}$  of infinite  $\mathbb{k}$ -rank, as desired.

Next if  $\deg \alpha = \deg \gamma = 3$ , let

$$\alpha = \beta_1 \longleftrightarrow \beta_2 \longleftrightarrow \cdots \longleftrightarrow \beta_{m-2} \longleftrightarrow \gamma = \beta_{m-1}$$

be the path between them, for suitable  $m$ . Also suppose  $\alpha$  is adjacent to  $\beta_2, \alpha_1, \alpha_2$ , and  $\gamma$  is adjacent to  $\beta_{m-2}, \gamma_1, \gamma_2$ . We again appeal to Figure 4.1 to generate a representation  $\mathcal{M}$  which has infinite  $\mathbb{k}$ -rank.

**Remark 4.1.** A brief remark for the reader, about checking that the relations hold in Figure 4.1 and the other figures in this paper: (a) On each commuting sub-diagram, one only needs to check the braid relations in  $J_{<k}$ , and account for the ‘+’ signs. (b) On each directed cycle in the diagrams, one needs to check that no braid word in  $J \setminus J_{<k}$  occurs, nor any word  $\mathbf{s}_i^{d_i}$ .

**Step 3.** Thus the Dynkin multigraph is in fact acyclic, with at most one vertex of degree 3 and no vertex of degree  $\geq 4$ . First, we use the first diagram in Figure 3.1 to show there exists a unique node  $i$  such that  $d_i \geq 3$ . Second, use Figure 4.2 with  $m = 1$  to show this node has degree 1 (i.e., is pendant).




FIGURE 4.2. Diagrammatic calculus for Step 3; here  $m' = m - 1$ .

Third, use the second diagram in Figure 3.1 to show that the Dynkin multigraph of  $W$  is in fact simply laced. Hence from above, it is a tree graph.

Next, suppose there is a vertex  $\beta$  of degree 3. Let

$$\alpha = \beta_1 \longleftrightarrow \beta_2 \longleftrightarrow \cdots \longleftrightarrow \beta_{m-2} \longleftrightarrow \beta = \beta_{m-1}$$

be a path from  $\alpha$  to  $\beta$ ; further, let  $\gamma_1$  and  $\gamma_2$  be two more vertices adjacent to  $\beta$ . Then the module pictured in Figure 4.2 again has infinite  $\mathbb{k}$ -rank, a contradiction. This shows that the adjacency multigraph is necessarily a path graph. As we have seen,  $\mathbf{d} = (d, 2, \dots, 2)$  for some  $d \geq 3$ , and so we obtain a quotient of the algebra  $NC_A(n, d)$  as desired.

**Step 4.** To finish the proof, we need to determine the  $\mathbb{k}$ -rank of  $NTL_W(\mathbf{d})$ , where  $W$  is of type  $A_n$  and  $\mathbf{d} = (d, 2, \dots, 2)$ . By Theorem 2.4, it suffices to enumerate all the monomials in  $x_1, \dots, x_n$  that are not equivalent (via commutation relations  $x_i x_j = x_j x_i$  for  $|i - j| > 1$ ) to any monomial containing the strings  $x_1^d, x_i^2$  for  $i > 1$ , or containing any of the length 3 braid words. We extend the argument in [20] to accomplish this.

Consider a nonzero monomial in  $NTL_W(\mathbf{d})$ , and in its equivalence class under the commutation relations ( $x_i x_j = x_j x_i$  for  $|i - j| > 1$ ), look at the lexicographically smallest monomial  $\mathbf{w}$  as in [20]. We claim that all the  $x_1$ 's occur in a contiguous string in  $\mathbf{w}$ . Indeed, suppose not. Then by the arguments in [20], the monomial  $\mathbf{w}$  can be assumed to be composed of decreasing runs, where in each run the indices go down by one, except possibly where  $x_1$  is followed by  $x_1$ . Since  $x_1$ 's can only occur at the ends of such runs, if there are two  $x_1$ 's that are not consecutive, a closest such pair must be part of a substring that looks like

$x_1x_jx_{j-1}\cdots x_2x_1$ . But this can be reduced to  $x_jx_{j-1}\cdots x_1x_2x_1$  and thus zero, proving our claim.

One can repeat the arguments in [20, Section 3] and verify that Lemmas 1, 3, and 4 there still hold, with the following exceptions:

- (1) There might be  $x_1$  followed by  $x_1$  in the monomial  $\mathbf{w}$ , contrary to Lemma 1; or
- (2) The indices of the peaks might not be strictly increasing, precisely when the first run starts with  $x_i$  for  $i \geq 2$ , ends with  $x_1^j$  for  $j \geq 2$ , and is followed by  $x_2$ , contrary to Lemma 2.

Let us set aside the monomials that come under exception (2) for the moment. Then the monomials we need to count are precisely those counted by [20], with the added caveat that there might be a contiguous block of up to  $d - 1$   $x_1$ 's. Now if the monomial contains no  $x_1$ , then this is simply a monomial in  $NTL_W((2, \dots, 2))$  on the generators  $x_2, \dots, x_n$ ; by the results of [20], there are  $C_n$  such monomials. If the monomial does contain a nonempty block of  $x_1$ 's, then deleting all but one  $x_1$  gives a monomial containing  $x_1$  in  $NTL_W((2, \dots, 2))$ ; conversely, given a monomial in  $NTL_W((2, \dots, 2))$  containing  $x_1$ , one can form  $d - 1$  different valid monomials from there by adding  $x_1$ 's. The number of  $NTL_W((2, \dots, 2))$ -monomials containing an  $x_1$  is simply  $C_{n+1} - C_n$  (since there are  $C_n$  monomials not having  $x_1$ ), so the total number is

$$C_n + (d - 1)(C_{n+1} - C_n) = (d - 1)C_{n+1} - (d - 2)C_n.$$

Finally, we count the monomials that exhibit exception (2). Such a monomial must start with  $x_i x_{i-1} \cdots x_2 x_1^j x_2$ , and then either the next term is  $x_3$ , or the remaining string is a non-zero monomial on the generators  $x_{i+1}, \dots, x_n$ . Indeed, the peak of the next run cannot be  $x_k$  for  $4 \leq k \leq i$ , else we can move it to the left by the commutation relations and the entire monomial reduces to zero. The next run cannot even contain such a  $x_k$ , else it would have to contain  $x_i$  and thus the initial segment of this run till  $x_i$  can be commuted leftward to reduce the monomial to zero, proving our claim. Continuing this argument, we see that the  $x_1^j$  in our monomial is followed by a string of the form  $x_2 x_3 \cdots x_\ell$ , with  $2 \leq \ell \leq i$ , and a nonzero (possibly empty) monomial in the generators  $x_{i+1}, \dots, x_n$ . Thus the number of such monomials is  $(i - 1)C_{n-i+1}$ . Summing over all possible  $i$  and  $j$ , the total number becomes

$$(d - 2) \sum_{i=2}^n (i - 1)C_{n-i+1} = (d - 2) \sum_{j=1}^{n-1} jC_{n-j}.$$

As in [20], we can verify these indeed correspond to non-zero monomials, so adding this to the previous count completes the proof.  $\square$

## 5. PROOF OF THEOREM C AND ITS COROLLARIES

We now consider the remaining cases  $4 \leq k \leq \infty$ , starting with  $k = 4$ .

*Proof of Theorem C.* Since  $n\mathcal{H}(W, \mathbf{d}, J_{<\infty})$  surjects onto  $n\mathcal{H}(W, \mathbf{d}, J_{<4})$ , it follows that  $n\mathcal{H}(W, \mathbf{d}, J_{<4})$  has finite  $\mathbb{k}$ -rank if  $n\mathcal{H}(W, \mathbf{d}, J_{<\infty})$  does; these cases are listed in Theorem 1.3. On the other hand,  $n\mathcal{H}(W, \mathbf{d}, J_{<4}) \twoheadrightarrow n\mathcal{H}(W, \mathbf{d}, J_{<3})$ , so  $n\mathcal{H}(W, \mathbf{d}, J_{<4})$  has infinite  $\mathbb{k}$ -rank whenever  $n\mathcal{H}(W, \mathbf{d}, J_{<3})$  does. The cases where this is not so are listed in Theorem B. Comparing these two lists, we see that only the cases where  $W$  is of type  $E_n (n \geq 9)$ ,  $F_n (n \geq 5)$  or  $H_n (n \geq 5)$  with  $\mathbf{d} = (2, \dots, 2)$  remain to be checked for finiteness.

If  $W$  has type  $E_n$ , there are no braid relations of length 4 or more, and so  $n\mathcal{H}(W, \mathbf{d}, J_{<4}) \cong n\mathcal{H}(W, \mathbf{d}, J_{<\infty})$ , which is known by [28] to have infinite  $\mathbb{k}$ -rank for  $n \geq 9$ . Next, assume  $W$  is

of type  $F_n$  with  $n \geq 5$ . We will prove that  $n\mathcal{H}(W, \mathbf{d}, J_{<4})$  has infinite  $\mathbb{k}$ -rank for  $n = 5$ , from which the  $n > 5$  cases would follow trivially.

Let  $W$  be of type  $F_5$ , and label the generators as in Figure 5.1.




FIGURE 5.1. Dynkin graphs for  $F_5$  and  $H_5$

Then the infinite rank module defined by the diagram in Figure 5.2 proves our claim.




FIGURE 5.2. An infinite rank module for  $F_5$

For the remaining case, it suffices to show a similar module for  $H_5$ . This is demonstrated in Figure 5.3.

Next, we compute the ranks of the finite rank nil-Hecke algebras. First suppose  $W$  is of type  $A$  and  $\mathbf{d} = (d, 2, \dots, 2)$  or  $(2, \dots, 2, d)$  for some  $d > 2$ . As all braid relations have length 3,  $n\mathcal{H}(W, \mathbf{d}, J_{<4}) = n\mathcal{H}(W, \mathbf{d}, J_{<\infty})$ , and so we are done by Theorem 1.3.

Otherwise, we fix  $\mathbf{d} = (2, \dots, 2)$  henceforth. Now if  $W$  is simply laced (of type  $ADE$ ) or of type  $I_2(m)$  with  $m < 4$ , then there are no braid relations of length 4 or more, so  $n\mathcal{H}(W, \mathbf{d}, J_{<k}) \cong n\mathcal{H}(W, \mathbf{d}, J_{<\infty}) = NC_W((2, \dots, 2))$ . This has  $\mathbb{k}$ -rank precisely  $|W|$ . If instead  $W$  is of type  $I_2(m)$ ,  $m \geq k = 4$ , then both sides of the braid relation (i.e., the longest word in  $NC_W((2, 2))$ ) get killed in  $n\mathcal{H}(W, \mathbf{d}, J_{<4})$ , while the remaining  $2m - 1$  words are not killed in the quotient. The result now follows by Theorem 2.3.

We tackle the case of  $B_n$  next. Let the generators of the corresponding Weyl group  $W(B_n)$  be  $s_0, s_1, \dots, s_{n-1}$ , where the labeling is chosen such that  $m_{01} = 4$ , and  $m_{i,i+1} = 3$  for all






FIGURE 5.3. An infinite rank module for  $H_5$

$i \in \{1, \dots, n - 2\}$ . By Theorem 2.3, we need to count the set  $B_{n,4}$  of elements  $w \in W(B_n)$ , for which no reduced expression contains the substring  $s_0s_1s_0s_1$  or  $s_1s_0s_1s_0$ . For this, we use the following well-known combinatorial description of  $W(B_n)$  (see, for example, [4]):

**Theorem 5.1.** *Let  $W$  be the group of all permutations  $w$  on  $S = \{\pm 1, \dots, \pm n\}$  such that  $w(-a) = -w(a) \forall a \in S$ . Then  $W = W(B_n)$  is the Coxeter group of type  $B_n$ , with generators*

$$s_i = [1, \dots, i - 1, i + 1, i, i + 2, \dots, n], \quad 1 \leq i \leq n - 1; \quad s_0 = [-1, 2, \dots, n],$$

where we write  $w = [a_1, a_2, \dots, a_n] \in W(B_n)$  if  $w(i) = a_i \forall 1 \leq i \leq n$ .

Further, given any  $w \in W$ , the length function  $\ell(\cdot)$  satisfies the following properties:

- (1)  $\ell(ws_0) = \ell(w) + 1$  if and only if  $w(1) < 0$ ; and
- (2)  $\ell(ws_i) = \ell(w) + 1$  if and only if  $w(i) < w(i + 1)$  for  $i \in \{1, \dots, n - 1\}$ .

Using this, we now prove the following lemma:

**Lemma 5.2.** *Let  $w = [a_1, \dots, a_n] \in W(B_n)$  be as above. Then  $w \in B_{n,4}$  if and only if the following holds: no pair of indices  $i < j$  satisfies  $a_i > a_j$  and  $a_i, a_j < 0$ .*

*Proof.* Suppose  $w \in W(B_n)$  does not satisfy the condition: thus, there exist  $i < j$  such that  $a_j < a_i < 0$ . Call such a pair of indices a *bad pair*. One can pick  $i, j$  with minimal  $|i - j|$ . Perform the following operations on  $w$ : if the leftmost negative number in  $w$  occurs in position  $i > 1$ , multiply by the transposition  $s_{i-1}$ . If it occurs in position 1, multiply by  $s_0$ . Each move decreases the length by 1, and after sufficiently many applications, this brings  $a_i$  to position 1. By another sequence of length-decreasing transpositions, we can bring  $a_j$

to 2. Now one verifies that multiplying by  $s_0s_1s_0s_1$  further reduces the length by 4. Thus  $w'w'' = w''$ , where  $w'$  contains the string  $s_0s_1s_0s_1$ , and  $\ell(w) - \ell(w') = \ell(w'')$ . This implies  $w'^{-1}w''$  is a reduced expression for  $w$  with the substring  $s_1s_0s_1s_0$ , and thus  $w \notin B_{n,4}$ .

For the other direction, suppose  $w$  satisfies the given condition. We will prove  $w \in B_{n,4}$  by induction on  $\ell(w)$ . The base cases of  $\ell(w) \leq 1$  are clear. For the induction step, suppose to the contrary that  $w$  has a reduced expression with  $s_0s_1s_0s_1$  in it, and let  $s_i$  be the rightmost generator in that expression. Then  $w = w's_i$ , with  $\ell(w') = \ell(w) - 1$ . There are two cases:

*Case 1:* First suppose  $s_0s_1s_0s_1$  does not occur at the rightmost position in this expression for  $w$ , so it has to occur in the expression for  $w'$ . Now  $ws_i$  has smaller length than  $s$ , so if  $i \geq 1$ ,  $w(i) > w(i+1)$ . Multiplying by  $s_i$  swaps these, so if  $w$  does not have a bad pair, swapping  $w(i)$  with  $w(i+1)$  cannot create further bad pairs. Thus by the induction hypothesis,  $ws_i = w'$  does not have a reduced expression containing  $s_0s_1s_0s_1$ , a contradiction. Similarly, if  $i = 0$ ,  $w(1)$  must be negative, and swapping its sign cannot create more bad pairs. By a similar reasoning as above, we are done in this case.

*Case 2:* Otherwise,  $w = w''s_0s_1s_0s_1$ , with  $\ell(w'') = \ell(w) - 4$ . As this is a reduced expression,  $\ell(ws_1) < \ell(w)$ , implying  $w(1) > w(2)$ . Now  $w_1 = ws_1 = [w(2), w(1), \dots]$ , so  $\ell(w_1s_0) < \ell(w_1)$  implies  $w_1(1) = w(2) < 0$ . Next,  $w_2 = w_1s_0 = [-w(2), w(1), \dots]$ , so  $w_3 = w_2s_1 = [w(1), -w(2), \dots]$ , and  $w_4 = w_3s_0 = [-w(1), -w(2), \dots]$ . Since  $\ell(w_3s_0) < \ell(w_3)$ , we have  $w_3(1) = w(1) < 0$ . Thus  $(1, 2)$  is a bad pair in  $w$ , a contradiction. This finishes the proof.  $\square$

Returning to the main proof, we want to enumerate the elements in  $W(B_n)$  without a bad pair. This is counted in [39] as the  $\overline{12}$ -avoiding signed permutations, and the answer is precisely  $\sum_{k=0}^n \binom{n}{k}^2 k!$ . This shows the result for type  $B_n$ .

The only remaining cases are  $F_4$ ,  $H_3$  and  $H_4$ : all of finite  $\mathbb{k}$ -rank. By computer checking and use of Theorem 2.3, one verifies the numbers stated in Theorem C. The proof is now complete.  $\square$

We conclude by proving the two corollaries to Theorem C.

*Proof of Corollary 1.8.* Since  $n\mathcal{H}(W, \mathbf{d}, J_{<5}) \twoheadrightarrow n\mathcal{H}(W, \mathbf{d}, J_{<4})$ , it suffices to check the cases in Theorem B. The dihedral case is easy to verify, and the case of type  $A$  with  $\mathbf{d} = (d, 2, \dots, 2)$  or  $(2, \dots, 2, d)$  is immediate since there are no braid relations of length 4 or more in this case. As the same fact holds for types  $A, B, D, E, F$ , hence for these types (with  $\mathbf{d} = (2, \dots, 2)$ ) it follows that  $n\mathcal{H}(W, \mathbf{d}, J_{<5}) = n\mathcal{H}(W, \mathbf{d}, J_{<\infty}) = NC_W((2, \dots, 2))$  is the usual nil-Coxeter algebra, and hence has  $\mathbb{k}$ -rank  $|W|$ .

Finally, suppose  $W$  is of type  $H_3$  or  $H_4$ , and  $\mathbf{d} = (2, \dots, 2)$ . As all braid words have length either 3 or 5, it follows that  $J_{<3} = J_{<4}$  in both cases, and so the algebras and hence their  $\mathbb{k}$ -ranks remain unchanged when passing from  $k = 4$  to  $k = 5$ .  $\square$

*Proof of Corollary 1.10.* Since  $n\mathcal{H}(W, \mathbf{d}, J_{<k}) \twoheadrightarrow n\mathcal{H}(W, \mathbf{d}, J_{<5})$ , to identify the finite  $\mathbb{k}$ -rank nil-Hecke algebras, we need only consider the cases in Corollary 1.8, i.e., for  $k = 5$ . Once again the case of type  $A$  with  $\mathbf{d} = (d, 2, \dots, 2)$  or  $(2, \dots, 2, d)$  is immediate, and the dihedral case of  $I_2(m)$  is again easily checked. Since none of the remaining cases have braid relations of length 6 or more, in all non-dihedral cases with  $\mathbf{d} = (2, \dots, 2)$ , we have  $n\mathcal{H}(W, \mathbf{d}, J_{<k}) = n\mathcal{H}(W, \mathbf{d}, J_{<\infty}) = NC_W((2, \dots, 2))$  as above. This completes the proof.  $\square$

## 6. PROOF OF THEOREM D: NILPOTENCE OF THE AUGMENTATION IDEAL

We conclude this work by proving the outstanding, final result.

*Proof of Theorem D.* The key equivalence to be shown here is (1)  $\iff$  (3); that (1)  $\iff$  (2) was shown in earlier sections. First assume  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  has infinite  $\mathbb{k}$ -rank. Recall from above that every such algebra was shown to have infinite rank by demonstrating an explicit cyclic  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$ -module of infinite  $\mathbb{k}$ -rank; moreover, every diagram above that described such a module has at least one directed cycle that is “accompanied” by a “+” symbol. Let the generators along the edges of any such fixed cycle be  $\alpha_1, \alpha_2, \dots, \alpha_m$ , and suppose a basis element  $v$  in that module is associated to the initial node of the edge corresponding to  $\alpha_1$ . Take an arbitrary positive integer  $N$ . Then the element  $\mathbf{s} = (\mathbf{s}_{\alpha_m} \cdots \mathbf{s}_{\alpha_2} \mathbf{s}_{\alpha_1})^j$  belongs to  $\mathfrak{m}^N$  for large enough  $j \gg 0$ , and  $\mathbf{s}v$  is non-zero by how the module action is defined. This implies  $\mathfrak{s} \in \mathfrak{m}^N$  is non-zero; and since  $N$  was arbitrary,  $\mathfrak{m}$  is not nilpotent.

For the other direction, assume  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  has finite  $\mathbb{k}$ -rank. Via an application of Theorem 2.4, one obtains a finite set  $S_{\text{fin}}$  of monomials corresponding to equivalence classes in  $S$  (as defined in Theorem 2.4) that gives a  $\mathbb{k}$ -basis of the algebra in question. Thus,  $S_{\text{fin}}$  has an element of maximal length, say  $\ell$ . We claim that  $\mathfrak{m}^{\ell+1} = 0$ , which would prove the desired nilpotency.

Indeed, consider a monomial  $\mathbf{s}'$  in  $\mathfrak{m}^{\ell+1}$ : this must be a product of strictly more than  $\ell$  generators. If  $\mathbf{s}'$  equals some monomial composed of  $\ell$  generators or less, one can reduce this via using the defining relations of  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$ . Since applying the braid relations does not change the number of monomials, this can be reduced to some monomial containing the string  $\mathbf{s}_i^{d_i}$  or a braid word of length  $\geq k$  at some point, and hence it is zero in  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$ .

On the other hand, if  $\mathbf{s}'$  is not equal in  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  to any monomial composed of  $\ell$  generators or less, the corresponding string of  $\mathbf{s}_i$ 's cannot belong to any equivalence class in  $S$ , and thus must reduce to 0 (as in the proof of Theorem 2.4). This completes the proof of the equivalence.

Finally, suppose these conditions hold and  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  is finite-dimensional, where  $\mathbb{k}$  is now a field. It is clear that if  $x \in \mathfrak{m}$  and  $\mathfrak{m}^{\ell+1} = 0$ , then  $(1 - x)^{-1} = 1 + x + \cdots + x^\ell$ , and so  $n\mathcal{H}(W, \mathbf{d}, J_{<k})$  is local.  $\square$

## REFERENCES

- [1] N. Andruskiewitsch and H.-J. Schneider, *On the classification of finite-dimensional pointed Hopf algebras*, Annals of Mathematics 171(1):375–417, 2010.
- [2] J. Assion, *A proof of a theorem of Coxeter*, C. R. Math. Rep. Acad. Sci. Canada 1(1):41–44, 1978/79.
- [3] S. Billey, W. Jockusch, and R.P. Stanley, *Some combinatorial properties of Schubert polynomials*, Journal of Algebraic Combinatorics 2(4):345–374, 1993.
- [4] A. Bjorner and F. Brenti, *Combinatorics of Coxeter groups*, Vol. 231. Springer Science & Business Media, 2006.
- [5] G. Benkart and J. Meinel, *The center of the affine nilTemperley–Lieb algebra*, Mathematische Zeitschrift 284(1):413–439, 2016.
- [6] C. Berg, F. Saliola, and L. Serrano, *Pieri operators on the affine nilCoxeter algebra*, Transactions of the American Mathematical Society 366(1):531–546, 2014.
- [7] J.N. Bernstein, I.M. Gelfand, and S.I. Gelfand, *Schubert cells and cohomology of the spaces  $G/P$* , Russian Mathematical Surveys 28(2):1–26, 1973.
- [8] D. Bessis, *Zariski theorems and diagrams for braid groups*, Inventiones Mathematicae 145(3):487–507, 2001.
- [9] E. Brieskorn, *Die Fundamentalgruppe des Raumes der regulären Orbits einer endlichen komplexen Spiegelungsgruppe*, Inventiones Mathematicae 12(1):57–61, 1971.
- [10] H.S.M. Coxeter, *Discrete groups generated by reflections*, Annals of Mathematics 35(3):588–621, 1934.
- [11] H.S.M. Coxeter, *Factor groups of the braid group*, Proceedings of the 4th Canadian Mathematical Congress (Banff, 1957), University of Toronto Press, 95–122, 1959.
- [12] V.G. Drinfeld, *Degenerate affine Hecke algebras and Yangians*, Functional Analysis and its Applications 20(1):58–60, 1986.

- [13] P. Etingof and V. Ginzburg, *Symplectic reflection algebras, Calogero-Moser space, and deformed Harish-Chandra homomorphism*, *Inventiones Mathematicae* 147(2):243–348, 2002.
- [14] P. Etingof and O. Schiffmann, *Lectures on Quantum Groups*, *Lectures in Mathematical Physics*, International Press, Somerville, MA, 1998.
- [15] C.K. Fan, *A Hecke algebra quotient and properties of commutative elements of a Weyl group*, Ph.D. thesis, MIT, 1995.
- [16] C.K. Fan, *A Hecke algebra quotient and some combinatorial applications*, *Journal of Algebraic Combinatorics* 5(3):175–189, 1996.
- [17] C.K. Fan, *Structure of a Hecke algebra quotient*, *Journal of the American Mathematical Society* 10:139–167, 1997.
- [18] S. Fomin and C. Greene, *Noncommutative schur functions and their applications*, *Discrete Mathematics* 193(1–3):179–200, 1998.
- [19] S. Fomin and R.P. Stanley, *Schubert polynomials and the nil-Coxeter algebra*, *Advances in Mathematics* 103(2):196–207, 1994.
- [20] N. Gowravaram and T. Khovanova, *On the Structure of nil-Temperley-Lieb Algebras of type A*, preprint, arXiv:1509.00462.
- [21] J.J. Graham, *Modular representations of Hecke algebras and related algebras*, Ph.D. thesis, University of Sydney, 1995.
- [22] M. Graña, I. Heckenberger, and L. Vendramin, *Nichols algebras of group type with many quadratic relations*, *Advances in Mathematics* 227(5):1956–1989, 2011.
- [23] S. Hart, *How many elements of a Coxeter group have a unique reduced expression?*, *Journal of Group Theory* 20(5):903–910, 2017.
- [24] I. Heckenberger and L. Vendramin, *A classification of Nichols algebras of semi-simple Yetter-Drinfeld modules over non-abelian groups*, *Journal of the European Mathematical Society* 19(2):299–356, 2017.
- [25] I. Heckenberger and L. Vendramin, *The classification of Nichols algebras over groups with finite root system of rank two*, *Journal of the European Mathematical Society* 19(7):1977–2017, 2017.
- [26] J.E. Humphreys, *Reflection groups and Coxeter groups*, *Cambridge studies in advanced mathematics no. 29*, Cambridge University Press, Cambridge-New York-Melbourne, 1990.
- [27] A. Khare, *Generalized nil-Coxeter algebras, cocommutative algebras, and the PBW property*, *AMS Contemporary Mathematics* 688:139–168, 2017. (Special issue in honor of Donald S. Passman.)
- [28] A. Khare, *Generalized nil-Coxeter algebras over discrete complex reflection groups*, *Transactions of the American Mathematical Society* 370(4):2971–2999, 2018.
- [29] A. Khare, *Generalized nil-Coxeter algebras*, *Sem. Lothar. Combin.* 80B, art. #29, 12 pp., 2018.
- [30] M. Khovanov, *NilCoxeter algebras categorify the Weyl algebra*, *Communications in Algebra* 29(11):5033–5052, 2001.
- [31] M. Khovanov and A. Lauda, *A diagrammatic approach to categorification of quantum groups I*, *Representation Theory* 13:309–347, 2009.
- [32] T. Kildetoft, M. Mackaay, V. Mazorchuk, and J. Zimmermann, *Simple transitive 2-representations of small quotients of Soergel bimodules*, *Transactions of the American Mathematical Society* 371(8):5551–5590, 2019.
- [33] C. Korff and C. Stroppel, *The  $\widehat{\mathfrak{sl}}(n)_k$ -WZNW fusion ring: A combinatorial construction and a realisation as quotient of quantum cohomology*, *Advances in Mathematics* 225(1):200–268, 2010.
- [34] B. Kostant and S. Kumar, *The nil Hecke ring and cohomology of  $G/P$  for a Kac-Moody group  $G$* , *Advances in Mathematics* 62(3):187–237, 1986.
- [35] A. Lascoux and M.-P. Schützenberger, *Fonctorialité des polynômes de Schubert*, *Invariant Theory (R. Fossum, W. Haboush, M. Hochster and V. Lakshmibai, Eds.)*, *Contemporary Mathematics Volume 88*, pages 585–598, American Mathematical Society, Providence, 1989.
- [36] G. Lusztig, *Affine Hecke algebras and their graded version*, *Journal of the American Mathematical Society* 2:599–635, 1989.
- [37] L. Paris, *Artin monoids inject into their groups*, *Commentarii Mathematici Helvetici* 77(3):609–637, 2002.
- [38] A. Postnikov, *Affine approach to quantum Schubert calculus*, *Duke Mathematical Journal* 128(3):473–509, 2005.
- [39] R. Simion, *Combinatorial statistics on type-B analogues of noncrossing partitions and restricted permutations*, *Electronic Journal of Combinatorics* 7(2), art. #R9, 27 pp., 2000.
- [40] G.C. Shephard and J.A. Todd, *Finite unitary reflection groups*, *Canadian Journal of Mathematics* 6:274–304, 1954.

- [41] A.V. Shepler and S. Witherspoon, *A Poincaré–Birkhoff–Witt theorem for quadratic algebras with group actions*, Transactions of the American Mathematical Society 366(12):6483–6506, 2014.
- [42] A.V. Shepler and S. Witherspoon, *Poincaré–Birkhoff–Witt Theorems*, Commutative Algebra and Non-commutative Algebraic Geometry, Volume I: Expository Articles (D. Eisenbud, S.B. Iyengar, A.K. Singh, J.T. Stafford, and M. Van den Bergh, Eds.), Mathematical Sciences Research Institute Proceedings Volume 67, pages 259–290, Cambridge University Press, Cambridge, 2015.
- [43] N.J.A. Sloane, *The On-line Encyclopedia of Integer Sequences*, Sequence A014143.
- [44] J.R. Stembridge, *On the fully commutative elements of Coxeter groups*, Journal of Algebraic Combinatorics 5(4):353–385, 1996.
- [45] J.R. Stembridge, *Some combinatorial aspects of reduced words in finite Coxeter groups*, Transactions of the American Mathematical Society 349(4):1285–1332, 1997.
- [46] J.R. Stembridge, *The enumeration of fully commutative elements of Coxeter groups*, Journal of Algebraic Combinatorics 7(3):291–320, 1998.
- [47] C. Walton and S. Witherspoon, *Poincaré–Birkhoff–Witt deformations of smash product algebras from Hopf actions on Koszul algebras*, Algebra & Number Theory 8(7):1701–1731, 2014.

(S. Bhattacharya) INDIAN INSTITUTE OF SCIENCE BANGALORE – 560012, INDIA; B.S.(MATH) STUDENT  
*Email address:* [sutanayb@iisc.ac.in](mailto:sutanayb@iisc.ac.in)

(A. Khare) DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE, INDIA AND  
ANALYSIS & PROBABILITY RESEARCH GROUP, BANGALORE, INDIA  
*Email address:* [khare@iisc.ac.in](mailto:khare@iisc.ac.in)