

Fast Goodstein Walks

David Fernández-Duque and Andreas Weiermann

December 1, 2021

Abstract

We define a variant of the Goodstein process based on fast-growing functions and show that it terminates, but this fact is not provable in Kripke-Platek set theory or other theories of strength the Bachmann-Howard ordinal. We moreover show that this Goodstein process is of maximal length, so that any alternative Goodstein process based on the same fast-growing functions will also terminate.

1 Introduction

Goodstein's classic principle [9] is perhaps the oldest example of a statement unprovable in Peano arithmetic (PA) [11] which does not involve direct coding of metamathematical objects. The proof itself does use the well-foundedness of ε_0 [8], the proof-theoretic ordinal of PA. Goodstein's principle is based on *hereditary exponential normal forms*, whereby each natural number is written in base k in the standard way, as is each exponent appearing in the expansion, and so forth. Thus for example we may write $18 = 2^{2^2} + 2$ in base-2 hereditary exponential normal form.

The Goodstein process then proceeds as follows: given a natural number m_0 , write m_0 in base-2 hereditary exponential normal form. Then, compute m_1 by replacing every 2 appearing in the normal form of m_0 by 3, then subtract one. We continue inductively in this fashion, defining m_{k+1} by writing m_k in base- $(k+2)$ normal form, replacing $k+2$ by $k+3$, and subtracting 1. The operation of replacing every instance of $k+2$ by $k+3$ is an instance of the *base change operation*. The process terminates if $m_\ell = 0$ for some ℓ : Goodstein's principle states that such an ℓ always exists.

One may wonder if the use of normal forms is essential. For example, we can write $18 = 2^{2+1} + 2^2 + 2^2 + 2$. This begs the question: will Goodstein processes terminate no matter how we write natural numbers? Surprisingly, the answer is *yes*. A *Goodstein walk* is a sequence m_0, m_1, \dots where each m_{k+1} is defined by writing m_k using *any* expression formed from $0, +, (k+2)^x$, then changing the base from $k+2$ to $k+3$ and subtracting one. The authors have recently shown that every Goodstein walk is finite [7], using the fact that hereditary exponential normal forms are *base change maximal*, meaning that applying

the base change operator to the normal form of m yields the maximal value among all the possible terms for m . This implies that Goodstein processes using such normal forms have maximal length, hence since these processes are finite, so is any Goodstein walk.

Goodstein's theorem may also be extended by considering notations based on functions other than the exponential. A parametrized version of the Ackermann function gives rise to independence results for theories between PA and the second-order arithmetical theory ATR_0 of arithmetical transfinite recursion [1]. This process is based on a notion of normal form based on a 'sandwiching' procedure, but other natural notions of normal forms can be considered [6]. More generally, Goodstein walks are also naturally defined in terms of the Ackermann function. The sandwiching normal forms are base-change maximal in this setting, meaning that every Ackermannian Goodstein walk is finite [7].

For writing much bigger numbers, one may switch to notation using even faster-growing functions. The ordinal ε_0 can be used to define very fast-growing functions by transfinite recursion. There are several prominent examples of this, e.g. the Hardy function [10]. Goodstein principles based on fast-growing functions [2] give rise to independence from theories with strength the Bachmann-Howard ordinal, including Kripke-Platek set theory (KP) (see e.g. [12]).

Goodstein walks may also be considered in the context of fast-growing functions. Our main goal is to show that these walks are finite, a fact independent of KP. We will use a function that grows slightly faster than standard fast-growing hierarchies and denote it $\mathbb{A}_k(\xi)$, where k is the number of iterations and $\xi < \varepsilon_0$ is an ordinal. The precise definition will be given in Section 3. Note that Peano arithmetic does not prove that these functions are total, but KP does (as do much weaker theories).

Let us give a brief description of KP; a more formal treatment can be found in e.g. [3]. The theory KP (with infinity) is axiomatized by all axioms of ZFC except for powerset, but with separation restricted to Δ_0 formulas and replacement restricted to Δ_0 -collection. This theory, while much more powerful than Peano arithmetic, does not prove that the real line exists as a set. It proves the same arithmetical formulas as other well-known theories such as the theory ID_1 of non-iterated inductive definitions, and $\Pi_1^1\text{-CA}_0^-$ of parameter-free Π_1^1 comprehension. Precise details of these theories are not needed, except that their proof-theoretic ordinal is $\vartheta(\varepsilon_{\Omega+1})$ (also denoted $\psi(\varepsilon_{\Omega+1})$), where Ω is the first uncountable ordinal and $\vartheta: \varepsilon_{\Omega+1} \rightarrow \Omega$ is a function that transforms possibly uncountable ordinals into countable ones. Most relevant to us is that this ordinal and its system of fundamental sequences can be used to bound the provably total computable functions of any of these three theories. This allows us to prove that the Goodstein principle based on the \mathbb{A}_k functions is also unprovable, by showing that the process reaches zero more slowly than stepping down these fundamental sequences.

2 Ordinals below $\varepsilon(\kappa)$

In this section, we review some elementary notions from ordinal arithmetic. We assume basic familiarity with ordinal addition, multiplication, and exponentiation. The successor of α will be denoted $\alpha + 1$, when it exists. We will use the following simple inequalities, proven by routine induction.

Lemma 2.1. *Let α, β be ordinals.*

1. *If $\alpha, \beta > 1$ then $\alpha + \beta \leq \alpha\beta$.*
2. *If $\alpha > 1$ then $\beta \leq \alpha^\beta$.*

Throughout the text we will use normal forms for ordinals based on either ω or Ω , the first uncountable ordinal. Let $\kappa \in \{\omega, \Omega\}$ and $\xi > 0$ be an ordinal. There exist unique ordinals α, β, γ with $\beta < \kappa$ such that $\xi = \kappa^\alpha \beta + \gamma$ and $\gamma < \kappa^\alpha$. This is the κ -normal form of ξ . Note that the ω -normal form of ξ is not precisely its Cantor normal form, as Cantor normal forms do not involve coefficients. For our purposes, we simply say that an expression $\alpha + \beta$ is in *Cantor normal form* if $\alpha + \beta > \alpha' + \beta$ for all $\alpha' < \alpha$.

The ordinal $\varepsilon(\kappa)$ is defined as the least $\varepsilon > \kappa$ such that $\varepsilon = \omega^\varepsilon$. If $\xi = \kappa^\alpha \beta + \gamma < \varepsilon(\kappa)$ is in κ -normal form, then we may also deduce that $\alpha < \xi$. We write $\varepsilon_0 = \varepsilon(\omega)$ and $\varepsilon_{\Omega+1} = \varepsilon(\Omega)$ as is standard. Define $\kappa_0(\alpha) = \alpha$ and $\kappa_{i+1}(\alpha) = \kappa^{\kappa_i(\alpha)}$. We set $\kappa_i = \kappa_i(1)$. Then, $\varepsilon(\kappa) = \sup_{n < \omega} \kappa_i$.

Next we define fundamental sequences. We will reserve the more standard notation $\xi[n]$ for arbitrary systems of fundamental sequences, whereas the notation $[\xi]_n$ refers exclusively to the operation defined below.

Definition 2.2. *Let κ be an ordinal and $\varepsilon = \varepsilon(\kappa)$ the least ε -number above κ . We define fundamental sequences for ordinals below ε and $\theta < \kappa$ recursively as follows.*

1. $[0]_\kappa = [1]_\kappa = 0$,
2. $[\kappa^\alpha \beta + \gamma]_\kappa = \kappa^\alpha \beta + [\gamma]_\kappa$ if $\gamma > 0$,
3. $[\kappa^\alpha \beta]_\kappa = \kappa^\alpha \theta$ if β is a limit,
4. $[\kappa^{\alpha+1}]_\kappa = \kappa^\alpha \theta$,
5. $[\kappa^\alpha(\beta + 1)]_\kappa = \kappa^\alpha \beta + [\kappa^\alpha]_\kappa$ if $\beta > 0$, and
6. $[\kappa^\alpha]_\kappa = \kappa^{[\alpha]_\kappa}$.

We write simply $[\alpha]_\kappa$ when $\kappa = \omega$. It is not hard to check that $[\alpha + \beta]_\kappa = \alpha + [\beta]_\kappa$ whenever $\alpha + \beta$ is in Cantor normal form. We define the set of coefficients of α in coefficient κ -normal form by $C_\kappa(0) = \{0\}$, $C_\kappa(\kappa^\alpha \theta + \beta) = C_\kappa(\alpha) \cup C_\kappa(\beta) \cup \{\theta\}$. We define the maximal coefficient of α by $\text{mc}_\kappa(\alpha) = \max C(\alpha)$, and omit the subindex κ when $\kappa = \omega$. The *terminal part* of ξ , denoted $\tau_\kappa(\xi)$ or $\tau(\xi)$ when clear from context, is given recursively by

1. $\tau(0) = 0$ and $\tau(\zeta + 1) = 1$,

2. $\tau(\kappa^\alpha\beta + \gamma) = \tau(\gamma)$ if $\gamma > 0$,
3. $\tau(\kappa^\alpha\beta) = \beta$ if β is a limit,
4. $\tau(\kappa^\alpha(\beta + 1)) = \tau(\alpha)$ if α is a limit, and
5. $\tau(\kappa^{\alpha+1}(\beta + 1)) = \kappa$.

Then it is readily checked that if $\tau = \tau(\xi)$ is a limit and $\theta < \tau$ then $[\xi|\theta]_\kappa < \xi$, and moreover $\xi = \lim_{\theta \rightarrow \tau} [\xi|\theta]_\kappa$. Note however that we have defined $[\xi|\theta]_\kappa$ even when $\theta \geq \tau$, but this is mostly because it is convenient to have $[\xi + 1|\theta]_\kappa = \xi$ for all θ .

The following is checked by an easy induction on ξ .

Lemma 2.3. *If $\xi < \varepsilon(\kappa)$ is a limit and $\theta < \kappa$ then*

$$\theta \leq \text{mc}([\xi|\theta]) \leq \max\{\text{mc}(\xi), \theta\}.$$

Lemma 2.4. *Let $\lambda < \varepsilon(\kappa)$ be a limit ordinal.*

1. *If $\xi < \lambda$ and $\text{mc}_\kappa(\xi) < \theta$ then $\xi < [\lambda|\theta]_\kappa$.*
2. *Whenever $\theta < \eta < \kappa$ it follows that $[\lambda|\theta]_\kappa < [\lambda|\eta]$.*

The fundamental sequences admit a sort of left inverse, given by the following operation.

Definition 2.5. *Define $[\xi]_\kappa$ where, $\xi = \omega^\alpha\beta + \gamma$ in κ -normal form, by*

1. $[0]_\kappa = 0$,
2. $[\xi]_\kappa = \kappa^\alpha\beta + [\gamma]_\kappa$ if $\gamma > 0$
3. $[\omega^\alpha\beta]_\kappa = \omega^{\alpha+1}$ if $\beta > 1$
4. $[\omega^\alpha]_\kappa = \omega^{[\alpha]_\kappa}$.

As before, we omit the subindex when $\kappa = \omega$.

Lemma 2.6. *If $\theta \in (1, \kappa)$ and ξ is a limit with $\tau(\xi) = \kappa$ then $[[\xi|\theta]_\kappa]_\kappa = \xi$.*

Proof. We sketch the proof, which proceeds by induction on ξ . The critical case is where $\xi = \kappa^\alpha$. If α is a limit then $[\kappa^\alpha|\theta]_\kappa = \kappa^{[\alpha|\theta]_\kappa}$, so that $[[\kappa^\alpha|\theta]_\kappa]_\kappa = \kappa^{[[\alpha|\theta]_\kappa]_\kappa}$. By induction hypothesis, $[[\alpha|\theta]_\kappa]_\kappa = \alpha$, so that $[[\kappa^\alpha|\theta]_\kappa]_\kappa = \kappa^\alpha$. Otherwise, $\alpha = \alpha' + 1$ and $[\kappa^{\alpha'+1}|\theta]_\kappa = \kappa^{\alpha'}\theta$. So, $[[\kappa^{\alpha'+1}|\theta]_\kappa]_\kappa = \kappa^{\alpha'+1}$. \square

As a corollary, we obtain that fundamental sequences are injective in the following sense.

Corollary 2.7. *If $\tau_\kappa(\alpha) = \tau_\kappa(\alpha') = \kappa$ and $\theta, \theta' \in (1, \kappa)$ are such that $[\alpha|\theta]_\kappa = [\alpha'|\theta']_\kappa$, then $\alpha = \alpha'$.*

Proof. By Lemma 2.6, we have that $\alpha = [[\alpha|\theta]_\kappa]_\kappa = [[\alpha'|\theta']_\kappa]_\kappa = \alpha'$. \square

Next we turn our attention to $\kappa = \omega$. Here, the *Bachmann property* holds for the system of fundamental sequences. It is convenient to define this notion with some generality.

Definition 2.8. Let Λ be a countable ordinal. A system of fundamental sequences on Λ is a function $[\cdot]: \Lambda \times \mathbb{N} \rightarrow \Lambda$ such that

1. $\alpha[n] \leq \alpha$ with equality holding if and only if $\alpha = 0$,
2. $\alpha[n] < \alpha[m]$ whenever $n < m$, and
3. $\lambda = \lim_{n \rightarrow \infty} \alpha[n]$ whenever λ is a limit.

The system of fundamental sequences has the Bachmann property if whenever $\alpha[n] < \beta < \alpha$, it follows that $\alpha[n] \leq \beta[1]$.

The fundamental sequences we have defined are known to enjoy the Bachmann property [?].

Lemma 2.9. If $\lambda < \varepsilon_0$ is a limit and $\xi \in ([\lambda|n], \lambda)$, it follows that $[\lambda|n] \leq [\xi|1]$.

Finally, we want to observe that if $[\beta|q] \leq \alpha \leq \beta$, then we can obtain information about the coefficients of β from those of α . To this end, we define a *truncation* of α to be any $\beta \leq \alpha$ such that $C(\beta) \subseteq \{0, 1\} \cup C(\alpha)$.

Lemma 2.10. Let $\alpha, \beta < \varepsilon_0$ with β a limit. If $[\beta|q] \leq \alpha \leq \beta$, there is a truncation α' of α and some $q' \geq q$ such that $\alpha' = [\beta|q']$.

Proof. By induction on β . Write $\beta = \omega^{\beta_1} b + \beta_0$ and $\alpha = \omega^{\alpha_1} a + \alpha_0$ in ω -normal form and consider the following cases.

CASE 1 ($\beta_0 > 0$). Then we must have $\alpha = \omega^{\beta_1} b + \alpha_0$ with $[\beta_0|q] \leq \alpha_0 < \beta_0$, which by the induction hypothesis yields a truncation γ of α_0 and $q' > q$ such that $\gamma = [\beta_0|q']$. The desired truncation of α is then $\omega^{\beta_1} b + \gamma$.

CASE 2 ($\beta_0 = 0$). Consider the following sub-cases.

CASE 2.1 ($b > 1$). Then $\alpha_1 = \beta_1$ and $b = a + 1$. Moreover, $[\omega^{\beta_1}|q] \leq \alpha_0 < \omega^{\beta_1}$, yielding a truncation γ of α_0 and $q' > q$ such that $\gamma = [\omega^{\beta_1}|q']$. Then, $\omega^{\alpha_1} a + \gamma$ is the desired truncation of α .

CASE 2.2 ($b = 1$). We consider two further sub-cases.

CASE 2.2.1 ($\beta_1 = \delta + 1$). Then, $[\beta|q] = \omega^\delta q$, and from $[\beta|q] \leq \alpha < \beta$ we see that $\alpha = \omega^{\beta_1} q' + \alpha_0$ for some $q' \geq q$. In this case, $\omega^{\beta_1} q'$ is a truncation of α and $\omega^{\beta_1} q' = [\beta|q']$.

CASE 2.2.2 (β_1 is a limit). We have that $[\omega^{\beta_1}|q] \leq \alpha < \omega^{\beta_1}$, so that $[\omega^{\beta_1}|q] \leq \alpha_1 < \omega^{\beta_1}$. Let γ and q' be the truncation and number given by the induction hypothesis for α_1 . Then, ω^γ is the corresponding truncation of α . \square

3 Parametrized fast-growing hierarchies

We may use fundamental sequences to define very large natural numbers in terms of ordinals below ε_0 . We introduce a version with an extra parameter k , which will serve as the ‘base’ in our fast Goodstein walks.

Definition 3.1. For $2 \leq \lambda \leq \omega$, we define $\mathbb{A}_\lambda: \varepsilon(\lambda^+) \rightarrow \lambda^+$ as follows. First, introduce the abuse of notation $\mathbb{A}_k(\alpha - 1) = \text{mc}(\alpha)$, when α is a limit or zero ($\alpha - 1$ itself remains undefined when α is a limit). Suppose inductively that $\mathbb{A}_\lambda(\xi)$ is defined for all $\xi < \alpha$ and define $\mathbb{A}_\lambda(\xi)$ according to the following cases.

CASE 1 ($\lambda < \omega$). We divide in two sub-cases.

CASE 1.1 ($\xi < \omega$). Set $\mathbb{A}(\xi) = \xi + 1$.

CASE 1.2 ($\xi \geq \omega$). Write $\xi = \alpha + b$ with α a limit. Define $\mathbb{A}_\lambda^{(i)}(\xi)$ recursively by

- i. $\mathbb{A}_\lambda^{(0)}(\xi) = \mathbb{A}_\lambda(\alpha - 1)$ and
- ii. $\mathbb{A}_\lambda^{(i+1)}(\xi) = \mathbb{A}_\lambda([\alpha | \mathbb{A}_\lambda^{(i)}(\xi)])$.

Then, set $\mathbb{A}_k(\xi) = \mathbb{A}_k^{(k)}(\xi)$.

CASE 1.3 ($\lambda = \omega$). $\mathbb{A}_\omega(\xi)$ is inductively defined to be the least $\theta > \text{mc}_\Omega(\xi)$ such that if $\zeta < \xi$ and $\text{mc}_\Omega(\zeta) < \theta$ then $\mathbb{A}_\omega(\zeta) < \theta$. We further define $\mathbb{A}_\omega^{(i)}(\xi) = \mathbb{A}_\omega(\xi)$ for all i .

We remark that \mathbb{A}_ω is a variant of the ϑ function [4], except that $\mathbb{A}_\omega(\xi)$ may take values that are not principal numbers. In particular, the proof that \mathbb{A}_ω is well-defined is essentially identical to the proof that ϑ is well-defined. The main characterization we use of the \mathbb{A}_ω function is the following.

Proposition 3.2. If $\zeta, \xi < \varepsilon_{\Omega+1}$, then $\mathbb{A}_\omega(\zeta) < \mathbb{A}_\omega(\xi)$ if and only if $\zeta < \xi$ and $\text{mc}_\Omega(\zeta) < \mathbb{A}_\omega(\xi)$.

Throughout the paper often write $\mathbb{A}(\xi) = \mathbb{A}_\lambda(\xi)$ whenever λ is fixed and clear from context. For $k \in [2, \omega)$ and $\xi = \alpha + b \in (0, \varepsilon_0)$ with α a limit, we define $[\xi]_k = [\alpha | \mathbb{A}^{(k-1)}(\xi)]$. It is readily checked that Definition 3.1 yields

$$\mathbb{A}_k(\xi) = \mathbb{A}_k([\xi]_k) \tag{1}$$

for all ξ .

The values $\mathbb{A}_k(\xi)$ grow rather quickly, but the following simple lower bound will be useful to us. Recall that if α is zero or a limit, then $\mathbb{A}(\alpha - 1) = \text{mc}(\alpha)$ by definition.

Lemma 3.3. Let $k \in [2, \omega)$ and $\xi < \varepsilon_0$. Write \mathbb{A} for \mathbb{A}_k

1. If $\xi > \omega$ and $i < j < \omega$, then $\mathbb{A}(\xi - 1) \leq \mathbb{A}^{(i)}(\xi) < \mathbb{A}^{(j)}(\xi)$.
2. $\text{mc}(\xi) \leq \mathbb{A}(\xi - 1) < \mathbb{A}(\xi)$.

3. If $\xi > \omega$ and $k \in [2, \omega)$, then $\mathbb{A}_k(\xi) > 2\text{mc}(\xi) + 2$.

Proof. Proceed to prove the three claims simultaneously by induction on $\xi = \alpha + b$, where α is zero or a limit.

Proof of Claim 1. We have that $\mathbb{A}^{(0)}(\xi) = \mathbb{A}(\xi - 1)$, and induction hypothesis applied to $[\alpha|\mathbb{A}^{(i)}(\xi)] < \xi$ yields

$$\mathbb{A}^{(i+1)}(\xi) = \mathbb{A}([\alpha|\mathbb{A}^{(i)}(\xi)]) > \text{mc}([\alpha|\mathbb{A}^{(i)}(\xi)]) \geq \mathbb{A}^{(i)}(\xi).$$

Induction on i, j then yields the claim.

Proof of Claim 2. If $\xi < \omega$, $\xi = \text{mc}(\xi) = \mathbb{A}(\xi - 1)$ and $\mathbb{A}(\xi) = \xi + 1 > \xi$. Otherwise, α is a limit. That $\text{mc}(\xi) \leq \mathbb{A}(\xi - 1)$ follows by definition when $b = 0$, otherwise we may apply the induction hypothesis to the third claim to see that $\text{mc}(\xi) < 2\text{mc}(\xi - 1) + 2 < \mathbb{A}(\xi - 1)$. That $\mathbb{A}(\xi - 1) < \mathbb{A}(\xi)$ follows from direct computation when $\xi = \omega$, as $\mathbb{A}(\xi - 1) = 1 < k + 1 = \mathbb{A}(\omega)$. If $\xi > \omega$, the first claim yields $\mathbb{A}(\xi - 1) = \mathbb{A}^{(0)}(\xi) \leq \mathbb{A}^{(k)}(\xi) = \mathbb{A}(\xi)$.

Proof of Claim 3. Consider two cases.

CASE 1 ($\alpha = \omega$). Proceed by induction on b . If $b = 1$ then $\mathbb{A}^{(0)}(\xi) = \mathbb{A}(\omega) = k + 1 \geq 3$, so inductively $\mathbb{A}^{(i)}(\xi) \geq 3 + i$ yielding $\mathbb{A}(\xi) = \mathbb{A}^{(k)}(\xi) = 3 + k \geq 5 = 2\text{mc}(\xi) + 3$. Otherwise, the induction hypothesis yields $\mathbb{A}^{(0)}(\xi) \geq 2(b - 1) + 3 = 2b + 1$, so $\mathbb{A}^{(k)}(\xi) \geq 2b + k + 1 \geq 2b + 3 = 2\text{mc}(\xi) + 3$.

CASE 2 ($\alpha > \omega$). We have by the second claim that $\mathbb{A}^{(1)}(\xi) > \text{mc}(\xi) \geq 1 = \text{mc}(\omega)$. Since $\alpha > \omega$ and $\mathbb{A}^{(1)}(\xi) > \text{mc}(\omega)$, we have that $[\alpha|\mathbb{A}^{(1)}(\xi)] > \omega$, so we may apply the induction hypothesis to conclude that

$$\mathbb{A}(\xi) \geq \mathbb{A}^{(2)}(\xi) = \mathbb{A}([\alpha|\mathbb{A}^{(1)}(\xi)]) > 2\text{mc}([\alpha|\mathbb{A}^{(1)}(\xi)]) + 2 > 2\text{mc}(\xi) + 2. \quad \square$$

Note that $\mathbb{A}_\omega^{(i)}(\zeta)$ has not been defined. We set $\mathbb{A}_\omega^{(i)}(\zeta) = \mathbb{A}_\omega(\zeta)$ for all i , which allows us to give a uniform statement in the following lemma.

Lemma 3.4. *Let $2 \leq k < \lambda \leq \omega$ and $\alpha < \beta < \varepsilon(\lambda^+)$. If $\text{mc}_{\lambda^+}(\alpha) < \mathbb{A}_\lambda^{(k-1)}(\beta)$, then $\mathbb{A}_\lambda(\alpha) < \mathbb{A}_\lambda(\beta)$.*

Proof. By induction on β . First assume $\lambda < \omega$. Write $\beta = \lambda + t$ for λ a limit. If $\alpha = \lambda + t'$ for some natural number t' , then $t' < t$ and the claim is established by an easy induction on t . Otherwise, using Lemma 2.4.1, we see that $\alpha \leq \lfloor \beta \rfloor_k$, and moreover Lemma 3.3.1 yields

$$\mathbb{A}^{(k-1)}(\lfloor \beta \rfloor_k) > \text{mc}(\lfloor \beta \rfloor_k) \geq \mathbb{A}^{(k-1)}(\beta) > \text{mc}(\alpha).$$

Thus we may apply the induction hypothesis to see that $\mathbb{A}(\alpha) < \mathbb{A}(\lfloor \beta \rfloor_k) = \mathbb{A}(\beta)$.

If $\lambda = \omega$, this simply repeats the characterization we have given for establishing $\mathbb{A}_\omega(\alpha) < \mathbb{A}_\omega(\beta)$, recalling that we have set $\mathbb{A}^{(k)}(\beta) = \mathbb{A}(\beta)$. \square

Lemma 3.5. Fix $k \geq 0$ and let $\mathbb{A} = \mathbb{A}_k$. Let $\lambda < \varepsilon_0$ be a limit. Then, if $[\lambda|q] < \zeta < \lambda$, it follows that $\mathbb{A}(\zeta) > \mathbb{A}([\lambda|q])$.

Proof. Write $\zeta = \alpha + d$ and $[\lambda|q] = \beta + c$ with α, β limits or zero and proceed by induction on ζ . Note that either $\alpha = \beta$ or else $[\lambda|q] < \alpha < \lambda$, from which it follows using the Bachmann property that $[\lambda|q] \leq [\alpha|1] < [\zeta]_k$. With this in mind, consider two cases.

CASE 1 ($\alpha = \beta$). Then,

$$\alpha + c = [\lambda|q] < \zeta = \alpha + d,$$

so $c < d$ and $\mathbb{A}(\alpha + c) < \mathbb{A}(\alpha + d)$ by Lemma 3.4.

CASE 2 ($[\zeta]_k > [\lambda|q]$). In this case, $[\lambda|q] < [\zeta]_k < \zeta < \lambda$, so we see by the induction hypothesis that

$$\mathbb{A}([\lambda|q]) < \mathbb{A}([\zeta]_k) = \mathbb{A}(\zeta). \quad \square$$

Corollary 3.6. Fix $k \geq 0$ and let $\mathbb{A} = \mathbb{A}_k$. Let $\xi = \lambda + b < \varepsilon_0$ with λ a limit. Then, if $[\xi]_k < \zeta < \lambda$, it follows that $\mathbb{A}(\zeta) > \mathbb{A}(\xi)$.

Proof. Note that $[\xi]_k = [\lambda|q]$ with $q = \mathbb{A}^{(k-1)}(\xi)$. Thus Lemma 3.5 yields $\mathbb{A}(\xi) = \mathbb{A}([\xi]_k) < \mathbb{A}(\zeta)$. \square

Lemma 3.7. Given $\xi < \zeta < \varepsilon_0$, $\mathbb{A}(\zeta) \notin (\mathbb{A}(\xi), \mathbb{A}(\xi + 1))$.

Proof. Write $\xi = \alpha + a$ and $\zeta = \beta + b$ with α, β limits. Proceed by induction on ζ and consider three cases.

CASE 1 ($\xi \geq \beta$). Then, since $\xi < \zeta$ we have that $\alpha = \beta$ and $a < b$, so that $\mathbb{A}(\xi + 1) = \mathbb{A}(\alpha + a + 1) \leq \mathbb{A}(\alpha + b) = \mathbb{A}(\zeta)$.

CASE 2 ($[\zeta]_k \leq \xi$). By Corollary 3.6, $\mathbb{A}(\zeta) \leq \mathbb{A}(\xi)$.

CASE 3 ($[\zeta]_k > \xi$). In this case the claim is immediate from the induction hypothesis. \square

4 Normal forms based on the fast-growing hierarchy

In order to define maximal Goodstein processes for the \mathbb{A} functions, we must identify base-change maximal normal forms. We define them below, with maximality proven in subsequent sections. Below, for an expression $\varphi(\xi)$ we set $\xi_* = \nu\xi.\varphi$ if either ξ_* is the maximum element of ε_0 satisfying $\varphi(\xi_*)$, or $\xi_* = 0$ and no such maximum exists.

Definition 4.1. Fix $k \in [2, \omega)$ and let $\mathbb{A}(\xi) = \mathbb{A}_k(\xi)$. Given $m \in \mathbb{N}$, we define $\mathbb{A}(\xi)$ to be the k -normal form of m , in symbols $m =_k \mathbb{A}(\xi)$, if $m = \mathbb{A}(\xi)$ and there exist sequences $\xi_0, \xi_1, \dots, \xi_n$ such that:

1. $\xi_0 = 0$.
2. Given $i < n$, $\xi_{i+1} = \nu\zeta \cdot (\mathbb{A}(\zeta) \leq m \text{ and } \text{mc}(\zeta) \geq \mathbb{A}(\xi_i))$.
3. $\mathbb{A}(\xi_n) = m$.

We call $(\xi_i)_{i \leq n}$ the normal form sequence of m .

Let us introduce some notation for normal form sequences. Suppose that m has normal form sequence $(\mathbb{A}(\xi_i))_{i \leq n}$. Then, we define $\dot{m} = m_{n-1}$, and $\dot{\xi} = \xi_{n-1}$. It is easy to see that $\dot{m} =_k \mathbb{A}(\dot{\xi})$, with the same normal form sequence as m but truncated at the second to last element.

We often need results of the form *If $\mathbb{A}(\xi)$ is in normal form, then $\mathbb{A}(\zeta)$ is also in normal form*, where ζ is ‘similar’ to ξ in some way. The following general principle will be useful for establishing this.

Lemma 4.2. *If $\mathbb{A}(\xi)$ is in normal form and ζ is such that*

- (a) $\mathbb{A}(\zeta) < \mathbb{A}(\xi + 1)$,
- (b) $\mathbb{A}(\xi) \leq \text{mc}(\zeta)$, and
- (c) *if θ is such that $\text{mc}(\zeta) \geq \mathbb{A}(\theta)$ and $\mathbb{A}(\theta) \leq \mathbb{A}(\zeta)$, then $\theta \leq \zeta$,*

then $\mathbb{A}(\zeta)$ is in normal form.

Proof. Let $(\xi_n)_{i \leq n}$ be the normal form sequence for $\mathbb{A}(\xi)$. Note that $\mathbb{A}(\xi) \leq \text{mc}(\zeta) < \mathbb{A}(\zeta) < \mathbb{A}(\xi + 1)$ implies that $\xi \geq \omega$ (otherwise $\mathbb{A}(\xi + 1) = \mathbb{A}(\xi) + 1$), so $n > 0$. Note also that the maximality of ξ_i yields $\xi_{i+1} \leq \xi_i$ for $0 < i < n$, and $\xi_{i+1} = \xi_i$ is impossible since $\mathbb{A}(\xi_{i+1}) > \text{mc}(\xi_{i+1}) \geq \mathbb{A}(\xi)$. It follows that $\xi_{i+1} > \xi_n$ for all $i < n - 1$.

Let $(\zeta_r)_{i \leq r}$ be the normal form sequence for $\mathbb{A}(\zeta)$. We claim that $r = n + 1$, $\zeta_i = \xi_i$ for $i \leq n$, and $\zeta_r = \zeta$, witnessing that $\mathbb{A}(\zeta)$ is in normal form. If $i < n$, $\mathbb{A}(\xi_{i+1}) \leq \mathbb{A}(\xi) < \mathbb{A}(\zeta)$. If $\delta > \xi_{i+1}$ and $\text{mc}(\delta) \geq \mathbb{A}(\xi_i)$, the maximality of ξ_{i+1} yields $\mathbb{A}(\delta) > \mathbb{A}(\xi)$. Moreover, $\delta > \xi_{i+1} > \xi$, so Lemma 3.7 yields $\mathbb{A}(\delta) \geq \mathbb{A}(\xi + 1) > \mathbb{A}(\zeta)$. Thus by induction on i we see that $\zeta_i = \xi_i$, since ξ_{i+1} is maximal with $\mathbb{A}(\xi_{i+1}) \leq \mathbb{A}(\zeta)$ and $\text{mc}(\xi_{i+1}) \geq \mathbb{A}(\xi_i) \stackrel{\text{IH}}{=} \mathbb{A}(\zeta_i)$. Conditions (b) and (c) then guarantee that $\zeta_{n+1} = \zeta$, as needed. \square

Below we provide some applications of this general lemma.

Lemma 4.3. *If ξ is a successor and $m = \mathbb{A}(\xi)$ is in normal form, then either*

- (a) $n = \mathbb{A}(\xi - 1)$ *is in normal form, or*
- (b) $\xi = \alpha + \mathbb{A}(\dot{\xi})$, *where α is a limit or zero and $\text{mc}(\alpha) < \mathbb{A}(\dot{\xi})$.*

Proof. We have that $\mathbb{A}(\xi - 1) < \mathbb{A}(\xi) < \mathbb{A}(\dot{\xi} + 1)$, and if $\mathbb{A}(\theta) \leq \mathbb{A}(\xi - 1)$ and $\text{mc}(\theta) \geq \mathbb{A}(\dot{\xi})$, then we also have that $\mathbb{A}(\theta) \leq \mathbb{A}(\xi)$, which by maximality of ξ implies that $\theta \leq \xi$. But $\mathbb{A}(\xi) > \mathbb{A}(\xi - 1)$, so $\theta \neq \xi$, hence $\theta \leq \xi - 1$. Thus $\mathbb{A}(\xi - 1)$ is in normal form unless $\text{mc}(\xi - 1) < \mathbb{A}(\dot{\xi})$. The only way to have that $\text{mc}(\xi) \geq \mathbb{A}(\dot{\xi})$ but $\text{mc}(\xi - 1) < \mathbb{A}(\dot{\xi})$ is that $\xi = \alpha + \mathbb{A}(\dot{\xi})$ with $\text{mc}(\alpha) < \mathbb{A}(\dot{\xi})$, as needed. \square

Lemma 4.4. *Let $k \in [2, \omega)$ and write \mathbb{A} for \mathbb{A}_k . Suppose that $\mathbb{A}(\xi)$ is in normal form and let $i < k - 1$. Then, $\mathbb{A}([\xi | \mathbb{A}^{(i)}(\xi)])$ is in normal form.*

Proof. Let $m = \mathbb{A}(\xi)$, $q = \mathbb{A}^{(i)}(\xi)$, and $m' = \mathbb{A}([\xi | \mathbb{A}^{(i)}(\xi)])$. Consider two cases.

CASE 1 ($b = 0$ or $\text{mc}(\xi - 1) < \dot{m}$). In order to apply Lemma 4.2, we note that $\mathbb{A}([\alpha | q]) < \mathbb{A}(\xi) < \mathbb{A}(\dot{\xi} + 1)$, where the first inequality uses Lemma 3.3.1. Lemma 3.3.1 also yields $q \geq \text{mc}(\xi) \geq \dot{m}$. Thus it suffices to show that if θ is such that $\text{mc}(\theta) \geq \dot{m}$ and $\mathbb{A}(\theta) \leq m'$, then $\theta \leq [\alpha | q]$. If $\theta \geq \xi$ then $\mathbb{A}(\theta) > m > m'$ by maximality of ξ . If $\theta = \alpha + p$ then $p < b$. Consider two sub-cases. If $b = 0$, no such p exists. If $\text{mc}(\xi - 1) < \dot{m}$ holds, this implies that $\text{mc}(\theta) \leq \dot{m}$, contrary to our choice of θ . Finally, we note that if $\theta \in ([\alpha | q], \alpha)$, Lemma 3.5 yields $\mathbb{A}(\theta) > m'$. We thus conclude that $\mathbb{A}([\alpha | q])$ is in normal form.

CASE 2 ($b > 0$ and $\text{mc}(\xi - 1) \geq \dot{m}$). Lemma 4.3 shows that $\mathbb{A}(\xi - 1)$ is in normal form. We use this to check that $[\alpha | q]$ is in normal form, once again using Lemma 4.2. Using Lemma 3.3, we can see that $\mathbb{A}(\xi - 1) \leq q \leq \text{mc}([\alpha | q])$ and $\mathbb{A}([\alpha | q]) < \mathbb{A}(\xi)$. Thus it remains to check that if χ is such that $\text{mc}(\chi) \geq \mathbb{A}(\xi - 1)$ and $\mathbb{A}(\chi) \leq m'$, then $\chi \leq [\alpha | q]$. Since $\mathbb{A}(\xi - 1)$ is in normal form, we must have that $\chi \leq \xi - 1$. If $p < b$ then $\text{mc}(\alpha + p) < \mathbb{A}(\alpha + p) \leq \mathbb{A}(\xi - 1)$. If $\chi \in ([\alpha | q], \alpha)$, Lemma 3.5 yields $\mathbb{A}(\chi) > m'$. Hence indeed $\mathbb{A}([\alpha | q])$ is in normal form. \square

5 Base change

The last ingredient we need in order to define our maximal Goodstein process is the base change operator. As in the classical Goodstein process, $\uparrow_k^\lambda m$ replaces every instance of k by λ in the normal form of m .

Definition 5.1. *Given $2 \leq k < \lambda \leq \omega$ and $m \in \mathbb{N}$, we define the base change operation $\uparrow_k^\lambda m = \uparrow m$ inductively as follows. Let $\kappa = \lambda^+$, and set:*

1. $\uparrow_k^\lambda 0 = 0$
2. For $m = {}_k \mathbb{A}_k(\xi)$ we set $\uparrow m = \mathbb{A}(\uparrow \xi)$, and
3. For $\xi = \omega^\alpha b + \gamma$ in ω -normal form, we set $\uparrow \xi = \kappa^{\uparrow \alpha} \uparrow b + \uparrow \gamma$.

Base-change maximal normal forms lead to monotone base-change operators, in the following sense.

Proposition 5.2. *Let $2 \leq k < \lambda \leq \omega$ and write \uparrow for \uparrow_k^λ . Suppose that m has the property that for all $n < m$, if $n = \mathbb{A}_k(\zeta)$, then $\uparrow n \geq \mathbb{A}_\lambda(\uparrow \zeta)$. Then, whenever $2 \leq k < \lambda$ and $i < j < m$, it follows that $\uparrow_k^\lambda i < \uparrow_k^\lambda j$.*

Proof. Working inductively, we may assume that $j = i + 1$. Then, we have that $i = i + 1 = |\mathbb{A}_k(i)|$, and by base-change maximality,

$$\uparrow j \geq \uparrow \mathbb{A}_k(i) = \mathbb{A}_\ell(\uparrow i) = \uparrow i + 1 > \uparrow i.$$

□

In fact, this monotonicity property is crucial for proving that Goodstein processes terminate, and Proposition 5.2 tells us that we have this property for free, if we prove base-change maximality.

Remark 5.3. *Our goal is to prove that base-change maximality indeed holds. If we are to prove this by induction on m , then Proposition 5.2 tells us that we may assume that the base-change operator is monotone below m . Thus this will be used as a hypothesis in many of the following results. Once Theorem 6.4 and the subsequent Corollary 6.5 (which states that the base change is monotone) have been established, we may drop this assumption wherever it was used.*

Lemma 5.4. *Assume monotonicity of base change below m and let $2 \leq k < \lambda \leq \omega$. If $\text{mc}(\xi), \text{mc}(\zeta) < m$, then $\uparrow_k^\lambda \xi < \uparrow_k^\lambda \zeta$, and if $\text{mc}(\xi) < \text{mc}(\zeta) < m$, then $\uparrow \text{mc}(\xi) < \uparrow \text{mc}(\zeta)$.*

Proof sketch. Both claims are easily verified using the fact that the base change is applied to the coefficients of ξ and ζ , and we may apply monotonicity to them given that they are bounded by m . □

Lemma 5.5. *Assume monotonicity of base change up to $\text{mc}(\alpha)$, where $\alpha < \varepsilon_0$ is a limit. Let $n \in \mathbb{N}$, $2 \leq k < \lambda \leq \omega$, and $2 \leq k < \lambda \leq \omega$. Write \uparrow for \uparrow_k^λ . Then,*

1. $\text{mc}(\uparrow[\alpha|n]) \leq \text{mc}([\uparrow\alpha|\uparrow n]_{\lambda+})$, and
2. $\uparrow[\alpha|n] \leq [\uparrow\alpha|\uparrow n]_{\lambda+}$.

Proof. By induction on α , where the critical cases are $\alpha = \omega^\beta c$ with $c > 1$ and $\alpha = \omega^{\beta+c}$ with β a limit and $c > 0$. In the first case, we have that

$$\uparrow[\alpha|n] = \uparrow(\omega^\beta(c-1) + [\omega^\beta|n]) = \omega^{\uparrow\beta} \uparrow(c-1) + \uparrow[\omega^\beta|n],$$

while

$$[\uparrow\alpha|\uparrow n]_{\lambda+} = [\omega^{\uparrow\beta} \uparrow c | \uparrow n]_{\lambda+} = \omega^{\uparrow\beta} (\uparrow c - 1) + [\omega^{\uparrow\beta} | \uparrow n]_{\lambda+}.$$

Monotonicity below $\text{mc}(\alpha)$ yields $\uparrow(c-1) \leq \uparrow c - 1$, while the induction hypothesis yields $\uparrow[\omega^\beta|n] \leq [\omega^{\uparrow\beta}|\uparrow n]_{\lambda+}$. From this it readily follows that $\uparrow[\alpha|n] \leq [\uparrow\alpha|\uparrow n]_{\lambda+}$, establishing the first item, and inspection on the coefficients involved establishes the second. The case for $\alpha = \omega^{\beta+c}$ is similar. □

Note that as an immediate corollary we obtain that $\mathbb{A}_\lambda(\uparrow[\alpha|n]) \leq \mathbb{A}_\lambda([\uparrow\alpha|\uparrow n])$. The following claims are easily verified simultaneously by induction on n and α .

Lemma 5.6. *Let $2 \leq k < \lambda \leq \omega$ and assume monotonicity of base change below n .*

1. *If $n \in \mathbb{N}$ then $n \leq \uparrow_k^\lambda n$, and if $n = \mathbb{A}(\xi)$ with $\xi \geq \omega$, then $2n + 2 \leq \uparrow_k^\lambda n$.*
2. *If $\alpha < \varepsilon_0$ and $\text{mc}(\alpha) < n$ then $\alpha \leq \uparrow_k^\lambda \alpha$, and if $\lambda < \omega$ then $\uparrow_k^\lambda \alpha \in \text{Lim}$ if and only if $\alpha \in \text{Lim}$.*

Proof. Write \uparrow instead of \uparrow_k^ℓ , \mathbb{A} for \mathbb{A}_k and \mathbb{B} for \mathbb{A}_λ . For the first claim, proceed by induction on n . Clearly, $0 \leq \uparrow 0$. Otherwise, write $n =_k \mathbb{A}(\xi)$, so that $\uparrow n = \mathbb{B}(\uparrow \xi)$. If ξ is finite, then $\xi = n - 1$ and the induction hypothesis yields $\uparrow \xi \geq \xi$, so that $\uparrow n = \uparrow \xi + 1 \geq n$. Otherwise, proceed by a secondary induction on ξ to show that if $\xi \geq \omega$ then $2\mathbb{A}(\xi) + 2 < \mathbb{B}(\uparrow \xi)$. By induction on i , we show that $\mathbb{A}^{(i)}(\xi) \leq \mathbb{B}^{(i)}(\uparrow \xi)$. For $i = 0$ we have if ξ is a limit that $\mathbb{A}^{(0)}(\xi) = \text{mc}(\xi) \leq \uparrow \text{mc}(\xi) = \mathbb{B}^{(0)}(\uparrow \xi)$, where we have used Lemma 3.3 to see that $\text{mc}(\xi) < n$, so that we may apply the induction hypothesis. Otherwise,

$$\mathbb{A}^{(0)}(\xi) = \mathbb{A}(\alpha + b - 1) \leq \mathbb{B}(\uparrow \alpha + \uparrow(b - 1)) \leq \mathbb{B}(\uparrow \alpha + \uparrow b - 1) = \mathbb{B}^{(0)}(\uparrow \xi),$$

where we have used monotonicity of base change below n to conclude that $\uparrow(b - 1) \leq \uparrow b - 1$.

Now assume that $\mathbb{A}^{(i)}(\xi) \leq \mathbb{B}^{(i)}(\uparrow \xi)$. We see that

$$\begin{aligned} \mathbb{A}^{(i+1)}(\xi) &= \mathbb{A}([\alpha|\mathbb{A}^{(i)}(\xi)]) \stackrel{\text{IH}(\xi)}{\leq} \mathbb{B}(\uparrow[\alpha|\mathbb{A}^{(i)}(\xi)]) \\ &\leq \mathbb{B}([\uparrow\alpha|\uparrow\mathbb{A}^{(i)}(\xi)]) \stackrel{\text{IH}^{(i)}}{\leq} \mathbb{B}([\uparrow\alpha|\uparrow\mathbb{B}^{(i)}(\xi)]) = \mathbb{B}^{(i+1)}(\uparrow \xi), \end{aligned}$$

where first we use the induction hypothesis on $[\alpha|\mathbb{A}^{(i)}(\xi)] < \xi$, then the secondary induction hypothesis on i , and we have used Lemma 5.5 to see that $\mathbb{B}(\uparrow[\alpha|\mathbb{A}^{(i)}(\xi)]) \leq \mathbb{B}([\uparrow\alpha|\uparrow\mathbb{A}^{(i)}(\xi)])$. Lemma 3.3 then yields

$$2\mathbb{A}(\xi) + 2 = 2\mathbb{A}^{(k)}(\xi) + 2 \leq 2\mathbb{B}^{(k)}(\uparrow \xi) + 2 < \mathbb{B}^{(k+1)}(\uparrow \xi) = \mathbb{B}(\uparrow \xi).$$

The second claim follows by an easy induction on α using the first claim. \square

In general we have that $\uparrow[\alpha|b] \leq [\uparrow\alpha|\uparrow b]$. On occasion it will be useful for this to be an equality, and we may define a variant of base change such that $\uparrow[\alpha|b] = [\uparrow\alpha|\uparrow b]$.

Definition 5.7. *Let $2 \leq k < \lambda \leq \omega$, and write \uparrow for \uparrow_k^λ and set $\kappa = \lambda^+$. For $\xi < \varepsilon_0$, define $\uparrow \xi = \uparrow_k^\lambda \xi$ recursively by*

1. $\uparrow 0 = 0$,
2. $\uparrow(\omega^\alpha b + \gamma) = \kappa^\alpha + \uparrow \gamma$ if $\omega^\alpha b + \gamma$ is in ω -normal form,

3. $\zeta\omega^\alpha(b+1) = \kappa^{\uparrow\alpha}\uparrow b + \zeta\omega^\alpha$ if $b > 0$,
4. $\zeta\omega^{\alpha+1} = \kappa^{\uparrow\alpha+1}$,
5. $\zeta\omega^\alpha = \kappa^{\zeta\alpha}$ if α is not a successor.

Lemma 5.8. Let $\xi < \varepsilon_0$ be a limit ordinal, $t \in (1, \omega)$ and $2 \leq k < \lambda \leq \omega$. Write ζ for ζ_k^λ and \uparrow for \uparrow_k^λ . Then,

1. $\zeta\xi = \lceil \uparrow[\xi|t] \rceil$.
2. $\uparrow[\xi|t] = \lceil \zeta\xi \uparrow t \rceil$.
3. If $t > 1$ and ζ is a limit with $\uparrow[\xi|t] = \lceil \zeta \uparrow t \rceil$, then $\zeta = \zeta\xi$.
4. $\mathbb{A}_\lambda(\zeta\xi + t) \leq \mathbb{A}_\lambda(\uparrow\xi + t)$.
5. ξ is a limit iff $\zeta\xi$ is a limit.
6. $\tau_{\lambda^+}(\zeta\xi) = \lambda^+$.

Proofsketch. Let $\kappa = \lambda^+$. The first two items follow by induction on ξ . The critical case is where $\xi = \omega^{\alpha+1}$. For the first claim, $[\xi|t] = \omega^\alpha t$, so that $\uparrow[\xi|t] = \kappa^{\uparrow\alpha}\uparrow t$, and

$$\lceil \uparrow[\xi|t] \rceil_\kappa = \lceil \kappa^{\uparrow\alpha}\uparrow t \rceil_\kappa = \kappa^{\uparrow\alpha+1} = \zeta\xi.$$

For the second, we have that

$$\uparrow[\xi|t] = \uparrow\omega^\alpha t = \kappa^{\uparrow\alpha}\uparrow t = \lceil \kappa^{\uparrow\alpha+1} \uparrow t \rceil_\kappa = \lceil \zeta\omega^\alpha t \uparrow t \rceil.$$

Other cases follow by applying the induction hypothesis to the relevant sub-terms.

The third claim follows from Corollary 2.7, since from $\lceil \zeta \uparrow t \rceil = \uparrow[\xi|t] = \lceil \zeta\xi \uparrow t \rceil$ and injectivity of the fundamental sequences we obtain $\zeta = \zeta\xi$. The fourth follows from the fact that $\zeta\xi + t \leq \uparrow\xi + t$ and $\text{mc}(\zeta\xi + t) \leq \text{mc}(\uparrow\xi + t)$, which are verified by a routine induction, and the fifth by induction and case-by-case inspection.

The sixth item also proceeds by induction. We treat the cases where $\xi = \omega^{\alpha+1}$ and $\xi = \omega^\alpha(b+1)$ with $b > 0$. We have that $\zeta\omega^{\alpha+1} = \kappa^{\uparrow\alpha+1}$, and $\tau_\kappa(\kappa^{\uparrow\alpha+1}) = \kappa$ by definition. Similarly, $\zeta\omega^\alpha(b+1) = \kappa^{\uparrow\alpha}\uparrow b + \zeta\omega^\alpha$, and by the induction hypothesis, $\tau_\kappa(\zeta\omega^\alpha) = \kappa$, so that $\tau_\kappa(\xi) = \kappa$ as well. \square

Lemma 5.9. Let $2 \leq k < \lambda \leq \omega$ and write \uparrow for \uparrow_k^λ and ζ for ζ_k^λ . If $c > 1$, $\gamma < \varepsilon_0$, and $\theta < \varepsilon(\lambda^+)$ is such that $\tau_\kappa(\theta) = \kappa$ and $[\theta \uparrow c]_{\lambda^+} = \uparrow\gamma$, then $[[\gamma] \uparrow c] = \gamma$ and $\zeta[\gamma] = \theta$.

Proof. Let $\kappa = \lambda^+$. By Lemma 5.8.2, $\uparrow[[\gamma] \uparrow c] = \lceil \zeta[\gamma] \uparrow c \rceil_\kappa$, so that $[[\gamma] \uparrow c] = \gamma$ iff $\lceil \zeta[\gamma] \uparrow c \rceil_\kappa = \gamma$ iff $\lceil \zeta[\gamma] \uparrow c \rceil_\kappa = [\theta \uparrow c]_\kappa$ iff $\zeta[\gamma] = \theta$, where the last equivalence follows from Corollary 2.7 and the fact that $\tau_\kappa(\zeta[\gamma]) = \kappa$ by Lemma 5.8.6. Thus we prove that $\zeta[\gamma] = \theta$ by induction on θ . Write $\theta = \kappa^\alpha \delta + \beta$ in κ -normal form and $\gamma = \omega^{\tilde{\alpha}} d + \tilde{\beta}$ in ω -normal form, so that $\uparrow\gamma = \kappa^{\uparrow\tilde{\alpha}} \uparrow d + \uparrow\tilde{\beta}$.

CASE 1 ($\beta > 0$). Then $[\theta|\uparrow c]_\kappa = \kappa^\alpha d + [\beta|\uparrow c]_\kappa$, so that the assumption that $[\theta|\uparrow c]_\kappa = \uparrow\gamma$ yields $\uparrow\check{\alpha} = \alpha$, $\uparrow d = \delta$, and $\uparrow\check{\beta} = [\beta|\uparrow c]$. Since $\uparrow\check{\beta} = [\beta|\uparrow c]$, we may apply the induction hypothesis to obtain $\beta = \check{\zeta}[\check{\beta}]$, and hence

$$\check{\zeta}[\gamma] = \uparrow\omega^{\check{\alpha}}d + \check{\zeta}[\check{\beta}] = \kappa^\alpha\delta + \beta = \theta.$$

CASE 2 ($\beta = 0$ and $\delta = \eta + 1 > 1$). Then $[\theta|\uparrow c]_\kappa = \kappa^\alpha\eta + [\kappa^\alpha|\uparrow c]_\kappa$, and $\uparrow\check{\alpha} = \alpha$, $\uparrow d = \eta$, and $\uparrow\check{\beta} = [\omega^\alpha|\uparrow c]$. The induction hypothesis yields $\omega^\alpha = \check{\zeta}[\check{\beta}]$, hence

$$\check{\zeta}[\gamma] = \uparrow\omega^{\check{\alpha}}d + \check{\zeta}[\check{\beta}] = \omega^\alpha\eta + \omega^\alpha = \theta.$$

CASE 3 ($\beta = 0$, $\delta = 1$, and $\alpha = \chi + 1$). Then $[\theta|\uparrow c] = \omega^\delta\uparrow c$, and $\uparrow\check{\alpha} = \chi$, $\uparrow d = \uparrow c$ so that $d = c > 1$, and $\uparrow\check{\beta} = 0$, so $\check{\beta} = 0$. Then, $[\gamma] = \omega^{\check{\alpha}+1}$, so that

$$\check{\zeta}[\gamma] = \kappa^{\uparrow\check{\alpha}+1} = \kappa^{\chi+1} = \theta.$$

CASE 4 ($\beta = 0$, $\delta = 1$, and α is a limit). Then $[\theta|\uparrow c] = \omega^{[\alpha|\uparrow c]}$, and $\uparrow\check{\alpha} = [\alpha|\uparrow c]$, $d = 1$, and $\check{\beta} = 0$. The induction hypothesis yields $\alpha = \check{\zeta}[\check{\alpha}]$, hence since α is a limit, so is $[\check{\alpha}]$, and

$$\check{\zeta}[\gamma] = \check{\zeta}\omega^{[\check{\alpha}]} = \kappa^{\check{\zeta}[\check{\alpha}]} = \kappa^\alpha = \theta. \quad \square$$

We have noted that in general $\uparrow[\xi|n] \neq [\uparrow\xi|\uparrow n]_{\lambda^+}$, but there is an important case where this equality does hold. Below, $\tau = \tau_\Omega$.

Lemma 5.10. *Write \uparrow for \uparrow_k^ω and let $\xi < \varepsilon_{\Omega+1}$. If $\tau(\uparrow\xi) = \Omega$, then $\uparrow[\xi|n] = [\uparrow\xi|\uparrow n]_{\lambda^+}$.*

Proof sketch. This follows by induction, where the critical case is when $\xi = \omega^\zeta b$. We cannot have that $\uparrow b \in \text{Lim}$, since this would imply that $\tau(\uparrow\xi) = \uparrow b < \Omega$. Hence $\uparrow b = \beta + 1$ for some β , which means that $b =_k \mathbb{A}(b-1)$ (as $\mathbb{A}(\zeta)$ is a limit for any $\zeta \geq \omega$), and thus $\uparrow b = \uparrow(b-1) + 1$, so that $\uparrow(b-1) = \beta$. Now, write $\zeta = \gamma + a$ with γ a limit, and as above we see that a is either zero or else $\uparrow a = \uparrow(a-1) + 1$. If $a = 0$, we have that

$$\begin{aligned} \uparrow[\xi|n] &= \uparrow(\omega^\gamma(b-1) + \omega^{[\gamma|n]}) = \Omega^{\uparrow\gamma}\uparrow(b-1) + \Omega^{\uparrow[\gamma|n]} \\ &\stackrel{\text{IH}}{=} \Omega^{\uparrow\gamma}\beta + \Omega^{\uparrow[\gamma|\uparrow n]} = [\Omega^{\uparrow\gamma}(\beta+1)|\uparrow n]. \end{aligned}$$

If $\uparrow a = \alpha + 1$, we have that

$$\begin{aligned} \uparrow[\xi|n] &= \uparrow(\omega^\zeta(b-1) + \omega^\alpha n) = \Omega^{\uparrow\gamma+\uparrow a}\beta + \Omega^{\uparrow(a-1)}\uparrow n \\ &= \omega^{\uparrow\gamma+\alpha+1}\beta + \omega^\alpha\uparrow n = [\Omega^{\uparrow\gamma+\alpha+1}(\beta+1)|\uparrow n]_\Omega = [\uparrow\xi|\uparrow n]_\Omega. \quad \square \end{aligned}$$

6 Maximality of Base Change

Our strategy for proving that every fast Goodstein walk is finite proceeds by showing that the normal forms we have given provide the maximal value after base change, thus yield the longest possible Goodstein processes. In this section we prove this maximality property. We begin with some useful lemmas.

Lemma 6.1. *Assume monotonicity of base change below m . Suppose that $m =_k \mathbb{A}(\xi)$ with $\xi = \alpha + b$, where α is a limit and $b > 0$. Then,*

1. $\uparrow \mathbb{A}(\alpha - 1 + b) < 1 + \mathbb{B}(\uparrow \alpha - 1 + \uparrow b)$, and
2. if $\mathbb{A}(\alpha - 1 + b)$ is not in normal form, then $\uparrow \mathbb{A}(\alpha - 1 + b) < 1 + \mathbb{B}(\uparrow \alpha - 1 + \uparrow b)$.

Proof. We consider two cases.

CASE 1 ($\mathbb{A}(\xi - 1)$ is in normal form). Then, monotonicity below m yields $\uparrow(-1 + b) \leq -1 + \uparrow b$, hence

$$\uparrow \mathbb{A}(\xi - 1) = \mathbb{B}(\uparrow(\xi - 1)) \leq \mathbb{B}(\uparrow \alpha - 1 + \uparrow b).$$

If $\uparrow b$ is infinite, then the inequality is strict, since $-1 + \uparrow b = \uparrow b$.

CASE 2 ($\mathbb{A}(\xi - 1)$ is not in normal form). By Lemma 4.3, we have that $mc(\alpha) < b$ and $b =_k \mathbb{A}(\zeta)$ for some $\zeta > \xi$, which in particular implies that $\zeta > \omega$. Note that in cases where $\mathbb{A}(\xi - 1)$ is not in normal form, it suffices to show that $\uparrow \mathbb{A}(\xi - 1) \leq \mathbb{B}(\uparrow \alpha - 1 + \uparrow b)$, as $\mathbb{B}(\uparrow \alpha - 1 + \uparrow b) \leq \mathbb{B}(\uparrow \alpha - 1 + \uparrow b)$. We prove the more general claim that for all $t \in [1, b + 1]$,

$$\uparrow \mathbb{A}(\alpha - t + b) \leq \mathbb{B}(\uparrow \alpha - t + \uparrow b).$$

Consider two sub-cases and proceed by induction on $b - t$.

CASE 2.1 ($\mathbb{A}(\xi - t) \leq b$). If $\uparrow b$ is infinite then $-t + \uparrow b = \uparrow b$, and

$$\uparrow \mathbb{A}(\xi - t) \leq \uparrow b < \mathbb{B}(\uparrow \alpha + \uparrow b) = \mathbb{B}(\uparrow \alpha - t + \uparrow b).$$

If $\uparrow b$ is finite, $\uparrow b > 2b + 2$ by Lemma 5.6, and we have that

$$\begin{aligned} \mathbb{B}(\uparrow \alpha - t + \uparrow b) &\geq \mathbb{B}(\uparrow \alpha - b - 1 + \uparrow b) \geq \mathbb{B}(\uparrow \alpha + \lceil \uparrow b/2 \rceil) \\ &> 2(\uparrow b/2) = \uparrow b \geq \uparrow \mathbb{A}(\xi - t) \end{aligned}$$

(note that here $\lceil \uparrow b/2 \rceil$ is the standard integer ceiling function).

CASE 2.2 ($\mathbb{A}(\xi - t) > b$). By the induction hypothesis, $\uparrow \mathbb{A}(\xi - t - 1) < 1 + \mathbb{B}(\uparrow \alpha - t - 1 - \uparrow b)$. We claim moreover that $\mathbb{A}(\xi - t - 1) \geq b$. Note that $mc(\alpha) < b$, so that $t < b - 1$ (otherwise $\mathbb{A}(\xi - t) = mc(\alpha)$). Since $\zeta > \xi$, by Lemma 3.5,

$$b = \mathbb{A}(\zeta) \notin (\mathbb{A}(\xi - t - 1), \mathbb{A}(\xi - t)),$$

from which it follows that $b \leq \mathbb{A}(\xi - t - 1)$. Note moreover that if λ is infinite this implies that $\uparrow b$ is infinite.

We proceed by induction on i to show that, for $i \leq k$, $\mathbb{A}^{(i)}(\xi - t) < 1 + \mathbb{B}^{(i)}(\uparrow\alpha - t + \uparrow b)$. By the induction hypothesis for $b - t - 1 < b - t$ we have that

$$\begin{aligned} \uparrow\mathbb{A}^{(0)}(\xi - t) &= \uparrow\mathbb{A}(\xi - t - 1) < 1 + \mathbb{B}(\uparrow\alpha - t - 1 - \uparrow b) \\ &\leq 1 + \mathbb{B}^{(0)}(\uparrow\alpha - t - \uparrow b). \end{aligned}$$

We have that $\mathbb{A}^{(i)}(\xi - t) \geq \mathbb{A}(\xi - t - 1) \geq b$, so that Lemma 4.4 yields $\mathbb{A}^{(i+1)}(\xi - t) =_k \mathbb{A}([\alpha|\mathbb{A}^{(i)}(\xi - t)])$. Hence,

$$\uparrow\mathbb{A}^{(i+1)}(\xi - t) = \mathbb{B}(\uparrow[\alpha|\mathbb{A}^{(i)}(\xi - t)]) = \mathbb{B}([\uparrow\alpha|\uparrow\mathbb{A}^{(i)}(\xi - t)]).$$

If λ is finite then

$$\mathbb{B}([\uparrow\alpha|\uparrow\mathbb{A}^{(i)}(\xi - t)]) \stackrel{\text{IH}}{\leq} \mathbb{B}([\uparrow\alpha|\mathbb{B}^{(i)}(\uparrow\xi - t)]) = \mathbb{B}^{(i+1)}(\uparrow\xi - t).$$

If λ is infinite then $\uparrow\mathbb{A}^{(i)}(\xi - t) < \mathbb{B}^{(i)}(\uparrow\alpha - t + \uparrow b)$ yields

$$\mathbb{B}([\uparrow\alpha|\uparrow\mathbb{A}^{(i)}(\xi - t)]) < \mathbb{B}(\uparrow\xi) = \mathbb{B}^{(i+1)}(\uparrow\alpha - t + \uparrow b).$$

The claim follows by setting $i = k$, since

$$\uparrow\mathbb{A}^{(k)}(\xi - t) < 1 + \mathbb{B}^{(k)}(\uparrow\alpha - t + \uparrow b) \leq 1 + \mathbb{B}(\uparrow\alpha - t + \uparrow b),$$

as desired. \square

Lemma 6.2. *Let $2 \leq k < \lambda \leq \omega$ and write \uparrow for \uparrow_k^λ , \uparrow for \uparrow_k^λ . Suppose that $\mathbb{A}(\xi)$ is in normal form, where $\xi = \alpha + b$ with α a limit, and let $i < k$. Then,*

1. $\uparrow\mathbb{A}^{(i)}(\xi) < 1 + \mathbb{B}^{(i)}(\uparrow\xi)$, and
2. $\mathbb{B}(\uparrow[\alpha|\mathbb{A}^{(i)}(\xi)]) < 1 + \mathbb{B}^{(i+1)}(\uparrow\xi)$.

Proof. We prove both claims simultaneously by induction on i . When $i = 0$, the first claim follows from Lemma 6.1, or by definition when $\lambda < \omega$ and ξ is a limit (in which case this becomes $\uparrow\text{mc}(\xi) < 1 + \uparrow\text{mc}(\xi)$). For $i + 1$, the induction hypothesis on the second claim and Lemma 4.4 yield

$$\uparrow\mathbb{A}^{(i+1)}(\xi) = \mathbb{B}(\uparrow[\alpha|\mathbb{A}^{(i)}(\xi)]) < 1 + \mathbb{B}^{(i+1)}(\uparrow\xi).$$

For the second, Lemma 5.5 yields $\mathbb{B}(\uparrow[\alpha|\mathbb{A}^{(i)}(\xi)]) \leq \mathbb{B}([\uparrow\alpha|\uparrow\mathbb{A}^{(i)}(\xi)])$. If λ is finite, then using Lemma 3.4 we see that

$$\mathbb{B}([\uparrow\alpha|\uparrow\mathbb{A}^{(i)}(\xi)]) \stackrel{\text{IH}}{\leq} \mathbb{B}([\uparrow\alpha|\mathbb{B}^{(i)}(\uparrow\xi)]) = \mathbb{B}^{(i+1)}(\uparrow\xi).$$

If λ is infinite, from the first claim we see that $\uparrow\mathbb{A}^{(i)}(\xi) < 1 + \mathbb{B}^{(i)}(\uparrow\xi) = \mathbb{B}(\uparrow\xi)$, therefore

$$\mathbb{B}([\uparrow\alpha|\uparrow\mathbb{A}^{(i)}(\xi)]) < \mathbb{B}(\uparrow\xi) = 1 + \mathbb{B}^{(i+1)}(\uparrow\xi). \quad \square$$

Corollary 6.3. *Assume monotonicity of base change below $m = {}_k \mathbb{A}(\xi)$, and let j_* be the unique integer such that*

$$0 < \lfloor \xi \rfloor_k^{j_*} < \omega.$$

Then for all $i \leq k$ and $j < j_$,*

$$\uparrow(m-1) < \mathbb{B}(\uparrow \lfloor \xi \rfloor_k^j) < 1 + \mathbb{B}^{(k)}(\uparrow \xi) \leq \uparrow m.$$

Proof. Lemma 6.2.2 and induction on $j < j_*$ yield

$$\mathbb{B}(\uparrow \lfloor \xi \rfloor_k^{j+1}) \leq \mathbb{B}(\uparrow \lfloor \xi \rfloor_k^j) < 1 + \mathbb{B}^{(k)}(\uparrow \xi).$$

Write $\lfloor \xi \rfloor_k^{j_*-1} = \gamma + c$ with γ a limit. Since $0 < \lfloor \xi \rfloor_k^{j_*} < \omega$ and $\lfloor \xi \rfloor_k^{j_*-1} = [\gamma|q]$ for some q , we must have that $\gamma = \omega$ and $\lfloor \xi \rfloor_k^{j_*-1} = \omega + c$. For all i ,

$$\begin{aligned} \mathbb{A}^{(i+1)}(\lfloor \xi \rfloor_k^{j_*-1}) &= \mathbb{A}([\omega|\mathbb{A}^{(i)}(\lfloor \xi \rfloor_k^{j_*-1})]) \\ &= \mathbb{A}(\mathbb{A}^{(i)}(\lfloor \xi \rfloor_k^{j_*-1})) = \mathbb{A}^{(i)}(\lfloor \xi \rfloor_k^{j_*-1}) + 1, \end{aligned}$$

and since $m = \mathbb{A}^{(k)}(\lfloor \xi \rfloor_k^{j_*-1})$, we see that $m-1 = \mathbb{A}^{(k-1)}(\lfloor \xi \rfloor_k^{j_*-1})$. Lemma 6.2 then yields

$$\uparrow(m-1) = \uparrow \mathbb{A}^{(k-1)}(\lfloor \xi \rfloor_k^{j_*-1}) < 1 + \mathbb{B}^{(k-1)}(\uparrow \lfloor \xi \rfloor_k^{j_*-1}) \leq 1 + \mathbb{B}(\uparrow \lfloor \xi \rfloor_k^{j_*-1}).$$

□

We are now ready to prove that our normal forms are base-change maximal.

Theorem 6.4. *If $2 \leq k < \lambda \leq \omega$ and $\mathbb{A}(\zeta) = {}_k \mathbb{A}(\xi)$, then $\mathbb{A}(\uparrow_k^\lambda \zeta) \leq \mathbb{A}(\uparrow_k^\lambda \xi)$.*

Proof. Write \uparrow for \uparrow_k^λ . Let $m = \mathbb{A}(\xi)$. Note that $\zeta < \dot{\xi}$. This is because $m \in (\mathbb{A}(\dot{\xi}), \mathbb{A}(\dot{\xi} + 1))$, but by Lemma 3.7, if $\zeta \geq \dot{\xi}$ then $\mathbb{A}(\zeta) \notin (\mathbb{A}(\dot{\xi}), \mathbb{A}(\dot{\xi} + 1))$. Thus we write $\xi = \alpha + a$ and $\zeta = \beta + b$ with α, β limits and consider two cases.

CASE 1 ($\zeta < \xi$). Let i_* be the unique number such that $0 < \lfloor \xi \rfloor_k^{i_*} < \omega$. We claim that there is $i \leq i_*$ such that $\zeta = \lfloor \xi \rfloor_k^i$. If not, there is $i \leq i_*$ such that $\lfloor \xi \rfloor_k^{i+1} < \zeta \leq \lfloor \xi \rfloor_k^i$. Write $\lfloor \xi \rfloor_k^i = \gamma + c$ with γ a limit or zero. If $\zeta < \gamma$, then Lemma 3.5 yields $\mathbb{A}(\xi) > \mathbb{A}(\lfloor \xi \rfloor_k^i) = \mathbb{A}(\xi)$, contrary to assumption. Thus $\zeta = \gamma + c'$, but then $\mathbb{A}(\zeta) = \mathbb{A}(\xi)$ yields $c = c'$ and $\zeta = \lfloor \xi \rfloor_k^i$. But by Corollary 6.3, $\mathbb{A}(\uparrow \lfloor \xi \rfloor_k^i) < \mathbb{A}(\uparrow \xi)$, as needed.

CASE 2 $\zeta \in (\xi, \dot{\xi})$. Using the induction hypothesis (which yields monotonicity below m) and Lemma 5.4, we see that $\uparrow \zeta < \uparrow \dot{\xi}$. Maximality of ξ yields $\text{mc}(\zeta) \leq \dot{m} - 1$, so Corollary 6.3 yields $\text{mc}(\uparrow \beta) < \mathbb{B}^{(k)}(\uparrow \dot{\xi})$. Lemma 3.4 then yields $\mathbb{B}(\uparrow \zeta) < \mathbb{B}(\uparrow \dot{\xi}) < \mathbb{B}(\uparrow \xi)$.

□

Corollary 6.5. *If $n < m$ and $2 \leq k < \lambda \leq \omega$, then $\uparrow_k^\lambda n < \uparrow_k^\lambda m$.*

Proof. Write $n = {}_k \mathbb{A}(\xi)$ and without loss of generality assume $m = n + 1$. Since $m = \mathbb{A}(\mathbb{A}(\xi))$, maximality of the base change yields

$$\uparrow_k^\lambda m \geq \mathbb{B}(\uparrow_k^\lambda \mathbb{A}(\xi)) = \uparrow_k^\lambda n + 1. \quad \square$$

7 Normal form preservation

The next step in showing that the Goodstein process for our normal forms terminates is to show that normal forms are preserved after base-change, which is needed to show that the ordinal assignment gives a decreasing sequence of ordinals.

Lemma 7.1. *Let $2 \leq k < \lambda \leq \omega$ and write \uparrow for \uparrow_k^λ . If $m =_k \mathbb{A}_k(\xi)$ and $2 \leq k < \ell < \omega$, then $\uparrow m =_k \mathbb{A}_\ell(\uparrow \xi)$.*

Proof. Write \mathbb{A} for \mathbb{A}_k and \mathbb{B} for \mathbb{A}_ℓ , and proceed by induction on m . By induction hypothesis we have that $\uparrow m =_\lambda \mathbb{B}(\uparrow \xi)$. By monotonicity (Corollary 6.5), $\text{mc}(\uparrow \xi) > \text{mc}(\uparrow \xi)$. So, in view of Lemma 4.2, it suffices to show that $\uparrow m < \mathbb{B}(\uparrow \xi)$ and that if θ is such that $\mathbb{B}(\theta) \leq \uparrow m$ and $\text{mc}(\theta) \geq \text{mc}(\uparrow \xi)$, then $\theta \leq \uparrow \xi$.

To see that $\mathbb{B}(\uparrow \xi) < \mathbb{B}(\uparrow \xi + 1)$, write $\lfloor \xi + 1 \rfloor_k^i = \gamma_i + c_i$ with γ_i a limit. Then, $\xi = \gamma_i + d$ for some i and $d < c_i$, since otherwise we would have $\mathbb{A}(\xi) \geq \mathbb{A}(\xi + 1)$. From this it is readily checked that $\mathbb{B}(\uparrow \xi) < \mathbb{B}(\uparrow \lfloor \xi + 1 \rfloor_k^i) < \mathbb{B}(\uparrow \xi + 1)$.

With this, it remains to show that $\uparrow \xi$ is maximal. Towards a contradiction, assume that $\hat{\theta}$ is minimal with the property that $\hat{\theta} > \xi$, $\text{mc}(\hat{\theta}) \geq \uparrow m$, and $\mathbb{B}(\hat{\theta}) \leq \uparrow m$. Since $\mathbb{B}(\hat{\theta}) = \mathbb{B}(\lfloor \hat{\theta} \rfloor_\ell)$, we must have that $\lfloor \hat{\theta} \rfloor_\ell \leq \uparrow \xi$. Write $\xi = \alpha + a$ and $\hat{\theta} = \hat{\beta} + \hat{b}$ with $\alpha, \hat{\beta}$ limits, so that $\uparrow \xi = \uparrow \alpha + \uparrow a$. We cannot have $\uparrow \alpha = \hat{\beta}$, since in this case $\uparrow b > \hat{a}$, and $\mathbb{B}(\uparrow \xi) < \mathbb{B}(\hat{\theta})$. Thus $\hat{\beta} > \uparrow \alpha$. By minimality of $\hat{\theta}$, either $\text{mc}(\hat{\beta}) \geq \uparrow m$ and $\hat{b} = 0$, or else $\text{mc}(\beta) < \uparrow m$ and $\hat{b} = \uparrow m$. In either case, there is b such that $\uparrow b = \hat{b}$.

By Lemma 2.10, there is a truncation $\hat{\zeta}$ of $\uparrow \xi$ and some $\hat{c} \geq \mathbb{B}^{(k)}(\hat{\theta})$ such that $\hat{\zeta} = \lfloor \hat{\beta} \rfloor \hat{c}$. Since $\hat{\zeta}$ is a truncation of $\uparrow \xi$, we have that $\hat{\zeta} = \uparrow \zeta$ for some $\zeta < \xi$; in particular, $\hat{c} = \uparrow c$ for some c .

Let $\theta = \lceil \zeta \rceil + b$. Lemma 5.9 implies that $\lceil \lceil \zeta \rceil \rceil c = \zeta$ and $\hat{\beta} = \uparrow \beta$. We aim to prove that $\mathbb{A}(\theta) \leq m$, and proceed by assuming $\mathbb{A}(\theta) > m$ toward a contradiction. We claim that for all $i < k$,

- (a) $\uparrow \mathbb{A}^{(i)}(\theta) \leq \mathbb{B}^{(i)}(\hat{\theta})$,
- (b) $\mathbb{A}^{(i)}(\theta) < c$,
- (c) $\mathbb{A}^{(i+1)}(\theta) < m$, and
- (d) $\mathbb{A}(\lfloor \theta \rfloor \mathbb{A}^{(j)}(\theta))$ is in normal form for $j < k - 1$.

The third item for $i = k - 1$ will yield our desired claim, since it becomes $\mathbb{A}(\theta) < m$, contradicting that $\mathbb{A}(\xi)$ is in normal form. However, we must prove the four claims simultaneously for induction on i to go through.

Assume that all four claims hold for $i' < i$. Let $e = \mathbb{A}^{(i)}(\theta)$.

Proof of (a). When $i = 0$, we have by Lemma 5.6 that

$$\uparrow e = \uparrow \mathbb{A}(\theta - 1) \leq \mathbb{B}(\hat{\theta} - 1) = \mathbb{B}^{(0)}(\hat{\theta}),$$

establishing the first claim when $i = 0$. For $i > 0$, we use the induction hypothesis on the fourth claim to see that

$$\begin{aligned}\uparrow e &= \uparrow \mathbb{A}([\beta|A^{(i-1)}(\theta)]) \\ &= \mathbb{B}(\uparrow[\beta|A^{(i-1)}(\theta)]) = \mathbb{B}([\xi|\beta|\uparrow \mathbb{A}^{(i-1)}(\theta)]) \\ &\leq \mathbb{B}([\xi|\beta|\mathbb{B}^{(i-1)}(\hat{\theta})]) = \mathbb{B}^{(i)}(\hat{\theta}),\end{aligned}\tag{2}$$

where in (2) we have used the induction hypothesis to see that $\uparrow \mathbb{A}^{(i-1)}(\theta) \leq \mathbb{B}^{(i-1)}(\hat{\theta})$. This establishes the first claim when $i > 0$.

Proof of (b). Using the previous claim we see that

$$\uparrow e = \uparrow \mathbb{A}^{(i)}(\theta) \leq \mathbb{B}^{(i)}(\hat{\theta}) < \mathbb{B}^{(k)}(\hat{\theta}) < \uparrow c,$$

and by monotonicity of the base change operator, $e < c$, establishing the second claim.

Proof of (c). Since $e < c$, we have that $[\beta|e] < \xi$ and $\text{mc}([\beta|e]) \leq \text{mc}(\zeta) \leq \text{mc}(\xi)$. By Lemma 3.4, $\mathbb{A}^{(i+1)}(\theta) = \mathbb{A}([\beta|e]) < \mathbb{A}(\xi) = m$.

Proof of (d). Let $j < k - 1$ and $\chi = [\beta|e]$. Note that the previous item yields $\mathbb{A}(\chi) < m < \mathbb{A}(\xi + 1)$. Since $\text{mc}(\chi) \geq \dot{m}$, by Lemma 4.2, it suffices to show that if $\text{mc}(\eta) \geq \dot{m}$ and $\eta > \chi$, then $\mathbb{A}(\eta) > \mathbb{A}(\chi)$. If $\eta > \xi$, then the maximality of ξ yields $\mathbb{A}(\eta) > \mathbb{A}(\xi) > \mathbb{A}(\chi)$. If $\eta \leq \xi$, then $\eta \in (\chi, \theta)$, which by Lemma 3.5, implies that $\mathbb{A}(\eta) > \mathbb{A}(\chi)$. Thus $\mathbb{A}(\chi)$ is in normal form, as required.

Applying (c) with $i = n - 1$, we conclude that ξ is not maximal with $\text{mc}(\xi) \geq \dot{m}$ and $\mathbb{A}(\xi) \leq m$, contradicting the original assumption that $\mathbb{A}(\xi)$ is in normal form. Thus we conclude that $\mathbb{B}(\uparrow \xi)$ is in normal form as well. \square

With this and a simple induction, we obtain the following useful property.

Corollary 7.2. *If $2 \leq k < \ell < \lambda \leq \omega$ and $m \in \mathbb{N}$ then*

$$\uparrow_k^\lambda m = \uparrow_\ell^\lambda \uparrow_k^\ell m.$$

8 Fast Goodstein Walks

Now we are ready to define our fast Goodstein processes and prove that they terminate. We also show that Goodstein walks based on the \mathbb{A} function always terminate, even if normal forms are not used.

Definition 8.1. *Given a natural number m , we define a sequence $(\mathcal{G}_i m)_{i < \alpha}$, where $\alpha \leq \omega$, by the following recursion.*

1. $\mathcal{G}_0 m = m$;
2. if $\mathcal{G}_i m > 0$, then $\mathcal{G}_{i+1} m = \uparrow_{k+2}^{k+3} \mathcal{G}_i m - 1$;

3. if $\mathcal{G}_i m = 0$, then $\alpha = i + 1$ and the sequence terminates.

The sequence $(\mathcal{G}_i m)_{i < \alpha}$ is the Fast Goodstein sequence starting on m .

Theorem 8.2. Given any $m \in \mathbb{N}$, the Fast Goodstein sequence starting on m terminates on finite time.

Proof. Let $(\mathcal{G}_i m)_{i < \alpha}$ be the fast Goodstein sequence starting on m . Let $i < \alpha$. Then,

$$\begin{aligned} \uparrow_{i+3}^\omega \mathcal{G}_{i+1} m &= \uparrow_{i+3}^\omega (\uparrow_{i+2}^{i+3} \mathcal{G}_i m - 1) \\ &< \uparrow_{i+3}^\omega \uparrow_{i+2}^{i+3} \mathcal{G}_i m \end{aligned} \quad (3)$$

$$= \uparrow_{i+2}^\omega \mathcal{G}_i m, \quad (4)$$

where (3) follows from Corollary 6.5 and (4) from Corollary 7.2. Hence $(\uparrow_{i+2}^\omega \mathcal{G}_i m)_{i < \alpha}$ is a decreasing sequence of ordinals, so α must be finite. \square

It is not needed to write numbers in normal form in order for the process to terminate. Natural numbers may be represented using the \mathbb{A}_k functions in various ways. To make this precise, we build terms for numbers and ordinals out of this function. Given fixed $k \geq 2$, the set of k -terms and ordinal k -terms are defined inductively as follows:

1. 0 is both a k -term and an ordinal k -term.
2. If τ is an ordinal term, then $\mathbb{A}_k(\tau)$ is both a k -term and an ordinal k -term.
3. If τ, σ are ordinal terms and ν is a number term, $\omega^\tau \nu + \sigma$ is an ordinal term.

The set of k -terms will be denoted \mathbb{T}_k , and we set $\mathbb{T} = \bigcup_{k < \omega} \mathbb{T}_k$. The *value* of a term is defined inductively in the obvious way by $|0| = 0$, $|\mathbb{A}_k(\tau)| = \mathbb{A}_k(|\tau|)$, and $|\omega^\tau \nu + \sigma| = \omega^{|\tau|} |\nu| + |\sigma|$. We remark that we use the notation for function symbols and the functions they represent, but whether an expression should be treated as a term or as a number will be made clear.

The base change operator can be applied to arbitrary terms, even those in normal form. Given $k \leq \ell$ and $\tau \in \mathbb{T}_k$, we define $\uparrow^\ell \tau$ recursively by

1. $\uparrow^\ell 0 = 0$,
2. $\uparrow^\ell \mathbb{A}_k(\tau) = \mathbb{A}_\ell(\uparrow^\ell \tau)$,
3. $\uparrow^\ell (\omega^\tau \nu + \sigma) = \omega^{\uparrow^\ell \tau} \uparrow^\ell \nu + \uparrow^\ell \sigma$.

Then, normal forms give maximal base change in the following sense.

Proposition 8.3. If $2 \leq k < \ell < \omega$ and τ is any k -term, then $\uparrow^\ell \tau \leq \uparrow_k^\ell |\tau|$.

Proof. Follows from Theorem 6.4 using induction on term complexity. \square

With this, we may define Goodstein walks, in which natural numbers are written using *any* term.

Definition 8.4. A fast Goodstein walk is a sequence $(m_i)_{i=0}^\alpha$, where $\alpha \leq \omega$, such that for every $i < \alpha$, there is an $(i+2)$ -term τ_i with $|\tau_i| = m_i$ and $m_{i+1} = \uparrow^{i+3} \tau - 1$.

Theorem 8.5. Every fast Goodstein walk is finite.

Proof. Let $(m_i)_{i=0}^\alpha$ be a Goodstein walk for \mathcal{F} . Let $m = m_0$. By induction on i , we check that $m_i \leq \mathcal{G}_i m$. For the base case this is clear. Otherwise, $m_{i+1} = |\uparrow^{i+3} \tau_i| - 1$ for some $(i+2)$ -term τ_i , and thus

$$m_{i+1} = |\uparrow^{i+3} \tau_i| - 1 \leq \uparrow_{i+2}^{i+3} m_i - 1 \stackrel{\text{IH}}{\leq} \uparrow_{i+2}^{i+3} \mathcal{G}_i m - 1 = \mathcal{G}_{i+1} m,$$

where the second inequality uses Corollary 6.5 along with the induction hypothesis for i . Thus if we choose i such that $\mathcal{G}_i m = 0$, we must have $\alpha \leq i$. \square

Example 8.6. Consider alternative normal forms obtained by writing $m \simeq_k \mathbb{A}_k(\xi)$, where ξ is maximal so that $\mathbb{A}(\xi) = m$. Such normal forms give alternative Goodstein sequences, which are terminating by Theorem 8.5.

9 Fundamental sequences for the ϑ function

In order to prove that Theorem 8.2 is not provable in KP, we must review fundamental sequences for the ϑ function, as defined by Buchholz [4]. The ϑ function is essentially defined as \mathbb{A}_ω , but with the additional condition that all values must be additively indecomposable, i.e. for $\xi < \varepsilon_{\Omega+1}$, $\vartheta(\xi)$ is defined recursively by

$$\vartheta(\xi) = \min \{ \theta : \forall \zeta < \xi (\text{mc}_\Omega(\zeta) < \theta \rightarrow \vartheta(\zeta) < \theta) \}.$$

In this section, $\mathbb{B} = \mathbb{A}_\omega$.

Definition 9.1. For $\xi < \varepsilon_{\Omega+1}$, we define $(\xi)_\vartheta$ inductively as follows.

- (a) $(0)_\vartheta = 0$.
- (b) If α is countable, then $(\mathbb{B}(\alpha))_\vartheta = \alpha + \vartheta(0)$ (recall that $\vartheta(0) = 1$).
- (c) Let α, β be countable and $\beta = \gamma + n$ with γ a limit or zero. Define

$$x_\alpha^\beta = \begin{cases} 1 & \text{if } \gamma = \omega^{\omega^\alpha} \gamma, \\ 0 & \text{otherwise.} \end{cases} \quad y_\alpha = \begin{cases} 1 & \text{if } \alpha = \omega^{\omega^\alpha}, \\ 0 & \text{otherwise.} \end{cases}$$

Then,

$$(\mathbb{B}(\Omega(1 + \alpha) + \beta))_\vartheta = \omega^{\omega^\alpha} (y_\alpha + \beta + x_\alpha^\beta).$$

- (d) $(\mathbb{B}(\Omega^2 + \beta))_\vartheta = \vartheta(\Omega + \beta)$.

We claim that $\xi = (\xi)_\vartheta$ for all $\xi \in \mathbb{A}(\varepsilon_{\Omega+1})$. This will follow from the next few lemmas.

Lemma 9.2. *If $\xi < \mathbb{A}(\varepsilon_{\Omega+1})$ then $\xi \leq (\xi)_\vartheta$.*

Proof. Proceed by induction on ξ and consider several cases.

CASE 1 ($\Omega \leq \xi < \Omega^2$). Write $\xi = \Omega(1 + \alpha) + \beta$ and $\beta = \gamma + n$, where γ is a limit or zero, and define $x = x_{\alpha}^{\beta}$, $y = y_{\alpha}$ as in the definition of $(\xi)_\vartheta$. The claim is that

$$\begin{aligned} \mathbb{A}(\Omega(1 + \alpha) + \beta) &\leq (\mathbb{A}(\Omega(1 + \alpha) + \beta))_\vartheta \\ &= \omega^{\omega^\alpha}(y + \beta + x). \end{aligned}$$

Let $\theta = \omega^{\omega^\alpha}(y + \beta + x)$. To show that $\mathbb{A}(\xi) \leq \theta$, we check that if $\zeta < \xi$ and $\text{mc}_\Omega(\zeta) < \theta$, then $\mathbb{A}(\zeta) < \theta$. If $\zeta < \Omega$, then $\mathbb{A}(\zeta) = \zeta + 1 < \theta$, as θ is a limit since it is a multiple of ω^{ω^α} . Otherwise, write $\zeta = \Omega(1 + \alpha') + \beta'$. For $x' = x_{\alpha'}^{\beta'}$ and $y' = y_{\alpha'}$, we have by induction that $\mathbb{A}(\zeta) \leq \omega^{\omega^{\alpha'}}(y' + \beta' + x')$. Consider the following cases.

CASE 1.1 ($\beta' < \gamma$). Then $y' + \beta' + x' < \gamma$, since γ must be a limit. The induction hypothesis yields

$$\mathbb{A}(\zeta) < \omega^{\omega^\alpha} \gamma < \theta.$$

CASE 1.2 ($\gamma \leq \beta' < \beta$). Then $x' = x$ and β is a successor. Consider two sub-cases.

CASE 1.2.1 ($\alpha' = \alpha$). Then also $y' = y$ and the induction hypothesis yields $\mathbb{A}(\zeta) \leq \omega^{\omega^\alpha}(y + \beta + x - 1) < \theta$.

CASE 1.2.2 ($\alpha' < \alpha$). The induction hypothesis yields

$$\begin{aligned} \mathbb{A}(\zeta) &\leq \omega^{\omega^{\alpha'}}(y' + \beta + x - 1) = \omega^{\omega^{\alpha'}} y' + \omega^{\omega^{\alpha'}}(\beta + x - 1) \\ &< \omega^{\omega^{\alpha'}} y' + \omega^{\omega^\alpha}(y + \beta + x) = \omega^{\omega^\alpha}(y + \beta + x). \end{aligned}$$

CASE 1.3 ($\beta' \geq \beta$). In this case we have that $\beta' < \theta$ and $\alpha' < \alpha$. Then, $\beta' = \omega^{\omega^\alpha} \delta + \eta$ for some $\delta < (y + \beta + x)$ and some $\eta < \omega^{\omega^\alpha}$. The induction hypothesis yields

$$\begin{aligned} \mathbb{A}(\zeta) &\leq \omega^{\omega^{\alpha'}}(y' + \omega^{\omega^\alpha} \delta + \eta + x') \\ &= \omega^{\omega^{\alpha'}} y' + \omega^{\omega^{\alpha'}}(\omega^{\omega^\alpha} \delta) + \omega^{\omega^{\alpha'}}(\eta + x') \\ &= \omega^{\omega^{\alpha'}} y' + \omega^{\omega^\alpha} \delta + \omega^{\omega^{\alpha'}}(\eta + x') \\ &< \omega^{\omega^{\alpha'}} y' + \omega^{\omega^\alpha} \delta + \omega^{\omega^\alpha} = \omega^{\omega^{\alpha'}} y' + \omega^{\omega^\alpha}(\delta + 1) \\ &= \omega^{\omega^\alpha}(\delta + 1) \leq \omega^{\omega^\alpha}(y + \beta + x) = \theta. \square \end{aligned}$$

We need some auxiliary definitions before giving the fundamental sequences for the ϑ function. We will use the function $\tau_\Omega(\xi)$ defined in Section 2, and write τ for τ_Ω . If α is a limit and $\beta < \Omega$, we define $\vartheta^{(i)}(\alpha + \beta)$ recursively by $\vartheta^{(0)}(\alpha + \beta) = \beta$ and $\vartheta^{(i+1)}(\alpha + \beta) = \vartheta([\alpha|\vartheta^{(i)}(\alpha + \beta)]_\Omega$.

Definition 9.3. For an ordinal $\alpha < \varepsilon_{\Omega+1}$, we define

$$\text{FIX}(\alpha) = \{\vartheta(\gamma) : \alpha + \Omega \leq \gamma \text{ and } \text{mc}_\Omega(\alpha) < \vartheta(\gamma)\}.$$

For $\alpha = \Omega\alpha'$ and $\beta < \Omega$, we set

1. $\vartheta_\alpha^*(0) = \begin{cases} \tau(\alpha) & \text{if } \text{mc}_\Omega([\alpha|1]_\Omega) < \text{mc}_\Omega(\alpha) \\ = \tau(\alpha) = \vartheta(\alpha_1) \text{ with } \alpha_1 > \alpha, \\ 0 & \text{otherwise,} \end{cases}$
2. $\vartheta_\alpha^*(\beta + 1) = \vartheta(\alpha + \beta)$, and
3. $\vartheta_\alpha^*(\beta) = \beta$ if β is a limit.

Then we set

1. $\{\alpha + \beta|n\} = \alpha + \{\beta|n\}$ if $\beta > 0$ and $\alpha + \beta$ is in Cantor normal form.
2. $\{\vartheta(\Omega\alpha + \beta)|n\} = \vartheta(\Omega\alpha + \{\beta|n\})$ if $\beta \in \text{Lim} \setminus \text{FIX}(\alpha)$.
3. In other cases, $\{\vartheta(\Omega\alpha + \beta)|n\}$ is given by
 - (a) $\vartheta_\alpha^*(\beta) \cdot n$ if $\alpha = 0$,
 - (b) $\vartheta^{(n)}(\alpha + \vartheta_\alpha^*(\beta))$ if $\tau(\alpha) = \Omega$, and
 - (c) $\vartheta([\alpha|\{\tau(\alpha)|n\}] + \vartheta_\alpha^*(\beta))$ if $\tau(\alpha) \in (1, \Omega)$.

Recall from the introduction that KP is a restriction of ZFC with proof-theoretic ordinal $\vartheta(\varepsilon_{\Omega+1})$. This ordinal can be used to bound the provably total computable functions of KP.

Definition 9.4. For $i < \omega$ and $\alpha < \vartheta(\varepsilon_{\Omega+1})$, define $\{\alpha|i\}^*$ recursively by $\{\alpha|0\}^* = \alpha$ and $\{\alpha|i+1\}^* = \{\{\alpha|i\}^*|i+1\}$. Define $F(n)$ to be the least ℓ such that $\{\vartheta(\Omega_n)|\ell\}^* = 0$.

The function F is total since $\{\vartheta(\Omega_n)|i+1\}^* < \{\vartheta(\Omega_n)|i\}^*$ whenever the right-hand side is not zero, but this fact is not provable in KP. The following is a standard result from proof theory; see e.g. [5].

Theorem 9.5. Let φ be Σ_1^0 formula and suppose that $\text{KP} \vdash \forall x \exists y \varphi(x, y)$. Let $f_\varphi(n)$ be the least m such that $\varphi(n, m)$. Then, $\exists m \forall n > m f_\varphi(n) < F(n)$.

In order to compare $F(n)$ to the length of our fast Goodstein walks, the following property will be useful. It is shown to hold in general for any system of fundamental sequences with the Bachmann property in [6].

Proposition 9.6. Let $(\xi_n)_{n \in \mathbb{N}}$ be a sequence of ordinals below $\vartheta(\varepsilon_{\Omega+1})$ such that, for all n , $\xi_n[n+1] \leq \xi_{n+1} \leq \xi_n$. Then, for all n , $\xi_n \geq \{\xi|n\}^*$.

10 Independence

Our strategy to prove that Theorems 8.2 and 8.5 are not provable in KP is to show that they grow at least as fast as the function F of Definition 9.4. By Proposition 9.6, it suffices to show that $\{\uparrow_{k+2}^\omega m|k\} \leq \uparrow_{k+2}^\omega(m-1)$ for all k , as this will allow us to use Proposition 9.6 to prove that $\{\uparrow_2^\omega m|k\}^* \leq \uparrow_{k+2}^\omega \mathcal{G}_k m$. By choosing suitable $m = m(n)$, this will show that the termination time for the fast Goodstein process is bounded below by $F(n)$. We begin with a useful lemma, which we use within an inductive argument later.

Lemma 10.1. *Let $k \geq 2$ and write \uparrow for \uparrow_k^ω , $\mathbb{A} = \mathbb{A}_k$, and $\mathbb{B} = \mathbb{A}_\omega$. Suppose that $\xi \in [\omega^2, \varepsilon_0)$ is such that $\uparrow n = (\uparrow n)_\vartheta$ for all $n < \mathbb{A}(\xi)$. Write $\xi = \alpha + b$ with α zero or a limit. Then,*

$$\vartheta_{\Omega - \Omega^2 + \uparrow \alpha}^*(\uparrow b) \leq \uparrow \mathbb{A}(\xi - 1).$$

Proof. Let $\alpha' = \Omega - \Omega^2 + \uparrow \alpha$ and consider the following cases.

CASE 1 ($b = 0$ and $\text{mc}_\Omega([\alpha'|1]_\Omega) < \text{mc}_\Omega(\alpha') = \tau(\alpha') = \vartheta(\tilde{\alpha})$ with $\tilde{\alpha} > \alpha'$). Then, $\vartheta_{\alpha'}^*(0) = \text{mc}_\Omega(\alpha')$. We claim that $\text{mc}_\Omega(\alpha') = \text{mc}_\Omega(\uparrow \alpha)$. Write $\uparrow a = \Omega^2 \alpha_1 + \Omega \alpha_0$ and $\alpha' = \Omega^2 \alpha'_1 + \Omega \alpha'_0$ with $\alpha_0, \alpha'_0 < \Omega$. Note that $\alpha'_1 = -1 + \alpha_1$ and $\alpha'_0 \in \{\alpha_0, 1 + \alpha_0\}$. If $\alpha'_0 > 0$, then $\tau(\alpha') = \alpha'_0$. From $\tau(\alpha') = \vartheta(\tilde{\alpha})$, we see that α'_0 is a limit, hence $\alpha'_0 = \alpha_0$. But then, Lemma 3.3 yields $\alpha'_0 \leq \text{mc}_\Omega(\uparrow a) < \mathbb{A}(\xi - 1)$. If $\alpha'_0 = 0$, then similar reasoning shows that α'_1 cannot be finite, hence $\alpha'_1 = \alpha_1$. It follows that $\tau(\alpha') = \tau(\Omega^2 \alpha'_1) = \tau(\Omega^2 \alpha_1) \leq \text{mc}_\Omega(\uparrow a) < \mathbb{A}(\xi - 1)$.

CASE 2 (β is a successor). Then, $\beta - 1 = \uparrow(b - 1)$. By Theorem 6.4 and the assumption, we see that

$$\begin{aligned} \uparrow \mathbb{A}(\xi - 1) &\geq \mathbb{B}(\uparrow \alpha + \uparrow(b - 1)) \\ &= \vartheta(\Omega - \Omega^2 + \uparrow \alpha + \uparrow(b - 1)) = \vartheta_{\Omega - \Omega^2 + \uparrow \alpha}^*(\uparrow b). \end{aligned}$$

CASE 3 (other cases). Otherwise, $\vartheta_{\Omega - \Omega^2 + \uparrow \alpha}^*(\uparrow b) \in \{0, \uparrow b\}$, so $\vartheta_{\Omega - \Omega^2 + \uparrow \alpha}^*(\uparrow b) \leq \uparrow b$. Since $\mathbb{A}(\xi - 1) > b$ by Lemma 3.3, it follows that $\uparrow \mathbb{A}(\xi - 1) > \vartheta_{\Omega - \Omega^2 + \uparrow \alpha}^*(\uparrow b)$. \square

Lemma 10.2. *If $m \in \mathbb{N}$ and $k < \omega$, then*

- (1) $\{\uparrow_{k+2}^\omega m|k\} \leq \uparrow_{k+2}^\omega(m-1)$, and
- (2) $\uparrow_{k+2}^\omega m = (\uparrow_{k+2}^\omega m)_\vartheta$.

Proof. We prove the claims simultaneously by induction on m . Write \uparrow for \uparrow_{k+2}^ω , \mathbb{A} for \mathbb{A}_{k+2} , \mathbb{B} for \mathbb{A}_ω , and let $m =_k \mathbb{A}(\xi)$. For the induction to go through, we need to additionally show the side condition that if $\uparrow m = \gamma + \delta$ in Cantor normal form and $\delta = \omega^\delta$, then $\uparrow(m-1) \geq \gamma + \omega^{\{\delta|k\}}$.

Proof of Claim (1). Write $m =_{k+2} \mathbb{A}(\xi)$ and consider the following cases.

CASE 1 ($\xi < \omega$). Then $m - 1 = \xi$ and $\uparrow m = \mathbb{B}(\uparrow \xi) = \uparrow \xi + 1$, so

$$\uparrow(m - 1) \geq \uparrow \xi = \{\uparrow m | k\}.$$

We do not check the side condition as $1 < \omega^1$.

CASE 2 ($\omega \leq \xi < \omega^2$). Write $\xi = \omega a + b$ with $a > 0$ and let $\alpha = -1 + \uparrow a$, $\beta = \uparrow b$. Letting $x = x_\alpha^\beta$ and $y = y_\alpha$, we have that $(\mathbb{A}(\xi))_\vartheta = \omega^{\omega^\alpha}(y + \beta + x)$. In this case, we claim that $\{\uparrow m | i\} \leq \uparrow \mathbb{A}^{(i+1)}(\xi)$ for all $i < k$, which suffices as $\mathbb{A}^{(k+1)}(\xi) < m - 1$. Consider the following cases.

CASE 2.1 ($y + \beta + x = 0$). This case is impossible since $\beta = 0$ implies that $\omega^{\omega^\alpha} \beta = \beta$, so $x = 1$.

CASE 2.2 ($y + \beta + x$ is a successor). We first claim that

$$\omega^{\omega^\alpha}(y + \beta + x - 1) \leq \uparrow \mathbb{A}^{(0)}(\xi - 1).$$

First assume that β is a successor. Note that in this case, $\beta - 1 = \uparrow(b - 1)$, and it is not hard to check that $x_\alpha^{\beta-1} = x$, so that

$$\begin{aligned} \omega^{\omega^\alpha}(y + \beta + x - 1) &= \mathbb{B}(\uparrow(\xi - 1)) \\ &\leq \uparrow \mathbb{A}(\xi - 1) = \uparrow \mathbb{A}^{(0)}(\xi - 1). \end{aligned}$$

If β is a limit then we must have $x = 1$ and $\beta = \gamma$, so

$$\begin{aligned} \beta &= \gamma = \omega^{\omega^\alpha} \gamma = \omega^{\omega^\alpha} \beta \\ &= \omega^{\omega^\alpha}(y + \beta + x - 1). \end{aligned}$$

However, $\mathbb{A}(\xi - 1) > b$, so $\uparrow \mathbb{A}^{(0)}(\xi - 1) = \uparrow \mathbb{A}(\xi - 1) > \uparrow b = \beta$. To continue, consider two sub-cases.

CASE 2.2.1 ($\alpha = 0$). In this case we claim that $\{\omega^{\omega^\alpha}(y + \beta + x) | i\} \leq \mathbb{A}^{(i)}(\xi)$. We have that

$$\omega^{\omega^\alpha}(y + \beta + x) = \omega(y + \beta + x) = \omega(y + \beta + x - 1) + \omega$$

and $\{\omega^{\omega^\alpha}(y + \beta + x) | i\} = \omega(y + \beta + x - 1) + i$. For $i = 0$, we already have that $\omega(y + \beta + x - 1) \leq \mathbb{A}^{(0)}(\xi - 1)$, and for $i + 1$, we see that

$$\begin{aligned} \uparrow \mathbb{A}^{(i+1)}(\xi) &= \uparrow \mathbb{A}([\omega | \mathbb{A}^{(i+1)}(\xi)]) = \uparrow \mathbb{A}(\mathbb{A}^{(i)}(\xi)) \\ &\leq \mathbb{B}(\uparrow \mathbb{A}^{(i)}(\xi)) \stackrel{\text{IH}}{\leq} \mathbb{B}(\omega(y + \beta + x - 1) + i) \\ &= \omega(y + \beta + x - 1) + i + 1. \end{aligned}$$

CASE 2.2.2 ($\alpha > 0$). We claim that $\{\omega^{\omega^\alpha}(y + \beta + x) | i\} \leq \mathbb{A}^{(i+1)}(\xi)$ if $i \leq k$. In this case it is possible that α is an epsilon-number, so we prove the stronger claim that

$$\omega^{\omega^\alpha}(y + \beta + x - 1) + \omega^{\{\omega^\alpha | i\}} \leq \mathbb{A}^{(i+1)}(\xi).$$

Note that $\{\alpha|i\} \leq \{\alpha|k\} \leq \uparrow(a-1)$, the latter being by the induction hypothesis. Moreover, $\{\omega^\alpha|i\} \leq \omega^{\{\alpha|k\}i}$ if $i \geq 1$. So, define $\alpha' = \{\alpha|k\}$; we will show that

$$\omega^{\omega^\alpha}(y + \beta + x - 1) + \omega^{\omega^{\alpha'}i} \leq \mathbb{A}^{(i+1)}(\xi). \quad (5)$$

For $i = 0$, we have that

$$\begin{aligned} \uparrow\mathbb{A}^{(1)}(\xi) &\geq \uparrow\mathbb{A}^{(0)}(\xi) + 1 \geq \omega^{\omega^\alpha}(y + \beta + x - 1) + 1 \\ &= \omega^{\omega^\alpha}(y + \beta + x - 1) + \omega^{\omega^{\alpha'}0}. \end{aligned}$$

For $x' = x_{\alpha'}^{\mathbb{A}^{(i)}(\xi)}$ and $y' = y_{\alpha'}$, the induction hypothesis yields

$$\begin{aligned} \uparrow\mathbb{A}^{(i+1)}(\xi) &= \uparrow\mathbb{A}(\omega(a-1) + \mathbb{A}^{(i)}(\xi)) \\ &\geq \mathbb{B}(\Omega\alpha' + \mathbb{A}^{(i)}(\xi)) \\ &= \omega^{\omega^{\alpha'}}(y' + \mathbb{A}^{(i)}(\xi) + x') \\ &\geq \omega^{\omega^{\alpha'}}(y' + \omega^{\omega^\alpha}(y + \beta + x - 1) + \omega^{\omega^{\alpha'}i}) \\ &= \omega^{\omega^{\alpha'}}y' + \omega^{\omega^\alpha}(y + \beta + x - 1) + \omega^{\omega^{\alpha'}(i+1)} \\ &\geq \omega^{\omega^\alpha}(y + \beta + x - 1) + \omega^{\omega^{\alpha'}(i+1)}. \end{aligned}$$

Setting $i = k$, we see that

$$\omega^{\omega^\alpha}(y + \beta + x - 1) + \omega^{\{\omega^\alpha|k\}} \leq \mathbb{A}^{(k+1)}(\xi) \leq \uparrow(m-1),$$

as needed.

CASE 2.3 ($y + \beta + x$ is a limit). Then forcibly $x = 0$ and β is a limit, so $y + \beta = \beta$. Write $\beta = \beta_0 + \omega^\delta$ in Cantor normal form. Then,

$$(\mathbb{A}(\xi))_\vartheta = \omega^{\omega^\alpha}\beta = \omega^{\omega^\alpha}(\beta_0 + \omega^\delta) = \omega^{\omega^\alpha}\beta_0 + \omega^{\omega^\alpha+\delta},$$

and

$$\{(\mathbb{A}(\xi))_\vartheta|k\} = \omega^{\omega^\alpha}\beta_0 + \{\omega^{\omega^\alpha+\delta}|k\}.$$

The induction hypothesis yields $\uparrow(b-1) = \beta_0 + \eta$ for some $\eta \geq \{\omega^\delta|k\}$, such that if $\omega^\delta = \delta$ then $\eta \geq \omega^{\{\omega^\delta|k\}}$. For suitable x' and y' ,

$$\begin{aligned} \uparrow\mathbb{A}^{(0)}(\xi - 1) &= \uparrow\mathbb{A}(\xi - 1) = (\uparrow(\xi - 1))_\vartheta = \omega^{\omega^\alpha}(y' + \uparrow(b-1) + x') \\ &\geq \omega^{\omega^\alpha}\uparrow(b-1) = \omega^{\omega^\alpha}(\beta_0 + \eta) = \omega^{\omega^\alpha}\beta_0 + \omega^{\omega^\alpha}\eta. \end{aligned}$$

Note that from $x = 0$ we see that $\omega^\alpha + \delta > \delta$, hence $\omega^\alpha + \delta$ is in Cantor normal form and $\omega^{\omega^\alpha+\delta} > \omega^\alpha + \delta$ as the latter is not additively indecomposable (hence it is not an epsilon-number). To proceed, consider the following cases.

CASE 2.3.1 ($\delta < \omega^\delta$). By a case-by-case analysis we see that $\{\omega^{\omega^\alpha+\delta}|k\} = \omega^{\omega^\alpha}\{\omega^\delta|k\}$. Since $\eta \geq \{\omega^\delta|k\}$, we readily see that

$$\begin{aligned} \uparrow\mathbb{A}^{(0)}(\xi - 1) &= \omega^{\omega^\alpha}\beta_0 + \omega^{\omega^\alpha}\eta \\ &\geq \omega^{\omega^\alpha}\beta_0 + \omega^{\omega^\alpha}\{\omega^\delta|k\} = \{(\uparrow m)_\vartheta|k\}. \end{aligned}$$

CASE 2.3.2 ($\delta = \omega^\delta$). Then the side condition for ε -numbers yields $\eta \geq \omega^{\{\delta|k\}}$, and $\{\omega^{\omega^\alpha + \delta}|k\} = \omega^{\omega^\alpha + \{\delta|k\}}$, so

$$\uparrow \mathbb{A}^{(0)}(\xi - 1) \geq \omega^{\omega^\alpha} \beta_0 + \omega^{\omega^\alpha + \{\delta|k\}} = \{(\uparrow m)_{\vartheta}|k\}.$$

CASE 3 ($\omega^2 \leq \xi$). Write $\xi = \alpha + b$ with $\alpha = \omega \tilde{\alpha}$, so that $\uparrow \xi = \uparrow \alpha + \uparrow b$. Then, we claim that

$$\begin{aligned} \uparrow m &= \mathbb{B}(\uparrow \xi) = \mathbb{B}(\uparrow \alpha + \uparrow b) \\ &= \vartheta(\Omega - \Omega^2 + \uparrow \alpha + \uparrow b) = \vartheta(\zeta), \end{aligned}$$

where $\zeta = \gamma + \uparrow b$ with $\gamma = \Omega - \Omega^2 + \uparrow \alpha$. Note that $\zeta \geq \Omega$, so $\vartheta(\zeta) = \omega^{\vartheta(\zeta)}$, meaning that we must prove that $\uparrow(m - 1) \geq \omega^{\{\vartheta(\zeta)|k\}}$. Consider three subcases.

CASE 3.1 ($\uparrow b \in \text{Lim} \setminus \text{FIX}(\gamma)$). Then,

$$\begin{aligned} \uparrow(m - 1) &= \uparrow(\mathbb{A}(\xi) - 1) \geq \uparrow \mathbb{A}^{(0)}(\xi) = \uparrow \mathbb{A}(\alpha + (b - 1)) \\ &\geq \mathbb{A}(\uparrow \alpha + \uparrow(b - 1)) \stackrel{\text{IH}}{\geq} \uparrow \mathbb{A}(\uparrow \alpha + \{\uparrow b|k\}) \\ &\stackrel{\text{IH}}{=} \vartheta(\gamma + \{\uparrow b|k\}) = \omega^{\vartheta(\gamma + \{\uparrow b|k\})} = \omega^{\{\vartheta(\gamma + \uparrow b)|k\}} = \omega^{\{\vartheta(\zeta)|k\}}. \end{aligned}$$

CASE 3.2 ($\alpha = \omega^2$ and $\uparrow b \notin \text{Lim} \setminus \text{FIX}(\Omega)$). Then $\xi = \omega^2 + b$, so that $\uparrow \xi = \omega^2 + \uparrow b$, and $\zeta = \Omega + \uparrow b$. We have that $\{\vartheta(\zeta)|k\} = \vartheta^{(k)}(\zeta)$ and $m - 1 \geq \mathbb{A}^{(k+1)}(\xi)$, so $\uparrow(m - 1) \geq \uparrow \mathbb{A}^{(k+1)}(\xi)$. But induction on i shows that $\mathbb{A}^{(i+1)}(\uparrow \xi) > \omega_i(\vartheta_\Omega^*(\uparrow b) + 1)$. Note that

$$\uparrow \mathbb{A}^{(i+1)}(\xi) = \uparrow \mathbb{A}([\Omega^2 | \mathbb{A}^{(i)}(\xi)]) = \mathbb{B}([\Omega^2 | \uparrow \mathbb{A}^{(i)}(\xi)]) = \mathbb{B}(\Omega \uparrow \mathbb{A}^{(i)}(\xi)).$$

If $i = 0$, we see that $\uparrow \mathbb{A}^{(0)}(\xi) = \uparrow \mathbb{A}(\xi - 1)$. By Lemma 10.1 and the induction hypothesis applied to the second claim, $\uparrow \mathbb{A}(\xi - 1) \geq \vartheta_\Omega^*(\uparrow b)$. With this, we see that

$$\mathbb{B}(\Omega \uparrow \mathbb{A}^{(0)}(\xi)) \geq \mathbb{B}(\Omega \vartheta_\Omega^*(\uparrow b)) > \vartheta_\Omega^*(\uparrow b) + 1,$$

and for the inductive step,

$$\begin{aligned} \mathbb{B}(\Omega \uparrow \mathbb{A}^{(i)}(\xi)) &\stackrel{\text{IH}}{\geq} \mathbb{B}(\Omega \omega_{i-1}(\vartheta_\Omega^*(\uparrow b) + 1)) \\ &= \omega_{i+1}(\vartheta_\Omega^*(\uparrow b) + 1) \geq \omega^{\omega_i(\vartheta_\Omega^*(\uparrow b) + 1)}. \end{aligned}$$

Setting $i = k + 1$, we see that

$$\uparrow(m - 1) \geq \omega^{\{\vartheta(\zeta)|k\}} = \omega^{\{\uparrow m|k\}},$$

as needed.

CASE 3.3 ($\alpha > \omega^2$ and $\uparrow b \notin \text{Lim} \setminus \text{FIX}(\gamma)$). Write $\alpha = \omega^2 \alpha_1 + \alpha_0$ and let $v = 1$ if $\alpha_1 \leq k$, otherwise $v = 0$, and let $w = 1$ if $\alpha_1 = 1$, otherwise $w = 0$. Clearly $0 \leq w \leq v \leq 1$. Letting $\gamma = -\omega^2 v + \alpha$, we see that $\omega^2 v + \gamma$ is in Cantor normal

form, so that $[\omega^2 v + \gamma|i] = \omega^2 v + [\gamma|i]$ for all i . Moreover, $\uparrow\alpha = \Omega^2 v + \uparrow\gamma$ is in Cantor normal form, as is $\Omega w + \uparrow\gamma$, from which we see that $[\Omega w + \uparrow\gamma|\chi]_\Omega = \Omega w + [\uparrow\gamma|\chi]_\Omega$ for all χ .

Let $\theta = \vartheta_{\Omega w + \uparrow\gamma}^*(\uparrow b)$ and $A = \mathbb{A}(\alpha + b - 1)$. As before, Lemma 10.1 yields $\theta \leq \uparrow A$. Let $\tau = \tau(\Omega w + \uparrow\gamma)$ and consider two sub-cases.

CASE 3.3.1 ($0 < \tau < \Omega$). We prove by induction on ρ that if $\omega\rho \leq \gamma$ and $\tau(\uparrow\omega\rho) = \tau$, then

$$[\uparrow\omega\rho|\{\tau|k\}]_\Omega + \theta \leq \uparrow[\omega\rho|A]. \quad (6)$$

The claim will then follow, as from this we also obtain that $\Omega w + [\uparrow\omega\rho|\{\tau|k\}]_\Omega + \theta \leq \Omega w + \uparrow[\omega\rho|A]$. Moreover,

$$\text{mc}_\Omega(\Omega w + [\uparrow\omega\rho|\{\tau|k\}]_\Omega + \theta) = \theta < A = \text{mc}_\Omega(\Omega w + \uparrow[\omega\rho|A]),$$

so that the induction hypothesis on claim (2) yields

$$\begin{aligned} \{\vartheta(\Omega w + \uparrow\gamma + \uparrow b)|k\}_\Omega &= \vartheta(\Omega w + [\uparrow\gamma|\{\tau|k\}]_\Omega + \theta) \leq \vartheta(\Omega w + \uparrow[\gamma|A]) \\ &\stackrel{\text{IH}}{=} \mathbb{B}(\Omega^2 v + \uparrow[\gamma|A]) = \uparrow\mathbb{A}(\omega^2 v + \uparrow[\gamma|A]) \\ &= \uparrow\mathbb{A}([\alpha|A]) = \uparrow\mathbb{A}^{(1)}(\alpha + b) < \uparrow\mathbb{A}(\alpha + b). \end{aligned}$$

Moreover, $\Omega w + [\uparrow\gamma|\{\tau|k\}]_\Omega + \theta \geq \Omega$, so $\{\vartheta(\Omega w + \uparrow\gamma + \uparrow b)|k\}_\Omega = \omega^{\{\vartheta(\Omega w + \uparrow\gamma + \uparrow b)|k\}_\Omega}$, meaning that we do not need to check the side-condition for epsilon-numbers separately.

So, it remains to prove (6). Write $\omega\rho = \omega^{\rho_1} c + \rho_0$ in coefficient ω -normal form and consider the following cases.

CASE (i) ($\rho_0 > 0$). Then we have inductively that

$$\begin{aligned} [\uparrow\omega\rho|\{\tau|k\}]_\Omega + \theta &= \Omega^{\uparrow\rho_1} \uparrow c + [\uparrow\rho_0|\{\tau|k\}]_\Omega + \theta \\ &\leq \Omega^{\uparrow\rho_1} \uparrow c + \uparrow[\rho_0|A] = \uparrow[\omega\rho|A]. \end{aligned}$$

CASE (ii) ($\rho_0 = 0$ and $\uparrow c \in \text{Lim}$). Then, $\uparrow c = \tau$ by the definition of τ , and $[\omega\rho|A] = \omega^{\rho_1}(c - 1) + \delta$ for some δ with $\text{mc}(\delta) \geq A$. From this we see that

$$\begin{aligned} \uparrow[\omega\rho|A] &= \uparrow(\omega^{\rho_1}(c - 1) + \delta) = \Omega^{\uparrow\rho_1} \uparrow(c - 1) + \uparrow\delta \\ &\stackrel{\text{IH}}{\geq} \Omega^{\uparrow\rho_1} \{\tau|k\} + \theta = [\uparrow\omega\rho|\{\tau|k\}]_\Omega + \theta. \end{aligned}$$

CASE (iii) ($\rho_0 = 0$, $\uparrow c \in \text{Succ}$, and $\rho_1 = \omega\rho'$). Write $\uparrow(c - 1) = \eta$, so that $\uparrow c = \eta + 1$. Then, $\tau(\uparrow\rho_1) = \tau(\uparrow\omega\rho)$, so the induction hypothesis and Lemma 2.1 yield

$$\begin{aligned} [\uparrow\omega^{\rho_1} c|\{\tau|k\}]_\Omega + \theta &= \uparrow\omega^{\rho_1}(c - 1) + \Omega^{[\uparrow\rho_1|\{\tau|k\}]_\Omega} + \theta \\ &\leq \uparrow\omega^{\rho_1}(c - 1) + \Omega^{[\uparrow\rho_1|\{\tau|k\}]_\Omega + \theta} \\ &\stackrel{\text{IH}}{\leq} \Omega^{\uparrow\rho_1} \eta + \Omega^{\uparrow[\rho_1|A]} = \uparrow[\omega^{\rho_1}|A]. \end{aligned}$$

CASE (iv) ($\rho_0 = 0, \uparrow c, \rho_1 \in \text{Succ}$). Then, $\rho_1 = \omega\rho' + t$ for some finite $t > 0$. The assumption that $\tau(\omega\rho) = \tau$ yields $\uparrow t = \tau$, and the main induction hypothesis yields $\uparrow(t-1) \geq \{\tau|k\}$.

If $\theta = 0$, note that $A \geq 1$, so

$$\begin{aligned} \uparrow[\omega\rho|A] &= \uparrow[\omega^{\omega\rho'+t}c|A] = \uparrow(\omega^{\omega\rho'+t}(c-1) + \omega^{\omega\rho'+t-1}A) \\ &\geq \uparrow(\omega^{\omega\rho'+t}(c-1) + \omega^{\omega\rho'+t-1}) = \Omega^{\uparrow\rho_1}\eta + \Omega^{\uparrow\omega\rho'+\uparrow(t-1)} \\ &\stackrel{\text{IH}}{\geq} \Omega^{\uparrow\rho_1}\eta + \Omega^{\uparrow\omega\rho'+\{\tau|k\}} = [\uparrow\omega^{\rho_1}c|\{\tau|k\}]_{\Omega} + \theta. \end{aligned}$$

Otherwise, θ is a limit ordinal, so that $A \geq \theta > 1$. Since also $\omega^{\omega\rho'+t-1} > 1$, using Lemma 2.1 we see that

$$\begin{aligned} \uparrow[\omega\rho|A] &= \uparrow[\omega^{\omega\rho'+t}c|A] = \uparrow(\omega^{\omega\rho'+t}(c-1) + \omega^{\omega\rho'+t-1}A) \\ &\leq \uparrow(\omega^{\omega\rho'+t}(c-1) + \omega^{\omega\rho'+t-1} + A) \\ &= \Omega^{\uparrow\rho_1}\eta + \Omega^{\uparrow\omega\rho'+\uparrow(t-1)} + \uparrow A \stackrel{\text{IH}}{\geq} \Omega^{\uparrow\rho_1}\eta + \Omega^{\uparrow\omega\rho'+\{\tau|k\}} + \uparrow A \\ &= [\uparrow\omega^{\rho_1}c|\{\tau|k\}]_{\Omega} + \uparrow A \geq [\uparrow\omega^{\rho_1}c|\{\tau|k\}]_{\Omega} + \theta. \end{aligned}$$

CASE 3.3.2 ($\tau = \Omega$). Define $\theta_i = \vartheta^{(i)}(\Omega w + \uparrow\gamma + \theta)$. We claim that $\theta_i \leq \uparrow\mathbb{A}^{(i)}(\alpha + b)$ for all i . Since $\uparrow(m-1) \leq \uparrow\mathbb{A}^{(k+1)}(\alpha + b)$ and $\{\vartheta(\Omega w + \uparrow\gamma)|k\} = \theta_k$, this yields the desired result.

For $i = 0$, we already have that $\theta \leq \uparrow\mathbb{A}(\alpha + b - 1) = \uparrow\mathbb{A}^{(0)}(\alpha + b)$. For $i + 1$, we see that

$$\begin{aligned} \theta_{i+1} &= \vartheta(\Omega w + [\uparrow\gamma|\theta_i]_{\Omega}) \\ &\leq \mathbb{B}(\Omega^2 v + [\uparrow\gamma|\theta_i]_{\Omega}) && \text{by main IH} \\ &\leq \mathbb{B}(\Omega^2 v + [\uparrow\gamma|\uparrow\mathbb{A}^{(i)}(\alpha + b)]_{\Omega}) && \text{by induction on } i \\ &= \mathbb{B}(\Omega^2 v + \uparrow[\gamma|\mathbb{A}^{(i)}(\alpha + b)]) && \text{by Lemma 5.10 and } \tau(\uparrow\gamma) = \Omega \\ &= \uparrow\mathbb{A}(\omega^2 v + [\gamma|\mathbb{A}^{(i)}(\alpha + b)]) \\ &= \uparrow\mathbb{A}^{(i+1)}(\alpha + b). \end{aligned}$$

As above, $\{\vartheta(\Omega w + \uparrow\gamma)|k\}$ is an epsilon-number, so we obtain the side condition for free.

Proof of Claim (2). By Lemma 9.2, $\uparrow_{k+2}^{\omega} m \leq (\uparrow_{k+2}^{\omega} m)_{\vartheta}$. Now, let $\theta < (\uparrow_{k+2}^{\omega} m)_{\vartheta}$. Let ℓ be such that $\theta < \{(\uparrow m)_{\vartheta}|\ell\}$; since fundamental sequences are increasing, we may assume $\ell \geq k$. By Corollary 7.2, $\uparrow_{k+2}^{\omega} \uparrow_{k+2}^{\ell+2} m = \uparrow_{k+2}^{\omega} m$, and by the first claim,

$$\uparrow_{k+2}^{\omega} (\uparrow_{k+2}^{\ell+2} m - 1) \geq \{(\uparrow m)_{\vartheta}|\ell\} > \theta.$$

Since $\theta < (\uparrow m)_{\vartheta}$ was arbitrary, we conclude that $\uparrow_{k+2}^{\omega} m \geq (\uparrow_{k+2}^{\omega} m)_{\vartheta}$, as needed. \square

The following corollary, while not needed for our main results, provides a direct comparison of the ϑ and \mathbb{B} functions.

Corollary 10.3. *For all $\xi < \vartheta(\varepsilon_{\Omega+1})$,*

1. $\xi = (\xi)_{\vartheta}$, and
2. there exist $k \geq 2$ and $\xi < \varepsilon_0$ such that $\zeta = \uparrow_k^{\omega} \xi$.

Proof. The first claim is a consequence of the second, since then $\xi = \uparrow_k^{\omega} m$ for suitable m, k , and Lemma 10.2 yields

$$\xi = \uparrow_k^{\omega} m = (\uparrow_k^{\omega} m)_{\vartheta} = (\xi)_{\vartheta}.$$

Thus we focus on the second claim. Then, $\xi < \vartheta(\Omega_i)$ for some i , and

$$\uparrow_2^{\omega} \mathbb{A}(\omega_i) \geq \mathbb{B}(\uparrow_2^{\omega} \omega_i) = \mathbb{B}(\Omega_i) > \xi.$$

Thus there exist m' and $k' \geq 2$ such that $\uparrow_{k'}^{\omega} m' \geq \xi$. Let $\xi' \geq \xi$ be least with the property that $\uparrow_{k'}^{\omega} m' \geq \xi'$ for some m' and $k' \geq 2$.

We claim that $\xi' = \xi$. If not, let $k \geq k'$ be large enough so that $\{\xi'|k\} \geq \xi$, and let $m = \uparrow_{k'}^{k'+2} m'$. By Lemma 10.2, we have that

$$\xi \leq \{\xi'|k\} \leq \uparrow_{k+2}^{\omega}(m-1) < \uparrow_{k+2}^{\omega} m = \xi'.$$

But $\zeta := \uparrow_{k+2}^{\omega}(m-1)$ contradicts the minimality of ξ' . Thus we conclude that $\xi' = \xi$, as desired. \square

Corollary 10.4. *Given $m, k \in \mathbb{N}$, $\uparrow_{k+2}^{\omega} \mathcal{G}_k m \geq \{\uparrow_2^{\omega} m|k\}^*$.*

Proof. We have that

$$\uparrow_{k+3}^{\omega} \mathcal{G}_{k+1} m = \uparrow_{k+3}^{\omega} (\mathcal{G}_k m - 1) \geq \{\mathcal{G}_k m|k\},$$

where the inequality is an instance of Lemma 10.2.(1). Proposition 9.6 then yields $\uparrow_{k+2}^{\omega} \mathcal{G}_k m \geq \{\uparrow_2^{\omega} m|k\}^*$ for all k . \square

Theorem 10.5. *Theorem 8.2 (and hence Theorem 8.5) is not provable in KP.*

Proof. Let $m(n) := \mathbb{A}_2(\omega_{n+2})$. It is not hard to check that $m(n)$ is in normal form, as $\xi > \omega_n$ implies also that $\text{mc}(\xi) \geq \text{mc}(\omega_n)$ and hence $\mathbb{A}(\xi) > \mathbb{A}(\omega_n)$ by Lemma 3.4. Then, $\uparrow_2^{\omega} m(n) = \mathbb{A}(\Omega_{n+2}) = \vartheta(\Omega_{n+2})$, the last inequality being by Lemma 10.2. Let $G(n)$ be the least value of k such that $\mathcal{G}_k m(n) = 0$. By Corollary 10.4,

$$\uparrow_{k+2}^{\omega} \mathcal{G}_k m(n) \geq \{\uparrow_2^{\omega} m(n)|k\}^* = \{\vartheta(\Omega_{n+2})|k\}^*.$$

Hence, $G(n) \geq F(n)$, as any value of k with $\mathcal{G}_k m(n) = 0$ also has $\{\vartheta(\Omega_{n+2})|k\}^* = 0$. The function $G(n)$ is clearly computable, hence definable by a Σ_1^0 formula. It follows that $\forall x \exists y (G(n) = y)$ is not provable in KP, hence neither is Theorem 6.4. \square

11 Concluding remarks

We have proven that Goodstein processes based on the \mathbb{A} function always terminate, leading to independence results of strength the Bachmann-Howard ordinal. This opens various lines of research. The \mathbb{A} function is not the only fast-growing function based on transfinite recursion below ε_0 . A natural question is how sensitive the termination and independence results presented here are to the precise choice of fast-growing functions used (e.g. the aforementioned Hardy function [10]), and in particular if the normal forms based on successive maximization we used yield maximality of base change in a more general context.

Variation of normal forms also leads to questions regarding independence. Note that the alternative normal forms of Example 8.6 give rise to a terminating Goodstein process, but our methods do not establish any lower bounds on such processes, so it is not immediately obvious whether termination is provable in KP. We conjecture that it is not, and remark that independence for this alternative Goodstein principle would also imply Theorem 10.5. Thus a different strategy for the current work would have been to prove that the maximal Goodstein principle terminates, and that the alternative one leads to independence, thus obtaining two independence results at once. The drawback of such a ‘dual’ approach is that, at least in this case, the two Goodstein processes would have to be studied separately, for example providing different ordinal assignments for each. We thus leave the analysis of this alternative Goodstein process for future work, but such a ‘dual’ approach may be interesting when studying less involved Goodstein processes.

Another direction involves Goodstein processes based on fast-growing hierarchies up to ordinals $\Lambda > \varepsilon_0$, perhaps on the Bachmann-Howard ordinal, or even larger ordinals where suitable systems of fundamental sequences are known, e.g. the ordinal of Π_1^1 -CA₀. The challenge is in identifying the base-change maximal normal forms in these contexts, and the proof-theoretic strength of termination.

References

- [1] T. Arai, D. Fernández-Duque, S. Wainer, and A. Weiermann. Predicatively unprovable termination of the Ackermannian Goodstein principle. *Proceedings of the American Mathematical Society*, 2019. Accepted for publication.
- [2] Toshiyasu Arai, Stanley S. Wainer, and Andreas Weiermann. Goodstein sequences based on a parametrized ackermann-péter function. *Bull. Symb. Log.*, 27(2):168–186, 2021.
- [3] J. Barwise. *Admissible sets and structures: An approach to definability theory*. Perspectives in mathematical logic. Springer-Verlag, Berlin, New York, 1975.

- [4] W. Buchholz. Ordinal notations and fundamental sequences, 2003. Manuscript.
- [5] E.A. Cichon, W. Buchholz, and A. Weiermann. A uniform approach to fundamental sequences and hierarchies. *Mathematical Logic Quarterly*, 40:273–286, 1994.
- [6] David Fernández-Duque and Andreas Weiermann. Ackermannian goodstein sequences of intermediate growth. In Marcella Anselmo, Gianluca Della Vedova, Florin Manea, and Arno Pauly, editors, *Beyond the Horizon of Computability - 16th Conference on Computability in Europe, CiE 2020, Fisciano, Italy, June 29 - July 3, 2020, Proceedings*, volume 12098 of *Lecture Notes in Computer Science*, pages 163–174. Springer, 2020.
- [7] David Fernández-Duque and Andreas Weiermann. A walk with goodstein. *CoRR*, arXiv:2004.09110, 2020.
- [8] R.L. Goodstein. On the restricted ordinal theorem. *Journal of Symbolic Logic*, 9(2):33–41, 1944.
- [9] R.L. Goodstein. Transfinite ordinals in recursive number theory. *Journal of Symbolic Logic*, 12(4):123–129, 12 1947.
- [10] G.H. Hardy. A theorem concerning the infinite cardinal numbers. *Quarterly Journal of Mathematics*, 35:87, 1904.
- [11] L. Kirby and J. Paris. Accessible independence results for Peano arithmetic. *Bulletin of the London Mathematical Society*, 14(4):285–293, 1982.
- [12] W. Pohlers. *Proof Theory, The First Step into Impredicativity*. Springer-Verlag, Berlin Heidelberg, 2009.