

# Pick's Theorem in Two-Dimensional Subspace of $\mathbb{R}^3$

Lin Si

College of Science, Beijing Forestry University, Beijing, 100083, P.R.China

E-mail: silind@163.com

**Abstract.** In this note, we given a version of Pick's theorem for the simple lattice polygon in two-dimensional subspace of  $\mathbb{R}^3$ .

**Keywords.** Pick's theorem; lattice polygon; two-dimensional subspace.

**2010 Mathematics Subject Classification.** 51M25, 52C05.

## 1. Introduction

In the Euclidean plane  $\mathbb{R}^2$ , a lattice point is one whose coordinates are both integers. A lattice polygon is a polygon with all vertices on integer coordinates. The area  $A(P)$  of a simple lattice polygon  $P$  can be given by the celebrated Pick's theorem[5]

$$A(P) = I(P) + \frac{1}{2}B(P) - 1,$$

where  $B(P)$  is the number of lattice points on the boundary of  $P$  and  $I(P)$  is the number of lattice points in the interior of  $P$ .

Pick's formula can be used to compute the area of a lattice polygon conveniently.

For example, on Figure 1,  $I(P) = 60$ ,  $B(P) = 15$ . Then, the area of the polygon is  $A(P) = 60 + 15 - 1 = 74$ .




Figure 1

There are many papers concerning Pick's theorem and it's generalizations[1,2,3,4], which mostly be discussed in two dimension.

Unfortunately, Pick theorem is failed in three dimensions. In 1957, John Reeve found a class of tetrahedra, named as Reeve tetrahedra later, whose vertices are

$$(0, 0, 0)^T, (1, 0, 0)^T, (0, 1, 0)^T, (1, 1, r)^T,$$

where  $r$  is a positive integer.

All Reeve tetrahedra contain the same number of lattice points, but their volumes are different.

In this note, we discussed Pick's theorem in two-dimensional subspace of  $\mathbb{R}^3$ . For any  $(a, b, c)^T \in \mathbb{Z}^3$  with  $(a, b, c) = 1$ , i.e. the greatest common factor of  $a, b, c$  is one, denote by  $K$ ,  $ax + by + cz = 0$ , the two-dimensional subspace of  $\mathbb{R}^3$ . Then we established the following theorem.

**Theorem.** *If  $P$  is simple lattice polygon in the  $K$ , then the area of  $P$  is*

$$k(I(P) + \frac{1}{2}B(P) - 1),$$

where  $B(P)$  is the number of lattice points on the boundary of  $P$  in  $\mathbb{Z}^3$ ,  $I(P)$  is the number of lattice points in the interior of  $P$  in  $\mathbb{Z}^3$  and  $k$  is the constant  $(a^3 + ab^2)\sqrt{a^2 + b^2 + c^2}$ .

**Remark.** *Although the simple lattice polygon  $P$  is in the two-dimensional subspace  $K$ , the lattice points in  $P$  belong to  $\mathbb{Z}^3$ .*

Let  $(a, b, c)^T = (1, 0, 0)^T$  in the Theorem, then we can get Pick's theorem in some coordinate plane of  $\mathbb{R}^3$ .

**Corollary.** *If  $P$  is simple lattice polygon in the  $K$ , whose normal vector is  $(1, 0, 0)^T$ , then the area of  $P$  is*

$$I(P) + \frac{1}{2}B(P) - 1.$$

## 2. Proof of Main Result

For any  $(a, b, c)^T \in \mathbb{Z}^3$  with  $(a, b, c) = 1$ , there is two-dimensional subspace of  $\mathbb{R}^3$

$$ax + by + cz = 0, \tag{1}$$

whose normal vector is just  $(a, b, c)^T$ . We denote this two-dimensional subspace by  $K$ .

By the theory of linear equations system,  $(-b, a, 0)^T$  and  $(-c, 0, a)^T$  are two linearly independent solutions of (1). We denote  $(-b, a, 0)^T$  by  $\alpha$  and  $(-c, 0, a)^T$  by  $\beta$ . Obviously,  $\alpha$  and  $\beta$  are also the basis of  $K$ .

**Lemma 1.** *For any  $(a, b, c)^T \in \mathbb{Z}^3$  with  $(a, b, c) = 1$ , there exists the lattice basis with the minimal area in the two-dimensional subspace  $K$ .*

**Proof.** The area of parallelogram generated by  $\alpha$  and  $\beta$  is

$$\frac{1}{\sqrt{a^2 + b^2 + c^2}} \begin{vmatrix} a & -b & -c \\ b & a & 0 \\ c & 0 & a \end{vmatrix} = \frac{a^3 + ab^2 + ac^2}{\sqrt{a^2 + b^2 + c^2}} = a\sqrt{a^2 + b^2 + c^2}.$$

Denote  $(a, b, c)^T$  by  $n$ . For any lattice basis in  $K$ ,  $k_1\alpha + k_2\beta$  and  $l_1\alpha + l_2\beta$ , where  $k_i, l_i \in \mathbb{Z}(i = 1, 2)$  and  $\begin{vmatrix} k_1 & l_1 \\ k_2 & l_2 \end{vmatrix} = 0$ . The area of parallelogram generated by  $k_1\alpha + k_2\beta$  and  $l_1\alpha + l_2\beta$  is

$$\frac{1}{\sqrt{a^2 + b^2 + c^2}} \left| n \begin{vmatrix} k_1 & l_1 \\ k_2 & l_2 \end{vmatrix} \right| = \frac{1}{\sqrt{a^2 + b^2 + c^2}} \left| (n, \alpha, \beta) \times \begin{pmatrix} 1 & 0 & 0 \\ 0 & k_1 & l_1 \\ 0 & k_2 & l_2 \end{pmatrix} \right|,$$

where  $\begin{vmatrix} k_1 & l_1 \\ k_2 & l_2 \end{vmatrix}$  denote the determinant of  $n, k_1\alpha + k_2\beta$  and  $l_1\alpha + l_2\beta$ .

Thus the lattice basis  $k_1\alpha + k_2\beta$  and  $l_1\alpha + l_2\beta$  have the minimal area if and only if  $\begin{vmatrix} k_1 & l_1 \\ k_2 & l_2 \end{vmatrix} = 1$ .

Let  $k_1 = 1, k_2 = 0, l_1 = 0, l_2 = 1$ ,  $\alpha, \beta$  are the lattice basis with the minimal area in the two-dimensional subspace  $K$ .  $\square$

**Lemma 2.** For any  $(a, b, c)^T \in \mathbb{Z}^3$  with  $(a, b, c) = 1$ , there exists the orthogonal lattice basis in the two-dimensional subspace  $K$ .

**Proof.** By Lemma 1,  $\alpha, \beta$  are the lattice basis with the minimal area in the two-dimensional subspace  $K$ . By Schmidt orthogonalization, let

$$\begin{aligned}\gamma_1 &= \alpha = (-b, a, 0)^T \\ \gamma_2 &= \beta - \frac{(\beta, \gamma_1)}{(\gamma_1, \gamma_1)}\gamma_1 \\ &= (-c, 0, a)^T - \frac{bc}{a^2 + b^2}\gamma_1 \\ &= (-c, 0, a)^T - \left(\frac{-b^2c}{a^2 + b^2}, \frac{abc}{a^2 + b^2}, 0\right)^T \\ &= \left(\frac{-a^2c}{a^2 + b^2}, \frac{-abc}{a^2 + b^2}, \frac{a^3 + ab^2}{a^2 + b^2}\right)^T,\end{aligned}$$

where  $(\beta, \gamma_1)$  denote the usual inner product of  $\beta, \gamma_1$  in  $\mathbb{R}^3$ .

Thus

$$\begin{aligned}\eta_1 &= \gamma_1 = \alpha = (-b, a, 0)^T \\ \eta_2 &= (a^2 + b^2)\gamma_2 = (-a^2c, -abc, a^3 + ab^2)^T\end{aligned}$$

are the orthogonal lattice basis in the two-dimensional subspace  $K$ .  $\square$

**Proof of Theorem** By Lemma 2,  $\eta_1, \eta_2$  are the orthogonal lattice basis in the two-dimensional subspace  $K$ .

The area of parallelogram generated by  $\eta_1, \eta_2$  is

$$\begin{aligned}& \frac{1}{\sqrt{a^2 + b^2 + c^2}} \begin{vmatrix} a & -b & -a^2c \\ b & a & -abc \\ c & 0 & a^3 + ab^2 \end{vmatrix} \\ &= \frac{a^5 + a^3b^2 + ab^2c^2 + a^3c^2 + a^3b^2 + ab^4}{\sqrt{a^2 + b^2 + c^2}} \\ &= \frac{a(a^4 + 2a^2b^2 + b^4 + (a^2 + b^2)c^2)}{\sqrt{a^2 + b^2 + c^2}} \\ &= \frac{a(a^2 + b^2)(a^2 + b^2 + c^2)}{\sqrt{a^2 + b^2 + c^2}} \\ &= (a^3 + ab^2)\sqrt{a^2 + b^2 + c^2},\end{aligned}$$

which just is the constant  $k$  in the Theorem.  $\square$

### Acknowledgements.

This work was supported by the Beijing Higher Education Young Elite Teacher Project (Granted No.YETP0770), the National Natural Science Foundation of China (Grant No.11001014), and the Young Teachers Domestic Visiting Scholars Program of Beijing Forestry University.

## REFERENCES

- [1] Ding R, Kolodziejczyk K, Reay J, A new Pick-type theorem on the hexagonal lattice, *Discrete Math.* **68** (1988)171-177
- [2] Funkenbusch W W, From Euler's formula to Pick's formula using an edge theorem, *The Amer. Math. Monthly* **81** (1974)647-648
- [3] Grunbaum B, Shephard G C, Pick's theorem, *The Amer. Math. Monthly* **100** (1993) 150-161
- [4] Liu A C F, Lattice points and Pick's theorem, *Math. Mag.* **52** (1979)232-235
- [5] Pick G A, Geometrisches zur Zahlentheorie, Sitzungber. Lotos, Naturwissen Zeitschrift **19** (1899) 311-319