

Black box tests for algorithmic stability

Byol Kim and Rina Foygel Barber

May 26, 2022

Abstract

Algorithmic stability is a concept from learning theory that expresses the degree to which changes to the input data (e.g., removal of a single data point) may affect the outputs of a regression algorithm. Knowing an algorithm’s stability properties is often useful for many downstream applications—for example, stability is known to lead to desirable generalization properties and predictive inference guarantees. However, many modern algorithms currently used in practice are too complex for a theoretical analysis of their stability properties, and thus we can only attempt to establish these properties through an empirical exploration of the algorithm’s behavior on various data sets. In this work, we lay out a formal statistical framework for this kind of *black box testing* without any assumptions on the algorithm or the data distribution, and establish fundamental bounds on the ability of any black box test to identify algorithmic stability.

1 Introduction

Consider an algorithm \mathcal{A} that maps data sets to fitted regression functions:

$$\mathcal{A} : \underbrace{(X_1, Y_1), \dots, (X_n, Y_n)}_{\text{Training data points in } \mathbb{R}^d \times \mathbb{R}} \mapsto \underbrace{\hat{\mu} : \mathbb{R}^d \rightarrow \mathbb{R}}_{\text{Fitted function}}.$$

In many practical settings where regression algorithms are used, the distribution P of the data points (X_i, Y_i) is potentially complex and cannot be assumed to satisfy strong simplifying conditions (such as a parametric model or a high degree of smoothness). Guarantees bounding the predictive error of the fitted model $\hat{\mu}$ can still be obtained by testing its predictive accuracy on additional labeled data that was not used for training (i.e., a holdout set). Of course, in settings where the available training data is limited, we may not want to sacrifice sample size in order to obtain this independent holdout set—that is, if n is the number of available labeled data points drawn from the unknown data distribution P , we may want to run the regression algorithm \mathcal{A} on the entire set of n data points. In this type of setting, where no holdout set is available, it is nonetheless possible to give guarantees on predictive error by assuming a condition of *stability* on the algorithm \mathcal{A} [Bousquet and Elisseeff, 2002]. The stability condition essentially requires

that slight perturbations of the training data set $((X_i, Y_i))_{i=1, \dots, n}$, such as removing or replacing a single data point, can alter the fitted model $\hat{\mu}$ only slightly.

Unfortunately, verifying the stability of a given regression algorithm is itself challenging. Some algorithms are known to satisfy stability by construction—for example, if \mathcal{A} is a k -nearest neighbor algorithm, then removing a single data point in the training set will only perturb predictions locally near that point. However, in many settings, particularly with modern machine learning algorithms, it is not feasible to analyze an algorithm’s stability properties theoretically, since the algorithm is too complex for this to be tractable. Instead, we can only examine its properties through empirical testing, to try to determine based on its output whether it appears to be stable or unstable. We may ask, therefore, whether stability of an algorithm can instead be validated empirically, to ensure that the algorithms used in practice also enjoy the theoretical guarantees that accompany the stability property.

In this work, our aim is to examine the question of whether it is possible to infer the stability of an algorithm through *black box testing* that does not peer inside the construction of the algorithm but instead tries to determine the stability property by observing the empirical behavior of the algorithm on various data sets. In particular, we will establish fundamental bounds on our ability to test for the stability property in the absence of assumptions on the distribution P or the algorithm \mathcal{A} .

2 Background and framework

2.1 Algorithmic stability

We define an *algorithm* \mathcal{A} as any function taking any finite collection of (x, y) -pairs to a regression function (i.e., a function $\hat{\mu}$ that maps each x to some y). Formally, as a map between spaces, \mathcal{A} can be written as

$$\mathcal{A} : \left(\bigcup_{n \geq 0} (\mathbb{R}^d \times \mathbb{R})^n \right) \times [0, 1] \rightarrow \mathcal{M}_d,$$

where $\mathcal{M}_d = \{\text{Measurable functions } \mu : \mathbb{R}^d \rightarrow \mathbb{R}\}$. Given n points $(x_1, y_1), \dots, (x_n, y_n)$, the fitted regression function $\hat{\mu}$ obtained from \mathcal{A} is

$$\hat{\mu} = \mathcal{A}[(x_1, y_1), \dots, (x_n, y_n); \xi], \quad \xi \sim \text{Uniform}[0, 1].$$

Here, the role of the last argument ξ is to encode the random behavior of \mathcal{A} , if needed—for example, if \mathcal{A} uses stochastic gradient descent, ξ determines the random sequence of data points seen by the algorithm. Effectively, we can think of ξ as the random seed with which we initialize the procedure.

Throughout this paper, we will assume that \mathcal{A} is *symmetric*, i.e., given any training data set $(x_1, y_1), \dots, (x_n, y_n)$ and its permuted version $(x_{\sigma(1)}, y_{\sigma(1)}), \dots, (x_{\sigma(n)}, y_{\sigma(n)})$,

$$\mathcal{A}[(x_1, y_1), \dots, (x_n, y_n); \xi] \stackrel{d}{=} \mathcal{A}[(x_{\sigma(1)}, y_{\sigma(1)}), \dots, (x_{\sigma(n)}, y_{\sigma(n)}); \xi],$$

where $\stackrel{d}{=}$ denotes equality in distribution with respect to the random ξ . For example, this property holds for stochastic gradient descent, since ξ is used to draw a uniformly random ordering of the n data points.

In a typical statistical learning problem, an algorithm \mathcal{A} is applied to training data that is assumed to arise from some unknown random process. In this work, we focus on the independent and identically distributed (i.i.d.) setting, i.e.,

$$(X_i, Y_i) \stackrel{\text{iid}}{\sim} P, \quad i = 1, 2, \dots$$

We are interested in the stability of the predictions made by a model obtained by applying \mathcal{A} to a training data set from P . That is to say, we would like to establish that slight perturbations to the training data set will not substantially affect the resulting fitted function $\hat{\mu}$.

Definition 1. *Let \mathcal{A} be a symmetric algorithm. Let $\epsilon \geq 0$ and $\delta \in [0, 1)$. We say that \mathcal{A} is (ϵ, δ) -stable with respect to training data sets of size n from data distribution P —or, equivalently, the triple (\mathcal{A}, P, n) is (ϵ, δ) -stable—if*

$$\mathbb{P} \{ |\hat{\mu}_n(X_{n+1}) - \hat{\mu}_{n-1}(X_{n+1})| > \epsilon \} \leq \delta, \quad (1)$$

where $\hat{\mu}_n$ and $\hat{\mu}_{n-1}$ are the fitted models obtained from independent runs of the algorithm \mathcal{A} using the full training data set and the data set after removing the last data point, i.e.,

$$\hat{\mu}_n = \mathcal{A}[(X_1, Y_1), \dots, (X_n, Y_n); \xi], \quad \hat{\mu}_{n-1} = \mathcal{A}[(X_1, Y_1), \dots, (X_{n-1}, Y_{n-1}); \xi'],$$

where $\xi, \xi' \stackrel{\text{iid}}{\sim} \text{Uniform}[0, 1]$.

In this definition, the probability calculated in (1) is taken with respect to both the data $(X_1, Y_1), \dots, (X_n, Y_n) \stackrel{\text{iid}}{\sim} P$ and the algorithmic randomness $\xi, \xi' \stackrel{\text{iid}}{\sim} \text{Uniform}[0, 1]$.

2.1.1 Other notions of algorithmic stability

The earliest works on algorithmic stability date back to at least the late 1970s [Rogers and Wagner, 1978, Devroye and Wagner, 1979a,b]. Since then, various notions of algorithmic stability has been proposed, all describing different ways an algorithm may exhibit “continuity” as a map from the data space $\bigcup_{n \geq 0} (\mathbb{R}^d \times \mathbb{R})^n$ to the model space \mathcal{M}_d . Among many such competing definitions, some of the most influential in the fields of learning theory and statistics are due to Bousquet and Elisseeff [2002], where certain forms of algorithmic stability are shown to imply good generalization behavior (see also Kearns and Ron [1999]).

The formulation of stability that we use in this paper (Definition 1) is perhaps most similar to the *hypothesis stability* condition of Bousquet and Elisseeff [2002], which instead requires a bound on $\mathbb{E} [|\ell(\hat{\mu}_n; (X_{n+1}, Y_{n+1})) - \ell(\hat{\mu}_{n-1}; (X_{n+1}, Y_{n+1}))|]$ for a loss function ℓ . Definition 1 has appeared in Barber et al. [2021] under the name of *out-of-sample stability*, where it was shown that this condition is sufficient for predictive coverage guarantees of

the jackknife (i.e., leave-one-out cross-validation) and jackknife+ methods. Earlier work by Steinberger and Leeb [2020] also establishes coverage properties for the jackknife under a similar stability property.

Many recent works have also proposed extensions and modifications of existing notions of stability Kutin and Niyogi [2002a,b], Mukherjee et al. [2006], Shalev-Shwartz et al. [2010] proposed further relaxations and variants of Bousquet and Elisseeff [2002]’s definitions in the context of characterizing sufficient and necessary conditions for learning problems. Interestingly, Shalev-Shwartz et al. [2010, Section 6.2] give a generic but infeasible meta-algorithm in which stability plays the role of regularization, which they use to establish that a statistical problem is asymptotically *learnable* only if it can be learned with a stable algorithm. More recently, Deng et al. [2021] advocated a functional description of algorithmic stability. It must be noted that only some of the definitions considered in these works define algorithmic stability as a property that holds jointly for an algorithm \mathcal{A} and a data distribution P . For example, in Bousquet and Elisseeff [2002]’s work, while *hypothesis stability* is defined for each algorithm-distribution pair, *uniform stability* is instead required to hold uniformly over all possible data sets and hence, is unsuitable as a target for statistical inference.

2.2 Testing for stability

The focus of this work is the following statistical inference question: given a fixed $\epsilon \geq 0$ and $\delta \in [0, 1)$ and a desired confidence level $1 - \alpha$, we would like to test whether (\mathcal{A}, P, n) is (ϵ, δ) -stable. Writing \mathcal{D}_ℓ and \mathcal{D}_u to denote the available labeled and unlabeled data, respectively, we will write $\widehat{T}_{\epsilon, \delta} = \widehat{T}_{\epsilon, \delta}(\mathcal{A}, \mathcal{D}_\ell, \mathcal{D}_u) \in \{0, 1\}$ for the data-dependent outcome of such a hypothesis test, where $\widehat{T}_{\epsilon, \delta} = 1$ indicates that we believe stability holds. Define the “ground truth” as

$$T_{\epsilon, \delta}^* = T_{\epsilon, \delta}^*(\mathcal{A}, P, n) = \mathbb{1} \{(\mathcal{A}, P, n) \text{ is } (\epsilon, \delta)\text{-stable}\}.$$

Given this target, we seek a test $\widehat{T}_{\epsilon, \delta}$ that satisfies the following notion of *assumption-free* validity:

$$\text{For all } (\mathcal{A}, P, n), \text{ if } T_{\epsilon, \delta}^*(\mathcal{A}, P, n) = 0 \text{ then } \mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta} = 1 \right\} \leq \alpha. \quad (2)$$

In other words, our chance of falsely labeling a (\mathcal{A}, P, n) triple as (ϵ, δ) -stable using this test is bounded by α , uniformly over all (\mathcal{A}, P, n) .

2.3 Black box tests of stability

For some simple algorithms such as nearest neighbor methods or ridge regression, their stability properties have been analyzed and are hence known [Devroye and Wagner, 1979a,b, Bousquet and Elisseeff, 2002, Steinberger and Leeb, 2020]. For many modern learning algorithms and data sets, however, stability cannot be determined the same way, as they are too complex for this to be a tractable problem. Such algorithms are rightly regarded as “black boxes” in the sense that our only practical means of studying them is by observing

their empirical behavior, i.e., by sending different data sets through an algorithm and taking note of the changes in the output. Often these outputted fitted functions $\hat{\mu}$ will also be “black boxes” and be only accessible by evaluating them at various test points.

This “black box” setting is the framework we are interested in here. In this setting, for a test $\hat{T}_{\epsilon,\delta}$ to be admissible, the computations the test makes can only rely on observable outcomes. The situation closely resembles that of black box testing in software development where the performance of a code is tested by pushing different inputs through the code and observing the results assuming no knowledge of the inner structure [Myers et al., 2011]. We essentially perform the same task for algorithmic stability, but with the crucial difference that we also seek assumption-free validity.

To formalize this setting, we define a *black box test* of algorithmic stability. Let $\mathfrak{D}_\ell = \cup_{k \geq 0} (\mathbb{R}^d \times \mathbb{R})^k$ denote the space of labeled data sets of any size, and let $\mathfrak{D}_u = \cup_{k \geq 0} (\mathbb{R}^d)^k$ denote the space of unlabeled data sets of any size.

Definition 2 (Black box test). *Consider any test \hat{T} that takes as input an algorithm $\mathcal{A} : \mathfrak{D}_\ell \rightarrow \mathcal{M}_d$, a labeled data set $\mathcal{D}_\ell \in \mathfrak{D}_\ell$, and an unlabeled data set $\mathcal{D}_u \in \mathfrak{D}_u$, and returns a (possibly randomized) binary output $\hat{T}(\mathcal{A}, \mathcal{D}_\ell, \mathcal{D}_u) \in \{0, 1\}$. Then we say that \hat{T} is a black box test if for some measurable functions $f^{(1)}, f^{(2)}, \dots$ and T , it can be defined in the following way:*

1. At the initial stage $r = 1$,

(a) Generate a labeled data set $\mathcal{D}_\ell^{(1)} \in \mathfrak{D}_\ell$ and an unlabeled data set $\mathcal{D}_u^{(1)} \in \mathfrak{D}_u$ as a function of the input data sets:

$$\left(\mathcal{D}_\ell^{(1)}, \mathcal{D}_u^{(1)} \right) = f^{(1)} \left[\mathcal{D}_\ell, \mathcal{D}_u, \zeta^{(1)} \right],$$

where $\zeta^{(1)} \sim \text{Uniform}[0, 1]$.

(b) Fit a model and evaluate it using the generated data $(\mathcal{D}_\ell^{(1)}, \mathcal{D}_u^{(1)})$:

$$\hat{\mu}^{(1)} = \mathcal{A} \left[\mathcal{D}_\ell^{(1)}; \xi^{(1)} \right], \quad \hat{\mathcal{Y}}^{(1)} = \hat{\mu}^{(1)} \left[\mathcal{D}_u^{(1)} \right],$$

where $\xi^{(1)} \sim \text{Uniform}[0, 1]$ and the operation of $\hat{\mu}^{(1)}$ on $\mathcal{D}_u^{(1)}$ is understood in the pointwise sense.¹

2. At each stage $r = 2, 3, \dots$,

(a) Generate a labeled data set $\mathcal{D}_\ell^{(r)} \in \mathfrak{D}_\ell$ and an unlabeled data set $\mathcal{D}_u^{(r)} \in \mathfrak{D}_u$ as a function of all the data sets and predictions observed so far:

$$\begin{aligned} & \left(\mathcal{D}_\ell^{(r)}, \mathcal{D}_u^{(r)} \right) \\ &= f^{(r)} \left[\mathcal{D}_\ell, \mathcal{D}_u, \left(\mathcal{D}_\ell^{(s)} \right)_{s=1}^{r-1}, \left(\mathcal{D}_u^{(s)} \right)_{s=1}^{r-1}, \left(\hat{\mathcal{Y}}^{(s)} \right)_{s=1}^{r-1}, \left(\zeta^{(s)} \right)_{s=1}^{r-1}, \left(\xi^{(s)} \right)_{s=1}^{r-1}, \zeta^{(r)} \right], \end{aligned}$$

where $\zeta^{(r)} \sim \text{Uniform}[0, 1]$.

¹For example, if $\mathcal{D}_u^{(1)}$ contains two unlabeled points X_1 and X_2 , then $\hat{\mu}^{(1)} \left[\mathcal{D}_u^{(1)} \right] = (\hat{\mu}^{(1)}(X_1), \hat{\mu}^{(1)}(X_2))$.

(b) Fit a model and evaluate it using the generated data $(\mathcal{D}_\ell^{(r)}, \mathcal{D}_u^{(r)})$:

$$\widehat{\mu}^{(r)} = \mathcal{A} \left[\mathcal{D}_\ell^{(r)}; \xi^{(r)} \right], \quad \widehat{\mathcal{Y}}^{(r)} = \widehat{\mu}^{(r)} \left[\mathcal{D}_u^{(r)} \right],$$

where $\xi^{(r)} \sim \text{Uniform}[0, 1]$ and the operation of $\widehat{\mu}^{(r)}$ on $\mathcal{D}_u^{(r)}$ is understood in the pointwise sense.

3. Compute the output $\widehat{T} \in \{0, 1\}$ as a function of the sequence of generated data sets and observed predictions:

$$\widehat{T} = T \left[\mathcal{D}_\ell, \mathcal{D}_u, \left(\mathcal{D}_\ell^{(r)} \right)_{r \geq 1}, \left(\mathcal{D}_u^{(r)} \right)_{r \geq 1}, \left(\widehat{\mathcal{Y}}^{(r)} \right)_{r \geq 1}, \left(\zeta^{(r)} \right)_{r \geq 1}, \left(\xi^{(r)} \right)_{r \geq 1}, \zeta \right],$$

where $\zeta \sim \text{Uniform}[0, 1]$.

In the above, the role of the $\text{Uniform}[0, 1]$ random variables $\zeta^{(1)}, \xi^{(1)}, \zeta^{(2)}, \xi^{(2)}, \dots, \zeta$ is to allow for randomization at each step of the procedure, if desired.

2.3.1 Examples of black box tests

To make the notion of black box test concrete, we give several examples of procedures we might use in practice to estimate an algorithm's stability. Throughout, we will write $\mathcal{D}_\ell = ((X_1, Y_1), \dots, (X_{N_\ell}, Y_{N_\ell}))$ to denote the available labeled data and $\mathcal{D}_u = (X_{N_\ell+1}, \dots, X_{N_\ell+N_u})$ for the available unlabeled data.

First, we can consider splitting the available data and measuring stability on each split. We shall revisit this specific test in Section 3 when we study some fundamental bounds.

Example 1 (A simple Binomial test). *Define*

$$\kappa = \kappa(n, N_\ell, N_u) = \min \left\{ \frac{N_\ell}{n}, \frac{N_\ell + N_u}{n + 1} \right\}. \quad (3)$$

Then $\lfloor \kappa \rfloor$ is the number of copies of independent data sets—each consisting of n labeled training points and one unlabeled test point—that can be constructed from \mathcal{D}_ℓ and \mathcal{D}_u .

1. For $k = 1, \dots, \lfloor \kappa \rfloor$, construct the k -th data set with n labeled data points

$$(X_{(k-1)n+1}, Y_{(k-1)n+1}), \dots, (X_{kn}, Y_{kn})$$

and one unlabeled data point $X_{\lfloor \kappa \rfloor n + k}$.

2. For each $k = 1, \dots, \lfloor \kappa \rfloor$, train models

$$\begin{aligned} \widehat{\mu}_n^{(k)} &= \mathcal{A} \left[(X_{(k-1)n+1}, Y_{(k-1)n+1}), \dots, (X_{kn}, Y_{kn}); \xi^{(2k-1)} \right], \\ \widehat{\mu}_{n-1}^{(k)} &= \mathcal{A} \left[X_{(k-1)n+1}, Y_{(k-1)n+1}, \dots, (X_{kn-1}, Y_{kn-1}); \xi^{(2k)} \right] \end{aligned}$$

with $\xi^{(2k-1)}, \xi^{(2k)} \stackrel{\text{iid}}{\sim} \text{Uniform}[0, 1]$, and compute the difference in predictions

$$\Delta^{(k)} = \left| \widehat{\mu}_n^{(k)} (X_{\lfloor \kappa \rfloor n + k}) - \widehat{\mu}_{n-1}^{(k)} (X_{\lfloor \kappa \rfloor n + k}) \right|. \quad (4)$$

3. Compute how many of the $\Delta^{(k)}$'s exceed ϵ ,

$$B = \sum_{k=1}^{\lfloor \kappa \rfloor} \mathbb{1} \{ \Delta^{(k)} > \epsilon \},$$

and compare this test statistic against the $\text{Binomial}(\lfloor \kappa \rfloor, \delta)$ distribution, returning $\widehat{T}_{\epsilon, \delta} = 1$ if B is sufficiently small or $\widehat{T}_{\epsilon, \delta} = 0$ otherwise (precise details for the test are given in Section 3.1).

Alternatively, instead of partitioning the available data, we can use bootstrapped samples to run the algorithm and generate the empirical differences $\Delta^{(k)}$:

Example 2 (Bootstrapped sample method).

1. For $r = 1, \dots, R$, construct a bootstrapped training sample $(X_1^{(r)}, Y_1^{(r)}), \dots, (X_n^{(r)}, Y_n^{(r)})$ by sampling with replacement from \mathcal{D}_ℓ . Sample a test point $X_{n+1}^{(r)}$ from \mathcal{D}_u .
2. As for Example 1, train models $\widehat{\mu}_n^{(r)}$ and $\widehat{\mu}_{n-1}^{(r)}$, and compute the difference in predictions

$$\Delta^{(r)} = \left| \widehat{\mu}_n^{(r)} \left(X_{n+1}^{(r)} \right) - \widehat{\mu}_{n-1}^{(r)} \left(X_{n+1}^{(r)} \right) \right|.$$

3. As in Example 1, compute how many of the $\Delta^{(r)}$'s exceed ϵ ,

$$\sum_{r=1}^R \mathbb{1} \{ \Delta^{(r)} > \epsilon \},$$

and return $\widehat{T}_{\epsilon, \delta} = 1$ if this value is sufficiently small or $\widehat{T}_{\epsilon, \delta} = 0$ otherwise.

Or, we may want to use the available data to estimate P first, and then test the algorithm with simulated data from the estimated distribution \widehat{P} .

Example 3 (Simulated sample method).

1. Use \mathcal{D}_ℓ and \mathcal{D}_u to obtain an estimate \widehat{P} of P via any method, then for $r = 1, \dots, R$, draw a simulated training set $(X_1^{(r)}, Y_1^{(r)}), \dots, (X_n^{(r)}, Y_n^{(r)}) \stackrel{\text{iid}}{\sim} \widehat{P}$ and a simulated test point $X_{n+1}^{(r)} \sim \widehat{P}_X$, the marginal distribution of X under \widehat{P} .
2. Proceed as in Examples 1 and 2.

Furthermore, the black box testing framework also allows us to use a testing procedure that learns from the earlier stages to improve the design of future queries. For instance, after running any of the above methods for some number of stages, a procedure may try to narrow down the properties of the training set and/or the test point that appear to lead to a lack of stability based on the observed outcomes up to that stage. In the subsequent stages, the data set generation may be made to target these types of training and/or test data to learn more about the algorithm's stability properties before outputting the final answer $\widehat{T}_{\epsilon, \delta} \in \{0, 1\}$.

2.3.2 Inadmissible tests

Tests that require uncountably infinitely many executions of a black box are *not* admissible under Definition 2. An example is a test \widehat{T} that relies on evaluating

$$\sup_{y \in \mathbb{R}} \{ \mathcal{A} [(X_1, Y_1), \dots, (X_{n-1}, Y_{n-1}), (X_n, y); \xi] (X_{n+1}) \}, \quad (5)$$

since computing this supremum requires uncountably infinitely many calls to \mathcal{A} . Similarly,

$$\sup_{x \in \mathbb{R}^d} \{ \mathcal{A} [(X_1, Y_1), \dots, (X_n, Y_n); \xi] (x) \} \quad (6)$$

is also not allowed in general, as it requires uncountably infinitely many evaluations of the fitted regression function $\widehat{\mu}_n = \mathcal{A}[(X_1, Y_1), \dots, (X_n, Y_n); \xi]$.

2.3.3 Black box algorithms and black box functions

Definition 2 above treats both the algorithm \mathcal{A} itself as a black box, but also the resulting fitted functions $\widehat{\mu}$. That is, this definition allows for a setting where even the fitted functions themselves may be so complex that they can be studied only empirically—if we run the model fitting algorithm \mathcal{A} to produce functions $\widehat{\mu}_n$ and $\widehat{\mu}_{n-1}$, we might only be able to compare these functions by comparing $\widehat{\mu}_n(x)$ against $\widehat{\mu}_{n-1}(x)$ on various test points x , but might not be able to provide a meaningful bound on the maximum difference $\sup_x |\widehat{\mu}_n(x) - \widehat{\mu}_{n-1}(x)|$.

In some settings, however, we might assume that the black-box algorithm \mathcal{A} returns functions $\widehat{\mu}$ that can be studied analytically. As an example, \mathcal{A} might perform feature selection via some complex and intractable procedure, but might always output a linear model. In this scenario the resulting fitted model $\widehat{\mu}$ can therefore be studied theoretically as well as empirically. In particular, this means that calculations of the form (5) would continue to not be allowed for any black box test, but calculations of the form (6) would be permitted.

We will discuss this alternative definition of the black box setting later on in Section 5.2, and will see how this alternative formulation affects the results that we obtain.

3 Bounds for black box tests

In this section, we will first study the simple Binomial test defined in Example 1, and will analyze its performance as a black-box test that achieves valid inference regardless of the algorithm \mathcal{A} and distribution P . We will then prove that the performance of this simple procedure is in fact essentially optimal among all black box tests.

3.1 A simple Binomial test

Recall the simple Binomial test described in Example 1: after constructing $\lfloor \kappa \rfloor$ many independent data sets, each consisting of n labeled points and one unlabeled point, we

compute the difference in predictions $\Delta^{(k)}$ as in (4) induced by comparing $\widehat{\mu}_n$ and $\widehat{\mu}_{n-1}$ for each independent data set $k = 1, \dots, \lfloor \kappa \rfloor$, and then construct the test statistic

$$B = \sum_{k=1}^{\lfloor \kappa \rfloor} \mathbb{1} \{ \Delta^{(k)} > \epsilon \},$$

so that the empirical proportion $B/\lfloor \kappa \rfloor$ is an estimate of $\mathbb{P}\{|\widehat{\mu}_n(X_{n+1}) - \widehat{\mu}_{n-1}(X_{n+1})| > \epsilon\}$. Comparing to the bound (1) in the definition of stability (Definition 1), we see that we should return $\widehat{T}_{\epsilon, \delta} = 1$ if the empirical proportion $B/\lfloor \kappa \rfloor$ is sufficiently below δ .

To make this precise, we first observe that since the $\Delta^{(k)}$'s are constructed on independent subsets of the data, we have $B \sim \text{Binomial}(\lfloor \kappa \rfloor, \delta_\epsilon^*)$, where

$$\delta_\epsilon^* = \mathbb{P} \{ |\widehat{\mu}_n(X_{n+1}) - \widehat{\mu}_{n-1}(X_{n+1})| > \epsilon \} \quad (7)$$

is the unknown true probability (that is, δ_ϵ^* is the smallest possible value of δ' so that (\mathcal{A}, P, n) is (ϵ, δ') -stable). Therefore, we simply need to perform a one-tailed Binomial test on B to test whether $\delta_\epsilon^* \leq \delta$, in which case $T_{\epsilon, \delta}^* = 1$, or $\delta_\epsilon^* > \delta$, in which case $T_{\epsilon, \delta}^* = 0$. To perform this test, define $k_{\kappa, \delta}^* \in \{0, \dots, \lfloor \kappa \rfloor - 1\}$ and $a_{\kappa, \delta}^* \in (0, 1]$ as the unique values satisfying²

$$\mathbb{P} \{ \text{Binomial}(\lfloor \kappa \rfloor, \delta) < k \} + a_{\kappa, \delta}^* \cdot \mathbb{P} \{ \text{Binomial}(\lfloor \kappa \rfloor, \delta) = k \} = \alpha,$$

and then define

$$\widehat{T}_{\epsilon, \delta} = \begin{cases} 1, & \text{if } B < k_{\kappa, \delta}^*, \\ \mathbb{1} \{ \zeta \leq a_{\kappa, \delta}^* \}, \text{ where } \zeta \sim \text{Uniform}[0, 1], & \text{if } B = k_{\kappa, \delta}^*, \\ 0, & \text{if } B > k_{\kappa, \delta}^*. \end{cases} \quad (8)$$

Clearly, this test is a black box test in the sense of Definition 2. We now verify the assumption-free validity property (2) to ensure bounded error for detecting that an algorithm is *not* stable, and calculate this test's power to detect when an algorithm *is* stable.

Theorem 1. *Fix any parameters $\epsilon \geq 0$ and $\delta \in [0, 1)$, and any Type I error level $\alpha \in (0, 1)$. Fix any integers $n \geq 2$ and $N_\ell, N_u \geq 0$, and define $\lfloor \kappa \rfloor$ as in (3). Then the black box test $\widehat{T}_{\epsilon, \delta}$ defined in (8) above satisfies the assumption-free validity (2) at level $1 - \alpha$, i.e., for any (\mathcal{A}, P, n) that is not (ϵ, δ) -stable (i.e., $T_{\epsilon, \delta}^* = 0$), it holds that*

$$\mathbb{P} \{ \widehat{T}_{\epsilon, \delta} = 1 \} \leq \alpha.$$

Moreover, for any (\mathcal{A}, P, n) that is (ϵ, δ) -stable (i.e., $T_{\epsilon, \delta}^* = 1$), if either $\delta_\epsilon^* = 0$ or $\delta \leq 1 - \alpha^{1/\lfloor \kappa \rfloor}$ then the power of the test is given by

$$\mathbb{P} \{ \widehat{T}_{\epsilon, \delta} = 1 \} = \left\{ \alpha \cdot \left(\frac{1 - \delta_\epsilon^*}{1 - \delta} \right)^{\lfloor \kappa \rfloor} \right\} \wedge 1, \quad (9)$$

where δ_ϵ^* is defined as in (7).

²Abusing notation, the $\text{Binomial}(0, \delta)$ distribution should be interpreted as a point mass at zero. In this way, the defined procedure is able to include the case that $\lfloor \kappa \rfloor = 0$ (i.e., $N_\ell < n$ or $N_\ell + N_u < n + 1$).

These results are straightforward consequences of the fact that $B \sim \text{Binomial}(\lfloor \kappa \rfloor, \delta_\epsilon^*)$ for any (\mathcal{A}, P, n) , i.e., the distribution of B depends on (\mathcal{A}, P, n) only through the parameter δ_ϵ^* . We give the full proof in Appendix A.

To understand the setting in which the power calculation applies, the two special cases are exactly the settings where the outcome $\widehat{T}_{\epsilon, \delta} = 1$ can occur only when $B = 0$, i.e., when we observe $\Delta^{(k)} \leq \epsilon$ for all $k = 1, \dots, \lfloor \kappa \rfloor$ data sets. In particular, the condition $\delta \leq 1 - \alpha^{1/\lfloor \kappa \rfloor}$ is not very restrictive in practical scenarios. For example, taking $\delta = 0.1$ (i.e., a 10% chance of instability) and $\alpha = 0.1$ (i.e., a 10% Type I error rate), this condition is satisfied for any $\lfloor \kappa \rfloor < 21$ (i.e., the amount of data available for testing is less than 21 times than the sample size n we are interested in studying).

3.2 A bound on power for all black box tests

The simple black box test defined in (8) above appears to be a very naive and inefficient proposal—in particular, we only use a single split of the available data into $\lfloor \kappa \rfloor$ data sets of size $n + 1$ (i.e., n labeled training points and one unlabeled test point), and the algorithm \mathcal{A} is only called $2\lfloor \kappa \rfloor$ many times. We might expect to increase the power of the test by re-splitting or by resampling as in Example 2, or perhaps by creating new data through simulation or perturbation as in Example 3, so that it may be possible to observe the algorithm’s stability behavior on a wider range of different data sets.

However, our next result establishes that, despite its simplicity, the simple Binomial test proposed in (8) is optimal in some regimes, in the sense that no universally valid black box test can improve its power.

Theorem 2. *Fix any parameters $\epsilon \geq 0$ and $\delta \in [0, 1)$, any Type I error level $\alpha \in (0, 1)$, and any integers $n \geq 2$ and $N_\ell, N_u \geq 0$. Let $\widehat{T}_{\epsilon, \delta}$ be any black box test as in Definition 2, which satisfies the assumption-free validity (2) at level $1 - \alpha$. Then, for any (\mathcal{A}, P, n) that is (ϵ, δ) -stable (i.e., $T_{\epsilon, \delta}^* = 1$), the power of $\widehat{T}_{\epsilon, \delta}$ is bounded as*

$$\mathbb{P}\left\{\widehat{T}_{\epsilon, \delta} = 1\right\} \leq \left\{\alpha \cdot \left(\frac{1 - \delta_\epsilon^*}{1 - \delta}\right)^\kappa\right\} \wedge 1, \quad (10)$$

where $\kappa = \kappa(n, N_\ell, N_u)$ is defined as in (3).

We prove this theorem in Appendix B. The key idea is that for any triple (\mathcal{A}, P, n) that is (ϵ, δ) -stable, we can perturb the algorithm \mathcal{A} and the data distribution P very slightly so that the perturbed triple, call it (\mathcal{A}', P', n) , is *no longer* (ϵ, δ) -stable, but with only N_ℓ labeled data points and N_u unlabeled data points available, it is difficult to tell apart the outcomes of running the original algorithm \mathcal{A} on data from P from the outcomes of running the perturbed algorithm \mathcal{A}' on data from P' . Since validity of the test ensures that $\mathbb{P}\{\widehat{T}_{\epsilon, \delta} = 1\}$ must be $\leq \alpha$ for the triple (\mathcal{A}', P', n) , it therefore follows that this probability cannot be much larger than α for the original triple (\mathcal{A}, P, n) .

3.2.1 Matching upper and lower bounds

To understand the implications of Theorem 2, let us compare the upper bound on power (10), which holds universally for any black box test of stability with assumption-free validity, to the power achieved by our very simple test (9). In particular, in the case where κ is an integer, we see that the simple Binomial test defined in (8) is in fact optimal over all black box tests in the regime $\delta_\epsilon^* = 0$ or $\delta \leq 1 - \alpha^{1/\lfloor \kappa \rfloor}$. That is to say, there is no possibility of power gain by running a more complicated procedure in this scenario. For example, in this regime, if $N_\ell = n$ (i.e., the number of available labeled data points matches the sample size at which we want to test stability), then $\lfloor \kappa \rfloor = 1$ and we can therefore test the stability of \mathcal{A} with maximum power using only two runs of \mathcal{A} (on the n points to fit $\hat{\mu}_n$, and on the first $n - 1$ points to fit $\hat{\mu}_{n-1}$). This is a surprising result—we would intuitively expect that additional runs of \mathcal{A} , on more carefully curated data sets obtained via some more sophisticated resampling procedure, might yield useful information about stability beyond simply calling \mathcal{A} on the single available data set. However, by comparing the power achieved in Theorem 1 and the universal upper bound shown in Theorem 2, we see that any additional calls to \mathcal{A} are not adding any information to the test.

3.2.2 Power and sample size

Is Theorem 2 a pessimistic result? The answer depends on *how* one wishes to use a stability test. Broadly speaking, we can imagine two different types of goals.

First, we might simply be interested in characterizing the algorithm, by studying its behavior and its properties on different data distributions. In this setting, it is plausible that we might be interested in learning about the algorithm’s stability at a sample size n that is far smaller than the available data set size $N_\ell + N_u$. Here, the power of a black box test can be quite high as long as the sample size ratio κ is large.

Alternatively, it may be the case that we are not interested in learning about the algorithm’s stability for its own sake, but rather that we need to verify that stability holds in order to check assumptions for other procedures. For example, for the task of predicting plausible values of Y given a new value of X , methods based on cross-validation can be used to construct predictive intervals around $\hat{\mu}_n(X)$, but their coverage guarantees typically require algorithmic stability [Steinberger and Leeb, 2016, 2020, Barber et al., 2021]. In this case, we would probably want to test the algorithm’s stability at the same sample size as the number of available labeled data points (i.e., $n = N_\ell$), since we would expect that using the largest possible sample size for training would yield the most accurate fitted model. This is the scenario for which the implications of Theorem 2 are quite discouraging. When $\lfloor \kappa \rfloor = 1$, the maximum possible power is upper-bounded by $\alpha/(1 - \delta)$, which is not much higher than the type-I error level α if δ is small. In other words, it is impossible to be substantially better than random if we need to check the stability assumption at a sample size $n \approx N_\ell$.

4 Alternative targets for inference

So far, we have focused on finding an answer to the binary question “Does (\mathcal{A}, P, n) satisfy (ϵ, δ) -stability?”, with a fixed ϵ and δ . However, this is only one of many possible ways to formulate this inference problem, and here we list two additional questions that are also of interest.

Given a fixed $\epsilon \geq 0$ and a desired confidence level $1 - \alpha$, is it possible to compute a data-dependent estimate $\widehat{\delta}_\epsilon$ such that (\mathcal{A}, P, n) is $(\epsilon, \widehat{\delta}_\epsilon)$ -stable with probability at least $1 - \alpha$?

Given a fixed $\delta \in [0, 1)$ and a desired confidence level $1 - \alpha$, is it possible to compute a data-dependent estimate $\widehat{\epsilon}_\delta$ such that (\mathcal{A}, P, n) is $(\widehat{\epsilon}_\delta, \delta)$ -stable with probability at least $1 - \alpha$?

Similarly to how we defined the “ground truth” $T_{\epsilon, \delta}^*$ for the data-dependent outcome $\widehat{T}_{\epsilon, \delta}$, we define the associated targets as

$$\delta_\epsilon^* = \inf \{ \delta \geq 0 : T_{\epsilon, \delta}^* = 1 \}, \quad \epsilon_\delta^* = \inf \{ \epsilon \geq 0 : T_{\epsilon, \delta}^* = 1 \}.$$

Examining the definition of $T_{\epsilon, \delta}^*$, we can observe that $\delta_\epsilon^* = \mathbb{P}\{|\widehat{\mu}_n(X_{n+1}) - \widehat{\mu}_{n-1}(X_{n+1})| > \epsilon\}$ (as considered earlier in (7)), while ϵ_δ^* is the $(1 - \delta)$ -quantile of $|\widehat{\mu}_n(X_{n+1}) - \widehat{\mu}_{n-1}(X_{n+1})|$. By definition of (ϵ, δ) -stability, we see that for any (ϵ, δ) , it holds that

$$T_{\epsilon, \delta}^* = 1 \iff \delta \geq \delta_\epsilon^* \iff \epsilon \geq \epsilon_\delta^*. \quad (11)$$

Next, we define notions of assumption-free validity for inference on each of the new targets. For the problem of estimating δ_ϵ^* given a fixed $\epsilon \geq 0$, we seek an estimator $\widehat{\delta}_\epsilon$ such that

$$\mathbb{P} \left\{ \widehat{\delta}_\epsilon \geq \delta_\epsilon^* \right\} \geq 1 - \alpha \text{ for all } (\mathcal{A}, P, n). \quad (12)$$

(Equivalently, $\mathbb{P}\{T_{\epsilon, \widehat{\delta}_\epsilon}^* = 1\} \geq 1 - \alpha$.) In other words, given an algorithm \mathcal{A} and data drawn from P , we conclude with $1 - \alpha$ level confidence that (\mathcal{A}, P, n) is $(\epsilon, \widehat{\delta}_\epsilon)$ -stable. Similarly, for the problem of estimating ϵ_δ^* given a fixed $\delta \in [0, 1)$, we seek an estimator $\widehat{\epsilon}_\delta$ such that

$$\mathbb{P} \left\{ \widehat{\epsilon}_\delta \geq \epsilon_\delta^* \right\} \geq 1 - \alpha \text{ for all } (\mathcal{A}, P, n). \quad (13)$$

(Equivalently, $\mathbb{P}\{T_{\widehat{\epsilon}_\delta, \delta}^* = 1\} \geq 1 - \alpha$.) In other words, given an algorithm \mathcal{A} and data drawn from P , we conclude with $1 - \alpha$ level confidence that (\mathcal{A}, P, n) is $(\widehat{\epsilon}_\delta, \delta)$ -stable.

It turns out that these inference targets are highly interconnected—any procedure that is able to achieve one can be adapted to achieve the other two. The following proposition says that any assumption-free estimation procedure (for either parameter) can be turned into an assumption-free test $\widehat{T}_{\epsilon, \delta}$, and vice versa.

Proposition 1. *The following equivalence holds for inference procedures $\widehat{T}_{\epsilon, \delta}$, $\widehat{\delta}_\epsilon$, and $\widehat{\epsilon}_\delta$:*

- (a) For any fixed $\epsilon \geq 0$, suppose there is an estimator $\widehat{\delta}_\epsilon$ satisfying (12). Then for any $\delta \in [0, 1)$, the test $\widehat{T}_{\epsilon, \delta} = \mathbb{1}\{\widehat{\delta}_\epsilon \leq \delta\}$ satisfies (2).
- (b) For any fixed $\delta \in [0, 1)$, suppose there is an estimator $\widehat{\epsilon}_\delta$ satisfying (13). Then for any $\epsilon \geq 0$, the test $\widehat{T}_{\epsilon, \delta} = \mathbb{1}\{\widehat{\epsilon}_\delta \leq \epsilon\}$ satisfies (2).
- (c) Suppose $\{\widehat{T}_{\epsilon, \delta} : \epsilon \geq 0, \delta \in [0, 1]\}$ is a family of tests such that each $\widehat{T}_{\epsilon, \delta}$ satisfies (2). Then, for any fixed $\epsilon \geq 0$, the estimator $\widehat{\delta}_\epsilon = \inf\{\delta : \widehat{T}_{\epsilon, \delta'} = 1 \forall \delta' \geq \delta\}$ satisfies (12).
- (d) Similarly, under the same assumptions as in (c), for any fixed $\delta \in [0, 1)$, the estimator $\widehat{\epsilon}_\delta = \inf\{\epsilon : \widehat{T}_{\epsilon', \delta} = 1 \forall \epsilon' \geq \epsilon\}$ satisfies (13).

The proof is in Appendix C.

Proposition 1 implies that the problems of assumption-free inference for any of these three targets are essentially equivalent. Therefore, although we have exclusively focused on the question of testing stability at a fixed (ϵ, δ) in the earlier sections, our conclusions apply to the other two inference problems as well to bound the power of any black box procedure. Specifically, assuming that $\widehat{\delta}_\epsilon$ (respectively, $\widehat{\epsilon}_\delta$) is a black box procedure in the same sense as Definition 2, combining Theorem 2 with part (a) (respectively, part (b)) of Proposition 1 will yield an upper bound on the probability of the event $\widehat{\delta}_\epsilon \leq \delta$ (respectively, $\widehat{\epsilon}_\delta \leq \epsilon$).

5 Discussion

In this work, we have shown a universal bound on the power achievable by any black box test of stability (Theorem 2), and have seen that, surprisingly, in some regimes the maximum power is achieved by a simple Binomial test that only calls the algorithm once for each batch of n labeled data points and one unlabeled test point (Theorem 1). To conclude, we will now discuss some implications and extensions of these results, which we leave up to future work.

5.1 The role of uncountability

Our optimality result (Theorem 2) applies to a test with *assumption-free* validity as defined by (2), i.e., the bound on power applies to a test that controls the Type I error for *any* pair of algorithm and data distribution. Thus, to prove the theorem, given an arbitrary (ϵ, δ) -stable triple (\mathcal{A}, P, n) , we constructed a non- (ϵ, δ) -stable triple (\mathcal{A}', P', n) such that the resulting predictions are hard to distinguish from one another given a limited amount of available data. Our proof relies on finding a point x_* in the feature space \mathbb{R}^d , and/or a point y_* in the response space \mathbb{R} , with zero probability of being observed at any point throughout the course of the black box testing procedure (i.e., at any stage of Definition 2). In particular, establishing the existence of such a point relies on the uncountability of \mathbb{R}^d (for finding x_*) or of \mathbb{R} (for finding y_*).

However, in some settings either X and/or Y may be a discrete random variable that is known to take values in a *countable* space $\mathbb{X} \subsetneq \mathbb{R}^d$ and/or $\mathbb{Y} \subsetneq \mathbb{R}$. In this type of setting, we may be more interested in a test that is guaranteed to be valid *only* for P supported on the relevant space (e.g., $\mathbb{X} \times \mathbb{R}$ or $\mathbb{R}^d \times \mathbb{Y}$), as opposed to over all possible P on $\mathbb{R}^d \times \mathbb{R}$. For this more restricted definition of validity, the proof technique of finding x_* and/or y_* can no longer be carried out, and consequently the upper bound on power (10) no longer applies. Assumption-free testing in the discrete setting was studied by Lee and Barber [2021] for a different problem—namely, for inference on error in regression—and it seems likely that testing stability with higher power may be achievable using similar techniques.

Note that this setting is different than simply assuming that X and/or Y is discrete but not assuming a known support. In particular, if we assume X and Y are discrete but any values in $\mathbb{R}^d \times \mathbb{R}$ might appear in its support, then we may require the validity statement (2) to hold only for triples (\mathcal{A}, P, n) where P is discrete; nonetheless, the same upper bound on power given in Theorem 2 will still hold.

5.2 Black box algorithms and black box functions, revisited

In Section 2.3.3 above, we noted that there may be some settings where the algorithm \mathcal{A} is sufficiently complex that it can only be thought of as a “black box” in any analysis, but a fitted function $\hat{\mu}$ produced by the algorithm is simple enough to analyze theoretically. As mentioned earlier, an example of this type of setting is where fitted functions $\hat{\mu}$ are always linear models (and therefore, easy to analyze theoretically), but the model fitting algorithm \mathcal{A} is itself can only be studied empirically (for instance, some complex procedure for variable selection whose theoretical properties are not well understood). In this setting, we can use similar arguments as in Theorem 2 to establish an analogous bound on power, with $\kappa = N_\ell/n$ in place of $\kappa = \min\{N_\ell/n, (N_\ell + N_u)/(n + 1)\}$. That is, if we are able to study the fitted $\hat{\mu}$ ’s analytically, this is essentially equivalent to having an unlimited amount of unlabeled data, but we are still constrained by the number N_ℓ of available data points.

5.3 A different target: data-conditional stability

We have just seen that it is essentially impossible to test whether \mathcal{A} is stable at sample size n , if $N_\ell = n$ is the number of labeled data points available for the test. However, if the reason that we want to test stability is for verifying that predictive inference methods such as jackknife can be applied to \mathcal{A} , testing whether (\mathcal{A}, P, n) is (ϵ, δ) -stable may be too strong of a goal. Specifically, given a training data set $(X_1, Y_1), \dots, (X_n, Y_n)$ and a test point (X_{n+1}, Y_{n+1}) (where Y_{n+1} is of course unobserved, since it is the target of predictive inference), examining the proof of Barber et al. [2021, Theorem 5] we see that the predictive coverage of the jackknife method relies on bounding

$$|\hat{\mu}_{n;-i}(X_i) - \hat{\mu}_{n+1;-i}(X_i)| \tag{14}$$

where $\hat{\mu}_{n;-i}$ denotes the model fitted by running \mathcal{A} on data points $((X_j, Y_j) : 1 \leq j \leq n, j \neq i)$ and $\hat{\mu}_{n+1;-i}$ denotes the model fitted by running \mathcal{A} on data points $((X_j, Y_j) :$

$1 \leq j \leq n + 1, j \neq i$).

Requiring stability of (\mathcal{A}, P, n) , as in Definition 1, ensures that the quantity (14) is small with high probability *over a random draw of the entire data set*. On the other hand, when using stability in the proof of the jackknife’s predictive coverage, we only need to know that the quantity (14) is small *on the actual data set*, which we are able to observe with the exception of the last point (X_{n+1}, Y_{n+1}) . In other words, we only need to test whether \mathcal{A} is stable with respect to a random draw of this single last data point (and we can condition on the training set). This notion of “data-conditional stability” appears to be a weaker property than the (unconditional) stability property of Definition 1, and therefore it may be possible to test this conditional property with higher power than the bounds established in Theorem 2.

Acknowledgements

B.K. acknowledge the support from the National Institutes of Health via grants R01GM114029 and R01GM133848. R.F.B. was supported by the National Science Foundation via grants DMS-1654076 and DMS-2023109, and by the Office of Naval Research via grant N00014-20-1-2337.

References

- Rina Foygel Barber, Emmanuel J. Candès, Aaditya Ramdas, and Ryan J. Tibshirani. Predictive inference with the jackknife+. *The Annals of Statistics*, 49(1):486–507, 2021. doi: 10.1214/20-AOS1965. URL <https://doi.org/10.1214/20-AOS1965>.
- Olivier Bousquet and André Elisseeff. Stability and generalization. *Journal of Machine Learning Research*, 2:499–526, 2002.
- Zhun Deng, Hangfeng He, and Weijie Su. Toward better generalization bounds with locally elastic stability. In Marina Meila and Tong Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pages 2590–2600. PMLR, 2021. URL <https://proceedings.mlr.press/v139/deng21b.html>.
- L. Devroye and T. Wagner. Distribution-free inequalities for the deleted and holdout error estimates. *IEEE Transactions on Information Theory*, 25(2):202–207, 1979a. doi: 10.1109/TIT.1979.1056032.
- L. Devroye and T. Wagner. Distribution-free performance bounds for potential function rules. *IEEE Transactions on Information Theory*, 25(5):601–604, 1979b. doi: 10.1109/TIT.1979.1056087.
- Michael Kearns and Dana Ron. Algorithmic stability and sanity-check bounds for leave-one-out cross-validation. *Neural Computation*, 11:1427–1453, 1999.

Samuel Kutin and Partha Niyogi. Almost-everywhere algorithmic stability and generalization error. Technical Report TR-2002-03, Department of Computer Science, The University of Chicago, 2002a.

Samuel Kutin and Partha Niyogi. Almost-everywhere algorithmic stability and generalization error. In *Proceedings of the Eighteenth Conference on Uncertainty in Artificial Intelligence*, UAI'02, pages 275–282, San Francisco, CA, USA, 2002b. Morgan Kaufmann Publishers Inc. ISBN 1558608974.

Yonghoon Lee and Rina Foygel Barber. Distribution-free inference for regression: discrete, continuous, and in between, 2021. arXiv preprint.

Sayan Mukherjee, Partha Niyogi, Tomaso Poggio, and Ryan Rifkin. Learning theory: Stability is sufficient for generalization and necessary and sufficient for consistency of empirical risk minimization. *Advances in Computational Mathematics*, 25(1):161–193, 2006. doi: 10.1007/s10444-004-7634-z. URL <https://doi.org/10.1007/s10444-004-7634-z>.

Glenford J. Myers, Corey Sandler, and Tom Badgett. *The Art of Software Testing*. Wiley Publishing, 3rd edition, 2011. ISBN 1118031962.

W. H. Rogers and T. J. Wagner. A finite sample distribution-free performance bound for local discrimination rules. *The Annals of Statistics*, 6(3):506–514, 5 1978. doi: 10.1214/aos/1176344196. URL <https://doi.org/10.1214/aos/1176344196>.

Shai Shalev-Shwartz, Ohad Shamir, Nathan Srebro, and Karthik Sridharan. Learnability, stability and uniform convergence. *Journal of Machine Learning Research*, 11(90): 2635–2670, 2010. URL <http://jmlr.org/papers/v11/shalev-shwartz10a.html>.

Lukas Steinberger and Hannes Leeb. Leave-one-out prediction intervals in linear regression models with many variables, 2016. arXiv preprint.

Lukas Steinberger and Hannes Leeb. Conditional predictive inference for high-dimensional stable algorithms, 2020. arXiv preprint.

A Proof of Theorem 1

As calculated in Section 3.1, we have shown that, for any algorithm \mathcal{A} and distribution P , it holds that $B \sim \text{Binomial}(\lfloor \kappa \rfloor, \delta_\epsilon^*)$ (note that this distribution depends on \mathcal{A} and on P implicitly via the parameter δ_ϵ^*).

Next we check validity, meaning that we need to show $\mathbb{P}\{\widehat{T}_{\epsilon,\delta} = 1\} \leq \alpha$. If $T_{\epsilon,\delta}^* = 0$. By definition, $T_{\epsilon,\delta}^* = 0$ implies that $\delta_\epsilon^* > \delta$, and so we have

$$\begin{aligned} \mathbb{P}\{\widehat{T}_{\epsilon,\delta} = 1\} &= \mathbb{P}\{B < k_{\kappa,\delta}^*\} + a_{\kappa,\delta}^* \cdot \mathbb{P}\{B = k_{\kappa,\delta}^*\} \\ &= (1 - a_{\kappa,\delta}^*) \cdot \mathbb{P}\{B < k_{\kappa,\delta}^*\} + a_{\kappa,\delta}^* \cdot \mathbb{P}\{B \leq k_{\kappa,\delta}^*\} \\ &\leq (1 - a_{\kappa,\delta}^*) \cdot \mathbb{P}\{\text{Binomial}(\lfloor \kappa \rfloor, \delta) < k_{\kappa,\delta}^*\} + a_{\kappa,\delta}^* \cdot \mathbb{P}\{\text{Binomial}(\lfloor \kappa \rfloor, \delta) \leq k_{\kappa,\delta}^*\} \\ &= \alpha. \end{aligned}$$

In the above, the inequality of the next-to-last step is because $\delta_\epsilon^* > \delta$, and hence, the $\text{Binomial}(\lfloor \kappa \rfloor, \delta)$ distribution is stochastically smaller than the distribution of $B \sim \text{Binomial}(\lfloor \kappa \rfloor, \delta_\epsilon^*)$. The equality of the last step is a direct consequence of the definitions of $k_{\kappa,\delta}^*$ and of $a_{\kappa,\delta}^*$.

Finally, we calculate power in the case that $T_{\epsilon,\delta}^* = 1$, for the two special cases $\delta_\epsilon^* = 0$ and $\delta \leq 1 - \alpha^{1/\lfloor \kappa \rfloor}$. For the second special case, $\delta \leq 1 - \alpha^{1/\lfloor \kappa \rfloor}$ implies that $k_{\kappa,\delta}^* = 0$, and so we have

$$\begin{aligned} \mathbb{P}\{\widehat{T}_{\epsilon,\delta} = 1\} &= a_{\kappa,\delta}^* \cdot \mathbb{P}\{B = 0\} = a_{\kappa,\delta}^* \cdot \mathbb{P}\{\text{Binomial}(\lfloor \kappa \rfloor, \delta_\epsilon^*) = 0\} \\ &= \alpha \cdot \frac{\mathbb{P}\{\text{Binomial}(\lfloor \kappa \rfloor, \delta_\epsilon^*) = 0\}}{\mathbb{P}\{\text{Binomial}(\lfloor \kappa \rfloor, \delta) = 0\}} = \alpha \cdot \left(\frac{1 - \delta_\epsilon^*}{1 - \delta}\right)^{\lfloor \kappa \rfloor} = \left\{ \alpha \cdot \left(\frac{1 - \delta_\epsilon^*}{1 - \delta}\right)^{\lfloor \kappa \rfloor} \right\} \wedge 1, \end{aligned}$$

where the third equality holds by definition of $a_{\kappa,\delta}^*$, and the last equality holds since $\delta \leq 1 - \alpha^{1/\lfloor \kappa \rfloor}$ implies $\alpha \cdot \left(\frac{1 - \delta_\epsilon^*}{1 - \delta}\right)^{\lfloor \kappa \rfloor} \leq \frac{\alpha}{(1 - \delta)^{\lfloor \kappa \rfloor}} \leq 1$. If instead only the first special case holds, i.e., $\delta_\epsilon^* = 0$ and $\delta > 1 - \alpha^{1/\lfloor \kappa \rfloor}$, then this implies that $B = 0$ almost surely and that $k_{\kappa,\delta}^* > 0$. Therefore,

$$\mathbb{P}\{\widehat{T}_{\epsilon,\delta} = 1\} = \mathbb{P}\{B < k_{\kappa,\delta}^*\} + a_{\kappa,\delta}^* \cdot \mathbb{P}\{B = k_{\kappa,\delta}^*\} = 1 = \left\{ \alpha \cdot \left(\frac{1 - \delta_\epsilon^*}{1 - \delta}\right)^{\lfloor \kappa \rfloor} \right\} \wedge 1,$$

where the last step holds since $\delta_\epsilon^* = 0$ and $\delta > 1 - \alpha^{1/\lfloor \kappa \rfloor}$ implies $\alpha \cdot \left(\frac{1 - \delta_\epsilon^*}{1 - \delta}\right)^{\lfloor \kappa \rfloor} = \frac{\alpha}{(1 - \delta)^{\lfloor \kappa \rfloor}} > 1$.

B Proof of Theorem 2

To prove (10), by definition of κ , it is equivalent to show that

$$\mathbb{P}\{\widehat{T}_{\epsilon,\delta}(\mathcal{A}, P, n) = 1\} \leq \alpha \cdot \left(\frac{1 - \delta_\epsilon^*}{1 - \delta}\right)^{N_\ell/n} \quad (15)$$

and that

$$\mathbb{P}\{\widehat{T}_{\epsilon,\delta}(\mathcal{A}, P, n) = 1\} \leq \alpha \cdot \left(\frac{1 - \delta_\epsilon^*}{1 - \delta}\right)^{(N_\ell + N_u)/(n+1)}. \quad (16)$$

The first inequality (15) shows the bound on power that is due to the limited amount of labeled data, while the second inequality (16) shows the bound on power that is due to the limited amount of unlabeled data (since the total available number of draws of X is equal to $N_\ell + N_u$).

The proofs of (15) and (16) both follow the same general recipe. The key idea of each proof is to construct an algorithm \mathcal{A}' and a data distribution P' where (\mathcal{A}', P', n) is not (ϵ, δ) -stable, but with \mathcal{A}' and P' sufficiently similar to \mathcal{A} and P so that the two are difficult to distinguish using the limited available data. The constructions will follow a shared structure:

- We will construct \mathcal{A}' and P' so that

$$(\mathcal{A}', P', n) \text{ does not satisfy } (\epsilon, \delta)\text{-stability}, \quad (17)$$

and therefore, we must have

$$\mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}'_\ell, \mathcal{D}'_u) = 1 \right\} \leq \alpha, \quad (18)$$

where $(\mathcal{D}'_\ell, \mathcal{D}'_u)$ are drawn from P' (i.e., \mathcal{D}'_ℓ consists of N_ℓ many labeled points drawn i.i.d. from P' , and \mathcal{D}'_u consists of N_u many unlabeled points drawn i.i.d. from the marginal distribution P'_X).

- Distribution P' is constructed to be similar to P , by defining it as a mixture model of the form

$$P' = (1 - c) \cdot P + c \cdot P_1$$

for some other distribution P_1 on $\mathbb{R}^d \times \mathbb{R}$.

With this definition, we can couple the processes of drawing data from P and drawing data from P' . Specifically, suppose data sets $(\mathcal{D}_\ell, \mathcal{D}_u)$ are drawn from P and data sets $(\mathcal{D}'_\ell, \mathcal{D}'_u)$ are drawn from P' . Then by definition of P' , we can construct a coupling of $(\mathcal{D}_\ell, \mathcal{D}_u)$ and $(\mathcal{D}'_\ell, \mathcal{D}'_u)$ such that

$$\mathbb{P} \{ (\mathcal{D}'_\ell, \mathcal{D}'_u) = (\mathcal{D}_\ell, \mathcal{D}_u) \mid (\mathcal{D}_\ell, \mathcal{D}_u) \} \geq (1 - c)^{N_\ell + N_u}. \quad (19)$$

Moreover, if P_1 and P have the same marginal distribution of X , then the two distributions differ only on the labeled samples, and so we will instead have

$$\mathbb{P} \{ (\mathcal{D}'_\ell, \mathcal{D}'_u) = (\mathcal{D}_\ell, \mathcal{D}_u) \mid (\mathcal{D}_\ell, \mathcal{D}_u) \} \geq (1 - c)^{N_\ell}. \quad (20)$$

- Algorithm \mathcal{A}' is constructed to be similar to \mathcal{A} , in the following sense. For data sets $(\mathcal{D}_\ell, \mathcal{D}_u)$ drawn from P , we will design \mathcal{A}' to satisfy

$$\mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta}(\mathcal{A}, \mathcal{D}_\ell, \mathcal{D}_u) = \widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}_\ell, \mathcal{D}_u) \right\} = 1, \quad (21)$$

i.e., running the test $\widehat{T}_{\epsilon, \delta}$ using algorithm \mathcal{A} or using algorithm \mathcal{A}' will result in the same answer almost surely. (On the other hand, \mathcal{A}' might not yield the same result as \mathcal{A} for the test $\widehat{T}_{\epsilon, \delta}$ for some adversarial choices of the data.)

Combining (21) with (18) and (19) yields

$$\begin{aligned}
\alpha &\geq \mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}'_\ell, \mathcal{D}'_u) = 1 \right\} \\
&\geq \mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}_\ell, \mathcal{D}_u) = 1, (\mathcal{D}'_\ell, \mathcal{D}'_u) = (\mathcal{D}_\ell, \mathcal{D}_u) \right\} \\
&= \mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}_\ell, \mathcal{D}_u) = 1 \right\} \cdot \mathbb{P} \left\{ (\mathcal{D}'_\ell, \mathcal{D}'_u) = (\mathcal{D}_\ell, \mathcal{D}_u) \mid \widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}_\ell, \mathcal{D}_u) = 1 \right\} \\
&\geq \mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta}(\mathcal{A}, \mathcal{D}_\ell, \mathcal{D}_u) = 1 \right\} \cdot (1 - c)^{N_\ell + N_u}.
\end{aligned}$$

Here the last step holds since

$$\begin{aligned}
&\mathbb{P} \left\{ (\mathcal{D}'_\ell, \mathcal{D}'_u) = (\mathcal{D}_\ell, \mathcal{D}_u) \mid \widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}_\ell, \mathcal{D}_u) = 1 \right\} \\
&= \mathbb{E} \left[\mathbb{P} \left\{ (\mathcal{D}'_\ell, \mathcal{D}'_u) = (\mathcal{D}_\ell, \mathcal{D}_u) \mid (\mathcal{D}_\ell, \mathcal{D}_u), \zeta^{(1)}, \xi^{(1)}, \zeta^{(2)}, \xi^{(2)}, \dots, \zeta \right\} \mid \widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}_\ell, \mathcal{D}_u) = 1 \right] \\
&= \mathbb{E} \left[\mathbb{P} \left\{ (\mathcal{D}'_\ell, \mathcal{D}'_u) = (\mathcal{D}_\ell, \mathcal{D}_u) \mid (\mathcal{D}_\ell, \mathcal{D}_u) \right\} \mid \widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}_\ell, \mathcal{D}_u) = 1 \right] \\
&\geq \mathbb{E} \left[(1 - c)^{N_\ell + N_u} \mid \widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}_\ell, \mathcal{D}_u) = 1 \right] \\
&= (1 - c)^{N_\ell + N_u},
\end{aligned}$$

where the second step holds since the noise terms $\zeta^{(1)}, \xi^{(1)}, \zeta^{(2)}, \xi^{(2)}, \dots, \zeta$ used in defining the (potentially randomized) test $\widehat{T}_{\epsilon, \delta}$ are generated independently of the data, while the third step applies (19). Rearranging terms in our work above, we have therefore established that

$$\mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta}(\mathcal{A}, \mathcal{D}_\ell, \mathcal{D}_u) = 1 \right\} \leq \alpha \cdot (1 - c)^{-(N_\ell + N_u)}. \quad (22)$$

For the case that P_1 and P have the same marginal distribution of X , combining (21) with (18) and (20) yields

$$\alpha \geq \mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}'_\ell, \mathcal{D}'_u) = 1 \right\} \geq \mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta}(\mathcal{A}, \mathcal{D}_\ell, \mathcal{D}_u) = 1 \right\} \cdot (1 - c)^{N_\ell}$$

with an analogous calculation, and so for this case, we have established that

$$\mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta}(\mathcal{A}, \mathcal{D}_\ell, \mathcal{D}_u) = 1 \right\} \leq \alpha \cdot (1 - c)^{-N_\ell}. \quad (23)$$

Finally, to prove the bound (15), by examining (23) we see that it suffices to prove that

For any $c > 1 - \left(\frac{1 - \delta}{1 - \delta_\epsilon^*} \right)^{1/n}$, we can choose P_1 and \mathcal{A}' such that

$$(P_1)_X = P_X \text{ and both (17) and (21) hold.} \quad (24)$$

Similarly, to prove the bound (16), by examining (22) we see that it suffices to prove that

$$\text{For any } c > 1 - \left(\frac{1 - \delta}{1 - \delta_\epsilon^*} \right)^{1/(n+1)}, \text{ we can choose } P_1 \text{ and } \mathcal{A}' \text{ such that}$$

$$\text{both (17) and (21) hold.} \quad (25)$$

To complete the proof, we will now give explicit constructions of P_1 and \mathcal{A}' to verify each of these claims.

B.1 Proof of (24)

Consider the data sets $(\mathcal{D}_\ell, \mathcal{D}_u)$ drawn from P , and the data sets $(\mathcal{D}_\ell^{(1)}, \mathcal{D}_u^{(1)}), (\mathcal{D}_\ell^{(2)}, \mathcal{D}_u^{(2)}), \dots$ generated by running the test $\widehat{T}_{\epsilon, \delta}$ on input data sets $(\mathcal{D}_\ell, \mathcal{D}_u)$ using algorithm \mathcal{A} . Write

$$\mathcal{D}_\ell = ((X_1, Y_1), \dots, (X_{N_\ell}, Y_{N_\ell})), \quad \mathcal{D}_u = (X_{N_\ell+1}, \dots, X_{N_\ell+N_u}),$$

and for each $r \geq 1$ write

$$\mathcal{D}_\ell^{(r)} = ((X_1^{(r)}, Y_1^{(r)}), \dots, (X_{N_\ell^{(r)}}^{(r)}, Y_{N_\ell^{(r)}}^{(r)})), \quad \mathcal{D}_u^{(r)} = (X_{N_\ell^{(r)}+1}^{(r)}, \dots, X_{N_\ell^{(r)}+N_u^{(r)}}^{(r)}).$$

(Note that $N_\ell^{(r)}, N_u^{(r)}$ might be random, but are required to be finite almost surely.)

Define

$$\mathcal{Y} = \left\{ Y_1, \dots, Y_{N_\ell}, Y_1^{(1)}, \dots, Y_{N_\ell^{(1)}}^{(1)}, Y_1^{(2)}, \dots \right\} \subset \mathbb{R},$$

the set of all Y values that are observed at any point when running the test. We claim that there exists some $y_* \in \mathbb{R}$ such that

$$\mathbb{P} \{y_* \in \mathcal{Y}\} = 0. \quad (26)$$

To see why, consider any $y \in \mathbb{R}$. Then by the union bound,

$$\mathbb{P} \{y \in \mathcal{Y}\} \leq \sum_{i=1}^n \mathbb{P} \{Y_i = y\} + \sum_{r \geq 1} \sum_{i \geq 1} \mathbb{P} \left\{ N_\ell^{(r)} \geq i \text{ and } Y_i^{(r)} = y \right\}.$$

Clearly, for each term in the sum, there are at most countably infinitely many values $y \in \mathbb{R}$ for which the corresponding probability term is positive. Therefore in total, there are only countably many $y \in \mathbb{R}$ for which the sum is positive. This implies that some y_* satisfying (26) must exist.

Next we define

$$P_1 = P_X \times \delta_{y_*},$$

where δ_{y_*} is the point mass at y_* . In other words, P_1 is defined by drawing X from P_X (the marginal distribution of X under P), and then setting $Y = y_*$ deterministically. Clearly, P_1 and P have the same marginal distribution of X by definition.

Next we define the algorithm \mathcal{A}' . Note that since $c > 1 - \{(1 - \delta)/(1 - \delta_\epsilon^*)\}^{1/n}$, we have that $c > 1 - \{(1 - \delta - a)/(1 - \delta_\epsilon^* - a)\}^{1/n}$ for sufficiently small $a > 0$. Specifically, this is true for any $a \in (0, \{(1 - \delta) - (1 - \delta_\epsilon^*)(1 - c)^n\}/\{1 - (1 - c)^n\})$. Fix any such a . For a labeled data set $\mathcal{D} = ((x_1, y_1), \dots, (x_m, y_m))$, write $\mathcal{D}_{-i} = ((x_1, y_1), \dots, (x_{i-1}, y_{i-1}), (x_{i+1}, y_{i+1}), \dots, (x_m, y_m))$ to denote the data set with the i -th point removed, and let $|\mathcal{D}| = m$ be its cardinality. The fitted model $\mathcal{A}'[\mathcal{D}; \xi]$ is then defined as follows: for any test point $x \in \mathbb{R}^d$, the outputted prediction is given by

$$\mathcal{A}'[\mathcal{D}; \xi](x) = \begin{cases} 1 + \epsilon + \mathcal{Q}_{1-a}[\mathcal{D}](x) & \text{if } |\mathcal{D}| = n \text{ and } \sum_{i=1}^n \mathbb{1}\{y_i = y_*\} \geq 1, \\ \mathcal{A}[\mathcal{D}; \xi](x) & \text{otherwise,} \end{cases}$$

where $\mathcal{Q}_{1-a}[\mathcal{D}](x)$ is the smallest value such that

$$\mathbb{P} \left\{ \max_{i=1, \dots, n} \mathcal{A}[\mathcal{D}_{-i}; \xi'](x) \leq \mathcal{Q}_{1-a}[\mathcal{D}](x) \mid \mathcal{D} \right\} \geq 1 - a,$$

i.e., $\mathcal{Q}_{1-a}[\mathcal{D}](x)$ is the $(1 - a)$ -quantile of the random variable $\max_{i=1, \dots, n} \mathcal{A}[\mathcal{D}_{-i}; \xi'](x)$ where we condition on \mathcal{D} so that the randomness is over the draw of $\xi' \sim \text{Uniform}(0, 1)$ only. This ensures that if $y_* \notin \mathcal{D}$ (i.e., the label y_* is not observed in the data), then the two algorithms yield the same output. On the other hand, if the label y_* is observed at least once in a sample of size n , then \mathcal{A}' exhibits instability.

Let $(X'_1, Y'_1), (X'_2, Y'_2), \dots \stackrel{\text{iid}}{\sim} P'$. Then defining

$$\widehat{\mu}'_n = \mathcal{A}'[(X'_1, Y'_1), \dots, (X'_n, Y'_n); \xi], \quad \widehat{\mu}'_{n-1} = \mathcal{A}'[(X'_1, Y'_1), \dots, (X'_{n-1}, Y'_{n-1}); \xi'],$$

we see that by definition of \mathcal{A}' , if $\sum_{i=1}^n \mathbb{1}\{Y'_i = y_*\} \geq 1$ then $|\widehat{\mu}'_n(X'_{n+1}) - \widehat{\mu}'_{n-1}(X'_{n+1})| \geq 1 + \epsilon > \epsilon$ with probability at least $1 - a$ over the draw of $\xi' \sim \text{Uniform}(0, 1)$, i.e.,

$$\mathbb{P} \left\{ |\widehat{\mu}'_n(X'_{n+1}) - \widehat{\mu}'_{n-1}(X'_{n+1})| > \epsilon \mid \sum_{i=1}^n \mathbb{1}\{Y'_i = y_*\} \geq 1 \right\} \geq 1 - a.$$

Moreover, by definition of P' together with the fact that $\mathbb{P}\{Y = y_*\} = 0$ for $Y \sim P$ (since $\mathbb{P}\{y_* \in \mathcal{D}_\ell\} = 0$), we see that

$$\mathbb{P} \left\{ \sum_{i=1}^n \mathbb{1}\{Y'_i = y_*\} \geq 1 \right\} = 1 - \prod_{i=1}^n \mathbb{P}\{Y'_i \neq y_*\} = 1 - (1 - c)^n,$$

as well as

$$\begin{aligned} & \mathbb{P} \left\{ |\widehat{\mu}'_n(X'_{n+1}) - \widehat{\mu}'_{n-1}(X'_{n+1})| > \epsilon, \sum_{i=1}^n \mathbb{1}\{Y'_i = y_*\} = 0 \right\} \\ &= \mathbb{P} \left\{ |\widehat{\mu}_n(X_{n+1}) - \widehat{\mu}_{n-1}(X_{n+1})| > \epsilon, \sum_{i=1}^n \mathbb{1}\{Y'_i = y_*\} = 0 \right\} \\ &= \mathbb{P} \{ |\widehat{\mu}_n(X_{n+1}) - \widehat{\mu}_{n-1}(X_{n+1})| > \epsilon \} \mathbb{P} \left\{ \sum_{i=1}^n \mathbb{1}\{Y'_i = y_*\} = 0 \right\} \\ &= \delta_\epsilon^* (1 - c)^n. \end{aligned}$$

(Here δ_ϵ^* is defined relative to the original triple (\mathcal{A}, P, n) .) Combining these calculations, we have

$$\begin{aligned} \mathbb{P} \{ |\widehat{\mu}'_n(X'_{n+1}) - \widehat{\mu}'_{n-1}(X'_{n+1})| > \epsilon \} \\ \geq (1-a) \{1 - (1-c)^n\} + \delta_\epsilon^* (1-c)^n \\ = (1-a) - \{(1-a) - \delta_\epsilon^*\} (1-c)^n > \delta, \end{aligned}$$

where the last step holds by how c and a are related to δ , δ_ϵ^* , and n . Therefore, (\mathcal{A}', P', n) is not (ϵ, δ) -stable, which verifies (17).

Finally, we verify (21), i.e., we need to check that $\widehat{T}_{\epsilon, \delta}$ will return the same answer using \mathcal{A}' as using \mathcal{A} when the input data is drawn from P .

By Definition 2, we know that

$$\widehat{T}_{\epsilon, \delta}(\mathcal{A}, \mathcal{D}_\ell, \mathcal{D}_u) = T \left[\mathcal{D}_\ell, \mathcal{D}_u, \left(\mathcal{D}_\ell^{(r)} \right)_{r \geq 1}, \left(\mathcal{D}_u^{(r)} \right)_{r \geq 1}, \left(\widehat{\mathcal{Y}}^{(r)} \right)_{r \geq 1}, \left(\zeta^{(r)} \right)_{r \geq 1}, \left(\xi^{(r)} \right)_{r \geq 1}, \zeta \right]$$

for some function T , where for each $r \geq 1$,

$$\left(\mathcal{D}_\ell^{(r)}, \mathcal{D}_u^{(r)} \right) = f^{(r)} \left[\mathcal{D}_\ell, \mathcal{D}_u, \left(\mathcal{D}_\ell^{(s)} \right)_{s=1}^{r-1}, \left(\mathcal{D}_u^{(s)} \right)_{s=1}^{r-1}, \left(\widehat{\mathcal{Y}}^{(s)} \right)_{s=1}^{r-1}, \left(\zeta^{(s)} \right)_{s=1}^{r-1}, \left(\xi^{(s)} \right)_{s=1}^{r-1}, \zeta^{(r)} \right]$$

and

$$\widehat{\mu}^{(r)} = \mathcal{A} \left[\mathcal{D}_\ell^{(r)}; \xi^{(r)} \right], \quad \widehat{\mathcal{Y}}^{(r)} = \widehat{\mu}^{(r)} \left[\mathcal{D}_u^{(r)} \right].$$

Similarly, we have

$$\widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}_\ell, \mathcal{D}_u) = T \left[\mathcal{D}_\ell, \mathcal{D}_u, \left(\widetilde{\mathcal{D}}_\ell^{(r)} \right)_{r \geq 1}, \left(\widetilde{\mathcal{D}}_u^{(r)} \right)_{r \geq 1}, \left(\widetilde{\mathcal{Y}}^{(r)} \right)_{r \geq 1}, \left(\zeta^{(r)} \right)_{r \geq 1}, \left(\xi^{(r)} \right)_{r \geq 1}, \zeta \right]$$

for the same function T and the same noise terms $\zeta^{(1)}, \xi^{(1)}, \zeta^{(2)}, \xi^{(2)}, \dots, \zeta \stackrel{\text{iid}}{\sim} \text{Uniform}[0, 1]$, where for each $r \geq 1$,

$$\left(\widetilde{\mathcal{D}}_\ell^{(r)}, \widetilde{\mathcal{D}}_u^{(r)} \right) = f^{(r)} \left[\mathcal{D}_\ell, \mathcal{D}_u, \left(\widetilde{\mathcal{D}}_\ell^{(s)} \right)_{s=1}^{r-1}, \left(\widetilde{\mathcal{D}}_u^{(s)} \right)_{s=1}^{r-1}, \left(\widetilde{\mathcal{Y}}^{(s)} \right)_{s=1}^{r-1}, \left(\zeta^{(s)} \right)_{s=1}^{r-1}, \left(\xi^{(s)} \right)_{s=1}^{r-1}, \zeta^{(r)} \right],$$

and

$$\widetilde{\mu}^{(r)} = \mathcal{A} \left[\widetilde{\mathcal{D}}_\ell^{(r)}; \xi^{(r)} \right], \quad \widetilde{\mathcal{Y}}^{(r)} = \widetilde{\mu}^{(r)} \left[\widetilde{\mathcal{D}}_u^{(r)} \right].$$

We will now verify that, for all $r \geq 1$, it holds almost surely that

$$\left(\mathcal{D}_\ell^{(r)}, \mathcal{D}_u^{(r)} \right) = \left(\widetilde{\mathcal{D}}_\ell^{(r)}, \widetilde{\mathcal{D}}_u^{(r)} \right), \quad \widehat{\mu}^{(r)} = \widetilde{\mu}^{(r)}, \quad \widehat{\mathcal{Y}}^{(r)} = \widetilde{\mathcal{Y}}^{(r)}. \quad (27)$$

First consider $r = 1$. Then we have

$$\left(\mathcal{D}_\ell^{(1)}, \mathcal{D}_u^{(1)} \right) = f^{(1)} \left[\mathcal{D}_\ell, \mathcal{D}_u, \zeta^{(1)} \right] = \left(\widetilde{\mathcal{D}}_\ell^{(1)}, \widetilde{\mathcal{D}}_u^{(1)} \right),$$

and therefore

$$\widehat{\mu}^{(1)} = \mathcal{A} \left[\mathcal{D}_\ell^{(1)}; \xi^{(1)} \right] = \mathcal{A}' \left[\mathcal{D}_\ell^{(1)}; \xi^{(1)} \right] = \mathcal{A}' \left[\widetilde{\mathcal{D}}_\ell^{(1)}; \xi^{(1)} \right] = \widetilde{\mu}^{(1)},$$

where the second equality holds whenever $N_\ell^{(1)} = n$ and $Y_i^{(1)} \neq y_*$ for all $i = 1, \dots, n$ by definition of \mathcal{A}' (recall that $\mathbb{P}\{Y_i^{(1)} \neq y_* \text{ for all } i = 1, \dots, N_\ell^{(1)}\} = 1$ by definition of y_*), and so we have

$$\widehat{\mathcal{Y}}^{(1)} = \widehat{\mu}^{(1)} \left[\mathcal{D}_u^{(1)} \right] = \widetilde{\mu}^{(1)} \left[\widetilde{\mathcal{D}}_u^{(1)} \right] = \widetilde{\mathcal{Y}}^{(1)}.$$

Next consider any $r \geq 2$. Suppose that (27) holds for all $s = 1, \dots, r-1$. Then,

$$\begin{aligned} & \left(\mathcal{D}_\ell^{(r)}, \mathcal{D}_u^{(r)} \right) \\ &= f^{(r)} \left[\mathcal{D}_\ell, \mathcal{D}_u, \left(\mathcal{D}_\ell^{(s)} \right)_{s=1}^{r-1}, \left(\mathcal{D}_u^{(s)} \right)_{s=1}^{r-1}, \left(\widehat{\mathcal{Y}}^{(s)} \right)_{s=1}^{r-1}, \left(\zeta^{(s)} \right)_{s=1}^{r-1}, \left(\xi^{(s)} \right)_{s=1}^{r-1}, \zeta^{(r)} \right] \\ &= f^{(r)} \left[\mathcal{D}_\ell, \mathcal{D}_u, \left(\widetilde{\mathcal{D}}_\ell^{(s)} \right)_{s=1}^{r-1}, \left(\widetilde{\mathcal{D}}_u^{(s)} \right)_{s=1}^{r-1}, \left(\widetilde{\mathcal{Y}}^{(s)} \right)_{s=1}^{r-1}, \left(\zeta^{(s)} \right)_{s=1}^{r-1}, \left(\xi^{(s)} \right)_{s=1}^{r-1}, \zeta^{(r)} \right] \\ &= \left(\widetilde{\mathcal{D}}_\ell^{(r)}, \widetilde{\mathcal{D}}_u^{(r)} \right), \end{aligned}$$

where the second equality holds by induction. Therefore we have

$$\widehat{\mu}^{(r)} = \mathcal{A} \left[\mathcal{D}_\ell^{(r)}; \xi^{(r)} \right] = \mathcal{A}' \left[\mathcal{D}_\ell^{(r)}; \xi^{(r)} \right] = \mathcal{A}' \left[\widetilde{\mathcal{D}}_\ell^{(r)}; \xi^{(r)} \right] = \widetilde{\mu}^{(r)},$$

where the second equality holds whenever $N_\ell^{(r)} = n$ and $Y_i^{(r)} \neq y_*$ for all $i = 1, \dots, n$ by definition of \mathcal{A}' (recall that $\mathbb{P}\{Y_i^{(r)} \neq y_* \text{ for all } i = 1, \dots, N_\ell^{(r)}\} = 1$ by definition of y_*), and so we have

$$\widehat{\mathcal{Y}}^{(r)} = \widehat{\mu}^{(r)} \left[\mathcal{D}_u^{(r)} \right] = \widetilde{\mu}^{(r)} \left[\widetilde{\mathcal{D}}_u^{(r)} \right] = \widetilde{\mathcal{Y}}^{(r)}.$$

Therefore, (27) holds almost surely for all $r \geq 1$, and so

$$\begin{aligned} & \widehat{T}_{\epsilon, \delta}(\mathcal{A}, \mathcal{D}_\ell, \mathcal{D}_u) \\ &= T \left[\mathcal{D}_\ell, \mathcal{D}_u, \left(\mathcal{D}_\ell^{(r)} \right)_{r \geq 1}, \left(\mathcal{D}_u^{(r)} \right)_{r \geq 1}, \left(\widehat{\mathcal{Y}}^{(r)} \right)_{r \geq 1}, \left(\zeta^{(r)} \right)_{r \geq 1}, \left(\xi^{(r)} \right)_{r \geq 1}, \zeta \right] \\ &= T \left[\mathcal{D}_\ell, \mathcal{D}_u, \left(\widetilde{\mathcal{D}}_\ell^{(r)} \right)_{r \geq 1}, \left(\widetilde{\mathcal{D}}_u^{(r)} \right)_{r \geq 1}, \left(\widetilde{\mathcal{Y}}^{(r)} \right)_{r \geq 1}, \left(\zeta^{(r)} \right)_{r \geq 1}, \left(\xi^{(r)} \right)_{r \geq 1}, \zeta \right] \\ &= \widehat{T}_{\epsilon, \delta}(\mathcal{A}', \mathcal{D}_\ell, \mathcal{D}_u) \end{aligned}$$

holds almost surely. This proves the claim (21), and thus completes our proof of (24).

B.2 Proof of (25)

Define

$$\mathcal{X} = \left\{ X_1, \dots, X_{N_\ell + N_u}, X_1^{(1)}, \dots, X_{N_\ell^{(1)} + N_u^{(1)}}^{(1)}, X_1^{(2)}, \dots \right\} \subset \mathbb{R}^d,$$

the set of all X values that are observed at any point when running the test. (Here the X_i 's and $X_i^{(k)}$'s are defined as in Section B.1). Then there exists some $x_* \in \mathbb{R}$ such that

$$\mathbb{P}\{x_* \in \mathcal{X}\} = 0$$

(the proof of this claim is analogous to the proof of (26) in Section B.1, so we do not repeat the details here). Define

$$P_1 = \delta_{(x_*, 0)},$$

i.e., the point mass at $(x_*, 0)$.

Next we define the algorithm \mathcal{A}' . Similarly to Section B.1, if $c > 1 - \{(1 - \delta)/(1 - \delta_\epsilon^*)\}^{1/(n+1)}$, then we have $c > 1 - \{(1 - \delta - a)/(1 - \delta_\epsilon^* - a)\}^{1/(n+1)}$ for any $a \in (0, \{(1 - \delta) - (1 - \delta_\epsilon^*)(1 - c)^{n+1}\}/\{1 - (1 - c)^{n+1}\})$. As before, fix any such a . Using the same notation as in the definition of \mathcal{A}' for Section B.1, for any labeled data set $\mathcal{D} = ((x_1, y_1), \dots, (x_m, y_m))$ and any test point $x \in \mathbb{R}^d$ define

$$\mathcal{A}'[\mathcal{D}; \xi](x) = \begin{cases} 1 + \epsilon + \mathcal{Q}_{1-a}[\mathcal{D}](x) & \text{if } |\mathcal{D}| = n, \text{ and } \sum_{i=1}^n \mathbb{1}\{x_i = x_*\} \geq 1 \text{ or } x = x_*, \\ \mathcal{A}[\mathcal{D}; \xi](x) & \text{otherwise,} \end{cases}$$

where, as before, $\mathcal{Q}_{1-a}[\mathcal{D}](x)$ is the smallest value such that

$$\mathbb{P}\left\{\max_{i=1, \dots, n} \mathcal{A}[\mathcal{D}_{-i}; \xi'](x) \leq \mathcal{Q}_{1-a}[\mathcal{D}](x) \mid \mathcal{D}\right\} \geq 1 - a \text{ for an arbitrary small } a > 0,$$

Now we need to verify (17), i.e., (\mathcal{A}', P', n) is not (ϵ, δ) -stable. This proof is similar to the analogous proof in Section B.1. First, we can observe that if $\sum_{i=1}^{n+1} \mathbb{1}\{X'_i = x_*\} \geq 1$ then $|\widehat{\mu}'_n(X'_{n+1}) - \widehat{\mu}'_{n-1}(X'_{n+1})| \geq 1 + \epsilon > \epsilon$ with probability at least $1 - a$ over the draw of $\xi' \sim \text{Uniform}(0, 1)$, i.e.,

$$\mathbb{P}\left\{|\widehat{\mu}'_n(X'_{n+1}) - \widehat{\mu}'_{n-1}(X'_{n+1})| > \epsilon \mid \sum_{i=1}^n \mathbb{1}\{Y'_i = y_*\} \geq 1\right\} \geq 1 - a.$$

We can also calculate

$$\mathbb{P}\left\{\sum_{i=1}^{n+1} \mathbb{1}\{X'_i = x_*\} \geq 1\right\} = 1 - \prod_{i=1}^{n+1} \mathbb{P}\{X'_i \neq x_*\} = 1 - (1 - c)^{n+1}.$$

Next, exactly as in Section B.1, we have

$$\begin{aligned} & \mathbb{P}\left\{|\widehat{\mu}'_n(X'_{n+1}) - \widehat{\mu}'_{n-1}(X'_{n+1})| > \epsilon, \sum_{i=1}^{n+1} \mathbb{1}\{X'_i = x_*\} = 0\right\} \\ &= \mathbb{P}\left\{|\widehat{\mu}_n(X_{n+1}) - \widehat{\mu}_{n-1}(X_{n+1})| > \epsilon, \sum_{i=1}^{n+1} \mathbb{1}\{X'_i = x_*\} = 0\right\} \\ &= \mathbb{P}\{|\widehat{\mu}_n(X_{n+1}) - \widehat{\mu}_{n-1}(X_{n+1})| > \epsilon\} \mathbb{P}\left\{\sum_{i=1}^{n+1} \mathbb{1}\{X'_i = x_*\} = 0\right\} \\ &= \delta_\epsilon^*(1 - c)^{n+1}. \end{aligned}$$

Combining these calculations, we obtain

$$\begin{aligned} \mathbb{P} \left\{ \left| \widehat{\mu}'_n(X'_{n+1}) - \widehat{\mu}'_{n-1}(X'_{n+1}) \right| > \epsilon \right\} \\ \geq (1-a) \left\{ 1 - (1-c)^{n+1} \right\} + \delta_\epsilon^* (1-c)^{n+1} \\ = (1-a) - \{(1-a) - \delta_\epsilon^*\} (1-c)^{n+1} > \delta, \end{aligned}$$

where the last step holds by how c and a are related to δ , δ_ϵ^* , and n . Therefore, (\mathcal{A}', P', n) is not (ϵ, δ) -stable, which verifies (17) for this setting.

It remains to verify (21). The proof of (21) for this setting is essentially identical to the analogous statement in Section B.1, so we omit it here.

C Proof of Proposition 1

Assume $T_{\epsilon, \delta}^* = 0$, i.e., $\delta < \delta_\epsilon^*$ and $\epsilon < \epsilon_\delta^*$. To prove part (a), we have

$$\begin{aligned} \mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta} = 1 \right\} &= \mathbb{P} \left\{ \widehat{\delta}_\epsilon \leq \delta \right\} \quad \text{by definition of } \widehat{T}_{\epsilon, \delta} \\ &\leq \mathbb{P} \left\{ \widehat{\delta}_\epsilon < \delta_\epsilon^* \right\} \quad \text{since } \delta < \delta_\epsilon^* \\ &= \mathbb{P} \left\{ T_{\epsilon, \widehat{\delta}_\epsilon} = 0 \right\} \quad \text{by definition of } T_{\epsilon, \widehat{\delta}_\epsilon} = 0 \\ &\leq \alpha \quad \text{by assumption-free validity of } \widehat{\delta}_\epsilon \text{ (12)}. \end{aligned}$$

Similarly, to prove (b), we have

$$\begin{aligned} \mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta} = 1 \right\} &= \mathbb{P} \left\{ \widehat{\epsilon}_\delta \leq \epsilon \right\} \quad \text{by definition of } \widehat{T}_{\epsilon, \delta} \\ &\leq \mathbb{P} \left\{ \widehat{\epsilon}_\delta < \epsilon_\delta^* \right\} \quad \text{since } \epsilon < \epsilon_\delta^* \\ &= \mathbb{P} \left\{ T_{\widehat{\epsilon}_\delta, \delta} = 0 \right\} \quad \text{by definition of } T_{\widehat{\epsilon}_\delta, \delta} \\ &\leq \alpha \quad \text{by assumption-free validity of } \widehat{\epsilon}_\delta \text{ (13)}. \end{aligned}$$

Next, to prove part (c), we have

$$\begin{aligned} \mathbb{P} \left\{ \widehat{\delta}_\epsilon < \delta_\epsilon^* \right\} &= \sup_{\delta < \delta_\epsilon^*} \mathbb{P} \left\{ \widehat{\delta}_\epsilon \leq \delta \right\} \\ &\leq \sup_{\delta < \delta_\epsilon^*} \mathbb{P} \left\{ \widehat{T}_{\epsilon, \delta} = 1 \right\} \quad \text{by definition of } \widehat{\delta}_\epsilon \\ &\leq \sup_{\delta < \delta_\epsilon^*} \alpha \quad \text{by assumption-free validity of } \widehat{T}_{\epsilon, \delta} \text{ (2)} \\ &= \alpha. \end{aligned}$$

Similarly, to verify (d), we have

$$\begin{aligned}\mathbb{P}\{\widehat{\epsilon}_\delta < \epsilon_\delta^*\} &= \sup_{\epsilon < \epsilon_\delta^*} \mathbb{P}\{\widehat{\epsilon}_\delta \leq \epsilon\} \\ &\leq \sup_{\epsilon < \epsilon_\delta^*} \mathbb{P}\{\widehat{T}_{\epsilon,\delta} = 1\} \quad \text{by definition of } \widehat{\epsilon}_\delta \\ &\leq \sup_{\epsilon < \epsilon_\delta^*} \alpha \quad \text{by assumption-free validity of } \widehat{T}_{\epsilon,\delta} \text{ (2)} \\ &= \alpha.\end{aligned}$$