

# Nonlocal pseudospin dynamics in a quantum Ising chain

K. L. Zhang and Z. Song\*

*School of Physics, Nankai University, Tianjin 300071, China*

The existence of Majorana zero modes in nontrivial phase of quantum Ising chain results in not only the Kramers-like degeneracy spectrum, but also dynamic response for non-Hermitian perturbation in the ordered phase [Phys. Rev. Lett. **126**, 116401 (2021)]. In this work, we investigate the possible response of the degeneracy spectrum for Hermitian perturbations. We propose a single-particle description of the model in the ordered phase, associating with an internal degree of freedom characterized as a pseudospin. The effective magnetic field, arising from both local and nonlocal perturbations in terms of string operators, exerts on the pseudospin. We show that the action of string operator can be realized via a quench under the local perturbations. As an application, any ground states and excited states for the Hamiltonian with perturbation can be selected to identify the quantum phase, by adding the other perturbations to trigger a quench and measuring the Loschmidt echo.

## I. INTRODUCTION

Identifying the quantum phase diagram of a physical system is of vital importance in both condensed matter physics and quantum information science. In the past few decades, a large number of theoretical and experimental studies in this field have emerged [1–12]. The transverse field Ising chain [13–15], which is one of the paradigmatic model to explore quantum phase transition (QPT) and quantum information science, play a key role in this realm. In this model, the integrability has possibly attracted the attention of the researchers the most. In spite of the simplicity, it possesses all the basic elements of the QPT. The competition between the nearest neighbor interaction and the external magnetic field leads to an ordered phase and a disordered phase, which are separated by a quantum critical point, accompanied by a symmetry spontaneous breaking.

Methodologically, triggering the quantum quench dynamics [16–20] by suddenly changing system parameters is frequently used to study the QPT. After the quench, the system undergoes nonequilibrium dynamics [16], and one can measure the Loschmidt echo (LE) [21–26] to quantify the deviation of the evolved state from the initial state. Generally speaking, since the evolved state containing information of both the initial state and the postquench Hamiltonian, the behavior of LE can reflect the physical properties of the system. Besides the application in QPT, it is interesting to employ the quench protocol to manipulating the spin degrees of freedom in quantum information science.

In this paper, we investigate the possible response of the degeneracy spectrum of the transverse field Ising chain to Hermitian perturbations. We focus on the model with open boundary condition. As a result, the Majorana zero modes [27] in nontrivial phase leads to the Kramers-like degeneracy spectrum, demonstrating the dynamic response for non-Hermitian perturbation in the

ordered phase [28]. Our motivation is to provide a quench protocol for this model with Hermitian perturbations, which is more convenient for experimental implementation. To this end, we propose a single-particle description of the model in the ordered phase, associating with an internal degree of freedom characterized as a pseudospin. The effective magnetic field exerts on the pseudospin, which arises from both local and nonlocal perturbations in terms of string operators. As an application, a quantum state lives in the twofold degenerate subspace of the system in the ordered phase can be considered as a single qubit, and with the effective magnetic field, the time evolution operator can achieve two quantum gates—phase gate and Hadamard gate. We further show that the string operator action can be realized via a quench under the local perturbations. Any ground states and excited states for the Hamiltonian with perturbation can be selected for identifying the quantum phase, by adding the other perturbations to trigger a quench and measuring the LE. A possible application in the thermal state is also discussed. In addition, numerical simulations for a finite-size system are provided to support our results.

The remainder of this paper is organized as follows. In Sec. II, we present the transverse field Ising chain and its symmetries. In Sec. III, we introduce the pseudospin description for the model in the ordered phase and investigate the response of the degenerate spectrum to perturbations. In Sec. IV, we discuss the realization of the nonlocal operator, and the possible applications in the QPT and thermal state with numerical results of LEs. Finally, we summarize our results in Sec. V.

## II. MODEL AND SYMMETRIES

In this section, we present the Hamiltonian and a brief review on its basic properties, based on which we perform our investigations in this work. The model considered is the transverse field Ising chain with open boundary

---

\* songtc@nankai.edu.cn

condition, defined by the Hamiltonian

$$H_0 = -J \sum_{j=1}^{N-1} \sigma_j^x \sigma_{j+1}^x + g \sum_{j=1}^N \sigma_j^z, \quad (1)$$

where  $\sigma_j^\alpha$  ( $\alpha = x, y, z$ ) are the Pauli operators on site  $j$  and parameter  $g$  ( $g > 0$ ) is the transverse field strength. For simplicity, the following discussion assumes that  $J = 1$ . It can be checked that the model respects two symmetries. The first one is the parity symmetry

$$[p, H_0] = 0, \quad (2)$$

where the parity operator is defined as  $p = \prod_{j=1}^N (-\sigma_j^z)$ . The second one is a little subtle and is crucial to our main conclusion [28, 29]. The model with periodic boundary condition is exactly solvable and has been well studied [13]. At zero temperature, the QPT at  $g = 1$  separates an ordered phase of the system ( $g < 1$ ) from a disordered phase ( $g > 1$ ). However, when we consider the model with open boundary condition, it possesses an exclusive symmetry in the ordered phase  $g < 1$  in thermodynamic limit. Defining a nonlocal operator

$$D = \frac{1}{2} \sqrt{1 - g^2} \sum_{j=1}^N g^{j-1} D_j, \quad (3)$$

with a position-dependent component

$$D_j = \prod_{l < j} (-\sigma_l^z) \sigma_j^x - i \prod_{l < N-j+1} (-\sigma_l^z) \sigma_{N-j+1}^y, \quad (4)$$

(where  $i = \sqrt{-1}$ ), we have the commutation relation

$$[D, H_0] = [D^\dagger, H_0] = 0, \quad (5)$$

which is referred to as edge-spin symmetry, since it is the outcome of the edge operator of the Kitaev chain [27]. Operator  $D$  is a fermion operator, obeying the relations  $\{D, D^\dagger\} = 1$  and  $D^2 = (D^\dagger)^2 = 0$ . It should be noted that it is contingent on the following conditions:  $g < 1$ , a large  $N$  limit, and open boundary. In addition, operator  $D$  is non-universal and  $g$ -dependent.

From these symmetries, we have following implications: the complete eigenstates  $\{|\psi_n^+\rangle, |\psi_n^-\rangle\}$  of  $H_0$  with eigenenergy  $\varepsilon_n^\pm$ ,  $H_0 |\psi_n^\pm\rangle = \varepsilon_n^\pm |\psi_n^\pm\rangle$ , span two invariant subspaces with

$$p |\psi_n^\pm\rangle = \pm |\psi_n^\pm\rangle, \quad (6)$$

for any value of  $g$ . Importantly, within the region  $g < 1$ , we have the relations

$$\begin{aligned} D |\psi_n^+\rangle &= |\psi_n^-\rangle, D^\dagger |\psi_n^-\rangle = |\psi_n^+\rangle, \\ D^\dagger |\psi_n^+\rangle &= D |\psi_n^-\rangle = 0, \end{aligned} \quad (7)$$

which guarantee the existence of eigenstates degeneracy  $\varepsilon_n^+ = \varepsilon_n^- = \varepsilon_n$ , referred to as Kramers-like degeneracy. Accordingly, we also have the relations

$$i (D^\dagger - D) (|\psi_n^+\rangle \pm |\psi_n^-\rangle) = \pm i (|\psi_n^+\rangle \mp |\psi_n^-\rangle), \quad (8)$$

and

$$p (|\psi_n^+\rangle \pm |\psi_n^-\rangle) = |\psi_n^+\rangle \mp |\psi_n^-\rangle, \quad (9)$$

which play an important role in the quench dynamics, as demonstrated in the following section.

### III. NONLOCAL PSEUDOSPIN AND LOSCHMIDT ECHO

It is not surprising that a Hermitian perturbation can lift the degeneracy. However, it should lead to a fascinating dynamic phenomenon if an identical split in each level in the spectrum is obtained. Moreover, hybridizing two robust degenerate states on demand is a central task of quantum information processing, since these states are immune to weak local perturbations. We first focus on the ordered quantum phase  $0 < g < 1$ , considering a perturbed Hamiltonian

$$H = H_0 + H', \quad (10)$$

with  $H'$  being the combination of three types of actions,

$$H' = \kappa_x (D^\dagger + D) + i\kappa_y (D^\dagger - D) + \kappa_z p, \quad (11)$$

or explicitly,

$$\begin{aligned} H' &= \kappa_x \sqrt{1 - g^2} \sum_{j=1}^N g^{j-1} \prod_{l < j} (-\sigma_l^z) \sigma_j^x \\ &\quad - \kappa_y \sqrt{1 - g^2} \sum_{j=1}^N g^{j-1} \prod_{l < N-j+1} (-\sigma_l^z) \sigma_{N-j+1}^y \\ &\quad + \kappa_z \prod_{j=1}^N (-\sigma_j^z). \end{aligned} \quad (12)$$

We note that the perturbation is nonlocal, containing the string operators  $\prod_l (-\sigma_l^z)$ . Any pair of degenerate eigenstates  $(|\psi_n^+\rangle, |\psi_n^-\rangle)$  with energy  $\varepsilon_n$  spans a diagonal block in the form

$$\mathbf{B} \cdot \boldsymbol{\sigma} = \begin{pmatrix} \kappa_z & \kappa_x - i\kappa_y \\ \kappa_x + i\kappa_y & -\kappa_z \end{pmatrix}, \quad (13)$$

with parameter vector  $\mathbf{B} = (\kappa_x, \kappa_y, \kappa_z)$  and Pauli matrix  $\boldsymbol{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ . Therefore, under the basis  $(|\psi_1^+\rangle, |\psi_1^-\rangle, |\psi_2^+\rangle, |\psi_2^-\rangle, \dots)$ , we get an equivalent Hamiltonian for  $H$

$$H_{\text{eq}} = \bigoplus_{n=1}^{2^{N-1}} (\mathbf{B} \cdot \boldsymbol{\sigma} + \varepsilon_n I_2), \quad (14)$$

where  $I_2$  denotes the  $2 \times 2$  identity matrix. This is a single-particle description with the Pauli matrix presents a nonlocal pseudo spin, due to the robustness of the edge-spin symmetry against the random variations on the uniform distribution of parameters  $(J, g)$  in  $H_0$ .

The introduction of the perturbation term  $H'$  allows two possible applications through time evolution. First, the operation on the robust degenerate state can be realized. In the degenerate subspace with index  $n$ , an arbitrary state  $|\psi_n\rangle = \alpha|\psi_n^+\rangle + \beta|\psi_n^-\rangle$  acts as a single qubit, where  $\alpha$  and  $\beta$  are complex numbers encoding quantum information. An arbitrary unitary operation on this state can be realized by the time evolution operator

$$U(t) = e^{-i\varepsilon_n t} e^{-i\mathbf{B}\cdot\boldsymbol{\sigma}t} = e^{-i\varepsilon_n t} \left[ \cos(|\mathbf{B}|t) - i \frac{\mathbf{B}\cdot\boldsymbol{\sigma}}{|\mathbf{B}|} \sin(|\mathbf{B}|t) \right], \quad (15)$$

by choosing appropriate parameters  $\mathbf{B} = (\kappa_x, \kappa_y, \kappa_z)$  and evolved time  $t$ . For example, when choosing  $\mathbf{B} = (0, 0, \kappa_z)$ , it realizes the action of the phase gate

$$\mathcal{P} = e^{i\varepsilon_n t + i\kappa_z t} U(t) = \begin{pmatrix} 1 & 0 \\ 0 & e^{2i\kappa_z t} \end{pmatrix}. \quad (16)$$

When  $\mathbf{B} = (\kappa_x, 0, \kappa_x)$  and  $t' = \pi/(2\mathbf{B})$ , we get the Hadamard gate

$$\mathcal{H} = i e^{i\varepsilon_n t'} U(t') = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}. \quad (17)$$

Second, taking advantage of the perturbation term  $H'$ , the ordered and disordered quantum phases can be identified by the quench dynamics. In the following, we only give an analytical analysis of the expected results with parameter in the ordered quantum phase. since  $H'$  is  $g$ -dependent and is only defined in the region  $0 < g < 1$ . Based on a simplified version of  $H'$ , numerical results with parameters in both phases are presented in the next section. It can be checked that the solution of  $H_{\text{eq}}$  is simply given by the eigenvectors of  $\mathbf{B}\cdot\boldsymbol{\sigma}$ , i.e.,

$$|\phi_n^+\rangle = \cos\frac{\theta}{2} |\psi_n^+\rangle + \sin\frac{\theta}{2} e^{i\varphi} |\psi_n^-\rangle, \quad (18)$$

$$|\phi_n^-\rangle = \sin\frac{\theta}{2} |\psi_n^+\rangle - \cos\frac{\theta}{2} e^{i\varphi} |\psi_n^-\rangle, \quad (19)$$

with corresponding eigenenergies

$$E_n^\pm = \varepsilon_n \pm |\mathbf{B}| = \varepsilon_n \pm \sqrt{\kappa_x^2 + \kappa_y^2 + \kappa_z^2}, \quad (20)$$

where  $\theta$  and  $\varphi$  are defined as  $\cos\theta = \kappa_z/|\mathbf{B}|$  and  $\tan\varphi = \kappa_y/\kappa_x$ . The level spacing  $2\sqrt{\kappa_x^2 + \kappa_y^2 + \kappa_z^2}$  is  $n$  and  $g$  independent, resulting in a periodic dynamics among all the spectrum. It is exclusive and then is the manifestation of the ordered quantum phase.

To capture the effect of perturbation  $H'$  on the dynamics, we introduce the LE, which is a measure of reversibility and sensitivity to perturbation of quantum evolution. An initial quantum state  $|\Phi(0)\rangle$  evolves during a time  $t$  under a postquench Hamiltonian  $H_{\text{Pos}}$  reaching state  $|\Phi(t)\rangle$ . The behavior of  $\langle\Phi(0)|\Phi(t)\rangle$  is the basis of LE measurement, in which  $|\Phi(0)\rangle$  is an eigenstate of prequench Hamiltonian  $H_{\text{Pre}}$ . Two Hamiltonian  $H_{\text{Pre}}$  and

$H_{\text{Pos}}$  can be taken by two different sets of parameters  $(\kappa_x, \kappa_y, \kappa_z)$  for  $H$ . The LE is defined as

$$L(t) = |\langle\Phi(0)| e^{iH_{\text{Pos}}t} e^{-iH_{\text{Pre}}t} |\Phi(0)\rangle|^2, \quad (21)$$

where  $|\Phi(0)\rangle$  is usually an easily prepared state, such as the ground state of  $H_{\text{Pre}}$ . The dynamics in each subspace is clearly a rotation of a Bloch state with a fixed axis and frequency. Then a maximum oscillating amplitude is achieved when the initial Bloch state is perpendicular to the axis. In this sense,  $H_{\text{Pre}}$  cannot be taken as  $H_0$  with parameters  $(\kappa_x, \kappa_y, \kappa_z) = (0, 0, 0)$ , since  $|\Phi(0)\rangle$  is uncertain in practice due to the degeneracy. However, we can consider the setup as  $H_{\text{Pre}} = H_0 + \kappa_x(D^\dagger + D)$  and  $H_{\text{Pos}} = H_0 + \kappa_x(D^\dagger + D) + i\kappa_y(D^\dagger - D)$ , and the initial state is given as the ground state of the prequench Hamiltonian:  $|\Phi(0)\rangle = (|\psi_n^+\rangle - |\psi_n^-\rangle)/\sqrt{2}$ , under which we have

$$L(t) = |\langle\Phi(0)| e^{iH_{\text{Pos}}t} |\Phi(0)\rangle|^2 = \frac{2\kappa_x^2 + \kappa_y^2 + \kappa_y^2 \cos\left(2\sqrt{\kappa_x^2 + \kappa_y^2}t\right)}{2(\kappa_x^2 + \kappa_y^2)}. \quad (22)$$

We note that  $L(t)$  oscillates with period  $\tau = \pi/\sqrt{\kappa_x^2 + \kappa_y^2}$  and amplitude  $\kappa_y^2/(\kappa_x^2 + \kappa_y^2)$ , which tends to the maximum 1 in the limit of  $\kappa_x^2 \ll \kappa_y^2$ . The same conclusion can be obtained when we consider the case with replacing  $i\kappa_y(D^\dagger - D)$  by  $\kappa_z p$  in  $H_{\text{Pos}}$ .

#### IV. IDENTIFICATION OF THE QUANTUM PHASE

In this section, we analyze the characters of three types of perturbation and try to propose a practical scheme to demonstrate the dynamic detection of the phase diagram in experiment. We note that  $p$  is a typical string operator, which is a challenge to realized in experiment. We first consider the possible realization of the perturbation in practice. The perturbation term  $H'$  in Eq. (11) commutes to the unperturbed Hamiltonian  $H_0$ , then the whole Hamiltonian  $H_0 + H'$  is exactly solvable. However, the operators in  $H'$  are  $g$  dependent and need to be deliberately designed in practice.

##### A. Simplified perturbations

First, we consider a simplification of the perturbation in Eq. (11), that is, only the dominant terms of  $H'$  are taken into account:

$$H'_S = \kappa_x \sigma_1^x - \kappa_y \prod_{l=1}^{N-1} (-\sigma_l^z) \sigma_N^y + \kappa_z \prod_{j=1}^N (-\sigma_j^z) = \kappa_x \sigma_1^x - i\kappa_y p \sigma_N^x + \kappa_z p, \quad (23)$$

which is equal to a small  $g$  limit of  $H'$ . We find that the local operator  $\sigma_{1(N)}^x$  and the nonlocal operator  $p$  are two




FIG. 1. Spectrum of the low-lying eigenstates for Hamiltonian  $H = H_0 + H'_S$  as a function of  $g$  with parameters  $(\kappa_x, \kappa_y, \kappa_z)$  (a)  $(0, 0, 0)$ , (b)  $(0.1, 0, 0)$ , (c)  $(0, 0.1, 0)$  and (d)  $(0, 0, -0.1)$ , obtained numerically through exact diagonalization. The red and black colors of the lines in (a) and (d) denote even and odd parities of the corresponding eigenstates, respectively, while the eigenstates of spectrum (b) and (c) are not the eigenstates of parity operator  $p$ . Here  $E_g$  is the ground-state energy. System parameters are  $N = 10$  and  $J = 1$ .

elemental actions of the perturbations. The advantages of considering this perturbation are two folds: (i)  $H'_S$  is independent of the system parameter  $g$ , allowing us to implement the LE detection when the system parameter is unknown. (ii) The form of  $H'_S$  is simpler and thus is more convenient for experimental implementation. To see the effects of the perturbation  $H'_S$  on the spectrum of  $H_0$ , in Fig. 1, we present the spectrum of the low-lying eigenstates of the Hamiltonian  $H = H_0 + H'_S$  for different parameters  $(\kappa_x, \kappa_y, \kappa_z)$  and  $g$ . We can see that in the ordered phase, the perturbations with different  $(\kappa_x, \kappa_y, \kappa_z)$  all lead to almost equal level splitting for a fixed  $g$  and as  $g$  varying, which suggest that perturbation  $H'_S$  has the same effect as that in Eq. (11). As a local perturbation,  $\sigma_1^x$  can lift the degeneracy [see Fig. 1(b)], which has the same effect as the nonlocal case in Figs. 1(c) and 1(d). We also note that perturbations  $\kappa_x \sigma_1^x$  and  $-i\kappa_y p \sigma_N^x$  both break the parity symmetry, while  $\kappa_z p$  preserves it [see the red (even parity) and black (odd parity) lines in Figs. 1(a) and 1(d)].

In contrast, we investigate the effect of another type of perturbation, containing operator  $\sigma_N^y$ . Without loss of generality, we consider the case with Hamiltonian

$$H = H_0 + \sum_{j=1}^N \gamma_j \sigma_j^y, \quad (24)$$

where  $\gamma_j$  is an arbitrary set of real numbers. Applying a

set of local transformation [29]

$$\begin{aligned} \tau_j^x &= \sigma_j^x, \\ \tau_j^y &= \eta_j^+ \sigma_j^y - \eta_j^- \sigma_j^z, \\ \tau_j^z &= \eta_j^+ \sigma_j^z + \eta_j^- \sigma_j^y, \end{aligned} \quad (25)$$

with the factors  $\eta_j^+ = 1/\sqrt{1+\gamma_j^2}$  and  $\eta_j^- = \gamma_j/\sqrt{1+\gamma_j^2}$ , we have

$$H = -J \sum_{j=1}^{N-1} \tau_j^x \tau_{j+1}^x + \sum_{j=1}^N \sqrt{g^2 + \gamma_j^2} \tau_j^z, \quad (26)$$

which is still an Ising model since the new spin operators still satisfy the Lie algebra commutation relations

$$[\tau_j^\mu, \tau_j^\nu] = \sum_{\lambda=x,y,z} 2i\epsilon^{\mu\nu\lambda} \tau_j^\lambda. \quad (27)$$

Then weak perturbation  $\kappa \sigma_N^y$  cannot lift the degeneracy of  $H_0$ , since small derivation from uniform  $g$  does not affect the Majorana zero modes [27].

## B. Realization of string operator action

Second, we consider to realize the action of string operator  $p$  by the time evolution under a time-dependent local Hamiltonian

$$H_p(t) = \mathbf{g}(t) \sum_{l=1}^N \sigma_l^z, \quad (28)$$

which describes the action of an extra time-dependent transverse field. Here the coefficient is defined as

$$\mathbf{g}(t) = \begin{cases} \frac{\pi}{2\Delta}, & 0 < t \leq \Delta \\ 0, & \text{otherwise} \end{cases}. \quad (29)$$

After time  $\Delta$ , the effect of  $H_p(t)$  on the degenerate state  $|\psi_n\rangle = \alpha |\psi_n^+\rangle + \beta |\psi_n^-\rangle$  can be expressed as the time evolution operator

$$\begin{aligned} U(\Delta) &= \exp \left[ -i \int_0^\Delta H_p(t) dt \right] \\ &= \prod_{l=1}^N \exp \left( -i \int_0^\Delta \frac{\pi}{2\Delta} \sigma_l^z dt \right) \\ &= (i)^N \prod_{l=1}^N (-\sigma_l^z). \end{aligned} \quad (30)$$

It indicates that the time evolution operator takes the role of the operator  $p$ , i.e.,

$$U(\Delta) (\alpha |\psi_n^+\rangle + \beta |\psi_n^-\rangle) = (i)^N (\alpha |\psi_n^+\rangle - \beta |\psi_n^-\rangle). \quad (31)$$




FIG. 2. (a) Numerical results of LEs for the quench processes under the prequench and postquench Hamiltonians Eq. (32), with parameters  $g = 0.5$  and  $1.5$ . Other parameters are  $N = 12$ ,  $\kappa = 0.05$  and  $\Delta = 0.10$ . (b) Numerical results of LEs for the quench processes under Hamiltonians Eq. (35) for different  $g$  values. The gray dashed line denotes the minimum value of LE obtained from Eq. (22). The system size is  $N = 12$ . (c) Numerical results of average LEs as functions of  $g$  [Eq. (36)] for  $N = 8, 10$  and  $12$ . The gray dashed line represents the ideal average LEs expected for large  $N$  limits, and we set  $T = 500$ .

To verify this result, we perform numerical simulation for a quench process defined as

$$\begin{aligned} H_{\text{Pre}} &= H_0 + \kappa_x \sigma_1^x, \\ H_{\text{Pos}} &= H_0 + H_p. \end{aligned} \quad (32)$$

The initial state is taken as the ground state of the prequench Hamiltonian

$$|\Phi(0)\rangle = \sin \frac{\theta}{2} |\psi_g^+\rangle - \cos \frac{\theta}{2} e^{i\varphi} |\psi_g^-\rangle. \quad (33)$$

Then in the ordered phase, the expected final state is

$$\begin{aligned} |\Phi(\Delta)\rangle &= U(\Delta) |\Phi(0)\rangle \\ &= (i)^N \left( \sin \frac{\theta}{2} |\psi_g^+\rangle + \cos \frac{\theta}{2} e^{i\varphi} |\psi_g^-\rangle \right). \end{aligned} \quad (34)$$

It is expected that the LE obeys  $L(\Delta) = \cos^2 \theta = \kappa_z^2 / |\mathbf{B}|^2 = 0$ . While in the disordered phase, the ground state of  $H_0$  is non-degenerate and is separated from the excited state by an energy gap, then it is expected that  $L(\Delta) \approx 1$ . The numerical results of LEs obtained by exact diagonalization with system parameters  $g = 0.5$  and  $1.5$  are presented in Fig. 2(a), which are in accord with our analysis. Here the evolved state is calculated as  $|\Phi(t)\rangle = \exp(-iH_{\text{Pos}}t) |\Phi(0)\rangle$ , and the numerical computation is performed using a uniform mesh in the time discretization.

### C. Quantum phase transition

According to the conclusions in Sec. III and Sec. IV A, it is expected that the dynamics behavior of LE can be utilized to identify different quantum phases when we consider a simplified version of the perturbation in Eq. (23). It should lead to the similar oscillating behavior of LE described in Eq. (22) in the ordered phase if one implement the quench protocol. In contrast, such an oscillating behavior of  $L(t)$  in the region of  $g > 1$  is absent, since the non-degenerate eigenstates are not sensitive to

the perturbation of  $H'_S$ . These features allow us to observe significantly different dynamical behaviors in different quantum phase when the initial state is chosen as any eigenstate of  $H_{\text{Pre}}$ .

In the following, we consider the numerical simulation of quench process under the Hamiltonian

$$\begin{aligned} H_{\text{Pre}} &= H_0 + H'_S(\kappa_x = 0.05, \kappa_y = 0, \kappa_z = 0), \\ H_{\text{Pos}} &= H_0 + H'_S(\kappa_x = 0.05, \kappa_y = 0.1, \kappa_z = 0), \end{aligned} \quad (35)$$

where  $H'_S(\kappa_x, \kappa_y, \kappa_z)$  is defined in Eq. (23). The initial state is prepared as the ground state of  $H_{\text{Pre}}$ . In Fig. 2(b), we presented the LEs for different  $g$ , which are calculated by exact diagonalization. We can see that the results are in accord with our predictions for both phase. For small  $g$ , the minimum value of LE is the same as that of Eq. (22), which is estimated under the complex version of quench term in Eq. (11).

This verify that the proposed quench protocol can be utilized to identify different quantum phase of the transverse field Ising chain. To compare with the theoretical phase diagram, we introduce the average LE in the time interval  $[0, T]$ , which is defined as

$$\overline{L(g)} = \frac{1}{T} \int_0^T L(t) dt, \quad (36)$$

the value of which in the ordered phase can be estimated from Eq. (22) in long-time limit, that is

$$\overline{L(g < 1)} = \frac{2\kappa_x^2 + \kappa_y^2}{2(\kappa_x^2 + \kappa_y^2)}. \quad (37)$$

While in the disordered phase, it is expected that  $\overline{L(g > 1)} = 1$ . The numerical results of average LEs for different  $g$  and  $N$  are presented in Fig. 2(c), which are obtained by exact diagonalization. We can see that when the system size is larger, the average LE is closer to the ideal values (gray dashed line) that are expected in the thermodynamic limit.




FIG. 3. Numerical results of LEs for the quench processes under Hamiltonian Eq. (44) and three different initial thermal states with  $\beta = 1$ . The parameters of the system are  $g = 0.4$ ,  $N = 10$  and  $\kappa_x = 0.1$ .

#### D. Thermal state

Now we discuss a possibility of applying the quench protocol to the thermal state when  $g < 1$ . In the previous section, we have known that for the Ising chain with parameters  $g < 1$ , the robust degeneracy occurs not only in the ground states, but in all energy levels [see Eq. (7) and Fig. 1(a)]. In general, a thermal state of system  $H_0$  with temperature  $\beta^{-1}$  can be written as  $\rho_0 = e^{-\beta H_0} / \text{Tr} e^{-\beta H_0}$ , which preserves no quantum information. However, the robust degeneracy of the spectrum may enable a thermal state to preserve the quantum information in each degenerate subspace, in the case that the thermalization is induced by local perturbation from the environment.

Consider such a state as an initial state, with density matrix

$$\rho = \frac{\sum_{n=1}^{2^{N-1}} e^{-\beta E_n} |\Phi_n\rangle \langle \Phi_n|}{\sum_{n=1}^{2^{N-1}} e^{-\beta E_n}}, \quad (38)$$

where

$$|\Phi_n\rangle = \sin \frac{\theta_n}{2} |\psi_n^+\rangle - \cos \frac{\theta_n}{2} e^{i\varphi_n} |\psi_n^-\rangle. \quad (39)$$

At first, we estimate the dynamics of thermal state  $\rho$  under the quenched Hamiltonian

$$H_{\text{Pos}} = H_0 + \kappa_x (D^\dagger + D). \quad (40)$$

In fact, we have

$$\begin{aligned} & |\langle \Phi_n | e^{-iH_{\text{Pos}}t} | \Phi_n \rangle|^2 \\ &= \cos^2(\kappa_x t) + \sin^2(\kappa_x t) \sin^2 \theta_n \cos^2 \varphi_n, \end{aligned} \quad (41)$$

which is periodic function of time. We consider two types of distribution of  $\{\theta_n, \varphi_n\}$ , which are encoded with different information. (i) Random distribution:  $\{\theta_n\}$  and  $\{\varphi_n\}$  are respectively taken as the random samples that are uniformly distributed over the intervals  $[0, \pi)$  and

$[0, 2\pi)$ . Then the LE can be estimated by ignoring the Boltzmann factor in high temperature limit, that is

$$\begin{aligned} \bar{L}(t) &= \frac{1}{2^{N-1}} \sum_{n=1}^{2^{N-1}} |\langle \Phi_n | e^{-iH_{\text{Pos}}t} | \Phi_n \rangle|^2 \\ &\approx \frac{1}{2\pi^2} \int_0^\pi d\theta \int_0^{2\pi} d\varphi |\langle \Phi_n | e^{-iH_{\text{Pos}}t} | \Phi_n \rangle|^2 \\ &= \frac{1}{4} + \frac{3}{4} \cos^2(\kappa_x t). \end{aligned} \quad (42)$$

(ii) Fixed parity: the thermal state consists of the levels of the same parity, that is,  $\theta_n = \pi$  and  $\{\varphi_n\}$  is taken as a random sample in  $[0, 2\pi)$ . Then we have

$$\begin{aligned} \bar{L}(t) &\approx \frac{1}{2\pi} \int_0^{2\pi} d\varphi \cos^2(\kappa_x t) \\ &= \cos^2(\kappa_x t). \end{aligned} \quad (43)$$

For both cases,  $\bar{L}(t)$  are periodic functions but with different amplitudes, and note that the minimal value for the latter is zero.

In practice, based on the above analysis, one can consider the following quenched Hamiltonian instead of Eq. (40)

$$H_{\text{Pos}} = H_0 + \kappa_x \sigma_1^x, \quad (44)$$

and the definition of LE for density matrix is

$$L(t) = \left[ \text{Tr} \sqrt{\sqrt{\rho(0)} \rho(t) \sqrt{\rho(0)}} \right]^2, \quad (45)$$

which is also known as the Uhlmann fidelity [30, 31], characterizing the similarity between the initial state  $\rho(0)$  and evolved state  $\rho(t) = e^{-iH_{\text{Pos}}t} \rho(0) e^{iH_{\text{Pos}}t}$ . The numerical results for the two types of random initial states described above are presented in Fig. 3. As a comparison, the numerical result for initial thermal state  $\rho_0 = e^{-\beta H_0} / \text{Tr} e^{-\beta H_0}$  is also given. We can see that for initial states  $\rho$  of two cases, the LEs are close to the results in Eqs. (42) and (43), although the definitions of LEs and the forms of the quench Hamiltonian are different. For the initial thermal state  $\rho_0$ , the dynamics is not sensitive to the perturbation in Eq. (44).

#### V. SUMMARY

In summary, we have studied the consequence of the nonlocal perturbation term on the transverse field Ising chain. We proposed a pseudospin description for the Hamiltonian with perturbation term. In this description, the perturbation acts as an effective magnetic field, which lift the degenerate spectrum of the Hamiltonian in the ordered phase. We have shown that the string operator action can be realized via a quench process under the local perturbations. As an application, it is demonstrated that any ground states and excited states for the

Hamiltonian with perturbation can be selected for identifying the quantum phase, by adding the other perturbation to trigger a quench and measuring the LE. Our work, including the numerical results of LEs for a small-size system, provides a possible realization of the nonlocal operation as well as alternative quench protocol to detect the QPT. In addition, the result may shed light on the protocol of quantum information processing based

on nonlocal pseudospin as qubit.

## ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (under Grant No. 11874225).

- 
- [1] M. P. A. Fisher, Quantum phase transitions in disordered two-dimensional superconductors, *Phys. Rev. Lett.* **65**, 923 (1990).
- [2] D. Bitko, T. F. Rosenbaum, and G. Aeppli, Quantum critical behavior for a model magnet, *Phys. Rev. Lett.* **77**, 940 (1996).
- [3] M. Vojta, Y. Zhang, and S. Sachdev, Quantum phase transitions in  $d$ -wave superconductors, *Phys. Rev. Lett.* **85**, 4940 (2000).
- [4] Q. Si, S. Rabello, K. Ingersent, and J. L. Smith, Locally critical quantum phase transitions in strongly correlated metals, *Nature* **413**, 804 (2001).
- [5] D. Porras and J. I. Cirac, Effective quantum spin systems with trapped ions, *Phys. Rev. Lett.* **92**, 207901 (2004).
- [6] M. Uhlarz, C. Pfeiderer, and S. M. Hayden, Quantum phase transitions in the itinerant ferromagnet  $\text{ZrNi}_2$ , *Phys. Rev. Lett.* **93**, 256404 (2004).
- [7] H. M. Rønnow, R. Parthasarathy, J. Jensen, G. Aeppli, T. Rosenbaum, and D. McMorrow, Quantum phase transition of a magnet in a spin bath, *Science* **308**, 389 (2005).
- [8] R. Coldea, D. Tennant, E. Wheeler, E. Wawrzynska, D. Prabhakaran, M. Telling, K. Habicht, P. Smeibidl, and K. Kiefer, Quantum criticality in an ising chain: experimental evidence for emergent  $e_8$  symmetry, *Science* **327**, 177 (2010).
- [9] K. Kim, S. Korenblit, R. Islam, E. Edwards, M. Chang, C. Noh, H. Carmichael, G. Lin, L. Duan, C. J. Wang, *et al.*, Quantum simulation of the transverse ising model with trapped ions, *New J. Phys.* **13**, 105003 (2011).
- [10] J. Simon, W. S. Bakr, R. Ma, M. E. Tai, P. M. Preiss, and M. Greiner, Quantum simulation of antiferromagnetic spin chains in an optical lattice, *Nature* **472**, 307 (2011).
- [11] A. Trenkwalder, G. Spagnolli, G. Semeghini, S. Coop, M. Landini, P. Castilho, L. Pezze, G. Modugno, M. Inguscio, A. Smerzi, *et al.*, Quantum phase transitions with parity-symmetry breaking and hysteresis, *Nat. phys.* **12**, 826 (2016).
- [12] B. S. Rem, N. Käming, M. Tarnowski, L. Asteria, N. Fläschner, C. Becker, K. Sengstock, and C. Weitenberg, Identifying quantum phase transitions using artificial neural networks on experimental data, *Nat. Phys.* **15**, 917 (2019).
- [13] P. Pfeuty, The one-dimensional ising model with a transverse field, *Ann. Phys. (NY)* **57**, 79 (1970).
- [14] S. Sachdev, *Quantum phase transitions* (Cambridge university press, 2011).
- [15] A. Dutta, G. Aeppli, B. K. Chakrabarti, U. Divakaran, T. F. Rosenbaum, and D. Sen, *Quantum phase transitions in transverse field spin models: from statistical physics to quantum information* (Cambridge University Press, 2015).
- [16] A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, Colloquium: Nonequilibrium dynamics of closed interacting quantum systems, *Rev. Mod. Phys.* **83**, 863 (2011).
- [17] F. H. Essler and M. Fagotti, Quench dynamics and relaxation in isolated integrable quantum spin chains, *J. Stat. Mech.* **2016**, 064002 (2016).
- [18] N. O. Abeling and S. Kehrein, Quantum quench dynamics in the transverse field ising model at nonzero temperatures, *Phys. Rev. B* **93**, 104302 (2016).
- [19] P. Jurcevic, H. Shen, P. Hauke, C. Maier, T. Brydges, C. Hempel, B. P. Lanyon, M. Heyl, R. Blatt, and C. F. Roos, Direct observation of dynamical quantum phase transitions in an interacting many-body system, *Phys. Rev. Lett.* **119**, 080501 (2017).
- [20] E. Granet, M. Fagotti, and F. Essler, Finite temperature and quench dynamics in the transverse field ising model from form factor expansions, *SciPost Phys.* **9**, 033 (2020).
- [21] F. Andraschko and J. Sirker, Dynamical quantum phase transitions and the loschmidt echo: A transfer matrix approach, *Phys. Rev. B* **89**, 125120 (2014).
- [22] H. T. Quan, Z. Song, X. F. Liu, P. Zanardi, and C. P. Sun, Decay of loschmidt echo enhanced by quantum criticality, *Phys. Rev. Lett.* **96**, 140604 (2006).
- [23] M. Cozzini, P. Giorda, and P. Zanardi, Quantum phase transitions and quantum fidelity in free fermion graphs, *Phys. Rev. B* **75**, 014439 (2007).
- [24] M. Heyl, A. Polkovnikov, and S. Kehrein, Dynamical quantum phase transitions in the transverse-field ising model, *Phys. Rev. Lett.* **110**, 135704 (2013).
- [25] R. Jafari and H. Johannesson, Loschmidt echo revivals: Critical and noncritical, *Phys. Rev. Lett.* **118**, 015701 (2017).
- [26] B. Mera, C. Vlachou, N. Paunković, V. R. Vieira, and O. Viyuela, Dynamical phase transitions at finite temperature from fidelity and interferometric loschmidt echo induced metrics, *Phys. Rev. B* **97**, 094110 (2018).
- [27] A. Y. Kitaev, Unpaired majorana fermions in quantum wires, *Phys. Usp.* **44**, 131 (2001).
- [28] K. L. Zhang and Z. Song, Quantum phase transition in a quantum ising chain at nonzero temperatures, *Phys. Rev. Lett.* **126**, 116401 (2021).
- [29] K. L. Zhang and Z. Song, Ising chain with topological degeneracy induced by dissipation, *Phys. Rev. B* **101**, 245152 (2020).
- [30] A. Uhlmann, The “transition probability” in the state space of  $a^*$ -algebra, *Rep. Math. Phys.* **9**, 273 (1976).
- [31] N. T. Jacobson, L. C. Venuti, and P. Zanardi, Unitary equilibration after a quantum quench of a thermal state,

Phys. Rev. A **84**, 022115 (2011).