

L-spaces, taut foliations and the Whitehead link

Diego Santoro

Abstract

We prove that if M is a rational homology sphere that is a Dehn surgery on the Whitehead link, then M is not an L -space if and only if M supports a coorientable taut foliation. The left orderability of some of these manifolds is also proved, by determining which of the constructed taut foliations have vanishing Euler class.

We also present some more general results about the structure of the L -space surgery slopes for links whose components are unknotted and with pairwise linking number zero, and about the existence of taut foliations on the fillings of a k -holed torus bundle over the circle with some prescribed monodromy. Our results, combined with some results from [RSS03], also imply that all the rational homology spheres that arise as integer surgeries on the Whitehead link satisfy the L -space conjecture.

1 Introduction

In this paper we study the rational homology spheres obtained as surgery on the Whitehead link, motivated by the following conjecture.

Conjecture 1 (L-space conjecture). For an irreducible oriented rational homology 3-sphere M , the following are equivalent:

- 1) M supports a cooriented taut foliation;
- 2) M is not an L-space, i.e. its Heegaard Floer homology is not minimal;
- 3) M is left orderable, i.e. $\pi_1(M)$ is left orderable.

The equivalence between 1) and 2) was conjectured by Juhász in [Juh15], while the equivalence between 2) and 3) was conjectured by Boyer, Gordon and Watson in [BGW13].

Even if the properties involved in the conjecture are very different in flavour and nature it follows by the works of Ozsváth-Szábo, Bowden and Kazez-Roberts ([OS04], [Bow16], [KR17]) that 1) implies 2). Moreover it is now known that the conjecture holds for all the graph manifolds ([Ras17], [BC17], [HRRW20]).

It is therefore interesting to study the conjecture in the case of hyperbolic manifolds. In this direction, in [Zun20] the conjecture is proved for some manifolds obtained by considering mapping tori of pseudo-Anosov diffeomorphisms of closed surfaces and then by surgering on some collections of closed orbits.

In addition, in [Dun20], the conjecture is tested on a census of more than 300,000 hyperbolic rational homology spheres and proved for more than 60% of these manifolds.

One way of producing rational homology spheres is via Dehn surgery on knots or links in S^3 . When dealing with surgeries on knots, the different aspects of this conjecture have been studied separately in several papers. For example it has been proved that if a knot K admits a positive surgery that is an L-space then K is fibered, strongly quasipositive and the r -framed surgery along K is an L-space if and only if $r \geq 2g(K) - 1$, where $g(K)$ denotes the genus of the knot K ([OS05], [Ni07], [Hed10], [OS10]).

Taut foliations on surgeries on knots are constructed, for example, in [Rob01], [LR14], [DR19], [DR20], [Kri20] and it is possible to prove the left orderability of some of these manifolds by determining which of these foliations have vanishing Euler class, as done in [Hu19]. Another approach to study the left orderability of surgeries on knots is via representation theoretic methods, as presented in [CD18].

On the other hand, not much is known when it comes to the study of surgeries on links. Some results regarding integer L-space surgeries on links in S^3 are presented in [GN16], [GH17], [GN18], [Liu17] and [GLM20], while in [Ras20] rational L-space surgeries on satellites by algebraic links are studied. Concerning foliations, in [KR14] Kalelkar and Roberts construct coorientable taut foliations on some fillings of 3-manifolds that fiber over the circle. In particular, their methods can also be applied to surgeries on fibered links.

As far as we know, in this paper we provide the first example of the equivalence between conditions 1) and 2) of the conjecture for manifolds obtained via Dehn surgery on a hyperbolic link with at least 2 components.

We focus our attention on the Whitehead link, that is depicted in Figure 1.




Figure 1: The Whitehead link

We will denote the Whitehead link with WL and the $(p_1/q_1, p_2/q_2)$ -surgery on the Whitehead link with $S_{p_1/q_1, p_2/q_2}^3(\text{WL})$. Notice that since the Whitehead link has linking number zero, the homology of this manifold is isomorphic to $\mathbb{Z}_{p_1} \oplus \mathbb{Z}_{p_2}$. In particular, the manifold $S_{p_1/q_1, p_2/q_2}^3(\text{WL})$ is a rational homology sphere if and only if $p_1 \neq 0$ and $p_2 \neq 0$.

Recall that the Whitehead link exterior supports a complete hyperbolic structure and therefore, by virtue of the hyperbolic Dehn filling theorem, "most" of its fillings are hyperbolic.

The main result of this paper is the following:

Theorem 1.1. *Let p_1, q_1 and p_2, q_2 be two pairs of non vanishing coprime integers. Let $S_{p_1/q_1, p_2/q_2}^3(\text{WL})$ be the $(p_1/q_1, p_2/q_2)$ -surgery on the Whitehead link. Then*

- $S_{p_1/q_1, p_2/q_2}^3(\text{WL})$ is an L-space if and only if $p_1/q_1 \geq 1$ and $p_2/q_2 \geq 1$.
- $S_{p_1/q_1, p_2/q_2}^3(\text{WL})$ supports a cooriented taut foliation if and only if $p_1/q_1 < 1$ or $p_2/q_2 < 1$.

In particular, for all these manifolds we have 1) \Leftrightarrow 2) in Conjecture 1.

Notice that when some of the parameters p_1, p_2, q_1, q_2 vanish the only rational homology spheres that can be obtained are S^3 and lens spaces and it is known that they satisfy the L-space conjecture.

Also notice that the statement of Theorem 1.1 is invariant under the switch $p_1/q_1 \leftrightarrow p_2/q_2$. This is a consequence of the fact that the Whitehead link is symmetric, i.e. that there exists an isotopy exchanging its two components.

In the proof of the theorem we study the conditions of being an L-space and of supporting a taut foliation separately.

The key idea in the proof of the first part of the theorem is to use the results of J. Rasmussen and S.D. Rasmussen in [RR17] and Gorsky, Liu and Moore in [GLM20] to prove the following more general fact:

Theorem 1.2. *Suppose that \mathcal{L} is a non trivial link in S^3 with two unknotted components and linking number zero. Suppose that there exist rationals $p_1/q_1 > 0$ and $p_2/q_2 > 0$ such that $S_{p_1/q_1, p_2/q_2}^3(\mathcal{L})$ is an L-space. Then there exist non-negative integer numbers b_1, b_2 such that*

$$L(\mathcal{L}) = ([2b_1 + 1, \infty] \times [2b_2 + 1, \infty]) \cup (\{\infty\} \times \mathbb{Q}^*) \cup (\mathbb{Q}^* \times \{\infty\}).$$

In the previous statement the symbol $L(\mathcal{L})$ denotes the set of the *L-space filling slopes* of the exterior of \mathcal{L} . This is the set of slopes such that filling the exterior of \mathcal{L} with such slopes yields L-spaces.

Remark 1.3. We will not make use of this fact, but the integers b_1 and b_2 can be explicitly computed with the H function associated to \mathcal{L} , see [GLM20].

The foliations, on the other hand, are obtained by constructing branched surfaces without sink discs, using the results of Li ([Li02],[Li03]) and inspired by the works of Roberts and Kalelkar-Roberts ([Rob00],[Rob01],[KR14]). Also in this case, part of the proof of Theorem 1.1 will follow from a more general result, that allows to construct taut foliations on some fillings of manifolds that fiber over the circle with fiber a k -holed

torus and some particular type of monodromy. This is the content of Theorem 3.18, which seems to be interesting in itself.

We are also able to determine which of the taut foliations of Theorem 1.1 have zero Euler class, by adapting the ideas of Hu in [Hu19] to our case. This implies that the manifolds supporting such taut foliations have left orderable first fundamental group.

More precisely we have:

Theorem 1.4. *Let $S_{p_1/q_1, p_2/q_2}^3(\text{WL})$ be the $(p_1/q_1, p_2/q_2)$ -surgery on the Whitehead link, with $q_1, q_2 \neq 0$ and $p_1, p_2 > 0$.*

Then the foliations constructed in the proof of the Theorem 1.1 have vanishing Euler class if and only if $|q_i| \equiv 1 \pmod{p_i}$ for $i = 1, 2$.

In particular, for all these manifolds the L-space conjecture holds.

Moreover, Theorem 1.1, together with some results proved in [RSS03], implies the following

Theorem 1.5. *All the rational homology spheres obtained by integer surgery on WL satisfy the L-space conjecture.*

We refer to the last section of this paper for a more detailed statement of Theorem 1.5, which also combines results from [Zun20].

Structure of the paper. The paper is organised as follows. In Section 2 we recall the result of [RR17] and we prove Theorem 1.2 and the first part of Theorem 1.1. In Section 3 we focus our attention on taut foliations. In Subsection 3.1 we recall some basic notions on branched surfaces and the main result of [Li03]. In Subsection 3.2 we prove Theorem 3.18 and start the proof of the second part of Theorem 1.1, that is concluded in Subsection 3.3. In the last section we prove Theorem 1.4 and collect from [Dun20], [RSS03] and [Zun20] some other results about orderability, and non orderability, of some surgeries on the Whitehead link.

Acknowledgments. I warmly thank my advisors Bruno Martelli and Paolo Lisca for having presented this problem to me, for their support and for their useful comments on this paper. I also thank Alice Merz and Ludovico Battista for the several helpful discussions.

2 L-spaces

In this section we prove the first part of Theorem 1.1.

We start by recalling some definitions and the main result of [RR17]. Let Y be a rational homology solid torus, i.e. Y is a compact oriented 3-manifold with toroidal boundary such that $H_*(Y; \mathbb{Q}) \cong H_*(\mathbb{D}^2 \times S^1; \mathbb{Q})$.

We are interested in the study of the Dehn fillings on Y . We define the **set of slopes in Y** as

$$Sl(Y) = \{\alpha \in H_1(\partial Y; \mathbb{Z}) \mid \alpha \text{ is primitive}\} / \pm 1.$$

It is a well known fact that each element $[\alpha] \in Sl(Y)$ determines a Dehn filling on Y , that we will denote with $Y(\alpha)$.

Notice that since Y is a rational homology solid torus, there is a distinguished slope in $Sl(Y)$ that we call the *homological longitude* of Y and that is defined in the following way. We denote with $i : H_1(\partial Y; \mathbb{Z}) \rightarrow H_1(Y; \mathbb{Z})$ the map induced by the inclusion $\partial Y \subset Y$ and we consider a primitive element $l \in H_1(\partial Y; \mathbb{Z})$ such that $i(l)$ is torsion in $H_1(Y; \mathbb{Z})$. The element l is unique up to sign, and its equivalence class $[l] \in Sl(Y)$ is the homological longitude of Y . This definition, that may seem to be counterintuitive, is given so that when Y is the complement of a knot in S^3 , the homological longitude of Y coincides with the slope defined by the longitude the knot.

We want to study the fillings on Y that are L-spaces. For this reason we define the set of the **L-space filling slopes**:

$$L(Y) = \{[\alpha] \in Sl(Y) \mid Y(\alpha) \text{ is an L-space}\}$$

and we say that Y is **Floer simple** if Y admits multiple L-space filling slopes, i.e. if $|L(Y)| > 1$.

It turns out that if Y is Floer simple then the set $L(Y)$ has a simple structure, and this can be computed by knowing the **Turaev torsion** of Y . We only recall some properties of the Turaev torsion and we refer the reader to [Tur12] for the precise definitions.

Fix an identification $H_1(Y; \mathbb{Z}) = \mathbb{Z} \oplus T$, where T is the torsion subgroup, and denote by $\phi : H_1(Y; \mathbb{Z}) \rightarrow \mathbb{Z}$ the projection induced by this identification. Then the Turaev torsion of Y can be normalised to be written as a formal sum

$$\tau(Y) = \sum_{\substack{h \in H_1(Y; \mathbb{Z}) \\ \phi(h) \geq 0}} a_h h$$

where a_h is an integer for each h , $a_0 \neq 0$ and $a_h = 1$ for all but finitely many h . For example if $H_1(Y; \mathbb{Z}) = \mathbb{Z}$ the Turaev torsion of Y can be written as

$$\tau(Y) = \frac{\Delta(Y)}{1-t} \in \mathbb{Z}[[t]]$$

where $(1-t)^{-1}$ is expanded as an infinite sum in positive powers of t and $\Delta(Y)$ is the Alexander polynomial of Y normalised so that $\Delta(Y) \in \mathbb{Z}[t]$, $\Delta(Y)(0) \neq 0$ and $\Delta(Y)(1) = 1$. In fact, in this case the coefficients of $\tau(Y)$ are eventually constant and equal to the sum of all the coefficients of $\Delta(Y)$, and this value is exactly $\Delta(Y)(1) = 1$.

We define $S[\tau(Y)] = \{h \in H_1(Y; \mathbb{Z}) \mid a_h \neq 0\}$ to be the **support** of $\tau(Y)$. Notice that if $h \in S[\tau(Y)]$ then $\phi(h) \geq 0$ and that there exists a positive constant $c \in \mathbb{Z}$ such that if $h' \notin S[\tau(Y)]$ then $\phi(h') \leq c$.

We also define the following subset of $H_1(Y; \mathbb{Z})$:

$$D_{>0}^\tau(Y) = \{x - y \mid x \notin S[\tau(Y)], y \in S[\tau(Y)] \text{ and } \phi(x) > \phi(y)\} \cap i(H_1(\partial Y; \mathbb{Z}))$$

where $i : H_1(\partial Y; \mathbb{Z}) \rightarrow H_1(Y; \mathbb{Z})$ is induced from the inclusion.

Remark 2.1. The set $D_{>0}^\tau$ is always finite. In fact:

- if $x \notin S[\tau(Y)]$ is such that $\phi(x) \leq 0$ then there are no y in the support of $\tau(Y)$ that satisfy $\phi(y) < \phi(x)$;
- since the torsion subgroup of $H_1(Y; \mathbb{Z})$ is finite there are only finitely many $x \notin S[\tau(Y)]$ such that $\phi(x) > 0$ and for each such x there are only finitely many y in the support of $\tau(Y)$ such that $\phi(y) < \phi(x)$.

We are now ready to state the main result of [RR17] :

Theorem 2.2. ([RR17]) *If Y is Floer simple, then either*

- $D_{>0}^\tau(Y) = \emptyset$ and $L(Y) = Sl(Y) \setminus [l]$, or
- $D_{>0}^\tau(Y) \neq \emptyset$ and $L(Y)$ is a closed interval whose endpoints are consecutive elements in $i^{-1}(D_{>0}^\tau(Y))$.

We explain more precisely the second part of the statement of this theorem. Once we fix a basis (μ, λ) for $H_1(\partial Y; \mathbb{Z})$ we can associate to each element $a\mu + b\lambda \in H_1(\partial Y; \mathbb{Z})$ the element $a/b \in \overline{\mathbb{Q}} = \mathbb{Q} \cup \{\infty\} \subset S^1$. This association defines a map onto $\overline{\mathbb{Q}}$ that yields an identification between $Sl(Y)$ and $\overline{\mathbb{Q}}$.

If the set $D_{>0}^\tau$ is not empty, then we can apply this map to the set $i^{-1}(D_{>0}^\tau) \subset H_1(\partial Y; \mathbb{Z})$ and Theorem 2.2 states that if Y is Floer simple then $L(Y)$ is a closed interval in $Sl(Y) = \overline{\mathbb{Q}}$ whose endpoints are consecutive elements in the image of $i^{-1}(D_{>0}^\tau)$ in $\overline{\mathbb{Q}}$.

In the case of our interest we consider a link $\mathcal{L} = K_1 \sqcup K_2 \subset S^3$ with the following properties:

- \mathcal{L} has two unknotted components;
- \mathcal{L} has linking number zero.

By analogy with the definitions given for rational homology solid tori we denote with $Sl(\mathcal{L})$ the set of slopes of the exterior of \mathcal{L} and with $L(\mathcal{L})$ the set of L-space filling slopes of the exterior of \mathcal{L} .

We fix an orientation of the components of \mathcal{L} and in this way we obtain canonical meridian-longitude bases $(\mu_i, \lambda_i)_{i=1,2}$ of the first homology groups of the boundary tori of its exterior. The choice of these bases also determines an identification $Sl(\mathcal{L}) = \overline{\mathbb{Q}} \times \overline{\mathbb{Q}}$.

We denote with $S_{p_1/q_1, \bullet}^3(\mathcal{L})$ the manifold obtained by drilling the second component of \mathcal{L} and by performing (p_1/q_1) -surgery on the first. Analogously we denote with $S_{\bullet, p_2/q_2}^3(\mathcal{L})$ the manifold obtained by drilling the first component of \mathcal{L} and by performing (p_2/q_2) -surgery on the second. When it will not be important which component we are drilling and on which component we are surgering, we will simply use the symbol $S_{p/q}^3(\mathcal{L})$.

Notice that since \mathcal{L} has linking number zero, we have an isomorphism

$$H_1\left(S_{p_1/q_1, \bullet}^3(\mathcal{L}); \mathbb{Z}\right) \cong \mathbb{Z}_{p_1} \oplus \mathbb{Z}$$

where the image of the meridian μ_1 in $H_1\left(S_{p_1/q_1, \bullet}^3(\mathcal{L}); \mathbb{Z}\right)$ is mapped to $(1, 0)$ and the image of the meridian μ_2 in $H_1\left(S_{p_1/q_1, \bullet}^3(\mathcal{L}); \mathbb{Z}\right)$ is mapped to $(0, 1)$. An analogous result holds for $S_{\bullet, p_2/q_2}^3(\mathcal{L})$.

Lemma 2.3. *Fix $p \neq 0$ and q coprime integers. Suppose that $S_{p/q}^3(\mathcal{L})$ is Floer simple. Then the set $L\left(S_{p/q}^3(\mathcal{L})\right)$ has one of the following forms:*

- $L\left(S_{p/q}^3(\mathcal{L})\right) = \overline{\mathbb{Q}} \setminus \{0\}$, or
- there exists a natural number $k > 0$ such that either $L\left(S_{p/q}^3(\mathcal{L})\right) = [k, \infty]$ or $L\left(S_{p/q}^3(\mathcal{L})\right) = [\infty, -k]$.

Proof. We suppose that $S_{p/q}^3(\mathcal{L}) = S_{\bullet, p/q}^3(\mathcal{L})$, the case $S_{p/q}^3(\mathcal{L}) = S_{p/q, \bullet}^3(\mathcal{L})$ being analogous. We denote $S_{p/q}^3(\mathcal{L})$ with M .

The lemma follows from Theorem 2.2 together with a simple inspection on the possible forms of the set $D_{>0}^\tau(M)$:

- $D_{>0}^\tau(M)$ is empty: in this case we have that $L(M) = \overline{\mathbb{Q}} \setminus \{0\}$.
- $D_{>0}^\tau(M)$ is not empty: the projection ϕ associated to the identification

$$H_1(M; \mathbb{Z}) = \mathbb{Z} \oplus \mathbb{Z}_p$$

is simply the map $\phi(x, y) = x$. Since $i_*(H_1(\partial M; \mathbb{Z})) = \mathbb{Z} \times \{0\}$ we have that $D_{>0}^\tau(M)$ is by definition a subset of $\mathbb{Z}_{>0} \times \{0\}$, and we denote with $S = \{n_1, \dots, n_h\} \subset \mathbb{Z}_{>0}$ the first coordinates of the elements in $D_{>0}^\tau(M)$, listed in ascending order. Recall from Remark 2.1 that $D_{>0}^\tau(M)$ is always a finite set.

We have that

$$i^{-1}(D_{>0}^\tau(M)) = \{(n_i, m) \in \mathbb{Z} \times \mathbb{Z} \mid n_i \in S \text{ and } m \in \mathbb{Z}\}$$

and we know by Theorem 2.2 that $L(M)$ is a closed interval in $\overline{\mathbb{Q}}$ whose endpoints are consecutive elements in the set $\{n_i/m \mid n_i \in S \text{ and } m \in \mathbb{Z}\}$. Since the components of \mathcal{L} are unknotted we know that ∞ belongs to $L(M)$, and therefore we can conclude that either $L(M) = [n_h, \infty]$ or $L(M) = [\infty, -n_h]$.

This concludes the proof. \square

As we already anticipated, the first part of Theorem 1.1 will be a corollary of the following, more general, result:

Theorem 1.2. *Suppose that \mathcal{L} is a non trivial link with two unknotted components and linking number zero. Suppose that there exist rationals $p_1/q_1 > 0$ and $p_2/q_2 > 0$ such that $S_{p_1/q_1, p_2/q_2}^3(\mathcal{L})$ is an L-space. Then there exist non-negative integer numbers b_1, b_2 such that*

$$L(\mathcal{L}) = ([2b_1 + 1, \infty] \times [2b_2 + 1, \infty]) \cup (\{\infty\} \times \mathbb{Q}^*) \cup (\mathbb{Q}^* \times \{\infty\}).$$

As a corollary of Theorem 1.2 we have:

Corollary 2.4 (First part of Theorem 1.1). *The 3-manifold $S_{p_1/q_1, p_2/q_2}^3(\text{WL})$ is an L-space if and only if $p_1/q_1 \geq 1$ and $p_2/q_2 \geq 1$.*

Proof. The $(1, 1)$ -surgery on the Whitehead link is the Poincaré homology sphere, that has finite fundamental group and is therefore an L-space. In other words $(1, 1)$ belongs to $L(\text{WL})$. The Whitehead link also has unknotted components and linking number zero and we can therefore apply Theorem 1.2, that immediately implies the thesis. \square

To prove Theorem 1.2 we recall the definition of L-space link, as given by Gorsky and Némethi in [GN16]. We give the definition for a 2-components link, but it is obviously generalisable to links with more components.

Definition 2.5. A link $\mathcal{L} \subset S^3$ is an **L-space link** if all sufficiently large integer surgeries are L-spaces, i.e. if there exist integers p_1, p_2 such that $S_{d_1, d_2}^3(\mathcal{L})$ is an L-space for all integers $d_1 > p_1$ and $d_2 > p_2$.

For knots, the existence of a positive rational L-space surgery implies the existence of arbitrarily large L-space surgeries, but this fails in the case of links, as the Example 2.4 in [Liu17] shows.

The following lemma shows that such generalisation holds if \mathcal{L} has unknotted components and linking number zero. We will use the symbols $\lfloor x \rfloor$ and $\lceil x \rceil$, where x is a rational number, to denote the integers

$$\lfloor x \rfloor = \max\{k \in \mathbb{Z} | k \leq x\}$$

$$\lceil x \rceil = \min\{k \in \mathbb{Z} | k \geq x\}.$$

Lemma 2.6. *Let \mathcal{L} be a n -components link whose components are unknotted and have pairwise linking number zero. Suppose that there exist rationals $r_1 > 0, r_2 > 0, \dots, r_n > 0$ such that $S_{r_1, \dots, r_n}^3(\mathcal{L})$ is an L-space. Then $S_{s_1, \dots, s_n}^3(\mathcal{L})$ is an L-space for all (s_1, \dots, s_n) satisfying*

$$\begin{cases} s_i \geq \lfloor r_i \rfloor & \text{if } r_i \geq 1 \\ s_i > \lfloor r_i \rfloor = 0 & \text{if } 0 < r_i < 1. \end{cases}$$

In particular \mathcal{L} is an L-space link.

Proof. We suppose that \mathcal{L} has two components, the proof being analogous in the general case. We have the following cases:

- $r_1 \geq 1$ and $r_2 \geq 1$: we start by considering the set $L(S_{r_1, \bullet}^3(\mathcal{L}))$. We know by hypothesis that this set contains r_2 and since r_2 is positive it follows from Lemma 2.3 that $L(S_{r_1, \bullet}^3(\mathcal{L}))$ must be either $\overline{\mathbb{Q}} \setminus \{0\}$ or $[k, \infty]$ for some positive natural number k . In both of these cases, since $r_2 \geq 1$ we can deduce that

$$[\lfloor r_2 \rfloor, \infty] \subset L(S_{r_1, \bullet}^3(\mathcal{L})).$$

We now consider the set $L(S_{\bullet, \lfloor r_2 \rfloor}^3(\mathcal{L}))$. We have just proved that it contains r_1 and with the same argument as before we can deduce that

$$[\lfloor r_1 \rfloor, \infty] \subset L(S_{\bullet, \lfloor r_2 \rfloor}^3(\mathcal{L})).$$

By applying the same reasoning it follows that for every $s_1 \geq \lfloor r_1 \rfloor$ we have that

$$[\lfloor r_2 \rfloor, \infty] \subset L(S_{s_1, \bullet}^3(\mathcal{L}))$$

and this is exactly what we wanted. A pictorial sketch of the proof is showed in Figure 2.

- $r_1 \geq 1$ and $r_2 < 1$: since $0 < r_2 < 1$ we have $L(S_{r_1, \bullet}^3(\mathcal{L})) = \overline{\mathbb{Q}} \setminus \{0\}$ and in particular

$$(0, \infty] = (\lfloor r_2 \rfloor, \infty] \subset L(S_{r_1, \bullet}^3(\mathcal{L})).$$

This implies that for any $s_2 > 0$ we have that $S_{r_1, s_2}^3(\mathcal{L})$ is an L-space and therefore by applying again Lemma 2.3, since $r_1 \geq 1$, we have that

$$[\lfloor r_1 \rfloor, \infty] \subset L(S_{\bullet, s_2}^3(\mathcal{L}))$$

and this is exactly what we wanted.

- $r_1 < 1$ and $r_2 \geq 1$: this case is completely analogous to the previous one.
- $r_1 < 1$ and $r_2 < 1$: also in this case we have that

$$(0, \infty] = (\lfloor r_2 \rfloor, \infty] \subset L(S_{r_1, \bullet}^3(\mathcal{L})) = \overline{\mathbb{Q}} \setminus \{0\}.$$

As a consequence, for any $s_2 > 0$ we have that $S_{r_1, s_2}^3(\mathcal{L})$ is an L-space and therefore by applying again Lemma 2.3, since $0 < r_1 < 1$, we have that

$$(0, \infty] = (\lfloor r_1 \rfloor, \infty] \subset L(S_{\bullet, s_2}^3(\mathcal{L})) = \overline{\mathbb{Q}} \setminus \{0\}.$$

This concludes the proof. □




Figure 2

Remark 2.7. It follows from Theorem 1.2 that if \mathcal{L} has two components, then in the previous lemma the case $0 < r_1 < 1$ or $0 < r_2 < 1$ cannot occur.

Before proving Theorem 1.2 we recall the following theorem from [GLM20].

Theorem 2.8. ([GLM20]) *Assume that \mathcal{L} is a non trivial L-space link with unknotted components and linking number zero. Then there exist non-negative integers b_1, b_2 such that for $p_1, p_2 \in \mathbb{Z}$ we have that $S_{p_1, p_2}^3(\mathcal{L})$ is an L-space if and only if $p_1 > 2b_1$ and $p_2 > 2b_2$.*

Proof of Theorem 1.2. We know from Lemma 2.6 that \mathcal{L} is an L-space link. Therefore we can apply Theorem 2.8 and deduce that there exist non-negative integers b_1, b_2 such that

$$L(\mathcal{L}) \cap \mathbb{Z}^2 = [2b_1 + 1, \infty) \times [2b_2 + 1, \infty) \cap \mathbb{Z}^2$$

Exactly as in the proof of Lemma 2.6, we can use Lemma 2.3 to deduce that

$$L(\mathcal{L}) \supset ([2b_1 + 1, \infty) \times [2b_2 + 1, \infty]) \cup (\{\infty\} \times \mathbb{Q}^*) \cup (\mathbb{Q}^* \times \{\infty\}).$$

and therefore we only have to prove that this inclusion is an equality.

Suppose on the contrary that there exists an L-space surgery slope (r_1, r_2) , with r_1, r_2 rationals, such that

$$(r_1, r_2) \notin ([2b_1 + 1, \infty) \times [2b_2 + 1, \infty]) \cup (\{\infty\} \times \mathbb{Q}^*) \cup (\mathbb{Q}^* \times \{\infty\}).$$

We suppose that $r_1 < 2b_1 + 1$. The case $r_2 < 2b_2 + 1$ can be solved in the same way. We have the following cases:

- $1 \leq r_1 < 2b_1 + 1$.

By virtue of Lemma 2.3 we have that $[[r_1], \infty]$ is contained in $L(S_{\bullet, r_2}^3)$. This implies that $S_{[r_1], \bullet}^3$ is Floer simple and therefore, by applying again Lemma 2.3, we deduce that it admits integral L-space filling slopes. In this way we produce a point in

$$(L(\mathcal{L}) \cap \mathbb{Z}^2) \setminus \left(([2b_1 + 1, \infty) \times [2b_2 + 1, \infty)) \cap \mathbb{Z}^2 \right)$$

contradicting Theorem 2.8.

- $r_1 \in (-1, 1)$.

As a consequence of Lemma 2.3 we have that $L(S_{\bullet, r_2}^3) = \overline{\mathbb{Q}} \setminus \{0\}$. Therefore if we fix any negative integer $-m < 0$ we have that $r_2 \in L(S_{-m, \bullet}^3)$ and by applying again Lemma 2.3 we deduce that there exist integral L-space filling slopes on $S_{k, \bullet}^3$, contradicting Theorem 2.8.

- $r_1 \leq -1$.

By applying the same argument used in the first case we have that $L(S_{\bullet, r_2}^3)$ contains $[\infty, [r_1]]$. Therefore $S_{[r_1], \bullet}^3$ admits integral L-space filling slopes, contradicting Theorem 2.8.

The proof is complete. □

3 Coorientable Taut Foliations

In this section we study the existence of taut foliations on the Dehn fillings on the Whitehead link exterior. The main theorem in this section is the following:

Theorem 3.1. *Let p_1, q_1 and p_2, q_2 be two pairs of non vanishing coprime integers. Let $S_{p_1/q_1, p_2/q_2}^3(\text{WL})$ be the $(p_1/q_1, p_2/q_2)$ -surgery on the Whitehead link. Then $S_{p_1/q_1, p_2/q_2}^3(\text{WL})$ supports a cooriented taut foliation if and only if $p_1/q_1 < 1$ or $p_2/q_2 < 1$.*

In this paper the term *foliation* will refer to codimension-1 foliations of class $C^{\infty, 0}$, as defined for example in [CC00] and [KR17]. We recall the definition here. We denote with \mathbb{H}^k the k -dimensional Euclidean closed half space

$$\mathbb{H}^k = \{(x_1, \dots, x_k) \in \mathbb{R}^k \mid x_k \geq 0\}.$$

Definition 3.2. A $C^{\infty, 0}$ **codimension-1 foliation** \mathcal{F} of a smooth 3-manifold M with (possibly empty) boundary is a decomposition of M into the union of disjoint smoothly injectively immersed surfaces, called the **leaves** of \mathcal{F} , together with a collection of charts $(U_i, \phi_i)_{i \in \mathcal{I}}$ covering M such that:

- $\phi_i : U_i \rightarrow \mathcal{X}$ is a homeomorphism, where \mathcal{X} is either $\mathbb{R}^2 \times \mathbb{R}$ or $\mathbb{R}^2 \times \mathbb{H}^1$ or $\mathbb{H}^2 \times \mathbb{R}$, with the property that the image of each component of a leaf intersected with U_i is a slice $\mathbb{R}^2 \times \{\text{point}\}$ or $\mathbb{H}^2 \times \{\text{point}\}$;

- all partial derivatives of any order in the variables x and y on the domain of each transition function $\phi_j\phi_i^{-1}$ are continuous; here we have fixed coordinates (x, y, z) on \mathcal{X} .

The three local models for a foliation are depicted in Figure 3, where $\partial\mathcal{X}$ is shaded.




Figure 3

Remark 3.3. The tangent planes to the leaves of a foliation \mathcal{F} of a 3-manifold M define a continuous plane subbundle of TM , that we denote with $T\mathcal{F}$.

Definition 3.4. A foliation \mathcal{F} of a 3-manifold M is **orientable** if the plane bundle $T\mathcal{F}$ is orientable and is **coorientable** if the line bundle $TM/T\mathcal{F}$ is orientable.

Definition 3.5. A foliation \mathcal{F} of a 3-manifold M is **taut** if every leaf of \mathcal{F} intersects a closed transversal, i.e. a smooth simple closed curve in M that is transverse to \mathcal{F} .

There are several definitions of tautness and in general they are not equivalent. For details we refer to [CKR19], where also the relations among these different notions are discussed.

We recall that in order to prove Theorem 3.1 it is enough to prove the “if” part, since L-spaces do not support taut foliations (see [OS04], [Bow16], [KR17]), and we have already proved in the previous section that if $p_i/q_i \geq 1$ for $i = 1, 2$, then $S_{p_1/q_1, p_2/q_2}^3(\text{WL})$ is an L-space.

This section is organised as follows. In Section 3.1 we recall the required background regarding branched surfaces and we state the theorem of [Li03]. In Section 3.2 we prove Theorem 3.18, regarding the existence of taut foliations on Dehn fillings on manifolds that fiber over the circle with fiber a k -holed torus and with some prescribed monodromy. This theorem will be useful to prove that many of the manifolds of Theorem 3.1 support taut foliations. In Section 3.3 we conclude the proof of Theorem 3.1.

3.1 Background

In this and in the next sections we will assume familiarity with the basic notions of the theory of train tracks; see [PH16] for reference. We only point out that in the cases of our interest, train tracks can also have bigons as complementary regions.

We now recall some basic facts about branched surfaces. We refer to [FO84] and [Oer84] for more details.

Definition 3.6. A **branched surface with boundary** in a 3-manifold M is a closed subset $B \subset M$ that is locally diffeomorphic to one of the models in \mathbb{R}^3 of Figure 4a) or to one of the models in the closed half space of Figure 4b), where $\partial B := B \cap \partial M$ is represented with a bolded line:




Figure 4

Branched surfaces generalise the concept of train tracks from surfaces to 3-manifolds and when the boundary of B is non empty it defines a train track ∂B in ∂M .

If B is a branched surface it is possible to identify two subsets of B : the **branch locus** and the set of **triple points** of B . The branch locus is defined as the set of points where B is not locally homeomorphic to a surface. It is self-transverse and intersects itself in double points only. The set of triple points of B can be defined as the points where the branch locus is not locally homeomorphic to an arc. For example, the rightmost model of Figure 4a) contains a triple point.

The complement of the branch locus in B is a union of connected surfaces. The abstract closures of these surfaces under any path metric on M are called the **branch sectors** of B . Analogously the complement of the set of the triple points inside the branch locus is a union of 1-dimensional connected manifolds. Moreover to each of these manifolds we can associate an arrow in B pointing in the direction of the smoothing, as we did in Figure 5. We call these arrows **branch directions**, or also **cuspid directions**.



Figure 5

If B is a branched surface in M , we denote with N_B a fibered regular neighbourhood of B constructed as suggested in Figure 6.




Figure 6

The boundary of N_B decomposes naturally into the union of three compact subsurfaces $\partial_h N_B$, $\partial_v N_B$ and $N_B \cap \partial M$. We call $\partial_h N_B$ the **horizontal boundary** of N_B and $\partial_v N_B$ the **vertical boundary** of B . The horizontal boundary is transverse to the interval fibers of N_B while the vertical boundary intersects, if at all, the fibers of N_B in one or two proper closed subintervals contained in their interior. If we collapse each interval fiber of N_B to a point, we obtain a branched surface in M that is isotopic to B , and the image of $\partial_v N_B$ coincides with the branch locus of such a branched surface.

We also recall the definition of *splitting*¹.

Definition 3.7. Given two branched surfaces B_1 and B_2 in M we say that B_2 is obtained by **splitting** B_1 if N_{B_1} can be obtained as $N_{B_2} \cup J$, where J is a $[0, 1]$ -bundle such that $\partial_h J \subset \partial_h N_{B_2}$, $\partial_v J \cap \partial N_{B_2} \subset \partial_v N_{B_2}$ and ∂J meets ∂N_{B_2} so that the fibers agree.

Figure 7 shows two examples of splittings, depicted for the case of 1-branched manifolds, i.e. train tracks. The coloured region is the interval bundle J .




Figure 7

¹This operation is referred to as *restriction* in [Oer84]

Branched surfaces are useful to define partial foliations on 3-manifolds. To be more precise, the right notion of partial foliation is the following:


Definition 3.8. Let B be a branched surface in a 3-manifold M . A **lamination carried by B** is a closed subset Λ of some regular neighbourhood N_B of B such that Λ is a disjoint union of smoothly injectively immersed surfaces, called leaves, that intersect the fibers of N_B transversely. We say that Λ is **fully carried** by B if Λ is carried by B and intersects every fiber of N_B .

Remark 3.9. Analogously as in Definition 3.8, if S is a closed oriented surface and τ is a train track in S we can define what is a lamination (fully) carried by τ . In this case we say that an oriented simple closed curve γ is **realised** by τ if τ fully carries a union of finitely many disjoint curves that are parallel to γ inside S .


In [Li02], Li introduces the notion of *sink disc*.

Definition 3.10. Let B be a branched surface in M and let S be a branch sector in B . We say that S is a **sink disc** if S is a disc, $S \cap \partial M = \emptyset$ and the branch direction of any smooth curve or arc in its boundary points into S . We say that S is a **half sink disc** if S is a disc, $S \cap \partial M \neq \emptyset$ and the branch direction of any smooth arc in $\partial S \setminus \partial M$ points into S .

In Figure 8 some examples of sink discs and half sink discs are depicted. The bolded lines represent the intersection of the branched surface with ∂M . Notice that if S is a half sink disc the intersection $\partial S \cap \partial M$ can also be disconnected.



a) Examples of sink discs.



b) Examples of half sink discs.

Figure 8

If B contains a sink disc or a half sink disc there is a very simple way to eliminate it, namely it is enough to blow an air bubble in its interior, as in the following figure.




Figure 9: How to eliminate a sink disc by blowing an air bubble.

We want to avoid this situation. We say that a connected component of $M \setminus \text{int}(N_B)$ is a $\mathbb{D}^2 \times [0, 1]$ **region** if it is homeomorphic to a ball and its boundary can be subdivided into an annular region, corresponding to a component of $\partial_v N_B$, and two \mathbb{D}^2 regions corresponding to components of $\partial_h N_B$. We say that a $\mathbb{D}^2 \times [0, 1]$ region is **trivial** if the map collapsing the fibers of N_B is injective on $\text{int}(\mathbb{D}^2) \times \{0, 1\}$. In this case the image of $\mathbb{D}^2 \times \{0, 1\}$ via the collapsing map is called a **trivial bubble** in B . Trivial bubbles and trivial $\mathbb{D}^2 \times [0, 1]$ regions occur when we eliminate sink discs as in Figure 9.

When M and B have boundary these definitions generalise straightforwardly to the relative case, see [Li03].

In [Li02], Li introduces the definition of laminar branched surface and shows that laminar branched surfaces fully carry essential laminations². In [Li03] he generalises this definition to branched surfaces with boundary as follows:

Definition 3.11. Let B be a branched surface in a 3-manifold M . We say that B is **laminar** if B has no trivial bubbles and the following hold:

1. $\partial_h N_B$ is incompressible and ∂ -incompressible in $M \setminus \text{int}(N_B)$, and no component of $\partial_h N_B$ is a sphere or a properly embedded disc in M ;
2. there is no monogon in $M \setminus \text{int}(N_B)$, i.e. no disc $D \subset M \setminus \text{int}(N_B)$ such that $\partial D = D \cap N_B = \alpha \cup \beta$, where α is in an interval fiber of $\partial_v N_B$ and β is an arc in $\partial_h N_B$;
3. $M \setminus \text{int}(N_B)$ is irreducible and $\partial M \setminus \text{int}(N_B)$ is incompressible in $M \setminus \text{int}(N_B)$;
4. B contains no Reeb branched surfaces (see [GO89] for the definition);
5. B has no sink discs or half sink discs.

Since $\partial_h N_B$ is not properly embedded in $M \setminus \text{int}(N_B)$ we explain more precisely the request of ∂ -incompressibility in 1. : we require that if D is a disc in $M \setminus \text{int}(N_B)$ with $\text{int}(D) \subset M \setminus N_B$ and $\partial D = \alpha \cup \beta$ where α is an arc in $\partial_h N_B$ and β is an arc in ∂M , then there is a disc $D' \subset \partial_h N_B$ with $\partial D' = \alpha \cup \beta'$ where $\beta' = \partial D' \cap \partial M$.

The following theorem of [Li03] will be used profusely in this section.

²For the definition of essential lamination see [GO89], but we will not need their properties for our purposes.

Theorem 3.12. [Li03] *Let M be an irreducible and orientable 3-manifold whose boundary is union of k incompressible tori T_1, \dots, T_k . Suppose that B is a laminar branched surface in M such that $\partial M \setminus \partial B$ is a union of bigons. Then for any multislope $(s_1, \dots, s_k) \in \overline{\mathbb{Q}}^k$ that is realised by the train track ∂B , if B does not carry a torus that bounds a solid torus in $M(s_1, \dots, s_k)$, there exists an essential lamination Λ in M fully carried by B that intersects ∂M in parallel simple curves of multislope (s_1, \dots, s_k) . Moreover this lamination extends to an essential lamination of the filled manifold $M(s_1, \dots, s_k)$.*

Remark 3.13. The statement of the Theorem 3.12 is slightly more detailed than the version of [Li03]. The details we have added come from the proof of Theorem 3.12. In fact the idea of the proof is to split the branched surface B in a neighbourhood of ∂M so that it intersects T_i in parallel simple closed curves of slopes s_i , for $i = 1, \dots, k$. In this way, when gluing the solid tori, we can glue meridional discs of these tori to B so to obtain a branched surface $B(s_1, \dots, s_k)$ in $M(s_1, \dots, s_k)$ that is laminar and that therefore by virtue of [Li02, Theorem 1] fully carries an essential lamination. In particular, this essential lamination is obtained by gluing the meridional discs of the solid tori to an essential lamination in M that intersects T_i in parallel simple closed curves of slopes s_i , for $i = 1, \dots, k$.

Remark 3.14. In [Li03] the statement of the theorem is given for M with connected boundary, but as already noticed in [KR14] if M has multiple boundary components we can split B in a neighbourhood of each boundary tori T_i and the same proof of [Li03] works.

3.2 Constructing taut foliations

The aim of this subsection is to prove Theorem 3.18, that concerns the existence of taut foliations on Dehn fillings on manifolds that fiber over the circle with fiber a k -holed torus and with some prescribed monodromy. To do this we will recall a very simple, yet useful, way to build branched surfaces. First of all we fix some notations and recall the definition of fibered link.

Given an oriented surface S with (possibly empty) boundary and $h : S \rightarrow S$ an orientation preserving homeomorphism of S fixing ∂S pointwise we denote with M_h the mapping torus of h

$$M_h = \frac{S \times [0, 1]}{(h(x), 0) \sim (x, 1)}.$$

We orient $S \times [0, 1]$ as a product and we orient M_h with the orientation induced by $S \times [0, 1]$. We also identify S with its image in M_h via the map

$$\begin{aligned} S &\rightarrow S \times \{0\} \subset M_h \\ x &\mapsto (x, 0). \end{aligned}$$

The homeomorphism h is called the monodromy of M_h .

Definition 3.15. Let L be an oriented link in S^3 . We say that L is **fibered** if there exist a Seifert surface S for L , an orientation preserving homeomorphism h of S fixing ∂S pointwise and an orientation preserving homeomorphism

$$\chi : S^3 \setminus \text{int}(N_L) \rightarrow M_h,$$

where N_L denotes a tubular neighbourhood of L in S^3 , so that

- $\chi|_S$ is the inclusion $S \subset M_h$;
- $\chi(m_i) = \{x_i\} \times [0, 1]$, where m_i is a meridian for the i -th component of L and $x_i \in \partial S$ is a point.

We want to apply Theorem 3.12 to construct laminations on $S^3_{p_1/q_1, p_2/q_2}$ (WL) when $p_i/q_i < 1$ for at least one $i \in \{1, 2\}$ and then promote them to taut foliations. To do this we will define some branched surfaces in the exterior of the Whitehead link and then study their boundary train tracks.

The construction of these branched surfaces relies on the fact that the Whitehead link is a fibered link. In fact, as Figure 10 shows, the Whitehead link can be obtained as the boundary of a surface F that is a torus with two open discs removed. This torus is obtained by a sequence of three Hopf plumbings and this implies by standard results (see [Gab86], [Sta78]) that F is a fiber surface for WL with monodromy h given by $h = \tau_0 \tau_1 \tau_2^{-1}$, where τ_i denotes the positive Dehn twist along the curve γ_i .




Figure 10: The Whitehead link is the union of the red curve and the blue curve. The curves $\gamma_0, \gamma_1, \gamma_2$ lie on the Seifert surface that should be evident from the picture.

Before focusing on the specific example of the Whitehead link exterior, we work in a more general setting.

Let S be an oriented surface with boundary and let h be an orientation preserving homeomorphism of S fixing ∂S pointwise. We consider some pairwise disjoint properly embedded arcs $\alpha_1, \dots, \alpha_k$ in S and the discs $\overline{D}_i = \alpha_i \times [0, 1] \subset S \times [0, 1]$. Each of these discs has a “bottom” boundary, $\alpha_i \times \{0\}$, and a “top” boundary, $\alpha_i \times \{1\}$. When we project these discs to the mapping torus M_h , the bottom and top boundaries become respectively $\cup_i \alpha_i \subset S$ and $\cup_i h(\alpha_i) \subset S$.

We can isotope simultaneously the discs \overline{D}_i 's in a neighbourhood of $S \times \{1\} \subset S \times [0, 1]$ so that when projected to M_h their top boundaries define a family of arcs $\{\widehat{h(\alpha_i)}\}_{i=1, \dots, k}$ in S such that for each $i, j \in \{1, \dots, k\}$ the intersection between α_i and $\widehat{h(\alpha_j)}$ is transverse and minimal. Notice that each arc $\widehat{h(\alpha_i)}$ is isotopic as a properly embedded arc to $h(\alpha_i)$. We also denote with D_i the projected perturbed disc contained in M_h .

If we assign a (co)orientation to these discs, since S is (co)oriented, we can smoothen $S \cup D_1 \cup \dots \cup D_k$ to a branched surface B by imposing that the smoothing preserves the coorientation of S and of the discs. This operation is suggested Figure 11, where S is a torus with two open discs removed.




Figure 11

We prove the following lemma, that is only implicit in [KR14].

Lemma 3.16. *Let S be a connected and oriented surface with boundary and let h be an orientation preserving homeomorphism of S fixing ∂S pointwise. Let $\{\alpha_i\}_{i=1, \dots, k} \subset S$ be pairwise disjoint properly embedded arcs in S and suppose that $S \setminus \cup_{i=1}^k \alpha_i$ has no disc components. Denote with D_i 's the discs in M_h associated to the arcs α_i 's in the way described above and fix a coorientation for these discs. Let $B = S \cup D_1 \cup \dots \cup D_k$ denote the branched surface in M_h obtained by smoothing according to these coorientations. Then B has no trivial bubbles and satisfies conditions 1, 2, 3 and 4 of Definition 3.11.*

Proof. We denote the mapping torus M_h with M . We fix for each arc α_i a tubular neighbourhood N_{α_i} in S and we denote with S' the surface $S \setminus \cup_{i=1}^k \text{int}(N_{\alpha_i})$. The first observation is that by construction we have

$$M \setminus \text{int}(N_B) \cong S' \times [0, 1]$$

with a homeomorphism that identifies

$$\partial_h N_B = S' \times \{0, 1\}.$$

and

$$\partial_v N_B = \partial' S' \times [0, 1]$$

where $\partial' S'$ denotes the closure of $\partial S' \setminus \partial M$.

Basically, the proof follows from the fact that $M \setminus \text{int}(N_B)$ is homeomorphic to $S' \times [0, 1]$ and that S' has no discs components.

First of all, we notice that since by hypothesis $S \setminus \cup_{i=1}^k \alpha_i$ has no discs components, there are no $\mathbb{D}^2 \times [0, 1]$ regions in $M \setminus \text{int}(N_B)$ and in particular no trivial bubbles. We now verify that conditions 1 – 4 of Definition 3.11 hold.

1.
 - *The horizontal boundary $\partial_h N_B$ is incompressible in $M \setminus \text{int}(N_B)$* : this follows from the fact that the inclusions of $S' \times \{0\}$ and $S' \times \{1\}$ in $M \setminus \text{int}(N_B)$ are homotopy equivalences. In particular, if a simple closed curve in $\partial_h N_B$ bounds a disc in $M \setminus \text{int}(N_B)$ then it must be nullhomotopic in $\partial_h N_B$ and nullhomotopic simple closed curves in surfaces always bound embedded discs.
 - *The horizontal boundary $\partial_h N_B$ is ∂ -incompressible in $M \setminus \text{int}(N_B)$* : suppose that there is a disc $\Delta \subset M \setminus \text{int}(N_B)$ such that $\text{int}(\Delta) \subset M \setminus N_B$ and $\partial\Delta = a \cup b$, where a is an arc in $\partial_h N_B$ and $b = \partial\Delta \cap \partial M$. We have to find a disc $\Delta' \subset \partial_h N_B$ with $\partial\Delta' = a \cup b'$ where $b' = \partial\Delta' \cap \partial M$.

Without loss of generality we can suppose that $a \subset S' \times \{0\}$. The arc b is an arc in $\partial M \setminus \text{int}(N_B)$ with both endpoints in $S' \times \{0\}$ and since the connected components of $\partial M \setminus \text{int}(N_B)$ are either discs or annuli, there exists a homotopy in $\partial M \setminus \text{int}(N_B)$, relative to the boundary, from the arc b to an arc $b' \subset (S' \times \{0\}) \cap \partial M$. In particular since the simple closed curve $a \cup b$ is nullhomotopic in $M \setminus \text{int}(N_B)$, the curve $a \cup b'$ is nullhomotopic as well.

To conclude it is enough to observe that since the inclusion of $S' \times \{0\}$ in $M \setminus \text{int}(N_B)$ is a homotopy equivalence, the simple closed curve $a \cup b'$ bounds a disc Δ' in $S' \times \{0\}$.

- *No component of the horizontal boundary is a sphere or a properly embedded disc*: this follows by our hypotheses.
2. *there is no monogon in $M \setminus \text{int}(N_B)$* : this is a consequence of the fact that the branched surface B admits a coorientation.
 3.
 - *$M \setminus \text{int}(N_B)$ is irreducible*: this is a consequence of the fact that each component of $M \setminus \text{int}(N_B)$ is the product of a surface with boundary with $[0, 1]$.
 - *$\partial M \setminus \text{int}(N_B)$ is incompressible in $M \setminus \text{int}(N_B)$* : consider any boundary component T of M . By construction $T \setminus \text{int}(N_B)$ is a union of discs or an annulus (in case there are no endpoints of the arcs α_i on T). In the former case, $T \setminus \text{int}(N_B)$ is obviously incompressible in $M \setminus N_B$, while in the latter it is compressible if and only if it is the boundary of $S' \times [0, 1]$ and $S' \times [0, 1]$ is diffeomorphic to $\mathbb{D}^2 \times [0, 1]$, but this would contradict our hypotheses.
 4. *B contains no Reeb branched surfaces*: the presence of a Reeb branched surface would imply that some of the complementary regions of $\text{int}(N_B)$ are $\mathbb{D}^2 \times [0, 1]$ regions (see [GO89]) and we have already observed that there are no such regions.

The proof is complete. □

All the branched surfaces we will use are obtained with the previous construction. One problem that one has to face is that there could be sink discs in such branched surfaces. In [KR14] the authors present a useful procedure to build splittings of branched surfaces constructed in this way that are without sink discs, and part of the results that we are going to obtain can also be proved with the methods there presented (see Remark 3.24). However to be able to construct taut foliations on all the manifolds of the statement of Theorem 3.1 we will need to find a way to build our branched surfaces that is slightly different from the one presented in [KR14].

We fix some notation. We suppose that S is a torus with k open discs removed. We consider the curves $\gamma_0, \gamma_1, \dots, \gamma_k$ and we label the boundary components of S with numbers in $\{1, \dots, k\}$ as in Figure 12. We also orient S so that the orientation induced on the boundary components is the one of the figure.




Figure 12

We denote with τ_i the positive Dehn twist along the curve γ_i . Notice that since $\gamma_i \cap \gamma_j = \emptyset$ for $i, j = 1, \dots, k$ we have that $\tau_i \tau_j = \tau_j \tau_i$ for $i, j = 1, \dots, k$.

We focus on homeomorphisms of S of the following type:

$$h = \tau_0^{a_0} \tau_1^{a_1} \dots \tau_k^{a_k} \quad a_0 \in \mathbb{Z}, a_i \in \mathbb{Z} \setminus \{0\} \text{ for } i \neq 0.$$

We fix the following convention:

Convention: the indices $1, \dots, k$ have to be considered ordered ciclically; so we set $a_{k+1} = a_1$ and think of a_1 as consecutive to a_k . In particular when $k = 1$, *i.e.* S has only one boundary component, we set $a_{1+1} = a_1$ and think of a_1 as consecutive to itself.

Let $\partial_i S$ denote the boundary component of S labelled with i . Given such a homeomorphism h we assign to $\partial_i S$ a label with the following rule:

- we assign to $\partial_i S$ the label p_+ if a_i and a_{i+1} are both positive;
- we assign to $\partial_i S$ the label p_- if a_i and a_{i+1} are both negative;
- we assign to $\partial_i S$ the label n if a_i and a_{i+1} have different signs.

Figure 13 shows an example.




Figure 13: Example with $h = \tau_0 \tau_1^5 \tau_2^{10} \tau_3^{-5}$.

Finally we assign to each boundary component $\partial_i S$ two intervals I_i and J_i in $\overline{\mathbb{Q}}$ in the following way:

- if $\partial_i S$ has label p_+ we set $I_i = J_i = (\infty, 1)$;
- if $\partial_i S$ has label p_- we set $I_i = J_i = (-1, \infty)$;
- let $i_1 < \dots < i_{2c}$ the indices of the boundary components labelled with n . We set $I_{i_a} = (\infty, 0)$ when $a \in \{1, \dots, 2c\}$ is odd and we set $I_{i_a} = (0, \infty)$ when $a \in \{1, \dots, 2c\}$ is even.

Therefore we have $I_{i_1} = (\infty, 0)$, $I_{i_2} = (0, \infty)$, $I_{i_3} = (\infty, 0)$ and so on.

On the contrary, we set $J_{i_a} = (0, \infty)$ when $a \in \{1, \dots, 2c\}$ is odd and $J_{i_a} = (\infty, 0)$ when $a \in \{1, \dots, 2c\}$ is even.

Example 3.17. In the example of Figure 13 we have

$$\begin{aligned} I_1 &= J_1 = (\infty, 1) \\ I_2 &= (\infty, 0) \quad J_2 = (0, \infty) \\ I_3 &= (0, \infty) \quad J_3 = (\infty, 0). \end{aligned}$$

We are now ready to state the theorem. In the statement of the theorem, for each boundary torus T_i of the manifold M_h we have fixed as longitude the oriented curve $\partial_i S$ and as meridian the image in M_h of the curve $\{x_i\} \times [0, 1]$, oriented as $[0, 1]$, where $x_i \in \partial_i S$.

Theorem 3.18. *Let S be a k -holed torus as in Figure 12 and let h be a homeomorphism of S of the following form:*

$$h = \tau_0^{a_0} \tau_1^{a_1} \dots \tau_k^{a_k}$$

where $a_0 \in \mathbb{Z}$ and $a_i \in \mathbb{Z} \setminus \{0\}$ for $i = 1, \dots, k$. Then:

1. if $a_0 > 0$ (resp. $a_0 < 0$) then $M_h(s_1, \dots, s_k)$ supports a coorientable taut foliation for each multislope $(s_1, \dots, s_k) \in (\infty, 1)^k$ (resp. $(-1, \infty)^k$);
2. for any multislope $(s_1, \dots, s_k) \in (I_1 \times \dots \times I_k) \cup (J_1 \times \dots \times J_k)$, the filled manifold $M_h(s_1, \dots, s_k)$ supports a coorientable taut foliation, where the intervals I_i 's and J_i 's are the ones described above.

Remark 3.19. Notice that if h' is conjugated in $\text{MCG}(S, \partial S)$ to a homeomorphism h that satisfies the hypotheses of Theorem 3.18, then the conclusion of the theorem holds also for $M_{h'}$.

Remark 3.20. Notice that the first part of the theorem does not cover the case $a_0 = 0$. We did not investigate further this case, but for our purpose the statement of Theorem 3.18 will be sufficient.

3.2.1 Proof of the first part of Theorem 3.18

To prove Theorem 3.18 we will build branched surfaces in M_h satisfying the hypotheses of Theorem 3.12 by following the construction presented before Lemma 3.16.

We start by proving the first part of the theorem. We define a branched surface as follows. We consider the parallel arcs $\alpha_1, \dots, \alpha_k$ depicted in Figure 14.




Figure 14

We consider the discs $\alpha_i \times [0, 1]$, perturb them in a neighbourhood of $S \times \{1\}$ as explained in the discussion before Lemma 3.16, and project them to the mapping torus M_h . We consider the (co)oriented branched surface B in M_h obtained by adding these discs D_i to the surface S . The discs D_i 's are oriented so that the orientation on their boundary induces the given orientation on the arcs α_i 's. For an example, see Figure 15, where also some cusp directions are showed.




Figure 15: In this example, $a_0 = 2$ and $k = 3$.

Lemma 3.21. *The branched surface B is laminar and satisfies the hypotheses of Theorem 3.12.*

Proof. Thanks to Lemma 3.16 to prove that B is laminar it is enough to prove that B contains no sink discs or half sink discs. We prove that:


- *B contains no sink discs or half sink discs:* there are k sectors of B that are half discs and that coincide with the discs D_i 's; these sectors are never sink by construction (see Figure 11).

The other sectors coincide with the abstract closures of the connected components of $S \setminus (\cup_i \alpha_i \cup h(\alpha_i))$. Being a_0 non-zero, these sectors are discs and half discs³. We can organise these sectors in the following way. We refer to Figure 15 to visualise the situation. If we cut S along the arcs α_i 's we obtain k oriented annuli A_1, \dots, A_k , so that $\partial A_i \supset -\alpha_i \cup \alpha_{i+1}$ for $i \in \{1, \dots, k\}$, where $-\alpha_i$ denotes the arc α_i with the opposite orientation. Also notice that the cusp directions along α_i point inside A_i and the cusp directions along α_{i+1} point outside A_i .

It follows by the definition of the arcs α_i that $h(\alpha_i) = \tau_0^{a_0}(\alpha_i)$. Each of these annuli intersects $h(\alpha_j)$ in a_0 subarcs, for each $j = 1, \dots, k$. Therefore, when we cut along the $h(\alpha_i)$'s we subdivide each of this annuli in $k|a_0|$ discs and these discs coincide with the sectors of B in S . By construction each of these discs is contained in an annulus, say A_i , and intersects both α_i and α_{i+1} and therefore there is a cusp direction pointing outside it.

- *The only connected compact surface properly embedded in M_h carried by B is S :* if Σ is a compact surface properly embedded in M_h carried by B , then Σ induces an integral weight system on B ; that is to say, Σ defines a way to assign to each branch sector of B a non-negative integer such that along each connected component of the branch locus minus the set of triple points of B the weights sum according to the cusp directions, as represented in the following figure.

³When the product $k|a_0|$ satisfies $k|a_0| \leq 3$ there are only half disc sectors.



$$a = b + c$$

Figure 16

The way Σ induces a weight system is the following: we fix a point in the interior of each sector and we assign to each sector the number of intersections between Σ and the fiber of N_B over this fixed point. We denote with ω_i 's the weights of the half disc sectors D_i 's.

We have already observed that the other sectors of B are organised so that each of the annuli A_1, \dots, A_k contains $k|a_0|$ among discs and half discs sectors. When $a_0 > 0$ (resp. $a_0 < 0$), for each $i = 1, \dots, k$, we order in each annulus these sectors by following the direction of α_i (resp. $-\alpha_i$) and denote with $\Delta_{i,j}$ the discs contained in the annulus A_i , with $j = 1, \dots, ka_0$. We denote the weight of the disc $\Delta_{i,j}$ with $\omega_{i,j}$.

Let us fix an annulus A_{i_0} . Each pair of consecutive discs $\Delta_{i_0,j}, \Delta_{i_0,j+1}$ is separated by a subarc of $h(\alpha_l)$, for some $l = 1, \dots, k$. Since the arcs α_i 's (and therefore also the $h(\alpha_i)$'s) are parallel it follows by the orientation of the discs D_i 's that we have:

$$\omega_{i_0,j} + \omega_l = \omega_{i_0,j+1}.$$

This implies the following chain of inequality:

$$\omega_{i_0,1} \leq \omega_{i_0,2} \leq \dots \leq \omega_{i_0,k|a_0|} \leq \omega_{i_0,1}.$$

Therefore the weights $\omega_{i_0,j}$ are all equal and since all the arcs $h(\alpha_i)$'s intersect the annulus A_{i_0} we have that $\omega_l = 0$ for each $l = 1, \dots, k$. Therefore the sectors of B contained in S all have the same weight and the discs D_i 's have weight zero. This means that Σ is a finite number of parallel copies of S .

By construction, $\partial M_h \setminus \partial B$ is a union of bigons (see Figure 17). Moreover B does not carry any closed surface, and therefore it does not carry tori bounding a solid torus in $M_h(s_1, \dots, s_k)$, for any multislote (s_1, \dots, s_k) . Since M_h is irreducible and its boundary is union of incompressible tori, the hypotheses of Theorem 3.12 are fulfilled. \square

Proposition 3.22. *If $a_0 > 0$, for any multislote $(s_1, \dots, s_k) \in (\infty, 1)^k$ the branched surface B fully carries an essential lamination intersecting the boundary of M_h in parallel curves of multislote (s_1, \dots, s_k) . If $a_0 < 0$ the same happens for any multislote $(s_1, \dots, s_k) \in (-1, \infty)^k$.*

Proof. We study the multislopes realised by the boundary train tracks of B . In order to do this we assign rational weight systems to our boundary train tracks. Since our train tracks are oriented, we can associate to such a weight system the rational number w_{μ_i}/w_{λ_i} , where w_{μ_i} and w_{λ_i} are the *weighted* intersections of the train tracks with our fixed meridians μ_i and longitudes λ_i , as we would do with oriented simple closed curves. This quotient can be interpreted as a *slope* in the i -th boundary component of M_h . In fact it is easy to show that each slope p/q obtained in this way is realised by the train track. Since we want to study slopes *fully* carried by these train tracks, we have to require that each weight is strictly positive: if the weight of an arc is zero, the associated slope will not intersect the fibers over that arc. For details, see [PH16].

The boundary train tracks of B are all the same for each boundary tori, and only depend on the sign of a_0 . The two possible types of boundary train tracks are depicted in Figure 17.




Figure 17

We also endowed the two train tracks with weight systems. The slopes of these weight systems are always $x - y$, but since we have to impose that each sector of the train tracks has positive weight we have that:

- if $a_0 > 0$, x can vary in $(0, 1)$ and y can vary in $(0, +\infty)$;
- if $a_0 < 0$, x can vary in $(0, +\infty)$ and y can vary in $(0, 1)$.

By letting x, y vary we have that when $a_0 > 0$ the boundary train tracks realise all multislopes in $(\infty, 1)^k$ and when $a_0 < 0$ the boundary train tracks realise all slopes in $(-1, \infty)^k$. Thanks to Lemma 3.21 we can apply Theorem 3.12 to obtain the desired essential laminations. \square

Lemma 3.23. *All the laminations constructed in the previous proposition extend to taut foliations of the filled manifolds.*

Proof. Let Λ be one of the laminations constructed in Proposition 3.22 and suppose that Λ intersect ∂M_h in parallel curves of multislope (s_1, \dots, s_k) . We can also suppose that $\partial_h N_B \subset \Lambda$. First of all we notice that if the multislope is different from $(0, \dots, 0)$ then Λ does not have any compact leaves. In fact any leaf of Λ is carried by B and we have showed in the proof of Lemma 3.21 that the only connected compact surface carried by

B is the k -holed torus S . Therefore if Λ has a compact leaf then it should intersect ∂M_h in parallel curves of multislope $(0, \dots, 0)$.

We now consider the abstract closures (in a path metric on M_h) of the complementary regions of Λ . These closures are $[0, 1]$ -bundles; in fact they are unions, along $\partial_v N_B$, of:

- components of $M_h \setminus \text{int}(N_B)$, that are products of the type $F \times [0, 1]$, where F is a surface, with

$$\partial_h N_B \cap (F \times [0, 1]) = F \times \{0, 1\}$$

and

$$\partial_v N_B \cap (F \times [0, 1]) = \partial' F \times [0, 1]$$

where $\partial' F$ is the closure of $\partial F \setminus \partial M_h$;

- abstract closures of the components of $N_B \setminus \Lambda$. Since Λ intersects transversely the fibers of N_B also these closures are products with the same properties of the components of $M_h \setminus \text{int}(N_B)$.

Each component of the vertical boundary of N_B is an annulus $S^1 \times [0, 1]$ or a disc $[0, 1] \times [0, 1]$, where each interval $\{*\} \times [0, 1]$ is contained in a fiber of N_B . Both the product structures of the components of $M_h \setminus \text{int}(N_B)$ and of the abstract closures of the components of $N_B \setminus \Lambda$ define a foliation of the vertical boundary transverse to the interval fibers. Any of two such foliations are isotopic and therefore also the abstract closures of the complementary regions of Λ are products.

In particular, since the horizontal boundary of the closures of these complementary regions are leaves of Λ we can foliate these bundles with parallel leaves to obtain a foliation of the whole M_h . This foliation has no compact leaves and intersects the boundary of M in parallel curves of multislope (s_1, \dots, s_k) . Therefore the leaves of this foliation can be capped with the meridional discs of the solid tori to obtain a foliation of the filled manifold $M_h(s_1, \dots, s_k)$ that has no compact leaves as well, and that is therefore taut (see [Cal07, Example 4.23.]). \square

Remark 3.24. In the terminology of [KR14], if $|a_0| = 1$ then the pair of parallel k -uples

$$(\widetilde{h(\alpha)}, \alpha)$$

is good and oriented, where $\alpha = (\alpha_1, \dots, \alpha_k)$ and $\widetilde{h(\alpha)} = (\widetilde{h(\alpha_1)}, \dots, \widetilde{h(\alpha_k)})$. In this case the branched surface constructed in the previous discussion coincides with the branched surface associated to the sequence $(\widetilde{h(\alpha)}, \alpha)$ by Kalelkar and Roberts in [KR14].

3.2.2 Proof of the second part of Theorem 3.18

We now focus our attention on the second part of Theorem 3.18 and we define a new branched surface. We fix a new set of arcs $\alpha_1, \dots, \alpha_k$ in the following way. We consider arcs β_1, \dots, β_k as in Figure 18 and choose α_i so that $\widetilde{h(\alpha_i)} = \beta_i$. One example is depicted in Figure 19.




Figure 18

We now give orientations to the arcs α_i 's in order to build our branched surfaces. It will be simpler to state how to assign orientations to the β_i 's and we will orient each α_i as isotopic to $h^{-1}(\beta_i)$, for $i = 1, \dots, k$.

Definition 3.25. We say that an orientation of the arcs β_i 's is **coherent** if the following hold:

- if a boundary component $\partial_i S$ has label p_+ or p_- the arcs intersecting $\partial_i S$ are oriented so that one starts at $\partial_i S$ and the other ends at $\partial_i S$;
- if a boundary component $\partial_i S$ has label n the arcs intersecting $\partial_i S$ are oriented so that both start or both end at $\partial_i S$. In case the arcs both start at $\partial_i S$ we say that the component is of type $n_{\mathbf{o}}$ (the subscript \mathbf{o} stands for “out”) and if both end at $\partial_i S$ we say that is of type $n_{\mathbf{i}}$ (\mathbf{i} standing for “in”).

It is easy to see that since there is always an even number of boundary components of S with label n it is always possible to find a coherent orientation of the arcs β_i 's. In particular, there are exactly two of such orientations. Moreover the boundary components with label n are alternately of type $n_{\mathbf{o}}$ and $n_{\mathbf{i}}$. See Figure 19 for an example.




Figure 19

We fix a coherent orientation and as usual we consider the branched surface B that is the union of S and the images in M_h of the discs $\alpha_i \times [0, 1] \subset S \times [0, 1]$. We denote the image of $\alpha_i \times [0, 1]$ with D_i and orient the discs D_i 's so that the orientation on their boundary induces the given orientation on the α_i 's. Exactly as before we have:

Lemma 3.26. *The branched surface B is laminar and satisfies the hypotheses of Theorem 3.12.*

Proof. The proof is analogous to the one of Lemma 3.21. We only need to prove that B contains no sink discs or half sink discs, and that the only connected surface properly embedded in M_h carried by B is S .

- *B contains no sink discs or half sink discs:* there are k sectors of B that coincide with the discs D_i 's and they always have cusp directions pointing outside. We focus our attention on the sectors contained in S . We consider the k annuli A_i obtained by cutting S along the arcs of Figure 14. Each of these annuli contains in its interior some disc and half disc sectors and intersects two other half disc sectors. The former are never sink because each of these sectors has in its boundary two parallel subarcs of some arc of the α_i 's, as for example Figure 20 shows.




Figure 20

The sectors belonging to two consecutive annuli are never sink because each of these sectors has in its boundary two subarcs of two consecutive arcs of the α_i 's and, since we have fixed a coherent orientation of the arcs β_i 's, the cusp directions along the arcs α_i 's all point in the same direction. For an example, see Figure 21.




Figure 21

- *The only connected compact surface properly embedded in M_h carried by B is S :* suppose that Σ is a compact surface carried by B . Σ induces an integral weight

system on B . We denote with ω_i the weight of the discs D_i . The number of sectors contained in S is equal to $N = \sum_{i=1}^k |a_i|$. Since we have fixed a coherent orientation of the arcs β_i , the cusp directions along the arcs α_i 's all point in the same direction. We order the sectors in S according to this direction; we denote them with Δ_l and we denote their weights with δ_l , where $1 \leq l \leq N$. As in the proof of the first part of the theorem, we have that

$$\delta_1 \leq \delta_2 \leq \dots \leq \delta_N \leq \delta_1.$$

Therefore these weights are all equal and this implies that the discs D_i 's all have weight zero; that is to say, Σ is a finite number of parallel copies of S .

□

Proposition 3.27. *Let I_1, \dots, I_k and J_1, \dots, J_k be the intervals defined in the discussion before the statement of Theorem 3.18 and let B denote the branched surface associated to a coherent orientation of the arcs β_i 's. Then for one choice of coherent orientation of the arcs β_i 's, B fully carries essential laminations intersecting the boundary of M_h in parallel curves of multislope (s_1, \dots, s_k) , for $(s_1, \dots, s_k) \in I_1 \times \dots \times I_k$. Choosing the other coherent orientation yields B that fully carries essential laminations intersecting the boundary of M_h in parallel curves of multislope (s_1, \dots, s_k) , for $(s_1, \dots, s_k) \in J_1 \times \dots \times J_k$. Moreover these laminations extend to taut foliations of the filled manifold $M_h(s_1, \dots, s_k)$.*

Proof. We focus our attention on the boundary train tracks of B . For a fixed boundary component of S we have the four possible configurations showed in Figure 22 and for each of this configurations we have two possible way to fix a coherent orientation.




Figure 22

If the boundary component has label p_+ or p_- the type of the boundary train track does not depend on the choice of the coherent orientation, and is described in the following figure.




Figure 23

By assigning weights to these train tracks as we have already done in the proof of Proposition 3.22 we have that if the label is p_+ the train track realises all the slopes in the interval $(\infty, 1)$, while if the label is p_- the slopes realised are those in the interval $(-1, \infty)$.

On the other hand if the boundary component has label n the choice of the orientation yields two different train tracks. We represent in Figure 24 the two train tracks that can occur when we have the configuration of type $a)$.




Figure 24

The train track on the left realises all the slopes in the interval $(0, \infty)$, while the one on the right realises the slopes in the interval $(\infty, 0)$. Analogously, we have that if a boundary component has label n_i of type $b)$ we obtain a train track that realises all the slopes in $(0, \infty)$, and if it has label n_o of type $b)$ we obtain a train track that realises all the slopes in $(\infty, 0)$.

By fixing one or the other of the two possible coherent orientations, we have that the boundary train tracks of B realise all the multislopes in $I_1 \times \cdots \times I_k$ and $J_1 \times \cdots \times J_k$. By virtue of Lemma 3.26, we can apply Theorem 3.12 to obtain the desired essential laminations and Lemma 3.23 implies that these laminations extends to taut foliations of the filled manifolds. \square

Proof of Theorem 3.18. The first part of the theorem is the content of Proposition 3.22 and Lemma 3.23. The second part is the content of Proposition 3.27. \square

Example 3.28. For each natural number n we consider the n -component oriented link \mathcal{L}_n in figure.




Figure 25

We can represent the link \mathcal{L}_n in a different way, as in Figure 26. With this representation, it is evident that \mathcal{L}_n is a fibered link, with fiber surface a torus with n open discs removed. Moreover the monodromy associated to this fiber is

$$h = \tau_0^{-1} \tau_1 \dots \tau_n$$

where τ_i is the positive Dehn twist along the curve γ_i .




Figure 26

It follows by a theorem of Penner (see [Pen88]), that \mathcal{L}_n is a hyperbolic link for each n . Moreover Theorem 3.18 applies to these links and we can deduce that for any multi-slope $(s_1, \dots, s_n) \in (\infty, 1)^n \cup (-1, \infty)^n$, the filling of the exterior of \mathcal{L}_n with multi-slope (s_1, \dots, s_n) supports a coorientable taut foliation. Recall that these slopes are referred to the meridian-longitude bases given by the mapping torus; since the components of the link \mathcal{L}_n do not have pairwise linking number zero, the longitudes of these bases do not coincide with the canonical longitudes of the link.

3.3 The Whitehead link case

We now return to the Whitehead link exterior. Recall from the discussion preceding Figure 10 that the Whitehead link is a fibered link with fiber surface a 2-holed torus and monodromy $h = \tau_0 \tau_1 \tau_2^{-1}$, where the curves γ_i are represented in Figure 27. In what follow we will identify the exterior of the Whitehead link with the mapping torus M_h .




Figure 27

As a consequence of Theorem 3.18 and of the fact that the components of the Whitehead link have linking number zero, we have

Corollary 3.29. *Let $(p_1/q_1, p_2/q_2)$ be a multislope in*

$$(\infty, 1)^2 \cup (0, \infty) \times (\infty, 0) \cup (\infty, 0) \times (0, \infty).$$

Then $S^3_{p_1/q_1, p_2/q_2}(\text{WL})$ supports a coorientable taut foliation.

We have been able to prove that for slopes in the region depicted in Figure 28, the corresponding filling on the Whitehead exterior supports a coorientable taut foliation.




Figure 28

We now cover the remaining regions of Figure 28. We define a branched surface by considering the pair of arcs α, β shown in Figure 29-(left). We apply the monodromy h and we obtain what is depicted in Figure 29.




Figure 29

As usual, we consider the branched surface B associated to the arcs α and β .

Lemma 3.30. *The branched surface B is laminar and satisfies the hypotheses of Theorem 3.12.*

Proof. Since $S \setminus (\alpha \cup \beta)$ has no disc components, by virtue of Lemma 3.16 we only need to prove that B contains no sink disc or half sink discs. The sectors of B are five half discs. The cusp directions of B are depicted in Figure 30 and it is easy to check that none of these half discs is sink.




Figure 30

To prove that B satisfies the hypotheses of Theorem 3.12 we have to show that $\partial M_h \setminus \partial B$ is a union of bigons and that B does not carry a torus. By construction $\partial M_h \setminus \partial B$ is union of bigons (see Figure 31) and since each sector of B intersect ∂M_h it follows that any surface carried by B must intersect ∂M_h . Therefore B does not carry any closed surfaces and in particular it does not carry tori. \square

Corollary 3.31. *Let $(p_1/q_1, p_2/q_2)$ be slopes in*

$$(0, \infty) \times (-1, 1) \cup (-1, 1) \times (0, \infty)$$

Then the filled manifold $S^3_{p_1/q_1, p_2/q_2}(\text{WL})$ supports a coorientable taut foliation.

Proof. By virtue of Theorem 3.12 for any multislope (s_1, s_2) realised by the boundary train tracks of B there exists an essential lamination Λ fully carried by B intersecting the boundary of the exterior of the Whitehead link in parallel curves of multislope (s_1, s_2) . The boundary train tracks of B are the following:




Figure 31

By assigning weights to these train tracks as in Figure 32 it follows that these train tracks realise all the slopes in $(0, \infty) \times (-1, 1)$.




Figure 32

Let Λ be an essential lamination intersecting the boundary of M_h in parallel curves of one of these multislopes (s_1, s_2) . Since each leaf of Λ is carried by B and each sector of B intersects ∂M_h it follows that all the leaves of Λ intersect ∂M_h . It follows by the proof of Lemma 3.23 that we can construct a foliation of M_h such that each leaf of this foliation is parallel to some leaf of Λ . Therefore all the leaves of these foliation intersect ∂M and as a consequence when we cap these leaves with the meridional discs of the solid tori we obtain a foliation of $M_h(s_1, s_2)$ with the property that the cores of the tori are transversals intersecting all the leaves.

We have proved that for any $(p_1/q_1, p_2/q_2) \in (0, \infty) \times (-1, 1)$ the manifold $S^3_{p_1/q_1, p_2/q_2}(\text{WL})$ supports a coorientable taut foliation. Since the Whitehead link is symmetric we deduce that the same result holds also for any $(p_1/q_1, p_2/q_2) \in (-1, 1) \times (0, \infty)$. \square

Proof of Theorem 3.1. It is a consequence of Corollary 3.29 and Corollary 3.31. \square

4 Orderability

In this last section we will prove Theorem 1.4 and discuss some results about the orderability, and non orderability, of some surgeries on the Whitehead link. We recall the following definition:

Definition 4.1. Let G be a group. G is **left orderable** if there exists a total order $<$ on G that is invariant for the left multiplication by elements in G , i.e. such that for any $g, g' \in G$ we have that $g < g'$ if and only if $hg < hg' \forall h \in G$.

If G is the fundamental group of a closed, orientable, irreducible 3-manifold, left orderability translates in the following dynamical property.

Theorem 4.2. [BRW05] *Let N be a closed, irreducible, orientable 3-manifold. Then $\pi_1(N)$ is left orderable if and only if there exists a non trivial homomorphism $\varphi : \pi_1(N) \rightarrow \text{Homeo}^+(\mathbb{R})$.*

This result yields us a theoretical way to connect taut foliations to left orderability in the following way. Suppose that \mathcal{F} is a cooriented taut foliation on a rational homology 3-sphere N . We can associate to \mathcal{F} its tangent bundle $T\mathcal{F}$, that is a plane bundle over N . Being a plane bundle, we can associate to $T\mathcal{F}$ its Euler class $e(T\mathcal{F}) \in H^2(N; \mathbb{Z})$. Moreover, by a construction of Thurston (see [CD03]), it is possible to associate to \mathcal{F} a non trivial homomorphism

$$\varphi : \pi_1(N) \rightarrow \text{Homeo}^+(S^1).$$

Since there is an injective homomorphism from the universal cover $\widetilde{\text{Homeo}}^+(S^1)$ of $\text{Homeo}^+(S^1)$ into $\text{Homeo}^+(\mathbb{R})$, one would like to lift φ to a homomorphism

$$\tilde{\varphi} : \pi_1(N) \rightarrow \widetilde{\text{Homeo}}^+(S^1).$$

The obstruction to find such a lift is again a comohomology class in $H^2(N; \mathbb{Z})$, and it turns out that it coincides with $e(T\mathcal{F})$. For more details we refer to [BH19].

The upshot of the previous discussion is the following

Theorem 4.3. [BH19] *Let N be a rational homology sphere and let \mathcal{F} be a coorientable taut foliation on N . If the Euler class of \mathcal{F} vanishes then $\pi_1(N)$ is left orderable.*

We now consider the taut foliations obtained in the previous section and determine which of them have vanishing Euler class. To do this we will adapt part of the content of [Hu19] to our context.

We fix some notation. We denote with M the exterior of WL and we denote with S the 2-holed torus of Figure 10 that is a Seifert surface for WL. We fix a multislope $(p_1/q_1, p_2/q_2)$, with $p_1/q_1 < 1$ or $p_2/q_2 < 1$, and we denote with \mathcal{F} the foliation in M intersecting ∂M in parallel curves of multislope $(p_1/q_1, p_2/q_2)$, as constructed in the proof of Theorem 3.1. This foliation extends to a foliation $\hat{\mathcal{F}}$ of the filled manifold $S^3_{p_1/q_1, p_2/q_2}(\text{WL})$ so that in the glued solid tori N_1 and N_2 the foliation $\hat{\mathcal{F}}$ restricts to the standard foliations by meridional discs \mathcal{D}_1 and \mathcal{D}_2 . We can suppose without loss of generality that $p_1, p_2 > 0$. We orient the meridional disc D_i of N_i so that the gluing map identifies ∂D_i with the oriented curve $p_i\mu_i + q_i\lambda_i$ in ∂M .

The second homology group $H_2(M, \partial M; \mathbb{Z})$ is isomorphic to \mathbb{Z}^2 and in particular we can fix as generators two properly embedded surfaces S_1 and S_2 that are duals to the meridians of the two components of the Whitehead link. Since the Whitehead link has linking number zero, these surfaces can be taken to be Seifert surfaces for the components

of the link. In particular, it is easy to see that these can be chosen to be tori with one disc removed, so that $\partial S_i = \lambda_i$. One of these tori is showed in Figure 33 and the other can be obtained by an isotopy of S^3 exchanging the two components of WL.




Figure 33

We fix a nowhere vanishing section σ of $(T\mathcal{F})|_{\partial M}$ that is everywhere pointing outside of M . Hence the restrictions of σ to the boundary components of M also define nowhere vanishing sections σ_i of $(T\mathcal{D}_i)|_{\partial N_i}$ everywhere pointing inside N_i , for $i = 1, 2$. These sections yield us relative Euler classes in $H^2(M, \partial M; \mathbb{Z})$, $H^2(N_1, \partial N_1; \mathbb{Z})$ and $H^2(N_2, \partial N_2; \mathbb{Z})$, that we denote respectively with $e_\sigma(T\mathcal{F})$, $e_{\sigma_1}(T\mathcal{D}_1)$ and $e_{\sigma_2}(T\mathcal{D}_2)$. See [Hu19] for details.

Finally, we set $a_i = \langle e_\sigma(T\mathcal{F}), [S_i] \rangle$ and $b_i = \langle e_{\sigma_i}(T\mathcal{D}_i), [D_i] \rangle$, where D_i is a meridional disc in N_i .

Remark 4.4. Notice that since \mathcal{D}_i is the standard foliation of the solid torus by meridional discs, we have that b_i coincides with $\pm \langle e_{\sigma_i}(T\mathcal{D}_i), [D_i] \rangle = \pm \chi(D_i) = \pm 1$, where the sign depends on the orientation of the foliation \mathcal{D}_i .

We are interested in knowing when $e(T\hat{\mathcal{F}})$ vanishes. The following proposition tells us exactly when this happens. Recall that without loss of generality we are supposing $p_1, p_2 > 0$, whereas the signs of q_1 and q_2 are arbitrary.

Proposition 4.5. *We have that $e(T\hat{\mathcal{F}}) = 0$ if and only if $e(T\mathcal{F}) = 0$ and $a_i q_i \equiv b_i \pmod{p_i}$.*

Proof. The statement of this proposition is the generalisation to our case of the statements of Lemma 3.1 and Theorem 1.4 in [Hu19] and the proof that is presented there adapts almost unaltered. We give a brief sketch of the proof and refer to [Hu19] for the details. In what follows the cohomology and homology groups are all implicitly assumed with integer coefficients and we will denote with \overline{M} the $(p_1/q_1, p_2/q_2)$ -surgery on M . Since \overline{M} is a rational homology sphere as a consequence of the long exact sequence of the pair $(\overline{M}, \partial M)$ we have

$$0 \rightarrow H^1(\partial M) \xrightarrow{\delta} H^2(\overline{M}, \partial M) \xrightarrow{\iota} H^2(\overline{M}) \rightarrow 0. \quad (1)$$

Moreover there is an isomorphism

$$H^2(M, \partial M) \oplus H^2(N_1, \partial N_1) \oplus H^2(N_2, \partial N_2) \cong H^2(\overline{M}, \partial M) \quad (2)$$

defined by mapping the relative classes (c_M, c_{N_1}, c_{N_2}) to the sum $\overline{c_M} + \overline{c_{N_1}} + \overline{c_{N_2}}$, where each of these cohomology classes is obtained by extending to \overline{M} the corresponding relative class by the zero map.

By using the identification given by the isomorphism in (2) we obtain a short exact sequence:

$$0 \rightarrow H^1(\partial M) \xrightarrow{\psi} H^2(M, \partial M) \oplus H^2(N_1, \partial N_1) \oplus H^2(N_2, \partial N_2) \xrightarrow{\varphi} H^2(\overline{M}) \rightarrow 0 \quad (3)$$

where

$$\psi(\beta) = (\delta\beta, (\delta \circ f_1^*)(\beta), (\delta \circ f_2^*)(\beta))$$

with $f_1 : \partial N_1 \hookrightarrow \partial M$ and $f_2 : \partial N_2 \hookrightarrow \partial M$ denoting the gluing maps of the solid tori.

We suppose now that $e(T\hat{\mathcal{F}}) = 0$. By naturality of the euler class, $e(T\mathcal{F})$ is the image of $e(T\hat{\mathcal{F}}) = 0$ under the map induced by the inclusion $M \hookrightarrow \overline{M}$ and therefore we have that $e(T\mathcal{F}) = 0$.

Moreover it is easy to show that it also holds that

$$\varphi(e_\sigma(T\mathcal{F}), e_{\sigma_1}(T\mathcal{D}_1), e_{\sigma_2}(T\mathcal{D}_2)) = e(\mathcal{F}) = 0$$

and therefore there exists $\beta \in H^1(\partial M)$ such that $\psi(\beta) = (e_\sigma(T\mathcal{F}), e_{\sigma_1}(T\mathcal{D}_1), e_{\sigma_2}(T\mathcal{D}_2))$; in other words β satisfies

$$\begin{cases} \delta\beta = e_\sigma(T\mathcal{F}) \\ (\delta \circ f_1^*)(\beta) = e_{\sigma_1}(T\mathcal{D}_1) \\ (\delta \circ f_2^*)(\beta) = e_{\sigma_2}(T\mathcal{D}_2) \end{cases}$$

and this implies that $a_i q_i \equiv b_i \pmod{p_i}$. In fact we have

$$\begin{aligned} b_i &= \langle e_{\sigma_i}(T\mathcal{D}_i), [D_i] \rangle = \langle (\delta \circ f_i^*)(\beta), [D_i] \rangle = \langle \beta, [f(\partial D_i)] \rangle = \\ &= \langle \beta, p_i \mu_i + q_i \lambda_i \rangle = p_i \langle \beta, \mu_i \rangle + q_i \langle \beta, \lambda_i \rangle = p_i \langle \beta, \mu_i \rangle + q_i a_i \end{aligned}$$

where in the last equality we have used that

$$\langle \beta, \lambda_i \rangle = \langle \beta, [\partial S_i] \rangle = \langle \delta\beta, [S_i] \rangle = \langle e_\sigma(T\mathcal{F}), [S_i] \rangle = a_i.$$

We now prove that if $e(T\mathcal{F}) = 0$ and $a_i q_i \equiv b_i \pmod{p_i}$ for $i = 1, 2$, then $e(T\hat{\mathcal{F}}) = 0$.

We consider again the short exact sequence in (1). The nowhere vanishing section σ defines an element $e_\sigma(T\hat{\mathcal{F}}) \in H^2(\overline{M}, \partial M)$ that satisfies $\iota(e_\sigma(T\hat{\mathcal{F}})) = e(T\hat{\mathcal{F}})$ and therefore if we prove that $e_\sigma(T\hat{\mathcal{F}})$ belongs to the image of $\delta : H^1(\partial M) \rightarrow H^2(\overline{M}, \partial M)$ we obtain the thesis. Moreover under the isomorphism (2) the element $(e_\sigma(T\mathcal{F}), e_{\sigma_1}(T\mathcal{D}_1), e_{\sigma_2}(T\mathcal{D}_2))$ corresponds to $e_\sigma(T\hat{\mathcal{F}})$ and therefore it is enough to prove that $(e_\sigma(T\mathcal{F}), e_{\sigma_1}(T\mathcal{D}_1), e_{\sigma_2}(T\mathcal{D}_2))$ belongs to the image of ψ in the short exact sequence (3).

If we consider the long exact sequence of the pair $(M, \partial M)$ we have the following

$$H^1(M; \mathbb{Z}) \xrightarrow{\iota} H^1(\partial M) \xrightarrow{\delta} H^2(M, \partial M) \xrightarrow{\iota} H^2(M)$$

and since $\iota(e_\sigma(T\mathcal{F})) = e(T\mathcal{F}) = 0$ we deduce that there exists $\beta_0 \in H^1(\partial M)$ such that $\delta(\beta_0) = e_\sigma(T\mathcal{F}) \in H^2(M, \partial M)$. We now want to modify β_0 in order to find $\beta \in H^1(\partial M)$ that satisfies

$$\psi(\beta) = (e_\sigma(T\mathcal{F}), e_{\sigma_1}(T\mathcal{D}_1), e_{\sigma_2}(T\mathcal{D}_2))$$

that is to say, such that

$$\begin{cases} \delta\beta = e_\sigma(T\mathcal{F}) \\ (\delta \circ f_1^*)(\beta) = e_{\sigma_1}(T\mathcal{D}_1) \\ (\delta \circ f_2^*)(\beta) = e_{\sigma_2}(T\mathcal{D}_2). \end{cases}$$

We denote with $\mu_i^* \in H^1(M)$ the dual of $\mu_i \in H_1(M)$ and we define

$$\beta = \beta_0 + n_1\mu_1^* + n_2\mu_2^* \quad \text{where } n_i = -\langle \beta_0, \mu_i \rangle - \frac{a_i q_i - b_i}{p_i}.$$

Since $a_i q_i \equiv b_i \pmod{p_i}$ for $i = 1, 2$ it follows that n_i is an integer number. Moreover, since $\beta - \beta_0 \in \iota(H^1(M))$ we have that $\delta(\beta_0) = \delta(\beta) = e_\sigma(T\mathcal{F}) \in H^2(M, \partial M)$. We have to prove that $(\delta \circ f_i^*)(\beta) = e_{\sigma_i}(T\mathcal{D}_i)$ for $i = 1, 2$. Since

$$H^2(N_i, \partial N_i) \cong \text{Hom}(H_2(N_i, \partial N_i, \mathbb{Z}))$$

it is enough to prove that $\langle (\delta \circ f_i^*)(\beta), [D_i] \rangle = \langle e_{\sigma_i}(T\mathcal{D}_i), [D_i] \rangle$ and this follows by direct computation. □

Theorem 1.4. *Let $S_{p_1/q_1, p_2/q_2}^3(\text{WL})$ be the $(p_1/q_1, p_2/q_2)$ -surgery on the Whitehead link, with $q_1, q_2 \neq 0$ and $p_1, p_2 > 0$.*

Then the taut foliations constructed in the proof of the Theorem 1.1 have vanishing Euler class if and only if $|q_i| \equiv 1 \pmod{p_i}$ for $i = 1, 2$.

In particular, for all these manifolds the L-space conjecture holds.

Proof. First of all we observe that $e(T\mathcal{F}) = 0$. In fact, let T denote one of the boundary components of M ; the inclusion of T in M induces an isomorphism $\iota : H^2(M; \mathbb{Z}) \rightarrow H^2(T; \mathbb{Z})$. Therefore we have

$$e(T\mathcal{F}) = 0 \Leftrightarrow \iota(e(T\mathcal{F})) = 0.$$

By naturality of the Euler class we have that

$$\iota(e(T\mathcal{F})) = e(T(\mathcal{F}|_T))$$

and since $\mathcal{F}|_T$ admits a nowhere vanishing section we have that the last quantity is zero.

We now want to compute the numbers a_i . As a consequence of the proof of Theorem 1.7 in [Hu19] we have that

- $b_i = 1$ if $q_i < 0$, for $i = 1, 2$;
- $b_i = -1$ if $q_i > 0$, for $i = 1, 2$;

- $\langle e_\sigma(T\mathcal{F}), [S] \rangle = \chi(S) = -2$.

Since by construction S intersects positively in one point the meridians of the components of the Whitehead link, we have the equality $[S] = [S_1 + S_2]$ in $H_2(M, \partial M; \mathbb{Z})$ and hence

$$a_1 + a_2 = \langle e_\sigma(T\mathcal{F}), [S_1] \rangle + \langle e_\sigma(T\mathcal{F}), [S_2] \rangle = \chi(S) = -2.$$

As a consequence of [Thu86, Corollary 1, p. 118] for any $[F] \in H_2(M, \partial M; \mathbb{Z})$ we have the inequality

$$|\langle e_\sigma(T\mathcal{F}), [F] \rangle| \leq |\chi(F)|$$

and since S_1 and S_2 are 1-holed tori, this implies that $a_i = \langle e_\sigma(T\mathcal{F}), [S_i] \rangle = -1$ for $i = 1, 2$. Therefore, by virtue of Proposition 4.5 we have $e(T\hat{\mathcal{F}}) = 0$ if and only if for each $i = 1, 2$ it holds one of the following:

- q_i is positive and $q_i \equiv 1 \pmod{p_i}$;
- q_i is negative and $q_i \equiv -1 \pmod{p_i}$.

In other words $e(T\hat{\mathcal{F}}) = 0$ if and only if

$$|q_i| \equiv 1 \pmod{p_i} \quad \text{for } i = 1, 2$$

that is exactly what we wanted. □

We point out the following straightforward consequence of Theorem 1.4.

Corollary 4.6. *Let d_1, d_2 be two integers such that $d_1 < 0$ or $d_2 < 0$. Then the manifold $S_{d_1, d_2}^3(\text{WL})$ satisfies the L-space conjecture.*

We conclude by collecting from the literature some results regarding the orderability (or non orderability) of some surgeries on the Whitehead link, obtaining a generalisation of Corollary 4.6.

Theorem 1.5. *Let $m \neq 0$ be an integer.*

- *If $m \leq -1$ then the manifolds $S_{m, p/q}^3(\text{WL})$ and $S_{p/q, m}^3(\text{WL})$ have left orderable fundamental group for all rationals p/q .*
- *If $m \geq 1$ then the manifolds $S_{m, p/q}^3(\text{WL})$ and $S_{p/q, m}^3(\text{WL})$ have non left orderable fundamental group for all rationals $p/q \geq 1$.*

In particular, all the rational homology spheres obtained by integer surgery on WL satisfy the L-space conjecture.

Proof. The manifold $S_{m, \bullet}^3(\text{WL})$ fibers over the circle if and only if m is an integer (see [HMW11]). Moreover, in this case the fiber is a punctured torus.

- When $m \leq -1$ the monodromy of this fibration satisfies the hypotheses of [Zun20, Theorem 1] and therefore all the non trivial fillings of $S_{m, \bullet}^3(\text{WL})$ have left orderable fundamental group. Since WL is symmetric, the same result holds for $S_{\bullet, m}^3(\text{WL})$.

- When $m \geq 3$ and $p/q \geq 1$, the fundamental group of the manifold $S_{m,p/q}^3(\text{WL})$ was studied by Roberts, Shareshian and Stein in [RSS03], where they prove that it is not orderable. Their proof also works in the case $m = 2$. When $m = 1$ the manifold $S_{1,\bullet}^3(\text{WL})$ is the exterior of the right-handed trefoil knot and its surgeries are well known. In fact all surgeries but one yield Seifert fibered manifolds and we have already proved that these are L -spaces; since the L -space conjecture holds for Seifert fibered manifolds we deduce that these surgeries are all non orderable. The remaining surgery is a connected sum of two lens spaces and since its fundamental group has torsion it is not orderable as well.

This concludes the proof. □

Remark 4.7. Notice that even if the content of Corollary 4.6 is generalised by Theorem 1.5, the statement of Theorem 1.4 is not; in fact it is easy to see that there are also non-integer rationals p_1/q_1 and p_2/q_2 that satisfy the hypotheses of Theorem 1.4.

In [Dun20] Dunfield considers a census of more than 300,000 hyperbolic rational homology spheres, testing the conjecture for this census. These manifolds are obtained by filling 1-cusped hyperbolic 3-manifolds that can be triangulated with at most 9 ideal tetrahedra, see [Bur14]. We checked whether some of these manifolds studied by Dunfield arise as Dehn surgery on the Whitehead link and obtained the following

Proposition 4.8. *Among the 307,301 rational homology spheres studied in [Dun20] at least 625 are obtained as Dehn surgery on the Whitehead link. In [Dun20] it is proved that:*

- 300 of these manifolds are orderable;
- 250 are non orderable.

It follows from Theorem 1.5 that 16 of the remaining 75 manifolds are orderable and 10 are non orderable.

The code of the program can be found at [Cod]. The surgery coefficients yielding these manifolds are plotted in Figure 34, where the red dots represent the coefficients whose corresponding surgery is non orderable and the blue dots represent the coefficients whose corresponding surgery is orderable.




Figure 34

The examples of Theorem 1.5 and Proposition 4.8 are consistent with the conjecture, and together with Theorem 1.1 confirm that the L-space conjecture holds also for all these manifolds.

References

- [BC17] Steven Boyer and Adam Clay. Foliations, orders, representations, L-spaces and graph manifolds. *Advances in Mathematics*, 310:159–234, 2017.
- [BGW13] Steven Boyer, Cameron McA Gordon, and Liam Watson. On L-spaces and left-orderable fundamental groups. *Mathematische Annalen*, 356(4):1213–1245, 2013.
- [BH19] Steven Boyer and Ying Hu. Taut foliations in branched cyclic covers and left-orderable groups. *Transactions of the American Mathematical Society*, 372(11):7921–7957, 2019.
- [Bow16] Jonathan Bowden. Approximating C^0 -foliations by contact structures. *Geometric and Functional Analysis*, 26(5):1255–1296, 2016.
- [BRW05] Steven Boyer, Dale Rolfsen, and Bert Wiest. Orderable 3-manifold groups. In *Annales de l’institut Fourier*, volume 55, pages 243–288, 2005.
- [Bur14] Benjamin A. Burton. The cusped hyperbolic census is complete. *arXiv preprint arXiv:1405.2695*, 2014.
- [Cal07] Danny Calegari. *Foliations and the geometry of 3-manifolds*. Oxford University Press on Demand, 2007.
- [CC00] A. Candel and L. Conlon. *Foliations I*. Number v. 1 in Foliations I. American Mathematical Society, 2000.

- [CD03] Danny Calegari and Nathan M. Dunfield. Laminations and groups of homeomorphisms of the circle. *Inventiones mathematicae*, 152(1):149–204, 2003.
- [CD18] Marc Culler and Nathan Dunfield. Orderability and Dehn filling. *Geometry & Topology*, 22(3):1405–1457, 2018.
- [CKR19] Vincent Colin, William H. Kazez, and Rachel Roberts. Taut foliations. *Communications in Analysis and Geometry*, 27(2):357–375, 2019.
- [Cod] Code available at <https://sites.google.com/view/santoro/research>.
- [DR19] Charles Delman and Rachel Roberts. Persistently foliar composite knots. *arXiv preprint arXiv:1905.04838*, 2019.
- [DR20] Charles Delman and Rachel Roberts. Taut foliations from double-diamond replacements. *Characters in Low-Dimensional Topology*, 760:123, 2020.
- [Dun20] Nathan M. Dunfield. Floer homology, group orderability, and taut foliations of hyperbolic 3-manifolds. *Geometry & Topology*, 24(4):2075–2125, 2020.
- [FO84] William Floyd and Ulrich Oertel. Incompressible surfaces via branched surfaces. *Topology*, 23(1):117–125, 1984.
- [Gab86] David Gabai. Detecting fibred links in S^3 . *Commentarii Mathematici Helvetici*, 61(1):519–555, 1986.
- [GH17] Eugene Gorsky and Jennifer Hom. Cable links and L-space surgeries. *Quantum Topology*, 8(4):629–666, 2017.
- [GLM20] Eugene Gorsky, Beibei Liu, and Allison H Moore. Surgery on links of linking number zero and the Heegaard Floer d -invariant. *Quantum Topology*, 11(2):323–378, 2020.
- [GN16] Eugene Gorsky and András Némethi. Links of plane curve singularities are L-space links. *Algebraic & Geometric Topology*, 16(4):1905–1912, 2016.
- [GN18] Eugene Gorsky and András Némethi. On the set of l-space surgeries for links. *Advances in Mathematics*, 333:386–422, 2018.
- [GO89] David Gabai and Ulrich Oertel. Essential laminations in 3-manifolds. *Annals of Mathematics*, 130(1):41–73, 1989.
- [Hed10] Matthew Hedden. Notions of positivity and the Ozsváth–Szabó concordance invariant. *Journal of Knot Theory and its Ramifications*, 19(05):617–629, 2010.
- [HMW11] Craig D. Hodgson, G. Robert Meyerhoff, and Jeffrey R. Weeks. Surgeries on the Whitehead link yield geometrically similar manifolds. In *Topology'90*, pages 195–206. de Gruyter, 2011.

- [HRRW20] Jonathan Hanselman, Jacob Rasmussen, Sarah Dean Rasmussen, and Liam Watson. L-spaces, taut foliations, and graph manifolds. *Compositio Mathematica*, 156(3):604–612, 2020.
- [Hu19] Ying Hu. Euler class of taut foliations and Dehn filling. *arXiv preprint arXiv:1912.01645*, 2019.
- [Juh15] András Juhász. A survey of Heegaard Floer homology. In *New ideas in low dimensional topology*, pages 237–296. World Scientific, 2015.
- [KR14] Tejas Kalelkar and Rachel Roberts. Taut foliations in surface bundles with multiple boundary components. *Pacific Journal of Mathematics*, 273(2):257–275, 2014.
- [KR17] William Kazez and Rachel Roberts. C^0 approximations of foliations. *Geometry & Topology*, 21(6):3601–3657, 2017.
- [Kri20] Siddhi Krishna. Taut foliations, positive 3-braids, and the L-space conjecture. *Journal of Topology*, 13(3):1003–1033, 2020.
- [Li02] Tao Li. Laminar branched surfaces in 3-manifolds. *Geometry & Topology*, 6(1):153–194, 2002.
- [Li03] Tao Li. Boundary train tracks of laminar branched surfaces. In *Proceedings of symposia in pure mathematics*, volume 71, pages 269–286. Providence, RI; American Mathematical Society; 1998, 2003.
- [Liu17] Yajing Liu. L-space surgeries on links. *Quantum Topology*, 8(3):505–570, 2017.
- [LR14] Tao Li and Rachel Roberts. Taut foliations in knot complements. *Pacific Journal of Mathematics*, 269(1):149–168, 2014.
- [Ni07] Yi Ni. Knot Floer homology detects fibred knots. *Inventiones mathematicae*, 170(3):577–608, 2007.
- [Oer84] Ulrich Oertel. Incompressible branched surfaces. *Inventiones mathematicae*, 76(3):385–410, 1984.
- [OS04] Peter Ozsváth and Zoltán Szabó. Holomorphic disks and genus bounds. *Geometry & Topology*, 8(1):311–334, 2004.
- [OS05] Peter Ozsváth and Zoltán Szabó. On knot Floer homology and lens space surgeries. *Topology*, 44(6):1281–1300, 2005.
- [OS10] Peter S Ozsváth and Zoltán Szabó. Knot Floer homology and rational surgeries. *Algebraic & Geometric Topology*, 11(1):1–68, 2010.

- [Pen88] Robert C. Penner. A construction of pseudo-Anosov homeomorphisms. *Transactions of the American Mathematical Society*, 310(1):179–197, 1988.
- [PH16] Robert C. Penner and John L. Harer. *Combinatorics of Train Tracks. (AM-125), Volume 125*. Princeton University Press, 2016.
- [Ras17] Sarah Dean Rasmussen. L-space intervals for graph manifolds and cables. *Compositio Mathematica*, 153(5):1008–1049, 2017.
- [Ras20] Sarah Dean Rasmussen. Rational L-space surgeries on satellites by algebraic links. *Journal of Topology*, 13(4):1333–1387, 2020.
- [Rob00] Rachel Roberts. Taut foliations in punctured surface bundles, I. *Proceedings of the London Mathematical Society*, 82(3):747–768, 2000.
- [Rob01] Rachel Roberts. Taut foliations in punctured surface bundles, II. *Proceedings of the London Mathematical Society*, 83(2):443–471, 2001.
- [RR17] Jacob Rasmussen and Sarah Dean Rasmussen. Floer simple manifolds and L-space intervals. *Advances in Mathematics*, 322:738–805, 2017.
- [RSS03] Rachel Roberts, John Shareshian, and Melanie Stein. Infinitely many hyperbolic 3-manifolds which contain no reebless foliation. *Journal of the American Mathematical Society*, 16(3):639–679, 2003.
- [Sta78] John Stallings. Constructions of fibred knots and links. In *Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part*, volume 2, pages 55–60, 1978.
- [Thu86] William P. Thurston. A norm for the homology of 3-manifolds. *Memoirs of the American Mathematical Society*, 59(339):99–130, 1986.
- [Tur12] Vladimir Turaev. *Torsions of 3-dimensional manifolds*, volume 208. Birkhäuser, 2012.
- [Zun20] Jonathan Zung. Taut foliations, left-orders, and pseudo-Anosov mapping tori. *arXiv preprint arXiv:2006.07706*, 2020.

SCUOLA NORMALE SUPERIORE, PIAZZA DEI CAVALIERI 7, PISA
 Email address: `diego.santoro@sns.it`