

AN ALGEBRAIC PROPERTY OF REIDEMEISTER TORSION

TERUAKI KITANO AND YUTA NOZAKI

ABSTRACT. For a 3-manifold M and an acyclic $SL(2, \mathbb{C})$ -representation ρ of its fundamental group, the $SL(2, \mathbb{C})$ -Reidemeister torsion $\tau_\rho(M) \in \mathbb{C}^\times$ is defined. If there are only finitely many conjugacy classes of irreducible representations, then the Reidemeister torsions are known to be algebraic numbers. Furthermore, we prove that the Reidemeister torsions are not only algebraic numbers but also algebraic integers for most Seifert fibered spaces and infinitely many hyperbolic 3-manifolds. Also, for a knot exterior $E(K)$, we discuss the behavior of $\tau_\rho(E(K))$ when the restriction of ρ to the boundary torus is fixed.

CONTENTS

1. Introduction	1
2. Preliminaries	5
3. Reidemeister torsion and algebraic integers	9
4. Reidemeister torsion of Seifert fibered spaces	16
5. Continuous variation of Reidemeister torsion	19
References	21

1. INTRODUCTION

Let M be a connected compact n -manifold and let ρ be an acyclic $SL(2, \mathbb{C})$ -representation, namely the chain complex $C_*(M; \mathbb{C}_\rho^2)$ is acyclic. Then the $SL(2, \mathbb{C})$ -Reidemeister torsion $\tau_\rho(M) \in \mathbb{C}^\times$ is defined to be the alternative product of determinants (see Section 2.1). When ρ is not acyclic, we set $\tau_\rho(M) = 0$. Then $\tau_\rho(M)$ defines a \mathbb{C} -valued function on the $SL(2, \mathbb{C})$ -representation variety $R(M) = \text{Hom}(\pi_1(M), SL(2, \mathbb{C}))$, which factors through the $SL(2, \mathbb{C})$ -character variety $X(M)$ of M . In this paper, we mainly consider a 3-manifold M and the subspace $X^{\text{irr}}(M)$ of irreducible characters. Under the assumption that $X^{\text{irr}}(M)$ is a finite set, Johnson [8] defined the

2020 *Mathematics Subject Classification*. Primary 57K31, 57Q10, Secondary 11R04, 13P15, 57M05.

Key words and phrases. Reidemeister torsion, algebraic integer, resultant, Chebyshev polynomial, character variety, A -polynomial.

torsion polynomial $\sigma_M(t)$ of M by

$$\sigma_M(t) = \prod_{[\rho] \in X^{\text{irr}}(M), \text{ acyclic}} (t - \tau_\rho(M)) \in \mathbb{C}[t].$$

He mentioned that $\sigma_M(t)$ lies in $\mathbb{Q}[t]$ by considering the action of a Galois group. As a consequence, $\tau_\rho(M)$ is an algebraic number when $X^{\text{irr}}(M)$ is a finite set. The first author and Tran [15] described the torsion polynomial of the Brieskorn homology 3-sphere $\Sigma(p, q, r)$ in terms of the normalized Chebyshev polynomial (see also [13]). Moreover, we show that $\sigma_{\Sigma(p, q, r)}(t)$ has integral coefficients in Section 4. In other words, $\tau_\rho(\Sigma(p, q, r))$ is not only an algebraic number but also an algebraic integer for every irreducible representation ρ . Recall that $\alpha \in \mathbb{C}$ is called an *algebraic integer* if there is a monic polynomial over \mathbb{Z} such that α is a root of the polynomial. As a consequence, for the homology 3-sphere obtained by Dehn surgery along the (right-handed) (p, q) -torus knot $T_{p, q}$, its Reidemeister torsion $\tau_\rho(S_{-1/n}^3(T_{p, q}))$ is also an algebraic integer since $\Sigma(p, q, pqn + 1) = S_{-1/n}^3(T_{p, q})$ for $n > 0$.

Also, such a phenomenon was numerically observed for the figure-eight knot 4_1 in [12]. It is worth mentioning that while the Brieskorn homology 3-spheres are not hyperbolic, $S_{p/q}^3(4_1)$ is hyperbolic unless p/q is an integer with $|p/q| \leq 4$ or $p/q = \infty$. In this paper, we rigorously prove the observation in [12].

Theorem 1.1. *Let p, q be non-zero integers and either p or q is 1. For an $SL(2, \mathbb{C})$ -representation ρ of $\pi_1(S_{p/q}^3(4_1))$, the $SL(2, \mathbb{C})$ -Reidemeister torsion $\tau_\rho(S_{p/q}^3(4_1))$ is an algebraic integer.*




FIGURE 1. The twist knot $J(2, 2m)$ and $J(2, 2) = 3_1$, where $2m$ half-twists mean $|2m|$ negative half-twists if $m < 0$.

Furthermore, an analogous result holds for the complement of a twist knot $J(2, 2m)$ in Figure 1 ($m \in \mathbb{Z}$), which suggests that the Reidemeister torsions of the closed 3-manifold $S_{p/q}^3(J(2, 2m))$ might be algebraic integers as observed in Examples 3.10 and 3.11. Throughout this paper, for a knot K , let $E(K)$ denote the complement of an open tubular neighborhood of K . We also write μ and λ for a meridian and a preferred longitude of K , respectively.

Theorem 1.2. *Let K be a twist knot $J(2, 2m)$ and let $p = \pm 1$ and q an odd integer. For an irreducible $SL(2, \mathbb{C})$ -representation ρ of $\pi_1(E(K))$ satisfying $\rho(\mu^p \lambda^q) = I_2$, the $SL(2, \mathbb{C})$ -Reidemeister torsion $\tau_\rho(E(K))$ is an algebraic integer.*

Note that $J(2, 2)$ is a trefoil knot $3_1 = T_{-3,2}$, and $\tau_\rho(E(3_1))$ was computed by Johnson [8]. In fact, he expressed $\tau_\rho(E(T_{p,q}))$ in terms of cosine functions. In Section 4, we see that these are algebraic integers as well. We give the proofs of Theorems 1.1 and 1.2 in Section 3, which is based on the resultant of polynomials and the A -polynomial of knots introduced by Cooper, Culler, Gillet, Long, and Shalen [1].

Here we refer to some related results. First recall that the Reidemeister torsion $\tau_\rho(E(K))$ coincides with $\Delta_{K,\rho}(1)$, where $\Delta_{K,\rho}(t) \in \mathbb{C}[t^{\pm 1}]$ denotes the twisted Alexander polynomial associated with ρ (see [22], [11]). When K is hyperbolic and ρ is the holonomy representation, Dunfield, Friedl, and Jackson [4] called $\Delta_{K,\rho}(t)$ the hyperbolic torsion polynomial. They observed that the coefficients of hyperbolic torsion polynomial are often algebraic integers. This observation and Theorems 1.1 and 1.2 suggest that, under some conditions, Reidemeister torsions have an interesting algebraic property that is quite non-trivial from its definition. Recently, Yoon [23] proved a vanishing identity on the adjoint Reidemeister torsions of 2-bridge knots. This result also implies that Reidemeister torsions are imposed on strong algebraic restrictions. Moreover, we observe that the maximal $SL(2, \mathbb{C})$ -Reidemeister torsion in absolute value is often a Perron number in Examples 3.10 and 3.11. Here, a real algebraic number $\alpha > 1$ is called a *Perron number* if it is larger than the absolute values of the Galois conjugates of α . Such numbers appear, for example, in the study of the stretch factor of pseudo-Anosov homeomorphisms (see [5, Section 14.2.1]).




FIGURE 2. The knot K_0 .

We now turn our attention to the case where $X^{\text{irr}}(M)$ is an infinite set. For instance, if $M = E(K)$ for a knot K in S^3 , then $X^{\text{irr}}(M)$ has a positive dimension. The first author [10] gave the formula $\tau_\rho(E(4_1)) = 2 - 2 \operatorname{tr} \rho(\mu)$, where $\operatorname{tr} \rho(\mu) \neq 2$ for a meridian μ . Note that $\tau_\rho(E(4_1))$ is no longer an algebraic integer since $\operatorname{tr} \rho(\mu)$ can vary continuously. On the other hand,

$\tau_\rho(E(4_1))$ is determined by the restriction $r(\rho)$ of ρ to the boundary torus. Here, when $\partial M \neq \emptyset$, we denote by r the regular map $X(M) \rightarrow X(\partial M)$ between algebraic sets induced by the inclusion $\partial M \hookrightarrow M$. It is natural to ask whether the function $\tau_\rho(M)$ on $X(M)$ varies continuously while $r(\rho)$ is fixed. If K is a 2-bridge knot, then $\dim_{\mathbb{C}} X(E(K)) = 1$ and $r^{-1}(\rho_0)$ is a finite set, and thus $\tau_\rho(M)$ cannot vary continuously. We need to consider the case $\dim_{\mathbb{C}} X(E(K)) \geq 2$. An easy example of such a knot is the connected sum $K = K_1 \sharp K_2$ of non-trivial knots K_1 and K_2 . However, if $r(\rho)$ is fixed, then $\tau_\rho(E(K))$ depends only on $\tau_\rho(E(K_1))$ and $\tau_\rho(E(K_2))$ because of the multiplicativity of the Reidemeister torsions. In this paper, we focus on the knot K_0 in Figure 2, which is obtained from $4_1 \sharp 4_1$ by a construction given by Cooper and Long [2]. Then we can find a family C of representations of $\pi_1(E(K_0))$ and obtain the following result.

Theorem 1.3. *Let $\rho' \in C$. Then the function $\tau_\rho(E(K_0))$ varies continuously on the preimage $r^{-1}(r(\rho')) \subset X(E(K_0))$.*

Note that the knot K_0 is alternating, hyperbolic, and fibered (see Section 5). Theorem 1.3 is motivated by the authors' previous work [14]. They investigated the set

$$RT(M) = \{\tau_\rho(M) \mid [\rho] \in X^{\text{irr}}(M), \text{ acyclic}\} \subset \mathbb{C}$$

of all values of the Reidemeister torsion for irreducible representations. In [14], the authors proved that for 2-bridge knots K_1 and K_2 , the set $RT(\Sigma(K_1, K_2))$ is finite, while $X^{\text{irr}}(\Sigma(K_1, K_2))$ has positive dimension. Here the splice $\Sigma(K_1, K_2)$ is the closed 3-manifold $E(K_1) \cup_h E(K_2)$, where h is an orientation-reversing homeomorphism sending a meridian (resp. longitude) of K_1 to a longitude (resp. meridian) of K_2 . The proof is based on the fact that $\tau_\rho(E(K_j))$ cannot vary continuously if $r_j(\rho)$ is fixed ($j = 1, 2$) as mentioned before. In contrast, Theorem 1.3 implies the following consequence.

Corollary 1.4. *Let K be a 2-bridge knot such that polynomials $f_C(L, M)$ and $A_K(M, L) \in \mathbb{Z}[L, M]$ have a common zero (L_0, M_0) with $L_0, M_0 \neq 0$ and with $L_0 \neq \pm 1$ or $M_0 \neq \pm 1$, where A_K is the A -polynomial of K . Then the set $RT(\Sigma(K_0, K))$ is an infinite set.*

Here, $f_C(L, M) \in \mathbb{Z}[L, M]$ is the polynomial

$$\begin{aligned} &L^2 M^{16} - L((M^{32} + 1) - 4(M^{30} + M^2) - 2(M^{28} + M^4) + 16(M^{26} + M^6) \\ &+ 13(M^{24} + M^8) - 32(M^{22} + M^{10}) - 46(M^{20} + M^{12}) + 20(M^{18} + M^{14}) + 70M^{16}) + M^{16}, \end{aligned}$$

which is derived from $r(C) \subset X(\partial E(K_0))$. Note that the condition on K in Corollary 1.4 is generic enough. For instance, 3_1 and 4_1 satisfy the condition.

Acknowledgments. This work was supported by JSPS KAKENHI Grant Numbers JP19K03505 and JP20K14317.

2. PRELIMINARIES

2.1. $SL(2, \mathbb{C})$ -Reidemeister torsion and the torsion polynomial. Let M be a connected compact 3-manifold whose boundary is empty or a disjoint union of tori and let $\rho: \pi_1(M) \rightarrow SL(2, \mathbb{C})$ be a representation, namely a group homomorphism. Suppose ρ is acyclic, that is, $H_*(M; \mathbb{C}_\rho^2) = 0$. We first endow M with a cell decomposition so that M is a CW-complex. This gives the cellular chain complex $\{C_*(M; \mathbb{C}_\rho^2), \partial_*\}$ with an (ordered) basis \mathbf{c}_i of $C_i(M; \mathbb{C}_\rho^2)$ coming from i -cells. We next choose a basis \mathbf{b}_i of $\text{Im } \partial_{i+1}$ and its lift $\tilde{\mathbf{b}}_i$ to $C_{i+1}(M; \mathbb{C}_\rho^2)$. It follows from $H_*(M; \mathbb{C}_\rho^2) = 0$ that the union $\mathbf{b}_i \tilde{\mathbf{b}}_{i-1}$ is a basis of $C_i(M; \mathbb{C}_\rho^2)$. Let $[\mathbf{b}_i \tilde{\mathbf{b}}_{i-1} / \mathbf{c}_i]$ denote the change of basis matrix from \mathbf{c}_i to $\mathbf{b}_i \tilde{\mathbf{b}}_{i-1}$. Now, the $SL(2, \mathbb{C})$ -Reidemeister torsion (or simply *Reidemeister torsion*) $\tau_\rho(M)$ of M associated to ρ is defined by

$$\tau_\rho(M) = \prod_{i=0}^3 \det[\mathbf{b}_i \tilde{\mathbf{b}}_{i-1} / \mathbf{c}_i]^{(-1)^{i+1}} \in \mathbb{C}^\times.$$

See [8] and Section 1 of [9, 10] for details. For instance, when M is the complement of a 2-bridge knot K , the fundamental group $\pi_1(M)$ has a presentation with generators x, y and a relator r . Then, for an acyclic representation ρ , one can derive the formula

$$\tau_\rho(M) = \frac{\det(\rho(\partial r / \partial y))}{\det(\rho(x) - I_2)},$$

where $\partial r / \partial y$ denotes a Fox derivative of r .

We next consider the $SL(2, \mathbb{C})$ -character variety $X(M)$, which is the GIT quotient of the $SL(2, \mathbb{C})$ -representation variety $R(M) = \text{Hom}(\pi_1(M), SL(2, \mathbb{C}))$. Let $X^{\text{irr}}(M)$ denote the subspace of $X(M)$ consisting of conjugacy classes of irreducible representations. In [8, p. 54], when $X^{\text{irr}}(M)$ is a finite set, the *torsion polynomial* $\sigma_M(t)$ of M is defined by

$$\sigma_M(t) = \prod_{[\rho] \in X^{\text{irr}}(M), \text{acyclic}} (t - \tau_\rho(M)) \in \mathbb{Q}[t].$$

Remark 2.1. Reidemeister torsion is sometimes defined to be $\prod_{i=0}^3 \det[\mathbf{b}_i \tilde{\mathbf{b}}_{i-1} / \mathbf{c}_i]^{(-1)^i}$. Our $\tau_\rho(M)$ is the same as Reidemeister torsion in [8, 12, 15], but the inverse of Reidemeister torsion in [9, 10]. This difference is crucial for the results in this paper. For instance, $\tau_\rho(S_1^3(4_1))$ is an algebraic integer, but its inverse is not since the constant term of $\sigma_{S_1^3(4_1)}(t) = t^3 - 12t^2 + 20t - 8$ is not ± 1 .

2.2. The A -polynomial of knots. Let K be a knot in S^3 and recall that $E(K)$ denotes the complement of an open tubular neighborhood of K . We consider the algebraic subset U of $R(E(K))$ consisting of representations ρ such that $\rho(\lambda)$ and $\rho(\mu)$ are upper triangular matrices, where λ and μ denote a preferred longitude and a meridian, respectively. Now we define a map $\xi: U \rightarrow \mathbb{C}^2$ by $\xi(\rho) = (\rho(\lambda)_{11}, \rho(\mu)_{11})$, where X_{ij} denotes the (i, j) -entry of a matrix X . The Zariski closure of the image of ξ is some algebraic curves

and points. The product of the defining polynomials of the algebraic curves is known to be defined over \mathbb{Z} so that the coefficients are relatively prime. The resulting polynomial $A_K(L, M) \in \mathbb{Z}[L, M]$ is called the *A-polynomial* of K , which is defined up to sign. Following [1], we drop the factor $L - 1$ corresponding to abelian representations from the *A-polynomial*, that is, $A_{\text{unknot}}(L, M) = 1$.

For 2-bridge knots, their *A-polynomials* are computed from Riley polynomials (see [1, Section 7] and [2, Section 3]). Here we recall the definition of the Riley polynomial introduced in [21]. Let K be a 2-bridge knot. Then $\pi_1(E(K))$ has a presentation of the form $\langle x, y \mid wx = yw \rangle$, where w is a word in x and y . For $s, t \in \mathbb{C}$ with $s \neq 0$, we consider the representation $\rho_{s,t}$ of the free group $\langle x, y \rangle$ by

$$\rho_{s,t}(x) = \begin{pmatrix} s & 1 \\ 0 & 1/s \end{pmatrix}, \quad \rho_{s,t}(y) = \begin{pmatrix} s & 0 \\ -t & 1/s \end{pmatrix}$$

and define the *Riley polynomial* $\phi(s, t)$ of K by

$$\phi(s, t) = \rho_{s,t}(w)_{11} + (s - s^{-1})\rho_{s,t}(w)_{12} \in \mathbb{Z}[s^{\pm 1}, t].$$

It is shown in [21, Theorem 1] that $\rho_{s,t}$ factors through $\pi_1(E(K))$ if and only if $\phi(s, t) = 0$ holds. Moreover, every non-abelian representation is conjugate to some $\rho_{s,t}$. Note that $\rho_{s,t}$ is irreducible if and only if $t \neq 0$.

The *A-polynomial* of a 2-bridge knot is obtained from its Riley polynomial by eliminating t . More precisely, we consider the resultant

$$\text{res}_t(L - \rho_{M,t}(\lambda)_{11}, \phi(M, t)) \in \mathbb{Z}[L, M^{\pm 1}]$$

of polynomials in t , and the *A-polynomial* is a factor of this resultant. See Section 2.3 for the definition of the resultant.

Remark 2.2. The variable M in $A_K(L, M)$ can be taken as the variable s in $\phi(s, t)$ for a 2-bridge knot K . In this paper, however, we use a pair (L, M) for $A_K(L, M)$ and (s, t) for $\phi(s, t)$ under the conventions of these polynomials.

Example 2.3. Let us consider the case $K = J(2, 4) = 5_2$. Then $w = [y, x^{-1}]^2$, $\mu = x$, and $\lambda = [x, y^{-1}]^2[y, x^{-1}]^2$. Thus, we have

$$\begin{aligned} \phi(s, t) &= (2(s^2 + s^{-2}) - 3)t^2 + (-(s^4 + s^{-4}) + 3(s^2 + s^{-2}) - 6)t + 2(s^2 + s^{-2}) - t^3 - 3, \\ A_{5_2}(L, M) &= -L^3 + L^2(1 - 2M^2 - 2M^4 + M^8 - M^{10}) \\ &\quad + LM^4(-1 + M^2 - 2M^6 - 2M^8 + M^{10}) - M^{14}. \end{aligned}$$

Let us give an observation. Since $\phi(i, t) = -t^3 - 7t^2 - 14t - 7$ and $A_{5_2}(L, i) = (L - 1)^3$, there are three irreducible representations of $\pi_1(E(5_2))$ such that the restrictions to $\pi_1(\partial E(5_2))$ are identical. However, one can check that $\tau_{\rho_{i,t}}(E(5_2)) = (-t^3 - 3t^2 + 2t + 9)/2$, and thus $\tau_{\rho_{i,t}}(E(5_2))$'s are distinct for the three representations. That is, the Reidemeister torsion $\tau_\rho(E(5_2))$ is not necessarily determined by the restriction $r(\rho) = \rho|_{\partial E(5_2)}$. On the other hand, $\tau_\rho(E(5_2))$ can only take finitely many values while $r(\rho)$ is fixed. In particular, it does not vary continuously.

times the i th row to the $(i-1)$ st row for $i = 2, 3, \dots, n, n+2, n+3, \dots, m+n$:

$$\begin{aligned} \operatorname{res}_x(f, g) &= \begin{vmatrix} a_0 & a_0\zeta + a_1 & \cdots & & f(\zeta) & \cdots & \zeta^{n-1}f(\zeta) \\ & a_0 & & \cdots & & f(\zeta) & \\ & & \ddots & \ddots & & \ddots & \ddots \\ b_0 & b_0\zeta + b_1 & \cdots & a_0 & a_0\zeta + a_1 & \cdots & f(\zeta) \\ & b_0 & & \cdots & g(\zeta) & \cdots & \zeta^{m-1}g(\zeta) \\ & & \ddots & \ddots & & \ddots & \ddots \\ & & & b_0 & b_0\zeta + b_1 & \cdots & g(\zeta) \end{vmatrix} \\ &= \begin{vmatrix} a_0 & a_1 & \cdots & a_{m-1} & a_m & & & \\ & a_0 & a_1 & \cdots & a_{m-1} & a_m & & \\ & & \ddots & \ddots & & \ddots & \ddots & \\ & & & a_0 & a_0\zeta + a_1 & \cdots & & f(\zeta) \\ b_0 & b_1 & \cdots & b_{n-1} & b_n & & & \\ & b_0 & b_1 & \cdots & b_{n-1} & b_n & & \\ & & \ddots & \ddots & & \ddots & \ddots & \\ & & & b_0 & b_0\zeta + b_1 & \cdots & & g(\zeta) \end{vmatrix}. \end{aligned}$$

Since $g(\zeta)^m \mid f(\zeta)$, one can divide the $(m+n)$ th column by $g(\zeta)$. We then sweep out the bottom row by 1 at the bottom left:

$$\begin{aligned} \operatorname{res}_x(f, g) &= g(\zeta) \begin{vmatrix} a_0 & a_1 & \cdots & a_{m-1} & a_m & & & \\ & a_0 & a_1 & \cdots & a_{m-1} & a_m & & \\ & & \ddots & \ddots & & \ddots & \ddots & \\ & & & a_0 & a_0\zeta + a_1 & \cdots & & f(\zeta)g(\zeta)^{-1} \\ b_0 & b_1 & \cdots & b_{n-1} & b_n & & & \\ & b_0 & b_1 & \cdots & b_{n-1} & b_n & & \\ & & \ddots & \ddots & & \ddots & \ddots & \\ & & & b_0 & b_0\zeta + b_1 & \cdots & & 1 \end{vmatrix} \\ &= g(\zeta) \begin{vmatrix} a_0 & a_1 & \cdots & a_{m-1} & a_m & & & \\ & \ddots & \ddots & & \ddots & \ddots & & \\ & & a_0 & a_1 & \cdots & a_{m-1} & a_m & \\ b_0 & b_1 & \cdots & b_{n-1} & b_n & & & f(\zeta)g(\zeta)^{-1} \\ & \ddots & \ddots & & \ddots & \ddots & & \\ & & b_0 & b_1 & \cdots & b_{n-1} & b_n & \\ & & & 0 & 0 & \cdots & 0 & 1 \end{vmatrix}, \end{aligned}$$

Proof. First recall that $\phi(s, t) \in \mathbb{Z}[s^{\pm 1}, t]$ denotes the Riley polynomial of K . Since $A_K(L, M)$ is a factor of the resultant

$$f(L, M) = \text{res}_t(\phi(M, t), L - \rho_{M,t}(\lambda)_{11}) \in \mathbb{Z}[L, M^{\pm 1}],$$

it suffices to prove that the leading term of $f(L, M)$ with respect to L is a unit. By [21, Lemma 2], the coefficient of the leading term of $\phi(M, t)$ with respect to t is ± 1 . It follows from the definition of the resultant that the leading term of $f(L, M)$ with respect to L is a power of L up to sign.

Now, the latter follows from the equality $A_K(L, M) = A_K(L^{-1}, M^{-1})$ up to units in $\mathbb{Z}[L, M]$ (see [2, Proposition 4.2(1)]). \square

Before the next lemma, we recall a boundary slope of a knot K . For an incompressible surface S in $E(K)$, its boundary consists of parallel simple closed curves on $\partial E(K)$. Thus, S defines an element $p[\mu] + q[\lambda] \in H_1(\partial E(K); \mathbb{Z})$ up to sign, and then $p/q \in \mathbb{Q} \cup \{1/0\}$ is called a *boundary slope* of K . It is proved in [1, Theorem 3.4] that the slope of a side of the Newton polygon $N(A_K)$ of $A_K(L, M)$ is a slope of K . Combining this theorem and [6, Theorem 1(b)], we have that slopes of sides of $N(A_K)$ are even integers for a 2-bridge knot K .

Lemma 3.2. *Let K be a 2-bridge knot and let $p = \pm 1$ and $q > 0$. Then the coefficients of the highest and lowest degree terms of $\text{res}_L(A_K(L, M), M^p L^q - 1) \in \mathbb{Z}[M^{\pm 1}]$ are ± 1 .*

Proof. It suffices to discuss only the highest degree term of the resultant since $A_K(L, M) = A_K(L^{-1}, M^{-1})$ up to units in $\mathbb{Z}[L, M]$. Let $d = \deg_L A_K(L, M)$ and suppose $p = -1$. Let $cL^i M^j$ be the highest degree term of $A_K(L, M)$ with respect to the lexicographic order of (\deg_M, \deg_L) . Since this monomial corresponds to a corner v of the Newton polygon $N(A_K)$, [2, Theorem 11.3] implies $c = \pm 1$. When we choose cM^j 's in the first $|q|$ rows of the matrix appearing in the resultant, the highest degree term is $\pm (cM^j)^q (M^{-1})^{d-i} 1^i = \pm M^{jq-d+i}$. Now, let l be the line through v with slope $p = -1$. Since the slopes of the sides of $N(A_K)$ are even, $N(A_K)$ lies in the lower left half space. Therefore, all terms in the resultant except $\pm M^{jq-d+i}$ have a lower degree.

In the case $p = 1$, we can show the statement in a similar way by considering the lexicographic order of $(\deg_M, -\deg_L)$. \square

As a consequence of Lemma 3.2, if ρ_{s_0, t_0} is a representation satisfying the condition of Theorem 1.2, then s_0 and s_0^{-1} are algebraic integers.

Lemma 3.3. *For a root (s_0, t_0) of $\phi(s, t)$, if $s_0, s_0^{-1} \in \mathbb{A}$, then $t_0 \in \mathbb{A}$.*

Proof. By [21, Lemma 2], $\phi(s_0, t) \in \mathbb{A}[t]$ is monic. Since the ring \mathbb{A} of algebraic integers is integrally closed in \mathbb{C} , we conclude $t_0 \in \mathbb{A}$. \square

In the rest of this section, let $K = J(2, 2m)$ ($m \neq 0$). Define $d(m) = \deg_t \phi(1, t)$, then by [21, Lemma 2], we have

$$d(m) = \deg_t \phi(s, t) = \begin{cases} 2m - 1 & \text{if } m > 0, \\ -2m & \text{if } m < 0. \end{cases}$$

Lemma 3.4. $\frac{1}{n!} \frac{\partial^n A_K}{\partial L^n}(-1, M)$ is divisible by $(M^2 - 1)^{d(m)-n}$ for $0 \leq n < d(m)$.

Proof. The proof is by induction on m . The cases $m = -1, 0, 1, 2$ are directly confirmed by [7, Theorem 1]. For instance, in the case $m = 2$, we have

$$\begin{aligned} A_K(-1, M) &= (M^2 - 1)^3 (2M^8 + 4M^6 + 5M^4 + 4M^2 + 2), \\ \frac{1}{1!} \frac{\partial A_K}{\partial L}(-1, M) &= -(M^2 - 1)^2 (M^{10} - M^6 - 4M^4 - 6M^2 - 5), \\ \frac{1}{2!} \frac{\partial^2 A_K}{\partial L^2}(-1, M) &= (M^2 - 1) (M^8 + 2M^2 + 4). \end{aligned}$$

Suppose $m > 2$. Following [7], let

$$\begin{aligned} x(L, M) &= -L + L^2 + 2LM^2 + M^4 + 2LM^4 + L^2M^4 + 2LM^6 + M^8 - LM^8, \\ y(L, M) &= M^4(L + M^2)^4. \end{aligned}$$

Then one can check that $(M^2 - 1)^{2-k} \mid \frac{1}{k!} \frac{\partial^k x}{\partial L^k}(-1, M)$ for $0 \leq k \leq 2$ and $(M^2 - 1)^{4-k} \mid \frac{1}{k!} \frac{\partial^k y}{\partial L^k}(-1, M)$ for $0 \leq k \leq 4$. By the recursive relation in [7, Theorem 1] and Leibniz's rule, we have

$$\frac{1}{n!} \frac{\partial^n A_K}{\partial L^n} = \sum_{k=0}^2 \frac{1}{k!} \frac{\partial^k x}{\partial L^k} \frac{1}{(n-k)!} \frac{\partial^{n-k} A_{J(2, 2m-2)}}{\partial L^{n-k}} - \sum_{k=0}^4 \frac{1}{k!} \frac{\partial^k y}{\partial L^k} \frac{1}{(n-k)!} \frac{\partial^{n-k} A_{J(2, 2m-4)}}{\partial L^{n-k}}.$$

When $L = -1$, by the induction hypothesis, each term in the sums is divisible by $(M^2 - 1)^{2m-1-n}$.

The cases $m < -1$ also follow from the recursive relation in a similar way. \square

Proposition 3.5. Let $p = \pm 1$. Then, $\text{res}_L(A_K(L, M), M^p L^q - 1)$ is divisible by $(M^p(-1)^q - 1)^{d(m)}$.

Proof. Note that $M^p(-1)^q - 1 \mid M^2 - 1$ when $p = \pm 1$. Now, Lemma 3.4 allows us to apply Lemma 2.4 to $\zeta = -1 \in \mathbb{C}[M]$ and

$$f(L, M) = A_K(L, M), \quad g(L, M) = ML^q - 1 \in \mathbb{C}[M][L] = \mathbb{C}[L, M].$$

This completes the proof. \square

Here recall that $\tau_\rho(E(K))$ is written of the form

$$\frac{\det(\rho_{s_0, t_0}(\partial r / \partial y))}{\det(\rho_{s_0, t_0}(x) - I_2)} = -s_0(s_0 - 1)^{-2} F(s_0, t_0),$$

where $F(s, t) \in \mathbb{Z}[s^{\pm 1}, t]$. In the situation of Theorem 1.2, s_0 and $F(s_0, t_0)$ are algebraic integers by Lemmas 3.2 and 3.3. Hence, it suffices for proving the theorem to consider whether $(s_0 - 1)^{-1}$ is an algebraic integer.

Lemma 3.6. *Let $f(x) \in \mathbb{Z}[x]$ be a monic polynomial and let $\alpha \in \mathbb{C} \setminus \{1\}$ be a root of f . If $f(1) = \pm 1$, then $(\alpha - 1)^{-1}$ is an algebraic integer.*

Proof. Let $\beta = (\alpha - 1)^{-1}$, namely $\alpha = \beta^{-1} + 1$. Since f is monic, we have

$$0 = f(\alpha) = f(\beta^{-1} + 1) = \beta^{-\deg f} + \dots + f(1).$$

Here, by $f(1) = \pm 1$, there exists a monic polynomial $g \in \mathbb{Z}[x]$ such that $g(\beta) = 0$. \square

Proof of Theorem 1.2. Recall that q is odd. Let us show that $(s_0 - 1)^{-1} \in \mathbb{A}$. By Lemma 3.6, it suffices to find a monic polynomial $f(s)$ such that $f(s_0) = 0$ and $f(1) = \pm 1$. Proposition 3.5 implies that

$$f(s) := (s + 1)^{-d(m)} \operatorname{res}_L(A_K(L, s), sL^q - 1)$$

lies in $\mathbb{Z}[s^{\pm 1}]$ and it is monic. By [7, Corollary 3(i)], we have

$$\begin{aligned} f(1) &= (1 + 1)^{-d(m)} \operatorname{res}_L(\pm(L + 1)^{d(m)}, L^q - 1) \\ &= \pm 2^{-d(m)} \operatorname{res}_L(L + 1, L^q - 1)^{d(m)} = \pm 1. \end{aligned}$$

Therefore, $(s_0 - 1)^{-1}$ is an algebraic integer. \square

Remark 3.7. When q is even, $(s_0 - 1)^{-1}$ is not necessarily an algebraic integer. For example, in the case where $J(2, -2) = 4_1$ and $(p, q) = (1, 2)$, s_0 's are the roots of the polynomial

$$f(x) = x^{14} + 2x^{13} + x^{12} - 4x^{10} - 8x^9 - 10x^8 - 13x^7 - 10x^6 - 8x^5 - 4x^4 + x^2 + 2x + 1.$$

Here, $f(x)$ is irreducible since it is the product of two irreducible polynomials of degree 7 over the prime field \mathbb{F}_2 of order 2 and is the product of two irreducible polynomials of degree 4 and 10 over \mathbb{F}_3 . Thus, if $(s_0 - 1)^{-1} \in \mathbb{A}$, then $f(1) = \pm 1$. However, we have $f(1) = -49$.

On the other hand, $\tau_\rho(E(4_1)) = -2(s_0 + s_0^{-1} - 1)$ is an algebraic integer since $s_0^{\pm 1} \in \mathbb{A}$. The authors predict that Theorem 1.2 holds for every integer q .

3.2. The proof of Theorem 1.1. This subsection is devoted to proving that $\tau_\rho(S_{p/q}^3(4_1))$ is an algebraic integer when $p = 1$ or $q = 1$. We also observe that it is an algebraic integer when $p/q = 2/3$ in Example 3.10. Moreover, $\tau_\rho(S_{1/2}^3(5_2))$ is an algebraic integer as well (see Example 3.11 and Problem 3.9).

Let $q \neq 0$ and let ρ be an acyclic representation of $\pi_1(S_{p/q}^3(4_1))$. Then

$$\tau_\rho(S_{p/q}^3(4_1)) = \frac{-2(s + s^{-1} - 1)}{\det(\rho(\mu^{p'} \lambda^{q'}) - I_2)} = 2(s + s^{-1} - 1) s^{p'} L^{q'} (s^{p'} L^{q'} - 1)^{-2},$$

where $pq' - qp' = 1$ and $L = \rho(\lambda)_{11}$. We divide the proof of Theorem 1.1 into two cases: (I) $p = 1, q \neq 0$ and (II) $p \neq 0, q = 1$. Recall that

$$\begin{aligned} A_{4_1}(L, M) &= L^2 M^4 + L(-M^8 - 1 + M^6 + M^2 + 2M^4) + M^4 \\ &= (L^2 + L(-(M + M^{-1})^4 + 5(M + M^{-1})^2 - 2) + 1) M^4. \end{aligned}$$

Proof of Theorem 1.1(I). Let $p = 1, q \neq 0$. Then one can take $(p', q') = (0, 1)$. By Lemmas 3.2 and 3.3, it suffices to show that $(L_0 - 1)^{-1}$ is an algebraic integer. First, by direct computations, we have $\text{tr } \rho(\mu) \neq -2$. Indeed, if $s = -1$, then

$$\rho(\lambda) = \begin{pmatrix} -1 & \pm 2i\sqrt{3} \\ 0 & -1 \end{pmatrix},$$

and hence $\rho(x)\rho(\lambda)^q \neq I_2$. See also [15, Lemma 6.2].

Let $f(L) = A_{4_1}(L, L^{-q})$ and $g(L) = L^{-q} + L^q \in \mathbb{Z}[L^{\pm 1}]$. Then, we have $f(-1) = 0$ and

$$\begin{aligned} f'(L) &= (2L + (-g(L)^4 + 5g(L)^2 - 2) + L(-4g'(L)g(L)^3 + 10g'(L)g(L))) L^{-4q} \\ &\quad - 4q(L^2 + L(-g(L)^4 + 5g(L)^2 - 2) + 1) L^{-4q-1}. \end{aligned}$$

It follows from $g'(-1) = 0$ that $f'(-1) = 0$, and hence $f(L) = (L + 1)^2 h(L)$ for some $h(L) \in \mathbb{Z}[L^{\pm 1}]$. Here, $h(L_0) = 0$ and $h(1) = 1$ since $L_0 \neq -1$ and $f(1) = 4$. Therefore, Lemma 3.6 implies $(L_0 - 1)^{-1} \in \mathbb{A}$. \square

Next, we discuss the case (II), that is, $q = 1$. Then one can take $(p', q') = (-1, 0)$. If $(s - 1)^{-1}$ is an algebraic integer, we can obtain the desired result. However, the case $p = 4$, we have $A_{4_1}(s^{-4}, s) = s^{-2}(s^2 + 1)^2$, and hence $s = \pm i$. The minimal polynomial of $(\pm i - 1)^{-1}$ is $2x^2 + 2x + 1$, and thus they are not algebraic integers. This observation suggests that one needs to show that $\tau_\rho(S_p^3(4_1)) = 2(s^2 - s + 1)/(s - 1)^2$ is an algebraic integer directly. Note that

$$A_{4_1}(s^{-p}, s) = s^{2-p} + 2s^{4-p} + s^{6-p} - s^{8-p} - s^{-p} + s^{4-2p} + s^4.$$

Lemma 3.8. *The coefficient of the leading term of*

$$\text{res}_s(\tau(s - 1)^2 - 2(s^2 - s + 1), A_{4_1}(s^{-p}, s)) \in \mathbb{Z}[\tau]$$

is equal to ± 16 . Moreover, it is divisible by $4(2\tau - 3)^2$ if p is odd and by 16 if p is even.

Proof. Let $n = \deg A_{4_1}(s^{-p}, s)$ and

$$\begin{aligned} h(\tau) &= \text{res}_s(\tau(s - 1)^2 - 2(s^2 - s + 1), A_{4_1}(s^{-p}, s)) \\ &= \begin{vmatrix} \tau - 2 & -2\tau + 2 & \tau - 2 & & & \\ & \ddots & & \ddots & & \\ & & \tau - 2 & -2\tau + 2 & \tau - 2 & \\ * & \cdots & & * & & \\ & * & \cdots & & * & \end{vmatrix}. \end{aligned}$$

Then, by the definition of the resultant,

$$\begin{aligned} \frac{1}{n!} \frac{d^n}{d\tau^n} h(\tau) &= \text{res}_s((s-1)^2, A_{4_1}(s^{-p}, s)) \\ &= \pm A_{4_1}(1, 1)^2 = \pm 16. \end{aligned}$$

When p is odd, we first note that $(s+1)^2 \mid A_{4_1}(s^{-p}, s)$. Thus, $h(\tau)$ is a multiple of $\text{res}_s(\tau(s-1)^2 - 2(s^2 - s + 1), (s+1)^2) = \pm(4\tau - 6)^2$.

We next consider the case p even. For the matrix in the resultant $h(\tau)$, we add the j th column to the $(j+1)$ st column for $j = 1, 2, \dots, n+1$ in this order twice. Then one obtains

$$\begin{vmatrix} \tau - 2 & -2 & -4 & \cdots & & -2n - 2 \\ & \tau - 2 & -2 & -4 & \cdots & -2n \\ & & \ddots & & & \vdots \\ & & & \tau - 2 & -2 & -4 \\ * & \cdots & & * & *' & *' \\ & & * & \cdots & * & *' \end{vmatrix},$$

where four $*'$'s are multiples of 4. Since τ 's appear only in the diagonal and the entries in the upper triangle are even, the coefficient of τ^k is a multiple of 16 unless $k = n-1, n-2, n-3$. Here note that there are even number of odd integers in the $(n+i)$ th row for $i = 1, 2$ since $2 \mid A_{4_1}(1, 1)$. Thus, the coefficient of τ^k is a multiple of 16 when $k = n-1, n-2$.

Now, the coefficients in $h(\tau)$ are multiples of 16 except the coefficient of τ^{n-3} . Hence, it suffices to see that $16 \mid h(1) = \text{res}_s(-s^2 - 1, A_{4_1}(s^{-p}, s))$. By a property of the resultant, we have $h(1) = \pm A_{4_1}(i^{-p}, i) A_{4_1}((-i)^{-p}, -i)$, which is equal to 0 or ± 16 . \square

Proof of Theorem 1.1(II). By Lemma 3.8, when p is odd, $4^{-1}(2\tau - 3)^{-2}h(\tau)$ is in $\mathbb{Z}[\tau]$ and monic. When p is even, $16^{-1}h(\tau)$ is in $\mathbb{Z}[\tau]$ and monic as well. Hence, we conclude that $\tau_\rho(S_p^3(K))$ is an algebraic integer. \square

We now pose the following problem.

Problem 3.9. It is natural to ask whether $\tau_\rho(S_{p/q}^3(4_1))$ is always an algebraic integer. More generally, whether $\tau_\rho(S_{p/q}^3(K))$ is an algebraic integer for any knot K with $\dim_{\mathbb{C}} X^{\text{irr}}(E(K)) = 1$.

We confirm that $\tau_\rho(S_{p/q}^3(K))$ is an algebraic integer for every irreducible representation ρ in the cases of $S_{2/3}^3(4_1)$ and $S_{1/2}^3(5_2)$.

Example 3.10. Let us consider the case $p/q = 2/3$. Then there are 12 representations of $\pi_1(S_{2/3}^3(4_1))$ as follows.

s	t	$\tau_\rho(S_{2/3}^3(4_1))$
$-0.200325 + 0.979729i$	-3.42754	0.738094
$0.200325 + 0.979729i$	-3.42754	5.85638
$-0.490393 + 0.871501i$	-2.21504	2.38654
$-1.30664 + 0.0498758i$	$-0.392004 + 0.724199i$	$5.8872 - 0.943648i$
$-0.264802 + 0.964303i$	-1.43838	8.4028
$0.490393 + 0.871501i$	-2.21504	0.0164217
$0.264802 + 0.964303i$	-1.43838	0.258749
$1.30664 + 0.0498758i$	$-0.392004 - 0.724199i$	$-0.717017 + 0.0236658i$
$-0.764207 + 0.0291705i$	$-0.392004 - 0.724199i$	$5.8872 + 0.943648i$
$0.764207 + 0.0291705i$	$-0.392004 + 0.724199i$	$-0.717017 - 0.0236658i$
-0.615146	-0.135036	1.60981
0.615146	-0.135036	94.3908

Using the resultant, one finds a polynomial $f(x)^2g(x)$ which is zero at these exact values of the Reidemeister torsion, where

$$f(x) = x^{12} - 124x^{11} + 3142x^{10} - 34792x^9 + 196796x^8 - 561760x^7 + 627280x^6 + 254848x^5 - 866240x^4 + 153088x^3 + 253696x^2 - 66560x + 1024$$

and $g(x)$ is a certain polynomial. We can confirm that $g(x) \neq 0$ for the above 12 values $\tau_\rho(S_{2/3}^3(4_1))$. Thus, they must be the zeros of f , and hence they are algebraic integers.

Since $f(94) < 0 < f(95)$, the zero of f approximated by 94.3908 must be a real number. This value is larger than the absolute values of the other 11 zeros. Hence, this zero is a Perron number. Here the corresponding representation is an $SL(2, \mathbb{R})$ -representation. Indeed, s 's and t 's are respectively zeros of the polynomials

$$s^{24} - 3s^{22} - 3s^{20} + 8s^{18} + 12s^{16} - 7s^{14} - 20s^{12} - 7s^{10} + 12s^8 + 8s^6 - 3s^4 - 3s^2 + 1,$$

$$t^6 + 8t^5 + 23t^4 + 31t^3 + 23t^2 + 10t + 1.$$

They have zeros between 0.6 and 0.7 and between -0.2 and -0.1 , respectively. Therefore, one of the $SL(2, \mathbb{R})$ -representations gives a Perron number. Such a phenomenon is numerically seen in [12] as well. Also, the Reidemeister torsions of the Brieskorn homology 3-sphere $\Sigma(p, q, r)$ are real numbers (see Section 4). Thus, the maximal value $(4 \sin \frac{\pi}{2p} \sin \frac{\pi}{2q} \sin \frac{\pi}{2r})^{-2}$ is a Perron number. Note that $\pi_1(\Sigma(2, 3, 5))$ does not admit any $SL(2, \mathbb{R})$ -representation and the maximal Reidemeister torsion comes from an $SU(2)$ -representation (see [16, Remark 3.4]).

Example 3.11. There are 17 representations of $\pi_1(S_{1/2}^3(5_2))$ and three of them are not acyclic:

s	t	$\tau_\rho(S_{1/2}^3(5_2))$
$-0.471842 + 0.881683i$	-2.87048	2.81243
$0.165381 + 0.98623i$	-3.68825	12.3508
$-0.200082 + 0.979779i$	-2.2146	0.490587
$0.0681942 + 0.997672i$	-2.41933	0.313753
$1.29286 + 1.35876i$	$-0.182462 - 0.461334i$	$9.57556 - 0.520417i$
$-1.26355 + 0.363134i$	$0.158863 + 1.07159i$	$3.6856 - 0.147423i$
$-0.348139 + 0.937443i$	-1.1692	4.5522
1	$0.21508 - 1.30714i$	0
1	$0.21508 + 1.30714i$	0
1	0.56984	0
$-0.859034 + 0.511919i$	-0.62449	7.42456
$0.313791 + 0.949492i$	-1.80681	0.0670363
$0.942666 + 0.333737i$	0.0622582	148.658
$0.709501 + 0.704704i$	-1.66753	1.06479
$-0.731039 + 0.210094i$	$0.158863 - 1.07159i$	$3.6856 + 0.147423i$
$0.367529 + 0.386263i$	$-0.182462 + 0.461334i$	$9.57556 + 0.520417i$
$-0.986232 + 0.16537i$	0.445642	5.74328

In the same manner as Example 3.10, we can show that 14 values of $\tau_\rho(S_{1/2}^3(5_2))$ are the zeros of the monic polynomial

$$\begin{aligned}
& x^{14} - 210x^{13} + 10760x^{12} - 269160x^{11} + 3993232x^{10} \\
& - 38203808x^9 + 245006784x^8 - 1067441024x^7 + 3141232640x^6 - 6091473408x^5 \\
& + 7422475264x^4 - 5260713984x^3 + 1942106112x^2 - 314212352x + 13778944.
\end{aligned}$$

Hence they are algebraic integers.

As similar to the previous example, the zero of the above polynomial approximated by 148.658 is a Perron number. The corresponding representation is numerically an $SL(2, \mathbb{R})$ -representation by considering the conjugate

$$P^{-1}\rho(-)P, \text{ where } P \approx \begin{pmatrix} -0.950489i & -1.97479 \\ 0.341848 & 0.341848i \end{pmatrix}.$$

4. REIDEMEISTER TORSION OF SEIFERT FIBERED SPACES

In this section, we show that the $SL(2, \mathbb{C})$ -Reidemeister torsion $\tau_\rho(M)$ is an algebraic integer for a Seifert fibered space M under a mild condition, which any Brieskorn homology 3-sphere satisfies.

4.1. Chebyshev polynomials and variants. We use Chebyshev polynomials and variants to prove the algebraic integrality. The next lemma follows from the fact that $\cos \frac{2k-1}{2n}\pi = \sin \frac{n-2k+1}{2n}\pi$ ($k = 1, 2, \dots, n$) are the roots of the Chebyshev polynomial of the first kind $T_n(x)$ defined by $T_n(\cos \theta) = \cos n\theta$. Note that $T_{2n}(x)$ has the form of $2^{2n-1}x^{2n} + \dots + (-1)^n \in \mathbb{Z}[x]$.

Lemma 4.1. *Let a be a positive even integer. Then $(\sin \frac{k}{2a}\pi)^{-1}$ is an algebraic integer for $k = -a + 1, -a + 3, \dots, a - 1$.*

Proof. As mentioned above, $\sin \frac{k}{2a}\pi$ gives a zero of $T_a(x)$. Then $(\sin \frac{k}{2a}\pi)^{-1}$ is a zero of $x^a T_a(1/x) = \pm x^a + \dots + 2^{a-1}$. \square

Here let us introduce a variant of the normalized Chebyshev polynomial of the second kind.

Definition 4.2. Define a polynomial $V_n(x) \in \mathbb{Z}[x]$ inductively by $V_0(x) = 1$, $V_1(x) = x - 1$, and $V_n(x) = xV_{n-1}(x) - V_{n-2}(x)$ for $n \geq 2$.

By definition, $V_n(x)$ is of the form $x^n + \dots \pm 1$.

Lemma 4.3. *Suppose $\theta \neq (2k - 1)\pi$ for any $k \in \mathbb{Z}$. Then $V_n(2 \cos \theta) = \cos(n + \frac{1}{2})\theta / \cos \frac{\theta}{2}$.*

Proof. Use induction on n . The cases $n = 0, 1$ follow from the equalities $\cos \frac{3}{2}\theta = \cos \theta \cos \frac{1}{2}\theta - \sin \theta \sin \frac{1}{2}\theta$ and $\sin \theta = 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}$. Suppose $n \geq 2$. Then we have

$$\begin{aligned} V_n(2 \cos \theta) \cos \frac{\theta}{2} &= (2 \cos \theta V_{n-1}(2 \cos \theta) - V_{n-2}(2 \cos \theta)) \cos \frac{\theta}{2} \\ &= 2 \cos \theta \cos \left(n - \frac{1}{2}\right) \theta - \cos \left(n - \frac{3}{2}\right) \theta \\ &= \cos \left(n - \frac{1}{2}\right) \theta \cos \theta - \sin \left(n - \frac{1}{2}\right) \theta \sin \theta \\ &= \cos \left(n + \frac{1}{2}\right) \theta, \end{aligned}$$

where the second equality follows from the induction hypothesis. \square

Therefore, the roots of $V_n(x)$ are $2 \cos \frac{2k-1}{2n+1}\pi = 2 \sin \frac{2n+3-4k}{2(2n+1)}\pi$ ($k = 1, 2, \dots, n$). Since $V_n(0) = \pm 1$, we obtain the next consequence.

Corollary 4.4. *Let a be a positive odd integer. Then $(2 \sin \frac{a-2k}{2a}\pi)^{\pm 1}$ is an algebraic integer for $k = -a + 4, -a + 8, \dots, a - 2$.*

4.2. Seifert fibered spaces. Let M be an orientable Seifert fibered space with Seifert index

$$\{b, (\epsilon = o, g), (a_1, b_1), \dots, (a_m, b_m)\},$$

whose base orbifold is a closed oriented surface with m singular points. Here b is an integer, g is a non-negative integer and a_i, b_i are coprime integers.

Now any non-trivial value of the Reidemeister torsion $\tau_\rho(M)$ for an irreducible $SL(2, \mathbb{C})$ -representation $\rho: \pi_1(M) \rightarrow SL(2, \mathbb{C})$ is given as follows. See [9, Main Theorem] and Remark 2.1.

Proposition 4.5.

$$\tau_\rho(M)^{-1} = 2^{4-m-g} \prod_{i=1}^m \left(1 - (-1)^{s_i} \cos \frac{r_i k_i \pi}{a_i} \right),$$

where

- (1) $r_i, s_i \in \mathbb{Z}$ such that $a_i s_i - b_i r_i = -1$ ($i = 1, \dots, m$),
- (2) $k_i \in \mathbb{Z}$ such that $0 \leq k_i \leq a_i$ and $k_i \equiv b_i \pmod{2}$ ($i = 1, \dots, m$).

Remark 4.6. A pair (r_i, s_i) is not unique, but it depends on only (a_i, b_i) . On the other hand k_i does on a representation ρ .

We can see the following by Lemmas 4.1 and 4.3. We write m_o for the number of odd a_i 's.

Proposition 4.7. *For a Seifert fibered space M with an irreducible representation ρ , if $2m_o + g \geq 4$, then the Reidemeister torsion $\tau_\rho(M)$ is an algebraic integer.*

Proof. Here we deform each factor, which contains a cosine function, of the formula in Lemma 4.5 to a more simple form.

- Case (I): a_i is odd and b_i is even.

In this case we see s_i must be odd since $a_i s_i - b_i r_i = -1$. Now,

$$\begin{aligned} 1 - (-1)^{s_i} \cos \frac{r_i k_i \pi}{a_i} &= 1 + \cos \frac{r_i k_i \pi}{a_i} \\ &= 2 \cos^2 \left(\frac{r_i k_i \pi}{2a_i} \right) \\ &= 2 \sin^2 \left(\frac{r_i k_i \pi}{2a_i} + \frac{\pi}{2} \right) \\ &= 2 \sin^2 \left(\frac{2a_i \pi}{2a_i} - \frac{(r_i k_i + a_i) \pi}{2a_i} \right) \\ &= 2 \sin^2 \left(\frac{a_i - r_i k_i}{2a_i} \pi \right). \end{aligned}$$

- Case (II): a_i is odd and b_i is odd.

In this case s_i may be odd and r_i even by changing s_i to $s_i + b_i$ and r_i to $r_i + a_i$. Similarly, we have

$$1 - (-1)^{s_i} \cos \frac{r_i k_i \pi}{a_i} = 2 \sin^2 \left(\frac{a_i - r_i k_i}{2a_i} \pi \right).$$

- Case (III): a_i is even.

In this case b_i must be odd and s_i may be odd by changing s_i to $s_i + b_i$. Similarly to Cases (I) and (II), we see

$$1 - (-1)^{s_i} \cos \frac{r_i k_i \pi}{a_i} = 2 \sin^2 \left(\frac{a_i - r_i k_i}{2a_i} \pi \right).$$

Now we see

$$\tau_\rho(M) = 2^{2m_o + g - 4} \prod_{a_i: \text{odd}} \left(2 \sin \frac{a_i - r_i k_i}{2a_i} \pi \right)^{-2} \cdot \prod_{a_i: \text{even}} \left(\sin \frac{a_i - r_i k_i}{2a_i} \pi \right)^{-2}.$$

Recall that m_o is the number of odd a_i 's. By Lemma 4.1 and Corollary 4.4, it can be seen that all factors

$$2^{2m_o+g-4}, \left(2 \sin \frac{a_i - r_i k_i}{2a_i} \pi\right)^{-2} \quad (a_i: \text{ odd}), \left(\sin \frac{a_i - r_i k_i}{2a_i} \pi\right)^{-2} \quad (a_i: \text{ even}),$$

are algebraic integers. This completes the proof. \square

Consider a Brieskorn homology 3-sphere $\Sigma(a_1, a_2, a_3)$. This is a Seifert fibered space with $m = 3$, $g = 0$ and $m_o \geq 2$. Since there exist finitely many conjugacy classes of irreducible representations, the torsion polynomial can be defined.

Corollary 4.8. *The torsion polynomial $\sigma_{\Sigma(a_1, a_2, a_3)}(t) \in \mathbb{Q}[t]$ lies in $\mathbb{Z}[t]$.*

Recall that we write $T(p, q)$ for the (p, q) -torus knot. By similar arguments and Johnson's computation [8], we see the following.

Proposition 4.9. *For any irreducible representation ρ , the Reidemeister torsion $\tau_\rho(E(T_{p,q}))$ is an algebraic integer.*

5. CONTINUOUS VARIATION OF REIDEMEISTER TORSION

In this final section, we prove Theorem 1.3 which asserts that for a certain knot K_0 the Reidemeister torsion $\tau_\rho(E(K_0))$ can vary continuously while the restriction $r(\rho)$ is fixed. We construct the knot K_0 in Figure 2 following [2, Section 8].




FIGURE 3. The knot $4_1\#4_1$ with a braid axis.

Remark 5.1. Since Figure 2 is an alternating diagram, K_0 is an alternating knot. Then, we confirm that K_0 is prime by [17, Theorem 4.4]. The Alexander polynomial $\Delta_{K_0}(t)$ of K_0 is computed as follows:

$$t^{12} - 11t^{11} + 55t^{10} - 169t^9 + 358t^8 - 551t^7 + 635t^6 - 551t^5 + 358t^4 - 169t^3 + 55t^2 - 11t + 1.$$

In particular, K_0 is not a torus knot, and thus it is hyperbolic by [19, Corollary 2]. Moreover, since $\Delta_{K_0}(t)$ is monic, K_0 is a fibered knot of genus 6 by [20, Theorem 1.2].

Let $\varpi: E(K_0) \rightarrow E(4_1 \sharp 4_1)$ denote a double branched cover whose branch set is the braid axis. The map ϖ induces a homomorphism $\varpi_*: \pi_1(E(K_0)) \rightarrow \pi_1(E(4_1 \sharp 4_1))$ and a regular map $\varpi^*: X(E(4_1 \sharp 4_1)) \rightarrow X(E(K_0))$.

Recall that irreducible representations of

$$\pi_1(E(4_1)) = \langle x, y \mid [y, x]^{-1}x = y[y, x]^{-1} \rangle$$

can be given by

$$\rho(x) = \begin{pmatrix} s & 1 \\ 0 & s^{-1} \end{pmatrix}, \quad \rho(y) = \begin{pmatrix} s & 0 \\ -t & s^{-1} \end{pmatrix},$$

up to conjugate, where $s, t \in \mathbb{C}^\times$ satisfies $\phi(s, t) = 0$. Suppose $s \neq \pm 1$. Consider irreducible representations $\rho_{s,t} * (P_u^{-1} \rho_{s,t} P_u)$ of $\pi_1(E(4_1 \sharp 4_1))$, where

$$P_u = \begin{pmatrix} u & (u - u^{-1})/(s - s^{-1}) \\ 0 & u^{-1} \end{pmatrix} \in SL(2, \mathbb{C})$$

is an element of the centralizer of $\rho_{s,t}(x)$. Then we define an irreducible representation $\rho_{s,t,u}$ of $\pi_1(E(K_0))$ by $\rho_{s,t,u} = \varpi^*(\rho_{s,t} * (P_u^{-1} \rho_{s,t} P_u))$.

Proof of Theorem 1.3. Define

$$C = \{\rho_{s,t,u} \mid s, t, u \in \mathbb{C}^\times, s \neq \pm 1, \phi(s, t) = 0\}.$$

There is a Wirtinger representation of $\pi_1(E(K_0))$ with 16 generators and 15 relations. Then, by Mathematica, we compute the Reidemeister torsion as follows:

$$\tau_{\rho_{s,t,u}}(E(K_0)) = \frac{f_2(s, t)u^2 + f_0(s, t) + f_{-2}(s, t)u^{-2}}{s^{70}(s-1)^{19}(s+1)^{18}},$$

where $f_j(s, t) \in \mathbb{Z}[s, t]$ ($j = 2, 0, -2$) are certain complicated polynomials. Moreover, $\text{res}_t(f_j(s, t), \phi(s, t)) \in \mathbb{Z}[s]$ is the product of six factors $s, s \pm 1, s^2 \pm s - 1$, and $g_j(s)$ with some multiplicities, where $g_j(s) \in \mathbb{Z}[s]$ for $j = \pm 2$.

Now, it suffices to show that $f_2(s, t)$ and $f_{-2}(s, t)$ do not vanish simultaneously. By the definition of C , we have $s \neq 0, \pm 1$. It follows from $t \neq 0$ and $\phi(s, t) = 0$ that $s^2 \pm s - 1 \neq 0$ since $s^2 + s^{-2} - 3 = (s - s^{-1} + 1)(s - s^{-1} - 1)$. Finally, by a computer calculation, we can check that $\text{res}_s(g_2(s), g_{-2}(s)) \in \mathbb{Z}$ is non-zero, namely, they have no common zeros. \square




FIGURE 4. Generators x_1, x_2, \dots, x_{16} of $\pi_1(E(K_0))$.

Example 5.2. When $(s, t) = (i, \frac{-5+\sqrt{5}}{2})$, the Reidemeister torsion $\tau_{\rho_{s,t,u}}(E(K_0))$ is equal to

$$\frac{1}{16} \left(\left((6765 + 7695i) + (6383 + 2015i)\sqrt{5} \right) u^2 + (73122 + 21422\sqrt{5}) \right. \\ \left. + \left((6765 - 7695i) + (6383 - 2015i)\sqrt{5} \right) u^{-2} \right)$$

Also, when $(s, t) = (2, \frac{5-\sqrt{105}}{8})$, $\tau_{\rho_{s,t,u}}(E(K_0))$ is equal to

$$\left((8505120805 - 233834087\sqrt{105})u^2 + 16(633637427\sqrt{105} + 6409291542) \right. \\ \left. + (58019711838\sqrt{105} + 594558745850)u^{-2} \right) / 6291456.$$

Remark 5.3. Let $s_{\pm} = \frac{1 \pm \sqrt{5}}{2}$. Then $s_{\pm}^{\pm 1}$ are the roots of $s^2 - s^{-2} - 3$ and we have four reducible representations $\rho_{s_{\pm}^{\pm 1}, 0, u}$. First, $\rho_{s_{\pm}^{-1}, 0, u}$ is equivalent to $\rho_{s_{\pm}, 0, u}$. Next, as elements of $X(E(K_0))$, they are independent of u since

$$P_u^{-1} \begin{pmatrix} s & 0 \\ 0 & s^{-1} \end{pmatrix} P_u = \begin{pmatrix} s & 1 - u^{-2} \\ 0 & s^{-1} \end{pmatrix}.$$

Moreover, for any $u \in \mathbb{C}^{\times}$, we have

$$\tau_{\rho_{s_{\pm}, 0, u}}(E(K_0)) = 256(8222 \mp 3677\sqrt{5}).$$

Finally, let us prove Corollary 1.4 stated at the end of Section 1.

Proof of Corollary 1.4. First, the longitude λ of K_0 is written as

$$\lambda = x_2 x_7^{-1} x_1 x_{12} x_5^{-1} x_{11} x_{16}^{-1} x_6^{-1} x_{10} x_{15}^{-1} x_9 x_4 x_{13}^{-1} x_3 x_8^{-1} x_{14}^{-1}.$$

By a computer calculation, we have $\rho_{s,t,u}(\lambda)_{11} = f(s, t)/g(s)$. Then the resultant $\text{res}_t(g(s)L - f(s, t), \phi(s, t))$ is the product of four factors s , $s \pm 1$, and $f_C(L, s)$, where $f_C(L, s)$ is a polynomial written after Corollary 1.4. Since $s \neq 0, \pm 1$, we conclude that $\rho_{s,t,u}(x)_{11} = s$ and $\rho_{s,t,u}(\lambda)_{11} = L$ must satisfy $f_C(L, s) = 0$.

Now, we have a 2-bridge knot such that $f_C(L, M)$ and $A_K(M, L)$ have a common zero (L_0, M_0) with $L_0, M_0 \neq 0$ and $\{L_0, M_0\} \not\subset \{1, -1\}$. It follows from the definition of the A -polynomial and [2, Theorem 3.1] that there exist $\{\rho_u\}_u \subset C$ corresponding to (L_0, M_0) and $\rho_K \in X(E(K))$ corresponding to (M_0, L_0) . Moreover, $\{L_0, M_0\} \not\subset \{1, -1\}$ implies that ρ_u coincides with ρ_K on the boundary torus, and thus we have a family $\{\rho_u * \rho_K\}_u$ of representations of $\pi_1(\Sigma(K_0, K))$. By the multiplicativity of the Reidemeister torsion and Theorem 1.3, the set $RT(\Sigma(K_0, K))$ is an infinite set. \square

REFERENCES

- [1] D. Cooper, M. Culler, H. Gillet, D. D. Long, and P. B. Shalen. Plane curves associated to character varieties of 3-manifolds. *Invent. Math.*, 118(1):47–84, 1994.
- [2] D. Cooper and D. D. Long. Remarks on the A -polynomial of a knot. *J. Knot Theory Ramifications*, 5(5):609–628, 1996.

- [3] D. A. Cox, J. Little, and D. O’Shea. *Using algebraic geometry*, volume 185 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2005.
- [4] N. M. Dunfield, S. Friedl, and N. Jackson. Twisted Alexander polynomials of hyperbolic knots. *Exp. Math.*, 21(4):329–352, 2012.
- [5] B. Farb and D. Margalit. *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2012.
- [6] A. Hatcher and W. Thurston. Incompressible surfaces in 2-bridge knot complements. *Invent. Math.*, 79(2):225–246, 1985.
- [7] J. Hoste and P. D. Shanahan. A formula for the A-polynomial of twist knots. *J. Knot Theory Ramifications*, 13(2):193–209, 2004.
- [8] D. Johnson. A geometric form of Casson’s invariant and its connection to Reidemeister torsion. preprint, 1988. available at https://celebratio.org/media/docs/Johnson_GeomFormCassonInvar.1988.pdf.
- [9] T. Kitano. Reidemeister torsion of Seifert fibered spaces for $SL(2; \mathbf{C})$ -representations. *Tokyo J. Math.*, 17(1):59–75, 1994.
- [10] T. Kitano. Reidemeister torsion of the figure-eight knot exterior for $SL(2; \mathbf{C})$ -representations. *Osaka J. Math.*, 31(3):523–532, 1994.
- [11] T. Kitano. Twisted Alexander polynomial and Reidemeister torsion. *Pacific J. Math.*, 174(2):431–442, 1996.
- [12] T. Kitano. Some numerical computations on Reidemeister torsion for homology 3-spheres obtained by Dehn surgeries along the figure-eight knot. arXiv:1603.03728v2, 2016.
- [13] T. Kitano. A polynomial defined by the $SL(2; \mathbf{C})$ -Reidemeister torsion for a homology 3-sphere obtained by a Dehn surgery along a $(2p, q)$ -torus knot. *Tohoku Math. J. (2)*, 69(4):571–583, 2017.
- [14] T. Kitano and Y. Nozaki. Finiteness of the image of the Reidemeister torsion of a splice. *Ann. Math. Blaise Pascal*, 27(1):19–36, 2020.
- [15] T. Kitano and A. T. Tran. A torsion polynomial defined by $SL(2; \mathbf{C})$ -Reidemeister torsion of 3-manifolds. unpublished notes.
- [16] T. Kitano and Y. Yamaguchi. $SL(2; \mathbb{R})$ -representations of a Brieskorn homology 3-sphere. arXiv:1602.07503v3, 2016.
- [17] W. B. R. Lickorish. *An introduction to knot theory*, volume 175 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1997.
- [18] C. Maclachlan and A. W. Reid. *The arithmetic of hyperbolic 3-manifolds*, volume 219 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2003.
- [19] W. Menasco. Closed incompressible surfaces in alternating knot and link complements. *Topology*, 23(1):37–44, 1984.
- [20] K. Murasugi. On a certain subgroup of the group of an alternating link. *Amer. J. Math.*, 85:544–550, 1963.
- [21] R. Riley. Nonabelian representations of 2-bridge knot groups. *Quart. J. Math. Oxford Ser. (2)*, 35(138):191–208, 1984.
- [22] M. Wada. Twisted Alexander polynomial for finitely presentable groups. *Topology*, 33(2):241–256, 1994.
- [23] S. Yoon. Adjoint Reidemeister torsions of two-bridge knots. *Proc. Amer. Math. Soc.*, to appear.

DEPARTMENT OF INFORMATION SYSTEMS SCIENCE, FACULTY OF SCIENCE AND ENGINEERING, SOKA UNIVERSITY, TANGI-CHO 1-236, HACHIOJI, TOKYO 192-8577, JAPAN
Email address: `kitano@soka.ac.jp`

GRADUATE SCHOOL OF ADVANCED SCIENCE AND ENGINEERING, HIROSHIMA UNIVERSITY, 1-3-1 KAGAMIYAMA, HIGASHI-HIROSHIMA CITY, HIROSHIMA, 739-8526, JAPAN
Email address: `nozakiy@hiroshima-u.ac.jp`