

# Portfolio selection models based on interval-valued conditional value at risk (ICVaR) and empirical analysis

Jinping Zhang<sup>a,\*</sup>, Keming Zhang<sup>a</sup>

<sup>a</sup>*School of Mathematics and Physics, North China Electric Power University, Beijing, 102206, P. R. China*

---

## Abstract

Risk management is very important for individual investors or companies. There are many ways to measure the risk of investment. Prices of risky assets vary rapidly and randomly due to the complexity of finance market. Random interval is a good tool to describe uncertainty with both randomness and imprecision. Considering the uncertainty of financial market, we employ random intervals to describe the returns of a risk asset and consider the tail risk, which is called the interval-valued Conditional Value at Risk (ICVaR, for short). Such an ICVaR is a risk measure and satisfies subadditivity. Under the new risk measure ICVaR, as a manner similar to the classical portfolio model of Markowitz, optimal interval-valued portfolio selection models are built. Based on the real data from mainland Chinese stock market, the case study shows that our models are interpretable and consistent with the practical scenario.

*Keywords:* Portfolio Selection, Interval-Valued Random Variable, Value at Risk (VaR), Conditional Value at Risk (CVaR)

---

## 1. Introduction

A portfolio is a collection of stocks, bonds, and financial derivatives held by investors or financial institutions to spread risk. Since H. Markowitz published the pioneering paper on mean-variance model [13] in 1952, there have been many works on the applications and extension of the classical models.

---

\*Corresponding author

*Email addresses:* zhangjinping@ncepu.edu.cn (Jinping Zhang),  
k.m.zhang@ncepu.edu.cn (Keming Zhang)

Risk measurement is very important for investors or financial analysts and must be considered in portfolio selection models. Besides variance, there are many other ways to measure the risk of assets. For example, Konnor and Yamazaki [8] (1991) studied the optimal portfolio problem based on absolute deviation or semi-variance, which is easy to calculate than variance of portfolio. Value at risk (VaR, for short) is a financial risk measurement that has been developed and widely used (See e.g. [1, 2, 14]. Hamel et al. [4] (2013) extended real-valued VaR to set-valued case. VaR is convenient, easy to calculate. But VaR has no nice mathematical properties such as convexity, sub-additivity, which are crucial to solve the optimal problem. In order to overcome the drawback of VaR, one can consider the conditional value at risk (CVaR, for short), the coherent risk measurement with nice mathematical properties [12, 15]. Because of the asymmetry of information and the incompleteness of the market, the price of risky assets shows a more complex form than a scalar variable. In order to describe the complexity, randomness and inaccuracy of the financial market, the tool of interval-valued variables is a good choice. For example, one can consider interval-valued return, interval-valued VaR (IVaR, for short), interval-valued CVaR (See e.g. [10], 2017). Based on interval-valued return, it is natural to study interval-valued portfolio selection. Ida [6] (2003) studied portfolio selection with interval coefficients. [7] (2004) considered portfolio problem with both interval and fuzzy coefficients. Giove et al. [3] (2006) studied the interval portfolio model based on regret function. Zhang and Li [18] (2009) considered the portfolio selection models based on interval-valued semi-variance. Also, there are some references deal with the portfolio selection without using the usual risk measure such as variance, semi-variance, VaR CVaR. For example, Mansini and Speranza [11] (1999) built the portfolio model with minimum transaction lots, which is independent of the risk function.

In this paper, we collect the closing price, the highest price and the lowest price on each trade day. Then the range of return of risky asset can be obtained as an interval-valued random variable. The interval-valued CVaR is used to describe the risk instead of the variance. Then we build the interval-valued optimal portfolio selection models. The optimal portfolio maximizes the return for a given level of risk and the optimal portfolio minimizes the risk for a given level of return. The interval-valued case is a natural extension of the classic portfolio model, which can comprehensively reflect the complexity of the financial market and the risk appetite of investors.

The rest of the paper is organized as follows: Section 2 is the preliminaries of interval-valued variables and interval-valued programming. Section 3 contributes

to the interval-valued portfolio selection models based on ICVaR. Section 4 is devoted to the application of our models in real financial market. Section 5 is the concluding remarks.

## 2. Preliminaries

In this section, we list the knowledge of interval-valued variables and interval-valued programming needed later.

### 2.1. Operations and order of intervals

Let  $\mathbb{R}$  be the set of all real numbers,  $A, B$  real closed intervals. Let  $A = [a^L, a^U] = \{a \in \mathbb{R} : a^L \leq a \leq a^U\}$ ,  $B = [b^L, b^U] = \{b \in \mathbb{R} : b^L \leq b \leq b^U\}$ . If  $a^L = a^U = a$ , the interval  $A = [a, a]$  degenerates to a real number.

$$A + B := [a^L + b^L, a^U + b^U],$$

$$\lambda \cdot A = \lambda[a^L, a^U] := \begin{cases} [\lambda a^L, \lambda a^U], & \text{if } \lambda \geq 0, \\ [\lambda a^U, \lambda a^L], & \text{if } \lambda < 0. \end{cases}$$

Let  $m(A) := \frac{1}{2}(a^L + a^U)$ ,  $w(A) := \frac{1}{2}(a^U - a^L)$ , which are called the mean and width of the interval  $A$ , respectively. Then  $A$  also can be determined by  $m(A)$  and  $w(A)$ .

Ranking intervals is a crucial step to study interval-valued optimization problems. There are infinitely many ways to rank intervals [17]. The popular and natural way is using the endpoints or their functions such as mean and width. In this paper, we use the rule “mean-first, left-second” to rank intervals for calculating the IVaR and ICVaR. E.g. for two intervals  $A = [a^L, a^U]$  and  $B = [b^L, b^U]$ , define  $A < B$  if  $m(A) < m(B)$ . If  $m(A) = m(B)$  and  $a^L < b^L$ , then  $A < B$ .  $A = B$  if and only if  $m(A) = m(B)$ ,  $a^L = b^L$ . Multiple intervals can be ranked well according to this rule.

### 2.2. Interval-valued optimization

We shall employ the method introduced in [16] (2001) to solve interval-valued optimization portfolio. In [16], at first the authors defined an index  $\Gamma$  to describe the acceptability for ranking two intervals. Let  $\otimes$  be a partial order of intervals. For any closed intervals  $A$  and  $B$ , define

$$\Gamma(A \otimes B) = \frac{m(B) - m(A)}{w(B) + w(A)}, \quad (1)$$

where  $w(B) + w(A) \neq 0$ . The value of function  $\Gamma(A \otimes B)$ , denoted by  $\gamma$ , may be interpreted as the grade of acceptability of ‘the interval  $A$  is less than the interval  $B$ ’.

The grade of acceptability of  $A \otimes B$  may be classified and interpreted further on the basis of comparative position of mean and width of interval  $B$  with respect to those of interval  $A$  as follows:

$$\Gamma(A \otimes B) = \begin{cases} \leq 0, & \text{if } m(B) \leq m(A), \\ > 0, < 1, & \text{if } m(A) < m(B) \text{ and } a^U > b^L \\ \geq 1, & \text{if } m(A) < m(B) \text{ and } a^U \leq b^L. \end{cases} \quad (2)$$

If  $\Gamma(A \otimes B) \leq 0$ , then the premise ‘ $A$  is less than  $B$ ’ is not accepted. If  $0 < \Gamma(A \otimes B) < 1$ , then the interpreter accepts the premise ‘ $A$  is less than  $B$ ’ with different grades of satisfaction ranging from zero to one (excluding zero and one). If  $\Gamma(A \otimes B) \geq 1$ , the interpreter is absolutely satisfied with the premise ‘ $A \otimes B$ ’.

### 2.3. A satisfactory crisp equivalent system of $Ax \leq B$

Let  $A = [a^L, a^U]$ ,  $B = [b^L, b^U]$  and  $x$  be a nonnegative singleton variable. In order to solve linear programming problems with interval coefficients, Sengupta et al. [16] proposed a satisfactory crisp equivalent form for the interval inequality relations  $Ax \leq B$  or  $Ax \geq B$ .

The satisfactory crisp equivalent form of  $Ax \leq B$  is given as follows ([16]):

$$Ax \leq B \text{ iff } \begin{cases} a^U x \leq b^U, \\ \Gamma(B \otimes Ax) \leq \gamma \in [0, 1], \end{cases} \quad (3)$$

where  $\gamma$  may be interpreted as an optimistic threshold given and fixed by the decision maker. Similarly, for  $Ax \geq B$ , the satisfactory crisp equivalent form is:

$$Ax \geq B \text{ iff } \begin{cases} a^L x \geq b^L, \\ \Gamma(Ax \otimes B) \leq \gamma \in [0, 1]. \end{cases} \quad (4)$$

Now we consider the maximizing interval-valued linear programming problem:

$$\begin{aligned} & \text{Maximize}_{\otimes} && Z = \sum_{i=1}^n [c_i^L, c_i^U] x_i, \\ & \text{subject to} && \sum_{i=1}^n [a_{ij}^L, a_{ij}^U] x_i \leq [b_j^L, b_j^U], \\ & && j = 1, 2, \dots, k \\ & && x_i \geq 0, i = 1, \dots, n. \end{aligned} \quad (5)$$

According to [16], the principle of  $\gamma$ -index indicates that for the maximization (minimization) problem, an interval with a higher midpoint is superior (inferior) to an interval with a lower mid-value. Therefore, in order to obtain the max (min) of the interval-valued objective function, considering the midpoint is our primary concern. We reduce the interval objective function its central value and use conventional LP (Linear Programming) techniques for its solution. We also consider the width but as a secondary attribute, only to confirm whether it is within the acceptable limit of the decision maker. If it is not, one has to reduce the extent of width (uncertainty) according to his satisfaction and thus to obtain a less wide interval from the non-dominated alternatives accordingly. So by the satisfactory crisp equivalent form (3), the following LP problem (6) is the necessary equivalent form of the model (5):

$$\begin{aligned}
\text{Maximize} \quad & m(Z) = \frac{1}{2} \sum_{i=1}^n (c_i^L + c_i^U) x_i, \\
\text{subject to} \quad & \sum_{i=1}^n a_{ij}^U x_i \leq b_j^U, \\
& \sum_{i=1}^n [(a_{ij}^L + a_{ij}^U) - \gamma(a_{ij}^U - a_{ij}^L)] x_i \leq b_j^L + b_j^U + \gamma(b_j^U - b_j^L), \\
& j = 1, \dots, k \\
& x_i \geq 0, \text{ for } i = 1, \dots, n.
\end{aligned} \tag{6}$$

Similarly, the minimizing interval linear programming can be constructed as follows:

$$\begin{aligned}
\text{Minimize}_{\ominus} \quad & Z = \sum_{i=1}^n [c_i^L, c_i^U] x_i, \\
\text{subject to} \quad & \sum_{i=1}^n [a_{ij}^L, a_{ij}^U] x_i \geq [b_j^L, b_j^U], \text{ for } j = 1, \dots, k \\
& x_i \geq 0, i = 1, \dots, n.
\end{aligned} \tag{7}$$

According to (4), the interval-valued programming problem (7) can be transformed to the following real valued linear programming model (8):

$$\begin{aligned}
\text{Minimize} \quad & m(Z) = \frac{1}{2} \sum_{i=1}^n (c_i^L + c_i^U) x_i, \\
\text{subject to} \quad & \sum_{i=1}^n a_{ij}^L x_i \geq b_j^L, \\
& \sum_{i=1}^n [(a_{ij}^L + a_{ij}^U) + \gamma(a_{ij}^U - a_{ij}^L)] x_i \geq b_j^L + b_j^U - \gamma(b_j^U - b_j^L), \\
& j = 1, \dots, k \\
& x_i \geq 0, \text{ for } i = 1, \dots, n.
\end{aligned} \tag{8}$$

### 3. Portfolio selection based on ICVaR

We consider the portfolio problem with  $n$  risky assets. Due to the volatility and complexity of financial market, the profit is uncertain. There are several tools to describe uncertainty, for example, random variable, fuzzy variable, set-valued variable etc. Here the return is considered as a random interval, which includes not only randomness but also imprecision.

#### 3.1. Interval-valued risk measure

##### 3.1.1. Interval-valued random variable

Let  $(\Omega, \mathcal{F}, P)$  be a complete probability space.  $\mathbb{R}$  be the real space equipped with the Borel sigma-algebra  $\mathcal{B}(\mathbb{R})$ .  $K(\mathbb{R})$  ( $K_c(\mathbb{R})$ ) denotes the family of all nonempty closed (nonempty, closed and convex, respectively) subsets of  $\mathbb{R}$ . The mapping  $V : \Omega \rightarrow K(\mathbb{R})$  is called *measurable set-valued mapping* or *random set* if for any element  $B \in \mathcal{B}(\mathbb{R})$ ,  $\{\omega \in \Omega : V(\omega) \cap B \neq \emptyset\} \in \mathcal{F}$ .

$V$  is called an *interval-valued random variable* or *random interval* if  $V$  is measurable and takes value in  $K_c(\mathbb{R})$ .

For any  $X \in K_c(\mathbb{R})$ , the *distribution function* of  $V$  is defined by

$$F(X) := P(\omega \in \Omega : V(\omega) < X),$$

where  $<$  is the partial order according to the rule ‘mean-first, left-second’.  $V$  is called *integrably bounded* if  $\int_{\Omega} \sup_{x \in V(\omega)} |x| dP < \infty$ . The real valued random variable  $f$  is called an *integrable selection* of  $V$  if  $f(\omega) \in V(\omega)$  a.s. and  $\int_{\Omega} f dP < \infty$ .

For an integrably bounded random set  $V$ ,

$$S_V^1 := cl\{f \in L^1(\Omega, \mathbb{R}) : f(\omega) \in V(\omega) \text{ a.s.}\},$$

where  $L^1(\Omega, \mathbb{R})$  denotes the family of all integrable real valued random variables, the closure  $cl$  is taken in  $L^1(\Omega, \mathbb{R})$ . For two random sets  $V$  and  $U$ , it is well known that  $V = U$  a.s. if and only if  $S_V^1 = S_U^1$  ([5]).

The *expectation* of  $V$  is defined by

$$E(V) = \{E(f), f \in S_V^1\},$$

which is closed in  $\mathbb{R}$ .

By Lemma 3.1 of [19], a random set  $V$  is an interval-valued random variable if and only if  $V = [f, g]$ , where both  $f$  and  $g$  are real valued random variables.

For an integrably bounded interval-valued random variable  $V = [f, g]$ , obviously it holds that  $E(V) = [E(f), E(g)]$

### 3.1.2. IVaR and ICVaR

As a manner similar to the usual value at risk and conditional value at risk, we define the interval value at risk (IVaR) and interval conditional value at risk (ICVaR) as following:

**Definition 3.1.** Let  $(\Omega, \mathcal{F}, P)$  be a complete probability space.  $R$  is the random interval-valued return of a risky asset. Given confidence level  $1 - \alpha$  ( $0 < \alpha < 1$ ), the value at risk is defined by

$$IVaR := -\inf\{X \in K_c(\mathbb{R}) : P(R < X) = 1 - \alpha\},$$

where  $\inf$  is taken according to the order  $<$ .

**Definition 3.2.** Let  $(\Omega, \mathcal{F}, P)$  be a complete probability space.  $R$  is the random interval-valued return of a risky asset. Given confidence level  $1 - \alpha$  ( $0 < \alpha < 1$ ),  $IVaR$  is the value at risk. The conditional value at risk is defined by

$$ICVaR := -E(R|R < -IVaR).$$

Similar to the real-valued CVaR, ICVaR describes the tail risk.

**Proposition 3.1.** *ICVaR is a coherent risk measure. I.e. it has the following properties:*

- (1) *Transition-equivariant*
- (2) *Positively homogeneous*
- (3) *Convex*
- (4) *Monotonic w.r.t stochastic dominance of order 1, monotonic w.r.t monotonic dominance of order 2.*

*Proof.* The expectation of random interval is linear for positive coefficients (c.f.[9]). Then it is easy to prove the result as a manner similar to the real-valued case.  $\square$

### 3.2. Portfolio Models

Let the rate of return of asset  $R = [r^L, r^U]$  be an interval-valued random variable,  $R_{ij} = [r_{ij}^L, r_{ij}^U]$  be the rate of return of the  $i$ -th security at the  $j$ -th period,  $i = 1, \dots, n$ ,  $j = 1, \dots, k$ .

Given the confidence level  $1 - \alpha$ ,  $IVaR_{ij}$  ( $ICVaR_{ij}$ ) is the value at risk (conditional value at risk, resp.) of the  $i$ -th risky asset at the  $j$ -th period,  $i = 1, \dots, n$ ,  $j = 1, \dots, k$ . Let  $x_i$  ( $i = 1, \dots, n$ ) be the proportion of the  $i$ -th asset in the portfolio. By Proposition 3.1, the  $ICVaR$  of portfolio is less than the linear composition of  $ICVaR_i$  with weights  $x_i \geq 0$  ( $i = 1, \dots, n$ ).

Now we build two models of portfolio selection as follows:

- Model 1

We consider the optimal portfolio problem under given acceptable maximum of risk level:

$$\begin{aligned}
 & \text{Maximize}_{\otimes} & E(R_P) &= \sum_{i=1}^n x_i E(R_i), \\
 & \text{subject to} & \sum_{i=1}^n [ICVaR_{ij}^L, ICVaR_{ij}^U] x_i &\leq [ICVaR_{0j}^L, ICVaR_{0j}^U], \\
 & & j &= 1, \dots, k, \\
 & & \sum_{i=1}^n x_i &= 1, \quad x_i \geq 0, \quad i = 1, \dots, n.
 \end{aligned} \tag{9}$$

where  $E(R_i)$  is the expectation return of the  $i$ -th asset,  $ICVaR_{0j} = [ICVaR_{0j}^L, ICVaR_{0j}^U]$  is the acceptable maximum risk in the  $j$ -period,  $j = 1, \dots, m$ , which are subjective and given by the investors.

By (6), the corresponding real-valued linear programming problem of (9) is represented as (10) below

$$\begin{aligned}
 & \text{Maximize} & m(E(R_P)) &= \sum_{i=1}^n x_i m(E(R_i)), \\
 & \text{subject to} & \sum_{i=1}^n x_i ICVaR_{ij}^U &\leq ICVaR_{0j}^U, \\
 & & \sum_{i=1}^n x_i [m(ICVaR_{ij}) - \gamma w(ICVaR_{ij})] &\leq m(ICVaR_{0j}) + \gamma w(ICVaR_{0j}), \\
 & & j &= 1, \dots, k \\
 & & x_i &\geq 0, \text{ for } i = 1, \dots, n,
 \end{aligned} \tag{10}$$

where  $m(A)$  is the midpoint of interval  $A$  and  $w(A)$  the semi-width of  $A$ .  $\gamma \in (0, 1)$  is a given index in advance, which can describe the degree of risk appetite of investors. The larger  $\gamma$ , the lower the risk aversion.

- Model 2

$$\begin{aligned}
 & \text{Maximize}_{\otimes} & \sum_{i=1}^n x_i ICVaR_i, \\
 & \text{subject to} & \sum_{i=1}^n x_i E(R_{ij}) &\geq R_{0j}, \\
 & & j &= 1, \dots, k \\
 & & \sum_{i=1}^n x_i &= 1, \quad x_i \geq 0, \quad i = 1, \dots, n.
 \end{aligned} \tag{11}$$



Correspondingly, by (8), the interval-valued linear programming problem (11) is represented as the following real-valued linear programming model (12) below

$$\begin{aligned}
& \text{Minimize} && \frac{1}{2} \sum_{i=1}^n x_i (ICVaR_i^L + ICVaR_i^U), \\
& \text{subject to} && \sum_{i=1}^n x_i E(r_{ij}^L) \geq r_{0j}^L, \\
& && \sum_{i=1}^n x_i [m(E(R_{ij})) + \gamma w(E(R_{ij}))] \geq m(R_{0j}) - \gamma w(R_{0j}), \\
& && j = 1, \dots, k \\
& && x_i \geq 0, \text{ for } i = 1, \dots, n,
\end{aligned} \tag{12}$$

where  $m$ ,  $w$  and  $\gamma$  have the same meaning as that in (10).

#### 4. Case Study

This paper selects the stocks of ten listed companies of financial markets in Chinese mainland as samples for a case study. The listed companies include oil, bank, coal, liquor, security, medicine, technology industry and so on. The ticker symbols of stocks are listed in Table 1. Daily prices including the daily closing price  $S_j$ , the highest price  $S_j^U$  and the lowest price  $S_j^L$  are collected from Jan. 2016 to Sept. 2020 (Data source: <http://www.finance.sina.cn>). The software R is employed to make the empirical analysis.

Table 1: Sample Assets

Number	ST01	ST02	ST03	ST04	ST05
Ticker	SH600028	SH600085	SH600188	SH600536	SH601939
Number	ST06	ST07	ST08	ST09	ST10
Ticker	SZ000333	SZ6000735	SZ000776	SZ000858	SZ002428

According to the daily closing price, the highest price and the lowest price of the stock, we calculate the exact value and the interval value of return rate. Let the real valued return  $r_j = \ln S_j - \ln S_{j-1}$ ,  $r_j^L = \ln S_j^L - \ln S_{j-1}$ ,  $r_j^U = \ln S_j^U - \ln S_{j-1}$ ,  $R_j = [r_j^L, r_j^U]$  be the interval return. As usual, the real valued return is not normal distributed, see Table 2. Then it is not suitable to compute the real valued VaR by using the variance-covariance method. The historical simulation method is employed to estimate the real-valued VaR, CVaR and IVaR, ICVaR. There are many methods to rank intervals. Here the ‘‘mean-first, left-second’’ of

Table 2: Jarque-Bera test

Number	ST01	ST02	ST03	ST04	ST05
$\chi^2$	843.55	1372.1	611.96	123.05	1285.1
Number	ST06	ST07	ST08	ST09	ST10
$\chi^2$	313.1	292.48	1883.1	205.07	138.27
$df = 2, p < 2.2 \times 10^{-16}$					

intervals ranking method is used to estimate IVaR and ICVaR under the confidence level  $1 - \alpha = 0.95$ . By the satisfactory crisp equivalent system, interval-valued programming problems are transformed into real-valued ones. Then the optimal solutions of the corresponding portfolio models are obtained.

Table 3: Portfolio, Model 1 with the same  $ICVaR_{0j}, j = 1, \dots, 5$ 

$ICVaR_{0j} = [0.008, 0.08], j = 1, \dots, 5$												
	ST01	ST02	ST03	ST04	ST05	ST06	ST07	ST08	ST09	ST10	$ST_{Op}$	$\gamma$
[1]	0	0	0	0	0	0	0	0	1	0	0.0009	0.15
[2]	0	0	0	0	0	0.0349	0	0	0.9651	0	0.0009	0.05
[3]	0	0	0	0	0	0.0909	0	0	0.9091	0	0.0009	0.04
[4]	0	0	0	0	0	0.1462	0	0	0.8538	0	0.0009	0.03
[5]	0	0	0	0	0	0.1736	0	0	0.8264	0	0.0009	0.025
[6]	0	0	0	0	0	0.2008	0	0	0.7992	0	0.0009	0.02
[7]	0	0	0	0	0	0.2869	0	0	0.7131	0	0.0009	0.01
$ICVaR_{0j} = [0.003, 0.07], j = 1, \dots, 5$												
	ST01	ST02	ST03	ST04	ST05	ST06	ST07	ST08	ST09	ST10	$ST_{Op}$	$\gamma$
[1]	0	0	0	0	0.1537	0.7908	0	0.0000	0.0554	0	0.0007	0.15
[2]	0	0	0	0	0.1875	0.1268	0	0.3520	0.3337	0	0.0005	0.05
[3]	0	0	0	0	0.2136	0.0973	0	0.3935	0.2956	0	0.0004	0.04
[4]	0	0	0	0	0.2395	0.0680	0	0.4348	0.2577	0	0.0004	0.03
[5]	0	0	0	0	0.2523	0.0535	0	0.4552	0.2389	0	0.0004	0.025
[6]	0	0	0	0	0.2652	0.0389	0	0.4755	0.2203	0	0.0003	0.02
[7]	0	0	0	0	0.3270	0	0	0.4792	0.1938	0	0.0002	0.01

In Table 3, when the maximum risk level  $ICVaR_0$  is different, the selection of assets is different. When taking the lower acceptable risk level  $ICVaR_{j0} = [0.003, 0.07]$ , the investor prefers to the stock ST05 (CCB, China Construction Bank) and ST08 (GF Securities Co., Ltd) since these two stocks have stable returns and lower risk compared with the stock ST06 and ST09, which implies the investor is more conservative than one who take  $ICVaR_{0j} = [0.008, 0.08]$ . Table 3 also shows that the less  $\gamma$ , the more allocation to assets with lower risk. Correspondingly, the weights of ST06 (Mediea Group) and ST09 (Wuliangye Group) increase with respect to  $\gamma$ . The result implies that ST06 and ST09 have higher return but accompanying with higher risk, which is consistent with the actual situation of the market.

Table 4: Portfolio, Model 2 with the same  $R_{0j}$ ,  $j = 1, \dots, 5$

$R_{0j} = [-0.025, 0.025], j = 1, \dots, 5$												
	ST01	ST02	ST03	ST04	ST05	ST06	ST07	ST08	ST09	ST10	$ST_{OP}$	$\gamma$
[1]	0	0	0	0	1	0	0	0	0	0	0.0243	0.04
[2]	0	0	0	0	0.9045	0	0.0955	0	0	0	0.0265	0.03
[3]	0	0	0	0	0.8332	0	0.1668	0	0	0	0.0281	0.025
[4]	0	0	0	0	0.7253	0.0285	0.2462	0	0	0	0.0302	0.02
[5]	0	0	0	0	0.3844	0.1797	0.4358	0	0	0	0.0361	0.01
$R_{0j} = [-0.02, 0.02], j = 1, \dots, 5$												
	ST01	ST02	ST03	ST04	ST05	ST06	ST07	ST08	ST09	ST10	$ST_{OP}$	$\gamma$
[1]	0	0	0	0	0.9717	0	0.0283	0	0	0	0.0250	0.04
[2]	0	0	0	0	0.8534	0	0.1466	0	0	0	0.0277	0.03
[3]	0	0	0	0	0.7825	0.0062	0.2113	0	0	0	0.0292	0.025
[4]	0	0	0	0	0.6458	0.0665	0.2877	0	0	0	0.0316	0.02

Table 5: Portfolio, Model 1 with different  $ICVaR_{0j}$  for different  $j = 1, \dots, 5$

$ICVaR_{01} = [0.004, 0.08] \quad ICVaR_{02} = [0.0045, 0.06] \quad ICVaR_{03} = [0.007, 0.08]$												
$ICVaR_{04} = [0.007, 0.06] \quad ICVaR_{05} = [0.018, 0.07]$												
	ST01	ST02	ST03	ST04	ST05	ST06	ST07	ST08	ST09	ST10	$ST_{OP}$	$\gamma$
[1]	0	0	0	0	0	0.1700	0	0	0.8300	0	0.0009	0.05
[2]	0	0	0	0	0	0.1700	0	0	0.8300	0	0.0009	0.04
[3]	0	0	0	0	0	0.1734	0	0	0.8266	0	0.0009	0.03
[4]	0	0	0	0	0	0.1962	0	0	0.8038	0	0.0009	0.025
[5]	0	0	0	0	0	0.2189	0	0	0.7811	0	0.0009	0.02
[6]	0	0	0	0	0	0.8222	0	0	0.1778	0	0.0009	0.01

Table 6: Portfolio, Model 2 with different  $R_{0j}$  for different  $j = 1, \dots, 5$

$R_{01} = [-0.020, 0.025] \quad R_{02} = [-0.020, 0.020] \quad R_{03} = [-0.025, 0.025]$												
$R_{04} = [-0.025, 0.020] \quad R_{05} = [-0.021, 0.016]$												
	ST01	ST02	ST03	ST04	ST05	ST06	ST07	ST08	ST09	ST10	$ST_{OP}$	$\gamma$
[1]	0	0	0	0	1	0	0	0	0	0	0.0243	0.15
[2]	0	0	0	0	0.5045	0.4955	0	0	0	0	0.0295	0.05
[3]	0	0	0	0	0.3512	0.6155	0.0333	0	0	0	0.0315	0.04
[4]	0	0	0	0	0.0533	0.7534	0.1933	0	0	0	0.0366	0.03

In Table 4, given the minimum return constraint  $R_{0j} = [-0.025, 0.025]$  and  $[-0.02, 0.02]$ , to minimize the risk, it is obvious that ST05 has the highest proportion in each scenario, very different from the case in Model 1. Its lower risk property explains this result. Also one can figure out that the weight of stock ST05 increases with respect to  $\gamma$  however the proportion of ST07 behaves in the contrary. In a word, the solution is sensitive to the constraints and also depends on the index  $\gamma$ . One also can figure out the law from Table 5 and Table 6.

Different risk preference has different weights for portfolio selection. In the process of solving the model, we can reasonably allocate the investment share according to the degree of risk preference of investors. So that it is important to select appropriate constraints and  $\gamma$  for making a decision of portfolio selection for the investors.

## 5. Concluding remarks

This paper contributes to the optimal selection under interval-valued environment. At first, the interval-valued conditional value at risk ICVaR was defined, which is a coherent risk measure. An then, based on the risk measure, two interval-valued portfolio selection models were constructed, where both the return and risk of a risky asset are random intervals. The “mean-first, left-second” method was employed to rank the intervals for estimating IVaR and ICVaR by historical simulation method. Of course, one can use other methods to rank intervals, such as “mean-first, right-second” or possibility degree etc. Different ranking method will lead to different IVaR and ICVaR for the risky asset. On solving the maximization and minimization problem of interval values, the satisfactory crisp equivalent system of  $Ax \leq B$  ([16]) was used. In Model 1 and Model 2, the  $\Gamma$ -index can describe investors’ subjective preference or aversion to risk, which is an innovation different from the classical portfolio model. The smaller the index, the higher the risk aversion of investors. The result of case study also shows it. But the real finance market is very complicated. The risk of asset is not only related to the industry, but also closely related to the size of the company, trading time, national policy, and so on. It is expected that the models have certain guiding significance for investors’ choice.

## Acknowledgment

The research of Jinping Zhang was supported by Natural Science Foundation of Beijing Municipality (No.1192015).

## References

- [1] Castellacci, G. and Siclari, M. The practice of Delta-Gamma VaR: Implementing the quadratic portfolio model. *European Journal of Operational Research*, 2003, 150(3): 529-545.
- [2] Costello, A., Asem, E. and Gardner E. Comparison of historically simulated VaR: Evidence from oil prices. *Energy Economics*, 2008, 30(5): 2154-2166.
- [3] Giove, S., Funari, S. and Nardelli, C. An interval portfolio selection problem based on regret function. *European Journal of Operational Research*, 2006, 170(1): 253-264.
- [4] Hamel, A. H., Rudloff, B. and Yankova, M. Set-valued average value at risk and its computation. *Mathematics and Financial Economics*, 2013, 7(2): 229-246.
- [5] F. Hiai, and H. Umegaki. Integrals, conditional expectations and martingales of multivalued functions. *J. Multivar. Anal.*, 1977, 7: 149-182.
- [6] Ida, M. Portfolio selection problem with Interval Coefficients, *Applied Mathematics Letters*, 2003, 16: 709-713.
- [7] Ida, M. Solutions for the portfolio selection problem with Interval and fuzzy Coefficients, *Reliable Computing*, 2004, 1: 389-400.
- [8] Konno, H. and Yamazaki, H. Mean-absolute deviation portfolio optimization model and its applications to Tokyo stock market. *Management Science*, 1991, 37(5): 519-531
- [9] Li, S., Ogura, Y. and Kreinovich, V. *Limit Theorems and Applications of Set-Valued and Fuzzy Sets-Valued Random Variables*, Kluwer Academic Publishers, 2002
- [10] Li, Z., Zhang, J. and Wang, X. Interval-valued risk measure models and empirical analysis. Fuzzy Systems Association, International Conference on Soft Computing & Intelligent Systems. IEEE, 2017.
- [11] Mansini, R. and Speranza, M. G. Heuristic algorithms for the portfolio selection problem with minimum transaction lots. *European Journal of Operational Research*, 1999, 114: 219-233.

- [12] Mansini, R., Ogryczak, W. and Speranza, M. G. Conditional value at risk and related linear programming models for portfolio optimization. *Ann Oper Res*, 2007, 152: 227-256. DOI 10.1007/s10479-006-0142-4
- [13] Markowitz, H. Portfolio Selection. *Journal of Finance*, 1952, 7(1): 77-91.
- [14] Ramazan, G. and Faruk, S. High volatility, thick tails and extreme value theory in value-at-risk estimation. *Insurance Mathematics & Economics*, 2003, 33(2): 337-356.
- [15] Rockafellar, R. T. and Uryasev, S. Optimization of conditional value-at-risk. *J. Risk*, 2000, 2: 21-42.
- [16] Sengupta, A., Pal, T. K. and Chakraborty, D. Interpretation of inequality constraints involving interval coefficients and a solution to interval linear programming, *Fuzzy Sets and Systems*, 2001, 119(1): 129-138.
- [17] Wang, W. and Wang, Z. Total orderings defined on the set of all fuzzy numbers. *Fuzzy Sets and Systems*, 2014, 243: 131-141.
- [18] Zhang, J. and Li, S. The portfolio selection problem with random interval-valued return rates. *International Journal of Innovative Computing, Information and Control*, 2009, 5: 47-56.
- [19] Zhang, J., Li, S., Mitoma, I. and Okazaki, Y. On set-valued stochastic integrals in an M-type 2 Banach space, *J. Math. Anal. Appl.*, 2009, 350: 216-233.