

# CONGRUENCES FOR ODD CLASS NUMBERS OF QUADRATIC FIELDS WITH ODD DISCRIMINANT

JIGU KIM AND YOSHINORI MIZUNO

**Abstract.** For any distinct two primes  $p_1 \equiv p_2 \equiv 3 \pmod{4}$ , let  $h(-p_1)$ ,  $h(-p_2)$  and  $h(p_1p_2)$  be the class numbers of the quadratic fields  $\mathbb{Q}(\sqrt{-p_1})$ ,  $\mathbb{Q}(\sqrt{-p_2})$  and  $\mathbb{Q}(\sqrt{p_1p_2})$ , respectively. Let  $\omega_{p_1p_2} := (1 + \sqrt{p_1p_2})/2$  and let  $\Psi(\omega_{p_1p_2})$  be the Hirzebruch sum of  $\omega_{p_1p_2}$ . We show that  $h(-p_1)h(-p_2) \equiv h(p_1p_2)\Psi(\omega_{p_1p_2})/n \pmod{8}$ , where  $n = 6$  (respectively,  $n = 2$ ) if  $\min\{p_1, p_2\} > 3$  (respectively, otherwise). We also consider the real quadratic order with conductor 2 in  $\mathbb{Q}(\sqrt{p_1p_2})$ .

## 1. INTRODUCTION AND RESULTS

In this paper  $\Delta$  denotes a *quadratic discriminant*, which means that it is a non-square integer such that  $\Delta \equiv 0$  or  $1 \pmod{4}$ . It is called *fundamental* if it is the discriminant of a quadratic number field. Every quadratic discriminant  $\Delta$  is a unique square multiple of a fundamental discriminant and we write  $\Delta = f^2d$ , where  $f$  is a positive integer and  $d$  is the fundamental discriminant of the field  $K = \mathbb{Q}(\sqrt{d})$ . By  $h(\Delta)$  (respectively,  $h^+(\Delta)$ ) we denote the *wide* (respectively, *narrow*) *class number* of  $\mathcal{O}_\Delta$  the quadratic order with *conductor*  $f$  in  $K$ . Let  $\omega_\Delta := (\sigma_\Delta + \sqrt{\Delta})/2$ , where  $\sigma_\Delta = 0$  if  $\Delta$  is even and  $\sigma_\Delta = 1$  if  $\Delta$  is odd. For  $\Delta > 0$ , the *fundamental unit* of the quadratic order  $\mathcal{O}_\Delta$  is denoted by  $\varepsilon_\Delta$ . We have

$$\varepsilon_\Delta = \frac{t + u\sqrt{\Delta}}{2},$$

where  $t$  and  $u$  are positive integers satisfying  $t \equiv u \pmod{2}$ .

Any real quadratic irrational  $\xi$  has a continued fraction expansion of the form

$$\xi = \hat{v}_1 + \frac{1}{\hat{v}_2 + \frac{1}{\dots + \frac{1}{\hat{v}_k + \frac{1}{v_0 + \frac{1}{\dots + \frac{1}{v_{l-1} + \frac{1}{v_0 + \dots}}}}}}}$$

---

2010 *Mathematics Subject Classification.* Primary 11R29, Secondary 11A55, 11F20.

*Key words and phrases.* Class numbers, Quadratic fields, Hirzebruch sums.

with the *basic period*  $\{v_0, \dots, v_{l-1}\}$ . We write  $\xi = [\hat{v}_1, \dots, \hat{v}_k; \overline{v_0, \dots, v_{l-1}}]$ , and let

$$\Psi(\xi) := \begin{cases} \sum_{i=0}^{l-1} (-1)^{k+i} v_i & \text{if } l \text{ is even,} \\ 0 & \text{if } l \text{ is odd.} \end{cases}$$

This number  $\Psi(\xi)$  is said to be the *Hirzebruch sum* of  $\xi$ . When the *pre-period*  $\{\hat{v}_1, \dots, \hat{v}_k\}$  is empty, we understand  $k = 0$ . In the 1970s, Hirzebruch and Zagier [5, 13] obtained a beautiful formula on the class number of an imaginary quadratic field in terms of the Hirzebruch sum<sup>1</sup> (for the formula, see Remark 2). As an application of Zagier's formula, Chua et al. solved Richard Guy's conjecture (cf. [3, p. 1347]), which is equivalent to the following theorem.

**Theorem 1.1** ([3, Theorem 1.3]). *For any prime  $p \equiv 3 \pmod{4}$  with  $p > 3$ ,*

$$h(-p) \equiv h(4p) \frac{\Psi(\omega_{4p})}{3} \pmod{16}.$$

The result obtained by Zagier [13], however, does not work in the case of primes congruent to 1 modulo 4. Cheng and Guo [2] used results obtained by Lu [9], and they proved similar congruences modulo 8 (respectively, modulo 4) if  $p \equiv 1 \pmod{8}$  (respectively, if  $p \equiv 5 \pmod{8}$ ) and  $\varepsilon_p$  has integral coefficients).

The second named author and Kaneko [7] presented an explicit form of genus character  $L$ -functions of quadratic orders, and they generalized Zagier's results (for detail, see Section 2). As a consequence, the second named author showed a reasonable counterpart of Theorem 1.1.

**Theorem 1.2** ([10, Theorem 1]). *For any prime  $p \equiv 1 \pmod{4}$ ,*

$$h(-4p) \equiv h(16p) \frac{\Psi(\omega_{16p})}{3} \pmod{16}.$$

We remark that for a prime  $p \equiv 1 \pmod{4}$ ,  $h(16p)$  the class number of the quadratic order  $\mathcal{O}_{16p}$  equals to  $3h(p)$  (respectively,  $h(p)$ ) if  $p \equiv 5 \pmod{8}$  and  $\varepsilon_p$  has integral coefficients (respectively, otherwise) (cf. Lemma 3.8). The odd class number of  $\mathcal{O}_\Delta$  ( $\Delta > 0$ ) deserves special emphasis in both proofs of Theorems 1.1 and 1.2. For a fundamental discriminant  $d > 0$ , it is well-known that  $h(d)$  is odd if and only if

- (i) either  $d = 8$ ,
- (ii) or  $d = p$  for some prime  $p \equiv 1 \pmod{4}$ ,
- (iii) or  $d \in \{4p, 8p\}$  for some prime  $p \equiv 3 \pmod{4}$ ,
- (iv) or  $d = p_1 p_2$  for some distinct primes  $p_1 \equiv p_2 \equiv 3 \pmod{4}$

(cf. [6, Theorem 5.6.13]). In this paper, we study a similar problem regarding the odd class number of  $\mathbb{Q}(\sqrt{p_1 p_2})$  for distinct primes  $p_1 \equiv p_2 \equiv 3 \pmod{4}$ .

We state the main results of this paper as follows.

---

<sup>1</sup>In [3] and [10], the authors used the Zagier sum, which is defined by a *negative continued fraction*. We note that the Hirzebruch sum is three times the Zagier sum,  $3 \mid \Psi(\omega_{4p})$  in Theorem 1.1 and  $3 \mid \Psi(\omega_{16p})$  in Theorem 1.2 (for detail, see [2, p. 312], [10, p.353] and [8, Satz 2]).

**Theorem 1.3.** For distinct primes  $p_1 \equiv p_2 \equiv 3 \pmod{4}$ ,

$$h(-p_1)h(-p_2) \equiv h(p_1p_2) \frac{\Psi(\omega_{p_1p_2})}{n} \pmod{8},$$

where  $n = 6$  (respectively,  $n = 2$ ) if  $\min\{p_1, p_2\} > 3$  (respectively, otherwise).

**Theorem 1.4.** For distinct primes  $p_1 \equiv p_2 \equiv 3 \pmod{4}$ ,

$$h(-p_1)h(-p_2)\theta(-p_1, -p_2, 2) \equiv h(4p_1p_2) \frac{\Psi(\omega_{4p_1p_2})}{n} \pmod{8}.$$

Here  $n$  is defined as in Theorem 1.3 and  $\theta(-p_1, -p_2, 2)$  is defined by

$$\theta(-p_1, -p_2, 2) := (2 - \chi_{-p_1}(2))(2 - \chi_{-p_2}(2)) - (1 - \chi_{-p_1}(2))(1 - \chi_{-p_2}(2))$$

with the Kronecker symbols  $\chi_{-p_i} := \left(\frac{-p_i}{\cdot}\right)$ .

**Remark.** With the same notation as above, we see that  $h(-p_1)h(-p_2)$  is odd by Gauss' genus theory. We also note that  $n \mid \Psi(\omega_{p_1p_2})$  and  $n \mid \Psi(\omega_{4p_1p_2})$ , which can be inferred from [8, Satz 2] (see also Lemma 3.9 (i)). It is well-known that  $\chi_m(2) = (-1)^{(m^2-1)/8}$  for an odd integer  $m$ .

The outline of this paper is as follows. In Section 2, we recall the Kaneko-Mizuno-Zagier formula and the related notions. In Section 3, we review a calculation method of Hirzebruch sums via Dedekinds sums, and we show some facts on the fundamental units of  $\mathcal{O}_{p_1p_2}$  and  $\mathcal{O}_{4p_1p_2}$ . In Section 4, we prove Theorems 1.3 and 1.4 according to the cases whether  $\varepsilon_{p_1p_2}$  has half-integral coefficients and  $f = 1$  simultaneously or not.

## 2. THE KANEKO-MIZUNO-ZAGIER FORMULA

To state the results of the second named author and Kaneko [7], we recall the genus character on the narrow class group for a general discriminant (for detail, we refer to [1, Section 3] and [7, Appendix A.2]).

By  $F = [a, b, c]$  we denote the primitive quadratic form  $F(x, y) = ax^2 + bxy + cy^2$  with  $a, b, c \in \mathbb{Z}$  such that  $\gcd(a, b, c) = 1$ . We call  $\Delta := b^2 - 4ac$  the discriminant of  $F$ . As mentioned in Section 1, we assume that  $\Delta$  is not a square. The action  $F \mapsto AF$  of  $A = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$  is defined by  $(AF)(x, y) := F(\alpha x + \gamma y, \beta x + \delta y)$ . We denote by  $[F]$  the equivalence class of  $F$  with respect to the action  $F \mapsto AF$  of  $A \in \mathrm{SL}_2(\mathbb{Z})$ . Let  $\mathfrak{F}_\Delta$  be the set of all equivalence classes on “not negative-definite”<sup>2</sup> primitive forms of discriminant  $\Delta$ . By  $\mathcal{C}_\Delta^+$  (respectively,  $\mathcal{C}_\Delta$ ) we denote the narrow (respectively, wide) class group of a quadratic order with discriminant  $\Delta$ . We denote by  $[\mathfrak{a}]^+$  (respectively,  $[\mathfrak{a}]$ ) the narrow (respectively, wide) ideal class of an ideal  $\mathfrak{a}$  in  $\mathcal{O}_\Delta$ .

<sup>2</sup>It means “positive-definite” (respectively, “indefinite”) for  $\Delta < 0$  (respectively,  $\Delta > 0$ ).

**Proposition 2.1** ([6, Theorem 6.4.2]). *For a primitive form  $F = [a, b, c]$  of discriminant  $\Delta$ , let  $I(F) := (|a|, \frac{b+\sqrt{\Delta}}{2})\sqrt{\Delta}^{(1-\text{sign}(a))/2}$  be an ideal of  $\mathcal{O}_\Delta$ . The map  $\Phi_\Delta : \mathfrak{F}_\Delta \rightarrow \mathcal{C}_\Delta^+$  defined by*

$$\Phi_\Delta([F]) := [I(F)]^+$$

*is bijective. We define the composition  $[F_1] * [F_2] \in \mathfrak{F}_\Delta$  by*

$$[F_1] * [F_2] := \Phi_\Delta^{-1}(\Phi_\Delta([F_1])\Phi_\Delta([F_2])).$$

*With this composition,  $\mathfrak{F}_\Delta$  is an abelian group and  $\Phi_\Delta$  is an isomorphism.*

Using Proposition 2.1, we define the genus character as follows.

**Definition 2.2** ([7, Appendix A.2]). *Let  $\Delta = d_1 d_2 f^2$  be a discriminant, where  $d_1$  and  $d_2$  are distinct fundamental discriminants and  $f$  is a positive integer. We define the genus character  $\chi_{d_1, d_2}^{(\Delta)} : \mathfrak{F}_\Delta \rightarrow \{\pm 1\}$  as follows. We write  $d_1 = q_1^* q_2^* \cdots q_m^*$  with prime fundamental discriminants<sup>3</sup>  $q_i^*$ 's. For  $[F] \in \mathfrak{F}_\Delta$ , we define*

$$\chi_{d_1, d_2}^{(\Delta)}([F]) := \prod_{i=1}^m \chi^{(q_i^*)}([a, b, c]),$$

*where  $[a, b, c]$  is a representative of  $[F]$  and*

$$\chi^{(q_i^*)}([a, b, c]) := \begin{cases} \chi_{q_i^*}(a) & \text{if } \gcd(a, q_i^*) = 1, \\ \chi_{q_i^*}(c) & \text{if } \gcd(c, q_i^*) = 1, \end{cases}$$

*with  $\chi_{q_i^*}$  the quadratic character of  $\mathcal{O}_{q_i^*}$ . This map is a well-defined homomorphism and gives the genus character  $\chi_{d_1, d_2}^{(\Delta)} : \mathcal{C}_\Delta^+ \rightarrow \{\pm 1\}$  through  $\Phi_\Delta$  in Proposition 2.1.*

We review quadratic irrationals and their continued fractions (for detail, we refer to [6, Chapter 1 and Chapter 5.5]). Let  $\mathbb{X}_\Delta := \{\xi = \frac{b+\sqrt{\Delta}}{2a}; a \neq 0, b, c, \in \mathbb{Z}, \gcd(a, b, c) = 1, b^2 - 4ac = \Delta\}$  be the set of all *quadratic irrationals* of discriminant  $\Delta$ . The action  $\xi \mapsto A\xi$  of  $A = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \text{GL}_2(\mathbb{Z})$  is defined by  $A\xi := (\alpha\xi + \beta)/(\gamma\xi + \delta)$ . Two quadratic irrationals  $\xi_1, \xi_2 \in \mathbb{X}_\Delta$  are *properly equivalent* ( $\xi_1 \sim_+ \xi_2$ ) if there exists  $M \in \text{SL}_2(\mathbb{Z})$  such that  $\xi_1 = M\xi_2$ . Two quadratic irrationals  $\xi_1, \xi_2 \in \mathbb{X}_\Delta$  are *equivalent* ( $\xi_1 \sim \xi_2$ ) if there exists  $M \in \text{GL}_2(\mathbb{Z})$  such that  $\xi_1 = M\xi_2$ . We denote by  $\mathfrak{X}_\Delta^+$  (respectively,  $\mathfrak{X}_\Delta$ ) the set of all proper equivalence (respectively, equivalence) classes of  $\mathbb{X}_\Delta$ .

**Proposition 2.3** ([6, Theorem 5.5.8]). *For a quadratic irrational  $\xi = \frac{b+\sqrt{\Delta}}{2a}$ , let  $I(\xi) := (|a|, \frac{b+\sqrt{\Delta}}{2})\sqrt{\Delta}^{(1-\text{sign}(a))/2}$  be an ideal of  $\mathcal{O}_\Delta$ . The map  $\iota_\Delta^+ : \mathfrak{X}_\Delta^+ \rightarrow \mathcal{C}_\Delta^+$  defined by*

$$\iota_\Delta^+([\xi]_{\sim_+}) = [I(\xi)]^+$$

*is bijective. Also, the map  $\iota_\Delta : \mathfrak{X}_\Delta \rightarrow \mathcal{C}_\Delta$  defined by  $\iota_\Delta([\xi]_{\sim}) = [I(\xi)]$  is bijective.*

<sup>3</sup>For any odd prime  $q_i > 0$ ,  $q_i^* := (-1)^{\frac{q_i-1}{2}} q_i$ . If  $d_1$  is even,  $q_1^* = 2^* \in \{-4, \pm 8\}$  is defined by  $d_1 = q_1^* q_2^* \cdots q_m^*$ .

By  $\xi'$  we denote the *conjugate* of  $\xi \in \mathbb{X}_\Delta$ . For  $\Delta > 0$ , let  $\mathbb{X}_\Delta^0 := \{\xi \in \mathbb{X}_\Delta; -1 < \xi' < 0, 1 < \xi\}$  be the set of all *reduced quadratic irrationals* of discriminant  $\Delta$ . We denote the *continued fraction* of  $\xi \in \mathbb{R}$  by  $\xi = [u_0, u_1, u_2, \dots]$  with  $u_0 \in \mathbb{Z}$  and  $u_i \in \mathbb{N}$  for  $i \geq 1$ . We define the number  $\xi_n := [u_n, u_{n+1}, u_{n+2}, \dots]$  for  $n \geq 0$ . We denote by  $\mathcal{N} : \mathbb{Q}(\sqrt{\Delta}) \rightarrow \mathbb{Q}$  the field norm and by  $\varepsilon_\Delta$  the fundamental unit of  $\mathcal{O}_\Delta$ . The following proposition provides some properties of continued fractions.

**Proposition 2.4** ([6, Theorems 2.2.2, 2.2.9 and 2.3.3]). *Let  $\xi = [u_0, u_1, u_2, \dots]$  be a continued fraction of an irrational  $\xi \in \mathbb{R} \setminus \mathbb{Q}$ .*

(i) *The sequence  $(u_n)_{n \geq 0}$  is ultimately periodic if and only if  $\xi$  is a quadratic irrational, and it is periodic if and only if  $\xi$  is a reduced quadratic irrational.*

(ii) *If  $\xi \in \mathbb{X}_\Delta$ , then  $\xi_n \in \mathbb{X}_\Delta$  and  $\xi_n \sim_+ (-1)^n \xi$ .*

(iii) *Let  $\xi = [\hat{v}_1, \dots, \hat{v}_k; \overline{v_0, \dots, v_{l-1}}] \in \mathbb{X}_\Delta$ . The  $l$  distinct numbers  $\xi_k, \xi_{k+1}, \dots, \xi_{k+l-1}$  are all reduced quadratic irrationals which are equivalent to  $\xi$ .*

(iv) *In the notation above, the period length  $l$  is odd if and only if  $\mathcal{N}(\varepsilon_\Delta) = -1$ .*

(v) *(The inverse period) If  $\xi = [\overline{v_0, v_1, \dots, v_{l-1}}]$ , then  $-\xi'^{-1} = [\overline{v_{l-1}, v_{l-2}, \dots, v_0}]$ .*

Now we introduce the main results of [7].

**Theorem 2.5** ([7, Lemma 10 (2) and Theorem 3]). *Let  $\Delta = d_1 d_2 f^2$  be a discriminant with distinct negative fundamental discriminants  $d_1$  and  $d_2$ , and a positive integer  $f$ . Let  $\varepsilon_\Delta$  be the fundamental unit of  $\mathcal{O}_\Delta$  and  $\omega(d_i)$  the number of the roots of unity in  $\mathcal{O}_{d_i}$ . We define*

$$\theta(d_1, d_2, f) := \prod_{\substack{p|f \\ p: \text{prime}}} \frac{(1 - \chi_{d_1}(p))(1 - \chi_{d_2}(p)) - p^{m_p-1}(p - \chi_{d_1}(p))(p - \chi_{d_2}(p))}{1 - p},$$

where  $m_p$  is a positive integer such that  $p^{m_p}$  is the highest power of  $p$  dividing  $f$  if  $f > 1$ . The empty product is understood as being 1 if  $f = 1$ . Then we have  $\mathcal{N}(\varepsilon_\Delta) = 1$  and

$$24 \frac{h(d_1)h(d_2)}{\omega(d_1)\omega(d_2)} \theta(d_1, d_2, f) = \sum_{\xi \in \mathbb{X}_\Delta^0} \chi_{d_1, d_2}^{(\Delta)}([I(\xi)]^+) [\xi], \quad (1)$$

where  $\chi_{d_1, d_2}^{(\Delta)}$  is defined in Definition 2.2 and  $[I(\xi)]^+$  is defined in Proposition 2.3.

Under the same assumption as in Theorem 2.5, we will restate the equality (1) in terms of  $\mathfrak{X}_\Delta$  and the Hirzebruch sum  $\Psi$ . Let  $\eta := \frac{b+\sqrt{\Delta}}{2a} \in \mathbb{X}_\Delta$ . By Propositions 2.3, 2.1 and  $d_1 < 0$ , we have

$$\begin{aligned} \chi_{d_1, d_2}^{(\Delta)}([I(-\eta)]^+) &= \chi_{d_1, d_2}^{(\Delta)}([-a, b, -c]) \\ &= \chi_{d_1}(-1) \chi_{d_1, d_2}^{(\Delta)}([a, b, c]) \\ &= -\chi_{d_1, d_2}^{(\Delta)}([I(\eta)]^+). \end{aligned} \quad (2)$$

By Proposition 2.4 (ii), (iii) and (iv), for any  $\eta = [\hat{v}_1, \dots, \hat{v}_k; \overline{v_0, \dots, v_{2m-1}}] \in \mathbb{X}_\Delta$ , we have  $\{\xi \in \mathbb{X}_\Delta^0 \mid \xi \sim_+ (-1)^k \eta\} = \{\eta_k, \eta_{k+2}, \dots, \eta_{k+2m-4}, \eta_{k+2m-2}\}$  and  $\{\xi \in \mathbb{X}_\Delta^0 \mid \xi \sim_+ (-1)^{k+1} \eta\} =$

$\{\eta_{k+1}, \eta_{k+3}, \dots, \eta_{k+2m-3}, \eta_{k+2m-1}\}$ . Hence we have

$$\begin{aligned} \sum_{\substack{\xi \in \mathfrak{X}_{\Delta}^0 \\ \xi \sim \eta}} \chi_{d_1, d_2}^{(\Delta)}([I(\xi)]^+) [\xi] &= \chi_{d_1, d_2}^{(\Delta)}([I(\eta_k)]^+) \Psi(\eta_k) \\ &= \left( (-1)^k \chi_{d_1, d_2}^{(\Delta)}([I(\eta)]^+) \right) \left( (-1)^k \Psi(\eta) \right) \end{aligned}$$

since  $[\eta_{k+i}] = v_i$  for  $0 \leq i \leq 2m-1$  and (2). Therefore, we get

$$24 \frac{h(d_1)h(d_2)}{\omega(d_1)\omega(d_2)} \theta(d_1, d_2, f) = \sum_{[\eta] \sim \mathfrak{X}_{\Delta}} \chi_{d_1, d_2}^{(\Delta)}([I(\eta)]^+) \Psi(\eta), \quad (3)$$

which we call the *Kaneko-Mizuno-Zagier formula*.

**Remark.** Zagier [13, (23)] proved the equality (3) in the case where  $\gcd(d_1, d_2) = 1$  and  $f = 1$ . We remark that Lu [9] obtained a different kind of formula for  $h(d_1)h(d_2)$  in the case where  $d_1 \mid d_2$  and  $f = 1$  (cf. [2, Theorem 2.12]).

### 3. PRELIMINARY LEMMAS

In this section, we show some lemmas required in the proofs of Theorems 1.3 and 1.4.

**3.1. Hirzebruch sums via Dedekind sums.** To introduce one of the calculation methods of Hirzebruch sums, we review some properties of Dedekind sums (for detail, we refer to [11]).

**Definition 3.1.** Let  $h$  and  $k$  be integers such that  $\gcd(h, k) = 1$  and  $k \geq 1$ . Let

$$s(h, k) := \sum_{m=1}^k \left( \left( \frac{hm}{k} \right) \right) \left( \left( \frac{m}{k} \right) \right),$$

where

$$((x)) := \begin{cases} x - [x] - \frac{1}{2} & \text{if } x \notin \mathbb{Z}, \\ 0 & \text{if } x \in \mathbb{Z}. \end{cases}$$

We call this number  $s(h, k)$  the *Dedekind sum*.

**Proposition 3.2.** Let  $h$  and  $k$  be integers such that  $\gcd(h, k) = 1$  and  $k \geq 1$ .

- (i)  $s(-h, k) = -s(h, k)$ .
- (ii)  $s(h, k)$  depends only on  $h \pmod{k}$ .
- (iii) (The reciprocity law) If  $h \geq 1$ , then  $s(h, k) + s(k, h) = \frac{h^2+k^2+1}{12hk} - \frac{1}{4}$ .
- (iv) For an integer  $h'$  such that  $hh' \equiv 1 \pmod{k}$ ,  $s(h, k) = s(h', k)$ .
- (v)  $6ks(h, k)$  is an integer.
- (vi) If  $k$  is odd, then  $\left( \frac{h}{k} \right) = (-1)^{\frac{1}{2}(\frac{k-1}{2} - 6ks(h, k))}$ , where  $\left( - \right)$  is the Jacobi symbol.

*Proof.* (i) and (ii) follow from Definition 3.1. We refer (iii) to Theorem 1 in [11, Chapter 2]. We refer (iv), (v) and (vi) to the equation (33c), Theorem 2 and Theorem 4 in [11, Chapter 3], respectively.  $\square$

We define a subset  $\mathfrak{M}$  of  $\mathrm{SL}_2(\mathbb{Z})$  by

$$\mathfrak{M} := \left\{ M : M = \begin{pmatrix} x & y \\ z & w \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}) \text{ such that } z \neq 0 \right\}.$$

For any  $M = \begin{pmatrix} x & y \\ z & w \end{pmatrix} \in \mathfrak{M}$ , we define

$$n_M := \frac{x+w}{z} - \mathrm{sign}(z)(3 + 12s(w, |z|)),$$

where  $\mathrm{sign}(z) := z/|z|$ . The following three lemmas will be used to prove Lemmas 4.3 and 4.4 in Section 4.

**Lemma 3.3** ([7, Lemma 8]). *Let  $M = \begin{pmatrix} x & y \\ z & w \end{pmatrix} \in \mathfrak{M}$  such that  $x \neq 0$ , and let  $M' = \begin{pmatrix} k & -1 \\ 1 & 0 \end{pmatrix} M$  for some integer  $k$ . Then it follows that*

$$n_{M'} = \begin{cases} k + 9 + n_M & \text{if } x < 0 \text{ and } z > 0, \\ k - 3 + n_M & \text{if } x > 0 \text{ or } z < 0. \end{cases}$$

**Lemma 3.4.** *Let  $M = \begin{pmatrix} x & y \\ z & w \end{pmatrix} \in \mathfrak{M}$ , and let  $M' = \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix} M \begin{pmatrix} 1 & -k \\ 0 & 1 \end{pmatrix}$  for some integer  $k$ . Then one has  $n_{M'} = n_M$ .*

*Proof.* We have  $M' = \begin{pmatrix} x+kz & -kx+y-k^2z+kw \\ z & -kz+w \end{pmatrix} \in \mathfrak{M}$  and  $n_{M'} = \frac{x+w}{z} - \mathrm{sign}(z)(3 + 12s(-kz+w, |z|))$ . By Proposition 3.2 (ii) we have  $n_{M'} = n_M$ .  $\square$

**Lemma 3.5.** *Let  $M = \begin{pmatrix} x & y \\ z & w \end{pmatrix} \in \mathfrak{M}$  such that  $x > 0$  and  $y \neq 0$ . Let  $M' = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} M \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ . Then we have  $n_{M'} = -n_M$ .*

*Proof.* We have  $M' = \begin{pmatrix} w & z \\ y & x \end{pmatrix} \in \mathfrak{M}$ . By Proposition 3.2 (iv) we have  $s(w, |z|) = s(x, |z|)$ . By Proposition 3.2 (iii) we have  $s(x, |z|) = -s(|z|, x) + \frac{x^2+z^2+1}{12x|z|} - \frac{1}{4}$ . By Proposition 3.2 (i) we have  $n_M = \frac{x+w}{z} + 12s(z, x) - \frac{x^2+z^2+1}{xz}$ . Similarly, by Proposition 3.2 (iii) and (i), we have  $n_{M'} = \frac{x+w}{y} + 12s(y, x) - \frac{x^2+y^2+1}{xy}$ . By Proposition 3.2 (iv) and (i), we see that  $s(z, x) = s(-y, x) = -s(y, x)$ . Therefore, we have  $n_M + n_{M'} = \frac{(y+z)(xw-yz-1)}{xyz} = 0$ .  $\square$

Now we introduce a calculation method of Hirzebruch sums via Dedekind sums.  $\Delta$  always stands for a positive discriminant. Let  $\omega_\Delta := (\sigma_\Delta + \sqrt{\Delta})/2$ , where  $\sigma_\Delta = 0$  if  $\Delta$  is even and  $\sigma_\Delta = 1$  if  $\Delta$  is odd. We write the fundamental unit  $\varepsilon_\Delta = q + r\omega_\Delta$  with positive integers  $q$  and  $r$ . For  $\xi \in \mathbb{X}_\Delta$ , we define  $M_\xi$  by a unique integral matrix  $M_\xi := \begin{pmatrix} x & y \\ z & w \end{pmatrix}$  such that

$$\begin{cases} \varepsilon_\Delta \xi &= x\xi + y, \\ \varepsilon_\Delta &= z\xi + w. \end{cases}$$



From the definition, we easily see that

$$M_{\omega_\Delta} = \begin{pmatrix} q + r\sigma_\Delta & r(\Delta - \sigma_\Delta)/4 \\ r & q \end{pmatrix}. \quad (4)$$

For  $\xi = \frac{b+\sqrt{\Delta}}{2a} \in \mathbb{X}_\Delta$ , we see that  $\xi = \begin{pmatrix} 1 & (b-\sigma_\Delta)/2 \\ 0 & a \end{pmatrix} \omega_\Delta$  and

$$M_\xi = \begin{pmatrix} 1 & (b-\sigma_\Delta)/2 \\ 0 & a \end{pmatrix} M_{\omega_\Delta} \begin{pmatrix} 1 & (b-\sigma_\Delta)/2 \\ 0 & a \end{pmatrix}^{-1}. \quad (5)$$

We note that  $M_\xi \in \mathfrak{M}$  for any  $\xi \in \mathbb{X}_\Delta$ . For  $\xi \in \mathbb{X}_\Delta$ , we define  $n(\xi) := n_{M_\xi}$ .

**Lemma 3.6.** *Let  $\Delta$  be a positive discriminant such that  $\mathcal{N}(\varepsilon_\Delta) = 1$ . For any  $\xi \in \mathbb{X}_\Delta^0$ , we have*

$$\Psi(\xi) = n(\xi).$$

*Proof.* The above equality was proved in the middle of the proof of Theorem 2 in [7, Section 5]. The assumption of [7, Theorem 2] is that  $\Delta$  is of the form  $d_1 d_2 f^2$ , where  $d_1$  and  $d_2$  are distinct negative fundamental discriminants, and  $f$  is a positive integer. However, we can see that the above equality still holds under the weaker assumption, that is,  $\mathcal{N}(\varepsilon_\Delta) = 1$ .  $\square$

**Lemma 3.7.** *Let  $\Delta$  be a positive discriminant such that  $\mathcal{N}(\varepsilon_\Delta) = 1$ . Let  $\omega_\Delta = (\sigma_\Delta + \sqrt{\Delta})/2$ . Then we have*

$$\Psi(\omega_\Delta) = n(\omega_\Delta).$$

*Proof.* We have  $(\omega_\Delta)_1 = 1/(\omega_\Delta - \lfloor \omega_\Delta \rfloor) \in \mathbb{X}_\Delta$ , where  $(\omega_\Delta)_1$  denotes the shift by 1 as given in the notation of Proposition 2.4. Since  $0 < \omega_\Delta - \lfloor \omega_\Delta \rfloor < 1$  and  $\omega'_\Delta - \lfloor \omega_\Delta \rfloor < -1$ , we see that  $(\omega_\Delta)_1 \in \mathbb{X}_\Delta^0$ . By Lemma 3.6 we have  $\Psi((\omega_\Delta)_1) = n((\omega_\Delta)_1)$ . From the definition,  $\Psi(\omega_\Delta) = -\Psi((\omega_\Delta)_1)$ . Let  $A := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $B := \begin{pmatrix} 1 & -\lfloor \omega_\Delta \rfloor \\ 0 & 1 \end{pmatrix}$ . We see that  $(\omega_\Delta)_1 = AB\omega_\Delta$  and  $M_{(\omega_\Delta)_1} = ABM_{\omega_\Delta}B^{-1}A^{-1}$ . By (4) we have

$$BM_{\omega_\Delta}B^{-1} = \begin{pmatrix} q + r\sigma_\Delta - r\lfloor \omega_\Delta \rfloor & -\lfloor \omega_\Delta \rfloor^2 + \sigma_\Delta \lfloor \omega_\Delta \rfloor + (\Delta - \sigma_\Delta)/4 \\ r & q + r\lfloor \omega_\Delta \rfloor \end{pmatrix} \in \mathfrak{M}$$

and  $q + r\sigma_\Delta - r\lfloor \omega_\Delta \rfloor > \varepsilon'_\Delta > 0$ . Since  $\omega_\Delta$  and  $\omega'_\Delta$  are two distinct roots of  $X^2 - \sigma_\Delta X + (\sigma_\Delta - \Delta)/4$ , the  $(1, 2)$ -component of  $BM_{\omega_\Delta}B^{-1}$  is not equal to 0. By Lemma 3.4 and Lemma 3.5, we have  $n_{M_{\omega_\Delta}} = n_{BM_{\omega_\Delta}B^{-1}} = -n_{ABM_{\omega_\Delta}B^{-1}A^{-1}} = -n_{M_{(\omega_\Delta)_1}}$ . Therefore, we have  $\Psi(\omega_\Delta) = -\Psi((\omega_\Delta)_1) = -n((\omega_\Delta)_1) = n(\omega_\Delta)$ .  $\square$

**3.2. Fundamental units of quadratic orders.** The first two lemmas are standard facts on quadratic orders.

**Lemma 3.8** ([6, p. 328]). *Let  $\Delta = f^2 d$  be a discriminant with a fundamental discriminant  $d$  and a conductor  $f$ . Then the class numbers of  $\mathcal{O}_d$  and  $\mathcal{O}_\Delta$  have the following relation:*

$$h(f^2 d) = h(d) \frac{f}{[\mathcal{O}_d^\times : \mathcal{O}_{f^2 d}^\times]} \prod_{\substack{q|f \\ q: \text{prime}}} (1 - \chi_d(q)q^{-1}).$$



**Lemma 3.9.** *Let  $p_1$  and  $p_2$  be distinct primes such that  $p_1 \equiv p_2 \equiv 3 \pmod{4}$ . Let  $\varepsilon_{p_1 p_2} = \frac{t+u\sqrt{p_1 p_2}}{2}$  be the fundamental unit of  $\mathcal{O}_{p_1 p_2}$  such that  $t \equiv u \pmod{2}$ . Then we have*

- (i)  $h(p_1 p_2)$  is odd and  $\mathcal{N}(\varepsilon_{p_1 p_2}) = 1$ .
- (ii) If  $p_1 p_2 \equiv 1 \pmod{8}$ , then  $t \equiv u \equiv 0 \pmod{2}$ .
- (iii)  $(h(4p_1 p_2), \varepsilon_{4p_1 p_2})$  is equal to

$$\begin{cases} (h(p_1 p_2), \varepsilon_{p_1 p_2}) & \text{if } p_1 p_2 \equiv 1 \pmod{8}, \\ (3h(p_1 p_2), \varepsilon_{p_1 p_2}) & \text{if } p_1 p_2 \equiv 5 \pmod{8} \text{ and } t \equiv u \equiv 0 \pmod{2}, \\ (h(p_1 p_2), \varepsilon_{p_1 p_2}^3) & \text{if } p_1 p_2 \equiv 5 \pmod{8} \text{ and } t \equiv u \equiv 1 \pmod{2}. \end{cases}$$

*Proof.* (i) We refer to [6, Theorem 5.6.13 and Theorem 5.2.2].

(ii) We refer to [6, Theorem 5.2.3].

(iii) If  $t \equiv u \equiv 0 \pmod{2}$ , then  $\varepsilon_{4p_1 p_2} = \varepsilon_{p_1 p_2}$ . Now suppose that  $t \equiv u \equiv 1 \pmod{2}$ . By [6, Theorem 5.2.3] we have  $p_1 p_2 \equiv 5 \pmod{8}$  and  $\varepsilon_{4p_1 p_2} = \varepsilon_{p_1 p_2}^3$ . The results on  $h(4p_1 p_2)$  follow from Lemma 3.8 and  $\chi_d(2) = (-1)^{(d^2-1)/8}$  for odd  $d$ .  $\square$

By modifying the results of [12] and [14], we show the following properties on the fundamental integral units  $\varepsilon_{p_1 p_2}$  and  $\varepsilon_{4p_1 p_2}$ .

**Lemma 3.10.** *Let  $p_1$  and  $p_2$  be distinct primes such that  $p_1 \equiv p_2 \equiv 3 \pmod{4}$ . Let  $f = 2$  if  $p_1 p_2 \equiv 5 \pmod{8}$  and  $\varepsilon_{p_1 p_2}$  has half-integral coefficients, and let  $f = 1$  or  $2$  otherwise. Let  $\Delta = p_1 p_2 f^2$  and  $\varepsilon_\Delta = x + y\sqrt{p_1 p_2}$  be the fundamental unit of  $\mathcal{O}_\Delta$ . Then we have*

- (i)  $x, y$  are integers such that  $x \equiv 7 \pmod{8}$  and  $y \equiv 0 \pmod{4}$ . Moreover,  $y \equiv 4 \pmod{8}$  if  $(p_1, p_2) \equiv (3, 3) \pmod{8}$ , and  $y \equiv 0 \pmod{8}$  if  $(p_1, p_2) \equiv (7, 7) \pmod{8}$ .
- (ii)  $p_1 \varepsilon_\Delta = (X + Y\sqrt{p_1 p_2})^2$  for some integers  $X$  and  $Y$ .

*Proof.* (i) We refer to [12, Theorem 2].

(ii) By [14, Lemma 3.2] there exist  $X, Y \in \mathbb{Q}$  such that  $p_1(x + y\sqrt{p_1 p_2}) = (X + Y\sqrt{p_1 p_2})^2$ . We have  $p_1 x = X^2 + p_1 p_2 Y^2$  and  $p_1 y = 2XY$ ; so, we get

$$p_1 p_2 Y^4 - p_1 x Y^2 + \frac{p_1^2 y^2}{4} = 0.$$

Then  $Y^2 = (x + \alpha)/(2p_2)$  with  $\alpha \in \{\pm 1\}$  since  $x^2 - p_1 p_2 y^2 = 1$ . It follows that  $2 \mid (x + \alpha)$  and  $p_2 \mid (x + \alpha)$ ; thus we have that  $Y^2 \in \mathbb{Z}$  and  $X^2 = p_1 x - p_1 p_2 Y^2 \in \mathbb{Z}$ . Therefore,  $X$  and  $Y$  are integers.  $\square$

We use Dirichlet's method [4] (see also [12]) to prove the following lemma about the fundamental half-integral unit  $\varepsilon_{p_1 p_2}$ .

**Lemma 3.11.** *Let  $p_1$  and  $p_2$  be distinct primes such that  $p_1 \equiv p_2 \equiv 3 \pmod{4}$ . Assume that  $p_1 p_2 \equiv 5 \pmod{8}$  and  $\varepsilon_{p_1 p_2} = \frac{t+u\sqrt{p_1 p_2}}{2}$  with  $t \equiv u \equiv 1 \pmod{2}$ . Then we have  $t \equiv 1 \pmod{4}$ .*

*Proof.* We note that  $(t, u) \in \mathbb{N}^2$  is the smallest nontrivial solution of  $x^2 - p_1 p_2 y^2 = \pm 4$ , and  $t^2 - p_1 p_2 u^2 = 4$ . Since  $(t-2)(t+2) = p_1 p_2 u^2$  and  $(t-2, t+2) = 1$ , there exist odd positive

integers  $r$  and  $s$  satisfying one of the following:

$$(t-2, t+2) = \begin{cases} (p_1 r^2, p_2 s^2), \\ (p_2 r^2, p_1 s^2), \\ (p_1 p_2 r^2, s^2), \\ (r^2, p_1 p_2 s^2). \end{cases}$$

For the third case (resp., the fourth case) we have that  $s^2 - p_1 p_2 r^2 = 4$  (resp.,  $r^2 - p_1 p_2 s^2 = -4$ ), which is a contradiction with the minimality of  $(t, u)$ . Therefore, we have that  $t = 2 + p_i r^2 \equiv 1 \pmod{4}$ , where  $i = 1$  or  $2$ .  $\square$

#### 4. PROOFS OF THEOREMS 1.3 AND 1.4

Let  $p_1$  and  $p_2$  be distinct primes congruent to 3 modulo 4. Under the same notation as in Theorem 2.5, let  $d_1 = -p_1$  and  $d_2 = -p_2$ , and let  $f = 1$  or  $2$ . By the Kaneko-Mizuno-Zagier formula (3) we have

$$\begin{aligned} & h(-p_1)h(-p_2)\theta(-p_1, -p_2, f) \\ = & \begin{cases} \sum_{[\eta] \sim \mathfrak{X}_{p_1 p_2 f^2}} \chi_{-p_1, -p_2}^{(p_1 p_2 f^2)}([I(\eta)]^+) \frac{\Psi(\eta)}{6} & \text{if } \min\{p_1, p_2\} > 3, \\ \sum_{[\eta] \sim \mathfrak{X}_{p_1 p_2 f^2}} \chi_{-p_1, -p_2}^{(p_1 p_2 f^2)}([I(\eta)]^+) \frac{\Psi(\eta)}{2} & \text{if } \min\{p_1, p_2\} = 3. \end{cases} \end{aligned} \quad (6)$$

Moreover, we remark that if  $\min\{p_1, p_2\} > 3$  (resp.,  $\min\{p_1, p_2\} = 3$ ), then  $\Psi(\eta)/6$  (resp.,  $\Psi(\eta)/2$ ) is an integer (cf. [8, Satz 2] and Lemma 3.9 (i)).

To show Theorems 1.3 and 1.4, we prove the following four lemmas which are similar to those in [3] and [10].

**Lemma 4.1.** *Following the above notation, for any equivalence class  $[\eta]_{\sim}$  in  $\mathfrak{X}_{p_1 p_2 f^2}$ , there exists a representative  $\frac{\beta + \sqrt{p_1 p_2 f^2}}{2\alpha} \in \mathbb{X}_{p_1 p_2 f^2}^0$  of  $[\eta]_{\sim}$  such that  $(a, p_1) = 1$  and  $a > 0$ .*

*Proof.* By Proposition 2.4 (iii) there exists a reduced quadratic irrational  $\xi = \frac{\beta + \sqrt{p_1 p_2 f^2}}{2\alpha} \in \mathbb{X}_{p_1 p_2 f^2}^0$  such that  $\xi \sim \eta$ . Since  $\xi' < \xi$ ,  $\alpha$  is a positive integer. If  $(\alpha, p_1) = 1$ , we take  $(a, b) = (\alpha, \beta)$ . Now, we suppose that  $p_1$  divides  $\alpha$ . Let  $\gamma = \frac{\beta^2 - p_1 p_2 f^2}{4\alpha} \in \mathbb{Z}$ . We have that  $p_1$  divides  $\beta$  and  $(\gamma, p_1) = 1$  since  $\gcd(\alpha, \beta, \gamma) = 1$ . With the same notation as in Proposition 2.4, let  $\xi_1 = \frac{\beta_1 + \sqrt{p_1 p_2 f^2}}{2\alpha_1} \in \mathbb{X}_{p_1 p_2 f^2}^0$ . We see that  $\alpha_1 > 0$ ,  $\xi_1 \sim \xi \sim \eta$  and

$$\xi_1 = \frac{1}{\xi - [\xi]} = \frac{2\alpha}{\beta - 2\alpha[\xi] + \sqrt{p_1 p_2 f^2}} = \frac{-\beta + 2\alpha[\xi] + \sqrt{p_1 p_2 f^2}}{-2\gamma + 2\beta[\xi] - 2\alpha[\xi]^2}.$$

Therefore,  $\alpha_1 = -\gamma + \beta[\xi] - \alpha[\xi]^2$  is coprime to  $p_1$ . We take  $(a, b) = (\alpha_1, \beta_1)$ .  $\square$

**Lemma 4.2.** *Following the above notation, let  $\eta$  and  $\delta$  be two quadratic irrationals of discriminant  $p_1 p_2 f^2$  such that  $[I(\eta)] = [I(\delta)]^{-1}$  in  $\mathcal{C}_{p_1 p_2 f^2}$ . Then we have*

$$\chi_{-p_1, -p_2}^{(p_1 p_2 f^2)}([I(\eta)]^+) \Psi(\eta) = \chi_{-p_1, -p_2}^{(p_1 p_2 f^2)}([I(\delta)]^+) \Psi(\delta).$$

*Proof.* By Lemma 4.1 there exists  $\xi = \frac{b+\sqrt{p_1p_2f^2}}{2a} \in \mathbb{X}_{p_1p_2f^2}^0$  such that  $\xi \sim \eta$ ,  $(a, p_1) = 1$  and  $a > 0$ . Let  $\xi^{\text{op}} := [\xi] - \xi' = [\xi] + \frac{-b+\sqrt{p_1p_2f^2}}{2a}$ . Since  $I(\xi) = (a, \frac{b+\sqrt{p_1p_2f^2}}{2})$  and  $I(\xi^{\text{op}}) = (a, \frac{-b+\sqrt{p_1p_2f^2}}{2})$ , we have  $[I(\xi)] = [I(\xi^{\text{op}})]^{-1}$  in  $\mathcal{C}_{p_1p_2f^2}$  and  $\xi^{\text{op}} \sim \delta$ . By Lemma 3.8 (i), (iii) and Proposition 2.4 (iv), the period length of  $\xi$  is even. We write  $\xi = [\overline{v_0, v_1, \dots, v_{2m-1}}]$ . Then we have  $\xi^{\text{op}} = [\overline{v_0, v_{2m-1}, v_{2m-2}, \dots, v_2, v_1}]$  by Proposition 2.4 (v). We see that  $\chi_{-p_1, -p_2}^{(p_1p_2f^2)}([I(\xi)]^+) = \chi_{-p_1}(a) = \chi_{-p_1, -p_2}^{(p_1p_2f^2)}([I(\xi^{\text{op}})]^+)$  and  $\Psi(\xi) = \sum_{i=0}^{2m-1} (-1)^i v_i = \Psi(\xi^{\text{op}})$ . Therefore, we have

$$\chi_{-p_1, -p_2}^{(p_1p_2f^2)}([I(\eta)]^+) \Psi(\eta) = \chi_{-p_1}(a) \sum_{i=0}^{2m-1} (-1)^i v_i = \chi_{-p_1, -p_2}^{(p_1p_2f^2)}([I(\delta)]^+) \Psi(\delta)$$

as desired.  $\square$

**Lemma 4.3.** *Following the above notation, we further assume that  $f = 2$  if  $p_1p_2 \equiv 5 \pmod{8}$  and  $\varepsilon_{p_1p_2}$  has half-integral coefficients, and we assume that  $f = 1$  or  $2$  otherwise. For any  $\eta \in \mathbb{X}_{p_1p_2f^2}$ , we have*

$$\chi_{-p_1, -p_2}^{(p_1p_2f^2)}([I(\eta)]^+) \Psi(\eta) \equiv \Psi(\omega_{p_1p_2f^2}) \pmod{8}.$$

*Proof.* With the same notation as in the paragraph before Lemma 3.6, we write  $\sigma_\Delta$ ,  $\omega_\Delta$  and  $\varepsilon_\Delta = q + r\omega_\Delta$ , where  $\Delta = p_1p_2f^2$  and  $q, r$  are positive integers. By Lemma 4.1 there exists  $\xi = \frac{b+\sqrt{p_1p_2f^2}}{2a} \in \mathbb{X}_{p_1p_2f^2}^0$  such that  $\xi \sim \eta$ ,  $(a, p_1) = 1$  and  $a > 0$ . Let  $c = \frac{b^2 - p_1p_2f^2}{4a} \in \mathbb{Z}$ , and let  $m$  be an integer such that  $b = 2m + \sigma_\Delta$ . Let  $A := \begin{pmatrix} 1 & m \\ 0 & 1 \end{pmatrix}$  and  $B := \begin{pmatrix} 3 & -1 \\ 1 & 0 \end{pmatrix}$ . By Lemma 3.9 we have  $\mathcal{N}(\varepsilon_\Delta) = 1$ . By (4) we have

$$\begin{aligned} AM_{\omega_\Delta} A^{-1} &= \begin{pmatrix} 1 & m \\ 0 & 1 \end{pmatrix} \begin{pmatrix} q + r\sigma_\Delta & r(\Delta - \sigma_\Delta)/4 \\ r & q \end{pmatrix} \begin{pmatrix} 1 & -m \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} q + rm + r\sigma_\Delta & -rac \\ r & q - rm \end{pmatrix} \end{aligned}$$

and

$$BAM_{\omega_{p_1p_2}} A^{-1} = \begin{pmatrix} 3(q + rm + r\sigma_\Delta) - r & -3rac - (q - rm) \\ q + rm + r\sigma_\Delta & -rac \end{pmatrix}.$$

By (5) we have

$$\begin{aligned} M_\xi &= \begin{pmatrix} 1 & m \\ 0 & a \end{pmatrix} \begin{pmatrix} q + r\sigma_\Delta & r(\Delta - \sigma_\Delta)/4 \\ r & q \end{pmatrix} \begin{pmatrix} 1 & -\frac{m}{a} \\ 0 & \frac{1}{a} \end{pmatrix} \\ &= \begin{pmatrix} q + rm + r\sigma_\Delta & -rc \\ ra & q - rm \end{pmatrix} \end{aligned}$$

and

$$BM_\xi = \begin{pmatrix} 3(q + rm + r\sigma_\Delta) - ra & -3rc - (q - rm) \\ q + rm + r\sigma_\Delta & -rc \end{pmatrix}.$$

Since  $\xi = \frac{b + \sqrt{p_1 p_2 f^2}}{2a} \in \mathbb{X}_{p_1 p_2 f^2}^0$ , we have  $2m + \sigma_\Delta > -\sqrt{\Delta}$  and  $q + rm + r\sigma_\Delta > \varepsilon'_\Delta > 0$ . By Lemma 3.7, Lemma 3.4 and Lemma 3.3, we have

$$\Psi(\omega_\Delta) = n(\omega_\Delta) = n_{AM\omega_\Delta A^{-1}} = n_{BAM\omega_\Delta A^{-1}}. \quad (7)$$

By Lemma 3.6 and Lemma 3.3, we have

$$\Psi(\xi) = n(\xi) = n_{BM\xi}. \quad (8)$$

Since  $(a, p_1) = 1$  and  $\chi_{-p_1, -p_2}^{(p_1 p_2 f^2)}([I(\xi)]^+) = \chi_{-p_1}(a)$ , by (7) and (8), it suffice to show that

$$\begin{aligned} & \frac{-r - rac}{q + rm + r\sigma_\Delta} - 12s(-rac, q + rm + r\sigma_\Delta) \\ \equiv & \chi_{-p_1}(a) \left( \frac{-ra - rc}{q + rm + r\sigma_\Delta} - 12s(-rc, q + rm + r\sigma_\Delta) \right) \pmod{8}. \end{aligned}$$

By Lemma 3.10 (i) we have

$$q + rm + r\sigma_\Delta \equiv 3 \pmod{4}. \quad (9)$$

By Lemma 3.10 (i) if  $f = 1$ , then  $8 \mid r$ , and if  $f = 2$ , then  $4 \mid r$ ,  $2 \mid b$  and at least one of  $\{a, c\}$  is odd; so, we have  $8 \mid r(a-1)(c-1)$  and  $8 \mid r(a+1)(c+1)$ . By Proposition 3.2 (v) it suffices to show that

$$\begin{aligned} & 6(q + rm + r\sigma_\Delta)s(-rac, q + rm + r\sigma_\Delta) \\ \equiv & \chi_{-p_1}(a)6(q + rm + r\sigma_\Delta)s(-rc, q + rm + r\sigma_\Delta) \pmod{4}. \end{aligned} \quad (10)$$

By Proposition 3.2 (vi) we have

$$\left( \frac{-rac}{q + rm + r\sigma_\Delta} \right) = (-1)^{\frac{1}{2} \left( \frac{q+rm+r\sigma_\Delta-1}{2} - 6(q+rm+r\sigma_\Delta)s(-rac, q+rm+r\sigma_\Delta) \right)} \quad (11)$$

and

$$\left( \frac{-rc}{q + rm + r\sigma_\Delta} \right) = (-1)^{\frac{1}{2} \left( \frac{q+rm+r\sigma_\Delta-1}{2} - 6(q+rm+r\sigma_\Delta)s(-rc, q+rm+r\sigma_\Delta) \right)}. \quad (12)$$

By dividing or multiplying (11) by (12), we have

$$\left( \frac{a}{q + rm + r\sigma_\Delta} \right) = (-1)^{\frac{1}{2}Z}, \quad (13)$$

where

$$\begin{aligned} Z = & \frac{1 - \chi_{-p_1}(a)}{2} (q + rm + r\sigma_\Delta - 1) - 6(q + rm + r\sigma_\Delta)s(-rac, q + rm + r\sigma_\Delta) \\ & + 6\chi_{-p_1}(a)(q + rm + r\sigma_\Delta)s(-rc, q + rm + r\sigma_\Delta). \end{aligned}$$

By Lemma 3.10 (ii) there exist integers  $T$  and  $U$  such that  $p_1\varepsilon_\Delta = (T + U\omega_\Delta)^2$ ; so, we have that  $p_1(q + r\sigma_\Delta/2) = (T + U\sigma_\Delta/2)^2 + U^2\Delta/4$  and  $p_1r/2 = (T + U\sigma_\Delta/2)U$ . Then we have

$$\begin{aligned} p_1(q + rm + r\sigma_\Delta) &= \left( T + \frac{U\sigma_\Delta}{2} \right)^2 + \frac{U^2\Delta}{4} + \left( T + \frac{U\sigma_\Delta}{2} \right) Ub \\ &= \left( T + \frac{U\sigma_\Delta}{2} + \frac{Ub}{2} \right)^2 - U^2ac \end{aligned} \quad (14)$$

and  $T + U(\sigma_\Delta + b)/2 \in \mathbb{Z}$ . Let  $e \in \mathbb{Z}$  and  $a_1 \in \mathbb{Z} \setminus 2\mathbb{Z}$  such that  $a = 2^e a_1$ . By (14) we have

$$\left(\frac{p_1(q + rm + r\sigma_\Delta)}{a_1}\right) = 1,$$

where  $\left(\frac{\cdot}{\cdot}\right)$  is the *Kronecker symbol* (we refer the relation between the Kronecker symbol and the Jacobi symbol to [6, Section 3.5]). We note that  $(q + rm + r\sigma_\Delta, a) = 1$  by  $\det M_\xi = 1$ . By the law of quadratic reciprocity and (9), we have

$$\begin{aligned} \left(\frac{a}{q + rm + r\sigma_\Delta}\right) &= (-1)^{\frac{a_1-1}{2} \frac{q+rm+r\sigma_\Delta-1}{2}} \left(\frac{q + rm + r\sigma_\Delta}{a}\right) \\ &= (-1)^{\frac{a_1-1}{2}} \left(\frac{p_1}{a_1}\right) \left(\frac{q + rm + r\sigma_\Delta}{2^e}\right). \end{aligned} \quad (15)$$

We claim that

$$\left(\frac{q + rm + r\sigma_\Delta}{2^e}\right) = \left(\frac{p_1}{2^e}\right). \quad (16)$$

If  $e$  is even, it is clear. If  $e \geq 3$ , by (14), then  $p_1(q + rm + r\sigma_\Delta) \equiv 1 \pmod{8}$ ; thus we have  $p_1 \equiv q + rm + r\sigma_\Delta \pmod{8}$ . Now we assume that  $e = 1$ . If  $f = 2$ , we have that  $p_1 p_2 = m^2 - ac$ ,  $m$  is odd and  $4 \mid ac$ , which is a contradiction with  $\gcd(a, b, c) = 1$ . Therefore, we have that  $f = 1$  and  $p_1 p_2 \equiv 1 \pmod{8}$ . We write  $\varepsilon_{p_1 p_2} = q + r\omega_{p_1 p_2} = x + y\sqrt{p_1 p_2}$ , i.e.,  $q = x - y$  and  $r = 2y$ . By Lemma 3.10 (i) we have that  $q \equiv 3 \pmod{8}$  and  $r \equiv 0 \pmod{8}$  if  $(p_1, p_2) \equiv (3, 3) \pmod{8}$ , and that  $q \equiv 7 \pmod{8}$  and  $r \equiv 0 \pmod{8}$  if  $(p_1, p_2) \equiv (7, 7) \pmod{8}$ ; so, we have  $p_1 \equiv q + rm + r \pmod{8}$ . Therefore, (16) holds. By (15) and (16), we have

$$\left(\frac{a}{q + rm + r\sigma_\Delta}\right) = \chi_{-p_1}(a). \quad (17)$$

By (9), (13) and (17), we get (10) as desired.  $\square$

**Lemma 4.4.** *Following the above notation, we further assume that  $p_1 p_2 \equiv 5 \pmod{8}$  and  $\varepsilon_{p_1 p_2} = \frac{t+u\sqrt{p_1 p_2}}{2}$  is the fundamental unit of  $\mathcal{O}_{p_1 p_2}$  such that  $t \equiv u \equiv 1 \pmod{2}$ . Then for any  $\eta \in \mathbb{X}_{p_1 p_2}$ , we have*

$$\chi_{-p_1, -p_2}^{(p_1 p_2)}([I(\eta)]^+) \Psi(\eta) \equiv \Psi(\omega_{p_1 p_2}) \pmod{8}.$$

*Proof.* We write  $\varepsilon_{p_1 p_2} = \frac{t+u\sqrt{p_1 p_2}}{2} = q + r\omega_{p_1 p_2}$ , where  $q, r$  are positive integers and  $r$  is odd. We note that  $t = 2q + r$  and  $u = r$ . By Lemma 4.1 there exists  $\xi = \frac{b+\sqrt{p_1 p_2}}{2a} \in \mathbb{X}_{p_1 p_2}^0$  such that  $\xi \sim \eta$ ,  $(a, p_1) = 1$  and  $a > 0$ . Let  $c = \frac{b^2 - p_1 p_2}{4a} \in \mathbb{Z}$ , and let  $m$  be an integer such that  $b = 2m + 1$ .

It follows that  $a$  and  $c$  are odd because  $p_1 p_2 \equiv 5 \pmod{8}$ . Let  $A := \begin{pmatrix} 1 & m \\ 0 & 1 \end{pmatrix}$ ,  $B := \begin{pmatrix} 3 & -1 \\ 1 & 0 \end{pmatrix}$

and  $C := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ . By (4) we have

$$BAM_{\omega_{p_1 p_2}} A^{-1} = \begin{pmatrix} 3(q + rm + r) - r & -3rac - (q - rm) \\ q + rm + r & -rac \end{pmatrix}$$

and

$$BCAM_{\omega_{p_1 p_2}} A^{-1} C^{-1} = \begin{pmatrix} 3(q - rm) + rac & 3r - (q + rm + r) \\ q - rm & r \end{pmatrix}.$$

By (5) we have

$$BM_\xi = \begin{pmatrix} 3(q+rm+r) - ra & -3rc - (q-rm) \\ q+rm+r & -rc \end{pmatrix}$$

and

$$BCM_\xi C^{-1} = \begin{pmatrix} 3(q-rm) + rc & 3ra - (q+rm+r) \\ q-rm & ra \end{pmatrix}.$$

Since  $\xi = \frac{b+\sqrt{p_1 p_2}}{2a} \in \mathbb{X}_{p_1 p_2}^0$ , we have  $-\sqrt{p_1 p_2} < 2m+1 < \sqrt{p_1 p_2}$ ; so, we have  $q+rm+r > \varepsilon'_{p_1 p_2} > 0$  and  $q-rm > \varepsilon'_{p_1 p_2} > 0$ . By Lemma 3.7, Lemma 3.4, Lemma 3.5 and Lemma 3.3, we have

$$\Psi(\omega_{p_1 p_2}) = n(\omega_{p_1 p_2}) = n_{BAM_{\omega_{p_1 p_2}} A^{-1}} = -n_{BCAM_{\omega_{p_1 p_2}} A^{-1} C^{-1}}. \quad (18)$$

By Lemma 3.6, Lemma 3.5 and Lemma 3.3, we have

$$\Psi(\xi) = n(\xi) = n_{BM_\xi} = -n_{BCM_\xi C^{-1}}. \quad (19)$$

We note that only one of  $\{q+rm+r, q-rm\}$  is odd because  $r$  is odd. We have that

$$(q+rm+r-1)(q-rm-1) = \left(\frac{t}{2} - 1\right)^2 - \left(\frac{ub}{2}\right)^2 = 2 - t - u^2 ac$$

since  $t^2 - p_1 p_2 u^2 = 4$  and  $p_1 p_2 = b^2 - 4ac$ . Therefore, by Lemma 3.11 we have that

$$\begin{aligned} (q+rm+r-1)(q-rm-1) \equiv 0 \pmod{4} &\iff ac \equiv 1 \pmod{4} \\ &\iff \chi_{-4}(a) = \chi_{-4}(c). \end{aligned} \quad (20)$$

**Case 1.** Suppose that  $q+rm+r$  is odd. Since  $(a, p_1) = 1$  and  $\chi_{-p_1, -p_2}^{(p_1 p_2)}([I(\xi)]^+) = \chi_{-p_1}(a)$ , by (18) and (19), it suffices to show that

$$\begin{aligned} &\frac{-r - rac}{q+rm+r} - 12s(-rac, q+rm+r) \\ &\equiv \chi_{-p_1}(a) \left( \frac{-ra - rc}{q+rm+r} - 12s(-rc, q+rm+r) \right) \pmod{8}. \end{aligned}$$

It suffices to show that

$$\begin{aligned} &\frac{r(\chi_{-4}(a) - \chi_{-p_1}(a))(\chi_{-4}(c) - \chi_{-p_1}(a))}{2} \\ &\equiv -6(q+rm+r)s(-rac, q+rm+r) \\ &\quad + \chi_{-p_1}(a)6(q+rm+r)s(-rc, q+rm+r) \pmod{4} \end{aligned} \quad (21)$$

by Proposition 3.2 (v),  $4 \mid (a - \chi_{-p_1}(a))(c - \chi_{-p_1}(a))$ , and  $\alpha \equiv \chi_{-4}(\alpha) \pmod{4}$  for any odd  $\alpha$ .

By Proposition 3.2 (vi) we have

$$\left( \frac{-rac}{q+rm+r} \right) = (-1)^{\frac{1}{2} \left( \frac{q+rm+r-1}{2} - 6(q+rm+r)s(-rac, q+rm+r) \right)} \quad (22)$$

and

$$\left( \frac{-rc}{q+rm+r} \right) = (-1)^{\frac{1}{2} \left( \frac{q+rm+r-1}{2} - 6(q+rm+r)s(-rc, q+rm+r) \right)}. \quad (23)$$

By dividing or multiplying (22) by (23), we have

$$\left( \frac{a}{q+rm+r} \right) = (-1)^{\frac{1}{2} Z}, \quad (24)$$

where

$$Z = \frac{1 - \chi_{-p_1}(a)}{2}(q + rm + r - 1) - 6(q + rm + r)s(-rac, q + rm + r) \\ + 6\chi_{-p_1}(a)(q + rm + r)s(-rc, q + rm + r).$$

By Lemma 3.9 (iii), Lemma 3.10 (ii) and  $t \equiv u \equiv 1 \pmod{2}$ , there exist integers  $X, Y, T$  and  $U$  such that  $p_1\varepsilon_{p_1p_2} = \left(\frac{(t-u\sqrt{p_1p_2})(X+Y\sqrt{p_1p_2})}{2}\right)^2 = (T + U\omega_{p_1p_2})^2$ ; so, we have that  $p_1(q + r/2) = (T + U/2)^2 + U^2p_1p_2/4$  and  $p_1r/2 = (T + U/2)U$ . Then we have

$$p_1(q + rm + r) = \left(T + \frac{U}{2}\right)^2 + \frac{U^2p_1p_2}{4} + \left(T + \frac{U}{2}\right)Ub = \left(T + \frac{U}{2} + \frac{Ub}{2}\right)^2 - U^2ac$$

and

$$\left(\frac{p_1(q + rm + r)}{a}\right) = 1.$$

We note that  $(q + rm + r, a) = 1$  by  $\det M_\xi = 1$ . By the law of quadratic reciprocity, we have

$$\begin{aligned} \left(\frac{a}{q + rm + r}\right) &= (-1)^{\frac{a-1}{2}\frac{q+rm+r-1}{2}} \left(\frac{q + rm + r}{a}\right) \\ &= \begin{cases} \left(\frac{p_1}{a}\right) & \text{if } q + rm + r - 1 \equiv 0 \pmod{4}, \\ (-1)^{\frac{a-1}{2}} \left(\frac{p_1}{a}\right) & \text{if } q + rm + r - 1 \equiv 2 \pmod{4}, \end{cases} \\ &= \begin{cases} \chi_{-4}(a)\chi_{-p_1}(a) & \text{if } q + rm + r - 1 \equiv 0 \pmod{4}, \\ \chi_{-p_1}(a) & \text{if } q + rm + r - 1 \equiv 2 \pmod{4}. \end{cases} \end{aligned} \quad (25)$$

By (20), (24) and (25), we get (21) as desired.

**Case 2.** Suppose that  $q - rm$  is odd. By (18) and (19), it suffices to show that

$$\frac{rac + r}{q - rm} - 12s(r, q - rm) \equiv \chi_{-p_1}(a) \left(\frac{ra + rc}{q - rm} - 12s(ra, q - rm)\right) \pmod{8}.$$

It suffices to show that

$$\begin{aligned} &\frac{r(\chi_{-4}(a) - \chi_{-p_1}(a))(\chi_{-4}(c) - \chi_{-p_1}(a))}{2} \\ &\equiv 6(q - rm)s(r, q - rm) - \chi_{-p_1}(a)6(q - rm)s(ra, q - rm) \pmod{4} \end{aligned} \quad (26)$$

by Proposition 3.2 (v),  $4 \mid (a - \chi_{-p_1}(a))(c - \chi_{-p_1}(a))$ , and  $\alpha \equiv \chi_{-4}(\alpha) \pmod{4}$  for any odd  $\alpha$ .

By Proposition 3.2 (vi) we have

$$\left(\frac{r}{q - rm}\right) = (-1)^{\frac{1}{2}\left(\frac{q-rm-1}{2} - 6(q-rm)s(r, q-rm)\right)} \quad (27)$$

and

$$\left(\frac{ra}{q - rm}\right) = (-1)^{\frac{1}{2}\left(\frac{q-rm-1}{2} - 6(q-rm)s(ra, q-rm)\right)}. \quad (28)$$

By dividing or multiplying (27) by (28), we have

$$\left(\frac{a}{q - rm}\right) = (-1)^{\frac{1}{2}\left(\frac{1-\chi_{-p_1}(a)}{2}(q-rm-1) - 6(q-rm)s(r, q-rm) + 6\chi_{-p_1}(a)(q-rm)s(ra, q-rm)\right)}. \quad (29)$$



By Lemma 3.9 (iii), Lemma 3.10 (ii) and  $t \equiv u \equiv 1 \pmod{2}$ , there exist integers  $X, Y, T$  and  $U$  such that  $p_1 \varepsilon_{p_1 p_2} = \left( \frac{(t-u\sqrt{p_1 p_2})(X+Y\sqrt{p_1 p_2})}{2} \right)^2 = (T + U\omega_{p_1 p_2})^2$ ; so, we have that  $p_1(q + r/2) = (T + U/2)^2 + U^2 p_1 p_2 / 4$  and  $p_1 r / 2 = (T + U/2)U$ . Then we have

$$p_1(q - rm) = \left(T + \frac{U}{2}\right)^2 + \frac{U^2 p_1 p_2}{4} - \left(T + \frac{U}{2}\right)Ub = \left(T + \frac{U}{2} - \frac{Ub}{2}\right)^2 - U^2 ac$$

and

$$\left(\frac{p_1(q - rm)}{a}\right) = 1.$$

We note that  $(q - rm, a) = 1$  by  $\det M_\xi = 1$ . By the law of quadratic reciprocity, we have

$$\begin{aligned} \left(\frac{a}{q - rm}\right) &= (-1)^{\frac{a-1}{2} \frac{q-rm-1}{2}} \left(\frac{q - rm}{a}\right) \\ &= \begin{cases} \left(\frac{p_1}{a}\right) & \text{if } q - rm - 1 \equiv 0 \pmod{4}, \\ (-1)^{\frac{a-1}{2}} \left(\frac{p_1}{a}\right) & \text{if } q - rm - 1 \equiv 2 \pmod{4}, \end{cases} \\ &= \begin{cases} \chi_{-4}(a) \chi_{-p_1}(a) & \text{if } q - rm - 1 \equiv 0 \pmod{4}, \\ \chi_{-p_1}(a) & \text{if } q - rm - 1 \equiv 2 \pmod{4}. \end{cases} \end{aligned} \quad (30)$$

By (20), (29) and (30), we get (26) as desired.  $\square$

*Proofs of Theorems 1.3 and 1.4.* Let  $f = 1$  or  $2$ , and let  $n = 6$  if  $\min\{p_1, p_2\} > 3$  and  $n = 2$  otherwise. By (6) we have

$$\begin{aligned} &h(-p_1)h(-p_2)\theta(-p_1, -p_2, f) - \frac{h(p_1 p_2 f^2)\Psi(\omega_{p_1 p_2 f^2})}{n} \\ &= \frac{1}{n} \sum_{[\eta] \sim \mathfrak{X}_{p_1 p_2 f^2}} \left( \chi_{-p_1, -p_2}^{(p_1 p_2 f^2)}([I(\eta)]^+) \Psi(\eta) - \Psi(\omega_{p_1 p_2 f^2}) \right). \end{aligned}$$

For  $\xi \in \mathbb{X}_{p_1 p_2 f^2}^0$ , let  $\xi^{\text{op}} := [\xi] - \xi'$ . In the proof of Lemma 4.2, we see that  $[I(\xi)] = [I(\xi^{\text{op}})]^{-1}$  in  $\mathcal{C}_{p_1 p_2 f^2}$ . By Lemma 3.9  $h(p_1 p_2 f^2)$  is odd; so, for any  $\xi \in \mathbb{X}_{p_1 p_2 f^2}^0$  such that  $\xi$  is not equivalent to  $\omega_{p_1 p_2 f^2}$ , we have  $[I(\xi)] \neq [I(\xi^{\text{op}})]$ . Therefore, there exist representatives  $\omega_{p_1 p_2 f^2}, \xi_{(1)}, \xi_{(2)}, \dots, \xi_{(m)}$  such that

$$\mathfrak{X}_{p_1 p_2 f^2} = \{[\omega_{p_1 p_2 f^2}] \sim, [\xi_{(1)}] \sim, [\xi_{(1)}^{\text{op}}] \sim, \dots, [\xi_{(m)}] \sim, [\xi_{(m)}^{\text{op}}] \sim\},$$

where  $m = \frac{h(p_1 p_2 f^2) - 1}{2}$ . By Lemma 4.2 we have

$$\begin{aligned} &h(-p_1)h(-p_2)\theta(p_1, p_2, f) - \frac{h(p_1 p_2 f^2)\Psi(\omega_{p_1 p_2 f^2})}{n} \\ &= \frac{2}{n} \sum_{i=1}^m \left( \chi_{-p_1, -p_2}^{(p_1 p_2 f^2)}([I(\xi_{(i)})]^+) \Psi(\xi_{(i)}) - \Psi(\omega_{p_1 p_2 f^2}) \right). \end{aligned}$$

Therefore, by Lemma 4.3 or Lemma 4.4 and  $2 \parallel n$ , we have

$$h(-p_1)h(-p_2)\theta(-p_1, -p_2, f) - h(p_1 p_2 f^2) \frac{\Psi(\omega_{p_1 p_2 f^2})}{n} \equiv 0 \pmod{8}$$

as desired.  $\square$

APPENDIX

We denote odd primes by  $p_1$  and  $p_2$ . For the small discriminant of  $\mathbb{Q}(\sqrt{p_1 p_2})$  having the class number 1, 3 or 5, we list the factorizations of

$$H(f) := H(-p_1, -p_2, f) := h(p_1 p_2 f^2) \frac{\Psi(\omega_{p_1 p_2 f^2})}{n} - h(-p_1)h(-p_2)\theta(-p_1, -p_2, f).$$

Here  $f$  is 1 or 2,  $h(\Delta)$  is the class number of the quadratic order with discriminant  $\Delta$ ,  $\Psi$  is the Hirzebruch sum,  $\omega_{p_1 p_2} = (1 + \sqrt{p_1 p_2})/2$ ,  $\omega_{4p_1 p_2} = \sqrt{p_1 p_2}$ ,  $n = 6$  (resp.,  $n = 2$ ) if  $\min\{p_1, p_2\} > 3$  (resp., if  $\min\{p_1, p_2\} = 3$ ),  $\theta(1) := \theta(-p_1, -p_2, 1) := 1$  and  $\theta(2) := \theta(-p_1, -p_2, 2) := (2 - (\frac{-p_1}{2}))(2 - (\frac{-p_2}{2})) - (1 - (\frac{-p_1}{2}))(1 - (\frac{-p_2}{2}))$  with the Kronecker symbol  $(-)$ . We denote the fundamental unit of  $\mathbb{Q}(\sqrt{p_1 p_2})$  by  $x_{p_1 p_2} + y_{p_1 p_2} \omega_{p_1 p_2}$ .

Table A1. Factorizations of  $H(-p_1, -p_2, f)$  for  $h(p_1 p_2) = 1$  and  $p_1 p_2 \leq 161$

$\Delta = p_1 p_2 f^2$	$\Psi(\omega_\Delta)$	$(x_{p_1 p_2}, y_{p_1 p_2})$	$h(\Delta), h(-p_1), h(-p_2)$	$\theta(f)$	$H(f)$
$21 = 3 \cdot 7$	2	(2,1)	1,1,1	1	0
$21 \cdot 2^2$	6		1,1,1	3	0
$33 = 3 \cdot 11$	2	(19,8)	1,1,1	1	0
$33 \cdot 2^2$	10		1,1,1	5	0
$57 = 3 \cdot 19$	2	(131,40)	1,1,1	1	0
$57 \cdot 2^2$	10		1,1,1	5	0
$69 = 3 \cdot 23$	6	(11,3)	1,1,3	1	0
$69 \cdot 2^2$	18		1,1,3	3	0
$77 = 7 \cdot 11$	6	(4,1)	1,1,1	1	0
$77 \cdot 2^2$	18		1,1,1	3	0
$93 = 3 \cdot 31$	6	(13,3)	1,1,3	1	0
$93 \cdot 2^2$	18		1,1,3	3	0
$129 = 3 \cdot 43$	2	(15371,2968)	1,1,1	1	0
$129 \cdot 2^2$	10		1,1,1	5	0
$133 = 7 \cdot 19$	6	(79,15)	1,1,1	1	0
$133 \cdot 2^2$	18		1,1,1	3	0
$141 = 3 \cdot 47$	10	(87,16)	1,1,5	1	0
$141 \cdot 2^2$	26		3,1,5	3	$2^3 \cdot 3$
$161 = 7 \cdot 23$	18	(10847,1856)	1,1,3	1	0
$161 \cdot 2^2$	18		1,1,3	1	0

Table A2. Factorizations of  $H(-p_1, -p_2, f)$  for  $h(p_1p_2) = 3$  and  $p_1p_2 \leq 1509$ 

$\Delta = p_1p_2f^2$	$\Psi(\omega_\Delta)$	$(x_{p_1p_2}, y_{p_1p_2})$	$h(\Delta), h(-p_1), h(-p_2)$	$\theta(f)$	$H(f)$
$321 = 3 \cdot 107$	18	(203,24)	3,1,3	1	$2^3 \cdot 3$
$321 \cdot 2^2$	42		3,1,3	5	$2^4 \cdot 3$
$469 = 7 \cdot 67$	18	(31,3)	3,1,1	1	$2^3$
$469 \cdot 2^2$	54		3,1,1	3	$2^3 \cdot 3$
$473 = 11 \cdot 43$	18	(83,8)	3,1,1	1	$2^3$
$473 \cdot 2^2$	42		3,1,1	5	$2^4$
$993 = 3 \cdot 331$	18	(2563,168)	3,1,3	1	$2^3 \cdot 3$
$993 \cdot 2^2$	42		3,1,3	5	$2^4 \cdot 3$
$1101 = 3 \cdot 367$	22	(177,11)	3,1,9	1	$2^3 \cdot 3$
$1101 \cdot 2^2$	66		3,1,9	3	$2^3 \cdot 3^2$
$1257 = 3 \cdot 419$	38	(98539,5720)	3,1,9	1	$2^4 \cdot 3$
$1257 \cdot 2^2$	94		3,1,9	5	$2^5 \cdot 3$
$1509 = 3 \cdot 503$	46	(246,13)	3,1,21	1	$2^4 \cdot 3$
$1509 \cdot 2^2$	138		3,1,21	3	$2^4 \cdot 3^2$

Table A3. Factorizations of  $H(-p_1, -p_2, f)$  for  $h(p_1p_2) = 5$  and  $p_1p_2 \leq 3997$ 

$\Delta = p_1p_2f^2$	$\Psi(\omega_\Delta)$	$(x_{p_1p_2}, y_{p_1p_2})$	$h(\Delta), h(-p_1), h(-p_2)$	$\theta(f)$	$H(f)$
$817 = 19 \cdot 43$	30	(331,24)	5,1,1	1	$2^3 \cdot 3$
$817 \cdot 2^2$	42		5,1,1	5	$2^4$
$1393 = 7 \cdot 199$	30	(3487,192)	5,1,9	1	$2^4$
$1393 \cdot 2^2$	78		5,1,9	1	$2^3 \cdot 7$
$1641 = 3 \cdot 547$	30	(4267,216)	5,1,3	1	$2^3 \cdot 3^2$
$1641 \cdot 2^2$	54		5,1,3	5	$2^3 \cdot 3 \cdot 5$
$1897 = 7 \cdot 271$	42	(42895,2016)	5,1,11	1	$2^3 \cdot 3$
$1897 \cdot 2^2$	90		5,1,11	1	$2^6$
$3997 = 7 \cdot 571$	54	(280,9)	5,1,5	1	$2^3 \cdot 5$
$3997 \cdot 2^2$	162		5,1,5	3	$2^3 \cdot 3 \cdot 5$

## REFERENCES

- [1] W. Bosma and P. Stevenhagen, *On the computation of quadratic 2-class groups*, J. Théor. Nr. Bordx. **8** (1996), 283–313. [3](#)
- [2] W. Cheng and X. Guo, *Some congruences connection quadratic class numbers with continued fractions*, Acta Arith. **191** (2019), 309–340. [2](#), [6](#)
- [3] L. Chua, B. Gunby, S. Park and A. Yuan, *Proof of a conjecture of Guy on class numbers*, Inter. J. Number Theory **11** (2015), 1345–1355. [2](#), [10](#)
- [4] P.G.L. Dirichlet, *Einige neue Sätze über unbestimmte Gleichungen*, Abh. Königlich Preussischen Akad. Wiss. (1834), 649–664; Gesammelte Werke, Chelsea Pub. Co., NY, 1969, pp. 219–236. [9](#)

- [5] F. Hirzebruch, *Hilbert modular surfaces*, L'Enseignement Math. **19** (1973), 183–281. [2](#)
- [6] F. Halter-Koch, *Quadratic irrationals: An introduction to classical number theory*, Pure and Applied Mathematics, CRC Press, Boca Raton, FL, 2013. [2](#), [4](#), [5](#), [8](#), [9](#), [13](#)
- [7] M. Kaneko and Y. Mizuno, *Genus character  $L$ -functions of quadratic orders and class numbers*, J. Lond. Math. Soc. **102** (2020), 69–98. [2](#), [3](#), [4](#), [5](#), [7](#), [8](#)
- [8] H. Lang, *Über einfache periodische Kettenbrüche und Vermutungen von P. Chowla und S. Chowla*, Acta Arith. **28** (1976), 419–428. [2](#), [3](#), [10](#)
- [9] H. Lu, *Hirzebruch sum and class number of the quadratic fields*, Chinese Sci. Bull. **36** (1991), 1145–1147. [2](#), [6](#)
- [10] Y. Mizuno, *Congruences relating class numbers of quadratic orders and Zagier's sums*, J. Number Theory **227** (2021), 352–370. [2](#), [10](#)
- [11] H. Rademacher and E. Grosswald, *Dedekind Sums*, Carus Mathematical Monographs, vol. 16, Mathematical Association of America, Washington, DC, 1972. [6](#)
- [12] K. S. Williams, *Historical remark on a theorem of Zhang and Yue*, J. Number Theory **147** (2015), 691–693. [9](#)
- [13] D. Zagier, *Nombres de classes et fractions continues*, Astérisque **24-25** (1975), 81–97. [2](#), [6](#)
- [14] Z. Zhang and Q. Yue, *Fundamental unit of real quadratic fields of odd class number*, J. Number Theory **137** (2014), 122–129. [9](#)

Institute of Mathematical Sciences,  
Ewha Womans University,  
Seoul, Korea  
E-mail: [jigu.kim.math@gmail.com](mailto:jigu.kim.math@gmail.com)

Graduate School of Technology Industrial and Social Sciences,  
Tokushima University,  
Tokushima, Japan  
E-mail: [mizuno.yoshinori@tokushima-u.ac.jp](mailto:mizuno.yoshinori@tokushima-u.ac.jp)