

FURTHER q -ANALOGUES OF THE (G.2) SUPERCONGRUENCE OF VAN HAMME

YUDONG LIU AND XIAOXIA WANG*

ABSTRACT. In 2015, Swisher generalized the (G.2) supercongruence of Van Hamme to the modulus p^4 . In this paper, we first propose two q -analogues of Swisher's supercongruence and then a new q -congruence with parameters is present. Furthermore, we prove a q -congruence modulo the fourth power of a cyclotomic polynomial, which was conjectured by the authors early.

1. INTRODUCTION

In 1997, Van Hamme [23] developed 13 mysterious p -adic analogues of Ramanujan-type π -formulas which were marked as (A.2)–(M.2), such as,

$$(G.2) \quad \sum_{k=0}^{(p-1)/4} (8k+1) \frac{(\frac{1}{4})_k^4}{k!^4} \equiv p \frac{\Gamma_p(\frac{1}{2})\Gamma_p(\frac{1}{4})}{\Gamma_p(\frac{3}{4})} \pmod{p^3}, \quad \text{if } p \equiv 1 \pmod{4}. \quad (1.1)$$

Here and throughout this paper, p is an odd prime and $\Gamma_p(x)$ is the p -adic Gamma function [17]. Then the research of the Ramanujan-type congruence and supercongruence has caught a lot of authors' attention. Later, Swisher [22, Theorem 1.2] proved a stronger version of (1.1):

$$\sum_{k=0}^{(p-1)/4} (8k+1) \frac{(\frac{1}{4})_k^4}{k!^4} \equiv -(-1)^{\frac{p-1}{4}} p \Gamma_p(\frac{1}{2})\Gamma_p(\frac{1}{4})^2 \pmod{p^4}, \quad \text{if } p \equiv 1 \pmod{4}. \quad (1.2)$$

Since $\Gamma_p(\frac{1}{4})\Gamma_p(\frac{3}{4}) = -(-1)^{\frac{p-1}{4}}$ for $p \equiv 1 \pmod{4}$, the right-hand side of (1.2) is the same as that of (1.1).

Recently, the authors [15] gave two q -analogues of (1.1) as follows: for positive integers $n \equiv 1 \pmod{4}$,

$$\sum_{k=0}^M [8k+1] \frac{(q; q^4)_k^4}{(q^4; q^4)_k^4} q^{2k} \equiv \frac{(q^2; q^4)_{(n-1)/4} [n] q^{(1-n)/4}}{(q^4; q^4)_{(n-1)/4}} \pmod{[n]\Phi_n(q^2)}; \quad (1.3)$$

2010 *Mathematics Subject Classification.* Primary 33D15; Secondary 11A07, 11B65.

Key words and phrases. basic hypergeometric series; supercongruences; q -congruences; q -analogue; cyclotomic polynomial.

This work is supported by National Natural Science Foundations of China (11661032).

* Corresponding author.

$$\sum_{k=0}^M [8k+1]_{q^2} [8k+1]^2 \frac{(q^2; q^8)_k^4}{(q^8; q^8)_k^4} q^{-4k} \equiv -\frac{2[n]_{q^2} (q^4; q^8)_{(n-1)/4}}{(1+q^2)(q^8; q^8)_{(n-1)/4}} q^{(3-n)/2} \pmod{[n]_{q^2} \Phi_n(q^2)^2}, \quad (1.4)$$

here and in what follows $M = (n-1)/4$ or $n-1$.

In fact, Guo and Schlosser [8, Theorem 2] have presented a q -supercongruence: for even $d \geq 4$ and integer $n \geq 1$ with $n \equiv -1 \pmod{d}$,

$$\sum_{k=0}^{n-1} [2dk+1] \frac{(q; q^d)_k^d}{(q^d; q^d)_k^d} q^{\frac{d(d-3)k}{2}} \equiv 0 \pmod{\Phi_n(q)^2}, \quad (1.5)$$

which is just q -analogue of a companion of Van Hamme's (G.2) for $p \equiv 3 \pmod{4}$ when $d = 4$. The authors [16] also proposed the following conjecture.

Conjecture 1. *Let $n \equiv 1 \pmod{4}$ be a positive integer. Then, modulo $[n] \Phi_n(q)^3$,*

$$\sum_{k=0}^{(n-1)/2} [6k+1] \frac{(q; q^4)_k (q; q^2)_k^3}{(q^2; q^2)_k (q^4; q^4)_k^3} q^{k^2+k} \equiv \sum_{k=0}^{(n-1)/4} [8k+1] \frac{(q; q^4)_k^4}{(q^4; q^4)_k^4} q^{2k}. \quad (1.6)$$

Here and in what follows, $(a; q)_n = (1-a)(1-aq) \cdots (1-aq^{n-1})$ is the q -shifted factorial and $(a_1, a_2, \dots, a_m; q)_n = (a_1; q)_n (a_2; q)_n \cdots (a_m; q)_n$ denotes the product of q -shifted factorials for simplicity. q -integer $[n]$ is defined as $[n] = [n]_q = (1-q^n)/(1-q)$ and $\Phi_n(q)$ is the n -th cyclotomic polynomial.

For some other recent progress on congruences and q -congruences, see [2–7, 9–12, 18, 19, 21, 24–28]. Especially, Guo and Zudilin [9] introduced the ‘creative microscoping’ method which is useful for proving q -congruences.

The first aim of this paper is to give the following two q -supercongruences modulo the fourth power of a cyclotomic polynomial which are the q -analogues of (1.2) and also the generalizations of the q -congruences (1.3) and (1.4).

Theorem 1. *Let $n \equiv 1 \pmod{4}$ be positive integer. Then, modulo $[n] \Phi_n(q)^3$, we have*

$$\sum_{k=0}^M [8k+1] \frac{(q; q^4)_k^4}{(q^4; q^4)_k^4} q^{2k} \equiv \frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} [n] q^{(1-n)/4} \{1 + [n]^2 B(n, q)\}; \quad (1.7)$$

$$\begin{aligned} & \sum_{k=0}^M [8k+1]_{q^2} [8k+1]^2 \frac{(q^2; q^8)_k^4}{(q^8; q^8)_k^4} q^{-4k} \\ & \equiv -\frac{2[n]_{q^2} (q^4; q^8)_{(n-1)/4}}{(1+q^2)(q^8; q^8)_{(n-1)/4}} q^{(5-n)/2} \{1 + [n]_{q^2}^2 B(n, q^2)\}, \end{aligned} \quad (1.8)$$

here and throughout this paper

$$B(n, q) = \frac{(n^2-1)(1-q)^2}{24} + \sum_{k=1}^{(n-1)/4} \frac{q^{4k-2}}{[4k-2]^2}.$$

The second purpose of this paper is to prove Conjecture 1 by comparing the following result with the q -supercongruence (1.7).

Theorem 2. *Let $n \equiv 1 \pmod{4}$ be a positive integer. Then, modulo $[n]\Phi_n(q)^3$,*

$$\sum_{k=0}^{(n-1)/2} [6k+1] \frac{(q; q^4)_k (q; q^2)_k^3}{(q^2; q^2)_k (q^4; q^4)_k^3} q^{k^2+k} \equiv \frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} [n] q^{(1-n)/4} \{1 + [n]^2 B(n, q)\}. \quad (1.9)$$

The rest of the paper is organized as follows. We shall prove Theorem 1 by establishing a parameter generalized q -congruence which based on Watson's ${}_8\phi_7$ transformation and the Chinese remainder theorem for coprime polynomials in the next section. In Section 3, we will give simple proof of Theorem 2. Finally, we will give another five q -analogues of (1.2) in Section 4.

2. PROOF OF THEOREM 1

The following Watson's ${}_8\phi_7$ transformation (cf. [1, Appendix (II.18)])

$$\begin{aligned} & {}_8\phi_7 \left[\begin{matrix} a, & qa^{\frac{1}{2}}, & -qa^{\frac{1}{2}}, & b, & c, & d, & e, & q^{-N} \\ & a^{\frac{1}{2}}, & -a^{\frac{1}{2}}, & aq/b, & aq/c, & aq/d, & aq/e, & aq^{N+1} \end{matrix} ; q, \frac{a^2 q^{N+2}}{bcde} \right] \\ &= \frac{(aq, aq/de; q)_N}{(aq/d, aq/e; q)_N} {}_4\phi_3 \left[\begin{matrix} aq/bc, & d, & e, & q^{-N} \\ & aq/b, & aq/c, & deq^{-N}/a \end{matrix} ; q, q \right] \end{aligned} \quad (2.1)$$

will play an important role in our proof. In fact, we can prove Theorem 1 by establishing the following parameter generalized q -congruence. Obviously, the special cases $c = d = q^{3/2}$ and $q \rightarrow q^2, c = d = q^{7/2}$ of the following q -congruence are just the two results in Theorem 1.

Theorem 3. *Let $n \equiv 1 \pmod{4}$ be positive integer. Then, modulo $[n]\Phi_n(q)^3$, we have*

$$\begin{aligned} & \sum_{k=0}^M [8k+1] \frac{(q; q^4)_k^4 (cq; q^4)_k (qd; q^4)_k}{(q^4; q^4)_k^4 (q^4/c; q^4)_k (q^4/d; q^4)_k} (cd)^{-k} q^{5k} \\ & \equiv \frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} [n] q^{(1-n)/4} \{1 + [n]^2 B(n, q)\} \\ & \quad \times \sum_{k=0}^M \frac{(q^3/cd; q^4)_k (q; q^4)_k^3}{(q^4; q^4)_k (q^4/c; q^4)_k (q^4/d; q^4)_k (q^2; q^4)_k} q^{4k}. \end{aligned} \quad (2.2)$$

In order to prove Theorem 3, we first recall the following two parametric q -congruences which the authors [14, Lemmas 1 and 2] have proved.

Lemma 1. *Let d, n be positive integers with $\gcd(d, n) = 1$. Let r be an integer and let a, b, c and e be indeterminates. Then, modulo $[n]$,*

$$\sum_{k=0}^{m_1} [2dk+r] \frac{(q^r, cq^r, eq^r, q^r/b, aq^r, q^r/a; q^d)_k}{(q^d, q^d/c, q^d/e, q^d b, q^d/a, aq^d; q^d)_k} \left(\frac{b}{ce}\right)^k q^{(2d-3r)k} \equiv 0; \quad (2.3)$$

$$\sum_{k=0}^{n-1} [2dk + r] \frac{(q^r, cq^r, eq^r, q^r/b, aq^r, q^r/a; q^d)_k}{(q^d, q^d/c, q^d/e, q^db, q^d/a, aq^d; q^d)_k} \left(\frac{b}{ce}\right)^k q^{(2d-3r)k} \equiv 0, \quad (2.4)$$

where $0 \leq m_1 \leq n-1$ and $dm_1 \equiv -r \pmod{n}$.

Lemma 2. *Let $n > 1$, $d \geq 2$, r be integers with $\gcd(r, d) = 1$ and $n \equiv r \pmod{d}$ such that $n + d - nd \leq r \leq n$. Then, modulo $\Phi_n(q)(1 - aq^n)(a - q^n)$,*

$$\begin{aligned} & \sum_{k=0}^{(n-r)/d} [2dk + r] \frac{(q^r, cq^r, eq^r, q^r/b, aq^r, q^r/a; q^d)_k}{(q^d, q^d/c, q^d/e, q^db, q^d/a, aq^d; q^d)_k} \left(\frac{b}{ce}\right)^k q^{(2d-3r)k} \\ & \equiv \frac{(q^{2r}/b; q^d)_{(n-r)/d}}{(bq^d; q^d)_{(n-r)/d}} [n] \left(\frac{b}{q^r}\right)^{(n-r)/d} \sum_{k=0}^M \frac{(q^{d-r}/ce, q^r/b, aq^r, q^r/a; q^d)_k}{(q^d, q^d/c, q^d/e, q^{2r}/b; q^d)_k} q^{dk}. \end{aligned} \quad (2.5)$$

We also need the following result.

Lemma 3. *Let $n \equiv 1 \pmod{4}$ be positive integer. Then, modulo $b - q^n$,*

$$\begin{aligned} & \sum_{k=0}^{(n-1)/4} [8k + 1] \frac{(q, q/b, qc, qd, aq, q/a; q^4)_k}{(q^4, q^4b, q^4/c, q^4/d, q^4/a, aq^4; q^4)_k} (cd)^{-k} q^{5k} b^k \\ & \equiv [n] \frac{(q, q^3; q^4)_{(n-1)/4}}{(aq^4, q^4/a; q^4)_{(n-1)/4}} \sum_{k=0}^{(n-1)/4} \frac{(q^3/cd, q/b, aq, q/a; q^4)_k}{(q^4, q^4/c, q^4/d, q^4, q^2/b; q^4)_k} q^{4k}. \end{aligned} \quad (2.6)$$

Proof. Letting $q \rightarrow q^4$, $a = q$, $b = qc$, $c = qd$, $d = q/a$, $e = aq$ and $N = (n-1)/4$ in Watson's formula (2.1), we get

$$\begin{aligned} & \sum_{k=0}^{(n-1)/4} [8k + 1] \frac{(q, q^{1-n}, qc, qd, aq, q/a; q^4)_k}{(q^4, q^{4-n}, q^4/c, q^4/d, q^4/a, aq^4; q^4)_k} (cd)^{-k} q^{(5+n)k} \\ & \equiv [n] \frac{(q, q^3; q^4)_{(n-1)/4}}{(aq^4, q^4/a; q^4)_{(n-1)/4}} \sum_{k=0}^{(n-1)/4} \frac{(q^3/cd, q^{1-n}, aq, q/a; q^4)_k}{(q^4, q^4/c, q^4/d, q^4, q^{2-n}; q^4)_k} q^{4k}, \end{aligned}$$

which is just the special case $b = q^n$ of (2.6). This establishes Lemma 3. \square

We now give a parametric version of Theorem 3.

Theorem 4. *Let $n \equiv 1 \pmod{4}$ be positive integer. Then, modulo $\Phi_n(q)^2(a - q^n)(1 - aq^n)$,*

$$\begin{aligned} & \sum_{k=0}^{(n-1)/4} [8k + 1] \frac{(q; q^4)_k^2 (cq, dq, qa, q/a; q^4)_k}{(q^4; q^4)_k^2 (q^4/c, q^4/d, q^4a, q^4/a; q^4)_k} (cd)^{-k} q^{5k} \\ & \equiv \left\{ q^{(1-n)/4} [n] \frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} + q^{(1-n)/4} [n] \frac{h_q(a)}{(1-a^2)(1-a^n)(q^2, q^4; q^4)_{(n-1)/4}} \right\} \end{aligned}$$

$$\times \sum_{k=0}^{(n-1)/4} \frac{(q, q^3/cd, aq, q/a; q^4)_k}{(q^2, q^4, q^4/c, q^4/d; q^4)_k} q^{4k}, \quad (2.7)$$

where

$$h_q(a) = (1-aq^n)(a-q^n) \left\{ (q^2; q^4)_{(n-1)/4}^2 (1-a^n) - n(aq^2, q^2/a; q^4)_{(n-1)/4} (1-a)a^{(n-1)/2} \right\}.$$

Proof. It is clear that $\Phi_n(q)$, $1-aq^n$, $a-q^n$ and $b-q^n$ are pairwise relatively prime polynomials and the following congruences can be easily verified

$$\frac{(b-q^n)(ab-1-a^2+aq^n)}{(a-b)(1-ab)} \equiv 1 \pmod{(1-aq^n)(a-q^n)}; \quad (2.8)$$

$$\frac{(1-aq^n)(a-q^n)}{(a-b)(1-ab)} \equiv 1 \pmod{(b-q^n)}. \quad (2.9)$$

Applying Lemma 2 for $d=4$, $r=1$, $e=d$ and Lemma 3 with the Chinese remainder theorem for coprime polynomials, we obtain, modulo $\Phi_n(q)(1-aq^n)(a-q^n)(b-q^n)$,

$$\begin{aligned} & \sum_{k=0}^{(n-1)/4} [8k+1] \frac{(q, cq, dq, qa, q/a, q/b; q^4)_k}{(q^4, q^4/c, q^4/d, q^4a, q^4/a, q^4b; q^4)_k} (cd)^{-k} q^{5k} b^k \\ & \equiv [n]A_q(a, b, n) \sum_{k=0}^{(n-1)/4} \frac{(q^3/cd, q/b, aq, q/a; q^4)_k}{(q^4, q^4/c, q^4/d, q^4, q^2/b; q^4)_k} q^{4k}, \end{aligned} \quad (2.10)$$

where

$$\begin{aligned} A_q(a, b, n) &= \frac{(b-q^n)(ab-1-a^2+aq^n)}{(a-b)(1-ab)} \frac{(b/q)^{(n-1)/4} (q^2/b; q^4)_{(n-1)/4}}{(bq^4; q^4)_{(n-1)/4}} \\ &+ \frac{(1-aq^n)(a-q^n)}{(a-b)(1-ab)} \frac{(q, q^3; q^4)_{(n-1)/4}}{(aq^4, q^4/a; q^4)_{(n-1)/4}}. \end{aligned} \quad (2.11)$$

It is easy to get

$$\begin{aligned} (q^4/a; q^4)_{(n-1)/4} &= (1-q^4/a)(1-q^8/a) \cdots (1-q^{n-1}/a) \\ &\equiv (1-q^{4-n}/a)(1-q^{8-n}/a) \cdots (1-q^{-1}/a) \\ &= (-1)^{(n-1)/4} (aq; q^4)_{(n-1)/4} \frac{q^{-(n-3)(n-1)/8}}{a^{(n-1)/4}} \pmod{\Phi_n(q)}. \end{aligned} \quad (2.12)$$

Similarly, we have

$$(aq^3; q^4)_{(n-1)/4} \equiv (-1)^{(n-1)/4} (q^2/a; q^4)_{(n-1)/4} q^{-(n-1)^2/8} a^{(n-1)/4} \pmod{\Phi_n(q)}. \quad (2.13)$$

Note that the following q -congruence which has been proposed by Guo [3, Equation (2.3)]

$$(aq; q)_{n-1} \equiv \sum_{k=0}^{n-1} a^k \pmod{\Phi_n(q)}. \quad (2.14)$$

Then $(aq^4, q^4/a; q^4)_{(n-1)/4}$ and $(q, q^3; q^4)_{(n-1)/4}$ can be transformed as

$$\begin{aligned} (aq^4, q^4/a; q^4)_{(n-1)/4} &\equiv (-1)^{(n-1)/4} \frac{q^{-(n-3)(n-1)/8}}{a^{(n-1)/4}} (aq, aq^4; q^4)_{(n-1)/4} \\ &= (-1)^{(n-1)/4} \frac{q^{-(n-3)(n-1)/8}}{a^{(n-1)/4}} \frac{(aq; q)_{n-1}}{(aq^2, aq^3; q^4)_{(n-1)/4}} \\ &\equiv q^{(1-n)/4} \frac{(1-a^n)}{(1-a)(aq^2, q^2/a; q^4)_{(n-1)/4} a^{(n-1)/2}} \pmod{\Phi_n(q)} \end{aligned} \quad (2.15)$$

and

$$(q, q^3; q^4)_{(n-1)/4} = \frac{(q; q)_{n-1}}{(q^2, q^4; q^4)_{(n-1)/4}} \equiv \frac{n}{(q^2, q^4; q^4)_{(n-1)/4}} \pmod{\Phi_n(q)}. \quad (2.16)$$

Combine the q -congruences (2.15) and (2.16), $A_q(a, b, n)$ reduces to the following result with $b = 1$,

$$A_q(a, 1, n) = q^{(1-n)/4} \frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} + q^{(1-n)/4} \frac{h_q(a)}{(1-a)^2(1-a^n)(q^2, q^4; q^4)_{(n-1)/4}}. \quad (2.17)$$

Thus we arrive at Theorem 4 instantly after taking $b = 1$ in both sides of (2.10). \square

Proof of Theorem 3. We first need to compute the 3rd derivative of $h_q(a)$ at $a = 1$ as

$$h_q'''(1) = (q^2; q^4)_{(n-1)/4}^2 (1-q^n)^2 \left\{ -\frac{n(n^2-1)}{4} - 6n \sum_{k=1}^{(n-1)/4} \frac{q^{4k-2}}{(1-q^{4k-2})^2} \right\}. \quad (2.18)$$

According to the L'Hospital rule, we have

$$\begin{aligned} \lim_{a \rightarrow 1} \frac{h_q(a)}{(1-a^2)(1-a^n)(q^2, q^4; q^4)_{(n-1)/4}} &= \frac{h_q'''(1)}{-6n(q^2, q^4; q^4)_{(n-1)/4}} \\ &= \frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} [n]^2 B(n, q). \end{aligned} \quad (2.19)$$

Owing to the fact that the denominators of both sides of (2.7) as $a \rightarrow 1$ are prime to $\Phi_n(q)^2$. Combine with the result (2.19), the q -congruence (2.2) for $M = (n-1)/4$ modulo $\Phi_n(q)^4$ is thus a direct conclusion from Theorem 4 by letting $a \rightarrow 1$. On the other hand, since $(q; q^4)_k \equiv 0 \pmod{\Phi_n(q)^4}$ for k in the range $(n-1)/4 \leq k \leq n-1$, Theorem 3 is also true for $M = n-1$ modulo $\Phi_n(q)^4$. At the same time, from Lemma 1 for $d = 4$, $r = a = b = 1$ and $e = d$, the q -congruence (2.2) is also valid modulo $[n]$. This completes the proof Theorem 3. \square

3. PROOF OF THEOREM 2

In order to prove Theorem 2, we first need to establish the following q -congruence, which is the case of [9, Theorem 4.5] with $b \rightarrow q/b$ and $r = 1$.

Lemma 4. *Let $n \equiv 1 \pmod{4}$ be positive integer. Then, modulo $[n](a - q^n)(1 - aq^n)$,*

$$\sum_{k=0}^{(n-1)/2} [6k+1] \frac{(aq, q/a, q; q^2)_k (q/b; q^4)_k q^{k^2+k} b^k}{(aq^4, q^4/a, q^4; q^4)_k (q^2b; q^2)_k} \equiv \frac{(q^2/b; q^4)_{(n-1)/4} [n] \left(\frac{b}{q}\right)^{(n-1)/4}}{(bq^4; q^4)_{(n-1)/4}}. \quad (3.1)$$

We also find a q -congruence for the left-hand side of (3.1) modulo $b - q^n$.

Lemma 5. *Let $n \equiv 1 \pmod{4}$ be positive integer. Then, modulo $b - q^n$,*

$$\sum_{k=0}^{(n-1)/2} [6k+1] \frac{(aq, q/a, q; q^2)_k (q/b; q^4)_k q^{k^2+k} b^k}{(aq^4, q^4/a, q^4; q^4)_k (q^2b; q^2)_k} \equiv [n] \frac{(q, q^3; q^4)_{(n-1)/4}}{(aq^4, q^4/a; q^4)_{(n-1)/4}}. \quad (3.2)$$

Sketch of proof. Using the following summation formula of Rahman [20, eq. (4.6)]

$$\sum_{k=0}^{\infty} \frac{(1 - aq^{3k}) (a, d, q/d; q)_k (b; q^2)_k a^k q^{\binom{k+1}{2}}}{(1-a)(q^2, aq^2/d, adq; q^2)_k (aq/b; q)_k b^k} = \frac{(aq, aq^2, adq/b, aq^2/bd; q^2)_{\infty}}{(aq/b, aq^2/b, aq^2/d, adq; q^2)_{\infty}} \quad (3.3)$$

with $q \rightarrow q^2$, $a = q$, $d = q/a$ and $b = q^{1-n}$, we just get the q -congruence (3.2) with $b = q^n$. \square

Proof of Theorem 2. Combining the above two lemmas and the q -congruences (2.8), (2.9) with the Chinese remainder theorem for coprime polynomials, we are led to the following result: modulo $[n](1 - aq^n)(a - q^n)(b - q^n)$,

$$\sum_{k=0}^{(n-1)/2} [6k+1] \frac{(aq, q/a; q^2)_k (q; q^2)_k (q/b; q^4)_k q^{k^2+k} b^k}{(aq^4, q^4/a; q^4)_k (q^4; q^4)_k (q^2b; q^2)_k} \equiv [n] A_q(a, b, n). \quad (3.4)$$

In the same manner as the proof of Theorem 3, taking $b \rightarrow 1$ and $a \rightarrow 1$ in sequence, we immediately arrive at Theorem 2 from (3.4). \square

4. THE OTHER q -ANALOGUES OF SWISHER'S SUPERCONGRUENCE (1.2)

From Theorem 3, we can get another five different q -analogues of Swisher's supercongruence (1.2). Fixing $c = d = -1$; $c = -1, d \rightarrow \infty$; $c = -1, d = 0$; $c \rightarrow \infty, d \rightarrow \infty$ and $c \rightarrow 0, d \rightarrow 0$ respectively in Theorem 3, we obtain the following results.

Corollary 5. *Let $n \equiv 1 \pmod{4}$ be positive integer. Then, modulo $[n]\Phi_n(q)^3$,*

$$\begin{aligned} & \sum_{k=0}^M [8k+1] \frac{(q; q^4)_k^2 (q^2; q^8)_k^2}{(q^4; q^4)_k^2 (q^8; q^8)_k^2} q^{5k} \\ & \equiv \frac{(q^2; q^4)_{(n-1)/4} [n] q^{(1-n)/4} \{1 + [n]^2 B(n, q)\}}{(q^4; q^4)_{(n-1)/4}} \sum_{k=0}^M \frac{(q^3; q^4)_k (q; q^4)_k^3}{(q^4; q^4)_k (-q^4; q^4)_k^2 (q^2; q^4)_k} q^{4k}; \end{aligned} \quad (4.1)$$

$$\sum_{k=0}^M [8k+1] \frac{(q; q^4)_k^3 (q^2; q^8)_k}{(q^4; q^4)_k^3 (q^8; q^8)_k} q^{2k^2+4k}$$

$$\equiv \frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} [n] q^{(1-n)/4} \{1 + [n]^2 B(n, q)\} \sum_{k=0}^M \frac{(q; q^4)_k^3}{(q^4; q^4)_k (-q^4; q^4)_k (q^2; q^4)_k} q^{4k}; \quad (4.2)$$

$$\sum_{k=0}^M [8k+1] \frac{(q; q^4)_k^3 (q^2; q^8)_k}{(q^4; q^4)_k^3 (q^8; q^8)_k} q^{3k-2k^2}$$

$$\equiv \frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} [n] q^{(1-n)/4} \{1 + [n]^2 B(n, q)\} \sum_{k=0}^M \frac{(q; q^4)_k^3}{(q^4; q^4)_k (-q^4; q^4)_k (q^2; q^4)_k} (-q)^{3k}; \quad (4.3)$$

$$\sum_{k=0}^M [8k+1] \frac{(q; q^4)_k^4}{(q^4; q^4)_k^4} q^{4k^2+3k}$$

$$\equiv \frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} [n] q^{(1-n)/4} \{1 + [n]^2 B(n, q)\} \sum_{k=0}^M \frac{(q; q^4)_k^3}{(q^4; q^4)_k (q^2; q^4)_k} q^{4k}; \quad (4.4)$$

$$\sum_{k=0}^M [8k+1] \frac{(q; q^4)_k^4}{(q^4; q^4)_k^4} q^{k-4k^2}$$

$$\equiv \frac{(q^2; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} [n] q^{(1-n)/4} \{1 + [n]^2 B(n, q)\} \sum_{k=0}^M \frac{(q; q^4)_k^3}{(q^4; q^4)_k (q^2; q^4)_k} (-q)^{2k-2k^2}. \quad (4.5)$$

Taking $q \rightarrow 1$ and $n = p \equiv 1 \pmod{4}$ be a prime in (4.1)–(4.5) and the two q -supercongruences (1.7)–(1.8) in Theorem 1, and applying the known congruence by Sun [21, p. 7]:

$$H_{(p-1)/2}^{(2)} \equiv 0 \pmod{p} \quad \text{for } p > 3,$$

we are led to the following supercongruence

$$\sum_{k=0}^m (8k+1) \frac{(\frac{1}{4})_k^4}{k!^4} \equiv \frac{(\frac{1}{2})_{(p-1)/4}}{(1)_{(p-1)/4}} p \left\{ 1 - \frac{p^2}{16} H_{(p-1)/4}^{(2)} \right\} \pmod{p^4}, \quad (4.6)$$

where $m = (p-1)/4$ or $p-1$ and the harmonic numbers of 2-order is defined as

$$H_n^{(2)} = \sum_{k=1}^n \frac{1}{k^2} \quad \text{with } n \in \mathbb{N}.$$

Recall that for prime $p \geq 5$ and rational number $a \in \mathbb{Z}_p$, the p -adic Gamma function Γ_p has the following basic properties [13],

$$\Gamma_p(1) = -1, \quad \Gamma_p\left(\frac{1}{2}\right)^2 = (-1)^{\frac{p+1}{2}} \quad (a)_n = (-1)^n \frac{\Gamma_p(a+n)}{\Gamma_p(a)}, \quad G_1(a) = G_1(1-a),$$

$$\Gamma_p(a+bp) \equiv \Gamma_p(a) \left(1 + G_1(a)bp + \frac{G_2(a)b^2p^2}{2} \right) \pmod{p^3},$$

$$\Gamma_p(x)\Gamma_p(1-x) = (-1)^{a_0(x)},$$

where $G_k(a) := \Gamma_p^{(k)}(a)/\Gamma_p(a)$, $\Gamma_p^{(k)}(a)$ is k -th derivative of $\Gamma_p(a)$ and $a_0(x) \in \{1, 2, \dots, p\}$ satisfies $x - a_0(x) \equiv 0 \pmod{p}$. We have the following result

$$\begin{aligned} \frac{\left(\frac{1}{2}\right)_{(p-1)/4}}{(1)_{(p-1)/4}} &= \frac{\Gamma_p((1+p)/4)\Gamma_p(1)}{\Gamma_p(1/2)\Gamma_p((3+p)/4)} \\ &= -(-1)^{(p-1)/4}\Gamma_p(1/2)\Gamma_p((1+p)/4)\Gamma_p((1-p)/4) \\ &\equiv -(-1)^{(p-1)/4}\Gamma_p(1/2) \left\{ \Gamma_p(1/4) + \Gamma'_p(1/4)\frac{p}{4} + \Gamma''_p(1/4)\frac{p^2}{2 \times 4^2} \right\} \\ &\times \left\{ \Gamma_p(1/4) - \Gamma'_p(1/4)\frac{p}{4} + \Gamma''_p(1/4)\frac{p^2}{2 \times 4^2} \right\} \pmod{p^3}. \end{aligned} \quad (4.7)$$

Recently, Wang and Pan [24, Page 6] have proved that

$$H_{(p-1)/4}^{(2)} \equiv \frac{\Gamma''_p(1/4)}{\Gamma_p(1/4)} - \left\{ \frac{\Gamma'_p(1/4)}{\Gamma_p(1/4)} \right\}^2 \pmod{p}. \quad (4.8)$$

Thus, the combination of (4.7) and (4.8) can bring out

$$p \frac{\left(\frac{1}{2}\right)_{(p-1)/4}}{(1)_{(p-1)/4}} \left\{ 1 - \frac{p^2}{16} H_{(p-1)/4}^{(2)} \right\} \equiv -(-1)^{(p-1)/4} p \Gamma_p(1/2) \Gamma_p(1/4)^2 \pmod{p^4}.$$

This shows that our q -congruences (1.7), (1.8) and (4.1)–(4.5) are indeed q -analogues of Swisher’s supercongruence (1.2).

REFERENCES

- [1] G. Gasper and M. Rahman, *Basic hypergeometric series*, second edition, Encyclopedia of Mathematics and Its Applications **96**, Cambridge University Press, Cambridge, 2004.
- [2] V.J.W. Guo, Proof of some q -supercongruences modulo the fourth power of a cyclotomic polynomial, *Results Math.* **75** (2020), Art. 77.
- [3] V.J.W. Guo, q -Supercongruences modulo the fourth power of a cyclotomic polynomial via creative microscoping, *Adv. Appl. Math.* **120** (2020), Art. 102078.
- [4] V.J.W. Guo, Proof of a generalization of the (C.2) supercongruence of Van Hamme, *Rev. R. Acad. Cienc. Exactas Fís. Nat., Ser. A Mat.* **115** (2021), Art. 45.
- [5] V.J.W. Guo, Some variations of a ‘divergent’ Ramanujan-type q -supercongruence, *J. Difference Equ. Appl.* in press; <https://doi.org/10.1080/10236198.2021.1900140>.
- [6] V.J.W. Guo and M.J. Schlosser, Proof of a basic hypergeometric supercongruence modulo the fifth power of a cyclotomic polynomial, *J. Difference Equ. Appl.* **25** (2019), 921–929.
- [7] V.J.W. Guo and M.J. Schlosser, Some new q -congruences for truncated basic hypergeometric series: even power, *Results Math.* **75** (2020), Art. 1.
- [8] V.J.W. Guo and M.J. Schlosser, A new family of q -supercongruences modulo the fourth power of a cyclotomic polynomial, *Results Math.* **75** (2020), Art. 155.
- [9] V.J.W. Guo and W. Zudilin, A q -microscope for supercongruences, *Adv. Math.* **346** (2019), 329–358.
- [10] V.J.W. Guo and W. Zudilin, Dwork-type supercongruences through a creative q -microscope, *J. Combin. Theory, Ser. A* **178** (2021), Art. 105362.
- [11] J.-C. Liu, On a congruence involving q -Catalan numbers, *C. R. Math. Acad. Sci. Paris* **358** (2020), 211–215.

- [12] J.-C. Liu and F. Petrov, Congruences on sums of q -binomial coefficients, *Adv. Appl. Math.* **116** (2020), Art. 102003.
- [13] L. Long and R. Ramakrishna, Some supercongruences occurring in truncated hypergeometric series, *Adv. Math.* **290** (2016), 773–808.
- [14] Y. Liu and X. Wang, q -Analogues of two Ramanujan-type supercongruences, *J. Math. Anal. Appl.* **502** (1) (2021), Art. 125238.
- [15] Y. Liu and X. Wang, q -Analogues of the (G.2) supercongruence of Van Hamme, *Rock. Mt. J. Math.* to appear.
- [16] Y. Liu and X. Wang, Some q -supercongruences from Rahman’s summation formula, preprint, 2021, arXiv: 2103.06416.
- [17] Y. Morita, A p -adic supercongruence of the Γ function, *J. Fac. Sci. Univ. Tokyo* **22** (1975), 255–266.
- [18] H.-X. Ni and H. Pan, On a conjectured q -congruence of Guo and Zeng, *Int. J. Number Theory* **14** (6) (2018), 1699–1707.
- [19] H.-X. Ni and H. Pan, Divisibility of some binomial sums, *Acta Arith.* **194** (2020), 367–381.
- [20] M. Rahman, Some quadratic and cubic summation formulas for basic hypergeometric series, *Canad. J. Math.* **45** (2) (1993), 394–411.
- [21] Z.-W. Sun, A new series for π^3 and related congruences, *Internat. J. Math.* **26** (8) (2019), Art. 1550055.
- [22] H. Swisher, On the supercongruence conjectures of Van Hamme, *Res. Math. Sci.* **2** (2015), Art. 18.
- [23] L. Van Hamme, Some conjectures concerning partial sums of generalized hypergeometric series, in: *p -Adic Functional Analysis* (Nijmegen, 1996), Lecture Notes in Pure and Appl. Math. **192**, Dekker, New York (1997), 223–236.
- [24] C. Wang and H. Pan, Supercongruences concerning truncated hypergeometric series, preprint, 2018, arXiv: 1806.02735.
- [25] X. Wang and M. Yue, Some q -supercongruences from Watson’s ${}_8\phi_7$ transformation formula, *Results Math.* **75** (2020), Art. 71.
- [26] X. Wang and M. Yue, A q -analogue of a Dwork-type supercongruence, *Bull. Aust. Math. Soc.* **103** (2) (2021), 303–310.
- [27] X. Wang and M. Yu, Some new q -congruences on double sums, *Rev. R. Acad. Cienc. Exactas Fís. Nat., Ser. A Mat.* **115**(2021), Art. 9.
- [28] W. Zudilin, Congruences for q -binomial coefficients, *Ann. Combin.* **23** (2019), 1123–1135.

DEPARTMENT OF MATHEMATICS, SHANGHAI UNIVERSITY, SHANGHAI 200444, P. R. CHINA
Email address: lydshdx@163.com (Y. Liu), xiaoxiawang@shu.edu.cn (X. Wang)