

HOMOTOPY RIBBON DISCS WITH A FIXED GROUP

ANTHONY CONWAY

ABSTRACT. In the topological category, the classification of homotopy ribbon discs is known when the fundamental group G of the complement is \mathbb{Z} and the Baumslag-Solitar group $BS(1, 2)$. We prove classification results under the assumption that the group G is good, geometrically 2-dimensional and satisfies the Farrell-Jones conjecture. As an application, for any knot $J \subset S^3$ whose knot group $G(J)$ is good, we classify the homotopy ribbon discs for $J\# - J$ whose complement has group $G(J)$. A similar application is obtained for $BS(m, n)$ when $|m - n| = 1$.

1. INTRODUCTION

Given a knot $K \subset S^3$, we consider the problem of classifying locally flat discs $D \subset D^4$ with boundary K , up to topological ambient isotopy rel. boundary. Naturally, K need not bound such a disc (i.e. K need not be *slice*) but if it does, then it is conjectured that it necessarily bounds one for which the inclusion induced map $\pi_1(S^3 \setminus K) \rightarrow \pi_1(D^4 \setminus D)$ is surjective; such discs are called *homotopy ribbon*. For this reason, and for technical purposes, we restrict our attention to homotopy ribbon discs with boundary K . Additionally, observe that if D_1 and D_2 are two ambiently isotopic slice discs with boundary K , then their groups must be isomorphic: $\pi_1(D^4 \setminus D_1) \cong \pi_1(D^4 \setminus D_2)$. Summarising, the goal of this article is to study the following question.

Question 1.1. *Given a knot $K \subset S^3$ and a ribbon group G , can one describe the set of homotopy ribbon discs for K with group G , considered up to topological ambient isotopy rel. boundary?*

Here, a group is called *ribbon* if it arises as $\pi_1(D^4 \setminus D)$ for some (smoothly embedded) ribbon disc $D \subset D^4$ ¹. We work with ribbon groups instead of fundamental groups of locally flat disc exteriors for convenience: the former admit an algebraic characterisation [FT05, Theorem 2.1], while no such description appears to be known for the latter [FT05, Question 1.7]. Examples of ribbon groups include $G = \mathbb{Z}$ and the Baumslag-Solitar group $G = BS(1, 2)$ and in those cases, Question 1.1 has been fully resolved [FT05, CP21]. The answers, which will be partially recalled in Remark 1.11 below, both rely on Freedman's 5-dimensional s-cobordism theorem [Fre82] and therefore make use of the fact that \mathbb{Z} and $BS(1, 2)$ are *good* groups. We refer to [BKK⁺21, Definition 12.12] for the precise definition of a good group and to [BKK⁺21, Chapter 19] for a survey, but note that the class of good groups contains all groups of subexponential growth as well as all elementary amenable groups (e.g. solvable groups). At the time of writing, it is unknown whether all groups are good: this is equivalent to the question of whether the free group F_2 is good [BKK⁺21, Proposition 19.7].

Remark 1.2. The only elementary amenable ribbon groups are \mathbb{Z} and $BS(1, 2)$, as can be seen by combining [Hil02, Corollary 2.6.1] with the fact that ribbon groups have deficiency one and abelianise to \mathbb{Z} . As a consequence, if the class of good ribbon groups were eventually shown to coincide with the class of elementary amenable ribbon groups, then the current article would contain no new result. Despite this shortcoming, we hope that the approach taken here will be of interest given the recent surge of activity around the topic of 2-discs in the 4-ball, both in the smooth and topological category [JZ20, CP21, Hay20, SS21, HKS⁺21, HS21, Hay21, DMS22].

1.1. Existence. From now on, a *G-ribbon disc* refers to a homotopy ribbon disc $D \subset D^4$ with $\pi_1(D^4 \setminus D) \cong G$. We recall and motivate a sufficient condition for the existence of a *G-ribbon disc* with boundary K , which is due to Friedl and Teichner [FT05, Theorem 1.9]. First,

¹ $D \subset D^4$ is *ribbon* if the restriction of the radial function $D^4 \rightarrow \mathbb{R}$ to D is Morse and admits no local maxima.

if K bounds a locally flat disc $D \subset D^4$, then $\partial N_D = M_K$, where $N_D := D^4 \setminus \nu D$ is the exterior of D and M_K denotes the 3-manifold obtained by 0-framed surgery on K . Next, if $D \subset D^4$ is a G -ribbon disc for a knot K , then there is an epimorphism $\pi_1(M_K) \twoheadrightarrow \pi_1(N_D) \cong G$ and (N_D, M_K) satisfies Poincaré duality or, using surgery theory jargon, is a (4-dimensional) *Poincaré pair*. If, additionally, the disc exterior $N_D = D^4 \setminus \nu D$ is aspherical, then we have a homotopy equivalence $N_D \simeq K(G, 1)$ and we deduce that $(K(G, 1), M_K)$ is a Poincaré pair.

Remark 1.3. It is expected that ribbon disc exteriors are aspherical [Gor81, Conjecture 6.5] (see also [How85]). As noted in [FT05, Section 2] this would imply the *ribbon group conjecture*: ribbon groups are geometrically 2-dimensional². Here recall that a group G is called *geometrically 2-dimensional* if $K(G, 1)$ is (homotopy equivalent to) a 2-complex. Both statements are in fact particular cases of the *Whitehead conjecture* which states that every connected subcomplex of a 2-dimensional aspherical CW complex is itself aspherical [Whi41]; see [Ros07] for a nice overview. Howie proved that locally indicable ribbon groups are geometrically 2-dimensional [How82, Theorem 5.2]. On the other hand, to the best of our knowledge, the Whitehead conjecture is not known to imply that exteriors of homotopy ribbon discs are aspherical; see also Remark 1.11 below.

We argued that if D is a G -ribbon disc with aspherical exterior and boundary a knot K , then $\pi_1(M_K) \twoheadrightarrow \pi_1(N_D) \cong G$ is an epimorphism and $(K(G, 1), M_K)$ is a Poincaré pair. On the other hand, if we start with an epimorphism $\pi_1(M_K) \twoheadrightarrow G$ onto a group G , then there is an embedding $\varphi: M_K \hookrightarrow K(G, 1) = BG$ that induces the given surjection on fundamental groups and, if G is geometrically 2-dimensional, then [FT05, Lemma 3.2] shows that $(K(G, 1), M_K)$ is a Poincaré pair if and only if the induced map

$$(FT) \quad \varphi^*: H^i(BG; \mathbb{Z}[G]) \rightarrow H^i(M_K; \mathbb{Z}[G]_\varphi) \text{ is an isomorphism for } i = 1, 2.$$

Under an additional condition on the group G , Friedl and Teichner prove that this leads to a sufficient condition for K to bound a G -ribbon disc [FT05, Theorem 1.9 and Lemma 3.2].

Theorem 1.4 (Friedl-Teichner). *Let $K \subset S^3$ be a knot and let G be a good geometrically 2-dimensional ribbon group such that $\tilde{L}_4^h(\mathbb{Z}[G]) = 0$. If $\varphi: \pi_1(M_K) \twoheadrightarrow G$ is an epimorphism that satisfies (FT), then there exists a G -ribbon disc $D \subset D^4$ with aspherical exterior and boundary K such that the composition $\pi_1(M_K) \twoheadrightarrow \pi_1(N_D) \cong G$ agrees with φ .*

Remark 1.5. We make a couple of remarks on this theorem.

- Friedl and Teichner actually prove a stronger result. Instead of asking for G to be geometrically 2-dimensional, they merely demand that $H_3(G) = 0$ and $H^i(G; \mathbb{Z}[G]) = 0$ for $i > 2$ and instead of assuming that G is ribbon, they only require that G be finitely presented and satisfy $H_1(G) = \mathbb{Z}$ and $H_2(G) = 0$. Finally, they do not require G to be good, only that the surgery sequence (with h -decorations) be exact for all 4-dimensional Poincaré pairs (X, M) with $\pi_1(X) = G$.
- The fact that the disc exterior is aspherical is implicit in [FT05, proof of Theorem 1.9]: their surgery theoretic argument yields a disc D whose exterior $N_D = D^4 \setminus \nu D$ is homotopy equivalent to $K(G, 1)$, which is aspherical.
- The groups \mathbb{Z} and $BS(1, 2)$ satisfy all the assumptions of Theorem 1.4. Additionally, for those groups, condition (FT) simplifies considerably. Indeed if G is poly-(torsion-free abelian) (or PTFA for short), then (FT) reduces to

$$(Ext) \quad \text{Ext}_{\mathbb{Z}[G]}^1(H_1(M_K; \mathbb{Z}[G]_\varphi), \mathbb{Z}[G]) = 0$$

and for $G = \mathbb{Z}$ it reduces further to the condition $\Delta_K = 1$; all of this is explained in [FT05, Sections 1 and 4 and Lemma 3.3].

1.2. Uniqueness and classification. We now return to the set $\mathcal{D}_G(K)$ of topological ambient isotopy classes of G -ribbon discs with boundary K . In fact, we will mostly be concerned with the subset $\mathcal{D}_G^a(K) \subset \mathcal{D}_G(K)$ of discs with aspherical exteriors. To that effect, inspired by [HKT09, Definition 1.2], we describe some assumptions on the group G that we will require.

²Friedl and Teichner refer to geometrically 2-dimensional groups as *aspherical* groups.

Definition 1.6. A group G satisfies properties *W-AA* if

- (W) the Whitehead group $\text{Wh}(G)$ vanishes;
- (A4) the assembly map $A_4: H_4(BG; \mathbf{L}_\bullet) \rightarrow L_4(\mathbb{Z}[G])$ is an isomorphism;³
- (A5) the assembly map $A_5: H_5(BG; \mathbf{L}_\bullet) \rightarrow L_5(\mathbb{Z}[G])$ is surjective.

We will mostly use these conditions as a blackbox, but note that thanks to extensive work on the Farrell-Jones conjecture (see [Lue21] for a survey) they should not be thought of as insurmountable restrictions. We discuss all of this in more detail in Remark 1.11 below and refer to [Ran92, CW21, Lue20, Lue21] for background on assembly maps in L -theory. Returning to our aim of describing $\mathcal{D}_G(K)$, we consider the set

$$(\text{Epi}) \quad \text{Epi}^{FT}(\pi_1(M_K), G) := \{\varphi: \pi_1(M_K) \rightarrow G \mid \varphi \text{ is an epimorphism that satisfies (FT)}\}$$

and observe that it is acted upon (by postcomposition) by the group $\text{Aut}(G)$ of automorphisms of G . Thanks to the discussion leading up to Theorem 1.4, note that sending a G -ribbon disc with aspherical exterior to an epimorphism $\pi_1(M_K) \twoheadrightarrow \pi_1(N_D) \cong G$ defines a map

$$\Phi: \mathcal{D}_G^a(K) \rightarrow \text{Epi}^{FT}(\pi_1(M_K), G) / \text{Aut}(G)$$

which does not depend on the choice of the isomorphism $\pi_1(N_D) \cong G$. If G is a good geometrically 2-dimensional ribbon group such that $\tilde{L}_4(\mathbb{Z}[G]) = 0$, then Theorem 1.4 ensures that Φ is surjective. Our main technical result gives conditions on G for Φ to be injective.

Theorem 1.7. *Let K be a knot and let G be a geometrically 2-dimensional good group that satisfies (W) and (A5). If D_1 and D_2 are two G -ribbon discs with aspherical exteriors and boundary K such that $\Phi(D_1) = \Phi(D_2)$, then D_1 and D_2 are ambiently isotopic rel. boundary.*

We note that this result can alternatively be stated with normal subgroups instead of epimorphisms as this is easier to verify in practice. To state this concisely, given a slice disc D for a knot K , we use $\iota_D: \pi_1(M_K) \rightarrow \pi_1(N_D)$ to denote the inclusion induced map.

Corollary 1.8. *Let K be a knot and let G be a geometrically 2-dimensional good group that satisfies (W) and (A5). If D_1 and D_2 are two G -ribbon discs with aspherical exteriors and boundary K such that $\ker(\iota_{D_1}) = \ker(\iota_{D_2})$, then D_1 and D_2 are ambiently isotopic rel. boundary.*

For smoothly embedded discs, the hypotheses of these results can be relaxed.

Remark 1.9. If D_1 and D_2 are ribbon discs with aspherical exteriors and $\pi_1(N_{D_i}) \cong G$ for $i = 1, 2$, then the assumption that G be geometrically 2-dimensional can be omitted in both Theorem 1.7 and Corollary 1.8: in this case $K(G, 1) \simeq N_{D_i}$ has the homotopy type of a 2-complex.

Combining Theorems 1.4 and 1.7, we obtain an answer to Question 1.1 provided we make some restrictions on the ribbon group G and require the ribbon disc exteriors to be aspherical.

Theorem 1.10. *Let $K \subset S^3$ be a knot and let G be a geometrically 2-dimensional good ribbon group that satisfies properties *W-AA*. Mapping a G -ribbon disc D to the epimorphism $\pi_1(M_K) \twoheadrightarrow \pi_1(N_D) \cong G$ defines a bijection Φ between the two following sets:*

- (1) the set $\mathcal{D}_G^a(K)$ of G -ribbon discs with aspherical exterior and boundary K , considered up to ambient isotopy rel. boundary;
- (2) the set $\text{Epi}^{FT}(\pi_1(M_K), G) / \text{Aut}(G)$ defined in (Epi).

Proof. We argue in Remark 2.1 below that since G is a geometrically 2-dimensional ribbon group with $\text{Wh}(G) = 0$, requiring G to satisfy condition (A4) is equivalent to asking for $\tilde{L}_4(\mathbb{Z}[G]) = 0$. Thus the hypotheses of Theorem 1.4 are satisfied and so Φ is surjective. The injectivity of Φ follows from Theorem 1.7 which we can apply since G satisfies properties *W-AA*. \square

Remark 1.11. We collect a couple of remarks on this result.

³In the work of Hambleton, Kreck and Teichner [HKT09] *W-AA* only requires A_4 to be injective.

- If the ribbon group conjecture (or more optimistically the Whitehead conjecture) were true, then requiring G to be geometrically 2-dimensional would be superfluous; recall Remark 1.3. It is also tempting to conjecture that exteriors of G -ribbon discs are aspherical and in this case we would have $\mathcal{D}_G^a(K) = \mathcal{D}_G(K)$. This latter conjecture holds when G is PTFA [CP21, Lemma 2.1] (e.g. when $G = \mathbb{Z}$ and $G = BS(1, 2)$) and is a consequence of the Whitehead conjecture if the disc exterior is homotopy equivalent to a 2-complex.
- The groups \mathbb{Z} and $BS(1, 2)$ satisfy the hypotheses of Theorem 1.10 and in this case, unpacking the definition of $\text{Epi}^{\text{FT}}(\pi_1(M_K), G)/\text{Aut}(G)$ recovers [CP21, Theorems 1.5 and 1.6]. Instead of repeating those statements, we note that for $G = \mathbb{Z}$, $\text{Epi}^{\text{FT}}(\pi_1(M_K), G)/\text{Aut}(G)$ has at most one element, while for $G = BS(1, 2)$ it has at most 2 [CP21, Section 4]. Estimating the cardinality of this set in general appears to be more challenging. Naturally, the set $\mathcal{D}_G(K)$ is often empty: for example, we refer to [FT05, Corollary 3.4] for an obstruction (based on the Alexander polynomial) to a knot K bounding a G -ribbon disc.
- As we alluded to in Corollary 1.8, the classification result of Theorem 1.10 can be stated in terms of normal subgroups of $\pi_1(M_K)$ instead of epimorphisms originating from $\pi_1(M_K)$: to a G -ribbon disc D , one associates the normal subgroup $\ker(\pi_1(M_K) \twoheadrightarrow \pi_1(N_D))$ of $\pi_1(M_K)$. This was the perspective taken in [CP21] where, using that $BS(1, 2)$ is metabelian, the results were then formulated using submodules of the Alexander module $H_1(M_K; \mathbb{Z}[t^{\pm 1}])$; the details are in [CP21, Section 3].
- The requirement that the group be good is hard to verify in practice. On the other hand G satisfies property W-AA if it is geometrically 2-dimensional and satisfies the Farrell-Jones conjecture: if a group G is geometrically 2-dimensional, then $K(G, 1)$ is a 2-complex and the claim now follows as in [KL20, Lemma 2.3] (the core of the argument will be recalled both in the proof of Theorem 1.7 and in Remark 2.1). We treat the Farrell-Jones conjecture as a blackbox, but refer the interested reader to [Lue21] for a survey and to [Lue21, Chapter 15] for a list of groups for which the conjecture is known to hold.

Example 1.12. We argue that if the knot group $G(J)$ of a classical knot $J \subset S^3$ is good, then it satisfies the hypotheses of Theorem 1.10; thus if $G(J)$ is good, this result classifies $G(J)$ -ribbon discs for $J\# - J$. The group of $J \subset S^3$ is ribbon (the ribbon knot $J\# - J$ bounds a smoothly embedded ribbon disc with group $G(J)$ as explained in [FT05, page 2135]). The sphere theorem ensures that $G(J)$ is geometrically 2-dimensional (the knot exterior is aspherical and has the homotopy type of a 2-complex; see e.g. [Lic97, Theorem 11.7]). The Farrell-Jones conjecture holds for $G(J)$ because it holds for the fundamental group of any 3-manifold with boundary [Lue21, Theorem 15.1 (e)]. Since knot groups are PTFA by work of Strebel [Str74], $G(J)$ -ribbon discs are aspherical by [CP21, Lemma 2.1] and thus $\mathcal{D}_{G(J)}(J\# - J) = \mathcal{D}_{G(J)}^a(J\# - J)$. Finally, as we noted in Remark 1.5, since $G(J)$ is PTFA, we can use condition (Ext) instead of condition (FT).

Example 1.13. For $m, n \in \mathbb{Z}$ with $|m - n| = 1$, we argue that if the Baumslag-Solitar group $BS(m, n)$ is good, then it satisfies the conditions of Theorem 1.10; thus if $BS(m, n)$ is good, this result classifies $BS(m, n)$ -ribbon discs with aspherical exterior. The fact that $BS(m, n)$ is ribbon when $|m - n| = 1$ can be seen by looking at the handle diagram depicted in Figure 1. Baumslag-Solitar groups are geometrically 2-dimensional: the universal cover of the presentation 2-complex for $\langle a, b \mid ba^mb^{-1} = b^n \rangle$ is homeomorphic to the product of \mathbb{R} with a tree; see e.g. [FKS11, Section 2]. Additionally, every Baumslag-Solitar group $BS(m, n)$ satisfies the Farrell-Jones conjecture [FW15, GMR15].

We conclude with a brief final remark concerning asphericity. The methods of this paper rely heavily on G -ribbon disc exteriors (conjecturally) being aspherical. Currently, non-aspherical 4-manifolds with boundary M_K and fundamental group G are poorly understood beyond the group $G = \mathbb{Z}$ [CP20]. This is the reason why we only work in D^4 instead of in other 4-manifolds.

Acknowledgments. I wish to thank Daniel Kasprowski and Markus Land for insightful correspondence related to [KL20] and for helpful comments on a draft of this paper. I am also grateful to Lisa Piccirillo for explaining to me why $BS(m, n)$ is ribbon when $|m - n| = 1$ and to Jonathan Hillman for pointing me towards [Hil02, Corollary 2.6.1].




FIGURE 1. Assuming that $|m - n| = 1$, this figure depicts a handle diagram of a ribbon disc exterior with fundamental for $BS(m, n)$. Indeed, since $|m - n| = 1$, the red and blue knots form a handle diagram for D^4 in which the green knot is sliced by a ribbon disc D with $\pi_1(N_D) = BS(m, n)$.

Conventions. Throughout this article, we work in the topological category. Manifolds are assumed to be compact and oriented. Homeomorphisms, homotopy equivalences and isotopies are *rel. boundary* if they fix the boundary pointwise. If M_1, M_2 are two n -manifolds with boundary Y , a cobordism between M_1 and M_2 is *relative Y* if, when restricted to Y , it is the product $Y \times [0, 1]$.

2. PROOF OF THE MAIN TECHNICAL RESULT

We recall the statement of Theorem 1.7 and prove it. Let K be a knot and let G be a geometrically 2-dimensional good group that satisfies (W) and (A5). The aim is to prove that if D_1 and D_2 are two G -ribbon discs with aspherical exteriors and boundary K such that $\Phi(D_1) = \Phi(D_2) \in \text{Epi}^{FT}(\pi_1(M_K), G)/\text{Aut}(G)$, then D_1 and D_2 are ambiently isotopic rel. boundary.

Proof of Theorem 1.7. Assume that D_1 and D_2 are two G -ribbon discs with aspherical exteriors and boundary K and that their epimorphisms agree in $\text{Epi}^{FT}(\pi_1(M_K), G)/\text{Aut}(G)$. We must show that D_1 and D_2 are ambiently isotopic rel. boundary. As noted in [CP21, Lemma 2.5], Alexander’s trick ensures that it is equivalent to prove that their exteriors N_{D_1} and N_{D_2} are homeomorphic rel. boundary. Our strategy decomposes into two steps. The first step uses the conditions on the epimorphisms to show that id_{M_K} extends to a homotopy equivalence $N_{D_1} \simeq N_{D_2}$. The second step uses surgery theory to improve this homotopy equivalence to a homeomorphism rel. boundary; here is where we rely on properties (W) and (A5) as well as on the fact that G is good.

We start with the first step. Since the epimorphisms of D_1 and D_2 agree, there exists an automorphism Ψ of G that makes the following diagram commute:

$$\begin{array}{ccc}
 \pi_1(M_K) & \xrightarrow{=} & \pi_1(M_K) \\
 \iota_{D_1} \downarrow & & \downarrow \iota_{D_2} \\
 \pi_1(N_{D_1}) & & \pi_1(N_{D_2}) \\
 \cong \downarrow & & \downarrow \cong \\
 G & \xrightarrow{\Psi, \cong} & G.
 \end{array}$$

Since the bottom vertical maps in this diagram are isomorphisms, we deduce that there exists an isomorphism $g: \pi_1(N_{D_1}) \cong \pi_1(N_{D_2})$ such that $g \circ \iota_{D_1} = \iota_{D_2}$; such isomorphisms were called *compatible* in [CP21, Section 2]. As the D_i have aspherical exteriors, the obstruction theory argument from [CP21, end of proof of Lemma 2.1] shows that the identity $\text{id}_{M_K}: M_K \rightarrow M_K$ extends to a homotopy equivalence $f: N_{D_1} \rightarrow N_{D_2}$ which induces g on fundamental groups.

We now move on to the second step: we use surgery theory to improve the homotopy equivalence f to a homeomorphism $N_{D_1} \cong N_{D_2}$ rel. boundary. We describe the argument very briefly for readers that are familiar with surgery theory before giving some more details. Consider the

surgery sequence, where we can ignore decorations thanks to condition **(W)**:

$$\mathcal{N}(N_{D_2} \times [0, 1], \partial(N_{D_2} \times [0, 1])) \xrightarrow{\sigma_5} L_5(\mathbb{Z}[G]) \rightarrow \mathcal{S}(N_{D_2}, \partial N_{D_2}) \xrightarrow{\eta} \mathcal{N}(N_{D_2}, \partial N_{D_2}) \xrightarrow{\sigma_4} L_4(\mathbb{Z}[G]).$$

We use that disc exteriors have trivial H_2 to deduce that η is the zero map. More concretely, we obtain a degree one normal map

$$(1) \quad (F', f, \text{id}_{N_{D_2}}): (W', N_{D_1}, N_{D_2}) \rightarrow (N_{D_2} \times [0, 1], N_{D_2}, N_{D_2})$$

that we can assume to be 2-connected by surgery below the middle dimension. We then use property **(A5)** and the fact that G is geometrically 2-dimensional to deduce that σ_5 is surjective. We infer that N_{D_1} and N_{D_2} are s -cobordant either by appealing to the exactness of the surgery sequence (which requires G to be good) or by using the surjectivity of σ_5 to replace F' by another degree one normal map with vanishing surgery obstruction (despite being slightly longer, this argument has the advantage of not requiring G to be good).⁴ The result then follows from Freedman's 5-dimensional relative s -cobordism theorem which we can apply because G is good.

We give more details. The set $\mathcal{N}(N_{D_2}, \partial N_{D_2})$ consists of equivalence classes of degree one normal maps $M \rightarrow N_{D_2}$ that restrict to a homeomorphism on the boundary. Two such degree one normal maps $f_i: M_i \rightarrow N_{D_2}$ for $i = 1, 2$ are equivalent if there exists a rel. boundary cobordism (W, M_1, M_2) and a degree one normal map

$$(W, M_1, M_2) \rightarrow (N_{D_2} \times [0, 1], N_{D_2}, N_{D_2})$$

that restricts to f_i on M_i for $i = 1, 2$. A homotopy equivalence $h: M \rightarrow N_{D_2}$ rel. boundary is in particular a degree one normal map that we denote by $\eta(h) \in \mathcal{N}(N_{D_2}, \partial N_{D_2})$.

We claim that η is the zero map. Under the isomorphism

$$(2) \quad \mathcal{N}(N_{D_2}, \partial N_{D_2}) \cong H^4(N_{D_2}, \partial N_{D_2}) \oplus H^2(N_{D_2}, \partial N_{D_2}; \mathbb{Z}_2) = H^4(N_{D_2}, \partial N_{D_2}) \cong \mathbb{Z}$$

we have $\eta(h) = \frac{1}{8}(\sigma(M) - \sigma(N_{D_2}))$; this fact is well known to surgeons but we refer to [CP21, Proposition 2.2] in case the reader is curious about the details. Since the signature of a disc exterior vanishes and h is a homotopy equivalence, we deduce that $\eta(h) = 0$, as claimed.

We assert that the map σ_5 from the surgery sequence is surjective. This relies on surgery spectra and the algebraic theory of surgery. We treat this largely as a blackbox but note that this part of surgery theory was developed by Quinn [Qui70, Qui71] and Ranicki [Ran79, Ran81]; we also refer to [CW21, Section 4.4] for a nice overview of these topics and to [CMR09, Section 4] for a helpful account of the rel. boundary case. Using the relation between the assembly map and the surgery obstruction (as mentioned for example in [CW21, page 158]) and the fact that N_{D_2} is a $K(G, 1)$, the following diagram commutes:

$$\begin{array}{ccc} \mathcal{N}(N_{D_2} \times [0, 1], \partial(N_{D_2} \times [0, 1])) & \xrightarrow{\sigma_5} & L_5(\mathbb{Z}[G]) \\ \downarrow \cong & & \downarrow = \\ H_5(N_{D_2}; \mathbf{L}\langle 1 \rangle_\bullet) & \xrightarrow{\cong} & H_5(N_{D_2}; \mathbf{L}_\bullet) \longrightarrow L_5(\mathbb{Z}[G]) \\ \downarrow \cong & & \downarrow \cong \\ H_5(BG; \mathbf{L}\langle 1 \rangle_\bullet) & \xrightarrow{\cong} & H_5(BG; \mathbf{L}_\bullet) \xrightarrow{A_5} L_5(\mathbb{Z}[G]). \end{array}$$

Here \mathbf{L}_\bullet denotes the L -theory spectrum of the integers and $\mathbf{L}\langle 1 \rangle_\bullet$ denotes its 1-connective cover. The fact that $H_5(BG; \mathbf{L}\langle 1 \rangle_\bullet) \rightarrow H_5(BG; \mathbf{L}_\bullet)$ is an isomorphism follows because $K(G, 1)$ admits a 2-dimensional CW-model (the Atiyah-Hirzebruch spectral sequence argument is the same as in [KL20, proof of Lemma 2.3]) and the fact that the top left vertical map is an isomorphism is a fact from algebraic surgery theory; see e.g. [CMR09, Equation (27)]. Using this commutative diagram and property **(A5)** (which stipulates that the assembly map A_5 is surjective), one deduces that σ_5 is surjective.

There are now two closely related ways to conclude that N_{D_1} and N_{D_2} are s -cobordant rel. boundary. The first way is shorter but uses that the group G is good: since η is the zero map, σ_5 is surjective and the surgery sequence is exact (because G is good), the structure set $\mathcal{S}(N_{D_2}, \partial N_{D_2})$ (to which f belongs) is trivial. The second argument (inspired by [KL20]) is slightly longer

⁴Thus the fact that N_{D_1} and N_{D_2} are s -cobordant rel. boundary can be proved without using that G is good.

but does not require that the group G be good: since $\eta \equiv 0$, there is a rel. boundary cobordism (W, N_{D_1}, N_{D_2}) and a degree one normal map

$$(F, f, \text{id}_{N_{D_2}}): (W, N_{D_1}, N_{D_2}) \rightarrow (N_{D_2} \times [0, 1], N_{D_2}, N_{D_2}).$$

Perform surgery below the middle dimension on the interior of W to obtain the 2-connected degree one normal map F' with surgery obstruction $x := \sigma(F') \in L_5(\mathbb{Z}[G])$ that we alluded to in (1). Using the surjectivity of σ_5 , one can find a degree one normal map

$$\Psi: (V, N_{D_2}, N_{D_2}) \rightarrow (N_{D_2} \times [0, 1], N_{D_2}, N_{D_2})$$

that restricts to the identity on both boundary components and with $-x$ as its surgery obstruction; stacking Ψ on top of F' leads to a degree one normal map F'' with vanishing surgery obstruction $\sigma(F'') \in L_5(\mathbb{Z}[G])$ and it follows that F'' is normal bordant rel. $M_K \times [0, 1]$ to a homotopy equivalence. Thus, we have two arguments for why N_{D_1} and N_{D_2} are s-cobordant rel. boundary.

Since G is good, we can apply Freedman's 5-dimensional relative s -cobordism theorem [FQ90, Theorem 7.1A] and it follows that N_{D_1} and N_{D_2} are homeomorphic rel. boundary. As we already mentioned, [CP21, Lemma 2.5] implies that the discs are ambiently isotopic rel. boundary. \square

We conclude by proving a statement that was used in the proof of Theorem 1.10.

Remark 2.1. Assume that G is a geometrically 2-dimensional ribbon group with vanishing Whitehead torsion (condition (W)). We claim that G satisfies $\tilde{L}_4(\mathbb{Z}[G]) = 0$ if and only if it satisfies (A4), which stipulates that the assembly map $A_4: H_4(BG; \mathbf{L}_\bullet) \rightarrow L_4(\mathbb{Z}[G])$ is an isomorphism. Since G is a ribbon group, there is a (smoothly embedded) ribbon disc $D \subset D^4$ with $\pi_1(N_D) \cong G$. This time, N_D might not be aspherical, but it is still a 2-complex with vanishing H_2 . An Atiyah-Hirzebruch spectral sequence argument therefore shows that $H_4(N_D; \mathbf{L}\langle 1 \rangle_\bullet) \rightarrow H_4(BG; \mathbf{L}\langle 1 \rangle_\bullet)$ is an isomorphism. Here, it is helpful to note that $H_2(G) = 0$: use $H_2(N_D) = 0$ together with the exact sequence $\pi_2(N_D) \rightarrow H_2(N_D) \rightarrow H_2(\pi_1(N_D)) \rightarrow 0$; see e.g. [Bro82, Equation (0.1)]. The same argument as above then produces the following commutative diagram:

$$\begin{array}{ccccc} \mathcal{N}(N_D, M_K) & \xrightarrow{\sigma_4} & & \xrightarrow{\quad} & L_4(\mathbb{Z}[G]) \\ \downarrow \cong & & & & \downarrow = \\ H_4(N_D; \mathbf{L}\langle 1 \rangle_\bullet) & \xrightarrow{\cong} & H_4(N_D; \mathbf{L}_\bullet) & \longrightarrow & L_4(\mathbb{Z}[G]) \\ \downarrow \cong & & \downarrow \cong & & \downarrow = \\ H_4(BG; \mathbf{L}\langle 1 \rangle_\bullet) & \xrightarrow{\cong} & H_4(BG; \mathbf{L}_\bullet) & \xrightarrow{A_4} & L_4(\mathbb{Z}[G]). \end{array}$$

As explained in (2) and [FQ90, Section 11.3B], the surgery obstruction σ_4 maps the set of normal invariants $\mathcal{N}(N_D, \partial N_D) \cong \mathbb{Z}$ isomorphically onto the $L_4(\mathbb{Z}) \cong \mathbb{Z}$ -summand of $L_4(\mathbb{Z}[G]) = L_4(\mathbb{Z}) \oplus \tilde{L}_4(\mathbb{Z}[G])$. The claim now follows by combining this fact with the commutativity of the diagram.

REFERENCES

- [BKK⁺21] Stefan Behrens, Boldizár Kalmaár, Min Hoon Kim, Mark Powell, and Arunima Ray. *The disc embedding theorem*. Oxford University press, 2021.
- [Bro82] Kenneth S. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1982.
- [CMR09] Matija Cencelj, Yuri Muranov, and Dušan Repovš. On structure sets of manifold pairs. *Homology Homotopy Appl.*, 11(2):195–222, 2009.
- [CP20] Anthony Conway and Mark Powell. Embedded surfaces with infinite cyclic knot group. *Preprint, available at arXiv:2009.13461*, 2020.
- [CP21] Anthony Conway and Mark Powell. Characterisation of homotopy ribbon discs. *Adv. Math.*, 391:Paper No. 107960, 29, 2021.
- [CW21] Stanley Chang and Shmuel Weinberger. *A course on surgery theory*, volume 211 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2021.
- [DMS22] Irving Dai, Abishek Mallick, and Matthew Stoffregen. Equivariant knots and knot floor homology. 2022. <https://arxiv.org/abs/2201.01875>.
- [FKS11] Eric M. Freden, Teresa Knudson, and Jennifer Schofield. Growth in Baumslag-Solitar groups I: subgroups and rationality. *LMS J. Comput. Math.*, 14:34–71, 2011.
- [FQ90] Michael H. Freedman and Frank Quinn. *Topology of 4-manifolds*, volume 39 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1990.

- [Fre82] Michael Hartley Freedman. The topology of four-dimensional manifolds. *J. Differential Geometry*, 17(3):357–453, 1982.
- [FT05] Stefan Friedl and Peter Teichner. New topologically slice knots. *Geom. Topol.*, 9:2129–2158, 2005.
- [FW15] Thomas Farrell and Xiaolei Wu. Isomorphism conjecture for Baumslag-Solitar groups. *Proc. Amer. Math. Soc.*, 143(8):3401–3406, 2015.
- [GMR15] Giovanni Gandini, Sebastian Meinert, and Henrik Rüping. The Farrell-Jones conjecture for fundamental groups of graphs of abelian groups. *Groups Geom. Dyn.*, 9(3):783–792, 2015.
- [Gor81] Cameron Gordon. Ribbon concordance of knots in the 3-sphere. *Math. Ann.*, 257(2):157–170, 1981.
- [Hay20] Kyle Hayden. Exotic ribbon disks and symplectic surfaces. 2020. <https://arxiv.org/abs/2003.13681>.
- [Hay21] Kyle Hayden. Corks, covers, and complex curves. 2021. <https://arxiv.org/abs/2107.06856>.
- [Hil02] Jonathan Hillman. *Four-manifolds, geometries and knots*, volume 5 of *Geometry & Topology Monographs*. Geometry & Topology Publications, Coventry, 2002.
- [HKS⁺21] Kyle Hayden, Alexandra Kjuchukova, Krishna Siddhi, Maggie Miller, Mark Powell, and Nathan Sunukjian. Brunnian exotic surface links in the 4-ball. 2021. <https://arxiv.org/abs/2106.13776>.
- [HKT09] Ian Hambleton, Matthias Kreck, and Peter Teichner. Topological 4-manifolds with geometrically two-dimensional fundamental groups. *J. Topol. Anal.*, 1(2):123–151, 2009.
- [How82] James Howie. On locally indicable groups. *Math. Z.*, 180(4):445–461, 1982.
- [How85] James Howie. On the asphericity of ribbon disc complements. *Trans. Amer. Math. Soc.*, 289(1):281–302, 1985.
- [HS21] Kyle Hayden and Isaac Sundberg. Khovanov homology and exotic surfaces in the 4-ball. 2021. <https://arxiv.org/abs/2108.04810>.
- [JZ20] András Juhász and Ian Zemke. Distinguishing slice disks using knot Floer homology. *Selecta Math. (N.S.)*, 26(1):Paper No. 5, 18, 2020.
- [KL20] Daniel Kasprowski and Markus Land. Topological 4-manifolds with 4-dimensional fundamental group. 2020. <https://arxiv.org/abs/2007.03399>.
- [Lic97] W. B. Raymond Lickorish. *An introduction to knot theory*, volume 175 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1997.
- [Lue20] Wolfgang Lueck. Assembly maps. In *Handbook of homotopy theory*, CRC Press/Chapman Hall Handb. Math. Ser., pages 851–890. CRC Press, Boca Raton, FL, [2020] ©2020.
- [Lue21] Wolfgang Lueck. *Isomorphism Conjectures in K- and L-Theory*. 2021. <https://www.him.uni-bonn.de/lueck/data/ic.pdf>.
- [Qui70] Frank Quinn. A geometric formulation of surgery. In *Topology of Manifolds (Proc. Inst., Univ. of Georgia, Athens, Ga., 1969)*, pages 500–511. Markham, Chicago, Ill., 1970.
- [Qui71] Frank Quinn. $B_{(\text{TOP}_n)^-}$ and the surgery obstruction. *Bull. Amer. Math. Soc.*, 77:596–600, 1971.
- [Ran79] Andrew Ranicki. The total surgery obstruction. In *Algebraic topology, Aarhus 1978 (Proc. Sympos., Univ. Aarhus, Aarhus, 1978)*, volume 763 of *Lecture Notes in Math.*, pages 275–316. Springer, Berlin, 1979.
- [Ran81] Andrew Ranicki. *Exact sequences in the algebraic theory of surgery*, volume 26 of *Mathematical Notes*. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1981.
- [Ran92] Andrew Ranicki. *Algebraic L-theory and topological manifolds*, volume 102 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1992.
- [Ros07] Stephan Rosebrock. The Whitehead conjecture—an overview. *Sib. Elektron. Mat. Izv.*, 4:440–449, 2007.
- [SS21] Isaac Sundberg and Jonah Swann. Relative khovanov-jacobsson classes. 2021. <https://arxiv.org/abs/2103.01438>.
- [Str74] Ralph Strebel. Homological methods applied to the derived series of groups. *Comment. Math. Helv.*, 49:302–332, 1974.
- [Whi41] John Henry Constantine Whitehead. On adding relations to homotopy groups. *Ann. of Math. (2)*, 42:409–428, 1941.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE MA 02139, UNITED STATES
 Email address: anthonyconway@gmail.com