

Polynomial Turing Compressions for Some Graph Problems Parameterized by Modular-Width

Weidong Luo ^{*†}

Abstract

In this paper we investigate the parameterized complexity for NP-hard graph problems parameterized by a structural parameter modular-width. We develop a recipe that is able to simplify the process of obtaining polynomial Turing compressions for a class of graph problems parameterized by modular-width. By using the recipe, we demonstrate that several problems, which include CHROMATIC NUMBER, INDEPENDENT SET, HAMILTONIAN CYCLE, etc. have polynomial Turing compressions parameterized by modular-width. In addition, under the assumption that $P \neq NP$, we provide tight kernels for a few problems such as STEINER TREE parameterized by modular-width. As a byproduct of the result of the tight kernels, new parameterized algorithms for these problems are obtained.

1 Introduction

Kernelization [14] is a theoretical formalization of efficient preprocessing for NP-hard problems. It is an important research direction in parameterized complexity [8, 11]. A *compression* or *generalized kernelization* for a parameterized problem $Q \subseteq \Sigma^* \times \mathbb{N}$ is a polynomial-time algorithm that, given any instance (x, k) , returns an instance (x', k') of an arbitrary parameterized problem Q' , such that $(x, k) \in Q$ if and only if $(x', k') \in Q'$, and $|x'| + k'$ is bounded by a function $f(k)$. We say that Q has a *polynomial compression* if $f(k)$ is a polynomial function of k . If we require the output to be an instance of Q , then the restricted compression is called *kernelization*. The output (x', k') of a kernelization is called *kernel*. Turing compression is a natural relaxation of compression. A *Turing compression* for a parameterized problem Q is a polynomial-time algorithm A that solves Q with the help of the oracle for an arbitrary parameterized problem Q' . More specifically, for an input instance (x, k) of Q , A is able to decide the instance by adaptively producing strings with lengths bounded by $f(k)$ bits and querying the oracle whether the strings are in Q' . We say that Q has a *polynomial Turing compression* if $f(k)$ is a polynomial function of k . We say that a parameterized problem is fixed-parameter tractable (FPT) if there is an $f(k) \cdot |x|^{\mathcal{O}(1)}$ time algorithm that can decide whether a string (x, k) is a yes instance of the problem, where f is a computable function. Note that a polynomial Turing compression for Q is not sufficient for an FPT algorithm for Q since Q' could be an undecidable problem. For compression, on the one hand, polynomial compressions for many parameterized problems have been obtained, on the other hand, many problems have been proved to have no polynomial compressions unless $\text{coNP} \subseteq \text{NP/poly}$. As the computational power of Turing compression is not weaker than that of compression, it is possible to obtain polynomial Turing compressions for some problems without polynomial compressions. More than ten years ago Binkele-Raible et al. [3] found the first problem, namely LEAF OUT-TREE(k), which has a polynomial Turing

^{*}Department of Computer Science, Humboldt-Universität zu Berlin, Germany

[†]Emails: luo@informatik.hu-berlin.de or weidong.luo@yahoo.com

compression but does not admit a polynomial compression unless $\text{coNP} \subseteq \text{NP}/\text{poly}$. However, by now, people only found limited problems of this kind [1, 4, 9, 18, 19, 20, 21, 24].

Modular-width is a structural graph parameter that was first proposed in [6]. Then, Gajarský et al. [15] introduce it into parameterized complexity. Modular-width is closely related to *modular decomposition* whose definition can be found in the preliminaries of this paper. A *module* of a graph $G = (V, E)$ is a set of vertices $M \subseteq V$ with the property that for each vertex $v \in V \setminus M$, either all vertices of M are adjacent to v or all vertices of M are not adjacent to v . Equivalently, M is a module if every vertex in M has the same neighbors among the vertices not in M . Obviously, the empty set, the vertex set V , and all one element sets $\{v\}$ for $v \in V$ are modules, furthermore, these are called the *trivial modules*. (We also call the empty set the empty module.) A graph is called *prime* if all modules of the graph are trivial modules. The modular-width of a graph G , which is denoted by $\text{mw} = \text{mw}(G)$, is the vertex number of a largest prime induced subgraph of G .

Our results. We study the polynomial Turing compressions for several graph problems parameterized by modular-width. We first regard the CLIQUE as an example to show the basic idea for getting a polynomial Turing compression parameterized by modular-width. Since the parameters of all the problems considered in this paper are modular-width, we observe that the ideas of constructing polynomial Turing compressions for these problems have some common place. So we make use of the properties of modular-width to develop a recipe that is able to assist us to design polynomial Turing compressions for these problems. In particular, our study gives rise to the following main result.

Theorem 1. *The following problems have polynomial Turing compressions parameterized by modular-width.*

- INDEPENDENT SET, CLIQUE, and VERTEX COVER.
- CHROMATIC NUMBER, DOMINATING SET, and NONBLOCKER.
- HAMILTONIAN CYCLE, HAMILTONIAN PATH, and PARTITIONING INTO PATHS.
- FEEDBACK VERTEX SET, ODD CYCLE TRANSVERSAL, and CONNECTED VERTEX COVER.
- LONGEST INDUCED PATH, MAXIMUM INDUCED FOREST, and INDUCED MATCHING.
- INDEPENDENT TRIANGLE PACKING and INDEPENDENT CYCLE PACKING.

In addition, we obtain linear kernels for some problems parameterized by modular-width. Moreover, the kernels for these problems are the best ones that can be obtained under the assumption that $\text{P} \neq \text{NP}$.

Theorem 2. STEINER TREE, CONNECTED DOMINATING SET, and MAX LEAF SPANNING TREE parameterized by modular-width have kernels with mw vertices and do not admit kernels with $\text{mw} - 1$ vertices unless $\text{P} = \text{NP}$.

For any graph G the connected domination number of G plus the number of leaves in the maximum leaf spanning tree of G equals the vertex number of G [10]. So the following corollary is obtained by applying the $\mathcal{O}(1.36^n)$ time exact exponential algorithm on the kernel of STEINER TREE [22], and the $\mathcal{O}(1.94^n)$ time exact exponential algorithm on the kernel of CONNECTED DOMINATING SET [12].

Corollary 1. STEINER TREE has an $\mathcal{O}^*(1.36^{\text{mw}})$ time algorithm. CONNECTED DOMINATING SET and MAX LEAF SPANNING TREE have $\mathcal{O}^*(1.94^{\text{mw}})$ time algorithms.

Note that Theorem 5.13 in [2] implies the best previous algorithm for each of the problems in Corollary 1 whose time complexity is $\mathcal{O}(c^{\text{mw}})$ for a large constant c .

2 Preliminaries

Throughout this paper, we only consider undirected graphs. Let Σ denote $\{0, 1\}$. For a graph G , we use \overline{G} to denote the complementary graph of G . Unless otherwise specified, $V(G)$ and $E(G)$ indicate the vertex set and the edge set of a graph G , respectively. For a vertex $v \in V(G)$, $N(v)$ represents the set of all neighbors of v . Vertex set $N[v] = N(v) \cup \{v\}$. Unless otherwise stated, a graph is denoted by a set whose elements are the vertices and edges of it. If M is a subset of the vertex set $V(G)$, then $G[M]$ denotes the subgraph induced in G by M . Vertex set $N(M)$ exactly includes all vertices that are not in M but are adjacent to some vertices of M . Vertex set $N[M] = N(M) \cup M$. The cardinality of a set S is denoted by $|S|$. Symbols \mathbb{G} and \mathbb{N} denote the set of undirected graphs and the set of natural numbers, respectively. Unless otherwise specified, $\text{mw}(G)$ (or mw) denotes the modular-width of a graph G . For any two vertex sets M and M' , the edges between M and M' are all edges (u, v) such that $u \in M$ and $v \in M'$. We use K_n to indicate a complete graph with n vertices and C_n to denote a cycle with n vertices. If X is a set of vertices, we write $G - X$ for the subgraph induced in G by $V(G) \setminus X$. In addition, we simply use $G - G'$ to represent $G - V(G')$. The \mathcal{O}^* -notation suppresses factors that are polynomial in the input size. The definitions of the problems discussed in this paper can be found in Appendix B.

Next, we give the formal definition of Turing compression as well as the definition of modular decomposition and its related concepts. First, recall the definition of Turing compression given by Binkele-Raible et al. [3].

Definition 1. A t -oracle for a parameterized problem Q is an oracle that takes as input (x, k) with $|x| \leq t$, $|k| \leq t$, and decides whether $(x, k) \in Q$ in constant time.

Definition 2. A Turing compression for a parameterized problem Q of size $g(k)$ is an algorithm that, given an input (x, k) together with a $g(k)$ -oracle for a parameterized problem Q' , decides whether a given instance $(x, k) \in Q$ in time polynomial in $|x| + k$.

We sometimes abuse terminology by saying that Q has a $g(k)$ Turing compression if there is a Turing compression for Q of size $g(k)$. We say Q has a polynomial Turing compression if Q has a $g(k)$ Turing compression, where $g(k)$ is a polynomial function of k . Note that if the oracle problem is restricted to Q , then the restricted Turing compression is called *Turing kernelization*.

Assume that $G = (V, E)$ is a graph. We call a module M of G a *strong module* if for any module M' of G , either M' is a subset of M , M is a subset of M' , or the intersection of M and M' is empty. This means that M is a strong module of G if M does not overlap with any other module of G . A module M of G is *maximal* if $M \subsetneq V$ and there is no modular M' such that $M \subsetneq M' \subsetneq V$. Let $P = \{M \mid M \subseteq V\}$ be a vertex partition of V . If every element M in P is a module of G , then P is a *modular partition*. Since P is a partition of V , any two modules M and M' of P are disjoint. Therefore, either each vertex of M is adjacent to all vertices of M' or no vertices of M are adjacent to a vertex of M' . Additionally, we say that M and M' are *adjacent* if each vertex of M is adjacent to all vertex of M' , and that M and M' are *non-adjacent* if no vertices of M are adjacent to a vertex of M' . A modular partition P that only contains maximal strong modules is a *maximal modular partition*. Notice that a graph has an unique maximal modular partition. For each modular partition P of V , the *quotient graph* $G_{/P} = (V_M, E_P)$ can be defined as follows. For a module M of P , let v_M represent its corresponding vertex. All vertices v_M constitute the vertex set V_M . If any two modules M and M' of P are adjacent, then the edge $(v_M, v_{M'})$ is a member of E_P . In other words, the vertices of the quotient graph $G_{/P}$ are the modules of P , and the adjacencies of these modules form the edges of $G_{/P}$.

For a module M of G , the subgraph $G[M]$ is called a *factor*. Suppose $P = \{M \mid M \subseteq V(G)\}$ is a modular partition of $V(G)$. Clearly, G can be reconstructed if all factors related to P and the quotient graph G/P are given. Furthermore, we consider every factor related to P . It can be recursively decomposed until one vertex factors are obtained. There is a quotient graph at each recursive level, and every bottom factor is a single vertex. Clearly, G can be reconstructed recursively in a bottom-up fashion by combining factors with quotient graph in each recursive level. Since modular partition of a graph may not be unique, a graph may have more than one way to recursively decompose. Next, we introduce a particular way to recursively decompose a graph, which is called *modular decomposition*. Recall that a module is strong if it does not overlap any other modules. Let us remark that all strong modules of a graph $G = (V, E)$ can be represented by an inclusion tree $MD(G)$. Each strong module M of G corresponds to a vertex v_M of $MD(G)$. The root vertex v_V of $MD(G)$ corresponds to the strong module V of G . Every leaf $v_{\{v\}}$ of $MD(G)$ corresponds to a vertex v of G . For any two strong modules M and M' of G , $v_{M'}$ is a descendant of v_M in $MD(G)$ if and only if M' is a proper subset of M . The inclusion tree $MD(G)$ is called the *modular decomposition tree* of G . Clearly, M' is a maximal strong module of $G[M]$ if and only if $v_{M'}$ is a child of v_M . We sometimes abuse terminology by saying that M' is a module of the vertex v_M , and $G[M']$ is a factor of the vertex v_M .

Modular decomposition theorem [16] proved that for any graph $G = (V, E)$, one of the following three conditions is satisfied: (1) G is not connected, (2) \overline{G} is not connected, (3) Both G and \overline{G} are connected and the quotient graph G/P is a prime graph, where P is the maximal modular partition of V . Consider an internal vertex v_M of $MD(G)$ and its corresponding module M of G . If $G[M]$ is disconnected, then the vertex set of each connected component is a maximal strong module of $G[M]$ and all these maximal strong modules is a maximal modular partition of $G[M]$. Thus, every child of v_M is corresponding to the vertex set of a connected component of $G[M]$ and the quotient graph of v_M is an independent set. Furthermore, the internal vertex v_M in this case is called *parallel*. If $\overline{G[M]}$ is disconnected, then the vertex set of each connected component of $\overline{G[M]}$ is a maximal strong module of $G[M]$ and all these maximal strong modules is a maximal modular partition of $G[M]$. Thus, every child of v_M is corresponding to the vertex set of a connected component of $\overline{G[M]}$ and the quotient graph of v_M is a complete graph. Furthermore, the internal vertex v_M in this case is called *series*. If both $G[M]$ and $\overline{G[M]}$ are connected. According to the modular decomposition theorem, $G[M]/P$ is a prime graph, where P is the maximal modular partition of M . Since every module in P is a maximal strong module, every child of v_M is corresponding to a maximal strong module in P and the quotient graph of v_M is the prime graph $G[M]/P$. Furthermore, the internal vertex v_M in this case is called *prime*. Collectively, the internal vertices of $MD(G)$ have three types, namely, parallel, series and prime. Since the maximal modular partition of a graph is unique, the children and the quotient graph of an internal vertex are unique. Therefore, the modular decomposition tree of a graph is unique. In addition, G can be reconstructed from $MD(G)$ if the parallel vertices and series vertices are labeled their types and the prime vertices are attached their quotient graphs. Given a graph G , the modular decomposition tree of G can be obtained in time $O(m + n)$ [23]. Additionally, the *modular-width* of a graph G can also be defined as the minimum number k such that the children number of any prime vertex in $MD(G)$ is at most k . More information about modular decomposition can be found in [17]. Note that we do not consider the null graph, which is the graph having no vertices, and the empty module in the proofs of this paper. (The results in this paper are trivial on the null graph or the empty module.)

Let $\text{cw}(G)$ denote the celebrated parameter *clique-width* of a graph G [7]. According to the definitions of clique-width and modular-width, we have $\text{cw}(G) \leq \text{mw}(G) + 2$. Clearly, the $(\text{mw} + 2)$ -expression for any graph G can be obtained in polynomial time. Thus, Theorem 5.13 in [2] implies that CONNECTED VERTEX COVER, CONNECTED DOMINATING SET, and STEINER

TREE have $2^{\mathcal{O}(\text{mw})}n^{\mathcal{O}(1)}$ time algorithms, and Theorem 5 in [5] implies that DOMINATING SET can be solved in $\mathcal{O}^*(4^{\text{mw}})$ time.

3 An example for Turing compression

To see how a modular decomposition and low modular width can be helpful for getting small Turing compressions, let us first consider the CLIQUE problem as an example. The *clique number* $\omega(G)$ is the number of vertices in a maximum clique of G . In this paper, we define that the *weighted clique number* of a weighted graph is the total weight of the vertices of a maximum weighted clique in the weighted graph.

Lemma 1. CLIQUE(mw) has a polynomial Turing compression.

Proof. Let (G, k) be an input of CLIQUE(mw). Suppose M is a module of $G = (V, E)$. Assume C is a subset of V such that $G[C]$ is a maximum clique of G . Assume S' is a subset of M such that $G[S']$ is a maximum clique of $G[M]$. Clearly, the intersection of $G[C]$ and $G[M]$ is a clique. More strongly, we claim that the intersection of $G[C]$ and $G[M]$ is either the empty set or a maximum clique of $G[M]$. If $C \cap M = S \neq \emptyset$ where $G[S]$ is not a maximum clique of $G[M]$, then each vertex of S is adjacent to all vertices of $C \setminus M = C \setminus S$ and $|S| < |S'|$. Since M is a module and S is a subset of M , each vertex of M is adjacent to all vertices of $C \setminus S$, so the subgraph induced by $(C \setminus S) \cup S'$ is a clique of G whose vertex number is larger than $|C|$. It contradicts that $G[C]$ is a maximum clique of G . Next, we will first compute $\omega(G)$ using a polynomial-time algorithm with an oracle for WEIGHTED CLIQUE problem, and then compare $\omega(G)$ with k to decide whether (G, k) is a yes instance of CLIQUE(mw).

Since CLIQUE(mw) can be solved in $1.74^{\text{mw}}|V|^{\mathcal{O}(1)}$ time [13], we can solve it in polynomial time if $|V| \geq 1.74^{\text{mw}}$. Without loss of generality, assume henceforth that $|V| < 1.74^{\text{mw}}$. We first obtain the modular decomposition tree $MD(G)$ in linear time. In order to obtain $\omega(G)$, we compute $\omega(G[M])$ for every vertex v_M of $MD(G)$ in a bottom-up fashion. If v_M is a leaf of $MD(G)$, then $\omega(G[M]) = 1$. If v_M is an internal vertex of $MD(G)$. Let $P = \{M' \mid M' \subseteq M\}$ be the maximal modular partition of M . Then every child $v_{M'}$ of v_M is corresponding to a module M' in P and the quotient graph of v_M is $G[M]_{/P}$. If v_M is a parallel vertex, then $\omega(G[M])$ is the maximum one of $\omega(G[M'])$ for all M' . If v_M is a series vertex, then $\omega(G[M])$ is the sum of $\omega(G[M'])$ for all M' . If v_M is a prime vertex. We solve a WEIGHTED CLIQUE problem: Take the quotient graph $G[M]_{/P}$ and assign a weight $\omega(G[M'])$ to each vertex $v_{M'}$, then query an oracle for WEIGHTED CLIQUE at most $|V|$ times to obtain the weighted clique number of the weighted quotient graph. The input of each query is the weighted quotient graph together with a parameter whose value is at most $|V|$. Clearly, the weighted clique number of the weighted quotient graph is equal to $\omega(G[M])$. Next, we analyze the size of each query. We know the weighted quotient graph has at most mw vertices and the integer weight of each vertex is not great than $|V|$. Moreover, since $|V| < 1.74^{\text{mw}}$, the binary encoding lengths of the parameter and the integer weight of each vertex are at most mw bits. Thus, the size of each query is $\mathcal{O}(\text{mw}^2)$ bits. Overall, we prove that CLIQUE(mw) has an $\mathcal{O}(\text{mw}^2)$ Turing compression. \square

By the well-known relations between CLIQUE, INDEPENDENT SET, and VERTEX COVER, and because modular-width does not change under graph complementation, we have the following corollary.

Corollary 2. INDEPENDENT SET(mw) and VERTEX COVER(mw) have polynomial Turing compressions.

4 Recipe

Given graphs $G = (V, E)$ and $H = (V_H, E_H)$, as well as a module M of G . Assume G_S is a supergraph of G , which is obtained as follows: (1) add all vertices and edges of both graphs G and H into G_S , (2) for every vertex v of $N(M)$ and every vertex u of H , add edge (v, u) to G_S . Let G' be the subgraph induced in G_S by $(V \setminus M) \cup V_H$, and we say that G' is obtained from G by replacing the factor $G[M]$ with H . Furthermore, we call the process of obtaining G' from G a *modular replacement*. Clearly, V_H is a module of G' .

Recall that, for each modular partition P of $V(G)$, the quotient graph $G_{/P} = (V_M, E_P)$ can be defined as follows. For each module M of P , let v_M be its corresponding vertex. All vertices v_M constitute the vertex set V_M . If any two modules M and M' of P are adjacent, then the edge $(v_M, v_{M'})$ is a member of E_P .

Lemma 2. *Let each F_i be a function from \mathbb{G} to \mathbb{N} , where $i \in \{1, \dots, r\}$. For any graphs G and H , as well as any module M of G , suppose the following statement holds: that $F_i(G[M]) = F_i(H)$ for all i implies that $F_i(G) = F_i(G')$ for all i , where G' is obtained from G by replacing the factor $G[M]$ with H . Then, for any graph G and any modular partition P of $V(G)$, we have a conclusion that the quotient graph $G_{/P}$ together with the function values $F_1(G[M]), \dots, F_r(G[M])$ for all modules M in P completely determine the function values $F_1(G), \dots, F_r(G)$.*

Proof. Let tuple $T(G)$ denote the function values $(F_1(G), \dots, F_r(G))$. For a graph G and a modular partition P of $V(G)$, we say X is a values-attached quotient graph that is generated from P if X is constructed as follows: (1) X includes all the vertices and edges of the quotient graph $G_{/P} = (V_M, E_P)$, moreover, (2) for each vertex v_M of X , the tuple $T(G[M])$ is attached to v_M . Suppose that \mathbb{X} includes all X . Assume that the binary relation R over sets \mathbb{X} and \mathbb{G} includes all pairs (X, G) such that X is generated from some modular partition of $V(G)$. For each $i \in \{1, \dots, r\}$, let f_i be the composition relation of R and F_i over \mathbb{X} and \mathbb{N} , which means that $f_i = R; F_i = \{(X, n) \mid \text{there exists a graph } G \text{ of } \mathbb{G} \text{ such that } (X, G) \in R \text{ and } (G, n) \in F_i\}$.

Consider each binary relation f_i . According to the definition of R , for any X of \mathbb{X} , there is at least one element G in \mathbb{G} such that $(X, G) \in R$. Furthermore, as F_i is a function, for any G of \mathbb{G} , there is an element n in \mathbb{N} such that $(G, n) \in F_i$. Therefore, for any X of \mathbb{X} , there is at least an element n in \mathbb{N} such that $(X, n) \in f_i$. This means that the binary relation f_i is left-total. For an X in \mathbb{X} , assume that \mathbb{G}' consists of all G such that $(X, G) \in R$. Next, we demonstrate that $F_i(G_1)$ equals $F_i(G_2)$ for any graphs $G_1, G_2 \in \mathbb{G}'$. Assume that G_1 and G_2 are in \mathbb{G}' . As (X, G_1) and (X, G_2) are elements of R , there exists a modular partition P_1 of G_1 such that X is generated from P_1 , and there exists a modular partition P_2 of G_2 such that X is also generated from P_2 . Moreover, the quotient graph G_{1/P_1} and the quotient graph G_{2/P_2} are isomorphic, and there exists an edge-preserving bijection g from the vertex set of G_{1/P_1} to the vertex set of G_{2/P_2} such that, for every vertex v_{M_1} of G_{1/P_1} and $v_{M_2} = g(v_{M_1})$ of G_{2/P_2} , the tuple $T(G[M_1])$ equals the tuple $T(G[M_2])$. Additionally, we say a factor $G[M_2]$ of G_2 is corresponding to a factor $G[M_1]$ of G_1 if $v_{M_2} = g(v_{M_1})$. Next, consider every module M_1 of P_1 . We replace each factor $G[M_1]$ of G_1 with its corresponding factor $G[M_2]$ of G_2 one by one. Since $T(G[M_1])$ equals to $T(G[M_2])$, according to the prerequisite of this lemma, the value of function F_i for the new obtained graph after every modular replacement does not change.¹ After the final modular replacement, G_2 is obtained, and we have that $F_i(G_1)$ equals $F_i(G_2)$. Therefore, $F_i(G)$ is a fixed number for any $G \in \mathbb{G}'$, and, for every $X \in \mathbb{X}$, there is a fixed number n such that $(X, n) \in f_i$. This means that the binary relation f_i is right-unique. Consequently, the binary relation f_i is a function from \mathbb{X} to \mathbb{N} , moreover, $F_i(G)$ is equal to $f_i(X)$, where $(X, G) \in R$. \square

¹The prerequisite of this lemma is the second sentence of Lemma 2.

Let each F_i be a function from \mathbb{G} to \mathbb{N} , where $i \in \{1, \dots, r\}$. For each i , we say an algorithm solves $F_i(G)$ if, for every input G , the algorithm produces the function value $F_i(G)$. The decision version of each $F_i(G)$ is as follows: given a graph G and an integer k , decide whether $F_i(G) \leq k$ (or $F_i(G) \geq k$). Recall that mw denotes the modular-width of G .

Lemma 3. *Let each F_i be a function from \mathbb{G} to \mathbb{N} , where $i \in \{1, \dots, r\}$ and r is a constant. If the following statements hold.*

(1) *For any graphs G and H , as well as any module M of G , that $F_i(H) = F_i(G[M])$ for all i implies that $F_i(G) \leq F_i(G')$ (or $F_i(G) \geq F_i(G')$) for all i , where G' is obtained from G by replacing the factor $G[M]$ with H .*

(2) *For each i , there is a $2^{\text{mw}^{\mathcal{O}(1)}}|G|^{\mathcal{O}(1)}$ time algorithm to solve $F_i(G)$.*

(3) *$F_i(G) \leq |G|^{\mathcal{O}(1)}$ for each i .*

Then each $F_i(G)$ can be solved by a polynomial-time algorithm together with an oracle for a language Q that is able to decide whether a string of length $\text{mw}^{\mathcal{O}(1)}$ is in Q . Moreover, the decision version of each $F_i(G)$ has a polynomial Turing compression parameterized by mw .

Proof. For all i , assume c is a large enough constant such that $F_i(G) \leq |G|^c$ and $F_i(G)$ can be solved by a $2^{\text{mw}^c}|G|^c$ time algorithm. Without loss of generality, suppose $\text{mw} \geq \log^{\frac{1}{c}}|G|$. (Otherwise, each $F_i(G)$ can be solved in polynomial time based on the second statement.) If the first statement hold. Based on the definition of G' , H is a factor of G' . Hence, for the graphs G' and $G[M]$, as well as the module $V(H)$ of G' , that $F_i(G[M]) = F_i(H)$ for all i implies that $F_i(G') \leq F_i(G)$ (or $F_i(G') \geq F_i(G)$) for all i , where G is obtained from G' by replacing the factor H with $G[M]$. As a result, we have that $F_i(G) = F_i(G')$ for all i . This means that functions F_1, \dots, F_r fulfill the conclusion of Lemma 2. Let the definitions of values-attached quotient graph X , set \mathbb{X} , binary relation R and functions f_1, \dots, f_r be the same as that in Lemma 2. For every i , since f_i is a function, $f_i(X)$ is a fixed number. Suppose that language Q includes all the strings $(X, f_1(X), \dots, f_r(X))$, where $X \in \mathbb{X}$.

Recall that every vertex v_M of the modular decomposition tree $MD(G)$ of a graph G is corresponding to the strong module M of G , where $MD(G)$ can be constructed in linear time. Here, we call $G[M]$ the corresponding graph of v_M . Roughly speaking, in order to solve the functions $F_1(G), \dots, F_r(G)$, we compute in a bottom-up fashion the values of all F_i for the graphs that are corresponding to the vertices of $MD(G)$. First, since the corresponding graph of every leaf of $MD(G)$ is the singleton graph K_1 , according to the second statement, every $F_i(K_1)$ can be solved in $\mathcal{O}(1)$ time. Secondly, consider an internal vertex v_M of $MD(G)$. Let $P = \{M' \mid M' \subseteq M\}$ be the maximal modular partition of M . Then every child $v_{M'}$ of v_M is corresponding to the module M' in P and the quotient graph of v_M is $G[M]_{/P}$. If v_M is a prime vertex. Assume $X \in \mathbb{X}$ is the values-attached quotient graph that is generated from P . Since the function values $F_1(G[M']), \dots, F_r(G[M'])$ for all modules M' in P and the quotient graph $G[M]_{/P}$ are given, X can be obtained immediately. Let k_1, \dots, k_r be non-negative integers all of which are less than $|G|^c$. Exhaustively generate string (X, k_1, \dots, k_r) and query the oracle whether the string is a member of Q . Based on Lemma 2, we have $F_i(G[M]) = f_i(X)$ for all i . Additionally, we also have $F_i(G[M]) \leq |G|^c$ for all i . Hence, we obtain the function values $F_1(G[M]), \dots, F_r(G[M])$ by finding the string $(X, k_1, \dots, k_r) = (X, f_1(X), \dots, f_r(X))$ after querying the oracle at most $(|G|^c + 1)^r = |G|^{\mathcal{O}(1)}$ times. Furthermore, the length of binary code of every string (X, k_1, \dots, k_r) is at most $\mathcal{O}(\text{mw}^2 \log |G|)$, which is $\text{mw}^{\mathcal{O}(1)}$ since $\text{mw} \geq \log^{\frac{1}{c}}|G|$. If v_M is a parallel vertex. Suppose P contains t modules, where $2 \leq t \leq |V(G)|$. The quotient graph $G[M]_{/P}$ is \overline{K}_t . Consider any two modules M'_1 and M'_2 of P . Let M'_{12} be $M'_1 \cup M'_2$ and $G[M'_{12}]$ be the subgraph induced by M'_{12} in $G[M]$. Then, $P' = \{M'_1, M'_2\}$ is a modular partition of M'_{12} in $G[M'_{12}]$. Since the values-attached quotient graph that is generated from P' are given, we can obtain the function values $F_1(G[M'_{12}]), \dots, F_r(G[M'_{12}])$ by using

the same method as that of the prime vertex. Now, consider the new modular partition $P_1 = \{M'_{12}\} \cup (P \setminus \{M'_1, M'_2\})$ of M , which contains only $t-1$ modules and the quotient graph $G[M]_{/P_1}$ is \bar{K}_{t-1} . Moreover, for every module M' in P_1 , the function values $F_1(G[M']), \dots, F_r(G[M'])$ are obtained. Thus, if we repeat the above-mentioned process $t-2$ times, then the function values $F_1(G[M]), \dots, F_r(G[M])$ can be obtained in the last time. If v_M is a series vertex. We can solve $F_1(G[M]), \dots, F_r(G[M])$ by using the same strategy as that of the parallel vertex. Therefore, for any internal vertex v_M of $MD(G)$, we obtain $F_1(G[M]), \dots, F_r(G[M])$ in $|G|^{\mathcal{O}(1)}$ time and the length of every string that is generated to query the oracle for Q is bounded by $\text{mw}^{\mathcal{O}(1)}$. In addition, the vertex number of $MD(G)$ is $O(|G|)$. As a result, functions $F_1(G), \dots, F_r(G)$ can be solved by a polynomial-time algorithm together with an oracle for Q that is able to decide whether a string of length $\text{mw}^{\mathcal{O}(1)}$ is in Q . Furthermore, it is obvious that the decision version of each $F_i(G)$ has a polynomial Turing compression parameterized by mw . \square

5 Polynomial Turing compression for problems

In this section, we demonstrate polynomial Turing compressions for several problems parameterized by modular-width with the help of the recipe provided in section 4.

Lemma 4. CHROMATIC NUMBER(mw) has a polynomial Turing compression.

Proof. Suppose that function $\chi(G)$ represents the chromatic number of a graph $G = (V, E)$. We demonstrate the problem has a polynomial Turing compression by proving $\chi(G)$ fulfills the statements of Lemma 3. Theorem 6 in [15] gives an $\mathcal{O}^*(2^{\text{mw}})$ algorithm for $\chi(G)$, so function $\chi(G)$ fulfills the statement 2 of Lemma 3. Since $\chi(G)$ is not greater than $|V|$, it satisfies the statement 3 of Lemma 3.

Next, we prove $\chi(G)$ fulfills the statement 1 of Lemma 3. Let $H = (V_H, E_H)$ be a graph and M be a module of G . Assume graph $G' = (V', E')$ is obtained from G by replacing the factor $G[M]$ with H . Let M' be $N(M)$. Suppose $\chi(G[M]) = \chi(H)$. Assume integer $q = \chi(G)$ and C is a color set with q different colors. Then, there exists a coloring $c : V \rightarrow C$ for G . Suppose C_M consists of all $c(v)$, where $v \in M$. Assume $C_{M'}$ consists of all $c(v)$, where $v \in M'$. As each vertex of M is adjacent to all vertices of M' , the intersection of C_M and $C_{M'}$ is empty. Consider the graph G' . Since $\chi(H) = \chi(G[M])$ and $\chi(G[M]) \leq |C_M|$, there exists a coloring $c_H : V_H \rightarrow C_M$ for H . Suppose $c' : V' \rightarrow C$ is a function such that $c'(v) = c(v)$ for all $v \in V' \setminus V_H$ and $c'(v) = c_H(v)$ for all $v \in V_H$. Clearly, for any two adjacent vertices $u, v \in V' \setminus V_H$ or $u, v \in V_H$, we have $c'(v) \neq c'(u)$. For any two adjacent vertices $u \in V' \setminus V_H$ and $v \in V_H$, since $N(V_H) = M'$ and the intersection of C_M and $C_{M'}$ is empty, we have $c'(v) \neq c'(u)$. Hence, function c' is a coloring for G' and G' is q -colorable. Consequently, $\chi(G') \leq q = \chi(G)$. \square

For a graph G , a partition of G into paths is a set of vertex disjoint paths of G whose union contains every vertex of G . Suppose L is a path of G and X is a subset of $V(G)$. We say L traverses $G[X]$ if the intersection of L and $G[X]$ is not empty. Assume function $c(G)$ is the number of the connected components of a graph G .

Let $L = (U, F)$ be a path with n vertices, where vertex set $U = \{u_1, \dots, u_n\}$ and edge set $F = \{(u_1, u_2), \dots, (u_{n-1}, u_n)\}$. We say L' is a sub-path of L if $L' = (U', F')$ such that $U' = \{u_i, \dots, u_j\}$ and $F' = \{(u_i, u_{i+1}), \dots, (u_{j-1}, u_j)\}$, where $1 \leq i \leq j \leq n$. Let $P = (W, D)$ be a path with m vertices, where vertex set $W = \{w_1, \dots, w_m\}$ and edge set $D = \{(w_1, w_2), \dots, (w_{m-1}, w_m)\}$. The operation of substituting the path P for the sub-path L' of L is as follows: (1) delete all vertices and edges of L' as well as edges $(u_{i-1}, u_i), (u_j, u_{j+1})$ from L . (2) add all

vertices and edges of P as well as edges $(u_{i-1}, w_1), (w_m, u_{j+1})$ into L .² Clearly, L is again a path with $n + m - (j - i + 1)$ vertices.

Lemma 5. PARTITIONING INTO PATHS(mw) and HAMILTONIAN CYCLE(mw) have polynomial Turing compressions.

Proof. Assume function $F_v(G)$ represents the vertex number of $G = (V, E)$. Suppose function $F_{pip}(G)$ denotes the least integer k such that G has a partition into k paths. Assume $F_{hc}(G) = 0$ if G has a Hamiltonian cycle, otherwise $F_{hc}(G) = 1$. We demonstrate that the problems have polynomial Turing compressions by proving $F_{hc}(G)$, $F_{pip}(G)$, and $F_v(G)$ fulfill all statements of Lemma 3. Theorem 7 in [15] gives $2^{\mathcal{O}(mw^2 \log mw)} |G|^{\mathcal{O}(1)}$ time algorithms for $F_{hc}(G)$ and $F_{pip}(G)$, so functions $F_{hc}(G)$ and $F_{pip}(G)$ fulfill the statement 2 of Lemma 3. Since $F_{hc}(G)$ and $F_{pip}(G)$ are not greater than $|V|$, they satisfy the statement 3 of Lemma 3. Obviously, $F_v(G)$ fulfills the statement 2 and statement 3 of Lemma 3.

Next, we prove that $F_{hc}(G)$, $F_{pip}(G)$, and $F_v(G)$ fulfill the statement 1 of Lemma 3. Let $H = (V_H, E_H)$ be a graph and M be a module of G . Assume graph $G' = (V', E')$ is obtained from G by replacing the factor $G[M]$ with H . Let M' be $N(M)$. Suppose $F_{hc}(G[M]) = F_{hc}(H)$, $F_{pip}(G[M]) = F_{pip}(H)$, and $F_v(G[M]) = F_v(H)$. Clearly, we have $F_v(G') \leq F_v(G)$.

First, consider $F_{pip}(G)$ and $F_{pip}(G')$. Assume \mathbb{L} is a partition of G into $F_{pip}(G)$ paths. Assume $\mathbb{O} \subseteq \mathbb{L}$ includes all the paths each of which is a subset of $G[V \setminus M]$, $\mathbb{P} \subseteq \mathbb{L}$ includes all the paths each of which traverse both $G[M]$ and $G[V \setminus M]$, and $\mathbb{Q} \subseteq \mathbb{L}$ includes all the paths each of which is a subset of $G[M]$. Obviously, \mathbb{L} is the union of \mathbb{P} , \mathbb{Q} , and \mathbb{O} . Suppose s equals the sum of $c(L \cap G[M])$ for all $L \in \mathbb{P}$. Then, we have

$$F_{pip}(G[M]) \leq s + |\mathbb{Q}| = \sum_{L \in \mathbb{P}} c(L \cap G[M]) + |\mathbb{Q}| \leq F_v(G[M])$$

In addition, we have $F_{pip}(G[M]) = F_{pip}(H)$ and $F_v(G[M]) = F_v(H)$. Hence, there exists a partition of H into paths \mathbb{R} whose cardinality is $s + |\mathbb{Q}|$. Now, we construct a set \mathbb{L}' as follows. (1) Add all elements of \mathbb{O} into \mathbb{L}' . (2) Consider every path L of \mathbb{P} . Suppose $L \cap G[M]$ contains r connected components L_1, \dots, L_r . Clearly, each L_i is a sub-path of L , where $i \in \{1, \dots, r\}$. For every i , substitute a path of \mathbb{R} for the sub-path L_i of L and delete the path from \mathbb{R} . After substituting for all the r sub-paths of L , we obtain a new path L of G' and add L to \mathbb{L}' . (3) Add all remaining paths of \mathbb{R} into \mathbb{L}' . Clearly, all elements of \mathbb{L}' are paths of G' and any two paths of \mathbb{L}' have no intersection. Next, we will demonstrate that the union of the paths of \mathbb{L}' contains every vertex of G' . First, \mathbb{R} is a partition of H into paths, and every path of \mathbb{R} is added into \mathbb{L}' either as an element of it in the third step or as a sub-path of an element of it in the second step. Hence, the union of the paths of \mathbb{L}' contains every vertex of V_H . Second, according to the definition of \mathbb{O} and \mathbb{P} , we have that $V \setminus M$ is a subset of $\bigcup_{L \in (\mathbb{O} \cup \mathbb{P})} V(L)$. Apart from the vertices of $\bigcup_{L \in \mathbb{P}} V(L \cap G[M])$, which are not vertices of $V \setminus M$, every vertex of $\bigcup_{L \in (\mathbb{O} \cup \mathbb{P})} V(L)$ is included in some path that is added into \mathbb{L}' in the first or the second step. Hence, the union of the paths of \mathbb{L}' contains every vertex of $V \setminus M$. Consider the cardinality of \mathbb{L}' . Obviously, $|\mathbb{O}|$ paths are added into \mathbb{L}' in the first step and $|\mathbb{P}|$ paths are added into \mathbb{L}' in the second step. As s paths are deleted from \mathbb{R} in the second step, $|\mathbb{Q}|$ paths are added into \mathbb{L}' in the third step. Thus, the cardinality of \mathbb{L}' equals that of \mathbb{L} . As a result, \mathbb{L}' is a partition of G' into $F_{pip}(G)$ paths, and $F_{pip}(G') \leq F_{pip}(G)$.

Then, consider Hamiltonian cycles of G and G' . Suppose C is a Hamiltonian cycle of G . Then, we have $c(C \cap G[M]) = c(C \cap G[V \setminus M])$. Since we have $F_{pip}(G[M]) = F_{pip}(H)$ and

²If $i - 1 \leq 0$, then vertex u_{i-1} does not exist and edges $(u_{i-1}, u_i), (u_{i-1}, w_1)$ are not taken into consideration. The explanation goes the same way if $j + 1 \geq n + 1$.

$F_v(G[M]) = F_v(H)$, there exists a partition of H into paths whose cardinality is $c(C \cap G[M])$. For the Hamiltonian cycle C , using the similar substitution operations as that for L in the proof for $F_{pip}(G') \leq F_{pip}(G)$, we can obtain a Hamiltonian cycle for G' . Thus, G has a Hamiltonian cycle implies G' has a Hamiltonian cycle, and $F_{hc}(G') \leq F_{hc}(G)$. \square

Corollary 3. HAMILTONIAN PATH(mw) has a polynomial Turing compression.

The domination number of a graph is the number of vertices in a smallest dominating set for the graph. Let function $\gamma(G)$ represent the domination number of a graph G .

Lemma 6. Let M be a module of a graph G . The intersection of M and a minimum dominating set of G is either \emptyset , $\{v\}$, or a minimum dominating set of $G[M]$, where v is a vertex of M .

Proof. Let D denote a minimum dominating set of G . The intersection of D and M is a minimum dominating set of $G[M]$ if $M' = \emptyset$. Assume henceforth $M' \neq \emptyset$.

If the intersection of D and M contains more than two vertices. Assume u is a vertex of M and v is a vertex of M' . As $N[M]$ is a subset of the union of $N[u]$ and $N[v]$, $D' = \{u, v\} \cup (D \setminus M)$ is a dominating set of G . It contradicts that D is a minimum dominating set of G .

If the intersection of D and M contains exact two vertices u, v . If $\gamma(G[M]) = 1$. Assume $\{w\}$ is a minimum dominating set of $G[M]$. Since $N[\{u, v\}]$ is a subset of $N[w]$, $D' = \{w\} \cup (D \setminus \{u, v\})$ is a dominating set of G . It contradicts that D is a minimum dominating set of G . If $\gamma(G[M]) \geq 3$, then the intersection of D and M' includes at least one vertex, otherwise there exists a vertex of M that is not adjacent to any vertex of D or in D . Thus, it is easy to verify that $D \setminus \{u\}$ (or $D \setminus \{v\}$) is a dominating set of G . It contradicts that D is a minimum dominating set of G . If $\gamma(G[M]) = 2$ and $\{u, v\}$ is not a dominating set of $G[M]$, then the intersection of D and M' contains at least one vertex, otherwise there exists a vertex of M that is not adjacent to any vertex of D or in D . Therefore, $D \setminus \{u\}$ (or $D \setminus \{v\}$) is a dominating set of G . It contradicts that D is a minimum dominating set of G . Thus, we have $\gamma(G[M]) = 2$ and $\{u, v\}$ is a dominating set of $G[M]$. This means that $\{u, v\}$ is a minimum dominating set of $G[M]$. \square

Lemma 7. DOMINATING SET(mw) has a polynomial Turing compression.

Proof. We demonstrate the problem has a polynomial Turing compression by proving $\gamma(G)$ fulfills all statements of Lemma 3. As we discussed in the preliminaries, Theorem 5 in [5] implies that $\gamma(G)$ can be solved in $\mathcal{O}^*(4^{mw})$ time. Thus, $\gamma(G)$ fulfills the statement 2 of Lemma 3. Since $\gamma(G)$ is not greater than $|V|$, it satisfies the statement 3 of Lemma 3.

Next, we prove $\gamma(G)$ fulfills the statement 1 of Lemma 3. Let $H = (V_H, E_H)$ be a graph and M be a module of G . Assume graph $G' = (V', E')$ is obtained from G by replacing the factor $G[M]$ with H . Let M' be $N(M)$. Suppose $\gamma(G[M]) = \gamma(H)$. Let D denote a minimum dominating set of G . Based on Lemma 6, the intersection of D and M is either \emptyset , $\{u\}$, or a minimum dominating set of $G[M]$, where u is a vertex in M . If the intersection of D and M is \emptyset , then the intersection of D and M' contains at least one vertex x . (Note that, as we have mentioned, we do not consider the empty module.) Furthermore, x is adjacent to all vertices of V_H in G' . Thus, D is a dominating set of G' and $\gamma(G') \leq \gamma(G)$. If the intersection of D and M is $\{u\}$. If $\gamma(G[M]) = 1$. Since $\gamma(G[M]) = \gamma(H)$, assume $\{v\}$ is a minimum dominating set of H . Clearly, $\{v\} \cup (D \setminus \{u\})$ is a dominating set of G' . Thus, we have $\gamma(G') \leq \gamma(G)$. If $\gamma(G[M]) \geq 2$, then the intersection of D and M' contains at least one vertex v , otherwise there exists a vertex in M that is not adjacent to any vertex of D or in D . Suppose w is a vertex of H . Clearly, $\{w\} \cup (D \setminus \{u\})$ is a dominating set of G' . Thus, we have $\gamma(G') \leq \gamma(G)$. If the intersection of D and M is a minimum dominating set of $G[M]$. Suppose D_H is a minimum dominating set of H . It is not hard to verify that $D_H \cup (D \setminus M)$ is a dominating set of G' . Additionally, since $\gamma(G[M]) = \gamma(H)$, we have $\gamma(G') \leq \gamma(G)$. \square

Since a nonblocker of a graph is the complement of a dominating set of the graph, and modular-width does not change under graph complementation, we have the following corollary.

Corollary 4. $\text{NONBLOCKER}(\text{mw})$ has a polynomial Turing compression.

Lemma 8. $\text{LONGEST INDUCED PATH}(\text{mw})$ has a polynomial Turing compression.

Proof. Let $G = (V, E)$ be a graph. Without loss of generality, suppose the vertex number of a longest induced path of G is not less than four. (Otherwise it can be solved in polynomial time.) Suppose functions $F_{lip}(G)$ and $F_v(G)$ represent the vertex number of a longest induced path and the vertex number of G , respectively. We demonstrate the problem has a polynomial Turing compression by proving $F_{lip}(G)$ and $F_v(G)$ fulfill all statements of Lemma 3. Since $F_{lip}(G)$ has an $\mathcal{O}^*(1.74^{\text{mw}})$ algorithm [13], it fulfills the statement 2 of Lemma 3. Since $F_{lip}(G)$ is not greater than $|V|$, it satisfies the statement 3 of Lemma 3. Obviously, $F_v(G)$ fulfills the statement 2 and statement 3 of Lemma 3.

Next, we prove $F_{lip}(G)$ and $F_v(G)$ fulfill the statement 1 of Lemma 3. Let $H = (V_H, E_H)$ be a graph and M be a module of G . Suppose graph $G' = (V', E')$ is obtained from G by replacing the factor $G[M]$ with H . Let M' be $N(M)$. Suppose $F_{lip}(G[M]) = F_{lip}(H)$ and $F_v(G[M]) = F_v(H)$. Obviously, we have $F_v(G) \leq F_v(G')$. Assume L is a longest induced path of G . Clearly, any vertex of L has at most two adjacent vertices of L in G . If $L \cap M$ is the empty set. Obviously, L is an induced path of G' and $F_{lip}(G) \leq F_{lip}(G')$. If $L \cap M$ contains exact one vertex u . There are at most two vertices v, w of L that are included in M' , otherwise u has more than two adjacent vertices of L . Furthermore, there is no edge between v and w , otherwise at least one vertex of v and w has three adjacent vertices in L since the vertex number of L is not less than four. Suppose, x is a vertex of H . Then, v and w are the only two vertices of L that are adjacent to x in G' . Therefore, $\{x, (x, v), (x, w)\} \cup (L \setminus \{u, (u, v), (u, w)\})$ is an induced path of G' . As a result, we have $F_{lip}(G) \leq F_{lip}(G')$. If $L \cap M$ contains exact two vertices $\{u, v\}$. Since the vertex number of L is not less than four, there is at least one vertex in $L \cap M'$. If $L \cap M'$ contains exact one vertex w , then there is at least one vertex $x \in L \cap (V \setminus N[M])$ and (w, x) is an edge of L . Thus, w is adjacent to three vertices u, v, x of L . If $L \cap M'$ has at least two vertices w and x , then there are at least four edges among the vertices u, v, w and x , which are (u, w) , (w, v) , (v, x) , and (x, u) . Therefore, the subgraph induced by vertex set $\{u, v, w, x\} \subseteq L$ contains a cycle with four vertices. If $3 \leq |L \cap M| < |V(L)|$. $L \cap M'$ contains at least one vertex v . Since v is adjacent to all vertices of M , at least three vertices of L are adjacent to v . If $L \cap M$ equals $|V(L)|$. It means that $G[M]$ contains an induced path with $|V(L)|$ vertices. Since $F_{lip}(G[M]) = F_{lip}(H)$, there exists an induced path with $|V(L)|$ vertices in H . Thus, we have $F_{lip}(G) \leq F_{lip}(G')$. \square

Let M be a module of a graph $G = (V, E)$, and S be a subset of $V \setminus M$. Assume graph G' is obtained from G by replacing the factor $G[M]$ with a graph $H = (V_H, E_H)$. Let $I_{G[M]}$ be an independent set of $G[M]$. Let I_H be an independent set of H . Let $G_1 = (V_1, E_1)$ be the subgraph in G induced by vertex set $S \cup I_{G[M]}$. Let $G_2 = (V_2, E_2)$ be the subgraph in G' induced by vertex set $S \cup I_H$.

Lemma 9. If the cardinality of I_H and $I_{G[M]}$ are equal, then G_1 and G_2 are isomorphic.

Proof. If $I_{G[M]}$ is empty. Clearly, I_H is empty. Hence, both G_1 and G_2 are the subgraph in $G[V \setminus M]$ induced by vertex set S . If S is empty. Both G_1 and G_2 are edgeless graph on the same vertices. If both $I_{G[M]}$ and S are not empty. Since the cardinality of I_H and $I_{G[M]}$ are equal, suppose $S = \{v_1, \dots, v_r\}$, $I_{G[M]} = \{u_1, \dots, u_s\}$, and $I_H = \{w_1, \dots, w_s\}$. Assume f is a bijection from V_1 to V_2 such that $f = \{[v_1, v_1], \dots, [v_r, v_r], [u_1, w_1], \dots, [u_s, w_s]\}$. According to the definition of G_1 and G_2 , it is easy to verify that f is an edge-preserving bijection. \square

Lemma 10. *Let $G = (V, E)$ be a graph. G has an edge if and only if $G[V \setminus S]$ has an edge, where S is a minimum feedback vertex set or a minimum odd cycle transversal of G .*

Proof. If S is a minimum feedback vertex set of G . For one direction, if G has an edge. $G[V \setminus S]$ has an edge if S is empty. If S is not empty. After deleting any $|S| - 1$ vertices of S from G , there is at least one cycle with at least 3 vertices in G , otherwise G has a feedback vertex set of $|S| - 1$ vertices. As the edges of a cycle with at least 3 vertices cannot be entirely removed by deleting one vertex, $G[V \setminus S]$ has at least one edge. The other direction is trivial.

If S is a minimum odd cycle transversal of G . The proof goes the same way. \square

Lemma 11. *Let M be a module of a graph $G = (V, E)$. The following statements hold:*

(1) *The intersection of M and a minimum vertex cover of G is either M or a minimum vertex cover of $G[M]$.*

(2) *The intersection of M and a minimum feedback vertex set of G is either M , $M \setminus \{v\}$, a minimum vertex cover of $G[M]$, or a minimum feedback vertex set of $G[M]$, where v is a vertex in M .*

(3) *The intersection of M and a minimum odd cycle transversal of G is either M , a minimum vertex cover of $G[M]$, or a minimum odd cycle transversal of $G[M]$.*

Proof. Let M' denote $N(M)$. Since M is a module, each vertex of M' is adjacent to all vertices of M , and each vertex of M is adjacent to all vertices of M' .

Let C represent a minimum vertex cover of G . If M' is empty. The intersection of M and C is a minimum vertex cover of $G[M]$. If M' is not empty. If the intersection of M and C does not equal M , then M' is a subset of C , otherwise there exists an edge between M and M' that is not covered by any vertex of C . Hence, the intersection of M and C is a minimum vertex cover of $G[M]$. As a result, the intersection of M and C is either M or a minimum vertex cover of $G[M]$.

Let F represent a minimum feedback vertex set of G . If M' is empty. The intersection of M and F is a minimum feedback vertex set of $G[M]$. If M' is not empty. If the subgraph induced by $M \setminus F$ has an edge, then M' is a subset of F , otherwise there exists a C_3 in $G[V \setminus F]$. Thus, the intersection of M and F is a minimum feedback vertex set of $G[M]$. If the subgraph induced by $M \setminus F$ has no edges and has at least two vertices, then $M' \setminus F$ contains at most one vertex, otherwise there exists a C_4 in $G[V \setminus F]$. Hence, the intersection of F and M is a minimum vertex cover of $G[M]$. If the subgraph induced by $M \setminus F$ is a single vertex v , then the intersection of F and M is $M \setminus \{v\}$, where v is a vertex of M . If $M \setminus F$ is empty, then the intersection of F and M is M . As a result, the intersection of M and F is either M , $M \setminus \{v\}$, a minimum vertex cover of $G[M]$, or a minimum feedback vertex set of $G[M]$.

Let O represent a minimum odd cycle transversal of G . If M' is empty, the intersection of M and O is a minimum odd cycle transversal of $G[M]$. If M' is not empty. If the subgraph induced by $M \setminus O$ contains an edge, then M' is a subset of O , otherwise there exists a C_3 in $G[V \setminus O]$. Hence, the intersection of M and O is a minimum odd cycle transversal of $G[M]$. If the subgraph induced by $M \setminus O$ contains a vertex but no edges, then the subgraph induced by $M' \setminus O$ contains no edges, otherwise there exists a C_3 in $G[V \setminus O]$. Therefore, the intersection of M and O is a minimum vertex cover of $G[M]$. If $M \setminus O$ is the empty set, then the intersection of M and O is M . As a result, the intersection of M and O is either M , a minimum vertex cover of $G[M]$, or a minimum odd cycle transversal of $G[M]$. \square

We know the vertex cover number of a graph is the vertex number of a minimum vertex cover of the graph. In this paper, connected vertex cover number, feedback vertex set number and odd cycle transversal number of a graph are defined in the analogous way.

Lemma 12. *For a connected graph $G = (V, E)$, let $M \subsetneq V$ be a module of it and C be a minimum connected vertex cover of it. If the vertex cover number of $G[M]$ is large than zero, then the intersection of M and C is either M or a minimum vertex cover of $G[M]$; if the vertex cover number of $G[M]$ equals zero, then the intersection of M and C is either M , a minimum vertex cover of $G[M]$, or $\{v\}$, where v is a vertex in M .*

Proof. Let M' denote $N(M)$. $M' \neq \emptyset$ since G is connected and $M \neq V$. Clearly, C always contains a vertex cover of $G[M]$ and a vertex cover of $G[M']$. If $M' \setminus C$ contains a vertex v , then the intersection of M and C is M , otherwise there exists a vertex u of M such that edge (v, u) is not covered by C . If $M' \setminus C$ is the empty set. Clearly, every edge between M' and M has at least one endpoint in C . If $G[C \setminus M]$ is connected, then the intersection of M and C is a minimum vertex cover of $G[M]$. If $G[C \setminus M]$ is not connected and the vertex cover number of $G[M]$ is larger than zero, then the intersection of M and C is a minimum vertex cover of $G[M]$. If $G[C \setminus M]$ is not connected and the vertex cover number of $G[M]$ is zero, then the intersection of M and C is a vertex of M to ensure the connectivity of $G[C]$. \square

If G has at least two connected components each of which contains at least one edge, then connected vertex covers of G do not exist and, in this case, we define the connected vertex cover number of G equals $|V(G)| + 1$.

Lemma 13. *CONNECTED VERTEX COVER(mw), FEEDBACK VERTEX SET(mw), and ODD CYCLE TRANSVERSAL(mw) have polynomial Turing compressions.*

Proof. Suppose functions $F_{oct}(G)$, $F_{fvs}(G)$, $F_{cvc}(G)$, $F_{vc}(G)$, and $F_v(G)$ represent the odd cycle transversal number, the feedback vertex set number, the connected vertex cover number, the vertex cover number and the vertex number of a graph $G = (V, E)$, respectively. We demonstrate the three problems have polynomial Turing compressions by proving all the functions fulfill the statements of Lemma 3. Clearly, all functions satisfy the statement 3 of Lemma 3, and $F_v(G)$ complies with the statement 2 of Lemma 3. As we discussed in the preliminaries, Theorem 5.13 in [2] implies a $2^{\mathcal{O}(\text{mw})}n^{\mathcal{O}(1)}$ algorithm for $F_{cvc}(G)$. There is an $\mathcal{O}^*(1.74^{\text{mw}})$ algorithm for both $F_{vc}(G)$ and $F_{fvs}(G)$ [13]. Furthermore, Lemma 18 in Appendix A demonstrates an $\mathcal{O}^*(3^{\text{mw}})$ algorithm for $F_{oct}(G)$. Hence, $F_{vc}(G)$, $F_{cvc}(G)$, $F_{fvs}(G)$, and $F_{oct}(G)$ fulfill the statement 2 of Lemma 3.

Next, we prove all functions fulfill the statement 1 of Lemma 3. Let $H = (V_H, E_H)$ be a graph and M be a module of G . Suppose graph $G' = (V', E')$ is obtained from G by replacing the factor $G[M]$ with H . Suppose $F_v(G[M]) = F_v(H)$, $F_{vc}(G[M]) = F_{vc}(H)$, $F_{cvc}(G[M]) = F_{cvc}(H)$, $F_{fvs}(G[M]) = F_{fvs}(H)$, and $F_{oct}(G[M]) = F_{oct}(H)$. Let vertex sets C , E , F , and O represent a minimum vertex cover, a minimum connected vertex cover, a minimum feedback vertex, and a minimum odd cycle transversal of G , respectively. Let vertex sets C_H , F_H , and O_H represent a minimum vertex cover, a minimum feedback vertex, and a minimum odd cycle transversal of H , respectively. Obviously, we have $F_v(G') \leq F_v(G)$.

First, consider the vertex cover numbers of G and G' . Based on Lemma 11, the intersection of C and M is either M or a minimum vertex cover of $G[M]$. Let S denote $V \setminus (C \cup M)$. If the intersection of C and M is M , then $(C \setminus M) \cup V_H$ is a vertex cover of G' . If the intersection of C and M is a minimum vertex cover of $G[M]$. According to Lemma 9, the subgraph induced by $S \cup (M \setminus C) = V \setminus C$ in G and the subgraph induced by $S \cup (V_H \setminus C_H)$ in G' are isomorphic. Therefore, the subgraph induced by $S \cup (V_H \setminus C_H)$ in G' is an independent set, and $(C \setminus M) \cup C_H$ is a vertex cover of G' . Consequently, we have $F_{vc}(G') \leq F_{vc}(G)$.

Secondly, consider the connected vertex cover numbers of G and G' . If G is connected. It is obvious that $F_{cvc}(G') \leq F_{cvc}(G)$ if $M = V$. Next, we assume $M \subsetneq V$. According to Lemma 12, the intersection of M and E is either M , a minimum vertex cover of $G[M]$, or $\{v\}$, where

v is a vertex of M . If the intersection of M and E is M . Clearly, $(E \setminus M) \cup V_H$ is a connected vertex cover of G' . If the intersection of M and E is a minimum vertex cover of $G[M]$. Let S denote $V \setminus (E \cup M)$. According to Lemma 9, the subgraph induced by $S \cup (V_H \setminus C_H)$ in G' and the subgraph induced by $S \cup (M \setminus E) = V \setminus E$ in G are isomorphic. Therefore, the subgraph induced by $S \cup (V_H \setminus C_H)$ in G' is an independent set, and $(E \setminus M) \cup C_H$ is a vertex cover of G' . If the subgraph in G induced by $E \setminus M$ is connected, then $(E \setminus M) \cup C_H$ is a connected vertex cover of G' . If the subgraph in G induced by $E \setminus M$ is not connected, then $E \cap M$ contains at least a vertex u such that every connected component of $G[E \setminus M]$ contains a vertex that is adjacent to u , otherwise the subgraph in G induced by E is not connected. Furthermore, since any vertex of C_H and u has the same neighbors in $E \setminus M$. Therefore, $(E \setminus M) \cup C_H$ is a connected vertex cover of G' . If the intersection of M and E contains exactly one vertex v . Without loss of generality, assume $\{v\} \subsetneq M$. (The case $M = \{v\}$ has been discussed.) Suppose u is also a vertex of M . Since G is connected, u is adjacent to every vertex of $N(M)$. Thus, $N(M)$ is a subset of E , otherwise there exists a vertex u' of $N(M)$ such that (u, u') is not covered by E . As every edge of $G[M]$ is covered by E , $\{v\}$ is a vertex cover of $G[M]$. Furthermore, since $F_{vc}(G[M]) = F_{vc}(H)$ and $F_v(G[M]) = F_v(H)$, there exists a vertex w of H such that w is a vertex cover of H . Therefore, $(E \setminus \{v\}) \cup \{w\}$ is a vertex cover of G' . In addition, according to Lemma 9, the subgraph induced by E in G and the subgraph induced by $(E \setminus \{v\}) \cup \{w\}$ in G' are isomorphic. Thus, the subgraph induced by $(E \setminus \{v\}) \cup \{w\}$ is connected. As a result, $(E \setminus \{v\}) \cup \{w\}$ is a connected vertex cover of G' . Consequently, we have $F_{cvc}(G') \leq F_{cvc}(G)$ if G is connected.

If G is not connected. If G has at least two connected components each of which contains at least one edge, then the connected vertex covers of G do not exist and $F_{cvc}(G) = |V| + 1$. In addition, we have $F_{cvc}(G') \leq |V'| + 1$ and $|V'| = |V|$. Thus, $F_{cvc}(G') \leq F_{cvc}(G)$. If G has one connected component K that contains at least one edge and the others are trivial connected components. (A trivial connected component is a singleton.) Suppose E is a minimum connected vertex cover of G . Clearly, E is a subset of K . If M contains no trivial connected components. Then M is a subset of K and the proof for this case goes the same as the proof for the case that G is connected. If M contains a trivial connected component and no vertices of K . Then, H has no edges as $F_{vc}(G[M]) = F_{vc}(H)$. Thus, E is a connected vertex cover of G' and $F_{cvc}(G') \leq F_{cvc}(G)$. If M contains a trivial connected component and a vertex of K . Then, M contains all vertices of K , otherwise M is not a module of G . Thus, $G[V \setminus M]$ are trivial connected components and $F_{cvc}(G) = F_{cvc}(G[M])$. In addition, we have $F_{cvc}(G[M]) = F_{cvc}(H)$. Moreover, $F_{cvc}(H) = F_{cvc}(G')$ as $G[V \setminus M]$ are trivial connected components. Thus, we have $F_{cvc}(G') \leq F_{cvc}(G)$. If G only contains trivial connected components. It is obvious that $F_{cvc}(G') \leq F_{cvc}(G)$.

Thirdly, consider the feedback vertex numbers of G and G' . Based on Lemma 11, the intersection of F and M is either M , $M \setminus \{v\}$, a minimum vertex cover of $G[M]$, or a minimum feedback vertex set of $G[M]$, where v is a vertex of M . Let S denote $V \setminus (F \cup M)$. If the intersection of F and M is M , then $(F \setminus M) \cup V_H$ is a feedback vertex set of G' . If the intersection of F and M is $M \setminus \{v\}$. Assume u is a vertex of H , based on Lemma 9, the subgraph induced by $S \cup \{v\}$ in G and the subgraph induced by $S \cup \{u\}$ in G' are isomorphic. Hence, the subgraph induced by $S \cup \{u\}$ in G' is a forest, and $(F \setminus M) \cup (V_H \setminus \{u\})$ is a feedback vertex set of G' . If the intersection of F and M is a minimum vertex cover of $G[M]$. Based on Lemma 9, the subgraph induced by $S \cup (M \setminus F) = V \setminus F$ in G and the subgraph induced by $S \cup (V_H \setminus C_H)$ in G' are isomorphic. So the subgraph induced by $S \cup (V_H \setminus C_H)$ in G' has no cycle, and $(F \setminus M) \cup C_H$ is a feedback vertex set of G' . If the intersection of F and M is a minimum feedback vertex set of $G[M]$. If the graph induced by $M \setminus F$ in G has no edges. Based on Lemma 10, $G[M]$ has no edges. Thus, H has no edges, moreover, G and G' are isomorphic

according to Lemma 9. If the graph induced by $M \setminus F$ in G has an edge, then $N(M)$ is a subset of F , otherwise there exists a triangle in $G[V \setminus F]$. Therefore, $(F \setminus M) \cup F_H$ is a feedback vertex set of G' . Consequently, we have $F_{fvs}(G') \leq F_{fvs}(G)$.

Finally, consider the odd cycle transversal numbers of G and G' . Based on Lemma 11, the intersection of O and M is either M , a minimum vertex cover of $G[M]$, or a minimum odd cycle transversal of $G[M]$. Let S denote $V \setminus (O \cup M)$. If the intersection of O and M is M , then $(O \setminus M) \cup V_H$ is an odd cycle transversal of G' . If the intersection of O and M is a minimum vertex cover of $G[M]$. According to Lemma 9, the subgraph induced by $S \cup (M \setminus O) = V \setminus O$ in G and the subgraph induced by $S \cup (V_H \setminus C_H)$ in G' are isomorphic. Therefore, the subgraph induced by $S \cup (V_H \setminus C_H)$ has no odd cycles, and $(O \setminus M) \cup C_H$ is an odd cycle transversal of G' . If the intersection of O and M is a minimum odd cycle transversal of $G[M]$. If the subgraph induced by $M \setminus O$ in G has no edges. Based on Lemma 10, $G[M]$ has no edges. Thus, H has no edges, moreover, G and G' are isomorphic according to Lemma 9. If the subgraph induced by $M \setminus O$ in G has an edge, then $N(M)$ is a subset of O , otherwise there exists a triangle in $G[V \setminus O]$. Hence, $(O \setminus M) \cup O_H$ is an odd cycle transversal of G' . Consequently, we have $F_{oct}(G') \leq F_{oct}(G)$. \square

By the well-known relation between FEEDBACK VERTEX SET and MAXIMUM INDUCED FOREST, since modular-width does not change under graph complementation, we have the following corollary.

Corollary 5. MAXIMUM INDUCED FOREST(mw) *has a polynomial Turing compression.*

Let the independent triangle packing number of a graph be the number of triangles of a maximum independent triangle packing of the graph. Furthermore, we define the independent cycle packing number and the induced matching number in the analogous way.

Lemma 14. INDUCED MATCHING(mw), INDEPENDENT TRIANGLE PACKING(mw), and INDEPENDENT CYCLE PACKING(mw) *have polynomial Turing compressions.*

Proof. Suppose functions $F_{im}(G)$, $F_{itp}(G)$, $F_{icp}(G)$, and $F_v(G)$ denote the induced matching number, the independent triangle packing number, the independent cycle packing number, and the vertex number of a graph $G = (V, E)$, respectively. We demonstrate the three problems have polynomial Turing compressions by proving $F_{im}(G)$, $F_{itp}(G)$, $F_{icp}(G)$, and $F_v(G)$ fulfill all the statements of Lemma 3. It is obvious that $F_v(G)$ fulfills the statement 2 of Lemma 3. Since $F_{im}(G)$, $F_{itp}(G)$, and $F_{icp}(G)$ can be solved in $\mathcal{O}^*(1.74^{\text{mw}})$ time [13], they fulfill the statement 2 of Lemma 3. Since all functions are not greater than $|V|$, they satisfy the statement 3 of Lemma 3.

Next, we prove all functions fulfill the statement 1 of Lemma 3. Let $H = (V_H, E_H)$ be a graph and M be a module of G . Graph $G' = (V', E')$ is obtained from G by replacing the factor $G[M]$ with H . Suppose $F_{im}(G[M]) = F_{im}(H)$, $F_{itp}(G[M]) = F_{itp}(H)$, $F_{icp}(G[M]) = F_{icp}(H)$, and $F_v(G[M]) = F_v(H)$. Let M' be $N(M)$. For vertices u, v, w of a graph, if (u, v) , (u, w) , and (v, w) are edges of the graph, then we use (u, v, w) to represent the triangle induced by vertices u, v , and w . Clearly, we have $F_v(G) \leq F_v(G')$.

First, consider the induced matching numbers of G and G' . Let edge set I represent a maximum induced matching of G . Let X denote the set of all vertices that are incident to the edges of I . Clearly, each vertex of X has exact one adjacent vertex in X . If $M' = \emptyset$, we have $F_{im}(G) \leq F_{im}(G')$ as $F_{im}(G[M]) = F_{im}(H)$. Suppose $M' \neq \emptyset$ henceforth. If the intersection of X and M is the empty set, then $F_{im}(G) = F_{im}(G[V \setminus M]) \leq F_{im}(G')$. If the intersection of X and M contains exact one vertex v , then there exists a vertex $u \in M'$ such that $(u, v) \in I$. Furthermore, the intersection of X and M' equals $\{u\}$, otherwise there is more than one vertex

in $X \cap M'$ that is adjacent to v . Therefore, for any vertex w in H , the intersection of $N(w)$ and X equals $\{u\}$. As a result, edge set $\{(w, u)\} \cup (I \setminus \{(v, u)\})$ is an induced matching of G' , and $F_{im}(G) \leq F_{im}(G')$. If the intersection of X and M contains more than one vertex, then the intersection of X and M' is the empty set, otherwise there is more than one vertex in $X \cap M$ that is adjacent to a vertex in $X \cap M'$. Thus, for every edge $e \in I$, both endpoints of e are either in M or in $V \setminus N[M]$. Suppose $I_{G[M]}$ is the intersection of I and $G[M]$. Assume I_H is a maximum induced matching of H . Then, $I_H \cup (I \setminus I_{G[M]})$ is an induced matching of G' . Moreover, since $F_{im}(G[M]) = F_{im}(H)$, the cardinality of I_H is not less than that of $I_{G[M]}$. Hence, $F_{im}(G) \leq F_{im}(G')$.

Secondly, consider the independent triangle packing numbers of G and G' . Let T represent a maximum independent triangle packing of G . Let Y be all vertices of T . Clearly, each vertex of Y has exact two adjacent vertices in Y . If $M' = \emptyset$, we have $F_{itp}(G) \leq F_{itp}(G')$ since $F_{itp}(G[M]) = F_{itp}(H)$. Suppose $M' \neq \emptyset$ hereafter. If the intersection of Y and M is the empty set, then $F_{itp}(G) = F_{itp}(G[V \setminus M]) \leq F_{itp}(G')$. If the intersection of Y and M contains exact one vertex u , then there exist two vertices $v, w \in M'$ such that (u, v, w) is a triangle of T . Moreover, the intersection of X and M' equals $\{v, w\}$, otherwise there are more than two vertices in $X \cap M'$ that are adjacent to u . Therefore, for any vertex x in H , the intersection of $N(x)$ and Y equals $\{v, w\}$. Thus, $(x, v, w) \cup (T \setminus (u, v, w))$ is an independent triangle packing of G' , and $F_{itp}(G) \leq F_{itp}(G')$. If the intersection of Y and M contains exact two vertices u and v . If u, v belong to different triangles of T , then the intersection of Y and M' has at least four vertices. This means that there are at least four vertices in $Y \cap M'$ that are adjacent to u (or v). If u, v are in the same triangle of T , then there is a vertex $w \in M'$ such that (u, v, w) is a triangle of T . Moreover, w is the only vertex of $Y \cap M'$, otherwise there are at least three vertices in $Y \cap N[M]$ that are adjacent to u (or v). Additionally, since $F_{im}(G[M]) = F_{im}(H)$ and $G[M]$ contains an edge (u, v) , there is at least one edge (x, y) in H . Thus, $(x, y, w) \cup (T \setminus (u, v, w))$ is an independent triangle packing of G' , and $F_{itp}(G) \leq F_{itp}(G')$. If the intersection of Y and M includes more than two vertices, then the intersection of Y and M' is the empty set, otherwise there are more than two vertices in $Y \cap M$ that are adjacent to a vertex in $Y \cap M'$. Thus, for every triangle (u, v, w) of T , all vertices u, v and w are either in M or in $V \setminus N[M]$. Suppose $T_{G[M]}$ is the intersection of T and $G[M]$. Assume T_H is a maximum independent triangle packing of H . Then, $T_H \cup (T \setminus T_{G[M]})$ is an independent triangle packing of G' . In addition, since $F_{itp}(G[M]) = F_{itp}(H)$, the number of triangles of T_H is not less than that of $T_{G[M]}$. As a result, we have $F_{itp}(G) \leq F_{itp}(G')$.

Finally, consider the independent cycle packing numbers of G and G' . Let P represent a maximum independent cycle packing of G . Let Z be all vertices of P . Clearly, each vertex of Z has exact two adjacent vertices in Z . If $M' = \emptyset$, we have $F_{icp}(G) \leq F_{icp}(G')$ since $F_{icp}(G[M]) = F_{icp}(H)$. Suppose $M' \neq \emptyset$ henceforth. If the intersection of Z and M is the empty set, then $F_{icp}(G) = F_{icp}(G[V \setminus M]) \leq F_{icp}(G')$. If the intersection of Z and M contains exact one vertex u . Assume C_1 is the cycle of P that contains vertex u . There exist two vertices $v, w \in M'$ such that C_1 contains v and w . Moreover, the intersection of Z and M' only contains two vertices v and w , otherwise there are more than two vertices in $Z \cap M'$ that are adjacent to u . Therefore, for any vertex x in H , the intersection of $N(x)$ and Z equals $\{v, w\}$. Suppose graph C_2 equals $\{x, (x, v), (x, w)\} \cup (C_1 \setminus \{u, (u, v), (u, w)\})$. Obviously, C_2 is a cycle of G' and $C_2 \cup (P \setminus C_1)$ is an independent cycle packing of G' . Thus, we have $F_{icp}(G) \leq F_{icp}(G')$. If the intersection of Z and M contains exact two vertices u and v . If vertices u, v belong to different cycles of P , then the intersection of Z and M' includes at least four vertices. This means that there are at least four vertices in $Z \cap M'$ that are adjacent to u (or v). If u, v are in the same cycle C of P , then there is only one vertex $w \in M' \cap Z$, otherwise u (or v) has more than two neighbor vertices in $N[M] \cap Z$. Thus, cycle C is the triangle (u, v, w) . Additionally, since

$F_{im}(G[M]) = F_{im}(H)$ and $G[M]$ contains an edge (u, v) , there is at least one edge (x, y) in H . Thus, $(x, y, w) \cup (P \setminus C)$ is an independent cycle packing of G' , and $F_{icp}(G) \leq F_{icp}(G')$. If the intersection of Z and M contains more than two vertices, then the intersection of Z and M' is the empty set, otherwise there are more than two vertices in $Z \cap M$ that are adjacent to a vertex in $Z \cap M'$. Thus, for every cycle C of P , all vertices of C are either in M or in $V \setminus N[M]$. Suppose $P_{G[M]}$ is the intersection of P and $G[M]$. Assume P_H is a maximum independent cycle packing of H . Then, $P_H \cup (P \setminus P_{G[M]})$ is an independent cycle packing of G' . In addition, since $F_{icp}(G[M]) = F_{icp}(H)$, the number of cycle of P_H is not less than that of $P_{G[M]}$. As a result, we have $F_{icp}(G) \leq F_{icp}(G')$. \square

6 Optimal kernels

In this section, we provide optimal kernels for STEINER TREE(mw), CONNECTED DOMINATING SET(mw), and MAX LEAF SPANNING TREE(mw) under the assumption that $P \neq NP$.

For a Steiner tree of a graph G , the size of the Steiner tree is the number of its edges, and the non-terminal vertices of the Steiner tree are called Steiner vertices. If G contains at least two connected components each of which includes a terminal, then the Steiner tree of G does not exist and we define the size of the Steiner tree of G is $|V(G)|$.

Lemma 15. STEINER TREE(mw) has a kernel with mw vertices and does not admit a kernel with $\text{mw} - 1$ vertices unless $P = NP$.

Proof. Let (G, K, k) be an input of STEINER TREE. Consider first that $G = (V, E)$ is not connected. If G has at least two connected components each of which includes a terminal of K , then return yes if $k \geq |V|$ and no otherwise. If G has only one connected component C that contains the terminals of K , then delete all vertices of G that are not in C as these vertices cannot be used for the Steiner tree. Without loss of generality, assume henceforth that G is connected. Obtain the modular decomposition tree $MD(G)$ in linear time. Then, the root vertex v_V of $MD(G)$ must be prime or series. Let $P = \{M \mid M \subseteq V\}$ be the maximal modular partition of V . Then every child v_M of v_V is corresponding to a module M of P and the quotient graph of v_V is $G_{/P}$. If $G[K]$ is connected, then the instance (G, K, k) is yes if $|K| \leq k + 1$ and no otherwise; assume henceforth that $G[K]$ is not connected. If all terminals of K are included in a module M of P . Since $G[K]$ is not connected, at least one Steiner vertex is needed to connect them. In addition, since v_V is prime or series, M has at least one adjacent module. Picking any vertex v of an adjacent module of M is sufficient to connect all vertices in K via edges to v . Thus, the size of the Steiner tree of G is $|K|$. This means that the instance (G, K, k) is yes if $|K| \leq k$ and no otherwise; assume henceforth that the terminals of K are distributed in more than one module of P . If v_V is series. As the terminals of K are distributed in more than one module of P , $G[K]$ is connected. It contradicts that $G[K]$ is not connected. Therefore, v_V must be prime and the vertex number of $G_{/P}$ is at most mw.

Suppose that terminal set K' contains all vertices v_M of $G_{/P}$ such that M contains at least one terminal of K . We have $|K'| \geq 2$ since the terminals of K are distributed in more than one module of P . We claim that (G, K, k) is a yes instance of STEINER TREE(mw) if and only if $(G_{/P}, K', k')$ is a yes instance of STEINER TREE(mw), where $k' = k - (|K| - |K'|)$.

For one direction, assume that (G, K, k) is a yes instance and that T is a Steiner tree of G of size at most k . Assume that X' includes all vertices v_M of $G_{/P}$ such that M contains at least one vertex of $V(T)$. K' is a subset of X' as K is a subset of $V(T)$. Moreover, based on the definition of X' and K' , vertex set $X' \setminus K'$ includes all vertices v_M of $G_{/P}$ such that M contains at least one vertex of $V(T)$ but contains no vertices of K . Thus, the cardinality of $X' \setminus K'$ is not more than $|V(T)| - |K|$. Hence, $|X'|$ is not greater than $|V(T)| - |K| + |K'|$.

Let H denote the subgraph induced by X' in $G_{/P}$. Then, H is connected as T is connected. Thus, we can obtain a Steiner tree of $G_{/P}$ by extracting a spanning tree of H . Moreover, the size of the Steiner tree of $G_{/P}$ is $|X'| - 1$, which is not greater than $|V(T)| - (|K| - |K'|) - 1$. In addition, $|V(T)| - 1 \leq k$ since (G, K, k) is a yes instance. As a result, the size of the Steiner tree of $G_{/P}$ is not greater than $k - (|K| - |K'|)$ and $(G_{/P}, K', k')$ is a yes instance of STEINER TREE(mw).

For another direction, assume that $(G_{/P}, K', k')$ is a yes instance and that T' is a Steiner tree of $G_{/P}$ of size at most k' . Next, we will construct a Steiner tree T of size at most $k' + (|K| - |K'|)$ for G . At the first step, for each v_M of $G_{/P}$, if v_M is a Steiner vertex of T' , then add a vertex of M into T ; if v_M is a terminal of T' , then add a vertex of $M \cap K$ into T . So far, every module M corresponding to a vertex v_M of T' contains exact one vertex of T . Moreover, these vertices could generate a tree T in G , which is the same as T' . At the second step, add each of the remaining vertices of K into T as leaf by adhering it to the vertex of T in its neighbor module. (The neighbor module must exist since $|K'| \geq 2$ and, in the graph $G_{/P}$, each vertex of K' is adjacent to at least one vertex of T' .) Thus, we obtain a new tree T that is a subgraph of G and contains all the vertices of K . Moreover, $|V(T')|$ vertices are added into T at the first step and $|K| - |K'|$ vertices are added into T at the second step. Additionally, $|V(T')| - 1 \leq k'$ since $(G_{/P}, K', k')$ is a yes instance. Thus, the vertex number of T is at most $k' + (|K| - |K'|) + 1$. Consequently, the size of the Steiner tree T is at most $k = k' + (|K| - |K'|)$, and (G, K, k) is a yes instance of STEINER TREE(mw).

Now, consider the lower bound for the size of the kernel. For any graph G , the modular-width of G is not greater than the vertex number of G . Therefore, if STEINER TREE(mw) has a kernel with $\text{mw} - 1$ vertices, then we can solve STEINER TREE(mw) in polynomial time by repeatedly using the kernelization at most mw times. \square

The connected domination number of $G = (V, E)$ is the number of vertices of a minimum connected dominating set of G . If G is not connected, then we define the connected domination number of G is $|V| + 1$.

Lemma 16. CONNECTED DOMINATING SET(mw) has a kernel with mw vertices and does not admit a kernel with $\text{mw} - 1$ vertices unless $P = NP$.

Proof. Let (G, k) be an input of CONNECTED DOMINATING SET(mw). If $G = (V, E)$ is not connected. Return yes if $k \geq |V| + 1$ and no otherwise. Assume G is connected henceforth. Without loss of generality, suppose the connected domination number of G is not less than three. (All instances with connected domination number less than three can be decided by a polynomial-time subroutine.) Thus, return no if $k \leq 2$. Suppose henceforth that $k \geq 3$. Then, obtain the modular decomposition tree $MD(G)$ in linear time. Let $P = \{M \mid M \subseteq V\}$ be the maximal modular partition of V . Then every child v_M of the root vertex v_V of $MD(G)$ is corresponding to a module M of P and the quotient graph of v_V is $G_{/P}$. v_V cannot be parallel as G is connected. If v_V is series, then any two vertices from different modules of P are a connected dominating set of G . It contradicts that the connected domination number of G is not less than three. Thus, the root vertex v_V must be prime and the vertex number of $G_{/P}$ is not greater than mw .

We claim that the intersection of a module M of P and a minimum connected dominating set of G is either \emptyset or $\{v\}$, where v is a vertex of M . Let M' indicate $N(M)$. Since M is a maximal strong module of G , we have $M \subsetneq V$. Furthermore, $M' \neq \emptyset$ as G is connected. Suppose D is a minimum connected dominating set of G . If the intersection of D and M contains more than two vertices. Assume u is a vertex of M and v is a vertex of M' . Then, $\{u, v\} \cup (D \setminus M)$ is a dominating set of G since $N[M]$ is a subset of $N[u] \cup N[v]$. In addition, for any vertex w of

$D \setminus M$, if w is adjacent to a vertex of M , then w is adjacent to u . Hence, the subgraph induced by $\{u\} \cup (D \setminus M)$ is connected. Moreover, u is adjacent to v . As a result, $\{u, v\} \cup (D \setminus M)$ is a connected dominating set of G . It contradicts that D is a minimum connected dominating set of G . If the intersection of D and M contains exact two vertices u and v . Since $|D| \geq 3$, the intersection of D and M' contains at least one vertex x . Since $N[u]$ is a subset of $N[v] \cup N[x]$, $D \setminus \{u\}$ is a dominating set of G . In addition, for any vertex w of $D \setminus M$, if w is adjacent to u , then w is adjacent to v . Thus, the subgraph induced by $D \setminus \{u\}$ is connected. As a result, $D \setminus \{u\}$ is a connected dominating set of G . It contradicts that D is a minimum connected dominating set of G .

Now, we claim that (G, k) is a yes instance of CONNECTED DOMINATING SET(mw) if and only if $(G_{/P}, k)$ is a yes instance of CONNECTED DOMINATING SET(mw).

For one direction, assume that (G, k) is a yes instance and that D is a minimum connected dominating set of G . Then, we have $|D| \leq k$. Suppose $D' \subseteq V(G_{/P})$ includes all vertices v_M such that the intersection of M and D contains exact one vertex. Since any module M contains at most one vertex of D , there are $|D|$ modules of P each of which contains a vertex of D . Hence, $|D'| = |D| \leq k$. Additionally, for any two vertices u, v of D , if (u, v) is an edge of $G[D]$, then the module M including u is adjacent to the module M' including v , moreover, vertices v_M and $v_{M'}$, which are vertices of D' , are adjacent in $G_{/P}$. Thus, the subgraph induced in $G_{/P}$ by D' is also connected. Clearly, for any module M' of P , if the intersection of M' and D is \emptyset , then M' has at least one adjacent module M such that $M \cap D$ contains exact one vertex, otherwise every vertex of M' is neither in D nor adjacent to any member of D . Thus, for any vertex of $G_{/P}$, it is adjacent to a vertex of D' if it is not in D' . As a result, D' is a connected dominating set of $G_{/P}$ and $(G_{/P}, k)$ is a yes instance of CONNECTED DOMINATING SET(mw).

For another direction, assume that $(G_{/P}, k)$ is a yes instance. It is easy to check that the smallest prime graph is the path on four vertices. Since $G_{/P}$ is prime, it contains at least four vertices. Additionally, we have $k \geq 3$. Thus, there exists a connected dominating set D of $G_{/P}$ such that $3 \leq |D| \leq k$. For every vertex v_M of $G_{/P}$, if v_M is in D , then add a vertex of M into vertex set D' . For each module M , if the intersection of M and D' contains a vertex, then add M to P' . Obviously, the cardinality of D' equals that of D and $3 \leq |D'| \leq k$. For any two vertices v_M and $v_{M'}$ of D , if v_M and $v_{M'}$ are adjacent in $G_{/P}$, then modules M and M' are adjacent, moreover, the vertex in $D' \cap M$ and the vertex in $D' \cap M'$ are adjacent in G . Thus, the subgraph induced in G by D' is also connected. Consider a module M' of $P \setminus P'$. Since D is a dominating set of $G_{/P}$, M' has at least one adjacent module M such that v_M is in D . This means that $D' \cap M$ contains a vertex u and all vertices of M' are adjacent to u . Consider a module M' of P' . Since $|D| \geq 3$ and the subgraph induced in $G_{/P}$ by D is connected, M' has at least one adjacent module M such that v_M is in D . This means that $D' \cap M$ contains a vertex u and all vertices of M' are adjacent to u . Therefore, any vertex of G is adjacent to a vertex of D' . As a result, D' is a connected dominating set of G and (G, k) is a yes instance of CONNECTED DOMINATING SET(mw).

Next, consider the lower bound for the size of the kernel. For any graph G , the modular-width of G is not greater than the vertex number of G . Therefore, if CONNECTED DOMINATING SET(mw) has a kernel with $\text{mw} - 1$ vertices, then we can solve CONNECTED DOMINATING SET(mw) in polynomial time by repeatedly using the kernelization at most mw times. \square

For any graph G the connected domination number of G plus the number of leaves in the maximum leaf spanning tree of G equals the vertex number of G [10]. Therefore, the following corollary holds.

Corollary 6. MAX LEAF SPANNING TREE(mw) has a kernel with mw vertices and does not admit a kernel with $\text{mw} - 1$ vertices unless $P = NP$.

7 Conclusions

In this paper we try to explore more examples for polynomial Turing compression. We find that when we solve a graph problem by using the modular decomposition tree of the graph, we need to investigate the quotient graphs of the vertices of the tree repeatedly. This mode is similar with the way that Turing compression solves problems. Based on this idea, polynomial Turing compressions are demonstrated for several graph problems parameterized by modular-width. Moreover, we construct a recipe to facilitate the process of obtaining these results. Additionally, we find conditional tight kernels for a few graph problems parameterized by modular-width, and new parameterized algorithms for these problems are obtained.

As a direction for further research, it would be interested to investigate if more problems have polynomial Turing compressions parameterized by modular-width such as k -PATH(mw). In addition, Fomin et al. [13] gives a meta-theorem proved a family of problems are FPT parameterized by modular-width. Can we also give a meta-theorem to prove that the family of problems have polynomial Turing compressions?

Acknowledgements

This work is supported by CSC 201708430114 fellowship.

References

- [1] A. M. Ambalath, R. Balasundaram, C. R. H., V. Koppula, N. Misra, G. Philip, and M. S. Ramanujan. On the kernelization complexity of colorful motifs. In *Parameterized and Exact Computation - 5th International Symposium, IPEC 2010, Chennai, India, December 13-15, 2010. Proceedings*, pages 14–25, 2010.
- [2] B. Bergougnoux and M. M. Kanté. Fast exact algorithms for some connectivity problems parameterized by clique-width. *Theor. Comput. Sci.*, 782:30–53, 2019.
- [3] D. Binkele-Raible, H. Fernau, F. V. Fomin, D. Lokshtanov, S. Saurabh, and Y. Villanger. Kernel(s) for problems with no kernel: On out-trees with many leaves. *ACM Trans. Algorithms*, 8(4):38:1–38:19, 2012.
- [4] H. L. Bodlaender, B. M. P. Jansen, and S. Kratsch. Kernelization lower bounds by cross-composition. *SIAM J. Discrete Math.*, 28(1):277–305, 2014.
- [5] H. L. Bodlaender, E. J. van Leeuwen, J. M. M. van Rooij, and M. Vatshelle. Faster algorithms on branch and clique decompositions. In P. Hlinený and A. Kucera, editors, *Mathematical Foundations of Computer Science 2010, 35th International Symposium, MFCS 2010, Brno, Czech Republic, August 23-27, 2010. Proceedings*, volume 6281 of *Lecture Notes in Computer Science*, pages 174–185. Springer, 2010.
- [6] B. Courcelle, J. A. Makowsky, and U. Rotics. Linear time solvable optimization problems on graphs of bounded clique-width. *Theory Comput. Syst.*, 33(2):125–150, 2000.
- [7] B. Courcelle and S. Olariu. Upper bounds to the clique width of graphs. *Discret. Appl. Math.*, 101(1-3):77–114, 2000.
- [8] M. Cygan, F. V. Fomin, L. Kowalik, D. Lokshtanov, D. Marx, M. Pilipczuk, M. Pilipczuk, and S. Saurabh. *Parameterized Algorithms*. Springer, 2015.

- [9] H. Donkers and B. M. P. Jansen. A Turing kernelization dichotomy for structural parameterizations of F-minor-free deletion. In *Graph-Theoretic Concepts in Computer Science - 45th International Workshop, WG 2019, Vall de Núria, Spain, June 19-21, 2019, Revised Papers*, pages 106–119, 2019.
- [10] R. J. Douglas. NP-completeness and degree restricted spanning trees. *Discrete Mathematics*, 105(1-3):41–47, aug 1992.
- [11] R. G. Downey and M. R. Fellows. *Fundamentals of Parameterized Complexity*. Texts in Computer Science. Springer, 2013.
- [12] F. V. Fomin, F. Grandoni, and D. Kratsch. Solving connected dominating set faster than 2^n . *Algorithmica*, 52(2):153–166, 2008.
- [13] F. V. Fomin, M. Liedloff, P. Montealegre, and I. Todinca. Algorithms parameterized by vertex cover and modular width, through potential maximal cliques. *Algorithmica*, 80(4):1146–1169, feb 2017.
- [14] F. V. Fomin, D. Lokshtanov, S. Saurabh, and M. Zehavi. *Kernelization: Theory of Parameterized Preprocessing*. Cambridge University Press, 2013.
- [15] J. Gajarský, M. Lampis, and S. Ordyniak. Parameterized algorithms for modular-width. In G. Z. Gutin and S. Szeider, editors, *Parameterized and Exact Computation - 8th International Symposium, IPEC 2013, Sophia Antipolis, France, September 4-6, 2013, Revised Selected Papers*, volume 8246 of *Lecture Notes in Computer Science*, pages 163–176. Springer, 2013.
- [16] T. Gallai. Transitiv orientierbare graphen. *Acta Mathematica Academiae Scientiarum Hungaricae*, 18(1-2):25–66, mar 1967.
- [17] M. Habib and C. Paul. A survey of the algorithmic aspects of modular decomposition. *comput. Sci. Rev.*, 4(1):41–59, 2010.
- [18] B. M. P. Jansen. Turing kernelization for finding long paths and cycles in restricted graph classes. *J. Comput. Syst. Sci.*, 85:18–37, 2017.
- [19] B. M. P. Jansen and D. Marx. Characterizing the easy-to-find subgraphs from the viewpoint of polynomial-time algorithms, kernels, and Turing kernels. In *Proceedings of the Twenty-Sixth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2015, San Diego, CA, USA, January 4-6, 2015*, pages 616–629, 2015.
- [20] B. M. P. Jansen, M. Pilipczuk, and M. Wrochna. Turing kernelization for finding long paths in graphs excluding a topological minor. In *12th International Symposium on Parameterized and Exact Computation, IPEC 2017, September 6-8, 2017, Vienna, Austria*, pages 23:1–23:13, 2017.
- [21] A. Schäfer, C. Komusiewicz, H. Moser, and R. Niedermeier. Parameterized computational complexity of finding small-diameter subgraphs. *Optimization Letters*, 6(5):883–891, 2012.
- [22] O. Suchý. Exact algorithms for steiner tree. *New Developments in Exact Algorithms and Lower Bounds, Pre-FSTTCS 2014 workshop, December 13-14, 2014, New Delhi, India*, 2014.

- [23] M. Tedder, D. Corneil, M. Habib, and C. Paul. Simpler linear-time modular decomposition via recursive factorizing permutations. In *Automata, Languages and Programming*, pages 634–645. Springer Berlin Heidelberg, 2008.
- [24] S. Thomassé, N. Trotignon, and K. Vuskovic. A polynomial Turing-kernel for weighted independent set in bull-free graphs. *Algorithmica*, 77(3):619–641, 2017.

8 Appendix A: an algorithm for odd cycle transversal

We provide an $\mathcal{O}^*(3^{\text{mw}})$ time algorithm for ODD CYCLE TRANSVERSAL(mw). Recall that, for a graph G , we can obtain the modular decomposition tree $MD(G)$ in linear time. Additionally, for every vertex v_M of $MD(G)$, we say $G[M]$ is the corresponding graph of v_M and M is the corresponding module of v_M .

Lemma 17. *Let M and O be a module and a minimum odd cycle transversal of $G = (V, E)$, respectively. The following statements hold:*

(1) *If the intersection of O and M is a minimum vertex cover of $G[M]$, then $O \setminus M$ together with any minimum vertex cover of $G[M]$ is a minimum odd cycle transversal of G .*

(2) *If the intersection of O and M is a minimum odd cycle transversal of $G[M]$, then $O \setminus M$ together with any minimum odd cycle transversal of $G[M]$ is a minimum odd cycle transversal of G .*

Proof. Consider statement (1). Let O' denote $O \setminus M$ together with a minimum vertex cover of $G[M]$. It is obvious that $V \setminus (O' \cup M)$ equals $V \setminus (O \cup M)$, and the cardinality of independent sets $M \setminus O$ and $M \setminus O'$ are equal. Thus, according to Lemma 9, the subgraph induced by $V \setminus O$ and the subgraph induced by $V \setminus O'$ are isomorphic. This means that the subgraph induced by $V \setminus O$ has an odd cycle if and only if the subgraph induced by $V \setminus O'$ has an odd cycle. Thus, O' is an odd cycle transversal of G . Additionally, the cardinality of O and O' are the same. Consequently, O' is a minimum odd cycle transversal of G .

Consider statement (2). Let O' denote $O \setminus M$ together with a minimum odd cycle transversal of $G[M]$. If $N(M) = \emptyset$. Obviously, O' is a minimum odd cycle transversal of G . Assume $N(M) \neq \emptyset$ henceforth. If $G[M]$ is a bipartite graph, then both $O \cap M$ and $O' \cap M$ are empty sets. Thus, O' equals O . If $G[M]$ is not a bipartite graph, then $G[M]$ has an edge. Based on Lemma 10, the remaining graph after deleting any minimum odd cycle transversal from $G[M]$ has an edge. Additionally, $N(M) \neq \emptyset$. Thus, $N(M) \setminus O$ is the empty set, otherwise there is a triangle in $G[V \setminus O]$. This means that there are no edges between vertex sets $M \setminus O'$ and $(V \setminus M) \setminus O'$. Moreover, both the subgraph induced by $M \setminus O'$ and the subgraph induced by $(V \setminus M) \setminus O'$ have no odd cycles. Therefore, O' is an odd cycle transversal of G . In addition, the cardinality of O and O' are equal. Consequently, O' is a minimum odd cycle transversal of G . \square

Lemma 18. ODD CYCLE TRANSVERSAL(mw) *has an $\mathcal{O}^*(3^{\text{mw}})$ algorithm.*

Proof. Let function $F_{\text{oct}}(G)$ and $F_{\text{vc}}(G)$ represent the odd cycle transversal number and the vertex cover number of a graph $G = (V, E)$, respectively. Since $F_{\text{vc}}(G)$ can be solved in $\mathcal{O}^*(1.74^{\text{mw}})$ time [13], we can obtain $F_{\text{vc}}(G[M])$ for all vertices v_M of $MD(G)$ in $\mathcal{O}^*(1.74^{\text{mw}})$ time. Next, we will compute $F_{\text{oct}}(G[M])$ for every vertex v_M of $MD(G)$ in a bottom-up fashion, and finally obtain $F_{\text{oct}}(G)$.

If v_M is a leaf of $MD(G)$, then $F_{\text{oct}}(G[M]) = 0$. Consider an internal vertex v_M of $MD(G)$. Let $P = \{M' \mid M' \subseteq M\}$ be the maximal modular partition of M . Then every child $v_{M'}$ of v_M is corresponding to a module M' in P and the quotient graph of v_M is $G[M]_{/P}$. If v_M is a parallel

vertex, then $F_{oct}(G[M])$ is the sum of $F_{oct}(G[M'])$ for all $G[M']$. If v_M is a series vertex, then, for any minimum odd cycle transversal $O_{G[M]}$ of $G[M]$, the vertices of $M \setminus O_{G[M]}$ are distributed in at most two modules of P , otherwise the subgraph in $G[M]$ induced by $M \setminus O_{G[M]}$ contains a triangle. Additionally, based on Lemma 11, the intersection of M' and $O_{G[M]}$ is either M' , a minimum vertex cover of $G[M']$, or a minimum odd cycle transversal of $G[M']$. Thus, we consider the following two cases. If all the vertices of $M \setminus O_{G[M]}$ are distributed in only one module of P , then we have

$$F_{oct}(G[M]) = |M| - \max_{M' \in P} \{|M'| - F_{oct}(G[M'])\}$$

If all the vertices of $M \setminus O_{G[M]}$ are distributed in exact two modules of P , then we have

$$F_{oct}(G[M]) = |M| - \max_{M', M'' \in P} \{|M'| + |M''| - F_{vc}(G[M']) - F_{vc}(G[M''])\}$$

Clearly, $F_{oct}(G[M])$ is the smaller one of the two cases.

From now on, we consider that v_M is a prime vertex. Suppose v_M has r children. Obviously, $r \leq \text{mw}$. Assume \mathbb{S} includes all vertex sets $S \subseteq M$ such that, for every $M' \in P$, the intersection of S and M' is either M' , a minimum vertex cover of $G[M']$, or a minimum odd cycle transversal of $G[M']$. For a vertex set $S \in \mathbb{S}$ and a module $M' \in P$, we say the pattern of $S \cap M'$ is pattern 0 if $S \cap M'$ is M' ; we say the pattern of $S \cap M'$ is pattern 1 if $S \cap M'$ is a minimum vertex cover of $G[M']$; we say the pattern of $S \cap M'$ is pattern 2 if $S \cap M'$ is a minimum odd cycle transversal of $G[M']$. In addition, we say $S \cap M'$ has the pattern x if the pattern of $S \cap M'$ is pattern x , where $x \in \{0, 1, 2\}$. Suppose $P = \{M_1, \dots, M_r\}$ and $i \in \{1, \dots, r\}$. Let $\overline{t_1 \dots t_r}$ be a ternary number with r trits, where $t_i \in \{0, 1, 2\}$ for all i . Clearly, ternary number $\overline{t_1 \dots t_r}$ can denote any integer between 0 and $3^r - 1$. Suppose $\mathbb{T}_{\overline{t_1 \dots t_r}}$ includes all $S \in \mathbb{S}$ such that the pattern of $S \cap M_i$ is pattern t_i for every i . For example, $\mathbb{T}_{\overline{2 \dots 2}}$ includes all S such that $S \cap M_i$ is a minimum odd cycle transversal of $G[M_i]$ for every i . Clearly, we have $\mathbb{S} = \mathbb{T}_0 \cup \dots \cup \mathbb{T}_{3^r - 1}$. Moreover, according to Lemma 11 and the definition of \mathbb{S} , every minimum odd cycle transversal of $G[M]$ is an element of \mathbb{S} . Therefore, there exists a j such that \mathbb{T}_j includes a minimum odd cycle transversal of $G[M]$, where $j \in \{0, \dots, 3^r - 1\}$.

Consider \mathbb{T}_j for each j . We claim that, for any two elements $S_0, S \in \mathbb{T}_j$, S_0 is a minimum odd cycle transversal of $G[M]$ if and only if S is a minimum odd cycle transversal of $G[M]$. Without loss of generality, it is enough to prove one direction to prove the claim. Assume S_0 is a minimum odd cycle transversal of $G[M]$. Suppose $S_i = (S_{i-1} \setminus M_i) \cup (S \cap M_i)$ for all i . Now, consider the relationship between S_{i-1} and S_i for each i . Both S_0 and S are elements of \mathbb{T}_j , so $S_{i-1} \cap M_i$, which equals $S_0 \cap M_i$, and $S \cap M_i$ have the same pattern. According to Lemma 17, that S_{i-1} is a minimum odd cycle transversal of $G[M]$ implies that S_i is a minimum odd cycle transversal of $G[M]$. Thus, each element of $\{S_0, S_1, \dots, S_r\}$ is a minimum odd cycle transversal of $G[M]$. Additionally, it is clear that S equals S_r . Consequently, S is a minimum odd cycle transversal of $G[M]$.

Now, we know that there exists a j such that every element of \mathbb{T}_j is a minimum odd cycle transversal of $G[M]$. Since the quotient graph $G[M]_P$ as well as $F_{vc}(G[M_i])$ and $F_{oct}(G[M_i])$ for all $M_i \in P$ are given, we can construct an element of \mathbb{T}_j in polynomial time. For every \mathbb{T}_j , obtain an element S from \mathbb{T}_j and reserve S only if it is an odd cycle transversal of $G[M]$. Then, select a minimum one, which must be a minimum odd cycle transversal of $G[M]$, from all reserved S . Since $0 \leq j \leq 3^r - 1$, this process can be done in $\mathcal{O}^*(3^r)$ time. \square

9 Appendix B: problem zoo

We provide problem statements to all problems discussed in this paper below.

CHROMATIC NUMBER

Input: A graph G and an integer k .

Question: Does there exist at most k colors to color the vertices of G such that no two adjacent vertices share the same color?

CLIQUE

Input: A graph G and an integer k .

Question: Does there exist a set X of at least k vertices of G such that any two vertices in X are adjacent?

CONNECTED DOMINATING SET

Input: A graph G and an integer k .

Question: Does there exist a set X of at most k vertices of G such that $G[X]$ is connected and every vertex not in X is adjacent to at least one member of X ?

CONNECTED VERTEX COVER

Input: A graph G and an integer k .

Question: Does there exist a set X of at most k vertices of G such that $G[X]$ is connected and every edge of G has at least one end point in X ?

DOMINATING SET

Input: A graph G and an integer k .

Question: Does there exist a set X of at most k vertices of G such that every vertex not in X is adjacent to at least one member of X ?

HAMILTONIAN CYCLE

Input: A graph G .

Question: Does there exist a cycle of G that visits each vertex of G exactly once?

HAMILTONIAN PATH

Input: A graph G .

Question: Does there exist a path of G that visits each vertex of G exactly once?

FEEDBACK VERTEX SET

Input: A graph G and an integer k .

Question: Does there exist a set X of at most k vertices of G such that $G - X$ is a forest?

INDEPENDENT CYCLE PACKING

Input: A graph G and an integer k .

Question: Does G contain an induced subgraph consisting of k pairwise vertex-disjoint cycles?

INDEPENDENT SET

Input: A graph G and an integer k .

Question: Does there exist a set X of at least k vertices of G such that any two vertices in X have no edge?

INDEPENDENT TRIANGLE PACKING

Input: A graph G and an integer k .

Question: Does there exist a set X of $3k$ vertices of G such that the subgraph induced by X is k pairwise vertex-disjoint triangles?

INDUCED MATCHING

Input: A graph G and an integer k .

Question: Does there exist a set X of $2k$ vertices of G such that the subgraph induced by X is a matching consisting of k edges?

LONGEST INDUCED PATH

Input: A graph G and an integer k .

Question: Does G contain the path on k vertices as an induced subgraph?

MAX INDUCED FOREST

Input: A graph G and an integer k .

Question: Does there exist a set X of k vertices of G such that the subgraph induced by X has no cycles?

MAX LEAF SPANNING TREE

Input: A graph G and an integer k .

Question: Does there exist a spanning tree of G with at least k leaves?

NONBLOCKER

Input: A graph G and an integer k .

Question: Does there exist a set X of at least k vertices of G such that every vertex in X is adjacent to a vertex outside of X ?

ODD CYCLE TRANSVERSAL

Input: A graph G and an integer k .

Question: Does there exist a set X of at most k vertices of G such that $G - X$ is a bipartite graph?

PARTITIONING INTO PATHS

Input: A graph G and an integer k .

Question: Does G contain k vertex disjoint paths whose union includes every vertex of G ?

STEINER TREE

Input: A graph G , a set $K \subseteq V(G)$, and an integer k .

Question: Does there exist a connected subgraph of G that contains at most k edges and contains all vertices of K ?

VERTEX COVER

Input: A graph G and an integer k .

Question: Does there exist a set X of at most k vertices of G such that every edge of G has at least one end point in X ?

WEIGHTED CLIQUE

Input: A graph G with vertex weight $w : V(G) \rightarrow \mathbb{Z}^+$ and an integer k .

Question: Does there exist a clique of G such that the total weight of the vertices of the clique is at least k ?