

Chambolle-Pock's primal-dual method with mismatched adjoint

Dirk A. Lorenz & Felix Schneppe*

January 14, 2022

Abstract

The primal-dual method of Chambolle and Pock is a widely used algorithm to solve various optimization problems written as convex-concave saddle point problems. Each update step involves the application of both the forward linear operator and its adjoint. However, in practical applications like computerized tomography, it is often computationally favourable to replace the adjoint operator by a computationally more efficient approximation. This leads to an adjoint mismatch in the algorithm.

In this paper, we analyze the convergence of Chambolle-Pock's primal-dual method under the presence of a mismatched adjoint. We present an upper bound on the error of the primal solution and derive step-sizes and mild conditions under which convergence to a fixed point is still guaranteed. Furthermore we present convergence rates similar to these of Chambolle-Pock's primal-dual method without the adjoint mismatch. Moreover, we illustrate our results both for an academic and a real-world application.

1 Introduction

Inverse problems occur whenever unknown quantities are measured indirectly. In many cases this process can be error-prone. Nevertheless, these inverse problems appear in many practical applications and are often attacked by solving minimization problems, which can be formulated as the minimization of an expression

$$\min_x F(Ax) + G(x). \quad (1)$$

Sometimes these kind of problems are hard to solve and it can be beneficial to examine the equivalent dual

$$\min_y G^*(-A^*y) + F^*(y) \quad (2)$$

*Dirk A. Lorenz, d.lorenz@tu-braunschweig.de, , Institute of Analysis and Algebra, TU Braunschweig, 38092 Braunschweig, Felix Schneppe, f.schneppe@tu-braunschweig.de, Institute of Analysis and Algebra, TU Braunschweig, 38092 Braunschweig

or the saddle point problem

$$\min_x \max_y G(x) + \langle Ax, y \rangle - F^*(y).$$

with the Fenchel conjugates G^*, F^* of G, F instead of the primal problem above.

If both G and F^* are proper, convex, lower semicontinuous functionals, the primal-dual algorithm

$$\begin{aligned} x^{i+1} &= \text{prox}_{\tau_i G}(x^i - \tau_i A^* y^i), \\ \bar{x}^{i+1} &= x^{i+1} + \omega_i(x^{i+1} - x^i), \\ y^{i+1} &= \text{prox}_{\sigma_{i+1} F^*}(y^i + \sigma_{i+1} A \bar{x}^{i+1}). \end{aligned} \tag{3}$$

of Chambolle and Pock [4] with stepsizes τ_i and σ_i and extrapolation parameter ω_i has proven to be a simple and effective solution method.

In practical applications, however, it can happen that the operator is given as a discretization of a continuous operator, while the backprojection operator is likewise only available as an often more coarsely discretized version of the adjoint operator. This discrepancy arises not only due to the limitations of the computers, but also to enable shorter computing time on the one hand and due to the physical importance of the operators on the other [2, 12, 16]. The influence of such a mismatch has been studied on various algorithms. [13, 5, 16, 9, 7, 8]

In this paper we examine the convergence of the Chambolle-Pock method in the case of a mismatched adjoint, i.e., we examine the algorithm

$$\begin{aligned} x^{i+1} &= \text{prox}_{\tau_i G}(x^i - \tau_i V^* y^i), \\ \bar{x}^{i+1} &= x^{i+1} + \omega_i(x^{i+1} - x^i), \\ y^{i+1} &= \text{prox}_{\sigma_{i+1} F^*}(y^i + \sigma_{i+1} A \bar{x}^{i+1}). \end{aligned} \tag{4}$$

with $V \neq A$ for convergence to a fixed point.

Example 1.1 (Counterexamples for convergence). Here are two simple examples that show that the mismatched iteration does not necessarily converge. In the first example, we consider the problem $\min_x \|x\|_1$ on \mathbb{R}^n and model this with $A = I$, $F(y) = \|y\|_1$, and $G \equiv 0$. We consider the most basic form of Chambolle-Pock's method with constant τ, σ and $\omega = 1$, i.e. the mismatched iteration is

$$\begin{aligned} x^{i+1} &= x^i - \tau V^T y^i \\ y^{i+1} &= \text{proj}_{[-1,1]}(y^i + \sigma A(2x^{i+1} - x^i)). \end{aligned}$$

If we consider the mismatch $V = -\alpha I$ with $\alpha > 0$ (instead of I), the iteration becomes

$$\begin{aligned} x^{i+1} &= x^i + \alpha \tau y^i \\ y^{i+1} &= \text{proj}_{[-1,1]}(y^i + \sigma(2x^{i+1} - x^i)). \end{aligned}$$

If we initialize with $x^0 > 0$ and $y^0 > 0$ (component-wise), we get that the entries in x^i are strictly increasing and hence, will not converge to the unique solution $x = 0$.

Another example where one observes non-convergence is the following. Consider the linear saddle point problem

$$\min_x \max_y \langle Ax, y \rangle$$

with $A \in \mathbb{R}^{m \times n}$, $x \in \mathbb{R}^n$, and $y \in \mathbb{R}^m$. We model with $G \equiv 0$ and $F^* \equiv 0$ and hence, the mismatched iteration reads as

$$\begin{aligned} x^{i+1} &= x^i - \tau V^T y^i \\ y^{i+1} &= y^i + \sigma A(2x^{i+1} - x^i). \end{aligned}$$

One can write this iteration in block form as

$$\begin{pmatrix} x^{i+1} \\ y^{i+1} \end{pmatrix} = \begin{pmatrix} I & 0 \\ -2\sigma A & I \end{pmatrix}^{-1} \begin{pmatrix} I & -\tau V^T \\ -\sigma A & I \end{pmatrix} \begin{pmatrix} x^i \\ y^i \end{pmatrix} =: M \begin{pmatrix} x^i \\ y^i \end{pmatrix}.$$

It is simple to create matrices A and small perturbations V of A such that the iteration matrix M has spectral radius larger than 1 (one can take a matrix A with uniformly distributed entries in $[0, 1]$ and increase a small number of entries by one to create V , for example).

Note that in both examples saddle points exist and are still fixed points of the mismatched iterations.

Our first result states that fixed points of the iteration (4) with mismatch are not too far away from fixed points of the original iteration (3) when the functional G is strongly convex.

Theorem 1.2. *If G is a γ_G -strongly convex function, (x^*, y^*) is the fixed point of the original Chambolle-Pock method (3) and (\hat{x}, \hat{y}) as the fixed point of the Chambolle-Pock method with mismatched adjoint (4), the difference between the primal solutions is bounded by*

$$\|x^* - \hat{x}\| \leq \frac{1}{\gamma_G} \|(V - A)^* \hat{y}\|.$$

Proof. Since ∂G and ∂F^* are monotone, we can conclude for $-A^* y^* \in \partial G(x^*)$, $-V^* \hat{y} \in \partial G(\hat{x})$, $Ax^* \in \partial F^*(y^*)$, $A\hat{x} \in \partial F^*(\hat{y})$ that

$$\langle x^* - \hat{x}, -A^* y^* + V^* \hat{y} \rangle \geq 0$$

and

$$\langle x^* - \hat{x}, A^*(y^* - \hat{y}) \rangle = \langle A(x^* - \hat{x}), y^* - \hat{y} \rangle \geq 0.$$

These sum up to

$$\langle x^* - \hat{x}, (V - A)^* \hat{y} \rangle \geq 0.$$

But since G is strongly convex with parameter γ_G , we are able to state

$$\langle x^* - \hat{x}, (V - A)^* \hat{y} \rangle \geq \gamma_G \|x^* - \hat{x}\|^2.$$

Furthermore, it is

$$\gamma_G \|x^* - \hat{x}\|^2 \leq \langle x^* - \hat{x}, (V - A)^* \hat{y} \rangle \leq \|x^* - \hat{x}\| \|(V - A)^* \hat{y}\|,$$

which shows

$$\gamma_G \|x^* - \hat{x}\| \leq \|(V - A)^* \hat{y}\|.$$

□

Notably, we can not show that the mismatched algorithm will converge to the original solution (a situation which possible for other mismatched iterations [9, 5]). However, since we can bound the difference of fixed points by the mismatch, we conclude that the analysis of the Chambolle-Pock method with mismatched adjoint can be of interest in practical applications.

The rest of the paper is structured in the following way. In the Section 2 we reformulate (4) and present easily verifiable properties, which allow convergences. Following to this section, we prove linear convergence, if both G and F^* are proper, convex, lower-semicontinuous functionals and provide step lengths, which allow linear convergence and are easy to compute. In Section 3, we present numerical examples and Section 4 concludes the paper.

Throughout this paper, we will use $\langle x, x' \rangle_T := \langle Tx, x' \rangle$ and the norm-like notation $\|x\|_T^2 := \langle x, x \rangle_T$ for an (not necessarily symmetric) positive semidefinite operators T .¹ Additionally, we will write $A \geq 0$ for a set A , if every element a satisfies $a \geq 0$.

2 Convergence analysis

This section is all about showing convergence of the algorithm, if $\|V - A\|$ is small enough, and establishing convergence rates that hold under certain circumstances. We do not proof the existence of a fixed point, but take its existence as granted. Since the original proof of Chambolle and Pock in [4] relies on having the exact adjoint, we make use of the ideas made in [15] and [6] in our proof and rely on the following, slightly modified version of [6, theorem 2.1].

Theorem 2.1. *On a Hilbert Space \mathcal{U} , let $\tilde{H}_{i+1} : \mathcal{U} \rightrightarrows \mathcal{U}$ and $M_{i+1}, W_{i+1}, Z_{i+1} \in \mathcal{L}(\mathcal{U}, \mathcal{U})$ for $i \in \mathbb{N}$. Suppose that*

$$0 \in W_{i+1} \tilde{H}_{i+1}(u^{i+1}) + M_{i+1}(u^{i+1} - u^i) \quad (5)$$

is solvable for $\{u^{i+1}\}_{i \in \mathbb{N}} \subset \mathcal{U}$ and let $\hat{u} \in \mathcal{U}$ be a stationary point of the iteration.

If $Z_{i+1}M_{i+1}$ is self-adjoint and for some $\Delta_{i+1} \in \mathbb{R}$ the condition

$$\langle \tilde{H}_{i+1}(u^{i+1}), u^{i+1} - \hat{u} \rangle_{Z_{i+1}W_{i+1}} \quad (6)$$

$$\geq \frac{1}{2} \|u^{i+1} - \hat{u}\|_{Z_{i+2}M_{i+2} - Z_{i+1}M_{i+1}}^2 - \frac{1}{2} \|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 - \Delta_{i+1} \quad (7)$$

holds for all $i \in \mathbb{N}$, then so does the descent inequality

$$\frac{1}{2} \|u^N - \hat{u}\|_{Z_{N+1}M_{N+1}}^2 \leq \frac{1}{2} \|u^0 - \hat{u}\|_{Z_1M_1}^2 + \sum_{i=0}^{N-1} \Delta_{i+1}.$$

While the maps \tilde{H}_i , M_i , and W_i define the iteration (5), the operator Z_i and the number Δ_i are only used to establish inequality (6) and the final descent inequality. The operator Z_i is called *test operator* and the Δ_i can be used to further quantify the descent.

¹In the case that T is not positive semi-definite, the expression $\|x\|_T$ will not be used, however, its square $\|x\|_T^2$ will (but may be a negative number.)

For obvious reasons we aim for non-positive Δ_{i+1} and want that the operator $Z_{i+1}M_{i+1}$ grow as fast as possible to obtain fast convergence.

Consequently, we now show that

- the algorithm can be expressed like (5).
- with the right step length choices an operator Z_{i+1} with $Z_{i+1}M_{i+1}$ being self-adjoint exists.
- for some non-positive Δ_{i+1} the inequality (6) can be obtained.

2.1 Reformulation

In order to use Theorem 2.1 to prove convergence of the iteration (4), we reformulate iteration (4) in terms of (5). First, recall that

$$v = \text{prox}_f(x) \Leftrightarrow x - v \in \partial f(v).$$

Moreover, it is easy to see that a stationary point $\hat{u} = \begin{pmatrix} \hat{x} \\ \hat{y} \end{pmatrix}$ of iteration (4) fulfills $\hat{u} \in H^{-1}(0)$ with

$$H(u) = H(x, y) = \begin{pmatrix} \partial G(x) + V^*y \\ \partial F^*(y) - Ax \end{pmatrix}. \quad (8)$$

With $\bar{x}^{i+1} = (1 + \omega_i)x^{i+1} - \omega_i x^i$, we rewrite the update steps of the algorithm (4) as

$$0 \in \begin{pmatrix} x^{i+1} + \tau_i \partial G(x^{i+1}) - x^i + \tau_i V^* y^i \\ y^{i+1} + \sigma_{i+1} \partial F^*(y^{i+1}) - y^i - \sigma_{i+1} A \bar{x}^{i+1} \end{pmatrix}. \quad (9)$$

Hence, we define

$$\tilde{H}_{i+1}(u) = \begin{pmatrix} \partial G(x) + V^*y \\ \partial F^*(y) - A[(1 + \omega_i)x - \omega_i x^i] - \omega_i V(x^i - x) \end{pmatrix} \quad (10)$$

as well as the *preconditioner*

$$M_{i+1} = \begin{pmatrix} I & -\tau_i V^* \\ -\omega_i \sigma_{i+1} V & I \end{pmatrix} \quad (11)$$

and the *step length operator*

$$W_{i+1} = \begin{pmatrix} \tau_i I & 0 \\ 0 & \sigma_{i+1} I \end{pmatrix} \quad (12)$$

and obtain, that the iteration (9) can be written as a preconditioned proximal-point method with

$$0 \in W_{i+1} \tilde{H}_{i+1}(u^{i+1}) + M_{i+1}(u^{i+1} - u^i). \quad (13)$$

Hence, the convergence of algorithm (4) can be analyzed by means of Theorem 2.1.

2.2 Test operator and step length bounds

Before we proceed with checking the conditions of Theorem 2.1, we have to make a choice regarding the test operator Z_{i+1} and this will induce some first conditions on the step length τ_i and σ_i . We define

$$Z_{i+1} := \begin{pmatrix} \varphi_i I & 0 \\ 0 & \psi_{i+1} I \end{pmatrix} \quad (14)$$

for $i \in \mathbb{N}$ and show that this choice is applicable in Theorem 2.1 for appropriate φ_i and ψ_{i+1} . Since we need that $Z_{i+1}M_{i+1}$ is self adjoint and it holds that

$$Z_{i+1}M_{i+1} = \begin{pmatrix} \varphi_i I & -\tau_i \varphi_i V^* \\ -\omega_i \sigma_{i+1} \psi_{i+1} V & \psi_{i+1} I \end{pmatrix}$$

we require that the values φ_i, ψ_{i+1} of the test operator fulfill

$$\omega_i \sigma_{i+1} \psi_{i+1} = \tau_i \varphi_i.$$

We introduce the “tested dual stepsize”

$$\eta_i = \psi_i \sigma_i \quad (15)$$

and get that the extrapolation constant fulfills

$$\omega_i = \frac{\eta_i}{\eta_{i+1}}. \quad (16)$$

Consequently, the “tested primal stepsize” also fulfills

$$\eta_i = \eta_{i+1} \omega_i = \omega_i \sigma_{i+1} \psi_{i+1} = \tau_i \varphi_i.$$

Furthermore, Theorem 2.1 needs $Z_{i+1}M_{i+1}$ to be positive semidefinite.

Lemma 2.2 ([6], Lemma 3.4). *Let $i \in \mathbb{N}$ and suppose that conditions (15) and (16) hold. Then we have that $Z_{i+1}M_{i+1}$ is self-adjoint and satisfies*

$$Z_{i+1}M_{i+1} = \begin{pmatrix} \varphi_i I & -\eta_i V^* \\ -\eta_i V & \psi_{i+1} I \end{pmatrix} \geq \begin{pmatrix} \delta \varphi_i I & 0 \\ 0 & \psi_{i+1} I - \frac{\eta_i^2}{\varphi_i(1-\kappa)} VV^* \end{pmatrix} \quad (17)$$

for all $\delta \in [0, \kappa]$ with $\kappa \in (0, 1)$.

Proof. The first statement follows from the computation

$$\begin{aligned} Z_{i+1}M_{i+1} &= \begin{pmatrix} \varphi_i I & 0 \\ 0 & \psi_{i+1} I \end{pmatrix} \begin{pmatrix} I & -\tau_i V^* \\ -\omega_i \sigma_{i+1} V & I \end{pmatrix} \\ &= \begin{pmatrix} \varphi_i I & -\varphi_i \tau_i V^* \\ -\omega_i \psi_{i+1} \sigma_{i+1} V & \psi_{i+1} I \end{pmatrix} \\ &= \begin{pmatrix} \varphi_i I & -\eta_i V^* \\ -\eta_i V & \psi_{i+1} I \end{pmatrix}, \end{aligned}$$

where we used (15) and (16) in the last step. For the second claim we observe

$$\begin{aligned} M &:= Z_{i+1}M_{i+1} - \begin{pmatrix} \delta \varphi_i I & 0 \\ 0 & \psi_{i+1} I - \frac{\eta_i^2}{\varphi_i(1-\kappa)} VV^* \end{pmatrix} \\ &= \begin{pmatrix} (1-\delta)\varphi_i I & -\eta_i V^* \\ -\eta_i V & (-\eta_i V)((1-\kappa)\varphi_i)^{-1}(-\eta_i V)^* \end{pmatrix}. \end{aligned}$$

Since $1 > \kappa \geq \delta$ and $(1-\kappa)\varphi_i > 0$, we derive the positive semidefiniteness of M . \square

Hence, we assume

$$\psi_{i+1} \geq \frac{\eta_i^2}{\varphi_i(1-\kappa)} \|V\|^2 \quad (18)$$

to ensure that $Z_{i+1}M_{i+1}$ is positive semi-definite.

Next we investigate the operator $Z_{i+1}M_{i+1} - Z_{i+2}M_{i+2}$ and show that it is easy to evaluate the respective semi-norm.

Lemma 2.3 ([6, Lemma 3.5]). *Let $i \in \mathbb{N}$ and assume that conditions (15), (16) and (18) are fulfilled and define*

$$\Xi_{i+1} := 2 \begin{pmatrix} \tau_i \mu_G I & \tau_i V^* \\ -\sigma_{i+1} V & \sigma_{i+1} \mu_{F^*} I \end{pmatrix}$$

If the constants μ_G and μ_{F^*} are chosen such that

$$\varphi_{i+1} = \varphi_i(1 + 2\tau_i \mu_G), \quad (19)$$

$$\psi_{i+1} = \psi_i(1 + 2\sigma_i \mu_{F^*}), \quad (20)$$

then it holds that

$$\|u\|_{Z_{i+1}M_{i+1} - Z_{i+2}M_{i+2}}^2 = \|u\|_{Z_{i+1}\Xi_{i+1}}^2.$$

Proof. We use the expression for $Z_{i+1}M_{i+1}$ from (17) and the conditions (15), (19), (20) and (16) to get

$$\begin{aligned} & Z_{i+1}M_{i+1} + Z_{i+1}\Xi_{i+1} - Z_{i+2}M_{i+2} \\ &= \begin{pmatrix} \varphi_i I & -\eta_i V^* \\ -\eta_i V & \psi_{i+1} I \end{pmatrix} + \begin{pmatrix} \varphi_i I & 0 \\ 0 & \psi_{i+1} I \end{pmatrix} \begin{pmatrix} 2\tau_i \mu_G I & 2\tau_i V^* \\ -2\sigma_{i+1} V & 2\sigma_{i+1} \mu_{F^*} I \end{pmatrix} - \begin{pmatrix} \varphi_{i+1} I & -\eta_{i+1} V^* \\ -\eta_{i+1} V & \psi_{i+2} I \end{pmatrix} \\ &= \begin{pmatrix} \varphi_i I & -\eta_i V^* \\ -\eta_i V & \psi_{i+1} I \end{pmatrix} + \begin{pmatrix} 2\varphi_i \tau_i \mu_G I & 2\varphi_i \tau_i V^* \\ -2\sigma_{i+1} \psi_{i+1} V & 2\sigma_{i+1} \psi_{i+1} \mu_{F^*} I \end{pmatrix} - \begin{pmatrix} \varphi_{i+1} I & -\eta_{i+1} V^* \\ -\eta_{i+1} V & \psi_{i+2} I \end{pmatrix} \\ &= \begin{pmatrix} \varphi_{i+1} I & \eta_i V^* \\ (-\eta_i - 2\eta_{i+1}) V & \psi_{i+2} I \end{pmatrix} - \begin{pmatrix} \varphi_{i+1} I & -\eta_{i+1} V^* \\ -\eta_{i+1} V & \psi_{i+2} I \end{pmatrix} \\ &= \begin{pmatrix} 0 & (\eta_i + \eta_{i+1}) V^* \\ -(\eta_i + \eta_{i+1}) V & 0 \end{pmatrix}. \end{aligned}$$

This shows that $Z_{i+1}M_{i+1} + Z_{i+1}\Xi_{i+1} - Z_{i+2}M_{i+2}$ is skew-symmetric, and hence, it holds for all u that $\|u\|_{Z_{i+1}(M_{i+1} + \Xi_{i+1}) - Z_{i+2}M_{i+2}}^2 = 0$ from which the statement follows. \square

2.3 Technical estimates

This finally brings us to a position, in which we can estimate the term

$$\begin{aligned} D &:= \langle \tilde{H}_{i+1}(u^{i+1}), u^{i+1} - \hat{u} \rangle_{Z_{i+1}W_{i+1}} - \frac{1}{2} \|u^{i+1} - \hat{u}\|_{Z_{i+2}M_{i+2} - Z_{i+1}M_{i+1}}^2 \\ &= \langle \tilde{H}_{i+1}(u^{i+1}), u^{i+1} - \hat{u} \rangle_{Z_{i+1}W_{i+1}} - \frac{1}{2} \|u^{i+1} - \hat{u}\|_{Z_{i+2}\Xi_{i+1}}^2, \end{aligned}$$

where the operator \tilde{H} is given by (10), the preconditioner is given by (11) and the step length operator is given by (12). In order to do so, we assume that both F^* and G are strongly convex functionals and choose the step length appropriately, and arrive at the following estimate of D .

Theorem 2.4. *Suppose that the conditions (15), (16), (19), and (20) hold, that G, F^* are γ_G/γ_{F^*} -strongly convex, respectively, with*

$$\gamma_G \geq \frac{\epsilon}{2\omega_i} \|A - V\| + \mu_G \quad (21)$$

for some $\epsilon > 0$. Then it holds that

$$D \geq \eta_{i+1}(\gamma_{F^*} - \mu_{F^*} - \frac{1 + \omega_i}{2\epsilon} \|A - V\|) \|y^{i+1} - \hat{y}\|^2 - \frac{\eta_i \epsilon \|A - V\|}{2} \|x^{i+1} - x^i\|^2.$$

Proof. We observe

$$\begin{aligned} -\frac{1}{2} \|u^{i+1} - \hat{u}\|_{Z_{i+1}\Xi_{i+1}}^2 &= (\eta_{i+1} - \eta_i) \langle V(x^{i+1} - \hat{x}), y^{i+1} - \hat{y} \rangle \\ &\quad - \eta_i \mu_G \|x^{i+1} - \hat{x}\|^2 - \eta_{i+1} \mu_{F^*} \|y^{i+1} - \hat{y}\|^2, \end{aligned}$$

which gives, by definition of \tilde{H}_{i+1} in (10) and H in (8),

$$\begin{aligned} D &= \langle \tilde{H}_{i+1}(u^{i+1}), u^{i+1} - \hat{u} \rangle_{Z_{i+1}W_{i+1}} - \frac{1}{2} \|u^{i+1} - \hat{u}\|_{Z_{i+1}\Xi_{i+1}}^2 \\ &= \langle H(u^{i+1}), u^{i+1} - \hat{u} \rangle_{Z_{i+1}W_{i+1}} \\ &\quad + \eta_{i+1} \langle (A - V)(x^{i+1} - \bar{x}^{i+1}), y^{i+1} - \hat{y} \rangle \\ &\quad + (\eta_{i+1} - \eta_i) \langle V(x^{i+1} - \hat{x}), y^{i+1} - \hat{y} \rangle \\ &\quad - \eta_i \mu_G \|x^{i+1} - \hat{x}\|^2 - \eta_{i+1} \mu_{F^*} \|y^{i+1} - \hat{y}\|^2. \end{aligned} \quad (22)$$

Now we estimate the first term on the right hand side: Since $\hat{u} \in H^{-1}(0)$ with $\hat{u} = \begin{pmatrix} \hat{x} \\ \hat{y} \end{pmatrix}$, we have

$$-V^* \hat{y} \in \partial G(\hat{x}) \quad \text{and} \quad A \hat{x} \in \partial F^*(\hat{y})$$

This yields to

$$\begin{aligned} \langle H(u^{i+1}), u^{i+1} - \hat{u} \rangle_{Z_{i+1}W_{i+1}} &= \langle H(u^{i+1}) - H(\hat{u}), u^{i+1} - \hat{u} \rangle_{Z_{i+1}W_{i+1}} \\ &= \eta_i \langle \partial G(x^{i+1}) - \partial G(\hat{x}), x^{i+1} - \hat{x} \rangle \\ &\quad + \eta_{i+1} \langle \partial F^*(y^{i+1}) - \partial F^*(\hat{y}), y^{i+1} - \hat{y} \rangle \\ &\quad + \eta_i \langle V^*(y^{i+1} - \hat{y}), x^{i+1} - \hat{x} \rangle \\ &\quad + \eta_{i+1} \langle A(\hat{x} - x^{i+1}), y^{i+1} - \hat{y} \rangle. \end{aligned}$$

Now the strong convexity of G and F^* with constants γ_G and γ_{F^*} , respectively, results in the inequality

$$\begin{aligned} \langle H(u^{i+1}), u^{i+1} - \hat{u} \rangle_{Z_{i+1}W_{i+1}} &\geq \eta_i \gamma_G \|x^{i+1} - \hat{x}\|^2 + \eta_{i+1} \gamma_{F^*} \|y^{i+1} - \hat{y}\|^2 \\ &\quad + \eta_i \langle V(x^{i+1} - \hat{x}), y^{i+1} - \hat{y} \rangle \\ &\quad + \eta_{i+1} \langle A(\hat{x} - x^{i+1}), y^{i+1} - \hat{y} \rangle. \end{aligned}$$

Plugging this into the definition of D in (22), collecting terms gives

$$\begin{aligned} D &\geq \eta_i (\gamma_G - \mu_G) \|x^{i+1} - \hat{x}\|^2 + \eta_{i+1} (\gamma_{F^*} - \mu_{F^*}) \|y^{i+1} - \hat{y}\|^2 \\ &\quad + \eta_i \langle V(x^{i+1} - \hat{x}), y^{i+1} - \hat{y} \rangle + \eta_{i+1} \langle A(\hat{x} - x^{i+1}), y^{i+1} - \hat{y} \rangle \\ &\quad + \eta_{i+1} \langle V(\bar{x}^{i+1} - x^{i+1}), y^{i+1} - \hat{y} \rangle + (\eta_{i+1} - \eta_i) \langle V(x^{i+1} - \hat{x}), y^{i+1} - \hat{y} \rangle \\ &= \eta_i (\gamma_G - \mu_G) \|x^{i+1} - \hat{x}\|^2 + \eta_{i+1} (\gamma_{F^*} - \mu_{F^*}) \|y^{i+1} - \hat{y}\|^2 \\ &\quad + \eta_{i+1} \langle (A - V)(\hat{x} - x^{i+1}), y^{i+1} - \hat{y} \rangle + \eta_{i+1} \langle (A - V)(x^{i+1} - \bar{x}^{i+1}), y^{i+1} - \hat{y} \rangle. \end{aligned}$$

We use the extrapolation $\bar{x}^{i+1} - x^{i+1} = \omega_i(x^{i+1} - x^i)$ to get

$$\begin{aligned} \langle (A - V)(x^{i+1} - \bar{x}^{i+1}), y^{i+1} - \hat{y} \rangle &= \omega_i \langle (A - V)(x^i - x^{i+1}), y^{i+1} - \hat{y} \rangle \\ &\geq -\omega_i \|A - V\| \|x^i - x^{i+1}\| \|y^{i+1} - \hat{y}\| \\ &\geq -\frac{\omega_i \|A - V\|}{2} \left(\epsilon \|x^i - x^{i+1}\|^2 + \frac{1}{\epsilon} \|y^{i+1} - \hat{y}\|^2 \right) \end{aligned}$$

by Young's inequality and similarly we derive

$$\begin{aligned} \langle (A - V)(\hat{x} - x^{i+1}), y^{i+1} - \hat{y} \rangle &\geq -\|A - V\| \|\hat{x} - x^{i+1}\| \|y^{i+1} - \hat{y}\| \\ &\geq -\frac{\|A - V\|}{2} \left(\epsilon \|\hat{x} - x^{i+1}\|^2 + \frac{1}{\epsilon} \|y^{i+1} - \hat{y}\|^2 \right). \end{aligned}$$

With these two estimate we continue to lower bound D and arrive at

$$\begin{aligned} D &\geq \eta_i (\gamma_G - \mu_G) \|x^{i+1} - \hat{x}\|^2 + \eta_{i+1} (\gamma_{F^*} - \mu_{F^*}) \|y^{i+1} - \hat{y}\|^2 \\ &\quad - \eta_{i+1} \epsilon \frac{\|A - V\|}{2} \|x^{i+1} - \hat{x}\|^2 - \eta_{i+1} \frac{\|A - V\|}{2\epsilon} \|y^{i+1} - \hat{y}\|^2 \\ &\quad - \eta_{i+1} \epsilon \frac{\omega_i \|A - V\|}{2} \|x^i - x^{i+1}\|^2 - \eta_{i+1} \frac{\omega_i \|A - V\|}{2\epsilon} \|y^{i+1} - \hat{y}\|^2 \\ &= \eta_{i+1} \left(\gamma_{F^*} - \mu_{F^*} - \frac{(1 + \omega_i) \|A - V\|}{2\epsilon} \right) \|y^{i+1} - \hat{y}\|^2 \\ &\quad + \left[\eta_i (\gamma_G - \mu_G) - \eta_{i+1} \epsilon \frac{\|A - V\|}{2} \right] \|x^{i+1} - \hat{x}\|^2 \\ &\quad - \eta_{i+1} \epsilon \frac{\omega_i \|A - V\|}{2} \|x^{i+1} - x^i\|^2 \\ &\geq \left[\eta_{i+1} (\gamma_{F^*} - \mu_{F^*} - \frac{1 + \omega_i}{2\epsilon} \|A - V\|) \right] \|y^{i+1} - \hat{y}\|^2 - \frac{\eta_i \epsilon \|A - V\|}{2} \|x^{i+1} - x^i\|^2. \end{aligned}$$

□

Additionally, we want to draw attention to the connection towards our estimate of $Z_{i+1}M_{i+1}$ in (17). Hence, we estimate $-\Delta_{i+1}$ in the following way.

Lemma 2.5. *Let the assumptions of Theorem 2.4 be fulfilled and assume furthermore that (18), (21) and*

$$\varphi_i \geq \frac{\epsilon \eta_i}{\delta} \|A - V\| \quad (23)$$

hold with $0 < \delta \leq \kappa < 1$. Define

$$S_{i+1} = \begin{pmatrix} (\delta \varphi_i - \epsilon \eta_i \|A - V\|)I & 0 \\ 0 & \psi_{i+1}I - \frac{\eta_i^2}{\varphi_i(1-\kappa)} VV^* \end{pmatrix}, \quad (24)$$

and assume that

$$\frac{1}{2} \|u^{i+1} - u^i\|_{S_{i+1}}^2 + \eta_{i+1} \left(\gamma_{F^*} - \mu_{F^*} - \frac{1 + \omega_i}{2\epsilon} \|A - V\| \right) \|y^{i+1} - \hat{y}\|^2 \geq -\Delta_{i+1},$$

is fulfilled. Then it holds that

$$-\Delta_{i+1} \leq \frac{1}{2} \|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 + D.$$

Proof. In the first step, we rewrite

$$\begin{aligned} S_{i+1} &= \begin{pmatrix} (\delta\varphi_i - \epsilon\eta_i\|A - V\|)I & 0 \\ 0 & \psi_{i+1}I - \frac{\eta_i^2}{\varphi_i(1-\kappa)}VV^* \end{pmatrix} \\ &= \underbrace{\begin{pmatrix} \delta\varphi_i I & 0 \\ 0 & \psi_{i+1}I - \frac{\eta_i^2}{\varphi_i(1-\kappa)}VV^* \end{pmatrix}}_{\leq Z_{i+1}M_{i+1}} - \begin{pmatrix} \epsilon\eta_i\|A - V\|I & 0 \\ 0 & 0 \end{pmatrix}, \end{aligned}$$

so Lemma 2.2 gives

$$\frac{1}{2}\|u^{i+1} - u^i\|_{S_{i+1}}^2 \leq \frac{1}{2}\|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 - \frac{\eta_i\epsilon\|A - V\|}{2}\|x^{i+1} - x^i\|^2.$$

Hence,

$$\begin{aligned} -\Delta_{i+1} &\leq \frac{1}{2}\|u^{i+1} - u^i\|_{S_{i+1}}^2 + \eta_{i+1}(\gamma_{F^*} - \mu_{F^*} - \frac{1 + \omega_i}{2\epsilon}\|A - V\|)\|y^{i+1} - \hat{y}\|^2 \\ &\leq \frac{1}{2}\|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 - \frac{\eta_i\epsilon\|A - V\|}{2}\|x^{i+1} - x^i\|^2 \\ &\quad + \eta_{i+1}(\gamma_{F^*} - \mu_{F^*} - \frac{1 + \omega_i}{2\epsilon}\|A - V\|)\|y^{i+1} - \hat{y}\|^2 \end{aligned}$$

which shows the claim since Theorem 2.4 shows that the last two terms are a lower bound for D . \square

To obtain a generally applicable estimate for the iteration (4) in terms of (5), we further recognize

$$\begin{aligned} -\Delta_{i+1} &\leq \frac{1}{2}\|u^{i+1} - u^i\|_{S_{i+1}}^2 + \eta_{i+1} \left(\gamma_{F^*} - \mu_{F^*} - \frac{1 + \omega_i}{2}\|A - V\| \right) \|y^{i+1} - \hat{y}\|^2 \\ &\leq \frac{1}{2}\|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 \\ &\quad + \langle \tilde{H}_{i+1}(u^{i+1}), u^{i+1} - \hat{u} \rangle_{Z_{i+1}M_{i+1}} - \frac{1}{2}\|u^{i+1} - \hat{u}\|_{Z_{i+1}\Xi_{i+1}}^2, \end{aligned}$$

which yields to

$$\begin{aligned} -\Delta_{i+1} &\leq \frac{1}{2}\|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 + \langle \tilde{H}_{i+1}(u^{i+1}), u^{i+1} - \hat{u} \rangle_{Z_{i+1}M_{i+1}} \\ &\quad - \frac{1}{2}\|u^{i+1} - \hat{u}\|_{Z_{i+2}M_{i+2} - Z_{i+1}M_{i+1}}^2 \end{aligned}$$

by the equality

$$\|u^{i+1} - \hat{u}\|_{Z_{i+1}\Xi_{i+1}}^2 = \|u^{i+1} - \hat{u}\|_{Z_{i+2}M_{i+2} - Z_{i+1}M_{i+1}}^2.$$

Since rearranging these terms yields to

$$\begin{aligned} \langle \tilde{H}_{i+1}(u^{i+1}), u^{i+1} - \hat{u} \rangle_{Z_{i+1}M_{i+1}} &\geq \frac{1}{2}\|u^{i+1} - \hat{u}\|_{Z_{i+2}M_{i+2} - Z_{i+1}M_{i+1}}^2 \\ &\quad - \frac{1}{2}\|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 - \Delta_{i+1} \end{aligned}$$

and thus all conditions of Theorem 2.1 are satisfied, we finally proved:

Theorem 2.6. *Suppose that the step length conditions (15), (16), (18)–(20) and (23) hold with $\epsilon > 0$ and $0 < \delta \leq \kappa < 1$. Additionally suppose that G, F^* are γ_G/γ_{F^*} -strongly convex, respectively, fulfill (21):*

$$\gamma_G \geq \frac{\epsilon}{2\omega_i} \|A - V\| + \mu_G.$$

Furthermore, let S_{i+1} be defined by (24) and $\hat{u} = \begin{pmatrix} \hat{x} \\ \hat{y} \end{pmatrix}$ fulfill $0 \in \tilde{H}(\hat{u})$. Then it holds

$$\frac{1}{2} \|u^N - \hat{u}\|_{Z_{N+1}M_{N+1}}^2 \leq \frac{1}{2} \|u^0 - \hat{u}\|_{Z_1M_1}^2 + \sum_{i=0}^{N-1} \Delta_{i+1}$$

for every Δ_i which fulfills

$$\frac{1}{2} \|u^{i+1} - u^i\|_{S_{i+1}}^2 + \eta_{i+1} (\gamma_{F^*} - \mu_{F^*} - \frac{1 + \omega_i}{2\epsilon} \|A - V\|) \|y^{i+1} - \hat{y}\|^2 \geq -\Delta_{i+1}.$$

2.4 Convergence rates

With the results from the previous section, we are in position to ensure convergence of (4) under easily verifiable conditions.

In this section we want to show that if both G as well as F^* are proper, strongly convex, lower-semicontinuous functions, then with a wise choice of the step sizes we get linear convergence. Even though this is a result, we can expect from a first-order method, it may come as a surprise, since there is an error in the adjoint. We proceed as follows: First we show the conditions under which we can expect linear convergence, then we select the parameters accordingly in order to obtain a choice of the step sizes that is easier for practice. Hence, we begin with the proof of linear convergence for strongly convex functions G and F^* .

Theorem 2.7. *Choose $\mu_G > 0, \mu_{F^*} > 0$ and suppose that G is γ_G -strongly convex and F^* is γ_{F^*} -strongly convex with*

$$\gamma_G \geq \frac{\epsilon}{2\omega_i} \|A - V\| + \mu_G, \quad \gamma_{F^*} \geq \frac{1 + \omega_i}{2\epsilon} \|A - V\| + \mu_{F^*} \quad (25)$$

for some $\epsilon > 0$. Furthermore suppose that (5) is solvable and let $\hat{u} \in \mathcal{U}$ be a stationary point of the iteration. Then the Chambolle-Pock method with mismatched adjoint given by (4) with constant step lengths

$$\begin{aligned} \tau_i = \tau &:= \min \left\{ \frac{\epsilon^{-1}\delta}{\|A - V\|}, \sqrt{\frac{(1 - \kappa)\mu_{F^*}}{\|V\|^2\mu_G}} \right\}, \\ \sigma_i = \sigma &:= \frac{\mu_G}{\mu_{F^*}} \tau, \\ \omega_i = \omega &:= (1 + 2\tau\mu_G)^{-1} \end{aligned}$$

for some $0 \leq \delta \leq \kappa < 1$ and all $i \in \mathbb{N}$ converges with $\|u^N - \hat{u}\|^2 = \mathcal{O}(\omega^N)$.

Proof. We take $\varphi_0 := \frac{1}{\tau}$ and $\psi_0 := \frac{1}{\sigma}$ and by (19) we get

$$\varphi_i \tau = \underbrace{\varphi_0 \tau}_{=1} (1 + 2\tau\mu_G)^i = (1 + 2\tau\mu_G)^i$$

By (15) and (20) we get

$$\begin{aligned} \psi_{i+1} \sigma &= \psi_i \sigma (1 + 2\sigma\mu_{F^*}) = \psi_i \sigma (1 + 2\tau\mu_G) \\ &= \underbrace{\psi_0 \sigma}_{=1} (1 + 2\tau\mu_G)^{i+1} = (1 + 2\tau\mu_G)^{i+1}. \end{aligned}$$

Hence, again by (15), we get

$$\eta_i = (1 + 2\tau\mu_G)^i$$

and, by (16)

$$\omega = \frac{\eta_i}{\eta_{i+1}} = \frac{1}{1 + 2\tau\mu_G}.$$

Now we claim, that the matrix S_{i+1} defined in (24) is positive semi-definite, which is equivalent to the conditions $\delta\varphi_i - \eta_i\epsilon\|A - V\| \geq 0$ and $\psi_{i+1} \geq \eta_i^2\|V\|^2/(\varphi_i(1 - \kappa))$. The first condition is fulfilled if $\tau \leq \delta/(\epsilon\|A - V\|)$ and the second condition is fulfilled if $\tau^2 \leq (1 - \kappa)\mu_{F^*}/(\|V\|^2\mu_G)$ which are fulfilled by assumption. Consequently, we can choose $\Delta_{i+1} = 0$ for $i \in \mathbb{N}$ and as a result, we get from Theorem 2.6 and Lemma 2.5

$$\begin{aligned} \|u^0 - \hat{u}\|_{Z_1 M_1}^2 &\geq \|u^N - \hat{u}\|_{Z_{N+1} M_{N+1}}^2 \\ &\geq \delta\varphi_N \|x^N - \hat{x}\|^2 + \left(\psi_{N+1} - \frac{\eta_N^2\|V\|^2}{\varphi_N(1 - \kappa)} \right) \|y^N - \hat{y}\|^2 \\ &= \varphi_N \left(\delta \|x^N - \hat{x}\|^2 + \left(\frac{\psi_{N+1}}{\varphi_N} - \frac{\eta_N^2\|V\|^2}{\varphi_N^2(1 - \kappa)} \right) \|y^N - \hat{y}\|^2 \right) \end{aligned}$$

Using the properties of φ_N, ψ_N and η_N we arrive at

$$\|u^0 - \hat{u}\|_{Z_1 M_1}^2 \geq (1 + 2\tau\mu_G)^N \left(\frac{\delta}{\tau} \|x^N - \hat{x}\|^2 + \left(\frac{\tau}{\sigma}(1 + 2\tau\mu_G) - \tau \right) \|y^N - \hat{y}\|^2 \right)$$

and the desired convergence rate. \square

Nevertheless, this theorem is not immediately practical, since it is unclear, if all parameters can be chosen such that all conditions are fulfilled. Hence, we derive a method that allows for a feasible choice of the parameters.

If we plug the definition of ω into conditions (25), we get

$$\gamma_G \geq \frac{\epsilon}{2}(1 + 2\tau\mu_G)\|A - V\| + \mu_G \quad (26)$$

$$\gamma_{F^*} \geq \frac{1 + \tau\mu_G}{\epsilon(1 + 2\tau\mu_G)}\|A - V\| + \mu_{F^*}. \quad (27)$$

Since $\frac{1}{2} \leq \frac{1+t}{1+2t} \leq 1$ for $t > 0$, we can fulfill (27) if we have

$$\gamma_{F^*} = \frac{\|A - V\|}{\epsilon} + \mu_{F^*}. \quad (28)$$

We express the quantities μ_G and μ_{F^*} by $\mu_{F^*} = a\gamma_{F^*}$, $\mu_G = b\gamma_G$ with $0 \leq a, b \leq 1$. Then, we can use (28) to get a valid value for ϵ , namely

$$\epsilon = \frac{\|A - V\|}{(1-a)\gamma_{F^*}}.$$

Furthermore, we observe that it is always beneficial to choose δ as large as possible, i.e. we set $\delta = \kappa$. Additionally, using this ϵ and μ_G in (26), we get

$$\gamma_G \geq \frac{1+2\tau b\gamma_G}{2(1-a)\gamma_{F^*}} \|A - V\|^2 + b\gamma_G,$$

which yields to the inequality

$$\tau \leq \frac{(1-b)(1-a)\gamma_{F^*}}{\|A-V\|^2 b} - \frac{1}{2b\gamma_G}. \quad (29)$$

On the other hand, we plug in $\delta = \kappa$, and the values for ϵ , μ_{F^*} and μ_G into the definition of τ in Theorem 2.7 and get

$$\tau = \min \left\{ \frac{\kappa(1-a)\gamma_{F^*}}{\|A-V\|^2}, \sqrt{\frac{(1-\kappa)a\gamma_{F^*}}{\|V\|^2 b\gamma_G}} \right\}.$$

and decide that

$$\frac{\kappa^2(1-a)^2\gamma_{F^*}^2}{\|A-V\|^4} \geq \frac{(1-\kappa)a\gamma_{F^*}}{\|V\|^2 b\gamma_G} \quad (30)$$

should hold in the definition. Hence, we have to find a, b and κ , such that

$$\frac{\kappa(1-a)\gamma_{F^*}}{\|A-V\|^2} \leq \frac{(1-b)(1-a)\gamma_{F^*}}{\|A-V\|^2 b} - \frac{1}{2b\gamma_G},$$

which is equivalent to

$$\kappa \leq \frac{1-b}{b} - \frac{\|A-V\|^2}{2b(1-a)\gamma_{F^*}\gamma_G} = \frac{1}{b} \left(1 - b - \frac{\|A-V\|^2}{2(1-a)\gamma_{F^*}\gamma_G} \right). \quad (31)$$

Clearly, the upper bound increases with decreasing the value of b . Hence, if we restrict ourselves to $b \leq \frac{1}{2}$ and use our degrees of freedom to set $a = \frac{1}{2}$. Now (30) turns into

$$\frac{\kappa^2\gamma_{F^*}^2}{4\|A-V\|^4} \geq \frac{1}{b} \frac{1-\kappa}{2\|V\|^2\gamma_G}, \quad (32)$$

which is equivalent to the condition

$$b \leq \frac{\kappa}{1-\kappa} \frac{\|A-V\|^4}{\|V\|^2} \frac{2}{\gamma_{F^*}^2\gamma_G} \quad (33)$$

on b . From (31) we have

$$\kappa \leq \frac{1}{b} \left(\frac{1}{2} - \frac{\|A-V\|^2}{\gamma_{F^*}\gamma_G} \right). \quad (34)$$

The later is positive, whenever

$$2\|A-V\|^2 < \gamma_{F^*}\gamma_G \quad (35)$$

is fulfilled. So by (34) we are able to find a small enough b for every $\kappa \in (0, 1)$, such that the corresponding inequality holds. In conclusion, we have proven:

Theorem 2.8. *Suppose that G, F^* are γ_G/γ_{F^*} -strongly convex functions fulfilling*

$$\gamma_{F^*}\gamma_G > 2\|A-V\|^2.$$

Define

$$b = \min \left\{ \frac{1}{2}, \frac{1}{\kappa} \left(\frac{1}{2} - \frac{\|A-V\|^2}{\gamma_{F^*}\gamma_G} \right), \frac{1-\kappa}{\kappa} \frac{\|A-V\|^4}{\|V\|^2} \frac{2}{\gamma_{F^*}\gamma_G} \right\}$$

for some arbitrary $\kappa \in (0, 1)$. Furthermore suppose that (5) is solvable and let $\hat{u} \in \mathcal{U}$ be a stationary point of the iteration. Then the Chambolle-Pock method with mismatched adjoint and constant step lengths

$$\begin{aligned} \tau_i = \tau &:= \sqrt{\frac{(1-\kappa)\gamma_{F^*}}{2b\|V\|^2\gamma_G}}, \\ \sigma_i = \sigma &:= 2b \frac{\gamma_G}{\gamma_{F^*}} \tau, \\ \omega_i = \omega &:= (1 + 2b\tau\gamma_G)^{-1} \end{aligned}$$

converges with $\|u^N - \hat{u}\|^2$ decaying to zero at the rate $\mathcal{O}(\omega^N)$.

Nevertheless, many problems involve only one strongly convex function while the other only remains to be convex. But even in this case, we are able to obtain strong convergence with a rate.

Theorem 2.9. *Suppose that G is a γ_G -strongly convex and F^* is convex function with $\gamma_G > 0$. Choose the initial step lengths τ_0, σ_0 as well as $\delta, \kappa, \epsilon > 0$ and $\mu_G > 0$, such that*

$$\gamma_G \geq \frac{\epsilon\sqrt{1+2\tau_0\mu_G}}{2} \|A-V\| + \mu_G \quad (36)$$

holds, $\delta \leq \kappa < 1$ is fulfilled and the initial step length bounds

$$\tau_0 \leq \frac{\delta}{\epsilon\|A-V\|} \text{ and } \sigma_0\tau_0 \leq \frac{1-\kappa}{\|V\|^2} \quad (37)$$

are satisfied. Furthermore suppose that (5) is solvable and let $\hat{u} \in \mathcal{U}$ be a stationary point of the iteration. Then the Chambolle-Pock method with mismatched adjoint and step lengths

$$\begin{aligned} \omega_i &:= \sqrt{1+2\tau_i\mu_G}^{-1} \\ \tau_{i+1} &:= \tau_i\omega_i \\ \sigma_{i+1} &:= \frac{\sigma_i}{\omega_i} \end{aligned}$$

converges with $\|x^N - \hat{x}\|^2$ decaying to zero at the rate $\mathcal{O}(N^{-2})$.

Proof. Since we want to make use of (2.6) and the only information on γ_{F^*} is its non-negativity, we have to choose $\mu_{F^*} = 0$. Hence $\psi_{i+1} = \psi_i(1 + 2\sigma_i\mu_{F^*}) = \psi$ in (20) remains a constant to choose. Furthermore we have $\omega_i \leq 1$, which yields to decreasing $\tau_{i+1} = \tau_i\omega_i$ and to $\gamma_G \leq \frac{\epsilon\sqrt{1+2\tau_i\mu_G}}{2} \|A-V\| + \mu_G$ for all $i \in \mathbb{N}$. Additionally it leads to

$$\varphi_i \geq \frac{\epsilon\eta_i}{\delta} \|A-V\|.$$

Likewise

$$\psi \geq \frac{\eta_i^2}{\varphi_i(1-\kappa)} \|V\|^2$$

holds by $\sigma_0\tau_0 \leq \frac{1-\kappa}{\|V\|^2}$. Still remaining is the update rule $\varphi_{i+1} = \varphi_i(1 + 2\tau_i\mu_G)$, which deviates from the definition of $\eta_i = \psi\sigma_i = \varphi_i\tau_i$, which also provides $\omega_i = \frac{\eta_i}{\eta_{i+1}}$. Thus, the conditions in Theorem 2.6 are attained and it is

$$\begin{aligned} -\Delta_{i+1} &\leq \frac{1}{2} \|u^{i+1} - u^i\|_{S_{i+1}}^2 + \eta_{i+1} \left(\gamma_{F^*} - \mu_{F^*} - \frac{1 + \omega_i}{2\epsilon} \|A - V\| \right) \|y^{i+1} - y^i\|^2 \\ &= \frac{1}{2} (\delta\varphi_i - \epsilon\eta_i \|A - V\|) \|x^{i+1} - x^i\|^2 \\ &\quad + \frac{1}{2} \left(\psi - \frac{\eta_i^2}{\varphi_i(1 - \kappa)} \|V\|^2 - \frac{1 + \omega_i}{2\epsilon} \|A - V\| \right) \|y^{i+1} - y^i\|^2. \end{aligned}$$

Choosing

$$\psi \geq \frac{\|A - V\|}{\epsilon \left(1 - \frac{\sigma_0\tau_0}{1-\kappa} \|V\|^2 \right)},$$

both coefficients are non-negative, which allows the choice of $\Delta_{i+1} = 0$. Consequently, we have

$$\frac{1}{2} \delta\varphi_N \|x^N - \hat{x}\|^2 \leq \frac{1}{2} \|u^N - \hat{u}\|_{Z_{N+1}M_{N+1}}^2 \leq \frac{1}{2} \|u^0 - \hat{u}\|_{Z_1M_1}^2.$$

Citing [4], we have $\tau_N \in \mathcal{O}(N^{-1})$ and consequently it is

$$\begin{aligned} \|x^N - \hat{x}\|^2 &\leq \frac{1}{\varphi_N} \frac{1}{\delta} \|u_0 - \hat{u}\|_{Z_1M_1}^2 \\ &= \frac{\tau_N^2}{\psi\sigma_0\tau_0} \frac{1}{\delta} \|u_0 - \hat{u}\|_{Z_1M_1}^2 \in \mathcal{O}(N^{-2}). \end{aligned}$$

□

In the situation of Theorem 2.9, the choice of parameters is more straightforward: As soon as you choose

$$\sigma_0\tau_0 < \|V\|^{-2}$$

it is clear that $0 \leq \delta \leq \kappa < 1$, $\epsilon > 0$ and μ_G exist, such that all demands in Theorem 2.9 are met. Therefore, we are able to recover the result of Chambolle and Pock from [4], however, it might come surprising that only the norm of the “backward operator” V appears in the conditions and that the “forward operator” only enters through the difference $\|A - V\|$.

Corollary 2.10. *Suppose that G is a γ_G -strongly convex and F^* is a convex function with $\gamma_G > 0$. Choose the initial step lengths $\tau_0, \sigma_0 > 0$ as well as $\mu_G \in (0, \gamma_G)$, such that the initial step length bound*

$$\sigma_0\tau_0 < \frac{1}{\|V\|^2}$$

is satisfied. Furthermore suppose that (5) is solvable and let $\hat{u} \in \mathcal{U}$ be a stationary point of the iteration. Then the Chambolle-Pock method with mismatched adjoint and step lengths

$$\begin{aligned} \omega_i &:= \frac{1}{\sqrt{1+2\tau_i\mu_G}} \\ \tau_{i+1} &:= \tau_i\omega_i \\ \sigma_{i+1} &:= \frac{\sigma_i}{\omega_i} \end{aligned}$$

converges with $\|x^N - \hat{x}\|^2$ decaying to zero at the rate $\mathcal{O}(N^{-2})$.

3 Numerical examples

In this section we report some numerical experiments to illustrate the results.

3.1 Convex quadratic problems

As examples where all quantities and solutions can be computed exactly, we consider convex quadratic problems of the form

$$\min_{x \in \mathbb{R}^n} \frac{\alpha}{2} \|x\|_2^2 + \frac{1}{2\beta} \|Ax - b\|_2^2 \quad (38)$$

with $\alpha, \beta > 0$, $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$. With $G(x) = \frac{\alpha}{2} \|x\|_2^2$ and $F(\zeta) = \frac{1}{2\beta} \|\zeta - b\|_2^2$ this is of the form (1). The conjugate functions are $G^*(\xi) = \frac{1}{2\alpha} \|x\|_2^2$ and $F^*(y) = \frac{\beta}{2} \|y\|_2^2 + \langle y, b \rangle$ and the respective proximal operators are readily computed as

$$\begin{aligned} \text{prox}_{\tau G}(x) &= \frac{x}{1+\tau\alpha} \\ \text{prox}_{\sigma F^*}(y) &= \frac{y - \sigma b}{1+\sigma\beta} \end{aligned}$$

and the optimal primal solution is

$$x^* = \left(\alpha I + \frac{1}{\beta} A^T A \right)^{-1} \left(\frac{1}{\beta} A^T b \right).$$

Note that G is strongly convex with constant $\gamma_G = \alpha$ and F^* is strongly convex with constant $\gamma_{F^*} = \beta$ and hence, for $\alpha, \beta > 0$ we can use Theorem 2.8 to obtain valid stepsizes. For a numerical experiment we choose $n = 400$, $m = 200$, a random matrix $A \in \mathbb{R}^{m \times n}$ and a perturbation $V \in \mathbb{R}^{m \times n}$ by adding a small random matrix to A , i.e.

$$V = A + E \quad \text{with} \quad \|E\| \leq \eta.$$

The resulting algorithms is

$$\begin{aligned} x^{i+1} &= \frac{1}{1+\tau\alpha} (x^i - \tau V^T y^i) \\ y^{i+1} &= \frac{1}{1+\sigma\beta} \left(y^i + \sigma A(x^{i+1} + \omega(x^{i+1} - x^i)) - \sigma b \right). \end{aligned}$$

We check the condition $\gamma_G \gamma_{F^*} > 2\|A - V\|^2$ numerically and use Theorem 2.8 to obtain feasible step-sizes. If we consider this iteration with constants τ , σ and ω , we get as limit the unique fixed points

$$\begin{aligned} \hat{x} &= \left(\alpha I + \frac{1}{\beta} V^T A \right)^{-1} \left(\frac{1}{\beta} V^T b \right) = V^T (\alpha \beta I + AV^T)^{-1} b \\ \hat{y} &= -(\beta I + \frac{1}{\alpha} AV^T)^{-1} b = -\alpha (\alpha \beta I + AV^T)^{-1} b. \end{aligned}$$

The true primal solution is

$$x^* = \left(\alpha I + \frac{1}{\beta} A^T A \right)^{-1} \left(\frac{1}{\beta} A^T b \right) = A^T (\alpha \beta I + AA^T)^{-1} b$$

For our experiment we used $\alpha = \gamma_G = 0.15$ and $\beta = \gamma_{F^*} = 1$ and $\kappa = 0.01$ in Theorem 2.8. Figure 1 illustrates that the method with mismatched adjoint behaves as expected: We observe linear convergence towards the fixed point \hat{x} and the iterates reach the error to the true minimizer x^* that has been predicted by Theorem 1.2.




Figure 1: Convergence of the Chambolle-Pock algorithms (3) and (4).

In this example, we observe the typical behavior of the method. Since our algorithm is structurally the same as the original, the convergence is sustained by smaller step sizes. Hence, we notice slower convergence by a bit. Additionally, we want to draw attention to the tight upper bounds on the convergence rate and the final error in this example. Furthermore, it becomes obvious that the distortion of the small adjoint disturbs the classic Chambolle-Pock method. However, both converge to a fixed point in a similar radius around the true solution.

3.2 Computerized tomography

To illustrate a real-world application of our results, we observe the effect of them on computerized tomography (CT) [3]. In computerized tomography one aims to reconstruct a slice of an object but x-ray measurements taken in different directions. The data of the x-ray measurements are stored in the so-called sinogram and the map of the image of the slice to the sinogram is modeled by a linear map which is referred to as Radon transform or forward projection. The adjoint of the map is called backprojection. There exist various inversion formulas which express the inverse of the Radon transform explicitly but since the Radon transform is compact (when modeled as a map between the right function spaces [10]), any inversion formula has to be unstable. One popular stable, approximate inversion method is the so-called filtered backprojection (FBP) [3]. The method gives good approximate reconstruction when the number of projections is high. However, the quality of the reconstruction quickly gets worse when the number of projection decreases. There are numerous efforts

to increase reconstruction quality with only a few projection, as this lowers the x-ray dose for a CT scan, and a successful approach uses total variation (TV) regularization as a reconstruction method [14] and solves the respective minimization problem with the Chambolle-Pock method. Usually, the method takes a large number of iteration. Moreover, there are many ways to implement the forward and the backward projection. In applications it sometime happen that a pair of forward and backward projection are chose that are not adjoint to each other, either because this importance of adjointness is not noted, or on purpose (speed of computation, special choice of backprojector to achieve a certain reconstruction quality, see also [16, 11, 5, 9]). In this example we illustrate a slightly different effect: Choose a mismatched adjoint on purpose to lower the number of iteration until good reconstruction quality under TV regularization.

TV- L^2 : We exaggerate the mismatch and replace the adjoint by the filtered back projection, which is an approximation of both the adjoint and the inverse of the Radon transform. Hence, we aim to solve the TV- L^2 problem

$$\min_{x \in \mathcal{X}} \frac{1}{2} \|Rx - y\|^2 + \lambda \text{TV}(x),$$

where $\text{TV}(x) = \|\nabla x\|_1$. This can be expressed as the saddle point problem

$$\min_{x \in \mathbb{R}^{m \times n}} \max_{p \in \mathbb{R}^{s \times t}, q \in \mathbb{R}^{m \times n}} -\langle x, \text{div } p \rangle + \langle Rx - y, q \rangle - \delta_{\|\cdot\|_\infty \leq 1}(p) - \frac{1}{2\lambda} \|q\|^2.$$

In our experiment we want to recover some image \hat{x} with 112×112 pixels from just 20 equispaced tomographic measurements y and added 3% relative Gaussian noise. With $F(x_1, x_2) = \frac{1}{2} \|x_1 - g\|_2^2 + \lambda \|(|x_2|)\|_1$, $G(x) = 0$ and $A = \begin{pmatrix} R \\ \nabla \end{pmatrix}$ the saddle point formulation is exactly of the form (1). We use the algorithm from Corollary 2.10 where we used $V^* = (S^* \quad -\text{div})$ instead of A^* and S^* is the filtered backprojection. The iteration reads as

$$\begin{aligned} x^{i+1} &= x^i - \tau_i S^* p^i + \tau_i \text{div } q^i, \\ \omega_i &= \frac{1}{\sqrt{1+2\tau_i \mu_G}}, \\ \bar{x}^{i+1} &= x^{i+1} + \omega_i (x^{i+1} - x^i), \\ p^{i+1} &= (p^i + \sigma_i (R\bar{x}^{i+1} - g)) / (1 + \sigma_i), \\ q^{i+1} &= \lambda (q^i + \sigma_i \nabla \bar{x}^{i+1}) \odot \min(\lambda, |q^i + \sigma_i \nabla \bar{x}^{i+1}|), \\ \tau_{i+1} &= \tau_i \omega_i, \\ \sigma_{i+1} &= \frac{\sigma_i}{\omega_i}. \end{aligned}$$

where the update on q^{i+1} is considered element-wise. To achieve a good result, we take $\lambda = 0.15$ and set the initial step sizes as $\sigma_0 = \tau_0 = \frac{0.99}{\|\nabla\|}$. The implementation is done in Python 3.7 with the built-in forward operator as well as the backprojector from scikit-image. The operator norm of V is estimated by the norm of the adjoint operator S^* (which is computed numerically) plus 8 (cf. [1, Lemma 6.142]).

In Figure 2 we show the original image (the famous Shepp-Logan phantom), its noisy sinogram, the reconstruction by filtered backprojection and the results

of the Chambolle-Pock iteration for TV regularized reconstruction with the exact adjoint and with the filtered backprojection as adjoint. One notes that the use of the filtered backprojection leads to a good reconstruction and does so faster than the iteration with the exact adjoint. Especially, the reconstruction does not suffer from the strike artifacts in the FBP reconstruction (these are suppressed by TV regularization).




Figure 2: Reconstruction of the Shepp Logan phantom.

Figure 3 shows the distance of the iteration to the exact reconstruction (i.e. the true noise free Shepp-Logan phantom) and the objective value. Naturally, the Chambolle-Pock iteration does not drive this error to zero (for both exact and mismatched adjoint). There are at least three different reasons: There is some error in the sinogram, we only use few projections and there is TV regularization involved. However, we note that the iteration with mismatch gets closer to the original image. Moreover, it can be seen that, as expected, the use of a mismatched adjoint prevents the true minimization of the objective.





Figure 3: Left: Distance to exact reconstruction over iterations. Right: Decay of primal objective function value.

In conclusion of this experiment, we can observe some advantages of the proposed adjoint replacement. While being comparable in computational efficiency, the convergence behavior differs. Heuristically speaking, the filtered back projection as an approximation on the inverse Radon transform works as a warm start. Hence, we recognize that the shape of the Shepp-Logan phantom gets reconstructed early on. Furthermore, the smoothing effect of the ramp filter in the filtered backprojection allows larger stepsizes, which result in faster convergence. Additionally, we observe that after 200 iterations, our method suffers less from artifacts than both the reconstruction with exact adjoint and the filtered backprojection.

4 Conclusion

We have established step sizes, for which the Chambolle-Pock method converges, even if the adjoint is mismatched. Additionally, we presented results that showed, that not only strong convergence can be preserved under strong convexity assumptions, and also that the convergence rate is in a similar region. Furthermore, as a broad class of problems are in the scope of this paper, we established an upper bound on the distance between the original solution and the fixed point of iteration with mismatch. Thus, approximating the adjoint with a computationally more efficient algorithm can be done as long as the assumptions are respected. One of these assumptions is, that the iteration with mismatch still possesses a fixed point and more work is needed to understand when this is guaranteed. Furthermore, we discussed advantages and disadvantages of our analysis and presented a real-world example of the use of a willingly done adjoint mismatch.

Acknowledgement

This work has received funding from the European Union’s Framework Programme for Research and Innovation Horizon 2020 (2014-2020) under the Marie Skłodowska-Curie Grant Agreement No. 861137. 

References

- [1] K. Bredies and D. Lorenz. *Mathematical Image Processing*. Birkhäuser, Cham, 2018.
- [2] J. Buffiere, E. Maire, J. Adrien, J. Masse, and E. Boller. In situ experiments with X ray tomography: an attractive tool for experimental mechanics. *Experimental Mechanics*, 50:289–305, 2010.
- [3] T. M. Buzug. *Computed tomography*. Springer, Berlin, Heidelberg, 2008.
- [4] A. Chambolle and T. Pock. A first-order primal-dual algorithm for convex problems with applications to imaging. *Journal of Mathematical Imaging and Vision*, 40:120–145, 2011.
- [5] E. Chouzenoux, J.-C. Pesquet, C. Riddell, M. Savanier, and Y. Troussel. Convergence of proximal gradient algorithm in the presence of adjoint mismatch. *Inverse Problems*, 37(6):Paper No. 065009, 29, 2021.
- [6] C. Clason, S. Mazurenko, and T. Valkonen. Acceleration and global convergence of a first-order primal-dual method for nonconvex problems. *SIAM J. Optim.*, 29:933–963, 2019.
- [7] Y. Dong, P. Hansen, M. Hochstenbach, and N. A. B. Riis. Fixing nonconvergence of algebraic iterative reconstruction with an unmatched backprojector. *SIAM J. Sci. Comput.*, 41:A1822–A1839, 2019.
- [8] T. Elfving and P. Hansen. Unmatched projector/backprojector pairs: Perturbation and convergence analysis. *SIAM J. Sci. Comput.*, 40, 2018.
- [9] D. A. Lorenz, S. D. Rose, and F. Schöpfer. The randomized Kaczmarz method with mismatched adjoint. *BIT Numerical Mathematics*, 58:1079–1098, 2018.
- [10] F. Natterer. *The mathematics of computerized tomography*, volume 32 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2001. Reprint of the 1986 original.
- [11] W. J. Palenstijn, K. Batenburg, and J. Sijbers. Performance improvements for iterative electron tomography reconstruction using graphics processing units (GPUs). *Journal of structural biology*, 176 2:250–3, 2011.
- [12] C. Riddell, B. Bendriem, M. Bourguignon, and J. Kernévez. The approximate inverse and conjugate gradient: non-symmetrical algorithms for fast attenuation correction in SPECT. *Physics in medicine and biology*, 40 2:269–81, 1995.
- [13] M. Savanier, É. Chouzenoux, J. Pesquet, C. Riddell, and Y. Troussel. Proximal gradient algorithm in the presence of adjoint mismatch. *2020 28th European Signal Processing Conference (EUSIPCO)*, pages 2140–2144, 2021.
- [14] E. Y. Sidky, J. H. Jorgensen, and X. Pan. Convex optimization problem prototyping for image reconstruction in computed tomography with the Chambolle–Pock algorithm. *Physics in Medicine & Biology*, 57(10):3065, 2012.

- [15] T. Valkonen. Testing and non-linear preconditioning of the proximal point method. *Appl. Math. Optim.*, 82(2):591–636, 2020.
- [16] G. L. Zeng and G. T. Gullberg. Unmatched projector/backprojector pairs in an iterative reconstruction algorithm. *IEEE transactions on medical imaging*, 19(5):548–555, 2000.